


each R^2 is independently selected from hydroxyl, halogen, cyano, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, C_{1-4} alkoxy, hydroxy C_{1-4} alkyl, hydroxy C_{1-4} alkoxy, halo C_{1-4} alkyl, halo C_{1-4} alkoxy, hydroxyhalo C_{1-4} alkyl, hydroxyhalo C_{1-4} alkoxy, C_{1-4} alkoxy C_{1-4} alkyl, halo C_{1-4} alkoxy C_{1-4} alkyl, C_{1-4} alkoxy C_{1-4} alkyl wherein each C_{1-4} alkyl may optionally be substituted with one or two hydroxyl groups, hydroxyhalo C_{1-4} alkoxy C_{1-4} alkyl, R^{13} , C_{1-4} alkyl substituted with R^{13} , C_{1-4} alkyl substituted with $-C(=O)-R^{13}$, C_{1-4} alkoxy substituted with R^{13} , C_{1-4} alkoxy substituted with $-C(=O)-R^{13}$, $-C(=O)-R^{13}$, C_{1-4} alkyl substituted with $-NR^7R^8$, C_{1-4} alkyl substituted with $-C(=O)-NR^7R^8$, C_{1-4} alkoxy substituted with $-NR^7R^8$, C_{1-4} alkoxy substituted with $-C(=O)-NR^7R^8$, $-NR^7R^8$ and $-C(=O)-NR^7R^8$; or when two R^2 groups are attached to adjacent carbon atoms they may be taken together to form a radical of formula:



represents O or S;

Y represents $-CR^{18}=N-OR^{19}$ or $-E-D$;

D represents a 3 to 12 ring membered monocyclic or bicyclic carbocyclyl or a 3 to 12 ring membered monocyclic or bicyclic heterocyclyl containing at least one heteroatom selected from N, O or S, wherein said carbocyclyl and heterocyclyl may each be optionally substituted by one or more (e.g. 1, 2 or 3) R^1 groups;

E represents a bond;

R^1 represents hydrogen, halo, cyano, C_{1-6} alkyl, C_{1-6} alkoxy, $-C(=O)-O-C_{1-6}$ alkyl, C_{2-4} alkenyl, hydroxy C_{1-6} alkyl, halo C_{1-6} alkyl, hydroxyhalo C_{1-6} alkyl, cyano C_{1-4} alkyl, C_{1-6} alkoxy C_{1-6} alkyl wherein each C_{1-6} alkyl may optionally be substituted with one or two hydroxyl groups, $-NR^4R^5$, C_{1-6} alkyl substituted with $-O-C(=O)-C_{1-6}$ alkyl, C_{1-6} alkyl substituted with $-NR^4R^5$, $-C(=O)-NR^4R^5$, $-C(=O)-C_{1-6}$ alkyl- NR^4R^5 , C_{1-6} alkyl substituted with $-C(=O)-NR^4R^5$, $-S(=O)_2-C_{1-6}$ alkyl, $-S(=O)_2$ -halo C_{1-6} alkyl, $-S(=O)_2-NR^{14}R^{15}$, C_{1-6} alkyl substituted with $-S(=O)_2-C_{1-6}$ alkyl, C_{1-6} alkyl substituted with $-S(=O)_2$ -halo C_{1-6} alkyl, C_{1-6} alkyl substituted with $-S(=O)_2-NR^{14}R^{15}$, C_{1-6} alkyl substituted with $-NH-S(=O)_2-C_{1-6}$ alkyl, C_{1-6} alkyl substituted with $-NH-S(=O)_2$ -halo C_{1-6} alkyl, C_{1-6} alkyl substituted with $-NR^{12}-S(=O)_2-NR^{14}R^{15}$, R^6 , C_{1-6} alkyl substituted with R^6 , $-C(=O)-R^6$, C_{1-6} alkyl substituted with $-C(=O)-R^6$, hydroxy C_{1-6} alkyl substituted with R^6 , C_{1-6} alkyl

substituted with $-\text{Si}(\text{CH}_3)_3$, C_{1-6} alkyl substituted with $-\text{P}(=\text{O})(\text{OH})_2$ or C_{1-6} alkyl substituted with $-\text{P}(=\text{O})(\text{OC}_{1-6}\text{alkyl})_2$;

R^3 represents hydroxyl, C_{1-6} alkoxy, hydroxy C_{1-6} alkoxy, C_{1-6} alkoxy substituted with

- 5 $-\text{NR}^{10}\text{R}^{11}$, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, halo C_{1-6} alkyl optionally substituted with $-\text{O}-$
 $\text{C}(=\text{O})-\text{C}_{1-6}$ alkyl, hydroxy C_{1-6} alkyl optionally substituted with $-\text{O}-\text{C}(=\text{O})-\text{C}_{1-6}$ alkyl, hydroxy C_{2-6}
 alkenyl , hydroxy C_{2-6} alkynyl, hydroxyhalo C_{1-6} alkyl, cyano C_{1-6} alkyl, C_{1-6} alkyl substituted with
 carboxyl, C_{1-6} alkyl substituted with $-\text{C}(=\text{O})-\text{C}_{1-6}$ alkyl, C_{1-6} alkyl substituted with $-\text{C}(=\text{O})-\text{O}-\text{C}_{1-6}$
 10 alkyl , C_{1-6} alkyl substituted with C_{1-6} alkoxy C_{1-6} alkyl- $\text{O}-\text{C}(=\text{O})-$, C_{1-6} alkyl substituted with C_{1-6}
 $\text{alkoxyC}_{1-6}\text{alkyl}-\text{C}(=\text{O})-$, C_{1-6} alkyl substituted with $-\text{O}-\text{C}(=\text{O})-\text{C}_{1-6}$ alkyl, C_{1-6} alkoxy C_{1-6} alkyl
 wherein each C_{1-6} alkyl may optionally be substituted with one or two hydroxyl groups or with
 $-\text{O}-\text{C}(=\text{O})-\text{C}_{1-6}$ alkyl, C_{2-6} alkenyl substituted with C_{1-6} alkoxy, C_{2-6} alkynyl substituted with C_{1-6}
 alkoxy , C_{1-6} alkyl substituted with R^9 and optionally substituted with $-\text{O}-\text{C}(=\text{O})-\text{C}_{1-6}$ alkyl, C_{1-6}
 alkyl substituted with $-\text{C}(=\text{O})-\text{R}^9$, C_{1-6} alkyl substituted with hydroxyl and R^9 , C_{2-6} alkenyl
 15 substituted with R^9 , C_{2-6} alkynyl substituted with R^9 , C_{1-6} alkyl substituted with $-\text{NR}^{10}\text{R}^{11}$, C_{2-6}
 alkenyl substituted with $-\text{NR}^{10}\text{R}^{11}$, C_{2-6} alkynyl substituted with $-\text{NR}^{10}\text{R}^{11}$, C_{1-6} alkyl substituted
 with hydroxyl and $-\text{NR}^{10}\text{R}^{11}$, C_{1-6} alkyl substituted with one or two halogens and $-\text{NR}^{10}\text{R}^{11}$, $-\text{C}_{1-6}$
 $\text{alkyl}-\text{C}(\text{R}^{12})=\text{N}-\text{O}-\text{R}^{12}$, C_{1-6} alkyl substituted with $-\text{C}(=\text{O})-\text{NR}^{10}\text{R}^{11}$, C_{1-6} alkyl substituted with
 $-\text{O}-\text{C}(=\text{O})-\text{NR}^{10}\text{R}^{11}$, $-\text{S}(=\text{O})_2-\text{C}_{1-6}$ alkyl, $-\text{S}(=\text{O})_2$ -halo C_{1-6} alkyl, $-\text{S}(=\text{O})_2-\text{NR}^{14}\text{R}^{15}$, C_{1-6}
 20 alkyl substituted with $-\text{S}(=\text{O})_2-\text{C}_{1-6}$ alkyl, C_{1-6} alkyl substituted with $-\text{S}(=\text{O})_2$ -halo C_{1-6} alkyl, C_{1-6}
 alkyl substituted with $-\text{S}(=\text{O})_2-\text{NR}^{14}\text{R}^{15}$, C_{1-6} alkyl substituted with $-\text{NR}^{12}-\text{S}(=\text{O})_2-\text{C}_{1-6}$ alkyl, C_{1-6}
 alkyl substituted with $-\text{NH}-\text{S}(=\text{O})_2$ -halo C_{1-6} alkyl, C_{1-6} alkyl substituted with
 $-\text{NR}^{12}-\text{S}(=\text{O})_2-\text{NR}^{14}\text{R}^{15}$, R^{13} , C_{1-6} alkyl substituted with $-\text{P}(=\text{O})(\text{OH})_2$ or C_{1-6} alkyl substituted
 with $-\text{P}(=\text{O})(\text{OC}_{1-6}\text{alkyl})_2$;

25

R^4 and R^5 each independently represent hydrogen, C_{1-6} alkyl, C_{1-6} alkyl substituted with $-\text{NR}^{14}\text{R}^{15}$, hydroxy C_{1-6} alkyl, halo C_{1-6} alkyl, hydroxyhalo C_{1-6} alkyl, C_{1-6} alkoxy C_{1-6} alkyl wherein
 each C_{1-6} alkyl may optionally be substituted with one or two hydroxyl groups, $-\text{S}(=\text{O})_2-\text{C}_{1-6}$
 alkyl , $-\text{S}(=\text{O})_2$ -halo C_{1-6} alkyl, $-\text{S}(=\text{O})_2-\text{NR}^{14}\text{R}^{15}$, $-\text{C}(=\text{O})-\text{NR}^{14}\text{R}^{15}$, $-\text{C}(=\text{O})-\text{O}-\text{C}_{1-6}$ alkyl, $-\text{C}(=\text{O})-\text{R}^{13}$, C_{1-6} alkyl substituted with $-\text{S}(=\text{O})_2-\text{C}_{1-6}$ alkyl, C_{1-6} alkyl substituted with $-\text{S}(=\text{O})_2$ -
 30 halo C_{1-6} alkyl, C_{1-6} alkyl substituted with $-\text{S}(=\text{O})_2-\text{NR}^{14}\text{R}^{15}$, C_{1-6} alkyl substituted with $-\text{NH}-\text{S}(=\text{O})_2-\text{C}_{1-6}$ alkyl, C_{1-6} alkyl substituted with $-\text{NH}-\text{S}(=\text{O})_2$ -halo C_{1-6} alkyl, C_{1-6} alkyl substituted
 with $-\text{NH}-\text{S}(=\text{O})_2-\text{NR}^{14}\text{R}^{15}$, R^{13} or C_{1-6} alkyl substituted with R^{13} ;

35

R^6 represents C_{3-8} cycloalkyl, C_{3-8} cycloalkenyl, phenyl, 4 to 7-membered monocyclic
 heterocyclyl containing at least one heteroatom selected from N, O or S; said C_{3-8} cycloalkyl,

~~C₃₋₈cycloalkenyl, phenyl, 4 to 7-membered monocyclic heterocyclyl, optionally and each independently being substituted by 1, 2, 3, 4 or 5 substituents, each substituent independently being selected from cyano, C₁₋₆alkyl, cyanoC₁₋₆alkyl, hydroxyl, carboxyl, hydroxyC₁₋₆alkyl, halogen, haloC₁₋₆alkyl, hydroxyhaloC₁₋₆alkyl, C₁₋₆alkoxy, C₁₋₆alkoxyC₁₋₆alkyl, C₁₋₆alkyl-O-C(=O)-, NR¹⁴R¹⁵, -C(=O)-NR¹⁴R¹⁵, C₁₋₆alkyl substituted with NR¹⁴R¹⁵, C₁₋₆alkyl substituted with -C(=O)-NR¹⁴R¹⁵, -S(=O)₂-C₁₋₆alkyl, -S(=O)₂-haloC₁₋₆alkyl, -S(=O)₂-NR¹⁴R¹⁵, C₁₋₆alkyl substituted with -S(=O)₂-C₁₋₆alkyl, C₁₋₆alkyl substituted with -S(=O)₂-haloC₁₋₆alkyl, C₁₋₆alkyl substituted with -S(=O)₂-NR¹⁴R¹⁵, C₁₋₆alkyl substituted with -NH-S(=O)₂-C₁₋₆alkyl, C₁₋₆alkyl substituted with -NH-S(=O)₂-haloC₁₋₆alkyl or C₁₋₆alkyl substituted with -NH-S(=O)₂-NR¹⁴R¹⁵;~~

R⁷ and R⁸ each independently represent hydrogen, C₁₋₆alkyl, hydroxyC₁₋₆alkyl, haloC₁₋₆alkyl, hydroxyhaloC₁₋₆alkyl or C₁₋₆alkoxyC₁₋₆alkyl;

R⁹ represents C₃₋₈cycloalkyl, C₃₋₈cycloalkenyl, phenyl, naphthyl, or 3 to 12 membered monocyclic or bicyclic heterocyclyl containing at least one heteroatom selected from N, O or S, said C₃₋₈cycloalkyl, C₃₋₈cycloalkenyl, phenyl, naphthyl, or 3 to 12 membered monocyclic or bicyclic heterocyclyl each optionally and each independently being substituted with 1, 2, 3, 4 or 5 substituents, each substituent independently being selected from =O, C₁₋₄alkyl, hydroxyl, carboxyl, hydroxyC₁₋₄alkyl, cyano, cyanoC₁₋₄alkyl, C₁₋₄alkyl-O-C(=O)-, C₁₋₄alkyl substituted with C₁₋₄alkyl-O-C(=O)-, C₁₋₄alkyl-C(=O)-, C₁₋₄alkoxyC₁₋₄alkyl wherein each C₁₋₄alkyl may optionally be substituted with one or two hydroxyl groups, halogen, haloC₁₋₄alkyl, hydroxyhaloC₁₋₄alkyl, -NR¹⁴R¹⁵, -C(=O)-NR¹⁴R¹⁵, C₁₋₄alkyl substituted with -NR¹⁴R¹⁵, C₁₋₄alkyl substituted with -C(=O)-NR¹⁴R¹⁵, C₁₋₄alkoxy, -S(=O)₂-C₁₋₄alkyl, -S(=O)₂-haloC₁₋₄alkyl, -S(=O)₂-NR¹⁴R¹⁵, C₁₋₄alkyl substituted with -S(=O)₂-NR¹⁴R¹⁵, C₁₋₄alkyl substituted with -NH-S(=O)₂-C₁₋₄alkyl, C₁₋₄alkyl substituted with -NH-S(=O)₂-haloC₁₋₄alkyl, C₁₋₄alkyl substituted with -NH-S(=O)₂-NR¹⁴R¹⁵, R¹³, -C(=O)-R¹³, C₁₋₄alkyl substituted with R¹³, phenyl optionally substituted with R¹⁶, phenylC₁₋₆alkyl wherein the phenyl is optionally substituted with R¹⁶, a 5 or 6-membered aromatic monocyclic heterocyclyl containing at least one heteroatom selected from N, O or S wherein said heterocyclyl is optionally substituted with R¹⁶;

or when two of the substituents of R⁹ are attached to the same atom, they may be taken together to form a 4 to 7-membered saturated monocyclic heterocyclyl containing at least one heteroatom selected from N, O or S;

R¹⁰ and R¹¹ each independently represent hydrogen, carboxyl, C₁₋₆alkyl, cyanoC₁₋₆alkyl, C₁₋₆alkyl substituted with -NR¹⁴R¹⁵, C₁₋₆alkyl substituted with -C(=O)-NR¹⁴R¹⁵, haloC₁₋₆alkyl,

hydroxyC₁₋₆alkyl, hydroxyhaloC₁₋₆alkyl, C₁₋₆alkoxy, C₁₋₆alkoxyC₁₋₆alkyl wherein each C₁₋₆alkyl may optionally be substituted with one or two hydroxyl groups, R⁶, C₁₋₆alkyl substituted with R⁶, -C(=O)-R⁶, -C(=O)-C₁₋₆alkyl, -C(=O)-hydroxyC₁₋₆alkyl, -C(=O)-haloC₁₋₆alkyl, -C(=O)-hydroxyhaloC₁₋₆alkyl, C₁₋₆alkyl substituted with -Si(CH₃)₃, -S(=O)₂-C₁₋₆alkyl, -S(=O)₂-haloC₁₋₆alkyl, -S(=O)₂-NR¹⁴R¹⁵, C₁₋₆alkyl substituted with -S(=O)₂-C₁₋₆alkyl, C₁₋₆alkyl substituted with -S(=O)₂-haloC₁₋₆alkyl, C₁₋₆alkyl substituted with -S(=O)₂-NR¹⁴R¹⁵, C₁₋₆alkyl substituted with -NH-S(=O)₂-C₁₋₆alkyl, C₁₋₆alkyl substituted with -NH-S(=O)₂-haloC₁₋₆alkyl, C₁₋₆alkyl substituted with carboxyl, or C₁₋₆alkyl substituted with -NH-S(=O)₂-NR¹⁴R¹⁵;

R¹² represents hydrogen or C₁₋₄alkyl optionally substituted with C₄₋₄alkoxy;

R¹³ represents C₃₋₈cycloalkyl or a saturated 4 to 6-membered monocyclic heterocyclyl containing at least one heteroatom selected from N, O or S, wherein said C₃₋₈cycloalkyl or monocyclic heterocyclyl is optionally substituted with 1, 2 or 3 substituents each independently selected from halogen, hydroxyl, C₁₋₆alkyl, haloC₁₋₆alkyl, -O-, cyano, -C(=O)-C₁₋₆alkyl, C₄₋₆alkoxy, or -NR¹⁴R¹⁵;

R¹⁴ and R¹⁵ each independently represent hydrogen, or haloC₁₋₄alkyl, or C₁₋₄alkyl optionally substituted with a substituent selected from hydroxyl, C₄₋₄alkoxy, amino or mono- or di-(C₄₋₄alkyl)amino;

R¹⁶ represents hydroxyl, halogen, cyano, C₁₋₄alkyl, C₁₋₄alkoxy, -NR¹⁴R¹⁵ or -C(=O)NR¹⁴R¹⁵;

R¹⁸ represents hydrogen, C₁₋₆alkyl, C₃₋₈cycloalkyl, C₁₋₄alkyl substituted with C₃₋₈cycloalkyl;

R¹⁹ represents hydrogen; C₁₋₆alkyl; C₃₋₈cycloalkyl; C₁₋₆alkyl substituted with -O-R²⁰; -(CH₂)_r-CN; -(CH₂)_r-CONR²⁰R²¹; -(CH₂)_{r1}-NR²⁰R²¹; -(CH₂)_{r1}-NR²⁰COR²¹; -(CH₂)_{r1}-NR²⁰-(CH₂)_s-SO₂-R²¹; -(CH₂)_{r1}-NH-SO₂-NR²⁰R²¹; -(CH₂)_{r1}-NR²⁰CO₂R²¹; -(CH₂)_r-SO₂NR²⁰R²¹; phenyl optionally substituted with 1, 2, 3, 4 or 5 substituents each independently selected from halogen, C₁₋₄alkyl, C₄₋₄alkoxy, cyano or amine; a 5- or 6-membered aromatic monocyclic heterocycle containing at least one heteroatom selected from N, O or S; said heterocycle being optionally substituted with 1, 2, 3 or 4 substituents each independently selected from halogen, C₁₋₄alkyl, C₄₋₄alkoxy, cyano or amine; wherein said C₁₋₆alkyl and C₃₋₈cycloalkyl may be optionally substituted by one or more R²⁰-groups

R²⁰ and R²¹ independently represent hydrogen, C₁₋₆ alkyl, C₁₋₆ alkanol -(CH₂)_n-O-C₁₋₆alkyl, or when attached to a nitrogen atom R²⁰ and R²¹ can be taken together to form with the nitrogen atom to which they are attached a monocyclic saturated 4, 5 or 6-membered ring which optionally contains a further heteroatom selected from O, S or N;

5

~~R²² and R²³ independently represent hydrogen, C₁₋₆ alkyl, or hydroxyC₁₋₆alkyl;~~

~~m independently represents an integer equal to 0, 1 or 2;~~

~~n independently represents an integer equal to 0, 1, 2, 3 or 4;~~

10 ~~s independently represents an integer equal to 0, 1, 2, 3 or 4;~~

~~r independently represent an integer equal to 1, 2, 3, or 4;~~

~~r1 independently represent an integer equal to 2, 3 or 4;~~

~~a N-oxide thereof or, a pharmaceutically acceptable salt thereof or a solvate thereof.~~

15 According to a second aspect of the disclosure there is provided a pharmaceutical composition comprising a compound of formula (I-A) or (I-B) as defined above.

According to a third aspect of the disclosure there is provided a compound as defined above for use in therapy.

20

According to a fourth aspect of the disclosure there is provided a compound as defined above for use in the prophylaxis or treatment of a disease state or condition mediated by a FGFR kinase.

25 According to a fifth aspect of the disclosure there is provided a compound as defined above for use in the prophylaxis or treatment of cancer.

According to a sixth aspect of the disclosure there is provided use of a compound as defined above for the manufacture of a medicament for the prophylaxis or treatment of cancer.

30

Preferred compounds of the present invention are compounds having an IC_{50} value of less than $1\text{ }\mu\text{M}$, more preferably less than $0.1\text{ }\mu\text{M}$.

The ~~invention~~ disclosure provides compounds that have FGFR inhibiting or modulating activity, and which may be useful in preventing or treating disease states or conditions mediated by FGFR kinases.

In one ~~embodiment~~ example, there is provided a compound as defined herein for use in therapy, for use as a medicine. In a further ~~embodiment~~ example, there is provided a compound as defined herein for use in the prophylaxis or treatment, in particular in the treatment, of a disease state or condition mediated by a FGFR kinase.

Thus, for example, the compounds of the ~~invention~~ disclosure may be useful in alleviating or reducing the incidence of cancer. Therefore, in a further ~~example~~ embodiment, there is provided a compound as defined herein for use in the prophylaxis or treatment, in particular the treatment, of cancer. In one ~~example~~ embodiment, the compound as defined herein is for use in the prophylaxis or treatment of FGFR-dependent cancer. In one ~~example~~ embodiment, the compound as defined herein is for use in the prophylaxis or treatment of cancer mediated by FGFR kinases.

Accordingly, the ~~disclosure~~ invention provides *inter alia*:

- A method for the prophylaxis or treatment of a disease state or condition mediated by a FGFR kinase, which method comprises administering to a subject in need thereof a compound of the formula (I) as defined herein.
- A method for the prophylaxis or treatment of a disease state or condition as described herein, which method comprises administering to a subject in need thereof a compound of the formula (I) as defined herein.
- A method for the prophylaxis or treatment of cancer, which method comprises administering to a subject in need thereof a compound of the formula (I) as defined herein.
- ~~A method for alleviating or reducing the incidence of a disease state or condition mediated by a FGFR kinase, which method comprises administering to a subject in need thereof a compound of the formula (I) as defined herein.~~
- A method of inhibiting a FGFR kinase, which method comprises contacting the kinase with a kinase-inhibiting compound of the formula (I) as defined herein.


- A method of modulating a cellular process (for example cell division) by inhibiting the activity of a FGFR kinase using a compound of the formula (I) as defined herein.
- A compound of formula (I) as defined herein for use as a modulator of a cellular process (for example cell division) by inhibiting the activity of a FGFR kinase.
- A compound of formula (I) as defined herein for use in the prophylaxis or treatment of cancer, in particular the treatment of cancer.
- ~~– A compound of formula (I) as defined herein for use as a modulator (e.g. inhibitor) of FGFR.~~
- The use of a compound of formula (I) as defined herein for the manufacture of a medicament for the prophylaxis or treatment of a disease state or condition mediated by a FGFR kinase, the compound having the formula (I) as defined herein.
- The use of a compound of formula (I) as defined herein for the manufacture of a medicament for the prophylaxis or treatment of a disease state or condition as described herein.
- The use of a compound of formula (I) as defined herein for the manufacture of a medicament for the prophylaxis or treatment, in particular the treatment, of cancer.
- ~~– The use of a compound of formula (I) as defined herein for the manufacture of a medicament for modulating (e.g. inhibiting) the activity of FGFR.~~
- Use of a compound of formula (I) as defined herein in the manufacture of a medicament for modulating a cellular process (for example cell division) by inhibiting the activity of a FGFR kinase.
- The use of a compound of the formula (I) as defined herein for the manufacture of a medicament for prophylaxis or treatment of a disease or condition characterised by up-regulation of a FGFR kinase (e.g. FGFR1 or FGFR2 or FGFR3 or FGFR4).
- The use of a compound of the formula (I) as defined herein for the manufacture of a medicament for the prophylaxis or treatment of a cancer, the cancer being one which is characterised by up-regulation of a FGFR kinase (e.g. FGFR1 or FGFR2 or FGFR3 or FGFR4).

- ~~— The use of a compound of the formula (I) as defined herein for the manufacture of a medicament for the prophylaxis or treatment of cancer in a patient selected from a sub-population possessing a genetic aberrations of FGFR3 kinase.~~
- ~~— The use of a compound of the formula (I) as defined herein for the manufacture of a medicament for the prophylaxis or treatment of cancer in a patient who has been diagnosed as forming part of a sub-population possessing a genetic aberrations of FGFR3 kinase.~~
- A method for the prophylaxis or treatment of a disease or condition characterised by up-regulation of a FGFR kinase (e.g. FGFR1 or FGFR2 or FGFR3 or FGFR4), the method comprising administering a compound of the formula (I) as defined herein.
- A method for alleviating or reducing the incidence of a disease or condition characterised by up-regulation of a FGFR kinase (e.g. FGFR1 or FGFR2 or FGFR3 or FGFR4), the method comprising administering a compound of the formula (I) as defined herein.
- ~~— A method for the prophylaxis or treatment of (or alleviating or reducing the incidence of) cancer in a patient suffering from or suspected of suffering from cancer; which method comprises (i) subjecting a patient to a diagnostic test to determine whether the patient possesses a genetic aberrations of FGFR3 gene; and (ii) where the patient does possess the said variant, thereafter administering to the patient a compound of the formula (I) as defined herein having FGFR3 kinase inhibiting activity.~~
- ~~— A method for the prophylaxis or treatment of (or alleviating or reducing the incidence of) a disease state or condition characterised by up-regulation of an FGFR kinase (e.g. FGFR1 or FGFR2 or FGFR3 or FGFR4); which method comprises (i) subjecting a patient to a diagnostic test to detect a marker characteristic of up-regulation of a FGFR kinase (e.g. FGFR1 or FGFR2 or FGFR3 or FGFR4) and (ii) where the diagnostic test is indicative of up-regulation of a FGFR kinase, thereafter administering to the patient a compound of the formula (I) as defined herein having FGFR kinase inhibiting activity.~~


In one ~~embodiment~~example, the disease mediated by FGFR kinases is a oncology related disease (e.g. cancer). In one example~~embodiment~~, the disease mediated by FGFR kinases is a non-oncology related disease (e.g. any disease disclosed herein excluding cancer). In

CLAIMS

1. A compound of formula (I-A) or (I-B):





or



including any tautomeric or stereochemically isomeric form thereof, wherein

each R^2 is independently selected from hydroxyl, halogen, cyano, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, C_{1-4} alkoxy, hydroxy C_{1-4} alkyl, hydroxy C_{1-4} alkoxy, halo C_{1-4} alkyl, halo C_{1-4} alkoxy, hydroxyhalo C_{1-4} alkyl, hydroxyhalo C_{1-4} alkoxy, C_{1-4} alkoxy C_{1-4} alkyl, halo C_{1-4} alkoxy C_{1-4} alkyl, C_{1-4} alkoxy C_{1-4} alkyl wherein each C_{1-4} alkyl may optionally be substituted with one or two hydroxyl groups, hydroxyhalo C_{1-4} alkoxy C_{1-4} alkyl, R^{13} , C_{1-4} alkyl substituted with R^{13} , C_{1-4} alkyl substituted with $-C(=O)-R^{13}$, C_{1-4} alkoxy substituted with R^{13} , C_{1-4} alkoxy substituted with $-C(=O)-R^{13}$, $-C(=O)-R^{13}$, C_{1-4} alkyl substituted with $-NR^7R^8$, C_{1-4} alkyl substituted with $-C(=O)-NR^7R^8$, C_{1-4} alkoxy substituted with $-NR^7R^8$, C_{1-4} alkoxy substituted with $-C(=O)-NR^7R^8$, $-NR^7R^8$ and $-C(=O)-NR^7R^8$; or when two R^2 groups are attached to adjacent carbon atoms they may be taken together to form a radical of formula:



Y represents $-CR^{18}=N-OR^{19}$ or $-E-D$;

D represents a 3 to 12 ring membered monocyclic or bicyclic carbocyclyl or a 3 to 12 ring membered monocyclic or bicyclic heterocyclyl containing at least one heteroatom selected from N, O or S, wherein said carbocyclyl and heterocyclyl may each be optionally substituted by one or more (e.g. 1, 2 or 3) R¹ groups;

E represents a bond;

R¹ represents hydrogen, halo, cyano, C₁₋₆alkyl, C₁₋₆alkoxy, -C(=O)-O-C₁₋₆alkyl, C₂₋₆alkenyl, hydroxyC₁₋₆alkyl, haloC₁₋₆alkyl, hydroxyhaloC₁₋₆alkyl, cyanoC₁₋₄alkyl, C₁₋₆alkoxyC₁₋₆alkyl wherein each C₁₋₆alkyl may optionally be substituted with one or two hydroxyl groups, -NR⁴R⁵, C₁₋₆alkyl substituted with -O-C(=O)-C₁₋₆alkyl, C₁₋₆alkyl substituted with -NR⁴R⁵, -C(=O)-NR⁴R⁵, -C(=O)-C₁₋₆alkyl-NR⁴R⁵, C₁₋₆alkyl substituted with -C(=O)-NR⁴R⁵, -S(=O)₂-C₁₋₆alkyl, -S(=O)₂-haloC₁₋₆alkyl, -S(=O)₂-NR¹⁴R¹⁵, C₁₋₆alkyl substituted with -S(=O)₂-C₁₋₆alkyl, C₁₋₆alkyl substituted with -S(=O)₂-haloC₁₋₆alkyl, C₁₋₆alkyl substituted with -S(=O)₂-NR¹⁴R¹⁵, C₁₋₆alkyl substituted with -NH-S(=O)₂-C₁₋₆alkyl, C₁₋₆alkyl substituted with -NH-S(=O)₂-haloC₁₋₆alkyl, C₁₋₆alkyl substituted with -NR¹²-S(=O)₂-NR¹⁴R¹⁵, R⁶, C₁₋₆alkyl substituted with R⁶, -C(=O)-R⁶, C₁₋₆alkyl substituted with -C(=O)-R⁶, hydroxyC₁₋₆alkyl substituted with R⁶, C₁₋₆alkyl substituted with -Si(CH₃)₃, C₁₋₆alkyl substituted with -P(=O)(OH)₂ or C₁₋₆alkyl substituted with -P(=O)(OC₁₋₆alkyl)₂;

R³ represents hydroxyl, C₁₋₆alkoxy, hydroxyC₁₋₆alkoxy, C₁₋₆alkoxy substituted with -NR¹⁰R¹¹, C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, haloC₁₋₆alkyl optionally substituted with -O-C(=O)-C₁₋₆alkyl, hydroxyC₁₋₆alkyl optionally substituted with -O-C(=O)-C₁₋₆alkyl, hydroxyC₂₋₆alkenyl, hydroxyC₂₋₆alkynyl, hydroxyhaloC₁₋₆alkyl, cyanoC₁₋₆alkyl, C₁₋₆alkyl substituted with carboxyl, C₁₋₆alkyl substituted with -C(=O)-C₁₋₆alkyl, C₁₋₆alkyl substituted with -C(=O)-O-C₁₋₆alkyl, C₁₋₆alkyl substituted with C₁₋₆alkoxyC₁₋₆alkyl-O-C(=O)-, C₁₋₆alkyl substituted with C₁₋₆alkoxyC₁₋₆alkyl-C(=O)-, C₁₋₆alkyl substituted with -O-C(=O)-C₁₋₆alkyl, C₁₋₆alkoxyC₁₋₆alkyl wherein each C₁₋₆alkyl may optionally be substituted with one or two hydroxyl groups or with -O-C(=O)-C₁₋₆alkyl, C₂₋₆alkenyl substituted with C₁₋₆alkoxy, C₂₋₆alkynyl substituted with C₁₋₆alkoxy, C₁₋₆alkyl substituted with R⁹ and optionally substituted with -O-C(=O)-C₁₋₆alkyl, C₁₋₆alkyl substituted

with $-C(=O)-R^9$, C_{1-6} alkyl substituted with hydroxyl and R^9 , C_{2-6} alkenyl substituted with R^9 , C_{2-6} alkynyl substituted with R^9 , C_{1-6} alkyl substituted with $-NR^{10}R^{11}$, C_{2-6} alkenyl substituted with $-NR^{10}R^{11}$, C_{2-6} alkynyl substituted with $-NR^{10}R^{11}$, C_{1-6} alkyl substituted with hydroxyl and $-NR^{10}R^{11}$, C_{1-6} alkyl substituted with one or two halogens and $-NR^{10}R^{11}$, $-C_{1-6}$ alkyl- $C(R^{12})=N-O-R^{12}$, C_{1-6} alkyl substituted with $-C(=O)-NR^{10}R^{11}$, C_{1-6} alkyl substituted with $-O-C(=O)-NR^{10}R^{11}$, $-S(=O)_2-C_{1-6}$ alkyl, $-S(=O)_2$ -halo C_{1-6} alkyl, $-S(=O)_2-NR^{14}R^{15}$, C_{1-6} alkyl substituted with $-S(=O)_2-C_{1-6}$ alkyl, C_{1-6} alkyl substituted with $-S(=O)_2$ -halo C_{1-6} alkyl, C_{1-6} alkyl substituted with $-S(=O)_2-NR^{14}R^{15}$, C_{1-6} alkyl substituted with $-NR^{12}-S(=O)_2-C_{1-6}$ alkyl, C_{1-6} alkyl substituted with $-NH-S(=O)_2$ -halo C_{1-6} alkyl, C_{1-6} alkyl substituted with $-NR^{12}-S(=O)_2-NR^{14}R^{15}$, R^{13} , C_{1-6} alkyl substituted with $-P(=O)(OH)_2$ or C_{1-6} alkyl substituted with $-P(=O)(OC_{1-6}alkyl)_2$;

R^4 and R^5 each independently represent hydrogen, C_{1-6} alkyl, C_{1-6} alkyl substituted with $-NR^{14}R^{15}$, hydroxy C_{1-6} alkyl, halo C_{1-6} alkyl, hydroxyhalo C_{1-6} alkyl, C_{1-6} alkoxy C_{1-6} alkyl wherein each C_{1-6} alkyl may optionally be substituted with one or two hydroxyl groups, $-S(=O)_2-C_{1-6}$ alkyl, $-S(=O)_2$ -halo C_{1-6} alkyl, $-S(=O)_2-NR^{14}R^{15}$, $-C(=O)-NR^{14}R^{15}$, $-C(=O)-O-C_{1-6}$ alkyl, $-C(=O)-R^{13}$, C_{1-6} alkyl substituted with $-S(=O)_2-C_{1-6}$ alkyl, C_{1-6} alkyl substituted with $-S(=O)_2$ -halo C_{1-6} alkyl, C_{1-6} alkyl substituted with $-S(=O)_2-NR^{14}R^{15}$, C_{1-6} alkyl substituted with $-NH-S(=O)_2-C_{1-6}$ alkyl, C_{1-6} alkyl substituted with $-NH-S(=O)_2$ -halo C_{1-6} alkyl, C_{1-6} alkyl substituted with $-NH-S(=O)_2-NR^{14}R^{15}$, R^{13} or C_{1-6} alkyl substituted with R^{13} ;

R^6 represents C_{3-8} cycloalkyl, C_{3-8} cycloalkenyl, phenyl, 4 to 7-membered monocyclic heterocyclyl containing at least one heteroatom selected from N, O or S; said C_{3-8} cycloalkyl, C_{3-8} cycloalkenyl, phenyl, 4 to 7-membered monocyclic heterocyclyl, optionally and each independently being substituted by 1, 2, 3, 4 or 5 substituents, each substituent independently being selected from cyano, C_{1-6} alkyl, cyano C_{1-6} alkyl, hydroxyl, carboxyl, hydroxy C_{1-6} alkyl, halogen, halo C_{1-6} alkyl, hydroxyhalo C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} alkoxy C_{1-6} alkyl, C_{1-6} alkyl $O-C(=O)-$, $-NR^{14}R^{15}$, $-C(=O)-NR^{14}R^{15}$, C_{1-6} alkyl substituted with $-NR^{14}R^{15}$, C_{1-6} alkyl substituted with $-C(=O)-NR^{14}R^{15}$, $-S(=O)_2-C_{1-6}$ alkyl, $-S(=O)_2$ -halo C_{1-6} alkyl, $-S(=O)_2-NR^{14}R^{15}$, C_{1-6} alkyl substituted with $-S(=O)_2-C_{1-6}$ alkyl, C_{1-6} alkyl substituted with $-S(=O)_2$ -halo C_{1-6} alkyl, C_{1-6} alkyl substituted with $-S(=O)_2$ -

~~NR¹⁴R¹⁵, C₁₋₆alkyl substituted with -NH-S(=O)₂-C₁₋₆alkyl, C₁₋₆alkyl substituted with -NH-S(=O)₂-haloC₁₋₆alkyl or C₁₋₆alkyl substituted with -NH-S(=O)₂-NR¹⁴R¹⁵;~~

R⁷ and R⁸ each independently represent hydrogen, C₁₋₆alkyl, hydroxyC₁₋₆alkyl, haloC₁₋₆alkyl, hydroxyhaloC₁₋₆alkyl or C₁₋₆alkoxyC₁₋₆alkyl;

R⁹ represents C₃₋₈cycloalkyl, C₃₋₈cycloalkenyl, phenyl, naphthyl, or 3 to 12 membered monocyclic or bicyclic heterocyclyl containing at least one heteroatom selected from N, O or S, said C₃₋₈cycloalkyl, C₃₋₈cycloalkenyl, phenyl, naphthyl, or 3 to 12 membered monocyclic or bicyclic heterocyclyl each optionally and each independently being substituted with 1, 2, 3, 4 or 5 substituents, each substituent independently being selected from =O, C₁₋₄alkyl, hydroxyl, carboxyl, hydroxyC₁₋₄alkyl, cyano, cyanoC₁₋₄alkyl, C₁₋₄alkyl-O-C(=O)-, C₁₋₄alkyl substituted with C₁₋₄alkyl-O-C(=O)-, C₁₋₄alkyl-C(=O)-, C₁₋₄alkoxyC₁₋₄alkyl ~~wherein each C₁₋₄alkyl may optionally be substituted with one or two hydroxyl groups, halogen, haloC₁₋₄alkyl, hydroxyhaloC₁₋₄alkyl, -NR¹⁴R¹⁵, -C(=O)-NR¹⁴R¹⁵, C₁₋₄alkyl substituted with -NR¹⁴R¹⁵, C₁₋₄alkyl substituted with -C(=O)-NR¹⁴R¹⁵, C₁₋₄alkoxy, -S(=O)₂-C₁₋₄alkyl, -S(=O)₂-haloC₁₋₄alkyl, -S(=O)₂-NR¹⁴R¹⁵, C₁₋₄alkyl substituted with -S(=O)₂-NR¹⁴R¹⁵, C₁₋₄alkyl substituted with -NH-S(=O)₂-C₁₋₄alkyl, C₁₋₄alkyl substituted with -NH-S(=O)₂-haloC₁₋₄alkyl, C₁₋₄alkyl substituted with -NH-S(=O)₂-NR¹⁴R¹⁵, R¹³, -C(=O)-R¹³, C₁₋₄alkyl substituted with R¹³, phenyl optionally substituted with R¹⁶, phenylC₁₋₆alkyl wherein the phenyl is optionally substituted with R¹⁶, a 5 or 6-membered aromatic monocyclic heterocyclyl containing at least one heteroatom selected from N, O or S wherein said heterocyclyl is optionally substituted with R¹⁶;~~

or when two of the substituents of R⁹ are attached to the same atom, they may be taken together to form a 4 to 7-membered saturated monocyclic heterocyclyl containing at least one heteroatom selected from N, O or S;

R¹⁰ and R¹¹ each independently represent hydrogen, carboxyl, C₁₋₆alkyl, cyanoC₁₋₆alkyl, C₁₋₆alkyl substituted with -NR¹⁴R¹⁵, C₁₋₆alkyl substituted with -C(=O)-NR¹⁴R¹⁵, haloC₁₋₆alkyl, hydroxyC₁₋₆alkyl, hydroxyhaloC₁₋₆alkyl, C₁₋₆alkoxy, C₁₋₆alkoxyC₁₋₆alkyl ~~wherein each C₁₋₆alkyl may optionally be substituted with one or two hydroxyl groups,~~ R⁶, C₁₋₆alkyl substituted with R⁶, -C(=O)-R⁶, -C(=O)-C₁₋₆alkyl, -C(=O)-hydroxyC₁₋₆alkyl, -C(=O)-haloC₁₋₆alkyl, -C(=O)-hydroxyhaloC₁₋₆alkyl, C₁₋₆alkyl substituted with -Si(CH₃)₃, -S(=O)₂-C₁₋₆alkyl, -S(=O)₂-haloC₁₋₆alkyl, -S(=O)₂-NR¹⁴R¹⁵, C₁₋₆alkyl substituted with -S(=O)₂-C₁₋₆alkyl, C₁₋₆alkyl substituted with -S(=O)₂-haloC₁₋₆alkyl, C₁₋₆alkyl substituted

with $-S(=O)_2-NR^{14}R^{15}$, C_{1-6} alkyl substituted with $-NH-S(=O)_2-C_{1-6}$ alkyl, C_{1-6} alkyl substituted with $-NH-S(=O)_2$ -halo C_{1-6} alkyl, C_{1-6} alkyl substituted with carboxyl, or C_{1-6} alkyl substituted with $-NH-S(=O)_2-NR^{14}R^{15}$;

R^{12} represents hydrogen or C_{1-4} alkyl optionally substituted with C_{1-4} alkoxy;

R^{13} represents C_{3-8} cycloalkyl or a saturated 4 to 6-membered monocyclic heterocyclyl containing at least one heteroatom selected from N, O or S, wherein said C_{3-8} cycloalkyl or monocyclic heterocyclyl is optionally substituted with 1, 2 or 3 substituents each independently selected from halogen, hydroxyl, C_{1-6} alkyl, halo C_{1-6} alkyl, $=O$, cyano, $-C(=O)-C_{1-6}$ alkyl, C_{1-6} alkoxy, or $-NR^{14}R^{15}$;

R^{14} and R^{15} each independently represent hydrogen, or halo C_{1-4} alkyl, or C_{1-4} alkyl optionally substituted with a substituent selected from hydroxyl, C_{1-4} alkoxy, amino or mono- or di(C_{1-4} alkyl)amino;

R^{16} represents hydroxyl, halogen, cyano, C_{1-4} alkyl, C_{1-4} alkoxy, $-NR^{14}R^{15}$ or $-C(=O)NR^{14}R^{15}$;

R^{18} represents hydrogen, C_{1-6} alkyl, C_{3-8} cycloalkyl, C_{1-4} alkyl substituted with C_{3-8} cycloalkyl;

R^{19} represents hydrogen; C_{1-6} alkyl; C_{3-8} cycloalkyl; C_{1-6} alkyl substituted with $-O-R^{20}$; $-(CH_2)_r-CN$; $-(CH_2)_r-CONR^{20}R^{21}$; $-(CH_2)_{r1}-NR^{20}R^{21}$; $-(CH_2)_{r1}-NR^{20}COR^{21}$; $-(CH_2)_{r1}-NR^{20}-(CH_2)_s-SO_2-R^{21}$; $-(CH_2)_{r1}-NH-SO_2-NR^{20}R^{21}$; $-(CH_2)_{r1}-NR^{20}CO_2R^{21}$; $-(CH_2)_r-SO_2NR^{20}R^{21}$; phenyl optionally substituted with 1, 2, 3, 4 or 5 substituents each independently selected from halogen, C_{1-4} alkyl, C_{1-4} alkyloxy, cyano or amino; a 5- or 6-membered aromatic monocyclic heterocycle containing at least one heteroatom selected from N, O or S, said heterocycle being optionally substituted with 1, 2, 3 or 4 substituents each independently selected from halogen, C_{1-4} alkyl, C_{1-4} alkyloxy, cyano or amino; wherein said C_{1-6} alkyl and C_{3-8} cycloalkyl, may be optionally substituted by one or more R^{20} groups

R^{20} and R^{21} independently represent hydrogen, C_{1-6} alkyl, C_{1-6} alkanol $-(CH_2)_n-O-C_{1-6}$ alkyl, or when attached to a nitrogen atom R^{20} and R^{21} can be taken together to form

with the nitrogen atom to which they are attached a monocyclic saturated 4, 5 or 6-membered ring which optionally contains a further heteroatom selected from O, S or N;

R^{22} and R^{23} independently represent hydrogen, C_{1-6} alkyl, or hydroxy C_{1-6} alkyl;

m independently represents an integer equal to 0, 1 or 2;

n independently represents an integer equal to 0, 1, 2, 3 or 4;

s independently represents an integer equal to 0, 1, 2, 3 or 4;

r independently represent an integer equal to 1, 2, 3, or 4;

r1 independently represent an integer equal to 2, 3 or 4;

a N-oxide thereof, or a pharmaceutically acceptable salt thereof ~~or a solvate thereof~~.

2. A compound according to claim 1 wherein the compound is a compound of formula (I-A).


3. A compound according to claim 1 wherein the compound is a compound of formula (I-B).

4. A compound according to any one of the preceding claims wherein Y represents - $CR^{18}=N-OR^{19}$.


5. A compound according to any one of claims 1 to 3 wherein Y is -E-D.

6. A compound according to claim 5 wherein D is optionally substituted pyrazolyl.

7. A compound according to any one of the preceding claims wherein R^2 is independently selected from hydroxyl, halogen, cyano, C_{1-4} alkyl, C_{2-4} alkenyl, C_{1-4} alkoxy, hydroxy C_{1-4} alkyl, hydroxy C_{1-4} alkoxy, halo C_{1-4} alkoxy, C_{1-4} alkoxy C_{1-4} alkyl, R^{13} ,



or




including any tautomeric or stereochemically isomeric form thereof;

wherein n , R^2 and R^3 are as defined in any of the preceding claims;

a *N*-oxide thereof or, a pharmaceutically acceptable salt thereof ~~or a solvate thereof~~.


14. A compound according to claim 13, wherein R^2 represents C_{1-4} alkoxy (for example CH_3O-) or halogen (for example fluoro) or hydroxyl; R^3 represents C_{1-4} alkyl (e.g. $-CH_2-$ or $-CH_2-CH_2-CH_2-$) substituted with R^9 (e.g. wherein R^9 represents an optionally substituted aromatic 5 membered monocyclic heterocyclyl, for example unsubstituted imidazolyl (e.g. imidazol-2-yl), or unsubstituted triazolyl (e.g. triazol-3-yl) or R^9 represents an optionally substituted saturated 5 membered monocyclic heterocyclyl, for example 2-pyrrolidinonyl (e.g. 2-pyrrolidinon-5-yl or 2-pyrrolidinon-1-yl) or 2-oxazolidinonyl (e.g. 2-oxazolidinon-5-yl)) or R^3 represents C_{1-4} alkyl (e.g. $-CH_2CH_2-$) substituted with $-NR^{10}R^{11}$ wherein one of R^{10} and R^{11} represents hydrogen and the other represents C_{1-6} alkyl, for example $-CH_3$ or $-CH(CH_3)_2$ (e.g. R^3 represents $-CH_2CH_2NHCH_3$ or $-CH_2CH_2NHCH(CH_3)_2$), or $-C(=O)-C_{1-6}$ alkyl, for example $-C(=O)-CH_3$ (e.g. R^3 represents $-CH_2CH_2NH-C(=O)-CH_3$), or R^6 (e.g. wherein R^6 represents an optionally substituted 4 membered saturated heterocycle (e.g. oxetanyl)); n is 2, 3, or 4.

15. A compound according to claim 1 wherein the compound is




a *N*-oxide thereof or, a pharmaceutically acceptable salt thereof ~~or a solvate thereof~~.

16. A compound according to claim 1 wherein the compound is



a *N*-oxide thereof or, a pharmaceutically acceptable salt thereof ~~or a solvate thereof~~.

17. A compound according to claim 1 wherein the compound is



a *N*-oxide thereof or, a pharmaceutically acceptable salt thereof ~~or a solvate thereof~~.

18. A compound according to any one of claims 1 to 17 or a pharmaceutically acceptable salt ~~or solvate~~ thereof.

19. A pharmaceutical composition comprising a compound of formula (I-A) or (I-B) as claimed in any one of claims 1 to 18.

20. A compound as claimed in any one of claims 1 to 18 for use in therapy.

21. A compound as claimed in any one of claims 1 to 18 for use in the prophylaxis or treatment of a disease state or condition mediated by a FGFR kinase.