NOVEL FUSED PYRROLOCARBAZOLES

CROSS-REFERENCE TO RELATED APPLICATIONS

This application claims priority of U.S. Provisional Application No. 60/532,182, filed December 23, 2003, which is incorporated herein by reference in its entirety for all purposes.

FIELD OF THE INVENTION

The present invention relates generally to fused pyrrolocarbazoles, including pharmaceutical compositions, diagnostic kits, assay standards or reagents containing the same, and methods of using the same as therapeutics. The invention is also directed to intermediates and processes for making these novel compounds.


BACKGROUND OF THE INVENTION

Publications cited throughout this disclosure are incorporated in their entirety herein by reference.

Various synthetic small organic molecules that are biologically active and generally known in the art as "fused pyrrolocarbazoles" have been prepared (See U.S. Patent Nos. 5,475,110; 5,591,855; 5,594,009; 5,616,724; and 6,630,500). In addition, U.S. Patent 5,705,511 discloses fused pyrrolocarbazole compounds which possess a variety of functional pharmacological activities. The fused pyrrolocarbazoles were disclosed to be used in a variety of ways, including: enhancing the function and/or survival of cells of neuronal lineage, either singularly or in combination with neurotrophic factor(s) and/or indolocarbazoles; enhancing trophic factor-induced activity; inhibition of protein kinase C ("PKC") activity; inhibition of *trk* tyrosine kinase activity; inhibition of proliferation of a prostate cancer cell-line; inhibition of the cellular pathways involved in the inflammation process; and enhancement of the survival of neuronal cells at risk of dying. However, there remains a need for novel pyrrolocarbazole derivatives that possess beneficial properties. This invention is directed to this, as well as other important ends.

SUMMARY OF THE INVENTION

The present invention in one aspect is directed to fused pyrrolocarbazole compounds of Formula I:



I

and its stereoisomeric forms, mixtures of stereoisomeric forms, or pharmaceutically acceptable salt forms thereof, wherein the constituent members are defined *infra*.


The fused pyrrolocarbazoles of the present invention may be used in a variety of ways, including: for inhibition of angiogenesis; as antitumor agents; for enhancing the function and/or survival of cells of neuronal lineage, either singularly or in combination with neurotrophic factor(s) and/or indolocarbazoles; for enhancing trophic factor-induced activity; inhibition of kinase activity, such as *trk* tyrosine kinase ("*trk*"), vascular endothelial growth factor receptor ("VEGFR") kinase, preferably VEGFR1 and VEGFR2, mixed lineage kinase ("MLK"), dual leucine zipper bearing kinase ("DLK"), platelet derived growth factor receptor kinase ("PDGFR"), protein kinase C ("PKC"), Tie-2, or CDK-1, -2, -3, -4, -5, -6; for inhibition of NGF-stimulated *trk* phosphorylation; for inhibition of proliferation of a prostate cancer cell-line; for inhibition of the cellular pathways involved in the inflammation process; and for enhancement of the survival of neuronal cells at risk of dying. In addition, the fused pyrrolocarbazoles may useful for inhibition of c-met, c-kit, and mutated Flt-3 containing internal tandem duplications in the juxtamembrane domain. Because of these varied activities, the disclosed compounds find utility in a variety of settings, including research and therapeutic environments.

In other embodiments, the compounds of the present invention are useful for treating or preventing angiogenesis and angiogenic disorders such as cancer of solid tumors, endometriosis, retinopathy, diabetic retinopathy, psoriasis, hemangioblastoma, ocular disorders or macular degeneration. In another embodiment, the compounds of the present invention are useful for treating or preventing neoplasia, rheumatoid arthritis, chronic arthritis, pulmonary fibrosis, myelofibrosis, abnormal wound healing, atherosclerosis, or restenosis. In further embodiments, the compounds of the present invention are useful for treating or preventing neurodegenerative diseases and disorders, such as Alzheimer's disease, amyotrophic lateral sclerosis, Parkinson's disease, stroke, ischemia, Huntington's disease, AIDS dementia, epilepsy, multiple sclerosis, peripheral neuropathy, chemotherapy induced peripheral neuropathy, AIDS related peripheral neuropathy, or injuries of the brain or spinal chord. In additional embodiments, the compounds of the present invention are useful for treating or preventing prostate disorders such as prostate cancer or benign prostate hyperplasia. In still other embodiments, the compounds of the present invention are useful for treating or preventing multiple myeloma and leukemias including, but not limited to, acute myelogenous leukemia, chronic myelogenous leukemia, acute lymphocytic leukemia, and chronic lymphocytic leukemia.

In further aspect, the present invention is directed to pharmaceutical compositions which comprises one or more pharmaceutically acceptable excipients and a therapeutically effective amount of a compound of the present invention.

DETAILED DESCRIPTION

Thus, in a first embodiment, the present invention provides a novel compound of Formula I:



I

wherein:

ring A and ring B, independently, and each together with the carbon atoms to which they are attached, are selected from:

(a) a phenylene ring in which from 1 to 3 carbon atoms may be replaced by nitrogen atoms; and

(b) a 5-membered aromatic ring in which either

(1) one carbon atom may be replaced with an oxygen, nitrogen, or sulfur atom;

(2) two carbon atoms may be replaced with a sulfur and a nitrogen atom, an oxygen and a nitrogen atom, or two nitrogen atoms; or

(3) three carbon atoms may be replaced with three nitrogen atoms, one oxygen and two nitrogen atoms, or one sulfur and two nitrogen atoms;

A^1 and A^2 are independently selected from H, H; H, OR; H, SR; H, $N(R)_2$; and a group wherein A^1 and A^2 together form a moiety selected from =O, =S, and =NR;

B^1 and B^2 are independently selected from H, H; H, OR; H, SR; H, $N(R)_2$; and a group wherein B^1 and B^2 together form a moiety selected from =O, =S, and =NR;

provided that at least one of the pairs A^1 and A^2, or B^1 and B^2 forms =O;

R is independently selected from H, optionally substituted alkyl, $C(=O)R^{1a}$, $C(=O)NR^{1c}R^{1d}$, optionally substituted arylalkyl and optionally substituted

heteroarylalkyl, wherein said optional substituents are one to three R^{10} groups;

R^1 is independently selected from H, $C(=O)R^{1a}$, OR^{1b}, $C(=O)NHR^{1b}$, $NR^{1c}R^{1d}$, and optionally substituted alkyl, wherein said optional substituents are one to three R^{10} groups;

R^{1a} is independently selected from optionally substituted alkyl, optionally substituted aryl and optionally substituted heteroaryl, wherein said optional substituents are one to three R^{10} groups;

R^{1b} is independently selected from H and optionally substituted alkyl, wherein said optional substituents are one to three R^{10} groups;

R^{1c} and R^{1d} are each independently selected from H and an optionally substituted alkyl, or together with the nitrogen to which they are attached form an optionally substituted heterocycloalkyl, wherein said optional substituents are one to three R^{10} groups;

R^2 is selected from H, $C(=O)R^{2a}$, $C(=O)NR^{2c}R^{2d}$, SO_2R^{2b}, CO_2R^{2b}, (alkylene)-$OC(=O)$-(alkylene)CO_2R^{11}, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl, wherein said optional substituents are one to three R^{10} groups;

R^{2a} is independently selected from optionally substituted alkyl, optionally substituted aryl, OR^{2b}, and $NR^{2c}R^{2d}$, wherein said optional substituents are one to three R^{10} groups;

R^{2b} is selected from H and optionally substituted alkyl, wherein said optional substituents are one to three R^{10} groups;

R^{2c} and R^{2d} are each independently selected from H and optionally substituted alkyl, or together with the nitrogen to which they are attached form an optionally substituted heterocycloalkyl, wherein said optional substituents are one to three R^{10} groups;

at least one of R^3, R^4, R^5, and R^6 is selected from $(alkylene)_xOR^{13}$, $C(=O)R^{13}$, $(CH_2)_pOR^{22}$, O-(alkylene)-R^{27}, $OCH(CO_2R^{18})_2$, $OCH[(CH_2)_pOR^{20}]_2$, C(=O)-(alkylene)-R^{25}, $NR^{11}R^{32}$, $NR^{11}R^{33}$, (alkylene)-$NR^{18}R^{19}$,

$C(R^{12})$=N-R^{18}, CH=N-OR^{13}, $C(R^{12})$=N-OR^{20},
$C(R^{11})$=N-$NR^{11}C(=O)NR^{14A}R^{14B}$, $C(R^{11})$=N-$NR^{11}SO_2R^{18}$,
$OC(=O)NR^{11}$(alkylene)-R^{26}, $OC(=O)[N(CH_2CH_2)_2N]$-R^{21},
$NR^{11}C(=O)OR^{23}$, $NR^{11}C(=O)$S-R^{18}, $NR^{11}C(=O)NR^{11}R^{23}$,
$NR^{11}C(=S)NR^{11}R^{23}$, $NR^{11}S(=O)_2N(R^{15})_2$, $NR^{11}C(=O)NR^{11}$(alkylene)-R^{24},
$NR^{11}C(=O)N(R^{11})NR^{16A}R^{16B}$, substituted alkyl, wherein one of the substituents is a spirocycloalkyl group, optionally substituted $(alkylene)_x$-cycloalkyl, and optionally substituted -$(alkylene)_x$-heterocycloalkyl, wherein the heterocycloalkyl does not include unsubstituted N-morpholinyl, N-piperidyl, or N-thiomorpholinyl,

wherein any said alkylene group may be optionally substituted with one to three R^{10} groups;

provided that when R^3, R^4, R^5, or R^6 is $C(=O)R^{13}$, then R^{13} does not include a heterocycloalkyl group that contains a nitrogen bonded to the carbonyl moiety; and

the other R^3, R^4, R^5, or R^6 moieties can be selected independently from H, halogen, R^{10}, OR^{20}, optionally substituted alkyl, optionally substituted alkenyl, and optionally substituted alkynyl, wherein said optional substituents are one to three R^{10} groups;

Q is selected from an optionally substituted $C_{1\text{-}2}$ alkylene, wherein said optional substituents are one to three R^{10} groups, CR^{20}=CR^{20}, O, $S(O)_y$, $C(R^7)$=N, N=$C(R^7)$, CH_2-Z', and Z'-CH_2; wherein

Z' is selected from O, S, C=O, and $C(=NOR^{11})$;

R^7 is selected from H, optionally substituted alkyl, and OR^{11}, wherein said optional substituents are one to three R^{10} groups;

R^{10} is selected from alkyl, aryl, heteroaryl, cycloalkyl, spirocycloalkyl, heterocycloalkyl, arylalkoxy, F, Cl, Br, I, CN, CF_3, $NR^{31A}R^{31B}$, NO_2, OR^{30}, OCF_3, =O, $=NR^{30}$, =N-OR^{30}, =N-$NR^{31A}R^{31B}$, $OC(=O)R^{30}$, $OC(=O)NHR^{29}$, O-$Si(R^{29})_4$, O-tetrahydropyranyl, ethylene oxide, $NR^{29}C(=O)R^{30}$, $NR^{29}CO_2R^{30}$, $NR^{29}C(=O)NR^{31A}R^{31B}$, $NHC(=NH)NH_2$, $NR^{29}S(O)_2R^{30}$, $S(O)_yR^{18}$, CO_2R^{30}, $C(=O)NR^{31A}R^{31B}$, $C(=O)R^{30}$, $(CH_2)_pOR^{30}$,

$CH=NNR^{31A}R^{31B}$, $CH=NOR^{30}$, $CH=NR^{30}$, $CH=NNHCH(N=NH)NH_2$, $S(=O)_2NR^{31A}R^{31B}$, $P(=O)(OR^{30})_2$, OR^{28}, and a monosaccharide wherein each hydroxyl group of the monosaccharide is independently either unsubstituted or is replaced by H, alkyl, alkylcarbonyloxy, or alkoxy;

R^{11} is selected from H and optionally substituted alkyl, wherein said optional substituents are one to three R^{10} groups;

R^{12} is selected from optionally substituted alkyl, optionally substituted aryl, and optionally substituted heteroaryl, wherein said optional substituents are one to three R^{10} groups;

R^{13} is independently selected from optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl, wherein said optional substituents are one to three R^{10} groups;

R^{14A} and R^{14B} are each independently selected from H, optionally substituted alkyl, optionally substituted aryl, and optionally substituted heteroaryl, wherein said optional substituents are one to three R^{10} groups;

R^{15} is independently selected from optionally substituted alkyl, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl, wherein said optional substituents are one to three R^{10} groups;

R^{16A} and R^{16B} are each independently selected from H and an optionally substituted alkyl, or together with the nitrogen to which they are attached form an optionally substituted heterocycloalkyl, wherein said optional substituents are one to three R^{10} groups;

R^{17} selected from optionally substituted cycloalkyl, optionally substituted heterocycloalkyl, and optionally substituted heteroaryl, wherein said optional substituents are one to three R^{10} groups;

R^{18} is selected from H, optionally substituted alkyl, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl, wherein said optional substituents are one to three R^{10} groups;

R^{19} is selected from CN and triazole;

R^{20} is selected from H, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted aryl, optionally substituted arylalkyl, optionally substituted heteroaryl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl, wherein said optional substituents are one to three R^{10} groups;

R^{21} is selected from optionally substituted aryl, and optionally substituted heteroaryl, wherein said optional substituents are one to three R^{10} groups;

R^{22} is optionally substituted C_5-C_{10} alkyl, wherein said optional substituents are one to three R^{10} groups;

R^{23} is selected from optionally substituted aryl, optionally substituted heteroaryl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl, wherein said optional substituents are one to three R^{10} groups;

R^{24} is selected from optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted cycloalkyl, optionally substituted heterocycloalkyl, OR^{20}, $O(CH_2)_pOR^{20}$, $(CH_2)_pOR^{20}$, SR^{17}, SOR^{15}, SO_2R^{18}, CN, $N(R^{18})_2$, $C(=O)N(R^{18})_2$, $NR^{18}C(=O)R^{18}$, $NR^{18}C(=O)N(R^{18})_2$, $C(=NR^{18})OR^{18}$, $C(R^{12})=NOR^{18}$, $NHOR^{20}$, $NR^{18}C(=NR^{18})N(R^{18})_2$, NHCN, $CONR^{18}OR^{18}$, CO_2R^{18}, $OCOR^{15}$, $OC(=O)N(R^{18})_2$, $NR^{18}C(=O)OR^{15}$, and $C(=O)R^{18}$, wherein said optional substituents are one to three R^{10} groups;

R^{25} is selected from optionally substituted cycloalkyl, optionally substituted heterocycloalkyl, OR^{20}, $O(CH_2)_pOR^{20}$, $(CH_2)_pOR^{20}$, SR^{17}, SOR^{15}, SO_2R^{18}, CN, $N(R^{17})_2$, $C(=O)N(R^{18})_2$, $NR^{18}C(=O)R^{18}$, $NR^{18}C(=O)N(R^{18})_2$, $C(=NR^{18})OR^{18}$, $C(R^{12})=NOR^{18}$, $NHOR^{20}$, $NR^{18}C(=NR^{18})N(R^{18})_2$, NHCN, $CONR^{18}OR^{18}$, CO_2R^{18}, $OCOR^{15}$, $OC(=O)N(R^{18})_2$, $NR^{18}C(=O)OR^{15}$, and $C(=O)R^{18}$, wherein said optional substituents are one to three R^{10} groups;

R^{26} is selected from optionally substituted cycloalkyl, optionally substituted heterocycloalkyl, optionally substituted aryl, optionally substituted heteroaryl, OR^{11}, $O(CH_2)_pOR^{20}$, $(CH_2)_pOR^{20}$, SR^{17}, SOR^{15}, SO_2R^{18}, CN, $N(R^{18})_2$, $C(=O)N(R^{18})_2$, $NR^{18}C(=O)R^{18}$, $NR^{18}C(=O)N(R^{18})_2$,

$C(=NR^{18})OR^{18}$, $C(R^{12})=NOR^{18}$, $NHOR^{20}$, $NR^{18}C(=NR^{18})N(R^{18})_2$, NHCN, $CONR^{18}OR^{18}$, CO_2R^{18}, $OCOR^{15}$, $OC(=O)N(R^{18})_2$, $NR^{18}C(=O)OR^{15}$, and $C(=O)R^{18}$, wherein said optional substituents are one to three R^{10} groups;

R^{27} is selected from optionally substituted cycloalkyl, CN, $C(R^{12})=NOR^{18}$, and $C(=O)N(R^{18})_2$, wherein said optional substituents are one to three R^{10} groups;

R^{28} is the residue of an amino acid after the removal of the hydroxyl moeity from the carboxyl group thereof;

R^{29} is H or alkyl;

R^{30} is H, alkyl, aryl, arylalkyl, heteroaryl, cycloalkyl, or heterocycloalkyl;

R^{31A} and R^{31B} are each independently selected from H, alkyl, and arylalkyl, or together with the nitrogen to which they are attached form a heterocycloalkyl;

R^{32} is optionally substituted aryl, wherein said optional substituents are one to three R^{10} groups;

R^{33} is optionally substituted cycloalkyl, optionally substituted heteroaryl, or optionally substituted heterocycloalkyl, wherein said optional substituents are one to three R^{10} groups;


p is independently selected from 1, 2, 3, and 4;

x is 0 or 1;

y is independently selected from 0, 1 and 2; or


a stereoisomeric or pharmaceutically acceptable salt form thereof.

In other embodiments, the compounds of Formula I as defined herein are not intended to include any compounds disclosed in PCT Publ. No. WO 02/28861. In particular, when R^1 is H; A^1,A^2 and B^1, B^2 are each =O or =S; ring A is a phenylene, ring B is




, wherein R^A is H or C_1-C_4 alkyl; and Q is CR^{20}=CR^{20}, wherein R^{20} is H, C_1-C_4 alkyl or C_1-C_4 alkoxy; then neither R^3 or R^4 includes (alkylene)-NR^aR^b, wherein R^a and R^b combine with the nitrogen to which they are attached to form a heterocycloalkyl group.

In another embodiment, the compounds of Formula I as defined herein are not intended to include any compounds disclosed in PCT Publ. No. WO 02/30942. In particular, when A^1,A^2 is =O; B^1, B^2 are independently H or OH, or B^1, B^2 combine to form =O; rings A and B are each a phenylene; Q is O, S, or CH_2; and R^2 is H, or optionally substituted



; then any of R^3, R^4, R^5, and R^6 cannot include (alkylene)$_x$$OR^{13}$, wherein R^{13} is cycloalkyl; $(CH_2)_pOR^{22}$, wherein R^{22} is C_5-C_7 alkyl; O-(alkylene)-R^{27}, wherein R^{27} is CN; $OCH[(CH_2)_pOR^{20}]_2$; $NR^{11}R^{33}$, wherein R^{33} is cycloalkyl; or NR^aR^b, wherein R^a and R^b combine with the nitrogen to which they are attached to form a heterocycloalkyl group.


In a further embodiment, the compounds of Formula I as defined herein are not intended to include any compounds disclosed in PCT Publ. No. WO 02/28874. In particular, when A^1,A^2 and B^1, B^2 are each =O; rings A and B are each a phenylene; Q is O or S; and R^2 is optionally substituted




; then any of R^3, R^4, R^5, and R^6 cannot include (alkylene)$_x$$OR^{13}$, wherein R^{13} is cycloalkyl; $(CH_2)_pOR^{22}$, wherein R^{22} is C_5-C_7 alkyl; O-(alkylene)-R^{27}, wherein R^{27} is CN;

$OCH[(CH_2)_pOR^{20}]_2$; $NR^{11}R^{33}$, wherein R^{33} is cycloalkyl; or NR^aR^b, wherein R^a and R^b combine with the nitrogen to which they are attached to form a heterocycloalkyl group.

In an additional embodiment, the compounds of Formula I as defined herein are not intended to include any compounds disclosed in PCT Publ. No. WO 98/07433. In particular, when A^1,A^2 is =O; B^1, B^2 are independently H or OH, or B^1, B^2 combine to form =O; rings A and B are each a phenylene; Q is O, S, or CH-R^a; and one of R^2 or R^a is H and the other is optionally substituted




or



,

wherein W is optionally substituted C_1 alkyl, or $NR^{31A}R^{31B}$; then any of R^3, R^4, R^5, and R^6 cannot include OR^{13} or $C(=O)NR^cR^d$ wherein R^c and R^d combine with the nitrogen to which they are attached to form a heterocycloalkyl group.

In a further embodiment, preferred compounds of Formula I as defined herein exclude certain compounds in which any of R^3, R^4, R^5, and R^6 include $(CH_2)_pOR^{22}$. In particular, in this embodiment, when R^1 is H; A^1,A^2 are both H; B^1, B^2 taken together are =O; rings A and B are both a phenylene, wherein ring B is unsubstituted; R^2 is $CH_2CH_2CH_2OH$; and Q is CH_2, then R^3 and R^4 cannot include $(CH_2)_pOR^{22}$.

Other aspects of the present invention include the compounds of Formula I as defined herein wherein rings A and B are a phenylene; or one of rings A and B are a phenylene, and the other ring A or ring B, together with the carbon atoms to which they are attached, are selected from a 5-membered aromatic ring in which one or two carbon atoms may be replaced with a nitrogen atom, preferably a pyrazolylene, and more preferably




or


. Further aspects include those compounds wherein R^1 is selected from H, substituted alkyl, and unsubstituted alkyl. Another aspect includes those compounds wherein R^2 is H, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, or optionally substituted cycloalkyl, and preferably

where R^2 is H or optionally substituted alkyl. Additional aspects include those compounds wherein groups A^1, A^2 and B^1, B^2 are each independently selected from H, H; and a group wherein A^1, A^2 or B^1, B^2 together form =O, and preferably those wherein A^1A^2 are H, H; and B^1B^2 together form =O. In yet another aspect, the invention includes compounds wherein Q is selected from an optionally substituted C_{1-2} alkylene, or preferably Q is CH_2 or CH_2CH_2. Additional aspects of the present invention include any combination of the above preferred substituents, such as, for example, a compound of Formula I with the preferred moieties of groups R^1 and R^2; or R^1 and Q; or R^1, R^2; or Q; etc.

In another embodiment of the present invention, there are included compounds having a structure of Formula II:





II

wherein ring B together with the carbon atoms to which it is attached, is selected from:


(a) a phenylene ring in which from 1 to 3 carbon atoms may be replaced by nitrogen atoms; and

(b) a 5-membered aromatic ring in which from 1 to 2 carbon atoms may be replaced by nitrogen atoms.

In one aspect, there are included compounds of Formula II wherein the B ring is a phenylene, or ring B is a pyrazolylene, or preferably



 or . Another aspect includes those compounds wherein R^1 is selected from H, substituted alkyl, and unsubstituted alkyl. Further aspects include those compounds wherein R^2 is H, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, or optionally substituted cycloalkyl, and preferably where R^2 is H or optionally substituted alkyl. Additional aspects include those compounds wherein groups A^1, A^2 and B^1, B^2 are each independently selected from H, H; and a group wherein A^1, A^2 or B^1, B^2 together form =O, and preferably those wherein A^1A^2 are H, H; and B^1B^2 together form =O. In yet another aspect, the invention includes compounds wherein Q is selected from an optionally substituted $C_{1\text{-}2}$ alkylene, or preferably Q is CH_2 or CH_2CH_2. Additional aspects of the present invention include any combination of the above preferred substituents, such as, for example, a compound of Formula II with the preferred moieties of groups R^1 and R^2; or R^1 and Q; or R^1, R^2; or Q; etc.

In yet another embodiment of the present invention, there are included compounds having a structure of Formula III:




III ,

where preferrably ring B is a phenylene, or ring B is a pyrazolylene, preferably


 or , and where R^1 is selected from H and optionally substituted alkyl;

and Formula IV:




IV ;

and Formula V:



V ;

and Formula VI:



VI

In certain aspects of the present invention, there are included compounds of Formulas III-VI wherein R^2 is H, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, or optionally substituted cycloalkyl, or preferably where R^2 is H or optionally substituted alkyl. Other aspects include those compounds wherein Q is selected from an optionally substituted C_{1-2} alkylene, or preferably Q is CH_2 or CH_2CH_2. Additional aspects of the present invention include any combination of the above preferred substituents for each of Formulas III-VI.

The following paragraphs show additional aspects of the invention for at least one R^3, R^4, R^5, and R^6 for compounds of Formulas I-VI and their respective preferred embodiments described heretofore.

1. OR^{13}; especially those where R^{13} is optionally substituted cycloalkyl, and particularly those where the cycloalkyl is a 5 or 6 membered ring.
2. $C(=O)R^{13}$; especially those where R^{13} is optionally substituted cycloalkyl, and particularly those where the cycloalkyl is a 5 or 6 membered ring.
3. (alkylene)OR^{13}; especially those where R^{13} is optionally substituted cycloalkyl, and particularly those where the cycloalkyl is a 5 or 6 membered ring.
4. $(CH_2)_pOR^{22}$; especially those where R^{22} is a branched chain alkyl.
5. O-(alkylene)-R^{27}.
6. $OCH(CO_2R^{18})_2$; especially those where R^{18} is optionally substituted alkyl.
7. $OCH[(CH_2)_pOR^{20}]_2$; especially those where R^{20} is optionally substituted alkyl.
8. C(=O)-(alkylene)-R^{25}.

9. $NR^{11}R^{33}$; especially those where R^{33} is optionally substituted heteroaryl.

10. (alkylene)-$NR^{18}R^{19}$; especially those where R^{18} is H or optionally substituted alkyl.

11. $C(R^{12})$=N-R^{18}; especially those where R^{12} is alkyl, and those where R^{18} is optionally substituted heterocycloalkyl.

12. CH=N-OR^{13}; especially those where R^{13} is optionally substituted heterocycloalkyl.

13. $C(R^{12})$=N-OR^{20}; especially those where R^{12} and R^{30} are optionally substituted alkyl.

14. $C(R^{11})$=N-$NR^{11}C(=O)NR^{14A}R^{14B}$.

15. $C(R^{11})$=N-$NR^{11}SO_2R^{18}$.

16. $OC(=O)NR^{11}$(alkylene)-R^{26}; especially those where R^{26} is optionally substituted aryl or optionally substituted heteroaryl.

17. $OC(=O)[N(CH_2CH_2)_2N]$-R^{21}; especially those where R^{21} is optionally substituted heteroaryl.

18. $NR^{11}C(=O)OR^{23}$; especially those where R^{23} is optionally substituted aryl.

19. $NR^{11}C(=O)S$-R^{18}.

20. $NR^{11}C(=O)NR^{11}R^{23}$; especially those where R^{23} is optionally substituted aryl or optionally substituted heteroaryl.

21. $NR^{11}C(=S)NR^{11}R^{23}$; especially those where R^{23} is optionally substituted aryl.

22. $NR^{11}S(=O)_2N(R^{15})_2$.

23. $NR^{11}C(=O)NR^{11}$(alkylene)-R^{24}; especially those where R^{24} is optionally substituted heterocycloalkyl, or optionally substituted heteroaryl.

24. $NR^{11}C(=O)N(R^{11})NR^{16A}R^{16B}$.

25. substituted alkyl, wherein one of the substituents is an optionally substituted spirocycloalkyl group.

26. optionally substituted $(alkylene)_x$-cycloalkyl, especially optionally substituted (C_1-alkylene)-cycloalkyl and optionally substituted cycloalkyl.

27. optionally substituted -$(alkylene)_x$-heterocycloalkyl, wherein the heterocycloalkyl does not include unsubstituted N-morpholinyl, N-

piperidyl, or N-thiomorpholinyl; especially optionally substituted (C_1-alkylene)-heterocycloalkyl, optionally substituted heterocycloalkyl, and more especially optionally substituted heterocycloalkyl groups with two heteroatoms, optionally substituted tetrahydrofuranyl and optionally substituted tetrahydropyranyl.

28. $NR^{11}R^{32}$, especially those where R^{32} is a phenyl group and wherein the phenyl group is optionally substituted with one or more alkoxy groups, and in particular, with one or more methoxy groups.

The preceding paragraphs may be combined to further define additional prefered embodiments of compounds of Formulas I-VI. For example, one such combination for R^3, R^4, R^5, or R^6 can include OR^{13}, $C(=O)R^{13}$, $(CH_2)_pOR^{22}$, $(CH_2)_pOR^{22}$, O-(alkylene)-R^{27}, $OCH(CO_2R^{18})_2$, $OCH[(CH_2)_pOR^{20}]_2$, and $C(=O)$-(alkylene)-R^{25}.

Another such combination includes $NR^{11}R^{32}$, $NR^{11}R^{33}$, (alkylene)-$NR^{18}R^{19}$, $C(R^{12})$=N-R^{18}, CH=N-OR^{13}, $C(R^{12})$=N-OR^{20}, $C(R^{11})$=N-$NR^{11}C(=O)NR^{14A}R^{14B}$, $C(R^{11})$=N-$NR^{11}SO_2R^{18}$, $OC(=O)NR^{11}$(alkylene)-R^{26}, $OC(=O)[N(CH_2CH_2)_2N]$-R^{21}, $NR^{11}C(=O)OR^{23}$, $NR^{11}C(=O)$S-R^{18}, $NR^{11}C(=O)NR^{11}R^{23}$, $NR^{11}C(=S)NR^{11}R^{23}$, $NR^{11}S(=O)_2N(R^{15})_2$, $NR^{11}C(=O)NR^{11}$(alkylene)-R^{24}, and $NR^{11}C(=O)N(R^{11})NR^{16A}R^{16B}$.

A third such combination includes OR^{13}, $NR^{11}R^{32}$, $NR^{11}R^{33}$, (alkylene)-$NR^{18}R^{19}$, $C(R^{12})$=N-R^{18}, CH=N-OR^{13}, $C(R^{12})$=N-OR^{20}, $OC(=O)NR^{11}$(alkylene)-R^{26}, $NR^{11}C(=O)NR^{11}R^{23}$, $NR^{11}C(=S)NR^{11}R^{23}$, $NR^{11}C(=O)NR^{11}$(alkylene)-R^{24}, $NR^{11}C(=O)N(R^{11})NR^{16A}R^{16B}$, C(=O)-cycloalkyl, and optionally substituted -$(\text{alkylene})_x$-heterocycloalkyl.

A fourth such combination includes OR^{13}, $C(=O)R^{13}$, $(CH_2)_pOR^{22}$, $OCH(CO_2R^{18})_2$, $OCH[(CH_2)_pOR^{20}]_2$, $NR^{11}R^{32}$, $NR^{11}R^{33}$, (alkylene)-$NR^{18}R^{19}$, CH=N-OR^{13}, $C(R^{12})$=N-OR^{20}, $OC(=O)NR^{11}$(alkylene)-R^{26}, $NR^{11}C(=O)NR^{11}R^{23}$, $NR^{11}C(=O)NR^{11}$(alkylene)-R^{24}, substituted alkyl, wherein the alkyl is substituted with at least a spiroalkyl group, optionally substituted -$(\text{alkylene})_x$-cycloalkyl, and optionally substituted -$(\text{alkylene})_x$-heterocycloalkyl.

A fifth such combination includes $NR^{11}R^{32}$, $NR^{11}R^{33}$, (alkylene)-$NR^{18}R^{19}$, $C(R^{12})$=N-R^{18}, CH=N-OR^{13}, $C(R^{12})$=N-OR^{20}, $C(R^{11})$=N-$NR^{11}C(=O)NR^{14A}R^{14B}$, $C(R^{11})$=N-$NR^{11}SO_2R^{18}$, $OC(=O)NR^{11}$(alkylene)-R^{26}, $OC(=O)[N(CH_2CH_2)_2N]$-R^{21},

$NR^{11}C(=O)OR^{23}$, $NR^{11}C(=O)S\text{-}R^{18}$, $NR^{11}C(=O)NR^{11}R^{23}$, $NR^{11}C(=S)NR^{11}R^{23}$, $NR^{11}S(=O)_2N(R^{15})_2$, $NR^{11}C(=O)NR^{11}(\text{alkylene})\text{-}R^{24}$, $NR^{11}C(=O)N(R^{11})NR^{16A}R^{16B}$, substituted alkyl, wherein the alkyl is substituted with at least a spiroalkyl group, optionally substituted -(alkylene)$_x$-cycloalkyl, and optionally substituted -(alkylene)$_x$-heterocycloalkyl.

A sixth such combination includes $NR^{11}R^{33}$, and $C(R^{12})=N\text{-}OR^{20}$.

A seventh such combination includes $NR^{11}R^{32}$, optionally substituted -(alkylene)$_x$-cycloalkyl, and optionally substituted -(alkylene)$_x$-heterocycloalkyl.

Other embodiments of the present invention include compounds of formulas I-VI wherein at least one of R^3, R^4, R^5, or R^6 is $NR^{11}R^{32}$ wherein R^{32} is phenyl optionally substituted with one to three OR^{30} groups, particularly those wherein R^{30} is C_1-C_8 alkyl. Further embodiments include those wherein R^{32} is phenyl substituted with one to three methoxy groups, or those wherein R^{32} is phenyl substituted with two methoxy groups.

In other aspects of the present invention, there are included compounds of Formulas I-VI wherein at least one R^3, R^4, R^5, and R^6 is $NR^{11}R^{33}$, wherein R^{33} is optionally substituted heteroaryl, or those wherein R^{33} is optionally substituted cycloalkyl, or those wherein R^{33} is optionally substituted heterocycloalkyl. Other embodiments include those wherein R^{33} is 5-6 membered heteroaryl ring with one to three nitrogen atoms in the ring. Further embodiments include compounds wherein R^{33} is a 6 membered heteroaryl ring with two nitrogen atoms in the ring. Additional embodiments include those wherein R^{33} is pyrimidinyl, or those wherein R^{33} is pyridazinyl, or those wherein R^{33} is pyridyl.

The following terms and expressions used herein have the indicated meanings.

In the formulas described and claimed herein, it is intended that when any symbol appears more than once in a particular formula or substituent, its meaning in each instance is independent of the other.

As used herein, the term "about" refers to a range of values from ± 10% of a specified value. For example, the phrase "about 50 mg" includes ± 10% of 50, or from 45 to 55 mg.

As used herein, a range of values in the form "x-y" or "x to y", or "x through y", include integers x, y, and the integers therebetween. For example, the phrases "1-6", or "1

to 6" or "1 through 6" are intended to include the integers 1, 2, 3, 4, 5, and 6. Preferred embodiments include each individual integer in the range, as well as any subcombination of integers. For example, preferred integers for "1-6" can include 1, 2, 3, 4, 5, 6, 1-2, 1-3, 1-4, 1-5, 2-3, 2-4, 2-5, 2-6, etc.

As used herein "stable compound" or "stable structure" refers to a compound that is sufficiently robust to survive isolation to a useful degree of purity from a reaction mixture, and preferably capable of formulation into an efficacious therapeutic agent. The present invention is directed only to stable compounds.


As used herein, the term "alkyl" refers to a straight-chain, or branched alkyl group having 1 to 8 carbon atoms, such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, isoamyl, neopentyl, 1-ethylpropyl, 3-methylpentyl, 2,2-dimethylbutyl, 2,3-dimethylbutyl, hexyl, octyl, etc. The alkyl moiety of alkyl-containing groups, such as alkoxy, alkoxycarbonyl, and alkylaminocarbonyl groups, has the same meaning as alkyl defined above. Lower alkyl groups, which are preferred, are alkyl groups as defined above which contain 1 to 4 carbons. A designation such as "C_1-C_4 alkyl" refers to an alkyl radical containing from 1 to 4 carbon atoms.

As used herein, the term "alkenyl" refers to a straight chain, or branched hydrocarbon chains of 2 to 8 carbon atoms having at least one carbon-carbon double bond. A designation "C_2-C_8 alkenyl" refers to an alkenyl radical containing from 2 to 8 carbon atoms. Examples of alkenyl groups include ethenyl, propenyl, isopropenyl, 2,4-pentadienyl, etc.

As used herein, the term "alkynyl" refers to a straight chain, or branched hydrocarbon chains of 2 to 8 carbon atoms having at least one carbon-carbon triple bond. A designation "C_2-C_8 alkynyl" refers to an alkynyl radical containing from 2 to 8 carbon atoms. Examples include ethynyl, propynyl, isopropynyl, 3,5-hexadiynyl, etc.

As used herein, the term "alkylene" refers to a branched or straight chained hydrocarbon of 1 to 8 carbon atoms, which is formed by the removal of two hydrogen atoms. A designation such as "C_1-C_4 alkylene" refers to an alkylene radical containing from 1 to 4 carbon atoms. Examples include methylene (-CH_2-), propylidene (CH_3CH_2CH=), 1,2-ethandiyl (-CH_2CH_2-), etc.


As used herein, the term "phenylene" refers to a phenyl group with an additional hydrogen atom removed, ie. a moiety with the structure of:



As used herein, the terms "carbocycle", "carbocyclic" or "carbocyclyl" refer to a stable, saturated or partially saturated, monocyclic or bicyclic hydrocarbon ring system which is saturated, partially saturated or unsaturated, and contains from 3 to 10 ring carbon atoms. Accordingly the carbocyclic group may be aromatic or non-aromatic, and includes the cycloalkyl and aryl groups defined herein. The bonds connecting the endocyclic carbon atoms of a carbocyclic group may be single, double, triple, or part of a fused aromatic moiety.

As used herein, the term "cycloalkyl" refers to a saturated or partially saturated mono- or bicyclic alkyl ring system containing 3 to 10 carbon atoms. A designation such as "C_5-C_7 cycloalkyl" refers to a cycloalkyl radical containing from 5 to 7 ring carbon atoms. Preferred cycloalkyl groups include those containing 5 or 6 ring carbon atoms. Examples of cycloalkyl groups include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclohexenyl, cycloheptyl, cyclooctyl, etc.

As used herein, the term "spirocycloalkyl" refers to a cycloalkyl group bonded to a carbon chain or carbon ring moiety by a carbon atom common to the cycloalkyl group and the carbon chain or carbon ring moiety. For example, a C_3 alkyl group substituted with an R group wherein the R group is spirocycloalkyl containing 5 carbon atoms refers to:



As used herein, the term "aryl" refers to a mono- or bicyclic hydrocarbon aromatic ring system having 6 to 12 ring carbon atoms. Examples include phenyl and naphthyl. Preferred aryl groups include phenyl or naphthyl groups. Included within the definition of "aryl" are fused ring systems, including, for example, ring systems in which an aromatic ring is fused to a cycloalkyl ring. Examples of such fused ring systems include, for example, indane and indene.

As used herein, the terms "heterocycle", "heterocyclic" or "heterocyclyl" refer to a mono- di-, tri- or other multicyclic aliphatic ring system that includes at least one heteroatom such as O, N, or S. The nitrogen and sulfur heteroatoms may be optionally oxidized, and the nitrogen may be optionally substituted in non-aromatic rings. Heterocycles are intended to include heteroaryl and heterocycloalkyl groups.

Some heterocyclic groups containing one or more nitrogen atoms include pyrrolidine, pyrroline, pyrazoline, piperidine, morpholine, thiomorpholine, N-methylpiperazine, indole, isoindole, imidazole, imidazoline, oxazoline, oxazole, triazole, thiazoline, thiazole, isothiazole, thiadiazole, triazine, isoxazole, oxindole, pyrazole, pyrazolone, pyrimidine, pyrazine, quinoline, iosquinoline, and tetrazole groups. Some heterocyclic groups formed containing one or more oxygen atoms include furan, tetrahydrofuran, pyran, benzofurans, isobenzofurans, and tetrahydropyran groups. Some heterocyclic groups containing one or more sulfur atoms include thiophene, thianaphthene, tetrahydrothiophene, tetrahydrothiapyran, and benzothiophenes.

As used herein, the term "heterocycloalkyl" refers to a cycloalkyl group in which one or more ring carbon atoms are replaced by at least one hetero atom such as -O-, -N-, or -S-, and includes ring systems which contain a saturated ring group bridged or fused to one or more aromatic groups. Some heterocycloalkyl groups containing both saturated and aromatic rings include phthalamide, phthalic anhydride, indoline, isoindoline, tetrahydroisoquinoline, chroman, isochroman, and chromene.

As used herein, the term "heteroaryl" refers to an aryl group containing 5 to 10 ring carbon atoms in which one or more ring carbon atoms are replaced by at least one hetero atom such as -O-, -N-, or -S-. Some heteroaryl groups of the present invention include pyridyl, pyrimidyl, pyrrolyl, furanyl, thienyl, imidazolyl, triazolyl, tetrazolyl, quinolyl, isoquinolyl, benzoimidazolyl, thiazolyl, pyrazolyl, and benzothiazolyl groups.

As used herein, the term "arylalkyl" refers to an alkyl group that is substituted with an aryl group. Examples of arylalkyl groups include, but are not limited to, benzyl, phenethyl, benzhydryl, diphenylmethyl, triphenylmethyl, diphenylethyl, naphthylmethyl, etc.

As used herein, the term "heteroarylalkyl" refers to an alkyl group that is substituted with a heteroaryl group.

As used herein, the term "alkoxy" refers to an oxygen radical substituted with an alkyl group. Examples include methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, isobutoxy, sec-butoxy, t-butoxy, etc.

As used herein, the term "arylalkoxy" refers to an aryl-substituted alkoxy group, such as benzyloxy, diphenylmethoxy, triphenylmethoxy, phenylethoxy, diphenylethoxy, etc.

As used herein, the term "alkylcarbonyloxy" refers to an RC(=O)O- group, wherein R is an alkyl group.

As used herein, the term "monosaccharide" refers to a simple sugar of the formula $(CH_2O)_n$. The monosaccharides can be straight-chain or ring systems, and can include a saccharose unit of the formula -CH(OH)-C(=O)-. Examples of monosaccharides include erythrose, threose, ribose, arabinose, xylose, lyxose, allose, altrose, glucose, mannose, gulose, idose, galactose, talose, erythulose, ribulose, xyulose, psicose, fructose, sorbose, tagatose, erythropentulose, threopentulose, glycerotetrulose, glucopyranose, fructofuranose, etc.

As used herein, the term "amino acid" refers to a group containing both an amino group and a carboxyl group. Embodiments of amino acids include α-amino, β-amino, γ-amino acids. The α-amino acids have a general formula HOOC-CH(side chain)-NH_2. The amino acids can be in their D, L or racemic configurations. Amino acids include naturally-occurring and non-naturally occurring moieties. The naturally-occurring amino acids include the standard 20 α-amino acids found in proteins, such as glycine, serine, tyrosine, proline, histidine, glutamine, etc. Naturally-occurring amino acids can also include non-α-amino acids (such as β-alanine, γ-aminobutyric acid, homocysteine, etc.), rare amino acids (such as 4-hydroxyproline, 5-hydroxylysine, 3-methylhistidine, etc.) and non-protein amino acids (such as citrulline, ornithine, canavanine, etc.). Non-naturally occurring amino acids are well-known in the art, and include analogs of natural amino acids. See Lehninger, A. L. *Biochemistry*, 2nd ed.; Worth Publishers: New York, 1975; 71-77, the disclosure of which is incorporated herein by reference. Non-naturally occurring amino acids also include α-amino acids wherein the side chains are replaced with synthetic derivatives. In certain embodiments, substituent groups for the compounds of the present invention include the residue of an amino acid after removal of the hydroxyl

moiety of the carboxyl group thereof; i.e., groups of formula -C(=O)CH(side chain)-NH_2. Representative side chains of naturally occurring and non-naturally occurring α-amino acids include are shown below in Table A.

Table A


H


CH_3-


HO-CH_2-


C_6H_5-CH_2-


HO-C_6H_4-CH_2-











HS-CH_2-

HO_2C-CH(NH_2)-CH_2-S-S-CH_2-

CH_3-CH_2-

CH_3-S-CH_2-CH_2-

CH_3-CH_2-S-CH_2-CH_2-

HO-CH_2-CH_2-

C_5H_9-

C_6H_{11}-

C_6H_{11}-CH_2-

CH_3-CH(OH)-

HO_2C-CH_2-NHC(=O)-CH_2-

HO_2C-CH_2-

HO_2C-CH_2-CH_2-

NH_2C(=O)-CH_2-

NH_2C(=O)-CH_2-CH_2-

$(CH_3)_2$-CH-

$(CH_3)_2$-CH-CH_2-

CH_3-CH_2-CH_2-

H_2N-CH_2-CH_2-CH_2-

H_2N-C(=NH)-NH-CH_2-CH_2-CH_2-

H_2N-C(=O)-NH-CH_2-CH_2-CH_2-

CH_3-CH_2-CH(CH_3)-

CH_3-CH_2-CH_2-CH_2-

H_2N-CH_2-CH_2-CH_2-CH_2-

As used herein, the term "*trk*" refers to the family of high affinity neurotrophin receptors presently comprising trk A, trk B, and trk C, and other membrane associated proteins to which a neurotrophin can bind.

As used herein, the term "VEGFR" refers to the family of high affinity vascular endothelial growth factor receptors presently comprising VEGFR1, VEGFR2, VEGFR3, and other membrane associated proteins to which a VEGF can bind.

As used herein, the term "MLK" refers to the family of high affinity mixed lineage kinases presently comprising MLK1, MLK2, MLK3, MLK4α & β, DLK, LZK, ZAK α & β , and other serine/threonine kinases classified within this family.

As used herein, the terms "enhance" or "enhancing" when used to modify the terms "function" or "survival" means that the presence of a compound of the present invention has a positive effect on the function and/or survival of a trophic factor responsive cell compared with a cell in the absence of the compound. For example, and not by way of limitation, with respect to the survival of, e.g., a cholinergic neuron, a compound of the present invention would evidence enhancement of survival of a cholinergic neuronal population at risk of dying (due to, e.g., injury, a disease condition, a degenerative condition or natural progression) when compared to a cholinergic neuronal population not presented with such a compound, if the treated population has a comparatively greater period of functionality than the non-treated population. As a further example, and again not by way of limitation, with respect to the function of, e.g., a sensory neuron, a compound of the present invention would evidence enhancement of the function (e.g. neurite extension) of a sensory neuronal population when compared to a sensory neuronal population not presented with such a compound, if the neurite extension of the treated population is comparatively greater than the neurite extension of the non-treated population.

As used herein, the terms "inhibit" or "inhibition" refer to a specified response of a designated material (e.g., enzymatic activity) is comparatively decreased in the presence of a compound of the present invention.

As used herein, the terms "cancer" or "cancerous" refer to any malignant proliferation of cells in a mammal. Examples include prostate, benign prostate hyperplasia, ovarian, breast, brain, lung, pancreatic, colorectal, gastric, stomach, solid

tumors, head and neck, neuroblastoma, renal cell carcinoma, lymphoma, leukemia, other recognized malignancies of the hematopoietic systems, and other recognized cancers.

As used herein the terms "neuron", "cell of neuronal lineage" and "neuronal cell" refer to a heterogeneous population of neuronal types having singular or multiple transmitters and/or singular or multiple functions; preferably, these are cholinergic and sensory neurons. As used herein, the phrase "cholinergic neuron" means neurons of the Central Nervous System (CNS) and Peripheral Nervous System (PNS) whose neurotransmitter is acetylcholine; exemplary are basal forebrain and spinal cord neurons. As used herein, the phrase "sensory neuron" includes neurons responsive to environmental cues (e.g., temperature, movement) from, e.g., skin, muscle and joints; exemplary is a neuron from the DRG.

As used herein the term "trophic factor" refers to a molecule that directly or indirectly affects the survival or function of a trophic factor responsive cell. Exemplary trophic factors include Ciliary Neurotrophic Factor (CNTF), basic Fibroblast Growth Factor (bFGF), insulin and insulin-like growth factors (e.g., IGF-I, IGF-II, IGF-III), interferons, interleukins, cytokines, and the neurotrophins, including Nerve Growth Factor (NGF), Neurotrophin-3 (NT-3), Neurotrophin-4/5 (NT-4/5) and Brain Derived Neurotrophic Factor (BDNF).

As used herein the term "trophic factor-responsive cell" refers to a cell which includes a receptor to which a trophic factor can specifically bind; examples include neurons (e.g., cholinergic and sensory neurons) and non-neuronal cells (e.g., monocytes and neoplastic cells).

As used herein the terms "trophic factor activity" and "trophic factor-induced activity", refer to both endogenous and exogenous trophic factors, where "endogenous" refers to a trophic factor normally present and "exogenous" refers to a trophic factor added to a system. As defined, "trophic factor induced activity" includes activity induced by (1) endogenous trophic factors; (2) exogenous trophic factors; and (3) a combination of endogenous and exogenous trophic factors.

As used herein, the term "at risk of dying" in conjunction with a biological material, e.g., a cell such as a neuron, refers to a state or condition which negatively impacts the biological material such that the material has an increased likelihood of dying

due to such state or condition. For example, compounds disclosed herein can "rescue" or enhance the survival of motoneurons which are naturally at risk of dying in an *in ovo* model of programmed cell death. Similarly, for example, a neuron may be at risk of dying due to the natural aging process which occasions the death of a neuron, or due to an injury, such as a trauma to the head, which may be such that neurons and/or glia, for example, impacted by such trauma may be at risk of dying. Further, for example, a neuron may be at risk of dying due to a disease state or condition, as in the case of neurons at risk of dying as occasioned by the disease ALS. Thus, by enhancing the survival of a cell at risk of dying by use of a compound of the claimed invention is meant that such compound decreases or prevents the risk of the death of the cell.

As used herein the term "contacting" refers to directly or indirectly causing placement together of moieties, such that the moieties directly or indirectly come into physical association with each other, whereby a desired outcome is achieved. Thus, as used herein, one can "contact" a target cell with a compound as disclosed herein even though the compound and cell do not necessarily physically join together (as, for example, is the case where a ligand and a receptor physically join together), as long as the desired outcome is achieved (e.g., enhancement of the survival of the cell). Contacting thus includes acts such as placing moieties together in a container (e.g., adding a compound as disclosed herein to a container comprising cells for *in vitro* studies) as well as administration of the compound to a target entity (e.g., injecting a compound as disclosed herein into a laboratory animal for *in vivo* testing, or into a human for therapy or treatment purposes).

As used herein, a "therapeutically effective amount" refers to an amount of a compound of the present invention effective to prevent or treat the symptoms of a particular disorder.

As used herein, the term "subject" refers to a warm blooded animal such as a mammal, preferably a human, or a human child, which is afflicted with, or has the potential to be afflicted with one or more diseases and conditions described herein.

As used herein, the term "pharmaceutically acceptable" refers to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for contact with the tissues of human beings and animals

without excessive toxicity, irritation, allergic response, or other problem complications commensurate with a reasonable benefit/risk ratio.

As used herein, "pharmaceutically acceptable salts" refer to derivatives of the disclosed compounds wherein the parent compound is modified by making acid or base salts thereof. Examples of pharmaceutically acceptable salts include, but are not limited to, mineral or organic acid salts of basic residues such as amines; alkali or organic salts of acidic residues such as carboxylic acids; and the like. The pharmaceutically acceptable salts include the conventional non-toxic salts or the quaternary ammonium salts of the parent compound formed, for example, from non-toxic inorganic or organic acids. For example, such conventional non-toxic salts include those derived from inorganic acids such as hydrochloric, hydrobromic, sulfuric, sulfamic, phosphoric, nitric and the like; and the salts prepared from organic acids such as acetic, propionic, succinic, glycolic, stearic, lactic, malic, tartaric, citric, ascorbic, pamoic, maleic, hydroxymaleic, phenylacetic, glutamic, benzoic, salicylic, sulfanilic, 2-acetoxybenzoic, fumaric, toluenesulfonic, methanesulfonic, ethane disulfonic, oxalic, isethionic, and the like.

The pharmaceutically acceptable salts of the present invention can be prepared from the parent compound which contains a basic or acidic moiety by conventional chemical methods. Generally, such salts can be prepared by reacting the free acid or base forms of these compounds with a stoichiometric amount of the appropriate base or acid in water or in an organic solvent, or in a mixture of the two. Generally, nonaqueous media like ether, ethyl acetate, ethanol, isopropanol, or acetonitrile are preferred. Lists of suitable salts are found in *Remington's Pharmaceutical Sciences*, 17th ed., Mack Publishing Company, Easton, PA, 1985, p. 1418, the disclosure of which is hereby incorporated by reference.

As used herein, the term "unit dose" refers to a single dose which is capable of being administered to a patient, and which can be readily handled and packaged, remaining as a physically and chemically stable unit dose comprising either the active compound itself, or as a pharmaceutically acceptable composition, as described hereinafter.

As used herein, "prodrug" is intended to include any covalently bonded carriers which release the active parent compound as defined in the present invention *in vivo* when

such prodrug is administered to a mammalian subject. Since prodrugs are known to enhance numerous desirable qualities of pharmaceuticals (e.g., solubility, bioavailability, manufacturing, etc.) the compounds of the present invention may be delivered in prodrug form. Thus, the present invention contemplates prodrugs of the claimed compounds, compositions containing the same, and methods of delivering the same. Prodrugs of a compound of the present invention may be prepared by modifying functional groups present in the compound in such a way that the modifications are cleaved, either in routine manipulation or *in vivo*, to the parent compound. Accordingly, prodrugs include, for example, compounds of the present invention wherein a hydroxy, amino, or carboxy group is bonded to any group that, when the prodrug is administered to a mammalian subject, cleaves to form a free hydroxyl, free amino, or carboxylic acid, respectively. Examples include, but are not limited to, acetate, formate and benzoate derivatives of alcohol and amine functional groups; and alkyl, carbocyclic, aryl, and alkylaryl esters such as methyl, ethyl, propyl, iso-propyl, butyl, isobutyl, sec-butyl, tert-butyl, cyclopropyl, phenyl, benzyl, and phenethyl esters, and the like.

It is recognized that compounds of the present invention may exist in various stereoisomeric forms. As such, the compounds of the present invention include their respective diastereomers or enantiomers. The compounds are normally prepared as racemates and can conveniently be used as such, but individual diastereomers or enantiomers can be isolated or synthesized by conventional techniques if so desired. Such racemates and individual diastereomers or enantiomers and mixtures thereof form part of the present invention.

It is well known in the art how to prepare and isolate such optically active forms. Specific stereoisomers can be prepared by stereospecific synthesis using enantiomerically pure or enantiomerically enriched starting materials. The specific stereoisomers of either starting materials or products can be resolved and recovered by techniques known in the art, such as resolution of racemic forms, normal, reverse-phase, and chiral chromatography, recrystallization, enzymatic resolution, or fractional recrystallization of addition salts formed by reagents used for that purpose. Useful methods of resolving and recovering specific stereoisomers described in Eliel, E. L.; Wilen, S.H. *Stereochemistry of Organic Compounds*; Wiley: New York, 1994, and Jacques, J, et al. *Enantiomers,*













Racemates, and Resolutions; Wiley: New York, 1981, each incorporated by reference herein in their entireties.












It is further recognized that functional groups present on the compounds of the present invention may contain protecting groups. For example, the amino acid side chain substituents of the compounds of the present invention can be substituted with protecting groups such as benzyloxycarbonyl or t-butoxycarbonyl groups. Protecting groups are known per se as chemical functional groups that can be selectively appended to and removed from functionalities, such as hydroxyl groups and carboxyl groups. These groups are present in a chemical compound to render such functionality inert to chemical reaction conditions to which the compound is exposed. Any of a variety of protecting groups may be employed with the present invention. Preferred groups for protecting lactams include silyl groups such as t-butyldimethylsilyl ("TBDMS"), dimethoxybenzhydryl ("DMB"), acyl, benzyl ("Bn"), and methoxybenzyl groups. Preferred groups for protecting hydroxy groups include TBS, acyl, benzyl, benzyloxycarbonyl ("CBZ"), t-butyloxycarbonyl ("Boc"), and methoxymethyl. Many other standard protecting groups employed by one skilled in the art can be found in Greene, T.W. and Wuts, P.G.M., "Protective Groups in Organic Synthesis" 2d. Ed., Wiley & Sons, 1991.







Synthesis











The general routes to prepare the examples shown in Tables 1-3 of the present invention are shown in the Schemes 1-18. The intermediates used to prepare the examples and their mass spectral data are shown in the Table B. The reagents and starting materials are commercially available, or readily synthesized by well-known techniques by one of ordinary skill in the arts. All processes disclosed in association with the present invention are contemplated to be practiced on any scale, including milligram, gram, multigram, kilogram, multikilogram or commercial industrial scale. All substituents in the synthetic schemes, unless otherwise indicated, are as previously defined.

Table B

I -1 399 (M + 1)		I-2 485 (M + 1)	
I-3 448 (M + 1)		I-4 355 (M + 1)	
I-5 439 (M + 1)		I-6 427 (M + 1)	
I-7 428 (M + 1)		I-8 568 (M + 1)	
I-9 490 (M + 1)		I-10 461 (M+)	
I-11 390 (M + 1)		I-12 580 (M + 1)	

I-13 369 (M + 1)		I-14 399 (M + 1)	
I-15 414 (M + 1)		I-16 504 (M + 1)	
I-17 464 (M + 1)		I-18 383 (M - 1)	
I-19 339 (M - 1)		I-20 329 (M + 1)	
I-21 399 (M + 1)		I-22 504 (M + 1)	
I-24 371 (M + 1)			


I-25 399 (M + 1)		I-26 427 (M + 1) 413 (M + 1) 413 (M + 1) 427 (M + 1) 399 (M + 1) 385 (M +) 442 (M + 1) 468 (M + 1) 524 (M + 1)	 R 26-1: i-butyl 26-1: i-propyl 26-3: propyl 26-4: butyl 26-5: ethyl 26-6: methyl 26-7: $CH_2CH_2NMe_2$ 26-8: $CH_2CH_2NC_4H_8$ 26-9: $(CH_2)_6NC_4H_8$
I-27 411 (M + 1)		I-28	 R 28-1: i-propyl 28-2: propyl 28-3: cyclopentyl 28-4: i-butyl
I-29 386 (M + 1) 386 (M + 1) 400 (M + 1)	 R 29-1: i-propyl 29-2: propyl 29-3: i-butyl 29-4: butyl	I-30 370 (M + 1)	

I-31  R 31-1: cyclopentyl 31-2: i-butyl 464 (M + 1)		I-32  385 (M +1)	
I-33  R 373 (M + 1) 33-1: ethyl 401 (M + 1) 33-2: i-butyl 387 (M + 1) 33-3: i-propyl 387 (M + 1) 33-4: propyl		I-34  329 (M + 1)	
I-35  492 (M + 1)		I-38  427 (M + 1)	
I-37  400 (M + 1)		I-40  311 (M + 1)	
I-41  399 (M + 1)		I-42  411 (M + 1)	

The general procedures to prepare the pyrrolocarbazoles of the present invention are described in U.S. Pat. Nos. 5,705,511 ("the '511 patent") and 6,630,500, PCT Publ. No. WO 00/47583, *J. Heterocyclic Chemistry*, **2001**, *38*, 591, and *J. Heterocyclic Chemistry*, **2003**, *40*, 135. In general, the lactam nitrogen or intermediate alcohol groups of the intermediates outlined in Table B may be protected with such groups as acetyl, substituted silyl, benzyl, Boc, or dimethoxybenzhydrol.

Intermediate I-20 containing a 2-methyl-pyrazole F-ring, used to prepare examples in Table 2, was prepared from the β–ketone, 2-methyl-1,4,6,7-tetrahydro-5*H*-indazol-5-one (Peet, N. P.; LeTourneau, M. E.; *Heterocycles,* **1991,** *32,* 41) using methods described in the '511 patent and in *J. Heterocyclic Chemistry*, 2003, *40*, 135.


Scheme 1.



I-34

As shown in Scheme 1, the N1-methyl pyrazole derivatives in Table 3 were prepared from the 1-methyl α–ketone (*J. Chem. Res.*, **1986**, 1401). The N2-methyl pyrazole intermediates were prepared according to procedures in *J. Heterocyclic Chem.* **1992**, *19*, 1355.


Scheme 2.



I-1 Examples 1-6

As outlined in Scheme 2, Examples 1-6 may be prepared from intermediate I-1 by converting the hydroxymethyl group, using trifluoroacetic anhydride, to an activated per-trifluoroacetate intermediate. This intermediate may be further reacted *in situ* with an alcohol, amine or amide to produce the appropriate product. In certain cases the alcohol may serve as the solvent, otherwise suitable solvents also include methylene chloride, dichloroethane or toluene. The hydroxymethyl intermediates may be readily prepared by reduction of the corresponding aldehyde or ester intermediate as described in U.S. Pat. No. 6,630,500 and herein. Alternatively the derivatives may be prepared by treating the hydroxymethyl intermediate I-1 with an acid catalyst, such as camphor sulfonic acid, trifluoroacetic acid or p-toluene sulfonic acid, in the presence of the appropriate alcohol in a suitable solvent such as methylene chloride or dichloromethane.


Scheme 3.



I-2 Example 7

As outlined in Scheme 3, the heterocycloalkyl derivatives may be prepared by reacting the ester intermediate I-2 with ethanolamine. Longer chain amino alcohols, such as 3-aminopropanol and 4-aminobutanol, produce the corresponding ring-expanded six or seven-membered heterocycloalkyl groups. Ethylene diamine and longer alkyl chain diamines may be used to produce the analogous imidazoline and ring-expanded diamine heterocycloalkyl derivatives.


Scheme 4.



Examples 9-11, 328, 329

As outlined in Scheme 4, Examples 9-11 and 328-329 were prepared by reacting an appropriate keto-ester intermediate, prepared using standard Friedel-Crafts acylation reactions (described in U.S. Pat. No. 6,630,500 and herein) with a hydrazine or an N-substituted hydrazine derivative.

Scheme 5.




Intermediate I-3 Example 8

As outlined in Scheme 5, Example 8 was prepared from the α,β-unsaturated ester (prepared by a Heck reaction with ethyl acrylate, described in the '511 patent) with

hydrazine. An N-substituted hydrazine derivative may be used to produce the N-substituted derivative.

Scheme 6.




R = H, alkyl

Examples 12-28; 232-235


As outlined in Scheme 6, the cyclic acetal and cyclic thioacetal Examples 12-28 and 232-235 were prepared by reacting an aldehyde or alkyl-ketone intermediate with a ω-diol, di-thiol or hydroxyl-thiol reagent and an acid catalyst such as CSA or p-toluene sulfonic acid in a suitable aprotic solvent such as toluene, benzene, or NMP. The diol or dithiol reagent may have substitutions on the alkyl chain such as in Examples 14, 15, 18-24, and 28. The thioether derivatives may be converted to sulfoxides or sulfones using standard oxidizing reagents such as hydrogen peroxide or mCPBA.

Scheme 7.



As outlined in Scheme 7, cycloalkyl and heterocycloalkyl ketone derivatives and cycloalkyl alkyl and heterocycloalkyl alkyl derivatives, such as in examples 32-38 may be prepared by treating the appropriate pyrrolocarbazole with an appropriate acid chloride, carboxylic acid, mixed carboxylic sulfonic anhydride or mixed carboxylic acid anhydride, in the presence of a Lewis acid catalyst such as $AlCl_3$, or $FeCl_3$ in a solvent such as methylene chloride, dichloroethane, nitromethane, nitrobenzene or carbon disulfide. The resulting ketone can be reduced to the alcohol using standard reducing agents such as sodium borohydride (as in Example 37) or $LiAlH_4$. The hydroxy group of the cycloalkyl or heterocycloalkyl alcohols may be replaced with hydrogen by treatment with trifluoroacetic acid and triethyl silane (as in Example 38 and 309) in methylene chloride.

Scheme 8.




Intermediates I-9 - I-11

Examples 39-42

Scheme 8 outlines a route to heterocycloalkenyl derivatives, such as Examples 39-42. The alkene may be further reduced by standards chemical or catalytic hydrogenation methods to the corresponding heterocycloalkyl derivative.

Scheme 9.




Intermediate I-12

Example 43


Scheme 9 outlines a route for the preparation of substituted or unsubstituted 2-, 3-, or 4-piperdinyl derivatives, such as Example 43. From intermediate I-12, the pyridyl ring may be catalytically reduced to the saturated piperidyl using a Raney nickel catalyst. Alternatively, using methods described for Examples 44 and 45, the piperidinyl may be incorporated by Stille or Suzuki coupling reactions using the appropriate piperidinyl stannane or borate and followed by a reduction to the saturated piperidinyl. The piperidinyl nitrogen may be alkylated by standard alkylation reactions as described for Examples 46-48. Standard aryl or heteroaryl Stille and Suzuki coupling techniques may also be used to prepare fused heterocycloalkyl-aryl derivatives.

Scheme 10.




Scheme 10 outlines routes for the preparation of Examples 49-51 using standard methodologies.

Scheme 11.




Intermediate I-14 Examples 52-53




Scheme 11 outlines the route to prepare carbamate-type derivatives, such as Examples 52-53, and 404-422. An alternate method to preparing N, N-di-substituted carbamates utilized a nitrophenyl carbonate intermediate which may be treated with various primary or secondary amines. Similarly urea, O-carbamate, and N-carbamate derivatives may be prepared from reaction of the appropriate amine or phenol intermediate with an isocyanate or chloroformate or from the appropriate nitrophenyl carbonate, nitrophenyl carbamate, or trichloromethylcarbonyl (see *J. Org. Chem.* **2003**, 68, 3733-3735). Ether derivatives, such as Examples 60-67, may be prepared from the phenol intermediates using standard alkylation techniques known to those skilled in the art of organic synthesis.

Scheme 12.




Scheme 12 shows a process for preparation of heterocycloalkyl lactone derivatives, such as Examples 68, 70-74. The intermediate keto-ester was reduced to a hydroxyl-ester intermediate, which cyclizes to the lactone.

Scheme 13.




Scheme 14.



Example 80 Example 83


Scheme 15.



Intermediate I-42 Example 135

Schemes 13-15 outline preparations for cyclic ether derivatives. The cyclic ether group was installed by formation of a keto-ester intermediate prepared through a standard Friedel-Crafts type acylation reaction (described in U.S. Pat. No. 6,630,500 and herein). The alkyl spacer between the chlorocarbonyl and ester may be substituted or unsubstituted. The keto-ester was then reduced to the diol intermediate using $LiBH_4$ in THF or $NaBH_4$ in ethanol. The ring may be closed by treatment with an acid such as trifluoroacetic acid in dichloromethane or dichloroethane. As shown in Scheme 14, the R^2 group could be incorporated after installation of the cyclic ether using a base and an electrophile, or as shown in Scheme 15, the R^2 group may be installed first, followed by formation of the cyclic ether ring.

Scheme 16.




Intermediates I-24 - I-27


Examples 29, 169, 236-286, 277-300

Scheme 16 outlines preparation of oxime and hydrazone derivatives (such as Examples 29, 169, 236-286, and 277-300) from the appropriate aldehyde or ketone using standard procedures known in the art.

Scheme 17.




Intermediate I-28



Examples 287-306

Scheme 17 shows preparation of arylamine derivatives, such as Examples 287-306. The aryl amine reagent is coupled with the bromo intermediate followed by lactam formation using RaNi and hydrogen in DMF/MeOH.

Scheme 18.



Intermediate I-29


Example 309

Scheme 18 shows a route for the preparation of heteroarylamine derivatives, such as Examples 309-318. Amino intermediates I-29 and I-37 were prepared by alkylation of the appropriate cyano-esters with the appropriate alkyl iodide or bromide followed by nitration, and subsequent RaNi reduction to provide the amino-lactam. Reaction of the amino lactam intermediate with an appropriate heteroaryl bromide or chloride produced the desired compounds.











Examples














Other features of the invention will become apparent in the course of the following descriptions of exemplary embodiments as shown in the following Tables 1-3. The compounds of Tables 1-3 show activity in the targets described herein at concentrations ranging from 0.1 nM to 10 μM. These examples are given for illustration of the invention and are not intended to be limiting thereof.















Table 1






























*R^1 is H unless otherwise noted










Ex. No.	R^3	R^2	Q	R^5
1		$CH_2CH_2CH_2OH$	CH_2	H
2		$CH_2CH_2CH_2OH$	CH_2	H
3		$CH_2CH_2CH_2OH$	CH_2	H
4		$CH_2CH_2CH_2OH$	CH_2	H
5		$CH_2CH_2CH_2OH$	CH_2	H
6		$CH_2CH_2CH_2OH$	CH_2	H
7		H	CH_2CH_2	OMe
8		$CH_2CH_2CH_2OH$	CH_2	H
9		H	CH_2CH_2	OCH_3
10		H	CH_2CH_2	OCH_3
11		H	CH_2CH_2	OCH_3














Ex. No.	R^3	R^2	Q	R^5
12		$CH_2CH_2CH_2OH$	CH_2	H
13		$CH_2CH_2CH_2OH$	CH_2	H
14		$CH_2CH_2CH_2OH$	CH_2	H
15		$CH_2CH_2CH_2OH$	CH_2	H
16		$CH_2CH_2CO_2Et$	CH_2	H
17		$CH_2CH_2CH_2OH$	CH_2	H
18		$CH_2CH_2CO_2Et$	CH_2	H
19		$CH_2CH_2CH_2OH$	CH_2	H
20		$CH_2CH_2CH_2OH$	CH_2	H
21		$CH_2CH_2CO_2Et$	CH_2	H
22		$CH_2CH_2CH_2OH$	CH_2	H
23		$CH_2CH_2CO_2Et$	CH_2	H
24		$CH_2CH_2CH_2OH$	CH_2	H















Ex. No.	R^3	R^2	Q	R^5
25		$CH_2CH_2CO_2Et$	CH_2	H
26		$CH_2CH_2CH_2OH$	CH_2	H
27		$CH_2CH_2CH_2OH$	CH_2	H
28		H	CH_2CH_2	O^iPr
29		CH_2CH_2OH	CH_2CH_2	OCH_3
30	$N\equiv C\text{-}N(H)CH_2$-	CH_2CH_2OH	CH_2CH_2	OCH_3
31		CH_2CH_2OBn	CH_2CH_2	OCH_3
32		H	CH_2CH_2	OCH_3
33		H	CH_2	H
34		H	CH_2	H
35		$CH_2CH_2CO_2Et$	CH_2	H
36		$CH_2CH_2CH_2OC(=O)$-cyclopropyl	CH_2	H
37		$CH_2CH_2CH_2OH$	CH_2	H
38		$CH_2CH_2CH_2OH$	CH_2	H
39		$CH_2CH_2CH_2OC(=O)$-CH_3	CH_2	H

















Ex. No.	R^3	R^2	Q	R^5
40		$CH_2CH_2CH_2OH$	CH_2	H
41		H	CH_2CH_2	OiPr
42		H	CH_2	H
43		$CH_2CH_2CH_2OH$	CH_2CH_2	OCH_3
44		$CH_2CH_2CH_2OH$	CH_2	H
45		$CH_2CH_2CH_2OH$	CH_2	H
46		$CH_2CH_2CH_2OH$	CH_2	H
47		$CH_2CH_2CH_2OH$	CH_2	H
48		$CH_2CH_2CH_2OH$	CH_2	H
49		$CH_2CH_2CH_2OCO$-CH_3	CH_2	H
50		$CH_2CH_2CH_2OH$	CH_2	H
51		$CH_2CH_2CH_2OH$	CH_2	H
52		H	CH_2CH_2	OiPr
53		H	CH_2CH_2	OiPr
















Ex. No.	R^3	R^2	Q	R^5
54		CH_2CH_2OH	CH_2CH_2	OCH_3
55		CH_2CH_2OBn $*R^1 = CH(p\text{-methoxyphenyl})_2$	CH_2CH_2	OCH_3
56		CH_2CH_2OBn	CH_2CH_2	OCH_3
57		CH_2CH_2OBn $*R^1 = CH(p\text{-methoxyphenyl})_2$	CH_2CH_2	OCH_3
58		CH_2CH_2OH	CH_2CH_2	OCH_3
59	F	H	CH_2CH_2	
60		H	CH_2CH_2	OiPr
61	H	CH_2CH_2OH	CH_2CH_2	
62	H	CH_2CH_2OH	CH_2CH_2	
63	H	CH_2CH_2OH	CH_2CH_2	
64	H	CH_2CH_2OH	CH_2CH_2	O-$CH(CO_2Me)_2$
65	H	CH_2CH_2OH	CH_2CH_2	$OCH[(CH_2)_3\text{-}OEt]_2$
66	H		CH_2CH_2	
67	H	H	CH_2CH_2	
68		$(CH_2)_3O_2C(CH_2)_2\text{-}CO_2CH_3$	CH_2	H



















Ex. No.	R^3	R^2	Q	R^5
69		H	CH_2CH_2	OCH_3
70		H	CH_2CH_2	OCH_3
71		H	CH_2CH_2	OCH_3
72		H	CH_2CH_2	OCH_3
73		Et	CH_2CH_2	OCH_3
74		$CH_2CH_2CH_2OH$	CH_2	H
75		H	CH_2CH_2	OCH_3
76		$CH_2CH_2CH_2OH$	CH_2	H
77		$CH_2CH_2CH_2OH$	CH_2	H
78		H	CH_2	H
79		H	CH_2	H
80		H	CH_2CH_2	OCH_3
81		CH_2CH_2OH	CH_2CH_2	OCH_3
















Ex. No.	R^3	R^2	Q	R^5
82		CH_2CH_2OH	CH_2CH_2	OCH_3
83		CH_2CH_3	CH_2CH_2	OCH_3
84	 single isomer (-)	CH_2CH_3	CH_2CH_2	OCH_3
85	 single isomer (+)	CH_2CH_3	CH_2CH_2	OCH_3
86		H	CH_2	H
87		$CH_2CH_2CH_2OH$	CH_2	H
88		$CH_2CH_2CH_2OH$	CH_2	H
89		$CH_2CH_2CH_2OH$	CH_2	H
90		$CH_2CH_2CH_2OH$	CH_2	H
91		H	CH_2	H
92		CH_2CH_3	CH_2	H
93		$CH_2CH_2CH_3$	CH_2	H
94		CH_3	CH_2	H

Ex. No.	R^3	R^2	Q	R^5
95		$CH_2CH_2CH_2CH_2CH_3$	CH_2	H
96		$CH_2CH{=}CH_2$	CH_2	H
97		$CH_2CH_2CH_2CH_2CH_3$	CH_2	H
98		$CH_2CH_2CH_2CH_3$	CH_2	H
99		$CH_2CH_2NEt_2$	CH_2	H
100		H	CH_2CH_2	OCH_3
101		H	CH_2CH_2	OCH_3
102		H	CH_2CH_2	OCH_3
103		H	CH_2CH_2	OCH_3
104			CH_2CH_2	OCH_3
105		CH_3	CH_2CH_2	OCH_3
106		$(CH_2)_4CH_3$	CH_2CH_2	OCH_3
107		$(CH_2)_5CH_3$	CH_2CH_2	OCH_3

Ex. No.	R^3	R^2	Q	R^5
108		$(CH_2)_3CH_3$	CH_2CH_2	OCH_3
109		$(CH_2)_2CH(CH_3)_2$	CH_2CH_2	OCH_3
110		$CH_2CH=CH_2$	CH_2CH_2	OCH_3
111		$CH_2CH_2CH_2OTBS$	CH_2CH_2	OCH_3
112		$CH_2CH_2CH_2OTHP$	CH_2CH_2	OCH_3
113		$CH_2CH_2CH_2OH$	CH_2CH_2	OCH_3
114		$CH_2CH_2NEt_2$	CH_2CH_2	OCH_3
115			CH_2CH_2	OCH_3
116		 *and R^1 is 	CH_2CH_2	OCH_3
117		$CH_2CH=CHEt$	CH_2CH_2	OCH_3
118		$CH_2CH_2CH_2CN$	CH_2CH_2	OCH_3
119		$CH_2CH_2CH_3$	CH_2CH_2	OCH_3
120		$CH_2CH(CH_3)_2$	CH_2CH_2	OCH_3

Ex. No.	R^3	R^2	Q	R^5
121		CH_2OCH_3	CH_2CH_2	OCH_3
122			CH_2CH_2	OCH_3
123		CH_2CH_2F	CH_2CH_2	OCH_3
124		$(CH_2)_3OTBS$	CH_2CH_2	OCH_3
125		$(CH_2)_3OH$	CH_2CH_2	OCH_3
126		$CH_2CH_2CH_3$	CH_2CH_2	OCH_3
127		$(CH_2)_2OTBS$	CH_2CH_2	OCH_3
128		$(CH_2)_2OH$	CH_2CH_2	OCH_3
129		CH_2CH_3	CH_2CH_2	OCH_3
130		$CH_2CH_2CH_2CN$	CH_2CH_2	OCH_3
131		CH_3	CH_2CH_2	OCH_3
132			CH_2CH_2	OCH_3
133		$CH_2C{\equiv}CH$	CH_2CH_2	OCH_3

Ex. No.	R^3	R^2	Q	R^5
134		H	CH_2CH_2	OH
135		CH_2CH_3	CH_2CH_2	OH
136		H	CH_2CH_2	OiPr
137		$CH_2CH_2CH_3$	CH_2CH_2	O^nPr
138		H	CH_2CH_2	O^nPr
139		$CH(CH_3)_2$	CH_2CH_2	O^iPr
140		CH_2CH_3	CH_2CH_2	OEt
141		H	CH_2CH_2	
142		H	CH_2CH_2	
143		$CH_2CH_2CH_2CN$	CH_2CH_2	O-$CH_2CH_2CH_2$-CN
144		H	CH_2CH_2	$O(CH_2)_3CN$
145		$CH_2CH=CH_2$	CH_2CH_2	OCH_2-$CH=CH_2$
146		H	CH_2CH_2	OCH_2-$CH=CH_2$
147		H	CH_2CH_2	$O(CH_2)_4CN$
148		$(CH_2)_4CN$	CH_2CH_2	$O(CH_2)_4CN$
149		$CH_2CH_2CH_2CN$	CH_2CH_2	O^iPr

Ex. No.	R^3	R^2	Q	R^5
150		CH_3 *R^1 is Me	CH_2CH_2	O^iPr
151		CH_3	CH_2CH_2	O^iPr
152		$CH(CH_3)CH_2CH_3$	CH_2CH_2	O^iPr
153			CH_2CH_2	O^iPr
154		CH_2CH_3	CH_2CH_2	O^iPr
155		CH_2CH_2F	CH_2CH_2	O^iPr
156		$CH_2CH_2CH_2F$	CH_2CH_2	O^iPr
157		$CH_2CH_2CH_2CH_3$	CH_2CH_2	O^iPr
158			CH_2CH_2	O^iPr
159		H	CH_2CH_2	OH
160		CH_3	CH_2CH_2	OMe
161		CH_2CH_3	CH_2CH_2	OEt
162		H	CH_2CH_2	OEt
163		H	CH_2CH_2	OiPr
164		$CH(CH_3)_2$	CH_2CH_2	O^iPr
165		CH_2CH_3	CH_2CH_2	O^iPr
























Ex. No.	R^3	R^2	Q	R^5
166		H	CH_2CH_2	OCH_3
167		$CH_2CH_2CH_2OH$	CH_2	H
168		H	CH_2	H
169		$CH_2CH_2CH_2OH$	CH_2	H
170		CH_2CH_3	CH_2CH_2	OCH_3
171		CH_2CH_3	CH_2CH_2	OCH_3

















Table 2




































*R^1 is H unless otherwise noted















Ex. No.	R^3	R^2	Q
172		H	CH_2CH_2
173		H	CH_2CH_2

















Ex. No.	R^3	R^2	Q
174		H	CH_2CH_2
175		H	CH_2CH_2
176		H	CH_2CH_2
177		CH_2CH_3	CH_2CH_2
178		$(CH_2)_3CN$	CH_2CH_2
179		$CH_2CH_2CH_3$	CH_2CH_2
180		CH_2CH_3	CH_2CH_2
181		CH_3	CH_2CH_2
182		$(CH_2)_3OH$	CH_2CH_2
183		$CH_2CH{=}CH_2$	CH_2CH_2
184		CH_2CH_2OEt	CH_2CH_2
185		$CH_2CH_2CH_2CH_3$	CH_2CH_2
186		CH_2CH_2OH	CH_2CH_2
187			CH_2CH_2














Ex. No.	R^3	R^2	Q
188		$CH(CH_3)_2$	CH_2CH_2
189		$CH_2CH(CH_3)_2$	CH_2CH_2
190		$CH_2C{\equiv}CH$	CH_2CH_2
191		$CH_2CH_2NEt_2$	CH_2CH_2
192			CH_2CH_2
193		$CH_2CH_2CH(CH_3)_2$	CH_2CH_2
194		$CH_2CH_2CH_2CH_2$-CH_3	CH_2CH_2
195		$CH_2CH_2CH_2$-$N(CH_3)_2$	CH_2CH_2
196			CH_2CH_2
197		$CH_2CH_2CH_2CH$-$(CH_3)_2$	CH_2CH_2
198	 single isomer	$CH_2CH_2CH_2CH_3$	CH_2CH_2
199	 single isomer	$CH_2CH_2CH_2CH_3$	CH_2CH_2
200		$CH_2CH_2CH_2F$	CH_2CH_2
201		$CH_2CH_2CH_2CH_2F$	CH_2CH_2





















Ex. No.	R^3	R^2	Q
202	 single isomer	iPr	CH_2CH_2
203	 single isomer	iPr	CH_2CH_2
204	 single isomer		CH_2CH_2
205	 single isomer		CH_2CH_2
206		$CH_2CH=CMe_2$	CH_2CH_2
207			CH_2CH_2
208		CH_2CH_2SEt	CH_2CH_2
209		H *R^1 is $C(=O)CH_3$	CH_2CH_2
210			CH_2CH_2
211			CH_2CH_2
212			CH_2CH_2
213			CH_2CH_2
214			CH_2CH_2
215			CH_2CH_2













Ex. No.	R^3	R^2	Q
216			CH_2CH_2
217	 single isomer	$CH_2CH(CH_3)_2$	CH_2CH_2
218	 single isomer	$CH_2CH(CH_3)_2$	CH_2CH_2
219		$CH_2CH_2CH_2CN$	CH_2CH_2
220		CH_2CH_3	CH_2CH_2
221		$CH_2CH_2CH_2CH_3$	CH_2CH_2
222		CH_3	CH_2CH_2
223		$CH_2CH=CH_2$	CH_2CH_2
224		$CH_2CH_2CH_3$	CH_2CH_2
225			CH_2CH_2
226		$CH_2CH_2CH_2F$	CH_2CH_2
227		$CH_2CH_2CH_2CH_2F$	CH_2CH_2















Ex. No.	R^3	R^2	Q
228		CH_2CH_3	CH_2CH_2
229		$CH_2CH_2CH_3$	CH_2CH_2
230		CH_2CH_3	CH_2CH_2
231		$CH_2CH_2CH_3CN$	CH_2CH_2
232		H *R^1 is $C(=O)CH_3$	CH_2CH_2
233		H	CH_2CH_2
234		H *R^1 is $C(=O)CH_3$	CH_2CH_2
235		H	CH_2CH_2
236		H	CH_2CH_2
237		H	CH_2CH_2
238		H	CH_2CH_2
239		H	CH_2CH_2
240		H	CH_2CH_2
241		H	CH_2CH_2



















Ex. No.	R^3	R^2	Q
242		H	CH_2CH_2
243		H	CH_2CH_2
244		H	CH_2CH_2
245		H	CH_2CH_2
246		$CH_2CH(CH_3)_2$	CH_2CH_2
247		$CH_2CH(CH_3)_2$	CH_2CH_2
248		$CH_2CH(CH_3)_2$	CH_2CH_2
249		$CH_2CH(CH_3)_2$	CH_2CH_2
250		$CH_2CH(CH_3)_2$	CH_2CH_2
251		H	CH_2CH_2
252		$CH(CH_3)_2$	CH_2CH_2
253		$CH(CH_3)_2$	CH_2CH_2
254		$CH(CH_3)_2$	CH_2CH_2
255		$CH(CH_3)_2$	CH_2CH_2
256		$CH_2CH(CH_3)_2$	CH_2CH_2
257		$CH_2CH_2CH_3$	CH_2CH_2
















Ex. No.	R^3	R^2	Q
258		$CH_2CH_2CH_3$	CH_2CH_2
259		$CH_2CH_2CH_2CH_3$	CH_2CH_2
260		$CH_2CH_2CH_2CH_3$	CH_2CH_2
261		$CH_2CH_2CH_2CH_3$	CH_2CH_2
262		$CH_2CH_2CH_2CH_3$	CH_2CH_2
263		CH_2CH_3	CH_2CH_2
264		CH_2CH_3	CH_2CH_2
265		CH_2CH_3	CH_2CH_2
266		CH_2CH_3	CH_2CH_2
267		CH_2CH_3	CH_2CH_2
268		$CH_2CH(CH_3)_2$	CH_2CH_2
269		$CH_2CH_2CH_3$	CH_2CH_2
270		$CH_2CH_2CH_3$	CH_2CH_2
















Ex. No.	R^3	R^2	Q
271		CH_3	CH_2CH_2
272		CH_3	CH_2CH_2
273		CH_3	CH_2CH_2
274		$CH_2CH_2NMe_2$	CH_2CH_2
275		$CH_2CH_2NMe_2$	CH_2CH_2
276		$CH_2CH_2NMe_2$	CH_2CH_2
277		$CH_2CH(CH_3)_2$	CH_2CH_2
278		$CH_2CH(CH_3)_2$	CH_2CH_2
279			CH_2CH_2
280			CH_2CH_2
281			CH_2CH_2
282			CH_2CH_2
283			CH_2CH_2
284			CH_2CH_2
285			CH_2CH_2
286			CH_2CH_2















Ex. No.	R^3	R^2	Q
287		$CH(CH_3)_2$	CH_2CH_2
288		$CH(CH_3)_2$	CH_2CH_2
289		$CH(CH_3)_2$	CH_2CH_2
290		$CH(CH_3)_2$	CH_2CH_2
291		$CH(CH_3)_2$	CH_2CH_2
292		$CH(CH_3)_2$	CH_2CH_2
293		$CH_2CH_2CH_3$	CH_2CH_2
294		$CH_2CH_2CH_3$	CH_2CH_2
295		$CH_2CH_2CH_3$	CH_2CH_2
296		$CH(CH_3)_2$	CH_2CH_2
297		$CH(CH_3)_2$	CH_2CH_2
298		$CH_2CH_2CH_3$	CH_2CH_2















Ex. No.	R^3	R^2	Q
299			CH_2CH_2
300		$CH_2CH_2CH_3$	CH_2CH_2
301		$CH_2CH_2CH_3$	CH_2CH_2
302		$CH_2CH_2CH_3$	CH_2CH_2
303		$CH(CH_3)_2$	CH_2CH_2
304		$CH_2CH(CH_3)_2$	CH_2CH_2
305		$CH_2CH(CH_3)_2$	CH_2CH_2
306		$CH(CH_3)_2$	CH_2CH_2
307		$CH_2CH_2CH_3$	CH_2CH_2
308		$CH_2CH(CH_3)_2$	CH_2CH_2
309		$CH(CH_3)_2$	CH_2CH_2
310		$CH_2CH_2CH_3$	CH_2CH_2
311		$CH_2CH(CH_3)_2$	CH_2CH_2
312		$CH(CH_3)_2$	CH_2CH_2

















Ex. No.	R^3	R^2	Q
313		$CH_2CH(CH_3)_2$	CH_2CH_2
314		$CH(CH_3)_2$	CH_2CH_2
315		$CH_2CH(CH_3)_2$	CH_2CH_2
316		$CH_2CH(CH_3)_2$	CH_2CH_2
317		$CH_2CH(CH_3)_2$	CH_2CH_2
318		$CH_2CH_2CH_3$	CH_2CH_2
319			CH_2CH_2
320			CH_2CH_2
321		H	CH_2CH_2
322		H	CH_2CH_2
323		H	CH_2CH_2
324		H	CH_2CH_2
325		H	CH_2CH_2
326		H	CH_2CH_2
327		$CH(CH_3)_2$	CH_2CH_2
328		H	CH_2CH_2

















Ex. No.	R^3	R^2	Q
329		$CH_2CH(CH_3)_2$	CH_2CH_2
330		$CH_2CH(CH_3)_2$	CH_2CH_2
331		$CH_2CH_2CH_3$	CH_2CH_2
332		$CH_2CH_2CH_3$	CH_2CH_2
333		$CH_2CH_2CH_3$	CH_2CH_2
334		$CH_2CH_2CH_2CH_3$	CH_2CH_2
335		$CH_2CH_2CH_3$	CH_2CH_2
336		$CH_2CH_2CH_3$	CH_2CH_2
337		$CH_2CH_2CH_2CH_3$	CH_2CH_2
338		$CH_2CH_2CH_2CH_3$	CH_2CH_2
339		$CH_2CH(CH_3)_2$	CH_2CH_2
340		$CH_2CH(CH_3)_2$	CH_2CH_2
341		$CH_2CH(CH_3)_2$	CH_2CH_2
342		$CH_2CH(CH_3)_2$	CH_2CH_2
343		$CH_2CH(CH_3)_2$	CH_2CH_2

Ex. No.	R^3	R^2	Q
344		$CH_2CH(CH_3)_2$	CH_2CH_2
345		$CH_2CH_2CH_3$	CH_2CH_2
346		$CH_2CH(CH_3)_2$	CH_2CH_2
347		$CH_2CH_2CH_2CH_3$	CH_2CH_2
348		$CH_2CH_2CH_3$	CH_2CH_2
349		$CH_2CH_2CH_3$	CH_2CH_2
350		$CH_2CH_2CH_3$	CH_2CH_2
351		$CH_2CH_2CH_3$	CH_2CH_2
352		$CH_2CH_2CH_3$	CH_2CH_2
353		$CH_2CH_2CH_3$	CH_2CH_2
354		$CH_2CH_2CH_3$	CH_2CH_2
355		CH_2CH_3	CH_2CH_2
356		$CH(CH_3)_2$	CH_2CH_2
357		$CH(CH_3)_2$	CH_2CH_2
358		$CH_2CH_2CH_3$	CH_2CH_2

Ex. No.	R^3	R^2	Q
359		$CH(CH_3)_2$	CH_2CH_2
360		$CH_2CH_2CH_3$	CH_2CH_2
361		$CH_2CH_2CH_3$	CH_2CH_2
362		$CH(CH_3)_2$	CH_2CH_2
363		$CH_2CH_2CH_3$	CH_2CH_2
364		$CH_2CH_2CH_3$	CH_2CH_2
365		$CH(CH_3)_2$	CH_2CH_2
366		$CH(CH_3)_2$	CH_2CH_2
367		$CH(CH_3)_2$	CH_2CH_2
368		$CH(CH_3)_2$	CH_2CH_2
369		$CH_2CH_2CH_3$	CH_2CH_2
370		$CH_2CH_2CH_3$	CH_2CH_2
371		$CH_2CH_2CH_3$	CH_2CH_2
372		$CH_2CH_2CH_3$	CH_2CH_2

Ex. No.	R^3	R^2	Q
373		$CH_2CH_2CH_3$	CH_2CH_2
374		$CH_2CH_2CH_3$	CH_2CH_2
375		$CH_2CH_2CH_3$	CH_2CH_2
376		$CH_2CH_2CH_3$	CH_2CH_2
377		$CH_2CH_2CH_3$	CH_2CH_2
378		$CH_2CH_2CH_3$	CH_2CH_2
379		$CH_2CH_2CH_3$	CH_2CH_2
380		$CH(CH_3)_2$	CH_2CH_2
381		$CH(CH_3)_2$	CH_2CH_2
382		$CH(CH_3)_2$	CH_2CH_2
383		$CH(CH_3)_2$	CH_2CH_2
384		$CH_2CH(CH_3)_2$	CH_2CH_2
385		$CH_2CH(CH_3)_2$	CH_2CH_2
386		$CH_2CH(CH_3)_2$	CH_2CH_2

Ex. No.	R^3	R^2	Q
387		$CH(CH_3)_2$	CH_2CH_2
388		$CH(CH_3)_2$	CH_2CH_2
389		$CH(CH_3)_2$	CH_2CH_2
390		$CH(CH_3)_2$	CH_2CH_2
391		$CH(CH_3)_2$	CH_2CH_2
392		$CH(CH_3)_2$	CH_2CH_2
393		$CH(CH_3)_2$	CH_2CH_2
394		$CH(CH_3)_2$	CH_2CH_2
395		$CH(CH_3)_2$	CH_2CH_2
396		$CH_2CH_2CH_2CH_3$	CH_2CH_2
397		$CH_2CH_2CH_2CH_3$	CH_2CH_2
398		$CH_2CH_2CH_2CH_3$	CH_2CH_2
399		$CH_2CH_2CH_3$	CH_2CH_2
400		$CH_2CH_2CH_3$	CH_2CH_2
401		$CH_2CH_2CH_3$	CH_2CH_2
402		$CH_2CH_2CH_3$	CH_2CH_2

Ex. No.	R^3	R^2	Q
403		$CH(CH_3)_2$	CH_2CH_2
404		CH_2CH_3	CH_2CH_2
405		CH_2CH_3	CH_2CH_2
406		CH_2CH_3	CH_2CH_2
407		$CH_2CH(CH_3)_2$	CH_2CH_2
408		$CH_2CH(CH_3)_2$	CH_2CH_2
409		$CH_2CH(CH_3)_2$	CH_2CH_2
410		$CH_2CH(CH_3)_2$	CH_2CH_2
411		$CH_2CH(CH_3)_2$	CH_2CH_2
412		$CH_2CH(CH_3)_2$	CH_2CH_2
413		$CH_2CH(CH_3)_2$	CH_2CH_2
414		$CH_2CH(CH_3)_2$	CH_2CH_2
415		$CH_2CH(CH_3)_2$	CH_2CH_2
416		$CH_2CH(CH_3)_2$	CH_2CH_2
417		$CH_2CH(CH_3)_2$	CH_2CH_2
418		$CH_2CH(CH_3)_2$	CH_2CH_2
























Ex. No.	R^3	R^2	Q
419		$CH_2CH(CH_3)_2$	CH_2CH_2
420		$CH_2CH(CH_3)_2$	CH_2CH_2
421		$CH_2CH(CH_3)_2$	CH_2CH_2
422		$CH(CH_3)_2$	CH_2CH_2
423		$CH_2CH(CH_3)_2$	CH_2CH_2
424		$CH(CH_3)_2$	CH_2CH_2
425		$CH(CH_3)_2$	CH_2CH_2
426		$CH_2CH(CH_3)_2$	CH_2CH_2
427		$CH(CH_3)_2$	CH_2CH_2
428		$CH_2CH(CH_3)_2$	CH_2CH_2
429		$CH_2CH(CH_3)_2$	CH_2CH_2
430		$CH_2CH(CH_3)_2$	CH_2CH_2

Table 3



Ex. No.	R^3	R^2	Q
431		H	CH_2CH_2
432		$CH_2CH_2CH_2CH_3$	CH_2CH_2
433		$CH_2CH(CH_3)_2$	CH_2CH_2
434			CH_2CH_2
435		CH_2CH_3	CH_2CH_2
436		$CH(CH_3)_2$	CH_2CH_2
437		$CH_2CH_2CH_3$	CH_2CH_2
438		CH_3	CH_2CH_2
439			CH_2CH_2
440		$CH_2CH_2OCH_2CH_3$	CH_2CH_2
441		$CH_2CH_2CH_2OH$	CH_2CH_2




















Ex. No.	R^3	R^2	Q
442		CH_2CH_2OH	CH_2CH_2
443		$CH(CH_3)_2$	CH_2CH_2
444		$CH_2CH(CH_3)_2$	CH_2CH_2
445		$CH_2CH(CH_3)_2$	CH_2CH_2
446		$CH_2CH(CH_3)_2$	CH_2CH_2
447		$CH_2CH(CH_3)_2$	CH_2CH_2
448		CH_2CH_3	CH_2CH_2





Table 4





Example No.	Structure	MS m/e (M + 1)
449		578



450		508
451		552
452		480
453		524

454		538
455		566
456		568
457		582

458		610
459		596
460		568
461		554

462		596
463		582
464		540
465		554

466		568
467		582
468		582
469		610

470		540
471		554

General Procedure for Examples 1-3.

To a suspension of the diol intermediate I-1 (1 equivalent) in either the appropriate alcohol, or the appropriate alcohol with either methylene chloride or chloroform, at room temperature in a reaction tube was slowly added trifluoroacetic anhydride (1-2 equiv). The tube was flushed with nitrogen and sealed tightly. The mixture was stirred at room temperature for 1-2 hours then heated to 80 °C for 2-60 h and monitored for disappearance of starting material by HPLC. Upon completion the reaction was allowed to cool to room temperature, concentrated and worked up by either triturating the residue with ether or collecting the resulting precipitate by filtration or extraction of the product from the reaction mixture with a suitable organic solvent. The solid product was purified by

triturating with ether or by flash chromatography on silica gel using ethyl acetate or a mixture of ethyl acetate and hexane.

Example 1. white solid ^{1}H NMR (DMSO-d_6): δ 0.87 (t, 6H), 1.50 (m, 4H), 1.93 (m, 2H), 3.46 (m, 2H), 4.52 (s, 2H), 4.62 (s, 2H), 4.73 (m, 3H), 4.89 (s, 2H), 7.37 (m, 2H), 7.48 (m, 1H), 7.66 (m, 2H), 7.91 (s, 1H), 8.54 (s, 1H), 9.47 (d, 1H); MS (m/e) 469 (M + 1).

Example 2. ^{1}H NMR (DMSO-d_6): δ 0.89 (d, 6H), 1.47 (m, 2H), 1.70 (m, 1H), 1.96 (m, 2H), 3.52 (m, 4H), 4.56 (s, 2H), 4.64 (s, 2H), 4.77 (m, 2H), 4.93 (s, 2H), 7.40 (m, 2H), 7.51 (d, 1H), 7.69 (m, 2H), 7.94 (s, 1H), 8.58 (s, 1H), 9.48 (d, 1H); MS (m/e) 469 (M + Na).

Example 3. ^{1}H NMR (DMSO-d_6): δ 1.51 (m, 2H), 1.98 (m, 4H), 3.52 (m, 2H), 3.67 (m, 1H), 3.84 (m, 2H), 4.58 (s, 2H), 4.73 (s, 2H), 4.79 (m, 4H), 4.95 (s, 2H), 7.28-7.48 (m, 2H), 7.54 (d, 1H), 7.67-7.80 (m, 2H), 7.96 (s, 1H), 7.60 (s, 1H), 9.51 (d, 1H); MS (ESI): m/e 505 $(M+Na)^+$.

General method for Examples 4-6

To a well-stirred suspension of intermediate diol I-1 (0.150 g, 0.37 mmol) in 7 mL of methylene chloride were added sequentially trifluoroacetic anhydride (0.395 g, 1.88 mmol) and N-methyl morpholine (0.152 g, 1.5 mmol) at 5 °C and under argon atmosphere. The resulted suspension was further stirred at room temperature for 3 h and the low boiling solvents were removed under vacuum. The solid was suspended in 70 mL of acetonitrile and the excess amine or amide was added then refluxed for 18 h and acetonitrile was removed under vacuum. The crude material was dissolved in a mixture of THF and MeOH (1:1 ratio, 7 mL) and treated with 10 mL of 0.5 M solution of NaOMe in MeOH. The reaction mixture was stirred at room temperature for 1 h and the low boiling solvents were removed under vacuum and the reaction mixture was quenched with water. The solid was filtered, washed with water and dried to provide the crude solid. The crude material was purified by silica gel column chromatography using 5% MeOH in methylene chloride to obtain pure product.

Example 4. ^{1}H NMR $(CD_3)_2SO$ δ: 1.86-1.99 (m, 2H), 2.0-2.46 (m,m, 4H), 3.15 (t, 2H), 3.35-3.47 (m, 2H), 4.45-4.52 (m, 4H), 4.71-4.73 (m, 2H), 4.88 (s, 2H), 7.30-7.41 (m, 3H), 7.62-7.69 (m, 2H), 7.82 (s, 1H), 8.55 (s, 1H), 9.46 (d, 1H); MS: m/e 466 (M+1).

Example 5. ^{1}H NMR $(CD_3)_2SO$ δ: 1.68 (br s, 4H), 1.90-1.92 (m, 2H), 2.11-2.14 (m, 2H), 3.09-3.22 (m, 2H), 3.40-3.47 (m, 2H), 4.50 (s, 1H), 4.65-4.74 (m, 4H), 4.87 (s, 2H), 7.32-7.41 (m, 3H), 7.61-7.68 (m, 2H), 7.81 (s, 1H), 8.53 (s, 1H), 9.46 (d, 1H); MS: m/e 480 (M+1).

Example 6. ^{1}H NMR $(CD_3)_2SO$ δ: 1.91-193 (m, 2H), 3.42-3.46 (m, 2H), 4.50 (s, 2H), 4.69-4.74 (m, 2H), 4.88 (s, 2H), 5.70 (s, 2H), 7.32-7.41 (m, 2H), 7.57-7.63 (m, 2H), 7.73 (d, 1H), 8.14 (s, 1H), 8.59 (s, 1H), 9.26 (s, 1H), 9.46 (d, 1H), 10.76 (S, 1H).

Example 7. To a stirred solution of ethanolamine (43.9 uL, 0.729 mmol) in THF (4 mL) was added sodium hydride (17.5 mg, 0.729 mmol) at room temperature under nitrogen. The reaction mixture stirred for 1 hour and a solution of ester intermediate I-2 (166 mg, 0.243 mmol) in THF (4 mL) was added. The mixture was stirred at room temperature overnight. The mixture was quenched with water, extracted with methylene chloride and washed with water and brine. The organic phase was dried over magnesium sulfate, filtered, and concentrated *in vacuo* to give a yellow residue. The residue was dissolved in methylene chloride and cooled to 0 °C in an ice bath. Thionyl chloride (70.9 uL, 0.972 mmol) was slowly added dropwise. The mixture became red-orange in color and was warmed to room temperature for 1.5 hours. The mixture was diluted with ethyl acetate, poured over ice and neutralized with solid potassium carbonate to pH 9. The solid precipitate which formed and was collected by filtration and purified by preparative TLC on silica gel using 10% methanol/ chloroform to give a yellowish-tan solid (10.3 mg, 10% yield). ^{1}H NMR (DMSO-d_6): δ 2.85 (m, 2H), 3.03 (t, 2H), 3.82 (s, 3H), 4.00 (t, 2H), 4.45 (t, 2H), 4.84 (s, 2H), 6.83 (d, 1H), 8.00 (d, 1H), 8.19 (d, 1H), 8.39 (s, 1H), 11.91 (s, 1H); MS (ESI): m/e 424 $(M+1)^+$.

Example 8. 3-bromo intermediate I-3 (0.224 g), ethyl acrylate (170 uL), Pd(II)acetate (3.5 mg), tri(O)tolylphosphine (18.6 mg), and TEA (0.3 mL) in dry DMF (3.2 mL) were heated to 90 °C for 21 h. The grey colored powder that formed was collected and diluted with methylene chloride (15 mL), washed with 2% citric acid and NaCl solution and dried. Evaporation of the solvent gave a dark brown gum, triturating gave an orange solid. ^{1}H NMR showed the product. The vinyl ester intermediate (120 mg) was mixed with 60 uL of hydrazine, in 200 uL of ethanol and heated to 70 °C for 3 h. The brown gum obtained was diluted with ethyl acetate. A dark colored gum that settled out was discarded. The ethyl acetate solution was evaporated and a foamy solid was obtained. The HPLC showed the presence of 60% product; MS m/e 453 $(M+1)^+$.

Example 9.

Step A. To a stirred suspension of aluminum chloride (0.47 g, 0.00353 mol) in 1,2-dichloroethane (5 mL) was added 3-carbomethoxypropionyl chloride (0.43 mL, 0.00353 mol). The reaction gradually became homogeneous. A suspension of intermediate I-4 (0.5g, 0.00141 mol) was added to the mixture. The orange reaction mixture was heated to reflux overnight. The mixture was cooled to room temperature and poured over ice and concentrated HCl was added. The mixture stirred for 30 min. and the solid was collected by filtration (0.52 g, 79% yield).

Step B. To a stirred solution of the product from step A (40 mg, 0.08 mmol) in DMF (5 mL) under nitrogen was added hydrazine (11.7 mL, 0.242 mmol). The mixture was heated to reflux overnight and monitored for disappearance of starting material by HPLC. The reaction mixture was cooled to room temperature and concentrated *in vacuo* to an oil. The product was purified by preparative HPLC to give the product as a tan solid (22% yield). ^{1}H NMR (DMSO-d_6): δ 2.86 (m,2H), 3.02 (m, 2H), 3.14 (t, 2H), 3.41 (t, 2H), 3.82 (s, 3H), 4.85 (s, 2H), 6.85 (d, 1H), 6.91 (s, 1H), 7.59 (d, 1H), 7.94 (d, 1H), 8.19 (d, 1H), 8.26 (s, 1H), 8.34 (s, 1H), 10.86 (s, 1H), 11.78 (s, 1H); MS (ESI): m/e 451 $(M+1)^+$.

Example 10. The ester intermediate produced in step A Example 9, (110 mg, 0.222 mmol) dissolved in DMF (3 mL) was placed in a sealed glass reaction tube. Methyl hydrazine

(35.4 uL, 0.666 mmol) was added to the reaction mixture and the tube was sealed and heated to 80 °C for 96 h. The mixture was cooled to room temperature and the solvent removed *in vacuo* leaving a yellow oil. Water was added to the oil and a precipitate formed that was collected by filtration. The solid was purified by preparative TLC chromatography on silica gel using 5% methanol/ chloroform to give a tan solid (12% yield). ^{1}H NMR (DMSO-d_6): δ 2.50 (t, 2H), 2.86 (m, 2H), 3.03 (m, 2H), 3.17 (t, 2H), 3.37 (s, 3H), 3.82 (s, 3H), 4.86 (s, 2H), 6.83 (d, 1H), 6.91 (s, 1H), 7.60 (d, 1H), 7.98 (d, 1H), 8.19 (d, 1H), 8.27 (s, 1H), 8.41 (s, 1H), 11.8 (s, 1H); MS (ESI): m/e 465 $(M+1)^+$.

Example 11. This compound was prepared by a similar method in Example 9 using the ester from step A (110 mg, 0.259 mmol) and 2-hydroxyethylhydrazine (52.6 uL, 0.777 mmol). The reaction mixture was heated to 100 °C for 48 h. The reaction solvent was removed *in vacuo* leaving an orange oil that was purified by preparative TLC chromatography on silica gel using 10% methanol/ chloroform to give a yellow solid (38 mg, 30% yield). ^{1}H NMR (DMSO-d_6): δ 2.56 (dd, 2H), 2.85 (dd, 2H), 3.02 (dd, 2H), 3.16 (dd, 2H), 3.67 (dd, 2H), 3.85 (dd, 2H), 3.82 (s, 3H), 4.69 (t, 1H), 4.86 (s, 2H), 6.83 (d, 1H), 6.91 (s, 1H), 7.61 (d, 1H), 7.99 (d, 1H), 8.20 (d, 1H), 8.26 (s, 1H), 8.41 (s, 1H), 11.79 (s, 1H); MS (ESI): m/e 495 $(M+1)^+$.

Synthesis of aldehyde intermediate I-5.

To a suspension of the indenocarbazole intermediate I-40 (8.0 g, 0.258 mol) in acetonitrile (300 mL) at room temperature under nitrogen was added ethyl acrylate (4.19 mL, 0.387 mol) followed by 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU) (1.93mL, 0.013 mol). After addition of DBU, the reaction changed colors from orange to green. The reaction mixture was heated to reflux overnight. The mixture remained heterogeneous throughout the course of the reaction and became dark in color. A small aliquot was removed after 18 h and the solid was collected by filtration. ^{1}H NMR of the sample showed no starting material remaining. The reaction mixture was cooled to room temperature and the solid was collected by filtration. The solid was washed several times with cold acetonitrile and dried *in vacuo* at 55 °C to yield a light orange solid (5.4 g, 78% yield). ^{1}H NMR (DMSO-d_6): δ 9.72 (t, 3H, J = 6. 8), 2.87 (m, 2H), 3.89 (q, 2H, J = 6.8),

4.49 (s, 2H), 4.88 (s, 2H), 4.92 (m, 2H), 7.29-7.48 (m, 3H), 7.50-7.73 (m, 3H), 7.96 (d, 1H, J = 7.33), 8.56 (s, 1H), 9.47 (d, 1H, J = 7.33). To a well stirred suspension of the ester intermediate (10 g, 24.3 mmol) and 1,1-dichloromethyl methylether (55.6 g, 488 mmol) in a mixture of methylene chloride (250 mL) and toluene (40 mL) was added tin(IV) chloride (95.1 g, 365 mmol) (1 M solution in methylene chloride). After 4.5 h of reaction at room temperature, the reaction was quenched with aqueous 2N HCl (150 mL) and methylene chloride and toluene were removed under vacuum. To the crude residue, an additional amount of 2N HCl (350 mL) and *t*-butylmethyl methyl ether (200 mL) was added. The resulted suspension was further stirred at room temperature for 14 h and filtered, washed with *t*-butylmethyl methylether. The crude material was suspended in a mixture of ethyl acetate (125 mL) and tetrahydrofuran (125 mL) and stirred for 14 h at room temperature then filtered and washed with cold ethyl acetate to obtain 8.4 g of aldehyde. (79% yield). ^{1}H NMR: (DMSO-d_6) δ: 0.98 (t, 3H), 2.92 (t, 2H), 3.88 (q, 2H), 4.65 (s, 2H), 4.92-5.08 (m, 4H), 7.25-7.45 (m, 2H), 7.75 (d, 1H), 7.85 (d,1H), 8.05 (d, 1H), 8.52 (s, 1H), 8.69 (s, 1H), 8.95 (s, 1H), 10.18 (s, 1H); MS m/e 439 (M+1).

General procedure for cyclic acetals and thioacetal Examples 12-28.

A mixture of intermediate I-5 (0.2 g, 0.45 mmol), the diol or thiol (> 4.5 mmol), p-toluenesulfonic acid (0.13 g, 0.68 mmol) and amberlyst (1 g) in a mixture of 1-methyl-2-pyrrolidinone (3 mL) and toluene (25 mL) was refluxed using Dean-Stark apparatus for 2 days and under argon atmosphere. The reaction mixture was filtered, washed with 1-methyl-2-pyrrolidinone and THF then concentrated. The oily liquid was quenched with aq. saturated $NaHCO_3$ solution and the solid was filtered, washed with water to provide the crude material, which was purified by silica gel column chromatography to furnish pure products. The product (1 mmol) was dissolved in THF (4 mL) and was added lithium borohydride (5 mmol) (1M solution in THF) at 0 °C then stirred at room temperature for 14 h. The reaction was quenched with aqueous $NaHCO_3$ solution and extracted with ethyl acetate (3 X 20 mL). The combined organic layers were washed with brine, dried and concentrated to yield the crude product, which was purified by silica gel column chromatography to provide pure product.

Example 12: ^{1}H NMR, (DMSO-d_6) δ: 1.82-2.00 (m, 2H), 3.42-3.55 (m, 2H), 3.95 (t, 2H), 4.12 (t, 2H), 4.51 (s, 2H), 4.65-4.80 (m, 2H), 4.91 (s, 2H), 5.90 (s, 2H), 7.32-7.45 (m, 2H),7.55-7.70 (m, 2H), 7.80 (d, 1H), 7.98 (s, 1H), 8.58 (s, 1H), 9.48 (d, 1H); MS m/e : 441 (M+1).

Example 13: ^{1}H NMR, (DMSO-d_6) δ: 1.82-1.95 (m, 2H), 3.42-3.62 (m, 2H), 3.95 (t, 2H), 4.25-4.39 (d,d, 2H), 4.65 (s, 2H), 4.63-4.72 (m, 2H), 4.89 (s, 2H), 5.72 (s, 1H), 7.32-7.55, (s, 2H), 7.68 (d, 1H), 7.69-7.82 (m, 2H), 7.96 (s, 1H), 8.55 (s, 1H), 9.48 (d, 1H); MS m/e 455 (M+1).

Example 14: ^{1}H NMR, (DMSO-d_6) δ: 0.80 (s, 3H), 1.28 (s, 3H), 1.97-2.33 (m, 2H), 3.40-3.52 (m, 2H), 3.69-3.76 (d,d 4H), 4.55 (s, 2H), 4.73-4.77 (m, 2H), 4.93 (s, 2H), 5.62 (s, 1H), 7.35-7.45 (m, 2H), 7.62-7.68 (m, 2H), 7.74 (d, 1H), 8.01 (s, 1H), 8.51 (s, 1H), 9.51 (d, 1H); MS m/e 483 (M+1).

Example 15: ^{1}H NMR, (DMSO-d_6) δ: 1.19-1.38 (4xd, 6H), 1.96-1.99 (m, 2H), 3.41-3.53 (m, 2H), 3.82-3.87 (m, 2H), 4.55 (s, 2H), 4.72-4.79 (m, 2H), 4.94 (s, 2H), 6.10 (s, 1H), 7.35-7.44 (m, 2H), 7.61-7.68 (m, 2H), 7.75 (d, 1H), 8.02 (s, 1H), 8.53 (s, 1H), 9.51 (d, 1H); MS m/e 469 (M+1).

Example 16: MS m/e 529 (M+1).

Example 17: MS m/e 487 (M+1).

Example 18: MS m/e 609 (M+1).

Example 19: MS m/e 567 (M+1).

Example 20: MS m/e 511 (M+1).

Example 21: MS m/e 557 (M+1).

Example 22: MS m/e 515 (M+1).

Example 23: MS m/e 539 (M+1).

Example 24: MS m/e 497 (M+1).

Example 25: MS m/e 515 (M+1).

Example 26: MS m/e 473 (M+1).

Example 27: MS m/e 471 (M+1).

Example 28: MS m/e 497 (M+1).

Example 29. To intermediate I-6 (30mg, 0.07 mmol) in pyridine (0.5 mL) was added O-(tetrahydro-2H-pyran-2yl)hydroxylamine (33 mg, 0.282 mmol, 4eq) and the reaction was heated to 100 °C overnight. The reaction was concentrated in vacuo, stirred with water, the product collected and dried at 80 °C. ^{1}H NMR (400MHz, DMSO) 8.52(1H, s), 8.42(1H, s), 8.17(1H, s), 7.89(1H, d), 7.81(1H, d), 7.74(1H, d), 6.91(1H, s), 6.82(1H, d), 5.35(1H, s), 5.01(1H, t), 4.81(2H, s), 4.68(2H, d), 3.86(6H, m), 3.56(1H, m), 3.30(2H, m), 2.79(2H, m), 1.90(2H, m), 1.68(4H, m) MS m/e 526 $(M+1)^+$

Example 30. The amino methyl intermediate I-7 (50 mg, 0.11 mmol) was added to a stirred solution of 10.65 µl cyanogen bromide (0.11 mmol) in dry ether (1mL) at -20 °C containing anhydrous sodium carbonate (23 mg, 0.22 mmol, 2eq). The mixture is stirred for 2 hours then allowed to warm to 0°C, and excess cyanogen bromide and sodium carbonate were then added. The reaction was allowed to warm to room temperature with stirring overnight and then concentrated. The residue was dissolved in ethyl acetate, washed with water and brine, dried over magnesium sulfate, filtered and evaporated. The sample was dissolved in DMF, suspended on silica gel, concentrated and then chromatographed on silica gel with methanol/methylene (95/5 increasing to 9/1). The fractions containing product were concentrated and then the residue triturated with ether, collected and dried. ^{1}H NMR (DMSO-d_6) δ 8.40(1H, s), 7.95(1H, s), 7.85(1H, d), 7.72(1H, d), 7.50(1H, d), 7.29(1H, m), 6.90(1H, s), 6.75(1H, d), 4.98(1H, m), 4.78(2H, s), 4.65(3H, m), 4.25(2H, d), 3.83(3H, m), 3.32(1H, m), 2.75(2H, m), 1.30(6H, d) MS m/e = 481 $(M+1)^+$

Example 31. To a stirred solution of intermediate I-8 (3.46 g, 7.1 mmol) in DMF (30 mL) under nitrogen was added vinyl tributyltin (3.11 mL, 0.0113 mol) and dichlorobis(triphenylphosphine)-palladium (0.49 g, 0.00071 mol). The mixture was heated at reflux overnight then cooled to room temperature, filtered through celite and concentrated *in vacuo* to a dark oil. This oil was triturated with ether and the resulting

precipitate was collected and purified by flash chromatography on silica gel using hexane/ ethyl acetate (1:1) then ethyl acetate (100%) to give a yellow solid (2.3 g, 63% yield). MS (ESI): m/e 515 $(M+1)^+$. To a stirred solution of this product (0.44 g, 0.855 mmol) in THF (10 mL) at room temperature under nitrogen was added osmium tetroxide 8.55 mL, 0.855 mmol, 0.1 M solution) and pyridine (0.55 mL, 6.84 mmol). The reaction mixture was stirred at room temperature overnight. The mixture was diluted with methylene chloride and washed with aqueous sodium bicarbonate solution. The organic phase was dried over magnesium sulfate, filtered, and concentrated *in vacuo* to a yellow solid which was purified by flash chromatography on silica gel to give a yellow solid (0.33 g, 72% yield). MS (ESI): m/e 549 $(M+1)^+$. To a stirred suspension of this diol (83 mg, 0.151 mmol) in benzene (8 mL) was added 1,1'-carbonyldiimidazole (48.9 mg, 0.302 mmol). The reaction mixture was heated at reflux overnight then cooled to room temperature. The solvent was removed *in vacuo* to give a tan solid that was triturated with water and collected by filtration. The solid was purified by preparative TLC chromatography on silica gel using 5% methanol/ chloroform to give a light tan solid (25% yield). ^{1}H NMR (DMSO-d_6): δ 2.73 (m, 2H), 3.25 (m, 4H), 3.82 (s, 3H), 3.85 (m, 2H), 4.35 (s, 2H), 4.64 (t, 1H), 4.75 (s, 2H), 4.94 (t, 1H), 6.11 (t, 1H), 6.82 (d, 1H), 6.89 (s, 1H), 7.15 (d, 3H), 7.63 (d, 1H), 7.82 (d, 1H), 7.90 (d, 1H), 8.12 (s, 1H), 8.46 (s, 1H), MS (ESI): m/e 575 $(M+1)^+$, 597 $(M+Na)^+$.

Example 32. To a stirred suspension of aluminum chloride (0.23 g, 1.76 mmol) in 1,2-dichloroethane (5 mL) at room temperature under nitrogen was added 3-oxabicyclo[3.1.0]hexane-2,4-dione (0.19 g, 1.76 mmol) in 1,2-dichloroethane (3 mL). The mixture gradually became homogeneous and a suspension of I-4 (0.25 g, 0.705 mmol) in 1,2-dichloroethane (3 mL) was added. The reaction mixture was heated to 60 °C overnight. Additional acylating agent (2 equivalents) was prepared and added to the reaction mixture. The mixture continued to be heated at 60 °C for an additional hour. The mixture was poured over ice and concentrated HCl was added. The resulting precipitate was collected by filtration and purified by preparative TLC chromatography on silica gel using 10% methanol/chloroform to give a yellowish-tan solid (70% yield). ^{1}H NMR (DMSO-d_6): δ 1.24 (m, 1H), 1.58 (m, 1H), 2.31 (m, 1H), 2.41 (m, 1H), 2.86 (m, 2H), 3.03

(m, 2H), 3.82 (s, 3H), 4.86 (q, 2H), 6.83 (d, 1H), 6.92 (s, 1H), 7.62 (d, 1H), 8.13 (d, 1H), 8.20 (d, 1H), 8.45 (s, 1H), 8.62 (s, 1H), 12.02 (s, 1H); MS (ESI): m/e 467 $(M+1)^+$, 489 $(M+Na)^+$.

Examples 33-36 were prepared using the general method of Example 32.

Example 33. MS (ESI): m/e 467 $(M+1)^+$

Example 34. MS (ESI): m/e 407 $(M+1)^+$

Example 35. MS (ESI): m/e 479 $(M+1)^+$

Example 36. MS (ESI): m/e 505 $(M+1)^+$

Example 37. This compound was prepared by sodium borohydride reduction of Example 36. MS (ESI) m/e 439 $(M+1)^+$.

Example 38. This compound was prepared by treating Example 37 with TFA and triethylsilane in CH_2Cl_2. MS (ESI) m/e 423 $(M+1)^+$.

Example 39: Bromo intermediate I-9 (68 mg, 0.139 mmol), tributylstannyl dioxene (104 mg, 0.277 mmol), and bis(triphenylphosphine)palladium dichloride (5 mg) were combined in 2 mL anhydrous N,N-dimethylformamide and heated at 80 °C for 16 h. The mixture was concentrated onto 700 mg silica and applied to a bed of silica. Medium pressure liquid chromatography employing a gradient of 1-3% methanol:dichloromethane afforded 47 mg of a yellow solid. NMR (DMSO-d_6) δ 9.55 (d, 1H), 8.6 (s, 1H), 7.95 (s, 1H), 7.6-7.75 (m, 3H), 7.3-7.45 (m, 2H), 7.05 (s, 1H), 5.0 (s, 2H), 4.75 (t, 2H), 4.58 (s, 2H), 4.3 (br s, 2H), 4,15 (m, 4H), 2.15 (m, 2H), 2.05 (s, 3H). MS (ES+): 495 (M+1).

Example 40: Approximately 40 mg of Example 39 was stirred for three hours and then heated to reflux for two hours with 40 mg potassium carbonate in 50 mL methanol:water (25:1). The solution was concentrated to approximately 5 mL. 10 mL water was added and the solution was cooled to 0 °C and the product collected by filtration. The white-yellow solid was washed with water and dried in vacuo to afford 28 mg of the product. NMR (DMSO-d_6) δ 9.55 (d, 1H), 8.6 (s, 1H), 7.95 (s, 1H), 7.6-7.75 (m, 3H), 7.35-7.5 (m,

2H), 7.05 (s, 1H), 5.0 (s, 2H), 4.8 (br s, 2H), 4.6 (s, 2H), 4.35 (br s, 2H), 4.17 (br s, 2H), 3.53 (br s, 2H), 2.0 (m, 2H).

Example 41. In a sealed reaction tube, bromo intermediate I-10 (50 mg, 0.109 mmol) in DMF (1 mL) was added bis(triphenylphosphine)palladium(II) chloride (4 mg, 0.005 mmol, 5 mol%) and 2-(tributylstannyl)-5,6-dihydro-[1,4]-dioxin (82 mg, 0.218 mmol) and heated to 90 °C overnight. The DMF was removed at reduced pressure and the residue dissolved in ethyl acetate, washed with sodium bicarbonate, brine, and then dried over magnesium sulfate. The drying agent was removed by filtration and the solvent evaporated. The product was purified by chromatography on silica gel using ethyl acetate/hexanes (70% increasing to 100% ethyl acetate). The fractions containing product were concentrated and the solid obtained was dried at 80 °C for 12 hours. MS m/e 467 $(m+1)^+$.

Example 42. This compound was prepared in the same manner as Example 41 using bromo intermediate I-11. ^{1}H NMR (DMSO-d_6) 11.90(1H, s), 9.22(1H, d), 8.41(1H, s), 7.92(1H, m), 7.53(2H, s), 7.28(1H, s), 6.98(2H, m), 4.92(2H, s), 4.30(2H, m), 4.10(4H, m), 3.89(3H, s). MS m/e 425 $(m+1)^+$.

Example 43. To intermediate I-12 (100 mg) in DMF (1 mL) was added palladium hydroxide (5 mg) and one drop of concentrated HCl. The reaction mixture was hydrogenated on a Parr apparatus at 50 psi for 5 hours. The catalyst was removed by filtration and the solvent concentrated in vacuo. The solid obtained was triturated with ether, collected and dried. MS m/e 496 $(M+1)^+$.

Example 44. This compound was prepared in a similar manner as Example 40 using intermediate I-9 and 1,2,5,6-tetrahydropyridine-4-boronic acid. MS m/e 450 $(M+1)^+$.

Example 45. This compound was prepared from Example 44 by hydrogenation over palladium on carbon. MS m/e 452 $(M+1)^+$.

Example 46-48 were prepared from Example 45 by treatment with the appropriate electrophile.

Example 46. MS m/e 452 $(M+1)^+$.

Example 47. MS m/e 530 $(M+1)^+$.

Example 48. MS m/e 559 $(M+1)^+$.

Example 49.

Step A: To a well-stirred suspension of intermediate I-13 (4.4 g, 11.9 mmol) in a mixture of 1-methyl-2-pyrrolidinone (51 mL) and methylene chloride (51 mL) were added sequentially triethylamine (2.99 g, 29.6 mmol), a catalytic amount of 4-(dimethylamino)pyridine (0.2 g) and acetic anhydride (3.04 g, 29.8 mmol) at room temperature. After 16 hours at room temperature, the reaction was quenched with water (150 mL) and the methylene chloride was removed under vacuum. The solid was filtered, washed with water and dried to provide product (4.5 g, 90% yield). ^{1}H NMR, (DMSO-d_6) δ: 2.01 (s, 3H), 2.06-2.18 (m, 2H), 4.06-4.16 (m, 2H), 4,51 (s, 2H), 4.75-4.87 (m, 2H), 4.92 (s, 2H), 7.27-7.78 (m, 6H), 8.00 (d, 1H), 8.54 (s, 1H), 9.51 (d, 2H).

Step B: To a well stirred suspension of the product from step A (0.5 g, 1.21 mmol) and 1,1-dichloromethyl methylether (1.39 g, 12.19 mmol) in a mixture of methylene chloride (42 mL) and toluene (5 mL) was added tin(IV) chloride (2.37 g, 9.11 mmol), (1 M solution in methylene chloride) at room temperature. The reaction was further stirred at room temperature for 4 h and at 50 ^{0}C for 45 min., then quenched the reaction at room temperature with aqueous 2N HCl (25 mL). Methylene chloride was removed under vacuum and the aqueous layer was triturated with *t*-butylmethyl methyl ether (30 mL) at room temperature for 14 h. The solid was filtered, washed with water and *t*-butylmethyl methyl ether then dried to obtain the crude product, which was triturated with *t*-butylmethyl methyl ether (20 mL) to afford product, (0.43 g, 81% yield,). ^{1}H NMR, (DMSO-d_6) δ: 2.00 (s, 3H), 2.15-2.27 (m, 2H), 4.10-4.16 (m, 2H), 4.48 (s, 2H), 4.76-4.90 (m, 2H), 4.95 (s, 2H), 7.33-7.45 (m, 2H), 7.64 (d, 1H), 7.89 (d, 1H), 8.04 (d, 1H), 8.51 (s, 1H), 8.64 (s, 1H), 9.51 (d, 1H), 10.11 (s, 1H); MS: m/e 439 (M+1).

Step C: A mixture of the product from step B (0.7 g, 1.6 mmol), N-methylhydroxylamine. HCl (1.06 g, 12.7 mmol), and sodium carbonate (1.18 g, 11.1 mmol) in 1-methyl-2-pyrrolidinone (13 mL) was heated to 80 °C for 18h. The reaction

was quenched with water (80 mL) at room temperature and the solid was filtered, washed with water and dried to obtain product (0.68 g, 91% yield). ^{1}H NMR, (DMSO-d_6) δ: 2.01 (s, 3H), 2.18-2.50 (m, 2H), 3.82 (s, 3H), 4.10-4.13 (m, 2H), 4.53 (s, 2H) 4.78-4.79 (m, 2H), 4.91 (s, 2H), 7.39-7.44 (m, 2H), 7.66-7.68 (d, 1H), 7.79 (d, 1H), 8.00 (s, 1H), 8.31 (d, 1H), 8.59 (s, 1H), 9.16 (s, 1H), 9.50-9.52 (d, 1H); MS m/e 468 (M+1).

Step D: A mixture of the product from step C (0.1 g, 0.21 mmol) and allyl acetate (4 mL) in 1-methyl-2-pyrrolidinone (1.5 mL) was heated to 100 °C for 20 h and the allyl acetate was removed under vacuum. The reaction was quenched with water (74 mL) and the solid was filtered, washed with water, then dried to provide the crude product, which was triturated with a mixture of tetrahydrofuran (1 mL) and diethyl ether (20 mL) to furnish Example 49 (0.08 g, 66% yield); MS m/e 568 (M+1).

Example 50: To a well stirred solution of Example 49 (0.195 g, 0.34 mmol) in tetrahydrofuran (1 mL) was added methanolic ammonia (25 mL) then heated to 80 °C for 14 h. The reaction was concentrated to ~2 mL and ethyl ether was added slowly. The solid was filtered, washed with diethyl ether and dried to provide Example 50 (89 mg, 53% yield); MS m/e 484 (M+1).

Example 51: A mixture of the product from step C, Example 49 (0.171 g, 0.364 mmol) and acrylonitrile (5 mL) in 1-methyl-2-pyrrolidinone (1 mL) was heated to 100 °C for 14 h. Acrylonitrile was removed under vacuum and the crude product was purified by silica gel column chromatography to obtain the required product (0.150 g, 79% yield). The product was suspended in tetrahydrofuran (2 mL) and treated with methanolic ammonia (15 mL) then heated to 70 °C for 24 hour. Methanol was removed under vacuum and the crude material was purified by silica gel column chromatography to obtain product (25 mg, 18% yield). MS: m/e 479 (M+1).

General procedure for Examples 52-53.

A mixture of phenol intermediate I-14 (0.05 mmol.), isocyanate (0.05 mmol), cesium hydrogen carbonate (0.5 mg) and tetrahydrofuran (0.5 mL) was stirred at room temperature for 1 day. The solvent was evaporated and the residue stirred for 8 hours with

ethyl acetate and 3N HCl. The ethyl acetate was removed by evaporation and the aqueous solution was decanted from the solid. The residue was triturated with methanol and the product collected.

Example 52. (79%) MS m/e 546 (M+1); ^{1}H NMR (DMSO-d_6) 11.60 (s, 1H), 8.34 (s, 1H), 8.16 (d, 1H), 7.80 (t, 1H), 7.59 (s, 1H), 7.52 (d, 1H), 7.34 (m, H), 7.26 (m, H), 7.19 (m, H), 4.76 (s, 2H), 4.68 (m, 1H), 3.00 (m, 2H), 2.83 (t, 4H), 2.66 (t, 2H), 1.31 (d, 6H).

Example 53. MS m/e 511 (M+1); ^{1}H NMR (DMSO-d_6) 11.61 (s, 1H), 8.33 (s, 1H), 8.30 (t, 1H), 8.16 (d, 1H), 7.67 (s, 1H), 7.57 (d, 1H), 7.50 (d, 1H), 7.33 (d, 2H), 7.19 (d, 2H), 6.86 (s, 1H), 6.78 (d, 1H), 4.77 (s, 2H), 4.69 (m, 1H), 4.28 (d, 2H), 4.18 (d, 1H), 3.00 (t, 2H), 2.82 (t, 2H), 1.31 (d, 6H).

Synthesis of amine intermediate I-15: To a stirred suspension of bromo-intermediate I-8 (2.44 g, 6.27 mmol) in benzene (180 mL)/ NMP (15 mL) was added p-toluenesulfonic acid (1.19 g, 6.27 mmol) and 4,4'-dimethoxybenzhydrol (1.84 g, 7.5 mmol). The reaction mixture was heated to reflux for 18 h then cooled to room temperature. The reaction solvent was removed *in vacuo*. The solid was dissolved in ethyl acetate and washed with sodium bicarbonate, water and brine. The organic phase was dried over magnesium sulfate, filtered and concentrated *in vacuo* to give the product (4.44 g). The product (2.28 g, 3.7 mmol) was placed in a schlenk tube along with dichlorobistriphenylphosphine palladium (210 mg, 0.3 mmol), triethylamine (1 mL) and 2-methoxyethanol (100 mL). Carbon monoxide was added and the tube was sealed and heated to 150 °C. After 4 h, additional carbon monoxide was added and the mixture continued to be heated. This was repeated one more time. After a total reaction time of 24 h, the mixture was cooled to room temperature and filtered through celite. The celite was washed with THF and the filtrate concentrated *in vacuo*. The solid was purified by flash chromatography on silica gel to give product.

Preparation of acid: To a stirred solution of ester (1 equiv.) at room temperature was added lithium hydroxide (3 equiv.) in one portion. The reaction mixture was heated to 60 °C for 48 h. The mixture was cooled to room temperature, diluted with ethyl acetate,

and washed with 10% HCl and water. The organic phase was dried over magnesium sulfate, filtered and concentrated *in vacuo* to product. MS (ESI): m/e 687 $(M+1)^+$. To a stirred solution of acid (1 equiv.) in benzene was added triethylamine (1.96 equiv.) followed by diphenylphosphoryl azide (2 equiv.). The reaction mixture was heated to reflux for 3 h. The amine (2 mL) was added and the mixture cooled to room temperature. The reaction solvent was removed in vacuo leaving an oil. The material was purified by preparative plate chromatography on silica gel using 5% methanol/chloroform to give I-16. Debenzylation was accomplished using palladium hydroxide in methanol with catalytic concentrated HCl under 50 psi hydrogen for 48-96 h to give intermediate I-15.

Examples 54-58 were prepared using intermediates I-15 or I-16.

Example 54. (71% yield) ^{1}H NMR (DMSO-d_6): δ 1.2-1.35 (m, 5H), 1.57 (m, 1H), 1.73 (m, 2H), 1.86 (,. 2H), 2.82 (m, 2H), 3.50 (m, 2H), 3.84 (m, 5H), 4.64 (m, 2H), 4.73 (s, 2H), 6.06 (m, 1H), 6.80 (d, 1H), 6.89 (s, 1H), 7.41 (d, 1H), 7.58 (d, 1H), 7.91 (d, 1H), 8.08 (s, 1H), 8.32 (d, 2H).

Example 55. (92% yield) ^{1}H NMR (DMSO-d_6): δ 2.75 (m, 2H), 3.30 (m, 2H), 3.75 (s, 6H), 3.80 (m, 2H), 3.81 (s, 3H), 4.30 (d, 2H), 4.35 (s, 2H), 4.62 (s, 2H), 4.78 (m, 2H), 6.26 (s, 1H), 3.39 (s, 1H), 6.49 (t, 1H), 6.69 (s, 1H), 6.82 (d, 1H), 6.90 (s, 1H), 6.95 (d, 4H), 7.01 (m, 2H), 7.14 (d, 3H), 7.19 (d, 4H), 7.58 (m, 2H), 7.73 (s, 1H), 7.90 (d, 1H), 8.48 (s, 1H).

Example 56. MS (ESI) m/e 630 $(M+1)^+$.

Example 57. (52% yield) ^{1}H NMR (DMSO-d): δ. 2.75 (m, 2H), 3.74 (s, 6H), 3.29 (m, 2H), 3.82 (s, 5H), 3.84 (m, 2H), 4.36 (s, 2H), 4.65 (s, 2H), 4.83 (m, 2H), 6.70 (s, 1H), 6.84 (d, 1H), 6.91 (s, 1H), 6.96 (d, 4H), 7.01 (m, 2H), 7.15 (m, 4H), 7.22 (d, 4H), 7.32 (s, 1H), 7.67 (s, 1H), 7.70 (d, 1H), 7.80 (d, 1H), 7.91 (d, 1H), 8.04 (s, 1H), 10.1 (s, 1H).

Example 58. light gray-brown solid (51% yield). ^{1}H NMR (DMSO-d_6): δ 1.09 (t, 6H), 2.79 (m, 2H), 3.40 (m, 2H), 3.70 (m, 2H), 3.82 (s, 3H), 4.60 (m, 2H), 4.73 (s, 2H), 5.75 (s, 2H), 6.80 (d, 1H), 6.89 (s, 1H), 7.59 (d, 2H), 7.89 (d, 1H), 8.15 (s, 1H), 8.32 (s, 1H); MS (ESI): m/e 500 $(M+1)^+$.

Example 59. A mixture of intermediate I-17 (46.3 mg, 0.1 mmol), Pd(dibenzylideneacetone)$_2$ (2.87 mg, 0.005 mmol), P(t-Bu)$_3$ (9.9 uL, 0.04 mmol), sodium *t*-butoxide (14.4 mg, 0.15 mmol) and pyrrolidine (13 uL, 0.15 mmol) in 2.0 mL of xylene was refluxed under argon for 72 hour. The reaction was monitored by HPLC. The reaction was diluted with CH_2Cl_2, filtered through celite and washed with CH_2Cl_2. The solvent was concentration and the product purified by flash chromatography using 70% EtOAc in hexane to provided 7.5 mg (17%) of the 6-cyano-8-fluoro-2-pyrrolidin-1-yl-12,13-dihydro-11H-11-aza-indeno[2,1-a]phenanthrene-5-carboxylic acid ethyl ester. MS: 452 m/e $(M-H)^-$. A mixture of cyanoester intermediate (6 mg, 0.013 mmol), Raney-Ni (20 mg) in 1.5 mL of DMF and 0.15 mL of MeOH was hydrogenated under 50 psi H_2 on a Parr apparatus for 1 week at room temperature. The reaction was monitored by HPLC. The catalyst was removed by filtration and the solvent concentrated to afford 3.5 mg (66%) of the product lactam. MS m/e 412 $(M+1)^+$.

Example 60. A mixture of intermediate I-14 (16.5 mg, 0.041 mmol) and cesium carbonate (88 mg, 1.1 eq) in 2.0 mL of CH_3CN was added cyclopentyl bromide (8.0 uL, 2.0 eq.) under N_2. After stirred at 70 °C for 24 hours, the mixture was diluted with CH_2Cl_2 and filtered through celite and concentrated. Purification by preparative TLC plate with CH_2Cl_2/MeOH afforded the product (4.0 mg, 23%); MS: 467 m/e $(M+1)^+$.

General Methods for Synthesis of Examples 61-67.

Method A: To a mixture of hydroxy intermediate (0.2 mmol), potassium iodide (3.3 mg, 0.1 eq.), *N*-tetrabutylammonium bromide (0.1 eq), cesium hydroxide hydrate (3 eq) and 20 mg of 4Å sieves in 2.0 mL of CH_3CN was added the appropriate alkyl bromide or alkyl iodide under N_2. After the mixture was stirred at 50 °C for 14-72 hours, the reaction mixture was diluted with CH_3CN and filtered through celite and concentrated. The

residue was diluted with CH_2Cl_2 and washed with water and dried over magnesium sulfate. Purification by preparative TLC plate or crystallization with CH_2Cl_2/MeOH afforded the desired products.

Method B: To a mixture of hydroxy intermediate (0.2 mmol) and cesium carbonate (3 eq) in 2.0 mL of CH_3CN was added the appropriate alkyl bromide or iodide under N_2. After the mixture was stirred at 50-80 °C for 14-72 hours, the mixture was diluted with CH_3CN, filtered through celite and concentrated. The residue was diluted with CH_2Cl_2 and washed with water and dried over magnesium sulfate. Purification by preparative TLC plate or crystallization with CH_2Cl_2/MeOH afforded the desired product.

Method C: To a mixture of hydroxy intermediate (0.1 mmol), sodium hydroxide (1.5 eq.) and *N*-tetrabutylammonium bromide (0.1 eq) in 0.5 mL of CH_2Cl_2 and 0.5 mL of water was added the appropriate alkyl bromide under N_2. After the mixture was stirred at room temperature for 14-72 hours, the reaction mixture was concentrated and the residue was washed with water and dried over magnesium sulfate. Purification by preparative TLC plate with CH_2Cl_2/MeOH or crystallization afforded the desired product.

Example 61. Method A; phenol I-18 and cyclopentyl bromide; 14 hr; preparative TLC (10% MeOH in CH_2Cl_2); yield 10%; MS: m/e 453 m/e $(M+1)^+$.

Example 62. Method A; phenol I-18 and cyclohexyl iodide; 40 hr; preparative TLC (10% MeOH in CH_2Cl_2); yield 10%; MS: m/e 467 m/e $(M+1)^+$.

Example 63. Method B; phenol I-18 and 3-bromocyclohexene; 48 hr at 80 °C; preparative TLC (10% MeOH in CH_2Cl_2); yield 19%; MS: m/e 487 m/e $(M+Na)^+$.

Example 64. Method A phenol I-18 and dimethyl bromomalonate; 48 hr at 80 °C; preparative TLC (10% MeOH in CH_2Cl_2); yield 18%; MS: m/e 537 m/e $(M+Na)^+$.

Example 65. Method A; phenol I-18 and 4-bromo-1,7-diethoxy-heptane; 24 hr at 80 °C; preparative TLC (10% MeOH in CH_2Cl_2); yield 3%; MS: m/e 571 m/e $(M+1)^+$.

Example 66 and Example 67. Method A; phenol I-19 and cyclopentyl bromide; 20 hr at 50 °C; preparative TLC (10% MeOH in CH_2Cl_2); yield Example 66 (5%); MS: 499 m/e $(M+Na)^+$; Example 67 (20%); MS 431 m/e $(M+Na)^+$.

Example 68. To a stirred suspension of aluminum chloride (0.35 g, 0.0026 mol) in methylene chloride (8 mL) under nitrogen was added 3-carbomethoxypropionyl chloride (0.323 mL, 0.0026 mol). The mixture gradually became homogeneous and a suspension of intermediate I-13 (0.32 g, 0.00087 mol) in 1,2-dichloroethane (4 mL) was added. The mixture was heated to reflux overnight. Additional acylating agent (1 equiv.) was prepared and added to the reaction mixture. After heating an additional 3 h, the mixture was cooled to room temperature. The reaction mixture was poured over ice, concentrated HCl was added along with additional methylene chloride. The mixture was stirred for 30 minutes then extracted with methylene chloride, dried over magnesium sulfate, and concentrated *in vacuo* to give a tan solid (0.41 g, 78% yield). To a stirred solution of the above compound (100 mg, 0.167 mmol) in THF/H_2O (4 mL, 3:1) was added sodium borohydride (12.6 mg, 0.334 mmol) in one portion. The mixture was stirred at room temperature overnight. The reaction solvent was removed and additional water was added to the flask. The resulting precipitate was collected by filtration. Further purification was accomplished using preparative plate chromatography on silica gel using 5% methanol/ chloroform as the developing solvent yielding a white solid (52 mg, 52% yield). MS (ESI): m/e 599 $(M+1)^+$. To a stirred solution of the previous product (25 mg, 0.0418 mmol) in THF (6 mL) under nitrogen was added sodium hydride (5 mg, 0.125 mmol) in one portion. The mixture was stirred at 0 °C initially then was warmed to room temperature. The yellow reaction mixture was diluted with dichloromethane and washed with water. The organic layer was dried over magnesium sulfate, filtered and concentrated *in vacuo* to a light yellow solid. The solid was purified by preparative plate chromatography developed in 5% methanol/ chloroform (12.6 mg, 53% yield). ^1H NMR ($CDCl_3$): δ 0.93 (m, 2H), 2.16 (m, 2H), 2.38 (m, 1H), 2.64 (s, 5H), 2.77 (m, 3H),2.79 (m,

2H), 3.71 (s, 3H), 4.01 (s, 2H), 4.15 (m, 2H), 4.46 (s, 2H), 4.52 (m, 2H), 5.72 (m, 1H), 7.36 (t, 1H), 7.49 (m, 4H), 7.62 (s, 1H), 9.43 (broad s, 1H); MS (ESI): m/e 567 $(M+1)^+$.

Example 69. To a suspension of aluminum chloride (0.17 g, 1.24 mmol) in 1,2-dichloroethane (5 mL) was added 2,2-dimethylsuccinic anhydride (0.16 g, 1.98 mmol). The mixture was stirred for 20 min. and a suspension of intermediate I-4 (0.22 g, 9.88 mmol) in 1,2-dichloroethane (4 mL) was added. The mixture was heated to 70 °C in an oil bath overnight. Additional acylating agent (2 equiv.) was added and the continued to be heated for 1 h. The reaction mixture was cooled to room temperature and poured over ice. Concentrated HCl (2 mL) was added and a precipitate formed. The solid was collected by filtration to give a tan solid (0.25 g, 85% yield). MS (ESI): m/e 483 $(M+1)^+$. To a stirred solution of the previous product (183 mg, 0.38 mmol) in THF (2 mL) /methanol (1 mL) at room temperature under nitrogen was added trimethylsilyldiazomethane dropwise (0.5 mL). The reaction mixture became cloudy and vigorous evolution of gas was observed. The mixture was stirred at room temperature for 1 h. The reaction solvent was removed *in vacuo* leaving a yellow film. The film was triturated with methylene chloride/ methanol and the solid collected by filtration. The solid was purified by preparative plate chromatography on silica gel to give a pale yellow solid (180 mg, 96% yield). MS (ESI): m/e 497 $(M+1)^+$. To a stirred solution of the previous product (180 mg, 0.362 mmol) in THF (5 mL) under nitrogen was added lithium borohydride (1.45 mL, 1.45 mmol) dropwise. The mixture stirred at room temperature for 4 h. The mixture was cooled and quenched with water. Vigorous evolution of gas was observed. After stirring at room temperature for 1 h, the reaction solvent was removed *in vacuo* leaving a white solid which was triturated with water. The solid was collected by filtration and purified by preparative plate chromatography on silica gel using 10% methanol/chloroform to give a tan solid (90.3 mg, 53% yield). MS (ESI): m/e 471 $(M+1)^+$. To a stirred suspension of the previous product (40 mg, 0.085 mmol) in methylene chloride (5 mL) at room temperature under nitrogen was added trifluoroacetic acid (6.5 µl, 0.085 mmol) dropwise. The mixture was stirred at room temperature for 4 h. The reaction solvent was removed *in vacuo* leaving a tan-brown solid which was purified by preparative plate chromatography on silica gel using 5% methanol/chloroform to give a tan solid (28 mg, 74% yield). ^{1}H NMR

(DMSO-d_6): δ 1.19 (d, 6H), 1.73 (t, 1H), 2.18 (m, 1H), 2.84 (m, 2H), 3.01 (m, 2H), 3.59 (d, 1H), 3.75 (d, 1H), 3.82 (s, 3H), 4.79 (s, 2H), 5.16 (m, 1H), 6.82 (d, 1H), 6.89 (s, 1H), 7.44 (d, 1H), 7.53 (d, 1H), 7.87 (s, 1H), 8.19 (d, 1H), 8.33 (s, 1H), 11.56 (s, 1H); MS (ESI): m/e 453 $(M+1)^+$.

Example 70. This compound was prepared using the general procedure of Example 68. To a stirred solution of the acyl intermediate from Example 69 (0.15 g, 0.31 mmol) in THF/ H_2O (3:1, 8 mL) under nitrogen was added sodium borohydride (47 mg, 1.24 mmol) in one portion. An oily precipitate formed on the walls of the flask. The mixture was stirred at room temperature overnight. The reaction mixture was cooled to 0 °C in an ice bath and methanol was added slowly to quench the reaction. The reaction solvent was removed *in vacuo* leaving a white solid that was collected by filtration and purified by preparative plate chromatography on silica gel using 10% methanol/chloroform to give the product (25 mg, 17% yield). ^{1}H NMR (DMSO-d_6): δ 1.30 (s, 3H), 1.34 (s, 3H), 2.27 (t, 2H), 2.85 (m, 2H), 3.02 (m, 2H), 3.82 (s, 3H), 4.82 (s, 2H), 5.77 (m, 1H), 6.82 (d, 1H), 6.90 (s, 1H), 7.53 (d, 1H), 7.60 (d, 1H), 8.02 (s, 1H), 8.20 (s, 1H), 8.44 (s, 1H), 11.69 (s, 1H); MS (ESI): m/e 467 $(M+1)^+$.

Example 71. This compound was prepared using the general procedure of Example 70. The acylated product was prepared (70 mg, 0.141 mmol) and suspended in THF/ H_20 (8 mL, 3:1) and cooled to 0 °C. Sodium borohydride (16 mg, 0.423 mmol) was added to the reaction mixture in one portion. Vigorous evolution of gas was observed. The mixture was warmed to room temperature overnight. The mixture was extracted with ethyl acetate, dried over magnesium sulfate, filtered, and concentrated *in vacuo* to a white solid. The solid was purified by preparative plate chromatography on silica gel using 5% methanol/ chloroform to give an off-white solid (12 mg, 18% yield). ^{1}H NMR (DMSO-d_6): δ 2.37 (m, 2H), 2.64-2.80 (m, 4H), 2.84 (m, 2H), 3.02 (m, 2H), 3.82 (s, 3H), 4.71 (m, 2H), 4.82 (s, 2H), 5.71 (t, 1H), 6.82 (d, 1H), 6.90 (s, 1H), 7.50 (d, 1H), 7.60 (d, 1H), 7.99 (s, 1H), 8.20 (d, 1H), 8.38 (s, 1H), 11.7 (s, 1H); MS (ESI): m/e 439 $(M+1)^+$.

Example 72. This compound was prepared using the general procedure for Example 70 using 3,3-dimethyl glutaric anhydride. MS (ESI): m/e 481$(M+1)^+$.

Example 73. Prepared by reaction of Example 72 with ethyl iodide and cesium carbonate in acetonitrile. MS (ESI): m/e 509$(M+1)^+$.

Example 74. This compound was prepared using the general procedure for Example 70. MS (ESI): m/e 495 $(M+1)^+$.

Example 75. Following the procedure for Example 9, step A, the acylated product was prepared (100 mg, 0.201 mmol) and suspended in THF (4 mL) at room temperature under nitrogen. Lithium borohydride (0.4 mL, 0.806 mmol, 2 M solution) was added dropwise to the reaction mixture. After stirring at room temperature for 4 h, the mixture was cooled to 0 °C in an ice bath and water was added slowly dropwise to quench the reaction. Evolution of gas was observed and the mixture gradually became homogeneous. The reaction solvent was removed *in vacuo* leaving a white solid which was triturated with water and the resulting precipitate was collected by filtration to give an off-white solid (86 mg, 97% yield). MS (ESI): m/e 443 $(M+1)^+$. To a stirred suspension of the previously prepared diol (80 mg, 0.181 mmol) in 1,2 dichloroethane (4 mL) was added trifluoroacetic anhydride (25.5 µl, 0.181 mmol) dropwise. The mixture became homogeneous and was heated in a sealed tube at 70 °C overnight. The mixture was cooled to room temperature and the solvent removed *in vacuo* leaving a dark solid. The solid was triturated with ether and collected by filtration. The solid was purified by preparative plate chromatography on silica gel using 5% methanol/ chloroform to give an tan solid (12 mg, 16% yield). ^{1}H NMR (DMSO-d_6): δ 1.83 (m, 1H), 2.02 (m, 2H), 2.35 (m, 1H), 2.86 (m, 2H), 3.03 (m, 2H), 3.83 (s, 3H), 3.86 (m, 1H), 4.07 (m, 1H), 4.81 (s, 2H), 4.97 (t, 1H), 6.82 (d, 1H), 6.90 (s, 1H), 7.44 (d, 1H), 7.53 (d, 1H), 7.87 (s, 1H), 8.19 (d, 1H), 8.33 (s, 1H), 11.56 (s, 1H); MS (ESI): m/e 425 $(M+1)^+$.

Example 76. The compound was prepared using the same method as Example 75 using intermediate I-13 and methyl-5-chloro-5-oxovalerate. The product was isolated as an off-

white solid (17.4 mg, 32% yield). ^{1}H NMR (DMSO-d_6) 4.76 (m, 2H), 4.93 (s, 2H), 7.35-7.45 (m, 2H), 7.53 (m, 1H), 7.68 (t, 2H), 7.90 (s, 1H), 8.53 (s, 1H), 9.51 (d, 1H); MS (ESI): m/e 453 $(M+1)^+$.

Example 77. The compound was prepared using the same method as Example 75 using intermediate I-13 and 3-carbomethoxypropionyl chloride to give an off-white solid (8.2 mg, 11%). ^{1}H NMR (DMSO-d_6): δ 2.01 (m, 4H), 2.41 (m, 2H), 3.51 (m, 2H), 3.90 (m, 1H), 4.08 (m, 2H), 4.56 (s, 2H), 4.75 (m, 2H), 4.95 (s, 2H), 4.98 (m, 1H), 7.35 (m, 1H), 7.43 (m, 1H), 7.51 (d, 1H), 7.70 (m, 2H), 7.90 (s, 1H), 8.54 (s, 1H), 9.50 (d, 1H); MS (ESI): m/e 439 $(M+1)^+$.

Example 78. To a stirred suspension of aluminum chloride (0.57g, 4.3 mmol) in 1,2-dichloroethane (6 mL) was added carbomethoxypropionyl chloride (0.529 mL, 4.3 mmol). The reaction mixture became homogeneous and a suspension of intermediate I-40 (0.61g, 1.72 mol) in 1,2-dichloroethane (4 mL) was added to the reaction mixture. The heterogeneous orange mixture was heated to 60 °C in an oil bath overnight. The mixture was poured over ice and concentrated HCl (2 mL) was added. The mixture was stirred at room temperature for 30 minutes then the resulting solid collected by filtration and air-dried overnight. The solid was slurried in methanol/methylene chloride and collected by filtration to give a tan solid (0.53g, 73% yield). MS (ESI): m/e 425 $(M+1)^+$. To a suspension of the above product (260 mg, 0.613 mmol) in THF (5 mL) under nitrogen was added lithium borohydride (1.23 mL, 2M in THF) dropwise and vigorous evolution of gas was observed. When all of the starting material had been consumed by HPLC, the reaction mixture was cooled to 0 °C in an ice bath and water was added very slowly. Vigorous evolution of gas was observed. After the evolution of gas subsided, the reaction mixture became homogeneous. The reaction solvent was removed *in vacuo* and additional water was added. The precipitate was collected by filtration yielding a tan solid (0.18 g, 75% yield). MS (ESI): m/e 399 $(M+1)^+$, 381 $(M-H_2O)^+$. To a suspension of the previous compound (43 mg, 0.108 mmol) in methylene chloride (2.5 mL) in a glass tube was added trifluoroacetic anhydride (15.2 μL, 0.108 mmol). The tube was sealed and heated to 80 °C in an oil bath overnight. The reaction was cooled to room temperature and

the solvent removed *in vacuo* leaving a gray solid. The solid was triturated with ether, collected by filtration and purified by preparative plate chromatography on silica gel developed with 5% methanol/chloroform to give a tan solid (26 mg, 63% yield). ^{1}H NMR (DMSO-d_6): δ 1.86 (m, 1H), 2.02 (m, 2H), 2.36 (m, 1H), 3.85 (dd, 1H), 4.08 (dd, 1H), 4.17 (s, 2H), 4.94 (s, 2H), 5.0 (m, 1H) 7.34 (m, 1H), 7.36-7.56 (m, 2H), 7.57 (d, 1H), 7.68 (d, 1H), 7.90 (s, 1H), 8.50 (s, 1H), 9.39 (d, 1H), 11.87 (s, 1H); MS (ESI): m/e 381 $(M+1)^+$

Example 79. The compound was prepared using the same method as Example 78 with methyl 4-chloro-4-oxobutyrate as the acid chloride. The product was isolated as a tan-yellow solid (36 mg, 34% yield). ^{1}H NMR (DMSO-d_6): δ 1.65 (m, 4H), 1.91 (m, 2H), 3.60 (m, 1H), 4.08 (d, 1H), 4.17 (s, 2H), 4.50 (d, 1H), 4.94 (s, 2H), 7.34 (t, 1H), 7.40-7.47 (m, 2H), 7.56 (d, 1H), 7.68 (d, 1H), 7.90 (s, 1H), 8.50 (s, 1H), 9.39 (d, 1H), 11.86 (s, 1H); MS (ESI): m/e 395 $(M+1)^+$.

Example 80. To a suspension of aluminum chloride (0.27 g, 2.01 mmol) in 1,2-dichloroethane (5 mL) at room temperature under nitrogen was added methyl-5-chloro-5-oxovalerate (0.28 mL, 2.01 mmol) dropwise. The suspension gradually became homogeneous and was stirred for 30 minutes. A suspension of intermediate I-4 (0.31 g, 0.875 mmol) in 1,2-dichloroethane (3 mL) was added to the reaction mixture and then heated to reflux overnight. The mixture was cooled to room temperature and poured over ice. Concentrated HCl (2 mL) was added and the mixture was stirred for 30 min. The resulting precipitate was collected by filtration and dried *in vacuo* to give a light orange solid (37 g, 88% yield). MS (ESI): m/e 483 $(M+1)^+$. To a stirred suspension of the previous product (240 mg, 0.498 mmol) in THF (8 mL) under nitrogen was added lithium borohydride (1.48 mL, 2.97 mmol, 2M soln). Evolution of gas was observed and the mixture stirred at room temperature for 4 h. The mixture was cooled to 0 °C and quenched carefully with water. After stirring at room temperature for 1 h, the reaction solvent was removed *in vacuo* . The resulting solid was triturated with water and collected by filtration to give a white solid (0.21 g, 91% yield). MS (ESI): m/e 457 $(M+1)^+$. To a stirred suspension of the previous product (77 mg, 0.168 mmol) in 1,2-dichloroethane (3 mL) at 0 °C under nitrogen was added trifluoroacetic acid (23.9 µl, 0.169 mmol). The

reaction mixture was stirred at 0 °C for 1 h then warmed to room temperature for 5 h. The reaction solvent was removed *in vacuo* leaving a tan solid which was purified by preparative plate chromatography on silica gel using 10% methanol/chloroform to give a tan solid (46.7 mg, 63% yield). ^{1}H NMR (DMSO-d_6): δ 1.59-1.67 (m, 4H), 1.91 (m, 2H), 2.84 (m, 2H), 3.00 (m, 2H), 3.60 (t, 1H), 4.07 (d, 1H), 4.48 (d, 1H), 4.79 (s, 2H), 6.82 (d, 1H), 6.90 (s, 1H), 7.43 (d, 1H), 7.51 (d, 1H), 7.86 (s, 1H), 8.19 (d, 1H), 8.33 (s, 1H), 11.5 (s, 1H); MS (ESI): m/e 439 $(M+1)^+$.

Example 81. To a stirred suspension of aluminum chloride (0.36 g, 2.69 mmol) in 1,2-dichloroethane (10 mL) under nitrogen was added 3-(carbomethoxy)propionyl chloride (0.33 mL, 2.69 mmol) dropwise. The mixture was stirred at room temperature for 30 min. Intermediate I-41 (0.43 g, 1.08 mmol) was added and the mixture stirred at room temperature for 1 h then warmed to 60 °C overnight. Additional acylating reagent (2 equiv.) was added and the reaction mixture continued to stir at 60 °C overnight. The mixture was cooled to room temperature, poured over ice and concentrated HCl was added. The mixture was stirred for 1 h and the precipitate collected by filtration being washed several times with water. The solid was purified by preparative plate chromatography on silica gel using 5% methanol/ chloroform to give the product (0.58 g, 85% yield). MS (ESI): m/e 627 $(M+1)^+$. To a stirred solution of the previous product (0.39 g, 0.622 mmol) in THF (10 mL) at room temperature under nitrogen was added lithium borohydride (4.9 mL, 4.98 mmol, 1.0 M solution) dropwise. Vigorous evolution of gas was observed and a precipitate began to form. The reaction mixture was cooled to 0 °C and water was carefully added. Following the subsiding of evolution of gas, the reaction solvent was removed *in vacuo* to give a white solid. The solid was triturated with water and collected by filtration to give an off-white solid (0.23g, 93% yield). MS (ESI): m/e 487 $(M+1)^+$. To a suspension of the previous product (130 mg, 0.267 mmol) in 1,2-dichloroethane (4 mL) in a glass tube was added trifluoroacetic acid (37.3 µl, 0.267 mmol) at room temperature. The reaction mixture was stirred for 30 min. at room temperature then heated to reflux for 1h. The mixture was cooled to room temperature and the solvent removed *in vacuo* leaving a grayish-white solid. The solid was purified by preparative plate chromatography on silica gel using 5% methanol/chloroform to give an off-white

solid (60.9 mg, 49% yield). ^{1}H NMR (DMSO-d_6): δ 1.79-1.86 (m, 1H), 2.02 (m, 2H), 2.36 (m, 1H), 2.79 (m, 2H), 3.82 (s, 3H), 3.86 (m, 2H), 4.08 (q, 1H), 4.64 (t, 2H), 4.30 (s, 2H), 5.00 (t, 2H), 6.81 (d, 1H), 6.90 (s, 1H), 7.46 (d, 1H), 7.64 (d, 1H), 7.84 (s, 1H), 7.88 (d, 1H), 8.36 (s, 1H); MS (ESI): m/e 469 $(M+1)^+$.

Example 82. This compound was prepared by the same general procedure as Example 81 and methyl-5-chloro-5-oxovalerate. The product was isolated as an off-white solid (85 mg, 80% yield). ^{1}H NMR (DMSO-d_6): δ 1.59-1.66 (m, 4H), 1.91 (m, 2H), 2.79 (m, 2H), 3.60 (t, 1H), 3.82 (s, 5H), 4.08 (d, 1H), 4.50 (d, 1H), 4.63 (t, 2H), 4.79 (s, 2H), 4.98 (t, 1H), 6.80 (d, 1H), 6.90 (s, 1H), 7.47 (d, 1H), 7.63 (d, 1H), 7.84 (s, 1H), 7.88 (d, 1H), 8.36 (s, 1H); MS (ESI): m/e 483 $(M+1)^+$, 505 $(M+Na)^+$.

Example 83. To a stirred suspension of Example 80 (52 mg, 0.119 mmol) in acetonitrile (5 mL) at room temperature under nitrogen was added cesium carbonate (193 mg, 0.593 mmol) followed by ethyl bromide (44.3 µl, 0.593 mmol). The reaction mixture was heated in a sealed tube at 100 °C for 1 h. The mixture was cooled to room temperature, diluted with methylene chloride and washed with water. The organic phase was dried over magnesium sulfate, filtered and concentrated *in vacuo* to a yellow film. The film was triturated with ether and the solid collected by filtration. The solid was purified by preparative plate chromatography on silica gel using 10% methanol/chloroform to give a light-yellow solid (25 mg, 45% yield). ^{1}H NMR (DMSO-d_6) δ 1.38 (t, 3H), 1.53-1.77 (m, 4H), 1.91 (m, 2H), 2.81 (m, 2H), 3.38 (m, 1H), 3.82 (s, 3H), 4.08 (d, 1H), 4.50 (d, 1H), 4.60 (m, 2H), 4.79 (s, 2H), 6.81 (d, 1H), 6.90 (s, 1H), 7.49 (d, 1H), 7.62 (d, 1H), 7.85 (s, 1H), 7.90 (d, 1H), 8.37 (s, 1H); MS (ESI): m/e 467 $(M+1)^+$.

Examples 84 and Example 85. The enantiomers of Example 83 were separated by chiral HPLC using a Chiralpak AD column 4.6 x 250 mm with a flow rate of 0.5 mL/min. with ethanol as solvent. Example 84 eluted with a R_t of 19.1 minutes MS (ESI): m/e 467 $(M+1)^+$. Example 85 eluted with a R_t of 24.7 minutes MS (ESI): m/e 467 $(M+1)^+$.

Examples 86-91 were prepared using the general procedure for Example 69 using the

appropriate cyclic anhydride.

Example 86. MS (ESI): m/e 423 $(M+1)^+$.

Example 87. MS (ESI): m/e 481 $(M+1)^+$.

Example 88. MS (ESI): m/e 481 $(M+1)^+$.

Example 89. MS (ESI): m/e 507 $(M+1)^+$.

Example 90. MS (ESI): m/e 467 $(M+1)^+$.

Example 91. MS (ESI): m/e 423 $(M+1)^+$.

Examples 92-99 were prepared from Example 86 using the method of Example 83.

Example 92. MS (ESI): m/e 451 $(M+1)^+$.

Example 93. MS (ESI): m/e 465 $(M+1)^+$.

Example 94. MS (ESI): m/e 437 $(M+1)^+$.

Example 95. MS (ESI): m/e 493 $(M+1)^+$.

Example 96. MS (ESI): m/e 463 $(M+1)^+$.

Example 97. MS (ESI): m/e 507 $(M+1)^+$.

Example 98. MS (ESI): m/e 479 $(M+1)^+$.

Example 99. MS (ESI): m/e 522 $(M+1)^+$.

Examples 100-103 were prepared from intermediate I-4 and the appropriate cyclic anhydride using the general procedure for Example 69.

Example 100. MS (ESI): m/e 493 $(M+1)^+$.

Example 101. MS (ESI): m/e 481 $(M+1)^+$.

Example 102. MS (ESI): m/e 467 $(M+1)^+$.

Example 103. MS (ESI): m/e 467 $(M+1)^+$.

Examples 104-123 were prepared from Example 80 and the appropriate bromide or iodide using the method for Example 83.

Example 104. MS (ESI): m/e 495 $(M+1)^+$

Example 105. MS (ESI): m/e 453 $(M+1)^+$

Example 106. MS (ESI): m/e 509 $(M+1)^+$.

Example 107. MS (ESI): m/e 523 $(M+1)^+$.

Example 108. MS (ESI): m/e 495 $(M+1)^+$.

Example 109. MS (ESI): m/e 509 $(M+1)^+$.

Example 110. MS (ESI): m/e 479 $(M+1)^+$.

Example 111. MS (ESI): m/e 611 $(M+1)^+$.

Example 112. MS (ESI): m/e 581 $(M+1)^+$.

Example 113. MS (ESI): m/e 497 $(M+1)^+$.

Example 114. MS (ESI): m/e 538 $(M+1)^+$.

Example 115. MS (ESI): m/e 552 $(M+1)^+$.

Example 116. MS (ESI): m/e 665 $(M+1)^+$.

Example 117. MS (ESI): m/e 507 $(M+1)^+$.

Example 118. MS (ESI): m/e 506 $(M+1)^+$.

Example 119. MS (ESI): m/e 481 $(M+1)^+$.

Example 120. MS (ESI): m/e 495 $(M+1)^+$.

Example 121. MS (ESI): m/e 483 $(M+1)^+$.

Example 122. MS (ESI): m/e 493 $(M+1)^+$.

Example 123. MS (ESI): m/e 507 $(M+1)^+$.

Examples 124-133 were prepared by reacting Example 103 with the appropriate bromide or iodide using the method of Example 80.

Example 124. MS (ESI): m/e 639 $(M+1)^+$

Example 125. MS (ESI): m/e 525 $(M+1)^+$.

Example 126. MS (ESI): m/e 509 $(M+1)^+$.

Example 127. MS (ESI): m/e 625 $(M+1)^+$.

Example 128. MS (ESI): m/e 511 $(M+1)^+$.

Example 129. MS (ESI): m/e 495 $(M+1)^+$.

Example 130. MS (ESI): m/e 534 $(M+1)^+$.

Example 131. MS (ESI): m/e 481 $(M+1)^+$.

Example 132. MS (ESI): m/e 521 $(M+1)^+$.

Example 133. MS (ESI): m/e 505 $(M+1)^+$.

Example 134. The phenol intermediate I-19 under Friedel Craft conditions as described for Example 80 formed the C-,O- bis-acylated adduct which was reduced to the diol using lithium borohydride in a similar manner as described in Example 80. To a stirred suspension of diol intermediate (1.19 g, 2.69 mmol) in CH_2Cl_2 (40 mL) was added TFA (0.21 mL, 2.69 mmol) under nitrogen. The reaction mixture was stirred for 2 hours then additional TFA (2 equiv.) was added. The reaction solvent was removed in vacuo leaving an off-white solid that was triturated with ether and collected by filtration. ^{1}H NMR (DMSO-d_6) δ 1.59-1.69 (m, 4H), 1.87-1.91 (m, 2H), 2.78 (m, 2H), 2.97 (m, 2H), 3.59 (m, 1H), 4.07 (d, 1H), 4.48 (d, 1H), 4.78 (s, 2H), 6.63 (d, 1H), 6.70 (s, 1H), 7.42 (d, 1H), 7.50 (d, 1H), 7.85 (s, 1H), 8.06 (d, 1H), 8.31 (s, 1H), 9.44 (s, 1H), 11.51 (s, 1H).

Example 135.

Step 1. A solution of N-ethyl intermediate I-42 (510 mg, 1.24 mmol) in dichloroethane (DCE) (15 ml) was added dropwise to a preformed solution of $AlCl_3$ (1.37 g, 10.31 mmol) and methyl-4-(chloro-formyl)butyrate (1.42 ml, 10.31 mmol) in DCE (10 ml) at room temperature. The reaction was allowed to stir overnight at room temperature and ^{1}H NMR of a worked up and extracted aliquot showed conversion to product. To the stirring solution was slowly added 10 ml of 2N HCl and the reaction allowed to stir at room temperature for 60 minutes. The resulting off white solid was collected by vacuum filtration and washed with ether; 684 mg, 88% yield. MS (ESI): m/e 647 $(M+Na)^+$.

Step 2. To a 0 °C solution of the product from Step 1 (684 mg, 1.24 mmol) in 20 ml of anhydrous THF was added dropwise 6.20 ml (12.36 mmol) of a 2.0 M solution of $LiBH_4$ in THF. The reaction was allowed to stir at room temperature for 60 hours and HPLC of an aliquot of the reaction showed conversion to product. The reaction was evaporated to dryness and the resulting white solid triturated in water and collected by vacuum filtration and air dried; 540 mg, 93% yield. MS (ESI): m/e 477 $(M+H)^+$.

Step 3. To a 0 °C solution of the product from Step 2 (530 mg, 1.13 mmol) in 10 ml of anhydrous DCE was added 87 µl of trifluoroacetic acid dropwise with a syringe. The solution was allowed to warm to room temperature overnight. An HPLC of an aliquot of the reaction showed conversion to product. The reaction was evaporated to dryness and

the resulting solid was triturated with ether and collected by vacuum filtration; 611mg, contains salts from reaction sequence, MS (ESI): m/e 475 $(M+Na)^+$.

General alkylation procedure for Examples 136-146: To a stirred suspension of Example 134 in acetonitrile (0.04-0.06 mM) under nitrogen was added the alkylating agent (3-5 equiv.) followed by cesium carbonate (3-5 equiv.). The mixture was heated to 50-80 °C overnight. When complete, the reaction mixture was cooled to room temperature and concentrated in vacuo to a solid. The solid was triturated with water and collected by filtration. The products were purified by flash column chromatography on silica gel, preparative silica gel plates or preparative HPLC.

Example 136. white solid (14.3 mg, 37% yield). ^{1}H NMR (DMSO-d_6) δ 1.30 (d, 6H), 1.60-1.67 (m, 4H), 1.87-1.90 (m, 2H), 2.90 (m, 2H), 3.00 (m, 2H), 3.59 (m, 1H), 4.07 (d, 1H), 4.48 (d, 1H), 4.69 (m, 1H), 4.79 (s, 2H), 6.78 (d, 1H), 6.87 (s, 1H), 7.43 (d, 1H), 7.51 (d, 1H), 7.86 (s, 1H), 8.16 (d, 1H), 8.34 (s, 1H), 11.55 (s, 1H).

Example 137. (6.52 mg, 14% yield) MS (ESI): m/e 509 $(M+H)^+$.
Example 138. (15.48 mg, 28% yield). MS (ESI): m/e 467 $(M+H)^+$.
Example 139. (15.90 mg, 2.7% yield). MS (ESI): m/e 509 $(M+H)^+$.
Example 140. (9.16 mg, 20% yield). MS (ESI): m/e 481 $(M+H)^+$.
Example 141. (20.8 mg, 17% yield). MS (ESI): m/e 493 $(M+H)^+$.
Example 142. (35 mg, 22% yield). ^{1}H NMR (DMSO-d_6): δ 0.34 (m, 2H), 0.58 (m, 2H), 1.26 (m, 1H), 1.63 (m, 4H), 2.83 (m, 2H), 2.99 (m, 2H), 3.61 (m, 1H), 3.88 (d, 2H), 4.07 (d, 1H), 4.48 (d, 1H), 4.79 (s, 2H), 6.79 (d, 1H), 6.89 (s, 1H), 7.43 (d, 1H), 7.51 (d, 1H), 7.87 (s, 1H), 8.17 (d, 1H), 8.32 (s, 1H).

Example 143. (27.17 mg, 41% yield). MS (ESI): m/e 581 $(M+Na)^+$.
Example 144. MS (ESI): m/e 492 $(M+1)^+$.
Example 145. MS (ESI): m/e 505 $(M+1)^+$.
Example 146. MS (ESI): m/e 465 $(M+1)^+$.
Example 147. MS (ESI): m/e 506 $(M+1)^+$.

Example 148. MS (ESI): m/e 587 $(M + 1)^+$.

Using the general alkylation procedure described for Examples 136-143, Examples 149-158 were prepared from Example 136.

Example 149. (20.73 mg, 56% yield). MS (ESI) m/e 556 $(M+Na)^+$.

Example 150. (12.04 mg, 26% yield). MS (ESI) m/e 517 $(M+Na)^+$.

Example 151. (5.09 mg, 11% yield). MS (ESI) m/e 503 $(M+Na)^+$.

Example 152. (6.94 mg, 13.7% yield). MS (ESI) m/e 545 $(M+Na)^+$

Example 153. (10.5 mg, 21% yield). MS (ESI) m/e 521 $(M+H)^+$.

Example 154. (19.1 mg, 11.7% yield). MS (ESI): m/e 517 $(M+Na)^+$.

Example 155. MS (ESI): m/e 512 $(M + 1)^+$.

Example 156. MS (ESI): m/e 527 $(M + 1)^+$.

Example 157. MS (ESI): m/e 523 $(M + 1)^+$.

Example 158. MS (ESI): m/e 492 $(M + 1)^+$.

Example 159. This compound was prepared according to the procedure for Example 158 where the phenol intermediate I-19 was acylated under Friedel Craft conditions with methyl 4-chloro-4-oxobutyrate and reduced with lithium borohydride. To a stirred suspension of intermediate diol (0.39 g, 0.91 mmol) in methylene chloride (15 mL) under nitrogen was added TFA (0.13 mL, 0.91 mmol) dropwise. The reaction mixture was stirred at room temperature for 4 h. The reaction solvent was removed in vacuo leaving a white solid that was triturated with ether and collected by filtration (0.32 g, 86%). MS (ESI): m/e 441 $(M+H)^+$, 463 $(M+Na)^+$.

Using the general alkylation conditions described for Examples 136-143, Examples 160-165 were prepared from Example 159.

Example 160. (15.8 mg, 49%). ^{1}H NMR (DMSO-d_6): δ 1.84 (m, 1H), 2.01 (m, 2H), 2.36 (m, 1H), 2.78 (m, 2H), 3.39 (m, 2H), 3.80 (s, 3H), 3.85 (m, 1H), 4.08 (m, 1H), 4.13 (s, 3H), 4.79 (s, 2H), 4.98 (dd, 1H), 6.81 (d, 1H), 6.90 (s, 1H), 7.48 (d, 1H), 7.62 (d, 1H), 7.85 (s, 1H), 7.92 (d, 1H), 8.37 (s, 1H).

Example 161. (24.3 mg, 32%). ^{1}H NMR (DMSO-d_6): δ 1.35-1.4 (m, 6H), 1.83 (m, 1H), 2.01 (m, 2H), 2.36 (m, 1H), 2.81 (m, 2H), 3.28 (m, 2H), 3.85 (m, 1H), 4.10 (m, 3H), 4.61 (m, 2H), 4.79 (s, 2H), 4.98 (m, 1H), 6.79 (d, 1H), 6.89 (s, 1H), 7.48 (d, 1H), 7.65 (d, 1H), 7.85 (s, 1H), 7.89 (d, 1H), 8.36 (s, 1H).

Example 162. (14.6 mg, 21%). MS (ESI): m/e 467 $(M+H)^+$.

Example 163. (26 mg, 32%). MS (ESI): m/e 453 $(M+H)^+$.

Example 164. This compound was prepared starting from Example 163 using the general procedure for Examples 148-155 (9.77 mg, 20%). MS (ESI) m/e 517 $(M+Na)^+$.

Example 165. This compound was prepared by alkylation of Example 163 to give a yellow solid (17.8 mg, 89%). MS (ESI): m/e 481 $(M+H)^+$.

Example 166. This compound was prepared in step 3 in Example 100. The ketoester in THF under nitrogen was reduced to the diol using lithium borohydride to give a solid which was collected purified by preparative plate chromatography on silica gel using 10% methanol/chloroform to give a tan solid. MS (ESI): m/e 511 $(M+1)^+$.

Example 167. The keto-acid intermediate from Example 86 (0.3 g, 0.543 mmol) in DMF (3.5 mL) was added HOBt-NH_2 (0.413 g, 2.717 mmol), and EDCI (0.521 g, 2.727 mmol). The resulted mixture was stirred at room temperature for 5 h and quenched with water at 5 °C and the precipitate was further stirred at room temperature for 40 minutes then filtered, washed with water and dried to obtain the amide intermediate (0.28 g, 93% yield, 85% purity); MS m/e 552 (M+1), 574(M+Na).

This intermediate (0.13 g, 0.235 mmol) in tetrahydrofuran (5 mL) was added lithium borohydride (0.092 g, 4.2 mmol) at 5 °C then stirred at room temperature for 2 h and quenched with water at 0 °C. The aqueous layer was extracted with a mixture of tetrahydrofuran and ethyl acetate (1:1 ratio) and the combined organics were washed with

brine, dried and concentrated to obtain the crude product. The crude product was triturated with a mixture of tetrahydrofuran and ethyl acetate at room temperature for 1 h, filtered, washed with ether and dried to provide the hydroxy-amide (0.09 g, 75% yield, 79% purity); MS m/e 534 (M+1), 494 (M-18). To the amide in methylene chloride (25 mL) was added trifluoroacetic acid (8 drops). The resulted mixture was stirred at room temperature for 2.5 h then diluted with ethyl acetate and concentrated under vacuum to furnish the crude product. The crude product was purified by silica gel column chromatography (0.0074 g, 14% yield); MS m/e 494 (M+1).

Example 168. The keto-methyl ester intermediate prepared using the method of Example 78 (0.026 g, 0.057 mmol) in NMP was added hydrazine hydrate (20 drops). The resulted clear solution was heated to 100 °C for 4 hours and quenched with water and the solid was filtered, washed with water and dried, (14 mg, 58% yield); MS m/e 435 (M+1), 457 (M+Na).

Example 169. A mixture of intermediate I-27 (1 eq) and methoxylamine hydrochloride (4 eq) in a mixture of 1-methyl-2-pyrrolidinone and ethyl alcohol was refluxed for 2 h. Ethyl alcohol was removed under vacuum and the reaction mixture was quenched with water and the resulted solid was filtered, washed with water and dried to afford product. MS m/e 440 (M+1).

Example 170. A dried flask containing I-22 (100 mg, 0.20 mmol), Cs_2CO_3 (91 mg, 0.25 mmol), *p*-anisidine (117 mg, 0.95 mmol), *rac*-2,2'-bis(diphenylphosphino)-1,1'-binaphthyl (20 mg, 0.09 mmol), and $Pd(OAc)_2$ (13 mg, 0.06 mmol) was de-gassed three times. Dry xylenes (5 mL) were added and the mixture was again de-gassed three times. After heating to 125 °C for 20 h, the black solution was cooled and the solvent removed in vacuo. Methylene chloride (10 mL) was added, the mixture filtered through Celite, and the mother liquor was chromatographed on silica gel (5% MeOH/CH_2Cl_2). The intermediate yielded (22 mg) and Raney nickel (50 mg) was hydrogenated in 5 mL 9:1 DMF/MeOH at room temperature under 50 psi of hydrogen pressure for 4 days, filtered through Celite, and concentrated in vacuo. The resulting brown solid was triturated with

MeOH to yield 9 mg (9%) as a tan powder: mp>300 °C; ^{1}H NMR (DMSO-d6) δ 1.37 (t, *J* = 6.9 Hz, 3 H), 2.81 (t, *J* = 5.8 Hz, 2 H), 3.27 (t, *J* = 5.8 Hz, 2 H), 3.72 (s, 3 H), 3.82 (s, 3 H), 4.56 (q, *J* = 6.8 Hz, 2 H), 4.70 (s, 2 H), 6.80 (dd, *J* = 2.6, 8.7 Hz, 1 H), 6.86 (d, *J* = 9.1 Hz, 2 H), 6.90 (d, *J* = 2.7 Hz, 1 H), 7.1 (d, *J* = 9.1 Hz, 2 H), 7.21 (dd, *J* = 2.0, 8.7 Hz, 1 H), 7.52 (d, *J* = 2.0 Hz 1 H), 7.56 (d, *J* = 8.8 Hz, 1 H), 7.78 (s, 1 H), 7.89 (d, *J* = 8.6 Hz, 1 H), 8.26 (s, 1 H); MS (*m/e*) 504 (M+).

Example 171. A dried flask containing I-22 (75 mg, 0.15 mmol), Cs_2CO_3 (68 mg, 0.19 mmol), 2,5-dimethoxyaniline (100 mg, 0.71 mmol), *rac*-2,2'-bis(diphenylphosphino)-1,1'-binaphthyl (15 mg, 0.07 mmol), and $Pd(OAc)_2$ (10 mg, 0.04 mmol) was de-gassed three times. Dry xylenes (5 mL) were added and the mixture was again de-gassed three times. After heating to 125 °C for 20 h, the black solution was cooled and the solvent removed in vacuo. Methylene chloride (10 mL) was added, the mixture filtered through Celite, and the mother liquor was chromatographed on silica gel (5% MeOH/CH_2Cl_2). The intermediate yielded (22 mg) and Raney nickel (50 mg) was hydrogenated in 5 mL 9:1 DMF/MeOH at room temperature under 50 psi of hydrogen pressure for 4 days, filtered through Celite, and concentrated in vacuo. The resulting brown solid was triturated with MeOH to yield 6 mg (8%) as a tan powder: mp>300 °C; ^{1}H NMR (DMSO-d6) δ 1.39 (t, *J* = 7.1 Hz, 3 H), 2.81 (t, *J* = 6.7 Hz, 2 H), 3.27 (t, *J* = 6.7 Hz, 2 H), 3.63 (s, 3 H), 3.82 (s, 6 H), 4.57 (q, *J* = 6.5 Hz, 2 H), 4.71 (s, 2 H), 6.30 (dd, *J* = 2.9, 8.7 Hz, 1 H), 6.72 (d, *J* = 2.8 Hz, 1 H), 6.81 (dd, *J* = 2.7, 8.7 Hz, 1 H), 6.89 (d, *J* = 8.8 Hz, 1 H), 6.90 (d, *J* = 3.3 Hz, 1 H), 7.28 (s, 1 H), 7.40 (dd, *J* = 2.1, 8.7 Hz, 1 H), 7.61 (d, *J* = 8.8 Hz, 1 H), 7.73 (d, *J* = 1.8 Hz, 1 H), 7.90 (d, *J* = 8.7 Hz, 1 H), 8.31 (s, 1 H); MS (m/e) 534 (M+).

Example 172. This compound was prepared by the general method for Example 80 except using nitromethane and methylene chloride as a solvent in step 1 and starting from the dihydroindazole intermediate I-20 and methyl-5-chloro-5-oxovalerate. MS (ESI): m/e 413 $(M+1)^+$.

Example 173. This compound was prepared by the general method for Example 75 starting from intermediate I-20. MS (ESI): m/e 399 $(M+1)^+$.

Example 174. This compound was prepared by the general method for Example 100 starting from intermediate I-20. MS (ESI): m/e 467 $(M+1)^+$.

Example 175. This compound was prepared by the general method for Example 91 and Example 102 starting from intermediate I-20. MS (ESI): m/e 441 $(M+1)^+$.

Example 176. This compound was prepared by the general method for Example 103 and Example 86 starting from intermediate I-20. MS (ESI): m/e 441 $(M+1)^+$.

Example 177. This compound was prepared from Example 175 and ethyl iodide by the general method for Example 83. MS (ESI): m/e 469 $(M+1)^+$.

Example 178. This compound was from Example 175 and bromobutyronitrile by the general method for Example 83. MS (ESI): m/e 508 $(M+1)^+$.

Examples 179-242 were prepared from Example 172 and the appropriate bromide or iodide by the general method described from Example 83.

Example 179. MS (ESI): m/e 455 $(M+1)^+$

Example 180. MS (ESI): m/e 441 $(M+1)^+$.

Example 181. MS (ESI): m/e 427 $(M+1)^+$.

Example 182. MS (ESI): m/e 471 $(M+1)^+$.

Example 183. MS (ESI): m/e 453 $(M+1)^+$.

Example 184. MS (ESI): m/e 485 $(M+1)^+$.

Example 185. MS (ESI): m/e 469 $(M+1)^+$.

Example 186. MS (ESI): m/e 457 $(M+1)^+$.

Example 187. MS (ESI): m/e 481 $(M+1)^+$.

Example 188. MS (ESI): m/e 455 $(M+1)^+$.

Example 189. MS (ESI): m/e 469 $(M+1)^+$.

Example 190. MS (ESI): m/e 451 $(M+1)^+$.

Example 191. MS (ESI): m/e 512 $(M+1)^+$.

Example 192. MS (ESI): m/e 467 $(M+1)^+$.

Example 193. MS (ESI): m/e 483 $(M+1)^+$.

Example 194. MS (ESI): m/e 483 $(M+1)^+$.

Example 195. MS (ESI): m/e 498 $(M+1)^+$.

Example 320. MS (ESI): m/e 483 $(M+1)^+$.

Example 196. MS (ESI): m/e 550 $(M+1)^+$.

Example 197. MS (ESI): m/e 497 $(M+1)^+$.

Example 198. MS (ESI): m/e 469 $(M+1)^+$.

Example 199. MS (ESI): m/e 469 $(M+1)^+$.

Example 200.MS (ESI): m/e 473 $(M+1)^+$.

Example 201. MS (ESI): m/e 487 $(M+1)^+$.

Example 202. MS (ESI): m/e 455 $(M+1)^+$.

Example 203. MS (ESI): m/e 455 $(M+1)^+$.

Example 204. MS (ESI): m/e 481 $(M+1)^+$.

Example 205. MS (ESI): m/e 481 $(M+1)^+$.

Example 206. MS (ESI): m/e 481 $(M+1)^+$.

Example 207. MS (ESI): m/e 509 $(M+1)^+$.

Example 208. MS (ESI): m/e 501 $(M+1)^+$.

Example 210. MS (ESI): m/e 497 $(M+1)^+$.

Example 211. MS (ESI): m/e 523 $(M+1)^+$.

Example 212. MS (ESI): m/e 487 $(M+1)^+$.

Example 213. MS (ESI): m/e 497 $(M+1)^+$.

Example 214. MS (ESI): m/e 469 $(M+1)^+$.

Example 215. MS (ESI): m/e 495 $(M+1)^+$.

Example 216. MS (ESI): m/e 466 $(M+1)^+$.

Example 217. MS (ESI): m/e 469 $(M+1)^+$.

Example 218. MS (ESI): m/e 469 $(M+1)^+$.

Example 209. To example 172 (50 mg, 0.121 mmol) was added Ac_2O (2 mL), and the reaction mixture was heated to 140 °C for 2 h. The solution was cooled to room temperature, Et_2O was added, and the yellow solid was collected and dried (33 mg, 60%):

^{1}H NMR (DMSO-d_6) δ 1.75 (m, 9 H), 2.62 (s, 3 H), 2.88 (m, 2 H), 3.25 (m, 2 H), 3.89 (s, 3 H), 5.17 (s, 2 H), 7.44 (d, 1 H), 7.54 (d, 1 H), 7.87 (s, 1 H), 8.72 (s, 1 H), 11.72 (s, 1 H) ppm; MS (m/e) 455 (M^+).

Example 219. This compound was prepared from Example 176 and bromobutyronitrile by the general method for Example 83. MS (ESI): m/e 508 $(M+1)^+$.

Example 220. This compound was prepared from Example 176 and ethyl iodide by the general method for Example 83. MS (ESI): m/e 469 $(M+1)^+$.

Examples 221-227 were prepared from Example 176 using the method described for Example 83.

Example 221. MS (ESI) m/e 497 (M + 1).

Example 222. MS (ESI): m/e 455 $(M+1)^+$.

Example 223. MS (ESI): m/e 481 $(M+1)^+$.

Example 224. MS (ESI): m/e 483 $(M+1)^+$.

Example 225. MS (ESI): m/e 509 $(M+1)^+$.

Example 226. MS (ESI): m/e 501 $(M+1)^+$.

Example 227. MS (ESI): m/e 515 $(M+1)^+$.

Example 228. This compound was prepared from Example 173 and ethyl iodide by the general method for Example 83. MS (ESI): m/e 427 $(M+1)^+$.

Example 229. This compound was prepared from Example 174 and propyl iodide by the general method for Example 86. MS (ESI): m/e 509 $(M+1)^+$.

Example 230. This compound was prepared from Example 174 and ethyl iodide by the general method for Example 83. MS (ESI): m/e 495 $(M+1)^+$.

Example 231. This compound was prepared from Example 174 and bromobutyronitrile by the general method for Example 83. MS (ESI): m/e 534 $(M+1)^+$.

Example 232. This compound was prepared from N-acyl aldehyde intermediate I-21 using the general procedure described for Example 12-28. MS (ESI): m/e 485 $(M+1)^+$.

Example 233. This compound was prepared from Example 232 using the deacylation method described . MS (ESI): m/e 443 $(M+1)^+$.

Example 234. This compound was prepared using the general method described for Example 231. MS (ESI): m/e 513 $(M+1)^+$.

Example 235. This compound was prepared from Example 234 using the method described for Example 233. MS (ESI): m/e 471 $(M+1)^+$.

General procedure for Examples 236-244.

A mixture of intermediate I-24 or I-25 (1 eq) and hydroxylamine hydrochloride or O-alkyl hydroxylamine hydrochloride in a mixture of 1-methyl-2-pyrrolidinone and ethyl alcohol was refluxed for 4 h. Ethyl alcohol was removed under vacuum then quenched with water and the resulted solid was filtered, washed with water and dried to obtain the product.

Example 236. MS m/e 400 (M+1)

Example 237. MS m/e 442 (M+1)

Example 238. MS m/e 453 (M+1)

Example 239. MS m/e 414 (M+1)

Example 240. MS m/e 442 (M+1)

Example 241. MS m/e 426 (M+1)

Example 242. MS m/e 426 (M+1)

Example 243. MS m/e 442 (M+1)

Example 244. MS m/e 428 (M+1).

Example 245. A mixture of intermediate I-24 (1 eq), methanesulfonyl hydrazide (5 eq) and *p*-toluenesulfonic acid monohydrate (1 eq) in a mixture of 1-methyl-2-pyrrolidinone and benzene was refluxed using a Dean-stark trap for 2.5 h. The reaction mixture was

quenched with saturated aqueous sodium bicarbonate solution and extracted thrice with a mixture of ethyl acetate and tetrahydrofuran (1:1 ratio). The combined organic layers were washed with brine, dried and concentrated, and the crude product was triturated with a mixture of tetrahydrofuran, ether and hexane to yield product. MS m/e 463 (M+1)

5

General procedure for Examples 246-250.

A mixture of intermediate methyl ketone intermediate I-26 (1 eq) and O-alkyl hydroxylamine hydrochloride (4 eq) in a mixture of 1-methyl-2-pyrrolidinone and ethyl
10 alcohol was refluxed for 2 h. Potassium carbonate (17 eq) was added at room temperature and heated to 65 °C for 1h. Ethyl alcohol was removed under vacuum, the reaction mixture was then quenched with water and extracted thrice with a mixture of tetrahydrofuran and ethyl acetate (1:1 ratio). The combined organics was washed with brine, dried and concentrated to provide a crude product, which was triturated with a
15 mixture of tetrahydrofuran, ether and hexane to furnish a pure product.

Example 246. MS m/e 470 (M+1)

Example 247. MS m/e 456 (M+1)

Example 248. MS m/e 484 (M+1)

Example 249. MS m/e 498 (M+1)

20 Example 250. MS m/e 509 (M + 1)

Example 251. This compound was prepared from methyl ketone intermediate I-25 using the general procedure for Example 246-250. MS m/e 428 (M + 1)

25 Examples 252-286 were prepared using the appropriate N-alkyl ketone intermediate I-26 by the general oxime synthesis methods described for Examples 246-250.

Example 252. MS m/e 442 (M+1)

Example 253. MS m/e 484 (M+1)

Example 254. MS m/e 470 (M+1)

30 Example 255. MS m/e 456 (M+1)

Example 256. To the intermediate I-32 (25mg, 0.07 mmol) in nitromethane (5 mL) was added 2-thiophene carbonyl chloride (75 µl 0.7 mmol, 10 eq) followed by aluminum chloride (94 mg, 0.7 mmol, 10 eq) added in small portions. The reaction mixture was stirred at room temperature overnight. The reaction was then concentrated, stirred with water and a few drops of 1 N HCl added. The product was collected by filtration, dissolved in methylene chloride/methanol and purified by preparative TLC eluting with 10% methanol/methylene chloride. The desired band was collected, stirred with methylene chloride/methanol, filtered, and concentrated. The sample was dried at 80°C under vacuum overnight. MS m/e 495 (M+1). Example 256 was prepared by from thienyl ketone intermediate and hydroxylamine HCl using the general procedure described for Examples 246-249. MS m/e 544 (M+1)

Example 257. MS m/e 456 (M+1)
Example 258. MS m/e 442 (M+1)
Example 259. MS m/e 456 (M+1)
Example 260. MS m/e 470 (M+1)
Example 261. MS m/e 484 (M+1)
Example 262. MS m/e 498 (M + 1)
Example 263. MS m/e 428 (M+1)
Example 264. MS m/e 442 (M+1)
Example 265. MS m/e 456 (M+1)
Example 266. MS m/e 470 (M+1)
Example 267. MS m/e 490 (M+1)
Example 268. MS m/e 499 (M+1)
Example 269. MS m/e 470 (M+1)
Example 270. MS m/e 484 (M+1)
Example 271. MS m/e 414 (M+1)
Example 272. MS m/e 428 (M+1)
Example 273. MS m/e 456 (M+1)
Example 274. MS m/e 471 (M+1)
Example 275. MS m/e 499 (M+1)

Example 276. MS m/e 485 (M+1)

Example 277. MS m/e 546 (M+1)

Example 278. MS m/e 562 (M+1)

Example 279. MS m/e 511 (M+1)

Example 280. MS m/e 497 (M+1)

Example 281. MS m/e 539 (M+1)

Example 282. MS m/e 525 (M+1)

Example 283. MS m/e 567 (M+1)

Example 284. MS m/e 553 (M+1)

Example 285. MS m/e 595 (M+1)

Example 286. MS m/e 581 (M+1)

Example 287. A mixture of intermediate I-28-1 (98.2 mg, 0.2 mmol), $Pd(OAc)_2$ (13.4 mg, 0.06 mmol), BINAP (56 mg, 0.09 mmol), Cs_2CO_3 (91 mg, 0.28 mmol) and *p*-anisidine (118 mg, 0.96 mmol) in 5.0 ml of o-xylene was refluxed under N_2 for 18 hr. The reaction was followed by HPLC. The reaction was cooled to room temperature and diluted with CH_2Cl_2, filtered through celite and washed with CH_2Cl_2. Concentration and purification by flash chromatography with 5% MeOH in CH_2Cl_2 and preparative TLC provided 35 mg (33%) of the 3-substituted cyano ester. MS: 534 m/e $(M+H)^+$. A mixture of this 3-substituted cyano ester (25 mg, 0.046 mmol), Raney-Ni (excess) and 5.0 mL of DMF and 0.5 mL of MeOH was hydrogenated under 50 psi H_2 on a Parr apparatus overnight at room temperature. The reaction was monitored by HPLC to completion. The RaNi was removed by filtration, the solvent concentrated and the product purified by silica chromatography (5% MeOH in CH_2Cl_2) to afford 16.0 mg (71%) of product. MS: 492 m/e $(M+H)^+$.

Example 288. This compound was prepared by the same general procedure as Example 287 with intermediate I-28-1 (98.2 mg, 0.2 mmol), $Pd(OAc)_2$ (13.4 mg, 0.06 mmol), BINAP (56 mg, 0.09 mmol), Cs_2CO_3 (91 mg, 0.28 mmol) and N-methylaniline (104 uL, 0.96 mmol) in 5.0 ml of o-xylene. The 3-substituted cyano ester (60 mg, 58%) was obtained. MS:518 m/e $(M+H)^+$. Hydrogenation of the 3-substituted cyano ester (50 mg,

0.097 mmol) as described for Example 287 afforded 33 mg (72%) of product. MS: 476 m/e $(M+H)^+$.

Example 289. This compound was prepared by the same general procedure as Example 287 with intermediate I-28-1 (98.2 mg, 0.2 mmol), $Pd(OAc)_2$ (13.4 mg, 0.06 mmol), BINAP (56 mg, 0.09 mmol), Cs_2CO_3 (91 mg, 0.28 mmol) and 2,5-dimethoxyaniline (147 mg, 0.96 mmol) in 5.0 ml of o-xylene. The 3-substituted cyano ester (45 mg, 40%) intermediate was obtained. MS: 564 m/e $(M+H)^+$. Hydrogenation of this 3-substituted cyano ester (40.0 mg, 0.07 mmol) as described for Example 287 afforded 31 mg (85%) of the product. MS: 522 m/e $(M+H)^+$.

Examples 290-306 were prepared using general procedures described for Examples 287-289 starting with the cyano-ester intermediate I-28 and the appropriate aniline.

Example 290. MS m/e 492 (M+1)

Example 291. MS m/e 522 (M+1)

Example 292. MS m/e 522 (M+1)

Example 293. MS m/e 492 (M+1)

Example 294. MS m/e 492 (M+1)

Example 295. MS m/e 521 (M+1)

Example 296. MS m/e 520 (M+1)

Example 297. MS m/e 560 (M+1)

Example 298. MS m/e 522 (M+1)

Example 299. MS m/e 548 (M+1)

Example 300. MS m/e 522 (M+1)

Example 301. MS m/e 506 (M+1)

Example 302. MS m/e 480 (M+1)

Example 303. MS m/e 492 (M+1)

Example 304. MS m/e 506 (M+1)

Example 305. MS m/e 506 (M+1)

Example 306. MS m/e 546 (M+1)

Example 307. This compound was prepared using the N-propyl cyano-ester intermediate I-28-2 and butyrolactam using the coupling and reductive cyclization procedure described for Examples 287-289. MS m/e 454 (M+1)

Example 308. This compound was prepared using the N-iso-butyl cyano-ester intermediate I-28-4 and 2-oxo-3,4,5-trihydro-imidazole using the coupling and reductive cyclization procedure described for Examples 287-289. MS m/e 469 (M+1).

Synthesis of intermediate I-29-1.

Step 1: To 1g (2.7 mmol) intermediate I-30 in 130 mL acetone was added 3.7 mL 10N NaOH, followed by 1.22 mL (12.2 mmol) 2-iodopropane. The reaction was heated to reflux for 48 hours then concentrated. Water was added and the product extracted with methylene chloride and washed with brine. The organic layer was dried over sodium sulfate, filtered, concentrated, and dried under vacuum at 80 °C to give 830 mg (75% yield) of the N-isopropyl product; MS m/e 413(M + H).

Step 2: To 1.32g (3.2 mmol) of intermediate from step 1 in 15 mL AcOH was added 409µl (6.41 mmol, 2 eq) 70% HNO_3 drop wise at r.t. The reaction was heated to 80°C for 2 hours and let stand at r.t. overnight. The solid was collected by filtration, washed with water and ether, and dried under vacuum at 80°C to give 1.1g (77% yield) of the nitro intermediate; MS m/e 458 (M + H).

Step 3: To 1.1g (2.47 mmol) of the nitro intermediate from step 3 in 40 ml DMF/13 mL MeOH was added a few spatula tips of RaNi under nitrogen atmosphere. The reaction mixture was stirred at 50 psi on a Parr Apparatus for 48 hours. The solution was filtered through celite, concentrated under vacuum, suspended and stirred with ether overnight to give 800 mg of intermediate I-29-1 as an off-white solid (86% yield); MS m/e 386(M + H).

Example 309. To 20 mg (0.52 mmol) intermediate I-29-1 in 1 mL n-butanol was added 7 mg (0.6) 2-chloropyrimidine, and the reaction was heated to reflux overnight. The reaction mixture was then concentrated, stirred with water, filtered, and eluted on a silica gel preparative TLC plate using 9% MeOH/CH_2Cl_2. The product was collected, stirred

with methylene chloride/acetone, filtered, and concentrated. The sample was dried at 80 °C under high vacuum overnight to give 11.4 mg as a yellow solid, 47% yield MS m/e 464 (M + H).

Examples 310-318 were prepared using the general procedure for Example 309.

Example 310. MS m/e 464 (M+1)

Example 311. MS m/e 478 (M+1)

Example 312. MS m/e 464 (M+1)

Example 313. MS m/e 508 (M+1)

Example 314. MS m/e 463 (M+1)

Example 315. MS m/e 546 (M+1)

Example 316. MS m/e 506 (M+1)

Example 317. MS m/e 506 (M+1)

Example 318. MS m/e 492 (M+1)

Example 319. A mixture of cyano-ester intermediate I-30 (500 mg, 1.4 mmol), 10 N NaOH (3 mL), 3-bromo cyclohexene (1 mL) in acetone (30 mL) was stirred at reflux 12 hours. The acetone was removed at reduced pressure, excess water was added and the product collected by filtration. MS (ES+) m/e 451 (M + 1). The above intermediate in DMF/MeOH (50 mL, 9:1) and 2 scoops of RaNi was hydrogenated on a Parr apparatus for 12 hours. The catalyst was removed by filtration and the solvent removed at reduced pressure. MeOH was added to the residue and the product was precipitated by addition of water to give 580 mg of intermediate product. MS (ES+) m/e 411 (M + 1). To a stirred suspension of the previous intermediate product (450 mg, 1.1 mmol) in methylene chloride (100 mL) was added 3-(carbomethoxy)propionyl chloride (1.5 mL, 11 mmol) then aluminum chloride (1.5 g, 11 mmol) under nitrogen atmosphere. The reaction was stirred over night at room temperature. The solid was collected, added to ice water, triturated and collected. The product was chromatographed on silica gel (CH_2Cl_2/MeOH; 95/5). MS (ES+) m/e 539 (M + 1). The THP ring was formed by the general procedure described for Example 93 using sodium borohydride for the reduction and TFA in dichloromethane for cyclization.

MS (ES+) m/e 495 (M + 1).

Example 321. To a stirred solution of aluminum chloride (0.49 g, 3.69 mmol, 10 equiv.) in methylene chloride (3 mL)-nitromethane (2 mL) at room temperature under nitrogen, was added cyclopentylcarbonyl chloride (0.45 mL, 3.69 mmol, 10 equiv.). The solution was stirred at room temperature for 15 min. The dihydroindazole template (0.12 g, 3.69 mmol, 1 equiv.) was added as a solid to the reaction mixture. After 24 h at room temperature, the reaction mixture was cooled to 0 °C and quenched with 10% HCl. The product precipitated out of solution and was collected by filtration (0.15 g, 97% yield). ^{1}H NMR (DMSO-d_6): δ 1.55 (m, 1H), 1.67 (m, 2H), 1.83 (m, 3H), 1.97 (m, 2H), 2.88 (dd, 2H), 3.26 (dd, 2H), 3.87 (s, 3H), 4.004 (m, 1H), 4.89 (s, 2H).7.62 (d, 1H), 8.09 (s, 1H), 8.51 (s, 1H), 8.91 (s, 1H), 11.91 (s, 1H); MS (ESI): m/e 425 $(M+H)^+$.

Example 322. To a stirred suspension of product from Example 321 (126 mg, 0.297 mmol, 1 equiv.) in THF (8 mL) under nitrogen was added lithium borohydride (1.5 mL, 2.97 mmol, 10 equiv.) dropwise. Vigorous evolution of gas was observed and gradually the reaction mixture became homogeneous. Additional lithium borohydride (1.5 mL, 2.97 mmol, 10 equiv.) was added and the mixture was stirred overnight. When no starting material was observed by HPLC, the mixture was cooled to 0 °C and quenched slowly and carefully with water. The solvent was removed in vacuo leaving a tan solid (110 mg, 87% yield). A sample of the solid (40 mg) was purified by preparatory plate chromatography on silica gel using 10% methanol/ methylene chloride to yield the product as an off-white solid (11.6 mg, 29% yield). ^{1}H NMR (DMSO-d_6): δ 1.23 (m, 2H), 1.43 (m, 1H), 1.55 (m, 3H), 1.73 (m, 1H), 2.21 (m, 1H), 2.86 (dd, 2H), 3.23 (dd, 2H), 3.88 (s, 3H), 4.42 (m, 1H), 4.77 (s, 2H), 5.07 (d, 1H), 7.40 (d, 1H), 7.48 (d, 1H), 7.82 (s, 8.34 (s, 1H), 8.88 (s, 1H), 11.41 (s, 1H); MS (ESI): m/e 427 $(M+H)^+$.

Example 323. To a stirred suspension of product from Example 322 (70 mg, 0.164 mmol, 1 equiv.) in methylene chloride (8 mL) was added trifloroacetic acid (63.2 μL, 0.820 mmol, 5 equiv.) followed by triethylsilane (0.13 mL, 0.820 mmol, 5 equiv.). The reaction mixture became homogeneous and was stirred at room temperature overnight. Reaction

solvent was removed in vacuo leaving a tan solid. The solid was purified by preparatory plate chromatography using 10% methanol/ methylene chloride to yield a tan solid (12.3 mg, 18% yield). ^{1}H NMR (DMSO-d_6): δ 1.25 (m, 2H), 1.50 (m, 2H), 1.65 (m, 4H), 2.19 (m, 1H), 2.74 (d, 2H), 2.86 (dd, 2H), 3.22 (dd, 2H), 3.86 (s, 3H), 4.77 (s, 2H), 7.26 (d, 1H), 7.45 (d, 1H), 7.69 (s, 1H), 8.34 (s, 1H), 8.88 (s, 1H), 11.38 (s, 1H); MS (ESI): m/e 411 $(M+H)^+$.

Example 324. This compound was prepared using the general procedure described for Example 321. MS m/e 439 (M + 1).

Example 325. This compound was prepared using the general procedure described for Example 322-323. MS m/e 425 (M + 1).

Example 326. This compound was prepared using 1-tributylstannyl cyclopentene and the 3-bromo intermediate using the same general procedure as Example 39. MS m/e 395 (M + 1).

Example 327. This compound was prepared using the same general procedure as Example 39, using 1-tributylstannyl cyclopentene and intermediate I-28-1. The product formed was subjected to RaNi/H_2 hydrogenation and the product purified by silica gel chromatography. MS m/e 439 (M + 1).

Example 328. This compound was prepared using the same method as described for Example 9 using intermediate I-31-1. MS m/e 425 (M + 1).

Example 329. This compound was prepared using the same method as described for Example 328 using the N-sec-butyl intermediate. MS m/e 481 (M + 1).

Example 330. Step 1: To a solution of intermediate I-32 (95 mg, 0.28 mmol) in dry methylene chloride (15 mL) and nitromethane (10 mL) was added 3-chloropropionyl chloride (0.24 mL, 2.51 mmol) and then aluminum chloride (335 mg, 2.52 mmol). The

reaction was stirred for 20 minutes, when LC-MS and HPLC indicated that the reaction was complete. The reaction was concentrated via rotary evaporation, and water (50 mL) and 1 N HCl (3 mL) were added to the remaining solid and the mixture was stirred for 1 hour at room temperature. The solid was filtered and washed with water (150 mL). The remaining solid was dissolved in methylene chloride and transferred to a sep. funnel, where it was washed with saturated $NaHCO_3$ solution, then brine, before drying with $MgSO_4$. After filtration, the organic layer was concentrated via rotary evaporation. The remaining solid was purified by silica gel chromatography (95/5 CH_2Cl_2/MeOH), yielding a light yellow solid; (14 mg, .029 mmol). ^1H NMR (DMSO-d_6) 8.8998 (s, 1H), 8.5374 (s, 1H), 8.4863 (s, 1H), 8.1163 (d, 1H, J=6.57 Hz), 7.8673 (d, 1H, J=8.84 Hz), 4.9332 (s, 2 H), 4.5450 (d, 2H, J=7.07), 4.0058 (t, 2H, J=6.315), 3.8758 (s, 3H), 3.7388 (t, 2H, J=5.81), 3.4572 (t, 2H, J=6.32), 2.8429 (t, 2H, J=7.075), 2.1434 (m, 1H), 0.8167 (d, 6H, J=6.82); MS m/e 475 ($M+H^+$).

To a solution of the intermediate from step 1 (9 mg, 0.019 mmol) in pyridine (1 mL) was added hydroxylamine hydrochloride (10 mg, 0.144 mmol). Additional pyridine (5 mL) was added, and the reaction was stirred for 20 hours at room temperature before heating the reaction to 116 °C. After two hours at 116 °C, HPLC indicated that the reaction was complete. The reaction was concentrated via rotary evaporation, then purified by silica gel chromatography (97/3 CH_2Cl_2/MeOH), yielding a solid (9.4 mg, 0.021 mmol, 100% yield). ^1H NMR ($CHCl_3$) 8.8935 (s, 1H), 8.1584 (s, 1H), 7.7629 (dd, 1H, J=1.51, 8.58), 7.4800 (d, 1H, J=8.59), 6.1555 (s, 1H), 4.8954 (s, 2H), 4.5418 (t, 2H, J=9.98), 4.4112 (d, 2H, J=7.57), 3.9774 (m, 4H), 2.9837 (t, 2H, J=7.325), 2.2552 (m, 1H), 0.8938 (d, 6H, J=6.82); MS m/e 454 ($M+H^+$).

Example 332. To a stirring solution of 3-amino intermediate I-29-2 (25 mg, 0.065 mmol) in CH_2Cl_2 (5 mL) was added phenyl isothiocyanate (12 μL, 0.098 mmol) and pyridine (8 μL, 0.095 mmol). The reaction was heated to 50 °C for 24 h. After cooled to room temperature, the resulted precipitate was collected by filtration and dried to give 19 mg (56%) of the desired product. ^1H NMR (DMSO-d_6) δ 9.83 (s, 1H), 9.72 (s, 1H), 8.86 (s, 1H), 8.39 (s, 1H), 7.99 (s, 1H), 7.66 (m, 1H), 7.50 (m, 3H), 7.34 (m, 2H), 7.12 (m, 1H),

4.73 (m, 2H), 4.56 (m, 2H), 3.86 (s, 3H), 3.47 (m, 2H), 2.84 (m, 2H), 1.84 (m, 2H), 0.93 (m, 3H); MS (m/e) 521 (M + 1).

Example 333. This compound was prepared using the same general procedure as Example 332 using 4-methoxyphenyl isothiocyanate. MS (m/e) 551 (M + 1).

Example 369. To a stirred solution of 3-amino intermediate I-29-2 (25 mg, 0.065 mmol) in CH_2Cl_2 (5 mL) was added methyl chlorothiolformate (11 μL, 0.130 mmol), pyridine (30 μL, 0.371 mmol). After stirring for 24 h at room temperature, the resulting precipitate was filtered, dried to give 26 mg (87%) of the desired product. ^{1}H NMR (DMSO-d_6) δ 10.35 (s, 1H), 8.85 (s, 1H), 8.38 (s, 1H), 8.13 (s, 1H), 7.64-7.54 (m, 2H), 4.71 (s, 2H), 4.52 (m, 2H), 3.86 (s, 3H), 3.45 (m, 2H), 3.44 (s, 3H), 2.83 (m, 2H), 1.81 (m, 2H), 0.89 (m, 3H); MS (m/e) 460 (M + 1).

Example 370. This compound was prepared using the general method for Example 369 using 3-amino intermediate I-29-2 and ethyl chlorothiolformate. MS (m/e) 474 (M + 1).

Compounds 331, 334-357, 359-368, 371-380, 384-386 were prepared using one of methods A-D as outlined below.

Method A. Example 356. To intermediate I-29-1 (20mg, 0.052 mmol) in 2 mL methylene chloride/6.3 μl pyridine was added 10μl (0.078 mmol, 1.5 eq) of 4-methoxyphenyl isocyanate. The reaction was stirred at r.t. overnight. The solid was filtered off and washed with ether and the sample was dried at 80 °C under high vacuum overnight. To give 17 mgs (59% yield) as an off-white solid, MS m/e 535 (M + H).

The following Examples were prepared using method A.

Example 331. MS m/e 535 (M + H).

Example 334. MS m/e 549 (M + H).

Example 335. MS m/e 505 (M + H).

Example 336. MS m/e 535 (M + H).

Example 337. MS m/e 519 (M + H).

Example 338. MS m/e 549 (M + H).

Example 339. MS m/e 519 (M + H).

Example 340. MS m/e 549 (M + H).

Example 341. MS m/e 549 (M + H).

Example 342. MS m/e 537 (M + H).

Example 343. MS m/e 537 (M + H).

Example 344. MS m/e 537 (M + H).

Example 345. MS m/e 535 (M + H).

Example 346. MS m/e 549 (M + H).

Example 347. MS m/e 549 (M + H).

Example 348. MS m/e 523 (M + H).

Example 349. MS m/e 523 (M + H).

Example 350. MS m/e 523 (M + H).

Example 351. MS m/e 537 (M + H).

Example 352. MS m/e 537 (M + H).

Example 353. MS m/e 537 (M + H).

Example 354. MS m/e 537 (M + H).

Example 355. MS m/e 521 (M + H).

Example 356. MS m/e 535 (M + H).

Example 357. MS m/e 505 (M + H).

Example 359. MS m/e 539 (M + H).

Example 360. MS m/e 539 (M + H).

Example 361. MS m/e 583 (M + H).

Example 362. MS m/e 523 (M + H).

Example 363. MS m/e 551 (M + H).

Example 364. MS m/e 519 (M + H).

Example 365. MS m/e 548 (M + H).

Example 366. MS m/e 523 (M + H).

Example 367. MS m/e 539 (M + H).

Example 368. MS m/e 537 (M + H).

Example 371. MS m/e 548 (M + H).

Example 372. MS m/e 539 (M + H).

Example 373. MS m/e 584 (M + H).

Example 374. MS m/e 519 (M + H).

Example 375. MS m/e 511 (M + H).

Example 376. MS m/e 524 (M + H).

Example 377. MS m/e 591 (M + H).

Example 378. MS m/e 591 (M + H).

Example 379. MS m/e 591 (M + H).

Example 380. MS m/e 551 (M + H).

Example 384. MS m/e 551 (M + H).

Example 385. MS m/e 553 (M + H).

Example 386. MS m/e 598 (M + H).

Method B: Example 358. To a stirred solution of 3-amino intermediate I-29-2 (25 mg, 0.0649 mmol) in CH_2Cl_2 (5 mL) was added *N*-methyl-*N*-phenylcarbamoyl chloride (66 mg, 0.389 mmol) and pyridine (80 μL, 0.989 mmol). The reaction was heated to 48 °C for 58 h. After cooling to room temperature, the reaction was concentrated in vacuo. The residue was purified by preparative thin layer chromatography (CH_2Cl_2:MeOH 10:1) to give 24 mg (71%) of the desired product. ^{1}H NMR (DMSO-d_6) δ 8.45 (s, 1H), 8.36 (s, 1H), 8.16 (s, 1H), 7.97 (s, 1H), 7.57-7.24 (m, 7H), 4.68 (s, 2H), 4.50 (m, 2H), 3.86 (s, 3H), 3.45 (m, 2H), 3.32 (s, 3H), 2.83 (m, 2H), 1.80 (m, 2H), 0.89 (m, 3H); MS (m/e) 519 (M + 1).

Examples 397-401 and 403 were prepared using the method for Example 396.

Example 396. To a stirred solution of intermediate I-29-4 (25 mg, 0.063 mmol) in CH_2Cl_2 (5 mL) was added phenyl chloroformate (0.125 mmol) and pyridine (20 μL, 0.247 mmol). After stirring 3 h at room temperature, the resulting precipitate was filtered and dried to give 28 mg (91%) of the desired product. MS m/e 520 (M + H).

Example 397. MS m/e 550 (M + H).

Example 398. MS m/e 538 (M + H).

Example 399. MS m/e 506 (M + H).

Example 400. MS m/e 536 (M + H).

Example 401. MS m/e 524 (M + H).

Example 403. MS m/e 506 (M + H).

Method C: Trichloroacetamide intermediate: To 300 mg (0.78 mmol) amine intermediate I-29-1 in 30 mL CH_2Cl_2/95 µl pyridine, was added 174 µl (1.56 mmol, 2 eq) trichloroacetyl chloride at r.t. The reaction was stirred overnight, concentrated, stirred with water, filtered, and the solid was washed with ether. The sample was collected and dried at 80 °C under high vacuum to give 392 mg of product (95% yield) ^{1}H NMR (DMSO-d_6) δ 10.86 (s, 1H), 8.79 (s, 1H), 8. 43 (s, 1H), 8.21 (s, 1H), 7.86 (d, 1H), 7.78 (d, 1H), 5.27 (m, 1H), 4.75 (m, 2H), 3.87 (s, 3H), 3.43 (m, 2H), 2.81 (m, 2H), 1.63 (d, 6H). MS m/e 530(M+1).

Example 381. To 20 mg (0.038 mmol) trichloroacetamide intermediate in 1 mL DMF was added 8 mg (0.06 mmol, 1.5 eq) potassium carbonate, followed by, 7.2 µl (0.06 mmol, 1.5 eq) 2-(aminoethyl)pyrrolidine. The reaction was heated to 80 °C overnight, concentrated in vacuo, stirred with water, filtered and collected. The product was purified by preparative silica gel TLC plate eluting with 10% MeOH/CH_2Cl_2. The pure product was collected, stirred with solvent, filtered, concentrated, and dried at 80 °C under high vacuum. ^{1}H NMR (DMSO-d_6) δ 8.77 (s, 1H), 8.72 (s, 1H), 8.35 (s, 1H), 8.03 (s, 1H), 7.68 (d, 1H), 7.34 (d, 1H), 6.16 (m, 1H), 5.20 (m, 1H), 4.69 (s, 2H), 3.86 (s, 3H), 3.30 (m, 11H), 2.78 (m, 2H), 1.71 (m, 3H), 1.58 (d, 6H). MS m/e 526 (M+1).

The following Examples were prepared using method C as described for Example 381.

Example 382. MS m/e 540 (M+1).

Example 383. MS m/e 500 (M+1).

Example 387. MS m/e 569 (M+1).

Example 388. MS m/e 534 (M+1).

Example 389. MS m/e 534 (M+1).

Example 390. MS m/e 533 (M+1).

Example 391. MS m/e 523 (M+1).

Example 392. MS m/e 547 (M+1).

Example 393. MS m/e 548 (M+1).

Method D: N-p-Nitrophenylcarbonate Intermediate: To 484 mg (1.26 mmol) amine intermediate I-29-1 in 50 mL CH_2Cl_2/152µl pyridine was added 254 mg (1.26 mmol, 1.5eq) 4-nitrophenyl chloroformate. The reaction was stirred at r.t. overnight then concentrated *in vacuo* and washed with sodium bicarbonate. The product was collected washed with water and dried at 80 °C under high vacuum to give 616 mg of product (89% yield).
^{1}H NMR (DMSO-d_6) δ 10.48 (s, 1H), 8.78 (s, 1H), 8.38 (m, 1H), 8.12 (s, 1H), 7.82 (d, 1H), 7.75 (m, 2H), 7.58 (m, 3H), 5.24 (m, 1H), 4.69 (s, 2H), 3.87 (s, 3H), 3.39 (m, 2H), 2.81 (m, 2H), 1.60 (d, 6H). MS m/e 551 (M+1).

Example 394. To 30 mg (0.055 mmol) of the N-Nitrophenylcarbonate Intermediate in 2 mL THF was added 12 µl (0.082 mmol, 1.5eq) N,N,N-trimethyl-1,3-propanediamine, and the reaction was heated to 40 °C for 2 hours and concentrated. The crude product was purified by preparative TLC plate eluting with 10-20% MeOH/CH_2Cl_2. The pure product was collected, stirred with solvent, filtered, and concentrated. The sample was dried at 80 °C under high vacuum. ^{1}H NMR (DMSO-d_6) δ 9.07 (s, 1H), 8.77 (s, 1H), 8.35 (s, 1H), 7.97 (s, 1H), 7.68 (d, 1H), 7.45 (d, 1H), 5.20 (m, 1H), 4.69 (s, 2H), 3.86 (s, 3H), 2.91 (m, 3H), 2.78 (m, 2H), 2.3 (m, 11H), 1.72 (m, 3H), 1.59 (d, 6H). MS m/e 528 (M+1).

Example 395. This compound was prepared using p-nitrophenylcarbonate intermediate and N,N,N, trimethyl ethylenediamine by the method described for Example 394. MS m/e 514 (M+1).

Example 396. To 25 mg (0.045 mmol) of the N-p-nitrophenyl intermediate was added 500 µl N-piperidinylethanol. The reaction was stirred at r.t for ~5 hours, diluted with methylene chloride, washed with water/brine and dried over sodium sulfate. The crude product was purified by preparative TLC plate eluting with 8-10% MeOH/ CH_2Cl_2. The

pure product was collected, stirred with solvent, filtered, and concentrated. The sample was dried at 80 °C under high vacuum. ^{1}H NMR (DMSO-d_6) δ 9.80 (s, 1H), 8.77 (s, 1H), 8.36 (s, 1H), 8.10 (s, 1H), 7.72 (d, 1H), 7.50 (d, 1H), 5.20 (m, 1H), 4.78 (s, 2H), 4.19 (m, 2H), 3.86 (s, 3H), 2.78 (m, 2H), 2.41 (m, 4H), 1.59 (d, 6H), 1.40 (m, 10H). MS m/e 541 (M+1).

Example 402. To a stirring solution of 3-amino intermediate I-29-2 (25 mg, 0.0649 mmol) in CH_2Cl_2 (5 mL) was added dimethylsulfamoyl chloride (35 μL, 0.325 mmol), pyridine (100 μL, 1.24 mmol). The reaction was heated to 48 °C for 58 h. After cooled to room temperature, the reaction was concentrated *in vacuo*. The residue was purified by preparative thin layer chromatography (CH_2Cl_2:MeOH 10:1) to give 22 mg (69%) of the desired product. ^{1}H NMR (DMSO-d_6) δ 9.74 (s, 1H), 8.86 (s, 1H), 8.38 (s, 1H), 7.72 (s, 1H), 7.64 (d, 1H), 7.37 (d, 1H), 4.71 (s, 2H), 4.52 (m, 2H), 3.86 (s, 3H), 3.45 (m, 2H), 2.84 (m, 2H), 2.70 (s, 6H), 1.82 (m, 2H), 0.91 (m, 3H); MS (m/e) 493 (M + 1).

General procedures for Examples 404-430.

O-Nitrophenylcarbonate intermediate: Step 1: A mixture of the phenol intermediate I-33-1 (192 mg, 0.525 mmol) and p-nitrophenyl carbonate (314 mg, 1.03 mmol) in DMF (4 mL) was heated to 100 °C for 20 h. Solvent was removed by rotary evaporation and the residue was extracted into CH_2Cl_2 and washed with aqueous $NaHCO_3$. The organic layer was dried over $MgSO_4$, filtered, and evaporated. The resulting residue was purified by column chromatography (silica gel, 3% MeOH in CH_2Cl_2) to afford product (156 mg, 56%). ^{1}H NMR ($CDCl_3$) δ 8.86 (s, 1H), 8.34 (d, 2H, J=9.1), 7.69 (d, 1H, J=2.1), 7.53 (d, 2H, J=9.1), 7.49, (d, 1H, J=8.8), 7.41, (d, 1H, J=8.8), 6.01 (s, 1H), 4.84 (s, 2H), 4.62 (q, 2H, J=7.1), 3.96 (s, 3H), 3.55 (t, 2H, 8.0), 3.01 (t, 2H, J=8.0), 1.55 (t, 3H, J=7.1). MS m/e 538 (M + H).

A suspension of the carbonate intermediate in THF (2 mL) was treated with the amine (2.5 eq) and warmed to 40 °C for 2 h. Solvent was removed by rotary evaporation, and the residue was extracted into CH_2Cl_2 and washed with dilute aqueous NaOH. The organic layer was dried over $MgSO_4$, filtered, and evaporated. The resulting residue was purified by triturating with water (2 x 1 mL) and ether (2 x 1 mL).

Example 404. MS m/e 541 (M + H).

Example 405. MS m/e 529 (M + H).

Example 406. MS m/e 521 (M + H).

Example 407. MS m/e 569(M + H).

Example 408. MS m/e 548 (M + H).

Example 409. MS m/e 549 (M + H).

Example 410. MS m/e 538 (M + H).

Example 411. MS m/e 535 (M + H).

Example 412. MS m/e 537 (M + H).

Example 413. MS m/e 541 (M + H).

Example 414. MS m/e 563 (M + H).

Example 415. MS m/e 539 (M + H).

Example 416. MS m/e 560 (M + H).

Example 417. MS m/e 543 (M + H).

Example 418. MS m/e 552 (M + H).

Example 419. MS m/e 537 (M + H).

Example 420. MS m/e 551 (M + H).

Example 421. MS m/e 563 (M + H).

Example 422. MS m/e 524 (M + H).

Example 423. MS m/e 590 (M + H).

Example 424. MS m/e 535 (M + H).

Example 425. MS m/e 549 (M + H).

Example 426. MS m/e 591 (M + H).

Example 427. MS m/e 535 (M + H).

Example 428. MS m/e 577 (M + H).

Example 429. MS m/e 619 (M + H).

Example 430. MS m/e 617 (M + H).

Example 431. This compound was prepared using the same procedure as Example 172, starting with intermediate I-34. MS m/e 413 (M + 1).

The following Examples 432-442 were prepared from Example 431 using the appropriate bromide or iodide by the general method described for Examples 179-195.

Example 432. MS m/e 469 (M + H).

Example 433. MS m/e 469 (M + H).

Example 434. MS m/e 481 (M + H).

Example 435. MS m/e 441 (M + H).

Example 436. MS m/e 455 (M + H).

Example 437. MS m/e 455 (M + H).

Example 438. MS m/e 427 (M + H).

Example 439. MS m/e 467 (M + H).

Example 440. MS m/e 485 (M + H).

Example 441. MS m/e 471 (M + H).

Example 442. MS m/e 457 (M + H).

Example 443 was prepared using the same method as Example 289 starting with N-isopropyl intermediate I-35 and 2,5-dimethoxyaniline; MS m/e 522 (M + H).

Example 444. This compound was prepared using the intermediate I-37 by the general procedure described for Example 309. MS m/e 478 (M + H).

Example 445. This compound was prepared using the intermediate I-38 by the general procedure described for Examples 236-244. MS m/e 442 (M + H).

Example 446. This compound was prepared using the intermediate I-38 by the general procedure described for Examples 236-244. MS m/e 456 (M + H).

Example 447. This compound was prepared using the intermediate I-37 by the general procedure described for Example 287. MS m/e 477 (M + H).

Example 448. This compound was prepared using the intermediate I-37 by the general procedure described for Example 447. MS m/e 450 (M + H).

Utility

The compounds of the present invention are useful, *inter alia*, as therapeutic agents. Particularly, the compounds are useful for kinase inhibition, such as, for example, *trk*, VEGFR, PDGFR, PKC, MLK, DLK, Tie-2, FLT-3, and CDK1-6. Various compounds of the present invention show enhanced pharmaceutical properties over those disclosed in the art and improved pharmacokinetic properties in mammals. The compounds of the present invention show enhanced pharmaceutical properties over those disclosed in the art, including increased MLK and DLK dual inhibition activity, or increased VEGFR and Tie-2 dual inhibition activity, along with improved pharmacokinetic properties in mammals. For example, compounds of the present invention, including those in which at least one of R^3, R^4, R^5, and R^6 is $NR^{11}R^{32}$ or optionally substituted -(alkylene)$_x$-heterocycloalkyl, wherein the heterocycloalkyl does not include unsubstituted N-morpholinyl, N-piperidyl, or N-thiomorpholinyl, wherein said alkylene group is optionally substituted with one to three R^{10} groups, have been shown to increase MLK and DLK inhibition activity. In another example, compounds of the present invention, including those in which at least one of R^3, R^4, R^5, and R^6 is $NR^{11}R^{33}$, show increased VEGFR and Tie-2 inhibition activity.

In one embodiment, the present invention provides a method for treating or preventing diseases and disorders, such as those disclosed herein, which comprises administering to a subject in need of such treatment or prevention a therapeutically effective amount of a compound of the present invention.

In an additional embodiment, the present invention provides a method for inhibiting *trk* kinase activity comprising providing a compound of the present invention in an amount sufficient to result in effective inhibition. Particularly, inhibition of *trk* implies utility in, for example, diseases of the prostate such as prostate cancer and benign prostate hyperplasia, as well as for the treatment of inflammation, such as neurological inflammation and chronic arthritis inflammation. In a preferred embodiment, the *trk* kinase receptor is *trk* A.

The majority of cancers have an absolute requirement for angiogenesis, the process by which new blood vessels are formed. The most potent angiogenic cytokine is vascular endothelial growth factor (VEGF) and there has been substantial research into the development of VEGF/VEGF receptor (VEGFR) antagonists. Receptor tyrosine kinase (RTK) inhibitors could have broad spectrum antitumor activity in patients with advanced pre-treated breast and colorectal carcinoma and Kaposi's sarcoma. Potentially these agents may play a role in the treatment of both early (adjuvant) and advanced cancer. The importance of angiogenesis for the progressive growth and viability of solid tumors is well established. Emerging data suggest an involvement of angiogenesis in the pathophysiology of hematologic malignancies as well. Recently, authors have reported increased angiogenesis in the bone marrow of patients with acute myeloid leukemia (AML) and normalization of bone marrow microvessel density when patients achieved a complete remission (CR) after induction chemotherapy. Tumor angiogenesis depends on the expression of specific mediators that initiate a cascade of events leading to the formation of new microvessels. Among these, VEGF (vascular endothelial growth factor), FGF (fibroblast growth factor) play a pivotal role in the induction of neovascularization in solid tumors. These cytokines stimulate migration and proliferation of endothelial cells and induce angiogenesis in vivo. Recent data suggest an important role for these mediators in hematologic malignancies as well. Isolated AML blasts overexpress VEGF and VEGF receptor 2. Thus, the VEGF/VEGFR-2 pathway can promote the growth of leukemic blasts in an autocrine and paracrine manner. Therefore, neovascularization and angiogenic mediators/receptors may be promising targets for anti-angiogenic and anti-leukemic treatment strategies. Thus, in other embodiments, the present invention provides a method for treating or preventing angiogenic disorders where VEGFR kinase activity contributes to pathological conditions, the method comprising providing a compound of the present invention in an amount sufficient to result in the vascular endothelial growth factor receptor being contacted with an effective inhibitory amount of the compound. Inhibition of VEGFR implies utility in, for example, angiogenic disorders such as cancer of solid tumors, endometriosis, macular degeneration, retinopathy, diabetic retinopathy, psoriasis, hemangioblastoma, as well as other ocular diseases and cancers.

FLT3, a member of the receptor tyrosine kinase (RTK) class III, is preferentially expressed on the surface of a high proportion of acute myeloid leukemia (AML) and B-lineage acute lymphocytic leukemia (ALL) cells in addition to hematopoietic stem cells, brain, placenta and liver. An interaction of FLT3 and its ligand has been shown to play an important role in the survival, proliferation and differentiation of not only normal hematopoetic cells but also leukemia cells. Mutations of the FLT3 gene was first reported as an internal tandem duplication (ITD) of the juxtamembrane (JM) domain-coding sequence, subsequently as a missense mutation of D835 within a kinase domain. ITD- and D835-mutations are essentially found in AML and their frequencies are approximately 20 and 6% of adults with AML, respectively. Thus, mutation of the FLT3 gene is so far the most frequent genetic alteration reported to be involved in AML. Several large-scale studies in well-documented patients published to date have demonstrated that ITD-mutation is strongly associated with leukocytosis and a poor prognosis. An inhibitor compound of FLT3 tyrosine kinase have an application in treatment of leukemia. The present invention provides a method for treating disorders characterized by responsiveness to FLT3 inhibition, the method comprising providing a compound of the present invention in an amount sufficient to result in the inhibition of FLT3.

Platelet-derived growth factor (PDGF) was one of the first polypeptide growth factors identified that signals through a cell surface tyrosine kinase receptor (PDGF-R) to stimulate various cellular functions including growth, proliferation, and differentiation. Since then, several related genes have been identified constituting a family of ligands (primarily PDGF A and B) and their cognate receptors (PDGF-R alpha and beta). To date, PDGF expression has been shown in a number of different solid tumors, from glioblastomas to prostate carcinomas. In these various tumor types, the biologic role of PDGF signaling can vary from autocrine stimulation of cancer cell growth to more subtle paracrine interactions involving adjacent stroma and even angiogenesis. Thus, in additional embodiments, the present invention provides a method for treating or preventing disorders where PDGFR activity contributes to pathological conditions, the method comprising providing a compound of the present invention in an amount sufficient to result in the platelet derived growth factor receptor being contacted with an effective inhibitory amount of the compound. Inhibition of PDGFR implies utility in, for example,

various forms of neoplasia, rheumatoid arthritis, chronic arthritis, pulmonary fibrosis, myelofibrosis, abnormal wound healing, diseases with cardiovascular end points, such as atherosclerosis, restenosis, post-angioplasty restenosis, and the like.

In further embodiments, the present invention provides a method for treating or preventing disorders where MLK activity contributes to pathological conditions, such as those listed above, wherein the method comprises providing a compound of the present invention in an amount sufficient to result in the MLK receptor being contacted with an effective inhibitory amount of the compound. Inhibition of MLK implies utility in, for example, forms of cancer where MLKs play a pathological role as well as in neurological disorders.

In still other embodiments, the present invention provides a method for treating disorders characterized by the aberrant activity of trophic factor responsive cells, the method comprising providing a compound of the present invention in an amount sufficient to result in the trophic factor cell receptor being contacted with an effective activity inducing amount of the compound. In certain preferred embodiments, the activity of the trophic factor responsive cells is ChAT activity.

Fibroblast growth factor receptors (FGFR) are members of a family of polypeptides synthesized by a variety of cell types during the processes of embryonic development and in adult tissues. FGFR have been detected in normal and malignant cells and are involved in biological events that include mitogenic and angiogenic activity with a consequent crucial role in cell differentiation and development. To activate signal transduction pathways, FGFR are coupled to fibroblast growth factors (FGF) and heparan sulfate (HS) proteoglycans to form a biologically fundamental ternary complex. Based on these considerations, inhibitors able to block the signaling cascade through a direct interaction with FGFR could have antiangiogenesis and subsequent antitumor activity. Accordingly, the present invention provides a method for treating disorders characterized by the aberrant activity of FGF responsive cells, the method comprising providing a compound of the present invention in an amount sufficient to result in the FGFR being contacted with an effective activity inducing amount of the compound.

The compounds of the present invention can also have positive effects on the function and survival of trophic factor responsive cells by promoting the survival of

neurons. With respect to the survival of a cholinergic neuron, for example, the compound may preserve the survival of a cholinergic neuronal population at risk of dying (due to, e.g., injury, a disease condition, a degenerative condition or natural progression) when compared to a cholinergic neuronal population not presented with such compound, if the treated population has a comparatively greater period of functionality than the non-treated population.

A variety of neurological disorders are characterized by neuronal cells which are dying, injured, functionally compromised, undergoing axonal degeneration, at risk of dying, etc. These neurodegenerative diseases and disorders include, but are not limited to, Alzheimer's disease; motor neuron disorders (e.g. amyotrophic lateral sclerosis); Parkinson's disease; cerebrovascular disorders (e.g., stroke, ischemia); Huntington's disease; AIDS dementia; epilepsy; multiple sclerosis; peripheral neuropathies including diabetic neuropathy and chemotherapy induced peripheral neuropathy, AID related peripheral neuropathy; disorders induced by excitatory amino acids; and disorders associated with concussive or penetrating injuries of the brain or spinal cord.

In other preferred embodiments, the compounds of the present invention are useful for treating or preventing multiple myeloma and leukemias including, but not limited to, acute myelogenous leukemia, chronic myelogenous leukemia, acute lymphocytic leukemia, and chronic lymphocytic leukemia.

In additional embodiments, the present compounds are also useful in the treatment of disorders associated with decreased ChAT activity or the death, injury to spinal cord motoneurons, and also have utility in, for example, diseases associated with apoptotic cell death of the central and peripheral nervous system, immune system and in inflammatory diseases. ChAT catalyzes the synthesis of the neurotransmitter acetylcholine, and it is considered an enzymatic marker for a functional cholinergic neuron. A functional neuron is also capable of survival. Neuron survival is assayed by quantification of the specific uptake and enzymatic conversion of a dye (e.g., calcein AM) by living neurons. The compounds described herein may also find utility in the treatment of disease states involving malignant cell proliferation, such as many cancers.

Additional embodiments of the invention are directed to the use of any compound described herein, and stereoisomers or pharmaceutically acceptable salts thereof, in the

treatment and/or prevention of any of the conditions, diseases and disorders described above. Further embodiments are directed to the use of the compounds described herein, and stereoisomers or pharmaceutically acceptable salts thereof, in the manufacture of a medicament for treating and/or preventing said conditions, disorders and diseases.

The compounds of the present invention have important functional pharmacological activities which find utility in a variety of settings, including both research and therapeutic arenas. For ease of presentation, and in order not to limit the range of utilities for which these compounds can be characterized, the activities of the compounds of the present invention can be generally described as follows:

A. Inhibition of enzymatic activity

B. Effect on the function and/or survival of trophic factor responsive cells

C. Inhibition of inflammation-associated responses

D. Inhibition of cell growth associated with hyperproliferative states

E. Inhibition of developmentally programmed motoneuron death

Inhibition of enzymatic activity can be determined using, for example, VEGFR inhibition (e.g., VEGFR2 inhibition), MLK inhibition (e.g., MLK1, MLK2 or MLK3 inhibition), PDGFR kinase inhibition, NGF-stimulated *trk* phosphorylation, PKC inhibition, or *trk* tyrosine kinase inhibition assays. Effect on the function and/or survival of trophic factor responsive cells, e.g., cells of a neuronal lineage, can be established using any of the following assays: (1) cultured spinal cord choline acetyltransferase ("ChAT") assay; (2) cultured dorsal root ganglion ("DRG") neurite extension assay; (3) cultured basal forebrain neuron ("BFN") ChAT activity assay. Inhibition of inflammation-associated response can be established using an indoleamine 2,3-dioxygenase ("IDO") mRNA assay. Inhibition of cell growth associated with hyperproliferative states can be

determined by measuring the growth of cell lines of interest, such as an AT2 line in the case of prostate cancer. Inhibition of developmentally programmed motoneuron death can be assessed *in ovo* using embryonic chick somatic motoneurons, which cells undergo naturally occurring death between embryonic days 6 and 10, and analyzing inhibition of such naturally occurring cell death as mediated by the compounds disclosed herein.

The inhibition of enzymatic activity by the compounds of the present invention can be determined using, for example, the following assays:

VEGFR Inhibition Assay
MLK Inhibition Assay
PKC Activity Inhibition Assay
trkA Tyrosine Kinase Activity Inhibition Assay
Tie-2 Inhibition Assay
CDK1-6 Inhibition Assay
Inhibition of NGF-stimulated trk phosphorylation in a whole cell preparation
Platelet Derived Growth Factor Receptor (PDGFR) inhibition assay

A description of assays that may be used in connection with the present invention are set forth below. They are not intended, nor are they to be construed, as limiting the scope of the disclosure.

Inhibition of trkA Tyrosine Kinase Activity

Selected compounds of the present invention were tested for their ability to inhibit the kinase activity of baculovirus-expressed human trkA cytoplasmic domain using an ELISA-based assay as previously described (Angeles et al., Anal. Biochem. 236: 49-55, 1996). Briefly, the 96-well microtiter plate was coated with substrate solution (recombinant human phospholipase C- γ1/glutathione S-transferase fusion protein (Rotin et al., EMBO J., 11: 559-567, 1992). Inhibition studies were performed in 100 µl assay mixtures containing 50 mM Hepes, pH 7.4, 40µM ATP, 10 mM $MnCl_2$, 0.1% BSA, 2% DMSO, and various concentrations of inhibitor. The reaction was initiated by addition of trkA kinase and allowed to proceed for 15 minutes at 37°C. An antibody to

phosphotyrosine (UBI) was then added, followed by a secondary enzyme-conjugated antibody, alkaline phosphatase-labelled goat anti-mouse IgG (Bio-Rad). The activity of the bound enzyme was measured via an amplified detection system (Gibco-BRL). Inhibition data were analyzed using the sigmoidal dose-response (variable slope) equation in GraphPad Prism. The concentration that resulted in 50% inhibition of kinase activity is referred to as "IC_{50}".

Inhibition of Vascular Endothelial Growth Factor Receptor Kinase Activity

Selected compounds of the present invention were examined for their inhibitory effects on the kinase activity of baculovirus-expressed VEGF receptor (human flk-1, KDR, VEGFR2) kinase domain using the procedure described for the trkA kinase ELISA assay described above. The kinase reaction mixture, consisting of 50 mM Hepes, pH 7.4, 40 μM ATP, 10 mM $MnCl_2$, 0.1% BSA, 2% DMSO, and various concentrations of inhibitor, was transferred to PLC-γ/GST-coated plates. VEGFR kinase was added and the reaction was allowed to proceed for 15 min. at 37 °C. Detection of phosphorylated product was accomplished by addition of anti-phosphotyrosine antibody (UBI). A secondary enzyme-conjugated antibody was delivered to capture the antibody-phosphorylated PLC-γ/GST complex. The activity of the bound enzyme was measured via an amplified detection system (Gibco-BRL). Inhibition data were analyzed using the sigmoidal dose-response (variable slope) equation in GraphPad Prism.

Inhibition of Mixed Lineage Kinase-1 Activity

The kinase activity of MLK1 was assessed using the Millipore Multiscreen TCA "in-plate" format as described for protein kinase C (Pitt & Lee, *J. Biomol. Screening*, 1: 47-51, 1996). Briefly, each 50-μl assay mixture contained 20 mM Hepes, pH 7.0, 1 mM EGTA, 10 mM $MgCl_2$, 1 mM DTT, 25 mM β-glycerophosphate, 60 μM ATP, 0.25 μCi [γ-^{32}P]ATP, 0.1% BSA, 500 μg/ml myelin basic protein (UBI #13-104), 2% DMSO, 1 μM of test compound, and 1 μg/ml of baculoviral GST-$MLK1_{KD}$. Samples were incubated for 15 min at 37°C. The reaction was stopped by adding ice cold 50% TCA and the proteins were allowed to precipitate for 30 min at 4°C. The plates were then washed with ice cold

25% TCA. Supermix scintillation cocktail was added, and the plates were allowed to equilibrate for 1-2 hours prior to counting using the Wallace MicroBeta 1450 PLUS scintillation counter.

Dual Leucine Zipper Bearing Kinase Assay

Compounds were tested for their ability to inhibit the kinase activity of recombinant baculoviral human DLK, containing the kinase domain and leucine zipper. Activity was measured in 384-well FluoroNunc plates (Cat#460372) using a time-resolved fluorescence readout (PerkinElmer Application Note 1234-968). Plates were coated with 30 µl of the protein substrate MKK7 (Merritt et al. 1999) at a concentration of 20 µg/ml in Tris buffered saline (TBS). Each 30 µl assay contained 20 mM MOPS (pH 7.2), 15 mM $MgCl_2$, 0.1 mM Na_3VO_4, 1 mM DTT, 5 mM EGTA, 25 mM β-glycerophosphate, 0.1% BSA, 100 µM ATP, and 2.5% DMSO. Reactions were started by the addition of 10 ng/ml GST-$hDLK_{KD/LZ}$. For IC_{50} determinations, a 10-point dose response curve was generated for each compound. Plates were incubated at 37°C for 30 minutes, and the reactions stopped by the addition of 100 mM EDTA. Product was detected using Europium-labeled anti-phosphothreonine (Wallac#AD0093; diluted 1:10000 in 3%BSA/T-TBS). Following overnight capture at 4°C, 50 µl enhancement solution (Wallac #1244-105) was added and the plate gently agitated for 5 min. The fluorescence of the resulting solution was then measured using the time-resolved fluorescence (TRF) mode in the Multilabel Reader (Victor2 Model # 1420-018 or Envision Model # 2100). Inhibition data was analyzed using GraphPad PRISM. See also Merritt, S.E., Mata, M., Nihalani, D., Zhu, C., Hu, X., and Holzman, L.B. (1999) The Mixed Lineage Kinase DLK utilizes MKK7 and not MKK4 as Substrate. *J. Biol. Chem.* 274, 10195-10202.

Tie-2 Tyrosine Kinase Assay

Compounds were tested for their ability to inhibit the kinase activity of recombinant baculoviral human His_6-Tie2 cytoplasmic domain using a modification of the ELISA described for trkA (Angeles *et al.*, 1996). A 384-well plate format was used for single-point screening while IC_{50}s were performed on 96-well plates. For single-point screening, each barcoded 384-well Costar High Binding plate (Cat # 3703) was coated

with 50 µl/well of 10 µg/ml substrate solution (recombinant human GST- PLC-γ; Rotin *et al.*, 1992) in Tris-buffered saline (TBS). The Tie2 activity was measured in 50-µl assay mixtures containing 50 mM HEPES (pH 7.2), 40 µM ATP, 10 mM $MnCl_2$, 2.5% DMSO, 0.05% BSA, and 200 ng/ml His_6-$Tie2_{CD}$. For IC_{50} determinations, the assays were run as described above but in 96-well Costar High Binding plates (Cat # 3703) and with the volumes doubled. A 10-point dose response curve was generated for each compound. The kinase reaction was allowed to proceed at 37°C for 20 minutes. The detection antibody, N1-Eu anti-phosphotyrosine (PT66) antibody (Wallac #AD0041), was added at 1:2000 diluted in block buffer [3% BSA in TBS with 0.05% Tween-20 (TBST)]. After one-hour incubation at 37°C, 50 µl of enhancement solution (Wallac #1244-105) was added and the plate was gently agitated. The fluorescence of the resulting solution was then measured using the time-resolved fluorescence (TRF) mode in the Multilabel Reader (Victor2 Model # 1420-018 or Envision Model # 2100). Inhibition data were analyzed using ActivityBase and IC_{50} curves were generated using XLFit. The cited references are as follows:

1. Angeles, T. S., Steffler, C., Bartlett, B. A., Hudkins, R. L., Stephens, R. M., Kaplan, D. R., and Dionne, C. A. (1996) Enzyme-linked immunosorbent assay for trkA tyrosine kinase activity. *Anal. Biochem.* 236, 49-55.

2. Rotin, D., Margolis, B., Mohammadi, M., Daly, R.J., Daum, G., Li, N., Fischer, E.H., Burgess, W.H., Ullrich, A., Schlessinger, J. (1992) SH2 domains prevent tyrosine dephosphorylation of the EGF receptor: identification of Tyr992 as the high-affinity binding site for SH2 domains of phospholipase C-γ. *EMBO J.* 11, 559-567.

Dosage and Formulation

For therapeutic purposes, the compounds of the present invention can be administered by any means that results in the contact of the active agent with the agent's site of action in the body of the subject. The compounds may be administered by any conventional means available for use in conjunction with pharmaceuticals, either as individual therapeutic agents or in combination with other therapeutic agents, such as, for example, analgesics. The compounds of the present invention are preferably administered

in therapeutically effective amounts for the treatment of the diseases and disorders described herein to a subject in need thereof.

A therapeutically effective amount can be readily determined by the attending diagnostician, as one skilled in the art, by the use of conventional techniques. The effective dose will vary depending upon a number of factors, including the type and extent of progression of the disease or disorder, the overall health status of the particular patient, the relative biological efficacy of the compound selected, the formulation of the active agent with appropriate excipients, and the route of administration. Typically, the compounds are administered at lower dosage levels, with a gradual increase until the desired effect is achieved.

Typical dose ranges are from about 0.01 mg/kg to about 100 mg/kg of body weight per day, with a preferred dose from about 0.01 mg/kg to 10 mg/kg of body weight per day. A preferred daily dose for adult humans includes about 25, 50, 100 and 200 mg, and an equivalent dose in a human child. The compounds may be administered in one or more unit dose forms. The unit dose ranges from about 1 to about 500 mg administered one to four times a day, preferably from about 10 mg to about 300 mg, two times a day. In an alternate method of describing an effective dose, an oral unit dose is one that is necessary to achieve a blood serum level of about 0.05 to 20 μg/ml in a subject, and preferably about 1 to 20 μg/ml.

The compounds of the present invention may be formulated into pharmaceutical compositions by admixture with one or more pharmaceutically acceptable excipients. The excipients are selected on the basis of the chosen route of administration and standard pharmaceutical practice, as described, for example, in *Remington: The Science and Practice of Pharmacy*, 20th ed.; Gennaro, A. R., Ed.; Lippincott Williams & Wilkins: Philadelphia, PA, 2000. The compositions may be formulated to control and/or delay the release of the active agent(s), as in fast-dissolve, modified-release, or sustained-release formulations. Such controlled-release, or extended-release compositions may utilize, for example biocompatible, biodegradable lactide polymers, lactide/glycolide copolymers, polyoxyethylene-polyoxypropylene copolymers, or other solid or semisolid polymeric matrices known in the art.

The compositions can be prepared for administration by oral means; parenteral means, including intravenous, intramuscular, and subcutaneous routes; topical or transdermal means; transmucosal means, including rectal, vaginal, sublingual and buccal routes; ophthalmic means; or inhalation means. Preferably the compositions are prepared for oral administration, particularly in the form of tablets, capsules or syrups; for parenteral administration, particularly in the form of liquid solutions, suspensions or emulsions; for intranasal administration, particularly in the form of powders, nasal drops, or aerosols; or for topical administration, such as creams, ointments, solutions, suspensions aerosols, powders and the like.

For oral administration, the tablets, pills, powders, capsules, troches and the like can contain one or more of the following: diluents or fillers such as starch, or cellulose; binders such as microcrystalline cellulose, gelatins, or polyvinylpyrrolidones; disintegrants such as starch or cellulose derivatives; lubricants such as talc or magnesium stearate; glidants such as colloidal silicon dioxide; sweetening agents such as sucrose or saccharin; or flavoring agents such as peppermint or cherry flavoring. Capsules may contain any of the afore listed excipients, and may additionally contain a semi-solid or liquid carrier, such as a polyethylene glycol. The solid oral dosage forms may have coatings of sugar, shellac, or enteric agents. Liquid preparations may be in the form of aqueous or oily suspensions, solutions, emulsions, syrups, elixirs, etc., or may be presented as a dry product for reconstitution with water or other suitable vehicle before use. Such liquid preparations may contain conventional additives such as surfactants, suspending agents, emulsifying agents, diluents, sweetening and flavoring agents, dyes and preservatives.

The compositions may also be administered parenterally. The pharmaceutical forms acceptable for injectable use include, for example, sterile aqueous solutions, or suspensions. Aqueous carriers include mixtures of alcohols and water, buffered media, and the like. Nonaqueous solvents include alcohols and glycols, such as ethanol, and polyethylene glycols; oils, such as vegetable oils; fatty acids and fatty acid esters, and the like. Other components can be added including surfactants; such as hydroxypropylcellulose; isotonic agents, such as sodium chloride; fluid and nutrient replenishers; electrolyte replenishers; agents which control the release of the active compounds, such as aluminum monostearate, and various co-polymers; antibacterial


agents, such as chlorobutanol, or phenol; buffers, and the like. The parenteral preparations can be enclosed in ampules, disposable syringes or multiple dose vials. Other potentially useful parenteral delivery systems for the active compounds include ethylene-vinyl acetate copolymer particles, osmotic pumps, implantable infusion systems, and liposomes.

Other possible modes of administration include formulations for inhalation, which include such means as dry powder, aerosol, or drops. They may be aqueous solutions containing, for example, polyoxyethylene-9-lauryl ether, glycocholate and deoxycholate, or oily solutions for administration in the form of nasal drops, or as a gel to be applied intranasally. Formulations for topical use are in the form of an ointment, cream, or gel. Typically these forms include a carrier, such as petrolatum, lanolin, stearyl alcohol, polyethylene glycols, or their combinations, and either an emulsifying agent, such as sodium lauryl sulfate, or a gelling agent, such as tragacanth. Formulations suitable for transdermal administration can be presented as discrete patches, as in a reservoir or microreservoir system, adhesive diffusion-controlled system or a matrix dispersion-type system. Formulations for buccal administration include, for example lozenges or pastilles and may also include a flavored base, such as sucrose or acacia, and other excipients such as glycocholate. Formulations suitable for rectal administration are preferably presented as unit-dose suppositories, with a solid based carrier, such as cocoa butter, and may include a salicylate.

As those skilled in the art will appreciate, numerous modifications and variations of the present invention are possible in light of the above teachings. It is therefore understood that within the scope of the appended claims, the invention may be practiced otherwise than as specifically described herein, and the scope of the invention is intended to encompass all such variations.

CLAIMS

1. A compound of Formula I:



I

wherein:

ring A and ring B, independently, and each together with the carbon atoms to which they are attached, are selected from:

(a) a phenylene ring in which from 1 to 3 carbon atoms may be replaced by nitrogen atoms; and

(b) a 5-membered aromatic ring in which either

(1) one carbon atom may be replaced with an oxygen, nitrogen, or sulfur atom;

(2) two carbon atoms may be replaced with a sulfur and a nitrogen atom, an oxygen and a nitrogen atom, or two nitrogen atoms; or

(3) three carbon atoms may be replaced with three nitrogen atoms, one oxygen and two nitrogen atoms, or one sulfur and two nitrogen atoms;

A^1 and A^2 are independently selected from H, H; H, OR; H, SR; H, $N(R)_2$; and a group wherein A^1 and A^2 together form a moiety selected from =O, =S, and =NR;

B^1 and B^2 are independently selected from H, H; H, OR; H, SR; H, $N(R)_2$; and a group wherein B^1 and B^2 together form a moiety selected from =O, =S, and =NR;

provided that at least one of the pairs A^1 and A^2, or B^1 and B^2 forms =O;

R is independently selected from H, optionally substituted alkyl, $C(=O)R^{1a}$, $C(=O)NR^{1c}R^{1d}$, optionally substituted arylalkyl and optionally substituted heteroarylalkyl, wherein said optional substituents are one to three R^{10} groups;

R^1 is independently selected from H, $C(=O)R^{1a}$, OR^{1b}, $C(=O)NHR^{1b}$, $NR^{1c}R^{1d}$, and optionally substituted alkyl, wherein said optional substituents are one to three R^{10} groups;

R^{1a} is independently selected from optionally substituted alkyl, optionally substituted aryl and optionally substituted heteroaryl, wherein said optional substituents are one to three R^{10} groups;

R^{1b} is independently selected from H and optionally substituted alkyl, wherein said optional substituents are one to three R^{10} groups;

R^{1c} and R^{1d} are each independently selected from H and an optionally substituted alkyl, or together with the nitrogen to which they are attached form an optionally substituted heterocycloalkyl, wherein said optional substituents are one to three R^{10} groups;

R^2 is selected from H, $C(=O)R^{2a}$, $C(=O)NR^{2c}R^{2d}$, SO_2R^{2b}, CO_2R^{2b}, (alkylene)-$OC(=O)$-(alkylene)CO_2R^{11}, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl, wherein said optional substituents are one to three R^{10} groups;

R^{2a} is independently selected from optionally substituted alkyl, optionally substituted aryl, OR^{2b}, and $NR^{2c}R^{2d}$, wherein said optional substituents are one to three R^{10} groups;

R^{2b} is selected from H and optionally substituted alkyl, wherein said optional substituents are one to three R^{10} groups;

R^{2c} and R^{2d} are each independently selected from H and optionally substituted alkyl, or together with the nitrogen to which they are attached form an optionally substituted heterocycloalkyl, wherein said optional substituents are one to three R^{10} groups;

at least one of R^3, R^4, R^5, and R^6 is selected from $(alkylene)_xOR^{13}$, $C(=O)R^{13}$,

$(CH_2)_pOR^{22}$, O-(alkylene)-R^{27}, $OCH(CO_2R^{18})_2$, $OCH[(CH_2)_pOR^{20}]_2$, $C(=O)$-(alkylene)-R^{25}, , $NR^{11}R^{32}$, $NR^{11}R^{33}$, (alkylene)-$NR^{18}R^{19}$, $C(R^{12})$=N-R^{18}, CH=N-OR^{13}, $C(R^{12})$=N-OR^{20}, $C(R^{11})$=N-$NR^{11}C(=O)NR^{14A}R^{14B}$, $C(R^{11})$=N-$NR^{11}SO_2R^{18}$, $OC(=O)NR^{11}$(alkylene)-R^{26}, $OC(=O)[N(CH_2CH_2)_2N]$-R^{21}, $NR^{11}C(=O)OR^{23}$, $NR^{11}C(=O)S$-R^{18}, $NR^{11}C(=O)NR^{11}R^{23}$, $NR^{11}C(=S)NR^{11}R^{23}$, $NR^{11}S(=O)_2N(R^{15})_2$, $NR^{11}C(=O)NR^{11}$(alkylene)-R^{24}, $NR^{11}C(=O)N(R^{11})NR^{16A}R^{16B}$, substituted alkyl, wherein one of the substituents is a spirocycloalkyl group, optionally substituted (alkylene)$_x$-cycloalkyl, and optionally substituted -(alkylene)$_x$-heterocycloalkyl, wherein the heterocycloalkyl does not include unsubstituted N-morpholinyl, N-piperidyl, or N-thiomorpholinyl;

wherein any said alkylene group may be optionally substituted with one to three R^{10} groups;

provided that when R^3, R^4, R^5, or R^6 is $C(=O)R^{13}$, then R^{13} does not include a heterocycloalkyl group that contains a nitrogen bonded to the carbonyl moiety; and

the other R^3, R^4, R^5, or R^6 moieties can be selected independently from H, halogen, R^{10}, OR^{20}, optionally substituted alkyl, optionally substituted alkenyl, and optionally substituted alkynyl, wherein said optional substituents are one to three R^{10} groups;

Q is selected from an optionally substituted $C_{1\text{-}2}$ alkylene, wherein said optional substituents are one to three R^{10} groups; CR^{20}=CR^{20}; O; $S(O)_y$; $C(R^7)$=N; N=$C(R^7)$; CH_2-Z'; and Z'-CH_2; wherein

Z' is selected from O, S, C=O, and $C(=NOR^{11})$;

R^7 is selected from H, optionally substituted alkyl, and OR^{11}, wherein said optional substituents are one to three R^{10} groups;

R^{10} is selected from alkyl, aryl, heteroaryl, cycloalkyl, spirocycloalkyl, heterocycloalkyl, arylalkoxy, F, Cl, Br, I, CN, CF_3, $NR^{31A}R^{31B}$, NO_2, OR^{30}, OCF_3, =O, $=NR^{30}$, =N-OR^{30}, =N-$NR^{31A}R^{31B}$, $OC(=O)R^{30}$, $OC(=O)NHR^{29}$, O-$Si(R^{29})_4$, O-tetrahydropyranyl, ethylene oxide, $NR^{29}C(=O)R^{30}$, $NR^{29}CO_2R^{30}$, $NR^{29}C(=O)NR^{31A}R^{31B}$, $NHC(=NH)NH_2$, $NR^{29}S(O)_2R^{30}$,

$S(O)_yR^{18}$, CO_2R^{30}, $C(=O)NR^{31A}R^{31B}$, $C(=O)R^{30}$, $(CH_2)_pOR^{30}$, $CH=NNR^{31A}R^{31B}$, $CH=NOR^{30}$, $CH=NR^{30}$, $CH=NNHCH(N=NH)NH_2$, $S(=O)_2NR^{31A}R^{31B}$, $P(=O)(OR^{30})_2$, OR^{28}, and a monosaccharide wherein each hydroxyl group of the monosaccharide is independently either unsubstituted or is replaced by H, alkyl, alkylcarbonyloxy, or alkoxy;

R^{11} is selected from H and optionally substituted alkyl, wherein said optional substituents are one to three R^{10} groups;

R^{12} is selected from optionally substituted alkyl, optionally substituted aryl, and optionally substituted heteroaryl, wherein said optional substituents are one to three R^{10} groups;

R^{13} is independently selected from optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl, wherein said optional substituents are one to three R^{10} groups;

R^{14A} and R^{14B} are each independently selected from H, optionally substituted alkyl, optionally substituted aryl, and optionally substituted heteroaryl, wherein said optional substituents are one to three R^{10} groups;

R^{15} is independently selected from optionally substituted alkyl, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl, wherein said optional substituents are one to three R^{10} groups;

R^{16A} and R^{16B} are each independently selected from H and an optionally substituted alkyl, or together with the nitrogen to which they are attached form an optionally substituted heterocycloalkyl, wherein said optional substituents are one to three R^{10} groups;

R^{17} is selected from optionally substituted cycloalkyl, optionally substituted heterocycloalkyl, and optionally substituted heteroaryl, wherein said optional substituents are one to three R^{10} groups;

R^{18} is selected from H, optionally substituted alkyl, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl, wherein said optional substituents are one to three R^{10} groups;

R^{19} is selected from CN and triazole;

R^{20} is selected from H, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted aryl, optionally substituted arylalkyl, optionally substituted heteroaryl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl, wherein said optional substituents are one to three R^{10} groups;

R^{21} is selected from optionally substituted aryl, and optionally substituted heteroaryl, wherein said optional substituents are one to three R^{10} groups;

R^{22} is optionally substituted C_5-C_{10} alkyl, wherein said optional substituents are one to three R^{10} groups;

R^{23} is selected from optionally substituted aryl, optionally substituted heteroaryl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl, wherein said optional substituents are one to three R^{10} groups;

R^{24} is selected from optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted cycloalkyl, optionally substituted heterocycloalkyl, OR^{20}, $O(CH_2)_pOR^{20}$, $(CH_2)_pOR^{20}$, SR^{17}, SOR^{15}, SO_2R^{18}, CN, $N(R^{18})_2$, $C(=O)N(R^{18})_2$, $NR^{18}C(=O)R^{18}$, $NR^{18}C(=O)N(R^{18})_2$, $C(=NR^{18})OR^{18}$, $C(R^{12})=NOR^{18}$, $NHOR^{20}$, $NR^{18}C(=NR^{18})N(R^{18})_2$, NHCN, $CONR^{18}OR^{18}$, CO_2R^{18}, $OCOR^{15}$, $OC(=O)N(R^{18})_2$, $NR^{18}C(=O)OR^{15}$, and $C(=O)R^{18}$, wherein said optional substituents are one to three R^{10} groups;

R^{25} is selected from optionally substituted cycloalkyl, optionally substituted heterocycloalkyl, OR^{20}, $O(CH_2)_pOR^{20}$, $(CH_2)_pOR^{20}$, SR^{17}, SOR^{15}, SO_2R^{18}, CN, $N(R^{17})_2$, $C(=O)N(R^{18})_2$, $NR^{18}C(=O)R^{18}$, $NR^{18}C(=O)N(R^{18})_2$, $C(=NR^{18})OR^{18}$, $C(R^{12})=NOR^{18}$, $NHOR^{20}$, $NR^{18}C(=NR^{18})N(R^{18})_2$, NHCN, $CONR^{18}OR^{18}$, CO_2R^{18}, $OCOR^{15}$, $OC(=O)N(R^{18})_2$, $NR^{18}C(=O)OR^{15}$, and $C(=O)R^{18}$, wherein said optional substituents are one to three R^{10} groups;

R^{26} is selected from optionally substituted cycloalkyl, optionally substituted heterocycloalkyl, optionally substituted aryl, optionally substituted heteroaryl, OR^{11}, $O(CH_2)_pOR^{20}$, $(CH_2)_pOR^{20}$, SR^{17}, SOR^{15}, SO_2R^{18}, CN,

$N(R^{18})_2$, $C(=O)N(R^{18})_2$, $NR^{18}C(=O)R^{18}$, $NR^{18}C(=O)N(R^{18})_2$, $C(=NR^{18})OR^{18}$, $C(R^{12})=NOR^{18}$, $NHOR^{20}$, $NR^{18}C(=NR^{18})N(R^{18})_2$, NHCN, $CONR^{18}OR^{18}$, CO_2R^{18}, $OCOR^{15}$, $OC(=O)N(R^{18})_2$, $NR^{18}C(=O)OR^{15}$, and $C(=O)R^{18}$, wherein said optional substituents are one to three R^{10} groups;

R^{27} is selected from optionally substituted cycloalkyl, CN, $C(R^{12})=NOR^{18}$, and $C(=O)N(R^{18})_2$, wherein said optional substituents are one to three R^{10} groups;

R^{28} is the residue of an amino acid after the removal of the hydroxyl moietyfrom the carboxyl group thereof;

R^{29} is H or alkyl;

R^{30} is H, alkyl, aryl, arylalkyl, heteroaryl, cycloalkyl, or heterocycloalkyl;

R^{31A} and R^{31B} are each independently selected from H, alkyl, and arylalkyl, or together with the nitrogen to which they are attached form a heterocycloalkyl;

R^{32} is optionally substituted aryl, wherein said optional substituents are one to three R^{10} groups;

R^{33} is selected from optionally substituted cycloalkyl, optionally substituted heteroaryl, and optionally substituted heterocycloalkyl, wherein said optional substituents are one to three R^{10} groups;


p is independently selected from 1, 2, 3, and 4;

x is 0 or 1;

y is independently selected from 0, 1 and 2; and

provided that:


1) when R^1 is H; A^1, A^2 and B^1, B^2 are each =O or =S; ring A is a phenylene; ring B is



, wherein R^A is H or C_1-C_4 alkyl; and Q is $CR^{20}=CR^{20}$, wherein R^{20} is H, C_1-C_4 alkyl or C_1-C_4 alkoxy; then neither R^3 or R^4 includes


(alkylene)-NR^aR^b, wherein R^a and R^b combine with the nitrogen to which they are attached to form heterocycloalkyl group; and

2) when A^1, A^2 is =O; B^1, B^2 are independently H or OH, or B^1, B^2 combine to form =O; rings A and B are each a phenylene; Q is O, S, or CH_2; and R^2 is H, or optionally substituted




then any of R^3, R^4, R^5, and R^6 cannot include $(alkylene)_xOR^{13}$, wherein R^{13} is cycloalkyl; $(CH_2)_pOR^{22}$, wherein R^{22} is C_5-C_7 alkyl; O-(alkylene)-R^{27}, wherein R^{27} is CN; $OCH[(CH_2)_pOR^{20}]_2$; $NR^{11}R^{33}$, wherein R^{33} is cycloalkyl; or NR^aR^b, wherein R^a and R^b combine with the nitrogen to which they are attached to form heterocycloalkyl group;

3) when A^1, A^2 and B^1, B^2 are each =O; rings A and B are each a phenylene; Q is O or S; and R^2 is optionally substituted



then any of R^3, R^4, R^5, and R^6 cannot include $(alkylene)_xOR^{13}$, wherein R^{13} is cycloalkyl; $(CH_2)_pOR^{22}$, wherein R^{22} is C_5-C_7 alkyl; O-(alkylene)-R^{27}, wherein R^{27} is CN; $OCH[(CH_2)_pOR^{20}]_2$; $NR^{11}R^{33}$, wherein R^{33} is cycloalkyl; or NR^aR^b, wherein R^a and R^b combine with the nitrogen to which they are attached to form heterocycloalkyl group;

4) when A^1, A^2 is =O; B^1, B^2 are independently H or OH, or B^1, B^2 combine to form =O; rings A and B are each a phenylene; Q is O, S, or CH-R^a; and one of R^2 or R^a is H and the other is optionally substituted



wherein W is optionally substituted C_1 alkyl, or $NR^{31A}R^{31B}$; then any of R^3, R^4, R^5, and R^6 cannot include OR^{13} or $C(=O)NR^cR^d$ wherein R^c and R^d combine with the nitrogen to which they are attached to form heterocycloalkyl group;

or a stereoisomer or pharmaceutically acceptable salt form thereof.

2. The compound of claim 1 wherein rings A and B are each a phenylene.

3. The compound of claim 1 wherein ring A is a phenylene, and ring B is a 5-membered aromatic ring in which one or two carbon atoms may be replaced with a nitrogen atom.

4. The compound of claim 3 wherein ring B is a pyrazolylene.

5. The compound of claim 1 wherein R^1 is selected from H, substituted alkyl, and unsubstituted alkyl.


6. The compound of claim 1 wherein groups A^1, A^2 and B^1, B^2 are each independently selected from H, H; and a group wherein A^1, A^2 or B^1, B^2 together form =O.

7. The compound of claim 6 wherein A^1, A^2 are H, H; and B^1, B^2 together form =O.

8. The compound of claim 1 wherein Q is optionally substituted C_{1-2} alkylene.

9. The compound of claim 8 wherein Q is CH_2 or CH_2CH_2.

10. The compound of claim 1 having a structure of Formula II:



II

wherein ring B together with the carbon atoms to which it is attached, is selected from:

(a) a phenylene ring in which from 1 to 3 carbon atoms may be replaced by nitrogen atoms; and

(b) a 5-membered aromatic ring in which from 1 to 2 carbon atoms may be replaced by nitrogen atoms; and


Q is selected from an optionally substituted C_{1-2} alkylene, wherein said optional substituents are one to three R^{10} groups.

11. The compound of claim 10 wherein A^1, A^2 are H, H; and B^1, B^2 together form =O.


12. The compound of claim 10 wherein the B ring is a phenylene.

13. The compound of claim 10 wherein the B ring is a pyrazolylene.

14. The compound of claim 13 wherein the B ring is




15. The compound of claim 13 wherein the B ring is



16. The compound of claim 10 wherein R^1 is H or unsubstituted alkyl.

17. The compound of claim 10 wherein Q is CH_2 or CH_2CH_2.


18. The compound of claim 10 having a structure of Formula III:



III


wherein ring B is a phenylene or a pyrazolylene, and R^1 is H or unsubstituted alkyl.

19. The compound of claim 18 having a structure of Formula IV:




IV

20. The compound of claim 18 having a structure of Formula V:



V.

21. The compound of claim 18 having Formula VI:



VI.

22. The compound of claims 18, 19, 20 or 21 wherein Q is CH_2 or CH_2CH_2.

23. The compound of claim 1 wherein at least one of R^3, R^4, R^5, or R^6 is selected from OR^{13}, $C(=O)R^{13}$, $(CH_2)_pOR^{22}$, O-(alkylene)-R^{27}, $OCH(CO_2R^{18})_2$, $OCH[(CH_2)_pOR^{20}]_2$, and C(=O)-(alkylene)-R^{25}.

24. The compound of any one of claims 1, 10, 18, 19, 20, or 21 wherein at least one of R^3, R^4, R^5, or R^6 is $NR^{11}R^{32}$..

25. The compound of claim 24 wherein at least one of R^3, R^4, R^5, or R^6 is NHR^{32}.

26. The compound of any one of claims 1, 10, 18, 19, 20, or 21 wherein at least one of R^3, R^4, R^5, or R^6 is optionally substituted -$(alkylene)_x$-heterocycloalkyl, wherein the heterocycloalkyl does not include unsubstituted N-morpholinyl, N-piperidyl, or N-thiomorpholinyl.

27. The compound of claim 26 wherein at least one of R^3, R^4, R^5, or R^6 is tetrahydropyranyl.

28. The compound of any one of claims 1, 10, 18, 19, 20, or 21 wherein at least one of R^3, R^4, R^5, or R^6 is $NR^{11}R^{33}$.

29. The compound of claim 28 wherein at least one of R^3, R^4, R^5, or R^6 is NHR^{33}.

30. The compound of claim 29 wherein R^{33} is optionally substituted heteroaryl.

31. The compound of any one of claims 1, 10, 18, 19, 20, or 21 wherein at least one of R^3, R^4, R^5, or R^6 is $C(R^{12})$=N-OR^{20}.


32. The compound of claim 31 wherein R^{12} and R^{20} are alkyl.




33. The compound of claim 1 wherein the compounds are selected in accordance with Table 1.

34. The compound of claim 1 wherein the compounds are selected in accordance with Table 2.


35. The compound of claim 1 wherein the compounds are selected in accordance with Table 3.

36. The compound of claim 1 wherein the compounds are selected in accordance with the following table:






























R^3	R^2	Q	R^5
	$CH_2CH_2CH_2OH$	CH_2	H
	CH_2CH_3	CH_2CH_2	OCH_3
	CH_2CH_3	CH_2CH_2	OCH_3





















37. The compound of claim 1 wherein the compounds are selected in accordance with the following table:





























R^3	R^2	Q
$CH_3ON=$	H	CH_2CH_2
tBuON=	H	CH_2CH_2
EtON=	H	CH_2CH_2
$Me_2HCCH_2ON=$	H	CH_2CH_2
$CH_2=CHCH_2ON=$	H	CH_2CH_2
HON=	H	CH_2CH_2
NOEt	H	CH_2CH_2
NOiPr	H	CH_2CH_2
$CH_3SO_2NH-N=$	H	CH_2CH_2
O-N=	$CH_2CH(CH_3)_2$	CH_2CH_2
O-N=	$CH_2CH(CH_3)_2$	CH_2CH_2
O-N=	$CH_2CH(CH_3)_2$	CH_2CH_2

R^3	R^2	Q
	$CH_2CH(CH_3)_2$	CH_2CH_2
	H	CH_2CH_2
	$CH(CH_3)_2$	CH_2CH_2
	$CH(CH_3)_2$	CH_2CH_2
	$CH(CH_3)_2$	CH_2CH_2
	$CH(CH_3)_2$	CH_2CH_2
	$CH_2CH(CH_3)_2$	CH_2CH_2
	$CH_2CH_2CH_3$	CH_2CH_2
	$CH_2CH_2CH_3$	CH_2CH_2
	$CH_2CH_2CH_2CH_3$	CH_2CH_2
	$CH_2CH_2CH_2CH_3$	CH_2CH_2
	$CH_2CH_2CH_2CH_3$	CH_2CH_2
	$CH_2CH_2CH_2CH_3$	CH_2CH_2
	CH_2CH_3	CH_2CH_2


R^3	R^2	Q
	CH_2CH_3	CH_2CH_2
	CH_2CH_3	CH_2CH_2
	CH_2CH_3	CH_2CH_2
	CH_2CH_3	CH_2CH_2
	$CH_2CH(CH_3)_2$	CH_2CH_2
	$CH_2CH_2CH_3$	CH_2CH_2
	$CH_2CH_2CH_3$	CH_2CH_2
	CH_3	CH_2CH_2
	CH_3	CH_2CH_2
	CH_3	CH_2CH_2
	$CH_2CH_2NMe_2$	CH_2CH_2
	$CH_2CH_2NMe_2$	CH_2CH_2
	$CH_2CH_2NMe_2$	CH_2CH_2
	$CH_2CH(CH_3)_2$	CH_2CH_2







R^3	R^2	Q
	$CH_2CH(CH_3)_2$	CH_2CH_2
		CH_2CH_2
		CH_2CH_2
		CH_2CH_2
		CH_2CH_2
		CH_2CH_2
		CH_2CH_2
		CH_2CH_2
		CH_2CH_2
	$CH(CH_3)_2$	CH_2CH_2
	$CH(CH_3)_2$	CH_2CH_2
	$CH(CH_3)_2$	CH_2CH_2
	$CH(CH_3)_2$	CH_2CH_2

R^3	R^2	Q
	$CH(CH_3)_2$	CH_2CH_2
	$CH(CH_3)_2$	CH_2CH_2
	$CH_2CH_2CH_3$	CH_2CH_2
	$CH_2CH_2CH_3$	CH_2CH_2
	$CH_2CH_2CH_3$	CH_2CH_2
	$CH(CH_3)_2$	CH_2CH_2
	$CH(CH_3)_2$	CH_2CH_2
	$CH_2CH_2CH_3$	CH_2CH_2
		CH_2CH_2
	$CH_2CH_2CH_3$	CH_2CH_2
	$CH_2CH_2CH_3$	CH_2CH_2
	$CH_2CH_2CH_3$	CH_2CH_2




R^3	R^2	Q
	$CH(CH_3)_2$	CH_2CH_2
	$CH_2CH(CH_3)_2$	CH_2CH_2
	$CH_2CH(CH_3)_2$	CH_2CH_2
	$CH(CH_3)_2$	CH_2CH_2
	$CH(CH_3)_2$	CH_2CH_2
	$CH_2CH_2CH_3$	CH_2CH_2
	$CH_2CH(CH_3)_2$	CH_2CH_2
	$CH(CH_3)_2$	CH_2CH_2
	$CH_2CH(CH_3)_2$	CH_2CH_2
	$CH(CH_3)_2$	CH_2CH_2
	$CH_2CH(CH_3)_2$	CH_2CH_2
	$CH_2CH(CH_3)_2$	CH_2CH_2
	$CH_2CH(CH_3)_2$	CH_2CH_2
	$CH_2CH_2CH_3$	CH_2CH_2

38. The compound of claim 1 wherein the compounds are selected in accordance with the following table:




R^3	R^2	Q
	$CH(CH_3)_2$	CH_2CH_2
	$CH_2CH(CH_3)_2$	CH_2CH_2
	$CH_2CH(CH_3)_2$	CH_2CH_2
	$CH_2CH(CH_3)_2$	CH_2CH_2
	$CH_2CH(CH_3)_2$	CH_2CH_2
	CH_2CH_3	CH_2CH_2

39. The compound of claim 1 wherein the compound is selected in accordance with the following table:


Structure




40. The compound of claim 1 wherein the compound is




5


41. The compound of claim 1 wherein the compound is




42. The compound of claim 1 wherein the compound is




43. The compound of claim 1 wherein the compound is



44. The compound of claim 1 wherein the compound is



45. The compound of claim 1 wherein the compound is



46. The compound of claim 45 wherein the compound is the (+)-isomer.

47. A pharmaceutical composition comprising a compound of claim 1 and at least one pharmaceutically acceptable excipient.

48. The pharmaceutical composition of claim 47, comprising a compound of claim 43.

49. The pharmaceutical composition of claim 47, comprising a compound of claim 44.

50. The pharmaceutical composition of claim 47, comprising a compound of claim 45.

51. The pharmaceutical composition of claim 47, comprising a compound of claim 46.

52. A method for treating a prostate disorder comprising administering to a subject in need of such treatment a therapeutically effective amount of a compound of claim 1.

53. The method of claim 52 wherein the prostate disorder is prostate cancer or benign prostate hyperplasia.

54. A method for treating an angiogenic disorder comprising administering to a subject in need of such treatment a therapeutically effective amount of a compound of claim 1.

55. The method of claim 54 wherein the angiogenic disorder is selected from the group consisting of cancer of solid tumors, hematologic tumors, macular degeneration, retinopathy of prematurity, diabetic retinopathy, rheumatoid arthritis, psoriasis, endometriosis, and restenosis.

56. A method for treating a pathological disorder comprising administering to a subject in need of such treatment a therapeutically effective amount of a compound of claim 1.

57. The method of claim 56 wherein the pathological disorder is selected from the group consisting of neoplasia, chronic arthritis, pulmonary fibrosis, myelofibrosis, abnormal wound healing, and atherosclerosis.

58. A method for treating a neurodegenerative disease or disorder comprising administering to a subject in need of such treatment a therapeutically effective amount of a compound of claim 1.

59. The method of claim 58 wherein the neurodegenerative disease or disorder is selected from the group consisting of Alzheimer's disease, amyotrophic lateral sclerosis, Parkinson's disease, stroke, ischemia, Huntington's disease, AIDS dementia, epilepsy,

multiple sclerosis, peripheral neuropathy, chemotherapy induced peripheral neuropathy, AIDS related peripheral neuropathy, and injury of the brain or spinal cord.

60. The method of claim 59 wherein the neurodegenerative disease is Alzheimer's disease.

61. The method of claim 60 wherein the subject is administered a compound of claim 44.

62. The method of claim 60 wherein the subject is administered a compound of claim 45.

63. The method of claim 60 wherein the subject is administered a compound of claim 46.

64. A method for treating multiple myeloma or leukemia which comprises administering to a subject in need of such treatment a therapeutically effective amount of a compound of claim 1.

65. The method of claim 64 wherein the leukemia is acute myelogenous leukemia, chronic myelogenous leukemia, acute lymphocytic leukemia, or chronic lymphocytic leukemia.

66. A method of treating cancer of solid tumors comprising administering to a subject in need of such treatment a therapeutically effective amount of a compound of claim 43.

67. A method of treating a hematologic tumor comprising administering to a subject in need of such treatment a therapeutically effective amount of a compound of claim 43.