

- 1 -

"Process for degumming a fatty substance and fatty
substance thus obtained"

The present invention relates to a process for degumming a fatty substance such as a crude or delecithinated, animal or vegetable oil or fat, and to the fatty substance thus obtained.

All fatty substances contain a certain number of impurities, liposoluble substances entrained during the bursting of the oil-bearing cells, which can make them unusable for consumption. Some of these impurities have a detrimental influence on the taste, odor and appearance of the product and on its storage.

The purpose of the refining of fatty substances is to remove the free fatty acids, oxidation products, unpleasant flavors, colorants and toxic products (such as pesticides and glycosides), as well as the phospholipids and the metals (such as iron and copper) which are present in trace form and generally bonded to organic compounds.

The presence of phospholipids in the crude oils entails a number of disadvantages. In the presence of water they hydrate and form deposits which decompose in the course of time. Experience also shows that a refined oil which is poorly freed from its phospholipids becomes acidic, oxidizes and rapidly takes on an unpleasant taste. Phospholipids are often bonded to heavy metals (such as calcium, magnesium, iron and copper), some of which have a detrimental influence on the conservation of the refined fatty substances, since they are oxidation catalysts. Phospholipids are also thermally unstable substances which darken the oil when they decompose at high temperature. Finally, since phospholipids are surfactants, their incomplete removal at the beginning of the refining entails the formation of foams and emulsions which result in abnormal losses of oil and deactivation of the decolorizing earths.

Among the fatty substances, some contain few phospholipids (for example palm fat, lauric and animal

fats); they can therefore be easily rid of these substances by dry degumming, that is to say by the addition of an acid in order to decompose them and of an earth in order to bind them thereto. Thus, these fatty substances
5 can be refined by neutralizing distillation or physical refining.

The oils obtained by pressing and/or by extraction with the aid of a solvent (for example soya, rapeseed and sunflower oils) are, on the other hand, very
10 rich in phospholipids and are therefore generally refined chemically. Refining of this type presents a number of disadvantages; one of these is that it gives rise to "soapstocks", mixtures of oil and of soaps, which have to be treated, and this involves losses of oil and
15 additional costs.

The total phospholipid content of these crude oils, expressed in the form of phosphorus, can be easily lowered from 800 ppm to 150-200 ppm by aqueous degumming or delecithination. The oil is stirred in the presence of
20 water at 80°C, resulting in hydration and flocculation of the phospholipids. They can therefore be separated by gravity separation or centrifuging. The 150-200 ppm of residual phosphorus represent chiefly the nonhydratable phospholipids, which are complexes of phosphatidic acid
25 and of phosphatidyl ethanolamine, which are combined with divalent ions (such as calcium, iron or magnesium). The removal of these nonhydratable phospholipids, which has become necessary for physical refining, can be carried out by virtue of a special degumming according to various
30 routes:

- An acidic degumming which consists in dissociating the phospholipid complex with the aid of an acid in order next to hydrate it. This superdegumming (see German Patents No. 2609705 and 132877), also including a special
35 cooling cycle, produces phosphorus contents which are much lower than with a conventional acidic degumming. However, the final result depends greatly on the quality of the crude oil. Finally, the removal of iron still requires much bleaching earth. This superdegumming has

therefore been supplemented (European Patent No. 0 348 004) by a second cooling cycle and the addition of water or of caustic soda to improve the purification. However, this results in a very long, very complex and
5 costly process.

- An acidic refining which, by virtue of an acid, dissociates the phospholipid complexes and then converts them in the presence of caustic soda into a completely hydratable sodium complex (see US Patent No. 4,698,185
10 and European Patent No. 0 349 178 and European Application No. 92200543.4). This process, requiring intense stirring, makes it possible to obtain oils with a low iron and phospholipid content; however, it requires 2 to 3 separations per centrifuging.

15 There are also known processes for refining oils and fats by treatment of the fatty substance firstly with an acid of the phosphoric acid type and then with a fatty acid salt or sodium or potassium carboxylate, but these processes involve two stages of treatment of the fatty
20 substance and do not enable a fine emulsion to be obtained.

One of the essential objectives of the present invention consists in overcoming the abovementioned disadvantages of the existing processes and in providing
25 an industrially and economically valid process making it possible to obtain fatty substances such as crude or delecithinated, animal or vegetable oils or fats which are completely degummed in order to permit their physical refining, making it possible especially to remove practically completely the phospholipids which they contain
30 and, more particularly, the nonhydratable phospholipids, when they contain them, and to reduce their iron content.

To this end, the degumming process of the invention consists in mixing the fatty substance to be treated
35 with a reactive aqueous solution of a complexing agent chosen from the group including citric acid, phosphoric acid, oxalic acid, tartaric acid, acids of the amino-carboxylic type, acids of the polyhydroxycarboxylic type, polycarboxylic acids, the salts of these acids and

mixtures of two or more of these substances and of an emulsifier of the anionic, cationic, zwitterionic or nonionic type or generated in situ by partial neutralization of the free fatty acids present in the fatty substance, the said solution making it possible of extracted [sic] the phospholipids present in the said fatty substance, the said mixing being carried out by adding, all at once, the aqueous solution of complexing agent and of emulsifier to the fatty substance or vice versa and by subjecting the whole to an intense stirring the rate of which lies between 500 and 15000 revolutions/minute, so as to form a fine emulsion.

According to a particular embodiment of the process of the invention the abovementioned stirring rate lies between 1200 and 10,000 revolutions/minute.

According to a particular embodiment of the process of the invention the abovementioned mixing is done at a temperature of 20 to 100°C, advantageously of 60 to 90°C.

According to a particularly advantageous embodiment of the invention the complexing agent is trisodium citrate or is an acid of the aminocarboxylic type, such as ethylenediaminetetraacetic acid or the disodium or trisodium salt of the latter, and the emulsifier is of the anionic type and consists of sodium lauryl sulfate, of the nonionic type and consists of one or more monoglycerides, or is generated in situ and is sodium and/or potassium carboxylate.

According to another embodiment of the invention the degumming process consists in dispersing the fatty substance in the form of fine droplets in a reactive aqueous solution of a complexing agent chosen from the group including citric acid, phosphoric acid, oxalic acid, tartaric acid, acids of the aminocarboxylic type, acids of the polyhydroxycarboxylic type, polycarboxylic acids, the salts of these acids and mixtures of two or more of these substances and of an emulsifier of the anionic, cationic, zwitterionic or nonionic type or generated in situ by partial neutralization of the free

fatty acids present in the fatty substance, the said solution making it possible to extract the phospholipids present in the said fatty substance.

5 A further subject of the invention is the degummed oils and fats obtained in accordance with the process described above.

Other details and special features of the invention will emerge from the description given below by way of nonlimiting example of some embodiments of the invention.
10

As already stated above, the present invention proposes to degum fatty substances such as crude or delecithinated, animal or vegetable oils or fats, by bringing the fatty substance to be treated into contact
15 with a reactive aqueous solution of a complexing agent and of an emulsifier making it possible to hydrate not only the hydratable phospholipids but above all and in particular the nonhydratable phospholipids if the fatty substance contains them. As already emphasized above, the
20 dissociation and the hydration of the nonhydratable phospholipids, such as phosphatidic acid and phosphatidyl ethanolamine, which are combined with divalent and trivalent metals (Ca^{++} , Mg^{++} , Fe^{++} or Fe^{+++}) is a difficult reaction. On the other hand, the phosphatidic acid and
25 phosphatidyl ethanolamine which are combined with monovalent metals (Na^+ , K^+) or even an H^+ cation are easily hydrated and removed from the fatty substance. Until now, complex ionization reactions in the presence of an acid, followed by a shift in equilibrium in the presence of
30 sodium hydroxide enabled this objective to be attained, but nevertheless required a number of separations by centrifuging in order to remove the nonhydratable phospholipids. In accordance with the invention, the oil or the fat to be degummed and the aqueous solution of
35 complexing agent and of emulsifier are mixed by adding, all at once, the aqueous solution to the oil or the fat or vice versa, and by subjecting the whole to an intense stirring the rate of which lies between 500 and 15,000 revolutions/minute and advantageously between

1,200 and 10,000 revolutions/minute, for a period generally of 10 seconds to 5 minutes. The purpose of this intense mixing is, in fact, to disperse the aqueous phase containing the reactants brought into contact (complexing agent and emulsifier) intensely in the oil or the fat so as to form a fine emulsion. The mixing of the fatty substance/aqueous solution of the reactants brought into contact is generally done at a temperature of the order of 20 to 100°C, but a temperature lying between 60 and 90°C is advantageously employed. A solution of sodium chloride whose concentration varies between 0.1 and 10 % may be added to the aqueous phase thus formed and the latter is then separated by gravity separation or centrifuging so as to obtain a degummed fatty substance essentially free from phospholipids. The degummed fatty substance is then either dried and then treated with a bleaching earth or treated directly without drying. The total content of phospholipids, expressed in the form of phosphorus, is much lower than 10 ppm after degumming. Furthermore, an iron content is obtained which is lower than 0.2 ppm, the value required for good conservation of the oil (A. J. Dijkstra, B. Cleenewerk F.S.T. 317-322, 1992). The physical refining of the fatty substance, which is performed after its degumming, therefore now requires only a small quantity of bleaching earth, of the same order as that employed for the chemical refining.

According to the invention, the complexing agents have a much higher affinity constant for the divalent cations than for the monovalent cations; as a result, they displace and preferentially complex the Ca^{++} , Mg^{++} , Fe^{++} and Fe^{+++} cations. The phosphatidic acid and the phosphatidyl ethanolamine which are thus released are therefore easily hydrated in a sodium form. This reaction of complexing of the divalent or trivalent cations (Mg, Ca, Fe) by the complexing agent requires the preliminary dissociation of the phospholipid-divalent cation complex. This dissociation requires both the presence of a complexing agent chosen from the group including citric acid, phosphoric acid, oxalic acid, tartaric acid, acids

of the aminocarboxylic type, acids of the polyhydroxy-carboxylic type, polycarboxylic acids, the salts of these acids and mixtures of two or more of these substances and of an emulsifier of the anionic, cationic, zwitterionic or nonionic type or generated in situ by partial neutralization of the free fatty acids present in the fatty substance and, as has just been stated, the use of an intense stirring and of a temperature which is preferably at least 60°C, advantageously from 60 to 90°C. Examples of preferred complexing agents employed within the scope of the present invention are trisodium citrate or acids of the aminocarboxylic type, such as ethylenediamine-tetraacetic acid or the disodium and trisodium salts of the latter. The complexing agent will be employed at least in a stoichiometric quantity in relation to the quantity of nonhydratable phospholipids or of total cations (Mg, Ca, Fe) which are present in the fatty substance to be treated. The emulsifier, for its part, is of the anionic, cationic, zwitterionic or nonionic type. The anionic emulsifier, such as sodium lauryl sulfate, is particularly suitable. The emulsifier may also be generated in situ by partial neutralization of the free fatty acids present in the fatty substance. Emulsifiers produced in this way are, for example, sodium and potassium carboxylates. Monoglycerides and their mixtures will be mentioned by way of nonlimiting examples of nonionic emulsifiers.

The quantity of water of the aqueous solution-fatty substance mixture may vary between 0.1 % and 99 % by weight according to the separation conditions employed. As already stated above, the reaction normally takes place between 10 seconds and 5 minutes but can be shortened or last longer if one of the parameters is modified, for example the quantity of water employed, the reaction temperature or the type of reactants brought into contact.

The degumming of soya oils as well as rapeseed, cotton, groundnut, sunflower and corn oils has been successfully carried out by employing the process of the

invention. As already mentioned, the process of the invention is particularly suitable for the degumming of fatty substances containing phospholipids consisting essentially of nonhydratable phospholipids, but is also
5 suitable for the degumming of fatty substances which are lean in nonhydratable phospholipids, so as better to remove some gums or mucilages. The degumming is carried out noncontinuously or continuously, followed by a separation by gravity separation or centrifuging. Washing
10 with water after the degumming of the fatty substance is beneficial but absolutely unnecessary.

The fatty substance such as oil may also be dispersed in the form of fine droplets in an aqueous solution containing the chemical reactants. This technique, described in Belgian Patent No. 595,219, employs
15 a column fitted with a jacket and a distribution system into which the fatty substance or the oil is injected continuously in an extremely divided form. An infinite number of oil droplets is thus formed, and these rise
20 slowly countercurrentwise in the aqueous solution. After coalescence at the top of the column, these oil droplets are separated continuously by gravity separation or centrifuging. In general, the reaction may be carried out in a countercurrent extractor or in a pulsed column for
25 liquid/liquid extraction. It is quite obvious that in the case of the use of this technique for dispersion of the fatty substance in the form of fine droplets in the aqueous solution of complexing agent and of emulsifier, the dispersion will also be done at a temperature of
30 between 20 and 100°C and advantageously between 60 and 90°C. The complexing agents and emulsifiers employed will be the same ones as those illustrated above.

Examples of degumming of fatty material, carried out on the basis of the process in accordance with the
35 invention are given below.

EXAMPLE 1

7 g of delecithinated soya oil whose phospholipid content, expressed in the form of phosphorus, is 80 ppm, and whose acidity, expressed as oleic acid, is 0.32 %, is

are heated to 75°C in a beaker. 21 ml of an aqueous solution made up of 5-millimolar di- or trisodium ethylenediaminetetraacetate salt and of 1.7-millimolar sodium lauryl sulfate are also heated to 75°C. The aqueous solution is added all at once to the oil. The mixture is stirred intensely for 45 seconds with the aid of an Ultra-Turax (type 725 = Janke & Kunkel KG) at 9,500 revolutions/minute.

The emulsion thus obtained is broken by adding 10 ml of a saturated sodium chloride solution or centrifuged directly at 5,000 revolutions/minute.

The phosphorus content, determined by the method of calorimetric [sic] determination of phosphorus (AOCS ca 12-55), is 6 ppm. The cation content, determined by atomic absorption according to the IUPAC method 2,631, is given in ppm.

CATION	BEFORE TREATMENT	AFTER TREATMENT
Magnesium	18	0.2
Calcium	46	1
Iron	0.55	0.04

By treating 7 g of delecithinated rapeseed oil in the same way as above, a phosphorus content of 5 ppm is obtained, determined by the same method of determination.

EXAMPLE 2

300 g of delecithinated soya oil are heated to 75°C in a beaker. 900 ml of an aqueous solution made up of 5-millimolar di- or trisodium ethylenediaminetetraacetate salt and of 1.7-millimolar sodium lauryl sulfate are also heated to 75°C. The aqueous solution is added all at once to the oil. The mixture is stirred intensely for 45 seconds with the aid of an Ultra-Turax (type T45 = Janke & Kunkel KG) at 10,000 revolutions/minute.

The emulsion thus obtained is broken by adding 400 ml of a saturated sodium chloride solution or centrifuged directly at 5,000 revolutions/minute.

The phosphorus content, determined by the method of calorimetric determination of phosphorus (AOCS ca

12-55), is 6 ppm. The cation content, determined by atomic absorption according to IUPAC method 2,631, is given in ppm.

5 The results below are given for two different soya oils.

SOYA OIL No. 1	BEFORE TREATMENT	AFTER TREATMENT
Phosphorus (ppm)	73	5.5
Acidity (oleic acid) %	0.32	
Calcium	46	1
Magnesium	16	< 0.2
Iron	0.55	0.04

SOYA OIL No. 2	BEFORE TREATMENT	AFTER TREATMENT
Phosphorus (ppm)	122	6.5
Acidity (oleic acid) %	4.24	
Calcium	68	1
Magnesium	36	< 0.2
Iron	4.9	0.05

EXAMPLE 3

300 g of delecithinated soya oil whose phospho-
lipid content, expressed in the form of phosphorus, is
25 80 ppm and whose acidity, expressed as oleic acid, is
0.32 %, are heated to 75°C in a beaker. 900 ml of an
aqueous solution made up of 10-millimolar trisodium
citrate and 1.7-millimolar sodium lauryl sulfate are also
heated to 75°C. The aqueous solution is added all at once
30 to the oil. The mixture is stirred intensely for
45 seconds with the aid of an Ultra-Turax (type
T 45 = Janke & Kunkel KG) at 10,000 revolutions/minute.

The emulsion thus obtained is broken by adding
10 ml of a saturated sodium chloride solution or centri-
35 fugal directly at 5,000 revolutions/minute.

The phosphorus content, determined by the method of colorimetric determination of phosphorus (AOCS ca 12-55), and the cation content, determined by atomic absorption according to the IUPAC method 2,631, is given below.

	BEFORE TREATMENT	AFTER TREATMENT
Phosphorus	80	2.2
Iron	0.55	0.03

EXAMPLE 4

According to the conditions described in Example 1, the test was performed in the presence of various emulsifiers the concentration of which, which is 1.7-millimolar, remains constant.

The Table below gives the phosphorus content after treatment with two different soya oils.

	SOYA OIL No. 1	SOYA OIL No. 2
Acidity % (oleic ac.) % Phosphorus ppm	0.32 73	4.24 122
EMULSIFIERS	RESIDUAL PHOSPHORUS AFTER TREATMENT	RESIDUAL PHOSPHORUS AFTER TREATMENT
ANIONIC		
Diocetyl sulfosuccinate	5.2	7.1
Na lauryl sulfate	6.1	7.3
CATIONIC		
Cetylpyridinium	4.0	6.3
Dodecyltrimethylammonium	5.5	5.2
Hexadecyltrimethylammonium	4.5	7.4
Tetradecyltrimethylammonium	4.9	5.1
ZWITTERIONIC		
Lauryl sulfobetaine	4.7	7.3
Tetramethylsulfobetaine	5.8	6.6
NONIONIC		
Triton X100	3.1	3.3
Triton X114	2.3	3.1
Tween 20	3.5	3.4

CLAIMS

1. Process for degumming a fatty substance such as a crude or delecithinated, animal or vegetable oil or fat, characterized in that it includes the mixing of this
5 fatty substance with a reactive aqueous solution of a complexing agent chosen from the group including citric acid, phosphoric acid, oxalic acid, tartaric acid, acids of the aminocarboxylic type, acids of the polyhydroxy-carboxylic type, polycarboxylic acids, the salts of these
10 acids and mixtures of two or more of these substances and of an emulsifier of the anionic, cationic, zwitterionic or nonionic type or generated in situ by partial neutralization of the free fatty acids present in the fatty substance, the said solution making it possible to
15 extract the phospholipids present in the said fatty substance, the said mixing being carried out by adding, all at once, the aqueous solution of complexing agent and of emulsifier to the fatty substance or vice versa and by subjecting the whole to an intense stirring the rate of
20 which lies between 500 and 15000 revolutions/minute, so as to form a fine emulsion.
2. Process in accordance with claim 1, characterized in that the rate of stirring is 1,200 to 10,000 revolutions/minute.
- 25 3. Process in accordance with either of claims 1 and 2, characterized in that the mixing is done at a temperature of 20 to 100°C.
4. Process in accordance with claim 3, characterized in that the temperature employed lies between 60 and
30 90°C.
5. Process in accordance with any one of claims 1 to 4, characterized in that, after the abovementioned mixing, the aqueous phase thus formed is separated off to obtain a degummed fatty substance essentially free from
35 phospholipids.
6. Process in accordance with any one of claims 1 to 5, characterized in that the abovementioned mixing is performed before the operation of physical refining of



the fatty substance.

7. Process for degumming a fatty substance such as a crude or delecithinated, animal or vegetable oil or fat, characterized in that it includes the dispersion of the fatty substance in the form of fine droplets in a reactive aqueous solution of a complexing agent chosen from the group including citric acid, phosphoric acid, oxalic acid, tartaric acid, acids of the aminocarboxylic type, acids of the polyhydroxycarboxylic type, polycarboxylic acids, the salts of these acids and mixtures of two or more of these substances and of an emulsifier of the anionic, cationic, zwitterionic or nonionic type or generated in situ by partial neutralization of the free fatty acids present in the fatty substance, the said solution making it possible to extract the phospholipids present in the said fatty substance.

8. Process in accordance with claim 7, characterized in that the dispersion is done at a temperature of 20 to 100°C.

9. Process in accordance with claim 8, characterized in that the temperature employed lies between 60 and 90°C.

10. Process in accordance with any one of claims 1 to 9, characterized in that the phospholipids which the fatty substance contains are essentially made of non-hydratable phospholipids.

11. Process in accordance with any one of claims 1 to 10, characterized in that the complexing agent is trisodium citrate or is of the aminocarboxylic type, such as ethylenediaminetetraacetic acid or the disodium or trisodium salt of the latter.

12. Process in accordance with any one of claims 1 to 11, characterized in that the complexing agent is employed at least in stoichiometric quantity relative to the quantity of nonhydratable phospholipids present in the fatty substance.

13. Process in accordance with any one of claims 1 to 12, characterized in that the emulsifier is of the anionic type and consists of sodium lauryl sulfate.



14. Process in accordance with any one of claims 1 to 12, characterized in that the emulsifier is generated in situ and is sodium and/or potassium carboxylate.

15. Process in accordance with any one of claims 1 to 5 12, characterized in that the emulsifier is nonionic and consists of one or more monoglycerides.

16. Fatty substance as obtained by the process in accordance with any one of claims 1 to 15.



* C L M S *

ABSTRACT

"Process for degumming a fatty substance and fatty
substance thus obtained"

Process for degumming a fatty substance such as a crude or delecithinated, animal or vegetable oil or fat, including bringing this fatty substance into contact with an aqueous solution of a complexing agent and of a detergent and/or emulsifier making it possible to hydrate the phospholipids present in the latter, and fatty substance thus obtained.

6-163132