


(I);

or a pharmaceutically acceptable salt thereof; and (b) radiation therapy; in amounts effective for treating the brain tumor.


[0012a] In yet another aspect, there is provided a use of a compound of formula (I):



(I);

or a pharmaceutically acceptable salt thereof; in the preparation of a medicament for treating glioma characterized by the presence of an IDH1 mutation or an IDH2 mutation, wherein the IDH1 or IDH2 mutation results in accumulation of R(-)-2-hydroxyglutarate, wherein the medicament is to be administered with radiation therapy in amounts effective for treating glioma; wherein said compound of formula (I), or a pharmaceutically acceptable salt thereof and the radiation are to be administered concurrently or wherein said compound of formula (I) is to be administered after radiation.


[0013] In another aspect, the invention provides a method for treating a brain tumor in a patient in need thereof comprising administering to the patient (a) a compound of formula (I)



(I);

or a pharmaceutically acceptable salt thereof; and (b) one or more additional therapeutic agents; in amounts effective for treating the brain tumor.

[0014] In another aspect, the invention provides a method for treating a brain tumor in a patient in need thereof comprising administering to the patient (a) a compound of formula (I)




(I);

or a pharmaceutically acceptable salt thereof; (b) radiation therapy; and (c) one or more additional therapeutic agents; in amounts effective for treating the brain tumor.

CLAIMS

1. Use of a compound of formula (I):



(I);

or a pharmaceutically acceptable salt thereof; in the preparation of a medicament for treating glioma characterized by the presence of an IDH1 mutation or an IDH2 mutation, wherein the IDH1 or IDH2 mutation results in accumulation of R(-)-2-hydroxyglutarate, wherein the medicament is to be administered with radiation therapy in amounts effective for treating glioma; wherein said compound of formula (I), or a pharmaceutically acceptable salt thereof and the radiation are to be administered concurrently or wherein said compound of formula (I) is to be administered after radiation.

2. The use of claim 1, wherein said compound of formula (I), or a pharmaceutically acceptable salt thereof, is to be administered in an amount of from 10 to 50 mg/day.

3. The use of claim 1, wherein said radiation therapy is to be administered in the form of x-ray radiation.

4. The use of claim 1, wherein said radiation therapy is to be administered in a cumulative dose of from 30 to 60 Gy.

5. The use of claim 1, wherein the IDH1 mutation is an R132X mutation.

6. The use of claim 5, wherein the IDH1 mutation is an R132H or R132C mutation.

7. The use of claim 1, wherein the IDH2 mutation is an R140X or R172X mutation.

8. The use of claim 7, wherein the IDH2 mutation is an R140Q, R140W, or R140L mutation.

9. The use of claim 7, wherein the IDH2 mutation is an R172K or R172G mutation.

10. The use of any one of claims 1 to 9, wherein said compound of formula (I) is to be administered in non-salt form.