

FUNCTIONALIZED GEL BEADS

CROSS REFERENCE

[0001] This application claims the benefit of U.S. Provisional Application No. 62/586,784, filed November 15, 2017, and U.S. Provisional Application No. 62/629,561, filed February 12, 2018, each of which applications is entirely incorporated herein by reference.

BACKGROUND

[0002] Samples may be processed for various purposes, such as identification of a type of moiety within the sample. The sample may be a biological sample. The biological samples may be processed for various purposes, such as detection of a disease (e.g., cancer) or identification of a particular species. There are various approaches for processing samples, such as polymerase chain reaction (PCR) and sequencing.

[0003] Biological samples may be processed within various reaction environments, such as partitions. Partitions may be wells or droplets. Droplets or wells may be employed to process biological samples in a manner that enables the biological samples to be partitioned and processed separately. For example, such droplets may be fluidically isolated from other droplets, enabling accurate control of respective environments in the droplets.

[0004] Biological samples in partitions may be subjected to various processes, such as chemical processes or physical processes. Samples in partitions may be subjected to heating or cooling, or chemical reactions, such as to yield species that may be qualitatively or quantitatively processed.

SUMMARY

[0005] The present disclosure provides methods for use in various sample processing and analysis applications. The methods provided herein may provide barcode molecules comprising one or more sequences such as a barcode sequence and a functional sequence. Barcode molecules may be attached to supports such as beads. Such methods may employ combinatorial (e.g., split pool) ligation reactions and may be useful, for example, in controlled analysis and processing of analytes such as biological particles, nucleic acid molecules, proteins, and perturbation agents.

[0006] In an aspect, the present disclosure provides a method for generating a plurality of barcode molecules, comprising: (a) providing a plurality of molecules coupled to a plurality of supports; and combinatorially assembling the plurality of barcode molecules coupled to the plurality of supports by coupling one or more molecules each comprising one or more segments to each of the plurality of molecules, wherein the plurality of barcode molecules comprises (i) a

first set of barcode molecules coupled to a support of the plurality of supports and (ii) a second set of barcode molecules coupled to the support, wherein barcode molecules of the first set of barcode molecules are different than barcode molecules of the second set of barcode molecules.

[0007] In some embodiments, the barcode molecules of the first set of barcode molecules and the barcode molecules of the second set of barcode molecules comprise barcode sequences that are different from barcode sequences of barcode molecules coupled to other supports of the plurality of supports.

[0008] In some embodiments, barcode sequences of the barcode molecules of the first set of barcode molecules and the barcode molecules of the second set of barcode molecules are identical.

[0009] In some embodiments, the plurality of supports is a plurality of beads, and wherein the barcode molecules of the first set of barcode molecules and the barcode molecules of the second set of barcode molecules are coupled to a bead of the plurality of beads. In some embodiments, at least a subset of the barcode molecules of the first set of barcode molecules and/or the barcode molecules of the second set of barcode molecules is coupled to an interior of the bead. In some embodiments, the barcode molecules of the first set of barcode molecules and the barcode molecules of the second set of barcode molecules are releasably coupled to the bead. In some embodiments, the barcode molecules of the first set of barcode molecules and the barcode molecules of the second set of barcode molecules are releasably coupled to the bead through chemical cross-linkers. In some embodiments, the plurality of beads is a plurality of gel beads. In some embodiments, the plurality of beads is dissolvable or disruptable.

[0010] In some embodiments, (b) comprises sequentially coupling multiple molecules to each of the plurality of molecules.

[0011] In some embodiments, in (b) the one or more molecules comprise a first molecule comprising a first segment and a second molecule comprising a second segment, and wherein the combinatorially assembling comprises coupling the first molecule to a molecule of the plurality of molecules and coupling the second molecule to the first molecule, wherein the molecule is coupled to the support. In some embodiments, the method further comprises coupling a third molecule comprising a third segment to the second molecule. In some embodiments, the third segment is different than the first segment or the second segment. In some embodiments, the third segment is different than the first segment and the second segment. In some embodiments, the first segment of the first molecule is different than the second segment of the second molecule.

[0012] In some embodiments, wherein the first molecule comprises a double-stranded region comprising the first segment and a single-stranded region. In some embodiments, the single-

stranded region of the first molecule has six or fewer nucleotides. In some embodiments, the single-stranded region of the first molecule has two nucleotides. In some embodiments, the sequence of the single-stranded region of the first molecule is selected from the group consisting of CA, GT, AC, and TG. In some embodiments, the first molecule further comprises an additional single-stranded region configured to couple to the molecule of the plurality of molecules. In some embodiments, the single-stranded region and the additional single-stranded region of the first molecule are included in the same strand of the first molecule. In some embodiments, the second molecule comprises a double-stranded region comprising the second segment and a single-stranded region. In some embodiments, the single-stranded region of the second molecule comprises a sequence complementary to the sequence of the single-stranded region of the first molecule. In some embodiments, the single-stranded region of the second molecule has six or fewer nucleotides. In some embodiments, the single-stranded region of the second molecule has two nucleotides. In some embodiments, the second molecule further comprises an additional single-stranded region. In some embodiments, the single-stranded region and the additional single-stranded region of the second molecule are included in different strands of the second molecule.

[0013] In some embodiments, in (b) the one or more molecules comprise a third molecule comprising a third segment and a fourth molecule comprising a fourth segment, and wherein the combinatorially assembling comprises coupling the third molecule to an additional molecule of the plurality of molecules and coupling the fourth molecule to the third molecule, wherein the additional molecule is coupled to the support. In some embodiments, (i) the first molecule comprises a double-stranded region comprising the first segment and a single-stranded region and (ii) the third molecule comprises a double-stranded region comprising the third segment and a single-stranded region. In some embodiments, the first segment of the first molecule and the third segment are the same. In some embodiments, the second segment of the second molecule and the fourth segment of the fourth molecule are the same. In some embodiments, the single-stranded region of the first molecule and the single-stranded region of the third molecule are the same. In some embodiments, the single-stranded region of the first molecule and the single-stranded region of the third molecule are different. In some embodiments, the single-stranded region of the first molecule and the single-stranded region of the third molecule each have six or fewer nucleotides. In some embodiments, the molecule and the additional molecule coupled to the support are the same. In some embodiments, the molecule and the additional molecule coupled to the support are different. In some embodiments, the second molecule comprises a first functional sequence and the fourth molecule comprises a second functional sequence, which first functional sequence is different than the second functional sequence.

[0014] In some embodiments, the one or more molecules comprise a plurality of first molecules comprising the first segment, a plurality of second molecules comprising the second segment, a plurality of third molecules comprising the third segment, and a plurality of fourth molecules comprising the fourth segment, and wherein the combinatorially assembling comprises (i) coupling first molecules of the plurality of first molecules to molecules of the plurality of molecules and coupling second molecules of the plurality of second molecules to the first molecules, and (ii) coupling third molecules of the plurality of third molecules to additional molecules of the plurality of molecules and coupling fourth molecules of the plurality of fourth molecules to the third molecules, wherein the molecules and the additional molecules of the plurality of molecules are coupled to the support. In some embodiments, among the one or more molecules, the plurality of first molecules comprises a greater number of molecules than the plurality of third molecules. In some embodiments, among the one or more molecules, the plurality of second molecules comprises a greater number of molecules than the plurality of fourth molecules. In some embodiments, the molecules of the plurality of molecules comprises a greater number of molecules than the additional molecules of the plurality of molecules.

[0015] In some embodiments, (b) comprises ligating the one or more molecules to each of the plurality of molecules coupled to the plurality of supports.

[0016] In some embodiments, (b) comprises partitioning the plurality of molecules coupled to the plurality of supports in separate partitions, and coupling at least a subset of the one or more molecules to molecules of the plurality of molecules in the separate partitions. In some embodiments, (b) further comprises partitioning the plurality of molecules coupled to the plurality of supports in one or more additional partitions, and coupling at least a subset of the one or more molecules to molecules of the plurality of molecules in the one or more additional partitions. In some embodiments, the separate partitions are wells. In some embodiments, the separate partitions are droplets.

[0017] In some embodiments, the barcode sequences are nucleic acid sequences. In some embodiments, the barcode sequences are amino acid sequences. In some embodiments, barcode molecules of the other supports of the plurality of supports comprise different barcode sequences. In some embodiments, each other support of the other supports of the plurality of supports comprises barcode molecules comprising a different barcode sequence.

[0018] In some embodiments, the one or more segments are at least one nucleotide in length.

[0019] In some embodiments, each of the barcode molecules of the first set of barcode molecules and the barcode molecules of the second set of barcode molecules comprises a unique identifier that is different than unique identifiers of other barcode molecules coupled to the support.

[0020] In some embodiments, the barcode molecules of the first set of barcode molecules and the barcode molecules of the second set of barcode molecules comprise at least 100,000 barcode molecules.

[0021] In some embodiments, each of the barcode molecules of the first set of barcode molecules and the barcode molecules of the second set of barcode molecules comprise a segment separated from the one or more segments by a sequence that is the same across at least a subset of the plurality of barcode molecules.

[0022] In some embodiments, the one or more segments are different for at least a subset of the plurality of barcode molecules.

[0023] In some embodiments, barcode molecules of the first set of barcode molecules include a first sequence for use with a first assay and barcode molecules of the second set of barcode molecules include a second sequence for use with a second assay, wherein the first assay is different than the second assay.

[0024] In some embodiments, barcode molecules of the first set of barcode molecules and barcode molecules of the second set of barcode molecules include both identical barcode sequences and different barcode sequences.

[0025] In some embodiments, barcode molecules of the first set of barcode molecules include first functional sequences and barcode molecules of the second set of barcode molecules include second functional sequences, wherein the first functional sequences and the second functional sequences are different. In some embodiments, the first set of barcode molecules coupled to the support comprises a greater number of barcode molecules than the second set of barcode molecules coupled to the support. In some embodiments, the first set of barcode molecules coupled to the support comprises at least 10,000 barcode molecules. In some embodiments, the second set of barcode molecules coupled to the support comprises at least 10,000 barcode molecules.

[0026] In some embodiments, the method further comprises contacting a solution comprising the first molecule and the second molecule coupled to the molecule of the plurality of molecules with an exonuclease. In some embodiments, the second molecule comprises a phosphorothioate moiety.

[0027] In another aspect, the present disclosure provides a method of processing a plurality of barcode molecules, comprising: (a) providing a plurality of molecules coupled to a plurality of supports; (b) subjecting a subset of molecules of the plurality of molecules to conditions sufficient to couple first molecules to the subset of molecules, wherein the subset of molecules are coupled to a support of the plurality of supports; (c) subjecting the first molecules coupled to the subset of molecules to conditions sufficient to couple second molecules to the first

molecules, thereby generating the plurality of barcode molecules and by-products, which plurality of barcode molecules comprise the second molecules coupled to the first molecules; and (d) contacting the products and by-products with a molecule capable of degrading the by-products.

[0028] In some embodiments, the molecule capable of degrading the by-products is an exonuclease.

[0029] In some embodiments, the second molecules comprise phosphorothioate moieties. In some embodiments, the phosphorothioate moieties are disposed at an end of the second molecules.

[0030] In some embodiments, the plurality of barcode molecules comprises a plurality of barcode sequences. In some embodiments, the plurality of barcode sequences of the plurality of barcode molecules coupled to the support are identical. In some embodiments, the plurality of barcode sequences comprises a plurality of nucleic acid barcode sequences.

[0031] In some embodiments, the plurality of supports comprises a plurality of beads, and wherein the plurality of barcode molecules are coupled to a bead of the plurality of beads. In some embodiments, the plurality of beads is a plurality of gel beads. In some embodiments, the plurality of beads is dissolvable or disruptable. In some embodiments, the plurality of barcode molecules is releasably coupled to the bead.

[0032] In some embodiments, (b) comprises ligating the first molecules to the molecules.

[0033] In some embodiments, (c) comprises ligating the second molecules to the first molecules.

[0034] In some embodiments, the method further comprises partitioning the plurality of molecules coupled to the plurality of supports in separate partitions. In some embodiments, (b) comprises coupling the first molecules to the subset of molecules within a partition of the separate partitions. In some embodiments, the method further comprises, subsequent to (b), partitioning the plurality of molecules coupled to the plurality of supports in one or more additional partitions. In some embodiments, (c) comprises coupling the second molecules to the first molecules within a partition of the one or more additional partitions. In some embodiments, the separate partitions are wells. In some embodiments, the separate partitions are droplets. In some embodiments, prior to (d), the plurality of barcode molecules coupled to the support and the by-products are recovered from the separate partitions.

[0035] In some embodiments, the first molecules and the second molecules are nucleic acid molecules, which nucleic acid molecules each comprise at least two nucleotides.

[0036] In some embodiments, the plurality of barcode molecules coupled to the support comprises at least 10,000 barcode molecules.

[0037] In some embodiments, the plurality of barcode molecules coupled to the support comprises a plurality of first barcode molecules and a plurality of second barcode molecules, which plurality of first barcode molecules and plurality of second barcode molecules are different. In some embodiments, the plurality of first barcode molecules comprise a plurality of first barcode sequences and the plurality of second barcode molecules comprise a plurality of second barcode sequences, which first barcode sequences are different than the second barcode sequences. In some embodiments, the plurality of first barcode molecules comprise first functional sequences and the plurality of second barcode molecules comprise second functional sequences, which first functional sequences are different than the second functional sequences.

[0038] In some embodiments, the plurality of barcode molecules comprise a plurality of functional sequences. In some embodiments, the functional sequences are poly(T) sequences.

[0039] In a further aspect, the present disclosure provides a composition comprising a support comprising a plurality of nucleic acid barcode molecules coupled thereto, which plurality of nucleic acid barcode molecules comprise one or more sequences selected from the sequences included in Table 1 or Table 2.

[0040] In some embodiments, the plurality of nucleic acid barcode molecules comprise one or more sequences from the sequences included in Table 1.

[0041] In some embodiments, the plurality of nucleic acid barcode molecules comprise one or more sequences from the sequences included in Table 2. In some embodiments, the plurality of nucleic acid barcode molecules comprise the sequence CCTTAGCCGCTAATAGGTGAGC. In some embodiments, the plurality of nucleic acid barcode molecules comprise the sequence TTGCTAGGACCGCCTTAAAGC.

[0042] In some embodiments, each nucleic acid barcode molecule of the plurality of nucleic acid barcode molecules comprises one or more sequences selected from the sequences included in Table 1 or Table 2.

[0043] In some embodiments, the plurality of nucleic acid barcode molecules comprise a first plurality of nucleic acid barcode molecules and a second plurality of nucleic acid barcode molecules, which first plurality of nucleic acid barcode molecules comprise one or more sequences selected from the sequences included in Table 1 or Table 2 and which second plurality of nucleic acid barcode molecules do not comprise a sequence included in Table 1 or Table 2. In some embodiments, the second plurality of nucleic acid barcode molecules are configured to interact with ribonucleic acid molecules. In some embodiments, the second plurality of nucleic acid barcode molecules comprise a poly(T) sequence. In some embodiments, the first plurality of nucleic acid barcode molecules are configured to interact with deoxyribonucleic acid molecules.

[0044] In some embodiments, the first plurality of nucleic acid barcode molecules comprises at least 10,000 nucleic acid barcode molecules. In some embodiments, the second plurality of nucleic acid barcode molecules comprises at least 10,000 nucleic acid barcode molecules. In some embodiments, among the plurality of nucleic acid barcode molecules, the first plurality of nucleic acid barcode molecules comprises a greater number of nucleic acid barcode molecules than the second plurality of nucleic acid barcode molecules. In some embodiments, among the plurality of nucleic acid barcode molecules, the second plurality of nucleic acid barcode molecules comprises a greater number of nucleic acid barcode molecules than the first plurality of nucleic acid barcode molecules.

[0045] In some embodiments, the support is a bead. In some embodiments, the bead is a gel bead. In some embodiments, the bead is dissolvable or disruptable. In some embodiments, the plurality of nucleic acid barcode molecules is releasably coupled to the bead.

[0046] In some embodiments, the plurality of nucleic acid barcode molecules coupled to the support comprise a plurality of barcode sequences. In some embodiments, the plurality of barcode sequences is the same for each of the plurality of nucleic acid barcode molecules. In some embodiments, the plurality of barcode sequences is a plurality of nucleic acid barcode sequences.

[0047] In some embodiments, the plurality of nucleic acid barcode molecules comprises at least 10,000 nucleic acid barcode molecules.

[0048] In another aspect, the present disclosure provides a kit comprising a plurality of barcode molecules, comprising: a plurality of supports; and a plurality of barcode molecules coupled to the plurality of supports, wherein the plurality of barcode molecules comprises (i) a first set of barcode molecules coupled to a support of the plurality of supports and (ii) a second set of barcode molecules coupled to the support, wherein first barcode molecules of the first set of barcode molecules are different than second barcode molecules of the second set of barcode molecules, and

wherein the first barcode molecules are configured to interact with different target molecules than the second barcode molecules.

[0049] In some embodiments, the first barcode molecules of the first set of barcode molecules and the second barcode molecules of the second set of barcode molecules comprise barcode sequences that are different from barcode sequences of barcode molecules coupled to other supports of the plurality of supports.

[0050] In some embodiments, the first barcode molecules of the first set of barcode molecules and the second barcode molecules of the second set of barcode molecules comprise barcode sequences, wherein the barcode sequences of the first barcode molecules of the first set

of barcode molecules and the second barcode molecules of the second set of barcode molecules are identical.

[0051] In some embodiments, the plurality of supports is a plurality of beads, and wherein the first barcode molecules of the first set of barcode molecules and the second barcode molecules of the second set of barcode molecules are coupled to a bead of the plurality of beads. In some embodiments, at least a subset of the first barcode molecules of the first set of barcode molecules and/or the second barcode molecules of the second set of barcode molecules is coupled to an interior of the bead. In some embodiments, the first barcode molecules of the first set of barcode molecules and the second barcode molecules of the second set of barcode molecules are releasably coupled to the bead. In some embodiments, the first barcode molecules of the first set of barcode molecules and the second barcode molecules of the second set of barcode molecules are releasably coupled to the bead through chemical cross-linkers. In some embodiments, the plurality of beads is a plurality of gel beads. In some embodiments, the plurality of beads is dissolvable or disruptable.

[0052] In some embodiments, the first barcode molecules or the second barcode molecules are configured to interact with deoxyribonucleic acid molecules. In some embodiments, the first barcode molecules or the second barcode molecules are configured to interact with ribonucleic acid molecules. In some embodiments, the first barcode molecules or the second barcode molecules are configured to interact with amino acids, polypeptides or proteins.

[0053] In some embodiments, the first barcode molecules of the first set of barcode molecules and the second barcode molecules of the second set of barcode molecules comprise barcode sequences, which barcode sequences are nucleic acid sequences. In some embodiments, the first barcode molecules of the first set of barcode molecules and the second barcode molecules of the second set of barcode molecules comprise barcode sequences, which barcode sequences are amino acid sequences.

[0054] In some embodiments, each of the first barcode molecules of the first set of barcode molecules and the second barcode molecules of the second set of barcode molecules comprises a unique identifier that is different than unique identifiers of other barcode molecules coupled to the support.

[0055] In some embodiments, the first barcode molecules of the first set of barcode molecules and the second barcode molecules of the second set of barcode molecules comprise at least 100,000 barcode molecules.

[0056] In some embodiments, the first barcode molecules of the first set of barcode molecules and the second barcode molecules of the second set of barcode molecules include both identical barcode sequences and different barcode sequences.

[0057] In some embodiments, the first barcode molecules of the first set of barcode molecules include a first sequence for use with a first assay and the second barcode molecules of the second set of barcode molecules include a second sequence for use with a second assay, wherein the first assay is different than the second assay. In some embodiments, the first sequences of the first barcode molecules of the first set of barcode molecules are configured to interact with first target molecules and the second sequences of the second barcode molecules of the second set of barcode molecules are configured to interact with second target molecules. In some embodiments, the first target molecules are deoxyribonucleic acid molecules and the second target molecules are ribonucleic acid molecules.

[0058] In some embodiments, the first set of barcode molecules coupled to the support comprises a greater number of barcode molecules than the second set of barcode molecules coupled to the support.

[0059] In a further aspect, the present disclosure provides a method for processing a plurality of analytes, comprising: (a) providing a plurality of barcode molecules coupled to a plurality of supports, wherein the plurality of barcode molecules comprises (i) a first set of barcode molecules coupled to a support of the plurality of supports and (ii) a second set of barcode molecules coupled to the support, wherein first barcode molecules of the first set of barcode molecules are different than second barcode molecules of the second set of barcode molecules; (b) partitioning the support of the plurality of supports in a partition, wherein subsequent to partitioning, the partition comprises the plurality of analytes; (c) using (i) a first barcode molecule from the first set of barcode molecules and a first analyte from the plurality of analytes to generate a first barcoded analyte, and (ii) a second barcode molecule from the second set of barcode molecules and a second analyte from the plurality of analytes to generate a second barcoded analyte; and (d) recovering (i) the first barcoded analyte or a derivative thereof and (ii) the second barcoded analyte or a derivative thereof from the partition.

[0060] In some embodiments, the first barcode molecules of the first set of barcode molecules and the second barcode molecules of the second set of barcode molecules comprise barcode sequences that are different from barcode sequences of barcode molecules coupled to other supports of the plurality of supports.

[0061] In some embodiments, the first barcode molecules of the first set of barcode molecules and the second barcode molecules of the second set of barcode molecules comprise barcode sequences, wherein the barcode sequences of the first barcode molecules of the first set of barcode molecules and the second barcode molecules of the second set of barcode molecules are identical. In some embodiments, the plurality of supports is a plurality of beads, and wherein the first barcode molecules of the first set of barcode molecules and the second barcode

molecules of the second set of barcode molecules are coupled to a bead of the plurality of beads. In some embodiments, at least a subset of the first barcode molecules of the first set of barcode molecules and/or the second barcode molecules of the second set of barcode molecules is coupled to an interior of the bead. In some embodiments, the first barcode molecules of the first set of barcode molecules and the second barcode molecules of the second set of barcode molecules are releasably coupled to the bead. In some embodiments, the first barcode molecules of the first set of barcode molecules and the second barcode molecules of the second set of barcode molecules are releasably coupled to the bead through chemical cross-linkers. In some embodiments, the plurality of beads is a plurality of gel beads. In some embodiments, the plurality of beads is dissolvable or disruptable.

[0062] In some embodiments, the plurality of analytes comprises a plurality of deoxyribonucleic acid molecules. In some embodiments, the plurality of analytes comprises a plurality of ribonucleic acid molecules. In some embodiments, the plurality of analytes comprises a plurality of amino acids, polypeptides or proteins.

[0063] In some embodiments, the first analyte is a deoxyribonucleic acid molecule. In some embodiments, the second analyte is a ribonucleic acid molecule.

[0064] In some embodiments, the first barcode molecules of the first set of barcode molecules and the second barcode molecules of the second set of barcode molecules comprise barcode sequences, which barcode sequences are nucleic acid sequences. In some embodiments, the first barcode molecules of the first set of barcode molecules and the second barcode molecules of the second set of barcode molecules comprise barcode sequences, which barcode sequences are amino acid sequences.

[0065] In some embodiments, each of the first barcode molecules of the first set of barcode molecules and the second barcode molecules of the second set of barcode molecules comprises a unique identifier that is different than unique identifiers of other barcode molecules coupled to the support.

[0066] In some embodiments, the first barcode molecules of the first set of barcode molecules and the second barcode molecules of the second set of barcode molecules comprise at least 100,000 barcode molecules.

[0067] In some embodiments, the first barcode molecules of the first set of barcode molecules and the second barcode molecules of the second set of barcode molecules include both identical barcode sequences and different barcode sequences.

[0068] In some embodiments, the first barcode molecules of the first set of barcode molecules include first functional sequences and the second barcode molecules of the second set of barcode molecules include second functional sequences, wherein the first functional

sequences and the second functional sequences are different. In some embodiments, the first functional sequences of the first barcode molecules of the first set of barcode molecules are configured to interact with first analytes of the plurality of analytes and the second functional sequences of the second barcode molecules of the second set of barcode molecules are configured to interact with second analytes of the plurality of analytes. In some embodiments, the first analytes are deoxyribonucleic acid molecules and the second analytes are ribonucleic acid molecules.

[0069] In some embodiments, the first set of barcode molecules coupled to the support comprises a greater number of barcode molecules than the second set of barcode molecules coupled to the support.

[0070] In some embodiments, barcode molecules of the other supports of the plurality of supports comprise different barcode sequences.

[0071] In some embodiments, the partition is a well. In some embodiments, the partition is a droplet.

[0072] In some embodiments, (c) comprises performing one or more nucleic acid extension reactions.

[0073] In some embodiments, (i) the first barcode molecule or a sequence thereof and (ii) the second barcode molecule or a sequence thereof are released from the support.

[0074] In some embodiments, the plurality of analytes is components of a cell. In some embodiments, subsequent to (b), the cell is lysed or permeabilized to provide access to the plurality of analytes.

[0075] In some embodiments, subsequent to (c), the first barcoded analyte and the second barcoded analyte are in solution in the partition.

[0076] Additional aspects and advantages of the present disclosure will become readily apparent to those skilled in this art from the following detailed description, wherein only illustrative embodiments of the present disclosure are shown and described. As will be realized, the present disclosure is capable of other and different embodiments, and its several details are capable of modifications in various obvious respects, all without departing from the disclosure. Accordingly, the drawings and description are to be regarded as illustrative in nature, and not as restrictive.

INCORPORATION BY REFERENCE

[0077] All publications, patents, and patent applications mentioned in this specification are herein incorporated by reference to the same extent as if each individual publication, patent, or patent application was specifically and individually indicated to be incorporated by reference. To

the extent publications and patents or patent applications incorporated by reference contradict the disclosure contained in the specification, the specification is intended to supersede and/or take precedence over any such contradictory material.

BRIEF DESCRIPTION OF THE DRAWINGS

[0078] The novel features of the invention are set forth with particularity in the appended claims. A better understanding of the features and advantages of the present invention will be obtained by reference to the following detailed description that sets forth illustrative embodiments, in which the principles of the invention are utilized, and the accompanying drawings (also “Figure” and “FIG.” herein), of which:

[0079] **FIG. 1** shows an example of a microfluidic channel structure for partitioning individual biological particles.

[0080] **FIG. 2** shows an example of a microfluidic channel structure for delivering barcode carrying beads to droplets.

[0081] **FIG. 3** shows an example of a microfluidic channel structure for co-partitioning biological particles and reagents.

[0082] **FIG. 4** shows an example of a microfluidic channel structure for the controlled partitioning of beads into discrete droplets.

[0083] **FIG. 5** shows an example of a microfluidic channel structure for increased droplet generation throughput.

[0084] **FIG. 6** shows another example of a microfluidic channel structure for increased droplet generation throughput.

[0085] **FIG. 7A** shows a cross-section view of another example of a microfluidic channel structure with a geometric feature for controlled partitioning. **FIG. 7B** shows a perspective view of the channel structure of **FIG. 7A**.

[0086] **FIG. 8** illustrates an example of a barcode carrying bead.

[0087] **FIG. 9A** schematically illustrates a combinatorial double ligation method. **FIG. 9B** shows the results of the combinatorial double ligation method.

[0088] **FIG. 10** schematically illustrates a combinatorial triple ligation method.

[0089] **FIG. 11** shows examples of beads comprising two or more different nucleic acid barcode molecules.

[0090] **FIG. 12** shows examples of differential functionalization of a bead comprising different starter sequences.

[0091] **FIG. 13** schematically illustrates a triple ligation reaction.

- [0092] FIG. 14 illustrates a double ligation reaction to provide a nucleic acid barcode molecule.
- [0093] FIG. 15 illustrates a double ligation reaction to provide two different nucleic acid barcode molecules.
- [0094] FIG. 16 illustrates a double ligation reaction to provide two different nucleic acid barcode molecules.
- [0095] FIG. 17 shows a nucleic acid barcode molecule generated by a triple ligation method.
- [0096] FIG. 18 shows a nucleic acid barcode molecule generated by a triple ligation method.
- [0097] FIG. 19 shows a comparison of the nucleic acid barcode molecules of FIGs. 17 and 18.
- [0098] FIG. 20 illustrates a triple ligation reaction to provide three different nucleic acid barcode molecules.
- [0099] FIG. 21 shows a combinatorial scheme for a triple ligation method.
- [00100] FIG. 22 shows a combinatorial scheme for a triple ligation method.
- [00101] FIG. 23 shows an example of the use of nucleic acid barcode molecules generated by the presently disclosed methods.
- [00102] FIG. 24 shows an example of the use of nucleic acid barcode molecules generated by the presently disclosed methods.
- [00103] FIG. 25 shows a computer system that is programmed or otherwise configured to implement methods provided herein.
- [00104] FIG. 26 shows the decrease in ligation efficiency throughout a triple ligation process.
- [00105] FIG. 27 shows an exonuclease treatment of barcode molecules coupled to gel beads.
- [00106] FIG. 28 shows an exonuclease treatment of barcode molecules coupled to gel beads.
- [00107] FIGs. 29A and 29B show a triple ligation process in which exonuclease treatment increases ligation efficiency.
- [00108] FIGs. 30A and 30B show a triple ligation process in which exonuclease treatment increases ligation efficiency.
- [00109] FIG. 31 shows a scheme relating to off-products in a ligation process.
- [00110] FIG. 32 shows a process in which fluorescent probes were used to examine the effects of exonuclease treatment.
- [00111] FIG. 33A shows a triple ligation process. FIGs. 33B and 33C show comparisons between exonuclease treatment and lack of exonuclease treatment for the triple ligation process shown in FIG. 33A.
- [00112] FIG. 34 shows a process involving both ExoI and ExoIII treatments.
- [00113] FIG. 35 shows a process involving both ExoI and HL-dsDNase treatment.

[00114] FIG. 36 shows comparisons of processes involving no exonuclease treatment, ExoI treatment, and ExoI and dsDNase treatment in which green and red probes are utilized as in FIG. 32.

DETAILED DESCRIPTION

[00115] While various embodiments of the invention have been shown and described herein, it will be obvious to those skilled in the art that such embodiments are provided by way of example only. Numerous variations, changes, and substitutions may occur to those skilled in the art without departing from the invention. It should be understood that various alternatives to the embodiments of the invention described herein may be employed.

[00116] Where values are described as ranges, it will be understood that such disclosure includes the disclosure of all possible sub-ranges within such ranges, as well as specific numerical values that fall within such ranges irrespective of whether a specific numerical value or specific sub-range is expressly stated.

[00117] The term “barcode,” as used herein, generally refers to a label, or identifier, that conveys or is capable of conveying information about an analyte. A barcode can be part of an analyte. A barcode can be independent of an analyte. A barcode can be a tag attached to an analyte (e.g., nucleic acid molecule) or a combination of the tag in addition to an endogenous characteristic of the analyte (e.g., size of the analyte or end sequence(s)). A barcode may be unique. Barcodes can have a variety of different formats. For example, barcodes can include: polynucleotide barcodes; random nucleic acid and/or amino acid sequences; and synthetic nucleic acid and/or amino acid sequences. A barcode can be attached to an analyte in a reversible or irreversible manner. A barcode can be added to, for example, a fragment of a deoxyribonucleic acid (DNA) or ribonucleic acid (RNA) sample before, during, and/or after sequencing of the sample. Barcodes can allow for identification and/or quantification of individual sequencing-reads.

[00118] The term “real time,” as used herein, can refer to a response time of less than about 1 second, a tenth of a second, a hundredth of a second, a millisecond, or less. The response time may be greater than 1 second. In some instances, real time can refer to simultaneous or substantially simultaneous processing, detection or identification.

[00119] The term “subject,” as used herein, generally refers to an animal, such as a mammal (e.g., human) or avian (e.g., bird), or other organism, such as a plant. For example, the subject can be a vertebrate, a mammal, a rodent (e.g., a mouse), a primate, a simian or a human. Animals may include, but are not limited to, farm animals, sport animals, and pets. A subject can be a healthy or asymptomatic individual, an individual that has or is suspected of having a

disease (e.g., cancer) or a pre-disposition to the disease, and/or an individual that is in need of therapy or suspected of needing therapy. A subject can be a patient. A subject can be a microorganism or microbe (e.g., bacteria, fungi, archaea, viruses).

[00120] The term “genome,” as used herein, generally refers to genomic information from a subject, which may be, for example, at least a portion or an entirety of a subject’s hereditary information. A genome can be encoded either in DNA or in RNA. A genome can comprise coding regions (e.g., that code for proteins) as well as non-coding regions. A genome can include the sequence of all chromosomes together in an organism. For example, the human genome ordinarily has a total of 46 chromosomes. The sequence of all of these together may constitute a human genome.

[00121] The terms “adaptor(s)”, “adapter(s)” and “tag(s)” may be used synonymously. An adaptor or tag can be coupled to a polynucleotide sequence to be “tagged” by any approach, including ligation, hybridization, or other approaches.

[00122] The term “sequencing,” as used herein, generally refers to methods and technologies for determining the sequence of nucleotide bases in one or more polynucleotides. The polynucleotides can be, for example, nucleic acid molecules such as deoxyribonucleic acid (DNA) or ribonucleic acid (RNA), including variants or derivatives thereof (e.g., single stranded DNA). Sequencing can be performed by various systems currently available, such as, without limitation, a sequencing system by Illumina®, Pacific Biosciences (PacBio®), Oxford Nanopore®, or Life Technologies (Ion Torrent®). Alternatively or in addition, sequencing may be performed using nucleic acid amplification, polymerase chain reaction (PCR) (e.g., digital PCR, quantitative PCR, or real time PCR), or isothermal amplification. Such systems may provide a plurality of raw genetic data corresponding to the genetic information of a subject (e.g., human), as generated by the systems from a sample provided by the subject. In some examples, such systems provide sequencing reads (also “reads” herein). A read may include a string of nucleic acid bases corresponding to a sequence of a nucleic acid molecule that has been sequenced. In some situations, systems and methods provided herein may be used with proteomic information.

[00123] The term “bead,” as used herein, generally refers to a particle. The bead may be a solid or semi-solid particle. The bead may be a gel bead. The gel bead may include a polymer matrix (e.g., matrix formed by polymerization or cross-linking). The polymer matrix may include one or more polymers (e.g., polymers having different functional groups or repeat units). Polymers in the polymer matrix may be randomly arranged, such as in random copolymers, and/or have ordered structures, such as in block copolymers. Cross-linking can be via covalent, ionic, or inductive, interactions, or physical entanglement. The bead may be a macromolecule.

The bead may be formed of nucleic acid molecules bound together. The bead may be formed via covalent or non-covalent assembly of molecules (e.g., macromolecules), such as monomers or polymers. Such polymers or monomers may be natural or synthetic. Such polymers or monomers may be or include, for example, nucleic acid molecules (e.g., DNA or RNA). The bead may be formed of a polymeric material. The bead may be magnetic or non-magnetic. The bead may be rigid. The bead may be flexible and/or compressible. The bead may be disruptable or dissolvable. The bead may be a solid particle (e.g., a metal-based particle including but not limited to iron oxide, gold or silver) covered with a coating comprising one or more polymers. Such coating may be disruptable or dissolvable.

[00124] The term “sample,” as used herein, generally refers to a biological sample of a subject. The biological sample may comprise any number of macromolecules, for example, cellular macromolecules. The sample may be a cell sample. The sample may be a cell line or cell culture sample. The sample can include one or more cells. The sample can include one or more microbes. The biological sample may be a nucleic acid sample or protein sample. The biological sample may also be a carbohydrate sample or a lipid sample. The biological sample may be derived from another sample. The sample may be a tissue sample, such as a biopsy, core biopsy, needle aspirate, or fine needle aspirate. The sample may be a fluid sample, such as a blood sample, urine sample, or saliva sample. The sample may be a skin sample. The sample may be a cheek swab. The sample may be a plasma or serum sample. The sample may be a cell-free or cell free sample. A cell-free sample may include extracellular polynucleotides. Extracellular polynucleotides may be isolated from a bodily sample that may be selected from the group consisting of blood, plasma, serum, urine, saliva, mucosal excretions, sputum, stool and tears.

[00125] The term “biological particle,” as used herein, generally refers to a discrete biological system derived from a biological sample. The biological particle may be a macromolecule. The biological particle may be a small molecule. The biological particle may be a virus. The biological particle may be a cell or derivative of a cell. The biological particle may be an organelle. The biological particle may be a rare cell from a population of cells. The biological particle may be any type of cell, including without limitation prokaryotic cells, eukaryotic cells, bacterial, fungal, plant, mammalian, or other animal cell type, mycoplasmas, normal tissue cells, tumor cells, or any other cell type, whether derived from single cell or multicellular organisms. The biological particle may be a constituent of a cell. The biological particle may be or may include DNA, RNA, organelles, proteins, or any combination thereof. The biological particle may be or may include a matrix (e.g., a gel or polymer matrix) comprising a cell or one or more constituents from a cell (e.g., cell bead), such as DNA, RNA, organelles, proteins, or any

combination thereof, from the cell. The biological particle may be obtained from a tissue of a subject. The biological particle may be a hardened cell. Such hardened cell may or may not include a cell wall or cell membrane. The biological particle may include one or more constituents of a cell, but may not include other constituents of the cell. An example of such constituents is a nucleus or an organelle. A cell may be a live cell. The live cell may be capable of being cultured, for example, being cultured when enclosed in a gel or polymer matrix, or cultured when comprising a gel or polymer matrix.

[00126] The term “macromolecular constituent,” as used herein, generally refers to a macromolecule contained within or from a biological particle. The macromolecular constituent may comprise a nucleic acid. In some cases, the biological particle may be a macromolecule. The macromolecular constituent may comprise DNA. The macromolecular constituent may comprise RNA. The RNA may be coding or non-coding. The RNA may be messenger RNA (mRNA), ribosomal RNA (rRNA) or transfer RNA (tRNA), for example. The RNA may be a transcript. The RNA may be small RNA that are less than 200 nucleic acid bases in length, or large RNA that are greater than 200 nucleic acid bases in length. Small RNAs may include 5.8S ribosomal RNA (rRNA), 5S rRNA, transfer RNA (tRNA), microRNA (miRNA), small interfering RNA (siRNA), small nucleolar RNA (snoRNAs), Piwi-interacting RNA (piRNA), tRNA-derived small RNA (tsRNA) and small rDNA-derived RNA (srRNA). The RNA may be double-stranded RNA or single-stranded RNA. The RNA may be circular RNA. The macromolecular constituent may comprise a protein. The macromolecular constituent may comprise a peptide. The macromolecular constituent may comprise a polypeptide.

[00127] The term “molecular tag,” as used herein, generally refers to a molecule capable of binding to a macromolecular constituent. The molecular tag may bind to the macromolecular constituent with high affinity. The molecular tag may bind to the macromolecular constituent with high specificity. The molecular tag may comprise a nucleotide sequence. The molecular tag may comprise a nucleic acid sequence. The nucleic acid sequence may be at least a portion or an entirety of the molecular tag. The molecular tag may be a nucleic acid molecule or may be part of a nucleic acid molecule. The molecular tag may be an oligonucleotide or a polypeptide. The molecular tag may comprise a DNA aptamer. The molecular tag may be or comprise a primer. The molecular tag may be, or comprise, a protein. The molecular tag may comprise a polypeptide. The molecular tag may be a barcode.

[00128] The term “partition,” as used herein, generally, refers to a space or volume that may be suitable to contain one or more species or conduct one or more reactions. A partition may be a physical compartment, such as a droplet or well. The partition may isolate space or volume from another space or volume. The droplet may be a first phase (e.g., aqueous phase) in a

second phase (e.g., oil) immiscible with the first phase. The droplet may be a first phase in a second phase that does not phase separate from the first phase, such as, for example, a capsule or liposome in an aqueous phase. A partition may comprise one or more other (inner) partitions. In some cases, a partition may be a virtual compartment that can be defined and identified by an index (e.g., indexed libraries) across multiple and/or remote physical compartments. For example, a physical compartment may comprise a plurality of virtual compartments.

[00129] Provided herein are methods that may be used for various sample processing and/or analysis applications. A method of the present disclosure may provide a support (e.g., a bead, such as a gel bead) attached thereto two or more different barcode molecules (e.g., nucleic acid barcode molecules) each comprising one or more sequences (e.g., priming sequences, barcode sequences, random N-mer sequences, capture sequences, poly(T) sequences, and/or other sequences). Two or more different barcode molecules attached to a support may comprise the same or different features. For example, a first barcode molecule attached to a support may comprise a first feature (e.g., a first sequence, such as a first nucleic acid sequence) and a second barcode molecule attached to the same support may comprise a second feature (e.g., a second sequence, such as a second nucleic acid sequence) that is the same as or different from said first feature. In some cases, a first barcode molecule attached to a support may comprise a first feature and a second feature and a second barcode molecule attached to the same support may comprise a third feature and a fourth feature, where the first and third features may be different from the second and fourth features. For example, different nucleic acid barcode molecules attached to the same bead may comprise the same barcode sequence and one or more different other sequences (e.g., functional or starter sequences). A barcode molecule may be releasably attached to a support (e.g., a bead). Methods of generating a plurality of barcode molecules coupled to a plurality of supports may comprise combinatorial (e.g., split pool) assembly and a series of hybridization and ligation processes. The methods may facilitate combinatorial construction of a bead library comprising beads comprising a plurality of different barcode molecules.

Methods of Generating Barcode Molecules

[00130] In an aspect, the present disclosure provides a method for generating a plurality of barcode molecules (e.g., nucleic acid barcode molecules) from a plurality of molecules (e.g., nucleic acid molecules, such as a nucleic acid molecule comprising a starter or functional flow cell sequence) coupled to a plurality of supports (e.g., beads). The method may comprise combinatorially assembling (e.g., using a split pool method) the plurality of barcode molecules by coupling one or more molecules to each of the plurality of molecules. Each of the molecules coupled to a molecule coupled to a support may comprise one or more features, such as one or

more nucleic acid sequences (e.g., barcode sequences, functional sequences, starter sequences, and/or overhang sequences, or complements thereof). A feature may comprise a barcode sequence or a portion of a barcode sequence. The plurality of molecules generated through this method may comprise (i) a first set of barcode molecules coupled to a support of a plurality of supports and (ii) a second set of barcode molecules coupled to the same support, where barcode molecules of the first set of barcode molecules are different than barcode molecules of the second set of barcode molecules. Barcode molecules of the first set of barcode molecules and barcode molecules of the second set of barcode molecules may comprise the same barcode sequence. The barcode molecules of the first set of barcode molecules and the barcode molecules of the second set of barcode molecules may comprise a barcode sequence that is different from other barcode sequences of other barcode molecules coupled to other supports of the plurality of supports. Accordingly, the method may provide a combinatorially constructed library of beads where each bead comprises two or more different barcode molecules, where each barcode molecule attached to a given bead comprises the same barcode sequence.

[00131] In some cases, methods of the present disclosure may be used to generate barcode molecules comprising one or more amino acids, peptides, proteins, PEG moieties, hydrocarbon chains, and/or other moieties. In some cases, methods of the present disclosure may be used to generate nucleic acid barcode molecules. In some cases, methods of the present disclosure may be used to generate barcode molecules that may be useful in a plurality of assays. A support may have multiple barcode molecule populations coupled thereto, each population of which may be configured for use in a different assay. For example, a first population of barcode molecules coupled to the support may comprise a first sequence (e.g., a poly(T) sequence) for use with a first assay (e.g., analysis of a messenger ribonucleic acid molecule) and a second population of barcode molecules coupled to the same support may comprise a second sequence (e.g., a capture sequence) for use with a second assay (e.g., analysis of a deoxyribonucleic acid molecule) that is different from the first assay.

[00132] The methods described herein may comprise multiple ligation and/or hybridization processes. For example, a method may comprise providing a bead (e.g., a gel bead) that may have a starter sequence coupled thereto; providing a first molecule comprising a first sequence; attaching the first sequence of the first molecule to the starter sequence, thereby generating a first product; providing a second molecule comprising a second sequence; and attaching the second sequence of the second molecule to the first product, thereby generating a second product. The second product may comprise a barcode molecule, and may comprise one or more barcode sequences, functional sequences, and/or other sequences. In some cases, the method may further comprise providing a third molecule comprising a third sequence and attaching the third

sequence of the third molecule to the second product, thereby generating a third product. In such a case, the third product may comprise a barcode molecule, and may comprise one or more barcode sequences, functional sequences, and/or other sequences. The starter sequence may comprise a functional sequence such as a flow cell functional sequence and/or a partial read sequence. The first, second, and third molecules may comprise first, second, and third barcode sequences, respectively. One or more of the first, second, and third molecules may comprise a functional sequence such as a poly(T) sequence or a DNA capture sequence. One or more of the first, second, and third molecules may also comprise an overhang sequence to facilitate attachment of a particular sequence at a specific location. The barcode molecules generated using the methods described herein may be releasably attached to a support. In some cases, a combinatorial (e.g., split pool) approach may be employed to provide a plurality of supports each comprising attached thereto two or more different barcode molecules.

[00133] A support used in a method of the present disclosure may be, for example, a well, matrix, rod, container, or bead(s). A support may have any useful features and characteristics, such as any useful size, fluidity, solidity, density, porosity, and composition. In some cases, a support may be a bead such as a gel bead. A bead may be solid or semi-solid. Additional details of beads are provided elsewhere herein.

[00134] A support (e.g., a bead) may comprise a starter sequence functionalized thereto (e.g., as described herein). A starter sequence may be attached to the support via, for example, a disulfide linkage. A starter sequence may comprise a partial read sequence and/or flow cell functional sequence. Such a sequence may permit sequencing of nucleic acid molecules attached to the sequence by a sequencer (e.g., an Illumina sequencer). Different starter sequences may be useful for different sequencing applications. A starter sequence may comprise, for example, a TruSeq or Nextera sequence. A starter sequence may have any useful characteristics such as any useful length and nucleotide composition. For example, a starter sequence may comprise 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or more nucleotides. In some cases, a starter sequence may comprise 15 nucleotides. Nucleotides of a starter sequence may be naturally occurring or non-naturally occurring (e.g., as described herein). A bead may comprise a plurality of starter sequences attached thereto. For example, a bead may comprise a plurality of first starter sequences attached thereto. In some cases, a bead may comprise two or more different starter sequences attached thereto. For example, a bead may comprise both a plurality of first starter sequences (e.g., Nextera sequences) and a plurality of second starter sequences (e.g., TruSeq sequences) attached thereto. For a bead comprising two or more different starter sequences attached thereto, the sequence of each different starter sequence may be distinguishable from the sequence of each other starter sequence at an end distal to the bead. For

example, the different starter sequences may comprise one or more nucleotide differences in the 2, 3, 4, 5, 6, 7, 8, 9, 10, or more nucleotides furthest from the bead.

[00135] A first molecule (e.g., a first nucleic acid molecule) may be provided that is capable of attaching to a starter sequence attached to a bead. A first molecule may be, for example, a nucleic acid molecule and/or may comprise an amino acid, peptide, polyethylene glycol (PEG) moiety, hydrocarbon chain, or another moiety. A first nucleic acid molecule may comprise a double-stranded region and one or more single-stranded regions. For example, a first nucleic acid molecule may comprise a central portion that is a double-stranded region and adjacent regions on either side that are single-stranded regions. Such a nucleic acid molecule may be referred to as a “splint oligonucleotide.” Both single-stranded regions of the first nucleic acid molecule may be on the same strand. The single-stranded region on a first end of a first nucleic acid molecule may comprise 1, 2, 3, 4, 5, 6, or more nucleotides. This single-stranded region may be referred to as an “overhang” sequence. The single-stranded region on a second end of a first nucleic acid molecule may comprise a sequence that is complementary to a starter sequence attached to the bead. For example, the bead may comprise a starter sequence comprising a TruSeq sequence, and a first nucleic acid molecule may comprise a single-stranded region comprising a sequence that is complementary to a distal portion of the TruSeq sequence. The complementary sequence may comprise any useful length and base composition. Similarly, the double-stranded region may comprise any useful length and composition. A strand of the double-stranded region of the first nucleic acid molecule may comprise a barcode sequence, and the corresponding strand may comprise a complement of the barcode sequence. As an example, the first nucleic acid molecule may comprise a first strand including a first barcode sequence and a second strand including a sequence complementary to a starter sequence or a portion thereof that is attached to a bead, a sequence complementary to the first barcode sequence of the first strand, and an overhang sequence.

[00136] A first molecule (e.g., a first nucleic acid molecule or segment) may be attached to the starter sequence attached to the support (e.g., bead, such as a gel bead). Attachment of the first molecule may comprise hybridization and/or ligation (e.g., as described herein). **FIG. 13** shows an example of this process. Bead 1300 comprises starter sequence 1302 attached thereto. As described herein, starter sequence 1302 may comprise a partial read sequence such as a TruSeq or Nextera sequence. In the second panel, first molecule 1304 is ligated to starter sequence 1302. First molecule 1304 comprises a first strand 1306 that comprises a first barcode sequence and a second strand 1308 that comprises a sequence complementary to starter sequence 1302, a sequence complementary to the first barcode sequence of first strand 1306, and an overhang sequence. Ligation of first molecule 1304 to starter sequence 1302 may comprise

hybridization of the sequence of second strand 1308 that is complementary to starter sequence 1302 to starter sequence 1302.

[00137] A second molecule (e.g., a second nucleic acid molecule or segment) may be provided that is capable of attaching to the overhang sequence of a first molecule attached to a starter sequence attached to a support (e.g., bead). A second molecule may be, for example, a nucleic acid molecule and/or may comprise an amino acid, peptide, PEG moiety, hydrocarbon chain, or another moiety. A second nucleic acid molecule may comprise a double-stranded region and one or more single-stranded regions. For example, a second nucleic acid molecule may comprise a central portion that is a double-stranded region and adjacent regions on either side that are single-stranded regions. The single-stranded regions of the second nucleic acid molecule may be on the same or different strands. In some cases, a first strand of a second nucleic acid molecule comprises a single-stranded region comprising 1, 2, 3, 4, 5, 6, or more nucleotides. This single-stranded region may be referred to as a “complementary overhang” sequence. This sequence may be complementary to the overhang sequence of a first nucleic acid molecule. The first strand may comprise a second single-stranded region at a second end that comprises one or more additional sequences such as a unique molecular identifier or a functional sequence. The single-stranded region of a second end of the first strand of the second nucleic acid molecule may comprise 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 45, 50, or more nucleotides. The first strand may further comprise a second barcode sequence in the double stranded region of the second nucleic acid molecule. The second strand may comprise a sequence complementary to the second barcode sequence. Accordingly, in some cases, a second nucleic acid molecule comprises a first strand comprising a complementary overhang sequence, a second barcode sequence, and one or more additional sequences such as a unique molecular identifier or a functional sequence and a second strand comprising a sequence that is complementary to the second barcode sequence. In other cases, the first strand may include a single-stranded region and the second strand may include a single-stranded region. The single-stranded region of a first strand of a second nucleic acid molecule may comprise a “complementary overhang” sequence that is complementary to the overhang sequence of a first nucleic acid molecule. The single-stranded region of the second strand of the second nucleic acid molecule may comprise 1, 2, 3, 4, 5, 6, or more nucleotides. This single-stranded region may be referred to as a “second overhang” sequence. The double-stranded region may comprise any useful length and composition. A strand of the double-stranded region of the second nucleic acid molecule may comprise a second barcode sequence, and the corresponding strand may comprise a complement of the second barcode sequence. As an example, the second nucleic acid molecule may comprise a first strand including a second barcode sequence and a complementary

overhang sequence and a second strand including a sequence complementary to the second barcode sequence of the first strand, and a second overhang sequence. A second nucleic acid molecule may be ligated to the first nucleic acid molecule (e.g., as described herein). In **FIG. 13**, second molecule 1310 comprises first strand 1312 comprising a complementary overhang sequence and a second barcode sequence and second strand 1314 comprising a sequence complementary to the second barcode sequence and a second overhang sequence. Ligation of second molecule 1310 to first molecule 1304 may comprise hybridization of the sequence of first strand 1312 that is complementary to the overhang sequence of strand 1308 of first molecule 1304 to strand 1308.

[00138] In some cases, a third molecule (e.g., a third nucleic acid molecule or segment) may be provided that is capable of attaching to the overhang sequence of a second molecule attached to a first molecule attached to a starter sequence attached to a support (e.g., bead). A third molecule may be, for example, a nucleic acid molecule and/or may comprise an amino acid, peptide, PEG moiety, hydrocarbon chain, or another moiety. A third nucleic acid molecule may comprise a double-stranded region and one or more single-stranded regions. For example, a third nucleic acid molecule may comprise a central portion that is a double-stranded region and adjacent regions on either side that are single-stranded regions. The single-stranded regions of the second nucleic acid molecule may be the same strand. The single-stranded region of a first end of a first strand of a third nucleic acid molecule may comprise 1, 2, 3, 4, 5, 6, or more nucleotides. This single-stranded region may be referred to as a “second complementary overhang” sequence. This sequence may be complementary to the second overhang sequence of a second nucleic acid molecule. The single-stranded region of a second end of the first strand of the third nucleic acid molecule may comprise 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 45, 50, or more nucleotides. This single-stranded region may comprise one or more sequences. For example, this region may comprise a unique molecular identifier, a third barcode sequence, and/or a functional sequence. The double-stranded region may comprise any useful length and composition. A strand of the double-stranded region of the second nucleic acid molecule may comprise a third barcode sequence, a unique molecular identifier, or another sequence and the corresponding strand may comprise a complement of the third barcode sequence, unique molecular identifier, or other sequence. As an example, the third nucleic acid molecule may comprise a first strand including a third barcode sequence, a second complementary overhang sequence that is complementary to the second overhang sequence of a second nucleic acid molecule, and one or more additional sequences such as a unique molecular identifier and a functional sequence. The second strand of the third nucleic acid molecule may include a sequence complementary to the third barcode sequence of the first strand. A third

nucleic acid molecule may be ligated to the second nucleic acid molecule (e.g., as described herein). In **FIG. 13**, third molecule 1316 comprises first strand 1318 comprising a second complementary overhang sequence, a third barcode sequence, and one or more additional sequences and second strand 1320 comprising a sequence complementary to the third barcode sequence. Ligation of third molecule 1316 to second molecule 1310 may comprise hybridization of the sequence of first strand 1318 that is complementary to the overhang sequence of strand 1314 of second molecule 1310 to strand 1314.

[00139] **FIG. 14** shows an example of a double ligation method involving a bead comprising a starter sequence comprising a TruSeq sequence. In the second panel, a first nucleic acid molecule comprising a first strand (top) and a second strand (bottom) is ligated to the TruSeq sequence. The second strand of the first nucleic acid molecule comprises a sequence complementary to a portion of the TruSeq sequence at one end and a two-nucleotide overhang sequence (CA) at a second end. The first strand comprises a first barcode sequence (indicated by the sequence XXXXXX, where X represents a variable nucleotide), and the second strand comprises a sequence complementary to the first barcode sequence. In the third panel of **FIG. 14**, a second nucleic acid molecule comprising a first strand (top) and a second strand (bottom) is ligated to the first nucleic acid molecule. The first strand comprises a barcode sequence (indicated by the sequence XXXXXXXX, where X represents a variable nucleotide), and the second strand comprises a sequence complementary to the second barcode sequence. The first strand also comprises a two-nucleotide sequence (GT) that is complementary to the overhang sequence of the first nucleic acid molecule as well as additional sequences indicated by N_{10} and $T_{30}VN$. The sequence N_{10} represents a random N-mer that represents a unique molecular identifier while the sequence $T_{30}VN$ represents a functional sequence. The functional sequence included in **FIG. 14** is a poly(T) sequence. Following ligation of the second nucleic acid molecule to the first nucleic acid molecule, the bead comprises a double-stranded nucleic acid barcode molecule attached thereto. The double-stranded nucleic acid barcode molecule may be denatured to remove the bottom strand and retain a single-stranded nucleic acid barcode molecule attached to the bead. This single-stranded nucleic acid barcode molecule may also be referred to as a “nucleic acid barcode molecule”.

[00140] Successive attachment of a first molecule, second molecule, and, in some cases, a third molecule to a bead or a sequence attached thereto may result in the generation of a bead comprising a barcode molecule (e.g., a nucleic acid barcode molecule). The barcode molecule may be useful in various analysis and processing applications. Barcode molecules attached to beads may be useful for capturing and/or labeling, for example, nucleic acid sequences. A barcode molecule (e.g., a nucleic acid barcode molecule) may comprise a functional sequence

designed to interact with a particular nucleic acid sequence or type of sequence. For example, a nucleic acid barcode molecule may comprise a functional sequence selected from the group consisting of capture sequences (e.g., random N-mers), poly(T) sequences, poly(C) sequences, primer sequences, universal primer sequences, primer annealing sequences, or other sequences. The nucleic acid barcode molecule may comprise one or more barcode sequences and/or unique molecular identifiers and be useful for indexing nucleic acid sequences deriving from one or a variety of sources. If a plurality of beads each comprising one or more barcode molecules comprising a unique barcode and/or unique molecular identifier is partitioned amongst a plurality of partitions (e.g., droplets or wells, as described herein) each comprising an analyte, the barcode molecules may be useful for indexing the analytes corresponding to each partition.

[00141] Multiple barcode molecules (e.g., nucleic acid barcode molecules) may be generated on the same support (e.g., bead). For example, a bead may comprise a plurality of starter sequences attached thereto, and barcode molecules may be generated using all or a portion of the plurality of starter sequences. The barcode molecules attached to a given support may be the same (e.g., comprising the same barcode sequences, functional sequences, and/or other sequences). In some cases, barcode molecules attached to the same support may differ in unique molecular identifiers included in each barcode molecule. Such barcode molecules may otherwise include the same components.

[00142] In some cases, multiple different barcode molecules (e.g., nucleic acid barcode molecules) may be generated on the same support (e.g., bead). For example, two different barcode molecules may be generated on the same support. Alternatively, three or more different barcode molecules may be generated on the same support. Different barcode molecules attached to the same support may comprise one or more different sequences. For example, different barcode molecules may comprise one or more different barcode sequences, functional sequences, and/or other sequences (e.g., starter sequences). In some cases, different barcode molecules attached to the same support may comprise the same barcode sequences and different functional sequences. For example, a first barcode molecule attached to a support may comprise a first functional sequence (e.g., a random N-mer) that is a capture sequence (e.g., for capturing a deoxyribonucleic acid (DNA) molecule), while a second barcode molecule attached to the same support may comprise a second functional sequence that is a poly(T) sequence (e.g., for capturing a messenger ribonucleic acid (mRNA) molecule which may comprise a poly(A) sequence). Different barcode molecules attached to the same support may comprise barcode sequences that are the same or different. Similarly, different barcode molecules may comprise unique molecular identifiers (UMIs) that are the same or different. Different barcode molecules attached to the same support may have several different configurations. A first barcode molecule

attached to a support may comprise a first functional sequence, a first barcode sequence, and a first starter sequence, and a second barcode molecule attached to the same support may comprise a second functional sequence, a second barcode sequence, and a second starter sequence. In a first example, the first and second barcode sequences may be the same, the first and second functional sequences may be the same, and the first and second starter sequences may be different. Such a configuration may facilitate performing different assays (e.g., sequencing methods) on the same target molecules. In a second example, the first and second barcode sequences may be the same, the first and second functional sequences may be different, and the first and second starter sequences may be the same. Such a configuration may facilitate performing different assays on different target molecules (e.g., RNA and DNA molecules). In a third example, the first and second barcode sequences may be different, the first and second functional sequences may be the same, and the first and second starter sequences may be the same. Such a configuration may facilitate sample preparation for multiple different subsequent assays (e.g., a first population processed using the first barcode molecules can be separated from a second population processed using the second barcode molecules). In other examples, any first pair of first and second sequences (e.g., functional sequence, barcode sequence, or starter sequence) in first and second barcode molecules may be the same, and second and third pairs of first and second sequences (not the first pair) may be different. In other examples, any first and second pairs of first and second sequences (e.g., functional sequence, barcode sequence, or starter sequence) in first and second barcode molecules may be the same, and a third pair of first and second sequences (not the first or second pair) may be different.

[00143] FIG. 15 shows an example of a double ligation method involving a bead comprising a plurality of TruSeq sequences. The TruSeq sequences attached to the bead comprise the same sequence. Accordingly, the same first nucleic acid molecule can be ligated to each TruSeq sequence. In the third panel, a second nucleic acid molecule comprising a functional sequence comprising a poly(T) sequence attaches to a first nucleic acid molecule, and a second nucleic acid molecule comprising a functional sequence comprising a capture sequence attached to another first nucleic acid molecule. The second barcode sequences of each second nucleic acid molecule may be the same. Accordingly, FIG. 15 demonstrates double ligation to generate multiple different nucleic acid barcode molecules attached to the same bead.

[00144] Different overhang sequences may be used to direct the formation of different nucleic acid barcode molecules associated with the same bead. FIG. 16 shows an example of a double ligation method involving a bead comprising a first starter sequence that is a TruSeq sequence and a second starter sequence that is a Nextera sequence. The first and second starter sequences comprise different nucleic acid sequences. Accordingly, different first nucleic acid molecules

ligate to each different starter sequence. In the second panel of **FIG. 16**, different first nucleic acid molecules are shown ligated to the different starter sequences (e.g., the TruSeq and Nextera sequences). The first nucleic acid molecule ligated to the TruSeq sequence comprises a sequence that is complementary to a portion of the TruSeq sequence, a first barcode sequence, and a first overhang sequence. The first overhang sequence comprises two nucleotides, CA. The first nucleic acid molecule ligated to the Nextera sequence comprises a sequence that is complementary to a portion of the Nextera sequence, a first barcode sequence that is the same or different from the first barcode sequence of the other first nucleic acid molecule, and a first overhang sequence that is different from the first overhang sequence of the other first nucleic acid molecule. The first overhang sequence corresponding to the Nextera sequence is a two nucleotide sequence, GT. By using first nucleic acid molecules with different overhang sequences, the second nucleic acid molecules that ligate to first nucleic acid molecules attached to the different starter sequences can be maintained. Accordingly, the overhang sequences prove useful as indices for controlling the generation of barcode molecules attached to the bead. In the bottom panel of **FIG. 16**, different second nucleic acid molecules are shown ligated to the different first nucleic acid molecules ligated to TruSeq and Nextera sequences. The second nucleic acid molecule corresponding to the TruSeq sequence comprises a first complementary overhang sequence with a sequence complementary to the first overhang sequence of the first nucleic acid molecule corresponding to the TruSeq sequence. Accordingly, the complementary overhang sequence has a sequence GT. The second nucleic acid molecule corresponding to the Nextera sequence comprises a complementary overhang sequence with a sequence complementary to the first overhang sequence of the first nucleic acid molecule corresponding to the Nextera sequence. Accordingly, the complementary overhang sequence has a sequence CA. Each second nucleic acid molecule further comprises a second barcode sequence and a functional sequence. The second barcode sequences of the second nucleic acid molecules may be the same or different. The functional sequences of the second nucleic acid molecules may be different. For example, the functional sequence of the second nucleic acid molecule corresponding to the TruSeq sequence may comprise a poly(T) sequence, while the functional sequence of the second nucleic acid molecule corresponding to the Nextera sequence may comprise a capture sequence. Each second nucleic acid molecule may further comprise a unique molecular identifier. The unique molecular identifiers of each second nucleic acid molecules may be the same or different. **FIG. 12** displays another example of differential functionalization of a bead comprising different starter sequences. The first nucleic acid molecule associated with each starter sequence comprises Parts A and S, and the second nucleic acid molecule associated with each starter sequence comprises Parts B and D. During the first attachment step, a portion

of Part S hybridizes to its corresponding starter sequence and an end of Part A ligates to the starter sequence. During the second attachment step, a portion of Part B hybridizes to an overhang sequence of Part S and an end of Part B ligates to an end of Part A. The resultant product is a partial double-stranded nucleic acid barcode molecule. The nucleic acid barcode molecule may be denatured to provide a single-stranded nucleic acid barcode molecule (e.g., as described herein). The first and second nucleic acid molecules of **FIG. 12** share the same first and second barcode sequences (shown highlighted). As indicated in the key, the overhang sequence associated with the TruSeq starter sequences is referred to as an “alpha” overhang, while the overhang sequence associated with the Nextera starter sequence is referred to as a “beta” overhang.

[00145] Triple ligation may also be employed to generate barcode molecules attached to beads, as described above. **FIG. 17** shows an example of functionalization of a bead comprising a TruSeq starter sequence using triple ligation. The first nucleic acid molecule comprises Parts A and S, the second nucleic acid molecule comprises Parts B and D, and the third nucleic acid molecule comprises Parts C and T. Part S comprises a sequence that is complementary to a portion of the TruSeq starter sequence on one end and an overhang sequence CA on the second end. The overhang sequence CA may be referred to as an “alpha” overhang. Part S also comprises a sequence complementary to a first barcode sequence of Part A. Part B comprises a complementary overhang sequence GT that is complementary to the alpha overhang. Part B further comprises a second barcode sequence and Part D comprises a sequence that is complementary to the second barcode sequence. Part D further comprises a second overhang sequence comprising the sequence AC. The second overhang sequence AC may be referred to as a “theta” overhang. Part C comprises a second complementary overhang sequence that is complementary to the theta overhang. Part C further comprises a third barcode sequence and a functional sequence, while Part T comprises a sequence that is complementary to the third barcode sequence. Similarly, **FIG. 18** shows an example of functionalization of a bead comprising a Nextera starter sequence using triple ligation. The first nucleic acid molecule comprises Parts A and S, the second nucleic acid molecule comprises Parts B and D, and the third nucleic acid molecule comprises Parts C and T. Part S comprises a sequence that is complementary to a portion of the Nextera starter sequence on one end and an overhang sequence GT on the second end. The overhang sequence GT may be referred to as a “beta” overhang. Part S also comprises a sequence complementary to a first barcode sequence of Part A. Part B comprises a complementary overhang sequence CA that is complementary to the beta overhang. Part B further comprises a second barcode sequence and Part D comprises a sequence that is complementary to the second barcode sequence. Part D further comprises a second

overhang sequence comprising the sequence TG. The second overhang sequence TG may be referred to as an “eta” overhang. Part C comprises a second complementary overhang sequence that is complementary to the eta overhang. Part C further comprises a third barcode sequence and a functional sequence, while Part T comprises a sequence that is complementary to the third barcode sequence. **FIG. 19** compares triple ligation for TruSeq and Nextera sequences, highlighting the overhang and complementary overhang sequences therein.

[00146] **FIG. 20** shows an example of triple ligation method involving a bead comprising a starter sequence comprising a TruSeq sequence and a starter sequence comprising a Nextera sequence. In the second panel, a first nucleic acid molecule comprising a first strand (top strand, indicated as “Part A”) and a second strand (bottom strand) is ligated to the TruSeq sequence. The second strand of the first nucleic acid molecule comprises a sequence complementary to a portion of the TruSeq sequence at one end and a two-nucleotide overhang sequence (CA) at a second end. Similarly, a first nucleic acid molecule comprising a first strand (top strand, indicated as “Part A”) and a second strand (bottom strand) is ligated to each Nextera sequence. The second strand of the first nucleic acid molecule comprises a sequence complementary to a portion of the Nextera sequence at one end and a two-nucleotide overhang sequence (GT) at a second end. The first strand of each first nucleic acid molecule comprises a first barcode sequence (indicated by the sequence XXXXXX, where X represents a variable nucleotide), and the second strand comprises a sequence complementary to the first barcode sequence. In the third panel, a second nucleic acid molecule comprising a first strand (top strand, indicated as “Part B”) and a second strand (bottom strand) is ligated to each first nucleic acid molecule. The first strand of each second nucleic acid molecule comprises a complementary overhang sequence that is complementary to the corresponding overhang sequence of a first nucleic acid molecule, and the second strand of each second nucleic acid molecule comprises an overhang sequence that differs based on the associated starter sequence. Accordingly, the first strand of the second nucleic acid molecule associated with the TruSeq sequence comprises a complementary overhang sequence GT, while the first strand of the second nucleic acid molecule associated with the Nextera sequences comprises a complementary overhang sequence CA. Similarly, the second strand of the second nucleic acid molecule associated with the TruSeq sequence comprises an overhang sequence AC, while the second strand of the second nucleic acid molecule associated with the Nextera sequence comprises an overhang sequence TG. Each second nucleic acid molecule comprises a second barcode sequence (indicated by the sequence XXXXXXXX, where X represents a variable nucleotide) in its first strand, and the complement of the second barcode sequence in its second strand.

[00147] In the fourth panel, a third nucleic acid molecule comprising a first strand (top strand, indicated as “Part C”) and a second strand (bottom strand) is ligated to each second nucleic acid molecule. The first strand of each third nucleic acid molecule comprises a complementary overhang sequence that is complementary to the corresponding overhang sequence of a second nucleic acid molecule, and the second strand of each third nucleic acid molecule comprises an overhang sequence that differs based on the associated starter sequence. Accordingly, the first strand of the third nucleic acid molecule associated with the TruSeq sequence comprises a complementary overhang sequence TG, while the first strand of the third nucleic acid molecule associated with the Nextera sequences comprises a complementary overhang sequence AC. Each third nucleic acid molecule comprises a third barcode sequence (indicated by the sequence XXXXXXXX, where X represents a variable nucleotide) in its first strand, and the complement of the third barcode sequence, or a portion thereof, in its second strand. The first strand of each third nucleic acid molecule also includes a unique molecular identifier (N_{10}) and a functional sequence. The functional sequence associated with the TruSeq sequence comprises a poly(T) sequence while the functional sequences associated with the Nextera sequences comprise different capture sequences.

[00148] FIG. 11 shows examples of beads including two or more different nucleic acid barcode molecules attached thereto. Each nucleic acid barcode molecule comprises a starter sequence (e.g., a partial read sequence such as a TruSeq or Nextera sequence), a barcode sequence, a unique molecular identifier, and a functional sequence. The barcode sequence may comprise two or more barcode sequences from two or more nucleic acid molecules, as well as one or more overhangs. For example, a double ligation method may be used to generate the nucleic acid barcode molecules such that the barcode sequence may comprise a first barcode sequence from a first nucleic acid molecule, a first overhang sequence or complement thereof, and a second barcode sequence from a second nucleic acid molecule. Alternatively, a triple ligation method may be used to generate the nucleic acid barcode molecules such that the barcode sequence may comprise a first barcode sequence from a first nucleic acid molecule, a first overhang sequence or complement thereof, a second barcode sequence from a second nucleic acid molecule, a second overhang sequence or complement thereof, and a third barcode sequence from a third nucleic acid molecule. Nucleic acid barcode molecules may be differently functionalized (e.g., comprise different functional sequences suited for different applications, such as interaction with different analytes). The functional sequence of a nucleic acid barcode molecule may be included in the second (in the case of double ligation) or third (in the case of triple ligation) nucleic acid molecule used to construct the nucleic acid barcode molecule. The starter sequence to which other components of the nucleic acid barcode molecule attach may

dictate the functional sequence of the nucleic acid barcode molecule after double or triple ligation. For example, the middle panel of **FIG. 11** shows a bead comprising a TruSeq and a Nextera starter sequence. Because the different starter sequences have different nucleic acid sequences, a different first nucleic acid molecule will ligate to the first (e.g., TruSeq) sequence than will ligate to the second (e.g., Nextera) starter sequence. By designing the first nucleic acid molecules associated with each starter sequence to include different overhang sequences, the second nucleic acid molecule that ligates to each different first nucleic acid molecule may also be controlled. For example, a first nucleic acid molecule that ligates to a first starter sequence may include a first overhang sequence, while a first nucleic acid molecule that ligates to a second starter sequence may include a different first overhang sequence. The second nucleic acid molecule that ligates to the first nucleic acid molecule ligated to the first starter sequence will include a sequence complementary to its overhang sequence, while the second nucleic acid molecule that ligates to the first nucleic acid molecule ligated to the second starter sequence will include a sequence complementary to its overhang sequence which is different from the other overhang sequence. Accordingly, different second nucleic acid molecules may ligate to different first nucleic acid molecules ligated to different starter sequences attached to the same bead based on different overhang sequences. This so-called “overhang control” construction method may facilitate construction of different nucleic acid barcode molecules including different functional sequences on the same bead. Additional nucleic acid molecules (e.g., third nucleic acid molecules, fourth nucleic acid molecules, etc.) may be selectively ligated in a similar fashion. The barcode sequences of the different first, second, third, etc. nucleic acid molecules may be the same such that different nucleic acid barcode molecules comprising different starter sequences may comprise the same barcode sequences.

[00149] Alternatively, concentration and other reaction parameters may be used to control the nucleic acid barcode molecules formed in a so-called “concentration control” construction method. For example, the top panel of **FIG. 11** shows two different nucleic acid barcode molecules attached to a bead, where each nucleic acid barcode molecule comprises the same starter sequence (e.g., a TruSeq sequence). The barcode sequence associated with each nucleic acid barcode molecule may be the same. In order to achieve multiple different nucleic acid barcode molecules attached to a bead, different concentrations of a second or third nucleic acid molecule comprising a first functional sequence and a second or third nucleic acid molecule comprising a second functional sequence may be added to a ligation reaction mixture. For example, for a double ligation reaction, equal concentrations of a second nucleic acid molecule comprising a first functional sequence and a second nucleic acid molecule comprising a second functional sequence may be added to the ligation reaction mixture, which may result in equal

concentrations of the first and second functional sequences attached to a bead. Alternatively, a higher concentration of second nucleic acid molecules including a first functional sequence may be used to provide a bead comprising a higher concentration of the first functional sequence than a second functional sequence. A mixture comprising second nucleic acid molecules comprising first functional sequences and second nucleic acid molecules comprising second functional sequences may comprise greater than 50% second nucleic acid molecules comprising first functional sequences, such as 51%, 52%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or more second nucleic acid molecules comprising first functional sequences. A mixture comprising second nucleic acid molecules comprising first functional sequences and second nucleic acid molecules comprising second functional sequences may be designed to provide a bead comprising greater than 50% nucleic acid barcode molecules comprising first functional sequences, such as 51%, 52%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or more nucleic acid barcode molecules comprising first functional sequences. Accordingly, different second nucleic acid molecules may ligate to the same first nucleic acid molecules ligated to the same starter sequences attached to the same bead based on different concentrations. The same method may be applied for triple ligation using beads comprising the same starter sequences, first nucleic acid molecules, and second nucleic acid molecules and different third nucleic acid molecules comprising different functional sequences. The “concentration control” construction method may thus facilitate construction of different nucleic acid barcode molecules including different functional sequences on the same bead. The barcode sequences of the first, second, third, etc. nucleic acid molecules may be the same such that different nucleic acid barcode molecules comprising different starter sequences may comprise the same barcode sequences.

[00150] The bottom panel of **FIG. 11** shows a bead comprising a first nucleic acid barcode molecule comprising a first starter sequence (e.g., a TruSeq sequence) and second and third nucleic acid barcode molecules comprising second starter sequences (e.g., Nextera sequences). The barcode sequence associated with each different nucleic acid barcode molecule may be the same. As described with regard to the top and middle panels of **FIG. 11**, overhang and concentration control construction methods may be used to construct different nucleic acid molecules attached to the same bead. For example, overhang sequences may be used to direct the generation of a particular first nucleic acid barcode molecule comprising the first starter sequence, while concentration control may be used to direct the generation of the second and third nucleic acid barcode molecules comprising the second starter sequences in a given ratio.

[00151] A concentration control scheme may be used to block a given barcode molecule from being produced (e.g., in a larger quantity) on the support (e.g., bead) by limiting the amount of necessary molecular segments or sequences or other reagents available for the reaction.

Alternatively or in addition, one or more molecules or portions thereof coupled to a bead may be blocked to prevent or lessen generation of a given barcode molecule on the support (e.g., bead).

[00152] A nucleic acid molecule and/or a starter sequence associated with a support (e.g., bead) may comprise natural and/or non-naturally occurring (e.g., modified) nucleotides. For example, a nucleotide of a nucleic acid molecule and/or starter sequence associated with a bead may comprise any number or concentration of guanine, cytosine, thymine, uracil, and adenine bases, as well as non-naturally occurring (e.g., modified) nucleosides. A modified nucleoside may comprise one or more modifications (e.g., alkylation, hydroxylation, oxidation, or other modification) in its nucleobase and/or sugar moieties. A nucleic acid molecule and/or a starter sequence associated with a bead may comprise a nucleotide comprising a modified phosphate linker moiety. A nucleotide of a nucleic acid molecule and/or starter sequence may comprise one or more detectable moieties such as one or more fluorophores.

[00153] A barcode sequence of a barcode molecule (e.g., a nucleic acid barcode molecule) may comprise a first barcode sequence, a second barcode sequence, and, in some cases, a third barcode sequence. A first overhang sequence or a complement thereof and, in some cases, a second overhang sequence or a complement thereof, may be inserted between two or more sequences of a barcode sequence. For example, a barcode sequence may comprise a first barcode sequence and a second barcode sequence flanking an overhang sequence or complement thereof. In some cases, a barcode sequence may comprise, in order, a first barcode sequence, a first overhang sequence or complement thereof, a second barcode sequence, a second overhang sequence or complement thereof, and a third barcode sequence. A barcode sequence may have any useful length and composition. In some cases, a barcode sequence may comprise 4 or more nucleotides, such as 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, or more nucleotides. For example, a barcode sequence may comprise a first barcode sequence comprising 6 nucleotides and a second barcode sequence comprising 8 nucleotides for a total of 14 nucleotides. In another example, a barcode sequence may comprise a first barcode sequence comprising 6 nucleotides, a second barcode sequence comprising 6 nucleotides, and a third barcode sequence comprising 8 nucleotides for a total of 20 nucleotides. A barcode sequence may comprise naturally occurring and/or non-naturally occurring nucleotides. A barcode sequence may comprise any useful sequence and composition of nucleotides. In some cases, all nucleic acid barcode molecules attached to a given bead comprise the same barcode sequences. In other cases, nucleic acid barcode molecules attached to a given bead may comprise different barcode sequences. For example, all nucleic acid barcode molecules may comprise the same first and second barcode sequences and different third barcode sequences. For beads for which all nucleic acid barcode molecules comprise the same barcode

sequence, each nucleic acid barcode molecule may comprise a different unique molecular identifier that identifies the nucleic acid barcode molecules attached to the beads.

[00154] Overhang sequences may help control the generation of different nucleic acid barcode molecules on a given bead (e.g., as described herein). A given overhang sequence may comprise 2, 3, 4, 5, 6, or more nucleotides. In some cases, the overhang sequence comprises 2 nucleotides. Examples of overhang sequences comprising 2 nucleotides include GT (referred to herein as “alpha”), CA (“beta”), AG (“gamma”), TC (“delta”), GA (“epsilon”), CT (“zeta”), AC (“eta”), and TG (“theta”). **FIG. 21** provides a scheme for triple ligation premised on differentiation between overhang sequences. The middle section provides overhang pairs (e.g., a first overhang and a second overhang) that may be used in triple ligation. As indicated therein, alpha and theta overhangs may be used in tandem; beta and eta overhangs may be used in tandem, delta and zeta overhangs may be used in tandem, and gamma and epsilon overhangs may be used in tandem.

[00155] Functional sequences may be useful for different applications and assays. For example, a functional sequence capable of hybridizing with an mRNA molecule (e.g., a functional sequence comprising a poly(T) sequence) may facilitate analysis of an mRNA analyte, while a functional sequence capable of hybridizing with a DNA molecule (e.g., a capture sequence, such as a capture sequence comprising a random N-mer) may facilitate analysis of a DNA analyte. A bead comprising a nucleic acid barcode molecule comprising a poly(T) sequence and a nucleic acid barcode molecule comprising a DNA capture sequence may be useful for analyzing an mRNA analyte and a DNA analyte in tandem (e.g., in a multi-assay method). Other examples of functional sequences include functional sequences capable of interacting with other nucleic acid molecules (e.g., transfer RNA (tRNA) molecules, ribosomal RNA (rRNA) molecules, and mitochondrial RNA (mtRNA) molecules), functional sequences capable of interacting with an oligonucleotide coupled to a labeling agent (e.g., an oligonucleotide coupled to a protein or antibody) and capable of detecting a an analyte (e.g., an intracellular protein, extracellular protein, transmembrane protein, or surface protein), and functional sequences capable of interacting with a nucleic acid sequence of or encoding a perturbation agent such as a clustered regularly interspersed short palindromic repeat (CRISPR) agent (e.g., crRNA or sgRNA), transcription-activator like effector nucleases (TALENs, e.g., TALEN mRNA sequence)), zinc finger nucleases (ZFN, e.g., ZFN mRNA sequence), or other perturbation agent. A functional sequence may comprise a random N-mer. In some cases, a functional sequence may be referred to as a “target sequence.” Examples of capture and primer sequences useful in nucleic acid barcode molecules of the claimed methods are provided in Table 1 below:

Table 1: Sequences for inclusion in nucleic acid barcode molecules

Sequence Number	Sequence	Sequence Number	Sequence
1	TGCCTTGTAACGCGAA	78	GAGCAACTATTTCGCGT
2	TATGGCCGCGCAATTA	79	TACCTACGATTGCGAG
3	TTCGAGCGCGCAATTA	80	TAATCGACCGGTTACG
4	ATTGCGCCGAACGTAT	81	GTTCACAATACGCGGT
5	GTTGCACGCGCAATTA	82	GCTAGCGCGCATATTA
6	TGCCATTGCGCGATAA	83	ACTGACTTACGCGGAT
7	AAGGATCGCGCCTATT	84	GCACGTTGCGCGTAATA
8	GTTACGCGCGCAATTA	85	AAGGTCCGCGCTATTA
9	AGCATGTCGCGCATT	86	CTAGTGAACGCGCTAT
10	TTCGCAACGGTCGAAT	87	GGCCATCGATTTCGTAA
11	ATTGCGCGCGAATTAC	88	ACCTTGCGCGATAGTA
12	TCTTAGCGGACGCAAT	89	CTAGGTCCGCGAATTA
13	ATCCATGGCGCGATTA	90	ATATCGACCGGTTACG
14	GTTTCGACGCGAATTA	91	ATAGCTTATGCGCGAC
15	CGATTGCGCGACATTA	92	ACGTTAATCGGTACGC
16	TGATCGCGCTACGAAT	93	CATCATCGATTGCGAG
17	CGCATTCAATTGGCGA	94	GTAACTCGACCGGATT
18	AACGTTGCGGATTGAC	95	TCGAACGCGTATTAGC
19	GCTTGACCGCGAATTA	96	TAGTCGACCGATTACG
20	ACTGCGCGATTTCGTAA	97	GATCACGCGATTTCGTA
21	TCCAATAATGCGCGGT	98	AGTTAGCGTTACGACC
22	AGTCATCGACCGGATT	99	GCTCAATAAGTCGCGT
23	AGACTTCGCGCGATTA	100	CTTAGGCGCGAATTAC
24	CTGAGTCGCGCAATTA	101	TAGGTCACGTTACGAC
25	TCGCTAACGGTCGAAT	102	ATACCTTATGCGCGAG
26	TATGCGCGCGAATTAC	103	GCACGATAGTTTCGCTA
27	TATGCGCGCTACGAAT	104	GTCATACAATTTCGCGG
28	AACTGCGCGATTTCGTA	105	CGACTATTATGCGCGA
29	TGGACCGCGCATATTA	106	TAAGCCGCGTATTAGC
30	TATCACAATGCGCGGT	107	CGAATTTCGACCGGTAT
31	GTCACGCGCGAATTAT	108	CGTCAATAATCGCGTG
32	GTCTAGCGCGCAATTA	109	GGAACCTTAATCGCGT
33	TCTGCAACGGTCGAAT	110	GCTCAATAATCGCGTG
34	CCAGTGCGCGAATTAT	111	CCAATGCGCGTTAGTA
35	ATCGTCACGCGATTAG	112	GAACCTTCGACCGGTAT
36	GCCAATCGACGTTAGT	113	CGCTTAATCGAACGGT
37	TTAGCGCGCGAATTAC	114	CCTTGAGTCGAACGAT
38	TCGATCAGTTACGCGA	115	ATCGAGTAACCGTTTCG
39	ACCTGAATACGCGGTT	116	TACCGTAACGTAGTCG
40	TGCGGTTCGAACCTAAT	117	GTTCTCATCGAACGGA
41	TGAACGCGCTACTATG	118	ATCTTGAATCGCGACG
42	TGGCTTAATCGCGACA	119	AGACCTTAATCGCGTG
43	TAGGTCCGCGACATTA	120	GACCAATATGTCGCGT
44	CCTTGCGCAACGATTA	121	CGTAATCGACCGGTAT
45	AAGTCCGCGCGATTAT	122	GTACACTAAGTCGCGT
46	GACTGTGCGCGCAATTA	123	TAGTACCGATTGACCG
47	TTAGGTCCGCTACGAA	124	AGTCTAATCGGTACGC

48	TCTGTGAACCGTCGAA	125	GTAAGTACGCGCTATC
49	GTCACAATACGCGGTT	126	CCTTGAATCGAACGGT
50	GAGACTTCGCGCATT	127	TCGACTAATCGGTACG
51	GAGCAATTCGCGCTAT	128	GCGGATTACGCTACTA
52	CGTTACGATTACGCGA	129	CCTTAGTAGTACGCGA
53	AGCACGTAATCGTTTCG	130	CCTTAAGTTACGCGAG
54	GCGTTACCGAACGTAT	131	GTAAGTACGCGCTATC
55	TTAACGACCGGTTACG	132	CTGTCGCGATCGATAA
56	ATTGCGCGCGATACTA	133	GGACAATCGCTCGTTA
57	CTGTTGACCGCGAATA	134	GTCCGTCGATCGATAA
58	GCAAGATTTCGCGCTAT	135	GACTTACGACCGGTAT
59	GGCCATCGCGAATTAT	136	CTGTTAATCGACCGGA
60	ATGGACCGCGCTATTA	137	ATGCGGTAACCTATCG
61	ACGCGATAATCGTTTCG	138	AGATAGTTACGCGTCC
62	TAGGCATTATCCGCGA	139	GAGTCCAACCTATCGGT
63	TCAGCTCGAACGGTTA	140	TTACAGTACTACGCGC
64	CATTCAATTGCGCGAG	141	GACTAATACGCGTTCG
65	GCTCAATTACGCGGAT	142	TGGTAACTATACCGGC
66	ACGGATCGCATCGTTA	143	CGTACGTAACCTATCGG
67	CTGCAATTACGCGGAT	144	GACCTTAATCGGTACG
68	TGTCGATTACGCGAAC	145	CATTACCGGATAGTCG
69	ACCTAGGCGCGATTAT	146	GATAGTTATCGCACCG
70	CGCGTAGCGCATATTA	147	ACTAGTCGTACGATGC
71	CAATCGTTACGCGGAT	148	GATCACTAATCGCGTG
72	TAACGCTTACGCGGAT	149	GCGTTACGCTAATACG
73	GCTAACGCGATTTCGTA	150	CACGATCGTACGGTAT
74	ATTCTTAATGCGCGAG	151	TGTACGTACGATCCGA
75	TTCCTTAATGCGCGAG	152	CTAGACTAATCGCGTG
76	CAAGGATTTCGCGCTAT	153	CATAGTCGTACGATGC
77	ACGATTTCGACCGGTAT	154	GTCGACTAATCGGTAT

[00156] Additional examples of capture and primer sequences that may be included in nucleic acid barcode molecules are included in Table 2 below:

Table 2: Sequences for inclusion in nucleic acid barcode molecules

Sequence Number	Sequence	Sequence Number	Sequence
155	/CCTTAGCCGCTAATAGGTGAGC	160	/ACGCTAGTTTCGCGTACGAAGC
156	/TTGCTAGGACCGGCCTTAAAGC	161	/GAGGATTGCGCACCTTACTAGC
157	/GAGGATTGCGCACCTTACTAGC	162	/TTGCTAGGACCGGCCTTAAAGC
158	/CAACTTTAGCGGTCCAAGGTGC	163	/GACAATTGTCGGCTCGACTAGC
159	/ACGCTAGTTTCGCGTACGAAGC		

[00157] In some cases, a barcode molecule (e.g., a nucleic acid barcode molecule) may comprise a sequence selected from the sequences included in Tables 1 and 2. In some cases, a barcode molecule may comprise a complement or reverse complement of a sequence selected from the sequences included in Tables 1 and 2. In some embodiments, a particular analyte of

interest (e.g., nucleic acid, cell surface protein, CRISPR RNA or other perturbation agent) may comprise, may be coupled to, or otherwise associated with an oligonucleotide comprising a sequence that is at least partially complementary to a sequence on a barcode molecule (e.g., a sequence in Table 1 or 2).

[00158] Multiple ligation reactions used in the preparation of barcode molecules (e.g., nucleic acid barcode molecules) according to the methods provided herein may be performed in a combinatorial fashion. Combinatorial (e.g., split pool) approaches permit the generation of a high diversity of barcode molecules (e.g., barcoded beads) using a reduced number of molecules (e.g., nucleic acid molecules). A combinatorial scheme involves assembling multiple molecular segments or sequences (e.g., a nucleic acid molecule or a portion thereof) to provide a larger molecule (e.g., a molecule comprising two or more molecular segments or sequences). **FIG. 9A** provides an example of a simplified combinatorial double ligation scheme (e.g., a split pool scheme). Partition 901 includes a plurality of beads comprising a starter sequence and a first molecule A1. Partition 902 includes a plurality of beads comprising a starter sequence and a first molecule A2. First molecules A1 are ligated to the beads of partition 901, and first molecules A2 are ligated to the beads of partition 902. Subsequent to this first ligation step, in process 903 beads of partition 901 and beads of partition 902 are pooled in container 904. Container 904 includes a pooled mixture comprising beads functionalized with first molecule A1 and beads functionalized with first n molecule A2. The pooled mixture is then partitioned between partitions 906 and 907. Partitions 906 and 907 therefore each include beads functionalized with first molecule A1 and beads functionalized with first molecule A2. Partition 906 includes second molecules B1 and partition 907 includes second molecules B2. Second molecules B1 are ligated to the beads of partition 906 to generate beads functionalized with A1 and B1 and beads functionalized with A2 and B1, and second molecules B2 are ligated to the beads of partition 907 to generate beads functionalized with A1 and B2 and beads functionalized with A2 and B2. Subsequent to this second ligation step, in process 908 beads of partition 906 and beads of partition 907 are pooled in container 909. Container 909 includes a pooled mixture comprising beads functionalized with A1 and B1, A1 and B2, A2 and B1, and A2 and B2. The differentially functionalized beads 910, 911, 912, and 913 are shown in **FIG. 9B**. These beads may be divided amongst different partitions and used for analysis of different analytes. For example, a single bead of the collection of beads may be co-partitioned with a single biological particle (e.g., a cell) comprising an analyte and one or more reagents. A plurality of partitions may be similarly generated (e.g., as described herein) such that each partition includes a bead having attached thereto a differently functionalized barcode molecule.

[00159] In some cases, a combinatorial ligation method such as that described with reference to **FIG. 9A** may comprise immobilizing a plurality of beads to a plurality of supports (e.g., wells of a well plate). The first ligation step may involve providing a plurality of first molecules (e.g., first nucleic acid molecules) to each partition and promoting a ligation reaction. Rather than pooling the plurality of beads from each partition, excess material may be washed out of each partition and a plurality of second molecules (e.g., second nucleic acid molecules) washed into each partition for a second ligation reaction. Accordingly, the methods of the present disclosure may comprise moving beads between partitions or moving fluids between partitions.

[00160] The scheme presented in **FIG. 9A** represents a simplified version of a combinatorial double ligation method (e.g., double ligation split pool method). The complexity of the scheme may be enhanced by providing different versions of the second molecule (e.g., a first version comprising a first functional sequence and a second version comprising a second functional sequence) in the same or different concentration for ligation to the same first molecule of a given bead (e.g., as described herein). This method may involve concentration control construction. Complexity may also be enhanced by providing a bead comprising two or more different starter sequences and employing molecules comprising overhang sequences to direct the generation of particular barcode molecules and provide different barcode molecules attached to the same bead.

[00161] **FIG. 10** provides an example of a simplified combinatorial triple ligation scheme (e.g., triple ligation split pool scheme). Partition 1001 includes a plurality of beads comprising a starter sequence and a first molecule A1. Partition 1002 includes a plurality of beads comprising a starter sequence and a first molecule A2. Partition 1003 includes a plurality of beads comprising a starter sequence and a first molecule A3. First molecules A1 are ligated to the beads of partition 1001, first molecules A2 are ligated to the beads of partition 1002, and first molecules A3 are ligated to the beads of partition 1003. Subsequent to this first ligation step, in process 1004 beads of partitions 1001, 1002, and 1003 are pooled in container 1005. Container 1005 includes a pooled mixture comprising beads functionalized with first molecule A1, beads functionalized with first molecule A2, and beads functionalized with first molecule A3. The pooled mixture is then partitioned in process 1006 between partitions 1007, 1008, and 1009. Partitions 1007, 1008, and 1009 therefore each include beads functionalized with first molecule A1, beads functionalized with first molecule A2, and beads functionalized with first molecule A3. Partition 1007 includes second molecules B1, partition 1008 includes second molecules B2, and partition 1009 includes second molecules B3. Second molecules B1 are ligated to the beads of partition 1007 to generate beads functionalized with A1 and B1, beads functionalized with A2 and B1, and beads functionalized with A3 and B1; second molecules B2 are ligated to the beads of partition 1008 to generate beads functionalized with A1 and B2, beads functionalized with A2

and B2, and beads functionalized with A3 and B2; and second molecules B3 are ligated to the beads of partition 1009 to generate beads functionalized with A1 and B3, beads functionalized with A2 and B3, and beads functionalized with A3 and B3. Subsequent to this second ligation step, in process 1010 beads of partitions 1007, 1008, and 1009 are pooled in container 1011. Container 1011 includes a pooled mixture comprising beads functionalized with A1 and B1, A1 and B2, A1 and B3, A2 and B1, A2 and B2, A2 and B3, A3 and B1, A3 and B2, and A3 and B3.

[00162] The pooled mixture is then partitioned in process 1012 between partitions 1013, 1014, and 1015. Partitions 1013, 1014, and 1015 therefore each include beads functionalized with A1 and B1, A1 and B2, A1 and B3, A2 and B1, A2 and B2, A2 and B3, A3 and B1, A3 and B2, and A3 and B3. Partition 1013 includes third molecules C1, partition 1014 includes third molecules C2, and partition 1015 includes third molecules C3. Third molecules C1 are ligated to the beads of partition 1013, third molecules C2 are ligated to the beads of partition 1014, and third molecules C3 are ligated to the beads of partition 1015. Subsequent to this third ligation step, beads of partitions 1013, 1014, and 1015 may be pooled and repartitioned for use in various processing and analysis applications (e.g., as described herein). The resultant pooled mixture comprises beads functionalized as follows: A1B1C1, A2B1C1, A3B1C1, A2B1C1, A2B2C1, A2B3C1, A3B1C1, A3B2C1, A3B3C1, A1B1C2, A2B1C2, A3B1C2, A2B1C2, A2B2C2, A2B3C2, A3B1C2, A3B2C2, A3B3C2, A1B1C3, A2B1C3, A3B1C3, A2B1C3, A2B2C3, A2B3C3, A3B1C3, A3B2C3, and A3B3C3. Accordingly, 9 different molecules (A1, A2, A3, B1, B2, B3, C1, C2, and C3) may be used to generate 27 different barcode molecules. As described elsewhere herein, the use of different starter sequences, overhangs, and controlled concentrations may further increase the number of different barcode molecules generated using the combinatorial (e.g., split pool) method.

[00163] Double and triple ligation reactions may be carried out in a series of well plates. For example, ligation of first molecules (e.g., first nucleic acid molecules) may be performed within wells of a 96 well plates using, e.g., 96 different first molecules (e.g., as described herein). Each well may also include a plurality of beads and one or more reagents. The plurality of beads may comprise one or more different starter sequences attached thereto. The plate may then be subjected to appropriate reaction conditions (e.g., appropriate pressure, concentration, and temperature conditions with appropriate reagents) to promote ligation between first molecules and starter sequences of beads within each well. Subsequent to this first ligation reaction, the contents of each well may be pooled and mixed. The contents of the resultant pooled mixture may then be redistributed between a plurality of wells in a separate well plate. For example, beads in the pooled mixture may be partitioned among 96 wells of a second 96 well plate. Each well may include a different second molecule (e.g., as described herein) and one or more

reagents. The plurality of beads in each well may include beads functionalized with each different first molecule. A second ligation reaction may be performed within each well to generate a plurality of beads comprising the same second molecule and a plurality of different first molecules within each well. The plurality of beads within each well may then be pooled to generate a second pooled mixture. If a third ligation reaction will be performed, the contents of the second pooled mixture may then be partitioned among 96 wells of a third 96 well plate. Each well may include a different third molecule (e.g., as described herein) and one or more reagents. The plurality of beads in each well may include beads functionalized with each different first molecule and each different second molecule. A third ligation reaction may be performed within each well to generate a plurality of beads comprising the same third molecule, a plurality of different first molecules, and a plurality of different second molecules within each well. The plurality of beads within each well may then be pooled to generate a third pooled mixture. Functionalized beads of the third pooled mixture may then be used in further processing or analysis applications (e.g., as described herein). **FIG. 21** provides additional details of a triple ligation scheme employing four different plates at each stage of the ligation reaction. **FIG. 22** demonstrates the capability of the scheme of **FIG. 21** to produce a large number of differently functionalized beads. As shown in **FIG. 22**, each triple ligation scheme performed using three 96 well plates provides nearly one million beads comprising different barcode molecules. By exploiting different overhang combinations (e.g., alpha/theta, beta/eta, delta/zeta, and gamma/epsilon), the number of different beads comprising different barcode molecules may be enhanced fourfold.

[00164] In any of the methods described herein, a double-stranded nucleic acid barcode molecule attached to a bead may be denatured to provide a single-stranded nucleic acid barcode molecule attached to the bead. Denaturation may be achieved controlling, for example, temperature, pressure, and/or pH conditions and/or by employing a chemical or biological agent such as a detergent.

[00165] In some cases, the methods described herein may be used to screen cells carrying mutations, e.g., mutations generated by gene editing such as CRISPR technology. For example, a bead comprising a first anchor oligonucleotide or nucleic acid molecule with a primer for CRISPR RNA (e.g., crRNA or guide RNA) or its complementary DNA and a second anchor oligonucleotide or nucleic acid molecule with a primer endogenous nucleic acid in the cell, e.g., total mRNA or a specific mRNA. The bead may be made into a partition with a cell transfected with CRISPR RNA or a plasmid expressing CRISPR RNA. In some cases, the expressed CRISPR RNA or the plasmid may have a barcode (CRISPR barcode) or a capture sequence. The primers (e.g., nucleic acid barcode molecules) on the bead may be used to amplify and sequence

the CRISPR RNA (e.g., using a nucleic acid barcode molecule comprising a sequence complementary to the CRISPR capture sequence, see **FIG. 23**) and endogenous mRNA (e.g., using a nucleic acid barcode molecule comprising an oligo(dT) sequence), thus determining the mutations generated by in the cell. In some cases, the methods may be used to perform single cell RNA sequencing, e.g., as described in Dixit, *et al.*, Perturb-Seq: Dissecting Molecular Circuits with Scalable Single-Cell RNA Profiling of Pooled Genetic Screens. *Cell*; Dec. 15, 2016;167(7):1853-1866.e17, which is incorporated herein by reference in its entirety.

[00166] As shown in **FIG. 24** and described elsewhere herein, beads comprising different nucleic acid barcode molecules generated by the methods described herein may be used to analyze mRNA and DNA sequences simultaneously. For example, mRNA and DNA sequences from the same biological particle (e.g., cell) within a partition may be attached to nucleic acid barcode molecules with the same barcode sequence.

[00167] A support (e.g., a bead such as a gel bead) may comprise a plurality of oligonucleotides or barcode molecules (e.g., nucleic acid barcode molecules) attached thereto. The support may comprise at least 10,000 barcode molecules attached thereto. For example, the support may comprise at least 100,000, 1,000,000, or 10,000,000 barcode molecules attached thereto. In any of the methods described herein, one or more oligonucleotides or barcode molecules may be releasably attached to a support. A barcode molecule may be releasable from a support upon application of a stimulus. Such a stimulus may be selected from the group consisting of a thermal stimulus, a photo stimulus, and a chemical stimulus. For example, the stimulus may be a reducing agent such as dithiothreitol. Application of a stimulus may result in one or more of (i) cleavage of a linkage between barcode molecules of the plurality of barcode molecules and the support, and (ii) degradation or dissolution of the support (e.g., bead) to release barcode molecules of the plurality of barcode molecules from the support. For a support comprising multiple different barcode molecules, different barcode molecules may be released from the bead upon application of different supports. For example, a first stimulus may be applied to release a first barcode molecule and may not release a second barcode molecule, and a second stimulus may be applied to release the second barcode molecule and may not release the first barcode molecule.

[00168] Different analytes capable of being analyzed by a support (e.g., a bead) comprising multiple different barcode molecules or a collection of such supports (e.g., beads) may be contained within or associated with a cell. A cell may be, for example, a human cell, an animal cell, or a plant cell. In some cases, the cell may be derived from a tissue or fluid, as described herein. The cell may be a prokaryotic cell or a eukaryotic cell. The cell may be a lymphocyte such as a B cell or T cell. Access to a plurality of molecules included in a cell may be provided

by lysing or permeabilizing the cell. Lysing the cell may release an analyte contained therein from the cell. A cell may be lysed using a lysis agent such as a bioactive agent. A bioactive agent useful for lysing a cell may be, for example, an enzyme (e.g., as described herein). Alternatively, an ionic or non-ionic surfactant such as TritonX-100, Tween 20, sarcosyl, or sodium dodecyl sulfate may be used to lyse a cell. Cell lysis may also be achieved using a cellular disruption method such as an electroporation or a thermal, acoustic, or mechanical disruption method. Alternatively, a cell may be permeabilized to provide access to a plurality of nucleic acid molecules included therein. Permeabilization may involve partially or completely dissolving or disrupting a cell membrane or a portion thereof. Permeabilization may be achieved by, for example, contacting a cell membrane with an organic solvent or a detergent such as Triton X-100 or NP-40.

[00169] A biological particle (e.g., a cell) including an analyte may be partitioned with a bead comprising one or more nucleic acid barcode molecules within a partition such as a well or droplet, e.g., as described herein. One or more reagents may be co-partitioned with a cell and bead. For example, a cell may be co-partitioned with one or more reagents selected from the group consisting of lysis agents or buffers, permeabilizing agents, enzymes (e.g., enzymes capable of digesting one or more nucleic acid molecules, extending one or more nucleic acid molecules, reverse transcribing an RNA molecule, permeabilizing or lysing a cell, or carrying out other actions), fluorophores, oligonucleotides, primers, barcodes, nucleic acid barcode molecules (e.g., nucleic acid barcode molecules comprising one or more barcode sequences), buffers, deoxynucleotide triphosphates, detergents, reducing agents, chelating agents, oxidizing agents, nanoparticles, beads, and antibodies. In some cases, a cell and bead may be co-partitioned with one or more reagents selected from the group consisting of temperature-sensitive enzymes, pH-sensitive enzymes, light-sensitive enzymes, reverse transcriptases, proteases, ligase, polymerases, restriction enzymes, nucleases, protease inhibitors, exonucleases, and nuclease inhibitors. For example, a cell and a bead may be co-partitioned with a reverse transcriptase and nucleotide molecules. Partitioning a cell, a bead, and one or more reagents may comprise flowing a first phase comprising an aqueous fluid, the cell, the bead, and the one or more reagents and a second phase comprising a fluid that is immiscible with the aqueous fluid toward a junction. Upon interaction of the first and second phases, a discrete droplet of the first phase comprising the cell, bead, and the one or more reagents may be formed. In some cases, the partition may comprise a single cell and a single bead. The cell may be lysed or permeabilized within the partition (e.g., droplet) to provide access to the plurality of molecules of the cell. Accordingly, molecules originating from the same cell may be isolated and barcoded within the same partition with the same barcode.

[00170] Combinatorial (e.g., split pool) ligation approaches such as those provided herein may generate significant diversity of barcode molecules. However, in some cases, unwanted off-products may also be generated. This may be due to ligation steps having less than 100% efficiency, such as between 95-97% efficiency. Off-products may not include the full length barcode sequence as the desired products and/or may lack a functional sequence. The overall efficiency of a combinatorial process may decrease for processes involving more ligation steps (e.g., triple ligation may be less efficient and generate more off-products than double ligation). Because off-products may interfere with biochemical reactions of interest, it may be beneficial to remove off-products (e.g., partial ligation products) without having other adverse effects on desired products. Removing the off-products may comprise degrading the off-products. Degrading these components may be accomplished by, for example, the use of one or more exonucleases. Exonucleases (e.g., ExoI) are typically inhibited by phosphorothioate bonds in the phosphate backbone of a nucleic acid. By incorporating a phosphorothioate bond between the final 3' base of a barcode molecule and the base immediately preceding it, a molecule can be created that is resistant to 3' → 5' exonuclease activity. This bond can be incorporated onto the final oligo that is ligated during a split pool ligation process. In this case, only molecules that are full length (e.g., desired products) can be protected from the exonuclease and partial ligation products (e.g., off-target products) may be digested by the exonuclease. This process may provide a population of molecules that include nearly 100% of the desired molecules (e.g., full length). Such a process may be particularly useful in a triple ligation approach where the initial efficiency may otherwise be undesirably low. This process may be used on any pool of molecules constructed in a combinatorial process where the final component (e.g., molecular segment or sequence) added includes a 3' phosphorothioate bond. The process may be performed within partitions (e.g., droplets or wells) and/or with molecules (e.g., barcode molecules) coupled to beads (e.g., gel beads) or a solid surface. Alternatively, the process may be performed with molecules (e.g., barcode molecules) in solution. A variety of exonucleases may be used in the process, including exonucleases unable to degrade phosphorothioate bonds. Other 3' modifications can be used to achieve the same effect including, for example, spacer molecules, biotin, and fluorophores. Blockers at the 3' end of a complete barcode molecule may also be employed. Such modifications may inhibit the ability of the molecule (e.g., barcode molecule) to participate in a given reaction, however. Accordingly, a modification may be removed or reversed subsequent to an exonuclease digestion process, e.g., using an enzymatic or chemical mechanism.

[00171] The use of exonucleases such as Exonuclease I (ExoI) to “clean-up” combinatorial processes performed on supports (e.g., gel beads) permits digestion of partially ligated

molecules, leaving only fully ligated barcode molecules coupled to the supports. Notably, exonucleases such as ExoI may be stored in solutions comprising one or more materials capable of degrading supports (e.g., gel beads). For example, a solution for storing ExoI may include dithiothreitol (DTT), which may degrade gel beads. Accordingly, an exonuclease may be purified before it is used according to the methods provided herein.

[00172] A method of improving efficiency of a combinatorial assembly process may comprise performing one or more ligation processes (e.g., as described herein), performing an additional ligation process using a molecular segment or sequence comprising a phosphorothioate moiety, and subjecting the resulting ligation products to conditions sufficient to remove or degrade products that do not comprise the molecular segment or sequence comprising the phosphorothioate moiety. These conditions may comprise combining the ligation products with an exonuclease such as ExoI. The exonuclease may be included in a buffered solution, such as a solution comprising Tris-HCl, MgCl₂, NaCl, Triton X-100, glycerol, and/or ATP. The conditions may comprise incubation at, e.g., about 37 degrees Celsius (°C) or higher for several minutes. In an example, the ligation products may be incubated with the exonuclease at about 37°C for about 1 hour and may then be subjected to approximately 15 minutes of incubation at a higher temperature such as about 65°C.

[00173] **FIG. 26** shows the decrease in ligation efficiency throughout a triple ligation process. The overall ligation efficiency is about 80%, wherein after the triple ligation process, about 80% are final products, with double ligation products remaining at about 15% and single ligation products remaining at about 4%.

[00174] **FIG. 27** shows an ExoI treatment of barcode molecules coupled to gel beads. In each panel (having four lanes), the left two lanes show results of a reaction without phosphorothioate, and the right two lanes show results of a reaction with phosphorothioate. As shown in each of the panels, the use of phosphorothioate successfully inhibits ExoI activity. Similarly, **FIG. 28** shows another ExoI treatment of barcode molecules coupled to gel beads. The figure illustrates three sets of results, a first set (left) of four lanes at 1.5 dilution, a second set (center) of four lanes at 1.25 dilution, and a third set (right) of four lanes at 1.125 dilution. In each set, the first two lanes shows results without ExoI treatment, and the latter two lanes shows results with ExoI treatment. Within each pair of lanes, the left lane shows results of a reaction without phosphorothioate and the right lane shows results of a reaction with phosphorothioate. As seen in each of the sets, phosphorothioate successfully inhibits ExoI activity, leaving the desired product about 50% intact. The exonuclease digests all DNA products that do not have a phosphorothioate at their respective 3' ends. Notably, ExoI purified to remove DTT was incubated with gel beads and minimal bead swelling was observed during the reaction process.

[00175] **FIGs. 29A and 30A** show a triple ligation process in which the overall ligation efficiency without exonuclease treatment is about 75%. As shown in **FIG. 29B**, the overall ligation efficiency increases to about 94% after incorporating an exonuclease treatment. **FIG. 30B** shows a process in which efficiency increases to about 82% after incorporating an exonuclease treatment. In this process, an unexpected band is observed which may include a phosphorothioate moiety and/or be a double-stranded product. **FIG. 31** shows a scheme relating to these scenarios.

[00176] **FIG. 32** shows a process in which fluorescent probes were used to examine the effects of exonuclease treatment. The plot on the left shows a plot of green fluorescence (y-axis) vs red fluorescence (x-axis), without ExoI treatment. The plot on the right shows a plot of green fluorescence (y-axis) vs red fluorescence (x-axis), with ExoI treatment.

[00177] **FIG. 33A** shows a triple ligation process. **FIG. 33B** shows a comparison between exonuclease treatment and lack of exonuclease treatment for the triple ligation process shown in **FIG. 33A**. In this example, the efficiency of the ligation process increased from about 90% to about 93% upon addition of an exonuclease process. **FIG. 33C** shows a process in which urea and Z1 washes are performed. With and without exonuclease treatment, ligation 2 products are not digested. Accordingly, this gel rules out the possibility of off target phosphorothioate products. Hairpin formation may block the exonuclease activity.

[00178] ExoIII may be used in addition to or instead of other exonucleases such as ExoI. ExoIII digests double-stranded DNA and is blocked by phosphorothioate bonds. As it may be provided in a solution comprising DTT, hydrogen peroxide may be added to an ExoIII solution to oxidize DTT and prevent unwanted digestion of gel beads. **FIG. 34** shows a process involving both ExoI and ExoIII treatments. Gel bead size was unaffected by the exonucleases and hydrogen peroxide.

[00179] Heat-labile dsDNase (arcticzymes) may also be used as alternatives to ExoIII. Such enzymes do not include DTT and thus do not necessitate the addition of hydrogen peroxide. The enzymes may also be deactivated at lower temperatures than ExoIII. **FIG. 35** shows a process involving both ExoI and HL-dsDNase treatment. **FIG. 36** shows comparisons of processes involving no exonuclease treatment, ExoI treatment, and ExoI and dsDNase treatment in which green and red probes are utilized as in **FIG. 32**. The combination of ExoI and HL-dsDNase may fully digest all partially ligated products while leaving about 50% of the desired product intact. This combination allows generation of functionalized gel beads with >99% desired barcode molecules and <1% off-target products without affecting the size of the gel beads (no swelling observed).

[00180] The present disclosure also provides kits comprising a plurality of barcode molecules (e.g., nucleic acid barcode molecules). A kit may comprise a plurality of supports (e.g., beads, such as gel beads) and a plurality of barcode molecules coupled to the plurality of supports. The plurality of barcode molecules may comprise (i) a first set of barcode molecules coupled to a support of the plurality of supports and (ii) a second set of barcode molecules coupled to the same support. First barcode molecules of the first set of barcode molecules may be different than second barcode molecules of the second set of barcode molecules. First barcode molecules of the first set of barcode molecules may be configured to interact with different target molecules than second barcode molecules of the second set of barcode molecules. First barcode molecules of the first set of barcode molecules and second barcode molecules of the second set of barcode molecules may comprise barcode sequences that are different from barcode sequences of barcode molecules coupled to other supports of the plurality of supports.

[00181] First barcode molecules and second barcode molecules may comprise barcode sequences (e.g., as described herein). The barcode sequences of first barcode molecules and second barcode molecules attached to the same support may be identical.

[00182] The plurality of supports may be a plurality of beads (e.g., as described herein), such that barcode molecules (e.g., first barcode molecules of the first set of barcode molecules and the second barcode molecules of the second set of barcode molecules) are coupled to a bead of the plurality of beads. At least a subset of the barcode molecules coupled to a bead may be coupled to an interior of the bead. Alternatively or in addition, barcode molecules may be coupled to a surface (e.g., an interior or exterior surface) of a bead. Barcode molecules may be releasably coupled to a bead, e.g., via chemical cross-linkers (e.g., as described herein). Beads may be gel beads. Beads may be dissolvable or disruptable (e.g., upon application of an appropriate stimulus).

[00183] Barcode molecules (e.g., first barcode molecules or said second barcode molecules) coupled to a support (e.g., bead) may be configured to interact with various target molecules. For example, some or all barcode molecules coupled to a given support (e.g., bead) may be configured to interact with DNA molecules. Alternatively or in addition, some or all barcode molecules coupled to a given support (e.g., bead) may be configured to interact with RNA molecules (e.g., mRNA molecules). For example, barcode molecules may comprise a barcode sequence comprising nucleotides (e.g., nucleic acid sequences) that may be configured to interact with one or more nucleic acid molecules (e.g., DNA and RNA molecules). Barcode molecules (e.g., some or all barcode molecules coupled to a given support) may also be configured to interact with amino acids, polypeptides or proteins. For example, barcode molecules may

comprise a barcode sequence comprising amino acids that may be configured to interact with one or more amino acids, polypeptides, or proteins.

[00184] Different barcode molecules coupled to a support (e.g., bead) may have the same and/or different components (e.g., as described herein). For example, different barcode molecules coupled to the same support may comprise the same or different barcode sequences, starter sequences, functional sequences, or other sequences. In an example, first barcode molecules and second barcode molecules coupled to the same support (e.g., bead) may comprise a unique identifier that is different than unique identifiers of other barcode molecules coupled to the support. The barcode sequences and/or other sequences of the barcode molecules may be identical. First and second barcode molecules coupled to the same support may have both identical and different barcode sequences. For example, first barcode molecules may have first and second barcode sequences and second barcode molecules may have third and fourth barcode sequences, where the first and third barcode sequences are the same but the second and fourth barcode sequences are different. Different barcode molecules coupled to the same support may have different functional sequences. For example, first barcode molecules coupled to a support may have a first sequence for use with a first assay and second barcode molecules coupled to the same support may have a second sequence for use with a second assay. The first assay may comprise, for example, analysis of first target molecules (e.g., DNA molecules) and the second assay may comprise analysis of second target molecules (e.g., RNA molecules).

[00185] Multiple different barcode molecules may be coupled to the same support. For example, two different barcode molecule populations may be coupled to the same support. In another example, three or more different barcode molecule populations may be coupled to the same support. Multiple thousands of barcode molecules may be coupled to the same support. For example, a support (e.g., bead) may comprise at least 100,000 barcode molecules. Different barcode molecules may be present on a support in equal or different portions. For example, a greater number of first barcode molecules may be coupled to the support than second barcode molecules.

[00186] The present disclosure also provides a method for processing a plurality of analytes using a plurality of supports comprising a plurality of barcode molecules (e.g., as described herein). The method may comprise providing a plurality of barcode molecules coupled to a plurality of supports (e.g., beads, such as gel beads, as described herein). The plurality of barcode molecules may comprise first and second barcode molecules coupled to the same support. The first and second barcode molecules may be different. The support may be partitioned into a partition (e.g., a droplet or well), such that the partition comprises the support comprising the plurality of barcode molecules as well as a plurality of analytes. The plurality of

analytes may comprise, for example, nucleic acid molecules (e.g., DNA and RNA molecules), amino acids, polypeptides, and/or proteins. The plurality of analytes may derive from the same source. For example, the plurality of analytes may be included in or derived from the same biological particle (e.g., cell). The biological particle may be lysed or permeabilized to provide access to the plurality of analytes. Subsequent to partitioning, a first barcode molecule of the first barcode molecules coupled to the support and a first analyte (e.g., a DNA molecule) of the plurality of analytes may be used to generate a first barcoded analyte. A second barcode molecule of the second barcode molecules coupled to the support and a second analyte (e.g., an RNA molecule) of the plurality of analytes may be used to generate a second barcoded analyte. The first and second barcoded analytes, or derivatives thereof, may be recovered from the partition. The first and second barcoded analytes, or derivatives thereof, may then be subjected to subsequent processing and analysis, such as nucleic acid sequencing. A derivative of a barcoded analyte may include one or more functional sequences in addition to those of the barcoded analyte, such as one or more sequencing adapters or primers or other functional sequences. In an example, the first and second barcoded analytes are generated and are subsequently functionalized with sequencing adapters suitable for performing a nucleic acid assay (e.g., nucleic acid sequencing). In another example, a derivative of a barcoded analyte may comprise a portion of the barcoded analyte. For example, a portion of the barcoded analyte may be separated from the barcoded analyte and, optionally, another molecular segment or functional sequence may be added to the barcoded analyte.

[00187] Barcode molecules (e.g., first barcode molecules or said second barcode molecules) coupled to a support (e.g., bead) may be configured to interact with various analytes. For example, some or all barcode molecules coupled to a given support (e.g., bead) may be configured to interact with DNA molecules. Alternatively or in addition, some or all barcode molecules coupled to a given support (e.g., bead) may be configured to interact with RNA molecules (e.g., mRNA molecules). For example, barcode molecules may comprise a barcode sequence comprising nucleotides (e.g., nucleic acid sequences) that may be configured to interact with one or more nucleic acid molecules (e.g., DNA and RNA molecules). Barcode molecules configured to interact with DNA molecules may comprise a functional sequence (e.g., a capture sequence, such as a random N-mer) configured to interact with DNA molecules. Barcode molecules configured to interact with RNA molecules may comprise a functional sequence (e.g., a poly(T) sequence) configured to interact with RNA molecules (e.g., mRNA molecules). Barcode molecules (e.g., some or all barcode molecules coupled to a given support) may also be configured to interact with amino acids, polypeptides or proteins. For example, barcode

molecules may comprise a barcode sequence comprising amino acids that may be configured to interact with one or more amino acids, polypeptides, or proteins.

[00188] Different barcode molecules (e.g., first and second barcode molecules) coupled to the same support (e.g., bead) may comprise barcode sequences that are different from barcode sequences of barcode molecules coupled to other supports of a plurality of supports (e.g., as described herein). First barcode molecules and second barcode molecules may comprise barcode sequences (e.g., nucleic acid sequences that may be configured to interact with nucleic acid molecules or amino acid sequences that may be configured to interact with amino acids, polypeptides, or proteins). Barcode sequences of different barcode molecules coupled to the same bead may be identical. In some cases, barcode sequences of different barcode molecules may be both the same and different (e.g., first barcode molecules may comprise first and second barcode sequences and second barcode molecules coupled to the same bead may comprise third and fourth barcode sequences, where the first and third barcode sequences are the same and the second and fourth barcode sequences are different). In an example, each barcode molecule coupled to a given support (e.g., each first barcode molecule and each second barcode molecule) may have a different unique identifier (e.g., unique molecular identifier). Different barcode molecules coupled to the same support may be coupled to the same bead, such as the same gel bead. The barcode molecules may be releasably coupled to the bead, e.g., via chemical cross-linkers. The bead may be dissolvable or disruptable.

[00189] Multiple different barcode molecules may be coupled to the same support (e.g., bead). For example, two different barcode molecule populations may be coupled to the same support. In another example, three or more different barcode molecule populations may be coupled to the same support. Multiple thousands of barcode molecules may be coupled to the same support. For example, a support (e.g., bead) may comprise at least 100,000 barcode molecules. Different barcode molecules may be present on a support in equal or different portions. For example, a greater number of first barcode molecules may be coupled to the support than second barcode molecules. Barcode molecules of other supports of the plurality of supports may comprise different barcode sequences.

[00190] Generating barcoded analytes may comprise performing one or more nucleic acid extension reactions. A nucleic acid extension reaction may comprise annealing an analyte (e.g., a DNA or RNA molecule) to a barcode molecule of a plurality of barcode molecules coupled to a support. For example, the analyte may hybridize to a functional sequence of the barcode molecule. An extension reaction (e.g., primer extension reaction) may then take place (e.g., using a polymerase molecule such as a DNA polymerase molecule). The hybridization and extension reactions may be isothermal or may occur at one or more elevated temperatures. The

extension product may then be separated from the barcode molecule. The extension product (e.g., barcoded analyte) may comprise one or more barcode sequences of the barcode molecule or complements thereof. The barcode molecule or sequences thereof may be coupled to the support (e.g., bead) throughout the extension reaction, or the barcode molecule or a sequence thereof may be released from the support during or after the extension reaction. In an example, the first and second barcode molecule or sequences thereof may be released from the support prior to the nucleic acid extension reaction (e.g., upon application of an appropriate stimulus, as described herein). The first and/or second barcoded analytes may be coupled to the bead. The first and/or second barcoded analytes may be in solution in the partition (e.g., droplet or well) subsequent to their generation. For example, the first and/or second barcoded analytes may be generated in solution. Alternatively, the first and/or second barcoded analytes may be generated using barcode molecules that are coupled to support and subsequently released from the support (e.g., by denaturing the extension products (barcoded analytes) from the barcode molecules coupled to the support and/or by releasing the first and/or second barcode molecules used to generate the first and second barcoded analytes or sequences thereof from the support).

Systems and methods for sample compartmentalization

[00191] In an aspect, the systems and methods described herein provide for the compartmentalization, depositing, or partitioning of one or more particles (e.g., biological particles, macromolecular constituents of biological particles, beads, reagents, etc.) into discrete compartments or partitions (referred to interchangeably herein as partitions), where each partition maintains separation of its own contents from the contents of other partitions. The partition can be a droplet in an emulsion. A partition may comprise one or more other partitions.

[00192] A partition may include one or more particles. A partition may include one or more types of particles. For example, a partition of the present disclosure may comprise one or more biological particles and/or macromolecular constituents thereof. A partition may comprise one or more gel beads. A partition may comprise one or more cell beads. A partition may include a single gel bead, a single cell bead, or both a single cell bead and single gel bead. A partition may include one or more reagents. Alternatively, a partition may be unoccupied. For example, a partition may not comprise a bead. A cell bead can be a biological particle and/or one or more of its macromolecular constituents encased inside of a gel or polymer matrix, such as via polymerization of a droplet containing the biological particle and precursors capable of being polymerized or gelled. Unique identifiers, such as barcodes, may be injected into the droplets previous to, subsequent to, or concurrently with droplet generation, such as via a microcapsule (e.g., bead), as described elsewhere herein. Microfluidic channel networks (e.g., on a chip) can be utilized to generate partitions as described herein. Alternative mechanisms may also be

employed in the partitioning of individual biological particles, including porous membranes through which aqueous mixtures of cells are extruded into non-aqueous fluids.

[00193] The partitions can be flowable within fluid streams. The partitions may comprise, for example, micro-vesicles that have an outer barrier surrounding an inner fluid center or core. In some cases, the partitions may comprise a porous matrix that is capable of entraining and/or retaining materials within its matrix. The partitions can be droplets of a first phase within a second phase, wherein the first and second phases are immiscible. For example, the partitions can be droplets of aqueous fluid within a non-aqueous continuous phase (e.g., oil phase). In another example, the partitions can be droplets of a non-aqueous fluid within an aqueous phase. In some examples, the partitions may be provided in a water-in-oil emulsion or oil-in-water emulsion. A variety of different vessels are described in, for example, U.S. Patent Application Publication No. 2014/0155295, which is entirely incorporated herein by reference for all purposes. Emulsion systems for creating stable droplets in non-aqueous or oil continuous phases are described in, for example, U.S. Patent Application Publication No. 2010/0105112, which is entirely incorporated herein by reference for all purposes.

[00194] In the case of droplets in an emulsion, allocating individual particles to discrete partitions may in one non-limiting example be accomplished by introducing a flowing stream of particles in an aqueous fluid into a flowing stream of a non-aqueous fluid, such that droplets are generated at the junction of the two streams. Fluid properties (e.g., fluid flow rates, fluid viscosities, etc.), particle properties (e.g., volume fraction, particle size, particle concentration, etc.), microfluidic architectures (e.g., channel geometry, etc.), and other parameters may be adjusted to control the occupancy of the resulting partitions (e.g., number of biological particles per partition, number of beads per partition, etc.). For example, partition occupancy can be controlled by providing the aqueous stream at a certain concentration and/or flow rate of particles. To generate single biological particle partitions, the relative flow rates of the immiscible fluids can be selected such that, on average, the partitions may contain less than one biological particle per partition in order to ensure that those partitions that are occupied are primarily singly occupied. In some cases, partitions among a plurality of partitions may contain at most one biological particle (e.g., bead, DNA, cell or cellular material). In some embodiments, the various parameters (e.g., fluid properties, particle properties, microfluidic architectures, etc.) may be selected or adjusted such that a majority of partitions are occupied, for example, allowing for at most a small percentage of unoccupied partitions. The flows and channel architectures can be controlled as to ensure a given number of singly occupied partitions, less than a certain level of unoccupied partitions and/or less than a certain level of multiply occupied partitions.

[00195] FIG. 1 shows an example of a microfluidic channel structure 100 for partitioning individual biological particles. The channel structure 100 can include channel segments 102, 104, 106 and 108 communicating at a channel junction 110. In operation, a first aqueous fluid 112 that includes suspended biological particles (or cells) 114 may be transported along channel segment 102 into junction 110, while a second fluid 116 that is immiscible with the aqueous fluid 112 is delivered to the junction 110 from each of channel segments 104 and 106 to create discrete droplets 118, 120 of the first aqueous fluid 112 flowing into channel segment 108, and flowing away from junction 110. The channel segment 108 may be fluidically coupled to an outlet reservoir where the discrete droplets can be stored and/or harvested. A discrete droplet generated may include an individual biological particle 114 (such as droplets 118). A discrete droplet generated may include more than one individual biological particle 114 (not shown in FIG. 1). A discrete droplet may contain no biological particle 114 (such as droplet 120). Each discrete partition may maintain separation of its own contents (e.g., individual biological particle 114) from the contents of other partitions.

[00196] The second fluid 116 can comprise an oil, such as a fluorinated oil, that includes a fluorosurfactant for stabilizing the resulting droplets, for example, inhibiting subsequent coalescence of the resulting droplets 118, 120. Examples of particularly useful partitioning fluids and fluorosurfactants are described, for example, in U.S. Patent Application Publication No. 2010/0105112, which is entirely incorporated herein by reference for all purposes.

[00197] As will be appreciated, the channel segments described herein may be coupled to any of a variety of different fluid sources or receiving components, including reservoirs, tubing, manifolds, or fluidic components of other systems. As will be appreciated, the microfluidic channel structure 100 may have other geometries. For example, a microfluidic channel structure can have more than one channel junction. For example, a microfluidic channel structure can have 2, 3, 4, or 5 channel segments each carrying particles (e.g., biological particles, cell beads, and/or gel beads) that meet at a channel junction. Fluid may be directed to flow along one or more channels or reservoirs via one or more fluid flow units. A fluid flow unit can comprise compressors (e.g., providing positive pressure), pumps (e.g., providing negative pressure), actuators, and the like to control flow of the fluid. Fluid may also or otherwise be controlled via applied pressure differentials, centrifugal force, electrokinetic pumping, vacuum, capillary or gravity flow, or the like.

[00198] The generated droplets may comprise two subsets of droplets: (1) occupied droplets 118, containing one or more biological particles 114, and (2) unoccupied droplets 120, not containing any biological particles 114. Occupied droplets 118 may comprise singly occupied droplets (having one biological particle) and multiply occupied droplets (having more than one

biological particle). As described elsewhere herein, in some cases, the majority of occupied partitions can include no more than one biological particle per occupied partition and some of the generated partitions can be unoccupied (of any biological particle). In some cases, though, some of the occupied partitions may include more than one biological particle. In some cases, the partitioning process may be controlled such that fewer than about 25% of the occupied partitions contain more than one biological particle, and in many cases, fewer than about 20% of the occupied partitions have more than one biological particle, while in some cases, fewer than about 10% or even fewer than about 5% of the occupied partitions include more than one biological particle per partition.

[00199] In some cases, it may be desirable to minimize the creation of excessive numbers of empty partitions, such as to reduce costs and/or increase efficiency. While this minimization may be achieved by providing a sufficient number of biological particles (e.g., biological particles **114**) at the partitioning junction **110**, such as to ensure that at least one biological particle is encapsulated in a partition, the Poissonian distribution may expectedly increase the number of partitions that include multiple biological particles. As such, where singly occupied partitions are to be obtained, at most about 95%, 90%, 85%, 80%, 75%, 70%, 65%, 60%, 55%, 50%, 45%, 40%, 35%, 30%, 25%, 20%, 15%, 10%, 5% or less of the generated partitions can be unoccupied.

[00200] In some cases, the flow of one or more of the biological particles (e.g., in channel segment **102**), or other fluids directed into the partitioning junction (e.g., in channel segments **104**, **106**) can be controlled such that, in many cases, no more than about 50% of the generated partitions, no more than about 25% of the generated partitions, or no more than about 10% of the generated partitions are unoccupied. These flows can be controlled so as to present a non-Poissonian distribution of single-occupied partitions while providing lower levels of unoccupied partitions. The above noted ranges of unoccupied partitions can be achieved while still providing any of the single occupancy rates described above. For example, in many cases, the use of the systems and methods described herein can create resulting partitions that have multiple occupancy rates of less than about 25%, less than about 20%, less than about 15%, less than about 10%, and in many cases, less than about 5%, while having unoccupied partitions of less than about 50%, less than about 40%, less than about 30%, less than about 20%, less than about 10%, less than about 5%, or less.

[00201] As will be appreciated, the above-described occupancy rates are also applicable to partitions that include both biological particles and additional reagents, including, but not limited to, microcapsules or beads (e.g., gel beads) carrying barcoded nucleic acid molecules (e.g., oligonucleotides) (described in relation to **FIG. 2**). The occupied partitions (e.g., at least about

10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, or 99% of the occupied partitions) can include both a microcapsule (e.g., bead) comprising barcoded nucleic acid molecules and a biological particle.

[00202] In another aspect, in addition to or as an alternative to droplet based partitioning, biological particles may be encapsulated within a microcapsule that comprises an outer shell, layer or porous matrix in which is entrained one or more individual biological particles or small groups of biological particles. The microcapsule may include other reagents. Encapsulation of biological particles may be performed by a variety of processes. Such processes may combine an aqueous fluid containing the biological particles with a polymeric precursor material that may be capable of being formed into a gel or other solid or semi-solid matrix upon application of a particular stimulus to the polymer precursor. Such stimuli can include, for example, thermal stimuli (e.g., either heating or cooling), photo-stimuli (e.g., through photo-curing), chemical stimuli (e.g., through crosslinking, polymerization initiation of the precursor (e.g., through added initiators)), mechanical stimuli, or a combination thereof.

[00203] Preparation of microcapsules comprising biological particles may be performed by a variety of methods. For example, air knife droplet or aerosol generators may be used to dispense droplets of precursor fluids into gelling solutions in order to form microcapsules that include individual biological particles or small groups of biological particles. Likewise, membrane based encapsulation systems may be used to generate microcapsules comprising encapsulated biological particles as described herein. Microfluidic systems of the present disclosure, such as that shown in **FIG. 1**, may be readily used in encapsulating cells as described herein. In particular, and with reference to **FIG. 1**, the aqueous fluid **112** comprising (i) the biological particles **114** and (ii) the polymer precursor material (not shown) is flowed into channel junction **110**, where it is partitioned into droplets **118**, **120** through the flow of non-aqueous fluid **116**. In the case of encapsulation methods, non-aqueous fluid **116** may also include an initiator (not shown) to cause polymerization and/or crosslinking of the polymer precursor to form the microcapsule that includes the entrained biological particles. Examples of polymer precursor/initiator pairs include those described in U.S. Patent Application Publication No. 2014/0378345, which is entirely incorporated herein by reference for all purposes.

[00204] For example, in the case where the polymer precursor material comprises a linear polymer material, such as a linear polyacrylamide, PEG, or other linear polymeric material, the activation agent may comprise a cross-linking agent, or a chemical that activates a cross-linking agent within the formed droplets. Likewise, for polymer precursors that comprise polymerizable monomers, the activation agent may comprise a polymerization initiator. For example, in certain cases, where the polymer precursor comprises a mixture of acrylamide monomer with a N,N'-

bis-(acryloyl)cystamine (BAC) comonomer, an agent such as tetraethylmethylenediamine (TEMED) may be provided within the second fluid streams **116** in channel segments **104** and **106**, which can initiate the copolymerization of the acrylamide and BAC into a cross-linked polymer network, or hydrogel.

[00205] Upon contact of the second fluid stream **116** with the first fluid stream **112** at junction **110**, during formation of droplets, the TEMED may diffuse from the second fluid **116** into the aqueous fluid **112** comprising the linear polyacrylamide, which will activate the crosslinking of the polyacrylamide within the droplets **118**, **120**, resulting in the formation of gel (e.g., hydrogel) microcapsules, as solid or semi-solid beads or particles entraining the cells **114**. Although described in terms of polyacrylamide encapsulation, other 'activatable' encapsulation compositions may also be employed in the context of the methods and compositions described herein. For example, formation of alginate droplets followed by exposure to divalent metal ions (e.g., Ca^{2+} ions), can be used as an encapsulation process using the described processes. Likewise, agarose droplets may also be transformed into capsules through temperature based gelling (e.g., upon cooling, etc.).

[00206] In some cases, encapsulated biological particles can be selectively releasable from the microcapsule, such as through passage of time or upon application of a particular stimulus, that degrades the microcapsule sufficiently to allow the biological particles (e.g., cell), or its other contents to be released from the microcapsule, such as into a partition (e.g., droplet). For example, in the case of the polyacrylamide polymer described above, degradation of the microcapsule may be accomplished through the introduction of an appropriate reducing agent, such as DTT or the like, to cleave disulfide bonds that cross-link the polymer matrix. See, for example, U.S. Patent Application Publication No. 2014/0378345, which is entirely incorporated herein by reference for all purposes.

[00207] The biological particle can be subjected to other conditions sufficient to polymerize or gel the precursors. The conditions sufficient to polymerize or gel the precursors may comprise exposure to heating, cooling, electromagnetic radiation, and/or light. The conditions sufficient to polymerize or gel the precursors may comprise any conditions sufficient to polymerize or gel the precursors. Following polymerization or gelling, a polymer or gel may be formed around the biological particle. The polymer or gel may be diffusively permeable to chemical or biochemical reagents. The polymer or gel may be diffusively impermeable to macromolecular constituents of the biological particle. In this manner, the polymer or gel may act to allow the biological particle to be subjected to chemical or biochemical operations while spatially confining the macromolecular constituents to a region of the droplet defined by the polymer or gel. The polymer or gel may include one or more of disulfide cross-linked

polyacrylamide, agarose, alginate, polyvinyl alcohol, polyethylene glycol (PEG)-diacrylate, PEG-acrylate, PEG-thiol, PEG-azide, PEG-alkyne, other acrylates, chitosan, hyaluronic acid, collagen, fibrin, gelatin, or elastin. The polymer or gel may comprise any other polymer or gel.

[00208] The polymer or gel may be functionalized to bind to targeted analytes, such as nucleic acids, proteins, carbohydrates, lipids or other analytes. The polymer or gel may be polymerized or gelled via a passive mechanism. The polymer or gel may be stable in alkaline conditions or at elevated temperature. The polymer or gel may have mechanical properties similar to the mechanical properties of the bead. For instance, the polymer or gel may be of a similar size to the bead. The polymer or gel may have a mechanical strength (e.g. tensile strength) similar to that of the bead. The polymer or gel may be of a lower density than an oil. The polymer or gel may be of a density that is roughly similar to that of a buffer. The polymer or gel may have a tunable pore size. The pore size may be chosen to, for instance, retain denatured nucleic acids. The pore size may be chosen to maintain diffusive permeability to exogenous chemicals such as sodium hydroxide (NaOH) and/or endogenous chemicals such as inhibitors. The polymer or gel may be biocompatible. The polymer or gel may maintain or enhance cell viability. The polymer or gel may be biochemically compatible. The polymer or gel may be polymerized and/or depolymerized thermally, chemically, enzymatically, and/or optically.

[00209] The polymer may comprise poly(acrylamide-co-acrylic acid) crosslinked with disulfide linkages. The preparation of the polymer may comprise a two-step reaction. In the first activation step, poly(acrylamide-co-acrylic acid) may be exposed to an acylating agent to convert carboxylic acids to esters. For instance, the poly(acrylamide-co-acrylic acid) may be exposed to 4-(4,6-dimethoxy-1,3,5-triazin-2-yl)-4-methylmorpholinium chloride (DMTMM). The polyacrylamide-co-acrylic acid may be exposed to other salts of 4-(4,6-dimethoxy-1,3,5-triazin-2-yl)-4-methylmorpholinium. In the second cross-linking step, the ester formed in the first step may be exposed to a disulfide crosslinking agent. For instance, the ester may be exposed to cystamine (2,2'-dithiobis(ethylamine)). Following the two steps, the biological particle may be surrounded by polyacrylamide strands linked together by disulfide bridges. In this manner, the biological particle may be encased inside of or comprise a gel or matrix (e.g., polymer matrix) to form a "cell bead." A cell bead can contain biological particles (e.g., a cell) or macromolecular constituents (e.g., RNA, DNA, proteins, etc.) of biological particles. A cell bead may include a single cell or multiple cells, or a derivative of the single cell or multiple cells. For example after lysing and washing the cells, inhibitory components from cell lysates can be washed away and the macromolecular constituents can be bound as cell beads. Systems and methods disclosed herein can be applicable to both cell beads (and/or droplets or other

partitions) containing biological particles and cell beads (and/or droplets or other partitions) containing macromolecular constituents of biological particles.

[00210] Encapsulated biological particles can provide certain potential advantages of being more storable and more portable than droplet-based partitioned biological particles.

Furthermore, in some cases, it may be desirable to allow biological particles to incubate for a select period of time before analysis, such as in order to characterize changes in such biological particles over time, either in the presence or absence of different stimuli. In such cases, encapsulation may allow for longer incubation than partitioning in emulsion droplets, although in some cases, droplet partitioned biological particles may also be incubated for different periods of time, e.g., at least 10 seconds, at least 30 seconds, at least 1 minute, at least 5 minutes, at least 10 minutes, at least 30 minutes, at least 1 hour, at least 2 hours, at least 5 hours, or at least 10 hours or more. The encapsulation of biological particles may constitute the partitioning of the biological particles into which other reagents are co-partitioned. Alternatively or in addition, encapsulated biological particles may be readily deposited into other partitions (e.g., droplets) as described above.

Beads

[00211] A partition may comprise one or more unique identifiers, such as barcodes. Barcodes may be previously, subsequently or concurrently delivered to the partitions that hold the compartmentalized or partitioned biological particle. For example, barcodes may be injected into droplets previous to, subsequent to, or concurrently with droplet generation. The delivery of the barcodes to a particular partition allows for the later attribution of the characteristics of the individual biological particle to the particular partition. Barcodes may be delivered, for example on a nucleic acid molecule (e.g., an oligonucleotide), to a partition via any suitable mechanism. Barcoded nucleic acid molecules can be delivered to a partition via a microcapsule. A microcapsule, in some instances, can comprise a bead. Beads are described in further detail below.

[00212] In some cases, barcoded nucleic acid molecules can be initially associated with the microcapsule and then released from the microcapsule. Release of the barcoded nucleic acid molecules can be passive (e.g., by diffusion out of the microcapsule). In addition or alternatively, release from the microcapsule can be upon application of a stimulus which allows the barcoded nucleic acid nucleic acid molecules to dissociate or to be released from the microcapsule. Such stimulus may disrupt the microcapsule, an interaction that couples the barcoded nucleic acid molecules to or within the microcapsule, or both. Such stimulus can include, for example, a thermal stimulus, photo-stimulus, chemical stimulus (e.g., change in pH

or use of a reducing agent(s)), a mechanical stimulus, a radiation stimulus; a biological stimulus (e.g., enzyme), or any combination thereof.

[00213] FIG. 2 shows an example of a microfluidic channel structure **200** for delivering barcode carrying beads to droplets. The channel structure **200** can include channel segments **201**, **202**, **204**, **206** and **208** communicating at a channel junction **210**. In operation, the channel segment **201** may transport an aqueous fluid **212** that includes a plurality of beads **214** (e.g., with nucleic acid molecules, oligonucleotides, molecular tags) along the channel segment **201** into junction **210**. The plurality of beads **214** may be sourced from a suspension of beads. For example, the channel segment **201** may be connected to a reservoir comprising an aqueous suspension of beads **214**. The channel segment **202** may transport the aqueous fluid **212** that includes a plurality of biological particles **216** along the channel segment **202** into junction **210**. The plurality of biological particles **216** may be sourced from a suspension of biological particles. For example, the channel segment **202** may be connected to a reservoir comprising an aqueous suspension of biological particles **216**. In some instances, the aqueous fluid **212** in either the first channel segment **201** or the second channel segment **202**, or in both segments, can include one or more reagents, as further described below. A second fluid **218** that is immiscible with the aqueous fluid **212** (e.g., oil) can be delivered to the junction **210** from each of channel segments **204** and **206**. Upon meeting of the aqueous fluid **212** from each of channel segments **201** and **202** and the second fluid **218** from each of channel segments **204** and **206** at the channel junction **210**, the aqueous fluid **212** can be partitioned as discrete droplets **220** in the second fluid **218** and flow away from the junction **210** along channel segment **208**. The channel segment **208** may deliver the discrete droplets to an outlet reservoir fluidly coupled to the channel segment **208**, where they may be harvested.

[00214] As an alternative, the channel segments **201** and **202** may meet at another junction upstream of the junction **210**. At such junction, beads and biological particles may form a mixture that is directed along another channel to the junction **210** to yield droplets **220**. The mixture may provide the beads and biological particles in an alternating fashion, such that, for example, a droplet comprises a single bead and a single biological particle.

[00215] Beads, biological particles and droplets may flow along channels at substantially regular flow profiles (e.g., at regular flow rates). Such regular flow profiles may permit a droplet to include a single bead and a single biological particle. Such regular flow profiles may permit the droplets to have an occupancy (e.g., droplets having beads and biological particles) greater than 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or 95%. Such regular flow profiles and devices that may be used to provide such regular flow profiles are provided in, for example, U.S. Patent Publication No. 2015/0292988, which is entirely incorporated herein by reference.

[00216] The second fluid **218** can comprise an oil, such as a fluorinated oil, that includes a fluorosurfactant for stabilizing the resulting droplets, for example, inhibiting subsequent coalescence of the resulting droplets **220**.

[00217] A discrete droplet that is generated may include an individual biological particle **216**. A discrete droplet that is generated may include a barcode or other reagent carrying bead **214**. A discrete droplet generated may include both an individual biological particle and a barcode carrying bead, such as droplets **220**. In some instances, a discrete droplet may include more than one individual biological particle or no biological particle. In some instances, a discrete droplet may include more than one bead or no bead. A discrete droplet may be unoccupied (e.g., no beads, no biological particles).

[00218] Beneficially, a discrete droplet partitioning a biological particle and a barcode carrying bead may effectively allow the attribution of the barcode to macromolecular constituents of the biological particle within the partition. The contents of a partition may remain discrete from the contents of other partitions.

[00219] As will be appreciated, the channel segments described herein may be coupled to any of a variety of different fluid sources or receiving components, including reservoirs, tubing, manifolds, or fluidic components of other systems. As will be appreciated, the microfluidic channel structure **200** may have other geometries. For example, a microfluidic channel structure can have more than one channel junctions. For example, a microfluidic channel structure can have 2, 3, 4, or 5 channel segments each carrying beads that meet at a channel junction. Fluid may be directed flow along one or more channels or reservoirs via one or more fluid flow units. A fluid flow unit can comprise compressors (e.g., providing positive pressure), pumps (e.g., providing negative pressure), actuators, and the like to control flow of the fluid. Fluid may also or otherwise be controlled via applied pressure differentials, centrifugal force, electrokinetic pumping, vacuum, capillary or gravity flow, or the like.

[00220] A bead may be porous, non-porous, solid, semi-solid, semi-fluidic, fluidic, and/or a combination thereof. In some instances, a bead may be dissolvable, disruptable, and/or degradable. In some cases, a bead may not be degradable. In some cases, the bead may be a gel bead. A gel bead may be a hydrogel bead. A gel bead may be formed from molecular precursors, such as a polymeric or monomeric species. A semi-solid bead may be a liposomal bead. Solid beads may comprise metals including iron oxide, gold, and silver. In some cases, the bead may be a silica bead. In some cases, the bead can be rigid. In other cases, the bead may be flexible and/or compressible.

[00221] A bead may be of any suitable shape. Examples of bead shapes include, but are not limited to, spherical, non-spherical, oval, oblong, amorphous, circular, cylindrical, and variations thereof.

[00222] Beads may be of uniform size or heterogeneous size. In some cases, the diameter of a bead may be at least about 10 nanometers (nm), 100 nm, 500 nm, 1 micrometer (μm), 5 μm , 10 μm , 20 μm , 30 μm , 40 μm , 50 μm , 60 μm , 70 μm , 80 μm , 90 μm , 100 μm , 250 μm , 500 μm , 1mm, or greater. In some cases, a bead may have a diameter of less than about 10 nm, 100 nm, 500 nm, 1 μm , 5 μm , 10 μm , 20 μm , 30 μm , 40 μm , 50 μm , 60 μm , 70 μm , 80 μm , 90 μm , 100 μm , 250 μm , 500 μm , 1mm, or less. In some cases, a bead may have a diameter in the range of about 40-75 μm , 30-75 μm , 20-75 μm , 40-85 μm , 40-95 μm , 20-100 μm , 10-100 μm , 1-100 μm , 20-250 μm , or 20-500 μm .

[00223] In certain aspects, beads can be provided as a population or plurality of beads having a relatively monodisperse size distribution. Where it may be desirable to provide relatively consistent amounts of reagents within partitions, maintaining relatively consistent bead characteristics, such as size, can contribute to the overall consistency. In particular, the beads described herein may have size distributions that have a coefficient of variation in their cross-sectional dimensions of less than 50%, less than 40%, less than 30%, less than 20%, and in some cases less than 15%, less than 10%, less than 5%, or less.

[00224] A bead may comprise natural and/or synthetic materials. For example, a bead can comprise a natural polymer, a synthetic polymer or both natural and synthetic polymers. Examples of natural polymers include proteins and sugars such as deoxyribonucleic acid, rubber, cellulose, starch (e.g., amylose, amylopectin), proteins, enzymes, polysaccharides, silks, polyhydroxyalkanoates, chitosan, dextran, collagen, carrageenan, ispaghula, acacia, agar, gelatin, shellac, sterculia gum, xanthan gum, Corn sugar gum, guar gum, gum karaya, agarose, alginic acid, alginate, or natural polymers thereof. Examples of synthetic polymers include acrylics, nylons, silicones, spandex, viscose rayon, polycarboxylic acids, polyvinyl acetate, polyacrylamide, polyacrylate, polyethylene glycol, polyurethanes, polylactic acid, silica, polystyrene, polyacrylonitrile, polybutadiene, polycarbonate, polyethylene, polyethylene terephthalate, poly(chlorotrifluoroethylene), poly(ethylene oxide), poly(ethylene terephthalate), polyethylene, polyisobutylene, poly(methyl methacrylate), poly(oxymethylene), polyformaldehyde, polypropylene, polystyrene, poly(tetrafluoroethylene), poly(vinyl acetate), poly(vinyl alcohol), poly(vinyl chloride), poly(vinylidene dichloride), poly(vinylidene difluoride), poly(vinyl fluoride) and/or combinations (e.g., co-polymers) thereof. Beads may also be formed from materials other than polymers, including lipids, micelles, ceramics, glass-ceramics, material composites, metals, other inorganic materials, and others.

[00225] In some instances, the bead may contain molecular precursors (e.g., monomers or polymers), which may form a polymer network via polymerization of the molecular precursors. In some cases, a precursor may be an already polymerized species capable of undergoing further polymerization via, for example, a chemical cross-linkage. In some cases, a precursor can comprise one or more of an acrylamide or a methacrylamide monomer, oligomer, or polymer. In some cases, the bead may comprise prepolymers, which are oligomers capable of further polymerization. For example, polyurethane beads may be prepared using prepolymers. In some cases, the bead may contain individual polymers that may be further polymerized together. In some cases, beads may be generated via polymerization of different precursors, such that they comprise mixed polymers, co-polymers, and/or block co-polymers. In some cases, the bead may comprise covalent or ionic bonds between polymeric precursors (e.g., monomers, oligomers, linear polymers), nucleic acid molecules (e.g., oligonucleotides), primers, and other entities. In some cases, the covalent bonds can be carbon-carbon bonds, thioether bonds, or carbon-heteroatom bonds.

[00226] Cross-linking may be permanent or reversible, depending upon the particular cross-linker used. Reversible cross-linking may allow for the polymer to linearize or dissociate under appropriate conditions. In some cases, reversible cross-linking may also allow for reversible attachment of a material bound to the surface of a bead. In some cases, a cross-linker may form disulfide linkages. In some cases, the chemical cross-linker forming disulfide linkages may be cystamine or a modified cystamine.

[00227] In some cases, disulfide linkages can be formed between molecular precursor units (e.g., monomers, oligomers, or linear polymers) or precursors incorporated into a bead and nucleic acid molecules (e.g., oligonucleotides). Cystamine (including modified cystamines), for example, is an organic agent comprising a disulfide bond that may be used as a crosslinker agent between individual monomeric or polymeric precursors of a bead. Polyacrylamide may be polymerized in the presence of cystamine or a species comprising cystamine (e.g., a modified cystamine) to generate polyacrylamide gel beads comprising disulfide linkages (e.g., chemically degradable beads comprising chemically-reducible cross-linkers). The disulfide linkages may permit the bead to be degraded (or dissolved) upon exposure of the bead to a reducing agent.

[00228] In some cases, chitosan, a linear polysaccharide polymer, may be crosslinked with glutaraldehyde via hydrophilic chains to form a bead. Crosslinking of chitosan polymers may be achieved by chemical reactions that are initiated by heat, pressure, change in pH, and/or radiation.

[00229] In some cases, a bead may comprise an acrydite moiety, which in certain aspects may be used to attach one or more nucleic acid molecules (e.g., barcode sequence, barcoded nucleic

acid molecule, barcoded oligonucleotide, primer, or other oligonucleotide) to the bead. In some cases, an acrydite moiety can refer to an acrydite analogue generated from the reaction of acrydite with one or more species, such as, the reaction of acrydite with other monomers and cross-linkers during a polymerization reaction. Acrydite moieties may be modified to form chemical bonds with a species to be attached, such as a nucleic acid molecule (e.g., barcode sequence, barcoded nucleic acid molecule, barcoded oligonucleotide, primer, or other oligonucleotide). Acrydite moieties may be modified with thiol groups capable of forming a disulfide bond or may be modified with groups already comprising a disulfide bond. The thiol or disulfide (via disulfide exchange) may be used as an anchor point for a species to be attached or another part of the acrydite moiety may be used for attachment. In some cases, attachment can be reversible, such that when the disulfide bond is broken (e.g., in the presence of a reducing agent), the attached species is released from the bead. In other cases, an acrydite moiety can comprise a reactive hydroxyl group that may be used for attachment.

[00230] Functionalization of beads for attachment of nucleic acid molecules (e.g., oligonucleotides) may be achieved through a wide range of different approaches, including activation of chemical groups within a polymer, incorporation of active or activatable functional groups in the polymer structure, or attachment at the pre-polymer or monomer stage in bead production.

[00231] For example, precursors (e.g., monomers, cross-linkers) that are polymerized to form a bead may comprise acrydite moieties, such that when a bead is generated, the bead also comprises acrydite moieties. The acrydite moieties can be attached to a nucleic acid molecule (e.g., oligonucleotide), which may include a priming sequence (e.g., a primer for amplifying target nucleic acids, random primer, primer sequence for messenger RNA) and/or one or more barcode sequences. The one more barcode sequences may include sequences that are the same for all nucleic acid molecules coupled to a given bead and/or sequences that are different across all nucleic acid molecules coupled to the given bead. The nucleic acid molecule may be incorporated into the bead.

[00232] In some cases, the nucleic acid molecule can comprise a functional sequence, for example, for attachment to a sequencing flow cell, such as, for example, a P5 sequence for Illumina® sequencing. In some cases, the nucleic acid molecule or derivative thereof (e.g., oligonucleotide or polynucleotide generated from the nucleic acid molecule) can comprise another functional sequence, such as, for example, a P7 sequence for attachment to a sequencing flow cell for Illumina sequencing. In some cases, the nucleic acid molecule can comprise a barcode sequence. In some cases, the primer can further comprise a unique molecular identifier (UMI). In some cases, the primer can comprise an R1 primer sequence for Illumina sequencing.

In some cases, the primer can comprise an R2 primer sequence for Illumina sequencing. Examples of such nucleic acid molecules (e.g., oligonucleotides, polynucleotides, etc.) and uses thereof, as may be used with compositions, devices, methods and systems of the present disclosure, are provided in U.S. Patent Pub. Nos. 2014/0378345 and 2015/0376609, each of which is entirely incorporated herein by reference.

[00233] FIG. 8 illustrates an example of a barcode carrying bead. A nucleic acid molecule **802**, such as an oligonucleotide, can be coupled to a bead **804** by a releasable linkage **806**, such as, for example, a disulfide linker. The same bead **804** may be coupled (e.g., via releasable linkage) to one or more other nucleic acid molecules **818**, **820**. The nucleic acid molecule **802** may be or comprise a barcode. As noted elsewhere herein, the structure of the barcode may comprise a number of sequence elements. The nucleic acid molecule **802** may comprise a functional sequence **808** that may be used in subsequent processing. For example, the functional sequence **808** may include one or more of a sequencer specific flow cell attachment sequence (e.g., a P5 sequence for Illumina® sequencing systems) and a sequencing primer sequence (e.g., a R1 primer for Illumina® sequencing systems). The nucleic acid molecule **802** may comprise a barcode sequence **810** for use in barcoding the sample (e.g., DNA, RNA, protein, etc.). In some cases, the barcode sequence **810** can be bead-specific such that the barcode sequence **810** is common to all nucleic acid molecules (e.g., including nucleic acid molecule **802**) coupled to the same bead **804**. Alternatively or in addition, the barcode sequence **810** can be partition-specific such that the barcode sequence **810** is common to all nucleic acid molecules coupled to one or more beads that are partitioned into the same partition. The nucleic acid molecule **802** may comprise a specific priming sequence **812**, such as an mRNA specific priming sequence (e.g., poly-T sequence), a targeted priming sequence, and/or a random priming sequence. The nucleic acid molecule **802** may comprise an anchoring sequence **814** to ensure that the specific priming sequence **812** hybridizes at the sequence end (e.g., of the mRNA). For example, the anchoring sequence **814** can include a random short sequence of nucleotides, such as a 1-mer, 2-mer, 3-mer or longer sequence, which can ensure that a poly-T segment is more likely to hybridize at the sequence end of the poly-A tail of the mRNA.

[00234] The nucleic acid molecule **802** may comprise a unique molecular identifying sequence **816** (e.g., unique molecular identifier (UMI)). In some cases, the unique molecular identifying sequence **816** may comprise from about 5 to about 8 nucleotides. Alternatively, the unique molecular identifying sequence **816** may compress less than about 5 or more than about 8 nucleotides. The unique molecular identifying sequence **816** may be a unique sequence that varies across individual nucleic acid molecules (e.g., **802**, **818**, **820**, etc.) coupled to a single bead (e.g., bead **804**). In some cases, the unique molecular identifying sequence **816** may be a

random sequence (e.g., such as a random N-mer sequence). For example, the UMI may provide a unique identifier of the starting mRNA molecule that was captured, in order to allow quantitation of the number of original expressed RNA. As will be appreciated, although **FIG. 8** shows three nucleic acid molecules **802**, **818**, **820** coupled to the surface of the bead **804**, an individual bead may be coupled to any number of individual nucleic acid molecules, for example, from one to tens to hundreds of thousands or even millions of individual nucleic acid molecules. The respective barcodes for the individual nucleic acid molecules can comprise both common sequence segments or relatively common sequence segments (e.g., **808**, **810**, **812**, etc.) and variable or unique sequence segments (e.g., **816**) between different individual nucleic acid molecules coupled to the same bead.

[00235] In operation, a biological particle (e.g., cell, DNA, RNA, etc.) can be co-partitioned along with a barcode bearing bead **804**. The barcoded nucleic acid molecules **802**, **818**, **820** can be released from the bead **804** in the partition. By way of example, in the context of analyzing sample RNA, the poly-T segment (e.g., **812**) of one of the released nucleic acid molecules (e.g., **802**) can hybridize to the poly-A tail of an mRNA molecule. Reverse transcription may result in a cDNA transcript of the mRNA, but which transcript includes each of the sequence segments **808**, **810**, **816** of the nucleic acid molecule **802**. Because the nucleic acid molecule **802** comprises an anchoring sequence **814**, it will more likely hybridize to and prime reverse transcription at the sequence end of the poly-A tail of the mRNA. Within any given partition, all of the cDNA transcripts of the individual mRNA molecules may include a common barcode sequence segment **810**. However, the transcripts made from the different mRNA molecules within a given partition may vary at the unique molecular identifying sequence **812** segment (e.g., UMI segment). Beneficially, even following any subsequent amplification of the contents of a given partition, the number of different UMIs can be indicative of the quantity of mRNA originating from a given partition, and thus from the biological particle (e.g., cell). As noted above, the transcripts can be amplified, cleaned up and sequenced to identify the sequence of the cDNA transcript of the mRNA, as well as to sequence the barcode segment and the UMI segment. While a poly-T primer sequence is described, other targeted or random priming sequences may also be used in priming the reverse transcription reaction. Likewise, although described as releasing the barcoded oligonucleotides into the partition, in some cases, the nucleic acid molecules bound to the bead (e.g., gel bead) may be used to hybridize and capture the mRNA on the solid phase of the bead, for example, in order to facilitate the separation of the RNA from other cell contents.

[00236] In some cases, precursors comprising a functional group that is reactive or capable of being activated such that it becomes reactive can be polymerized with other precursors to

generate gel beads comprising the activated or activatable functional group. The functional group may then be used to attach additional species (e.g., disulfide linkers, primers, other oligonucleotides, etc.) to the gel beads. For example, some precursors comprising a carboxylic acid (COOH) group can co-polymerize with other precursors to form a gel bead that also comprises a COOH functional group. In some cases, acrylic acid (a species comprising free COOH groups), acrylamide, and bis(acryloyl)cystamine can be co-polymerized together to generate a gel bead comprising free COOH groups. The COOH groups of the gel bead can be activated (e.g., via 1-Ethyl-3-(3-dimethylaminopropyl)carbodiimide (EDC) and N-Hydroxysuccinimide (NHS) or 4-(4,6-Dimethoxy-1,3,5-triazin-2-yl)-4-methylmorpholinium chloride (DMTMM)) such that they are reactive (e.g., reactive to amine functional groups where EDC/NHS or DMTMM are used for activation). The activated COOH groups can then react with an appropriate species (e.g., a species comprising an amine functional group where the carboxylic acid groups are activated to be reactive with an amine functional group) comprising a moiety to be linked to the bead.

[00237] Beads comprising disulfide linkages in their polymeric network may be functionalized with additional species via reduction of some of the disulfide linkages to free thiols. The disulfide linkages may be reduced via, for example, the action of a reducing agent (e.g., DTT, TCEP, etc.) to generate free thiol groups, without dissolution of the bead. Free thiols of the beads can then react with free thiols of a species or a species comprising another disulfide bond (e.g., via thiol-disulfide exchange) such that the species can be linked to the beads (e.g., via a generated disulfide bond). In some cases, free thiols of the beads may react with any other suitable group. For example, free thiols of the beads may react with species comprising an acrydite moiety. The free thiol groups of the beads can react with the acrydite via Michael addition chemistry, such that the species comprising the acrydite is linked to the bead. In some cases, uncontrolled reactions can be prevented by inclusion of a thiol capping agent such as N-ethylmaleimide or iodoacetate.

[00238] Activation of disulfide linkages within a bead can be controlled such that at most a small number of disulfide linkages are activated. Control may be exerted, for example, by controlling the concentration of a reducing agent used to generate free thiol groups and/or concentration of reagents used to form disulfide bonds in bead polymerization. In some cases, a low concentration (e.g., molecules of reducing agent:gel bead ratios of less than or equal to about 1:100,000,000,000, less than or equal to about 1:10,000,000,000, less than or equal to about 1:1,000,000,000, less than or equal to about 1:100,000,000, less than or equal to about 1:10,000,000, less than or equal to about 1:1,000,000, less than or equal to about 1:100,000, less than or equal to about 1:10,000) of reducing agent may be used for reduction. Controlling the

number of disulfide linkages that are reduced to free thiols may be useful in ensuring bead structural integrity during functionalization. In some cases, optically-active agents, such as fluorescent dyes may be coupled to beads via free thiol groups of the beads and used to quantify the number of free thiols present in a bead and/or track a bead.

[00239] In some cases, addition of moieties to a gel bead after gel bead formation may be advantageous. For example, addition of an oligonucleotide (e.g., barcoded oligonucleotide) after gel bead formation may avoid loss of the species during chain transfer termination that can occur during polymerization. Moreover, smaller precursors (e.g., monomers or cross linkers that do not comprise side chain groups and linked moieties) may be used for polymerization and can be minimally hindered from growing chain ends due to viscous effects. In some cases, functionalization after gel bead synthesis can minimize exposure of species (e.g., oligonucleotides) to be loaded with potentially damaging agents (e.g., free radicals) and/or chemical environments. In some cases, the generated gel may possess an upper critical solution temperature (UCST) that can permit temperature driven swelling and collapse of a bead. Such functionality may aid in oligonucleotide (e.g., a primer) infiltration into the bead during subsequent functionalization of the bead with the oligonucleotide. Post-production functionalization may also be useful in controlling loading ratios of species in beads, such that, for example, the variability in loading ratio is minimized. Species loading may also be performed in a batch process such that a plurality of beads can be functionalized with the species in a single batch.

[00240] A bead injected or otherwise introduced into a partition may comprise releasably, cleavably, or reversibly attached barcodes. A bead injected or otherwise introduced into a partition may comprise activatable barcodes. A bead injected or otherwise introduced into a partition may be degradable, disruptable, or dissolvable beads.

[00241] Barcodes can be releasably, cleavably or reversibly attached to the beads such that barcodes can be released or be releasable through cleavage of a linkage between the barcode molecule and the bead, or released through degradation of the underlying bead itself, allowing the barcodes to be accessed or be accessible by other reagents, or both. In non-limiting examples, cleavage may be achieved through reduction of di-sulfide bonds, use of restriction enzymes, photo-activated cleavage, or cleavage via other types of stimuli (e.g., chemical, thermal, pH, enzymatic, etc.) and/or reactions, such as described elsewhere herein. Releasable barcodes may sometimes be referred to as being activatable, in that they are available for reaction once released. Thus, for example, an activatable barcode may be activated by releasing the barcode from a bead (or other suitable type of partition described herein). Other activatable configurations are also envisioned in the context of the described methods and systems.

[00242] In addition to, or as an alternative to the cleavable linkages between the beads and the associated molecules, such as barcode containing nucleic acid molecules (e.g., barcoded oligonucleotides), the beads may be degradable, disruptable, or dissolvable spontaneously or upon exposure to one or more stimuli (e.g., temperature changes, pH changes, exposure to particular chemical species or phase, exposure to light, reducing agent, etc.). In some cases, a bead may be dissolvable, such that material components of the beads are solubilized when exposed to a particular chemical species or an environmental change, such as a change in temperature or a change in pH. In some cases, a gel bead can be degraded or dissolved at elevated temperature and/or in basic conditions. In some cases, a bead may be thermally degradable such that when the bead is exposed to an appropriate change in temperature (e.g., heat), the bead degrades. Degradation or dissolution of a bead bound to a species (e.g., a nucleic acid molecule, e.g., barcoded oligonucleotide) may result in release of the species from the bead.

[00243] As will be appreciated from the above disclosure, the degradation of a bead may refer to the disassociation of a bound or entrained species from a bead, both with and without structurally degrading the physical bead itself. For example, the degradation of the bead may involve cleavage of a cleavable linkage via one or more species and/or methods described elsewhere herein. In another example, entrained species may be released from beads through osmotic pressure differences due to, for example, changing chemical environments. By way of example, alteration of bead pore sizes due to osmotic pressure differences can generally occur without structural degradation of the bead itself. In some cases, an increase in pore size due to osmotic swelling of a bead can permit the release of entrained species within the bead. In other cases, osmotic shrinking of a bead may cause a bead to better retain an entrained species due to pore size contraction.

[00244] A degradable bead may be introduced into a partition, such as a droplet of an emulsion or a well, such that the bead degrades within the partition and any associated species (e.g., oligonucleotides) are released within the droplet when the appropriate stimulus is applied. The free species (e.g., oligonucleotides, nucleic acid molecules) may interact with other reagents contained in the partition. For example, a polyacrylamide bead comprising cystamine and linked, via a disulfide bond, to a barcode sequence, may be combined with a reducing agent within a droplet of a water-in-oil emulsion. Within the droplet, the reducing agent can break the various disulfide bonds, resulting in bead degradation and release of the barcode sequence into the aqueous, inner environment of the droplet. In another example, heating of a droplet comprising a bead-bound barcode sequence in basic solution may also result in bead degradation and release of the attached barcode sequence into the aqueous, inner environment of the droplet.

[00245] Any suitable number of molecular tag molecules (e.g., primer, barcoded oligonucleotide) can be associated with a bead such that, upon release from the bead, the molecular tag molecules (e.g., primer, e.g., barcoded oligonucleotide) are present in the partition at a pre-defined concentration. Such pre-defined concentration may be selected to facilitate certain reactions for generating a sequencing library, e.g., amplification, within the partition. In some cases, the pre-defined concentration of the primer can be limited by the process of producing nucleic acid molecule (e.g., oligonucleotide) bearing beads.

[00246] In some cases, beads can be non-covalently loaded with one or more reagents. The beads can be non-covalently loaded by, for instance, subjecting the beads to conditions sufficient to swell the beads, allowing sufficient time for the reagents to diffuse into the interiors of the beads, and subjecting the beads to conditions sufficient to de-swell the beads. The swelling of the beads may be accomplished, for instance, by placing the beads in a thermodynamically favorable solvent, subjecting the beads to a higher or lower temperature, subjecting the beads to a higher or lower ion concentration, and/or subjecting the beads to an electric field. The swelling of the beads may be accomplished by various swelling methods. The de-swelling of the beads may be accomplished, for instance, by transferring the beads in a thermodynamically unfavorable solvent, subjecting the beads to lower or high temperatures, subjecting the beads to a lower or higher ion concentration, and/or removing an electric field. The de-swelling of the beads may be accomplished by various de-swelling methods. Transferring the beads may cause pores in the bead to shrink. The shrinking may then hinder reagents within the beads from diffusing out of the interiors of the beads. The hindrance may be due to steric interactions between the reagents and the interiors of the beads. The transfer may be accomplished microfluidically. For instance, the transfer may be achieved by moving the beads from one co-flowing solvent stream to a different co-flowing solvent stream. The swellability and/or pore size of the beads may be adjusted by changing the polymer composition of the bead.

[00247] In some cases, an acrydite moiety linked to a precursor, another species linked to a precursor, or a precursor itself can comprise a labile bond, such as chemically, thermally, or photo-sensitive bond e.g., disulfide bond, UV sensitive bond, or the like. Once acrydite moieties or other moieties comprising a labile bond are incorporated into a bead, the bead may also comprise the labile bond. The labile bond may be, for example, useful in reversibly linking (e.g., covalently linking) species (e.g., barcodes, primers, etc.) to a bead. In some cases, a thermally labile bond may include a nucleic acid hybridization based attachment, e.g., where an oligonucleotide is hybridized to a complementary sequence that is attached to the bead, such that thermal melting of the hybrid releases the oligonucleotide, e.g., a barcode containing sequence, from the bead or microcapsule.

[00248] The addition of multiple types of labile bonds to a gel bead may result in the generation of a bead capable of responding to varied stimuli. Each type of labile bond may be sensitive to an associated stimulus (e.g., chemical stimulus, light, temperature, enzymatic, etc.) such that release of species attached to a bead via each labile bond may be controlled by the application of the appropriate stimulus. Such functionality may be useful in controlled release of species from a gel bead. In some cases, another species comprising a labile bond may be linked to a gel bead after gel bead formation via, for example, an activated functional group of the gel bead as described above. As will be appreciated, barcodes that are releasably, cleavably or reversibly attached to the beads described herein include barcodes that are released or releasable through cleavage of a linkage between the barcode molecule and the bead, or that are released through degradation of the underlying bead itself, allowing the barcodes to be accessed or accessible by other reagents, or both.

[00249] The barcodes that are releasable as described herein may sometimes be referred to as being activatable, in that they are available for reaction once released. Thus, for example, an activatable barcode may be activated by releasing the barcode from a bead (or other suitable type of partition described herein). Other activatable configurations are also envisioned in the context of the described methods and systems.

[00250] In addition to thermally cleavable bonds, disulfide bonds and UV sensitive bonds, other non-limiting examples of labile bonds that may be coupled to a precursor or bead include an ester linkage (e.g., cleavable with an acid, a base, or hydroxylamine), a vicinal diol linkage (e.g., cleavable via sodium periodate), a Diels-Alder linkage (e.g., cleavable via heat), a sulfone linkage (e.g., cleavable via a base), a silyl ether linkage (e.g., cleavable via an acid), a glycosidic linkage (e.g., cleavable via an amylase), a peptide linkage (e.g., cleavable via a protease), or a phosphodiester linkage (e.g., cleavable via a nuclease (e.g., DNAase)). A bond may be cleavable via other nucleic acid molecule targeting enzymes, such as restriction enzymes (e.g., restriction endonucleases), as described further below.

[00251] Species may be encapsulated in beads during bead generation (e.g., during polymerization of precursors). Such species may or may not participate in polymerization. Such species may be entered into polymerization reaction mixtures such that generated beads comprise the species upon bead formation. In some cases, such species may be added to the gel beads after formation. Such species may include, for example, nucleic acid molecules (e.g., oligonucleotides), reagents for a nucleic acid amplification reaction (e.g., primers, polymerases, dNTPs, co-factors (e.g., ionic co-factors), buffers) including those described herein, reagents for enzymatic reactions (e.g., enzymes, co-factors, substrates, buffers), reagents for nucleic acid modification reactions such as polymerization, ligation, or digestion, and/or reagents for

template preparation (e.g., tagmentation) for one or more sequencing platforms (e.g., Nextera® for Illumina®). Such species may include one or more enzymes described herein, including without limitation, polymerase, reverse transcriptase, restriction enzymes (e.g., endonuclease), transposase, ligase, proteinase K, DNase, etc. Such species may include one or more reagents described elsewhere herein (e.g., lysis agents, inhibitors, inactivating agents, chelating agents, stimulus). Trapping of such species may be controlled by the polymer network density generated during polymerization of precursors, control of ionic charge within the gel bead (e.g., via ionic species linked to polymerized species), or by the release of other species. Encapsulated species may be released from a bead upon bead degradation and/or by application of a stimulus capable of releasing the species from the bead. Alternatively or in addition, species may be partitioned in a partition (e.g., droplet) during or subsequent to partition formation. Such species may include, without limitation, the abovementioned species that may also be encapsulated in a bead.

[00252] A degradable bead may comprise one or more species with a labile bond such that, when the bead/species is exposed to the appropriate stimuli, the bond is broken and the bead degrades. The labile bond may be a chemical bond (e.g., covalent bond, ionic bond) or may be another type of physical interaction (e.g., van der Waals interactions, dipole-dipole interactions, etc.). In some cases, a crosslinker used to generate a bead may comprise a labile bond. Upon exposure to the appropriate conditions, the labile bond can be broken and the bead degraded. For example, upon exposure of a polyacrylamide gel bead comprising cystamine crosslinkers to a reducing agent, the disulfide bonds of the cystamine can be broken and the bead degraded.

[00253] A degradable bead may be useful in more quickly releasing an attached species (e.g., a nucleic acid molecule, a barcode sequence, a primer, etc) from the bead when the appropriate stimulus is applied to the bead as compared to a bead that does not degrade. For example, for a species bound to an inner surface of a porous bead or in the case of an encapsulated species, the species may have greater mobility and accessibility to other species in solution upon degradation of the bead. In some cases, a species may also be attached to a degradable bead via a degradable linker (e.g., disulfide linker). The degradable linker may respond to the same stimuli as the degradable bead or the two degradable species may respond to different stimuli. For example, a barcode sequence may be attached, via a disulfide bond, to a polyacrylamide bead comprising cystamine. Upon exposure of the barcoded-bead to a reducing agent, the bead degrades and the barcode sequence is released upon breakage of both the disulfide linkage between the barcode sequence and the bead and the disulfide linkages of the cystamine in the bead.

[00254] As will be appreciated from the above disclosure, while referred to as degradation of a bead, in many instances as noted above, that degradation may refer to the disassociation of a

bound or entrained species from a bead, both with and without structurally degrading the physical bead itself. For example, entrained species may be released from beads through osmotic pressure differences due to, for example, changing chemical environments. By way of example, alteration of bead pore sizes due to osmotic pressure differences can generally occur without structural degradation of the bead itself. In some cases, an increase in pore size due to osmotic swelling of a bead can permit the release of entrained species within the bead. In other cases, osmotic shrinking of a bead may cause a bead to better retain an entrained species due to pore size contraction.

[00255] Where degradable beads are provided, it may be beneficial to avoid exposing such beads to the stimulus or stimuli that cause such degradation prior to a given time, in order to, for example, avoid premature bead degradation and issues that arise from such degradation, including for example poor flow characteristics and aggregation. By way of example, where beads comprise reducible cross-linking groups, such as disulfide groups, it will be desirable to avoid contacting such beads with reducing agents, e.g., DTT or other disulfide cleaving reagents. In such cases, treatment to the beads described herein will, in some cases be provided free of reducing agents, such as DTT. Because reducing agents are often provided in commercial enzyme preparations, it may be desirable to provide reducing agent free (or DTT free) enzyme preparations in treating the beads described herein. Examples of such enzymes include, e.g., polymerase enzyme preparations, reverse transcriptase enzyme preparations, ligase enzyme preparations, as well as many other enzyme preparations that may be used to treat the beads described herein. The terms “reducing agent free” or “DTT free” preparations can refer to a preparation having less than about 1/10th, less than about 1/50th, or even less than about 1/100th of the lower ranges for such materials used in degrading the beads. For example, for DTT, the reducing agent free preparation can have less than about 0.01 millimolar (mM), 0.005 mM, 0.001 mM DTT, 0.0005 mM DTT, or even less than about 0.0001 mM DTT. In many cases, the amount of DTT can be undetectable.

[00256] Numerous chemical triggers may be used to trigger the degradation of beads. Examples of these chemical changes may include, but are not limited to pH-mediated changes to the integrity of a component within the bead, degradation of a component of a bead via cleavage of cross-linked bonds, and depolymerization of a component of a bead.

[00257] In some embodiments, a bead may be formed from materials that comprise degradable chemical crosslinkers, such as BAC or cystamine. Degradation of such degradable crosslinkers may be accomplished through a number of mechanisms. In some examples, a bead may be contacted with a chemical degrading agent that may induce oxidation, reduction or other chemical changes. For example, a chemical degrading agent may be a reducing agent, such as

dithiothreitol (DTT). Additional examples of reducing agents may include β -mercaptoethanol, (2S)-2-amino-1,4-dimercaptobutane (dithiobutylamine or DTBA), tris(2-carboxyethyl) phosphine (TCEP), or combinations thereof. A reducing agent may degrade the disulfide bonds formed between gel precursors forming the bead, and thus, degrade the bead. In other cases, a change in pH of a solution, such as an increase in pH, may trigger degradation of a bead. In other cases, exposure to an aqueous solution, such as water, may trigger hydrolytic degradation, and thus degradation of the bead. In some cases, any combination of stimuli may trigger degradation of a bead. For example, a change in pH may enable a chemical agent (e.g., DTT) to become an effective reducing agent.

[00258] Beads may also be induced to release their contents upon the application of a thermal stimulus. A change in temperature can cause a variety of changes to a bead. For example, heat can cause a solid bead to liquefy. A change in heat may cause melting of a bead such that a portion of the bead degrades. In other cases, heat may increase the internal pressure of the bead components such that the bead ruptures or explodes. Heat may also act upon heat-sensitive polymers used as materials to construct beads.

[00259] Any suitable agent may degrade beads. In some embodiments, changes in temperature or pH may be used to degrade thermo-sensitive or pH-sensitive bonds within beads. In some embodiments, chemical degrading agents may be used to degrade chemical bonds within beads by oxidation, reduction or other chemical changes. For example, a chemical degrading agent may be a reducing agent, such as DTT, wherein DTT may degrade the disulfide bonds formed between a crosslinker and gel precursors, thus degrading the bead. In some embodiments, a reducing agent may be added to degrade the bead, which may or may not cause the bead to release its contents. Examples of reducing agents may include dithiothreitol (DTT), β -mercaptoethanol, (2S)-2-amino-1,4-dimercaptobutane (dithiobutylamine or DTBA), tris(2-carboxyethyl) phosphine (TCEP), or combinations thereof. The reducing agent may be present at a concentration of about 0.1mM, 0.5mM, 1mM, 5mM, 10mM. The reducing agent may be present at a concentration of at least about 0.1mM, 0.5mM, 1mM, 5mM, 10mM, or greater than 10 mM. The reducing agent may be present at concentration of at most about 10mM, 5mM, 1mM, 0.5mM, 0.1mM, or less.

[00260] Any suitable number of molecular tag molecules (e.g., primer, barcoded oligonucleotide) can be associated with a bead such that, upon release from the bead, the molecular tag molecules (e.g., primer, e.g., barcoded oligonucleotide) are present in the partition at a pre-defined concentration. Such pre-defined concentration may be selected to facilitate certain reactions for generating a sequencing library, e.g., amplification, within the partition. In

some cases, the pre-defined concentration of the primer can be limited by the process of producing oligonucleotide bearing beads.

[00261] Although **FIG. 1** and **FIG. 2** have been described in terms of providing substantially singly occupied partitions, above, in certain cases, it may be desirable to provide multiply occupied partitions, e.g., containing two, three, four or more cells and/or microcapsules (e.g., beads) comprising barcoded nucleic acid molecules (e.g., oligonucleotides) within a single partition. Accordingly, as noted above, the flow characteristics of the biological particle and/or bead containing fluids and partitioning fluids may be controlled to provide for such multiply occupied partitions. In particular, the flow parameters may be controlled to provide a given occupancy rate at greater than about 50% of the partitions, greater than about 75%, and in some cases greater than about 80%, 90%, 95%, or higher.

[00262] In some cases, additional microcapsules can be used to deliver additional reagents to a partition. In such cases, it may be advantageous to introduce different beads into a common channel or droplet generation junction, from different bead sources (e.g., containing different associated reagents) through different channel inlets into such common channel or droplet generation junction (e.g., junction **210**). In such cases, the flow and frequency of the different beads into the channel or junction may be controlled to provide for a certain ratio of microcapsules from each source, while ensuring a given pairing or combination of such beads into a partition with a given number of biological particles (e.g., one biological particle and one bead per partition).

[00263] The partitions described herein may comprise small volumes, for example, less than about 10 microliters (μL), 5 μL , 1 μL , 900 picoliters (pL), 800 pL, 700 pL, 600 pL, 500 pL, 400pL, 300 pL, 200 pL, 100pL, 50 pL, 20 pL, 10 pL, 1 pL, 500 nanoliters (nL), 100 nL, 50 nL, or less.

[00264] For example, in the case of droplet based partitions, the droplets may have overall volumes that are less than about 1000 pL, 900 pL, 800 pL, 700 pL, 600 pL, 500 pL, 400pL, 300 pL, 200 pL, 100pL, 50 pL, 20 pL, 10 pL, 1 pL, or less. Where co-partitioned with microcapsules, it will be appreciated that the sample fluid volume, e.g., including co-partitioned biological particles and/or beads, within the partitions may be less than about 90% of the above described volumes, less than about 80%, less than about 70%, less than about 60%, less than about 50%, less than about 40%, less than about 30%, less than about 20%, or less than about 10% of the above described volumes.

[00265] As is described elsewhere herein, partitioning species may generate a population or plurality of partitions. In such cases, any suitable number of partitions can be generated or otherwise provided. For example, at least about 1,000 partitions, at least about 5,000 partitions,

at least about 10,000 partitions, at least about 50,000 partitions, at least about 100,000 partitions, at least about 500,000 partitions, at least about 1,000,000 partitions, at least about 5,000,000 partitions at least about 10,000,000 partitions, at least about 50,000,000 partitions, at least about 100,000,000 partitions, at least about 500,000,000 partitions, at least about 1,000,000,000 partitions, or more partitions can be generated or otherwise provided. Moreover, the plurality of partitions may comprise both unoccupied partitions (e.g., empty partitions) and occupied partitions.

Reagents

[00266] In accordance with certain aspects, biological particles may be partitioned along with lysis reagents in order to release the contents of the biological particles within the partition. In such cases, the lysis agents can be contacted with the biological particle suspension concurrently with, or immediately prior to, the introduction of the biological particles into the partitioning junction/droplet generation zone (e.g., junction **210**), such as through an additional channel or channels upstream of the channel junction. In accordance with other aspects, additionally or alternatively, biological particles may be partitioned along with other reagents, as will be described further below.

[00267] **FIG. 3** shows an example of a microfluidic channel structure **300** for co-partitioning biological particles and reagents. The channel structure **300** can include channel segments **301**, **302**, **304**, **306** and **308**. Channel segments **301** and **302** communicate at a first channel junction **309**. Channel segments **302**, **304**, **306**, and **308** communicate at a second channel junction **310**.

[00268] In an example operation, the channel segment **301** may transport an aqueous fluid **312** that includes a plurality of biological particles **314** along the channel segment **301** into the second junction **310**. As an alternative or in addition to, channel segment **301** may transport beads (e.g., gel beads). The beads may comprise barcode molecules.

[00269] For example, the channel segment **301** may be connected to a reservoir comprising an aqueous suspension of biological particles **314**. Upstream of, and immediately prior to reaching, the second junction **310**, the channel segment **301** may meet the channel segment **302** at the first junction **309**. The channel segment **302** may transport a plurality of reagents **315** (e.g., lysis agents) suspended in the aqueous fluid **312** along the channel segment **302** into the first junction **309**. For example, the channel segment **302** may be connected to a reservoir comprising the reagents **315**. After the first junction **309**, the aqueous fluid **312** in the channel segment **301** can carry both the biological particles **314** and the reagents **315** towards the second junction **310**. In some instances, the aqueous fluid **312** in the channel segment **301** can include one or more reagents, which can be the same or different reagents as the reagents **315**. A second fluid **316** that is immiscible with the aqueous fluid **312** (e.g., oil) can be delivered to the second junction

310 from each of channel segments **304** and **306**. Upon meeting of the aqueous fluid **312** from the channel segment **301** and the second fluid **316** from each of channel segments **304** and **306** at the second channel junction **310**, the aqueous fluid **312** can be partitioned as discrete droplets **318** in the second fluid **316** and flow away from the second junction **310** along channel segment **308**. The channel segment **308** may deliver the discrete droplets **318** to an outlet reservoir fluidly coupled to the channel segment **308**, where they may be harvested.

[00270] The second fluid **316** can comprise an oil, such as a fluorinated oil, that includes a fluorosurfactant for stabilizing the resulting droplets, for example, inhibiting subsequent coalescence of the resulting droplets **318**.

[00271] A discrete droplet generated may include an individual biological particle **314** and/or one or more reagents **315**. In some instances, a discrete droplet generated may include a barcode carrying bead (not shown), such as via other microfluidics structures described elsewhere herein. In some instances, a discrete droplet may be unoccupied (e.g., no reagents, no biological particles).

[00272] Beneficially, when lysis reagents and biological particles are co-partitioned, the lysis reagents can facilitate the release of the contents of the biological particles within the partition. The contents released in a partition may remain discrete from the contents of other partitions.

[00273] As will be appreciated, the channel segments described herein may be coupled to any of a variety of different fluid sources or receiving components, including reservoirs, tubing, manifolds, or fluidic components of other systems. As will be appreciated, the microfluidic channel structure **300** may have other geometries. For example, a microfluidic channel structure can have more than two channel junctions. For example, a microfluidic channel structure can have 2, 3, 4, 5 channel segments or more each carrying the same or different types of beads, reagents, and/or biological particles that meet at a channel junction. Fluid flow in each channel segment may be controlled to control the partitioning of the different elements into droplets. Fluid may be directed flow along one or more channels or reservoirs via one or more fluid flow units. A fluid flow unit can comprise compressors (e.g., providing positive pressure), pumps (e.g., providing negative pressure), actuators, and the like to control flow of the fluid. Fluid may also or otherwise be controlled via applied pressure differentials, centrifugal force, electrokinetic pumping, vacuum, capillary or gravity flow, or the like.

[00274] Examples of lysis agents include bioactive reagents, such as lysis enzymes that are used for lysis of different cell types, e.g., gram positive or negative bacteria, plants, yeast, mammalian, etc., such as lysozymes, achromopeptidase, lysostaphin, labiase, kitalase, lyticase, and a variety of other lysis enzymes available from, e.g., Sigma-Aldrich, Inc. (St Louis, MO), as well as other commercially available lysis enzymes. Other lysis agents may additionally or

alternatively be co-partitioned with the biological particles to cause the release of the biological particles' contents into the partitions. For example, in some cases, surfactant-based lysis solutions may be used to lyse cells, although these may be less desirable for emulsion based systems where the surfactants can interfere with stable emulsions. In some cases, lysis solutions may include non-ionic surfactants such as, for example, TritonX-100 and Tween 20. In some cases, lysis solutions may include ionic surfactants such as, for example, sarcosyl and sodium dodecyl sulfate (SDS). Electroporation, thermal, acoustic or mechanical cellular disruption may also be used in certain cases, e.g., non-emulsion based partitioning such as encapsulation of biological particles that may be in addition to or in place of droplet partitioning, where any pore size of the encapsulate is sufficiently small to retain nucleic acid fragments of a given size, following cellular disruption.

[00275] Alternatively or in addition to the lysis agents co-partitioned with the biological particles described above, other reagents can also be co-partitioned with the biological particles, including, for example, DNase and RNase inactivating agents or inhibitors, such as proteinase K, chelating agents, such as EDTA, and other reagents employed in removing or otherwise reducing negative activity or impact of different cell lysate components on subsequent processing of nucleic acids. In addition, in the case of encapsulated biological particles, the biological particles may be exposed to an appropriate stimulus to release the biological particles or their contents from a co-partitioned microcapsule. For example, in some cases, a chemical stimulus may be co-partitioned along with an encapsulated biological particle to allow for the degradation of the microcapsule and release of the cell or its contents into the larger partition. In some cases, this stimulus may be the same as the stimulus described elsewhere herein for release of nucleic acid molecules (e.g., oligonucleotides) from their respective microcapsule (e.g., bead). In alternative aspects, this may be a different and non-overlapping stimulus, in order to allow an encapsulated biological particle to be released into a partition at a different time from the release of nucleic acid molecules into the same partition.

[00276] Additional reagents may also be co-partitioned with the biological particles, such as endonucleases to fragment a biological particle's DNA, DNA polymerase enzymes and dNTPs used to amplify the biological particle's nucleic acid fragments and to attach the barcode molecular tags to the amplified fragments. Other enzymes may be co-partitioned, including without limitation, polymerase, transposase, ligase, proteinase K, DNase, etc. Additional reagents may also include reverse transcriptase enzymes, including enzymes with terminal transferase activity, primers and oligonucleotides, and switch oligonucleotides (also referred to herein as "switch oligos" or "template switching oligonucleotides") which can be used for template switching. In some cases, template switching can be used to increase the length of a

cDNA. In some cases, template switching can be used to append a predefined nucleic acid sequence to the cDNA. In an example of template switching, cDNA can be generated from reverse transcription of a template, e.g., cellular mRNA, where a reverse transcriptase with terminal transferase activity can add additional nucleotides, e.g., polyC, to the cDNA in a template independent manner. Switch oligos can include sequences complementary to the additional nucleotides, e.g., polyG. The additional nucleotides (e.g., polyC) on the cDNA can hybridize to the additional nucleotides (e.g., polyG) on the switch oligo, whereby the switch oligo can be used by the reverse transcriptase as template to further extend the cDNA. Template switching oligonucleotides may comprise a hybridization region and a template region. The hybridization region can comprise any sequence capable of hybridizing to the target. In some cases, as described elsewhere herein, the hybridization region comprises a series of G bases to complement the overhanging C bases at the 3' end of a cDNA molecule. The series of G bases may comprise 1 G base, 2 G bases, 3 G bases, 4 G bases, 5 G bases or more than 5 G bases. The template sequence can comprise any sequence to be incorporated into the cDNA. In some cases, the template region comprises at least 1 (e.g., at least 2, 3, 4, 5 or more) tag sequences and/or functional sequences. Switch oligos may comprise deoxyribonucleic acids; ribonucleic acids; modified nucleic acids including 2-Aminopurine, 2,6-Diaminopurine (2-Amino-dA), inverted dT, 5-Methyl dC, 2'-deoxyInosine, Super T (5-hydroxybutynl-2'-deoxyuridine), Super G (8-aza-7-deazaguanosine), locked nucleic acids (LNAs), unlocked nucleic acids (UNAs, e.g., UNA-A, UNA-U, UNA-C, UNA-G), Iso-dG, Iso-dC, 2' Fluoro bases (e.g., Fluoro C, Fluoro U, Fluoro A, and Fluoro G), or any combination.

[00277] In some cases, the length of a switch oligo may be at least about 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249 or 250 nucleotides or longer.

[00278] In some cases, the length of a switch oligo may be at most about 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249 or 250 nucleotides.

[00279] Once the contents of the cells are released into their respective partitions, the macromolecular components (e.g., macromolecular constituents of biological particles, such as RNA, DNA, or proteins) contained therein may be further processed within the partitions. In accordance with the methods and systems described herein, the macromolecular component contents of individual biological particles can be provided with unique identifiers such that, upon characterization of those macromolecular components they may be attributed as having been derived from the same biological particle or particles. The ability to attribute characteristics to individual biological particles or groups of biological particles is provided by the assignment of unique identifiers specifically to an individual biological particle or groups of biological particles. Unique identifiers, e.g., in the form of nucleic acid barcodes can be assigned or associated with individual biological particles or populations of biological particles, in order to tag or label the biological particle's macromolecular components (and as a result, its characteristics) with the unique identifiers. These unique identifiers can then be used to attribute the biological particle's components and characteristics to an individual biological particle or group of biological particles.

[00280] In some aspects, this is performed by co-partitioning the individual biological particle or groups of biological particles with the unique identifiers, such as described above (with reference to **FIG. 2**). In some aspects, the unique identifiers are provided in the form of nucleic acid molecules (e.g., oligonucleotides) that comprise nucleic acid barcode sequences that may be attached to or otherwise associated with the nucleic acid contents of individual biological particle, or to other components of the biological particle, and particularly to fragments of those nucleic acids. The nucleic acid molecules are partitioned such that as between nucleic acid

molecules in a given partition, the nucleic acid barcode sequences contained therein are the same, but as between different partitions, the nucleic acid molecule can, and do have differing barcode sequences, or at least represent a large number of different barcode sequences across all of the partitions in a given analysis. In some aspects, one nucleic acid barcode sequence can be associated with a given partition, although in some cases, two or more different barcode sequences may be present.

[00281] The nucleic acid barcode sequences can include from about 6 to about 20 or more nucleotides within the sequence of the nucleic acid molecules (e.g., oligonucleotides). The nucleic acid barcode sequences can include from about 6 to about 20, 30, 40, 50, 60, 70, 80, 90, 100 or more nucleotides. In some cases, the length of a barcode sequence may be about 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 nucleotides or longer. In some cases, the length of a barcode sequence may be at least about 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 nucleotides or longer. In some cases, the length of a barcode sequence may be at most about 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 nucleotides or shorter. These nucleotides may be completely contiguous, i.e., in a single stretch of adjacent nucleotides, or they may be separated into two or more separate subsequences that are separated by 1 or more nucleotides. In some cases, separated barcode subsequences can be from about 4 to about 16 nucleotides in length. In some cases, the barcode subsequence may be about 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16 nucleotides or longer. In some cases, the barcode subsequence may be at least about 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16 nucleotides or longer. In some cases, the barcode subsequence may be at most about 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16 nucleotides or shorter.

[00282] The co-partitioned nucleic acid molecules can also comprise other functional sequences useful in the processing of the nucleic acids from the co-partitioned biological particles. These sequences include, e.g., targeted or random/universal amplification primer sequences for amplifying the genomic DNA from the individual biological particles within the partitions while attaching the associated barcode sequences, sequencing primers or primer recognition sites, hybridization or probing sequences, e.g., for identification of presence of the sequences or for pulling down barcoded nucleic acids, or any of a number of other potential functional sequences. Other mechanisms of co-partitioning oligonucleotides may also be employed, including, e.g., coalescence of two or more droplets, where one droplet contains oligonucleotides, or microdispensing of oligonucleotides into partitions, e.g., droplets within microfluidic systems.

[00283] In an example, microcapsules, such as beads, are provided that each include large numbers of the above described barcoded nucleic acid molecules (e.g., barcoded oligonucleotides) releasably attached to the beads, where all of the nucleic acid molecules

attached to a particular bead will include the same nucleic acid barcode sequence, but where a large number of diverse barcode sequences are represented across the population of beads used. In some embodiments, hydrogel beads, e.g., comprising polyacrylamide polymer matrices, are used as a solid support and delivery vehicle for the nucleic acid molecules into the partitions, as they are capable of carrying large numbers of nucleic acid molecules, and may be configured to release those nucleic acid molecules upon exposure to a particular stimulus, as described elsewhere herein. In some cases, the population of beads provides a diverse barcode sequence library that includes at least about 1,000 different barcode sequences, at least about 5,000 different barcode sequences, at least about 10,000 different barcode sequences, at least about 50,000 different barcode sequences, at least about 100,000 different barcode sequences, at least about 1,000,000 different barcode sequences, at least about 5,000,000 different barcode sequences, or at least about 10,000,000 different barcode sequences, or more. Additionally, each bead can be provided with large numbers of nucleic acid (e.g., oligonucleotide) molecules attached. In particular, the number of molecules of nucleic acid molecules including the barcode sequence on an individual bead can be at least about 1,000 nucleic acid molecules, at least about 5,000 nucleic acid molecules, at least about 10,000 nucleic acid molecules, at least about 50,000 nucleic acid molecules, at least about 100,000 nucleic acid molecules, at least about 500,000 nucleic acids, at least about 1,000,000 nucleic acid molecules, at least about 5,000,000 nucleic acid molecules, at least about 10,000,000 nucleic acid molecules, at least about 50,000,000 nucleic acid molecules, at least about 100,000,000 nucleic acid molecules, at least about 250,000,000 nucleic acid molecules and in some cases at least about 1 billion nucleic acid molecules, or more. Nucleic acid molecules of a given bead can include identical (or common) barcode sequences, different barcode sequences, or a combination of both. Nucleic acid molecules of a given bead can include multiple sets of nucleic acid molecules. Nucleic acid molecules of a given set can include identical barcode sequences. The identical barcode sequences can be different from barcode sequences of nucleic acid molecules of another set.

[00284] Moreover, when the population of beads is partitioned, the resulting population of partitions can also include a diverse barcode library that includes at least about 1,000 different barcode sequences, at least about 5,000 different barcode sequences, at least about 10,000 different barcode sequences, at least at least about 50,000 different barcode sequences, at least about 100,000 different barcode sequences, at least about 1,000,000 different barcode sequences, at least about 5,000,000 different barcode sequences, or at least about 10,000,000 different barcode sequences. Additionally, each partition of the population can include at least about 1,000 nucleic acid molecules, at least about 5,000 nucleic acid molecules, at least about 10,000 nucleic acid molecules, at least about 50,000 nucleic acid molecules, at least about 100,000 nucleic acid

molecules, at least about 500,000 nucleic acids, at least about 1,000,000 nucleic acid molecules, at least about 5,000,000 nucleic acid molecules, at least about 10,000,000 nucleic acid molecules, at least about 50,000,000 nucleic acid molecules, at least about 100,000,000 nucleic acid molecules, at least about 250,000,000 nucleic acid molecules and in some cases at least about 1 billion nucleic acid molecules.

[00285] In some cases, it may be desirable to incorporate multiple different barcodes within a given partition, either attached to a single or multiple beads within the partition. For example, in some cases, a mixed, but known set of barcode sequences may provide greater assurance of identification in the subsequent processing, e.g., by providing a stronger address or attribution of the barcodes to a given partition, as a duplicate or independent confirmation of the output from a given partition.

[00286] The nucleic acid molecules (e.g., oligonucleotides) are releasable from the beads upon the application of a particular stimulus to the beads. In some cases, the stimulus may be a photo-stimulus, e.g., through cleavage of a photo-labile linkage that releases the nucleic acid molecules. In other cases, a thermal stimulus may be used, where elevation of the temperature of the beads environment will result in cleavage of a linkage or other release of the nucleic acid molecules from the beads. In still other cases, a chemical stimulus can be used that cleaves a linkage of the nucleic acid molecules to the beads, or otherwise results in release of the nucleic acid molecules from the beads. In one case, such compositions include the polyacrylamide matrices described above for encapsulation of biological particles, and may be degraded for release of the attached nucleic acid molecules through exposure to a reducing agent, such as DTT.

[00287] In some aspects, provided are systems and methods for controlled partitioning. Droplet size may be controlled by adjusting certain geometric features in channel architecture (e.g., microfluidics channel architecture). For example, an expansion angle, width, and/or length of a channel may be adjusted to control droplet size.

[00288] **FIG. 4** shows an example of a microfluidic channel structure for the controlled partitioning of beads into discrete droplets. A channel structure **400** can include a channel segment **402** communicating at a channel junction **406** (or intersection) with a reservoir **404**. The reservoir **404** can be a chamber. Any reference to “reservoir,” as used herein, can also refer to a “chamber.” In operation, an aqueous fluid **408** that includes suspended beads **412** may be transported along the channel segment **402** into the junction **406** to meet a second fluid **410** that is immiscible with the aqueous fluid **408** in the reservoir **404** to create droplets **416**, **418** of the aqueous fluid **408** flowing into the reservoir **404**. At the junction **406** where the aqueous fluid **408** and the second fluid **410** meet, droplets can form based on factors such as the hydrodynamic

forces at the junction **406**, flow rates of the two fluids **408**, **410**, fluid properties, and certain geometric parameters (e.g., w , h_0 , α , etc.) of the channel structure **400**. A plurality of droplets can be collected in the reservoir **404** by continuously injecting the aqueous fluid **408** from the channel segment **402** through the junction **406**.

[00289] A discrete droplet generated may include a bead (e.g., as in occupied droplets **416**). Alternatively, a discrete droplet generated may include more than one bead. Alternatively, a discrete droplet generated may not include any beads (e.g., as in unoccupied droplet **418**). In some instances, a discrete droplet generated may contain one or more biological particles, as described elsewhere herein. In some instances, a discrete droplet generated may comprise one or more reagents, as described elsewhere herein.

[00290] In some instances, the aqueous fluid **408** can have a substantially uniform concentration or frequency of beads **412**. The beads **412** can be introduced into the channel segment **402** from a separate channel (not shown in **FIG. 4**). The frequency of beads **412** in the channel segment **402** may be controlled by controlling the frequency in which the beads **412** are introduced into the channel segment **402** and/or the relative flow rates of the fluids in the channel segment **402** and the separate channel. In some instances, the beads can be introduced into the channel segment **402** from a plurality of different channels, and the frequency controlled accordingly.

[00291] In some instances, the aqueous fluid **408** in the channel segment **402** can comprise biological particles (e.g., described with reference to **FIGS. 1** and **2**). In some instances, the aqueous fluid **408** can have a substantially uniform concentration or frequency of biological particles. As with the beads, the biological particles can be introduced into the channel segment **402** from a separate channel. The frequency or concentration of the biological particles in the aqueous fluid **408** in the channel segment **402** may be controlled by controlling the frequency in which the biological particles are introduced into the channel segment **402** and/or the relative flow rates of the fluids in the channel segment **402** and the separate channel. In some instances, the biological particles can be introduced into the channel segment **402** from a plurality of different channels, and the frequency controlled accordingly. In some instances, a first separate channel can introduce beads and a second separate channel can introduce biological particles into the channel segment **402**. The first separate channel introducing the beads may be upstream or downstream of the second separate channel introducing the biological particles.

[00292] The second fluid **410** can comprise an oil, such as a fluorinated oil, that includes a fluorosurfactant for stabilizing the resulting droplets, for example, inhibiting subsequent coalescence of the resulting droplets.

[00293] In some instances, the second fluid 410 may not be subjected to and/or directed to any flow in or out of the reservoir 404. For example, the second fluid 410 may be substantially stationary in the reservoir 404. In some instances, the second fluid 410 may be subjected to flow within the reservoir 404, but not in or out of the reservoir 404, such as via application of pressure to the reservoir 404 and/or as affected by the incoming flow of the aqueous fluid 408 at the junction 406. Alternatively, the second fluid 410 may be subjected and/or directed to flow in or out of the reservoir 404. For example, the reservoir 404 can be a channel directing the second fluid 410 from upstream to downstream, transporting the generated droplets.

[00294] The channel structure 400 at or near the junction 406 may have certain geometric features that at least partly determine the sizes of the droplets formed by the channel structure 400. The channel segment 402 can have a height, h_0 and width, w , at or near the junction 406. By way of example, the channel segment 402 can comprise a rectangular cross-section that leads to a reservoir 404 having a wider cross-section (such as in width or diameter). Alternatively, the cross-section of the channel segment 402 can be other shapes, such as a circular shape, trapezoidal shape, polygonal shape, or any other shapes. The top and bottom walls of the reservoir 404 at or near the junction 406 can be inclined at an expansion angle, α . The expansion angle, α , allows the tongue (portion of the aqueous fluid 408 leaving channel segment 402 at junction 406 and entering the reservoir 404 before droplet formation) to increase in depth and facilitate decrease in curvature of the intermediately formed droplet. Droplet size may decrease with increasing expansion angle. The resulting droplet radius, R_d , may be predicted by the following equation for the aforementioned geometric parameters of h_0 , w , and α :

$$R_d \approx 0.44 \left(1 + 2.2 \sqrt{\tan \alpha} \frac{w}{h_0} \right) \frac{h_0}{\sqrt{\tan \alpha}}$$

[00295] By way of example, for a channel structure with $w = 21 \mu\text{m}$, $h = 21 \mu\text{m}$, and $\alpha = 3^\circ$, the predicted droplet size is $121 \mu\text{m}$. In another example, for a channel structure with $w = 25 \mu\text{m}$, $h = 25 \mu\text{m}$, and $\alpha = 5^\circ$, the predicted droplet size is $123 \mu\text{m}$. In another example, for a channel structure with $w = 28 \mu\text{m}$, $h = 28 \mu\text{m}$, and $\alpha = 7^\circ$, the predicted droplet size is $124 \mu\text{m}$.

[00296] In some instances, the expansion angle, α , may be between a range of from about 0.5° to about 4° , from about 0.1° to about 10° , or from about 0° to about 90° . For example, the expansion angle can be at least about 0.01° , 0.1° , 0.2° , 0.3° , 0.4° , 0.5° , 0.6° , 0.7° , 0.8° , 0.9° , 1° , 2° , 3° , 4° , 5° , 6° , 7° , 8° , 9° , 10° , 15° , 20° , 25° , 30° , 35° , 40° , 45° , 50° , 55° , 60° , 65° , 70° , 75° , 80° , 85° , or higher. In some instances, the expansion angle can be at most about 89° , 88° , 87° , 86° , 85° , 84° , 83° , 82° , 81° , 80° , 75° , 70° , 65° , 60° , 55° , 50° , 45° , 40° , 35° , 30° , 25° , 20° , 15° , 10° , 9° , 8° , 7° , 6° , 5° , 4° , 3° , 2° , 1° , 0.1° , 0.01° , or less. In some instances, the width, w , can be between a range of from about 100 micrometers (μm) to about 500 μm . In some instances, the

width, w , can be between a range of from about 10 μm to about 200 μm . Alternatively, the width can be less than about 10 μm . Alternatively, the width can be greater than about 500 μm . In some instances, the flow rate of the aqueous fluid **408** entering the junction **406** can be between about 0.04 microliters (μL)/minute (min) and about 40 $\mu\text{L}/\text{min}$. In some instances, the flow rate of the aqueous fluid **408** entering the junction **406** can be between about 0.01 microliters (μL)/minute (min) and about 100 $\mu\text{L}/\text{min}$. Alternatively, the flow rate of the aqueous fluid **408** entering the junction **406** can be less than about 0.01 $\mu\text{L}/\text{min}$. Alternatively, the flow rate of the aqueous fluid **408** entering the junction **406** can be greater than about 40 $\mu\text{L}/\text{min}$, such as 45 $\mu\text{L}/\text{min}$, 50 $\mu\text{L}/\text{min}$, 55 $\mu\text{L}/\text{min}$, 60 $\mu\text{L}/\text{min}$, 65 $\mu\text{L}/\text{min}$, 70 $\mu\text{L}/\text{min}$, 75 $\mu\text{L}/\text{min}$, 80 $\mu\text{L}/\text{min}$, 85 $\mu\text{L}/\text{min}$, 90 $\mu\text{L}/\text{min}$, 95 $\mu\text{L}/\text{min}$, 100 $\mu\text{L}/\text{min}$, 110 $\mu\text{L}/\text{min}$, 120 $\mu\text{L}/\text{min}$, 130 $\mu\text{L}/\text{min}$, 140 $\mu\text{L}/\text{min}$, 150 $\mu\text{L}/\text{min}$, or greater. At lower flow rates, such as flow rates of about less than or equal to 10 microliters/minute, the droplet radius may not be dependent on the flow rate of the aqueous fluid **408** entering the junction **406**.

[00297] In some instances, at least about 50% of the droplets generated can have uniform size. In some instances, at least about 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or greater of the droplets generated can have uniform size. Alternatively, less than about 50% of the droplets generated can have uniform size.

[00298] The throughput of droplet generation can be increased by increasing the points of generation, such as increasing the number of junctions (e.g., junction **406**) between aqueous fluid **408** channel segments (e.g., channel segment **402**) and the reservoir **404**. Alternatively or in addition, the throughput of droplet generation can be increased by increasing the flow rate of the aqueous fluid **408** in the channel segment **402**.

[00299] **FIG. 5** shows an example of a microfluidic channel structure for increased droplet generation throughput. A microfluidic channel structure **500** can comprise a plurality of channel segments **502** and a reservoir **504**. Each of the plurality of channel segments **502** may be in fluid communication with the reservoir **504**. The channel structure **500** can comprise a plurality of channel junctions **506** between the plurality of channel segments **502** and the reservoir **504**. Each channel junction can be a point of droplet generation. The channel segment **402** from the channel structure **400** in **FIG. 4** and any description to the components thereof may correspond to a given channel segment of the plurality of channel segments **502** in channel structure **500** and any description to the corresponding components thereof. The reservoir **404** from the channel structure **400** and any description to the components thereof may correspond to the reservoir **504** from the channel structure **500** and any description to the corresponding components thereof.

[00300] Each channel segment of the plurality of channel segments **502** may comprise an aqueous fluid **508** that includes suspended beads **512**. The reservoir **504** may comprise a second

fluid **510** that is immiscible with the aqueous fluid **508**. In some instances, the second fluid **510** may not be subjected to and/or directed to any flow in or out of the reservoir **504**. For example, the second fluid **510** may be substantially stationary in the reservoir **504**. In some instances, the second fluid **510** may be subjected to flow within the reservoir **504**, but not in or out of the reservoir **504**, such as via application of pressure to the reservoir **504** and/or as affected by the incoming flow of the aqueous fluid **508** at the junctions. Alternatively, the second fluid **510** may be subjected and/or directed to flow in or out of the reservoir **504**. For example, the reservoir **504** can be a channel directing the second fluid **510** from upstream to downstream, transporting the generated droplets.

[00301] In operation, the aqueous fluid **508** that includes suspended beads **512** may be transported along the plurality of channel segments **502** into the plurality of junctions **506** to meet the second fluid **510** in the reservoir **504** to create droplets **516**, **518**. A droplet may form from each channel segment at each corresponding junction with the reservoir **504**. At the junction where the aqueous fluid **508** and the second fluid **510** meet, droplets can form based on factors such as the hydrodynamic forces at the junction, flow rates of the two fluids **508**, **510**, fluid properties, and certain geometric parameters (e.g., w , h_0 , α , etc.) of the channel structure **500**, as described elsewhere herein. A plurality of droplets can be collected in the reservoir **504** by continuously injecting the aqueous fluid **508** from the plurality of channel segments **502** through the plurality of junctions **506**. Throughput may significantly increase with the parallel channel configuration of channel structure **500**. For example, a channel structure having five inlet channel segments comprising the aqueous fluid **508** may generate droplets five times as frequently than a channel structure having one inlet channel segment, provided that the fluid flow rate in the channel segments are substantially the same. The fluid flow rate in the different inlet channel segments may or may not be substantially the same. A channel structure may have as many parallel channel segments as is practical and allowed for the size of the reservoir. For example, the channel structure may have at least about 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 150, 500, 250, 300, 350, 400, 450, 500, 600, 700, 800, 900, 1000, 1500, 5000 or more parallel or substantially parallel channel segments.

[00302] The geometric parameters, w , h_0 , and α , may or may not be uniform for each of the channel segments in the plurality of channel segments **502**. For example, each channel segment may have the same or different widths at or near its respective channel junction with the reservoir **504**. For example, each channel segment may have the same or different height at or near its respective channel junction with the reservoir **504**. In another example, the reservoir **504** may have the same or different expansion angle at the different channel junctions with the plurality of channel segments **502**. When the geometric parameters are uniform, beneficially,

droplet size may also be controlled to be uniform even with the increased throughput. In some instances, when it is desirable to have a different distribution of droplet sizes, the geometric parameters for the plurality of channel segments **502** may be varied accordingly.

[00303] In some instances, at least about 50% of the droplets generated can have uniform size. In some instances, at least about 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or greater of the droplets generated can have uniform size. Alternatively, less than about 50% of the droplets generated can have uniform size.

[00304] **FIG. 6** shows another example of a microfluidic channel structure for increased droplet generation throughput. A microfluidic channel structure **600** can comprise a plurality of channel segments **602** arranged generally circularly around the perimeter of a reservoir **604**. Each of the plurality of channel segments **602** may be in fluid communication with the reservoir **604**. The channel structure **600** can comprise a plurality of channel junctions **606** between the plurality of channel segments **602** and the reservoir **604**. Each channel junction can be a point of droplet generation. The channel segment **402** from the channel structure **400** in **FIG. 2** and any description to the components thereof may correspond to a given channel segment of the plurality of channel segments **602** in channel structure **600** and any description to the corresponding components thereof. The reservoir **404** from the channel structure **400** and any description to the components thereof may correspond to the reservoir **604** from the channel structure **600** and any description to the corresponding components thereof.

[00305] Each channel segment of the plurality of channel segments **602** may comprise an aqueous fluid **608** that includes suspended beads **612**. The reservoir **604** may comprise a second fluid **610** that is immiscible with the aqueous fluid **608**. In some instances, the second fluid **610** may not be subjected to and/or directed to any flow in or out of the reservoir **604**. For example, the second fluid **610** may be substantially stationary in the reservoir **604**. In some instances, the second fluid **610** may be subjected to flow within the reservoir **604**, but not in or out of the reservoir **604**, such as via application of pressure to the reservoir **604** and/or as affected by the incoming flow of the aqueous fluid **608** at the junctions. Alternatively, the second fluid **610** may be subjected and/or directed to flow in or out of the reservoir **604**. For example, the reservoir **604** can be a channel directing the second fluid **610** from upstream to downstream, transporting the generated droplets.

[00306] In operation, the aqueous fluid **608** that includes suspended beads **612** may be transported along the plurality of channel segments **602** into the plurality of junctions **606** to meet the second fluid **610** in the reservoir **604** to create a plurality of droplets **616**. A droplet may form from each channel segment at each corresponding junction with the reservoir **604**. At the junction where the aqueous fluid **608** and the second fluid **610** meet, droplets can form based

on factors such as the hydrodynamic forces at the junction, flow rates of the two fluids **608**, **610**, fluid properties, and certain geometric parameters (e.g., widths and heights of the channel segments **602**, expansion angle of the reservoir **604**, etc.) of the channel structure **600**, as described elsewhere herein. A plurality of droplets can be collected in the reservoir **604** by continuously injecting the aqueous fluid **608** from the plurality of channel segments **602** through the plurality of junctions **606**. Throughput may significantly increase with the substantially parallel channel configuration of the channel structure **600**. A channel structure may have as many substantially parallel channel segments as is practical and allowed for by the size of the reservoir. For example, the channel structure may have at least about 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250, 300, 350, 400, 450, 500, 600, 700, 800, 900, 1000, 1500, 5000 or more parallel or substantially parallel channel segments. The plurality of channel segments may be substantially evenly spaced apart, for example, around an edge or perimeter of the reservoir. Alternatively, the spacing of the plurality of channel segments may be uneven.

[00307] The reservoir **604** may have an expansion angle, α (not shown in **FIG. 6**) at or near each channel junction. Each channel segment of the plurality of channel segments **602** may have a width, w , and a height, h_o , at or near the channel junction. The geometric parameters, w , h_o , and α , may or may not be uniform for each of the channel segments in the plurality of channel segments **602**. For example, each channel segment may have the same or different widths at or near its respective channel junction with the reservoir **604**. For example, each channel segment may have the same or different height at or near its respective channel junction with the reservoir **604**.

[00308] The reservoir **604** may have the same or different expansion angle at the different channel junctions with the plurality of channel segments **602**. For example, a circular reservoir (as shown in **FIG. 6**) may have a conical, dome-like, or hemispherical ceiling (e.g., top wall) to provide the same or substantially same expansion angle for each channel segments **602** at or near the plurality of channel junctions **606**. When the geometric parameters are uniform, beneficially, resulting droplet size may be controlled to be uniform even with the increased throughput. In some instances, when it is desirable to have a different distribution of droplet sizes, the geometric parameters for the plurality of channel segments **602** may be varied accordingly.

[00309] In some instances, at least about 50% of the droplets generated can have uniform size. In some instances, at least about 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or greater of the droplets generated can have uniform size. Alternatively, less than about 50% of the droplets generated can have uniform size. The beads and/or biological particle injected into the droplets may or may not have uniform size.

[00310] FIG. 7A shows a cross-section view of another example of a microfluidic channel structure with a geometric feature for controlled partitioning. A channel structure 700 can include a channel segment 702 communicating at a channel junction 706 (or intersection) with a reservoir 704. In some instances, the channel structure 700 and one or more of its components can correspond to the channel structure 100 and one or more of its components. FIG. 7B shows a perspective view of the channel structure 700 of FIG. 7A.

[00311] An aqueous fluid 712 comprising a plurality of particles 716 may be transported along the channel segment 702 into the junction 706 to meet a second fluid 714 (e.g., oil, etc.) that is immiscible with the aqueous fluid 712 in the reservoir 704 to create droplets 720 of the aqueous fluid 712 flowing into the reservoir 704. At the junction 706 where the aqueous fluid 712 and the second fluid 714 meet, droplets can form based on factors such as the hydrodynamic forces at the junction 706, relative flow rates of the two fluids 712, 714, fluid properties, and certain geometric parameters (e.g., Δh , etc.) of the channel structure 700. A plurality of droplets can be collected in the reservoir 704 by continuously injecting the aqueous fluid 712 from the channel segment 702 at the junction 706.

[00312] A discrete droplet generated may comprise one or more particles of the plurality of particles 716. As described elsewhere herein, a particle may be any particle, such as a bead, cell bead, gel bead, biological particle, macromolecular constituents of biological particle, or other particles. Alternatively, a discrete droplet generated may not include any particles.

[00313] In some instances, the aqueous fluid 712 can have a substantially uniform concentration or frequency of particles 716. As described elsewhere herein (e.g., with reference to FIG. 4), the particles 716 (e.g., beads) can be introduced into the channel segment 702 from a separate channel (not shown in FIG. 7). The frequency of particles 716 in the channel segment 702 may be controlled by controlling the frequency in which the particles 716 are introduced into the channel segment 702 and/or the relative flow rates of the fluids in the channel segment 702 and the separate channel. In some instances, the particles 716 can be introduced into the channel segment 702 from a plurality of different channels, and the frequency controlled accordingly. In some instances, different particles may be introduced via separate channels. For example, a first separate channel can introduce beads and a second separate channel can introduce biological particles into the channel segment 702. The first separate channel introducing the beads may be upstream or downstream of the second separate channel introducing the biological particles.

[00314] In some instances, the second fluid 714 may not be subjected to and/or directed to any flow in or out of the reservoir 704. For example, the second fluid 714 may be substantially stationary in the reservoir 704. In some instances, the second fluid 714 may be subjected to flow

within the reservoir **704**, but not in or out of the reservoir **704**, such as via application of pressure to the reservoir **704** and/or as affected by the incoming flow of the aqueous fluid **712** at the junction **706**. Alternatively, the second fluid **714** may be subjected and/or directed to flow in or out of the reservoir **704**. For example, the reservoir **704** can be a channel directing the second fluid **714** from upstream to downstream, transporting the generated droplets.

[00315] The channel structure **700** at or near the junction **706** may have certain geometric features that at least partly determine the sizes and/or shapes of the droplets formed by the channel structure **700**. The channel segment **702** can have a first cross-section height, h_1 , and the reservoir **704** can have a second cross-section height, h_2 . The first cross-section height, h_1 , and the second cross-section height, h_2 , may be different, such that at the junction **706**, there is a height difference of Δh . The second cross-section height, h_2 , may be greater than the first cross-section height, h_1 . In some instances, the reservoir may thereafter gradually increase in cross-section height, for example, the more distant it is from the junction **706**. In some instances, the cross-section height of the reservoir may increase in accordance with expansion angle, β , at or near the junction **706**. The height difference, Δh , and/or expansion angle, β , can allow the tongue (portion of the aqueous fluid **712** leaving channel segment **702** at junction **706** and entering the reservoir **704** before droplet formation) to increase in depth and facilitate decrease in curvature of the intermediately formed droplet. For example, droplet size may decrease with increasing height difference and/or increasing expansion angle.

[00316] The height difference, Δh , can be at least about 1 μm . Alternatively, the height difference can be at least about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500 μm or more. Alternatively, the height difference can be at most about 500, 400, 300, 200, 100, 90, 80, 70, 60, 50, 45, 40, 35, 30, 25, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, 1 μm or less. In some instances, the expansion angle, β , may be between a range of from about 0.5° to about 4° , from about 0.1° to about 10° , or from about 0° to about 90° . For example, the expansion angle can be at least about 0.01° , 0.1° , 0.2° , 0.3° , 0.4° , 0.5° , 0.6° , 0.7° , 0.8° , 0.9° , 1° , 2° , 3° , 4° , 5° , 6° , 7° , 8° , 9° , 10° , 15° , 20° , 25° , 30° , 35° , 40° , 45° , 50° , 55° , 60° , 65° , 70° , 75° , 80° , 85° , or higher. In some instances, the expansion angle can be at most about 89° , 88° , 87° , 86° , 85° , 84° , 83° , 82° , 81° , 80° , 75° , 70° , 65° , 60° , 55° , 50° , 45° , 40° , 35° , 30° , 25° , 20° , 15° , 10° , 9° , 8° , 7° , 6° , 5° , 4° , 3° , 2° , 1° , 0.1° , 0.01° , or less.

[00317] In some instances, the flow rate of the aqueous fluid **712** entering the junction **706** can be between about 0.04 microliters (μL)/minute (min) and about 40 $\mu\text{L}/\text{min}$. In some instances, the flow rate of the aqueous fluid **712** entering the junction **706** can be between about 0.01 microliters (μL)/minute (min) and about 100 $\mu\text{L}/\text{min}$. Alternatively, the flow rate of the

aqueous fluid **712** entering the junction **706** can be less than about 0.01 $\mu\text{L}/\text{min}$. Alternatively, the flow rate of the aqueous fluid **712** entering the junction **706** can be greater than about 40 $\mu\text{L}/\text{min}$, such as 45 $\mu\text{L}/\text{min}$, 50 $\mu\text{L}/\text{min}$, 55 $\mu\text{L}/\text{min}$, 60 $\mu\text{L}/\text{min}$, 65 $\mu\text{L}/\text{min}$, 70 $\mu\text{L}/\text{min}$, 75 $\mu\text{L}/\text{min}$, 80 $\mu\text{L}/\text{min}$, 85 $\mu\text{L}/\text{min}$, 90 $\mu\text{L}/\text{min}$, 95 $\mu\text{L}/\text{min}$, 100 $\mu\text{L}/\text{min}$, 110 $\mu\text{L}/\text{min}$, 120 $\mu\text{L}/\text{min}$, 130 $\mu\text{L}/\text{min}$, 140 $\mu\text{L}/\text{min}$, 150 $\mu\text{L}/\text{min}$, or greater. At lower flow rates, such as flow rates of about less than or equal to 10 microliters/minute, the droplet radius may not be dependent on the flow rate of the aqueous fluid **712** entering the junction **706**. The second fluid **714** may be stationary, or substantially stationary, in the reservoir **704**. Alternatively, the second fluid **714** may be flowing, such as at the above flow rates described for the aqueous fluid **712**.

[00318] In some instances, at least about 50% of the droplets generated can have uniform size. In some instances, at least about 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or greater of the droplets generated can have uniform size. Alternatively, less than about 50% of the droplets generated can have uniform size.

[00319] While **FIGs. 7A** and **7B** illustrate the height difference, Δh , being abrupt at the junction **706** (e.g., a step increase), the height difference may increase gradually (e.g., from about 0 μm to a maximum height difference). Alternatively, the height difference may decrease gradually (e.g., taper) from a maximum height difference. A gradual increase or decrease in height difference, as used herein, may refer to a continuous incremental increase or decrease in height difference, wherein an angle between any one differential segment of a height profile and an immediately adjacent differential segment of the height profile is greater than 90° . For example, at the junction **706**, a bottom wall of the channel and a bottom wall of the reservoir can meet at an angle greater than 90° . Alternatively or in addition, a top wall (e.g., ceiling) of the channel and a top wall (e.g., ceiling) of the reservoir can meet an angle greater than 90° . A gradual increase or decrease may be linear or non-linear (e.g., exponential, sinusoidal, etc.). Alternatively or in addition, the height difference may variably increase and/or decrease linearly or non-linearly. While **FIGs. 7A** and **7B** illustrate the expanding reservoir cross-section height as linear (e.g., constant expansion angle, β), the cross-section height may expand non-linearly. For example, the reservoir may be defined at least partially by a dome-like (e.g., hemispherical) shape having variable expansion angles. The cross-section height may expand in any shape.

[00320] The channel networks, e.g., as described above or elsewhere herein, can be fluidly coupled to appropriate fluidic components. For example, the inlet channel segments are fluidly coupled to appropriate sources of the materials they are to deliver to a channel junction. These sources may include any of a variety of different fluidic components, from simple reservoirs defined in or connected to a body structure of a microfluidic device, to fluid conduits that deliver fluids from off-device sources, manifolds, fluid flow units (e.g., actuators, pumps, compressors)

or the like. Likewise, the outlet channel segment (e.g., channel segment **208**, reservoir **604**, etc.) may be fluidly coupled to a receiving vessel or conduit for the partitioned cells for subsequent processing. Again, this may be a reservoir defined in the body of a microfluidic device, or it may be a fluidic conduit for delivering the partitioned cells to a subsequent process operation, instrument or component.

[00321] The methods and systems described herein may be used to greatly increase the efficiency of single cell applications and/or other applications receiving droplet-based input. For example, following the sorting of occupied cells and/or appropriately-sized cells, subsequent operations that can be performed can include generation of amplification products, purification (e.g., via solid phase reversible immobilization (SPRI)), further processing (e.g., shearing, ligation of functional sequences, and subsequent amplification (e.g., via PCR)). These operations may occur in bulk (e.g., outside the partition). In the case where a partition is a droplet in an emulsion, the emulsion can be broken and the contents of the droplet pooled for additional operations. Additional reagents that may be co-partitioned along with the barcode bearing bead may include oligonucleotides to block ribosomal RNA (rRNA) and nucleases to digest genomic DNA from cells. Alternatively, rRNA removal agents may be applied during additional processing operations. The configuration of the constructs generated by such a method can help minimize (or avoid) sequencing of the poly-T sequence during sequencing and/or sequence the 5' end of a polynucleotide sequence. The amplification products, for example, first amplification products and/or second amplification products, may be subject to sequencing for sequence analysis. In some cases, amplification may be performed using the Partial Hairpin Amplification for Sequencing (PHASE) method.

[00322] A variety of applications require the evaluation of the presence and quantification of different biological particle or organism types within a population of biological particles, including, for example, microbiome analysis and characterization, environmental testing, food safety testing, epidemiological analysis, e.g., in tracing contamination or the like.

Computer systems

[00323] The present disclosure provides computer systems that are programmed to implement methods of the disclosure. **FIG. 25** shows a computer system **2501** that is programmed or otherwise configured to, for example, (i) control a microfluidics system (e.g., fluid flow), (ii) sort occupied droplets from unoccupied droplets, (iii) polymerize droplets, (iv) perform sequencing applications, or (v) generate and maintain a library of barcode molecules. The computer system **2501** can regulate various aspects of the present disclosure, such as, for example, fluid flow rates in one or more channels in a microfluidic structure, polymerization application units, etc.. The computer system **2501** can be an electronic device of a user or a computer system that is

remotely located with respect to the electronic device. The electronic device can be a mobile electronic device.

[00324] The computer system **2501** includes a central processing unit (CPU, also “processor” and “computer processor” herein) **2505**, which can be a single core or multi core processor, or a plurality of processors for parallel processing. The computer system **2501** also includes memory or memory location **2510** (e.g., random-access memory, read-only memory, flash memory), electronic storage unit **2515** (e.g., hard disk), communication interface **2520** (e.g., network adapter) for communicating with one or more other systems, and peripheral devices **2525**, such as cache, other memory, data storage and/or electronic display adapters. The memory **2510**, storage unit **2515**, interface **2520** and peripheral devices **2525** are in communication with the CPU **2505** through a communication bus (solid lines), such as a motherboard. The storage unit **2515** can be a data storage unit (or data repository) for storing data. The computer system **2501** can be operatively coupled to a computer network (“network”) **2530** with the aid of the communication interface **2520**. The network **2530** can be the Internet, an internet and/or extranet, or an intranet and/or extranet that is in communication with the Internet. The network **2530** in some cases is a telecommunication and/or data network. The network **2530** can include one or more computer servers, which can enable distributed computing, such as cloud computing. The network **2530**, in some cases with the aid of the computer system **2501**, can implement a peer-to-peer network, which may enable devices coupled to the computer system **2501** to behave as a client or a server.

[00325] The CPU **2505** can execute a sequence of machine-readable instructions, which can be embodied in a program or software. The instructions may be stored in a memory location, such as the memory **2510**. The instructions can be directed to the CPU **2505**, which can subsequently program or otherwise configure the CPU **2505** to implement methods of the present disclosure. Examples of operations performed by the CPU **2505** can include fetch, decode, execute, and writeback.

[00326] The CPU **2505** can be part of a circuit, such as an integrated circuit. One or more other components of the system **2501** can be included in the circuit. In some cases, the circuit is an application specific integrated circuit (ASIC).

[00327] The storage unit **2515** can store files, such as drivers, libraries and saved programs. The storage unit **2515** can store user data, e.g., user preferences and user programs. The computer system **2501** in some cases can include one or more additional data storage units that are external to the computer system **2501**, such as located on a remote server that is in communication with the computer system **2501** through an intranet or the Internet.

[00328] The computer system **2501** can communicate with one or more remote computer systems through the network **2530**. For instance, the computer system **2501** can communicate with a remote computer system of a user (e.g., operator). Examples of remote computer systems include personal computers (e.g., portable PC), slate or tablet PC's (e.g., Apple® iPad, Samsung® Galaxy Tab), telephones, Smart phones (e.g., Apple® iPhone, Android-enabled device, Blackberry®), or personal digital assistants. The user can access the computer system **2501** via the network **2530**.

[00329] Methods as described herein can be implemented by way of machine (e.g., computer processor) executable code stored on an electronic storage location of the computer system **2501**, such as, for example, on the memory **2510** or electronic storage unit **2515**. The machine executable or machine readable code can be provided in the form of software. During use, the code can be executed by the processor **2505**. In some cases, the code can be retrieved from the storage unit **2515** and stored on the memory **2510** for ready access by the processor **2505**. In some situations, the electronic storage unit **2515** can be precluded, and machine-executable instructions are stored on memory **2510**.

[00330] The code can be pre-compiled and configured for use with a machine having a processor adapted to execute the code, or can be compiled during runtime. The code can be supplied in a programming language that can be selected to enable the code to execute in a pre-compiled or as-compiled fashion.

[00331] Aspects of the systems and methods provided herein, such as the computer system **2501**, can be embodied in programming. Various aspects of the technology may be thought of as “products” or “articles of manufacture” typically in the form of machine (or processor) executable code and/or associated data that is carried on or embodied in a type of machine readable medium. Machine-executable code can be stored on an electronic storage unit, such as memory (e.g., read-only memory, random-access memory, flash memory) or a hard disk. “Storage” type media can include any or all of the tangible memory of the computers, processors or the like, or associated modules thereof, such as various semiconductor memories, tape drives, disk drives and the like, which may provide non-transitory storage at any time for the software programming. All or portions of the software may at times be communicated through the Internet or various other telecommunication networks. Such communications, for example, may enable loading of the software from one computer or processor into another, for example, from a management server or host computer into the computer platform of an application server. Thus, another type of media that may bear the software elements includes optical, electrical and electromagnetic waves, such as used across physical interfaces between local devices, through wired and optical landline networks and over various air-links. The physical elements that carry

such waves, such as wired or wireless links, optical links or the like, also may be considered as media bearing the software. As used herein, unless restricted to non-transitory, tangible “storage” media, terms such as computer or machine “readable medium” refer to any medium that participates in providing instructions to a processor for execution.

[00332] Hence, a machine readable medium, such as computer-executable code, may take many forms, including but not limited to, a tangible storage medium, a carrier wave medium or physical transmission medium. Non-volatile storage media include, for example, optical or magnetic disks, such as any of the storage devices in any computer(s) or the like, such as may be used to implement the databases, etc. shown in the drawings. Volatile storage media include dynamic memory, such as main memory of such a computer platform. Tangible transmission media include coaxial cables; copper wire and fiber optics, including the wires that comprise a bus within a computer system. Carrier-wave transmission media may take the form of electric or electromagnetic signals, or acoustic or light waves such as those generated during radio frequency (RF) and infrared (IR) data communications. Common forms of computer-readable media therefore include for example: a floppy disk, a flexible disk, hard disk, magnetic tape, any other magnetic medium, a CD-ROM, DVD or DVD-ROM, any other optical medium, punch cards paper tape, any other physical storage medium with patterns of holes, a RAM, a ROM, a PROM and EPROM, a FLASH-EPROM, any other memory chip or cartridge, a carrier wave transporting data or instructions, cables or links transporting such a carrier wave, or any other medium from which a computer may read programming code and/or data. Many of these forms of computer readable media may be involved in carrying one or more sequences of one or more instructions to a processor for execution.

[00333] The computer system **2501** can include or be in communication with an electronic display **2535** that comprises a user interface (UI) **2540** for providing, for example, results of sequencing analysis, etc.. Examples of UIs include, without limitation, a graphical user interface (GUI) and web-based user interface.

[00334] Methods and systems of the present disclosure can be implemented by way of one or more algorithms. An algorithm can be implemented by way of software upon execution by the central processing unit **2505**. The algorithm can, for example, perform sequencing, etc.

[00335] Devices, systems, compositions and methods of the present disclosure may be used for various applications, such as, for example, processing a single analyte (e.g., RNA, DNA, or protein) or multiple analytes (e.g., DNA and RNA, DNA and protein, RNA and protein, or RNA, DNA and protein) from a single cell. For example, a biological particle (e.g., a cell or cell bead) is partitioned in a partition (e.g., droplet), and multiple analytes from the biological particle are processed for subsequent processing. The multiple analytes may be from the single cell. This

may enable, for example, simultaneous proteomic, transcriptomic and genomic analysis of the cell.

[00336] While preferred embodiments of the present invention have been shown and described herein, it will be obvious to those skilled in the art that such embodiments are provided by way of example only. It is not intended that the invention be limited by the specific examples provided within the specification. While the invention has been described with reference to the aforementioned specification, the descriptions and illustrations of the embodiments herein are not meant to be construed in a limiting sense. Numerous variations, changes, and substitutions will now occur to those skilled in the art without departing from the invention. Furthermore, it shall be understood that all aspects of the invention are not limited to the specific depictions, configurations or relative proportions set forth herein which depend upon a variety of conditions and variables. It should be understood that various alternatives to the embodiments of the invention described herein may be employed in practicing the invention. It is therefore contemplated that the invention shall also cover any such alternatives, modifications, variations or equivalents. It is intended that the following claims define the scope of the invention and that methods and structures within the scope of these claims and their equivalents be covered thereby.

CLAIMS

WHAT IS CLAIMED IS:

1. A method for generating a plurality of barcode molecules, comprising:
 - (a) providing a plurality of molecules coupled to a plurality of supports; and
 - (b) combinatorially assembling said plurality of barcode molecules coupled to said plurality of supports by coupling one or more molecules each comprising one or more segments to each of said plurality of molecules,wherein said plurality of barcode molecules comprises (i) a first set of barcode molecules coupled to a support of said plurality of supports and (ii) a second set of barcode molecules coupled to said support, wherein barcode molecules of said first set of barcode molecules are different than barcode molecules of said second set of barcode molecules.
2. The method of claim 1, wherein said barcode molecules of said first set of barcode molecules and said barcode molecules of said second set of barcode molecules comprise barcode sequences that are different from barcode sequences of barcode molecules coupled to other supports of said plurality of supports.
3. The method of claim 1, wherein barcode sequences of said barcode molecules of said first set of barcode molecules and said barcode molecules of said second set of barcode molecules are identical.
4. The method of claim 1, wherein said plurality of supports is a plurality of beads, and wherein said barcode molecules of said first set of barcode molecules and said barcode molecules of said second set of barcode molecules are coupled to a bead of said plurality of beads.
5. The method of claim 4, wherein at least a subset of said barcode molecules of said first set of barcode molecules and/or said barcode molecules of said second set of barcode molecules is coupled to an interior of said bead.
6. The method of claim 4, wherein said barcode molecules of said first set of barcode molecules and said barcode molecules of said second set of barcode molecules are releasably coupled to said bead.
7. The method of claim 6, wherein said barcode molecules of said first set of barcode molecules and said barcode molecules of said second set of barcode molecules are releasably coupled to said bead through chemical cross-linkers.
8. The method of claim 4, wherein said plurality of beads is a plurality of gel beads.
9. The method of claim 4, wherein said plurality of beads is dissolvable or disruptable.

10. The method of claim 1, wherein (b) comprises sequentially coupling multiple molecules to each of said plurality of molecules.
11. The method of claim 1, wherein in (b) said one or more molecules comprise a first molecule comprising a first segment and a second molecule comprising a second segment, and wherein said combinatorially assembling comprises coupling said first molecule to a molecule of said plurality of molecules and coupling said second molecule to said first molecule, wherein said molecule is coupled to said support.
12. The method of claim 11, further comprising coupling a third molecule comprising a third segment to said second molecule.
13. The method of claim 12, wherein said third segment is different than said first segment or said second segment.
14. The method of claim 12, wherein said third segment is different than said first segment and said second segment.
15. The method of claim 11, wherein said first segment of said first molecule is different than said second segment of said second molecule.
16. The method of claim 11, wherein said first molecule comprises a double-stranded region comprising said first segment and a single-stranded region.
17. The method of claim 16, wherein said single-stranded region of said first molecule has six or fewer nucleotides.
18. The method of claim 17, wherein said single-stranded region of said first molecule has two nucleotides.
19. The method of claim 18, wherein the sequence of said single-stranded region of said first molecule is selected from the group consisting of CA, GT, AC, and TG.
20. The method of claim 16, wherein said first molecule further comprises an additional single-stranded region configured to couple to said molecule of said plurality of molecules.
21. The method of claim 20, wherein said single-stranded region and said additional single-stranded region of said first molecule are included in the same strand of said first molecule.
22. The method of claim 16, wherein said second molecule comprises a double-stranded region comprising said second segment and a single-stranded region.
23. The method of claim 22, wherein said single-stranded region of said second molecule comprises a sequence complementary to the sequence of said single-stranded region of said first molecule.

38. The method of claim 28, wherein said one or more molecules comprise a plurality of first molecules comprising said first segment, a plurality of second molecules comprising said second segment, a plurality of third molecules comprising said third segment, and a plurality of fourth molecules comprising said fourth segment, and wherein said combinatorially assembling comprises (i) coupling first molecules of said plurality of first molecules to molecules of said plurality of molecules and coupling second molecules of said plurality of second molecules to said first molecules, and (ii) coupling third molecules of said plurality of third molecules to additional molecules of said plurality of molecules and coupling fourth molecules of said plurality of fourth molecules to said third molecules, wherein said molecules and said additional molecules of said plurality of molecules are coupled to said support.
39. The method of claim 38, wherein, among said one or molecules, said plurality of first molecules comprises a greater number of molecules than said plurality of third molecules.
40. The method of claim 38, wherein, among said one or more molecules, said plurality of second molecules comprises a greater number of molecules than said plurality of fourth molecules.
41. The method of claim 38, wherein said molecules of said plurality of molecules comprises a greater number of molecules than said additional molecules of said plurality of molecules.
42. The method of claim 1, wherein (b) comprises ligating said one or more molecules to each of said plurality of molecules coupled to said plurality of supports.
43. The method of claim 1, wherein (b) comprises partitioning said plurality of molecules coupled to said plurality of supports in separate partitions, and coupling at least a subset of said one or more molecules to molecules of said plurality of molecules in said separate partitions.
44. The method of claim 43, wherein (b) further comprises partitioning said plurality of molecules coupled to said plurality of supports in one or more additional partitions, and coupling at least a subset of said one or more molecules to molecules of said plurality of molecules in said one or more additional partitions.
45. The method of claim 43, wherein said separate partitions are wells.
46. The method of claim 43, wherein said separate partitions are droplets.
47. The method of claim 2, wherein said barcode sequences are nucleic acid sequences.
48. The method of claim 2, wherein said barcode sequences are amino acid sequences.

49. The method of claim 1, wherein said one or more segments are at least one nucleotide in length.
50. The method of claim 1, wherein each of said barcode molecules of said first set of barcode molecules and said barcode molecules of said second set of barcode molecules comprises a unique identifier that is different than unique identifiers of other barcode molecules coupled to said support.
51. The method of claim 1, wherein said barcode molecules of said first set of barcode molecules and said barcode molecules of said second set of barcode molecules comprise at least 100,000 barcode molecules.
52. The method of claim 1, wherein each of said barcode molecules of said first set of barcode molecules and said barcode molecules of said second set of barcode molecules comprise a segment separated from said one or more segments by a sequence that is the same across at least a subset of said plurality of barcode molecules.
53. The method of claim 1, wherein said one or more segments are different for at least a subset of said plurality of barcode molecules.
54. The method of claim 1, wherein barcode molecules of said first set of barcode molecules include a first sequence for use with a first assay and barcode molecules of said second set of barcode molecules include a second sequence for use with a second assay, wherein said first assay is different than said second assay.
55. The method of claim 1, wherein barcode molecules of said first set of barcode molecules and barcode molecules of said second set of barcode molecules include both identical barcode sequences and different barcode sequences.
56. The method of claim 1, wherein barcode molecules of said first set of barcode molecules include first functional sequences and barcode molecules of said second set of barcode molecules include second functional sequences, wherein said first functional sequences and said second functional sequences are different.
57. The method of claim 56, wherein said first set of barcode molecules coupled to said support comprises a greater number of barcode molecules than said second set of barcode molecules coupled to said support.
58. The method of claim 56, wherein said first set of barcode molecules coupled to said support comprises at least 10,000 barcode molecules.
59. The method of claim 56, wherein said second set of barcode molecules coupled to said support comprises at least 10,000 barcode molecules.
60. The method of claim 2, wherein barcode molecules of said other supports of said plurality of supports comprise different barcode sequences.

61. The method of claim 60, wherein each other support of said other supports of said plurality of supports comprises barcode molecules comprising a different barcode sequence.
62. The method of claim 11, further comprising contacting a solution comprising said first molecule and said second molecule coupled to said molecule of said plurality of molecules with an exonuclease.
63. The method of claim 62, wherein said second molecule comprises a phosphorothioate moiety.
64. A method of processing a plurality of barcode molecules, comprising:
 - (a) providing a plurality of molecules coupled to a plurality of supports;
 - (b) subjecting a subset of molecules of said plurality of molecules to conditions sufficient to couple first molecules to said subset of molecules, wherein said subset of molecules are coupled to a support of said plurality of supports;
 - (c) subjecting said first molecules coupled to said subset of molecules to conditions sufficient to couple second molecules to said first molecules, thereby generating said plurality of barcode molecules and by-products, which plurality of barcode molecules comprise said second molecules coupled to said first molecules; and
 - (d) contacting said products and by-products with a molecule capable of degrading said by-products.
65. The method of claim 64, wherein said molecule capable of degrading said by-products is an exonuclease.
66. The method of claim 64, wherein said second molecules comprise phosphorothioate moieties.
67. The method of claim 66, wherein said phosphorothioate moieties are disposed at an end of said second molecules.
68. The method of claim 64, wherein said plurality of barcode molecules comprises a plurality of barcode sequences.
69. The method of claim 68, wherein said plurality of barcode sequences of said plurality of barcode molecules coupled to said support are identical.
70. The method of claim 64, wherein said plurality of supports comprises a plurality of beads, and wherein said plurality of barcode molecules are coupled to a bead of said plurality of beads.
71. The method of claim 70, wherein said plurality of beads is a plurality of gel beads.
72. The method of claim 70, wherein said plurality of beads is dissolvable or disruptable.

73. The method of claim 70, wherein said plurality of barcode molecules is releasably coupled to said bead.
74. The method of claim 64, wherein (b) comprises ligating said first molecules to said molecules.
75. The method of claim 64, wherein (c) comprises ligating said second molecules to said first molecules.
76. The method of claim 64, further comprising partitioning said plurality of molecules coupled to said plurality of supports in separate partitions.
77. The method of claim 76, wherein (b) comprises coupling said first molecules to said subset of molecules within a partition of said separate partitions.
78. The method of claim 77, further comprising, subsequent to (b), partitioning said plurality of molecules coupled to said plurality of supports in one or more additional partitions.
79. The method of claim 78, wherein (c) comprises coupling said second molecules to said first molecules within a partition of said one or more additional partitions.
80. The method of claim 76, wherein said separate partitions are wells.
81. The method of claim 76, wherein said separate partitions are droplets.
82. The method of claim 76, wherein prior to (d), said plurality of barcode molecules coupled to said support and said by-products are recovered from said separate partitions.
83. The method of claim 68, wherein said plurality of barcode sequences comprises a plurality of nucleic acid barcode sequences.
84. The method of claim 64, wherein said first molecules and said second molecules are nucleic acid molecules, which nucleic acid molecules each comprise at least two nucleotides.
85. The method of claim 64, wherein said plurality of barcode molecules coupled to said support comprises at least 10,000 barcode molecules.
86. The method of claim 64, wherein said plurality of barcode molecules coupled to said support comprises a plurality of first barcode molecules and a plurality of second barcode molecules, which plurality of first barcode molecules and plurality of second barcode molecules are different.
87. The method of claim 86, wherein said plurality of first barcode molecules comprise a plurality of first barcode sequences and said plurality of second barcode molecules comprise a plurality of second barcode sequences, which first barcode sequences are different than said second barcode sequences.
88. The method of claim 86, wherein said plurality of first barcode molecules comprise first functional sequences and said plurality of second barcode molecules comprise second

- functional sequences, which first functional sequences are different than said second functional sequences.
89. The method of claim 64, wherein said plurality of barcode molecules comprise a plurality of functional sequences.
90. The method of claim 89, wherein said functional sequences are poly(T) sequences.
91. A composition comprising a support comprising a plurality of nucleic acid barcode molecules coupled thereto, which plurality of nucleic acid barcode molecules comprise one or more sequences selected from the sequences included in Table 1 or Table 2.
92. The composition of claim 91, wherein said plurality of nucleic acid barcode molecules comprise one or more sequences from the sequences included in Table 1.
93. The composition of claim 91, wherein said plurality of nucleic acid barcode molecules comprise one or more sequences from the sequences included in Table 2.
94. The composition of claim 93, wherein said plurality of nucleic acid barcode molecules comprise the sequence CCTTAGCCGCTAATAGGTGAGC.
95. The composition of claim 93, wherein said plurality of nucleic acid barcode molecules comprise the sequence TTGCTAGGACCGGCCTTAAAGC.
96. The composition of claim 91, wherein each nucleic acid barcode molecule of the plurality of nucleic acid barcode molecules comprises one or more sequences selected from the sequences included in Table 1 or Table 2.
97. The composition of claim 91, wherein said plurality of nucleic acid barcode molecules comprise a first plurality of nucleic acid barcode molecules and a second plurality of nucleic acid barcode molecules, which first plurality of nucleic acid barcode molecules comprise one or more sequences selected from the sequences included in Table 1 or Table 2 and which second plurality of nucleic acid barcode molecules do not comprise a sequence included in Table 1 or Table 2.
98. The composition of claim 97, wherein said second plurality of nucleic acid barcode molecules are configured to interact with ribonucleic acid molecules.
99. The composition of claim 98, wherein said second plurality of nucleic acid barcode molecules comprise a poly(T) sequence.
100. The composition of claim 97, wherein said first plurality of nucleic acid barcode molecules are configured to interact with deoxyribonucleic acid molecules.
101. The composition of claim 91, wherein said support is a bead.
102. The composition of claim 101, wherein said bead is a gel bead.
103. The composition of claim 101, wherein said bead is dissolvable or disruptable.

104. The composition of claim 101, wherein said plurality of nucleic acid barcode molecules is releasably coupled to said bead.
105. The composition of claim 91, wherein said plurality of nucleic acid barcode molecules coupled to said support comprise a plurality of barcode sequences.
106. The composition of claim 105, wherein said plurality of barcode sequences is the same for each of said plurality of nucleic acid barcode molecules.
107. The composition of claim 105, wherein said plurality of barcode sequences is a plurality of nucleic acid barcode sequences.
108. The composition of claim 91, wherein said plurality of nucleic acid barcode molecules comprises at least 10,000 nucleic acid barcode molecules.
109. The composition of claim 97, wherein said first plurality of nucleic acid barcode molecules comprises at least 10,000 nucleic acid barcode molecules.
110. The composition of claim 97, wherein said second plurality of nucleic acid barcode molecules comprises at least 10,000 nucleic acid barcode molecules.
111. The composition of claim 97, wherein, among said plurality of nucleic acid barcode molecules, said first plurality of nucleic acid barcode molecules comprises a greater number of nucleic acid barcode molecules than said second plurality of nucleic acid barcode molecules.
112. The composition of claim 97, wherein, among said plurality of nucleic acid barcode molecules, said second plurality of nucleic acid barcode molecules comprises a greater number of nucleic acid barcode molecules than said first plurality of nucleic acid barcode molecules.