9805471-1

IPOS Copy

PCT WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : A61K 31/505, C07D 471/04 — A1

(11) International Publication Number: **WO 97/44040**

(43) International Publication Date: 27 November 1997 (27.11.97)

(21) International Application Number: PCT/EP97/02710

(22) International Filing Date: 15 May 1997 (15.05.97)

(30) Priority Data:
96201450.2 23 May 1996 (23.05.96) EP
(34) Countries for which the regional or international application was filed: DE et al.

(71) Applicant *(for all designated States except US)*: JANSSEN PHARMACEUTICA N.V. [BE/BE]; Turnhoutseweg 30, B-2340 Beerse (BE).

(72) Inventors; and

(75) Inventors/Applicants *(for US only)*: KENNIS, Ludo, Edmond, Josephine [BE/BE]; Janssen Pharmaceutica N.V., Turnhoutseweg 30, B-2340 Beerse (BE). MERTENS, Josephus, Carolus [BE/BE]; Janssen Pharmaceutica N.V., Turnhoutseweg 30, B-2340 Beerse (BE).

(74) Agent: QUAGHEBEUR, Luc; Janssen Pharmaceutica N.V., Patent Dept., Turnhoutseweg 30, B-2340 Beerse (BE).


(81) Designated States: AL, AM, AU, BA, BB, BG, BR, CA, CN, CU, CZ, EE, GE, HU, IL, IS, JP, KG, KR, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, TR, TT, UA, US, UZ, VN, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).

Published
With international search report.
Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(54) Title: HEXAHYDRO-PYRIDO(4,3-B)INDOLE DERIVATIVES AS ANTIPSYCHOTIC DRUGS

(57) Abstract

This invention concerns the compounds of formula (I), the pharmaceutically acceptable addition salts and the stereochemically isomeric forms thereof, wherein Alk is C_{1-6}alkanediyl; R^1 is hydrogen, C_{1-6}alkyl, aryl or arylC_{1-6}alkyl; R^2, R^3 and R^4 are each independently selected from hydrogen, halo, hydroxy, nitro, cyano, C_{1-6}alkyl, C_{1-6}alkyloxy, trifluoromethyl, C_{1-6}alkylthio, mercapto, amino, mono- and di(C_{1-6}alkyl)amino, carboxyl, C_{1-6}alkyloxycarbonyl or C_{1-6}alkylcarbonyl; R^5 is hydrogen, C_{1-6}alkyl, phenyl or phenylC_{1-6}alkyl; R^6 is hydrogen, C_{1-6}alkyl, C_{1-6}alkyloxy, C_{1-6}alkylthio, or a radical of formula $-NR^8R^9$, wherein R^8 and R^9 are each independently selected from hydrogen, C_{1-6}alkyl, phenyl or phenylC_{1-6}alkyl; R^7 is hydrogen or C_{1-6}alkyl; or R^6 and R^7 taken together may form a bivalent radical of formula $-R^6-R^7-$; having central dopamine and serotonin antagonistic activity; their preparation, compositions containing them and their use as a medecine.



(I)