

Description

This invention relates to a novel antimicrobial composition which contains an isothiazolone complex or isothiazolone derivative complex of general formula (I) and 3,3,4,4-tetrachlorotetrahydrothiophene 1,1-dioxide (hereinafter sometimes referred to as "compound (II)") and in which synergism between both the active ingredients is made the best of. This antimicrobial composition is suited particularly for use in industrial water systems in pulp and paper manufacture.

In wastewater from the papermaking process in the field of paper and pulp industry and in various water systems in various fields of industry, such as in circulating cooling water, and further in water-based paints, sizing liquids for paper, latices, printing pastes, leathers and like products manufactured by using industrial water, harmful microorganisms can grow with ease to cause decrease in productivity and/or quality of products. In water systems in the paper and pulp industry, in particular, the growth and proliferation of bacteria, filamentous microorganisms and yeasts lead to the formation of slime. In watercourses in which the pulp slurry flows, in particular at places having a rough wall surface which comes into contact with the slurry and at places where the pulp slurry decreases in flow rate and becomes stagnant, for example in the chest, flow box and transportation pipe, slime formation and adherence take place. This slime often detaches and causes paper breakage and/or contamination of paper and pulp products. The growth of microorganisms in slime also causes various other troubles.

Where the papermaking machines are high-speed ones, these troubles present very serious problems, provoking marked decrease in productivity and significant economic losses.


Also in circulating cooling water systems for metal cutting oils, microorganisms can grow and proliferate to thereby affect the cooling power and/or emulsifiability of the cutting oils adversely or cause phenomena unfavorable from the public health viewpoint through bad odor emission and/or impairment of the working environment. Troubles caused by the growth of harmful microorganisms are also encountered in industrial products such as water-based paints, sizing liquids for paper, polymer latices, pulp for paper-making, pastes, leathers, and metal cutting oils.

Those agents which have been so far in use for the inhibition or prevention of growth of harmful microorganisms in the above-mentioned water systems or in the above-mentioned industrial products include, among others, organometallic compounds, organochlorine compounds, organosulfur compounds and quaternary ammonium compounds. These compounds, however, are toxic to humans, have unpleasant or offensive odor and, furthermore, may cause unfavorable phenomena such as foaming. In addition, water systems containing these agents, when discharged into ordinary rivers or watercourses or into the sea, may adversely affect aquatic animals and thus pose problems from the environmental protection viewpoint.

The present inventors conducted intensive investigations in an attempt to eliminate the above-mentioned drawbacks and, as a result, have now completed the present invention.

Thus, this invention provides an antimicrobial composition specified below:

An antimicrobial composition which contains, as combined active ingredients, an isothiazolone complex and or isothiazolone derivative complex of the general formula



wherein R¹ is a hydrogen atom or a substituted or unsubstituted monovalent hydrocarbyl group, R² and R³ are each independently a hydrogen atom or a halogen atom, M is a metal or ammonium cation, X is a complex-forming anion, a is 1 or 2 and b is an integer which is such that the anion X satisfies the valence of the cation M, and 3,3,4,4-tetrachlorotetrahydrothiophene 1,1-dioxide.

Some of the complexes of general formula (I) are the complexes described in United States Patent No. 4150026 whereas the compound (II) is also a bactericide described in United States Patent No. 2,957,887. The complexes of general formula (I) and the compound (II) can be produced in a manner analogous to the known method.

These bactericides separately are still unsatisfactory as antimicrobial agents, for example in that they are effective against only limited kinds of microorganism, hence they produce insufficient bactericidal effect, and that they fail to retain their effect for a prolonged period of time.

The antimicrobial composition according to the invention can produce a very high antimicrobial effect which could have never been expected from the experience with each individual constituent active ingre-

dient. Furthermore, it has a broad range of applications against harmful microorganisms, namely filamentous bacteria, other bacteria, yeasts, fungi, and so forth, irrespective of species thereof.

As for the complexes of the general formula (I), metal cations represented by M and complex-forming anions represented by X include respectively the corresponding metal cations and anions described in the above-mentioned U.S.P. No. 4150026, namely metal cations such as barium, cadmium, calcium, chromium, cobalt, copper, iron, lead, lithium, manganese, mercury, nickel, sodium, silver, strontium, tin, zinc or the like and anions such as chloride, bromide, iodide, sulfate, nitrate, acetate, perchlorate, bisulfate, bicarbonate, oxalate, maleate, p-toluene-sulfonate, carbonate, phosphate, and the like. As more preferred examples of M, there are mentioned cations of metals of valence of 2.

As the complex of general formula (I) which is to be used in accordance with the invention, there may be included, for example, compounds wherein R¹ is a C₁₋₁₀ alkyl group, R² and R³ are each independently a hydrogen atom or a halogen atom and M is zinc, calcium, copper or magnesium, and X is a mineral acid anion such as chloride, bromide, nitrate or sulfate. Preferably, there may be mentioned, for example, 5-chloro-2-methyl-3-isothiazolone-magnesium nitrate, 2-methyl-3-isothiazolone-magnesium nitrate, and 2-octyl-5-chloro-3-isothiazolone-calcium chloride and 2-octyl-4-chloro-3-isothiazolone-calcium chloride.

Such compounds of general formula (I) may be used either singly or in admixture of two or more of them. The compound of general formula (I) and the compound (II) are used generally in a weight ratio of 1/10 to 10/1, preferably 1/5 to 5/1.

A satisfactory effect as seen from the practical viewpoint can be achieved in the above ratio.

Basically, the antimicrobial composition according to the invention is prepared by admixing the above-mentioned two ingredients to a homogeneous composition. Generally, however, the composition is submitted to use in the form of a solution in solvents or an emulsion or dispersion, for instance.

As the solvent usable in that occasion, there may be mentioned alcohol type solvents (e.g. methyl alcohol, ethyl alcohol, ethylene glycol, diethylene glycol), ketone type solvents (e.g. acetone, methylethyl ketone), ether type solvents (e.g. dioxane, tetrahydrofuran) and hydrocarbon type solvents (e.g. benzene, toluene), among others. The antimicrobial composition according to the present invention may also be used in a carrier-borne form. The liquid carriers include, for example, the above-mentioned alcohols, ketones and hydrocarbons. As the solid carrier, for example, there may be mentioned clays (e.g. Kaolin, bentonite), talcs and silicas (e.g. diatomaceous soil). Thus the mode of application is not critical but various modes of application can be employed. The solvents and carriers may be conventional.

While the dose of the antimicrobial composition may vary depending on the concentration of the microorganisms, the composition is used generally in a dose of 0.01 to 100 ppm based on the amount of the composition in the case of water systems in the field of paper and pulp industry and the like, and 1 to 500 ppm based on the amount of the composition in the fields of water-based paints, pastes, leathers and the like.

The antimicrobial composition according to the invention may justifiably contain conventional additives or auxiliaries, such as stabilizers (e.g. magnesium nitrate, magnesium chloride, calcium chloride, sodium alginate) and surfactants (e.g. alkyl sulfates, polyethyleneglycol alkyl ethers), so long as they do not lead to failure to achieve the objects of the invention. While the amount of such additives and auxiliaries to be incorporated may vary depending on the amount of active ingredients (I) and (II) to be used, the additives and auxiliaries are preferably incorporated in a 0.5 to 10 weight percent based on the composition.

There will be mentioned Examples and Comparative Examples below. The examples are intended to illustrate this invention in further detail and should by no means be construed as limiting the scope of the present invention.

Antimicrobial compositions were prepared according to the following formulations. In the examples, "part(s)" invariably means "part(s)" by weight.

Example 1

Compound (II) 10 parts
3:1 (by weight) Mixture of 5-chloro-2-methyl-3-isothiazolone-magnesium nitrate and 2-methyl-3-isothiazolone-magnesium nitrate 10 parts
Diethylene glycol 78 parts
Sorpel 900A (Toho Chemical) 2 parts

The above-mentioned ingredients were admixed in the specified amount homogeneously to give a composition.

Example 2

An antimicrobial composition was prepared from the ingredients given below in the same manner as in Example 1.

Compound (II) 10 parts
5-Chloro-2-octyl-3-isothiazolone-calcium (II) chloride 10 parts
Diethylene glycol 78 parts
Sorpel 900A 2 parts,

Comparative Example 1

An antimicrobial composition was prepared from the ingredients given below in the same manner as in Example 1.

Compound (II) 20 parts
Diethylene glycol 78 parts
Sorpel 900A 2 parts,

Comparative Example 2

An antimicrobial composition was prepared from the ingredients given below in the same manner as in Example 1.

3:1 (by weight) Mixture of 5-chloro-2-methyl-3-isothiazolone-magnesium (II) nitrate and 2-methyl-3-isothiazolone-magnesium (II) nitrate 20 parts
Diethylene glycol 80 parts

Comparative Example 3

An antimicrobial composition was prepared from the ingredients given below in the same manner as in Example 1.

5-Chloro-2-octyl-3-isothiazolone-calcium (II) chloride 20 parts
Diethylene glycol 78 parts
Sorpel 900A 2 parts

The antimicrobial compositions prepared in the above Examples 1 to 2 and Comparative Examples 1 to 3 were tested for their effects in slime control and preservation.

Test Example 1Inhibition of microbial growth and control of slime formation in the papermaking process.

In the papermaking process in a certain paper mill, the antimicrobial compositions prepared in the above Examples 1 to 2 and comparative Examples 1 to 3 were separately added to a white water pit three times a day, each time for 2 hours, to a concentration in water of 20 ppm based on the amount of the composition. After 7 days of such addition, the number of the microorganisms in white water was counted.

For the counting, a white water sample taken was diluted with sterile water, a known amount of the dilution was placed in a dish, a Waksman agar medium solution was poured into the dish, and the mixture was homogenized and then allowed to solidify in a plate form. After 2 days of cultivation in an incubator (32°C), the microbial colonies formed were counted using a colony counter. The frequency of paper breakage during papermaking process was also recorded for further confirmation of the bactericidal effect. The results of these tests are shown below in Table 1.

Table 1

	Microbial count in white water (cells/ml)	Frequency paper breakage (times)
Example 1	$<10^2$	0
Example 2	$<10^2$	0
Comparative Example 1	2.1×10^4	7
Comparative Example 2	3.4×10^4	5
Comparative Example 3	6.8×10^3	6
No addition of bactericide	$\geq 10^8$	15

The data given in Table 1 indicate that the antimicrobial compositions according to the invention (Examples 1 to 2) have an excellent microbial growth inhibiting activity.

Test Example 2

Inhibition of microbial growth in sizing composition for papermaking

To a starch-based sizing liquid having a pH of 10.0, there were added a bouillon liquid medium and a preliminarily rotten sizing liquid. The mixture was stirred and each of the antimicrobial compositions prepared in the above Examples and Comparative Examples was added respectively to a concentration of 300 ppm based on the amount of the composition.


After 5 days' maintenance of the mixture in an incubator at 32°C, the viable count in each sizing liquid was conducted. The results thus obtained are shown below in Table 2.

Table 2

	Viable microbe cell number (cells/ml)
Example 1	$<10^2$
Example 2	$<10^2$
Comparative Example 1	2.5×10^6
Comparative Example 2	8.0×10^3
Comparative Example 3	7.6×10^3
No addition of bactericide	7.6×10^8

Claims

1. An antimicrobial composition characterized by containing, as combined active ingredients, an isothiazolone compound complex of the general formula



wherein R¹ is a hydrogen atom or a substituted or unsubstituted monovalent hydrocarbyl group, R² and R³ are each independently a hydrogen atom or a halogen atom, M is a metal or ammonium cation, X is a complex-forming anion, a is 1 or 2 and b is an integer which is such that the anion X satisfies the valence of the cation M, and 3,3,4,4-tetrachlorotetrahydrothiophene 1,1-dioxide.

2. An antimicrobial composition as claimed in Claim 1, wherein R¹ is a hydrogen atom or a C₁₋₁₀ alkyl, R² and R³ are each independently a hydrogen atom or a halogen atom, M is a metal of valence of 2 and X is chloride, bromide, nitrate or sulfate.

3. An antimicrobial composition as claimed in Claim 2, wherein the metal of valence of 2 represented by M is zinc, magnesium, calcium or copper.

4. An antimicrobial composition as claimed in Claim 1, wherein the isothiazolone compound complex is 5-chloro-2-methyl-3-isothiazolone-magnesium nitrate.

5. An antimicrobial composition as claimed in Claim 1, wherein the isothiazolone compound complex is 2-methyl-3-isothiazolone-magnesium nitrate.

6. An antimicrobial composition as claimed in Claim 1, wherein as the isothiazolone compound complex, an admixture of 5-chloro-2-methyl-3-isothiazolone-magnesium nitrate and 2-methyl-3-isothiazolone-magnesium nitrate in a weight ratio of 3 to 1 is used.

7. An antimicrobial composition as claimed in Claim 1 wherein the isothiazolone compound complex is 2-octyl-5-chloro-3-isothiazolone-calcium chloride.

8. An antimicrobial composition as claimed in Claim 1, wherein the isothiazolone compound complex and 3,3,4,4-tetrachlorotetrahydrothiophene 1,1-dioxide are contained in a weight ratio of 1/10 to 10/1.



Test Example 2

Inhibition of microbial growth in sizing composition for papermaking

To a starch-based sizing liquid having a pH of 10.0, there were added a bouillon liquid medium and a preliminarily rotten sizing liquid. The mixture was stirred and each of the antimicrobial compositions prepared in the above Examples and Comparative Examples was added respectively to a concentration of 300 ppm based on the amount of the composition.


After 5 days' maintenance of the mixture in an incubator at 32°C, the viable count in each sizing liquid was conducted. The results thus obtained are shown below in Table 2.

Table 2

	Viable microbe cell number (cells/ml)
Example 1	$<10^2$
Example 2	$<10^2$
Comparative Example 1	2.5×10^6
Comparative Example 2	8.0×10^3
Comparative Example 3	7.6×10^3
No addition of bactericide	7.6×10^8

Claims

1. An antimicrobial composition characterized by containing, as combined active ingredients, an isothiazolone compound complex of the general formula



wherein R¹ is a hydrogen atom or a substituted or unsubstituted monovalent hydrocarbyl group, R² and R³ are each independently a hydrogen atom or a halogen atom, M is a metal or ammonium cation, X is a complex-forming anion, a is 1 or 2 and b is an integer which is such that the anion X satisfies the valence of the cation M, and 3,3,4,4-tetrachlorotetrahydrothiophene 1,1-dioxide.

2. An antimicrobial composition as claimed in Claim 1, wherein R¹ is a hydrogen atom or a C₁₋₁₀ alkyl, R² and R³ are each independently a hydrogen atom or a halogen atom, M is a metal of valence of 2 and X is chloride, bromide, nitrate or sulfate.

3. An antimicrobial composition as claimed in Claim 2, wherein the metal of valence of 2 represented by M is zinc, magnesium, calcium or copper.

4. An antimicrobial composition as claimed in Claim 1, wherein the isothiazolone compound complex is 5-chloro-2-methyl-3-isothiazolone-magnesium nitrate.

5. An antimicrobial composition as claimed in Claim 1, wherein the isothiazolone compound complex is 2-methyl-3-isothiazolone-magnesium nitrate.

6. An antimicrobial composition as claimed in Claim 1, wherein as the isothiazolone compound complex, an admixture of 5-chloro-2-methyl-3-isothiazolone-magnesium nitrate and 2-methyl-3-isothiazolone-magnesium nitrate in a weight ratio of 3 to 1 is used.


7. An antimicrobial composition as claimed in Claim 1 wherein the isothiazolone compound complex is 2-octyl-5-chloro-3-isothiazolone-calcium chloride.

8. An antimicrobial composition as claimed in Claim 1, wherein the isothiazolone compound complex and 3,3,4,4-tetrachlorotetrahydrothiophene 1,1-dioxide are contained in a weight ratio of 1/10 to 10/1.



Patentansprüche

1. Antimikrobielle Zusammensetzung, dadurch gekennzeichnet, daß sie als kombinierte Wirkstoffe einen Komplex einer Isothiazolon-Verbindung mit der allgemeinen Formel



in der

R¹ ein Wasserstoff-Atom oder ein substituierter oder unsubstituierter einwertiger Kohlenwasserstoff-Rest ist,

R² und R³ unabhängig voneinander jeweils ein Wasserstoff-Atom oder ein Halogen-Atom sind,

M ein Metall- oder Ammonium-Kation ist,

X ein komplexbildendes Anion ist,

a 1 oder 2 ist und

b eine ganze Zahl derart ist, daß das Anion X die Valenz des Kations M absättigt, und 3,3,4,4-Tetrachlorotetrahydrothiophen-1,1-dioxid enthält.

2. Antimikrobielle Zusammensetzung nach Anspruch 1, worin

R¹ ein Wasserstoff-Atom oder ein C₁₋₁₀-Alkyl ist,

R² und R³ unabhängig voneinander jeweils ein Wasserstoff-Atom oder ein Halogen-Atom sind,

M ein Metall der Valenz 2 ist und

X Chlorid, Bromid, Nitrat oder Sulfat ist.

3. Antimikrobielle Zusammensetzung nach Anspruch 2, worin das durch M bezeichnete Metall der Valenz 2 Zink, Magnesium, Calcium oder Kupfer ist.

4. Antimikrobielle Zusammensetzung nach Anspruch 1, worin der Komplex einer Isothiazolon-Verbindung 5-Chloro-2-methyl-3-isothiazolon-Magnesiumnitrat ist.

5. Antimikrobielle Zusammensetzung nach Anspruch 1, worin der Komplex einer Isothiazolon-Verbindung 2-Methyl-3-isothiazolon-Magnesiumnitrat ist.


6. Antimikrobielle Zusammensetzung nach Anspruch 1, worin als Komplex einer Isothiazolon-Verbindung ein Gemisch aus 5-Chloro-2-methyl-3-isothiazolon-magnesiumnitrat und 2-Methyl-3-isothiazolon-Magnesiumnitrat in einem Gewichts-Verhältnis von 3 zu 1 eingesetzt wird.

7. Antimikrobielle Zusammensetzung nach Anspruch 1, worin der Komplex einer Isothiazolon-Verbindung 2-Octyl-5-chloro-3-isothiazolon-Calciumchlorid ist.

8. Antimikrobielle Zusammensetzung nach Anspruch 1, worin der Komplex einer Isothiazolon-Verbindung und 3,3,4,4-Tetrachlorotetrahydrothiophen-1,1-dioxid in einem Gewichts-Verhältnis von 1/10 bis 10/1 enthalten sind.

Revendications

1. Composition antimicrobienne caractérisée en ce qu'elle contient, en tant qu'ingrédients actifs combinés, un complexe d'un composé isothiazolone de formule générale



dans laquelle R¹ est un atome d'hydrogène ou un groupe hydrocarboné monovalent substitué ou non substitué, R² et R³ représentent chacun de manière indépendante un atome d'hydrogène ou un atome d'halogène, M est un cation d'un métal ou ammonium, X est un anion formant un complexe, a vaut 1 ou 2 et b est un nombre entier qui est tel que l'anion X satisfait la valence du cation M, et le 3,3,4,4-tétrachlorotétrahydrothiophène-1,1-dioxyde.



2. Composition antimicrobienne selon la revendication 1, dans laquelle R¹ est un atome d'hydrogène ou un groupe alkyle en C₁₋₁₀, R² et R³ représentent chacun de manière indépendante un atome d'hydrogène ou un atome d'halogène, M est un métal ayant une valence de 2, et X est chlorure, bromure, nitrate ou sulfate.

5 3. Composition antimicrobienne selon la revendication 2, dans laquelle le métal ayant la valence 2 représenté par M est le zinc, le magnésium, le calcium ou le cuivre.

4. Composition antimicrobienne selon la revendication 1, dans laquelle le complexe du composé isothiazolone est le 5-chloro-2-méthyl-3-isothiazolone-nitrate de magnésium.

10 5. Composition antimicrobienne selon la revendication 1, dans laquelle le complexe du composé isothiazolone est le 2-méthyl-3-isothiazolone-nitrate de magnésium.

6. Composition antimicrobienne selon la revendication 1, dans laquelle on utilise, à titre de complexe du composé isothiazolone, un mélange de 5-chloro-2-méthyl-3-isothiazolone-nitrate de magnésium et de 2-méthyl-3-isothiazolone-nitrate de magnésium, dans un rapport pondéral de 3 à 1.

15 7. Composition antimicrobienne selon la revendication 1, dans laquelle le complexe du composé isothiazolone est le 2-octyl-5-chloro-3-isothiazolone-chlorure de calcium.

8. Composition antimicrobienne selon la revendication 1 qui comprend le complexe du composé isothiazolone et le 3,3,4,4-tétrachlorotétrahydrothiophène-1,1-dioxyde, dans un rapport pondéral allant de 1/10 à 10/1.

20

25

30

35

40

45

50

55

60

65