

MANAGEMENT OF MEDICATION PREPARATION WITH DYNAMIC PROCESSING

CROSS-REFERENCE TO RELATED APPLICATION

5 This application claims priority from U.S. Provisional Application No. 62/077,759 filed
on November 10, 2014, entitled "MANAGEMENT OF MEDICATION PREPARATION WITH
DYNAMIC PROCESSING," the contents of which are incorporated by reference herein as if set
forth in full.

10 BACKGROUND

In many medical facilities, medication orders are provided to a pharmacy for preparation of doses corresponding to the medication orders for administration to a patient. In this regard, orders must be entered, received by the pharmacy, validated, and prepared according to manufacturer's specifications or established institutional guidelines. The preparation process involves the selection and, where required, preparation of drug products for administration to patients in compliance with the dose order. Once fulfilled, the resulting drug products (*i.e.*, doses) must be delivered to the patient for administration. One environment, by way of example, in which such processes occur is a hospital.

There are points in the foregoing process that may be susceptible to the loss or corruption of information due to miscommunication or other errors. This can be problematic in terms of logging and auditing of medication doses, which may be required by insurance and regulatory requirements. In turn, systems for management of pharmacy work flow have been proposed. For example, U.S. Provisional Patent Application No. 61/975,519 entitled "MANAGEMENT OF

MEDICATION DOSE ORDERS" filed on April 4, 2014, which is co-owned with the present application and incorporated by reference herein in its entirety, discloses certain embodiments of a pharmacy workflow management application. Accordingly, at least partially automated pharmacy workflow management applications have been devised to assist in management of the receipt, processing, organization, preparation, verification, and tracking of medication dose orders in the pharmacy or the like. However, further developments would be beneficial to assist in improvements to the pharmacy workflow management application to promote efficiency, reliability, and accuracy related to the preparation and management of medication dose orders.

SUMMARY

In view of the foregoing, the present disclosure is generally related to various embodiments of pharmacy workflow applications. In certain embodiments of the pharmacy workflow management application described herein, improvements in preparing a plurality of dose orders in a common batch according to a batch workflow are presented. Additionally, the present disclosure includes embodiments that relate to a dynamic determination of a final container for use in a dose order preparation protocol of the pharmacy workflow management application. Further still, enhancements to diluent volume accounting using the pharmacy workflow management application are discussed, that, for example, include dynamic diluent volume accounting in a dose order preparation protocol. The present disclosure also contemplates embodiments direct to adjusting the diluent ingredient volume (*e.g.*, dynamically for presentation in a dose preparation protocol). Additionally, improvements with respect to identifying and managing critical issues associated with the pharmacy workflow management application (*e.g.*, an error associated with a dose order record) are discussed.

The present disclosure presents an embodiment of the pharmacy workflow management application directed to preparing an aggregate plurality of dose orders via a batch workflow. In this regard, a plurality of dose orders having one or more identical constituent products associated with different dose order records may be prepared according to a batch workflow.

5 The batch workflow may proceed such that the plurality of dose orders may be prepared without initiating a new workflow process for each dose order. As such, the pharmacy workflow management application may aggregate the plurality of dose orders and display a corresponding dose preparation protocol (*e.g.*, for identifying at least one constituent product) applicable to each of the aggregate plurality of dose orders. Upon receipt of an indication indicative of
10 identification of the at least one constituent product, the pharmacy workflow management application may display one or more preparation steps (*e.g.*, those associated with the physical preparation of the dose) such that the aggregate plurality of dose orders may be prepared only upon the identification of the total volume of the at least one constituent product required to prepare the associated aggregate plurality of doses.

15 The present disclosure also presents an embodiment of the pharmacy workflow management application directed to dynamically determining a final container for use in a dose order preparation protocol. In this regard, the pharmacy workflow application may dynamically determine whether a formulary product indicated as being used in a dose preparation protocol is appropriate as a final container based on a received indication of a plurality of products
20 associated with the dose order. Upon the receipt of the received indication (*e.g.*, indicating the presence of a product as part of a product identification step), the pharmacy workflow management application may execute a series of logical steps in order to dynamically determine whether at least one product of the dose preparation protocol may constitute a selected product

corresponding to a final container. In turn, the pharmacy workflow management application may display a dose preparation protocol at least partially based on the dynamically determined final container (*e.g.*, associated with physical preparation steps, which may be modified based on the container in which the dose is prepared). Notably, if the given dose order analyzed by the

5 dynamically determined final container assessment fails to satisfy the series of logical steps, the pharmacy workflow management application may indicate that the given dose should be prepared in a new or empty container – *i.e.*, not a container of any of the products of the dose order.

The present disclosure presents an embodiment of the pharmacy workflow management

10 application that includes a preparation protocol where the diluent volume may be adjust based on the receipt of an indication of a product associated with a dose preparation protocol. That is, the received indication may correspond to, for example, the physical possession of a product for use in the preparation protocol as part of one or more product identification steps. The pharmacy workflow management application may dynamically determine a required diluent volume based

15 on the particular received indication. In this regard, the pharmacy workflow management application may communicate to a user (*e.g.*, via a display or other user interface), a dose preparation step associated with preparing a dose relative to the dynamically determined required diluent volume. In some embodiments, the foregoing described received indication may correspond to multiple discrete instances of containers of the same diluent product such that the

20 dose order may be prepared from multiple sources of diluent.

The present disclosure presents an embodiment of the pharmacy workflow management application directed to adjusting the diluent ingredient volume associated with the dose preparation protocol. In this regard, the pharmacy workflow application may calculate a QS

(*i.e.*, quantity sufficient) volume for each dose order item according to a series of logical steps based on various characteristics. This may include calculating the QS volume based in part on an accumulated dose volume for each ingredient, a final container determination, and/or a reused dose order status, among other criteria. In some instances, the pharmacy workflow management application may verify the determined QS volume via a QS volume sufficiency check such that the determined QS volume is analyzed with respect to various predetermined quality control measures (*e.g.*, a minimum measured volume, a maximum rounding error percentage, a maximum volume deviation, etc.). Notably, in some instances, based on the determination of the foregoing logical steps, the pharmacy workflow management application may determine that the QS is zero, which may indicate, for example, that a diluent need not be added to the dose in order to complete the preparation steps of the corresponding dose preparation protocol.

The present disclosure presents an embodiment of the pharmacy workflow management application directed to identifying and managing critical issues associated with the pharmacy workflow management application in a common location. In this regard, the pharmacy workflow management application may identify critical issues, which may include various errors such as an error regarding a dose order record, a password error, or other miscellaneous errors, and generate an alert corresponding to the identified error state. Based in part on the alert, the pharmacy workflow application may display the identified critical issues or error states in a common location, such as an attention tab of the pharmacy workflow management application. In some embodiments, the pharmacy workflow management application facilitates resolution of a dose order error within a pharmacy workflow queue. For example, the pharmacy workflow application may receive an input indicative of a resolution of the error state, which in turn may resolve the error by requeueing the dose error upon error resolution validation (*i.e.*, a

determination by the pharmacy workflow management application that the dose order no longer contains an error).

In this regard, a first aspect of the present disclosure includes a method of batch processing of a plurality of dose orders in a pharmacy workflow management application. The method includes accessing, by a processor executing said pharmacy workflow management application, a plurality of dose order records corresponding to dose orders received by the pharmacy workflow management application. The plurality of dose orders may include identical constituent products for use in preparing the plurality of dose orders, which have not been prepared. The method further includes aggregating the plurality of dose orders to create to an aggregated plurality of the dose orders. The method further includes displaying, at a user interface of the pharmacy workflow management application, a dose preparation protocol corresponding to the aggregated plurality of dose order records. The dose preparation protocol may include an aggregate amount of at least one of said identical constituent products needed to prepare each of the aggregated plurality of dose order records. The method may further include receiving, from a user by way of a user input of the pharmacy workflow management application, an indication of the aggregate amount of the at least one constituent product. The method may further include presenting to the user, via the user interface, one or more preparation steps regarding preparation of the aggregated plurality of doses upon receiving the aggregate amount of the at least one constituent product.

A number of feature refinements and additional features are applicable in the first aspect. These feature refinements and additional features may be used individually or in any combination. As such, each of the following features that will be discussed may be, but are not required to be, used with any other feature combination of features of the first aspect.

For example, in an embodiment, the method may further include determining, by the processor, the aggregate plurality of dose orders as suitable for preparation via a batch workflow. Notably, a batch workflow may correspond to preparing the aggregated plurality of dose orders in a common sequence such that a separate dose preparation protocol may not be required for each dose order of the aggregated plurality of dose orders. The method may further include initiating the dose preparation protocol corresponding to the aggregate plurality of dose orders for preparation via the batch workflow. The method may further include printing a dose label for each dose order of said aggregate plurality of dose orders. The foregoing noted indication of the aggregate amount of the at least one constituent product may correspond to one or more formulary products for use in preparation of the aggregate plurality of doses. In this regard, the method may further include printing a work-in-progress label upon receiving the aggregate amount of at least one constituent product such that the work-in-progress label is indicative of a remaining volume of the one or more formulary products after completion of said one or more preparation steps. As such, presenting the dose preparation steps may include presenting the one or more preparation steps free from printing a work-in-progress label for each use of the one or more formulary products such that the work-in-progress label is indicative of a remaining volume of said one or more formulary products after completion of the one or more preparation steps for a given dose of the aggregate plurality of doses.

In an embodiment, the indication of said aggregate amount of said at least one constituent product corresponds to one or more formulary products for use in preparation of said dose order. As such, the one or more formulary products may include a total formulary product volume such that the total formulary product volume equals the aggregate amount of said at least one constituent product. Furthermore, the accessing the plurality of dose order may include

identifying common associations among the plurality of dose order records such that the accessed plurality of dose order records comprise identical constituent products. Notably, the aggregating may include grouping the plurality of dose orders into a queue at least partially based on the identified common associations. In this regard, the method may further include
5 initiating the dose preparation protocol corresponding to the aggregate plurality of dose orders of the queue.

In some embodiments, the aggregate amount may include each of the identical constituent products of the dose order. Additionally, the foregoing noted indication may include a scan.

10 In this regard, a second aspect of the present disclosure includes a method for dynamic determination of a final container for use in a dose order preparation protocol of a pharmacy workflow management application. The method includes accessing, by a processor executing said pharmacy workflow management application, a dose order record corresponding to a dose order received by said pharmacy workflow management application. The accessed dose record,
15 in this instance, has not been prepared. The method further includes initiating a dose preparation protocol corresponding to the dose order. The method further includes receiving an indication of a plurality of products associated with the dose preparation protocol. The method further includes dynamically determining, in response to the receiving, at least one of the products as a selected product corresponding to a final container based at least in part on a final container
20 characteristic of the selected product. The method further includes displaying, at a user interface of the pharmacy workflow application, a dose preparation step associated with preparation of the dose relative to the final container.

A number of feature refinements and additional features are applicable in the second

aspect. These feature refinements and additional features may be used individually or in any combination. As such, each of the following features that will be discussed may be, but are not required to be, used with any other feature combination of features of the second aspect.

For example, in an embodiment, the final container characteristic may include a first data
5 field corresponding to the dose order record and a second data field corresponding to a formulary
data record associated with at least one of the plurality of indicated products. The first data field
and the second data field are comprised of a plurality of values, from which the dynamically
determined final container calculation may be based upon. In this regard, the dynamically
determining may include identifying a value of the first data field indicative of a final container
10 indicator of said dose order such that the final container indication indicates the dose order may
be suitable for preparation in the final container. The dynamically determining may also include
identifying a value of the second data field indicative of diluent product of the plurality of
products associated with the dose preparation protocol such that the diluent product indicates the
dose order may be suitable for preparation in the final container. The dynamically determining
15 may also include identifying a value of the first data field indicative of a dose order free of a
hazmat drug such that the dose order free of the hazmat drug indicates the dose order may be
suitable for preparation in the final container. The dynamically determining may also include
identifying a value of the first data field indicative of a diluent line item and identifying a value
of the second data field indicative of a QS drug field such that the diluent line item equals the QS
20 drug field such that the dose order may be suitable for preparation in said final container.

Moreover, the dynamically determining may also include identifying a value of the first
data field indicative of a total volume of the dose order such that the value is greater than a
minimum dose volume such that said dose order is suitable for preparation in said final

container. Notably, the minimum dose volume is configurable based on a site-specific preference. In some instances, the minimum dose volume may be configured to equal 24 ml.

The dynamically determining may also include identifying a value of the second data field indicative of a final container indication of the diluent product such that the final container

5 indication of the diluent product indicates that the diluent product may be suitable as the selected product corresponding to the final container. The dynamically determining may also include identifying a value of the second data field indicative of a dose order free of multiple selected products corresponding to the final container such that the dose order, which is free of multiple selected products corresponding to said final container, indicates that the dose order may be

10 suitable for preparation in the final container. The dynamically determining may also include identifying a value of the second data field indicative of a new container of the diluent product such that the new container of the diluent product indicates that the dose order may be suitable for preparation in the final container.

Additionally, the dynamically determining may also include identifying a value of the
15 first data field indicative of a total volume of the dose order and identifying a value of the second data field indicative of a volume of the diluent product such that the absolute percentage difference between the total volume of the dose order and the volume of the diluent product is less than a predetermined differential percentage such that said dose order may be suitable for preparation in the final container. Notably, the predetermined differential percentage is

20 configurable based on a site-specific preference. In some instances, the predetermined differential percentage may be configured to equal 100%. The dynamically determining may also include identifying a value of the first data field indicative of a total volume of the dose order and identifying a value of the second data field indicative of a carrier ingredient volume of

the dose order such that the difference between the carrier ingredient volume of the dose order and the total volume of the dose order is less than a predetermined waste volume such that the dose order may be suitable for preparation in said final container. Notably, the predetermined waste volume may be configurable based on a site-specific preference. In some instances, the
5 predetermined waste volume may be configured to equal 200 ml.

A third aspect includes a method for dynamic diluent volume accounting in a dose order preparation protocol of a pharmacy workflow management application. The method includes accessing, by a processor executing the pharmacy workflow management application, a dose order record corresponding to a dose order received by the pharmacy workflow management
10 application. In this instance, the dose order has not been prepared. The method may further include receiving a first indication of a product associated with a dose preparation protocol. The method may further include dynamically determining, in response to the receiving, a required diluent volume based on the indication. The method may further include displaying, at a user interface of the pharmacy workflow application, a dose preparation step associated with
15 preparation of the dose relative to the required diluent volume.

A number of feature refinements and additional features are applicable in the third aspect. These feature refinements and additional features may be used individually or in any combination. As such, each of the following features that will be discussed may be, but are not required to be, used with any other feature combination of features of the third aspect.

20 For example, the method may further include receiving a second indication of the dynamically determined required diluent volume. The indication of the dynamically determined required diluent volume may correspond to one or more formulary products for use in the preparation of the dose order. Moreover, the one or more formulary products may be associated

with a list of allowed diluent products relative to the dose preparation protocol. As such, each of the multiple indications of the one or more formulary products may be identical formulary products. The one or more formulary products may also include a total formulary product volume such that the total formulary product equals the dynamically determined required diluent
5 volume.

In an embodiment, the foregoing noted displaying may include displaying the dynamically determined required diluent volume. In this regard, the dynamically determined required diluent volume may be configurable based on receiving a second indication of the required diluent volume. The displaying may also include displaying at least one collapsible
10 area configured to hide information when collapsed. In some instances, the information may correspond to a second indication of the dynamically determined required diluent volume.

A fourth aspect includes a method for managing an error regarding a dose order record in a pharmacy workflow queue in a pharmacy workflow management application. The method includes accessing, at a processor executing the pharmacy workflow management application, a
15 dose order record corresponding to a dose order received by the pharmacy workflow management application. In some instances, the dose order has not been prepared. The method further includes first processing, by the processor, the dose order record to generate an alert corresponding to an error state related to the dose order. The method further includes displaying, at a user interface of the pharmacy workflow management application, the dose order in
20 corresponding relation to the alert. The method further includes receiving an input indicative of a resolution of the error. The method further includes second processing the dose order based on the input. The method further includes requeuing the dose order upon confirmation of the resolution.

A number of feature refinements and additional features are applicable in the fourth aspect. These feature refinements and additional features may be used individually or in any combination. As such, each of the following features that will be discussed may be, but are not required to be, used with any other feature combination of features of the fourth aspect.

5 For example, in an embodiment, the error state may include a parsing error. The parsing error may correspond to one or more unintelligible data attributes of the dose order record. The error state may also include a drug error. The drug error may correspond to one or more unrecognizable data attributes associated with a drug of the dose order record. The method may further include creating a formulary data record at least partially based on the unrecognizable
10 data attribute. The error state may also include an unknown unit error. The unknown unit error may correspond to an unrecognizable data attribute associated with a unit of the dose order record. The error state may also include an unknown product error. The unknown product error may correspond to an unrecognizable data attribute associated with a product of the dose order record. The method may further include creating a formulary data record at least partially based
15 on the unrecognizable data attribute.

 In some embodiments, the error state may include an unknown route error. The unknown route error may correspond to an unrecognizable data attribute associated with a route of the dose order. In this regard, the method may further include modifying a formulary data record at least partially based on the unrecognizable data attribute. The error state may also include a
20 passwords error. The passwords error may correspond to an expired password of a user account. Additionally, the passwords error may also correspond to a disabled user account. The error state may include a suspected discontinued dose error. The suspected discontinued dose error may correspond to a discontinued dose order. In this regard, the method may further include

deleting the discontinued dose order. The error state may also include a suspected duplicate error. The suspected duplicate error may correspond to a duplicate dose order. In this regard, the method may further include deleting the duplicate dose order.

A fifth aspect includes a system for batch processing of a plurality dose orders in a
5 pharmacy workflow management application. The system includes an access terminal remote from and in operative communication with the pharmacy workflow management application. The pharmacy workflow management application may be operable to access a plurality of dose order records corresponding to dose orders received by the pharmacy workflow management application. Moreover, the plurality of dose orders comprise identical constituent products. In
10 some instances, the plurality of received dose orders have not been prepared. The pharmacy workflow management application may also be operable to aggregate the plurality of dose orders to create an aggregated plurality of the dose orders. The system further includes a user interface provided at the access terminal for displaying a dose preparation protocol corresponding to the aggregated plurality of dose order records. The dose preparation protocol may include an
15 aggregate amount of at least one of the identical constituent products needed to prepare each of said aggregated plurality of dose orders. Furthermore, the pharmacy workflow application may also be operable to receive, from a user by way of a user input of the pharmacy workflow management application, an indication of the aggregate amount of the at least one constituent product. The user interface may also be operable to present to the user, via the user interface,
20 one or more preparation steps regarding preparation of the aggregated plurality of doses upon receiving the aggregate amount of the at least one constituent product.

A number of feature refinements and additional features are applicable in the fifth aspect. These feature refinements and additional features may be used individually or in any

combination. As such, each of the following features that will be discussed may be, but are not required to be, used with any other feature combination of features of the fifth aspect.

For example, in an embodiment, the pharmacy workflow management application may be operable to determine that the aggregate plurality of dose orders are suitable for preparation
5 via a batch workflow. In this regard, the batch workflow may correspond to preparing the aggregated plurality of dose orders in a common sequence such that a separate dose preparation protocol is not required for each dose order of the aggregated plurality of dose orders. The pharmacy workflow management application may also be operable to initiate the dose preparation protocol that corresponds to the aggregate plurality of dose orders for preparation via
10 the batch workflow. In this regard, the pharmacy workflow management application may be operable to print a dose label for each dose order of the aggregate plurality of dose orders. Notably, the indication of the aggregate amount of the at least one constituent product may correspond to one or more formulary products for use in preparation of the aggregate plurality of doses. The pharmacy workflow management application may also be operable to print a work-in-
15 progress label upon receiving the aggregate amount of at least one constituent product such that the work-in-progress label is indicative of a remaining volume of the one or more formulary products after completion of the one or more preparation steps. As such, the presenting of the pharmacy workflow management application may include presenting the one or more preparation steps free from printing a work-in-progress label for each use of the one or more
20 formulary products such that the work-in-progress label may also be indicative of a remaining volume of the one or more formulary products after completion of the one or more preparation steps for a given dose of the aggregate plurality of doses.

In an embodiment, the indication of the aggregate amount of the at least one constituent

product corresponds to one or more formulary products for use in the preparation of the dose order. The one or more formulary products may include a total formulary product volume such that the total formulary product volume equals the aggregate amount of the at least one constituent product. The accessing of the pharmacy workflow management application may also
5 include identifying common associations among the plurality of dose order records such that the accessed plurality of dose order records comprise identical constituent products. The aggregating of the pharmacy workflow management application may also include grouping the plurality of dose orders into a queue at least partially based on the identified common associations. In this regard, the pharmacy workflow management application may also be
10 operable to initiate the dose preparation protocol that corresponds to the aggregate plurality of dose orders of the queue. In some instances, the foregoing noted aggregate amount may include each of the identical constituent products of the dose order. Notably, the indication may include a scan.

A sixth aspect includes a system for dynamic determination of a final container for use in
15 a dose order preparation protocol of a pharmacy workflow management application. The system includes an access terminal remote from an in operative communication with the pharmacy workflow management application. The pharmacy workflow management application may be operable to access a dose order record corresponding to a dose order received by the pharmacy workflow management application. In some instances, the dose order has not been prepared.
20 Moreover, the pharmacy management application may also be operable to initiate a dose preparation protocol corresponding to the dose order. The pharmacy workflow application may also be operable to receive an indication of a plurality of products associated with the dose preparation protocol. Furthermore, the pharmacy workflow application may also be operable to

dynamically determine, in response to the received indication, at least one of the products as an elected product corresponding to a final container based at least in part on a final container characteristic of the selected product. The system further includes a user interface provided at the access terminal for displaying a dose preparation step associated with preparation of the dose
5 relative to the final container.

A number of feature refinements and additional features are applicable in the sixth aspect. These feature refinements and additional features may be used individually or in any combination. As such, each of the following features that will be discussed may be, but are not required to be, used with any other feature combination of features of the sixth aspect.

10 For example, in an embodiment, the final container characteristic may include a first data field corresponding to the dose order record and a second data field corresponding to a formulary data record associated with at least one of the plurality of indicated products. The first data field and the second data field may include a plurality of values. The pharmacy workflow management application may be operable to, in relation to the dynamically determining, identify
15 a value of the first data field indicative of a final container indicator of the dose order such that the final container indication indicates that the dose order may be suitable for preparation in the final container. The pharmacy workflow management application may also be operable to, in relation to the dynamically determining, identifying a value of the second data field indicative of a diluent product of the plurality of products associated with the dose preparation protocol such
20 that the diluent product indicates that the dose order may be suitable for preparation in the final container. The pharmacy workflow management application may also be operable to, in relation to the dynamically determining, identify a value of the first data field indicative of a dose order free of a hazmat drug such that the dose order, which is free of the hazmat drug, indicates that

the dose order may be suitable for preparation in the final container. The pharmacy workflow management application may also be operable to, in relation to the dynamically determining, identify a value of the first data field indicative of a diluent line item and identify a value of the second data field indicative of a QS drug field such that the diluent line item equals the QS drug field such that the dose order may be suitable for preparation in the final container.

In some embodiments, the pharmacy workflow management application may be operable to, in relation to the dynamically determining, identify a value of the first data field indicative of a total volume of the dose order such that the value is greater than a minimum dose volume such that the dose order may be suitable for preparation in the final container. Notably, the minimum dose volume may be configurable based on a site-specific preference. In one instance, the minimum dose volume may be configurable to equal 24 ml. The pharmacy workflow management application may be operable to, in relation to the dynamically determining, identify a value of the second data field indicative of a final container indication of the diluent product such that the final container indication of the diluent product indicates that the diluent product may be suitable as the selected product corresponding to the final container. The pharmacy workflow management application may also be operable to, in relation to the dynamically determining, identify a value of the second data field indicative of a dose order free of multiple selected products corresponding to the final container such that the dose order, which is free of multiple selected products, may correspond to the final container indication that the dose order may be suitable for preparation in the final container. The pharmacy workflow management application may also be operable to, in relation to the dynamically determining, identify a value of the second data field indicative of a new container of the diluent product such that the new container of the diluent product may indicate that the dose order may be suitable for preparation

in the final container.

Furthermore, in an embodiment, the pharmacy workflow management application is operable to, in relation to the dynamically determining, identify a value of the first data field indicative of a total volume of the dose order and identify a value of the second data field
5 indicative of a volume of the diluent product. In this regard, the absolute percentage difference between the total volume of the dose order and the volume of the diluent product is less than a predetermined differential percentage such that the dose order may be suitable for preparation in the final container. The predetermined differential percentage may be configurable based on a site-specific preference. In one embodiment, the predetermined differential percentage may be
10 configured to equal 100%. The pharmacy workflow management application may also be operable to, in relation to the dynamically determining, identify a value of the first data field indicative of a total volume of the dose order and identify a value of the second data field indicative of a carrier ingredient volume of the dose order. As such, the difference between the carrier ingredient volume of the dose order and the total volume of the dose order is less than a
15 predetermined waste volume such that the dose order may be suitable for preparation in the final container. The predetermined waste volume may be configurable based on a site-specific preference. In one instance, the predetermined waste volume may be configurable to equal 200 ml.

A seventh aspect includes a method for dynamic diluent volume accounting in a dose
20 order preparation protocol of a pharmacy workflow management application. The system includes an access terminal remote from an in operative communication with the pharmacy workflow management application. The pharmacy workflow management application may be operable to access a dose order record corresponding to a dose order received by the pharmacy

workflow management application. In some instances, the accessed dose order has not been prepared. Moreover, the pharmacy workflow management application may be operable to receive a first indication of a product associated with a dose preparation protocol. Furthermore, the pharmacy workflow management application may also be operable to dynamically
5 determine, in response to the received first indication, a required diluent volume based on the indication. The system further includes a user interface provided at the access terminal for displaying a dose preparation step associated with preparation of the dose relative to the required diluent volume.

A number of feature refinements and additional features are applicable in the seventh
10 aspect. These feature refinements and additional features may be used individually or in any combination. As such, each of the following features that will be discussed may be, but are not required to be, used with any other feature combination of features of the seventh aspect.

For example, the pharmacy workflow management application may be operable to receive a second indication of the dynamically determined required diluent volume. The
15 indication of the dynamically determined required diluent volume may correspond to one or more formulary products for use in preparation of the dose order. Notably, the one or more formulary products may be associated with a list of allowed diluent products relative to the dose preparation protocol. In some instances, each of the one or more formulary products may be identical formulary products. The one or more formulary products may include a total formulary
20 product volume such that the total formulary product equals the dynamically determined required diluent volume.

In an embodiment, the pharmacy workflow management application may be operable to, in relation to the displaying, display the dynamically determined required diluent volume. In this

regard, the pharmacy workflow management application may be operable to configure the dynamically determined required diluent volume based on receiving a second indication of the required diluent volume. In some instances, the user interface may be operable to display at least one collapsible area configured to hide information when collapsed. The information, in this
5 regard, may correspond to a second indication of the dynamically determined required diluent volume.

An eighth aspect includes a system for managing an error regarding a dose order record in a pharmacy workflow queue in a pharmacy workflow management application. The system includes an access terminal remote from an in operative communication with the pharmacy
10 workflow management application. The pharmacy workflow management application may be operable to access a dose order record corresponding to a dose order received by the pharmacy workflow management application. In some instances, the dose has not been prepared.

Moreover, the pharmacy workflow application may be operable to first process the dose order record to generate an alert corresponding to an error state related to the dose order. The system
15 further includes a user interface provided at the access terminal for displaying the dose order in corresponding relation to the alert. Furthermore, the pharmacy workflow management application may be operable to receive an input indicative of a resolution of the error. In this regard, the pharmacy management application may be operable to second process the dose order based on the input and requeue the dose order upon confirmation of the resolution.

20 A number of feature refinements and additional features are applicable in the eighth aspect. These feature refinements and additional features may be used individually or in any combination. As such, each of the following features that will be discussed may be, but are not required to be, used with any other feature combination of features of the eighth aspect.

For example, in an embodiment, the error state may include a parsing error. The parsing error may correspond to one or more unintelligible data attributes of the dose order record. The error state may also include a drug error. The drug error may correspond to one or more unrecognizable data attributes associated with a drug of the dose order record. In this regard, the pharmacy workflow application may be operable to create a formulary data record at least partially based on the unrecognizable data attribute. The error state may include an unknown unit error. The unknown unit error may correspond to an unrecognizable data attribute associated with a unit of the dose order record. The error state may also include an unknown product error. The unknown product error may correspond to an unrecognizable data attribute associated with a product of the dose order record. In this regard, the pharmacy workflow management application may be operable to create a formulary data record at least partially based on the unrecognizable data attribute.

Furthermore, in an embodiment, the error state may also include an unknown route error. The unknown route error may correspond to an unrecognizable data attribute associated with a route of the dose order. In this regard, the pharmacy workflow management application may be operable to modify a formulary data record at least partially based on the unrecognizable data attributes. The error state may also include a passwords error. The passwords error may correspond to an expired password of a user account. In other embodiments, the passwords error may correspond to a disabled user account.

In an embodiment, the error state may include a suspected discontinued dose error. The suspected discontinued dose error may correspond to a discontinued dose order. In this regard, the pharmacy workflow management application may be operable to delete the discontinued dose order. The error state may also include a suspected duplicate error. The suspected

duplicate error may correspond to a duplicate dose order. In this regard, the pharmacy workflow management application may be operable to delete the duplicate dose order.

BRIEF DESCRIPTION OF THE DRAWINGS

5 Fig. 1 is a schematic view of an embodiment of a system including a pharmacy workflow management application executed at a local system.

 Fig. 2 illustrates a flow diagram of an embodiment of a method of batch processing of a plurality of dose orders in a pharmacy workflow management application.

 Fig. 3 illustrates a flow diagram of an embodiment of a method for dynamic determination
10 of a final container for use in a dose order preparation protocol of a pharmacy workflow management application.

 Fig. 4a illustrates a flow diagram of an embodiment of a method for dynamic determination of a final container for use in a dose order preparation protocol of a pharmacy workflow management application.

15 Fig. 4b illustrates a flow diagram of an embodiment of a method for a QS volume determination for use in an embodiment of a method for dynamic determination of a final container for use in a dose order preparation protocol of a pharmacy workflow management application.

 Fig. 4c illustrates a flow diagram of an embodiment of a method for a QS volume sufficiency check for use in an embodiment of a method for dynamic determination of a final
20 container for use in a dose order preparation protocol of a pharmacy workflow management application.

 Fig. 4d illustrates a flow diagram of an embodiment of a method for the identification of active workstation products for use in an embodiment of a method for dynamic determination of a

final container for use in a dose order preparation protocol of a pharmacy workflow management application.

Fig. 4e illustrates a flow diagram of an embodiment of a method for dynamic determination of a final container for use in a dose order preparation protocol of a pharmacy workflow management application.

Fig. 5 illustrates a screenshot of a user interface of the pharmacy workflow application, according to an embodiment.

Fig. 6 illustrates a screenshot of a user interface of the pharmacy workflow application, according to an embodiment.

Fig. 7 illustrates a screenshot a user interface of the pharmacy workflow application, according to an embodiment.

Fig. 8 illustrates a flow diagram of an embodiment of a method for dynamic diluent volume accounting in a dose order preparation protocol of a pharmacy workflow management application.

Fig. 8a illustrates a screenshot of a user interface of the pharmacy workflow application, according to an embodiment.

Fig. 8b illustrates a screenshot of a user interface of the pharmacy workflow application, according to an embodiment.

Fig.8c illustrates a screenshot of a user interface of the pharmacy workflow application, according an embodiment.

Fig. 9 illustrates a screenshot of a user interface of the pharmacy workflow application, according to an embodiment.

Fig. 10 illustrates a flow diagram of an embodiment of a method for managing an error regarding a dose order record in a pharmacy workspace queue in a pharmacy management application.

Fig. 11 illustrates a screenshot of a user interface of the pharmacy workflow application,
5 according to an embodiment.

Fig. 12 illustrates a screenshot of a user interface of the pharmacy workflow application, according to an embodiment.

Fig. 13 illustrates a flow diagram of an embodiment of a method for identifying alerts on a dose order for use in an embodiment of a method for managing an error regarding a dose order
10 record in a pharmacy workspace queue in a pharmacy management application.

Fig. 14 illustrates a flow diagram of an embodiment of a method for identifying alerts on user accounts for use in an embodiment of a method for managing an error regarding a dose order record in a pharmacy workspace queue in a pharmacy management application.

Fig. 15 illustrates a flow diagram of an embodiment of a method for identifying alerts
15 scanned products barcodes for use in an embodiment of a method for managing an error regarding a dose order record in a pharmacy workspace queue in a pharmacy management application.

Fig. 16 illustrates a flow diagram of an embodiment of a method for reprocessing dose order for use in an embodiment of a method for managing an error regarding a dose order record in a pharmacy workspace queue in a pharmacy management application.

Fig. 17 illustrates a flow diagram of an embodiment of a method for requeueing dose
20 orders for use in an embodiment of a method for managing an error regarding a dose order record in a pharmacy workspace queue in a pharmacy management application.

DETAILED DESCRIPTION

The following description is not intended to limit the invention to the forms disclosed herein. Consequently, variations and modifications commensurate with the following teachings, skill and knowledge of the relevant art, are within the scope of the present invention. The
5 embodiments described herein are further intended to explain modes known of practicing the invention and to enable others skilled in the art to utilize the invention in such, or other embodiments and with various modifications required by the particular applications(s) or use(s) of the present invention.

The present disclosure contemplates a pharmacy workflow management application that
10 may facilitate pharmacy workflow management in relation to preparation of one or more doses at a pharmacy, hospital, or other facility at which doses are prepared for administration to a patient. The pharmacy workflow management application may be executed at a local system (also referred to herein as a local node). The local node may correspond to a facility such as a hospital, pharmacy, or other facility capable of preparing a dose for administration to a patient. Pharmacy workflow
15 management provided by the pharmacy workflow management application may include one or more activities within a pharmacy that may include management of a locally stored formulary, processing received medication dose orders at a pharmacy, creating corresponding dose order records for each received dose order, managing (e.g., viewing, sorting, prioritizing, etc.) dose order records, guiding a pharmacy technician in preparing the dose order medication, gathering
20 information regarding the preparation of a dose to fulfill a dose order, reviewing the medication dose order, and/or tracking of medication dose order records in the pharmacy.

The local node may comprise one or more devices that may include hardware and/or software that may execute machine readable instructions stored as machine readable non-transitory

data in a memory. The machine readable instructions, when executed by a processor may provide functionality related to the pharmacy workflow management application discussed herein. In this regard, the pharmacy workflow management application may include one or more processors operative to access machine-readable data stored in a storage medium (i.e., a memory) that provide
5 instructions to the one or more processors for execution to provide the functionality described herein.

The local node executing the pharmacy workflow management application may be in operative communication with a central server. The central server may provide support services to the local node to support the pharmacy workflow management application and/or may provide
10 additional services such as data backup, report generation, or access to data sets stored at the central server.

The present disclosure includes a description of an embodiment of dose order processing that may be performed by the pharmacy workflow management application. For instance, the pharmacy workflow management application may facilitate a method of batch processing of a
15 plurality of dose orders. In another embodiment, the pharmacy workflow application may facilitate the dynamic alteration of a final container in a dose order preparation protocol. In yet another embodiment, the pharmacy workflow application may facilitate dynamic diluent volume accounting and/or adjustment in a dose preparation protocol. In yet another embodiment, the pharmacy workflow application may facilitate managing an error regarding a dose order record.

20 With initial reference to Fig. 1, a system 100 is depicted that includes a plurality of local systems 110. The local systems 110 may each comprise a local node in the system 100. The local systems 110 may each execute a pharmacy workflow management application 114. For instance, each local system 110 may include at least one local device executing a client application for

providing functions related to the pharmacy workflow management application. As shown in Fig. 1, a plurality of local systems 110 may be provided that are each in operative communication with a central server 120. For example, the local systems 110 may be in operative communication the central server 120 by way of a wide area network 50 (e.g., the Internet).

5 Each local system 110 may be a healthcare facility such as a hospital, pharmacy, outpatient clinic, or the like, that prepares doses for administration to a patient. The respective pharmacy workflow management application 114 at each local system 110 may be operative to generate and/or capture medical information that may be provided to the central server 120 by way of the wide area network 50. As will come to be appreciated from the discussion below in relation to the
10 pharmacy workflow management application, the medical information may include medication dose order data that may include metadata regarding the dose order record and/or dose. The pharmacy workflow management application 114 at each local system 110 may allow for the pharmacy workflow management functions described in greater detail below. In any regard, any or all data generated by the pharmacy workflow management application may be provided to the
15 central server 120. In this regard, central server 120 may provide for data collection and/or data backup services in relation to the local systems 110.

Accordingly, in at least one embodiment the local systems 110 may comprise unaffiliated and discrete healthcare facilities capable of preparing medication doses for administration to patients. The central server 120 may be hosted by another discrete and unaffiliated third-party that
20 may be separate from any of entities of the local systems 110. For instance, the central server 120 may be hosted and/or executed by an application provider that provides one or more client applications for execution by the local systems 110 to facilitate the pharmacy workflow management application 114. Specifically, the central server 120 may be executed or hosted by

an application provider that provides the pharmacy workflow management application each local system 110.

A local system 110 may include a local database 112 for storage of information. As may be appreciated, the local database 112 may in fact include a collection of one or more databases collectively defining the local database 112. In an implementation of the present invention, the local system 110 may include a server that facilitates access to the database 112. The database may store data related to the current in process work and/or store archival data related to prior work completed at the pharmacy. The local database 112 may be a high-performance, highly scalable and SQL-compliant object database. In this regard, the database may scale easily to handle thousands of simultaneous users and potentially terabytes of data.

In addition to storing information related to pharmacy work (i.e., dose order records), the database 112 may include information in a variety of contexts including information related to formulary information, administrative information, or other information related to the pharmacy workflow management application 114 at the local system 110. In any regard, the local system 110 may execute a number of services (e.g., provided by modules executed by a processor). For instance, a data backup module may be provided that provides a rule-based approach to data backup from the local system 110 to the central server 120. The data backup module may define the interval at which the local system 110 provides backup data to the central server 120 and/or may dictate what information is provided to the central server. The backup module may also provide an administrator (e.g., at the central server or the local system 110) with information relating to the success or failure of data backup operations, system slowdowns, and the like. For instance, the backup module may facilitate selective backup of the local database 112 as described in U.S. Patent Application No. 62/019,227 incorporated by reference herein in its entirety.

Turning to Fig. 2, in one embodiment, the pharmacy workflow application 114 may facilitate preparation of a plurality of dose orders according to a batch workflow according generally to method 200. In this regard, a plurality of dose orders having identical constituent products associated with different dose order records may be prepared according to a batch workflow such that a plurality of such dose orders may be prepared without initiating a new workflow process for each of the plurality of dose orders.

In some instances, the preparation of dose orders in a batch workflow may facilitate efficient dose order preparation techniques and contribute to overall error reduction related to the dose fulfillment. Due at least in part to the fact that a new workflow is not being initiated for each of the plurality of doses, the batch workflow process may facilitate efficient dose order preparation by potentially eliminating repetitive preparation steps and organizing the steps relative to the plurality of doses such that like steps are performed together. In this regard, the batch workflow may comprise various product identification steps and various dose compounding steps. The product identification steps may establish, based on a received indication of an aggregate amount of at least one constituent product, the presence or possession of a product at, for example, a dose compounding station. The dose compounding steps may include one or more preparation steps regarding the physical preparation of the aggregate plurality of doses. Notably, the batch workflow may execute the various dose compounding steps after the completion of the various product identification steps for each of the aggregate plurality of doses of the batch. As such, the batch workflow may facilitate faster dose order preparation by grouping the product identification steps and the dose compounding steps such that, for example, the presence of at least one constituent product to fulfill the aggregate plurality of dose orders is established prior to the corresponding dose compounding step. For example, as discussed in greater detail below, the

batch workflow may be operable to identify an aggregate amount of at least one constituent product of the plurality of doses. In this regard, all the required products (*e.g.*, a diluent, a drug, or another product) may be identified prior to the physical preparation of the plurality of dose orders. This may eliminate repetitive steps related to identifying an amount of a product for each individual
5 dose. The elimination of this and other repetitive steps may save a substantial amount of time, which thereby increases the efficiency of the overall preparation of dose orders. Error rates may also be reduced. For example, each of the plurality of doses prepared via the batch workflow may comprise identical constituent components. In this regard, the products identified during the preparation during the batch workflow may be appropriately associated with any of the plurality
10 of dose orders. This may reduce errors by reducing the occurrence, for example, of inappropriate products being introduced into the prepared dose order.

The pharmacy workflow application 114 may access 204 a plurality of dose order records stored in, for example, local database 112. The dose order records may correspond to dose orders received by the pharmacy workflow application 114 that have not yet been prepared.
15 Furthermore, a dose order may include a request for medication that includes various constituent products. Each dose order accessed 204 by pharmacy workflow application 114 comprising the plurality of dose orders (which may include less than all dose orders at the pharmacy workflow application 114) may include identical constituent products such that each of the plurality of dose orders comprises a request for the same constituent products. In this regard, each dose order may
20 include a plurality of identical data attributes corresponding to, among other things, an identical ingredients listing, identical ingredient amount, identical ingredient concentration, and the like. Accordingly, based in at least in part on the plurality of identical data attributes, the plurality of dose orders may be identified as being eligible for being prepared via the batch workflow, as

described in greater detail below.

The pharmacy workflow application 114 may access 204 the plurality of dose orders based at least in part on identifying common associations (*e.g.*, one or more data attributes as described above) among the plurality of dose order records such that each dose order comprises a request for
5 identical constituent products, as described as above. In some embodiments, the common associations of the plurality of dose order records may be identified via a filter applied to the plurality of dose order records. That is, the pharmacy workflow application 114 may access a plurality of dose orders based on the identified common attributes such that the accessed dose order constitute dose orders that include identical constituent products. In some instances, the
10 pharmacy workflow application 114 may access the plurality of dose orders that include identical constituent products from a database of dose orders that include a second plurality of dose orders that do not include identical constituent products as the identical constituent products of the identified plurality. In this regard, the filter may facilitate accessing or identifying dose orders appropriate for preparation via the batch workflow by accessing the plurality of dose orders that
15 include identical constituent products.

The pharmacy workflow application 114 may aggregate 208 the accessed dose order records to create an aggregated plurality of dose orders (*e.g.*, a batch) to facilitate the batch workflow. This may be accomplished in terms of a queue such that each of the plurality of dose orders that share the common associations (*e.g.*, identical constituent products) are grouped
20 together to form the aggregated plurality of dose orders (*e.g.*, each identified dose order of “Cefazolin 1 mg in 10 ml NS syringe” may be grouped together in the queue as an aggregated plurality of dose orders to define a batch of orders). The pharmacy workflow application 114 may prepare the aggregate plurality of dose orders in a common batch according to a batch workflow.

The batch workflow may include a protocol for preparation of all dose orders in the batch.

In one embodiment, the queue may facilitate batch processing initiation by determining 212 the plurality of dose orders as suitable for batch processing and initiating 216 the dose preparation protocol corresponding to the aggregate plurality of dose orders for preparation via the batch workflow. In this regard, only those dose order records associated with the queue (*i.e.*, the dose order belongs to a batch of dose orders with identical constituent products) may be suitable for preparation according to the batch workflow. In this regard, the queue may facilitate the inclusion of only those dose orders with identical constituent products for preparation via the batch workflow. Accordingly, the pharmacy workflow application 114 may initiate 216 a dose preparation protocol corresponding to the aggregate plurality of dose orders for preparation via the batch workflow based at least in part on the presence of a dose order in of the queue. In some instances, the batch workflow is initiated directly from the queue, or the presence of a generated queue within the pharmacy workflow application 114. Initiating of the batch workflow from the queue may also provide a user safeguard, by ensuring that only the plurality of dose orders of identical constituent products may be selected for the batch workflow.

With returned reference to Fig. 1, local system 110 may also include a user interface 108. The user interface 108 may be generally operable to present and receive information associated with the batch workflow per the operation of the pharmacy workflow application 114. For instance, the pharmacy workflow application 114 may display 220 a dose preparation protocol corresponding to the aggregate plurality of dose order records. The dose preparation protocol may include information pertaining to the required aggregate amount of one or more of the identical constituent products needed to prepare each of the aggregated plurality of dose orders. For the sake of illustration, consider the example where the aggregate plurality of dose orders includes 10

dose orders, each for Cefazolin 1,000 mg, 0.9% Sodium Chloride 250 ml in 300 ml. In this illustration, the dose preparation protocol may include information pertaining to the volume of constituent products required to make all 10 dose orders (*i.e.*, 10,000 mg Cefazolin and 2,900 ml of Sodium Chloride). In this regard, the dose preparation protocol includes requirements
5 information pertaining to the preparation of all of the aggregate plurality of dose orders selected for batch preparation in the batch workflow.

The aggregate plurality of dose orders may be prepared by initially receiving 224 an indication related to products to be used to prepare the batch that identify the aggregate amount of at least one of the constituent products. In this regard, the pharmacy workflow management
10 application 114 may receive a user input indicative of the aggregate amount of the at least one constituent products. For example, the pharmacy workflow management application 114 may receive an indication via one or more product scans, NDC numbers, or other input that correspond to an indication of the possession of a product by a user for use in preparing the aggregate plurality of the dose orders. That is, the indication may correspond to the total amount (*e.g.*, volume or
15 mass) of the product required for preparing the aggregate plurality of dose orders. As such, the corresponding product of the indication may include a product volume such that the product volume is at least equal to the aggregate amount of the at least one constituent products. In some instances, the pharmacy workflow application 114 may require receipt of the indication identifying the aggregate amount of the at least one constituent product before presenting to the user one or
20 more preparation steps of the aggregate plurality of dose orders, as discussed in greater detail below. In this regard, a pharmacy technician, for example, may indicate the possession of the at least one constitute product to the pharmacy workflow application 114 before executing the one or more preparation steps.

In some embodiments, the user input may comprise a plurality of indications. For example, each of the plurality of indications may include indications related to different instances of a given product for use in preparing the aggregate plurality of dose orders. Accordingly, in this embodiment, the multiple instances of the given products, collectively, corresponding to the plurality of indications, may be at least equal to the aggregate amount of the at least one constituent product. In this regard, multiple, separate products may be used to prepare the aggregate plurality of dose orders. As the batch workflow may facilitate the preparation of any sized batch, including larger batches of 50, 100, or more, the indication corresponding to multiple separate products may support the preparation of the aggregate volume for larger batches.

For the sake of illustration, consider again the example where the aggregate plurality of dose orders includes 10 dose orders, each for Cefazolin 1,000 mg, 0.9% Sodium Chloride 250 ml in 300 ml. The pharmacy workflow application 114 may display the dose preparation protocol of the dose that includes information pertaining to the required volume of constituent products necessary to make all 10 dose orders (*i.e.*, 10,000 mg Cefazolin and 2,900 ml of Sodium Chloride), including the physical step associated with a user retrieving the products for use in the preparation of the aggregate plurality of dose orders. In turn, the pharmacy workflow application 114 may receive the user input corresponding to an indication identifying the amount of at least one of the constituent products that the user possesses for use in preparing the dose. For example, the user input may pertain to the 10,000 mg of Cefazolin or the 2,900 ml of Sodium Chloride – *i.e.*, the volume of at least one constituent product required to make all 10 dose orders, thus indicating the user has sufficient product to make dose order in the batch. In some instances, the user input may pertain to both the 10,000 mg of Cefazolin and the 2,900 ml of Sodium Chloride – *i.e.*, the volume of each different respective vials of the constituent products required to make all 10 dose orders.

In either case, the user input may include a plurality of indications, each of which may correspond to a different instance of the given product such that the plurality of indications corresponds to the aggregate amount of at least one constituent product. For example, the user may retrieve two 5,000 mg vials of Cefazolin. In turn, where the user scans each vial, each of the plurality of indications
5 may include separate indications for two 5,000 mg vials of Cefazolin. In this regard, the user input would include two indications of the 5,000 mg vial of Cefazolin such that the plurality of indication corresponds to the aggregate amount of the identical constituent product (*e.g.*, 2 vials of 5,000 mg Cefazolin equals the 10,000 mg of Cefazolin, the aggregate amount).

The pharmacy workflow application 114 may also present 228 preparation steps to the user
10 via user interface 108. As previously noted, a product preparation steps comprises the physical steps required to prepare a dose (*e.g.*, for the sake of illustration, remove 10 ml diluent volume from 0.9% Sodium Chloride – 300 ml, Acquire 10 ml from Cefazolin, etc.). Specifically, the pharmacy workflow application 114 may present 228 preparation steps to the user upon receipt of the user input indicative of the aggregate amount of the at least one constituent products. In this
15 regard, the pharmacy workflow application 114 may present preparation steps after all products for the aggregate amount of the at least one constituent product have been identified via the corresponding indication. Accordingly, the aggregate plurality of dose orders of the batch workflow may be prepared collectively such that any single dose order of the aggregate plurality of dose orders will not be prepared until the entire aggregate amount of the at least one constituent
20 product is indicated via the corresponding indication. This may facilitate a batch workflow by segmenting the preparation steps such that each preparation step is performed on each of the aggregated plurality of dose orders before proceeding to the next preparation step (*e.g.*, receiving an indication of an aggregate amount of product sufficient to fulfill the aggregate plurality of

orders, before physically preparing any of the aggregate plurality of dose orders, etc.).

In some embodiments, the pharmacy workflow application 114 may print 232 a label for each of the aggregate plurality of dose orders. A label may be applied to a prepared dose order and include information associated with the contents of the prepared dose order. Based on the
5 identified common associations of the queue filter, each of the prepared dose orders may contain identical constituent products. In this regard, printing 232 labels for each of the aggregate plurality of dose orders may be incorporated into the batch workflow process in a particular and separate processing segment of the foregoing. This may include, for example, printing the labels for each of the aggregate plurality of dose orders at a common time. For example, the pharmacy workflow
10 application 114 may print the labels after receiving an indication of the aggregate amount of the at least one constituent product. In other instances, the pharmacy workflow application 114 may print the labels after receiving an indication that the one or more steps of the preparation protocol have been completed. Accordingly, the labels may all be applied at a common time as a discrete step of the batch workflow process.

15 In another embodiment, the batch workflow may facilitate printing work-in-progress (“WIP”) labels such that the pharmacy workflow application 114 may print WIP labels after the completion of the one or more preparation steps. A WIP label may be applied, for example, to a partially-used diluent product to indicate the remaining diluent volume, expiration data, and the like. In this regard, to the extent that the preparation of a dose generates a partially-used diluent
20 product, the preparation of dose via batch workflow may require printing a WIP label only after the preparation of the aggregate plurality of dose orders of the batch.

For example, consider a batch comprised of a hypothetical 10 doses of Cefazolin 1,000 mg, 0.9% Sodium Chloride 250ml in 300 ml. A 3,000 ml bag of Sodium Chloride may satisfy the

required diluent to prepare the batch (*i.e.*, 3,000 ml > 2,900 ml required to prepare the batch). If the 10 doses of the hypothetical batch were prepared individually, 10 WIP labels may theoretically be printed (*e.g.*, a first WIP label after the first prepared dose indicating 2,700 ml remaining in the Sodium Chloride bag, a second WIP label after the second prepared dose indicating 2,400 ml
5 remaining in the Sodium Chloride bag, etc.). The multiple printing of WIP labels in this regard may contribute to additional processing steps that may result in process inefficiency. However, the same 10 doses prepared as a batch via the batch processing may result in printing only 1 WIP label, for example. That is, after presenting the one or more preparation steps regarding the aggregate plurality of doses, the pharmacy workflow application 114 may print one WIP label
10 indicating 100 ml remaining in the Sodium Chloride bag (*i.e.*, 3,000 ml bag – 2,900 ml required equals 100 ml).

Turning next to Fig. 3, according to another embodiment, the pharmacy workflow application 114 may facilitate the dynamic alteration of a final container in a dose order preparation protocol according generally to method 300. In this regard, a formulary product may be identified
15 as being capable of use as a final container – *i.e.*, the container which will ultimately contain the prepared dose. Accordingly, a dose order need not necessarily be prepared in a new or empty container. Additionally, a user need not guess or make ad hoc choices as to the final container. Rather, the pharmacy workflow application 114 dynamically determines whether a formulary product indicated as being used in a dose preparation protocol is appropriate as a final container
20 based on a received indication of a plurality of products associated with the dose order.

Considered in the context of preparing a dose order, for every dose order preparation there is a final container. The final container may either be a container associated with a formulary product used in the preparation of the dose or some other container (*e.g.*, a new or empty

container). Various rules and conditions, explained in greater detail below, executed by the pharmacy workflow application 114 may be applied to a dose order in view of a received indication of a product to be used to prepare the dose in order to dynamically determine a selected product as a final container during the dose preparation protocol. In this regard, based on the rules and conditions, the pharmacy workflow application 114 may receive an indication of a plurality of products to be used to prepare the dose and dynamically determine one of the plurality of products as the final container or whether to prepare a final container. Pharmacy workflow application 114 may also add a visual indicator to a user interface to specify the final container in relation to a listing of products of the dose preparation protocol. After each received indication, the pharmacy workflow application 114 may reevaluate the final container determination decision such that initially identified final container may not be valid until receiving an indication of the plurality of products.

In some instances, the dynamic alteration of a final container may facilitate efficient dose order preparation techniques and contribute to overall error reduction by substantially automating the final container determination. For example, absent the pharmacy workflow application 114 dynamically determining a final container, the user may be required to make a manual determination during the preparation of the dose order. In this regard, previous dose order preparation practices did not automate the final container determination. Rather, previous dose order preparation practice relied substantially on a user's judgment to determine the final container based in part on experiential knowledge. Due in part to the pharmacy workflow application 114 dynamically determining the final container based on a final container characteristic of a selected product, user judgment alone need not be relied upon to complete the determination. In this regard, error rates may be reduced by standardizing the final container determination (*e.g.*, by way of a

standard final container algorithm). In particular, dynamically determining the final container based on a final container characteristic of a selected product may reduce errors associated with waste produced during dose order preparation. For example, according to one embodiment, the dose order may only be prepared in a final container where the dose order exceeds a dose order minimum volume (*e.g.*, 20 ml, 24 ml, 20 ml, etc.)

Moreover, dose order preparation efficiency may also be increased because the standardized final container determination may be performed in the same manner in each circumstance. Notably, certain aspects of the standardized final determination may be customizable according to local preferences such as, and discussed in greater detail below, dose order volume rules, waste volume rules, and the like. The customizable aspects may facilitate system-wide consistency as the local preferences may be uniformly applied across a plurality of pharmacy workflow applications 114 operating in a common network, such as wide area network 50, for example. Accordingly, knowing the final container, the pharmacy workflow application 114 may display the correct dose preparation steps associated with preparation of the dose relative to the final container.

With further reference to Fig. 3, which illustrates the dynamic alteration of the final container, which includes the pharmacy workflow application 114 accessing 304 a dose order record corresponding to a dose order stored in, for example, local database 112. The dose order record may correspond to a dose order received by the pharmacy workflow application 114 that has not yet been prepared. The dose order may include a request for medication that is awaiting preparation via the pharmacy workflow application 114. In some instances, the dose order record of the corresponding dose order may be stored in a queue comprised of a plurality of dose order records with corresponding dose orders also awaiting preparation.

A dose preparation protocol may be initiated 308 that corresponds to the accessed dose order. The dose preparation protocol may include a number of product identification steps, which may include list of constituent products or a “recipe” for the dose order. This may include, among other things, a description of each constituent product for preparation of the dose order, the volume
5 or mass required of each constituent product, and the like for use in identify the various products required to prepare the dose order. For example, a dose preparation protocol for a hypothetical dose of Cefazolin 1,000 mg, 0.9% Sodium Chloride 250 ml in 300 ml may identify 1,000 mg of Cefazolin and 250 ml of 0.9% Sodium Chloride as within the dose preparation protocol. Moreover, the dose preparation protocol may include the various dose compounding steps, which may
10 communicate steps associated with the physical preparation of the dose order (*e.g.*, remove diluent, remove drug, apply label to dose, etc.).

The pharmacy workflow application 114 may then receive 312 an indication of a plurality of products associated with the dose preparation protocol. The received indication may be indicative of, for example, a pharmacy technician obtaining possession of the product for use in
15 preparing the dose. In this regard, the pharmacy workflow application 114 receives an indication that a product associated with a formulary data record of the same product has been identified for use in preparing the dose of the dose order. Each corresponding product of the indication may be associated with a plurality of corresponding data attributes per a formulary data record corresponding to the product that may describe, among other items, physical attributes of the
20 product, such as the product volume. In some instances, the plurality of corresponding data attributes may include a data field indicative of the suitability of a product as a final container. In any event, the totality of corresponding data attributes may correspond to a final container characteristic, such that the received indication facilitates a dynamic determination of the final

container.

Accordingly, the pharmacy workflow application 114 may dynamically determine 316, in response to the received indication, at least one of the products as a selected product corresponding to a final container based at least in part on analysis of the final container characteristic of the products indicated as being used to prepare the doses. That is, the pharmacy workflow application 114 may determine whether a product associated with the indication that the product is to be used in preparation of the dose order may constitute the final container, which will serve as a destination receptacle to contain the prepared dose. If the container of the indicated formulary product cannot be used as the final container, the dose may be prepared in a new, empty container. The dynamic determination may change for each received indication of the plurality of products associated with the dose preparation protocol during the product identification steps. For example, a first diluent product container may initially be determined to constitute the final container. A second diluent product container indication may be received that modifies the final container determination (*e.g.*, to the second diluent container, or into the empty container) based on receipt of the indicator associated with the second diluent product container. That is, receipt of the indication corresponding to the second diluent container may trigger a change in the final container determination.

The final container characteristic, as noted above, may correspond to a plurality of data attributes associated with the product for use in preparation of the dose order. As such, the dynamic determination may be at least partially based on the execution of a plurality of rules, as discussed in more detail below, operable to determine whether a particular product should constitute the final container (*e.g.*, a volume differential rule may be adopted that dictates that a formulary product should not be a final container if the percentage difference between product

total volume and dose order total volume is greater than or equal to a specified differential percentage). The dynamic determination may also be based on characteristics of the dose order itself, such as by requesting a final volume of the order (*e.g.*, the final container may be determined based on an adjusted diluent ingredient volume corresponding to the received product indications).

5 The previously noted interface 108 of local system 110 may also display 320 a dose preparation step associated with the preparation of the dose relative to the final container. That is, the preparation step of the preparation protocol may be at least partially based on the dynamically determined final container. For instance, the protocol may differ for a given dose order based on the final container determination (*e.g.*, a negative QS volume versus a positive QS volume, etc.).

10 Also, the preparation protocol may include displaying an indication to the user corresponding to the product that constitutes the final container. In some instances, the final container product may be denoted at the user interface with an insignia such as that of the encircled capital letters “FC,” for example. As such, the displaying may therefore facilitate communication to the user of which, if any (*i.e.*, in the case of requiring an empty container), container constitutes the final container

15 such that a user may use this information to prepare the dose of the dose order in the noted final container.

 The pharmacy workflow application 114 may at least in part facilitate dynamically determining the final container by adjusting the diluent ingredient volume associated with the dose preparation protocol. In particular, the pharmacy workflow application 114 may optimize the dose

20 preparation protocol by adjusting the request diluent ingredient volume before displaying the dose preparation step associated with preparation of the dose order relative to the final container. In order to calculate the exact required diluent volume for the dose order, after indication of a product to be used to prepare the dose (*e.g.*, a product scan), the pharmacy workflow application 114 may

identify the currently active workstation products for the dose order in the particular dose order. The pharmacy workflow application 114 may reevaluate this identification for each subsequent indication received. The pharmacy workflow application 114 may adjust the diluent ingredient volume, based in part on the volume provided by the drug ingredient and requested final volume of the dose order. The foregoing adjustment may therefore update the requested diluent ingredient volume of the dose order, subject to various exceptions, described in greater detail below.

Turning now to Figs. 4a–4e, the pharmacy workflow application 114 may dynamically determine a final container of the dose order based at least in part on a final container characteristic in relation to a plurality of logical steps according generally to the logical steps 400. Each of the plurality of logical steps may reference at least one data value or attribute of the final container characteristic in order to carry out the logical step. In some regards, the plurality of logical steps may be viewed as a checklist of requirements such that each requirement must be satisfied in order for the selected product of the dose order to constitute a final container. The failure of any one of the plurality of logical steps may result in the dose order being prepared in a new or empty container. The dose order may be prepared in the new or empty container even in the case where some of the logical steps are satisfied – *i.e.*, all logical steps must be satisfied in order for the selected container to constitute a final container. The logical steps may be performed in the illustrated sequences or in other sequences accordingly.

In one embodiment, the plurality of logical steps may comprise the logical steps 400 illustrated in Fig. 4. The logical steps may proceed such that a subsequent logical step may be executed upon the satisfaction of a previous logical step. It will be appreciated that the logical steps may include other logical steps, or omit one or more of the following described logical steps 400, as needed to dynamically determine the final container based on said final container

characteristic.

The logical steps 400 may reference various data attributes in dynamically determining the final container. In some instances, the data attribute may include a data field corresponding to a dose order record. In other instances, the data attribute may include a data field corresponding to a formulary record associated with at least one of the plurality of indicated products. In this regard, the formulary record associated with at least one of the plurality of indicated products may include a data field at least partially for the purpose of the pharmacy workflow application 114 dynamically determining the selected product as the final container. As a further example, each formulary product may be marked as a drug, diluent, or both a drug and diluent (*e.g.*, a pre-mix product). Additionally, chemotherapy or hazmat products may be flagged or indicated using a specific indication in the formulary record data field. Further, the formulary data record may indicate that only a product marked or indicated as a diluent or a drug/diluent can be used as a final container. As such, these products may be specifically indicated as a “final container” in the formulary to indicate that the product is eligible for use as such. Also, a product with a product volume less than or equal to a selectable, predetermined volume (*e.g.*, 5, 10, 20, or 25 ml or less, etc.) may be indicated as not eligible as a final container or a “non-final” container. Further still, products that are clinically deemed as not suitable to be final containers may be indicated as a non-final container. In some embodiments, other data fields may be included in the formulary data record for purposes of making the final container determination. Moreover, the specific values may be configurable for site-specific preferences. In any event, these data fields of the formulary data record may be accessed by one or more of the logical steps 400 to facilitate the pharmacy workflow application 114 dynamically determining the final container.

In this regard, pharmacy workflow application 114 may execute a logical step 404 to

determine whether the dose order is associated with a data attribute corresponding to an indicator indicative of the suitability of the dose order to be prepared in a final container. In some instances, logical step 404 may be referred to as a “Dose Logic Engine” rule or DLE rule. If the dose order does not include the indicator indicative of the suitability of the dose order to be prepared in a final container (e.g., expressly or by reference to a specific ingredient name in the dose order), the dose order will be prepared a new or empty container 448. That is, absent a positive indication of the suitability of the dose order for preparation in a final container, the selected dose order may not be prepared in a final container. This may be the case, for example, where the dose order is specifically indicated for preparation in a new or empty container. Upon the positive identification of the final container indication of the dose order, the pharmacy workflow application 114 may proceed to logical step 408. Otherwise, the pharmacy workflow application 114 may proceed to logical step 448 such that the dose order is prepared in the empty container.

In this regard, pharmacy workflow application 114 may execute a logical step 408 to determine whether the dose order is to include at least one diluent. That is, for a dose order record to be eligible to undergo the dynamic final container determination, at least one of the plurality products associated with preparation may correspond to a diluent product. Accordingly, it may be that only a product container including a diluent may be selected as a final container. That is, absent a diluent container being required for preparation of the dose order, the selected product container may not be used as a final container. As such, diluent-free dose orders may be prepared in a empty or new container 448. Upon the positive identification of a diluent in the dose order, however, the pharmacy workflow application 114 may proceed to logical step 412. Otherwise, the pharmacy workflow application 114 may proceed to logical step 448 such that the dose order is prepared in the empty container.

In this regard, pharmacy workflow application 114 may execute a logical step 412 to determine whether the dose order is associated with a formulary product having a data attribute corresponding to an indicator indicative of a “hazmat” drug. This may be the case, for example, where the dose order includes a drug intended for use in a chemotherapy treatment. Accordingly, a dose order associated with a hazmat drug may not be suitable for preparation in a final container. Rather, a dose order including a hazmat drug may be prepared in a new or empty container 448. In some embodiments, logical rule 412 may be configurable based on site-specific preferences. For example, the hazmat drug classification may be modified to include specific types of hazmat drugs or to distinguish between one category of hazmat drugs which may be prepared in a final container, and another category of hazmat drugs which may not be prepared in a final container. Furthermore, each product in a formulary may comprise an indication of either a hazmat drug or not a hazmat drug. Upon the positive identification of the suitability of the dose to undergo the final container determination based on a hazmat drug indicator, the pharmacy workflow application 114 may proceed to logical step 416. Otherwise, the pharmacy workflow application 114 may proceed to logical step 448 such that the dose order is prepared in the empty container.

In this regard, pharmacy workflow application 114 may execute a logical step 416 to determine whether the dose order is associated with a pre-populated “QS Drug” field equal to a diluent line item in the dose order. In this regard, the QS Drug field must equal the diluent line item in the dose order in order for the selected product to constitute a suitable final container. 448. That is, if the dose order includes a drug that requires a different carrier (*i.e.*, a QS Drug), than the diluent ingredient in the dose order to fulfill the dose, the dose may be prepared in to an empty container. As such logical step 416 verifies that the product contains the request volume of product to satisfy the requirements of the dose order. Absent this equality, the dose order will be prepared

in an empty or new container, absent any addition adjustments to the product (*e.g.*, volume addition or withdraw from the product). For example, a hypothetical preparation of Cefazolin 1000 mg, NS 250 ml in 250 ml, after scanning all products, may include a calculated QS volume of -10 ml (*i.e.*, inverse QS). In order to satisfy the dose order total volume (*i.e.*, final volume) 250 ml request, the pharmacy workflow application 114 will display a preparation step that instructs a user to remove 10 ml (QS volume) from the final container product. As such, the remove of the 10 ml may permit the 250 ml product to be used as a final container in the foregoing dose order example. Upon the positive identification of the noted equality, the pharmacy workflow application 114 may proceed to logical step 420. Otherwise, the pharmacy workflow application 114 may proceed to logical step 448 such that the dose order is prepared in the empty container.

The pharmacy workflow application 114 may execute a logical step 420 to determine whether the dose order comprises a total volume equal to or less than a minimum dose volume (*e.g.*, less than 5 ml, 10 ml, 24 ml, 30 ml, 35 ml etc.). If the total volume of the dose order is less than or equal to the minimum dose volume the dose may be prepared in an empty or new container 448. Accordingly, logical rule 420 may be configurable based on site-specific preferences. For example, the minimum dose volume may be selectable prior to dose preparation of a given dose. Upon the positive identification of the volume of the dose order exceeding the minimum dose volume, the pharmacy workflow application 114 may proceed to logical step 424. Otherwise, the pharmacy workflow application 114 may proceed to logical step 448 such that the dose order is prepared in the empty container.

In this regard, pharmacy workflow application 114 may execute a logical step 424 to determine whether the dose order includes a requirement for a diluent product associated with a formulary record data attribute corresponding to an indicator that the diluent product is suitable as

a final container. If the diluent product of the dose order is not indicated as being suitable as a final container (*e.g.*, as recorded in a formulary record corresponding to one diluent product), the dose order will be prepared a new or empty container 448. This may be the case, for example, where the diluent formulary product is in a stored in a vial and is indicated in a corresponding
5 formulary record as not being suitable as a final container. Upon the positive identification of the diluent final container indication, the pharmacy workflow application 114 may proceed to logical step 428. Otherwise, the pharmacy workflow application 114 may proceed to logical step 448 such that the dose order is prepared in the empty container.

In this regard, pharmacy workflow application 114 may execute a logical step 428 to
10 determine whether the dose order includes more than one indentified final container products. That is, if the dose order contains multiple products as indicated as being diluent products, any one of which could theoretically constitute a final container, the dose order will be prepared in a new or empty container 448. That may be due, in part, because only one diluent product may constituent the final container. Upon the positive identification of the existence of only one indicated diluent
15 product which may constitute a dynamically determined final container, the pharmacy workflow application 114 may proceed to logical step 432. Otherwise, the pharmacy workflow application 114 may proceed to logical step 448 such that the dose order is prepared in the empty container.

In this regard, pharmacy workflow application 114 may execute a logical step 432 to determine whether the selected product corresponding to the potential final container is not a
20 partially used diluent container. Accordingly, only a new or unused diluent product may constituent a final container. As such, where the indicated diluent formulary product corresponds to a partially used diluent formulary product (*e.g.*, a diluent formulary product used, in part, to prepare a prior dose order or a “work-in-progress” (“WIP”) product), the dose order may be

prepared in a new or empty container 448. Upon the positive identification that the selected product corresponds to a new or unused diluent formulary product, the pharmacy workflow application 114 may proceed to logical step 436. Otherwise, the pharmacy workflow application 114 may proceed to logical step 448 such that the dose order is prepared in the empty container.

5 In this regard, pharmacy workflow application 114 may execute a logical step 436 to determine whether the dose order satisfies a volume differential rule for a product being evaluated as a final container. That is, in order to satisfy the volume differential rule, the absolute difference between the diluent formulary product volume in its original container and the dose order total volume may not exceed a specified and predetermined differential percentage. The calculation of
10 the differential percentage may be represented according to the following expression: $[(\text{diluent product total volume} - \text{dose order total volume}) / (\text{dose order total volume})] * 100$. If the foregoing expression returns a value greater than or equal to the specified differential percentage, the dose order may be prepared in a new or empty container 448. Preparing the dose order in a new or empty container may also enable using the remaining volume of the indicated product for future
15 doses through assigning the product a work-in-process (WIP) label and designation in the formulary along with the appropriate expiration date as defined in the product formulary. In some embodiments, logical rule 436 may be configurable based on site-specific preferences. For example, the specified differential percentage may be selectable (*e.g.*, 50%, 75%, 100%, 125%, etc.). Upon the positive identification that the expression returns a value less than the specified
20 differential percentage, the pharmacy workflow application 114 may proceed to logical step 440. Otherwise, the pharmacy workflow application 114 may proceed to logical step 448 such that the dose order is prepared in the empty container.

 In this regard, pharmacy workflow application 114 may execute a logical step 440 to

determine whether the dose order satisfies the waste volume rule. That is, in order to satisfy the waste volume rule, the absolute difference between the dose order carrier ingredient volume and the diluent product total volume may not exceed a selectable and predetermined waste volume. That calculation of the waste volume may be represented according to the following expression:

5 (dose order carrier ingredient volume – diluent product total volume). If the foregoing expression returns a value greater than or equal to the waste volume, the dose order may be prepared in a new or empty container 448. Preparing the dose order in a new or empty container may also enable using the remaining volume of the product for future doses through assigning the product a work-in-process (WIP) label along with the appropriate expiration date as defined in the formulary data
10 record. In some embodiments, logical rule 440 may be configurable based on site-specific preferences. For example, the waste volume may be selectable (*e.g.*, 100 ml, 200 ml, 300 ml, etc.). For illustration, upon the positive identification that the expression returns a value less than 200 ml, the pharmacy workflow application 114 may determine that the selected product of the dose order may constitute the final container 444. Otherwise, the pharmacy workflow application 114
15 may proceed to logical step 448 such that the dose order is prepared in the empty container.

It will be appreciated that any or all of the foregoing logical rules may be applied in a multitude of use cases. That is, the logical rules may dynamically determine the final container in various circumstances based on the particular requirements of the initiated dose preparation protocol and to the received indication of the plurality of products to be used to prepare the dose.
20 In this regard, various use cases are presented for the sake of illustration below. The presented use cases are merely examples and should not be read to limit the applicability of the logical rules 400 to any particular use case or dose preparation protocol.

In this regard, with reference to Fig. 5, a user interface 500 illustrating an exemplary use

case may comprise dynamically determining a final container in a dose preparation protocol that includes a scenario wherein multiple diluent product formulary products suitable as a final container are indicated or scanned as part of a dose preparation protocol. For instance, the 0.9% Sodium Chloride 250 ml diluent product 504 and the 0.9% Sodium Chloride 500 ml diluent product 508 may both be utilized as a final container. In this instance, and with reference to the logical rule 428, neither of the scanned diluent formulary products may constitute the final container of the dose order. Rather, the dose order will be prepared in a new or empty container.

In this regard, with reference to Fig. 6, a user interface 600 illustrating another use case may comprise dynamically determining a final container in a dose preparation protocol that includes a negative QS volume with a final specified volume. A QS volume may reference the volume of diluent need to reconstitute the dose as described above in relation to Fig. 4b. With continued reference to the example in Fig. 6, a Cefazolin 1 g, 0.9% Sodium Chloride 200 ml in 200 ml dose order is specified 604. In this instance, a 250 ml bag of 0.9 % Sodium Chloride is scanned and dynamically determined to constitute the final container for the dose order 608. In this instance, the pharmacy workflow application 114 may calculate 1 negative QS volume of 60 ml based on the contributed diluent for the Cefazolin. With reference to logical rule 440, the waste volume of the dose order (*i.e.*, 60 ml) does not exceed a predetermined waste volume, in this case, 250 ml. As such, 60 ml of diluent is withdrawn from the foregoing 250 ml bag 612, such that the bag will constitute the final container after the addition of the Cefazolin drug.

In this regard, with reference to Fig. 7, a user interface 700 illustrating still another use case may comprise dynamically determining a final container in a dose preparation protocol that includes no diluent line item in the order. For example, and with continued reference to Fig. 7, the dose order may include a request for: "Cefazolin 250 mg in 10 ml," 704. In this regard, and with

reference to logical rule 408, because the dose order does not contain in indication corresponding to a diluent product, the dose order may be prepared in a new or empty container.

In this regard, another case may comprise dynamically determining a final container in a dose preparation protocol that includes a premix product. A premix product may include a
5 formulary product premixed with a drug and diluent to a specified concentration. In this regard, a dose order may, in some cases, comprise only a request for a premix product, and thus, the premix product need not be altered as part of a dose preparation protocol. Where the premix product is not altered (*i.e.*, the premix satisfied the dose order entirely), the premix product itself is the final container. In other instances, a dose order may comprise a request for medication which may be
10 prepared based at least in part on a premix product. Consider, with reference to Fig. 12, a user interface 1200 according to another use case, which displays a hypothetical dose order 1204 comprised of: Gentamicin 80 mg, 0.9 % Sodium Chloride 100 ml in 120 ml. In this regard, the pharmacy workflow application 114 may receive an indication corresponding to premix product 1208 comprised of a Gentamicin Sulfate in 0.9% Sodium Chloride Injection, 80 ml in 100 ml in
15 VIAFLEX Plus Container. In this regard, the premix product may be identified as the final container 1212 as noted by the encircled "FC." In this particular example, the premix product must be QS'd (*e.g.*, supplied with additional diluent) in order to reach the final volume specified of 120 ml, as depicted at 1216, showing 0.9% Sodium Chloride 20ml added to the premix product.

Turning next to Fig., 4b, the pharmacy workflow application 114 may adjusting the diluent
20 ingredient volume associated with the dose preparation protocol according generally to method 400b. In this regard, the method 400b may proceed generally in the following manner. Pharmacy workflow application 114 receives 404b a dose order awaiting preparation and initiates the dose order preparation protocol. The pharmacy workflow application may receive an indication, in the

form of a scan 408b of the dose order label corresponding to the required ingredients. For example, the received indication may correspond to a pharmacy technician taking possession of the product for use the preparation of the dose. Next, the preparation protocol is checked 412b for having a product selection method configured to verify product drugs and quantity. Upon a positive
5 indication of such, the pharmacy workflow application calculates 416b the total accumulated dose order volume of non-diluent ingredients. Next, the pharmacy workflow application 114 verifies 420b that the dose order is set for preparation in a final container. The system may then ascertain 424b whether the dose order is associated with a final specified volume. Notably, if the dose order does not both have final container product scanned and final volume of the dose order specified,
10 system will not request to determine a QS volume for the dose order 456b. If, however, the dose order does not have final container scanned, but the final volume of dose order is specified, the pharmacy workflow application 114 may QS 428b the dose order such that the QS volume is determined by reference to following expression: (Final Volume) – (Sum of the contributing dose volumes).

15 In some embodiments (*i.e.*, where the dose order is identified as being prepared in a final container), the pharmacy will ascertain 432b whether a reused dose order is selected for a final container. If so, the pharmacy workflow application 114 may not determine a QS volume for the dose order 456b. In other embodiments, the reused dose order may not be a final container for the dose order. In this regard, the dose order is checked to see whether the dose order is associated
20 with a final specified volume. As such, and according to step 436b, if reused dose order is not a final container for dose order, then the dose order is checked for having the final volume specified. If the final volume is not specified, system will request 436b to determine a QS volume for the dose order such that the QS volume is determined by reference to following expression: $[(-1) *]$

(total volume of the final container formulary product – dose order volume for the diluent ingredient)]. Conversely, if the reused dose order is not a final container for dose order and the final volume is specified, the pharmacy workflow application 114 may determine 440b a QS volume for the dose order such that the QS volume is determined by reference to following
5 expression: $(-1)*[(\text{sum of the contributing dose volumes that are not provided by the final container formulary product}) + (\text{total volume of the final container formulary product} - \text{dose order final volume})]$.

As a result of the pharmacy workflow application 114 determining that the QS volume is greater than zero, the pharmacy workflow application may perform one or more QS volume
10 sufficiency checks. Specifically, this may begin by pharmacy workflow application 114 determining 444b whether the dose order has a final volume specified. To the extent that the dose order does not have a final volume specified 448b, the dose order total volume may be determined by reference to the following equation: dose total volume = sum of accumulated volume by each dose order ingredients. However, if the dose order does have a final volume specified 452b, the
15 dose total volume is determined by reference to the following equation: dose total volume = specified dose order final volume.

Turning next to Fig. 4c, the pharmacy workflow application 114 may determine if the QS volume will be displayed in part by performing a QS volume sufficient check according generally to method 400c. Method 400 may proceed by initiating the dose preparation at step 404c. This
20 may correspond to the pharmacy workflow application 114 initiating the dose preparation protocol corresponding to the dose order. Next, the pharmacy workflow application 114 may receive an indication or a scan and prepare all required products for the dose order according to step 408c. The method 400c may continue at step 412c such that the product list is completed for a dose

order. The method 400c may continue at step 416c such that the user scans a dose order label to get the dose order procedure actions. The method 400c may continue at step 420c such that the QS volume is calculated (*e.g.*, by method 400b, as depicted in Fig. 4b). The method 400c may continue at step 424c such that the absolute value of the calculated QS volume is compared to the
5 minimum measureable volume (*e.g.*, according to the site configuration parameters). To the extent that the absolute volume of the calculated QS volume is less than the minimum measureable volume, the method 400c may continue at step 428c such that it is determined that the calculated QS volume is sufficient. In this regard the QS volume action will not be displayed in the dose procedure and the method 400c may conclude at step 432c such the preparation of the dose
10 procedure is complete.

However, with returned reference to step 424c, to the extent that the absolute value of the calculated QS volume is greater than the minimum measurable volume, the method 400c may continue at step 436c such that the absolute value volume is compared to the maximum volume deviation (*e.g.*, according to the site configuration parameters). To the extent that the absolute
15 value of the calculated QS volume is greater than the maximum volume deviation, the method 400c may continue at step 440c such that the pharmacy workflow application 114 determines that the calculated QS volume is not sufficient. In this regard, the dose preparation actions are displayed at step 432c. However, to the extent that the absolute value of the calculated QS volume is less than the maximum volume deviation, the method 400c may continue at step 444c such that
20 the pharmacy workflow application 114 calculates the deviation percentage according to the following expression: $(\text{calculated QS volume} / \text{dose total volume}) * 100$. The method 400c may continue at step 448c such the absolute value of the calculated QS volume is compared to a maximum rounding error percentage (*e.g.*, according to site configuration parameters). To the

extent that the absolute value of the calculated QS volume is greater than the maximum rounding error percentage, the method may continue at step 440 such the pharmacy workflow application 114 determines that the calculated QS volume is not sufficient. In this regard, the dose preparation actions are completed at step 432c. If, however, the absolute value of the calculated QS volume is less than the maximum rounding error percentage, the method 400c may continue at step 428c, such that the pharmacy workflow application 114 determines that the QS volume is sufficient such that the QS volume will not be displayed in the dose procedure. In this regard, the method 400c concludes at step 432.

Turning next to Fig. 4d, as previously noted, the pharmacy workflow application 114 may identify the currently active workstation products for the dose order in the particular dose order as part of the received indication according generally to method 400d. The method 400d may begin at step 404d such that the pharmacy workflow application 114 initiates the dose preparation protocol corresponding to the dose order. The method may continue at step 408d such that the system requires the user to scan a product label associated with the dose order. As such, step 408d may be performed relative to the foregoing received indication of a plurality of products associated with the dose preparation protocol. The method 400d may continue at step 426d such the labels of the plurality of products are scanned. The method 400d may continue by the pharmacy workflow application 114 determining if there are any active reuse dose orders at the workstation. If so, the method 400d may continue at step 432d by identifying active contributing reused doses containing drug ingredients from the workstation. This may be accomplished by sorting the active reused dose orders by dose order item count in ascending order. Regardless of any active reuse dose orders are present at the workstation, the method 400d may continue at step 436 by pharmacy workflow application 114 determining whether there are any active products at the workstation.

To the extent that there are active products at the workstation, the method 400d may identify active contributing products from the workstation.

Specifically, the pharmacy workflow application 114 may identify active contributing products from the workstation by executing step 440d of method 400d such that the active products
5 are sorted by product drug count in descending order. The method 400d may continue at step 444d such that the active products are sorted by activation date time with the earliest time being ranked first. The method 400d may continue at step 448d such the active drug products are sorted. The method 400d may continue at step 452d such that active products are sorted by activation date time with the earlier time being ranked first. The method 400d may continue at step 456d such
10 the active diluent products are sort according to the identified final container indication. The method 400d may continue at step 460 such that the active products are sorted by activation date time with the earlier time being ranked first. The method 400d may continue at step 464d such that the active diluent products are sorted according to the identified non-final container indication. The method 400d may continue at step 468d such the active products are sorted by activation date
15 time with the earlier time being ranked first. The method 400d may continue at step 472d, wherein the pharmacy workflow application 114 may determine if there are any active reuse dose orders at the workstation. To the extent that active reuse dose orders are at the workstation, the pharmacy workflow application 114 may identify active contributing reused dose order containing at least one diluent ingredient from the workstation. This may be accomplished by sorting the active
20 reused dose orders by dose order items count in ascending order.

The method 400d may continue at step 480d such that all active contributing products from the workstation are identified. With returned reference to step 436d, to the extent that there are not any active products at the workstation, the method 400d may also proceed to step 480d. In

either event, the method 400d may proceed to step 412d, such the pharmacy workflow application 114 determines if the product list is completed for the dose order. To the extent that the product list is not completed for the dose order, the method 400d may continue at previously discussed step 408d. However, to the extent that the product list is complete for the dose order, the method 400d may continue at step 416d such that the label of the dose order is scanned. The method 400d may continue at step 420d such that the dose procedure actions are completed. The method may continue at step 424d such that the dose preparation is completed.

Turning next to Fig. 4e, in other embodiments, the pharmacy workflow application 114 may execute a series of steps, according generally to method 400e, associated with an embodiment of the dynamically determined final container determination of the logical steps 400 depicted in Fig. 4. In this regard, method 400e may begin by initiating the dose order preparation. In some instances, this may correspond to initiating a dose preparation protocol corresponding to the dose order. The method 400e may continue at step 408e such that pharmacy workflow application 114 determines if the dose order will be prepared via a preparation mode that will verify the product drugs and quantity. To the extent that the dose order will not be prepared via a preparation mode that will verify the product drugs and quantity, the dose order may be prepared in an empty or new container 432e. However, to the extent that the dose order will be prepared via a preparation mode that will verify the product drugs and quantity, the method 400e may continue at step 412e such that the pharmacy workflow application 114 may determine if a reuse dose is active at the workstation. To the extent that the reuse dose is active at the workstation, the method 400e may continue to step 416e such that the pharmacy workflow application determines if the reused dose satisfies a complete dose. To the extent that the reused dose satisfies a complete dose the method 400e may continue at step 420e such that the final container is determined to be the container of

the reused dose. If, however, and with returned reference to steps 412e and 416e, the reused dose does not satisfy a completed dose or there is no reuse dose active at the workstation, the method 400e may proceed to step 424e such that the pharmacy workflow application 424e may determine if the dose order is partially prepared by a compounder. To the extent that the dose order is partially
5 prepared by the compounder, the method 400e may continue at step 428e such the final container is determined to be the compounder bag. If, however, the dose order is not partially prepared in a compounder the method 400e may continue according to the method 400.

Turning next to Fig. 8, according to another embodiment, the pharmacy workflow
10 application 114 may facilitate dynamic diluent volume accounting in a dose preparation protocol according generally to the method 800. In this regard, the diluent volume may be dynamically adjusted based on the receipt of an indication of a product associated with a dose preparation protocol. Accordingly, the pharmacy workflow application may display to a user a precise amount of required diluent volume for a given dose order selected for preparation. And in some
15 embodiments, the required diluent volume may be fulfilled by multiple separate diluent products such that the additive volume amount of the multiple separate diluent product containers is at least equal to the dynamically determined required diluent volume.

In some instances, the preparation of dose orders with dynamic diluent volume accounting may facilitate efficient dose order preparation techniques and contribute to overall error reduction.
20 In one embodiment, the dynamic diluent volume account allows the pharmacy workflow application 114 to receive an indication (*e.g.*, a scan) associated with multiple bag of the same diluent ingredient to prepare the required diluent volume for the dose preparation protocol. The received indication may establish the presence, for example, of the diluent products required to

prepare the dose at a preparation workstation. In this regard, a dose order may be prepared from multiple sources of diluent. This aspect may increase efficiency by facilitating a workflow that allows a user to prepare a dose order with various volumetric combinations of identical diluent products. Moreover, other aspects of the dynamic diluent volume accounting may reduce dose preparation errors. For example, according to another embodiment, the pharmacy workflow application 114, via the user interface 108, may display a running total of the volume of the diluent product required according to one or more preparation steps, which demonstrates the physical steps required to complete the dose preparation. In this regard, errors associated with using the wrong volume of diluent during dose preparation may be reduced.

The dynamic alteration of the volume includes the pharmacy workflow application 114 accessing 804 a dose order record corresponding to a dose order stored in, for example, local database 112. The dose order record may correspond to a dose order received by the pharmacy workflow application 114 that has not yet been prepared. Furthermore, the dose order may include a request for medication that is awaiting preparation in the pharmacy workflow application 114. In some instances, the corresponding dose order record of the dose order may be stored in a queue comprised of a plurality of dose order records with corresponding dose orders also awaiting preparation. In either event, in this embodiment, the pharmacy workflow application accessed a single dose record.

The pharmacy workflow application 114 may receive 808 an indication of a product associated with a dose preparation protocol. The dose preparation protocol may include, among other things, a list of ingredients or a “recipe” which communicates the constituent components of a dose order. In this embodiment, the preparation protocol may include at least one drug product and at least one diluent product. The at least one drug product may correspond to a pharmaceutical

drug or active medication ingredient for therapeutic effect. The at least one diluent product may correspond to a reconstitution solution for use in diluting the concentration of the product. In this regard, a dose order may specify a precise amount of product (*i.e.*, medication to be delivered) volume, and require the diluent to be added to the product in order to reach a final volume of the
5 dose order, for instance. The diluent added in this regard, may constitute a QS volume, or a volume of diluent (in ml) added to a product to complete the dose order. The QS volume, as discussed in more detail below, may change depending on a variety of factors, including the initial product concentration, requested final volume, and the like.

The received indication of the product associated with the dose preparation protocol may
10 facilitate the pharmacy workflow application 114 dynamically determining 812 a required diluent volume based on the received indication. That is, the indication may include data attributed of the drug corresponding to the drug product. The data attributes may include, among other things, the concentration and/or volume of the formulary drug product. Moreover, this indication may be received by the pharmacy workflow application 114 in the context of the dose preparation protocol
15 and associated dose order. In any event, the foregoing defined QS volume may be dynamically determined based on the data attributes of the indication.

It will be appreciated that the dynamic determination of the QS volume may be application in a variety of cases based on various data attributes. For the sake of illustration, the following discussion highlights the pharmacy workflow application 114 dynamic alteration of the QS volume
20 based on two specific data attributes: final volume of the dose order, and initial volume of the formulary drug product. Notably, the dynamic determination is not limited to either of the following two data attributes.

In this regard, with brief reference to Fig. 9, the QS volume may be dynamically determined

based on the final volume specified in the dose order as generally indicated by user interface 900. For example, a dose order may include a dose 904 of: Cefazolin 1 g, 0.9% sodium Chloride 200 ml in 200 ml. As such, the dose order of the example specified a final volume 908 of 200 ml. The received indication corresponding to a formulary record corresponding to the Cefazolin drug product notes a Cefazolin dose at a concentration of 100 mg/ml (not pictured). The dynamic determination will use this information to determine that the Cefazolin drug will constituent 10 ml of the 200 ml final volume (*i.e.*, $1000 \text{ mg} / (100 \text{ mg/ml}) = 10 \text{ ml}$) 912. As such, the pharmacy workflow application 114 may determine that the required diluent volume 916 for this particular dose order equate 190 ml (*i.e.*, $200 \text{ ml} - 10 \text{ ml} = 190 \text{ ml}$). For the sake of continued illustration, consider that the required diluent volume would change in the event that the final volumes specified in the dose order were different. That is, based on the logic of the foregoing determination, a Cefazolin 1 g, 0.9% sodium Chloride 200 ml in 300 ml dose, would result in a dynamic determination of a required diluent volume of 290 ml, assuming the received indication of the formulary drug product is the same as that above (*i.e.*, $1000 \text{ mg} / (100 \text{ mg/ml}) = 10 \text{ ml}$; $300 \text{ ml} - 10 \text{ ml} = 290 \text{ ml}$).

In this regard, the QS volume may be dynamically determined based on the concentration of the indicated formulary record for a drug product received in connection with the dose preparation protocol of the dose order. For example, a dose order may include the dose of: Cefazolin 1 g, 0.9% sodium Chloride 200 ml in 200 ml. As such, the dose order of the example specified a final volume of 200 ml. The received indication corresponding to a formulary drug product may note a Cefazolin dose at a concentration of 50 mg/ml. The dynamic determination will use this information to determine that the Cefazolin drug will constituent 20 ml of the 200 ml final volume (*i.e.*, $1000 \text{ mg} / (50 \text{ mg/ml}) = 20 \text{ ml}$). As such, the pharmacy workflow application

may determine that the required diluent volume for this particular dose order equate 180 ml (*i.e.*, 200 ml – 20 ml = 180 ml).

With return reference to Fig. 8, the previously noted interface 108 of local system 110 may also be operable to display 816 a dose preparation step associated with the preparation of the dose relative to the diluent volume. That is, the preparation step of the preparation protocol may be displayed at least partially based on the dynamically determined diluent volume. In this regard, the pharmacy workflow application 114 may communicate to a user the dynamically determined diluent volume such that the user may incorporate the dynamically determined diluent volume in the preparation of the dose order.

In some embodiments, the pharmacy workflow application 114 may receive one or more indications corresponding to the required diluent volume. In this regard, the pharmacy workflow application 114 may require the identification of the particular diluent product intended for use in the preparation of the dose order. Notably, in this embodiment, the required diluent volume may be satisfied by one or more diluent products. Each diluent product includes a diluent product volume. The pharmacy workflow application 114 may determine the diluent product volume of each diluent product in order to verify that the aggregate diluent formulary product volume is at least equal to the required diluent volume. In this regard, multiple diluent formulary products may be indicated in order to satisfy the dynamically determined required diluent volume.

The pharmacy workflow application 114 may, in the context of receiving an indication of the one or more diluent formulary products, display a running total of the volume of the diluent product yet needed to meet the required diluent volume. In this regard, the pharmacy workflow application 114 may communicate to the user the remaining volume of diluent formulary product required to satisfy the dynamically determined diluent volume. For example, a give dose order

may require 300 ml of 0.9 % Sodium Chloride. Upon receipt of an indication of a diluent formulary product of 0.9% Sodium Chloride in a 250 ml bag, the pharmacy workflow application may indicate to the user that 50 ml are required to satisfy the required diluent volume (*i.e.*, 300 ml – 250 ml = 50 ml). The remaining volume indicator may appear on the user interface 108
5 associated with the pharmacy workflow application 114. The user interface 108 may also include a collapsible area for selectively displaying information associated with the one or more indications of the diluent formulary product. In this regard, the one or more diluent formulary products, appearing on the user interface as space line items, may be collapsed into a single line that is indicative of the group of indicated diluent formulary products. Accordingly, the collapsible
10 areas may facilitate organizing information such that the user interface 108 displays information to the user pertinent to the particular dose preparation protocol step of the dose order record (*e.g.*, the collapsible area may hide the various line items of the one or more diluent formulary products after the required diluent volume has been reached).

The foregoing indications may be displayed by the pharmacy workflow application 114
15 via user interface 108. User interface 108 may be configurable to present various indicators at least partially based on the received first indication of a product associated with a dose preparation protocol relative to the dynamic diluent volume accounting. With collective reference to Figs. 8a, 8b, and 8c, the configuration may resemble one or more configurations 800, 804, and 808. With specific regard to configuration 800 of Fig. 8a, user interface 108 is operable to display running
20 total 816 indicative of the volume of the diluent product yet needed to meet the required diluent volume. The running total 816 may be calculated on the basis of one or more received indications of a product associated with a dose preparation protocol disposed on configuration 800 in region 820. For example, as illustrated in configuration 800, the dose preparation protocol required 7.5

ml of diluent Sodium Chloride (not pictured). Configuration 800, in this example, includes a first indication 824 of a product associated with a dose preparation protocol, in particular, corresponding to 0.9% Sodium Chloride in a 10 ml vial contributing 4 ml. In this regard, running total 816 indicates that 3.5 ml are yet required to meet the required diluent volume. As such, the pharmacy workflow application 114 may receive another indication 828, with reference to configuration 804 of Fig. 8b, to satisfied the running total amount. The user interface 108 may also include a collapsible area 832 for selectively displaying information associated with the indicated products. Notably, the information of the collapsible area may be temporality hidden from view upon selecting the collapsible area, as generally depicted by collapsed collapsible area 836 of configuration 812 of Fig. 8c. The collapsible area may facilitate dose order preparation by temporarily hiding information from the user when the information is not longer directly relevant to a particular step of the dose preparation protocol.

In another embodiment, the pharmacy workflow application 114 may determine whether an indicated diluent product is associated with a list of allowed diluent products relative to the dose preparation protocol. That is, various diluents may appropriately be used (*i.e.*, allowed diluents) in the preparation of a given dose preparation protocol (*e.g.*, Sodium Chloride, etc.). The pharmacy workflow application 114 may display a dose preparation step at least partially based on a received indication of a diluent product associated with the list of allowed diluent products of the dose preparation protocol. As discussed in greater detail above, in some cases, the pharmacy workflow application 114 may receive multiple indications of diluent products intended to satisfy the required diluent volume. In this regard, upon the receipt of one indication of a diluent product, the required diluent volume must be satisfied by subsequent indications associated with the same diluent product. In this regard, after an indication of one diluent product, rather than determine if

a subsequent diluent product is associated with a list of allowed diluent products of the dose preparation protocol, the pharmacy workflow application 114 may determine if the subsequent diluent product is the same diluent product as the first indicated diluent product. If any subsequently indicated diluent product is different than the first indicated diluent product, the subsequently indicated diluent product will not be able to satisfy the required diluent volume.

Turning next to Fig. 10, according to another embodiment, the pharmacy workflow application 114 may facilitate resolving critical issues associated with the pharmacy workflow management application 114 according generally to the method 1000. In this regard, the pharmacy workflow application 114 may be operable to identify various critical issues and display the identified critical issues in a common location to facilitate issue resolution. The identified critical issues may include various errors, such as an error regarding a dose order record, a password error, or other miscellaneous errors, as described in greater detail below. In this regard, according to one embodiment, errors associated with the dose order record may be identified and, in some instances, resolved within the pharmacy workflow queue. Accordingly, a dose order record with an error need not necessarily be prepared outside of the pharmacy workflow application.

In some instances, managing an error regarding a dose order in a pharmacy workflow queue may facilitate efficient dose order preparation techniques and contribute to error mitigation. Notably, in one embodiment, the pharmacy workflow application 114 may support user-resolution of errors within the pharmacy workflow application 114 framework. That is, an error associated with a dose order need not preclude the dose order from preparation via the pharmacy workflow application 114. Rather, the pharmacy workflow application 114 may identify the error and, in some cases, accept an input from an user indicative of a resolution of the identified error. In this regard, dose order efficiency may increase by facilitating the preparation of dose orders initially

associated with errors. Furthermore, managing errors in this regard may also mitigate the occurrence of additional errors. This may occur by providing a combined critical issues display (*i.e.*, the Attention Tab, discussed in greater detail below) by which users may learn of various types of errors. Moreover, dose orders with identified errors may not be reintroduced in to the dose order queue without verifying that the identified error has been resolved. In this regard, a
5 dose order with an error is not introduced in to the dose order queue.

Managing an error of a dose order record includes the pharmacy workflow application 114 accessing 1004 a dose order record corresponding to a dose order stored in, for example, local database 112. The dose order record may correspond to a dose order received by the pharmacy
10 workflow application 114 that has not yet been prepared. Furthermore, the dose order may include a request for medication that is awaiting preparation in the pharmacy workflow application 114. In some instances, the corresponding dose order record of the dose order may be stored in a queue comprised of a plurality of dose order records with corresponding dose orders also awaiting preparation. In either event, in this embodiment, the pharmacy workflow application accesses a
15 single dose record.

The pharmacy workflow application 114 may execute a first processing 1008 sequence in order to generate an alert corresponding to an error state related to the dose order. The error state, as will be discussed in more detail below, may correspond to a plurality of potential errors of the dose order records. The potential errors may include, for example, parsing errors, unknown unit
20 errors, unknown product errors, unknown route errors, and the like. The first processing 1008 may include attempting to initiate a dose preparation protocol such upon initial of the dose preparation protocol the pharmacy workflow application identifies the error state.

The previously noted interface 108 of local system 110 may also be operable to display

1012 a dose order in corresponding relation to the alert. That is, for example, the dose order may be displayed to the user in a manner that communicates to the user that the dose order is associated with an error state. The displayed dose order associated with the error state may, in some instances, be displayed with a plurality of other dose orders also associated an error state. The plurality of
5 dose order associated with an error state may be grouped according to error type via one or more tabs, and highlighted with a color indicative of the error state. In this regard, the user may view a queue of dose orders such that each dose order of the queue is associated with an error state. The ability to view a plurality of dose orders in a queue in this manner may facilitate resolution of errors across the plurality of dose orders associated with errors. In some instances this queue may
10 be disposed in an “Attention Tab,” as depicted in Fig. 11, of the pharmacy workflow application 114, as discussed in more detail below.

The Attention Tab may facilitate the resolution of the errors associated with the dose order via Attention Tab user interface 1100. In particular, the pharmacy workflow application 114 may be operable to receive 1016 an input from a user indicative of a resolution of an error. For example,
15 the input may correspond to a user-entered route to resolve an error of a dose order associated with an unknown route. In this regard, the Attention Tab 1100 may provide the user with the ability to remedy the error of the dose order record.

Furthermore, the pharmacy workflow application 114, via the attention tab 1100, may facilitate a second processing 1020 of the dose order based on the received input. In this regard,
20 the pharmacy workflow application 114 may verify that the error of the dose order record has been appropriately resolved. For example, a user-entered route may be verified to ensure that that dose order record is no longer associated with an unknown route error. The unknown route error may still exist, for example, where the user entered the wrong information for the route. As such, the

second processing 1020 may help ensure that a dose transferred from the Attention Tab 1100 is not subsequently sent back to the attention tab 1100 with additional errors – *i.e.*, only error-free dose order records may leave the attention tab. In this regard, the pharmacy workflow application 114 may requeue 1024 the dose order record upon confirmation of the successful resolution of the error of the dose order record.

As noted, the pharmacy workflow application 114 may be operable to generate an alert corresponding to a plurality of error states related to the dose order. For the sake of illustration, several error states associated with the foregoing attention tab 1100 are presented below. It will be appreciated that the several error states of the attention tab 1100 are not limiting, and that the pharmacy workflow application 114 may be operable to generate an alert corresponding to a plurality of other error states of the dose order record.

In this regard, the pharmacy workflow application 114 may be operable to generate an alert corresponding to an error state related to a parsing error. A dose order record associated with a parsing error may be disposed on a parsing errors tab 1104 of the attention tab 1100. A parsing error may include an error where, as a result of extracting the dose order from an independent data stream (*e.g.*, an HL7 data stream), various aspects of the alpha numeric characters became unintelligible. In other instances, a parsing error may correspond to a dose order with incorrectly inputted alpha numeric characters at the time of creating the dose order – *i.e.*, misspelling the name of a formulary drug product. In any event, a parsing error results at least when the pharmacy workflow application 114 cannot recognize one or data fields or categories of the dose order record (*e.g.*, the pharmacy workflow application 114 cannot determine which portion of a string of alpha numeric characters corresponds to a drug field – the drug field cannot be identified by reference to the string). This contrasts with other error types, discussed in greater below, such an unknown

drug error, where the pharmacy workflow application 114 may recognize the drug field, but not the contents of the field. Upon generating an alert corresponding to the parsing error, the user may view the parsing error.

In another aspect, the pharmacy workflow application 114 may be operable to generate an alert corresponding to an error state related to an unknown drug error. A dose order record associated with an unknown drug error may be disposed on an unknown drug error tab 1108 of the attention tab 1100. In this regard, the pharmacy workflow application 114 may identify a drug field for the dose order with an unrecognized value of data. As such, an unknown drug error may include an error where the drug type associated with the dose order record is unrecognizable by the pharmacy workflow application 114. The determination of the unknown drug error may be accomplished with reference to a formulary database comprised of a plurality of formulary data records corresponding to various products. If the drug type associated with the dose order record is not related to a similar formulary data record, the pharmacy workflow application 114 may generate an alert corresponding to the error state of the unknown drug error. Upon generating an alert corresponding to the unknown drug error, the user may view the unknown drug error or take steps to resolve the error. For example, the user may update the formulary record database by creating a new product definition consistent with the product information of the dose order record. In this regard, upon second processing, the pharmacy workflow application 114 may identify the newly created formulary record such that the dose order record is no longer associated with an unknown drug – *i.e.*, the error was resolved.

In this regard, the pharmacy workflow application 114 may be operable to generate an alert corresponding to an error state related to an unknown route error. A dose order record associated with an unknown route may be disposed on an unknown route tab 1112 of the attention tab 1100.

In this regard, the pharmacy workflow application 114 may identify a route field for the dose order with an unrecognized value of data. As such, an unknown route error may include an error where the route type (*i.e.*, how the drug is administered to a patient) associated with the dose order record is unrecognizable by the pharmacy workflow application 114. The determination of the unknown route error may be accomplished with reference to the formulary database. If the route type associated with the dose order record is not related to a similar formulary data record, the pharmacy workflow application 114 may generate an alert corresponding to the error state of the unknown route error. Upon generating an alert corresponding to the unknown route error, the user may view the unknown route error or take steps to resolve the error. For example, the user may update the formulary record database by creating a new route definition in an existing formulary record consistent with the route information of the dose order record. In this regard, upon second processing, the pharmacy workflow application 114 may identify the newly updated formulary record that contains the new route such that the dose order record is no longer associated with an unknown drug – *i.e.*, the error was resolved.

In this regard, the pharmacy workflow application 114 may be operable to generate an alert corresponding to an error state related to an unknown unit error. A dose order record associated with an unknown unit error may be disposed on an unknown unit errors tab 1116 of the attention tab 1100. In this regard, the pharmacy workflow application 114 may identify a unit field for the dose order with an unrecognized value of data. As such, an unknown unit error may include an error where the unit type (*i.e.*, the unit of measurement of the various products of the dose order) associated with the dose order record is unrecognizable by the pharmacy workflow application 114. Upon generating an alert corresponding to the parsing error, the user may view the unknown unit error.

In this regard, the pharmacy workflow application 114 may be operable to generate an alert corresponding to an error state related to an unknown products error. A dose order record associated with an unknown products error may be disposed on an unknown products errors tab 1120 of the attention tab 1100. In this regard, the pharmacy workflow application 114 may identify
5 a products field for the dose order with an unrecognized value of data. As such, an unknown products error may include an error where the product type (*e.g.*, a diluent) associated with the dose order record is unrecognizable by the pharmacy workflow application 114. The determination of the unknown product error may be accomplished with reference to a formulary database comprise of a plurality of formulary data records corresponding to various products. If
10 the product type associated with the dose order record is not related to a similar formulary data record, the pharmacy workflow application 114 may generate an alert corresponding to the error state of the unknown product error. Upon generating an alert corresponding to the unknown drug error, the user may view the unknown drug error or take steps resolute the error. For example, the user may update the formulary record database by creating a new product definition consistent
15 with the product information of the dose order record. In this regard, upon second processing, the pharmacy workflow application 114 may identify the newly created formulary record such that the dose order record is no longer associated with an unknown product – *i.e.*, the error was resolved.

Other critical issues may be addressed by the pharmacy workflow management application 114 via the attention tab 110. For example, the pharmacy workflow application 114 may be
20 operable to generate an alert corresponding to an error state related to a passwords error. A dose order record associated with a passwords error may be disposed on a passwords errors tab 1124 of the attention tab 1100. A passwords error may include an error where a user account password is expired or where a user account is disable. An administrator-level user may view alerts associated

with used accounts with expired passwords or that are otherwise disabled. In some embodiments, the pharmacy workflow application 114 may accept an input from an administrator-level user indicative of a resolution of the expired password alert or the account disablement alert.

The pharmacy workflow application 114 may also be operable to generate an alert
5 corresponding to an error state related to a suspected discontinued dose error. A dose order record associated with a suspected discontinued dose error may be disposed on a suspected discontinued dose errors tab 1128 of the attention tab 1100. A suspected discontinued dose error may include an error where the dose order record is suspected of being associated with a discontinued dose order. For example, the previously inputted dose order may no longer be required in the hospital
10 setting in which it was originally order. As such, the dose order may be cancelled or discontinued. According, the pharmacy workflow application may 114 may associate an alert with the identified suspected discontinued dose order. Upon generating an alert corresponding to the suspected discontinued dose error, the user may view the suspected discontinued dose error. In some embodiments, the pharmacy workflow application 114 may be operable to delete the suspected
15 discontinued dose order in response to a user input indicative of a resolution of the suspected discontinued dose error.

In this regard, the pharmacy workflow application 114 may be operable to generate an alert corresponding to an error state related to a suspected duplicated error. A dose order record associated with a suspected duplicated error may be disposed on a suspected duplicated errors tab
20 1132 of the attention tab 1100. A suspected duplicated error may include an error where the where the dose order record is suspected of being associated with a duplicate dose order. For example, a dose order may have inadvertently been inputted into the system twice. In such an instance, the duplicate dose order is not required and need not be prepared. According, the pharmacy workflow

application may 114 may associate an alert with the identified suspected duplicate dose order. Upon generating an alert corresponding to the suspected duplicate dose error, the user may view the suspected duplicate dose error. In some embodiments, the pharmacy workflow application 114 may be operable to delete the suspected duplicate dose order in response to a user input
5 indicative of a resolution of the suspected duplicate dose error.

In this regard, the pharmacy workspace application 114 may provide functionality associated with identifying suspect duplicate dose order and suspect discontinued dose orders as described in U.S. Provisional Patent Application No. 61/975,519 entitled "MANAGEMENT OF MEDICATION DOSE ORDERS" filed on April 4, 2014, which is co-owned with the present
10 application and is incorporated by reference herein in its entirety. In this regard, the pharmacy workflow application 114 may execute a duplicate order detection rule, such that the pharmacy workflow application 114 may be operative to review a first dose order record in view of other dose order records within the dose order queue to determine if the first dose order record corresponds to a potential duplicate dose order. In this regard, the pharmacy workflow application
15 114 may evaluate the metadata regarding the first dose order record to determine if other ones of the dose order records within the dose order queue match the identified metadata regarding the dose order to a predetermined correspondence. In some instances, the predetermined correspondence may be selectable by a user. It may be appreciated that the matching of the metadata between the dose order record being reviewed in the other dose order records in the queue
20 may not necessarily need to be identical. For example, the rule may be written such that if selected ones of the metadata fields are identical between the first dose order record being reviewed and the other dose order records in the dose order queue, the dose order record being reviewed may be flagged is a potential duplicate order. Such a rule may be applied to the first dose order record

when the record is initially populated the dose order queue.

In other embodiments, the pharmacy workflow application 114 may process incoming dose orders to determine if the order corresponds to a discontinuation of an order. That is, a common practice when processing dose orders is to indicate a discontinuation of a first order by subsequently sending a corresponding order to the first order at a later time with a discontinued status for the subsequently provided order. In prior approaches, the receipt of such a discontinued order may simply result in printing a label with the dose order details and a discontinuation status. This would in turn require a user to go through the printed dose order labels to locate the original dose order that was referenced in the discontinued dose order in order to replace the original dose order on a discontinued status. The pharmacy workflow application 114 may be utilized to automate this process pharmacy workflow application 114 by identifying receipt of a dose order in the dose order queue that contains a discontinued status and automatically matching the discontinued dose to one or more existing dose order records. Thus, receipt of a dose order with a discontinued status may trigger the pharmacy workflow application 114 to perform a query of the dose order queue to identify corresponding ones of the dose orders in the dose order queue that exceed a predetermined correspondence to the discontinued dose order. For example, one or more overlapping or identical pieces of metadata between the discontinued dose order and the identified discontinued order in the dose order queue may be determined. The predetermined correspondence may be customizable by a user. In any regard, upon identification of a dose order record from the dose order queue that corresponds to a received discontinued dose order, the original dose order may automatically change the status of the one or more identified dose order records to a discontinued status.

Turning next to Fig. 13, the pharmacy workflow application 114 may facilitate the

identification of alerts on a dose order of the dose record according generally to method 1300. Method 1300 may proceed by initiating the dose order initialization process 1304. This may correspond to the pharmacy workflow application 114 initiating the dose preparation protocol corresponding to the dose order. Next, the pharmacy workflow application 114 may obtain the XML backup string from the currently processed dose order 1308. The method 1500 may continue at step 1312 such that the dose status of the dose order is error and parse error data field on the dose order is non-empty. To the extent that the dose status of the dose order comprises an error value and the parse error data field of the dose order is nonempty, the method 1500 may continue at step 1316 such that pharmacy workflow application 114 adds an alert to the dose order indicative of a parsing error. However, to the extent that either the dose status of the dose order does not comprise an error, or the parse error data field of the dose order is empty, the method 1300 may continue at step 1320 such that the dose logic engine (“DLE”) initiates the dose logic engine execution sequence.

In this regard, regardless of whether the dose status of the dose order comprises an error, or the parse error data field of the dose order is nonempty, the method 1300 may proceed to step 1324 such that dose unit of the dose order is identified in the formulary. To the extent that the dose unit of the dose order is not identifiable in the formulary, the method 1300 may proceed to step 1328 such that pharmacy workspace application 114 adds an alert to the dose order record indicative of an unknown unit error. To the extent that a dose unit of the dose order is identifiable in the formulary, or the pharmacy workflow application 114 added an alert to a dose order item indicative of an unknown unit error, the method of 1300 may proceed to step 1332, such that pharmacy workflow application 114 may determine whether the dose drug of the dose order is identified in the formulary. To the extent that the dose drug of the dose order is not identified in

the formulary, the method 1500 may continue to step 1340 such that pharmacy workflow application 114 adds an alert to the dose order associated with an unknown drug error.

To the extent that the dose drug of the dose order is identified in the formulary, or the pharmacy workflow application 114 added an alert to the dose order indicative of an unknown
5 drug, the method 1300 may continue at step 1336 such that the pharmacy workflow application 114 may determine if the administration route of the dose order is found in the formulary. To the extent that the administration route of the dose order is not found in the formulary, the method 1300 may continue at step 1344 such that pharmacy workflow application 114 adds an alert to the dose order indicative of an unknown route. To the extent that an administration route of the dose
10 order is found in the formulary, or the pharmacy workflow application 114 added an alert to the dose order indicative of an unknown route, the method 1300 may proceed at step 1340 such that the pharmacy workflow application 114 determines if a dose order includes an unknown drug flag. If the dose order is includes an unknown drug flag, the method 1300 may proceed to step 1348 such that the pharmacy workflow application 114 may backup the dose order in a local server.

15 In this regard, method 1300 may proceed to step 1360 such that the dose order is successfully processed. If, however, the dose order is not enabled with the unknown drug flag, the method 1300 may continue at step 1352 such that pharmacy workflow application 114 determines whether there are any active alerts on the dose order or associated aspects of the dose order record. To the extent that there are active alerts on the dose order or dose order record, the method 1300
20 may continue at step 1356 such that all alerts added to the dose order or dose order record may be ignored. Regardless of whether there are any active alert on the dose order or dose order record, the method 1300 may continue at step 1360 such that the pharmacy workflow application 114 successfully processes the dose order.

In this regard, method 1300 may facilitate executing a dose order purge process (*e.g.*, after 45 days). In this regard, the method 1300 may continue at step 1364 such that the pharmacy workflow application 114 determines whether the dose order is purgeable. If the dose order is purgeable, the method of 1300 may continue at step 1368 such that the pharmacy workflow application 114 determines whether a dose order copy is stored in the local server. If the dose order copy is stored in the local server, the pharmacy workflow application 114 may remove the backup dose order from the local server.

Turning next to Fig. 14, the pharmacy workflow application 114 may facilitate the identification of alerts on a user account according generally to method 1400. Method 1400 may proceed upon the initiation of step 1404 such that the pharmacy workflow application 114 identifies an attempt by a user to log in to the pharmacy workflow application 114 via user interface 108, for example. The method 1400 may continue at step 1408 such that the pharmacy workflow application determines whether the password associated with the user account is expired. To the extent that the password associated with the user account is not expired, the method 1400 may continue at step 1412 such that the pharmacy workflow application 114 determines if the user account has been disabled in any regard. To the extent, that the user account has not been disabled, the method 1400 may continue at step 1416 such that the pharmacy workflow application 114 determines that the user successfully logged in to the pharmacy workflow application 114.

If, however, the password associated with the user account has been disabled, or the user account has otherwise been disabled, the method 1400 may continue at step 1420 such that the pharmacy workflow application 114 determines if the user account is a support user account. To the extent that the user account is a support user account, the method 1400 may proceed to step 1424 such that pharmacy workflow application 114 does not add any alert on the user account. To

the extent that the user account is not a support user account, the method 1400 may continue at step 1428 such that the pharmacy workflow application may determine if LDAP authentication is activated within the pharmacy workflow application 114. To the extent that LDAP authentication is activated within the pharmacy workflow application 114, the method 1400 may continue at step 1424 such that the pharmacy workflow application 114 does not add an alert to the user account. However to the extent that LDAP authentication is not activated within the pharmacy workflow application 114, the method 1400 may continue at step 1432 such that the pharmacy workflow application 114 determines if an alert of the same type has previously been added and active on this user account. To the extent that in alert of the same type has been previously added an active on this user account, the method 1400 may continue at step 1424 such that the pharmacy workflow application does not add an alert to the user account. However, to the extent that new alert of the same type has not been previously added and active on this user account, the method 1400 may continue at step 1436 such that the pharmacy workflow application 114 may at any alert on the user account indicative of in account locked status.

Turning next to Fig. 15, the pharmacy workflow application 114 may facilitate identification of alerts on scanned product barcodes according generally to a method 1500. Method 1500 may be comprised generally of steps associated with dose preparation 1501 and steps associated with product preparation 1502. In this regard method 1500 may begin at step 1504 such that the dose preparation is initiated. The method may continue according generally to dose preparation 1501 such that the method 1500 continues at step 1508 such that the pharmacy workflow application 114 prompts a user to scan a product. Notably, step 1508 may initiate the steps associated with product preparation steps 1502, discussed in greater detail below. The completion of steps generally associated with product preparation 1502 may facilitate the method

1500 to continue at step 1512 such that the product ingredient preparation is complete. The method 1500 may continue at step 1516 such that the pharmacy workflow application 114 determines whether the product list is completed for the dose order. To the extent that the product list is not completed for the dose order, the pharmacy workflow application 114 may return to step 1508
5 such that the pharmacy workflow application 114 prompts a user to scan a product. If, however, the product list is complete for the dose order, the method 1500 may continue by executing the steps generally associated with a dose procedure QS action 1518.

In this regard, the method 1500 may continue at step 1520 such that the dose procedure is displayed. The method 1500 may continue at step 1524 such that the pharmacy workflow
10 application 114 determines whether the dose order requires a positive QS (*e.g.*, diluent product). To the extent that the dose order requires a positive QS, the method 1500 may continue at step 1528 such that the user scans a diluent product for subsequent QS. The method 1500 may continue at step 1532 such that the pharmacy workflow application 114 determines whether the scanned diluent product is associated with an unknown barcode. To the extent that the scanned diluent
15 product is associated with an unknown barcode, the method 1500 may proceed at step 1536 such that the pharmacy workflow application 114 may add an alert associated with the scanned diluent product indicative of a new source product.

In this regard, the method 1500 may proceed with returned reference to step 1528 such that a user scans a diluent product for a subsequent QS determination. With returned reference to step
20 1532, to the extent that the scanned diluent product is not associated with an unknown barcode, the method 1500 may proceed to step 1540 such that the pharmacy workflow application 114 determines whether the scanned diluent product is associated with an inactive QS diluent product in the formulary. To the extent that the scanned diluent product is associated with an inactive QS

diluent product in the formulary, the method 1500 may proceed to step 1544 such that the pharmacy workflow application 114 may add an alert associated with the scanned diluent product indicative of an inactive formulary product.

In this regard, the method 1500 may proceed with returned reference to step 1528 such that
5 a user scans a diluent product for a subsequent QS determination. To the extent that the scanned diluent product is not associated with inactive QS diluent product in the formulary, the method 1500 may continue at step 1544 such that the dose procedure actions are completed. Notably, and with returned reference to step 1524 of method 1500, if the dose does not require a positive QS, the method 1500 may proceed to step 1544 too. In this regard, the method 1500 may proceed to
10 step 1548 such that the dose preparation is complete.

With returned reference to step 1508, the method 1500 may generally execute the steps associated with the product preparation 1502 such that a product may be scanned for product preparation at step 1552 after pharmacy workflow application 114 prompts a user to scan a product at step 1508. In some embodiments, step 1552 may initiate after a user starts the product
15 preparation at step 1550. The method 1500 may continue at step 1556 such that the pharmacy workflow application 114 determines whether the barcode associated with the scan product is unknown. If the barcode associated with the scan product is unknown, the method 1500 may continue at step 1560 such that pharmacy workflow application 114 adds an alert associated with the barcode of the scanned product indicative of a new product.

20 In this regard, the method 1500 may continue with returned reference to step 1552 such that a product may be scanned for product preparation. To the extent that the barcode associated with the scanned product is not unknown, the method 1500 may continue at step 1564 such that pharmacy workflow application 114 may determine whether the scanned product is in an active

formulary product. To the extent of the scan product is an inactive formulary product, the method 1500 may proceed to step 1580 such that pharmacy workflow application 114 may add an alert associated with the barcode of the scanned product indicative of an inactive formulary product.

In this regard, the method 1500 may continue with returned reference to step 1552 such
5 that a product may be scanned for product preparation. If, however, the scanned product is not associated with an inactive formulary product, the method 1500 may continue at step 1568 such that the user scans a diluent product for a reconstitution process. The method 1500 may continue at step 1572 such that pharmacy workflow application 114 determines if the barcode of the scanned reconstitution diluent product is unknown. To the extent that the barcode of the scanned
10 reconstitution diluent product is unknown, the method 1500 may continue with returned reference to step 1560 such that pharmacy workflow application 114 may add an alert associated with the barcode of the scanned reconstitution diluent product indicative of a new product. If, however, the barcode of the scanned reconstitution diluent product is not unknown, the method 1500 may continue at step 1576 such that pharmacy workflow application 114 may determine if the barcode
15 of the scanned reconstitution diluent product is inactive. To the extent that the barcode of the scanned reconstitution diluent product is inactive, the method 1500 may continue with returned reference to step 1580 such that the pharmacy workflow application 114 may add an alert associated with the barcode of the scanned reconstitution diluent product indicative of an inactive formulary product. If, however, the barcode of the scanned reconstitution diluent product is not
20 inactive, the method 1500 may continue at step 1584 such that the user may complete the product procedure actions. The method 1500 may continue at step 1588 such that the product preparation is complete. In this regard, the method 1500 may continue at step 1592 such that the pharmacy workflow application 114 indicates successful product preparation. Notably, the completion of

step 1588 may facilitate the continuation of method 1500 at step 1512 such that the product ingredient preparation is complete.

Turning next to Fig. 16, the pharmacy workflow application 114 may facilitate the reprocessing of dose orders according generally to method 1600. Method 1600 may generally
5 comprise steps associated with actions taken for each problematic dose order 1600a and steps associated with database transactions 1600b. In this regard, method 1600 may begin at step 1602 such pharmacy workflow application 114 may receive a list of problematic dose orders for reprocessing according to method 1600. The method 1600 may continue at step 1604 such that pharmacy workflow application 114 may determine if all active alerts of the problematic dose
10 order have been resolved. To the extent that all active alerts of the problematic dose order have not been resolved, method 1600 may continue at step 1606 such that the pharmacy workflow application 114 may generate an error message, which notes: “list of active alerts of the problematic dose order into error collection.”

In this regard, the method 1600 may continue at step 1613 such that pharmacy flow
15 application 114 may add the error message into in error collection associated with the dose order. If, however, the pharmacy workspace application 114 determines, with returned reference to step 1604, that all active alerts of the problematic dose order have been resolved, the method 1600 may continue by executing the steps generally associated with database transaction 1600b. In this regard, the method 1600 may continue at step 1608 such that the pharmacy workflow application
20 114 obtains a cloned copy of the dose order from a backup. The method 1600 may continue at step 1610 such that the pharmacy workflow application 114 determines if the problematic dose order has been successfully cloned from the backup. To the extent that the pharmacy workflow application 114 does not successfully obtain a clone from the backup, the method 1600 may

continue at step 1612 such that the pharmacy workflow application 114 may generate an error message, which notes: “unable to perform clone operation from backup.” In this regard, method 1600 may continue at step 1613 according to the discussion above.

If, however, the pharmacy workflow application 114 does successfully obtain a clone from
5 the backup, the method 1600 may continue at step 1614 such that the pharmacy workflow application 114 acquires a lock on the problematic dose order ID. The method 1600 may continue at step 1616 such that the pharmacy workflow application 114 determines whether a lock has been acquired on the problematic dose order ID. To the extent that a lock has not been acquired on the problematic dose order ID, the method the 1600 may continue at step 1618 such that the pharmacy
10 workflow application 114 may generate an error message, which notes: “unable to acquire lock on problematic dose order.” In this regard, method 1600 may continue at step 1613 according to the discussion above.

If, however, a lock is acquired on the problematic dose order ID, the method 1600 may continue at step 1620 such that the cloned dose order and problematic dose order may be linked to
15 each other. The method 1600 may continue at step 1622 such that the cloned dose order is saved with an empty dose order status. Method 1600 may continue at step 1624 such that the pharmacy workflow application 114 may determine whether the cloned dose order has been saved successfully. To the extent that the cloned dose order has not been saved successfully, the pharmacy workflow application 114 may generate an error message, which notes: “failed to save
20 the cloned dose order with empty dose status.” In this regard, method 1600 may continue at step 1613 according to the discussion above.

If, however, the cloned dose order has been saved successfully, the method 1600 may continue at step 1628 such that he cloned dose order is sent for processing. The method 1600 may

continue at step 1630 such that the pharmacy workflow application 114 may determine whether the problematic dose order has already been bypassed. To the extent that the problematic dose order has not already been bypassed, the method 1600 may proceed to step 1632 such that the pharmacy workflow application 114 may determine whether the problematic dose order is already discontinued. If the problematic dose order has already been bypassed or the problematic dose order is already discontinued, the method 1600 may continue at step 1634 such that the status of the problematic dose order is unchanged. The pharmacy work flow application 114 may save the dose status of the cloned dose order to a backup file.

With returned reference to step 1634, if the problematic dose order is not already discontinued, the method 1600 may continue at step 1636 such that the pharmacy workflow application 114 may discontinue the problematic dose order with a remake reason, which notes: “discontinue during reprocess of dose order.” The method 1600 may continue at step 1638 such that the pharmacy workflow application 114 may determine whether the problematic dose order has been successfully discontinued. To the extent that the problematic dose order has not been successfully discontinued, the method 1600 may continue at step 1640 such that the pharmacy workflow application 114 may generate an error message, which notes: “failed to discontinue the problematic dose order.” If, however, the problematic dose order has been successfully discontinued the method 1600 may proceed to step 1642 such that the pharmacy workflow application 114 may add a dose order note associated with the problematic dose order with the tag: “reprocess_cancel.”

In another aspect, the method 1600 may continue at step 1644 such that the pharmacy workflow application 114 makes the all alert status of the problematic dose order set to: “processed.” The method 1600 may continue at step 1646 such that the pharmacy workflow

application 114 may save the dose order status of the cloned dose order to a backup file. The method 1600 may continue at step 1648 such that the pharmacy workflow application 114 may set the dose status of the cloned dose order to “ignore.” The method 1600 may continue at step 1650 such that pharmacy workflow application 114 may add a dose order note associated with the cloned
5 dose order with the tag: “reprocess_ignore.” The method 1600 may continue at step 1652 such that the pharmacy workflow application 114 may remove the problematic dose order from the backup.

In this regard, the steps associated with database transaction 1600b may be substantially performed. The method 1600 may continue at step 1654 such that the pharmacy workflow application 114 may add the problematic dose order into one or more of the following lists:
10 reprocessed successfully and can be requeued, reprocessed successfully but discontinued, and reprocessed successfully but bypassed. The method 1600 may continue at step 1656 such that pharmacy workflow application 114 may unlock the reprocessed dose ID. Notably, step 1613 of method 1600, discussed in greater detail above, may proceed to step 1656. The method may continue at step 1658 such that the client device may receive a response from the pharmacy
15 workflow application 114 associated with the resolution of the problematic dose order.

Turning next to Fig. 17, the pharmacy work flow application 114 may facilitate the requeue of dose orders. In this regard, the method 1700 may begin at step 1704 such that the pharmacy workflow application 114 receives a list of problematic dose orders for the requeue process. The method 1700 may continue at step 1708 such that the pharmacy workflow application 114 acquires
20 a lock on the problematic dose ID. The method 1700 may continue at step 1712 such that the pharmacy workflow application determines if a lock has been acquired on the reprocessed dose IDs. To the extent that a lock has not been acquired on the reprocessed dose IDs, the method 1700 may continue and step 1716 such that the pharmacy workflow application 114 may generate an

error message, which notes: “unable to require lock on problematic dose order.” In this regard, the method 1700 may continue at step 1720 such that the pharmacy workflow application 114 adds an error message to an error collection of the problematic dose order. To the extent that the lock is acquired on the reprocessed dose ID, the method 1700 may continue at step 1724 such that the pharmacy workflow application 114 may determine if all active alerts of the problematic dose order have been resolved. If all active alert on the problematic dose order have not been resolved, the method 1700 may continue at step 1728 such that the pharmacy workflow application 114 may generate an error message, which notes: “this does order can’t be requeued until the problematic dose order is processed, which has following alerts that are unresolved.” In this regard, the method 1700 may continue at step 1720 as described above.

If, however, all active alerts on, or otherwise associated with, the problematic dose order have been resolved, the method 1700 may continue at step 1732 such that the pharmacy workflow application 114 may determine whether any other dose orders are linked to the problematic dose order. To the extent that there are not any dose orders in the system that are linked to the problematic dose order, the method 1700 may continue at step 1736 such that the pharmacy workflow application 114 may generate an error message, which notes: “cloned dose order is not found for the problematic dose.” In this regard, the method 1700 may continue at step 1720 as described above.

To the extent that there are dose orders in the system that are linked to the problematic dose order, the method 1700 may continue at step 1740 such that the pharmacy workflow application 114 determines whether the problematic dose order has already been discontinued outside of the reprocess functionality. If the problematic dose order has already been discontinued outside of the reprocess functionality, the method 1700 may continue at step 1744 such that the pharmacy

workflow application 114 may generate an error message, which notes: “problematic dose order has already been discontinued.” In this regard, the method 1700 may continue at step 1720 as described above.

If, however, the problematic dose order has not already been discontinued outside of the reprocess functionality, the method 1700 may continue at step 1748 such that the pharmacy workflow application 114 may determine whether the problematic dose order has already been bypassed. If the problematic dose order has not already been bypassed, the method 1700 may continue at step 1756 such that the pharmacy workflow application 114 may determine whether the cloned dose order has any active alerts associated with it. In this regard, if the problematic dose order has already been bypassed, or the cloned dose order has active alerts associated with it, the method 1700 may proceed to step 1752 and step 1760, respectively. That is, if the problematic dose order has already been bypassed, the method 1700 may proceed to step 1752 such that the pharmacy workflow application 114 may generate an error message that notes: “problematic dose order has already been bypassed.” And, if the cloned dose order has active alerts associated with it, the method 1700 may continue at step 1760 such that the pharmacy work flow application 114 may generate an error message, which notes: “cloned dose order has some active alerts associated with it.” In response to either step 1752 or step 1760, the method 1700 may continue at step 1720 as discussed above.

With returned reference to step 1756, to the extent that the cloned dose order does not have any active alerts associated with it, the method 1700 may continue at step 1764 such that the pharmacy workflow application 114 may find the dose status of the problematic dose order from the backup. The method 1700 may continue at step 1768 such that the pharmacy workflow application 114 may determine whether the dose status of the cloned dose order is found in the

backup. To the extent that the dose status of the cloned dose order is not found in the backup, the method 1700 may continue at step 1780 such that the pharmacy workflow application 114 may set the cloned dose order status to pending. If, however, the dose status of the cloned dose order is found in the backup, the method 1700 may continue at step 1772 such that the pharmacy workflow application 114 may set the cloned dose order dose status to: “dose status from backup of reprocess dose.” In response to the completion of either step 1772 or 1780, the method 1700 may continue at step 1776 such that the pharmacy workflow application 114 may determine whether the dose status of the cloned dose order has been successfully set. To the extent that the dose status of the cloned dose order has not been successfully set, the method 1700 may continue at step 1784 such that the pharmacy workflow application 114 may generate an error message, which notes: “failed to set the dose status as to cloned dose order.” In this regard, the method 1700 may continue at step 1720 as described above.

If, however, the dose status of the cloned dose order has been successfully set, the method 1700 may continue at step 1788 such that the pharmacy workflow application 114 may add the problematic dose order into a list of successfully requeued doses. The method 1700 may continue at step 1790 such that the pharmacy workflow application 114 may add a dose order note to the cloned dose order with the tag: “reprocess_queue.” The method 1700 may continue at step 1792 such that the pharmacy workflow application 114 may remove the cloned dose order from the backup in the case that the cloned dose order was associated with a back. The method 1700 may continue at step 1794 such that the pharmacy workflow application 114 may unlock the problematic dose ID. Notably, the completion of previously described steps 1720, may also facilitate the method 1700 continuing at step 1794. The method 1700 may continue at step 1796 such that the client receives a response from the pharmacy workflow application 114.

The foregoing description of the present invention has been presented for purposes of illustration and description. Furthermore, the description is not intended to limit the invention to the form disclosed herein. Consequently, variation and modifications commensurate with the above teachings, skill and knowledge of the relevant art, are within the scope of the present invention. The embodiments described hereinabove are further intended to explain best modes known of practicing the invention and to enable others skilled in the art to utilize the invention in such, or other embodiments and with various modification required by the particular application(s) or use(s) of the present invention. It is intended that the appended claims be construed to include alternative embodiments to the extent permitted by the prior art.

What is claimed is:

1. A method of batch processing of a plurality of dose orders in a pharmacy workflow management application, the method comprising:

5 accessing, by a processor executing said pharmacy workflow management application, a plurality of dose order records corresponding to dose orders received by said pharmacy workflow management application, wherein said plurality of dose orders comprise identical constituent products, and wherein said plurality of dose orders have not been prepared;

10 aggregating said plurality of dose orders to create to an aggregated plurality of said dose orders;

displaying, at a user interface of the pharmacy workflow management application, a dose preparation protocol corresponding to said aggregated plurality of dose order records, wherein said dose preparation protocol comprises an aggregate amount of at least one of said identical constituent products needed to prepare each of said aggregated plurality of dose order records;

15 receiving, from a user by way of a user input of the pharmacy workflow management application, an indication of said aggregate amount of said at least one constituent product; and

presenting to the user, via said user interface, one or more preparation steps regarding preparation of said aggregated plurality of doses upon said receiving said aggregate amount of said at least one constituent product.

20 2. The method of claim 1, further comprising:

determining, by said processor, said aggregate plurality of dose orders as suitable for preparation via a batch workflow, wherein said batch workflow corresponds to preparing said

aggregated plurality of dose orders in a common sequence such that a separate dose preparation protocol is not required for each dose order of said aggregated plurality of dose orders.

3. The method of claim 2, further comprising after said determining:

5 initiating said dose preparation protocol corresponding to said aggregate plurality of dose orders for preparation via said batch workflow.

4. The method of claim 3, further comprising:

printing a dose label for each dose order of said aggregate plurality of dose orders.

10

5. The method of claim 3, wherein said indication of said aggregate amount of said at least one constituent product corresponds to one or more formulary products for use in preparation of said aggregate plurality of doses.

15 6. The method of claim 5, further comprising:

printing a work-in-progress label upon said receiving said aggregate amount of at least one constituent product, wherein said work-in-progress label is indicative of a remaining volume of said one or more formulary products after completion of said one or more preparation steps.

20 7. The method of claim 5, wherein said presenting comprising presenting said one or more preparation steps free from printing a work-in-progress label for each use of said one or more formulary products, wherein said work-in-progress label is indicative of a remaining volume of said one or more formulary products after completion of said one or more preparation steps for a

given dose of said aggregate plurality of doses.

8. The method of claim 1, wherein said indication of said aggregate amount of said at least one constituent product corresponds to one or more formulary products for use in preparation of said dose order.

9. The method of claim 8, wherein said one or more formulary products includes a total formulary product volume such that said total formulary product volume equals said aggregate amount of said at least one constituent product.

10. The method of claim 1, wherein said accessing comprises identifying common associations among said plurality of dose order records such that said accessed plurality of dose order records comprise identical constituent products.

11. The method of claim 10, wherein said aggregating comprises grouping said plurality of dose orders into a queue at least partially based on said identified common associations.

12. The method of claim 11, further comprising:
initiating said dose preparation protocol corresponding to said aggregate plurality of dose orders of said queue.

13. The method of claim 1, wherein said aggregate amount includes each of said identical constituent products of said dose order.

14. The method of claim 1, wherein said indication comprises a scan.

15. A method for dynamic determination of a final container for use in a dose order preparation protocol of a pharmacy workflow management application, the method comprising:

5 accessing, by a processor executing said pharmacy workflow management application, a dose order record corresponding to a dose order received by said pharmacy workflow management application, wherein said dose order has not been prepared;

 initiating a dose preparation protocol corresponding to said dose order;

 receiving an indication of a plurality of products associated with said dose preparation
10 protocol;

 dynamically determining, in response to said receiving, at least one of the products as a selected product corresponding to a final container based at least in part on a final container characteristic of said selected product; and

 displaying, at a user interface of said pharmacy workflow application, a dose preparation
15 step associated with preparation of said dose relative to said final container.

16. The method of claim 15, wherein said final container characteristic is comprised of a first data field corresponding to said dose order record and a second data field corresponding to a formulary data record associated with at least one of said plurality of indicated products, and
20 wherein said first data field and said second data field are comprised of a plurality of values.

17. The method of claim 16, wherein said dynamically determining comprises identifying a value of said first data field indicative of a final container indicator of said dose order, and

wherein said final container indication indicates said dose order is suitable for preparation in said final container.

18. The method of claim 17, wherein said dynamically determining comprises identifying a
5 value of said second data field indicative of diluent product of said plurality of products
associated with said dose preparation protocol, and wherein said diluent product indicates said
dose order is suitable for preparation in said final container.

19. The method of claim 18, wherein said dynamically determining comprises identifying a
10 value of said first data field indicative of a dose order free of a hazmat drug, and wherein said
dose order free of said hazmat drug indicates said dose order is suitable for preparation in said
final container.

20. The method of claim 19, wherein said dynamically determining comprises identifying a
15 value of said first data field indicative of a diluent line item and identifying a value of said
second data field indicative of a QS drug field, and wherein said diluent line item equals said QS
drug field such that said dose order is suitable for preparation in said final container.

21. The method of claim 20, wherein said dynamically determining comprises identifying a
20 value of said first data field indicative of a total volume of said dose order, and wherein said
value is greater than a minimum dose volume such that said dose order is suitable for preparation
in said final container.

22. The method of claim 21, wherein said minimum dose volume is configurable based on a site-specific preference.

23. The method of claim 21, wherein said minimum dose volume is 24 ml.

5

24. The method of claim 21, wherein said dynamically determining comprises identifying a value of said second data field indicative of a final container indication of said diluent product, and wherein said final container indication of said diluent product indicates said diluent product is suitable as said selected product corresponding to said final container.

10

25. The method of claim 24, wherein said dynamically determining comprises identifying a value of said second data field indicative of a dose order free of multiple selected products corresponding to said final container, and wherein said dose order free of multiple selected products corresponding to said final container indicates said dose order is suitable for preparation in said final container.

15

26. The method of claim 25, wherein said dynamically determining comprises identifying a value of said second data field indicative of a new container of said diluent product, and wherein said new container of said diluent product indicates said dose order is suitable for preparation in said final container.

20

27. The method of claim 26, wherein said dynamically determining comprises identifying a value of said first data field indicative of a total volume of said dose order and identifying a

value of said second data field indicative of a volume of said diluent product, wherein the absolute percentage difference between said total volume of said dose order and said volume of said diluent product is less than a predetermined differential percentage such that said dose order is suitable for preparation in said final container.

5

28. The method of claim 27, wherein said predetermined differential percentage is configurable based on a site-specific preference.

29. The method of claim 27, wherein said predetermined differential percentage is 100%.

10

30. The method of claim 27, wherein said dynamically determining comprises identifying a value of said first data field indicative of a total volume of said dose order and identifying a value of said second data field indicative of a carrier ingredient volume of said dose order, wherein the difference between said carrier ingredient volume of said dose order and said total volume of said dose order is less than a predetermined waste volume such that said dose order is suitable for preparation in said final container.

15

31. The method of claim 30, wherein said predetermined waste volume is configurable based on a site-specific preference.

20

32. The method of claim 30, wherein said predetermined waste volume is 200 ml.

33. A method for dynamic diluent volume accounting in a dose order preparation protocol of a pharmacy workflow management application, the method comprising:

accessing, by a processor executing said pharmacy workflow management application, a dose order record corresponding to a dose order received by said pharmacy workflow

5 management application, wherein said dose order has not been prepared;

receiving a first indication of a product associated with a dose preparation protocol;

dynamically determining, in response to said receiving, a required diluent volume based on said indication; and

10 displaying, at a user interface of the pharmacy workflow application, a dose preparation step associated with preparation of said dose relative to said required diluent volume.

34. The method of claim 33, further comprising:

receiving a second indication of said dynamically determined required diluent volume.

15 35. The method of claim 34, wherein said indication of said dynamically determined required diluent volume corresponds to one or more formulary products for use in preparation of said dose order.

20 36. The method of claim 35, wherein said one or more formulary products is associated with a list of allowed diluent products relative to the dose preparation protocol.

37. The method of claim 36, wherein each of said one or more formulary products are identical formulary products.

38. The method of claim 35, wherein said one or more formulary products include a total formulary product volume such that said total formulary product equals said dynamically determined required diluent volume.

5

39. The method of claim 33, wherein said displaying comprises displaying said dynamically determined required diluent volume.

40. The method of claim 39, wherein said dynamically determined required diluent volume is
10 configurable based on receiving a second indication of said required diluent volume.

41. The method of claim 33, wherein said displaying comprises displaying at least one collapsible area configured to hide information when collapsed, and wherein said information corresponds to a second indication of said dynamically determined required diluent volume.

15

42. A method for managing an error regarding a dose order record in a pharmacy workflow queue in a pharmacy workflow management application, the method comprising:

accessing, at a processor executing said pharmacy workflow management application, a dose order record corresponding to a dose order received by said pharmacy workflow

5 management application, wherein said dose order has not been prepared;

first processing, by said processor, said dose order record to generate an alert corresponding to an error state related to said dose order;

displaying, at a user interface of said pharmacy workflow management application, said dose order in corresponding relation to said alert;

10 receiving an input indicative of a resolution of said error;

second processing said dose order based on said input; and

requeuing said dose order upon confirmation of said resolution.

43. The method of claim 42, wherein said error state comprises a parsing error.

15 44. The method of claim 43, wherein said parsing error corresponds to one or more unintelligible data attributes of said dose order record.

45. The method of claim 42, wherein said error state comprises a drug error.

20 46. The method of claim 45, wherein said drug error corresponds to one or more unrecognizable data attributes associated with a drug of said dose order record.

47. The method of claim 46, further comprising:

creating a formulary data record at least partially based on said unrecognizable data attribute.

5 48. The method of claim 42, wherein said error state comprises an unknown unit error.

49. The method of claim 48, wherein said unknown unit error corresponds to an unrecognizable data attribute associated with a unit of said dose order record.

10 50. The method of claim 42, wherein said error state comprises an unknown product error.

51. The method of claim 50, wherein said unknown product error corresponds to an unrecognizable data attribute associated with a product of said dose order record.

15 52. The method of claim 51, further comprising:

creating a formulary data record at least partially based on said unrecognizable data attribute.

53. The method of claim 42, wherein said error state comprises an unknown route error.

20

54. The method of claim 53, wherein said unknown route error corresponds to an unrecognizable data attribute associated with a route of said dose order.

55. The method of claim 54, further comprising:

modifying a formulary data record at least partially based on said unrecognizable data attribute.

5 56. The method of claim 42, wherein said error state comprises a passwords error.

57. The method of claim 56, wherein said passwords error corresponds to an expired password of a user account.

10 58. The method of claim 56, wherein said passwords error corresponds to a disabled user account.

59. The method of claim 42, wherein said error state comprises a suspected discontinued dose error.

15

60. The method of claim 59, wherein said suspected discontinued dose error corresponds to a discontinued dose order.

61. The method of claim 60, further comprising:

20 deleting said discontinued dose order.

62. The method of claim 42, wherein said error state comprises a suspected duplicate error.

63. The method of claim 62, wherein said suspected duplicate error corresponds to a duplicate dose order.

64. The method of claim 63, further comprising:

5 deleting said duplicate dose order.

65. A system for batch processing of a plurality dose orders in a pharmacy workflow management application, comprising:

an access terminal remote from and in operative communication with said pharmacy workflow management application, wherein said pharmacy workflow management application is operable to access a plurality of dose order records corresponding to dose orders received by said pharmacy workflow management application, and wherein said plurality of dose orders comprise identical constituent products, and wherein said plurality of dose orders have not been prepared, and wherein said pharmacy workflow management application is operable to aggregate said plurality of dose orders to create an aggregated plurality of said dose orders; and

10
15

a user interface provided at said access terminal for displaying a dose preparation protocol corresponding to said aggregated plurality of dose order records, wherein said dose preparation protocol comprises an aggregate amount of at least one of said identical constituent products need to prepare each of said aggregated plurality of dose orders, and wherein said pharmacy workflow application is operable to receive, from a user by way of a user input of said pharmacy workflow management application, an indication of said aggregate amount of said at least one constituent product, and where said user interface is operable to present to said user, via said user interface, one or more preparation steps regarding preparation of said aggregated

20

plurality of doses upon receiving said aggregate amount of said at least one constituent product.

66. The system of claim 65, wherein said pharmacy workflow management application is operable to determine said aggregate plurality of dose order as suitable for preparation via a batch workflow, wherein said batch workflow corresponds to preparing said aggregated plurality of dose orders in a common sequence such that a separate dose preparation protocol is not required for each dose order of said aggregated plurality of dose orders.

67. The system of claim 66, wherein said pharmacy workflow management application is operable to initiate said dose preparation protocol corresponding to said aggregate plurality of dose orders for preparation via said batch workflow.

68. The system of claim 67, wherein said pharmacy workflow management application is operable to print a dose label for each dose order of said aggregate plurality of dose orders.

69. The system of claim 67, wherein said indication of said aggregate amount of said at least one constituent product correspond to one or more formulary products for use in preparation of said aggregate plurality of doses.

70. The system of claim 69, wherein said pharmacy workflow management application is operable to print a work-in-progress label upon receiving said aggregate amount of at least one constituent product, and wherein said work-in-progress label is indicative of a remaining volume of said one or more formulary products after completion of said one or more preparation steps.

71. The system of claim 69, wherein said presenting comprises presenting said one or more preparation steps free from printing a work-in-progress label for each use of said one or more formulary products, and wherein said work-in-progress label is indicative of a remaining volume
5 of said one or more formulary products after completion of said one or more preparation steps for a given dose of said aggregate plurality of doses.

72. The system of claim 65, wherein said indication of said aggregate amount of said at least one constituent product corresponds to one or more formulary products for use in preparation of
10 said dose order.

73. The system of claim 72, wherein said one or more formulary products includes a total formulary product volume such that said total formulary product volume equals said aggregate amount of said at least one constituent product.
15

74. The system of claim 65, wherein said accessing comprises identifying common associations among said plurality of dose order records such that said accessed plurality of dose order records comprise identical constituent products.

20 75. The system of claim 74, wherein said aggregating comprises group said plurality of dose orders into a queue at least partially based on said identified common associations.

76. The system of claim 75, wherein said pharmacy workflow management application is

operable to initiate said dose preparation protocol corresponding to said aggregate plurality of dose orders of said queue.

77. The system of claim 65, wherein said aggregate amount includes each of said identical
5 constituent products of said dose order.

78. The system of claim 65, wherein said indication comprises a scan.

79. A system for dynamic determination of a final container for user in a dose order
10 preparation protocol of a pharmacy workflow management application, comprising:

an access terminal remote from an in operative communication with said pharmacy workflow management application, wherein said pharmacy workflow management application is operable to access a dose order record corresponding to a dose order received by said pharmacy workflow management application, and wherein said dose order has not been prepared, and
15 wherein said pharmacy management application is operable to initiate a dose preparation protocol corresponding to the dose order, and wherein said pharmacy workflow application is operable to receive an indication of a plurality of products associated with said dose preparation protocol, and wherein said pharmacy workflow application is operable to dynamically determine, in response to said received indication, at least one of said products as a elected product
20 corresponding to a final container based at least in part on a final container characteristic of said selected product; and

a user interface provided at said access terminal for displaying a dose preparation step associated with preparation of said dose relative to said final container.

80. The system of claim 79, wherein said final container characteristic is comprised of a first data field corresponding to said dose order record and a second data field corresponding to a formulary data record associated with at least one of said plurality of indicated products, and
5 wherein said first data field and said second data field are comprised of a plurality of values.

81. The system of claim 80, wherein said pharmacy workflow management application is operable to, in relation to said dynamically determining, identify a value of said first data field indicative of a final container indicator of said dose order, and wherein said final container
10 indication indicates said dose order is suitable for preparation in said final container.

82. The system of claim 81, wherein said pharmacy workflow management application is operable to, in relation to said dynamically determining, identifying a value of said second data field indicative of diluent product of said plurality of products associated with said dose
15 preparation protocol, and wherein said diluent product indicates said dose order is suitable for preparation in said final container.

83. The system of claim 82, wherein said pharmacy workflow management application is operable to, in relation to said dynamically determining, identify a value of said first data field
20 indicative of a dose order free of a hazmat drug, and wherein said dose order free of said hazmat drug indicates said dose order is suitable for preparation in said final container.

84. The system of claim 83, wherein said pharmacy workflow management application is

operable to, in relation to said dynamically determining, identify a value of said first data field indicative of a diluent line item and identifying a value of said second data field indicative of a QS drug field, and wherein said diluent line item equals said QS drug field such that said dose order is suitable for preparation in said final container.

5

85. The system of claim 84, wherein said pharmacy workflow management application is operable to, in relation to said dynamically determining, identify a value of said first data field indicative of a total volume of said dose order, and wherein said value is greater than a minimum dose volume such that said dose order is suitable for preparation in said final container.

10

86. The method of claim 85, wherein said minimum dose volume is configurable based on a site-specific preference.

87. The method of claim 85, wherein said minimum dose volume is 24 ml.

15

88. The system of claim 85, wherein said pharmacy workflow management application is operable to, in relation to said dynamically determining, identify a value of said second data field indicative of a final container indication of said diluent product, and wherein said final container indication of said diluent product indicates said diluent product is suitable as said selected

20 product corresponding to said final container.

89. The system of claim 88, wherein said pharmacy workflow management application is operable to, in relation to said dynamically determining, identify a value of said second data field

indicative of a dose order free of multiple selected products corresponding to said final container, and wherein said dose order free of multiple selected products corresponding to said final container indicates said dose order is suitable for preparation in said final container.

5 90. The system of claim 89, wherein said pharmacy workflow management application is operable to, in relation to said dynamically determining, identify a value of said second data field indicative of a new container of said diluent product, and wherein said new container of said diluent product indicates said dose order is suitable for preparation in said final container.

10 91. The system of claim 90, wherein said pharmacy workflow management application is operable to, in relation to said dynamically determining, identify a value of said first data field indicative of a total volume of said dose order and identifying a value of said second data field indicative of a volume of said diluent product, wherein the absolute percentage difference between said total volume of said dose order and said volume of said diluent product is less than
15 a predetermined differential percentage such that said dose order is suitable for preparation in said final container.

92. The system of claim 91, wherein said predetermined differential percentage is configurable based on a site-specific preference.

20

93. The system of claim 91, wherein said predetermined differential percentage is 100%.

94. The system of claim 91, wherein said pharmacy workflow management application is

operable to, in relation to said dynamically determining, identify a value of said first data field indicative of a total volume of said dose order and identifying a value of said second data field indicative of a carrier ingredient volume of said dose order, wherein the difference between said carrier ingredient volume of said dose order and said total volume of said dose order is less than
5 a predetermined waste volume such that said dose order is suitable for preparation in said final container.

95. The system of claim 94, wherein said predetermined waste volume is configurable based on a site-specific preference.

10

96. The system of claim 94, wherein said predetermined waste volume is 200 ml.

97. A system for dynamic diluent volume accounting in a dose order preparation protocol of a pharmacy workflow management application, comprising:

15 an access terminal remote from an in operative communication with said pharmacy workflow management application, wherein said pharmacy workflow management application is operable to access a dose order record corresponding to a dose order received by the pharmacy workflow management application, and wherein said dose order has not been prepared, and wherein said pharmacy workflow management application is operable to receive a first
20 indication of a product associated with a dose preparation protocol, and wherein said pharmacy workflow management application is operable to dynamically determine, in response to said received first indication, a required diluent volume based on said indication; and
a user interface provided at said access terminal for displaying a dose preparation step

associated with preparation of said dose relative to said required diluent volume.

98. The system of claim 97, wherein said pharmacy workflow management application is operable to receive a second indication of said dynamically determined required diluent volume.

5

99. The system of claim 98, wherein said indication of said dynamically determined required diluent volume corresponds to one or more formulary products for use in preparation of said dose order.

10 100. The system of claim 99, wherein said one or more formulary products is associated with a list of allowed diluent products relative to the dose preparation protocol.

101. The system of claim 100, wherein each of said one or more formulary products are identical formulary products.

15

102. The system of claim 99, wherein said one or more formulary products include a total formulary product volume such that said total formulary product equals said dynamically determined required diluent volume.

20 103. The system of claim 97, wherein said pharmacy workflow management application is operable to, in relation to said displaying, display said dynamically determined required diluent volume.

104. The system of claim 103, wherein said pharmacy workflow management application is operable to configure said dynamically determined required diluent volume based on receiving a second indication of said required diluent volume.

5 105. The system of claim 97, wherein said user interface is operable to display at least one collapsible area configured to hide information when collapsed, and wherein said information corresponds to a second indication of said dynamically determined required diluent volume.

106. A system for managing an error regarding a dose order record in a pharmacy workflow queue in a pharmacy workflow management application, comprising:

an access terminal remote from an in operative communication with said pharmacy workflow management application, wherein said pharmacy workflow management application is operable to access a dose order record corresponding to a dose order received by said pharmacy workflow management application, and wherein said dose has not been prepared, and wherein
15 said pharmacy workflow application is operable to first process said dose order record to generate an alert corresponding to an error state related to said dose order; and

a user interface provided at said access terminal for displaying said dose order in corresponding relation to said alert;

wherein said pharmacy workflow management application is operable to receive an input
20 indicative of a resolution of said error, and wherein said pharmacy management application is operable to second process said dose order based on said input, and wherein said pharmacy management application is operable to requeue said dose order upon confirmation of said resolution.

107. The system of claim 106, wherein said error state comprises a parsing error.

108. The system of claim 107, wherein said parsing error corresponds to one or more
5 unintelligible data attributes of said dose order record.

109. The system of claim 106, wherein said error state comprises a drug error.

110. The system of claim 109, wherein said drug error corresponds to one or more
10 unrecognizable data attributes associated with a drug of said dose order record.

111. The system of claim 110, wherein said pharmacy workflow application is operable to
create a formulary data record at least partially based on said unrecognizable data attribute.

15 112. The system of claim 106, wherein said error state comprises an unknown unit error.

113. The system of claim 112, wherein said unknown unit error corresponds to an
unrecognizable data attribute associated with a unit of said dose order record.

20 114. The system of claim 106, wherein said error state comprises an unknown product error.

115. The system of claim 114, wherein said unknown product error corresponds to an
unrecognizable data attribute associated with a product of said dose order record.

116. The system of claim 115, wherein said pharmacy workflow management application is operable to create a formulary data record at least partially based on said unrecognizable data attribute.

5

117. The system of claim 106, wherein said error state comprises an unknown route error.

118. The system of claim 117, wherein said unknown route error corresponds to an unrecognizable data attribute associated with a route of said dose order.

10

119. The system of claim 118, wherein said pharmacy workflow management application is operable to modify a formulary data record at least partially based on said unrecognizable data attributes.

15 120. The system of claim 106, wherein said error state comprises a passwords error.

121. The system of claim 120, wherein said passwords error corresponds to an expired password of a user account.

20 122. The system of claim 120, wherein said passwords error corresponds to a disabled user account.

123. The system of claim 106, said error state comprises a suspected discontinued dose error.

124. The system of claim 123, wherein said suspected discontinued dose error corresponds to a discontinued dose order.

5 125. The system of claim 124, wherein said pharmacy workflow management application is operable to delete said discontinued dose order.

126. The system of claim 106, wherein said error state comprises a suspected duplicate error.

10 127. The system of claim 126, wherein said suspected duplicate error corresponds to a duplicate dose order.

128. The system of claim 127, wherein said pharmacy workflow management application is operable to delete said duplicate dose order.

15