



1/13

Fig. 1




Leibovitz L-15 medium
 Hydrolysed gelatin
 Cell

Fig. 2




Leibovitz L-15 medium
 Hydrolysed gelatin
 Cell

Fig. 3



Hydrolysed gelatin
 Cell

Fig. 4



Hydrolysed gelatin
 Cell attached to the dish

2/13

Fig. 5





Fig. 6



3/13

Fig. 7






Fig. 8



4/13


Fig. 9a



- Cyt b5 expressed in terms of LDH activity
- Cyt b5 expressed in terms of protein
- Cyt P450 expressed in terms of LDH activity
- Cyt P450 expressed in terms of protein

5/13

Fig. 9b



—■— Cyt b5 expressed in
terms of LDH activity

—□— Cyt b5 expressed in
terms of protein

—●— Cyt P450 expressed in
terms of LDH activity

—○— Cyt P450 expressed in
terms of protein

6/13

Fig. 10





Fig. 13



7/13

Fig. 11a





Fig. 11b




8/13


Fig. 12



Newly attached cell.
Round shape and numerous microvilli



Flat shape and
distorted nucleus




Rounded
cell shape

Microvilli


9/13

Fig. 14




10/13

Fig. 15



11/13

Fig. 16



12/13

Fig. 17




Fig. 18

