

**GREMLIN-1 INHIBITOR FOR THE TREATMENT OF A BONE FRACTURE OR BONE
DEFECT.**

5 The present invention relates to methods for the treatment of a bone fracture or bone defect. The invention discloses the effective use of an anti-gremlin-1 antibody to accelerate the healing and bridging of bone tissue in segmental gap defects; and demonstrates that inhibitors of gremlin-1 activity may provide improved therapies for treating or preventing fracture non-union.

10 **BACKGROUND**

A bone fracture is a break or crack in bone tissue and may be the result of a traumatic injury, such as a fall or impact, but can also occur as a result of diseases that affect bone integrity.

15 Non-stabilised bone fractures heal through the process of endochondral ossification, which is initiated through the formation of a blood clot or haematoma. This is coupled with an inflammatory response that modulates immune cells and surrounding skeletal stem cell populations. The haematoma is subsequently replaced with a mineralised cartilaginous
20 callus through the action of various growth factors including transforming growth factor beta (TGF β) (Cho et al; 2002), fibroblastic growth factors (FGFs) (Schmid et al; 2009), and bone morphogenic proteins (BMPs) (Yu et al; 2010). Through the actions of osteoclasts and osteoblasts, the mineralised callus is replaced by woven bone. The final remodelling stage involves the replacement of woven bone with lamellar bone. The
25 completion of this process can take many years depending on the age and disease status of the patient.

Bone fractures are generally treated clinically through stabilisation, via the use of a support such as a splint, cast, or brace. In extreme cases involving complex fractures
30 surgical intervention may be required and involves the use of internal and external fixators that are attached directly to the bone. Even with these measures, in approximately 10% of patients the tissue repair process is deficient (Einhorn et al; 2014) resulting in delayed bone union (failure to reach union 6 months post-fracture) or non-union. A non-union is defined as incomplete healing within 9 months, combined with a lack of radiological
35 characteristics associated with fracture healing being observed over three consecutive months (Buza et al; 2016). Current surgical techniques for repairing non-union fractures and critical bone defects are often limited in terms of quantity and quality of the materials

available. Commonly used treatments involve the autologous or allogenic graft, however these carry the additional risk of donor site morbidity (Goulet et al; 1997) and infection (Bostrom et al; 2005), respectively.

- 5 A bone defect is a loss of bone, due to trauma or disease.

There is currently a great unmet medical need for improved treatment of bone fractures and bone defects. Accordingly, it is an object of the present invention to provide new methods for the treatment of a bone fracture or bone defect.

10

The present invention provides inhibitors of gremlin-1 activity for use in the treatment of a bone fracture or bone defect. The invention discloses the effective use of an anti-gremlin-1 antibody to accelerate the healing and bridging of bone tissue in segmental gap defects; and demonstrates that inhibitors of gremlin-1 activity may provide improved therapies for

15 treating or preventing fracture non-union.

DESCRIPTION OF THE INVENTION

- 20 Unless defined otherwise, all scientific and technical terms used herein have the same meaning as commonly understood by one of skill in the art. All publications referred to herein are incorporated by reference.

It will be appreciated that any of the embodiments described herein may be combined.

25

The present invention provides an inhibitor of gremlin-1 activity for use in the treatment of a bone fracture or bone defect. The invention also provides the use of an inhibitor of gremlin-1 activity for the manufacture of a medicament for the treatment of a bone fracture or bone defect. The invention further provides a method for the treatment of a bone

30 fracture or bone defect comprising administering a therapeutically effective amount of an inhibitor of gremlin-1 activity.

Gremlin-1 (also known as Dm and CKTSF1B1) is a 184 amino acid glycoprotein which forms part of the DAN family of cysteine-knot secreted proteins (along with Cerberus and

35 Dan amongst others). Gremlin binds and inhibits the ability of BMP-2, 4, and 7 to signal along with a documented pro-angiogenic role possibly through agonism of VEGFR2. The

main role of Gremlin-1 is during development, in which it is vital during kidney formation and during limb bud formation.

- Bone morphogenetic protein (BMP) signalling is known to control endochondral bone formation, with Gremlin 1 (GREM1) being one of the natural antagonists of this pathway through its binding to BMP2, BMP4 and BMP7 (Hsu et al; 1998). GREM1 conditional deletion in osteoblasts results in sensitisation of BMP signalling/activity and enhanced bone formation *in vivo* (Gazzerro et al; 2007), whilst conditional overexpression in the same cell type causes osteopenia and spontaneous fractures (Gazzerro et al; 2005).
- Furthermore, although the global knockout is embryonic lethal in a BL6 background, 49% of pups survived longer than 24hrs post birth in the C57BL/6/FVB mixed genetic background, and whilst developmental skeletal defects were abundantly present, elevated bone formation rates could be observed (Canalis et al; 2012). Despite this developmental function of GREM1 there is no data to suggest that inhibition of this protein alone will enhance postnatal bone fracture repair. Indeed, although endochondral bone formation is the main mechanism of skeletogenesis at embryonic stages, the mechanisms that regulate cell recruitment are distinct processes when compared to postnatal fracture repair (Ferguson et al; 1999). The role of inflammation has been indicated as a key factor in adult bone repair, thus developmental factors controlling skeletogenesis processes cannot simply be extrapolated to postnatal repair mechanisms.

The term Gremlin-1 as used in the present invention typically has the sequence as set out in the UniProt entry O60565 (SEQ ID NO: 1). The term Gremlin-1 may also refer to a Gremlin-1 polypeptide which:

- (a) comprises or consists of the amino acid sequence of SEQ ID NO: 1 with or without the N-terminal signal peptide, i.e. may comprise or consist of the mature peptide sequence as shown in SEQ ID NO: 21; or
- (b) is a derivative having one or more amino acid substitutions, modifications, deletions or insertions relative to the amino acid sequence of SEQ ID NO: 1 with or without the N-terminal signal peptide (as shown in SEQ ID NO: 21), which retains the activity of Gremlin-1, such as the amino acid sequence of SEQ ID NO: 20.
- (c) a variant thereof, such variants typically retain at least about 60%, 70%, 80%, 90%, 91%, 92%, 93%, 94% or 95% identity to SEQ ID NO: 1 (or SEQ ID NO: 20 or 21) (or even about 96%, 97%, 98% or 99% identity). In other words, such variants may retain about 60% - about 99% identity to SEQ ID NO: 1, suitably about 80% - about 99% identity to SEQ ID NO: 1, more suitably about 90% - about 99% identity to SEQ ID NO: 1 and

most suitably about 95% - about 99% identity to SEQ ID NO: 1. Variants are described further below.

- As discussed further below, residue numbers are typically quoted based on the sequence of SEQ ID NO: 1. However, residue numbering could readily be extrapolated by the skilled person to a derivative or variant sequence as discussed above. Where residue numbers are quoted, the invention also encompasses these residues on a variant or derivative sequence.
- 10 The present inventors have crystallised human Gremlin-1 alone, and in complex with an antibody termed Ab 7326 (Fab fragments). Crystallisation of Gremlin-1 has allowed putative residues in the BMP binding site to be determined. Furthermore, crystallisation with Ab 7326, which is an allosteric inhibitory antibody, has allowed residues in the antibody epitope to be determined. (WO 2018/115017 A2). Antibodies binding this
- 15 epitope have potential as therapeutic agents in the treatment of a bone fracture or bone defect.

Inhibitors of Gremlin-1 activity

- 20 An inhibitor of gremlin-1 activity according to the present invention is an agent that reduces or blocks the activity of gremlin-1. Inhibitors according to the present invention may partially or completely inhibit gremlin-1 activity. Inhibitors of use in the present invention include without limitation, inhibitors that are capable of binding to gremlin-1 or to
- 25 a nucleic acid molecule encoding gremlin-1, or are capable of inhibiting the expression of gremlin-1. Such inhibitors may be, without limitation, proteins, polypeptides, peptides, peptidomimetics, nucleic acids (e.g. DNA, RNA, antisense RNA and siRNA), carbohydrates, lipids, and small molecules.
- 30 In one embodiment, the inhibitor of gremlin-1 activity is an anti-gremlin-1 antibody or a functionally active fragment, variant or derivative thereof.

- The term "antibody" as referred to herein includes whole antibodies and any antigen binding fragment (*i.e.*, "antigen-binding portion") or single chains thereof. An antibody or
- 35 immunoglobulin typically refers to a glycoprotein comprising at least two heavy (H) chains and two light (L) chains inter-connected by disulfide bonds, or an antigen-binding portion thereof. Each heavy chain is comprised of a heavy chain variable region (abbreviated

herein as HCVR or V_H) and a heavy chain constant region. Each light chain is comprised of a light chain variable region (abbreviated herein as LCVR or V_L) and a light chain constant region. The variable regions of the heavy and light chains contain a binding domain that interacts with an antigen. The V_H and V_L regions can be further subdivided
5 into regions of hypervariability, termed complementarity determining regions (CDR), interspersed with regions that are more conserved, termed framework regions (FR).

The constant regions of the antibodies may mediate the binding of the immunoglobulin to host tissues or factors, including various cells of the immune system (*e.g.*, effector cells)
10 and the first component (C1q) of the classical complement system.

An antibody for use in the present invention may be a monoclonal antibody or a polyclonal antibody, and will typically be a monoclonal antibody. An antibody for use in the invention may be a chimeric antibody, a CDR-grafted antibody, a nanobody, a human or humanised
15 antibody or an antigen-binding portion of any thereof.

Polyclonal antibodies may be produced by routine methods, such as immunisation of a suitable animal with the antigen of interest. Blood may be subsequently removed from the animal and the immunoglobulin fraction purified.

20 Antibodies against Gremlin-1 may be obtained, where immunisation of an animal is necessary, by administering the polypeptides to an animal, *e.g.* a non-human animal, using well-known and routine protocols, see for example Handbook of Experimental Immunology, D. M. Weir (ed.), Vol 4, Blackwell Scientific Publishers, Oxford, England,
25 1986). Many warm-blooded animals, such as rabbits, mice, rats, sheep, goats, cows, camels, llamas or pigs may be immunised. However, rabbits, mice, and rats are generally most suitable.

Monoclonal antibodies may be prepared by any method known in the art such as the
30 hybridoma technique (Kohler & Milstein, 1975, *Nature*, 256:495-497), the trioma technique, the human B-cell hybridoma technique (Kozbor *et al.*, 1983, *Immunology Today*, 4:72) and the EBV-hybridoma technique (Cole *et al.*, *Monoclonal Antibodies and Cancer Therapy*, pp77-96, Alan R Liss, Inc., 1985).

35 Antibodies for use in the invention may also be generated using single lymphocyte antibody methods by cloning and expressing immunoglobulin variable region cDNAs generated from single lymphocytes selected for the production of specific antibodies by for

example the methods described by Babcook, J. *et al.*, 1996, Proc. Natl. Acad. Sci. USA 93(15): 7843-7848; WO 92/02551; WO 2004/051268 and WO 2004/106377.

The antibodies for use in the present invention can also be generated using various phage display methods known in the art and include those disclosed by Brinkman *et al.* (in J. Immunol. Methods, 1995, 182: 41-50), Ames *et al.* (J. Immunol. Methods, 1995, 184:177-186), Kettleborough *et al.* (Eur. J. Immunol. 1994, 24:952-958), Persic *et al.* (Gene, 1997 187 9-18), Burton *et al.* (Advances in Immunology, 1994, 57:191-280) and WO 90/02809; WO 91/10737; WO 92/01047; WO 92/18619; WO 93/11236; WO 95/15982; WO 95/20401; and US 5,698,426; US 5,223,409; US 5,403,484; US 5,580,717; US 5,427,908; US 5,750,753; US 5,821,047; US 5,571,698; US 5,427,908; US 5,516,637; US 5,780,225; US 5,658,727; US 5,733,743 and US 5,969,108.

Fully human antibodies are those antibodies in which the variable regions and the constant regions (where present) of both the heavy and the light chains are all of human origin, or substantially identical to sequences of human origin, but not necessarily from the same antibody. Examples of fully human antibodies may include antibodies produced, for example by the phage display methods described above and antibodies produced by mice in which the murine immunoglobulin variable and optionally the constant region genes have been replaced by their human counterparts e.g. as described in general terms in EP 0546073, US 5,545,806, US 5,569,825, US 5,625,126, US 5,633,425, US 5,661,016, US 5,770,429, EP 0438474 and EP 0463151.

Alternatively, an antibody according to the invention may be produced by a method comprising immunising a non-human mammal with a Gremlin-1 immunogen; obtaining an antibody preparation from said mammal; deriving therefrom monoclonal antibodies that recognise Gremlin-1.

The antibody molecules for use in the present invention may comprise a complete antibody molecule having full length heavy and light chains or a fragment or antigen-binding portion thereof. The term "antigen-binding portion" of an antibody refers to one or more fragments of an antibody that retain the ability to selectively bind to an antigen. It has been shown that the antigen-binding function of an antibody can be performed by fragments of a full-length antibody. The antibodies and fragments and antigen binding portions thereof may be, but are not limited to Fab, modified Fab, Fab', modified Fab', F(ab')₂, Fv, single domain antibodies (e.g. VH or VL or VHH), scFv, bi, tri or tetra-valent antibodies, Bis-scFv, diabodies, triabodies, tetrabodies and epitope-binding fragments of

any of the above (see for example Holliger and Hudson, 2005, Nature Biotech. 23(9):1126-1136; Adair and Lawson, 2005, Drug Design Reviews - Online 2(3), 209-217). The methods for creating and manufacturing these antibody fragments are well known in the art (see for example Verma et al., 1998, Journal of Immunological Methods, 216, 165-181). Other antibody fragments for use in the present invention include the Fab and Fab' fragments described in International patent applications WO 2005/003169, WO 2005/003170 and WO 2005/003171 and Fab-dAb fragments described in International patent application WO 2009/040562. Multi-valent antibodies may comprise multiple specificities or may be monospecific (see for example WO 92/22853 and WO 2005/113605). These antibody fragments may be obtained using conventional techniques known to those of skill in the art, and the fragments may be screened for utility in the same manner as intact antibodies.

In one example, the functionally active antibody fragment for use in the present invention is a Fab, Fab', F(ab')₂, Fv or scFv.

The constant region domains of the antibody molecule for use in the present invention, if present, may be selected having regard to the effector functions which may be required. For example, the constant region domains may be human IgA, IgD, IgE, IgG or IgM domains. In particular, human IgG constant region domains may be used, especially of the IgG1 and IgG3 isotypes when antibody effector functions are required. Alternatively, IgG2 and IgG4 isotypes may be used when antibody effector functions are not required. In one example, the isotype is IgG4P, as described by Angal S. et al, Mol Immunol, Vol 30(1), p105-108, 1993.

An antibody for use in the invention may be prepared, expressed, created or isolated by recombinant means, such as (a) antibodies isolated from an animal (e.g., a mouse) that is transgenic or transchromosomal for the immunoglobulin genes of interest or a hybridoma prepared therefrom, (b) antibodies isolated from a host cell transformed to express the antibody of interest, e.g., from a transfectoma, (c) antibodies isolated from a recombinant, combinatorial antibody library, and (d) antibodies prepared, expressed, created or isolated by any other means that involve splicing of immunoglobulin gene sequences to other DNA sequences.

An antibody for use in the invention may be a human antibody or a humanised antibody. The term "human antibody", as used herein, is intended to include antibodies having variable regions in which both the framework and CDR regions are derived from human

germline immunoglobulin sequences. Furthermore, if the antibody contains a constant region, the constant region also is derived from human germline immunoglobulin sequences. Human antibodies for use in the invention may include amino acid residues not encoded by human germline immunoglobulin sequences (e.g., mutations introduced
5 by random or site-specific mutagenesis *in vitro* or by somatic mutation *in vivo*). However, the term "human antibody", as used herein, is not intended to include antibodies in which CDR sequences derived from the germline of another mammalian species, such as a mouse, have been grafted onto human framework sequences.

10 Such a human antibody may be a human monoclonal antibody. Such a human monoclonal antibody may be produced by a hybridoma that includes a B cell obtained from a transgenic nonhuman animal, e.g., a transgenic mouse, having a genome comprising a human heavy chain transgene and a light chain transgene fused to an immortalized cell.

15 Human antibodies may be prepared by *in vitro* immunisation of human lymphocytes followed by transformation of the lymphocytes with Epstein-Barr virus.

The term "derivative" refers to any modified form of the antibody, for example a conjugate
20 of the antibody and another agent or effector molecule.

An effector molecule may comprise a single effector molecule or two or more such molecules so linked as to form a single moiety that can be attached to the antibodies for use in the present invention. Where it is desired to obtain an antibody fragment linked to
25 an effector molecule, this may be prepared by standard chemical or recombinant DNA procedures in which the antibody fragment is linked either directly or via a coupling agent to the effector molecule. Techniques for conjugating such effector molecules to antibodies are well known in the art (see, Hellstrom *et al.*, Controlled Drug Delivery, 2nd Ed., Robinson *et al.*, eds., 1987, pp. 623-53; Thorpe *et al.*, 1982, Immunol. Rev., 62:119-
30 58 and Dubowchik *et al.*, 1999, Pharmacology and Therapeutics, 83, 67-123). Particular chemical procedures include, for example, those described in WO 93/06231, WO 92/22583, WO 89/00195, WO 89/01476 and WO 2003/031581. Alternatively, where the effector molecule is a protein or polypeptide the linkage may be achieved using recombinant DNA procedures, for example as described in WO 86/01533 and EP
35 0392745.

The effector molecule may increase the half-life of the antibody *in vivo*, and/or reduce immunogenicity of the antibody and/or enhance the delivery of an antibody across an epithelial barrier to the immune system. Examples of suitable effector molecules of this type include polymers, albumin, albumin binding proteins or albumin binding compounds
5 such as those described in WO 2005/117984.

The term "humanised antibody" is intended to refer to CDR-grafted antibody molecules in which CDR sequences derived from the germline of another mammalian species, such as a mouse, have been grafted onto human framework sequences. Additional framework
10 region modifications may be made within the human framework sequences.

As used herein, the term 'CDR-grafted antibody molecule' refers to an antibody molecule wherein the heavy and/or light chain contains one or more CDRs (including, if desired, one or more modified CDRs) from a donor antibody (e.g. a murine or rat monoclonal
15 antibody) grafted into a heavy and/or light chain variable region framework of an acceptor antibody (e.g. a human antibody). For a review, see Vaughan *et al*, Nature Biotechnology, 16, 535-539, 1998. In one embodiment rather than the entire CDR being transferred, only one or more of the specificity determining residues from any one of the CDRs described herein above are transferred to the human antibody framework (see for
20 example, Kashmiri *et al.*, 2005, Methods, 36, 25-34). In one embodiment only the specificity determining residues from one or more of the CDRs described herein above are transferred to the human antibody framework. In another embodiment only the specificity determining residues from each of the CDRs described herein above are transferred to the human antibody framework.

25 When the CDRs or specificity determining residues are grafted, any appropriate acceptor variable region framework sequence may be used having regard to the class/type of the donor antibody from which the CDRs are derived, including mouse, primate and human framework regions. Suitably, the CDR-grafted antibody for use in the present invention
30 has a variable domain comprising human acceptor framework regions as well as one or more of the CDRs or specificity determining residues described above. Thus, provided in one embodiment is a neutralising CDR-grafted antibody wherein the variable domain comprises human acceptor framework regions and non-human donor CDRs.

35 Examples of human frameworks which can be used in the present invention are KOL, NEWM, REI, EU, TUR, TEI, LAY and POM (Kabat *et al.*, *supra*). For example, KOL and NEWM can be used for the heavy chain, REI can be used for the light chain and EU, LAY

and POM can be used for both the heavy chain and the light chain. Alternatively, human germline sequences may be used; these are available for example at:

<http://www.vbase2.org/> (see Retter *et al.*, Nucl. Acids Res. (2005) 33 (supplement 1), D671-D674).

5

In a CDR-grafted antibody for use in the present invention, the acceptor heavy and light chains do not necessarily need to be derived from the same antibody and may, if desired, comprise composite chains having framework regions derived from different chains.

- 10 Also, in a CDR-grafted antibody for use in the present invention, the framework regions need not have exactly the same sequence as those of the acceptor antibody. For instance, unusual residues may be changed to more frequently occurring residues for that acceptor chain class or type. Alternatively, selected residues in the acceptor framework regions may be changed so that they correspond to the residue found at the same
- 15 position in the donor antibody (see Reichmann *et al.*, 1998, Nature, 332, 323-324). Such changes should be kept to the minimum necessary to recover the affinity of the donor antibody. A protocol for selecting residues in the acceptor framework regions which may need to be changed is set forth in WO 91/09967.

- 20 It will also be understood by one skilled in the art that antibodies may undergo a variety of posttranslational modifications. The type and extent of these modifications often depends on the host cell line used to express the antibody as well as the culture conditions. Such modifications may include variations in glycosylation, methionine oxidation, diketopiperazine formation, aspartate isomerization and asparagine deamidation. A
- 25 frequent modification is the loss of a carboxy-terminal basic residue (such as lysine or arginine) due to the action of carboxypeptidases (as described in Harris, R.J. *Journal of Chromatography* 705:129-134, 1995).

- In one embodiment the antibody heavy chain comprises a CH1 domain and the antibody
- 30 light chain comprises a CL domain, either kappa or lambda.

- Biological molecules, such as antibodies or fragments, contain acidic and/or basic functional groups, thereby giving the molecule a net positive or negative charge. The amount of overall "observed" charge will depend on the absolute amino acid sequence of
- 35 the entity, the local environment of the charged groups in the 3D structure and the environmental conditions of the molecule. The isoelectric point (pI) is the pH at which a particular molecule or surface carries no net electrical charge. In one embodiment the

antibody or fragment according to the present disclosure has an isoelectric point (pI) of at least 7. In one embodiment the antibody or fragment has an isoelectric point of at least 8, such as 8.5, 8.6, 8.7, 8.8 or 9. In one embodiment the pI of the antibody is 8. Programs such as ** ExPASy http://www.expasy.ch/tools/pi_tool.html (see Walker, The Proteomics
5 Protocols Handbook, Humana Press (2005), 571-607) may be used to predict the isoelectric point of the antibody or fragment.

Antibodies for use in the invention may comprise at least one, at least two or all three heavy chain CDR sequences of SEQ ID NOS: 4 to 6 (HCDR1/HCDR2/HCDR3
10 respectively). These are the HCDR1/HCDR2/HCDR3 sequences of the Ab7326 antibody of the Examples as determined using Kabat methodology.

The Kabat and Chothia methods for determining CDR sequences are well known in the art (as well as other techniques). CDR sequences may be determined using any
15 appropriate method and in the present invention, whilst Kabat is typically employed, other techniques could be used as well. In the present instance, SEQ ID NO: 3 presents the Ab7326 HCDR1 sequence as determined using a combined Chothia & Kabat definition.

Antibodies for use in the invention may comprise at least one, at least two or all three light
20 chain CDR sequences of SEQ ID NOS: 7 to 9 (LCDR1/LCDR2/LCDR3 respectively). These are the LCDR1/LCDR2/LCDR3 sequences of Ab7326 using Kabat methodology.

In one embodiment, the antibody comprises at least a HCDR3 sequence of SEQ ID NO:
25 6.

Typically, the antibody comprises at least one heavy chain CDR sequence selected from SEQ ID NOS: 4 to 6 and at least one light chain CDR sequence selected from SEQ ID NOS 7 to 9. The antibody may comprise at least two heavy chain CDR sequences selected from SEQ ID NOS: 4 to 6 and at least two light chain CDR sequences selected
30 from SEQ ID NOS: 7 to 9. The antibody typically comprises all three heavy chain CDR sequences of SEQ ID NOS: 4 to 6 (HCDR1/HCDR2/HCDR3 respectively) and all three light chain CDR sequences SEQ ID NOS: 7 to 9 (LCDR1/LCDR2/LCDR3 respectively). The antibodies may be chimeric, human or humanised antibodies.

35 The antibody may comprise a heavy chain variable region (HCVR) sequence of SEQ ID NO: 10 or 12 (the HCVR of Ab7326 variants 1 and 2). The antibody may comprise a light chain variable region (LCVR) sequence of SEQ ID NO: 11 or 13 (the LCVR of Ab7326

variants 1 and 2). The antibody preferably comprises the heavy chain variable region sequence of SEQ ID NO: 10 or 12 and the light chain variable region sequence of SEQ ID NO: 11 or 13 (especially HCVR/LVCR pairs of SEQ ID NOs: 10/11 or 12/13).

- 5 Ab7326 variants 1 and 2 differ by a single amino acid in the heavy chain variable region, and by a single amino acid in the light chain variable region, as follows:
- Heavy chain variable region variant 1 has glutamic acid (E) at position 6. (SEQ ID NO:10)
 - Heavy chain variable region variant 2 has glutamine (Q) at position 6. (SEQ ID NO:12)
 - 10 • Light chain variable region variant 1 has serine (S) at position 7. (SEQ ID NO:11)
 - Light chain variable region variant 2 has threonine (T) at position 7. (SEQ ID NO:13)

Thus, in one embodiment, the antibody comprises a heavy chain variable region (HCVR) sequence of SEQ ID NO: 10, wherein the glutamic acid residue at position 6 is substituted
15 with a glutamine residue (E6Q); wherein the residue numbering is according to SEQ ID NO: 10.

In one embodiment, the antibody comprises a heavy chain variable region (HCVR) sequence of SEQ ID NO: 12, wherein the glutamine residue at position 6 is substituted
20 with a glutamic acid residue (Q6E); wherein the residue numbering is according to SEQ ID NO: 12.

In one embodiment, the antibody comprises a light chain variable region (LCVR) sequence of SEQ ID NO: 11, wherein the serine residue at position 7 is substituted with a
25 threonine residue (S7T); wherein the residue numbering is according to SEQ ID NO: 11.

In one embodiment, the antibody comprises a light chain variable region (LCVR) sequence of SEQ ID NO: 13, wherein the threonine residue at position 7 is substituted with a serine residue (T7S); wherein the residue numbering is according to SEQ ID NO:
30 13.

In one embodiment, the antibody comprises the sequence of SEQ ID NO: 3 or 4 for HCDR1, the sequence of SEQ ID NO: 5 for HCDR2, the sequence of SEQ ID NO: 6 for HCDR3, the sequence of SEQ ID NO: 7 for LCDR1, the sequence of SEQ ID NO: 8 for
35 LCDR2 and the sequence of SEQ ID NO: 9 for LCDR3; and wherein the heavy chain variable region comprises a sequence having at least 95% identity, (e.g. at least 95%, 96%, 97%, 98% or 99% identity), to the sequence of SEQ ID NO: 10 and the light chain

variable region comprises a sequence having at least 95% identity, (e.g. at least 95%, 96%, 97%, 98% or 99% identity), to the sequence of SEQ ID NO: 11.

In one embodiment, the antibody comprises the sequence of SEQ ID NO: 3 or 4 for
5 HCDR1, the sequence of SEQ ID NO: 5 for HCDR2, the sequence of SEQ ID NO: 6 for
HCDR3, the sequence of SEQ ID NO: 7 for LCDR1, the sequence of SEQ ID NO: 8 for
LCDR2 and the sequence of SEQ ID NO: 9 for LCDR3; and wherein the heavy chain
variable region comprises a sequence having at least 95% identity, (e.g. at least 95%,
96%, 97%, 98% or 99% identity), to the sequence of SEQ ID NO: 12 and the light chain
10 variable region comprises a sequence having at least 95% identity, (e.g. at least 95%,
96%, 97%, 98% or 99% identity) to the sequence of SEQ ID NO: 13.

The antibody may comprise a heavy chain (H-chain) sequence of
SEQ ID NO: 14 mouse full length IgG1 heavy chain variant 1, or
15 SEQ ID NO: 28 mouse full length IgG1 heavy chain variant 2, or
SEQ ID NO: 30 human full length IgG1 heavy chain variant 1, or
SEQ ID NO: 16 human full length IgG1 heavy chain variant 2, or
SEQ ID NO: 22 human full length IgG4P heavy chain variant 1, or
SEQ ID NO: 34 human full-length IgG4P heavy chain variant 2, or
20 SEQ ID NO: 18 Fab heavy chain variant 1, or
SEQ ID NO: 32 Fab heavy chain variant 2.

The antibody may comprise a light chain (L-chain) sequence of
SEQ ID NO: 15 mouse full length IgG1 light chain variant 1, or
25 SEQ ID NO: 29 mouse full length IgG1 light chain variant 2, or
SEQ ID NO: 31 human full length IgG1 light chain variant 1, or
SEQ ID NO: 17 human full length IgG1 light chain variant 2, or
SEQ ID NO: 23 human full length IgG4P light chain variant 1, or
SEQ ID NO: 35 human full-length IgG4P light chain variant 2, or
30 SEQ ID NO: 19 Fab light chain variant 1, or
SEQ ID NO: 33 Fab light chain variant 2

In one example, the antibody comprises a heavy chain / light chain sequence pair of
SEQ ID NOs: 14/15 mouse full length IgG1 variant 1, or
35 SEQ ID NOs: 28/29 mouse full length IgG1 variant 2, or
SEQ ID NOs: 30/31 human full length IgG1 variant 1, or
SEQ ID NOs: 16/17 human full length IgG1 variant 2, or

SEQ ID NOs: 22/23 human full length IgG4P variant 1, or
 SEQ ID NOs: 34/35 human full-length IgG4P variant 2, or
 SEQ ID NOs: 18/19 Fab light chain variant 1, or
 SEQ ID NOs: 32/33 Fab light chain variant 2

5

The variant forms of corresponding sequences may be interchanged. For example, the antibody may comprise a heavy chain / light chain sequence pair of

SEQ ID NOs: 14/29 mouse full length IgG1 heavy chain variant 1/light chain variant 2, or
 SEQ ID NOs: 28/15 mouse full length IgG1 heavy chain variant 2/light chain variant 1, or
 10 SEQ ID NOs: 30/17 human full length IgG1 heavy chain variant 1/light chain variant 2, or
 SEQ ID NOs: 16/31 human full length IgG1 heavy chain variant 2/light chain variant 1, or
 SEQ ID NOs: 22/35 human full length IgG4P heavy chain variant 1/light chain variant 2, or
 SEQ ID NOs: 34/23 human full-length IgG4P heavy chain variant 2/light chain variant 1, or
 SEQ ID NOs: 18/33 Fab heavy chain variant 1/light chain variant 2, or
 15 SEQ ID NOs: 32/19 Fab heavy chain variant 2/light chain variant 1.

The antibodies may be chimeric, human or humanised antibodies.

The antibody may alternatively be or may comprise a variant of one of the specific
 20 sequences recited above. For example, a variant may be a substitution, deletion or
 addition variant of any of the above amino acid sequences.

A variant antibody may comprise 1, 2, 3, 4, 5, up to 10, up to 20 or more (typically up to a
 maximum of 50) amino acid substitutions and/or deletions from the specific sequences
 25 discussed above. "Deletion" variants may comprise the deletion of individual amino acids,
 deletion of small groups of amino acids such as 2, 3, 4 or 5 amino acids, or deletion of
 larger amino acid regions, such as the deletion of specific amino acid domains or other
 features. "Substitution" variants typically involve the replacement of one or more amino
 acids with the same number of amino acids and making conservative amino acid
 30 substitutions. For example, an amino acid may be substituted with an alternative amino
 acid having similar properties, for example, another basic amino acid, another acidic
 amino acid, another neutral amino acid, another charged amino acid, another hydrophilic
 amino acid, another hydrophobic amino acid, another polar amino acid, another aromatic
 amino acid or another aliphatic amino acid. Some properties of the 20 main amino acids
 35 which can be used to select suitable substituents are as follows:

Table 1: Amino acid properties.

Ala	aliphatic, hydrophobic, neutral	Met	hydrophobic, neutral
Cys	polar, hydrophobic, neutral	Asn	polar, hydrophilic, neutral
Asp	polar, hydrophilic, charged (-)	Pro	hydrophobic, neutral
Glu	polar, hydrophilic, charged (-)	Gln	polar, hydrophilic, neutral
Phe	aromatic, hydrophobic, neutral	Arg	polar, hydrophilic, charged (+)
Gly	aliphatic, neutral	Ser	polar, hydrophilic, neutral
His	aromatic, polar, hydrophilic, charged (+)	Thr	polar, hydrophilic, neutral
Ile	aliphatic, hydrophobic, neutral	Val	aliphatic, hydrophobic, neutral
Lys	polar, hydrophilic, charged(+)	Trp	aromatic, hydrophobic, neutral
Leu	aliphatic, hydrophobic, neutral	Tyr	aromatic, polar, hydrophobic

"Derivatives" or "variants" generally include those in which instead of the naturally occurring amino acid the amino acid which appears in the sequence is a structural analog thereof. Amino acids used in the sequences may also be derivatized or modified, e.g. labelled, providing the function of the antibody is not significantly adversely affected.

Derivatives and variants as described above may be prepared during synthesis of the antibody or by post- production modification, or when the antibody is in recombinant form using the known techniques of site- directed mutagenesis, random mutagenesis, or enzymatic cleavage and/or ligation of nucleic acids.

Variant antibodies may have an amino acid sequence which has more than about 60%, or more than about 70%, e.g. 75 or 80%, typically more than about 85%, e.g. more than about 90 or 95% amino acid identity to the amino acid sequences disclosed herein (particularly the HCVR/LCVR sequences and the H- and L-chain sequences). Furthermore, the antibody may be a variant which has more than about 60%, or more than about 70%, e.g. 75 or 80%, typically more than about 85%, e.g. more than about 90 or 95% amino acid identity to the HCVR/LCVR sequences and the H- and L-chain sequences disclosed herein, whilst retaining the exact CDRs disclosed for these sequences. Variants may retain at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%,

97%, 98% or 99% identity to the HCVR/LCVR sequences and to the H- and L-chain sequences disclosed herein (in some circumstances whilst retaining the exact CDRs).

- 5 Variants typically retain about 60% - about 99% identity, about 80% - about 99% identity, about 90% - about 99% identity or about 95% - about 99% identity. This level of amino acid identity may be seen across the full length of the relevant SEQ ID NO sequence or over a part of the sequence, such as across about 20, 30, 50, 75, 100, 150, 200 or more amino acids, depending on the size of the full length polypeptide.
- 10 In connection with amino acid sequences, "sequence identity" refers to sequences which have the stated value when assessed using ClustalW (Thompson *et al.*, 1994, *supra*) with the following parameters:
Pairwise alignment parameters -Method: accurate, Matrix: PAM, Gap open penalty: 10.00, Gap extension penalty: 0.10;
- 15 Multiple alignment parameters -Matrix: PAM, Gap open penalty: 10.00, % identity for delay: 30, Penalize end gaps: on, Gap separation distance: 0, Negative matrix: no, Gap extension penalty: 0.20, Residue-specific gap penalties: on, Hydrophilic gap penalties: on, Hydrophilic residues: GPSNDQEKR. Sequence identity at a particular residue is intended to include identical residues which have simply been derivatized.

20

Antibodies having specific sequences and derivatives and variants which maintain the function or activity of these chains are therefore provided for use in the present invention.

- 25 "Derivatives" as used herein is intended to include reactive derivatives, for example thiol-selective reactive groups such as maleimides. The reactive group may be linked directly or through a linker segment to a polymer. It will be appreciated that the residue of such a group will in some instances form part of the product as the linking group between the antibody fragment and the polymer.

- 30 The polymer may be a synthetic or a naturally occurring polymer, for example an optionally substituted straight or branched chain polyalkylene, polyalkenylene or polyoxyalkylene polymer or a branched or unbranched polysaccharide, e.g. a homo- or hetero- polysaccharide.

- Specific optional substituents which may be present on a synthetic polymer include one or
35 more hydroxy, methyl or methoxy groups.

Specific examples of synthetic polymers include optionally substituted straight or branched chain poly(ethyleneglycol), poly(propyleneglycol) poly(vinylalcohol) or

derivatives thereof, especially optionally substituted poly(ethyleneglycol) such as methoxypoly(ethyleneglycol) or derivatives thereof.

Specific naturally occurring polymers include lactose, amylose, dextran, glycogen or derivatives thereof.

5

The size of the polymer may be varied as desired, but will generally be in an average molecular weight range from 500Da to 50000Da, for example from 5000 to 40000Da such as from 20000 to 40000Da. The polymer size may in particular be selected on the basis of the intended use of the product for example ability to localize to certain tissues or
10 extend circulating half-life (for review see Chapman, 2002, Advanced Drug Delivery Reviews, 54, 531-545). Thus, for example, where the product is intended to leave the circulation and penetrate tissue, it may be advantageous to use a small molecular weight polymer, for example with a molecular weight of around 5000Da. For applications where the product remains in the circulation, it may be advantageous to use a higher molecular
15 weight polymer, for example having a molecular weight in the range from 20000Da to 40000Da.

Suitable polymers include a polyalkylene polymer, such as a poly(ethyleneglycol) or, especially, a methoxypoly(ethyleneglycol) or a derivative thereof, and especially with a
20 molecular weight in the range from about 15000Da to about 40000Da.

In one example antibodies for use in the present invention are attached to poly(ethyleneglycol) (PEG) moieties. In one particular example the antibody is an antibody fragment and the PEG molecules may be attached through any available amino
25 acid side-chain or terminal amino acid functional group located in the antibody fragment, for example any free amino, imino, thiol, hydroxyl or carboxyl group. Such amino acids may occur naturally in the antibody fragment or may be engineered into the fragment using recombinant DNA methods (see for example US 5,219,996; US 5,667,425; WO98/25971, WO2008/038024). In one example the antibody molecule is a modified Fab
30 fragment wherein the modification is the addition to the C-terminal end of its heavy chain one or more amino acids to allow the attachment of an effector molecule. Suitably, the additional amino acids form a modified hinge region containing one or more cysteine residues to which the effector molecule may be attached. Multiple sites can be used to attach two or more PEG molecules.

35

Antibodies may compete for binding to Gremlin-1 with, or bind to the same epitope as, those defined above in terms of H-chain/L-chain, HCVR/LCVR or CDR sequences. In

particular, an antibody may compete for binding to Gremlin-1 with, or bind to the same epitope as, an antibody which comprises a HCDR1/HCDR2/HCDR3/LCDR1/LCDR2/LCDR3 sequence combination of SEQ ID NOs: 4/5/6/7/8/9. An antibody may compete for binding to Gremlin-1 with, or bind to the same epitope as, an antibody which comprises a HCVR and LCVR sequence pair of SEQ ID NOs: 10/11 or 12/13 or full length chains of SEQ ID Nos: 14/15 or 16/17.

An "epitope" is a region of an antigen that is bound by an antibody. Epitopes may be defined as structural or functional. Functional epitopes are generally a subset of the structural epitopes and have those residues that directly contribute to the affinity of the interaction. Epitopes may also be conformational, that is, composed of non-linear amino acids. In certain embodiments, epitopes may include determinants that are chemically active surface groupings of molecules such as amino acids, sugar side chains, phosphoryl groups, or sulfonyl groups, and, in certain embodiments, may have specific three-dimensional structural characteristics, and/or specific charge characteristics.

One can easily determine whether an antibody binds to the same epitope as, or competes for binding with, a reference antibody by using routine methods known in the art. For example, to determine if a test antibody binds to the same epitope as a reference antibody for use in the invention, the reference antibody is allowed to bind to a protein or peptide under saturating conditions. Next, the ability of a test antibody to bind to the protein or peptide is assessed. If the test antibody is able to bind to the protein or peptide following saturation binding with the reference antibody, it can be concluded that the test antibody binds to a different epitope than the reference antibody. On the other hand, if the test antibody is not able to bind to protein or peptide following saturation binding with the reference antibody, then the test antibody may bind to the same epitope as the epitope bound by the reference antibody of the invention.

To determine if an antibody competes for binding with a reference antibody, the above-described binding methodology is performed in two orientations. In a first orientation, the reference antibody is allowed to bind to a protein/peptide under saturating conditions followed by assessment of binding of the test antibody to the protein/peptide molecule. In a second orientation, the test antibody is allowed to bind to the protein/peptide under saturating conditions followed by assessment of binding of the reference antibody to the protein/peptide. If, in both orientations, only the first (saturating) antibody is capable of binding to the protein/peptide, then it is concluded that the test antibody and the reference antibody compete for binding to the protein/peptide. As will be appreciated by the skilled

person, an antibody that competes for binding with a reference antibody may not necessarily bind to the identical epitope as the reference antibody, but may sterically block binding of the reference antibody by binding an overlapping or adjacent epitope.

- 5 Two antibodies bind to the same or overlapping epitope if each competitively inhibits (blocks) binding of the other to the antigen. That is, a 1-, 5-, 10-, 20- or 100-fold excess of one antibody inhibits binding of the other by at least 50%, 75%, 90% or even 99% as measured in a competitive binding assay (see, e.g., Junghans et al., Cancer Res, 1990:50:1495-1502). Alternatively, two antibodies have the same epitope if essentially all
- 10 amino acid mutations in the antigen that reduce or eliminate binding of one antibody reduce or eliminate binding of the other. Two antibodies have overlapping epitopes if some amino acid mutations that reduce or eliminate binding of one antibody reduce or eliminate binding of the other.
- 15 Additional routine experimentation (e.g., peptide mutation and binding analyses) can then be carried out to confirm whether the observed lack of binding of the test antibody is in fact due to binding to the same epitope as the reference antibody or if steric blocking (or another phenomenon) is responsible for the lack of observed binding. Experiments of this sort can be performed using ELISA, RIA, surface plasmon resonance, flow cytometry or
- 20 any other quantitative or qualitative antibody-binding assay available in the art.

The anti-gremlin-1 antibody of the Examples, Ab7326, has been found to bind the following residues of Gremlin-1: Ile131, Lys147, Lys148, Phe149, Thr150, Thr151, Arg169, Lys174 and Gln175; where Lys147, Lys148, Phe149, Thr150, Thr151, Arg169,

25 Lys174 and Gln175 are present on one Gremlin-1 monomer and Ile131 is present on the second Gremlin-1 monomer. The numbering is based on the UniProt entry O60565 of SEQ ID NO: 1. As discussed in the Examples section, these epitope residues were identified using NCONT analysis at 4 Å from the Gremlin-1-Ab7326 Fab complex.

- 30 Antibodies for use in the invention may therefore bind to an epitope which comprises at least one residue selected from Ile131, Lys147, Lys148, Phe149, Thr150, Thr151, Arg169, Lys174 and Gln175 (with residue numbering based on SEQ ID NO: 1). Antibodies for use in the invention may bind an epitope which comprises 2, 3, 4, 5, 6, 7, 8 or all 9 of these residues (preferably at least 5 residues).

35

Antibodies for use in the invention may also recognise an epitope where Ile131 is present on a different Gremlin-1 monomer to the other residues.

Although these residues are provided for a particular sequence of human Gremlin-1, the skilled person could extrapolate the positions of these residues to other corresponding Gremlin sequences using routine techniques. Antibodies binding to epitopes comprising
5 the corresponding residues within these other Gremlin sequences are therefore also provided for use in the invention.

To screen for antibodies that bind to a particular epitope, a routine cross-blocking assay such as that described in Antibodies, Harlow and Lane (Cold Spring Harbor Press, Cold
10 Spring Harb., NY) can be performed. Other methods include alanine scanning mutants, peptide blots (Reineke (2004) Methods Mol Biol 248:443-63), or peptide cleavage analysis. In addition, methods such as epitope excision, epitope extraction and chemical modification of antigens can be employed (Tomer (2000) Protein Science 9: 487-496). Such methods are well known in the art.

15 Antibody epitopes may also be determined by x-ray crystallography analysis. Antibodies for use in the present invention may therefore be assessed through x-ray crystallography analysis of the antibody bound to Gremlin-1. Epitopes may, in particular, be identified in this way by determining residues on Gremlin-1 within 4Å of an antibody paratope residue.

20 Antibodies can be tested for binding to Gremlin-1 by, for example, standard ELISA or Western blotting. An ELISA assay can also be used to screen for hybridomas that show positive reactivity with the target protein. The binding selectivity of an antibody may also be determined by monitoring binding of the antibody to cells expressing the target protein,
25 for example by flow cytometry. Thus, a screening method may comprise the step of identifying an antibody that is capable of binding Gremlin-1 by carrying out an ELISA or Western blot or by flow cytometry.

Antibodies may selectively (or specifically) recognise Gremlin-1. An antibody, or other
30 compound, "selectively binds" or "selectively recognises" a protein when it binds with preferential or high affinity to the protein for which it is selective but does not substantially bind, or binds with low affinity, to other proteins. The selectivity of an antibody may be further studied by determining whether or not the antibody binds to other related proteins as discussed above or whether it discriminates between them. Antibodies for use in the
35 invention typically recognise human Gremlin-1.

Antibodies may also have cross-reactivity for related proteins, or for human Gremlin-1 and for Gremlin-1 from other species.

By specific (or selective), it will be understood that the antibody binds to the protein of interest with no significant cross-reactivity to any other molecule. Cross-reactivity may be assessed by any suitable method described herein. Cross-reactivity of an antibody may be considered significant if the antibody binds to the other molecule at least about 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90% or 100% as strongly as it binds to the protein of interest. An antibody that is specific (or selective) may bind to another molecule at less than about 90%, 85%, 80%, 75%, 70%, 65%, 60%, 55%, 50%, 45%, 40%, 35%, 30%, 25% or 20% the strength that it binds to the protein of interest. The antibody may bind to the other molecule at less than about 20%, less than about 15%, less than about 10% or less than about 5%, less than about 2% or less than about 1% the strength that it binds to the protein of interest.

Thus, antibodies suitable for use in the present invention may have a high affinity binding for (human) Gremlin-1. The antibody may have a dissociation constant (K_D) of less than <1 nM, and preferably <500 pM. In one example, the antibody has a dissociation constant (K_D) of less than 200pM. In one example, the antibody has a dissociation constant (K_D) of less than 100pM. A variety of methods can be used to determine the binding affinity of an antibody for its target antigen such as surface plasmon resonance assays, saturation assays, or immunoassays such as ELISA or RIA, as are well known to persons of skill in the art. An exemplary method for determining binding affinity is by surface plasmon resonance analysis on a BIAcore™ 2000 instrument (Biacore AB, Freiburg, Germany) using CM5 sensor chips, as described by Krinner et al., (2007) Mol. Immunol. February; 44 (5):916-25. (Epub 2006 May 11)).

The anti-Gremlin-1 antibody of the Examples, Ab7326, is an allosteric inhibitor of Gremlin-1 activity which binds to an epitope distal from the BMP binding site. (WO 2018/115017 A2) Ab7326 binds to Gremlin-1 with exceptionally high affinity with a K_d value <100pM, and is expected to be particularly useful for use in the present invention.

An inhibitor of gremlin-1 activity may have an effect on any of the functions of Gremlin-1, but typically reduces binding of Gremlin-1 to BMP (BMP 2, 4, and/or 7). Gremlin-1 is a negative regulator of BMP and so reduced binding increases signalling through BMP.

- BMP binding and signalling may be detected by any method known in the art. The Examples of the present application describe two functional assays for testing whether an agent reduces binding of gremlin-1 to BMP. Example 3 describes an Id1 reporter gene assay, where the Id1 gene is a target gene of BMP signalling. An increase in the signal in this assay may be used to determine if an agent reduces Gremlin-1 binding to BMP. Example 5 describes a SMAD phosphorylation assay. SMAD1, 5 and 8 are phosphorylated upon BMP signalling. An increase in SMAD phosphorylation may therefore be used to determine whether an agent reduces binding of Gremlin-1 to BMP.
- Once a suitable antibody has been identified and selected, the amino acid sequence of the antibody may be identified by methods known in the art. The genes encoding the antibody can be cloned using degenerate primers. The antibody may be recombinantly produced by routine methods.
- Examples of DNA sequences encoding full length heavy chains and light chains of Ab7326 are provided in the sequence listing:
- SEQ ID NO: 24 (Human IgG1 heavy chain DNA variant 1)
 - SEQ ID NO: 25 (Human IgG1 light chain DNA variant 1)
 - SEQ ID NO: 26 (Human IgG4P heavy chain DNA variant 1)
 - SEQ ID NO: 27 (Human IgG4P light chain DNA variant 1)

Pharmaceutical Compositions, Dosages and Dosage Regimes

- An inhibitor of gremlin-1 activity for use in the present invention may be provided in a pharmaceutical composition. The pharmaceutical composition will normally be sterile and will typically include a pharmaceutically acceptable carrier and/or adjuvant. A pharmaceutical composition for use in the invention may additionally comprise a pharmaceutically acceptable adjuvant and/or carrier.
- The pharmaceutical compositions for use in the invention may include one or more pharmaceutically acceptable salts. A "*pharmaceutically acceptable salt*" refers to a salt that retains the desired biological activity of the parent molecule and does not impart any undesired toxicological effects. Examples of such salts include acid addition salts and base addition salts.
- As used herein, "*pharmaceutically acceptable carrier*" includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption

delaying agents, and the like that are physiologically compatible. The carrier may be suitable for parenteral, e.g. intravenous, intramuscular, intradermal, intraocular, intraperitoneal, subcutaneous, spinal or other parenteral routes of administration, for example by injection or infusion. Alternatively, the carrier may be suitable for non-
5 parenteral administration, such as a topical, epidermal or mucosal route of administration. The carrier may be suitable for oral administration. Depending on the route of administration, the inhibitor may be coated in a material to protect it from the action of acids and other natural conditions that may inactivate the inhibitor.

10 Pharmaceutically acceptable carriers comprise aqueous carriers or diluents. Examples of suitable aqueous carriers that may be employed in the pharmaceutical compositions for use in the invention include water, buffered water and saline. Examples of other carriers include ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), and suitable mixtures thereof, vegetable oils, such as olive oil, and injectable organic
15 esters, such as ethyl oleate. In many cases, it will be desirable to include isotonic agents, for example, sugars, polyalcohols such as mannitol, sorbitol, or sodium chloride in the composition.

Pharmaceutical compositions typically must be sterile and stable under the conditions of
20 manufacture and storage. The composition can be formulated as a solution, micro-emulsion, liposome, or other ordered structure suited to high drug concentration.

Pharmaceutical compositions for use in the invention may comprise additional active ingredients.

25 Also envisaged are kits comprising an inhibitor of gremlin-1 activity and instructions for use in a method of treatment according to the invention.

Agents for use in the invention or formulations or compositions thereof may be
30 administered for therapeutic and/or prophylactic treatments.

In therapeutic applications, agents are administered to a subject already suffering from a disorder or condition, in an amount sufficient to cure, alleviate or partially arrest the condition or one or more of its symptoms. Such therapeutic treatment may result in a
35 decrease in severity of symptoms, or an increase in frequency or duration of symptom-free periods. An amount adequate to accomplish this is defined as a "*therapeutically effective amount*".

In prophylactic applications, agents are administered to a subject at risk of a disorder or condition, in an amount sufficient to prevent or reduce the subsequent effects of the condition or one or more of its symptoms. An amount adequate to accomplish this is
5 defined as a "*prophylactically effective amount*". Effective amounts for each purpose will depend on the severity of the disease or injury as well as the weight and general state of the subject.

A subject for administration may be a human or non-human animal. The term "*non-human animal*" includes all vertebrates, e.g., mammals and non-mammals, such as non-
10 human primates, dogs, cats, horses, sheep, cows, chickens, amphibians, reptiles, etc. Administration to humans is typical.

An agent or pharmaceutical composition for use in the invention may be administered via
15 one or more routes of administration using one or more of a variety of methods known in the art. As will be appreciated by the skilled artisan, the route and/or mode of administration will vary depending upon the desired results. Examples of routes of administration for agents or pharmaceutical compositions for use in the invention include parenteral routes, such as intravenous, intramuscular, intradermal, intraocular,
20 intraperitoneal, subcutaneous, or spinal routes of administration, for example by injection or infusion. Alternatively, an agent or pharmaceutical composition can be administered via a non-parenteral route, such as a topical, epidermal or mucosal route of administration. The agent or pharmaceutical composition may be for oral administration.

25 A suitable dosage of an inhibitory agent or pharmaceutical composition for use in the invention may be determined by a skilled medical practitioner. Actual dosage levels of the active ingredients in the pharmaceutical compositions for use in the present invention may be varied so as to obtain an amount of the active ingredient that is effective to achieve the desired therapeutic response for a particular patient, composition, and mode of
30 administration, without being toxic to the patient. The selected dosage level will depend upon a variety of pharmacokinetic factors including the activity of the particular compositions employed, the route of administration, the time of administration, the rate of excretion of the particular compound being employed, the duration of the treatment, the age, sex, weight, condition, general health and prior medical history of the patient being
35 treated, and like factors well known in the medical arts.

A suitable dose may be, for example, in the range of from about 0.01 µg/kg to about 1000 mg/kg body weight, typically from about 0.1 µg/kg to about 100 mg/kg body weight, of the patient to be treated. For example, a suitable dosage may be from about 1 µg/kg to about 10 mg/kg body weight per day or from about 10 µg/kg to about 5 mg/kg body weight per day.

Dosage regimens may be adjusted to provide the optimum desired response (e.g., a therapeutic response). For example, a single dose may be administered, several divided doses may be administered over time or the dose may be proportionally reduced or increased as indicated by the therapeutic situation. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subjects to be treated; each unit contains a predetermined quantity of active agent calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier.

Administration may be in single or multiple doses. Multiple doses may be administered via the same or different routes and to the same or different locations. Alternatively, doses can be via a sustained release formulation, in which case less frequent administration is required. Dosage and frequency may vary depending on the half-life of the inhibitory agent in the patient and the duration of treatment desired.

Agents, formulations or pharmaceutical compositions for use in the invention may be co-administered with one or other more other therapeutic agents. Combined administration of two or more agents may be achieved in a number of different ways. Both may be administered together in a single composition, or they may be administered in separate compositions as part of a combined therapy. For example, the one may be administered before, after or concurrently with the other.

Therapeutic indications

Inhibitors of gremlin-1 activity according to the present invention are provided for the treatment of a bone fracture or bone defect. A bone fracture is a break or crack in bone tissue and may be the result of a traumatic injury, such as a fall or impact, but can also occur as a result of diseases that affect bone integrity. A bone defect is a loss of bone, due to trauma or disease.

The fracture may be a fracture of any bone in the body.

The bone defect may be a bone defect in any bone of the body.

In one embodiment, the bone fracture is a delayed-union or non-union fracture. A delayed-union fracture is defined as a fracture which fails to reach union within 6-months post-fracture. A non-union fracture is defined as incomplete healing within 9 months, combined with a lack of radiological characteristics associated with fracture healing being observed over three consecutive months. (Einhorn et al; 2014; Buza et al; 2016). Examples of fractures that are that are prone to delayed-union or non-union development include tibia, distal radius, femoral neck and scaphoid.

In one embodiment, the bone fracture or bone defect occurs as a result of a disease that affects bone integrity. Examples of diseases that affect bone integrity include but are not limited to osteoporosis, osteogenesis imperfecta, diabetes, Paget's disease of bone, rheumatoid arthritis, ankylosing spondylitis, multiple myeloma, primary bone cancer (e.g. osteosarcoma, Ewing's sarcoma and chondrosarcoma), cancers that metastasise to the bone (e.g. breast cancer, prostate cancer and lung cancer), diffuse idiopathic skeletal hyperostosis, osteomyelitis, renal disease, Duchenne muscular dystrophy and thalassemia major.

FIGURES

Figure 1 shows percentage restoration of signal for the immunisation derived antibodies in the HEK-ID1 reporter gene assay.

Figure 2 shows percentage restoration of signal for library derived antibodies in the HEK-ID1 reporter gene assay.

Figure 3 shows results for the HEK-ID1 reporter gene assay with titrations of human Gremlin (Figure 3A) and mouse Gremlin (Figure 3B) and the effect of antibody 7326 (shown as antibody PB376) in restoring signalling of BMP.

Figure 4 shows a structural model of the Gremlin-Fab complex, with the possible BMP binding regions and the Fab epitope highlighted.

Figure 5 shows examination of the area devoid of callus/bone tissue during fracture repair in acquired X-Ray images. The area within the defect, which was devoid of tissue, was

quantified using definiens image analysis and subsequently compared between control and anti-gremlin 1 treated group. Results are presented as the mean $\pm$ SD of 10 rats/group. *P<0.05; **P< 0.01; ***P<0.001 as measured by Mann-Whitney U test.

5 Figure 6 shows examination of LMB (low mineral bone; newly formed bone) and HMB (high mineral bone; mature bone) within a 3mm femoral bone defect. Panel A: 3D μ Ct analysis of femoral bone defect region to detect newly formed bone or mature bone. Percentage of bone volume/tissue volume was measured and compared all subjects (total) between control v.s treatments with anti-gremlin 1. Comparisons between control
10 versus anti-gremlin 1 treated group in animals separated as low responders (LR- incomplete bridging) and high responders (HR complete bridging) were also performed. Results are presented as the mean $\pm$ SD. *P<0.05; **P< 0.01; ***P<0.001 as measured by Mann-Whitney U test. Panel B: representative μ Ct illustrating the 3D bone volume renderings of LR and HR groups in control and after anti-gremlin 1 treatment.

15 Figure 7 shows histomorphometric analysis of femoral bone defect. Percentage of bone volume/tissue volume (BV/TV (%)), trabecular number (Tb.N) and trabecular separation (Tb.Sp) was compared between control versus anti-gremlin 1 treated group. Results are presented as the mean $\pm$ SD of 10 rats/group. *P<0.05; **P< 0.01; ***P<0.001 as
20 measured by Mann-Whitney U test.

Figure 8 shows correlation of 3D μ CT analysis and 2D histomorphometry analysis of total BV/TV%. Correlations were performed in both groups on LMB (low mineral bone; newly formed bone) and HMB (high mineral bone; mature bone) within a 3mm femoral bone
25 defect and compared to 2D histomorphometry score data (n=20). The Pearson's score indicates significant correlation of BV/TV% between 3D μ CT analysis and 2D histomorphometry analysis.

The following Examples illustrate the invention.

30

EXAMPLES

Example 1- Protein expression, purification, refolding and structure determination.

35

Protein expression and inclusion body preparation

A truncated human Gremlin-1 coding sequence (SEQ ID NO: 20), optimised for expression in *E. coli*, was cloned into a modified pET32a vector (Merck Millipore) using BamHI/XhoI, generating a vector encoding the Gremlin sequence with an N-terminal 6His-TEV tag (pET-hGremlin1).

5

Expressed sequence:

MGSSHHHHHSSGENLYFQGSAMPGEEVLESSQEALHVTERKYLKRDWCKTQPLKQTI
HEEGCNSRTIINRFCYGQCNSFYIPRHIRKEEGSFQSCSFCKPKKFTTMMVTLNCPQLP
PTKKKRVTTRVKQCRCISIDLD; SEQ ID NO: 2 (with non-Gremlin residues of the 6His-TEV
tag shown in italics). Sequence numbering based on UniProt O60565 & SEQ ID NO: 1.

10

The pET-hGremlin1 plasmid DNA was used to transform BL21(DE3) cells. A single ampicillin resistant colony was picked from a LB/Amp agar plate and used to inoculate a 100 ml starter culture of LB/Amp. After shaking (200 rpm) for 16 hr at 37 °C, 25 ml of the starter culture was used to inoculate 500 mL of 2xTY/Amp media. The culture was shaken (250 rpm) at 37 °C until an OD₆₀₀ of 3 was achieved. Subsequently, the culture was supplemented with 20 mL of a MOPS + glycerol feed mix (1M MOPS pH 7.4, 40 % glycerol, 0.5 % MgSO₄, 0.42 % MgCl₂), induced with 300 µM IPTG and further incubated at 17 °C, 180 rpm for 16 hours. Cells were harvested in a centrifuge (4,000 g for 20 minutes at 4 °C).

15

20

Cell pellets were resuspended in Lysis Buffer (PBS pH 7.4, 0.35 mg/ml lysozyme, 10 µg/ml DNase and 3 mM MgCl₂) at 4 °C and the insoluble fraction was harvested by centrifugation at 3,500 g for 30 minutes at 4 °C. Pelleted inclusion bodies were washed three times by resuspending in wash buffer (50 mM Tris, 500 mM NaCl, 0.5 % Triton X-100, pH 8.0), followed by centrifugation at 21,000 g for 15 minutes. An additional two washes were performed using wash buffer without Triton X-100.

25

Solubilisation

30

Inclusion bodies were resuspended in denaturing buffer (8 M Urea, 100 mM Tris, 1 mM EDTA, 10 mM Na₂S₄O₆ and 100 mM Na₂SO₃, pH 8.5), stirred for 16 hrs at room-temperature and clarified by centrifugation at 21,000 g for 15 minutes.

Pre-refolding purification

35

The solubilized inclusion bodies were loaded onto a Sephacryl S-200 26/60 column (120

5 mL) equilibrated in 8 M Urea, 50 mM MES, 200 mM NaCl, 1 mM EDTA, pH 6.0. Fractions containing Gremlin-1 protein were diluted with 6 M Urea, 20 mM MES, pH 6.0 and loaded onto HiTrap SP HP cation exchange columns and eluted with a 1 M NaCl gradient over 10 column volumes (10 CVs). Fractions containing purified, denatured hGremlin-1 protein were pooled.

Refolding

10 Denatured purified Gremlin-1 protein was added drop-wise to re-folding buffer (50 mM Tris, pH 8.5, 150 mM NaCl, 5 mM GSH and 5 mM GSSG, 0.5 mM Cysteine, 5 mM EDTA, 0.5 M Arginine) to a final concentration of 0.1 mg/ml and incubated at 4 °C with constant stirring for 5 days. After 5 days the Gremlin-1 protein was dialysed against 20 mM HEPES, 100 mM NaCl, pH 7.5.

15 Following dialysis protein was applied to heparin HiTrap column and eluted using a gradient of 0-100 % heparin elution buffer (20 mM HEPES, 1 M NaCl, pH 7.5) over 20 CV. Correctly folded protein eluted at 1 M NaCl whereas any misfolded protein eluted at lower salt concentrations.

20 Protein eluting at 1 M NaCl was concentrated and purified further on a S75 26/60 column equilibrated with 20 mM Hepes, pH 7.5, 1 M NaCl.

Protein was characterised by SDS PAGE (shift in gel), demonstrated to have the expected molecular weight and correct arrangement of disulphide bonds using liquid
25 chromatography mass spectrometry (LC-MS) and to be active in a cell assay (ID1 reporter assay).

Gremlin 1 structure determination

30 Gremlin 1 protein crystals were grown using the hanging-drop method by mixing a solution of Gremlin 1 at 6.6 mg/ml and 0.1 M citric acid at pH 4, 1 M lithium chloride and 27 % polyethylene glycol (PEG) 6000 in a 1:1 ratio. Before data collection, crystals were cryo-protected by adding 20 % glycerol to the crystallization buffer. Diffraction data were collected at the Diamond Light Source and were processed using XDS (Kabsch, Wolfgang
35 (2010) Acta Crystallographica Section D 66, 125—132). Diffraction data statistics are summarized in the table below:

Table 2: Diffraction data statistics

Diffraction Statistics	
Wavelength (Å)	0.97949
Space group	C2
Cell dimensions	a= 84.55 Å, b=107.22 Å, c=77.09 Å; $\alpha=90.00^\circ$, $\beta=120.43^\circ$, $\gamma=90.00^\circ$
Resolution range* (Å)	26.19-2.72 (2.79-2.72)
Completeness (%)	98.5 (99.0)
Multiplicity	3.4 (3.4)
I/sigma	9.6 (2.0)
Rmerge	0.095 (0.622)
Refinement Statistics	
Resolution Range (Å)	26.19-2.72
R _{cryst}	0.24
R _{free}	0.29
R.m.s.d. bonds (Å)**	0.013
R.m.s.d. angles (°)	1.782

*values in parenthesis correspond to the highest resolution shell

5 **r.m.s.d root mean square deviation

Gremlin-1 structure was solved by molecular replacement using Phaser (McCoy *et al*, J Appl Cryst (2007), 40, 658-674) and a Gremlin-1 model available from proprietary Gremlin-1 / Fab complex coordinates. The resultant model of Gremlin-1 contained four
 10 copies of Gremlin 1 monomer organised as two dimers. Model corrections were made with Coot (Emsley *et al* Acta Crystallographica Section D: Biological Crystallography 66 (4), 486-501) and coordinates were refined using Refmac (Murshudov *et al* REFMAC5 for the refinement of macromolecular crystal structures. Acta Crystallographica Section D: Biological Crystallography. 2011;67(Pt 4):355-367). Final coordinates were validated with
 15 Molprobit (Chen et al. (2010) MolProbity: all-atom structure validation for macromolecular crystallography. Acta Crystallographica D66:12-21). A summary of model refinement statistics is shown in Table 2 above.

20 **Example 2 – BMP Binding residues on Gremlin-1**

As discussed above, Gremlin-1 belongs to the bone morphogenic protein (BMP) antagonist protein family within a sub-group known as the DAN family. Within the DAN family, Gremlin-1 shares greatest homology with Gremlin-2 (PRDC).

5

The 2.7 Å human Gremlin-1 structure resolved in Example 1 shares many features in common with the published mouse Gremlin-2 structure (Nolan *et al* (2013), Structure, 21, 1417–1429). The overall fold is very similar, with two copies of Gremlin-1 forming an antiparallel, non-covalent dimer, arranged in an arch. Each monomer adopts the

10 characteristic finger-wrist-finger arrangement with a cystine-knot motif towards the wrist end, opposite the fingers. Sequence identity between the proteins is 52 % rising to 67 % within the sequence visible in the two structures. The most highly conserved region lies in the extensive dimer interface where all the key contact residues are 100 % conserved.

15 Residues involved in BMP's 2, 4 & 7 binding to mouse Gremlin-2 (PRDC) and DAN (NBL1) have been identified using mutagenesis (Nolan *et al* (2013), Structure, 21, 1417–1429 and Nolan *et al* (2014) J. Biol. Chem. 290, 4759–4771). The predicted BMP binding epitope encompasses a hydrophobic patch spanning across both monomers on the convex surface of the dimer. Six residues were identified by mutagenesis; Trp72, Phe96,

20 Tyr98, Phe104, Tyr105 & Phe117 and are 100 % conserved in human Gremlin-1 (numbering based on the mouse Gremlin-2 sequence). The degree of homology extends to the positioning of the side chains which adopt an identical conformation in both proteins.

25 The amino acid numbering used in the Gremlin PDB file matches the numbering in the published mouse Gremlin-2 structure based on a structural alignment. This enables like for like comparison of amino acids when describing the structures. However, for clarity the key residues identified as playing a role in BMP binding are shown below with numbering based on the PDB file and UniProt file of SEQ ID NO: 1 in brackets:

30

Trp72(93), Phe96(117), Tyr98(119), Phe104(125), Tyr105(126) & Phe117(138).

In both mouse Gremlin-2 and human Gremlin-1 the hydrophobic BMP binding epitope is partially buried by an alpha helix formed by the N-terminal residues of each protein. A

35 model of BMP binding has been proposed whereby the N-terminus can flex, exposing the full BMP binding interface (Nolan *et al* (2013), Structure, 21, 1417–1429). In the present analysis, the N-terminal residues were removed from the human Gremlin-1 and mouse

Gremlin-2 structures before rendering a surface to reveal the similarity of the BMP binding faces on each protein.

The literature only describes mutagenesis of six residues that have an effect on BMP binding. It is possible that the actual BMP epitope covers a larger surface area, encompassing neighbouring amino acids. By highlighting all residues, within 6Å of those mutated, on the surface of Gremlin-1, a larger region of Gremlin-1 is revealed that could potentially be targeted by a therapeutic. This more extensive region encompasses the following amino acids of human Gremlin-1:

Asp92-Leu99
Arg116-His130
Ser137-Ser142
Cys176-Cys178

(Numbering based on SEQ ID NO: 1)

By combining published information with the crystal structure information of human Gremlin-1, regions of human Gremlin-1 that offer themselves as a potential route for therapeutic intervention blocking its interaction with BMP's have been identified.

Example 3 – Hek Id1 reporter gene assay

Background

The Hek Id1 reporter gene assay uses Clone 12 Hek293-Id1 reporter cells. This cell line was stably transfected with Id1 transcription factor. Id1 is a transcription factor in the BMP signalling pathway. Gremlin is known to bind BMPs prevent binding to their receptors reducing the luciferase signal from the reporter gene. Therefore, using this reporter assay, it is possible to screen anti-Gremlin antibodies and see if there are any that block the interaction of Gremlin with BMPs. A restoration of the luciferase signal is seen in these cells if there is a blocking of this interaction.

Method

Clone 12 cells were cultured in DMEM containing 10 % FCS, 1x L-Glutamine & 1x NEAA. Cells are also grown in the presence of Hygromycin B (200 µg/ml) to ensure cells do not lose Id1 gene expression. Cells were assayed in DMEM containing 0.5 % FCS, 1x L-Glutamine & 1x NEAA. Hygromycin B is not needed for the short time that the cells are in
5 the assay.

The cells were washed in PBS, lifted off using cell dissociation buffer, spun and counted before being seeded at 5x10⁴/well in 70 µl (Density of 7.14x10⁵/ml). Plates used were white, opaque Poly-D-Lysine coated 96-well sterile. Cells go in incubator for about 3-4
10 hours to settle down. BMP heterodimers were reconstituted to 200 µg/ml in 4 mM HCL. BMP was diluted to 10 µg/ml in assay media using a glass vial to give a new working stock.

In a polypropylene plate, Gremlin-1 was diluted 1:2 for an 8 point dose response curve
15 with a top final dose of 1 µg/ml.

An additional volume of 20 µl media was added per well and plates were incubated at 37 °C for 45 mins.

20 BMP prepared at 100x was added to all wells except wells containing cells only. All wells are made up to 60 µl with assay medium and incubated for a further 45 mins at 37 °C.

Post incubation, 30 µl of sample was transferred per well of assay plate and incubated for 20-24 hours before measuring luminescence signal.

25 Cell Steady Glo was thawed in advance at room temperature. Assay plates were cooled to room temperature for about 10-15 mins before adding the reagent. Luciferase signal was detected by addition of cell steady glo reagent (100 µl) for 20 minutes on shaker at room temperature and measuring luminescence using cell titre glo protocol on Synergy 2.

30 The maximum signal was generated from wells containing BMP and the minimum signal was generated from the wells containing cells only.

Results

35 Gremlin-1 full length and truncated forms were tested in the Hek-Id1 reporter gene assay to confirm the blocking activity against BMP4/7.

The percentage of inhibition from dose response assays was calculated based on the maximum and minimum signals in the assay and the data fitted using 4 parameter logistical fit. The IC₅₀ was calculated based on the inflexion point of the curve.

5

Table 3: Potency results for full length Gremlin-1 and truncated Gremlin-1 in the Hek-Id1 reporter gene assay.

Hek-Id1 Reporter gene assay	N	Geometric mean (nM)	95% CI (or range where N=<4)
Gremlin 1 Full length	2	1.6	1.3-1.9
Gremlin 1 truncated	2	1.7	1.1-2.5

15 Conclusion

Gremlin 1 was able to inhibit the BMP 4/7 signalling in the Hek-Id1 reporter gene assay.

20 **Example 4 – Production of anti-Gremlin-1 antibodies**

Anti-Gremlin-1 antibodies were derived by immunisation using purified gremlin-1 as described in Example 1, and by library panning. The library was generated in-house as a naïve human library with the V-regions amplified from blood donations.

25

Immunisation yielded 26 distinct antibodies binding Gremlin-1 from the first round of immunisation. These antibodies were scaled up and purified for testing in screening assays.

30 25 human and mouse cross-reactive antibodies from the library were panned using recombinant human Gremlin from R&D Systems. 10 antibodies were selected for scale up and purified as scFvs for testing in the screening assays.

35 **Example 5 – Screening of anti-Gremlin-1 antibodies**

Antibodies were screened using the Hek-Id1 reporter gene assay described in Example 3 and by measuring SMAD phosphorylation. SMAD1, 5 and 8 are phosphorylated upon BMP signalling. Inhibitors of Gremlin-1 therefore increase SMAD phosphorylation.

- 5 SMAD phosphorylation assays were conducted on A549 cells or on human lung fibroblasts. Phosphorylation levels were determined using MSD.

Results

- 10 In the Hek-Id1 reporter gene assay, there were no apparent hits with the immunisation derived antibodies (with a 10 fold excess of antibody tested against a BMP4/7 heterodimer). Results are shown in Figure 1.

- In contrast, a number of library derived antibodies were capable of restoring signal in the
15 Hek-Id1 reporter gene assay (50-fold excess of antibodies with a 50 % gremlin dose) (Figure 2). Of these, Ab2416 and Ab2417 contained high levels of endotoxin. Ab7326 maintained blocking ability at a 10-fold excess and 80 % inhibition Gremlin-1 concentration.

- 20 Additional results are presented in Figures 3A (human gremlin) and 3B (mouse Gremlin). These Figures show titrations of Ab7326 (labelled as PB376) up to 15 nM. Ab7326 was shown to restore signalling of BMP when blocked by either human (IC_{50} of 1.3 nM) or mouse (IC_{50} of 0.2 nM Gremlin). The antibody functions both as a human and mouse IgG1.

- 25 Sequences of the mouse and human full length IgG1 are presented below. In order to synthesise the mouse and human full length IgG1 proteins, the Ab7326 variable regions derived from the library were re-cloned into vectors comprising the appropriate antibody constant domains.

- 30 Because Ab7326 came from a naïve human library, where Abs are cloned as scFvs, in order to re-clone the 7326 variable regions as full length Abs or Fabs, it was necessary to PCR amplify the VH and VK using pools of primers/degenerate primers. The amplified PCR products were then digested and cloned simultaneously into mouse and human
35 vectors. As the VH and VK were amplified by pools of primers/degenerate primers, two variant forms of the products were obtained, differing by a single amino acid residue derived from subtly different primers annealing during the PCR process.

The two variant forms of heavy chain variable region differed by a single amino acid at position 6, and the two variant forms of the light chain variable region differed by a single amino acid at position 7, as shown below:

- 5
 - Heavy chain variable region variant 1 has glutamic acid (E) at position 6.
 - Heavy chain variable region variant 2 has glutamine (Q) at position 6.
 - Light chain variable region variant 1 has serine (S) at position 7.
 - Light chain variable region variant 2 has threonine (T) at position 7.

10 Mouse full length IgG1 – heavy chain variant 1 (SEQ ID NO: 14)

QVQLVESGAE VKKPGATVKI SCKVSGYTFT **DYYMH**WVQQA PGKGLEWMGL
VPEDGETIY AEKFQGRVTI TADTSTDYAY MELSSLRSED TAVYYCAT**DA**
RSGSYYPNH **FDY**WGQGLV TVSSAKTTP SVYPLAPGSA AQTNSMVTLG
 15 CLVKGYFPEP VTVWNSGSL SSGVHTFPAV LQSDLYTLSS SVTVPSSTWP
SETVTCNVAH PASSTKVDKK IVPRDCGCKP CICTVPEVSS VFIFPPKPKD
VTITLTPKV TCVVVDISKD DPEVQFSWFV DDVEVHTAQT QPREEQFNST
FRSVSELPIM HQDWLNGKEF KCRVNSAAFP APIEKTISK TGRPKAPQVY
TIPPPKEQMA KDKVSLTCMI TDFFPEDITV EWQWNGQPAE NYKNTQPIMD
 20 TDGSYFVYSK LNVQKSNWEA GNTFTCSVLH EGLHNHHTK SLSHSPGK

Mouse full length IgG1 – light chain variant 1 (SEQ ID NO: 15)

25 DIVMTQSPDS LAVSLGERAT INC**KSSQSVL YSSNNKN**YLA WYQQKPGQPP
 KLLIY**WASTR ESG**VPDRFSG SSGTDFTLT INSLQAEDVA VFYFC**QQYYDT**
PTFQGTRLE IKRTDAAPT VSIFFPSSEQL TSGGASVVCF LNNFYPKDIN
 VKWKIDGSER QNGVLNSWTD QDSKDSTYSM SSTLTTLTKDE YERHNSYTCE
 30 ATHKTSTSPI VKSFNRNEC

Mouse full length IgG1 – heavy chain variant 2 (SEQ ID NO: 28)

QVQLVQSGAE VKKPGATVKI SCKVSGYTFT **DYYMH**WVQQA PGKGLEWMGL
 35 **VPEDGETIY AEKFQ**GRVTI TADTSTDYAY MELSSLRSED TAVYYCAT**DA**
RSGSYYPNH **FDY**WGQGLV TVSSAKTTP SVYPLAPGSA AQTNSMVTLG
CLVKGYFPEP VTVWNSGSL SSGVHTFPAV LQSDLYTLSS SVTVPSSTWP
SETVTCNVAH PASSTKVDKK IVPRDCGCKP CICTVPEVSS VFIFPPKPKD
VTITLTPKV TCVVVDISKD DPEVQFSWFV DDVEVHTAQT QPREEQFNST
 40 FRSVSELPIM HQDWLNGKEF KCRVNSAAFP APIEKTISK TGRPKAPQVY
TIPPPKEQMA KDKVSLTCMI TDFFPEDITV EWQWNGQPAE NYKNTQPIMD
TDGSYFVYSK LNVQKSNWEA GNTFTCSVLH EGLHNHHTK SLSHSPGK

45 Mouse full length IgG1 – light chain variant 2 (SEQ ID NO: 29)

DIVMTQTPDS LAVSLGERAT INC**KSSQSVL YSSNNKN**YLA WYQQKPGQPP
 KLLIY**WASTR ESG**VPDRFSG SSGTDFTLT INSLQAEDVA VFYFC**QQYYDT**

PTFGQGTRLE IKRTDAAPT V SIFPPSSEQL TSGGASVVC F LNNFY PKDIN
 VKWKIDGSER QNGVLNSWTD QDSKDSTYSM SSTLT LTKDE YERHNSYTCE
 ATHKTSTSPI VKSFNRNEC

5

Human full length IgG1 – heavy chain variant 1 (SEQ ID NO: 30)

QVQLVESGAE VKKPGATVKI SCKVSGYTFT **DYYMH**WVQQA PGKGLEWMGL
VPEDGETIY AEKFQGRVTI TADTSTD TAY MELSSLRSED TAVYYCAT**DA**
 10 **RGSGSYYPNH FDY**WGQGLV TVSSASTKGP SVFPLAPSSK STSGGTAALG
 CLVKDYFPEP VTVSWNSGAL TSGVHTFPAV LQSSGLYSLS SVVTVPSSSL
 GTQTYICNVN HKPSNTKVDK KVEPKSCDKT HTCPCPCAPE LLGGPSVFLF
 PPKPKDTLMI SRTPEVTCVV VDVSHEDPEV KFNWYVDGVE VHNAKTKPRE
 EQYNSTYRVV SVLTVLHQDW LNGKEYKCKV SNKALPAPIE KTISKAKGQP
 15 **REPQ**VYTLPP SRDELTKNQV SLTCLVKGFY PSDIAVEWES NGQPENNYKT
 TPPVLDS DGS FFLYSKLTVD KSRWQQGNVF SCSVMHEALH NHYTQKSLSL
 SPGK

20 Human full length IgG1 – light chain variant 1 (SEQ ID NO: 31)

DIVMTQSPDS LAVSLGERAT INC**KSSQSVL YSSNNKNYLA** WYQQKPGQPP
 KLLIY**WASTR ESG**VPDRFSG SGSGTDFTLT INSLQAEDVA VFYFC**QQYYDT**
PTFGQGTRLE IKRTVAAPSV FIFPPSDEQL KSGTASVVCL LNNFY PREAK
 25 **VQ**WKVDNALQ SGNSQESVTE QDSKDSTYSL SSTLTLSKAD YEKHKVYACE
 VTHQGLSSPV TKSFNREGC

Human full length IgG1 – heavy chain variant 2 (SEQ ID NO: 16)

30 QVQLVQSGAE VKKPGATVKI SCKVSGYTFT **DYYMH**WVQQA PGKGLEWMGL
VPEDGETIY AEKFQGRVTI TADTSTD TAY MELSSLRSED TAVYYCAT**DA**
RGSGSYYPNH FDYWGQGLV TVSSASTKGP SVFPLAPSSK STSGGTAALG
 CLVKDYFPEP VTVSWNSGAL TSGVHTFPAV LQSSGLYSLS SVVTVPSSSL
 35 GTQTYICNVN HKPSNTKVDK KVEPKSCDKT HTCPCPCAPE LLGGPSVFLF
 PPKPKDTLMI SRTPEVTCVV VDVSHEDPEV KFNWYVDGVE VHNAKTKPRE
 EQYNSTYRVV SVLTVLHQDW LNGKEYKCKV SNKALPAPIE KTISKAKGQP
REPQVYTLPP SRDELTKNQV SLTCLVKGFY PSDIAVEWES NGQPENNYKT
 TPPVLDS DGS FFLYSKLTVD KSRWQQGNVF SCSVMHEALH NHYTQKSLSL
 40 SPGK

Human full length IgG1 – light chain variant 2 (SEQ ID NO: 17)

45 DIVMTQTPDS LAVSLGERAT INC**KSSQSVL YSSNNKNYLA** WYQQKPGQPP
 KLLIY**WASTR ESG**VPDRFSG SGSGTDFTLT INSLQAEDVA VFYFC**QQYYDT**
PTFGQGTRLE IKRTVAAPSV FIFPPSDEQL KSGTASVVCL LNNFY PREAK
VQWKVDNALQ SGNSQESVTE QDSKDSTYSL SSTLTLSKAD YEKHKVYACE
 VTHQGLSSPV TKSFNREGC

50

Antibody CDRs were determined using the Kabat method (highlighted in bold in the above sequences). Additional HCDR1 residues using the Chothia definition are in italics.

Constant region sequences are underlined.

- 5 Restoration of p-SMAD signalling with anti-Gremlin 1 antibodies is shown in Table 4 below.

Table 4: Restoration of p-SMAD signalling

	2417	2418	2419	2481	2482	2483	2484	7326	8427
BMP 2	109.1%	58.2%	32.6%	40.4%	35.3%	43.1%	104.0%	107.2%	51.3%
50ng/ml	+/- 2.8%	+/- 1.9%	+/- 1.4%	+/- 0.6%	+/- 0.8%	+/- 2.1%	+/- 2.7%	+/- 3.5%	+/- 1.4%
BMP 4	109.6%	71.3%	31.7%	60.1%	54.4%	72.5%	105.2%	110.0%	78.2%
25ng/ml	+/- 3.0%	+/- 3.1%	+/- 1.2%	+/- 2.2%	+/- 1.3%	+/- 2.1%	+/- 3.3%	+/- 3.8%	+/- 2.5%
BMP 7	111.5%	99.5%	53.8%	64.4%	52.3%	66.2%	105.2%	108.0%	72.6%
200	+/-	+/-	+/-	+/-	+/-	+/-	+/-	+/-	+/-
ng/ml	3.8%	3.2%	3.4%	1.3%	1.1%	1.2%	4.3%	3.2%	2.5%
BMP-2/7	119.3%	78.6%	50.8%	53.7%	47.6%	56.1%	120.4%	128.5%	62.8%
50ng/ml	+/- 2.6%	+/- 3.6%	+/- 2.7%	+/- 3.1%	+/- 1.5%	+/- 2.5%	+/- 4.4%	+/- 2.9%	+/- 2.5%
BMP4/7	113.7%	78.0%	61.4%	48.3%	41.7%	50.8%	112.4%	127.0%	63.3%
50ng/ml	+/- 3.1%	+/- 4.0%	+/- 4.0%	+/- 2.1%	+/- 1.7%	+/- 1.7%	+/- 2.5%	+/- 3.1%	+/- 2.1%

10

Results are shown as a percentage of SMAD phosphorylation by BMP alone (control BMP). Experiments were performed using lung fibroblasts from idiopathic pulmonary fibrosis patients. rhGremlin-1 and the anti-Gremlin-1 antibodies were preincubated for 45 minutes at room temperature. rhGremlin-1 and the anti-Gremlin-1 antibodies were then

15 added with BMP to the cells for 30 minutes.

Table 5 then shows further results in the SMAD phosphorylation assay, where displacement of BMP-2 or BMP4/7 from Gremlin 1-BMP complexes by anti-Gremlin-1 antibodies was investigated. Experiments were again performed using lung fibroblasts

20 from idiopathic pulmonary fibrosis patients. rhBMP-2 or rhBMP 4/7 were preincubated with rhGremlin-1 for 1 hour at room temperature. The BMP-2- or BMP4/7-Gremlin-1 complexes were incubated with different concentrations of the anti-Gremlin-1 antibodies

overnight at 4 °C. Antibody concentrations represent the final concentration on the plate.

Table 5: Displacement of BMP-2 or BMP4/7 from Gremlin 1-BMP complexes by anti-Gremlin-1 antibodies

5

		81.3 μg/ml	40.6 μg/ml	20.3 μg/ml	10.2 μg/ml	5.1 μg/ml	2.55 μg/ml	1.27 μg/ml	0.6 μg/ml
2484	BMP 2 50ng/ml	100.3% +/- 3.5%	98.8% +/- 2.7%	97.0% +/- 2.9%	93.5% +/- 2.6%	86.4% +/- 2.0%	79.9% +/- 1.9%	66.5% +/- 2.8%	54.8% +/- 0.3%
2484	BMP4/7 50ng/ml	136.4% +/- 4.2%	133.2% +/- 1.0%	121.4% +/- 1.4%	108.1% +/- 4.9%	86.6% +/- 4.4%	74.7% +/- 2.2%	65.8% +/- 0.6%	60.7% +/- 1.5%
7326	BMP 2 50ng/ml	103.7% +/- 1.1%	101.5% +/- 2.4%	99.4% +/- 3.8%	103.8% +/- 2.4%	100.3% +/- 2.2%	103.2% +/- 4.3%	102.8% +/- 2.8%	97.0% +/- 2.9%
7326	BMP4/7 50ng/ml	133.7% +/- 0.8%	132.3% +/- 1.8%	130.3% +/- 4.2%	125.6% +/- 10.0%	121.4% +/- 4.2%	120.9% +/- 3.3%	111.1% +/- 2.3%	102.0% +/- 4.5%

The results shown in Table 5 demonstrate that Ab7326 can displace already complexed BMP-2 or BMP4/7 from Gremlin 1-BMP complexes. Ab7326 can achieve this displacement at much lower concentrations than the comparison antibody 2484. This provides evidence that Ab7326 is an allosteric inhibitor, consistent with our finding that the binding site for Ab7326 is distal from the known BMP binding regions on gremlin-1. Thus Ab7326 is able to access the allosteric binding site even when BMP is complexed to gremlin-1, resulting in significantly improved inhibition of gremlin activity.

Example 6 – Obtaining the crystal structure of Gremlin-1 in complex with the 7326 Fab

The crystal structure of human Gremlin-1 in complex with Ab7326 Fab was solved at a resolution of 2.1 Å. Fab sequences are shown below:

20

Heavy chain: SEQ ID NO: 18

QVQLVESGAEVKKPGATVKISKVSGYTFDTYYMHVQQAAPGKGLEWMGLVDPEDGETIYA
EKFQGRVTITADTSTDYAMELSSLRSEDTAVYYCATDARGSGSYYPNHFYWGQGTLLVTV

SSASTKGPSVFPLAPSSKSTSGGTAAAGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQS
 SGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSC

5 Light chain: SEQ ID NO: 19

DIVMTQSPDLSAVSLGERATINCKSSQSVLYSSNNKNYLAWYQQKPGQPPKLLIYWASTRE
 SGVPDRFSGSGSTDFTLTINSLQAEDVAVYFCQQYYDTPTFGQGRLEIKRTVAAPSVFI
 FPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKSTYSLSSST
 10 LTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

The CCP4 software NCONT was then used to identify all contacts at 4 Å between
 Gremlin-1 and the Fab. The following residues were identified: Ile131, Lys147, Lys148,
 15 Phe149, Thr150, Thr151, Arg169, Lys174 and Gln175 (numbering based on the UniProt
 Sequence of SEQ ID NO: 1 (numbered as Ile110, Lys126, Lys127, Phe128, Thr129,
 Thr130, Arg148, Lys153 and Gln154 in the structure file which matches the numbering of
 mouse Gremlin-2).

20 Figure 4 shows structural models of the Gremlin-Fab complex, with the Fab epitope
 residues shown relative to the BMP binding regions.

Ab7326 is an inhibitory antibody which acts allosterically, i.e. it binds away from the BMP
 binding regions.

25

Example 7 – Affinity measurements for binding of anti-Gremlin-1 antibody Ab7326 to Gremlin-1.

Method

30 The affinity of anti-Gremlin mIgG for human Gremlin 1 was determined by biomolecular
 interaction analysis using surface plasmon resonance (SPR) technology on a Biacore
 T200 system, GE Healthcare Bio-Sciences AB. Anti-Gremlin mIgG was captured by an
 immobilised anti-mouse Fc surface and Gremlin 1 was titrated over the captured mIgG.
 The capture ligand (affinipure F(ab')₂ fragment of goat anti-mouse IgG, Fc fragment
 35 specific, 115-006-071, Jackson ImmunoResearch Inc.) was immobilised at 50 µg/ml in
 10mM NaAc, pH5.0 on flow cell 2 of a CM4 Sensor Chip via amine coupling chemistry,
 using 600s activation and deactivation injections, to a level of ~1600 response units (RU).
 HBS-EP+ buffer (0.01 M HEPES pH 7.4, 0.15 M NaCl, 3 mM EDTA, 0.05 % Surfactant
 P20) was used as the running buffer with a flow rate of 10 µl/min. A reference surface was
 40 prepared on flow cell 1 by activating and deactivating the surface as for flow cell 2 but

omitting the capture ligand.

The assay buffer was HBS-EP+ plus an extra 150mM NaCl to give a final NaCl concentration of 300mM plus 1% CMD40. A 60s injection of anti-Gremlin mIgG (at 5µg/ml in running buffer) was passed over flow cells 1 and 2 to give a capture level of

5 approximately 100 RU on the immobilised anti-mouse IgG, Fc surface. Recombinant human Gremlin 1 was titrated in running buffer from 5nM (using 2-fold dilutions) and injected over flow cells 1 and 2 at a flow rate of 30µl/min for 3min followed by a 5min dissociation phase. A buffer only control was also included. The surface was regenerated at a flow rate of 10µl/min by a 60s injection of 50mM HCl, a 30s injection of 5mM NaOH
10 and a 30s injection of 50mM HCl.

The kinetic data was determined using Biacore T200 evaluation software.

The affinity measurements were made at 25°C.

Results

Binding affinity, taken as the average K_D value for 5 determinations, was found to be

15 below 100 pM.

Example 8. Inhibition of gremlin-1 activity accelerates healing and bridging in an in vivo model of bone fracture repair.

20

8.1. Materials and Methods.

Rat fracture model and drug administration

Long bone segmental defect models have been widely used for the research of bone healing and regeneration (Sato et al; 2014). In the present study, a 3mm femoral defect
25 was created in 10-week old male rats and stabilised using an 8-hole PEEK plate (RIS. 602.105, RISystem, Switzerland). The plate was fixed to the bone with a forceps in the middle of the diaphysis, before the bone was drilled and fixed with screws. A 3mm fracture gap was created using a 0.44 mm Gigly saw. The defect size/consistency/fixation
30 was quality controlled by X-Ray imaging using Faxitron (MX-20-DC5, Faxitron Bioptics LLC, USA), this time point was defined as Day 0.

Weekly dosing was commenced on Day 1 for a period of 8 weeks as outlined in Table 1.

X-ray images were subsequently acquired during the in-life phase of the study at day 11, 25, 39 and 57 in order to assess the callus formation and the progress of healing.

35 Definiens image analysis was utilized to quantify the area of the defect that was devoid of bone tissue in the captured X-Ray images.

Table 6: Treatment Groups.

Group	Animal Number	Treatment	Dose	Dosing Regimen	Time
1	10	Vehicle	Vehicle: 1ml, s.c.	Once/week	Total of 57 days
2	10	Anti-Gremlin-1	30mg/kg, 1ml, s.c.	Once/week	

5 Micro-CT analysis of fracture healing

Femora (fractured side) were scanned at 17.2 μm resolution using micro-CT (SkyScan 1076). A region of approximately 15 mm of the callus with the fracture in the centre was acquired. The scans were reconstructed using the Skyscan NRECON software (1.7.10) and then the reconstructed slices were further segmented to exclude fixator pins at a 3mm defined region calculated from the mid-point of the femoral fracture defect.

Histomorphometric analysis of fracture callus in 3D was performed by SkyScan software (v. 1.13.1). The mid-point within the 3mm femoral fracture defect was determined and slices 1.5mm distal and proximal to the mid-point were segmented for each limb measured. Subsequently, the binarization of the reconstructed datasets and segmentation were performed following two defined thresholds, one to delineate the low mineralized callus (thus quantifying newly formed bone) and the other one to define mature bone. Further segmentation of these data was carried out on femora of animals classified as low responders based on satisfying the criteria of incomplete bridging of the femoral defect or high responders exhibiting bridging of the fracture site.

20

Histomorphometry analysis of fracture

Femora were fixed in 10 % neutral-buffered formalin for 24 h, dehydrated and embedded in methyl methacrylate (MMA) at low temperature. 50- μm -thick sections were stained with Toluidine Blue to quantify the bone elements of the healing gap defect.

Histomorphometric parameters were measured on the trabecular bone of the fracture defect site. Measurements were performed through image analysis.

Statistical analysis

The results were presented as mean values $\pm$ SD. Statistical analysis was performed using a two-tailed Mann Whitney U test with GraphPad Prism software unless otherwise stated.

30

8.2 Results.

Analysis of X-Ray images obtained during the in-life phase of the study indicated that the anti-gremlin-1 antibody accelerated fracture healing with the control and treated groups significantly diverging after 25 days ($P < 0.05$). This effect was apparent for the remainder of the study (Figure 5).

Micro-CT analysis of terminal samples revealed that treatment with anti-gremlin-1 antibody (30mg/kg/once weekly) led to an increase in newly formed bone within the fracture callus site ($P = 0.06$).

The incidence of fracture non-union in this model is approximately 60% with no intervention (Sato et al; 2014). To test whether gremlin-1 inhibition reduced the incidence of non-union development, the animals were classified as low responders (LR) and high responders (HR). Gremlin-1 inhibition resulted in a significant increase in the percentage of bone volume/tissue volume (BV/TV%) within LMB (low mineral bone; newly formed bone) ($P < 0.01$) and HMB (high mineral bone; mature bone) ($P < 0.01$) in the low responder group compared to controls, thus indicating progressive repair of the cohort likely to form non-union.

Additionally, there was a trend (non-significant) towards increased LMB and HMB BV/TV% in the high responder group in response to anti-gremlin-1 treatment (Figure 6A, representative images of LR and HR are shown in Figure 6B).

Two-dimensional histomorphometric analysis of bone parameters was performed on histological sections of the fracture site (Figure 7). Treatment with anti-gremlin-1 antibody significantly increased percentage of bone volume/tissue volume (BV/TV %) ($P < 0.05$) compared to control. Anti-gremlin-1 significantly increased trabecular number (Tb.N) ($P < 0.001$) and significantly decreased trabecular separation (Tb.Sp) ($P < 0.01$) indicating increased trabecular bone due to treatment with anti-gremlin-1.

Correlations were performed between two-dimensional histomorphometric analysis and three-dimensional μ CT analysis by comparing the LMB and HMB groups segmented in μ CT analysis and the two-dimensional histomorphometry analysis of fracture sections (Figure 8). Comparisons measured by Pearson's correlation revealed a positive and significant correlation between histomorphometry and μ CT analysis in the LMB ($P < 0.0001$) and HMB groups ($P < 0.0001$) thus validating the data from each data-set.

8.3. Conclusion.

Inhibition of gremlin-1 activity using a neutralising anti-gremlin-1 antibody resulted in accelerated fracture repair, with significant differences between control and treated groups evident after 25 days (3 doses of antibody). Additionally, terminal analysis of the fracture site indicated the enhanced formation of bone tissue in the low responder animals, which otherwise would likely form non-union. Therefore, inhibition of gremlin-1 activity is a promising therapy for the prevention or treatment of non-union fractures and may be of particular value for the treatment of fractures that are prone to non-union development, for example, tibia, distal radius, femoral neck and scaphoid.

SEQUENCE LISTING

SEQ ID NO: 1 (Human Gremlin-1; Uniprot ID: O60565)

MSRTAYTVGALLLLLTLLPAAEGKKKGSQGAI PPDKAQHNDSEQTQSPQQPGSRNRGRG
 QGRGTAMPGEEVLESSQEALHVTERKYLKRDWCKTQPLKQTIHEEGCNSRTIINRFCYGC
 NSFYIPRHIRKEEGSFQSCSFCKPKKFTTMMVTLNCPQLPPTKKKRVTRVKQCRCISIDL
 D

SEQ ID NO: 2 (Human truncated Gremlin-1 used in crystallography with N-terminal tag)

MGSSHHHHHSSGENLYFQGSAMPGEEVLESSQEALHVTERKYLKRDWCKTQPLKQTIHEE
 GCNSRTIINRFCYGCNSFYIPRHIRKEEGSFQSCSFCKPKKFTTMMVTLNCPQLPPTKK
 KRVTRVKQCRCISIDLD

SEQ ID NO: 3 (Ab7326 HCDR1 combined Kabat & Chothia)

GYTFTDYMH

SEQ ID NO: 4 (Ab7326 HCDR1 Kabat)

DYYMH

SEQ ID NO: 5 (Ab7326 HCDR2 Kabat)

LVDPEDGETIYAEKFQG

SEQ ID NO: 6 (Ab7326 HCDR3 Kabat)

DARGSGSYYPNHFDY

SEQ ID NO: 7 (Ab7326 LCDR1 Kabat)

KSSQSVLYSSNNKNYLA

5 SEQ ID NO: 8 (Ab7326 LCDR2 Kabat)

WASTRES

SEQ ID NO: 9 (Ab7326 LCDR3 Kabat)

QQYYDTPT

10

SEQ ID NO: 10 (Ab7326 Heavy chain variable region variant 1)QVQLVESGAIEVKKPGATVKISCKVSGYTFTDYIMHWVQQAPGKGLEWMGLVDPEDGETIYA
EKFQGRVTITADTSTDYIMELSSLRSEDTAVYYCATDARGSGSYYPNHFDYWGQGTLVTV
SS

15

SEQ ID NO: 11 (Ab7326 Light chain variable region variant 1)DIVMTQSPDSLAVSLGERATINCKSSQSVLYSSNNKNYLAWYQQKPGQPPKLLIYWASTRE
SGVPDRFSGSGSGTDFTLTINSLSQAEDVAVYFCQQYYDTPTFGQGRLEIK**20 SEQ ID NO: 12 (Ab7326 Heavy chain variable region variant 2)**QVQLVQSGAIEVKKPGATVKISCKVSGYTFTDYIMHWVQQAPGKGLEWMGLVDPEDGETIYA
EKFQGRVTITADTSTDYIMELSSLRSEDTAVYYCATDARGSGSYYPNHFDYWGQGTLVTV
SS**25 SEQ ID NO: 13 (Ab7326 Light chain variable region variant 2)**DIVMTQTPDSLAVSLGERATINCKSSQSVLYSSNNKNYLAWYQQKPGQPPKLLIYWASTRE
SGVPDRFSGSGSGTDFTLTINSLSQAEDVAVYFCQQYYDTPTFGQGRLEIK**SEQ ID NO: 14 (Mouse full length IgG1 heavy chain variant 1)**

30 QVQLVESGAIEVKKPGATVKISCKVSGYTFTDYIMHWVQQAPGKGLEWMGLVDPEDGETIYA
EKFQGRVTITADTSTDYIMELSSLRSEDTAVYYCATDARGSGSYYPNHFDYWGQGTLVTV
SSAKTTPPSVYPLAPGSAAQTNSMVTLGCLVKGYFPEPVTVTWNSGSLSSGVHTFPAVLQS
DLYTLSSSVTVPSSTWPSETVTCNVAHPASSTKVDKKIVPRDCGCKPCICTVPEVSSVFIF
PPKPKDVLITITLTPKVTCTVVDISKDDPEVQFSWFVDDVEVHTAQTQPREEQFNSTFRSVS

ELPIMHQDWLNGKEFKCRVNSAAFPAPIEKTISKTKGRPKAPQVYTI PPPKEQMAKDKVSL
TCMITDFFPEDITVEWQWNGQPAENYKNTQPIMDTDGSYFVYSKLVQKSNWEAGNTFTCS
VLHEGLHNHHTEKSLSHSPGK

5 **SEQ ID NO: 15 (Mouse full length IgG1 light chain variant 1)**

DIVMTQSPDSLAVSLGERATINCKSSQSVLYSSNNKNYLAWYQQKPGQPPKLLIYWASTRE
SGVPDRFSGSGSGTDFTLTINSLQAEDVAVYFCQQYYDTPTFGQGRLEIKRTDAAPT VSI
FPPSSEQLTSGGASVVCFLNNFYPKDINVWKIDGSERQNGVLNSWTDQDSKDSTYSMSST
LTLTKEDEYERHNSYTCEATHKTSTSPIVKSFNRNEC

10

SEQ ID NO: 16 (Human full length IgG1 heavy chain variant 2)

QVQLVQSGAEVKKPGATVKISCKVSGYTFTDYIMHWVQQAPGKGLEWMGLVDPEDGETIYA
EKFQGRVTITADTSTDYAMELSSLRSEDTAVYYCATDARGSGSYYPNHFDYWGQGT LVTV
SSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVT VSWNSGALTSGVHTFPAVLQS
15 SGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGG
PSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNS
TYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDEL T
KNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQG
NVFSCSVMEALHNHYTQKSLSLSPGK

20

SEQ ID NO: 17 (Human full length IgG1 light chain variant 2)

DIVMTQTPDSLAVSLGERATINCKSSQSVLYSSNNKNYLAWYQQKPGQPPKLLIYWASTRE
SGVPDRFSGSGSGTDFTLTINSLQAEDVAVYFCQQYYDTPTFGQGRLEIKRTVAAPSVFI
FPPSDEQLKSGTASVVCFLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSTYSLSST
25 LTLSKADYEEKHKVYACEVTHQGLSSPVTKSFNRGEC

SEQ ID NO: 18 (Fab heavy chain variant 1)

QVQLVESGAEVKKPGATVKISCKVSGYTFTDYIMHWVQQAPGKGLEWMGLVDPEDGETIYA
EKFQGRVTITADTSTDYAMELSSLRSEDTAVYYCATDARGSGSYYPNHFDYWGQGT LVTV
30 SSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVT VSWNSGALTSGVHTFPAVLQS
SGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSC

SEQ ID NO: 19 (Fab light chain variant 1)

DIVMTQSPDSLAVSLGERATINCKSSQSVLYSSNNKNYLAWYQQKPGQPPKLLIYWASTRE

SGVPDRFSGSGSGTDFTLTINSLQAEDVAVYFCQQYYDTPTFGQGRLEIKRTVAAPSVFI
 FPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSTYLSST
 LTLSKADYEEKHKVYACEVTHQGLSSPVTKSFNRGEC

5 **SEQ ID NO: 20 (Human truncated Gremlin-1 used in
 crystallography without N-terminal tag)**

AMPGEEVLESSQEALHVTERKYLKRDWCKTQPLKQTIHEEGCNSRTIINRFCYGCNSFYI
 PRHIRKEEGSFQSCSFCKPKKFTTMMVTLNCPQLPPTKKKRVTRVKQCRCSIDL

10 **SEQ ID NO: 21 (Mature Gremlin-1 sequence of SEQ ID NO: 1
 lacking the signal peptide of amino acids 1-21)**

KKKGSQGAIPPPDKAQHNDSEQTQSPQQPSRNRGRGQGRGTAMPGEEVLESSQEALHVTE
 RKYLKRDWCKTQPLKQTIHEEGCNSRTIINRFCYGCNSFYIPRHIRKEEGSFQSCSFCKP
 KKFTTMMVTLNCPQLPPTKKKRVTRVKQCRCSIDL

15

SEQ ID NO: 22 (Human IgG4P heavy chain variant 1)

QVQLVESGAIEVKKPGATVKISCKVSGYFTFDYYMHVWQQAPGKGLEWMGLVDPEDGETIYA
 EKFGQGRVTITADTSTDTAYMELSSLRSEDTAVYYCATDARGSGSYYPNHFDYWGQGTLLVTV
 SSASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQS
 20 SGLYSLSVVTVPSSSLGKTYTCNVDPKPSNTKVDKRVESKYGPPCPPCPAPEFLGGPSV
 FLFPPKPKDTLMISRTPEVTCVVVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYR
 VVSVLTIVLHQDWLNGKEYKCKVSNKGLPSSIEKTIKAKGQPREPQVYTLPPSQEEMTKNQ
 VSLTCLVKGFIYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVF
 SCSVMHEALHNHYTQKSLSLGLK

25

SEQ ID NO: 23 (Human IgG4P light chain variant 1)

DIVMTQSPDSLAVSLGERATINCKSSQSVLYSSNNKNYLAWYQQKPGQPPKLLIYWASTRE
 SGVPDRFSGSGSGTDFTLTINSLQAEDVAVYFCQQYYDTPTFGQGRLEIKRTVAAPSVFI
 FPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSTYLSST
 30 LTLSKADYEEKHKVYACEVTHQGLSSPVTKSFNRGEC

SEQ ID NO: 24 (Human IgG1 heavy chain DNA variant 1)

caagtgaactggtggaatccggggccgaagtgaaaaagcccggagccactgtgaagatct
 cttgcaaagtgtccggctacaccttcaccgactattacatgcactgggtccagcaggcacc

tgggaagggccttgagtggatgggtctggtcgatcccaggagcggcgaaactatctacgcc
gagaagtccagggctcgcgtcaccatcaccgccgacacttccaccgacaccgcgtacatgg
agctgtccagcttgaggtccgaggacacagccgtgtactactgcgccacggatgctcgggg
aagcggcagctactacccgaaccacttcgactactggggacagggcactctcgtgactgtc
5 tcgagcgtcttctacaaagggccccctccgtgttcccgcctcgcctccatcatcgaagtctacca
gcggaggcactgcggctctcggttgcctcgtgaaggactacttcccggagccggtgaccgt
gtcgtggaacagcggagccctgaccagcggggtgcacacctttccggccgtcttgacgtca
agcggcctttactccctgtcatcagtggtgactgtcccgtccagctcattgggaacccaaa
cctacatctgcaatgtgaatcacaaacctagcaacaccaaggttgacaagaaagtcgagcc
10 caaatcgtgtgacaagactcacacttgtccgccgtgcccggcaccggaactgctgggaggt
cccagcgtctttctgttccctccaaagccgaaagacacgctgatgatctcccgcaccccg
aggtcacttgctgtggtcgtggacgtgtcacatgaggacccagaggtgaagttcaattggta
cgtggatggcgtcgaagtccacaatgccaaaactaagcccagagaagaacagtacaattcg
acctaccgcgtcgtgtccgtgctcacggtgttgcatcaggattggctgaacgggaaggaat
15 acaagtgcaaagtgtccaacaaggcgtgccggcaccgatcgagaaaactatctccaaagc
gaagggacagcctagggaaacctcaagtctacacgctgccaccatcacgggatgaactgact
aagaatcaagtctcactgacttgtctggtgaaggggtttaccctagcgacattgccgtgg
agtgggaatccaacggccagccagagaacaactacaagactacccctccagtgtcgcactc
ggatggatcgttcttctttactcgaagctcacggtggataagtcccgggtggcagcagggga
20 aacgtgttctcctgctcgggtgatgcataagccctccataaccactatacccaaaagtcgc
tgtccctgtcgcgggaaag

SEQ ID NO: 25 (Human IgG1 light chain DNA variant 1)

gacattgtgatgaccagtcccccgattcgccttgccgtgtccctgggagaacgggccacca
25 ttaactgcaagagctcacagtcctgtattcatcgaacaacaagaattacctcgcatg
gtatcagcagaagcctggacagcctcccaagctgctcatctactgggctagcaccgcgaa
tccggggtgccggatagattctccggatcgggttcgggcactgacttcactctgactatca
actcactgcaagccgaggatgtcgcgggtgtacttctgtcagcagtactacgacacccccgac
ctttggacaaggcaccagactggagattaagcgtacgggtggccgctccctccgtgttcac
30 ttcccacccctccgacgagcagctgaagtcgggcaccgcctccgtcgtgtgctgctgaaca
acttctacccccgcgaggccaaggtgcagtggagggtggacaacgcctgcagtcgggcaa
ctcccaggaatccgtcaccgagcaggactccaaggacagcacctactccctgtcctccacc
ctgacctgtccaaggccgactacgagaagcacaaggtgtacgcctgcgaagtgaaccacc
agggcctgtccagccccgtgaccaagtccttcaaccggggcgagtgc

SEQ ID NO: 26 (Human IgG4P heavy chain DNA variant 1)

caagtgcaactggtggaatccggggccgaagtgaaaaagcccgagccactgtgaagatct
cttgcaaagtgtccggctacaccttcaccgactattacatgcactgggtccagcaggcacc
5 tgggaagggccttgagtggatgggtctggtcgatcccgaggacggcgaaactatctacgcc
gagaagttccagggtcgcgtcaccatcaccgccgacacttccaccgacaccgcgtacatgg
agctgtccagcttgaggtccgaggacacagccgtgtactactgcgccacggatgctcgggg
aagcggcagctactacccgaaccacttcgactactggggacagggcactctcgtgactgtc
tcgagcgttctacaaagggccctccgtgttcctcttgcccccttgctcccggtccacct
10 ccgagtctaccgccgtctggtgctgctcaaggactacttccccgagcccgtagcagt
gtcctggaactctggcgccctgacctccggcgtgcacaccttcctgccgtgctgcagtcc
tccggcctgtactccctgtcctccgtcgtgaccgtgccctcctccagcctgggcaccaaga
cctacacctgtaacgtggaccacaagccctccaacaccaaggtggacaagcgggtggaatc
taagtacggccctccctgccccccctgccctgccccctgaatttctgggcggaccttccgtg
15 ttctgttccccccaaagcccaaggacacctgatgatctcccgacccccgaagtgcct
gcgtgggtgggtggacgtgtccaggaagatcccgagggtccagttcaattggtagctggacgg
cgtggaagtgcacaatgccaagaccaagcccagagaggaacagttcaactccacctaccgg
gtggtgtccgtgctgaccgtgctgcaccaggactgggtgaacggcaaagagtacaagtgc
aggtgtccaacaaggcctgccctccagcatcgaaaagaccatctccaaggccaagggcc
20 gccccgcgagccccaggtgtacacctgccccctagccaggaagagatgaccaagaaccag
gtgtccctgacctgtctggtcaagggttctacctcccgacattgccgtggaatgggagt
ccaacggccagcccgagaacaactacaagaccacccccctgtgctggacagcgacggctc
cttcttccctgtactctcggtgaccgtggacaagtcccggtggcaggaaggcaacgtcttc
tctgtcctcgatgacagaggccctgcacaaccactacaccagaagtccctgtccctga
25 gcctgggcaag

SEQ ID NO: 27 (Human IgG4P light chain DNA variant 1)

gacattgtgatgaccagtcccccgattcgcttgccgtgtccctgggagaacgggccacca
ttaactgcaagagctcacagtcctgtattcatcgaacaacaagaattacctcgcatg
30 gtatcagcagaagcctggacagcctcccaagctgctcatctactgggctagcaccgcgaa
tccggggtgccgtagattctccggatcgggttcgggcactgacttcactctgactatca
actcactgcaagccgaggatgtcgcgggtgtacttctgtcagcagtactacgacacccccgac
ctttggacaaggcaccagactggagattaagcgtacgggtggccgctccctccgtgttcac
tccccacctccgacgagcagctgaagtccggcaccgcctccgtcgtgtgcctgctgaaca

acttctacccccgcgaggccaaggtgcagtggaaggtggacaacgccctgcagtcggcaa
ctcccaggaatccgtcaccgagcaggactccaaggacagcacctactccctgtcctccacc
ctgaccctgtccaaggccgactacgagaagcacaaggtgtacgcctgcgaagtgaccacc
agggcctgtccagccccgtgaccaagtccttcaaccggggcgagtgc

5

SEQ ID NO: 28 (Mouse full length IgG1 heavy chain variant 2)

QVQLVQSGAEVKKPGATVKISCKVSGYTFTDYMHVWQQAPGKGLEWMGLVDPEDGETIYA
EKFQGRVTITADTSTDYAMELSSLRSEDYAVYYCATDARGSGSYYPNHFDYWGQGTLLVTV
SSAKTTPPSVYPLAPGSAAQTNSMVTLGCLVKGYFPEPVTVTWNSGSLSSGVHTFPAVLQS
10 DLYTLSSSVTVPSSTWVPSSTVTCNVVHPASSTKVDKKIVPRDCGCKPCICTVPEVSSVFIF
PPKPKDVLTTITLTPKVTCTVVDISKDDPEVQFSWFVDDVEVHTAQTQPREEQFNSTFRSVS
ELPIMHQDWLNGKEFKCRVNSAAFPAPIEKTISKTKGRPKAPQVYTIPPPKEQMAKDKVSL
TCMITDFFPEDITVEWQWNGQPAENYKNTQPIMDTDGSYFVYSKLVQKSNWEAGNTFTCS
VLHEGLHNHHTKSLSHSPGK

15

SEQ ID NO: 29 (Mouse full length IgG1 light chain variant 2)

DIVMTQTPDSLAVSLGERATINCKSSQSVLYSSNNKNYLAWYQQKPGQPPKLLIYWASTRE
SGVPDRFSGSGSGTDFTLTINSLSQAEDVAVYFCQQYYDTPTFGQGRLEIKRTDAAPTQSI
FPPSSEQLTSGGASVVCFLNNFYPKDINVKWKIDGSEKQNGVLNSWTDQDSKSTYSMSST
20 LTTLTKDEYERHNSYTCEATHKTSTSPIVKSFNREK

SEQ ID NO: 30 (Human full length IgG1 heavy chain variant 1)

QVQLVESGAEVKKPGATVKISCKVSGYTFTDYMHVWQQAPGKGLEWMGLVDPEDGETIYA
EKFQGRVTITADTSTDYAMELSSLRSEDYAVYYCATDARGSGSYYPNHFDYWGQGTLLVTV
25 SSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQS
SGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGG
PSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNS
TYRVSIVLTIVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTK
KNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQG
30 NVFSCSVMEALHNHYTQKSLSLSPGK

SEQ ID NO: 31 (Human full length IgG1 light chain variant 1)

DIVMTQSPDSLAVSLGERATINCKSSQSVLYSSNNKNYLAWYQQKPGQPPKLLIYWASTRE
SGVPDRFSGSGSGTDFTLTINSLSQAEDVAVYFCQQYYDTPTFGQGRLEIKRTVAAPSVFI

FPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSTYLSLST
LTLSKADYEEKHKVYACEVTHQGLSSPVTKSFNRGEC

SEQ ID NO: 32 (Fab heavy chain variant 2)

5 QVQLVQSGAEVKKPGATVKISCKVSGYTFTDYMHVWVQAPGKGLEWMGLVDPEDGETIYA
EKFQGRVTITADTSTDYAMELSSLRSEDYAVYYCATDARGSGSYYPNHFDYWGQGTILVTV
SSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQS
SGLYSLSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSC

10 **SEQ ID NO: 33 (Fab light chain variant 2)**

DIVMTQTPDSLAVSLGERATINCKSSQSVLYSSNNKNYLAWYQQKPGQPPKLLIYWASTRE
SGVPDRFSGSGSGTDFTLTINSQAEDVAVYFCQQYYDTPTFGQGRLEIKRTVAAPSVFI
FPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSTYLSLST
LTLSKADYEEKHKVYACEVTHQGLSSPVTKSFNRGEC

15

SEQ ID NO: 34 (Human IgG4P heavy chain variant 2)

QVQLVQSGAEVKKPGATVKISCKVSGYTFTDYMHVWVQAPGKGLEWMGLVDPEDGETIYA
EKFQGRVTITADTSTDYAMELSSLRSEDYAVYYCATDARGSGSYYPNHFDYWGQGTILVTV
SSASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQS
20 SGLYSLSVVTVPSSSLGTQTYTCNVDHKPSNTKVDKRVESKYGPPCPPCPAPEFLGGPSV
FLFPPKPKDTLMISRTPEVTCVVVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYR
VVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTIKAKGQPREPQVYTLPPSQEEMTKNQ
VSLTCLVKGFIYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVF
SCSVMHEALHNHYTQKSLSLGLGK

25

SEQ ID NO: 35 (Human IgG4P light chain variant 2)

DIVMTQTPDSLAVSLGERATINCKSSQSVLYSSNNKNYLAWYQQKPGQPPKLLIYWASTRE
SGVPDRFSGSGSGTDFTLTINSQAEDVAVYFCQQYYDTPTFGQGRLEIKRTVAAPSVFI
FPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSTYLSLST
30 LTLSKADYEEKHKVYACEVTHQGLSSPVTKSFNRGEC

REFERENCES

Bostrom, M. P. & Seigerman, D. A. (2005), HSS journal: The Musculoskeletal Journal of Hospital for Special Surgery 1, 9-18. The clinical use of allografts, demineralized bone matrices, synthetic bone graft substitutes and osteoinductive growth factors: a survey study.

5

Buza, J. A., 3rd & Einhorn, T. (2016), Clinical cases in mineral and bone metabolism: The Official Journal of the Italian Society of Osteoporosis, Mineral Metabolism, and Skeletal Diseases 13, 101-105. Bone healing in 2016.

- 10 Canalis, E., Parker, K. & Zanotti, S. (2012), J.Cell Physiol 227, 269-277. Gremlin1 is required for skeletal development and postnatal skeletal homeostasis.

- Cho, T. J., Gerstenfeld, L. C. & Einhorn, T. A. (2002), Journal of Bone and Mineral Research: The Official Journal of the American Society for Bone and Mineral Research
15 17, 513-520. Differential temporal expression of members of the transforming growth factor beta superfamily during murine fracture healing.

Einhorn, T. A. & Gerstenfeld, L. C. (2015), Nat. Rev. Rheumatol. 11, 45-54. Fracture healing: mechanisms and interventions.

20

Ferguson, C., Alpern, E., Miclau, T. & Helms, J. A. (1999), Mechanisms of development 87, 57-66. Does adult fracture repair recapitulate embryonic skeletal formation?

- 25 Gazzo, E. et al. (2005), Endocrinology 146, 655-665. Skeletal overexpression of gremlin impairs bone formation and causes osteopenia.

Gazzo, E. et al. (2007), J.Biol.Chem. 282, 31549-31557. Conditional deletion of gremlin causes a transient increase in bone formation and bone mass.

- 30 Goulet, J. A., Senunas, L. E., DeSilva, G. L. & Greenfield, M. L. (1997), Clinical Orthopaedics and Related Research, 76-81. Autogenous iliac crest bone graft. Complications and functional assessment.

- 35 Hsu, D. R., Economides, A. N., Wang, X., Eimon, P. M. & Harland, R. M. (1998), Mol. Cell 1, 673-683. The Xenopus dorsalizing factor Gremlin identifies a novel family of secreted proteins that antagonize BMP activities.

Sato, K., Watanabe, Y., Harada, N., Abe, S., Matsushita, T., Yamanaka, K., Kaneko, T., & Sakai, Y. (2014) Tissue Eng Part C. Methods 20, 1037-1041.

- Schmid, G. J., Kobayashi, C., Sandell, L. J. & Ornitz, D. M. (2009), Developmental
5 Dynamics: An Official Publication of the American Association of Anatomists 238, 766-
774. Fibroblast growth factor expression during skeletal fracture healing in mice.

Yu, Y. Y. et al. (2010), Bone 46, 841-851. Immunolocalization of BMPs, BMP antagonists, receptors, and effectors during fracture repair.