

JATROPHA CURCAS CURCIN GENES, TISSUE-SPECIFIC PROMOTERS AND
GENERATION OF CURCIN-DEFICIENT TRANSGENIC JATROPHA PLANTS

CROSS-REFERENCE TO RELATED APPLICATION

[0001] The present application claims priority to U.S. provisional patent application Serial No. 61/184,416 filed 5 June 2009, which is incorporated herein by reference.

BACKGROUND OF THE INVENTION

[0002] The present invention relates to the isolation of *Jatropha curcas* curcin genes and tissue-specific promoters and to the production of curcin-deficient *Jatropha* plants. More specifically, the present invention relates to the isolation of *Jatropha curcas* *Curcin 1*, *Curcin 2* and *Curcin 2A*. The present invention further relates to of the *Curcin 1*, *Curcin 2A* and *Curcin 2* genes and more particularly to tissue specific promoters of the *Curcin 1* and *Curcin 2A* genes. The present invention further relates to production of curcin-deficient transgenic *jatropha* plants by using RNAi technology to suppress curcin gene expression.

[0003] The publications and other materials used herein to illuminate the background of the invention or provide additional details respecting the practice, are incorporated by reference, and for convenience are respectively grouped in the Bibliography.

[0004] Tissue specific promoters are strongly desired for plant biotechnology to express gene-of-interest in the particular plant tissue at right time (Mansoor et al., 2006). Plant seed endosperm accumulates storage materials, such as starch, proteins and lipids (Berger et al., 2006; Hannah and James, 2008). The endosperm-specific promoters have been used to drive the expression of genes that are involved in biosynthesis pathways of these storage materials (Roesler et al., 1997; Plant et al., 1994; Kuwano et al., 2009). The insecticidal toxins, such as *Bacillus thuringiensis* (Bt) δ -endotoxins, have been used to control insects (Christou et al., 2006; Roh et al., 2007). Although Bt toxin has high specificity as an insecticidal toxin and safety for the environment, it is still desirable to have Bt toxin specifically expressed in leaf tissues rather than in seeds and fruits (Datta et al., 1998). Therefore, there is an increasing demand of specific promoter to control the expression of gene of interest in specific tissues at particular developmental stages. In addition, the multi-gene transformation system was used for delivery several genes simultaneously constructed in one expression vector (Lin et al., 2003; Chen et al., 2006; Wakasa et al., 2006). Each of these multiple genes needs a different

promoter to drive them to avoid the gene silencing. However, a lack of suitable promoters is a critical limiting factor for such research (Qu et al., 2008).

[0005] *Jatropha* (*Jatropha curcas*) is a small tropical, woody plant belonging to the Euphorbiaceae family. It is a dicotyledonous plant and its seeds contain as much as 40% oil (Bringi, 1987). The oil from *jatropha* could be an efficient substitute for fossil fuel (Augustus et al., 2002; Azam et al., 2005; Forson et al., 2004; Pramanik, 2003). However, *Jatropha* suffers from several shortcomings that may limit its wide adoption. The productivity of the plant is constrained by the unfavourable male to female flower ratio and its oil content has not been optimized by breeding. This plant is also sensitive to biotic stresses such as viral (Narayanna et al., 2007), fungal and bacterium pathogens and abiotic stresses, especially cold and drought (<http://www.jatropha.org>). The presence of several toxic components (e.g. the protein toxin, curcin, and the cancer-causing agent phorbol esters) in seeds and leaves of the plant possess health hazards for farmers and bioprocess workers in the *Jatropha* industry.

[0006] Curcin is a toxin protein identified in *jatropha* seeds (Stirpe et al. 1976). The presence of curcin as well as other toxins in *jatropha* seeds prevents the use of *jatropha* seed meals as animal feed (Makkar et al., 1997). Curcin belongs to the type I of ribosome inactivating proteins (RIPs) that has RNA *N*-glycosidase activity and can irreversibly inactivate ribosomes (Barbieri, 1993, Lin et al., 2003a). Currently, five curcin proteins have been deposited to GenBank and their accession numbers are AAL58089 (Lin et al., 2003a), AAL86778 (Lin et al., 2003b), ABZ04128, AAR08395 and ABW17545. These curcin proteins can be divided into two types based on the length of their amino acid residues. The precursors of type 1 curcin proteins consist of 293-aa residues while the precursors of type 2 curcin proteins contain 309-aa residues. Lin et al. (2003a) identified a type 1 curcin protein from *jatropha* seeds that encodes a 32-kDa curcin precursor with a 42-amino acid signal peptide. Wei et al. (2005) cloned a type 2 curcin gene. The curcin gene, designated as *Curcin* 2, was found to be induced by stress. Members of curcin proteins within a type display 93% to 98% identity in amino acids while members between two different types show 87%.

[0007] An important strategy to improve agronomic and quality traits of *Jatropha curcas* is by genetic modification. Transgenic *Jatropha* plants can be generated expressing homologous or heterologous gene sequences. In many instances, over-expression or silencing by RNA interference (RNAi) of one or more homologous genes of defined function is desired. Gene sequences of *Jatropha* can be obtained from cDNA and genomic libraries and functions

of genes can be tentatively assigned by sequence homology with other plant genes of known function. Tissue-specific promoters are often utilized to express homologous or heterologous gene sequences in the desired tissues.

[0008] RNAi is a post-transcriptional, targeted gene-silencing technique that uses double-stranded RNA (dsRNA) to incite degradation of messenger RNA (mRNA) containing the same sequence as the dsRNA (Constans, 2002). Small interfering RNAs, or siRNAs are produced in at least two steps: an endogenous ribonuclease cleaves longer dsRNA into shorter, 21-23 nucleotide-long RNAs. The siRNA segments then mediate the degradation of the target mRNA (Zamore, 2001). RNAi has been used for gene function determination in a manner similar to antisense oligonucleotides (Constans, 2002). Expression vectors that continually express siRNAs in transiently and stably transfected have been engineered to express small hairpin RNAs (shRNAs), which get processed *in vivo* into siRNAs-like molecules capable of carrying out gene-specific silencing (Brummelkamp et al., 2002; Paddison et al., 2002). Post-transcriptional gene silencing by double-stranded RNA is discussed in further detail by Hutvagner and Zamore, 2002; Vaucheret et al., 2001; Hammond et al., 2001; Maine, 2000; Fire et al., 1998; and Timmons and Fire, 1998. Methods for using RNAi to inhibit the expression of a gene are known to those of skill in the art. See, e.g., U.S. Patent Nos. 5,034,323; 6,326,527; 6,452,067; 6,573,099; 6,753,139; and 6,777,588. See also International Patent Publications WO 97/01952, WO 98/36083, WO 98/53083, WO 99/32619 and WO 01/75164; and U.S. Patent Publications 2003/0175965, 2003/0175783, 2003/0180945, 2004/0214330, 2005/0244858, 2005/0277610, 2007/0265220 and 2010/0058498.

[0009] Thus, it is desired to isolate tissue-specific promoters for use in controlling the expression of a gene of interest in specific tissues at particular developmental stages in *Jatropha curcas* other *Jatropha* species as well as other plant species, such as for genetic engineering of such plants. It is also desired to isolated promoters that can be used in genetic engineering of *Jatropha curcas*, other *Jatropha* species as well as other plant species. It is also desired to produce curcin-deficient *jatropha* plants that are non-toxic to humans and animals.

SUMMARY OF THE INVENTION

[0010] The present invention relates to the isolation of *Jatropha curcas* curcin genes and tissue-specific promoters and to the production of curcin-deficient *Jatropha* plants. More

specifically, the present invention relates to the isolation of *Jatropha curcas* *Curcin 1*, *Curcin 2* and *Curcin 2A*. The present invention further relates to of the *Curcin 1*, *Curcin 2A* and *Curcin 2* genes and more particularly to tissue specific promoters of the *Curcin 1* and *Curcin 2A* genes. The present invention further relates to production of curcin-deficient transgenic *jatropha* plants by using RNAi technology to suppress curcin gene expression.

[0011] Thus in a first aspect, the present invention provides the sequences of three *Jatropha curcas* curcin genes. In one embodiment, the curin gene is *Curcin 1*. The nucleotide sequence of *Curcin 1* is set forth in SEQ ID NO:1. The coding sequence for Curcin 1 comprises nucleotides 474 - 1355 of SEQ ID NO:1. The protein sequence for Curcin 1 is set forth in SEQ ID NO:2.

[0012] In a second embodiment, the curin gene is *Curcin 2*. The nucleotide sequence of *Curcin 2* is set forth in SEQ ID NO:3. The coding sequence for Curcin 1 comprises nucleotides 439 - 1368 of SEQ ID NO:3. The protein sequence for Curcin 2 is set forth in SEQ ID NO:4.

[0013] In a third embodiment, the curin gene is *Curcin 2A*. The nucleotide sequence of *Curcin 2A* is set forth in SEQ ID NO:5. The coding sequence for Curcin 2A comprises nucleotides 475 - 1404 of SEQ ID NO:5. The protein sequence for Curcin 2A is set forth in SEQ ID NO:6.

[0014] In a second aspect, the present invention provides the tissue-specific promoters of two *Jatropha curcas* curcin genes and a promoter of a third *Jatropha curcas* curcin gene. In one embodiment, the promoter is derived from the *Curcin 1* gene. The nucleotide sequence of *Curcin 1* promoter is set forth in SEQ ID NO:7. The *Curcin 1* promoter is an endosperm-specific promoter. Fragments of this sequence are also active as tissue-specific promoters, i.e., as endosperm-specific promoters. Such fragments include the following: (a) nucleotides 1 to 2888 of SEQ ID NO:7, (b) nucleotides 1142 to 3181 of SEQ ID NO:7, (c) SEQ ID NO:7 with nucleotides 2944 to 3170 deleted, (d) nucleotides 1142 to 3181 of SEQ ID NO:7 with nucleotides 2944 to 3170 deleted and (e) nucleotides 2688 to 3181 of SEQ ID NO:7 with nucleotides 2944 to 3170 deleted.

[0015] In a second embodiment, the promoter is derived from the *Curcin 2A* gene. The nucleotide sequence of *Curcin 2A* promoter is set forth in SEQ ID NO:8. The *Curcin 2A* promoter is a leaf-specific promoter. Fragments of this sequence are also active as tissue specific promoters. However, the specificity changes from leaf-specific to non-tissue specific when some fragments are deleted. Such fragments that are active as non-tissue specific

promoters include the following: (a) nucleotides 912 to 2087 of SEQ ID NO:8, (b) nucleotides 1 to 2087 of SEQ ID NO:8 with nucleotides 1853 to 2076 deleted, (c) nucleotides 912 to 2087 of SEQ ID NO:8 with nucleotides 1853 to 2076 deleted and (d) nucleotides 1751 to 2087 of SEQ ID NO:8 with nucleotides 1853 to 2076 deleted.

[0016] In a third embodiment, the promoter is derived from the *Curcin 2* gene. The nucleotide sequence of *Curcin 2* promoter is set forth in SEQ ID NO:9. As described herein, this promoter is not expressed in endosperm or leaf tissues, but may be activated and expressed upon activation by jasmonic acid (JA) upon attack by insects and/or upon activation by ethylene during defense reaction or leaf senescence.

[0017] In a third aspect, the present invention provides transgenic *Jatropha* plants that are curcin-deficient and to the production of curcin-deficient transgenic *Jatropha* plants. In one embodiment, a transgenic plant that is curcin-deficient comprises a nucleic acid stably integrated in its genome, wherein the nucleic acid encodes a double stranded RNA (dsRNA) targeted to a curcin gene. In another embodiment, a transgenic plant that is curcin-deficient comprises a nucleic acid stably integrated in its genome, wherein the nucleic acid encodes short interfering RNA (siRNA) targeted to a curcin gene. In an additional embodiment, a curcin-deficient transgenic plant comprises a nucleic acid stably integrated in its genome, wherein the nucleic acid encodes a short hairpin RNA (shRNA) targeted to a curcin gene. In one embodiment, the curcin gene is *Curcin 1*. In another embodiment, the curcin gene is *Curcin 2*. In an additional embodiment, the curcin gene is *Curcin 2A*. In one embodiment, the nucleic acid includes a portion of the *Curcin 1* gene. In another embodiment, the nucleic acid includes a portion of the *Curcin 2* gene. In an additional embodiment, the nucleic acid includes a portion of the *Curcin 2A* gene. In one embodiment, the curcin-deficient transgenic plants are produced by transforming *Jatropha* plant tissue. In another embodiment, the transformation is *Agrobacterium*-mediated transformation.

BRIEF DESCRIPTION OF THE FIGURES

[0018] Figure 1 shows the isolation of BAC clones that contain curcin genes and the isolation of curcin genes. Seven curcin gene-containing BAC clones were digested by *HindIII* (left panel) and its Southern blot was hybridized with a probe comprising the *Curcin 1* coding region (right panel). The molecular weights of standard DNA markers (New England Biolabs, # N3012L) are indicated in kilobases (Kb). M, molecular marker.

[0019] Figure 2 shows the nucleotide sequence of *Curcin 1* gene (SEQ ID NO:1) and its deduced amino acid sequence (SEQ ID NO:2). The transcription initiation site of *Curcin 1* is indicated as position 1 on the nucleotide sequence. The nucleotide sequences of the intron in the 5' untranslated region (5'UTR) are shown in lower-case letters. The positions of oligo primers are highlighted in bold italics with arrows indicating forward or reverse direction. The two *Sau3AI* sites in the coding region are displayed in bold and underlined. Only the second poly(A) signal sequence is highlighted in bold.

[0020] Figure 3 shows the nucleotide sequence of *Curcin 2* gene (SEQ ID NO:3) and its deduced amino acid sequence (SEQ ID NO:4). The putative transcription initiation site of *Curcin 2* is indicated as position 1 on the nucleotide sequence. The putative nucleotide sequences of the intron in the 5' untranslated region (5'UTR) are shown in lower-case letters. The positions of oligo primers are highlighted in bold italics with arrows indicating forward or reverse direction. The two *Sau3AI* sites in the coding region are displayed in bold and underlined. Only the second poly(A) signal sequence is highlighted in bold. The annotation of *Curcin 2* was predicted based on the annotation of *Curcin 1*.

[0021] Figure 4 shows the nucleotide sequence of *Curcin 2A* gene (SEQ ID NO:5) and its deduced amino acid sequence (SEQ ID NO:6). The putative transcription initiation site of *Curcin 2A* is indicated as position 1 on nucleotide sequence. The nucleotide sequences of the intron in the 5' untranslated region (5'UTR) are shown in lower-case letters. The positions of oligo primers are highlighted in bold italics with arrows indicating forward or reverse direction. The *Sau3AI* site in the coding region is displayed in bold and underlined. Only the second poly(A) signal sequence is highlighted in bold.

[0022] Figure 5 shows detection of curcin gene polymorphism by cleaved amplified polymorphic sequence (CAPS) analysis. PCR fragments amplified with primers CF2 and CR2 from BAC clone 121E10 (lane B) or genomic DNA (lane G) were digested with *Sau3AI*. *Curcin 1* produced three bands (195bp, 243bp and 582bp), *Curcin 2* also produced three bands (195bp, 243bp and 580bp) whereas *Curcin 2A* produced two bands (243bp and 775bp). BAC clone 121E10 only contains *Curcin 1* and *Curcin 2*, therefore, the digested PCR products (lane B) have three bands (195-bp band, 243-bp band, 580-bp and 582-bp bands as they could not be separated in 1.2% agarose gel). Genomic DNA contains all of the three curcin genes. The digested PCR products of genomic DNA (lane G) had all of the bands as shown in lane B and an additional band with size at 775bp. M, 100-bp DNA ladder is indicated in base pairs (New England Biolabs, # N3231L).

[0023] Figure 6 shows an alignment of curcin proteins identified from *J. curcas*. Amino acid residues showing identity among eight curcin genes are shown in black background. *Dart* indicates the cleavage site of N-terminal signal peptide (Lin et al., 2003a). The amino acid sequences identifiers are as follows: Curcin 2A – SEQ ID NO:6; ABZ04128 – SEQ ID NO:10; AAR08395 – SEQ ID NO:11; ABW17545 – SEQ ID NO:12; Curcin 1 – SEQ ID NO:2; AAL58089 – SEQ ID NO:13; Curcin 2 – SEQ ID NO:4; and AAL86778 – SEQ ID NO:14.

[0024] Figure 7 shows a phylogenetic tree of the curcin proteins identified from *J. curcas*. The phylogenetic tree was created by unclustalW.

[0025] Figures 8A-8C show the expression of curcin genes in *J. curcas*. **Fig 8A:** Detection of the expression of *Curcin 1* and *Curcin 2A* genes in jatropha by RT-PCR. The primers for *Curcin 1* were CF and C1SR (Table 1), whereas the primers for *Curcin 2A* were CF and C2ASR (Table 1). L, RNA sample from leaves; E, RNA sample from endosperm. **Fig. 8B:** Expression of *Curcin 1* in endosperm at different developmental stages. Transcript levels were measured by real-time PCR and the identity of the amplified sequences was confirmed by sequencing. *Jatropha Actin* gene was used as an internal control, and relative values normalized to the 6-week-old seeds (6 wk) are shown. The primers for *Curcin 1* were C1SF and C1SR, and the primers for the *jatropha Actin* gene were Jc actin F2 and Jc actin R1 (Table 1). **Fig. 8C:** Expression of *Curcin 2* in leaves. Transcript levels were measured by real-time PCR and the identity of the amplified sequences was confirmed by sequencing. *Jatropha Actin* gene was used as an internal control, and relative values normalized to the young leaves (YL) are shown. The primers for *Curcin 2A* were C2ASF and C1SR (Table 1), and the primers for the *jatropha Actin* gene were Jc actin F2 and Jc actin R1 (Table 1). YL, young leaves; FL, full-expanded leaves; OL, old leaves; IL, full-expanded leaves infected with mealybuds (*Pseudococcidae hirsutus*).

[0026] Figure 9 shows detection of curcin proteins in *J. curcas* showing a Western blot probed with anti-Curcin 1N antibodies. A band with molecular size of about 34 to about 35 kDa (presumably the Curcin 2A protein) was detected in leaves. Another band with molecular size of about 28 kDa (presumably the mature Curcin 1 protein) was detected in the seeds of all the tested accessions. An unknown protein with molecular size of about 39 kDa was detected in leaves, calli, seeds but not in roots. Similarly, an unknown protein with molecular size of about 16 kDa was detected in seeds. The molecular weights of standard protein markers (Amersham Biosciences, RPN755) are shown in kilodaltons (kDa). L, leaves; C, Calli; R,

roots; S1, seeds from Indonesia accession; S2, seeds from India accession; S3, seeds from China accession; S4, seeds from South America accession.

[0027] Figures 10A-10D show subcellular immunogold localization of Curcin 1 proteins in endosperm cells of developing seeds. **Fig. 10A:** Endosperm cells of a six-week-old seed. Bar = 5 μm . **Fig. 10B:** Preimmune serum control showing no immunolabelling of protein body. Bar = 0.2 μm . **Fig. 10C:** Curcin 1 proteins immunolocalized to protein body. Bar = 0.2 μm . **Fig. 10D:** Oil bodies and cell walls of endosperm cells after immunolabelling with anti-Curcin 1C antibody. Bar = 0.5 μm . *Arrows* indicate plastids with starchy bodies in white colour. *Darts* indicate curcin proteins immunolabelled with anti-Curcin 1C antibody. cw, cell wall; ob, oil body; pb, protein body.

[0028] Figures 11A-11D show subcellular immunogold localization of Curcin 2A proteins in vacuolar contents of leaf mesophyll cells. **Fig. 11A:** Mesophyll cells after immunolabelling with preimmune serum. Bar = 2 μm . **Fig. 11B:** Preimmune serum control showing no immunolabelling of vacuolar contents (high magnification of area indicated with a frame in **Fig. 11A**). Bar = 0.2 μm . **Fig. 11C:** Mesophyll cells after immunolabelling with anti-Curcin 1C antibody. Bar = 2 μm . **Fig. 11D:** Curcin 2A proteins immunolocalized to vacuolar contents (high magnification of area indicated with a frame in **Fig. 11C**). Bar = 0.2 μm . *Darts* indicate curcin proteins immunolabelled with anti-Curcin 1C antibody. C, chloroplast; cw, cell wall; v, vacuole.

[0029] Figures 12A-12D show subcellular immunogold localization of Curcin 2A proteins in secondary cell walls of leaf tracheary elements. **Fig. 12A:** Leaf cross section after immunolabelling with anti-Curcin 1C antibody. Two adjacent tracheary elements, including an immature tracheary element (I) and a mature tracheary element (M), are surrounded by mesophyll cells. Bar = 5 μm . **Fig. 12B:** Preimmune serum control showing no immunolabelling of secondary cell wall of tracheary element. Bar = 0.2 μm . **Fig. 12C:** Curcin 2A proteins immunolocalized to the secondary cell wall of the immature tracheary element (high magnification of area indicated with a frame at lower left of **Fig. 12A**). Bar = 0.2 μm . **Fig. 12D:** Curcin 2A proteins immunolocalized to the secondary cell wall of the mature tracheary element (high magnification of area indicated with a frame at upper right of **Fig. 12A**). Bar = 0.2 μm . *Arrows* indicate pit area on the walls of tracheary elements. *Darts* indicate curcin proteins immunolabelled with anti-Curcin 1C antibody. mc, mesophyll cell; pcw, primary cell wall; scw, secondary cell wall; v, vacuole.

[0030] Figures 13A and 13B show the characterization of *Curcin 1* and *Curcin 2A* promoters. **Fig. 13A:** Transient expression of GFP in developing endosperm of *J. curcas* after bombardment with empty vector pC1300 (left panel), *P_{Ubi}:GFP:T_{Nos}* (middle panel) and *P_{CIP3}:GFP:T_{Nos}* (right panel) genes. GFP, fluorescence channel; Ph2, phase contrast channel; Merge, merge of the GFP and Ph2 channels. Bar = 10 μ m. **Fig. 13B:** Transient expression of GUS activity in the leaf tissues of *J. curcas* after bombardment with empty vector pC1300 (left panel), *P_{35S}:GUS:T_{Nos}* (middle panel) and *P_{C2AP3}:GUS:T_{Nos}* (right panel) genes. Cells showing GUS activities are in blue color (indicated with *darts*) after stained with X-Gluc solution.

[0031] Figure 14 shows the T-DNA region of the binary plasmid pANDA35HKC1 (not drawn to scale). NPT II, neomycin phosphotransferase gene; HPT, hygromycin phosphotransferase gene; 35S pro., 35S promoter; Curcin 1, partial sequence of *Curcin 1*; linker, *Gus* linker; T, terminator of nopaline synthase gene; RB, right border; LB, left border.

[0032] Figure 15 shows T₀ transgenic jatropha plants with knock-down expression of curcin genes. 8-month-old T₀ transgenic plants transformed with a binary vector pANDA35HKC1 that carries RNAi cassette designed for the curcin genes are shown.

[0033] Figure 16 shows the results of Southern blot analysis of the transgenic jatropha plants. **Fig. 16:** Southern blot analysis of the transgenic jatropha plants. Souther blots were probed with *Gus* linker (upper panel) or *Hpt* (lower panel) probes. Arrow indicates the position of RNAi cassette when the genomic DNAs of the transgenic plants were digested by restriction enzymes *KpnI* and *SpeI*. The primers for the *Gus* linker were Gus F and Gus R (Table 1). M, molecular maker; Wt, wild-type jatropha plants; L3 to L26, independent T₀ transgenic jatropha plants.

[0034] Figures 17A to 17B shows the expression of double-strand-RNA (dsRNA) from the RNAi cassette and *Curcin 2A*. **Fig. 17A:** Expression of dsRNA of *curcin 1* in young leaves. Transcript levels of dsRNA of *curcin 1* were measured by real-time PCR for the *Gus* linker and the identity of the amplified sequences was confirmed by sequencing. Jatropha *Actin* gene was used as an internal control, and relative values normalized to transgenic line 17 (L17) are shown. The primers for the *Gus* linker were Gus RT F1 and Gus RT R1, and the primers for the jatropha *Actin* gene were Jc actin F2 and Jc actin R1 (Table 1). Wt, wild-type plant; L17 to L46, independent T₀ plants. **Fig. 17B:** Expression of *Curcin 2A* in young leaves. Transcript levels were measured by real-time PCR and the identity of the amplified sequences was confirmed by sequencing. Jatropha *Actin* gene was used as an internal control, and relative

values normalized to control transgenic plant transformed with empty vector pC1300 are shown. The primers for *Curcin 2A* were C2ASF and C2ASR (Table 1), and the primers for the jatropha *Actin* gene were Jc actin F2 and Jc actin R1 (Table 1). Wt, wild-type plant; L17 to L46, independent T₀ plants.

[0035] Figure 18 shows the expression of Curcin 2A proteins in young leaves of transgenic jatropha plants. **Fig. 18:** Expression of Curcin 2A in young leaves of transgenic jatropha plants. The Curcin 2A proteins (indicated with arrow) were detected with anti-Curcin 1 antibodies. Wt, wild-type plant; L17 to L46 are independent transgenic plants.

[0036] Figures 19A-19D show the inheritance of transgenes to the T₁ generation. **Fig. 19A and 19B:** Shoots (A) and roots (B) of wild-type jatropha seedlings germinated and grown on half-strength MS medium containing 10 mg/L hygromycin for 12 days. **Fig. 19C and 19D:** Shoots (C) and roots (D) of T₁ seedlings of transgenic line L26 germinated and grown on half-strength MS medium containing 10 mg/L hygromycin for 12 days.

DETAILED DESCRIPTION OF THE INVENTION

[0037] The present invention relates to the isolation of *Jatropha curcas* curcin genes and tissue-specific promoters and to the production of curcin-deficient *Jatropha* plants. More specifically, the present invention relates to the isolation of *Jatropha curcas* *Curcin 1*, *Curcin 2* and *Curcin 2A*. The present invention further relates to of the *Curcin 1*, *Curcin 2A* and *Curcin 2* genes and more particularly to tissue specific promoters of the *Curcin 1* and *Curcin 2A* genes. The present invention further relates to production of curcin-deficient transgenic jatropha plants by using RNAi technology to suppress curcin gene expression.

[0038] Thus in a first aspect, the present invention provides the sequences of three *Jatropha curcas* curcin genes. In one embodiment, the curin gene is *Curcin 1*. The nucleotide sequence of *Curcin 1* is set forth in SEQ ID NO:1. The coding sequence for Curcin 1 comprises nucleotides 474 - 1355 of SEQ ID NO:1. The protein sequence for Curcin 1 is set forth in SEQ ID NO:2. As shown herein, *Curcin 1* is specifically expressed in the endosperm of developing seeds. Curcin 1 localizes to protein bodies of endosperm.

[0039] In a second embodiment, the curin gene is *Curcin 2*. The nucleotide sequence of *Curcin 2* is set forth in SEQ ID NO:3. The coding sequence for Curcin 2 comprises nucleotides 439 - 1368 of SEQ ID NO:3. The protein sequence for Curcin 2 is set forth in SEQ ID NO:4. As shown herein, *Curcin 2* is not expressed in developing seeds or leaf.

[0040] In a third embodiment, the curin gene is *Curcin 2A*. The nucleotide sequence of *Curcin 2A* is set forth in SEQ ID NO:5. The coding sequence for *Curcin 2A* comprises nucleotides 475 - 1404 of SEQ ID NO:5. The protein sequence for *Curcin 2A* is set forth in SEQ ID NO:6. As shown herein, *Curcin 2A* is specifically expressed in the leaves. *Curcin 2A* was detected in vacuoles of mesophyll cells and secondary cell wall of leaf tracheary elements.

[0041] In a second aspect, the present invention provides the tissue-specific promoters of two *Jatropha curcas* curin genes. In one embodiment, the promoter is derived from the *Curcin 1* gene. The nucleotide sequence of *Curcin 1* promoter is set forth in SEQ ID NO:7. Promoter deletion studies indicate that the *Curcin 1* promoter of 2888 base pairs is sufficient for its tissue-specific expression in the seed endosperm. Thus, the *Curcin 1* promoter is an endosperm-specific promoter. Fragments of this sequence are also active as tissue specific promoters, i.e., as endosperm-specific promoters. Such fragments include the following: (a) nucleotides 1 to 2888 of SEQ ID NO:7, (b) nucleotides 1142 to 3181 of SEQ ID NO:7, (c) SEQ ID NO:7 with nucleotides 2944 to 3170 deleted, (d) nucleotides 1142 to 3181 of SEQ ID NO:7 with nucleotides 2944 to 3170 deleted and (e) nucleotides 2688 to 3181 of SEQ ID NO:7 with nucleotides 2944 to 3170 deleted. The *Curcin 1* promoter or its fragments can be used to specifically express downstream DNA of interest in seed endosperm of transgenic plants. In some embodiments, the endosperm-specific *Curcin 1* promoter can be used to express DNA of interest, for example a target gene such as a gene encoding an enzyme or a protein, in endosperm of *jatropha* seeds as well as other crop seeds.

[0042] In a second embodiment, the promoter is derived from the *Curcin 2A* gene. The nucleotide sequence of *Curcin 2A* promoter is set forth in SEQ ID NO:8. Promoter deletion studies indicate that the *Curcin 2A* promoter of 1793 base pairs is sufficient for its tissue-specific expression in leaf tissues. Thus, the *Curcin 2A* promoter is a leaf-specific promoter. Fragments of this sequence are also active as tissue-specific promoters. However, the specificity changes from leaf-specific to non-tissue specific when some fragments are deleted. Such fragments that are active as non-tissue specific promoters include the following: (a) nucleotides 912 to 2087 of SEQ ID NO:8, (b) nucleotides 1 to 2087 of SEQ ID NO:8 with nucleotides 1853 to 2076 deleted, (c) nucleotides 912 to 2087 of SEQ ID NO:8 with nucleotides 1853 to 2076 deleted and (d) nucleotides 1751 to 2087 of SEQ ID NO:8 with nucleotides 1853 to 2076 deleted.

[0043] The *Curcin 2A* promoter can be used to specifically express downstream DNA of interest in leaf tissues of transgenic plants. In some embodiments, the leaf-specific *Curcin 2A* promoter can be used to express DNA of interest, for example a target gene such as a gene encoding an enzyme or a protein, in leaves or other green-tissues of *Jatropha* as well as other crops. In other embodiments, the non-tissue specific fragments of the *Curcin 2A* promoter can be used to direct the non-tissue specific expression of DNA of interest, such as enzymes or proteins, in *Jatropha*, as well as other crop seeds.

[0044] In a third embodiment, the promoter is derived from the *Curcin 2* gene. The nucleotide sequence of *Curcin 2* promoter is set forth in SEQ ID NO:9. As described herein, this promoter is not expressed in endosperm or leaf tissues, but may be activated and expressed upon activation by jasmonic acid (JA) upon attack by insects and/or upon activation by ethylene during defense reaction or leaf senescence.

[0045] The promoters of the present invention are particularly useful for preparing transgenic plants, including *Jatropha*, to contain a DNA of interest. Other plant species include, but are not limited to, castor bean, palm oil, coconut, peanut, rapeseeds, soybean, sunflower, cassava, tomato, beans, cotton, cucumber, cabbage, barley, oats, wheat, corn and rice. In one embodiment, the *Curcin 1* promoter or its fragments can be used to direct expression of the DNA of interest, such as a gene of interest, in the endosperm of seeds of *Jatropha* or other plant seeds. In another embodiment, the *Curcin 2A* promoter can be used to direct expression of the DNA of interest, such as a gene of interest, in leaves or other green tissue of *Jatropha* or other plant species. In a further embodiment, the *Curcin 2* promoter can be used to direct expression of the DNA of interest, such as a gene of interest, under inducible conditions, such as upon activation by JA upon attack by insects and/or upon activation by ethylene during defense reaction or leaf senescence.

[0046] The DNA that is inserted (the DNA of interest) into plants of the genera *Jatropha* or other plant species is not critical to the transformation process. Generally the DNA that is introduced into a plant is part of a construct. The DNA may be a gene of interest, e.g., a coding sequence for a protein, or it may be a sequence that is capable of regulating expression of a gene, such as an antisense sequence, a sense suppression sequence, a post-transcriptional gene silencing sequence (an RNAi sequence such as an siRNA, shRNA or dsRNA) or a micro-RNA (miRNA) sequence. The construct typically includes regulatory regions operatively linked to the 5' side of the DNA of interest and/or to the 3' side of the DNA of interest. A cassette containing all of these elements is also referred to herein as an expression

cassette. The expression cassettes may additionally contain 5' leader sequences in the expression cassette construct. The regulatory regions (i.e., promoters, transcriptional regulatory regions, and translational termination regions) and/or the polynucleotide encoding a signal anchor may be native/analogous to the host cell or to each other. The promoters and tissue-specific promoters identified herein are particularly useful for preparing constructions for the transformation of *Jatropha*, as well as for the transformation of other plant species. Alternatively, the regulatory regions and/or the polynucleotide encoding a signal anchor may be heterologous to the host cell or to each other. See, U.S. Patent No. 7,205,453 and U.S. Patent Application Publication Nos. 2006/0218670, 2006/0248616 and 20090100536, and the references cited therein. The expression cassettes may additionally contain 5' leader sequences in the expression cassette construct. Such leader sequences can act to enhance translation. Translation leaders are known in the art and include those described in International Publication No. WO 2008/094127 and the references cited therein.

[0047] In a preferred embodiment, the promoters and tissue-specific promoters identified herein are used in the practice of the invention. The promoters can be selected based on the desired outcome. That is, the nucleic acids can be combined with the tissue-specific promoters for expression in a specific host cell or host tissue of interest.

[0048] The DNA of interest that is under control of a promoter, such as a promoter described herein, may be any DNA as defined herein and may be used to alter any characteristic or trait of a plant species into which it is introduced, including *Jatropha* as a preferred embodiment. In one embodiment, the DNA of interest is introduced into a plant in order to enhance a trait of the plant. In another embodiment, an enhanced agronomic trait may be characterized by enhanced plant morphology, physiology, growth and development, yield, nutritional enhancement, disease or pest resistance, or environmental or chemical tolerance. In some aspects, the enhanced trait is selected from group of enhanced traits consisting of enhanced water use efficiency, enhanced temperature tolerance, increased yield, enhanced nitrogen use efficiency, enhanced seed protein enhanced seed oil and enhanced biomass. Increase yield may include increased yield under non-stress conditions and increased yield under environmental stress conditions. Stress conditions may include, for example, drought, shade, fungal disease, viral disease, bacterial disease, insect infestation, nematode infestation, extreme temperature exposure (cold or hot), osmotic stress, reduced nitrogen nutrient availability, reduced phosphorus nutrient availability and high plant density. In some embodiments, the DNA of interest may be used to modify metabolic pathways, such

as fatty acid biosynthesis or lipid biosynthesis pathways in seeds, or to modify resistance to pathogens in *Jatropha* or other plant species.

[0049] Generally, the expression cassette may additionally comprise a selectable marker gene for the selection of transformed cells. Selectable marker genes are utilized for the selection of transformed cells or tissues. Usually, the plant selectable marker gene will encode antibiotic resistance, with suitable genes including at least one set of genes coding for resistance to the antibiotic spectinomycin, the streptomycin phosphotransferase (spt) gene coding for streptomycin resistance, the neomycin phosphotransferase (nptII) gene encoding kanamycin or geneticin resistance, the hygromycin phosphotransferase (hpt or aphiv) gene encoding resistance to hygromycin, acetolactate synthase (als) genes. Alternatively, the plant selectable marker gene will encode herbicide resistance such as resistance to the sulfonylurea-type herbicides, glufosinate, glyphosate, ammonium, bromoxynil, imidazolinones, and 2,4-dichlorophenoxyacetate (2,4-D), including genes coding for resistance to herbicides which act to inhibit the action of glutamine synthase such as phosphinothricin or basta (e.g., the bar gene). See generally, International Publication No. WO 02/36782, U.S. Patent No. 7,205,453 and U.S. Patent Application Publication Nos. 2006/0218670, 2006/0248616, 2007/0143880 and 20090100536, and the references cited therein. This list of selectable marker genes is not meant to be limiting. Any selectable marker gene can be used. The selectable marker gene is also under control of a promoter operable in the plant species to be transformed. Such promoters include those described in International Publication No. WO 2008/094127 and the references cited therein.

[0050] Where appropriate, the DNA of interest may be optimized for increased expression in the transformed plant. That is, the coding sequences can be synthesized using plant-preferred codons for improved expression. Methods are available in the art for synthesizing plant-preferred genes. See, for example, U.S. Patent Nos. 5,380,831, 5,436,391, and 7,205,453 and U.S. Patent Application Publication Nos. 2006/0218670 and 2006/0248616.

[0051] In preparing the expression cassette, the various DNA fragments may be manipulated, so as to provide for the DNA sequences in the proper orientation and, as appropriate, in the proper reading frame. Toward this end, adapters or linkers may be employed to join the DNA fragments or other manipulations may be involved to provide for convenient restriction sites, removal of superfluous DNA, removal of restriction sites, or the like. For this purpose, *in vitro* mutagenesis, primer repair, restriction, annealing, resubstitutions, e.g. transitions and transversions may be involved.

[0052] Once a nucleic acid has been cloned into an expression vector, it may be introduced into a plant cell using conventional transformation procedures. The term "plant cell" is intended to encompass any cell derived from a plant including undifferentiated tissues such as callus and suspension cultures, as well as plant seeds, pollen or plant embryos. Plant tissues suitable for transformation include leaf tissues, root tissues, meristems, protoplasts, hypocotyls, cotyledons, scutellum, shoot apex, root, immature embryo, pollen, and anther. "Transformation" means the directed modification of the genome of a cell by the external application of recombinant DNA from another cell of different genotype, leading to its uptake and integration into the subject cell's genome. In this manner, genetically modified plants, plant cells, plant tissue, seed, and the like can be obtained.

[0053] DNA constructs containing the promoters of the present invention can be used to transform any plant and particularly *Jatropha* plants. The constructs may be introduced into the genome of the desired plant host by a variety of conventional techniques. Techniques for transforming a wide variety of higher plant species are well known and described in the technical and scientific literature. Transformation protocols may vary depending on the type of plant or plant cell, i.e., monocot or dicot, targeted for transformation, as is well known to the skilled artisan. For example, the DNA construct may be introduced directly into the genomic DNA of the plant cell using techniques such as electroporation and microinjection of plant cell protoplasts, or the DNA constructs can be introduced directly to plant tissue using ballistic methods, such as DNA particle bombardment. Alternatively, the DNA constructs may be combined with suitable T-DNA flanking regions and introduced into a conventional *Agrobacterium tumefaciens* host vector. The virulence functions of the *Agrobacterium tumefaciens* host will direct the insertion of the construct and adjacent marker into the plant cell DNA when the cell is infected by the bacteria. Thus, any method, which provides for effective transformation/transfection may be employed. See, for example, U.S. Patent Nos. 7,241,937, 7,273,966 and 7,291,765 and U.S. Patent Application Publication Nos. 2007/0231905 and 2008/0010704 and references cited therein. See also, International Published Application Nos. WO 2005/103271 and WO 2008/094127 and references cited therein.

[0054] In one embodiment, the explant tissue can be co-cultured with an *Agrobacterium* strain harboring a DNA construct containing a gene or nucleic acid of interest using techniques well known in the art. Transformed tissue can be selected using conventional techniques well known in the art. In another embodiment, the embryogenic liquid suspension

cultures can be co-cultured with an *Agrobacterium* strain harboring a DNA construct containing a gene or nucleic acid of interest using techniques well known in the art. Transformed tissue can be selected using conventional techniques well known in the art. In a further embodiment, the DNA can be introduced into the explant tissue or cells of the embryogenic liquid suspension culture using conventional techniques, such as particle bombardment. Transformed tissue can be selected using conventional techniques well known in the art. Transformed or transgenic plants can be regenerated using the methods described herein. In a further embodiment, *Jatropha* can be transformed with *Agrobacterium* as described in International Patent Application No. PCT/SG2009/000015 filed on 7 January 2009, incorporated herein by reference.

[0055] Transformed plant cells which are derived by any of the above transformation techniques can be cultured to regenerate a whole plant which possesses the transformed genotype and thus the desired phenotype, e.g., a transgenic plant. A "transgenic plant" is a plant into which foreign DNA has been introduced. A "transgenic plant" encompasses all descendants, hybrids, and crosses thereof, whether reproduced sexually or asexually, and which continue to harbor the foreign DNA. Regeneration techniques rely on manipulation of certain phytohormones in a tissue culture growth medium, typically relying on a biocide and/or herbicide marker which has been introduced together with the desired nucleotide sequences. See for example, International Published Application No. WO 2008/094127 and references cited therein.

[0056] The foregoing methods for transformation are typically used for producing a transgenic variety in which the expression cassette is stably incorporated. After the expression cassette is stably incorporated in transgenic plants, it can be transferred to other plants by sexual crossing. In one embodiment, the transgenic variety could then be crossed, with another (non-transformed or transformed) variety, in order to produce a new transgenic variety. Alternatively, a genetic trait which has been engineered into a particular cotton line using the foregoing transformation techniques could be moved into another line using traditional backcrossing techniques that are well known in the plant breeding arts. For example, a backcrossing approach could be used to move an engineered trait from a public, non-elite variety into an elite variety, or from a variety containing a foreign gene in its genome into a variety or varieties which do not contain that gene. As used herein, "crossing" can refer to a simple X by Y cross, or the process of backcrossing, depending on the context.

Any of a number of standard breeding techniques can be used, depending upon the species to be crossed.

[0057] Once transgenic plants of this type are produced, the plants themselves can be cultivated in accordance with conventional procedures. Transgenic seeds can, of course, be recovered from the transgenic plants. These seeds can then be planted in the soil and cultivated using conventional procedures to produce transgenic plants.

[0058] In a third aspect, the present invention provides transgenic jatropha plants that are curcin-deficient and to the production of curcin-deficient transgenic jatropha plants. In one embodiment, a curcin-deficient transgenic plant comprises a nucleic acid stably integrated in its genome, wherein the nucleic acid encodes a double stranded RNA (dsRNA) targeted to a curcin gene. In another embodiment, a curcin-deficient transgenic plant comprises a nucleic acid stably integrated in its genome, wherein the nucleic acid encodes a short interfering RNA (siRNA) targeted to a curcin gene. In an additional embodiment, a curcin-deficient transgenic plant comprises a nucleic acid stably integrated in its genome, wherein the nucleic acid encodes a short hairpin RNA (shRNA) targeted to a curcin gene. In one embodiment, the curcin gene is *Curcin 1*. In another embodiment, the curcin gene is *Curcin 2*. In an additional embodiment, the curcin gene is *Curcin 2A*. In one embodiment, the nucleic acid includes a portion of the *Curcin 1* gene. In another embodiment, the nucleic acid includes a portion of the *Curcin 2* gene. In an additional embodiment, the nucleic acid includes a portion of the *Curcin 2A* gene. In one embodiment, the curcin-deficient transgenic plants are produced by transforming jatropha plant tissue. The nucleic acid encoding the dsRNA, siRNA or shRNA is generally included in a construct, such as constructs described herein. The nucleic acid is under the operable control of a promoter. Suitable promoters that can be used include those described herein. The construct may further include other regulatory sequences, such as those described herein.

[0059] For example, a construct can be prepared that includes a sequence that is transcribed into an RNA that can anneal to itself, e.g., a double stranded RNA having a stem-loop structure. In some embodiments, one strand of the stem portion of a double stranded RNA comprises a sequence that is similar or identical to the sense coding sequence, or a fragment thereof, of the polypeptide of interest, and that is from about 10 nucleotides to about 2,500 nucleotides in length. The length of the sequence that is similar or identical to the sense coding sequence can be from 10 nucleotides to 500 nucleotides, from 15 nucleotides to 300 nucleotides, from 20 nucleotides to 100 nucleotides, or from 25 nucleotides to 100

nucleotides. The other strand of the stem portion of a double stranded RNA comprises a sequence that is similar or identical to the antisense strand, or a fragment thereof, of the coding sequence of the polypeptide of interest, and can have a length that is shorter, the same as, or longer than the corresponding length of the sense sequence. In some cases, one strand of the stem portion of a double stranded RNA comprises a sequence that is similar or identical to the 3' or 5' untranslated region, or a fragment thereof, of the mRNA encoding the polypeptide of interest, and the other strand of the stem portion of the double stranded RNA comprises a sequence that is similar or identical to the sequence that is complementary to the 3' or 5' untranslated region, respectively, or a fragment thereof, of the mRNA encoding the polypeptide of interest. In other embodiments, one strand of the stem portion of a double stranded RNA comprises a sequence that is similar or identical to the sequence of an intron or a fragment thereof in the pre-mRNA encoding the polypeptide of interest, and the other strand of the stem portion comprises a sequence that is similar or identical to the sequence that is complementary to the sequence of the intron or fragment thereof in the pre-mRNA.

[0060] The loop portion of a double stranded RNA can be from 3 nucleotides to 5,000 nucleotides, e.g., from 3 nucleotides to 25 nucleotides, from 15 nucleotides to 1,000 nucleotides, from 20 nucleotides to 500 nucleotides, or from 25 nucleotides to 200 nucleotides. The loop portion of the RNA can include an intron or a fragment thereof. A double stranded RNA can have zero, one, two, three, four, five, six, seven, eight, nine, ten, or more stem-loop structures.

[0061] In one embodiment described herein, a construct is prepared which comprises a constitutive promoter, such as the CaMV 35S promoter, a partial sequence of one of the *Curcin* genes described herein in a sense orientation, a linker, the partial sequence of the same *Curcin* gene in antisense orientation and terminator, such as the NOS terminator. In one embodiment, the *Curcin* gene is the *Curcin 1* gene and the partial sequence of the *Curcin 1* gene comprises 862 nucleotides and is set forth in SEQ ID NO:53. In another embodiment, the *Curcin* gene is the *Curcin 1* cDNA and the sequence of the *Curcin 1* cDNA comprises 1161 nucleotides and is set forth in SEQ ID NO:54. In an additional embodiment, *Curcin* gene is the *Curcin 2A* cDNA and the sequence of the *Curcin 2A* cDNA comprises 1176 nucleotides and is set forth in SEQ ID NO:55. In a further embodiment, *Curcin* gene is the putative *Curcin 2* cDNA and the sequence of the putative *Curcin 2* cDNA comprises 1140 nucleotides and is set forth in SEQ ID NO:56. In addition to the sequences set forth in SEQ ID NOs:53-56, partial sequences of these sequences can also be used. The partial sequences

may comprise 50 nucleotides, 100 nucleotides, 150 nucleotides, 200 nucleotides, 250 nucleotides, 300 nucleotides, 350 nucleotides, 400 nucleotides, 450 nucleotides, 500 nucleotides, 550 nucleotides, 600 nucleotides, 650 nucleotides, 700 nucleotides, 750 nucleotides, 800 nucleotides, 850 nucleotides or more up to less than the number of nucleotides in the full length sequences. It is understood the the partial sequences may comprise any number of nucleotides between the number of nucleotides illustrated, for example, 60 nucleotides, 175 nucleotides, etc. It is also understood that the partial sequences may comprise a range of nucleotides between the number illustrated or any number within those illustrated and the number of nucleotides in the full length sequence, for example, 50-100, 50-800, 100-800, 175-862 of SEQ ID NO:53, and the like.

[0062] An siRNA also can refer to an RNA duplex of nucleotides, or, in some alternative aspects, a single molecule of RNA (which can, in some embodiments, have secondary structure, such as loops so as to form a shRNA) that is targeted to a nucleic acid, e.g., a gene, of interest. A "RNA duplex" refers to the structure formed by the complementary pairing between at least two regions of a RNA molecule. Thus, the "RNA duplex" can comprise one, two, three or more RNA molecules. siRNA is "targeted" to a gene in that the nucleotide sequence of the duplex portion of the siRNA is complementary to a nucleotide sequence of the targeted gene. Thus, by using the sequence of a target gene, any siRNA can be routinely designed and made. In some embodiments, the length of the duplex of siRNAs is less than 30 nucleotides. In some embodiments, the length of the duplex of siRNAs is more than 30 nucleotides. In some embodiments, the duplex can be 40, 39, 38, 37, 36, 35, 34, 33, 32, 31, 30, 29, 28, 27, 26, 25, 24, 23, 22, 21, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11 or 10 or less nucleotides in length. In some embodiments, the length of the duplex is 19-25 nucleotides in length. The RNA duplex portion of the siRNA can be part of a hairpin structure. In one aspect, there is no hairpin structure in an siRNA of the invention. In addition to the duplex portion, the hairpin structure may contain a loop portion positioned between the two sequences that form the duplex. The loop can vary in length. In some embodiments the loop is 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12 or 13 or more nucleotides in length. The hairpin structure can also contain 3' or 5' overhang portions. In some embodiments, the overhang is a 3' or a 5' overhang 0, 1, 2, 3, 4 or 5 nucleotides in length. The siRNA can be encoded by a nucleic acid sequence, and the nucleic acid sequence can also include a promoter. The nucleic acid sequence can also include a polyadenylation signal. In some embodiments, the polyadenylation signal is a synthetic minimal polyadenylation signal. The siRNA can be

entirely, or in part, comprised of synthetic nucleotides, natural bases or modified bases. The siRNA molecule targeted to a *Curcin* gene of interest, such as the *Curcin 1* gene, the *Curcin 2* gene or the *Curcin 2A* gene described herein, can be designed using algorithms well known in the art.

[0063] For further description of RNAi techniques, see, e.g., U.S. Patent Nos. 5,034,323; 6,326,527; 6,452,067; 6,573,099; 6,753,139; and 6,777,588. See also International Patent Publications WO 97/01952, WO 98/36083, WO 98/53083, WO 99/32619 and WO 01/75164; and U.S. Patent Publications 2003/0175965, 2003/0175783, 2003/0180945, 2004/0214330, 2005/0244858, 2005/0277610, 2007/0265220, 2009/0215860, 2009/0308041 and 2010/0058498.

[0064] In accordance with this aspect of the invention, the curcin-deficient transgenic plants are produced by transforming jatropha plant tissue. In one embodiment, the transformation is *Agrobacterium*-mediated transformation. In another embodiment, the transformation is a ballistic method, such as DNA particle bombardment. In an additional embodiment, the transformation is direct delivery, such as electroporation or microinjection. In one embodiment, the *Agrobacterium*-mediated transformation is performed as described herein. In another embodiment, the *Agrobacterium*-mediated transformation is performed as described in International Patent Application No. PCT/SG2009/000015 filed on 7 January 2009, incorporated herein by reference. Plants cells are transformed as described herein. Transformed plant cells are selected as described herein. Transgenic plants are regenerated from the transformed plant cells as described herein. The transgenic plants are screened for stable incorporation of the RNAi construct and for curcin deficiency as described herein.

[0065] The practice of the present invention employs, unless otherwise indicated, conventional techniques of chemistry, molecular biology, microbiology, recombinant DNA, genetics, immunology, cell biology, cell culture and transgenic biology, which are within the skill of the art. See, e.g., Maniatis *et al.*, 1982, *Molecular Cloning* (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York); Sambrook *et al.*, 1989, *Molecular Cloning*, 2nd Ed. (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York); Sambrook and Russell, 2001, *Molecular Cloning*, 3rd Ed. (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York); Ausubel *et al.*, 1992), *Current Protocols in Molecular Biology* (John Wiley & Sons, including periodic updates); Glover, 1985, *DNA Cloning* (IRL Press, Oxford); Russell, 1984, *Molecular biology of plants: a laboratory course manual* (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.); Anand, *Techniques for the*

Analysis of Complex Genomes, (Academic Press, New York, 1992); Guthrie and Fink, *Guide to Yeast Genetics and Molecular Biology* (Academic Press, New York, 1991); Harlow and Lane, 1988, *Antibodies*, (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York); *Nucleic Acid Hybridization* (B. D. Hames & S. J. Higgins eds. 1984); *Transcription And Translation* (B. D. Hames & S. J. Higgins eds. 1984); *Culture Of Animal Cells* (R. I. Freshney, Alan R. Liss, Inc., 1987); *Immobilized Cells And Enzymes* (IRL Press, 1986); B. Perbal, *A Practical Guide To Molecular Cloning* (1984); the treatise, *Methods In Enzymology* (Academic Press, Inc., N.Y.); *Methods In Enzymology*, Vols. 154 and 155 (Wu et al. eds.), *Immunochemical Methods In Cell And Molecular Biology* (Mayer and Walker, eds., Academic Press, London, 1987); *Handbook Of Experimental Immunology*, Volumes I-IV (D. M. Weir and C. C. Blackwell, eds., 1986); Riott, *Essential Immunology*, 6th Edition, Blackwell Scientific Publications, Oxford, 1988; Fire et al., *RNA Interference Technology: From Basic Science to Drug Development*, Cambridge University Press, Cambridge, 2005; Schepers, *RNA Interference in Practice*, Wiley-VCH, 2005; Engelke, *RNA Interference (RNAi): The Nuts & Bolts of siRNA Technology*, DNA Press, 2003; Gott, *RNA Interference, Editing, and Modification: Methods and Protocols (Methods in Molecular Biology)*, Human Press, Totowa, NJ, 2004; Sohail, *Gene Silencing by RNA Interference: Technology and Application*, CRC, 2004.

EXAMPLES

[0066] The present invention is described by reference to the following Examples, which is offered by way of illustration and is not intended to limit the invention in any manner. Standard techniques well known in the art or the techniques specifically described below were utilized.

EXAMPLE 1

Materials and Methods

[0067] *Plant materials and growth condition: Jatropha curcas* Indonesia accession (JC-MD) was used for experiments except that it was specially mentioned. Plants were grown in experimental field in Singapore at temperatures ranging from 32°C during the day to 26°C at night, relative humidity averaged 85%, and the photoperiod was 12 to 13 h.

[0068] *Constructs*: To construct the expression vectors using GFP as the reporter gene, we amplified the *Pst*I-*Kpn*I fragment including GFP coding sequence from plasmid pSSZ41

(Kolesnik et al., 2004) using primer pairs GFPPstIF and GFPKpnIR (Table 1). PCR products were digested with restriction enzymes *PstI* and *KpnI* and then inserted into pCAMBIA1300 vector to generate pCGFP. The *Curcin 1* terminator was cloned into the region between *KpnI* and *EcoRI* sites of pCGFP to generate pCGFPT1. Promoters of curcin genes were cloned into pCGFPT1 at the region between *HindIII* and *PstI* sites to generate final expression constructs. The CaMV35S promoter in pC1305.1 was removed by digesting with *HindIII* and *NcoI* and the vector fragment was filled at ends and self-ligated to produce pC1305.1(-35S). Promoters of curcin genes were cloned into pC1305.1(-35S) at the region between *KpnI* and *PstI* to drive GUS reporter gene in the constructs. The RNAi construct for the curcin gene was made based on the pANDA vector pANDA35HK (Miki and Shimamoto 2004). Briefly, partial coding sequence of Curcin 1 was amplified using primers Curcin 1 TOPO-F and JCG-R (Table 1). The sequence of the amplified product is set forth in SEQ ID NO:53. The PCR product was cloned into pENTR D-TOPO (Invitrogen) and finally transferred into pANDA35HK vector to generate pANDA35HKC1 using Gateway[®] technology (Hartley et al., 2000).

TABLE 1
DNA Oligonucleotide Primers

Primer Name	Nucleotide Sequence (5' to 3') (SEQ ID NO:)
GFPPstIF	AACTGCAGATGGTGAGCAAGGGCGAGGAGCTG (15)
GFPKpnIR	GGGGTACCTTACTTGTACAGCTCGTCCATGCCGT (16)
CantigenF1	GCTGGTTCCACTCCAACCTTTAAC (17)
CantigenF2	CGGGATCCGGTACTTCTTCTTTAACG (18)
CantigenR1	TCAGACTTTGTATTTGACTGCATTC (19)
CantigenR2	CCCAAGCTTAATTATCTAATGCCTGCACCCC (20)
CantigenR3	CCCAAGCTTCACTAGGATATTGGTATAGTTTTC (21)
GSP1	GTCTGCTGGCTTTGAACTTT (22)
GSP1A	CACCCCTAAATCCACATCCT (23)
AAP	GGCCACGCGTCGACTAGTACGGGGGGGGGGGGGGGGG (24)
AUAP	GGCCACGCGTCGACTAGTAC (25)
C2RTF2	GGGCATCGGCTAGGGAAATA (26)
3'RACEF4	TCTCATCAAACCAAACTACAAA (27)
C2RTR2	AGAGGTCTCCCCAGTTGTTC (28)
5'RACEG4	TGAATCTTGCTGCCTCTG (29)
CTRTR2	GGGCATCGGCTAGGGAAATA (30)
3'RACEF1	GAGAGGATGTGGATTTAGG (31)
C2PF4	CTTCAAGACACTAGTTCAAAAA (32)
C1CDSR	TGATGAGAATGGACAACTAT (33)
C2AFAR3	CCATAACCAATGTATGATTTGGTA (34)
AD2	NGTCGASWGANAWGAA (35)
C2ACDSF	CCCAAGTAAAAGGTCTCA (36)

C2TR2EcoRI	CGGAATTCGTATTATTTGGATGGTAGAAAATT (37)
CF2	GATATTTGTGTTTCTTCAT (38)
CR2	TTTGTTGTCCTTTATTTATGCT (39)
CF	TTTACTTCCCCGTTTGCTCA (40)
C1SR	TATAGCATTAGCAACAATAAAT (41)
C2ASR	CATTAGCAACATGTTTGGACAAC (42)
Curcin 1 TOPO-F	CACCCCATTACTTATGACGCTACTAC (43)
JCG-R	GTAGGATTAAAGCCATGGCAGC (44)
Gus F	GTCAGTGGCAGTGAAGGGCGAAC (45)
Gus R	TTCCATACCTGTTACCGACGAC (46)
Gus RT F1	GTGGCAGTGAAGGGCGAAC (47)
Gus RT R1	AGGTACGGTAGGAGTTGGC (48)
C1SF	TGCAGTCAATTACAAAGTCTG (49)
C2ASF	AGTAAAAGGTCTCATGGGAGTC (50)
Jc actin F2	TAATGGTCCCTCTGGATGTG (51)
Jc actin R1	AGAAAAGAAAAGAAAAAAGCAGC (52)

[0069] *Jatropha transformation:* Cotyledons from 5-7 days old seedling were excised into 0.5×0.5 cm² discs and co-cultivated with agrobacteria (OD₅₉₅ = 0.25-0.35) suspended *Agrobacterium* suspension medium (liquid MS salts, B5 vitamins, 1.5 mg/L BA (6-benzylaminopurine), 0.1 mg/L NAA (α -naphthalene-acetic acid), 20 mg/L AS (Acetosyringone), 0.5 g/L MES (2-(N-morpholino)ethanesulfonic acid), 30 g/L sucrose, 10 g/L glucose, pH5.0-5.2) for 2-3 days at 22E C in the dark. The co-cultivated cotyledon discs were then rinsed several times with sterile water, following one wash with 300 mg/L cefotaxime. The cotyledon discs were then cultured on callus formation medium (MS salts, B5 vitamins, 1.5 mg/L 6-BA, 0.05 mg/L NAA, 3.5 mg/L hygromycin, 100 mg/L cefotaxime, 10 mg/L citric acid, 150 mg/L glutamine, 100 mg/L casein hydrolysate, 0.5 g/L MgCl₂, 30 g/L sucrose, 2.5 g/L phytigel, pH 5.8-6.0) at 25E C in darkness for 3 weeks. The newly emerged hygromycin-resistant calli were subcultured on shoot regeneration medium I (MS salts, B5 vitamins, 10 mg/L citric acid, 150 mg/L glutamine, 100 mg/L casein hydrolysate, 0.5 g/L MgCl₂, 30 g/L sucrose, 1.5 mg/L 6-BA, 0.05 mg/L IBA (indole-3-butyric acid), 2 mg/L adenine, 3.5 mg/L Hygromycin, 100 mg/L cefotaxime, 2.5 g/L phytigel, pH 5.8-6.0) at 25E C in 16h light/8h darkness photoperiod for 3 weeks. During this period, any shoots regenerated from calli were required to be subcultured on shoot regeneration medium II (MS salts, B5 vitamins, 10 mg/L citric acid, 150 mg/L glutamine, 100 mg/L casein hydrolysate, 30 g/L sucrose, pH 5.8-6.0, 1.5 mg /L 6-BA, 0.05 mg/L IBA, 0.5 mg/L GA (gibberellic acid), 4 mg/L hygromycin, 100 mg/L cefotaxime, 7 g/L agar). Calli with no regenerated shoots were

subcultured on shoot regeneration medium III (MS salts, B5 vitamins, 10 mg/L citric acid, 150 mg/L glutamine, 100 mg/L casein hydrolysate, 0.5 g/L MgCl_2 , 30 g/L sucrose, 1.5 mg/L 6-BA, 0.05 mg/L IBA, 3.5 mg/L hygromycin, 100 mg/L cefotaxime; 2.5 g/L phytigel, pH 5.8-6.0) for further regeneration of shoots. The regenerated shoots were subcultured on shoot elongation medium (MS salts, B5 vitamins, 10 mg/L citric acid, 150 mg/L glutamine, 100 mg/L casein hydrolysate, 30 g/L sucrose, 0.3 mg/L 6-BA, 0.1 mg/L GA, 7 g/L Agar, pH 5.8-6.0) at 25°C in 16h light/8h darkness photoperiod for 3 weeks. The elongated shoots at about 2.5 cm in length were rooted in root induction medium (MS salts, B5 vitamin, 0.07 mg/L IBA, 0.5 g/L MES, 30 g/L sucrose, 2.2 g/L phytigel, pH 5.8). Normally it takes more than one month to get roots.

[0070] *BAC library construction:* BAC library of *J. curcas* was made with BAC vector pIndigoBAC-5 as described by Peterson et al (2000). Briefly, the High Molecular Weight (HMW) DNA was isolated from the fresh leaves according to the method OPTION X described in the protocol. The HMW DNA was digested by *Hind*III and size-selected 100-150-kb fragments were ligated into pIndigoBAC-5 Vector. The product of ligation was transferred into *E. coli* DH10B competent cells and the transformants were arrayed in the 384-well plates. The high density hybridization membrane was prepared and used for BAC clone screening.

[0071] *Southern and northern blot analysis:* DNA gel blot analysis was carried out according to the standard procedures as previously described (Sambrook et al., 1989). Plant genomic DNA was isolated from leaves as described previously (Dellaporta et al. 1983). About 2 µg DNA was used for each lane in southern analysis. *Curcin 1* coding sequence amplified with primer pairs CantigenF1 and CantigenR1 (Table 1) was used as probes for BAC library screening and Southern blot analyses. Probes were labelled with [^{32}P]-dCTP by Rediprime II random prime labelling system (Amersham Biosciences, Piscataway, NJ, USA).

[0072] *Reverse transcription polymerase chain reaction (RT-PCR):* Total RNA was isolated from different tissues using RNeasy Plant Mini Kit (Qiagen, Hilden, Germany) according to manufacturer's instruction. The total RNA samples were treated with DNase I. First-strand cDNA was synthesized from 1 µg of total RNA using SMARTTM MMLV Reverse Transcriptase (Clontech) according to the manufacturer's instructions.

[0073] *Rapid amplification of cDNA ends (RACE) and thermal asymmetric interlaced (TAIL) PCR:* 5' RACE and 3' RACE were carried out according to the manufacturer's instruction (Invitrogen, Carlsbad, CA, USA). The primers for 5' RACE of *Curcin 1* were primer pairs GSP1 and AAP, and primer pairs GSP1A and AUAP (Table 1). The primers for

3'RACE of *Curcin 1* were primer pairs C2RTF2 and AUAP, and primer pairs 3'RACEF4 and AUAP (Table 1). The primers for 5' RACE of *Curcin 2A* were primer pairs C2RTR2 and AAP, and primer pairs 5'RACEG4 and AUAP (Table 1). The primers for 3'RACE of *Curcin 2A* were primer pairs CTRTF2 and AUAP, and primer pairs 3'RACEF1 and AUAP (Table 1). TAIL-PCR was performed according to the procedures published previously (Liu et al. 1995). The arbitrary degenerate (AD) primer used for TAIL-PCR was AD2 (Table 1).

[0074] *Real-time RT-PCR*: Quantitative real-time RT-PCR was followed the methods described by Chen et al (2007) and conducted on a Bio-Rad iCycler iQ5 real-time PCR system. Total RNA was extracted from jatropha seeds or leaves at developmental stages. The total RNA samples were first treated with DNase I, and then reverse transcribed to first-strand cDNA using iScript cDNA synthesis kit (Bio-Rad) according to the manufacturer's instructions. Oligo primers for real-time RT-PCR were designed based on the 3'UTR of each gene. To ensure maximum specificity and efficiency during quantitative PCR, oligo primers were further tested for linearity of response by constructing standard curves on five or six serial ten-fold dilutions. A standard reaction mixture (15 µl) consisted of 2 µl cDNA template, 1 x SsoFast EvaGreen supermix and 500 nM forward and reverse primers. The PCR reaction consisted of an initial denaturizing step of 95E C for 30 sec, followed by 40 cycles of 95E C for 5 sec, 60E C for 10 sec. A melting-curve reaction immediately followed the amplification with heating for 10 sec, starting at 65E C with 0.5E C increments. PCR product specificity was confirmed by melting-curve analysis and electrophoresis on 2% agarose gel to ensure that PCR reactions were free of primer dimmers and non-specific amplicons. The jatropha actin gene was used as internal reference to normalize the relative amount of total RNA for all samples. For each selected gene, triplicate sets of PCR reactions, including the actin controls and duplicate negative controls (reaction samples without cDNA templates), were prepared and run in a 96-well plate. The real-time RT-PCR experiments were repeated for each plate to ensure that similar results could be obtained.

[0075] *Particle bombardment and transient expression of GFP or GUS*: 1.0-µm gold particles were coated with plasmid DNA as directed by the manufacturer's instructions (Bio-Rad, CA, USA). Endosperm from immature jatropha seeds at six-week-old developmental stage were manually cut into thin slices of about 0.5-mm thickness. For each bombardment, five to six pieces of slices were bombarded at 1100 psi with a flight distance of 15 cm using a PDS-1000/He system (Bio-Rad). The bombarded samples were cultured overnight at 22-25°C

on wet filter paper in dishes. The bombarded samples were screened with a fluorescence stereomicroscope (Model SZX12, Olympus, Japan) for GFP expression and then imaged on LSM510 META inverted confocal microscope (Zeiss, Jena, Germany) at 488 nm with a band pass of 505-530 nm. The maize ubiquitin promoter and its fusion GFP gene in pSSZ41 (Kolesnik et al., 2004) were used as positive control for GFP assay. For transient GUS expression, leaf discs of 2x2 cm² in size were used for particle bombardment. GUS activity was detected according to the method as described previously (Hiei et al, 1994). X-Gluc staining of GUS activity was observed with a Nikon SMZ1500 stereo microscope and photographed with an attached Nikon DIGITAL SIGHT DS-SMC camera. The CaMV35S promoter with its fusion gene GUS in pC1305.1 was used as the positive control for GUS activity assay. For both GFP and GUS transient expression assays, empty vector pC1300 was used as the negative control.

[0076] Generation of curcin antibodies: The N- and C-terminal coding sequences of *Curcin 1* were amplified with primer pairs Cantigen F1 and Cantigen R2 for Curcin 1N, and primer pairs Cantigen F2 and Cantigen R3 for Curcin 1C, respectively (Table 1). The PCR products were cloned into vector pQE30 (Qiagen) to generate pQCF1R2 and pQCF2R3. The constructs were then introduced into *E. coli* BL21 strain for antigen expression. The antigen proteins were extracted and purified by polyhistidine-tagged protein purification kit (MACHEREY-NAGEL, Duren, Germany). The anti-Curcin 1N or anti-Curcin 1C antibodies were raised following standard procedure for polyclonal antibody production for rabbits.

[0077] Western blot analysis: Western blot analysis was carried out with 20 µg total proteins from different tissues of plants. Proteins were separated on 12% SDS polyacrylamide gels followed by blotting onto nitrocellulose membrane. Curcin proteins were detected with anti-Curcin 1N or anti-Curcin 1C antibody and a horseradish peroxidase-coupled secondary antibody (Bio-Rad) according to the product manual.

[0078] Immunogold electron microscopy: Immunogold electron microscopy was carried out according to the procedure described previously (Chye et al., 1999). Seed or leaf sections were fixed in a solution of 0.5% glutaraldehyde and 2% paraformaldehyde in 0.1 M phosphate buffer for 4 h under vacuum. Specimens were washed with 50 mM phosphate buffer for 45 min. After dehydration in a graded ethanol series, specimens were infiltrated in LR white resin (EMS, Hatfield, PA19440, USA) and embedded in gelatin capsules. Specimens were sectioned at 80 nm using a Leica Ultracut microtome and mounted on Formvar-coated slotted grids. Curcin proteins were detected with anti-curcin polyclonal

antibodies followed by labelling with 15 nm gold-conjugated goat anti-rabbit IgG antibody (EMS). Rabbit pre-immune serum was used in control experiments. Samples on grids were further stained with 2% uranyl acetate and 1% lead citrate. Samples were visualized with a transmission electron microscope (JEOL JEM-1230, JEO LTD, Tokyo 196-8558, Japan) operating at 120kv and photographed with a digital microphotography system (Gatan Inc., Pleasanton, CA94588, USA).

EXAMPLE 2

Isolation of *Curcin 1* and *Curcin 2*

[0079] To isolate curcin genes from *J. curcas*, a jatropha BAC library was constructed and seven BAC clones were identified from the BAC library to contain one or two members of curcin genes (Figure 1). BAC clone 121E10 carried two copies of curcin genes (Figure 1). BAC 121E10 was sequenced by shot-gun approach and the insert in the BAC was revealed to be 64,841 bp. The two curcin genes in the BAC insert were closely linked and arrayed in tandem manner. One curcin gene encodes a 293-aa curcin protein while another encodes a 309-aa curcin 2-like protein. The two curcin genes were thereafter designated as *Curcin 1* for the 293-aa curcin and *Curcin 2* for the 309-aa curcin, respectively. *Curcin 1* and *Curcin 2* are 8-kb from each other and *Curcin 1* locates at the 3' of *Curcin 2*. The curcin genes in the remaining six BAC clones were subcloned and sequenced. The curcin genes in BAC clones 127F22, 147I10 and 176I18 harboured both *Curcin 1* and *Curcin 2*, whereas the curcin genes in 213H9 and 221F16 carried *Curcin 1*, respectively. BAC clone 187G09 also carried *Curcin 1*, but its sequence showed mutation at -2463 bp downstream of the *Curcin 1* coding sequence.

[0080] We performed cDNA library screening and RT-PCR with 5'RACE and 3'RACE to isolate the full-length cDNA clones of the *Curcin 1* and *Curcin 2* genes. A cDNA corresponding to *Curcin 1* was isolated from the endosperm of immature jatropha seeds (Figure 2). *Curcin 1* cDNA consists of 1388 nucleotides excluding 3' poly(A) sequence (Figure 2). It contains a 66-bp 5' untranslated region (5'UTR) and a 213-bp 3'UTR (Figure 2). Comparing *Curcin 1* cDNA with its genomic sequence, a 227-bp intron was identified in 5'UTR (Figure 2). The intron is located at 12 bp upstream of start codon (Figure 2). There are two canonical polyadenylation signals (AATAAA) in the 3'-untranslated region of *Curcin 1* gene, which is typical for RIPs genes (Lin et al., 2003; Chow et al., 1999). The first polyadenylation signal locates at 80 bp downstream of the stop codon while the second one

resides at 20 bp upstream of the polyadenylation site (Figure 2). Although the genomic sequence of *Curcin 2* shows high identity to that *Curcin 1*, we failed to isolate cDNA for *Curcin 2* from either immature seeds or leaf tissues with the above-mentioned technologies. We reasoned that *Curcin 2* gene might not express under normal developmental condition. Figure 3 shows the annotation of *Curcin 2* with its genomic sequence, putative intron in the 5'UTR and its deduced amino acid residues.

EXAMPLE 3

Isolation of *Curcin 2A*

[0081] During our attempt to isolate *Curcin 2* cDNA from jatropha, we obtained a cDNA clone that showed homology to that of *Curcin 2*. There are fifty-nine single nucleotide polymorphisms (SNP) in the ORF regions between the cDNA and *Curcin 2*. The open reading frame (ORF) of the cDNA encodes a curcin protein with 309 amino acid residues which is identical to a type 2 curcin protein in GenBank (Acc. No.: ABZ04128). We designated this curcin gene as *Curcin 2A*. The genomic clone of *Curcin 2A* was isolated by PCR. Initially, a 1216-bp genomic region of *Curcin 2A* was isolated by PCR with primers C2PF4 derived from *Curcin 2* genomic clone and C1CDSR from *Curcin 2A* cDNA (Table 1). This 1216-bp fragment contains 881-bp *Curcin 2A* promoter. Secondly, a 982-bp fragment corresponding to the 5' regulation region of the *Curcin 2A* promoter at the upstream of the 881-bp region was isolated by TAIL-PCR with primer pairs C2AFAR3 and AD2 (Table 1). An 832-bp 3' genomic region of *Curcin 2A* was amplified with C2ACDSF derived from the *Curcin 2A* cDNA and C2TR2EcoRI based on the *Curcin 2* genomic sequence (Table 1). Totally, we have obtained a 3774-bp genomic clone of *Curcin 2A* including a 1793-bp promoter at the upstream of transcription initiation site and 758-bp 3' regulation region at the downstream of the stop codon. The gene organization of *Curcin 2A* is similar to those of *Curcin 1*. Figure 4 shows the annotation of *Curcin 2A* with its genomic sequence, intron in the 5'UTR and its deduced amino acid residues.

[0082] So far, we have isolated three curcin genes from *J. curcas*. Previously, we have obtained seven BAC clones from jatropha BAC library. However, DNA sequencing indicated that none of the seven BAC clones harbors *Curcin 2A*. BAC clones 121E10, 127F22, 147I10 and 176I18 carry *Curcin 1* and *Curcin 2*, BAC clone 187G09 contains *Curcin 1*, and BAC clones 213H9 and 221F16 harbor *Curcin 2*. To further verify the presence of *Curcin 2A* in the jatropha genome, we designated a pair of universal primers CF2 and CR2 (Table 1) to

amplify the three curcin genes from genomic DNA as well as *Curcin 1* and *Curcin 2* from BAC 121E10. The PCR products were digested with *Sau3AI* and separated on 2% agarose gel. Both *Curcin 1* and *Curcin 2* generate three fragments (195 bp, 243 bp and 582 bp for *Curcin 1*; 195 bp, 243 bp and 580 bp for *Curcin 2*), whereas *Curcin 2A* produces two fragments (243 bp and 775 bp) due to only one *Sau3AI* site present in the coding sequence of the gene (Figure 4). As shown in Figure 5, the PCR products amplified from BAC 121E10 gave three bands after digestion with *Sau3AI* (the 582-bp and the 580-bp fragments could not be separated on 1.2% agarose gel), while the products from jatropha genome had four bands including an additional 775-bp bands from *Curcin 2A*. The results indicate the presence of *Curcin1*, *Curcin 2* and *Curcin 2A* in jatropha genome.

EXAMPLE 4

Curcin proteins

[0083] The curcin proteins have been identified with Curcin 2A identical to ABZ04128. Total together, there are seven different kinds of curcin proteins (Figure 6). The first 42 amino acid residues of curcin AAL58089 encodes a signal peptide, which is cleaved in mature protein (Lin et al., 2003a). This N-terminal 42aa signal peptide is conserved among all curcin proteins (Figure 6). Among these curcin proteins, Curcin 1 identified in this study is more closed to AAL58089 and both belong to type 1 curcins with 293 AA (Figure 7). Curcin 2A (ABZ04128), AAR083395 and ABW17545 belong to type 2 curcins with 309 AA. These three type 2 curcins are closely related to each other in evolution (Figure 7). Interestingly, Curcin 2 is more closely related to type 1 curcins, such as Curcin 1 and AAL58089, than type 2 curcins (Figure 7).

EXAMPLE 5

Expression of *Curcin 1* and *Curcin 2A* in *Jatropha curcas*

[0084] Previous RT-PCR analysis indicated that *Curcin 2* did not express in jatropha seeds and leaves under normal developmental conditions. We then determined the expression of *Curcin 1* or *Curcin 2A* in the two tissues by RT-PCR with a common forward primer CF and either one of the two specific reverse primers C1SR for *Curcin 1* and C2ASR for *Curcin 2A* (Table 1). Primer CF covers the splicing junction region between exon 1 and exon 2 of *Curcin 1* or *Curcin 2A*. As shown in Figure 8A, the *Curcin 1* transcripts were detected in endosperm. On the contrary, the *Curcin 2A* transcripts were only present in leaves. We

further determined the temporal expression of *Curcin 1* in the endosperm of developing seeds (Figure 8B) as well as in leaves at different developmental stages or infected by insects (Figure 8C). The *Curcin 1* transcripts were detected at high level at 6 weeks after pollination (WAP) (Figure 8B). No transcript or very low level of *Curcin 1* transcript was detected at 2, 4 and 6 WAP (Figure 8B). Curcin 2A transcripts were only detected in young leaves (YL) but not in full-expanded leaves (FL), old leaves (OL) leaves infected with mealybuds (*Pseudococcidae hirsutus*) (Figure 8C).

EXAMPLE 6

Detection of curcin proteins in *Jatropha curcas*

[0085] Polyclonal antibodies against C-terminal region of Curcin 1 were raised. Western blot analysis indicated that Curcin 1 proteins were only detected in endosperm whereas Curcin 2A proteins were only detected in leaves (Figure 9). The detected Curcin 1 proteins should be the mature Curcin 1 proteins with the molecular size of about 28 kDa (MW = 27.8 kD) (Figure 9) (Lin et al., 2003a, 2003b). However, the detected Curcin 2A proteins had molecular size of about 34 to 35 kDa and they seemed to be the intact Curcin 2A proteins or precursors of Curcin 2A (MW = 34.9 kD) (Figure 9). It should be mentioned that two unknown proteins were detected by anti-Curcin 1 antibodies. One protein had molecular size of about 39 kDa and was detected in leaves, calli, seeds but not in roots (Figure 9). Another protein had molecular size of about 16 kDa and was detected in seeds (Figure 9).

EXAMPLE 7

Subcellular localization of Curcin 1 and Curcin 2A

[0086] Subcellular localization of curcin proteins was determined by immunogold electron microscopy. As *Curcin 1* is only expressed in endosperm at late developmental stage, we took 6-week-old endosperm tissues for subcellular immunogold localization assay of the Curcin 1 proteins. Oil bodies and protein bodies are two major organelles in 6-week-old endosperm cells (Figure 10A). A few plastids with starch bodies were also observed in endosperm cells at this stage (Figure 10A). The Curcin 1 proteins, which are indicated by gold particles in immunogold electron microscopy, localized to protein bodies only (Figure 10C). The cell walls between endosperm cells are primary cell walls, where no Curcin 1 protein was detected or only background labeling was detected (Figure 10D). No immunolabelling of protein body was detected when preimmune antibody control was used (Figure 10B). As the anti-Curcin 1

antibody also recognized a 16-kDa non-specific protein in jatropha seeds (Figure 9), the signal detected by immunogold electron microscopy in this study should include this non-specific recognition.

[0087] Although the anti-Curcin 1 antibody recognized both Curcin 2A and a 39-kDa non-specific protein from jatropha leaves (Figure 9), it was used to detect the subcellular localization of the two proteins in jatropha leaves by immunogold electron microscopy assay. Immunogold electron microscopy assay indicated that gold particles were immunolocalized to vacuolar contents of leaf mesophyll cells (Figure 11D) and secondary cell walls of leaf tracheary elements (Figure 12C and 12D). The localization of either the Curcin 2A proteins or the 39-kDa non-specific proteins or both to secondary cell walls of tracheary elements takes place in immature tracheary elements where the cells undergo program cell death (Figure 12A and 12C). In control experiments, no immunolabelling or only background signal was detected in vacuolar contents of leaf mesophyll cells (Figure 11A and 11B) or secondary cell walls of leaf tracheary elements (Figure 12B) when preimmune serum was used.

EXAMPLE 8

Characterization of *Curcin 1* and *Curcin 2A* Promoters

[0088] *Curcin 1* is specifically expressed in endosperm at late developmental stages. We performed PLACE analysis (<http://colon.backslash.backslash.www.dot.dna.dot.affrc.dot.go.dot.jp/backslash/PLACE/backslash/signalscan.dot.html>) for plant *cis*-acting regulation DNA elements in *Curcin 1* promoter. DNA elements that are involved in endosperm-specific or storage protein gene expression were identified. A DNA element called "RY repeat (CATGCAY)" or legumin box was identified at -1388 bp position (the transcription initiation site is designated at +1 position). The legumin box was found in seed-storage protein genes in legume such as soybean (*Glycine max*) (Fujiwara and Beachy, 1994). Another DNA element termed AACAA element was identified at -1534 bp. The core of AACAA element were found in rice (*Oryza sativa*) glutelin genes and involved in controlling the endosperm-specific expression (Wu et al., 2000). At upstream of the AACAA element, there are several E-box elements locate at from -2649 bp and -2273 bp. One of the E-box located at -2649 bp was also found in *napA* storage-protein gene of rapeseed (*Brassica napus*) (Stalberg et al., 1996). *Curcin 1* promoters containing all these *cis*-elements (C1P3 and C1P4) were sufficient to drive endosperm-specific gene expression (Table 2, Figure 13A). Deletion of E-box elements (C1P2) or E-box elements, legumin box and AACAA element (C1P1) attenuated or abolished

endosperm-specific gene expression (Table 2). No activity of *Curcin 1* promoters was detected in leaf tissues (Table 2).

TABLE 2
Characterization of *Curcin* Gene Promoters

Promoter ^a cells	Size of promoter ^b	Reporter	Tissue tested	Positive sections/ total sections bombarded	Positive or spots/ positive section ^c
Empty vector -		-	Endosperm	0/80	0
Empty vector -		-	Leaf	0/25	0
Ubiquitin	N.A.	GFP	Endosperm	64/81	10-20 cells
CaMV35S	N.A.	GUS	Leaf	5/25 ^d	5-20 spots
C1P1	-200 to +293	GFP	Endosperm	0/80	0
C1P2	-1746 to +293	GFP	Endosperm	5/80	1-3 cells
C1P3	-2888 to +293	GFP	Endosperm	42/80	5-10 cells
C1P4	-4956 to +293	GFP	Endosperm	41/75	5-10 cells
C1P1	-200 to +293	GUS	Leaf	0/25	0
C1P2	-1746 to +293	GUS	Leaf	0/25	0
C1P3	-2888 to +293	GUS	Leaf	0/25	0
C1P4	-4956 to +293	GUS	Leaf	0/25	0
C1P1D	-200 to +293 (Δ Intron)	GFP	Endosperm	2/60	1-3 cells
C1P2D	-1746 to +293 (Δ Intron)	GFP	Endosperm	38/80	5-10 cells
C1P3D	-2888 to +293 (Δ Intron)	GFP	Endosperm	41/83	5-10 cells
C1P4D	-4956 to +293 (Δ Intron)	GFP	Endosperm	43/86	5-10 cells
C1P1D	-200 to +293 (Δ Intron)	GUS	Leaf	0/25	0
C1P2D	-1746 to +293 (Δ Intron)	GUS	Leaf	0/25	0
C1P3D	-2888 to +293 (Δ Intron)	GUS	Leaf	0/25	0
C1P4D	-4956 to +293 (Δ Intron)	GUS	Leaf	0/25	0
C2AP1	-42 to +294	GFP	Endosperm	0/80	0
C2AP2	-881 to +294	GFP	Endosperm	20/80	3-5 cells
C2AP3	-1793 to +294	GFP	Endosperm	0/80	0
C2AP1	-42 to +294	GUS	Leaf	0/25	0
C2AP2	-881 to +294	GUS	Leaf	0/25	0
C2AP3	-1793 to +294	GUS	Leaf	5/25	2-10 spots
C2AP1D	-42 to +294 (Δ Intron)	GFP	Endosperm	3/80	1-3 cells
C2AP2D	-881 to +294 (Δ Intron)	GFP	Endosperm	4/75	1-3 cells
C2AP3D	-1793 to +294 (Δ Intron)	GFP	Endosperm	1/80	4 cells
C2AP1D	-42 to +294 (Δ Intron)	GUS	Leaf	0/25	0
C2AP2D	-881 to +294 (Δ Intron)	GUS	Leaf	0/25	0
C2AP3D	-1793 to +294 (Δ Intron)	GUS	Leaf	0/25	0
C2P1	-326 to +258	GFP	Endosperm	0/80	0
C2P2	-1981 to +258	GFP	Endosperm	0/80	0
C2P3	-3207 to +258	GFP	Endosperm	0/80	0
C2P1	-326 to +258	GUS	Leaf	0/25	0
C2P2	-1981 to +258	GUS	Leaf	0/25	0

C2P3	-3207 to +258	GUS	Leaf	0/25	0
------	---------------	-----	------	------	---

^aEmpty vector pC1300 was used as negative control. The CaMV35S promoter in pC1305.1 and the ubiquitin promoter in pSSZ41 (Kolesnik et al., 2004) were used as positive controls for endosperm and leaf tissues, respectively.

^bTranscription initiation site of each promoter is indicated as +1. "ΔIntron" indicates the intron at the 5'UTR of the curcin gene has been deleted.

^cFor transient GFP assay, usually individual cells showing transient GFP expression were observed. For transient GUS expression assay, we only count blue spots. One blue spot is regarded as one successful delivery case.

^dFor each bombardment, one piece of leaf disk at size of 2x2 cm² was used. Twenty-five leaf disks were bombarded for GUS transient assay. Even for positive control bombarded with pC1305.1, not all bombarded leaf disks gave blue spot for GUS activity.

[0089] *Curcin 2A* is specifically expressed in jatropha leaves. Totally, a *Curcin 2A* promoter of 1793 bp was isolated from *J. curcas*. A PLACE survey with the 1793-bp fragment indicated that several light-regulated *cis*-elements (Terzaghi and Cashmore, 1995) present in the *Curcin 1* promoter, which meets its function as a specific promoter with activity in green tissues like leaves. These light-regulated *cis*-elements include the GT-1 binding sites (Villain et al., 1996) at positions -64, -548, -590, -712, -768, -861, -916, -1315, -1316, -1329, -1411, -1539, -1683 and -1703, I boxes (Giuliano et al., 1988) at positions -548, -861, -916 and -1533, T boxes (Chan et al., 2001) at position -454 and -1581, and GATA boxes (Lam and Chua, 1989) at positions -548, -861, -916, -1093, -1533 and -1747. Although not all these putative *cis*-element may be functional, the presence of these elements in the *Curcin 2A* promoter supports its function as leaf- or green-tissue-specific promoter. The PLCAE survey also identified several stress-inducible *cis*-elements at *Curcin 2A* promoter. These stress-inducible *cis*-elements include NPR1-type W boxes (Yu et al., 2001) at positions -54 and -161, ERF3-type W boxes (Nishiuchi et al., 2004) at position at -702, TGAC-containing W-box (Eulgem et al., 1999) at positions -160, -261, -702 and -1453 and MYB binding sites (Urao et al., 1993) at positions -157 and -294. The presence of these putative stress-inducible *cis*-elements in the *Curcin 2A* promoter suggests that the *Curcin 2A* gene might be involved in plant defense. Promoter deletion studies indicated that the light-regulated and stress-inducible *cis*-elements at the *Curcin 2A* promoter between position -811 and position -1793 (C2AP3 in Table 2) is required for its leaf-specific activity (Table 2, Figure 13B). The *Curcin 2A* promoter carrying *cis*-elements at upstream to position -811 (C2AP2) still had activity but lost

its specificity as a leaf-specific promoter since it only had low level of activity in endosperm but not in leaf tissues (Table 2).

[0090] *Curcin 2* is not expressed in leaves and endosperm of *J. curcas* under normal developmental condition as no *Curcin 2* transcript was detected in these tissues. Transient GFP or GUS assay in endosperm or leaf tissues also did not detect any activity of the *Curcin 2* promoter in these tissues (Table 2). DNA alignment with the 3-kb DNA sequences of *Curcin 1* and *Curcin 2* promoters indicated that the two promoters just show homologous from positions -1 to -275 in *Curcin 2* promoter (or positions -1 to -286 in the *Curcin 1* promoter). The homologous sequences between *Curcin 2* and *Curcin 2A* promoters start from positions -1 to -1229 in the *Curcin 2* promoter (or positions -1 to -1345 in the *Curcin 2A* promoter). Although PLACE analysis with *Curcin 2* promoter identified many *cis*-elements, two of them which are unique to *Curcin 2* promoter are T/G-box (Boter *et al.*, 2004) at position -269 and ethylene responsive element (ERE) at position -782 (Tapia *et al.*, 2005). T/G-boxes, found in tomato proteinase inhibitor II (*pin2*) and leucine aminopeptidase (*LAP*) genes, are involved in jasmonate (JA) induction of these genes (Boter *et al.*, 2004), whereas the ERE motifs were found to mediate ethylene-induced activation of the U3 promoter region in the 5'-LTR region of TLC1.1 retrotransposon family in *Lycopersicon chilense* (Tapia *et al.*, 2005). The presence of T/G-box and ERE motif in the *Curcin 2* promoter suggests that the gene might be activated by JA upon attack by insects (Howe and Jander, 2008) and/or ethylene during defense reaction or leaf senescence (van Loon, *et al.*, 2006).

[0091] Both *Curcin 1* and *Curcin 2A* have introns in their 5'UTR with lengths of 227 bp and 228 bp, respectively (Figures 2 and 4). *Curcin 2* might also have the 227-bp intron (Figure 3). Deletion of the intron in the *Curcin 1* promoters with the length up to position -2888 (C1P3D and C1P4D) did not affect their activity and specificity (Table 2). However, deletion of the intron in the *Curcin 1* promoters with the lengths up to positions -200 (C1P1D) or -1746 (C1P2D) significantly increased their activity and the mutated promoters still retained specificity for endosperm (Table 2). The results indicate that the intron in the 5'UTR of *Curcin1* suppress its expression. Deletion of the intron in 5'UTR of *Curcin 2A* promoter with length up to positions -42 (C2AP1D) made the promoter have detectable activity in endosperm, whereas the same length of wild-type promoter (C2AP1) did not show any activity even in leaf tissues. Deletion of the intron in 5'UTR of *Curcin 2A* promoter C2AP2 (C2AP2D) did not significantly change its activity, which was only detected in endosperm (Table 2). Finally, deletion of the intron in 5'UTR of *Curcin 2A* promoter C2AP3 (C2AP3D)

changed its specificity and the activity was detected in endosperm rather than in leaf tissues (Table 3). The deletion studies with the *Curcin 2A* promoters suggest the intron in the 5'UTR of *Curcin 2A* affect both its activity and its specificity in leaf tissues.

EXAMPLE 9

Generation of Curcin-Deficient Transgenic Jatropha Plants

[0092] RNAi strategy was utilized to knock-down curcin gene expression in transgenic jatropha plants and produce curcin-deficient jatropha plants. Binary construct pANDA35HKC1 contains an RNAi cassette in its T-DNA region (Figure 14). The expression of the RNAi cassette generates double-strand RNA (dsRNA) of the *Curcin 1* gene. pANDA35HKC1 was used for jatropha transformation. Initially, ten independent T₀ transgenic plants (L3, L4, L11, L17, L19, L22, L23, L24, L26 and L46) were obtained after *Agrobacterium*-mediated transformation. After transplantation, only 4 plants (L17, L23, L26 and L46) were survived in greenhouse (Figure 15).

[0093] Southern blot analysis using *Gus* linker probe indicated that all of the 4 plants carried intact RNAi cassette (Figure 16, data not shown for L46). Southern blot analysis using *Hpt* probe showed that they all carried single copy of T-DNA (Figure 16, data not shown for L46).

[0094] The expression of the RNAi cassette and the *Curcin 2A* in the young leaves of the transgenic plants was detected by real-time PCR. Real-time PCR analysis with the *Gus* linker region indicated that the RNAi cassette was expressed in young leaves of the 4 transgenic plants (Figure 17A). Real-time PCR analysis showed that the *Curcin 2A* transcripts in young leaves of L17, L26 and L46 were 20% or less than the total *Curcin 2A* transcripts in the wild-type plants (Figure 17B). Although the expression of the RNAi cassette in L23 was comparable to that in other transgenic plants, the *Curcin 2A* transcripts in young leaves of L23 was higher than that in L17, L26 and L46, which was similar to that of wild-type plants.

[0095] The suppression of *Curcin 2A* gene in young leaves of the transgenic plants was further confirmed by western blot analysis using anti-Curcin 1 antibodies. In wild-type plants, Curcin 2A was only detected in young leaves but not in full-expanded leaves (Figure 18). This result was consistent with the fact that *Curcin 2A* transcripts were only detected in young leaves of wild-type plants (Figure 8). Curcin 2A was not or hardly detected in transgenic plants L17, L26 and L46 (Figure 18). Curcin 2A was detected in L23 but its expression level was lower than that of wild-type plants (Figure 18). The results indicate that the expression of

Curcin 2A in the transgenic plants was either partially or completely suppressed by the dsRNA derived from the expression of the RNAi cassette.

[0096] To further check the transmission of the T-DNA in the transgenic *jatropha* plants, we germinated T₁ seeds of L26 as well as seeds from wild-type plants on selection medium containing 10 mg/L hygromycin. The wild-type seeds were able to germinate on the selection medium, but the young seedlings died within one week (Figure 19). Fifty T₁ seeds of L26 were germinated on the selection medium and thirty-six seedlings were resistant to hygromycin (Figure 19). L26 carried one copy of T-DNA and the segregation of hygromycin-resistant and hygromycin-susceptible follow the Mendelian ratio for single gene inheritance ($\chi^2=0.240$; $0.5 < P < 0.8$; $df=1$). This result indicates that the T-DNA region in pANDA35HKC1 has stably integrated into *jatropha* genome and was transmitted to the progeny of the transgenic plants.

[0097] In conclusion, we have characterized the promoters of the three curcin genes. The wild-type *Curcin 1* promoter with length up to position -2888 is sufficient for its high and specific activity in endosperm, whereas the isolated *Curcin 2A* promoter with length up to position -1793 has specific activity in leaf tissues. The curcin-deficient transgenic *jatropha* plants were produced by using RNAi technology to suppress curcin gene expression. The curcin-deficient *jatropha* transgenic plants can be used for future breeding program.

[0098] The use of the terms “a” and “an” and “the” and similar referents in the context of describing the invention (especially in the context of the following claims) are to be construed to cover both the singular and the plural, unless otherwise indicated herein or clearly contradicted by context. The terms “comprising,” “having,” “including,” and “containing” are to be construed as open-ended terms (i.e., meaning “including, but not limited to,”) unless otherwise noted. Recitation of ranges of values herein are merely intended to serve as a shorthand method of referring individually to each separate value falling within the range, unless otherwise indicated herein, and each separate value is incorporated into the specification as if it were individually recited herein. For example, if the range 10-15 is disclosed, then 11, 12, 13, and 14 are also disclosed. All methods described herein can be performed in any suitable order unless otherwise indicated herein or otherwise clearly contradicted by context. The use of any and all examples, or exemplary language (e.g., “such as”) provided herein, is intended merely to better illuminate the invention and does not pose a

limitation on the scope of the invention unless otherwise claimed. No language in the specification should be construed as indicating any non-claimed element as essential to the practice of the invention.

[0099] It will be appreciated that the methods and compositions of the instant invention can be incorporated in the form of a variety of embodiments, only a few of which are disclosed herein. Embodiments of this invention are described herein, including the best mode known to the inventors for carrying out the invention. Variations of those embodiments may become apparent to those of ordinary skill in the art upon reading the foregoing description. The inventors expect skilled artisans to employ such variations as appropriate, and the inventors intend for the invention to be practiced otherwise than as specifically described herein. Accordingly, this invention includes all modifications and equivalents of the subject matter recited in the claims appended hereto as permitted by applicable law. Moreover, any combination of the above-described elements in all possible variations thereof is encompassed by the invention unless otherwise indicated herein or otherwise clearly contradicted by context.

BIBLIOGRAPHY

- [0100]** Augustus, GDPS, Jayabalan M and Seiler GJ. 2002. Evaluation and bioinduction of energy components of *Jatropha curcas*. Biomass and Bioenergy. 23:161-164.
- [0101]** Azam MM, Waris A and Nahar NM. 2005. Prospects and potential of fatty acid methyl esters of some non-traditional seed oils for use as biodiesel in India. Biomass and Bioenergy. 29:293-302.
- [0102]** Barbieri L, Battelli M and Stirpe F. 1993. Ribosome-inactivating protein from plants. Biochim Biophys Acta. 1154:237-282.
- [0103]** Berger F, Grini PE, Schnittger A. 2006. Endosperm: an integrator of seed growth and development. Curr Opin Plant Biol. 9:664-670.
- [0104]** Boter M, Ruiz-Rivero O, Abdeen A, Prat S. 2004. Conserved MYC transcription factors play a key role in jasmonate signaling both in tomato and *Arabidopsis*. Genes Dev. 18:1577-1591.
- [0105]** Bringi NY. 1987. Non-traditional oilseeds and oils of India. New Delhi: Oxford & IBH Publishing Co. Pvt. Ltd.
- [0106]** Brummelkamp, T.R. et al. (2002). A system for stable expression of short interfering RNAs in mammalian cells. Science 296:550-553.
- [0107]** Chan CS, Guo L, Shih MC. 2001. Promoter analysis of the nuclear gene encoding the chloroplast glyceraldehyde-3-phosphate dehydrogenase B subunit of *Arabidopsis thaliana*. Plant Mol Biol. 46:131-141.

- [0108] Chen GQ, Turner C, He X, Nguyen T, McKeon TA, Laudencia-Chingcuanco D. 2007. Expression profiles of genes involved in fatty acid and triacylglycerol synthesis in castor bean (*Ricinus communis* L.). *Lipids* 42: 263-274.
- [0109] Chen Q, Zhou H, Chen J, Wang X. 2006. A Gateway-based platform for multigene plant transformation. *Plant Molecular Biology*. 62: 927-936.
- [0110] Chow L, Chou M, Ho C, Chuang C, Pan F, Wu S, Lin J. 1999. Purification, characterization and molecular cloning of trichoanguin, a novel type I ribosome-inactivating protein from the seeds of *Trichosanthes anguina*. *Biochem J*. 338:211-219.
- [0111] Christou P, Capell T, Kohli A, Gatehouse JA, Gatehouse AM. 2006. Recent developments and future prospects in insect pest control in transgenic crops. *Trends Plant Sci*. 11:302-308.
- [0112] Chye ML, Huang BQ, Zee SY. 1999. Isolation of a gene encoding Arabidopsis membrane-associated acyl-CoA binding protein and immunolocalization of its gene product. *Plant J*. 18: 205-214.
- [0113] Constans A. (2002). RNAi for the masses. *The Scientist* 16:36.
- [0114] Datta, K., Vasquez, A., Tu, J., Torrizo, L., Alam M.F., Oliva, N., Abrigo, E., Khush, G.S., Datta, S.K. 1998. Constitutive and tissue-specific differential expression of the cryIA(b) gene in transgenic rice plants conferring resistance to rice insect pest. *Theor Appl Genet*. 97:20-30.
- [0115] Dellaporta SL, Wood J, Hicks JB. 1983. A plant DNA miniprep: version II. *Plant Mol. Biol. Rep.* 1:19-21.
- [0116] Eulgem T, Rushton PJ, Schmelzer E, Hahlbrock K and Somssich IE. 1999. Early nuclear events in plant defence signalling: rapid gene activation by WRKY transcription factors. *EMBO J*. 18:4689-4699.
- [0117] Fire, A. et al. (1998). Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*. *Nature* 391:806-811.
- [0118] Fujiwara T, Beachy RN. 1994. Tissue-specific and temporal regulation of a beta-conglycinin gene: roles of the RY repeat and other *cis*-acting elements. *Plant Mol Biol*. 24:261-272.
- [0119] Forson FK, Oduro EK and Hammond-Donkoh E. 2004. Performance of jatropha oil blends in a diesel engine. *Renewable Energy*. 29:1135-1145.
- [0120] Giuliano G, Pichersky E, Malik VS, Timko MP, Scolnik PA, Cashmore AR. 1988. An evolutionarily conserved protein binding sequence upstream of a plant light-regulated gene. *Proc Natl Acad Sci USA*. 85:7089-7093.
- [0121] Hammond, S.M et al. (2001). Post-transcriptional gene silencing by double-stranded RNA. *Nature Rev Gen* 2:110-119.
- [0122] Hannah LC, James M. 2008. The complexities of starch biosynthesis in cereal endosperms. *Curr Opin Biotechnol*. 19:160-165.
- [0123] Hartley JL, Temple GF, and Brasch MA. 2000. DNA cloning using in vitro site-specific recombination. *Genome Res*. 10:1788-1795.

- [0124] He WJ, Liu WY. 2004. Both N- and C-terminal regions are essential for cinnamomin A-chain to deadenylate ribosomal RNA and supercoiled double-stranded DNA. *Biochem J.* 377:17–23.
- [0125] Hiei Y, Ohta S, Komari T, Kumashiro T. 1994. Efficient transformation of rice (*Oryza sativa* L.) mediated by *Agrobacterium* and sequence analysis of the boundaries of the T-DNA. *Plant J.* 6: 271-282.
- [0126] Howe GA and Jander G. 2008. Plant immunity to insect herbivores. *Annu Rev Plant Biol.* 59:41-66.
- [0127] Hutvagner, G. and Zamore, P.D. (2002). RNAi: nature abhors a double-strand. *Curr Opin Genet Dev* 12(2):225-232.
- [0128] Kolesnik T, Szeverenyi I, Bachmann D, Kumar, CS, Jiang, S, Ramamoorthy, R, Cai, M, Ma ZG, Sundaresan V, Ramachandran S. 2004. Establishing an efficient Ac/Ds tagging system in rice: large-scale analysis of Ds flanking sequences. *Plant J.* 37:301-314.
- [0129] Kuwano M, Mimura T, Takaiwa F, Yoshida KT. 2009. Generation of stable 'low phytic acid' transgenic rice through antisense repression of the 1D-myo-inositol 3-phosphate synthase gene (RINO1) using the 18-kDa oleosin promoter. *Plant Biotechnol J.* 7:96-105.
- [0130] Lam E, Chua NH. 1989. ASF-2: A factor that binds to the cauliflower mosaic virus 35S promoter and a conserved GATA motif in cab promoters. *Plant Cell.* 1:1147-1156.
- [0131] Lin L, Liu Y, Xu X, Li B. 2003. Efficient linking and transfer of multiple genes by a multigene assembly and transformation vector system. *Proceedings of the National Academy of Sciences, USA.* 100, 5962–5967.
- [0132] Lin J, Chen Y, Xu Y, Yang F, Tang L, Chen F. 2003a. Cloning and expression of Curcin, a Ribosome-inactivating Protein from the seeds of *Jatropha curcas*. *Acta Botanica Sinica.* 45: 858-863.
- [0133] Lin J, Li Y, Zhou X, Tang K, Chen F. 2003b. Cloning and characterization of a Curcin gene encoding a Ribosome Inactivating Protein from *Jatropha curcas*. *DNA sequence.* 14: 311-317.
- [0134] Liu YG, Whittier RF. 1995. Thermal Asymmetric Interlaced PCR: Automatable Amplification and Sequencing of Insert End Fragments from PI and YAC Clones for Chromosome Walking. *Genomics.* 25: 674-681.
- [0135] Maine, E.M. (2000). A conserved mechanism for post-transcriptional gene-silencing? *Genome Biol.* 1(3):reviews1018.1–reviews1018.4.
- [0136] Makkar HPS, Becker K, Sporer F and Wink M. 1997. Studies on nutritive potential and toxic constituents of different provenances of *Jatropha curcas*. *J Agric Food Chem.* 45:3152-3157.
- [0137] Mansoor S, Amin I, Hussain M, Zafar Y, Briddon RW. 2006. Engineering novel traits in plants through RNA interference. *Trends Plant Sci.* 11:559-565.
- [0138] Miki D and Shimamoto K. 2004. Simple RNAi Vectors for Stable and Transient Suppression of Gene Function in Rice. *Plant and Cell Physiology.* 45:490-495.
- [0139] Moon YH, Song SK, Choi KW, Lee JS. 1997. Expression of a cDNA encoding *Phytolacca insularis* antiviral protein confers virus resistance on transgenic potato plants. *Mol Cells.* 7:807– 815.

- [0140] Narayana, DSA, Rangaswamy KT, Shankarappa KS, Maruthi MN, Lakshminarayana Reddy CN, Rekha AR, Keshava Murthy KV. 2007. Distinct Begomoviruses Closely Related to Cassava Mosaic Viruses cause Indian *Jatropha* Mosaic Disease. *Int'l J Virol* 3:1-11.
- [0141] Nishiuchi T, Shinshi H, Suzuki K. 2004. Rapid and transient activation of transcription of the ERF3 gene by wounding in tobacco leaves: Possible involvement of NtWRKYs and autorepression. *J Biol Chem*. 279: 55355-55361.
- [0142] Paddison, P.J. et al. (2002). Short hairpin RNAs (shRNAs) induce sequence specific silencing in mammalian cells. *Genes & Dev* 16:948-958.
- [0143] Peterson DG, Tomkins JP, Frisch DA, Wing RA, Paterson AH. 2000. Construction of plant bacterial artificial chromosome (BAC) libraries: An illustrated guide. *Journal of Agricultural Genomics* 5 (www dot ncgr dot org backslash research backslash jag).
- [0144] Plant AL, van Rooijen GJ, Anderson CP, Moloney MM. 1994. Regulation of an *Arabidopsis* oleosin gene promoter in transgenic *Brassica napus*. *Plant Mol Biol*. 25:193-205.
- [0145] Pramanik K. 2003. Properties and use of *jatropha curcas* oil and diesel fuel blends in compression ignition engine. *Renewable Energy*. 28:239-248.
- [0146] Qu L, Xing Y, Liu W, Xu X and Song Y. 2008. Expression pattern and activity of six glutelin gene promoters in transgenic rice. *J Exp. Bot*. 59:2417-2424.
- [0147] Roesler K, Shintani D, Savage L, Boddupalli S and Ohlrogge J. 1997. Targeting of the *Arabidopsis* homomeric acetyl-coenzyme A carboxylase to plastids of rapeseeds. *Plant Physiol*. 113: 75-81.
- [0148] Roh JY, Choi JY, Li MS, Jin BR, Je YH. 2007. *Bacillus thuringiensis* as a specific, safe, and effective tool for insect pest control. *J Microbiol Biotechnol*. 17:547-559.
- [0149] Sambrook J, Frisch E, Maniatis T. 1989. *Molecular Cloning: A Laboratory Manual* 2nd edn. Cold Spring Harbor: Cold Spring Harbor Laboratory Press.
- [0150] Stalberg K, Ellerstrom M, Ezcurra I, Ablov S, Rask L. 1996. Disruption of an overlapping E-box/ABRE motif abolished high transcription of the napA storage-protein promoter in transgenic *Brassica napus* seeds. *Planta*. 199:515-519.
- [0151] Stirpe F, Pession-Brizzia, Lorenzoni E, Strocchi P, Montanarol L, Sperti S. 1976. Studies on the Proteins from the Seeds of *Croton tiglium* and of *Jatropha curcas*. *Biochem J*. 156: 1-6.
- [0152] Tapia G, Verdugo I, Yanez M, Ahumada I, Theoduloz C, Cordero C, Poblete F, Gonzalez E, Ruiz-Lara S. 2005. Involvement of Ethylene in Stress-Induced Expression of the TLC1.1 Retrotransposon from *Lycopersicon chilense* Dun. *Plant Physiol*. 138:2075-2086.
- [0153] Terzaghi WB, Cashmore AR. 1995. Light-regulated transcription. *Annu Rev Plant Physiol Plant Mol Biol*. 46:445-474.
- [0154] Timmons, L. and Fire, A. (1998). Specific interference by ingested dsRNA. *Nature* 395:854.
- [0155] Urao T, Yamaguchi-Shinozaki K, Urao S, Shinozaki K. 1993. An *Arabidopsis* myb homolog is induced by dehydration stress and its gene product binds to the conserved MYB recognition sequence. *Plant Cell*. 5:1529-1539.

- [0156] van Loon LC, Rep M, Pieterse CM. 2006. Significance of inducible defense-related proteins in infected plants. *Annu Rev Phytopathol.* 44:135-162.
- [0157] Vaucheret, H. et al. (2001). Post-transcriptional gene silencing in plants. *J Cell Sci* 114:3083-3091.
- [0158] Vickers CE, Xue G, Gresshoff PM. 2006. A novel cis-acting element, ESP, contributes to high-level endosperm-specific expression in an oat globulin promoter. *Plant Mol Biol.* 62: 195-214.
- [0159] Villain P, Mache R, Zhou DX. 1996. The mechanism of GT element-mediated cell type-specific transcriptional control. *J Biol Chem.* 271:32593-32598.
- [0160] Wakasa Y, Yasuda H, Takaiwa F. 2006. High accumulation of bioactive peptide in transgenic rice seeds by expression of introduced multiple genes. *Plant Biotechnology Journal.* 4, 499–510.
- [0161] Wei Q, Huang M, Xu Y, Zhang X, Chen F. 2005. Expression of a ribosome inactivating protein (curcin 2) in *Jatropha curcas* is induced by stress. *J. Biosci.* 30:351-357.
- [0162] Wu, C., Washida, H., Onodera, Y., Harada, K., Takaiwa, F. 2000. Quantitative nature of the Prolamin-box, ACGT and AACA motifs in a rice glutelin gene promoter: minimal cis-element requirements for endosperm-specific gene expression. *Plant J.* 23:415-421.
- [0163] Yu D, Chen C, Chen Z. 2001. Evidence for an important role of WRKY DNA binding proteins in the regulation of NPR1 gene expression. *Plant Cell.* 13: 1527-1540.
- [0164] Zamore, P.D. (2001). RNA interference: listening to the sound of silence. *Nature Struct Biol* 8:746-750.

WHAT IS CLAIMED IS:

1. An isolated nucleic acid having endosperm-specific promoter activity in a plant selected from the group consisting of:
 - (a) a nucleic acid comprising nucleotides 1 to 3181 of SEQ ID NO:7;
 - (b) a nucleic acid comprising nucleotides 1 to 2888 of SEQ ID NO:7;
 - (c) a nucleic acid comprising nucleotides 1142 to 3181 of SEQ ID NO:7;
 - (d) a nucleic acid comprising nucleotides 1 to 3181 of SEQ ID NO:7 with nucleotides 2944 to 3170 deleted;
 - (e) a nucleic acid comprising nucleotides 1142 to 3181 of SEQ ID NO:7 with nucleotides 2944 to 3170 deleted;
 - (f) a nucleic acid comprising nucleotides 2688 to 3181 of SEQ ID NO:7 with nucleotides 2944 to 3170 deleted
2. A nucleic acid construct comprising the nucleic acid having endosperm-specific promoter activity of claim 1 operably linked to a second nucleic acid encoding a DNA of interest.
3. The nucleic acid construct of claim 2, wherein the DNA of interest encodes a double stranded RNA (dsRNA) molecule targeted to a *Curcin* gene and capable of reducing expression of a *Curcin* gene in a *Jatropha* plant by RNAi, wherein the *Curcin* gene is selected from the group consisting of *Curcin 1*, *Curcin 2* and *Curcin 2A*.
4. The nucleic acid construct of claim 3, wherein the *Curcin* gene is *Curcin 1*.
5. The nucleic acid construct of claim 3, wherein the *Curcin* gene is *Curcin 2*.
6. The nucleic acid construct of claim 3, wherein the *Curcin* gene is *Curcin 2A*.
7. The nucleic acid construct of any one of claims 3-6, wherein the dsRNA molecule is an siRNA molecule, a shRNA molecule or a micro-RNA molecule.

8. A transgenic plant cell containing within its genome the nucleic acid construct of claim 2.
9. The transgenic plant cell of claim 8, wherein the plant is selected from the group consisting of a *Jatropha* species, castor bean, palm oil, coconut, peanut, rapeseeds, soybean, sunflower, cassava, tomato, beans, cotton, cucumber, cabbage, barley, oats, wheat, corn and rice.
10. The transgenic plant cell of claim 9, wherein the *Jatropha* species is *Jatropha curcas*.
11. A transgenic plant containing within its genome the nucleic acid construct of claim 2.
12. The transgenic plant of claim 11, wherein the plant is selected from the group consisting of a *Jatropha* species, castor bean, palm oil, coconut, peanut, rapeseeds, soybean, sunflower, cassava, tomato, beans, cotton, cucumber, cabbage, barley, oats, wheat, corn and rice.
13. The transgenic plant of claim 12, wherein the *Jatropha* species is *Jatropha curcas*.
14. A method of producing a transgenic plant cell comprising transforming cells of the plant with the nucleic acid construct of claim 2.
15. A method of producing a transgenic plant comprising transforming cells of the plant with the nucleic acid construct of claim 2 and regenerating a transgenic plant from the transformed plant cells.
16. The method of claim 14 or 15, wherein the plant is selected from the group consisting of a *Jatropha* species, castor bean, palm oil, coconut, peanut, rapeseeds, soybean, sunflower, cassava, tomato, beans, cotton, cucumber, cabbage, barley, oats, wheat, corn and rice.
17. The method of claim 16, wherein the *Jatropha* species is *Jatropha curcas*.

18. A transgenic plant cell containing within its genome the nucleic acid construct of any one of claims 3-7.
19. The transgenic plant cell of claim 18, wherein the plant is selected from the group consisting of a *Jatropha* species, castor bean, palm oil, coconut, peanut, rapeseeds, soybean, sunflower, cassava, tomato, beans, cotton, cucumber, cabbage, barley, oats, wheat, corn and rice.
20. The transgenic plant cell of claim 19, wherein the *Jatropha* plant is *Jatropha curcas*.
21. A curcin deficient transgenic plant containing within its genome the nucleic acid construct of any one of claims 3-7.
22. The curcin deficient transgenic plant of claim 21, wherein the plant is selected from the group consisting of a *Jatropha* species, castor bean, palm oil, coconut, peanut, rapeseeds, soybean, sunflower, cassava, tomato, beans, cotton, cucumber, cabbage, barley, oats, wheat, corn and rice.
23. The curcin deficient transgenic plant of claim 22, wherein the *Jatropha* species is *Jatropha curcas*.
24. A method of producing a transgenic plant cell comprising transforming cells of the plant with the nucleic acid construct of any one of claims 3-7.
25. A method of producing a curcin deficient transgenic *Jatropha* plant comprising transforming cells of a plant with the nucleic acid construct of any one of claims 3-7 and regenerating a curcin deficient transgenic plant from the transformed plant cells.
26. The method of claim 24 or 25, wherein the plant is selected from the group consisting of a *Jatropha* species, castor bean, palm oil, coconut, peanut, rapeseeds, soybean, sunflower, cassava, tomato, beans, cotton, cucumber, cabbage, barley, oats, wheat, corn and rice.

27. The method of claim 26, wherein the *Jatropha* plant is *Jatropha curcas*.