


**IMIDAZOLE DERIVATIVES HAVING AN INHIBITORY
ACTIVITY FOR FARNESYL TRANSFERASE AND PROCESS FOR
PREPARATION THEREOF**

TECHNICAL FIELD

The present invention relates to a novel imidazole derivative represented by the following formula (1) which shows an inhibitory activity against farnesyl transferase:

[Formula 1]



in which A, n_1 and Y are defined as described below, or pharmaceutically acceptable salts or isomers thereof.

The present invention also relates to a process for preparation of the compound of formula (1), to intermediates which are used in the preparation of the compound of formula (1), and to a pharmaceutical composition comprising the compound of formula (1) as an active ingredient.

BACKGROUND ART

Mammalian Ras proteins act as molecular switches in the signalling events associated with cell growth and differentiation. The ras proto-oncogene family consists of three members, N-, K-, and H-ras, which code for highly homologous 4 types of proteins; i.e., H, N-ras proteins of 189 residues and two isomorphous K-ras-4B and K-ras-4A

proteins of 188 and 189 residues, respectively. The chemical basis for the switch mechanism involves cycling of the protein between the inactive (off) guanosine diphosphate (GDP) bound state and the active (on) guanosine triphosphate (GTP) bound state (Bourne, H. R.; Sanders, D. A.; McCormick, F.; *Nature*, 1991, 349, 117). Biochemical and structural studies have shown that point mutations of the residues 12, 13 and 61, positioned in the neighborhood of phosphoryl group of GTP, resulting in the decrease of guanosine triphosphatase activity are associated with many human cancers, particularly, pancreatic cancer, urinary bladder carcinoma, colon cancer, etc. (Bos, J. L., *Cancer Res.*, 1989, 49, 4682).

Ras protein is synthesized as a cytosolic precursor that ultimately localized to the cytoplasmic face of the plasma membrane after a series of posttranslational modification (Gibbs, J. B., *Cell* 1991, 65, 1). These series of biochemical modifications, by changing the electrical charge state or spacial structure to increase the hydrophobicity allow Ras protein to attach to cell membrane more easily. The first and obligatory step in the series is the addition of a farnesyl moiety to the cysteine residue of the C-terminal CAAX motif (C, cysteine; A, usually aliphatic residue; X, any other amino acid) in a reaction catalyzed by farnesyl protein transferase (FTase). This modification is essential for Ras function, as demonstrated by the inability of Ras mutants lacking the C-terminal cysteine to be farnesylated, to localize to the plasma, and to transform mammalian cells in culture (Hancock, J. F., Magee, A. I., Childs, J. E., Marshall, C. J., *Cell* 1989, 57, 1167). The subsequent posttranslational modifications, cleavage of the AAX residues, carboxyl methylation of the the farnesylated cysteine, and palmitoylation of the cysteines located upstream of the CAAX motif in H- and N-ras proteins are not obligatory for Ras membrane association or cellular transforming activity.

Interestingly, K-ras-4B, different from H- and N-ras, has a multiple lysine rich region named polybasic domain, instead of having cysteine required for palmitoylation, thereby facilitating the farnesylated ras protein to bind to anionic lipid layer of cell membrane. The inhibitors of FTase that catalyzes the obligatory modification have therefore been suggested as anticancer agents for tumors in which Ras oncogene contributes to transformation (Buses, J. E. et al., Chemistry & Biology, 1995, 2, 787). A number of FTase inhibitors recently identified demonstrated potent and specific ability to block Ras farnesylation, signalling and transformation in transformed cells and tumor cell lines both in vitro and in animal models (Kohl, N. E. et al., Proc. Natl. Acad. Sci. USA. 1994, 91, 9141; Kohl, N. E. et al., Nature Medicine, 1995, 1 792).

However, most of the inhibitors are related to CAAX motif as Ras substrate mimic and peptidic in nature or contain a sulfhydryl group (USP No. 5,141,851; Kohl, N. E. et al., Science, 1993, 260, 1934; PCT/US95/12224, Graham et al.; Sebt, S. M. et al., J. Biol. Chem., 1995, 270, 26802; James, G. L. et al., Science, 1993, 260, 1937; Bishop, W. R. et al., J. Biol. Chem., 1995, 270, 30611). Recently, a new type of peptidomimetic inhibitor imitating catalytic step of FTase has been reported (Poulter, C.D. et al., J. Am. Chem. Soc., 1996, 118, 8761). The chemical basis of the inhibitor design relates to the reaction mechanism. This is, transferring prenyl group by the enzyme is electrophilic displacement and the reaction requires (+) charge in a transition state.

These inhibitors previously described, however, possess limited activity and selectivity for inhibition of the oncogenic function of Ras proteins, particularly K-ras-4B, which is found to be most common in human cancer. Therefore, new inhibitor having the ability of effectively

inhibiting K-ras activity is required.

With regard to the restenosis and vascular proliferative diseases, it has been shown that inhibition of cellular ras prevents smooth muscle proliferation after vascular injury in vivo (Indolfi C. et al., Nature Med., 1995, 1(6), 541-545). This report definitively supports a role for farnesyl transferase inhibitors in this disease, showing inhibition of accumulation and proliferation of vascular smooth muscle.

DISCLOSURE OF INVENTION

The present inventors have performed studies for developing a compound having the structural characteristics imitating an intermediate state of catalytic reaction of FTase and as a result, found that imidazole derivatives according to the present invention can potently inhibit the enzyme.

Therefore, the object of the present invention is to provide an imidazole derivative of formula (1) which inhibits the activity of FTase, a process for preparation thereof, and an intermediate which can be used effectively for the preparation of the compound of formula (1).


It is another object of the present invention to provide a pharmaceutical composition comprising the compound of formula (1) as an active ingredient.

BEST MODE FOR CARRYING OUT THE INVENTION

It is the first object of the present invention to provide an imidazole derivative represented by the following formula (1) which

inhibit the activity of farnesyl transferase :


[Formula 1]



in which

n_1 represents an integer of 1 to 4,

A represents hydrogen; straight-chain or branched C_1 - C_{10} -alkyl which may be optionally substituted by C_3 - C_7 -cycloalkyl or lower alkoxy; or a radical selected from the following group:




wherein

R_1 and R_1' independently of one another represent hydrogen, halogen, cyano, nitro, hydroxycarbonyl, aminocarbonyl, aminothiocarbonyl, lower alkoxy, phenoxy, phenyl, benzyloxy, or lower alkyl which may be optionally substituted by C_3 - C_6 -cycloalkyl,

R_2 represents hydrogen or lower alkyl, or represents -E-F wherein E is $-CH_2-$, $-C(O)-$ or $-S(O)_2-$ and F is hydrogen; lower alkyl which may be optionally substituted by phenoxy or biphenyl; lower alkoxy which may be optionally substituted by aryl; phenyl; benzyl; benzyloxy; or amino which may be optionally substituted by lower alkyl, benzyl or C_5 - C_6 -cycloalkyl,

R_3 represents hydrogen, lower alkyl or phenyl,

R_4 represents a radical selected from the following group:



wherein

n_2 and n_3 independently of one another denote 0, 1, 2, 3 or 4,


R_5 and R_9 independently of one another represent hydrogen, lower alkyl, lower alkoxy, phenoxy, phenyl, hydroxy or halogen,

R_6 and R_8 independently of one another represent hydrogen, lower alkyl, lower alkoxy, phenoxy, phenyl, cyano, hydroxy or halogen,

R_7 represents hydrogen; lower alkyl which may be optionally substituted by C_3 - C_6 -cycloalkyl; lower alkoxy; hydroxy; C_3 - C_6 -cycloalkyl; di(lower alkyl)amino; phenyl; phenoxy; or halogen,

R_{10} represents hydrogen, lower alkyl or lower alkoxy,

Y represents a radical selected from the following group:




wherein

X represents O or S,

B represents hydrogen, or lower alkyl which may be optionally substituted by hydroxy, mercapto, lower alkoxy, lower alkylthio or aryl,

C represents hydrogen, or lower alkyl which may be optionally substituted by aryl; or represents a radical selected from the following group:




wherein

R_{11} and R_{12} independently of one another represent hydrogen, lower alkyl, lower alkoxy, halogen, cyano, hydroxycarbonyl, amino-carbonyl, aminothiocarbonyl, hydroxy, phenyl or phenoxy,

R_{13} and R_{14} independently of one another represent hydrogen, lower

alkyl, aryl or $-(CH_2)_{n_4}-X-R_{15}$ wherein X is defined as previously described, n_4 is an integer of 2 to 4 and R_{15} is lower alkyl, D represents amino acid residue or lower alkyl ester of amino acid residue; or represents a radical selected from the following group:



wherein

R_{10} is defined as previously described,

Q represents O, S, S=O or SO₂,

Z represents O, S, S=O, SO₂, C=O or C=S, or represents CH-R₂₀ or N-R₂₀ (wherein R₂₀ is hydrogen, lower alkyl or hydroxy),

n_5 denotes an integer of 1 to 3,


R_{16} and R_{17} independently of one another represents hydrogen; aryl;

lower alkyl which may be optionally substituted by aryl or

cyanoaryl; or $-\frac{1}{2}-(CH_2)_{n_4}-Q-R_{10}$ wherein n_4 , Q and R_{10} are defined as previously described,

R_{18} and R_{19} independently of one another represents hydrogen; halogen; hydroxy; cyano; lower alkyl; lower alkoxy; alkoxyalkyl; alkylthio; hydroxycarbonyl; aminocarbonyl; aminothiocarbonyl; alkylsulfonyl; alkylthioalkyl; alkylthioalkyloxy; aryl; or oxy, thio, sulfonyl or lower alkyl substituted by aryl,


G represents a radical selected by the following group:



wherein

R_{11} and R_{12} are defined as previously described,


I represents lower alkoxy, or represents a radical selected from the following group:



wherein

R_{16} , R_{17} and Z are defined as previously described,


L represents a radical selected from the following group:




wherein Z and Q are defined as previously described,

provided that (1) n_2 is other than 0 when R_3 is hydrogen, and

(2) Y is other than



, when A is



, or pharmaceutically acceptable salts or isomers

thereof.

Particularly, the compound according to the present invention has a quite different structure from the known inhibitors for farnesyl transferase, and furthermore it does never include the thiol moiety.

In the definitions for the substituents of the compound of formula (1), the term "lower alkyl" means a straight-chain or branched alkyl having 1 to 4 carbon atoms which includes methyl, ethyl, isopropyl, isobutyl and t-butyl.


Since the compound of formula (1) according to the present invention may have asymmetric carbon atoms depending on the substituents, it can be present in the form of R or S isomer, racemate, or mixtures thereof. Thus, the present invention also includes all of these stereoisomers and their mixtures.

Also, the compound of formula (1) according to the present invention can form a pharmaceutically acceptable salt. Such salt includes non-toxic acid addition salt containing pharmaceutically acceptable anion, for example a salt with inorganic acids such as hydrochloric acid,

sulfuric acid, nitric acid, phosphoric acid, hydrobromic acid, hydriodic acid, etc., a salt with organic carboxylic acids such as tartaric acid, formic acid, citric acid, acetic acid, trichloroacetic acid, trifluoroacetic acid, gluconic acid, benzoic acid, lactic acid, fumaric acid, maleic acid, asparagic acid, etc., or a salt with sulfonic acids such as methanesulfonic acid, benzenesulfonic acid, p-toluenesulfonic acid, naphthalenesulfonic acid, etc.; base addition salt for example a salt with pyridine or ammonia; and metal addition salt, for example, a salt with alkali metal or alkaline earth metal such as lithium salt. Further, the present invention includes a solvate of the compound of formula (1) such as alcoholate or hydrate thereof. They can be produced by conventional conversion methods.

Among the compound of formula (1) according to the present invention, the preferred compounds include those wherein

- n_1 represents an integer of 1 to 3,
- A represents hydrogen; straight-chain or branched C_1 - C_{10} -alkyl which may be optionally substituted by C_3 - C_7 -cycloalkyl or lower alkoxy; or a radical selected from the following group:




wherein

- R_1 and R_1' independently of one another represent hydrogen, halogen, cyano, nitro, hydroxycarbonyl, aminocarbonyl, aminothiocarbonyl, lower alkoxy, phenoxy, phenyl, benzyloxy, or lower alkyl which may be optionally substituted by C_3 - C_6 -cycloalkyl,
- R_2 represents hydrogen or lower alkyl, or represents -E-F wherein E is $-CH_2-$, $-C(O)-$ or $-S(O)_2-$ and F is hydrogen; lower alkyl which

may be optionally substituted by phenoxy or biphenyl; lower alkoxy which may be optionally substituted by aryl; phenyl; benzyl; benzyloxy; or amino which may be optionally substituted by lower alkyl, benzyl or C₅-C₆-cycloalkyl,

R₃ represents hydrogen or lower alkyl,

R₄ represents a radical selected from the following group:



wherein


n₂ and n₃ independently of one another denote 0, 1, 2, 3 or 4,

R₅, R₆, R₈ and R₉ independently of one another represent hydrogen, lower alkyl, lower alkoxy, hydroxy or halogen,

R₇ represents hydrogen; lower alkyl which may be optionally substituted by C₃-C₆-cycloalkyl; lower alkoxy; hydroxy; C₃-C₆-cycloalkyl; or halogen,

R₁₀ represents hydrogen, methyl or methoxy,

Y represents a radical selected from the following group:




wherein

X represents O or S,

B represents hydrogen, or lower alkyl which may be optionally substituted by lower alkoxy or aryl,

C represents hydrogen, or lower alkyl which may be optionally

substituted by aryl; or represents a radical selected from the following group:




wherein

R_{11} and R_{12} independently of one another represent hydrogen, lower alkyl, lower alkoxy, halogen, cyano, aminocarbonyl, phenyl or phenoxy,

R_{13} and R_{14} independently of one another represent hydrogen, lower

alkyl, aryl or $-(CH_2)_{n_4}-X-R_{15}$ wherein X is defined as previously described, n_4 is 2 and R_{15} is lower alkyl,

D represents amino acid residue or lower alkyl ester of amino acid residue; or represents a radical selected from the following group:



wherein

R_{10} is defined as previously described,

Q represents O, S, S=O or SO₂,

Z represents O, S, S=O, SO₂ or C=O, or represents CH-R₂₀ or N-R₂₀ (wherein R₂₀ is hydrogen, lower alkyl or hydroxy),


n₅ denotes an integer of 1 to 3,

R₁₆ and R₁₇ independently of one another represents hydrogen; aryl; lower alkyl which may be optionally substituted by aryl or

cyanoaryl; or $\text{---}\frac{1}{2}\text{---}(\text{CH}_2)_{n_4}\text{---Q---R}_{10}$ wherein n₄, Q and R₁₀ are defined as previously described,

R₁₈ and R₁₉ independently of one another represents hydrogen; halogen; hydroxy; cyano; lower alkyl; lower alkoxy; alkoxyalkyl; alkylthio; hydroxycarbonyl; aminocarbonyl; aminothiocarbonyl; alkylsulfonyl; alkylthioalkyl; alkylthioalkyloxy; aryl; or oxy, thio, sulfonyl or lower alkyl substituted by aryl,


G represents a radical selected by the following group:



wherein

R₁₁ and R₁₂ are defined as previously described,


I represents lower alkoxy, or represents a radical selected from the following group:





wherein

R₁₆, R₁₇ and Z are defined as previously described,


L represents a radical selected from the following group:



wherein Z and Q are defined as previously described,
provided that (1) n_2 is other than 0 when R_3 is hydrogen, and



Particularly preferred compounds include those wherein Y



Typical examples of the compound of formula (1) according to the present invention are presented in the following Table 1.

Table 1-1







COM. NO.	STRUCTURE	COM. NO.	STRUCTURE
1		2	
3		4	
5		6	

Table 1-2







COM. NO.	STRUCTURE	COM. NO.	STRUCTURE
7		8	
9		10	
11		12	

Table 1-3







COM. NO.	STRUCTURE	COM. NO.	STRUCTURE
13		14	
15		16	
17		18	

Table 1-4







COM. NO.	STRUCTURE	COM. NO.	STRUCTURE
19		20	
21		22	
23		24	

Table 1-5.







COM. NO.	STRUCTURE	COM. NO.	STRUCTURE
25		26	
27		28	
29		30	

Table 1-6







COM. NO.	STRUCTURE	COM. NO.	STRUCTURE
31		32	
33		34	
35		36	

Table 1-7







COM. NO.	STRUCTURE	COM. NO.	STRUCTURE
37		38	
39		40	
41		42	

Table 1-8


COM. NO.	STRUCTURE
43	 The chemical structure of compound 43 is a complex molecule. It features a central pyrazole ring. One nitrogen of the pyrazole is connected via a methylene group to a benzimidazole system. The benzimidazole system consists of a benzene ring fused to an imidazole ring, which is further fused to a pyrazole ring. The pyrazole ring of the benzimidazole system has a carbonyl group (C=O) attached to it, which is in turn connected to a piperidine ring. The piperidine ring has a methyl group attached to one of its nitrogens. The other nitrogen of the central pyrazole ring is connected via a methylene group to a 4-hydroxyphenyl group.

Table 1-9











COM. NO.	STRUCTURE	COM. NO.	STRUCTURE
44		45	
46		47	
48		49	
50		51	
52		53	

Table 1-10











COM. NO.	STRUCTURE	COM. NO.	STRUCTURE
54		55	
56		57	
58		59	
60		61	
62		63	

Table 1-11











COM. NO.	STRUCTURE	COM. NO.	STRUCTURE
64		65	
66		67	
68		69	
70		71	
72		73	

Table 1-12

COM. NO.	STRUCTURE	COM. NO.	STRUCTURE
74		75	
76		77	
78		79	
80		81	
82		83	

Table 1-13











COM. NO.	STRUCTURE	COM. NO.	STRUCTURE
84		85	
86		87	
88		89	
90		91	
92		93	

Table 1-14











COM. NO.	STRUCTURE	COM. NO.	STRUCTURE
94		95	
96		97	
98		99	
100		101	
102		103	

Table 1-15











COM. NO.	STRUCTURE	COM. NO.	STRUCTURE
104		105	
106		107	
108		109	
110		111	
112		113	

Table 1-16











COM. NO.	STRUCTURE	COM. NO.	STRUCTURE
114		115	
116		117	
118		119	
120		121	
122		123	

Table 1-17

COM. NO.	STRUCTURE	COM. NO.	STRUCTURE
124		125	
126		127	
128		129	
130		131	
132		133	

Table 1-18









COM. NO.	STRUCTURE	COM. NO.	STRUCTURE
134		135	
136		137	
138		139	
140		141	

Table 1-19









COM. NO.	STRUCTURE	COM. NO.	STRUCTURE
142		143	
144		145	
146		147	
148		149	

Table 1-20









COM. NO.	STRUCTURE	COM. NO.	STRUCTURE
150		151	
152		153	
154		155	
156		157	

Table 1-21

COM. NO.	STRUCTURE	COM. NO.	STRUCTURE
158		159	
160		161	
162		163	


It is another object of the present invention to provide processes for preparing the imidazole derivative of formula (1) as defined above.

According to the present invention, the imidazole derivative of formula (1) can be prepared by processes characterized in that

(a) a compound represented by the following formula (2) is


reacted in a solvent in the presence of a base with a compound represented by the following formula (3), then the trityl group in the product thus obtained is eliminated in the presence of trifluoroacetic acid to produce a compound represented by the following formula (1a); or

Reaction Scheme 1




(b) a compound represented by the following formula (4) is reacted in a solvent in the presence of a base with the compound of formula (3) to produce a compound represented by the following formula (1b); or


Reaction Scheme 2




(c) a compound represented by the following formula (5) is reacted in a solvent in the presence of a base with the compound of formula (3), the trityl group in the product thus obtained is eliminated in the presence of trifluoroacetic acid to produce a compound represented by the following formula (6), and then hydrogenation reaction is carried out to produce a compound represented by the following formula (1c); or

Reaction Scheme 3


(d) a compound represented by the following formula (7) is hydrolyzed to produce a compound represented by the following formula (8) which is then reacted with a compound represented by the following formula (9) in the presence of a coupling agent to produce a compound represented by the following formula (1d); or

Reaction Scheme 4


(e) the carbonyl group in a compound represented by the following formula (1e) is converted into the thiocarbonyl group in the presence of a sulfurizing agent to produce a compound represented by the following formula (1f); or

Reaction Scheme 5

(f) a compound represented by the following formula (1g) is coupled in a solvent with a compound represented by the following formula (10) to produce a compound represented by the following formula (1h); or


Reaction Scheme 6

(g) a compound represented by the following formula (11) is cyclized in an inert solvent to produce a compound represented by the following formula (1i); or

Reaction Scheme 7


(h) the amide group in the compound of formula (11) is converted into the thioamide group to produce a compound represented by the following formula (12) which is then cyclized in an inert solvent to produce a compound represented by the following formula (1j); or

Reaction Scheme 8




(i) a compound represented by the following formula (13) is reacted in a solvent with a compound represented by the following formula (14a) to produce the compound of formula (1j); or


Reaction Scheme 9




(j) the compound of formula (13) is reacted in a solvent with a compound represented by the following formula (14b) to produce a compound represented by the following formula (1k); or

Reaction Scheme 10

(k) a compound represented by the following formula (1l) is hydrolyzed in the presence of a base and the product thus obtained is reacted in a solvent in the presence of a coupling agent with a compound represented by the following formula (15) to produce a compound represented by the following formula (1m); or


Reaction Scheme 11

(l) a compound represented by the following formula (16) is reacted in a solvent in the presence of a base with a compound represented by the following formula (17) to produce a compound represented by the following formula (1n); or

Reaction Scheme 12

(m) a compound represented by the following formula (18) is reacted in a solvent in the presence of a base with the compound of formula (17) and deprotected to produce a compound represented by the following formula (10) which is then coupled with a compound represented by the following formula (19) to produce a compound represented by the following formula (1p):

Reaction Scheme 13



in the above reaction schemes

A, n₁, B, C, X, D, R₁₆, R₁₇, R₂, G, I, L, E and F are defined as previously described,

I' represents lower alkoxy,

I'' is identical with I except that lower alkoxy is not included,

T represents hydroxy or reactive leaving group, preferably halogen,

Tr represents trityl,

Cbz represents benzyloxycarbonyl and has the same meaning through the present specification.

However, the compound according to the present invention may be conveniently prepared by any methods designed by combining various synthetic ways known in the prior arts, and such combination can be easily performed by a person having ordinary skill in this art. The processes (a) to (m) will be more specifically explained in below.

In processes (a) to (e) for preparing the compound according to the present invention, any inert solvents which does not adversely affect to the reaction, preferably one or more selected from a group consisting of dimethylformamide, dimethylacetamide, ethanol, water, methylene chloride, chloroform, tetrahydrofuran and N-methylpyrrolidinone can be used. As the base, one or more selected from a group consisting of sodium hydride, potassium hydroxide, potassium carbonate, potassium t-butoxide, sodium amide, sodium bis(trimethylsilyl)amide and potassium bis(trimethylsilyl)amide, more preferably sodium hydride or potassium hydroxide can be mentioned. As the coupling agent used in the process for reacting the compound of formula (8) with the compound of formula (9), a mixture of 1-hydroxybenzotriazole and one or more substances selected from a group consisting of carbodiimides such as dicyclohexylcarbodiimide(DCC), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide(EDC), 1,1'-dicarbonyldiimidazole(CDI), etc., and inorganic dehydrating agent such as silicone tetrachloride can be mentioned. Among them, a mixture of 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide(EDC) and 1-hydroxybenzotriazole hydrate is particularly preferred.

The sulfurizing agent used in preparing the compound of formula (1f) from the compound of formula (1e) includes 2,4-bis(phenylthio)-1,3-dithia-2,4-diphosphatane-2,4-disulfide, Lawesson's Reagent and P_4S_{10} . 2,4-bis(phenylthio)-1,3-dithia-2,4-diphosphatane-2,4-disulfide can be used.

most preferably.

The compound of formula (1g) which is used as a starting material in process (f) can be prepared by deprotecting the corresponding compound which is protected by benzyloxycarbonyl group at position-1 of piperidine moiety. The deprotection reaction may be carried out by applying the conventional reaction conditions, preferably by using $\text{Pd}(\text{OH})_2/\text{C}$ or Pd/C in an alcohol solvent under hydrogen atmosphere. The compound of formula (1g) thus obtained is coupled with the compound of formula (10) in an inert solvent as mentioned above optionally in the presence of a tertiary amine base to produce the compound of formula (1h). Alternatively, the compound of formula (1g) can be reacted in the presence of a coupling agent as mentioned for process (d) with the carboxylic acid derivative ($\text{T}=\text{OH}$) to produce the compound of formula (1h) in the form of amide.

In the cyclization reactions of (g) and (h) for preparing the compounds (1i) and (1j), any inert solvents, preferably one or more selected from tetrahydrofuran and ethanol can be used. As the sulfurizing agent used in the conversion procedure of amide to thioamide group in process (h), 2,4-bis(phenylthio)-1,3-dithia-2,4-diphosphatane-2,4-disulfide, Lawesson's Reagent or P_4S_{10} , preferably Lawesson's Reagent can be mentioned.

In processes (i) and (j) for preparing the compounds (1j) and (1k) by reacting the compound of formula (13) with the compound of formula (14a) or (14b), one or more solvents selected from ethanol and isopropyl alcohol can be used. Also, ordinary inorganic base, such as for example, one or more selected from a group consisting of lithium hydroxide, sodium hydroxide and potassium hydroxide, preferably lithium


hydroxide can be used in the process (k) wherein the compound of formula (1l) is hydrolyzed and then reacted with the compound of formula (15) to produce the compound of formula (1m). As the coupling agent, those mentioned for process (d) can be used.

In processes (l) and (m), any inert solvents, preferably one or more selected from dimethylformamide and dimethylacetamide are used as the solvent, and one or more selected from a group consisting of sodium hydride, sodium amide, sodium bis(trimethylsilyl)amide and potassium bis(trimethylsilyl)amide are used as the base. The deprotection reaction in process (m) may be carried out under the conventional reaction conditions for deprotection, preferably in the presence of Pd/C or Pd(OH)₂/C under hydrogen atmosphere. Further, the coupling agent used for the coupling of the compound of formula (1o) with the compound of formula (19) may be the same with those mentioned for process (d).


The compound of formula (3) used as the key intermediate in processes (a) to (c) for preparing the compound of formula (1) according to the present invention is itself a novel compound. Therefore, it is another object of the present invention to provide the compound of formula (3). As depicted in the following Reaction Schemes 14 to 16, the compound of formula (3) can be prepared by a process characterized in that a compound represented by the following formula (20) is reacted in a solvent in the presence of a coupling agent with a compound represented by the following formula (21); the compound of formula (20) is reacted in a solvent in the presence of dimethylformamide(DMF) with thionyl chloride to produce a compound represented by the following formula (20a) and then the compound of formula (20a) thus obtained is reacted in a solvent with the compound of formula (21); or a compound

represented by the following formula (3a) is oxidized in a solvent to produce a compound represented by the following formula (3b).


Reaction Scheme 14



Reaction Scheme 15



Reaction Scheme 16



in the above Reaction Schemes 14, 15 and 16

B, C and D are defined as previously described,

Q_a represents S or S=O.

In the above processes according to Reaction Scheme 14 to 16


for preparing the compound (3), any inert solvents, preferably one or more selected from dimethylformamide, dimethylacetamide, methylene chloride, tetrahydrofuran and 1,2-dichloroethane are used as the solvent. As the coupling agent in Reaction Scheme 14, a mixture of 1-hydroxybenzotriazole and one or more substances selected from a group consisting of carbodiimides such as dicyclohexylcarbodiimide(DCC), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide(EDC), etc. can be mentioned. Among them, a mixture of 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide(EDC) and 1-hydroxybenzotriazole hydrate is particularly preferred. The dimethylformamide in the process of Reaction Scheme 15 is used in a catalytic amount. Also, excess metachloroperbenzoic acid is preferably used as the oxidant in the process according to the Reaction Scheme 16. However, the coupling agent, oxidant, solvent, catalyst, etc. may be appropriately selected beyond those as mentioned above as far as the purpose of the reaction can be accomplished. And the reaction conditions including the amount of reactants, reaction temperature, reaction time, etc. can easily be determined by a person skilled in this art depending on the specific reactants.

Since the compound of formula (8) which is used as an intermediate for preparing the compound of formula (1d) in process (d) is also a novel compound like the compound of formula (3), it is another object of the present invention to provide the intermediate compound of formula (8). It can be obtained by hydrolyzing the compound of formula (7).

On the other hand, the starting materials used in the above processes can be prepared according to the specific processes described in the following Reaction Schemes 17 to 29.


First, the compound of formula (2) can be obtained through protection and halogenation as depicted in the following Reaction Scheme 17.

Reaction Scheme 17




The compound of formula (4) wherein A is 4-cyanobenzyl may be synthesized through protection, acetylation, coupling, deprotection and halogenation as depicted in the following Reaction Scheme 18. More frequently, the compound (4) is prepared by a process wherein an amine compound is reacted with dihydroxyacetone to produce a mercaptoimidazole derivative, which is then desulfurized and halogenated as depicted in the following Reaction Scheme 19. *J. Med. Chem.*, 33, 1312-1329, 1990 in which a similar reaction is explained in detail can be referred to for the specific reaction conditions.


Reaction Scheme 18



Reaction Scheme 19




The amine compound used in the above Reaction Scheme 19 wherein A represents 1-(benzyloxycarbonyl)piperidine-4-ylmethyl may be synthesized from 4-aminomethylpiperidine through protection, benzyloxy-carbonylation and deprotection as depicted in Reaction Scheme 20.

Reaction Scheme 20

in the above Reaction Scheme 20

CbzCl represents benzylchloroformate and has the same meaning through the present specification.

The compound of formula (5) may be synthesized through esterification, protection, reduction and halogenation as depicted in the following Reaction Scheme 21.

Reaction Scheme 21

in the above Reaction Scheme 21


DIBAL represents diisobutylaluminumhydride.

Also, in the above Reaction Scheme 21, the alcohol compound obtained before preparing the final chloride compound may be reduced according to the conventional manner and then reacted with thionyl


chloride to produce the compound of formula (2) wherein n_1 is 3.


The compound of formula (20) used as a starting material in preparing the intermediate of formula (3) may be prepared, for example, according to a process described in the following Reaction Scheme 22, a process starting from 1-naphthaldehyde. Particularly, the intermediate of formula (3) wherein D is 1-naphthyl can be conveniently synthesized according to the following reactions of Schemes 23 and 24.

Reaction Scheme 22





Reaction Scheme 23





Reaction Scheme 24

The compound of formula (11) used as a starting material in process (g) can be prepared by coupling a hydrochloride salt of glycinate derivative with a hydrochloride salt of 4-imidazoleacetic acid, as represented in the following Reaction Scheme 25. As the coupling agent, those mentioned in process (d) can be used. While, the compound of formula (13) used in process (i) may be prepared according to the procedure described in the following Reaction Scheme 26 in which the chloride derivative obtained in the process of Reaction Scheme 19 is used as a starting material.

Reaction Scheme 25

Reaction Scheme 26


The compounds (14a) and (14b) used in processes (i) and (j) can be prepared according to the following Reaction Schemes 27 and 28, respectively. First, the compound of formula 14a can be synthesized by reacting an aldehyde derivative with methyl dichloroacetate in the presence of potassium *t*-butoxide. The compound of formula (14b) wherein I is I' can be synthesized by reacting a ketone derivative with a dialkylcarbonate in the presence of sodium hydride, then by reacting the product thus obtained with sulfuryl chloride.

Reaction Scheme 27Reaction Scheme 28

Finally, the reactant of formula (17) in processes (l) and (m) wherein G represents 1-naphthyl and L represents N-methyl-N-(2-

methoxyethyl)amino may be prepared from 1-naphthaldehyde according to the following Reaction Scheme 29. The other compounds (17) having different substituents may also be prepared by referring to Reaction Scheme 29.

Reaction Scheme 29



The compound of formula (1) prepared according to the processes above shows an inhibitory activity against farnesyl transferase, and thus can be effectively used as an anti-cancer agent. Therefore, the present invention also provides a pharmaceutical composition comprising the novel compound of formula (1), as defined above, or a pharmaceutically acceptable salt or an isomer thereof as an active ingredient together with

a pharmaceutically acceptable carrier. Particularly, the compound of formula (1) can be used very effectively for treating cancer, restenosis, atherosclerosis and infections from hepatitis delta and related viruses.

When the active compound according to the present invention is used for clinical purpose, it is preferably administered in an amount ranging from 10_{mg} to 100_{mg} per kg of body weight a day. The total daily dosage may be administered in one time or over several times. However, the specific administration dosage for the patient can be varied with the specific compound used, body weight of the subject patient, sex, hygienic condition, diet, time or method of administration, excretion rate, mixing ratio of the agent, severity of the disease to be treated, etc.

The compound of the present invention may be administered in the form of injections or oral preparations. Injections, for example, sterilized aqueous or oily suspension for injection, can be prepared according to the known procedure using suitable dispersing agent, wetting agent, or suspending agent. Solvents which can be used for preparing injections include water, Ringer's fluid and NaCl solution, and also sterilized fixing oil may be conveniently used as the solvent or suspending media. Any non-stimulative fixing oil including mono-, di-glyceride may be used for this purpose. Fatty acid such as oleic acid may also be used for injections.

As the solid preparation for oral administration, capsules, tablets, pills, powders and granules, etc., preferably capsules and tablets can be mentioned. It is also desirable for tablets and pills to be formulated into enteric-coated preparation. The solid preparations may be prepared by mixing the active compound of formula (1) according to the present invention with at least one carrier selected from a group consisting of

inactive diluents such as sucrose, lactose, starch, etc., lubricants such as magnesium stearate, disintegrating agent and binding agent.

The present invention will be more specifically explained in the following examples. However, it should be understood that the following examples are intended to illustrate the present invention but not in any manner to limit the scope of the present invention. Processes for preparing the starting substances used for obtaining the compound of formula (1) will be also explained in detail in the following Preparations.

Preparation 1

Synthesis of 1-(3,4-methylenedioxybenzyl)-5-chloromethyl-1H-imidazole hydrochloride

1-1) 1-(3,4-Methylenedioxybenzyl)-5-hydroxymethyl-1H-imidazole

A modified method from *J. Med. Chem.*, 33, 1312-1329, 1990 was carried out using dihydroxyacetone dimer and piperonylamine as starting materials. 1.37g(10 mmol) of piperonylamine, 1.08g(5.5 mmol) of dihydroxyacetone dimer and 1.15g(11 mmol) of potassium thiocyanide were introduced to 10_{ml} of isopropyl alcohol, and then 2_{ml} of acetic acid was added thereto and the mixture was reacted at room temperature for 48 hours. The reaction mixture was filtered and the residual solid thus obtained was washed with 5_{ml} of isopropyl alcohol(x2) and with 5_{ml} of water(x2). The filtered solid was introduced into 12.5_{ml} of 10% aqueous nitric acid solution and the resulting solution was cooled down to 0°C. After 10_{mg} of sodium nitrite was added portionwise to the reaction solution, the mixture was reacted at room temperature for 1 hour. The aqueous solution was washed with 10_{ml} of ethyl acetate, basified, and then recrystallized to obtain 1.16g (Yield 50%) of the title compound.

^1H NMR(CDCl_3) δ 4.45(s, 2H), 5.13(s, 2H), 5.97(s, 2H), 6.70(m, 2H), 6.78 (d, 1H), 6.95(s, 1H), 7.45(s, 1H)

FAB 233 (M+H), $\text{C}_{12}\text{H}_{12}\text{N}_2\text{O}_3$

1-2) 1-(3,4-Methylenedioxybenzyl)-5-chloromethyl-1H-imidazole hydrochloride

233_{mg}(1 mmol) of the compound prepared in Preparation 1-1) was dissolved in 3_{ml} of chloroform and then 355_{mg}(3 mmol) of thionyl chloride was slowly added dropwise thereto at 0°C. After stirring for 2 hours, the solvent was removed by distillation under reduced pressure and the remained hydrochloride was eliminated to obtain the title compound in a yield of 95%. The product thus obtained was directly used in the next reaction without purification.

Preparation 2

Synthesis of 1-(naphthalen-1-ylmethyl)-5-chloromethyl-1H-imidazole hydrochloride

2-1) 1-(Naphthalen-1-ylmethyl)-5-hydroxymethyl-1H-imidazole

The title compound was obtained in a yield of 65% according to the same procedure as Preparation 1-1) except that dihydroxyacetone dimer and (naphthalen-1-ylmethyl)amine were used as starting materials.

^1H NMR(CDCl_3) δ 4.44(s, 2H), 5.42(s, 2H), 6.78(d, 1H), 6.85(s, 1H), 7.25(m, 1H), 7.35(s, 1H), 7.43(m, 2H), 7.65(d, 1H), 7.68(d, 1H), 8.02(d, 1H)

FAB 239 (M+H), $\text{C}_{15}\text{H}_{14}\text{N}_2\text{O}$

2-2) 1-(Naphthalen-1-ylmethyl)-5-chloromethyl-1H-imidazole hydrochloride

The title compound was obtained in a yield of 90% according to the same procedure as Preparation 1-2) using the compound prepared in

Preparation 2-1). The product thus obtained was directly used in the next reaction without purification.

Preparation 3: Synthesis of 1-((R)- α -methylbenzyl)-5-chloromethyl-1H-imidazole hydrochloride

3-1) 1-((R)- α -methylbenzyl)-5-hydroxymethyl-1H-imidazole

The title compound was obtained in a yield of 60% according to the same procedure as Preparation 1-1) except that dihydroxyacetone dimer and (R)-(+)- α -methylbenzylamine were used as starting materials.

^1H NMR(CDCl_3) δ 1.73 (d, 3H), 4.28 (s, 1H), 4.43(d, 1H), 5.60(m, 1H), 6.75(s, 1H), 7.04(d, 2H), 7.23(m, 3H), 7.42(s, 1H)

FAB 203 (M+H), $\text{C}_{12}\text{H}_{14}\text{N}_2\text{O}$

3-2) 1-((R)- α -methylbenzyl)-5-chloromethyl-1H-imidazole hydrochloride

The title compound was obtained in a yield of 90% according to the same procedure as Preparation 1-2) using the compound prepared in Preparation 3-1). The product thus obtained was directly used in the next reaction without purification.

Preparation 4: Synthesis of 1-((S)- α -methylbenzyl)-5-chloromethyl-1H-imidazole hydrochloride

4-1) 1-((S)- α -methylbenzyl)-5-hydroxymethyl-1H-imidazole

The title compound was obtained in a yield of 55% according to the same procedure as Preparation 1-1) except that dihydroxyacetone dimer and (S)-(+)- α -methylbenzylamine were used as starting materials.

^1H NMR(CDCl_3) δ 1.73(d, 3H), 4.28(s, 1H), 4.43(d, 1H), 5.60(m, 1H), 6.75(s, 1H), 7.04(d, 2H), 7.23(m, 3H), 7.42(s, 1H)

FAB 203 (M+H), $\text{C}_{12}\text{H}_{14}\text{N}_2\text{O}$

4-2) 1-((S)- α -methylbenzyl)-5-chloromethyl-1H-imidazole hydrochloride

The title compound was obtained in a yield of 94% according to the same procedure as Preparation 1-2) using the compound prepared in Preparation 4-1). The product thus obtained was directly used in the next reaction without purification.

Preparation 5: Synthesis of 1-phenethyl-5-chloromethyl-1H-imidazole hydrochloride**5-1) 1-Phenethyl-5-hydroxymethyl-1H-imidazole**

The title compound was obtained in a yield of 70% according to the same procedure as Preparation 1-1) except that dihydroxyacetone dimer and phenethylamine were used as starting materials.

^1H NMR(CDCl_3) δ 3.08(t, 2H), 4.27(t, 2H), 4.47(s, 2H), 6.89(s, 1H), 7.05(d, 2H), 7.26(m, 3H), 7.44(s, 1H)

FAB 203 (M+H), $\text{C}_{12}\text{H}_{14}\text{N}_2\text{O}$

5-2) 1-Phenethyl-5-chloromethyl-1H-imidazole hydrochloride

The title compound was obtained in a yield of 90% according to the same procedure as Preparation 1-2) using the compound prepared in Preparation 5-1). The product thus obtained was directly used in the next reaction without purification.

Preparation 6: Synthesis of 3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole**6-1) 3-(Naphthalen-1-yl)-acrylic acid ethylester**

22.4g(0.10 mol) of triethylphosphonoacetate was dissolved in 500 ml of acetonitrile and 30.4g(0.2 mol) of 1,8-diazabicyclo[5.4.0]undec-7-

ene(1,5-5)(DBU) was slowly added thereto. To this solution was slowly added 15.6g(0.10 mol) of 1-naphthaldehyde dissolved in 20_{ml} of tetrahydrofuran and the mixture was stirred for 8 hours. The organic solvent was removed by distillation under reduced pressure. The resulting residue was dissolved in ethyl acetate, washed twice with water, dried over magnesium sulfate, concentrated and then subjected to silica gel column chromatography (eluent: n-hexane/ethyl acetate=95/5, v/v) to obtain 20.3g(0.090 mol, Yield 90%) of the title compound.

¹H NMR(CDCl₃) δ 1.33(t, 3H), 4.10(q, 2H), 6.75(q, 1H), 7.50(m, 3H), 7.73(d, 1H), 7.85(m, 2H), 8.10(d, 1H), 8.21(d, 1H)

FAB 227 (M+H)

6-2) 3-(Ethoxycarbonyl)-4-(naphthalen-1-yl)-1H-pyrrole

4.3g(18.9 mmol) of 3-(naphthalen-1-yl)-acrylic acid ethylester prepared in Preparation 6-1) and 3.68g(18.9 mmol) of tosylmethylisocyanide were dissolved in 100_{ml} of tetrahydrofuran. 2.55g(22.7 mmol) of potassium t-butoxide dissolved in 100_{ml} of tetrahydrofuran was slowly added thereto and the mixture was refluxed for 30 minutes. 100_{ml} of water was added to the reaction solution to stop the reaction and the solvent was removed under reduced pressure. The reaction solution was extracted with diethylether, washed with aqueous sodium chloride solution and then dried over magnesium sulfate. The solvent was removed under reduced pressure and the resulting residue was subjected to silica gel column chromatography(eluent: ethyl acetate/n-hexane=1/3, v/v) to obtain 3.85g(14.5 mmol, Yield 77%) of the title compound.

¹H NMR(CDCl₃) δ 1.27(t, 3H), 4.07(q, 2H), 6.76(s, 1H), 7.28-7.47(m, 5H), 7.59(s, 1H), 7.82(m, 2H), 9.99(s, 1H)

FAB 266 (M+H)

6-3) 3-Hydroxycarbonyl-4-(naphthalen-1-yl)-1H-pyrrole

2.64g(10 mmol) of the compound prepared in Preparation 6-2) was dissolved in 50_{ml} of 50% ethanol and 2.24g(40 mmol) of potassium hydroxide was added thereto. The reaction mixture was refluxed for 7 hours, cooled down to room temperature, adjusted to pH 4-5, extracted with ethyl acetate, dried over sodium sulfate. The solvent was removed under reduced pressure to obtain 1.90g(8.1 mmol, Yield 81%) of the title compound. The product thus obtained was directly used in the next reaction without purification.

¹H NMR(CDCl₃) δ 6.60(s, 1H), 7.32-7.49(m, 5H), 7.54(s, 1H), 7.84(m, 2H), 9.92(s, 1H)

FAB 238 (M+H)

6-4) 3-[N-(2-Methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole

234_{mg}(1 mmol) of the compound prepared in Preparation 6-3) was dissolved in 2_{ml} of dimethylformamide, and then 230_{mg}(1.2 mmol) of EDC, 101_{mg}(1 mmol) of triethylamine and 162_{mg}(1.2 mmol) of HOBT were added thereto. The resulting mixture was stirred at 0°C for 5 minutes. To the reaction solution was added 124_{mg}(1 mmol) of N-(2-methoxyethyl)-N-methylamine hydrochloride, which was then stirred at room temperature for 5 hours. The solvent was removed under reduced pressure and then 10_{ml} of saturated potassium carbonate solution was added to the residue. The resulting solution was extracted with 20 _{ml} of ethyl acetate, washed with 10_{ml} of 1N aqueous hydrochloric acid solution, washed with aqueous sodium chloride solution and water, dried over sodium sulfate and concentrated to give 246_{mg}(0.79 mmol, Yield 79%) of the title compound.

¹H NMR(CDCl₃) δ 2.46(s, 2H), 2.80-3.40(m, 8H), 3.40(s, 1H), 6.80(s, 1H), 7.00(s, 1H), 7.42(m, 4H), 7.73(d, 1H), 7.81(d, 1H), 8.17(d, 1H), 10.66 (s, 1H)

FAB 309 (M+H)

Preparation 7: Synthesis of 3-(morpholin-4-yl)carbonyl-4-(naphthalen-1-yl)-1H-pyrrole

234_{mg}(1 mmol) of the compound prepared in Preparation 6-3) was dissolved in 2_{ml} of dimethylformamide, and then 230_{mg}(1.2 mmol) of EDC and 162_{mg}(1.2 mmol) of HOBT were added thereto. The resulting mixture was stirred at 0°C for 5 minutes. To the reaction solution was added 87_{mg}(1 mmol) of morpholine, which was then stirred at room temperature for 5 hours. The solvent was removed under reduced pressure and then 10_{ml} of saturated potassium carbonate solution was added to the residue. The resulting solution was extracted with ethyl acetate, washed with 10_{ml} of 1N aqueous hydrochloric acid solution, washed with aqueous sodium chloride solution and water, dried over sodium sulfate and concentrated to give 243_{mg}(0.8 mmol, Yield 80%) of the title compound.

¹H NMR(CDCl₃) δ 2.13-3.52(br, 8H), 6.54(s, 1H), 7.31-7.51(m, 5H), 7.53 (s, 1H), 7.81(m, 2H), 9.93(s, 1H)

FAB 307 (M+H)

Preparation 8: Synthesis of 3-(4-methylpiperazin-1-yl)carbonyl-4-(naphthalen-1-yl)-1H-pyrrole

The title compound was obtained in a yield of 75% according to the same procedure as Preparation 6-4) except that the compound prepared in Preparation 6-3) and 4-methylpiperazine were used.

¹H NMR(CDCl₃) δ 1.15(br, 2H), 1.87(br, 2H), 1.92(s, 3H), 2.96(br, 2H), 3.41(br, 2H), 6.83(s, 1H), 7.09(s, 1H), 7.36-7.42(m, 4H), 7.73(d, 1H), 7.75 (d, 1H), 8.10(d, 1H), 10.52(s, 1H)

FAB (M+H): 320

Preparation 9: Synthesis of 3-{N-[2-(N,N-dimethylamino)ethyl]-N-methyl}carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole

The title compound was obtained in a yield of 82% according to the same procedure as Preparation 6-4) except that the compound prepared in Preparation 6-3) and N,N,N'-trimethyl-ethylenediamine were used.

^1H NMR(CDCl_3) δ 1.89(br, 3H), 2.18(br, 4H), 2.44(br, 2H), 2.75(s, 1H), 2.98(br, 1H), 3.36(br, 2H), 6.84(s, 1H), 7.07(s, 1H), 7.38-7.43(m, 4H), 7.74 (d, 1H), 7.83(d, 1H), 8.13(b, 1H), 10.14(br, 1H)

FAB (M+H): 322

Preparation 10: Synthesis of 4-(naphthalen-1-yl)-3-(thiomorpholin-4-yl) carbonyl-1H-pyrrole

234_{mg}(1 mmol) of the compound prepared in Preparation 6-3) was dissolved in 2_{ml} of dimethylformamide, and then 230_{mg}(1.2 mmol) of EDC and 162_{mg}(1.2 mmol) of HOBT were added thereto. The resulting mixture was stirred at 0°C for 5 minutes. To the reaction solution was added 87_{mg}(1 mmol) of thiomorpholine, which was then stirred at room temperature for 5 hours. The solvent was removed under reduced pressure and then 10_{ml} of saturated potassium carbonate solution was added to the residue. The resulting solution was extracted with ethyl acetate, washed with 10_{ml} of 1N aqueous hydrochloric acid solution, washed with saturated sodium chloride solution and water, dried over sodium sulfate and concentrated to give 258_{mg}(0.8 mmol, Yield 80%) of the title compound.

^1H NMR(CDCl_3) δ 1.35 (br, 2H), 2.14 (br, 2H), 3.21(br, 2H),

3.41(br, 2H), 6.91 (s, 1H), 7.21 (s, 1H), 7.31-7.51 (m, 4H), 7.80 (d, 1H), 7.87 (d, 1H), 8.11(d, 1H), 10.69(s, 1H)

FAB 323 (M+H)

Preparation 11: Synthesis of 3-(1,1-dioxothiomorpholin-4-yl)carbonyl-4-(naphthalen-1-yl)-1H-pyrrole

323mg(1 mmol) of the compound prepared in Preparation 10 was dissolved in 5_{ml} of dichloromethane, 430mg(1.5 mmol) of 60% 3-chloroperbenzoic acid(MCPBA) was added thereto, and then the mixture was stirred at room temperature for 1 hour. 3_{ml} of 10% sodium thiosulfite was added to the mixture in order to remove the excess 3-chloroperbenzoic acid and the resulting mixture was stirred at room temperature for 30 minutes. After adding 10_{ml} of saturated potassium carbonate solution thereto, the mixture was extracted with dichloromethane, washed with saturated sodium chloride solution and water, dried over sodium sulfate and concentrated to give 264mg(0.75 mmol, Yield 75%) of the title compound.

¹H NMR(CDCl₃) δ 1.50-2.30(br, 4H), 3.65 (br, 4H), 6.92 (s, 1H), 7.20 (s, 1H), 7.32-7.54 (m, 4H), 7.81 (d, 1H), 7.88 (d, 1H), 8.12(d, 1H), 10.69(s, 1H).

FAB 355 (M+H)

Preparation 12: Synthesis of 3-[N-(2-methylthioethyl)-N-methyl] carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole

234_{mg}(1 mmol) of the compound prepared in Preparation 6-3) was dissolved in 2_{ml} of dimethylformamide, and then 230_{mg}(1.2 mmol) of EDC, 101_{mg}(1 mmol) of triethylamine and 162_{mg}(1.2 mmol) of HOBT were added thereto. The resulting mixture was stirred at 0°C for 5

minutes. To the reaction solution was added 140_{mg}(1 mmol) of N-(2-methylthioethyl)-N-methylamine hydrochloride, which was then stirred at room temperature for 5 hours. The solvent was removed under reduced pressure and then 10_{mℓ} of saturated potassium carbonate solution was added to the residue. The resulting solution was extracted with 20_{mℓ} of ethyl acetate, washed with 10_{mℓ} of 1N aqueous hydrochloric acid solution, washed with saturated sodium chloride solution and water, dried over sodium sulfate and concentrated to give 243_{mg}(0.75 mmol, Yield 75%) of the title compound.

¹H NMR(CDCl₃) δ 1.98 (s, 3H), 2.13 (br, 2H), 2.46 (br, 2H), 2.65 (br, 1H), 2.95 (br, 1H), 3.29 (br, 1H), 6.81 (s, 1H), 7.02 (s, 1H), 7.43 (m, 4H), 7.72 (d, 1H), 7.82 (d, 1H), 8.18 (d, 1H), 10.65 (s, 1H)

FAB 325 (M+H)

Preparation 13: Synthesis of 3-hydroxycarbonyl-5-methyl-4-(naphthalen-1-yl)-1H-pyrrole

13-1) 3-ethoxycarbonyl-5-methyl-4-(naphthalen-1-yl)-1H-pyrrole

4.3g(18.9 mmol) of 3-(naphthalen-1-yl)-acrylic acid ethylester prepared in Preparation 6-1) and 3.95g(18.9 mmol) of α-methyltosylmethylisocyanide disclosed in A.M. van Leusen, et al., *Tetrahedron Letter*, 1975, 40, 3487 were dissolved in 100_{mℓ} of tetrahydrofuran. 2.55g(22.7 mmol) of potassium t-butoxide dissolved in 100_{mℓ} of tetrahydrofuran was slowly added thereto, which was then refluxed for 30 minutes. To the reaction solution was added 100_{mℓ} of water to stop the reaction and the solvent was removed under reduced pressure. The residue was extracted with diethylether, washed with saturated sodium chloride solution and dried over magnesium sulfate. The solvent was removed under reduced pressure and the residue was subjected to silica gel column chromatography using a solvent mixture of

ethyl acetate/n-hexane(1/3, v/v) as an eluent to give 3.50g(12.5 mmol, Yield 66%) of the title compound.

FAB 280 (M+H)

13-2) 3-Hydroxycarbonyl-5-methyl-4-(naphthalen-1-yl)-1H-pyrrole

2.80g(10 mmol) of the compound prepared in Preparation 13-1) was dissolved in 50_{ml} of 50% ethanol, 2.24g(40 mmol) of potassium hydroxide was added thereto, and the mixture was refluxed for 7 hours. The reaction solution was cooled down to room temperature, adjusted to pH 4-5, extracted with ethyl acetate and dried over sodium sulfate. The solvent was eliminated under reduced pressure to obtain 2.02g(8.1 mmol, Yield 81%) of the title compound.

FAB 252 (M+H)

Preparation 14: Synthesis of 5-methyl-3-(morpholin-4-yl)carbonyl-4-(naphthalen-1-yl)-1H-pyrrole

248_{mg}(1 mmol) of the compound prepared in Preparation 13-2) was dissolved in 2_{ml} of dimethylformamide, and then 230_{mg}(1.2 mmol) of EDC and 162_{mg}(1.2 mmol) of HOBT were added thereto. The resulting mixture was stirred at 0°C for 5 minutes. To the reaction solution was added 87_{mg}(1 mmol) of morpholine, which was then stirred at room temperature for 5 hours. The solvent was removed under reduced pressure and then 10_{ml} of saturated potassium carbonate solution was added to the residue. The resulting solution was extracted with ethyl acetate, washed with 10_{ml} of 1N aqueous hydrochloric acid solution, washed with saturated sodium chloride solution and water, dried over sodium sulfate and concentrated to give 224_{mg}(0.7 mmol, Yield 70%) of the title compound.

¹H NMR(CDCl₃) δ 2.12 (s, 3H), 2.80-3.40 (br, 8H), 7.01 (s,

1H), 7.30- 7.50 (m, 4H), 7.75-7.95 (m, 3H), 10.60 (br, 1H)

FAB 321 (M+H)

Example 1: Synthesis of 3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-1-[1-(3,4-methylenedioxybenzyl)-1H-imidazol-5-ylmethyl]-4-(naphthalen-1-yl)-1H-pyrrole(1)

62_{mg}(0.2 mmol) of the compound prepared in Preparation 6 was dissolved in 2_{ml} of dimethylformamide, 26.4_{mg}(0.66 mmol) of sodium hydride(60%) was added thereto at 0°C and then the mixture was stirred for 5 minutes. To the mixture was added 50_{mg}(2.2 mmol) of the compound prepared in Preparation 1 and the whole mixture was stirred at room temperature for 3 hours. The solvent was removed by distillation under reduced pressure and 3_{ml} of water was added to the residue. The mixture was then extracted twice with 10_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated, and subjected to silica gel column chromatography (eluent: dichloromethane/methanol=95/5, v/v) to obtain 78_{mg}(Yield 75%) of the title compound.

¹H NMR(CDCl₃) δ 2.40(m, 2H), 2.72(m, 1H), 2.91(s, 3H), 3.09(m, 2H), 3.32(br, 1H), 4.09(br, 1H), 4.89(s, 2H), 4.95(s, 2H), 5.89(s, 2H), 6.45(s, 1H), 6.62(d, 1H), 6.63(s, 1H), 6.70(d, 1H), 7.0(s, 1H), 7.16(s, 1H), 7.31(t, 1H), 7.41(m, 3H), 7.66(s, 1H), 7.73(d, 1H), 7.81(d, 1H), 8.03(d, 1H)

FAB (M+H) 523, C₃₁H₃₀N₄O₂

Example 2: Synthesis of 1-[1-(3,4-methylenedioxybenzyl)-1H-imidazol-5-ylmethyl]-3-(morpholin-4-yl)carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(2)

62_{mg}(0.2 mmol) of the compound prepared in Preparation 7 was dissolved in 2_{ml} of dimethylformamide, 26.4_{mg}(0.66 mmol) of sodium

hydride(60%) was added thereto at 0°C and then the mixture was stirred for 5 minutes. To the mixture was added 50_{mg}(2.2 mmol) of the compound prepared in Preparation 1 and the whole mixture was stirred at room temperature for 3 hours. The solvent was removed by distillation under reduced pressure and 3_{ml} of water was added to the residue. The mixture was then extracted twice with 10_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated, and subjected to silica gel column chromatography (eluent: dichloromethane/methanol=95/5, v/v) to obtain 70_{mg}(Yield 67%) of the title compound.

¹H NMR(CDCl₃) δ 2.36(br, 2H), 3.06(br, 4H), 3.33(br, 2H), 5.23(s, 2H), 5.33(s, 2H), 5.96(s, 2H), 6.65(s, 1H), 6.70-6.85(m, 3H), 7.18-7.50(m, 7H), 7.79(d, 1H), 7.81(d, 1H), 7.94(d, 1H)

FAB (M+H) 521, C₃₁H₂₈N₄O₄

Example 3: Synthesis of 1-[1-(3,4-methylenedioxybenzyl)-1H-imidazol-5-ylmethyl]-3-(4-methylpiperazin-1-yl)carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(3)

64_{mg}(0.2 mmol) of the compound prepared in Preparation 8 was dissolved in 2_{ml} of dimethylformamide, 26.4_{mg}(0.66 mmol) of sodium hydride(60%) was added thereto at 0°C and then the mixture was stirred for 5 minutes. To the mixture was added 50_{mg}(2.2 mmol) of the compound prepared in Preparation 1 and the whole mixture was stirred at room temperature for 3 hours. The solvent was removed by distillation under reduced pressure and 3_{ml} of water was added to the residue. The mixture was then extracted twice with 10_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated, and subjected to silica gel column chromatography (eluent: dichloromethane/methanol=90/10, v/v) to obtain 73_{mg}(Yield 67%) of the title compound.

¹H NMR(CDCl₃) δ 2.18(s, 3H), 2.30-2.60(br, 4H), 3.10-3.30(br,

4H), 4.98 (s, 2H), 5.05(s, 2H), 5.95(s, 2H), 6.44(s, 1H), 6.53(d, 1H), 6.70(d, 1H), 6.73(d, 1H), 7.14(d, 1H), 7.20-7.40(m, 3H), 7.50(m, 3H), 7.81(d, 1H), 7.83(d, 1H), 7.88(d, 1H)

FAB (M+H) 534, C₃₂H₃₁N₅O₃

Example 4: Synthesis of 3-[N-[2-(N,N-dimethylamino)ethyl]-N-methyl]carbamoyl-1-[1-(3,4-methylenedioxybenzyl)-1H-imidazol-5-ylmethyl]-4-(naphthalen-1-yl)-1H-pyrrole(4)

64_{mg}(0.2 mmol) of the compound prepared in Preparation 9 was dissolved in 2_{ml} of dimethylformamide, 26.4_{mg}(0.66 mmol) of sodium hydride(60%) was added thereto at 0°C and then the mixture was stirred for 5 minutes. To the mixture was added 50_{mg}(2.2 mmol) of the compound prepared in Preparation 1 and the whole mixture was stirred at room temperature for 3 hours. The solvent was removed by distillation under reduced pressure and 3_{ml} of water was added to the residue. The mixture was then extracted twice with 10_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated, and subjected to silica gel column chromatography (eluent: dichloromethane/methanol=90/10, v/v) to obtain 78_{mg}(Yield 71%) of the title compound.

¹H NMR(CDCl₃) δ 1.87(m, 1H), 2.01(m, 2H), 2.14(br, 6H), 2.36(br, 2H), 2.50-3.00(br, 1H), 3.29(br, 2H), 4.87(s, 2H), 4.95(s, 2H), 5.89(s, 2H), 6.45 (s, 1H), 6.50(d, 1H), 6.63(d, 1H), 6.72(d, 1H), 7.00(s, 1H), 7.18(s, 1H), 7.31(br, 1H), 7.35-7.47(m, 3H), 7.54(s, 1H), 7.73(d, 1H), 7.81(d, 1H), 8.01(br, 1H)

FAB (M+H) 536, C₃₂H₃₃N₅O₃

Example 5: Synthesis of 3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-naphthalen-1-yl)-1-[1-naphthalen-1-ylmethyl)-1H-imidazol-5-ylmethyl]-1H-pyrrole(5)

62_{mg}(0.2 mmol) of the compound prepared in Preparation 6 was dissolved in 2_{ml} of dimethylformamide, 26.4_{mg}(0.66 mmol) of sodium hydride(60%) was added thereto at 0°C and then the mixture was stirred for 5 minutes. To the mixture was added 58_{mg}(2.2 mmol) of the compound prepared in Preparation 2 and the whole mixture was stirred at room temperature for 3 hours. The solvent was removed by distillation under reduced pressure and 3_{ml} of water was added to the residue. The mixture was then extracted twice with 10_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated, and subjected to silica gel column chromatography (eluent: dichloromethane/methanol=95/5, v/v) to obtain 79_{mg}(Yield 75%) of the title compound.

¹H NMR(CDCl₃) δ 2.37(br, 2H), 2.72(br, 1H), 2.99(br, 3H), 3.00(br, 2H), 3.31(br, 1H), 3.71(br, 1H), 5.06(s, 2H), 5.48(s, 2H), 6.62(d, 1H), 6.91(d, 1H), 7.03(d, 1H), 7.27(d, 2H), 7.28-7.55(m, 6H), 7.58(s, 1H), 7.69(d, 1H), 7.75(d, 1H), 7.81(d, 2H), 7.87(d, 1H), 8.00(d, 1H)

FAB (M+H) 529, C₃₄H₃₂N₄O₂

Example 6: Synthesis of 3-(morpholin-4-yl)carbonyl-4-(naphthalen-1-yl)-1-[1-naphthalen-1-ylmethyl]-1H-imidazol-5-ylmethyl]-1H-pyrrole(6)

62_{mg}(0.2 mmol) of the compound prepared in Preparation 7 was dissolved in 2_{ml} of dimethylformamide, 26.4_{mg}(0.66 mmol) of sodium hydride(60%) was added thereto at 0°C and then the mixture was stirred for 5 minutes. To the mixture was added 58_{mg}(2.2 mmol) of the compound prepared in Preparation 2 and the whole mixture was stirred at room temperature for 3 hours. The solvent was removed by distillation under reduced pressure and 3_{ml} of water was added to the residue. The mixture was then extracted twice with 10_{ml} of ethyl

acetate, dried over anhydrous sodium sulfate, concentrated, and subjected to silica gel column chromatography (eluent: dichloromethane/methanol=95/5, v/v) to obtain 76_{mg} (Yield 72%) of the title compound.

¹H NMR(CDCl₃) δ 2.38(br, 2H), 3.06(br, 4H), 3.30(br, 2H), 4.99(s, 2H), 5.42(s, 2H), 6.58(d, 1H), 6.80(d, 1H), 7.00(s, 1H), 7.17(d, 1H), 7.25(s, 1H), 7.26-7.54(m, 6H), 7.69(d, 1H), 7.71-7.81(m, 3H), 7.85(d, 1H), 7.91(d, 1H)

FAB (M+H) 527, C₃₄H₃₀N₄O₂

Example 7: Synthesis of 3-(4-methylpiperazin-1-yl)carbonyl-4-(naphthalen-1-yl)-1-[1-(naphthalen-1-ylmethyl)-1H-imidazol-5-ylmethyl]-1H-pyrrole(7)

62_{mg} (0.2 mmol) of the compound prepared in Preparation 8 was dissolved in 2_{ml} of dimethylformamide, 26.4_{mg} (0.66 mmol) of sodium hydride (60%) was added thereto at 0°C and then the mixture was stirred for 5 minutes. To the mixture was added 58_{mg} (2.2 mmol) of the compound prepared in Preparation 2 and the whole mixture was stirred at room temperature for 3 hours. The solvent was removed by distillation under reduced pressure and 3_{ml} of water was added to the residue. The mixture was then extracted twice with 10_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated, and subjected to silica gel column chromatography (eluent: dichloromethane/methanol=90/10, v/v) to obtain 75_{mg} (Yield 69%) of the title compound.

¹H NMR(CDCl₃) δ 1.07(br, 2H), 1.77(d, 2H), 1.85(s, 3H), 2.84(br, 2H), 3.27(br, 2H), 4.99(s, 2H), 5.42(s, 2H), 6.58(d, 1H), 6.80(d, 1H), 7.01(d, 1H), 7.16(d, 1H), 7.25(s, 1H), 7.31-7.60(m, 6H), 7.68(d, 1H), 7.69-7.83(m, 3H), 7.85(d, 1H), 7.94(d, 1H)

FAB (M+H) 540, C₃₅H₃₃N₅O

Example 8: Synthesis of 3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-1-[1-((R)- α -methylbenzyl)-1H-imidazol-5-ylmethyl]-4-(naphthalen-1-yl)-1H-pyrrole(8)

62_{mg}(0.2 mmol) of the compound prepared in Preparation 6 was dissolved in 2_{ml} of dimethylformamide, 26.4_{mg}(0.66 mmol) of sodium hydride(60%) was added thereto at 0°C and then the mixture was stirred for 5 minutes. To the mixture was added 50_{mg}(2.2 mmol) of the compound prepared in Preparation 3 and the whole mixture was stirred at room temperature for 3 hours. The solvent was removed by distillation under reduced pressure and 3_{ml} of water was added to the residue. The mixture was then extracted twice with 10_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated, and subjected to silica gel column chromatography (eluent: dichloromethane/methanol=95/5, v/v) to obtain 70_{mg}(Yield 71%) of the title compound.

¹H NMR(CDCl₃) δ 1.78(d, 3H), 2.28(s, 1H), 2.40(br, 2H), 3.02(br, 3H), 3.09(br, 2H), 3.32(br, 2H), 4.71(d, 2H), 4.92(d, 2H), 5.12(q, 1H), 6.59(d, 1H), 7.00(m, 3H), 7.18(s, 1H), 7.20-7.39(m, 4H), 7.40-7.62(m, 3H), 7.74(m, 2H), 7.82(d, 1H), 8.04(d, 1H)

FAB (M+H) 493, C₃₁H₃₂N₄O₂

Example 9: Synthesis of 1-[1-((R)- α -methylbenzyl)-1H-imidazol-5-ylmethyl]-3-(morpholin-4-yl)carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(9)

62_{mg}(0.2 mmol) of the compound prepared in Preparation 7 was dissolved in 2_{ml} of dimethylformamide, 26.4_{mg}(0.66 mmol) of sodium hydride(60%) was added thereto at 0°C and then the mixture was stirred for 5 minutes. To the mixture was added 50_{mg}(2.2 mmol) of the compound prepared in Preparation 3 and the whole mixture was stirred at room temperature for 3 hours. The solvent was removed by

distillation under reduced pressure and 3_{ml} of water was added to the residue. The mixture was then extracted twice with 10_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated, and subjected to silica gel column chromatography (eluent: dichloromethane/methanol=95/5, v/v) to obtain 71_{mg} (Yield 72%) of the title compound.

¹H NMR(CDCl₃) δ 1.81(d, 3H), 2.28(br, 2H), 3.06(br, 4H), 3.29(br, 2H), 4.65(d, 1H), 4.96(d, 1H), 5.14(q, 1H), 6.62(d, 1H), 7.01(d, 2H), 7.04(s, 1H), 7.20(s, 1H), 7.23-7.36(m, 5H), 7.39-7.50(m, 3H), 7.76(s, 1H), 7.78(d, 1H), 7.84(d, 1H), 8.00(d, 1H)

FAB (M+H) 491, C₃₁H₃₀N₄O₂

Example 10: Synthesis of 1-[1-((R)-α-methylbenzyl)-1H-imidazol-5-ylmethyl]-3-(4-methylpiperazin-1-yl)carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(10)

64_{mg} (0.2 mmol) of the compound prepared in Preparation 8 was dissolved in 2_{ml} of dimethylformamide, 26.4_{mg} (0.66 mmol) of sodium hydride (60%) was added thereto at 0°C and then the mixture was stirred for 5 minutes. To the mixture was added 50_{mg} (2.2 mmol) of the compound prepared in Preparation 3 and the whole mixture was stirred at room temperature for 3 hours. The solvent was removed by distillation under reduced pressure and 3_{ml} of water was added to the residue. The mixture was then extracted twice with 10_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated, and subjected to silica gel column chromatography (eluent: dichloromethane/methanol=90/10, v/v) to obtain 73_{mg} (Yield 73%) of the title compound.

¹H NMR(CDCl₃) δ 1.09(br, 2H), 1.77(d, 3H), 1.83(s, 3H), 1.70-1.90(br, 2H), 2.90(br, 2H), 3.31(br, 2H), 4.73(d, 1H), 4.92(d, 1H), 5.14(q, 1H), 6.60(d, 1H), 7.01(m, 3H), 7.17(s, 1H), 7.20-7.35(m, 4H), 7.45(m, 3H), 7.73(m, 2H), 7.80(d, 1H), 8.00(d, 1H)

FAB (M+H) 504, $C_{32}H_{33}N_5O$

Example 11: Synthesis of 3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-1-[1-((S)- α -methylbenzyl)-1H-imidazol-5-ylmethyl]-4-(naphthalen-1-yl)-1H-pyrrole(11)

62_{mg}(0.2 mmol) of the compound prepared in Preparation 6 was dissolved in 2_{ml} of dimethylformamide, 26.4_{mg}(0.66 mmol) of sodium hydride(60%) was added thereto at 0°C and then the mixture was stirred for 5 minutes. To the mixture was added 50_{mg}(2.2 mmol) of the compound prepared in Preparation 4 and the whole mixture was stirred at room temperature for 3 hours. The solvent was removed by distillation under reduced pressure and 3_{ml} of water was added to the residue. The mixture was then extracted twice with 10_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated, and subjected to silica gel column chromatography (eluent: dichloromethane/methanol=95/5, v/v) to obtain 75_{mg}(Yield 75%) of the title compound.

¹H NMR(CDCl₃) δ 1.78(d, 3H), 2.28(s, 1H), 2.40(br, 2H), 3.02(br, 3H), 3.09(br, 2H), 3.32(br, 2H), 4.72(d, 2H), 4.93(d, 2H), 5.12(q, 1H), 6.59(d, 1H), 7.00(m, 3H), 7.18(s, 1H), 7.20-7.39(m, 4H), 7.40-7.62(m, 3H), 7.74(m, 2H), 7.82(d, 1H), 8.04(d, 1H)

FAB (M+H) 493, $C_{31}H_{32}N_4O_2$

Example 12: Synthesis of 1-[1-((S)- α -methylbenzyl)-1H-imidazol-5-ylmethyl]-3-(morpholin-4-yl)carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(12)

62_{mg}(0.2 mmol) of the compound prepared in Preparation 7 was dissolved in 2_{ml} of dimethylformamide, 26.4_{mg}(0.66 mmol) of sodium hydride(60%) was added thereto at 0°C and then the mixture was stirred for 5 minutes. To the mixture was added 50_{mg}(2.2 mmol) of the

compound prepared in Preparation 4 and the whole mixture was stirred at room temperature for 3 hours. The solvent was removed by distillation under reduced pressure and 3_{mℓ} of water was added to the residue. The mixture was then extracted twice with 10_{mℓ} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated, and subjected to silica gel column chromatography (eluent: dichloromethane/methanol=95/5, v/v) to obtain 73_{mg} (Yield 73%) of the title compound.

¹H NMR(CDCl₃) δ 1.81(d, 3H), 2.28(br, 2H), 3.06(br, 4H), 3.29(br, 2H), 4.64(d, 1H), 4.95(d, 1H), 5.14(q, 1H), 6.62(d, 1H), 7.01(d, 2H), 7.04(s, 1H), 7.20(s, 1H), 7.23-7.36(m, 5H), 7.39-7.50(m, 3H), 7.76(s, 1H), 7.78(d, 1H), 7.84(d, 1H), 8.00(d, 1H)

FAB (M+H) 491, C₃₁H₃₀N₄O₂

Example 13: Synthesis of 1-[1-((S)-α-methylbenzyl)-1H-imidazol-5-ylmethyl]-3-(4-methylpiperazin-1-yl)carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(13)

64_{mg} (0.2 mmol) of the compound prepared in Preparation 8 was dissolved in 2_{mℓ} of dimethylformamide, 26.4_{mg} (0.66 mmol) of sodium hydride(60%) was added thereto at 0°C and then the mixture was stirred for 5 minutes. To the mixture was added 50_{mg} (2.2 mmol) of the compound prepared in Preparation 4 and the whole mixture was stirred at room temperature for 3 hours. The solvent was removed by distillation under reduced pressure and 3_{mℓ} of water was added to the residue. The mixture was then extracted twice with 10_{mℓ} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated, and subjected to silica gel column chromatography (eluent: dichloromethane/methanol=90/10, v/v) to obtain 75_{mg} (Yield 75%) of the title compound.

¹H NMR(CDCl₃) δ 1.09(br, 2H), 1.77(d, 3H), 1.83(s, 3H), 1.70-1.90(br, 2H), 2.90(br, 2H), 3.31(br, 2H), 4.74(d, 1H), 4.93(d, 1H),

5.14(q, 1H), 6.60(d, 1H), 7.01(m, 3H), 7.17(s, 1H), 7.20-7.35(m, 4H), 7.45(m, 3H), 7.73(m, 2H), 7.80(d, 1H), 8.00(d, 1H)

FAB (M+H) 504, $C_{32}H_{33}N_5O$

Example 14: Synthesis of 3-[N-(2-methoxyethyl)-N-methyl]carbonyl-4-(naphthalen-1-yl)-1-[1-(phenethyl)-1H-imidazol-5-ylmethyl]-1H-pyrrole (14)

62_{mg}(0.2 mmol) of the compound prepared in Preparation 6 was dissolved in 2_{ml} of dimethylformamide, 26.4_{mg}(0.66 mmol) of sodium hydride(60%) was added thereto at 0°C and then the mixture was stirred for 5 minutes. To the mixture was added 50_{mg}(2.2 mmol) of the compound prepared in Preparation 5 and the whole mixture was stirred at room temperature for 3 hours. The solvent was removed by distillation under reduced pressure and 3_{ml} of water was added to the residue. The mixture was then extracted twice with 10_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated, and subjected to silica gel column chromatography (eluent: dichloromethane/methanol=95/5, v/v) to obtain 77_{mg}(Yield 78%) of the title compound.

¹H NMR(CDCl₃) δ 2.38(br, 2H), 2.70(m, 1H), 2.80(t, 2H), 2.90(m, 3H), 3.00(br, 2H), 3.31(br, 1H), 3.41(br, 1H), 4.03(t, 2H), 4.77(s, 2H), 6.66(d, 1H), 6.97(d, 1H), 7.06(d, 1H), 7.22(m, 3H), 7.30-7.60(m, 5H), 7.75(d, 1H), 7.80(d, 1H), 8.04(d, 1H)

FAB (M+H) 493, $C_{31}H_{32}N_4O_2$

Example 15: Synthesis of 3-(morpholin-4-yl)carbonyl-4-(naphthalen-1-yl)-1-[1-(phenethyl)-1H-imidazol-5-ylmethyl]-1H-pyrrole(15)

62_{mg}(0.2 mmol) of the compound prepared in Preparation 7 was dissolved in 2_{ml} of dimethylformamide, 26.4_{mg}(0.66 mmol) of sodium

hydride(60%) was added thereto at 0°C and then the mixture was stirred for 5 minutes. To the mixture was added 50_{mg}(2.2 mmol) of the compound prepared in Preparation 5 and the whole mixture was stirred at room temperature for 3 hours. The solvent was removed by distillation under reduced pressure and 3_{ml} of water was added to the residue. The mixture was then extracted twice with 10_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated, and subjected to silica gel column chromatography (eluent: dichloromethane/methanol=95/5, v/v) to obtain 79_{mg}(Yield 80%) of the title compound.

¹H NMR(CDCl₃) δ 2.28(br, 2H), 2.81(t, 2H), 2.83(br, 4H), 3.21(br, 2H), 4.07(t, 2H), 4.78(s, 2H), 6.68(d, 1H), 6.99(d, 1H), 7.10(d, 2H), 7.10(d, 2H), 7.23(m, 3H), 7.30(d, 1H), 7.50(m, 3H), 7.67(s, 1H), 7.77(d, 1H), 7.82(d, 1H), 8.00(d, 1H)

FAB (M+H) 491, C₃₁H₃₀N₄O₂

Example 16: Synthesis of 3-(4-methylpiperazin-1-yl)carbonyl-4-(naphthalen-1-yl)-1-[1-(phenethyl)-1H-imidazol-5-ylmethyl]-1H-pyrrole (16)

62_{mg}(0.2 mmol) of the compound prepared in Preparation 8 was dissolved in 2_{ml} of dimethylformamide, 26.4_{mg}(0.66 mmol) of sodium hydride(60%) was added thereto at 0°C and then the mixture was stirred for 5 minutes. To the mixture was added 50_{mg}(2.2 mmol) of the compound prepared in Preparation 5 and the whole mixture was stirred at room temperature for 3 hours. The solvent was removed by distillation under reduced pressure and 3_{ml} of water was added to the residue. The mixture was then extracted twice with 10_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated, and subjected to silica gel column chromatography (eluent: dichloromethane/methanol=90/10, v/v) to obtain 75_{mg}(Yield 75%) of the title compound.

^1H NMR(CDCl_3) δ 1.06(br, 2H), 1.90-2.00(br, 2H), 2.05(s, 3H), 2.80(t, 2H), 3.37(br, 4H), 4.04(t, 2H), 4.77(s, 2H), 6.69(d, 1H), 6.99(m, 2H), 7.09 (d, 2H), 7.20-7.56(m, 8H), 7.78(d, 1H), 7.83(d, 1H), 8.00(d, 1H)

FAB (M+H) 504, $\text{C}_{32}\text{H}_{33}\text{N}_5\text{O}$

Preparation 15: Synthesis of 1-(2-methoxy)phenethyl-5-chloromethyl-1H-imidazole hydrochloride

15-1) 1-(2-Methoxy)phenethyl-5-hydroxymethyl-1H-imidazole

The title compound was obtained in a yield of 65% according to the same procedure as Preparation 1-1) except that dihydroxyacetone dimer and 2-methoxyphenethylamine were used as starting materials.

^1H NMR(CDCl_3) δ 3.03(t, 2H), 3.75(s, 3H), 4.16(t, 2H), 4.47(s, 2H), 4.75(s, 1H), 6.74(s, 1H), 6.75-7.00(m, 3H), 7.13-7.30(m, 1H)

FAB 233 (M+H), $\text{C}_{13}\text{H}_{16}\text{N}_2\text{O}_2$ (M)

15-2) 1-(2-Methoxy)phenethyl-5-chloromethyl-1H-imidazole hydrochloride

The title compound was obtained in a yield of 89% according to the same procedure as Preparation 1-2) using the compound prepared in Preparation 15-1). The product thus obtained was directly used in the next reaction without purification.

Preparation 16: Synthesis of 1-(4-methoxy)phenethyl-5-chloromethyl-1H-imidazole hydrochloride

16-1) 1-(4-Methoxy)phenethyl-5-hydroxymethyl-1H-imidazole

The title compound was obtained in a yield of 60% according to the same procedure as Preparation 1-1) except that dihydroxyacetone dimer and 4-methoxyphenethylamine were used as starting materials.

^1H NMR(CDCl_3) δ 2.91(t, 2H), 3.68(s, 3H), 4.09(t, 2H), 4.36(s, 2H), 6.70(d, 2H), 6.77(s, 1H), 6.87(d, 2H), 7.13 (s, 1H)

FAB 233 (M+H), $\text{C}_{13}\text{H}_{16}\text{N}_2\text{O}_2$ (M)

16-2) 1-(4-Methoxy)phenethyl-5-chloromethyl-1H-imidazole hydrochloride

The title compound was obtained in a yield of 89% according to the same procedure as Preparation 1-2) using the compound prepared in Preparation 16-1). The product thus obtained was directly used in the next reaction without purification.

Preparation 17: Synthesis of 1-(2-fluoro)phenethyl-5-chloromethyl-1H-imidazole hydrochloride

17-1) 1-(2-Fluoro)phenethyl-5-hydroxymethyl-1H-imidazole

The title compound was obtained in a yield of 68% according to the same procedure as Preparation 1-1) except that dihydroxyacetone dimer and 2-fluorophenethylamine were used as starting materials.

^1H NMR(CDCl_3) δ 3.12(t, 2H), 3.50(br, 1H), 4.23 (t, 2H), 4.52(s, 2H), 6.82(s, 1H), 7.02(m, 3H), 7.20(m, 2H)

FAB 221 (M+H), $\text{C}_{12}\text{H}_{13}\text{N}_2\text{OF}$ (M)

17-2) 1-(2-Fluoro)phenethyl-5-chloromethyl-1H-imidazole hydrochloride

The title compound was obtained in a yield of 89% according to the same procedure as Preparation 1-2) using the compound prepared in Preparation 17-1). The product thus obtained was directly used in the next reaction without purification.

Preparation 18: Synthesis of 1-(2-chloro)phenethyl-5-chloromethyl-1H-imidazole hydrochloride

18-1) 1-(2-Chloro)phenethyl-5-hydroxymethyl-1H-imidazole

The title compound was obtained in a yield of 71% according to

the same procedure as Preparation 1-1) except that dihydroxyacetone dimer and 2-chlorophenethylamine were used as starting materials.

^1H NMR(CDCl_3) δ 3.13(t, 2H), 3.34(br, 1H), 4.18 (t, 2H), 4.42(s, 2H), 6.79(s, 1H), 6.94(d, 1H), 7.03-7.20(m, 3H), 7.29(d, 1H)

FAB 237 (M+H), $\text{C}_{12}\text{H}_{13}\text{N}_2\text{OCl}$ (M)

18-2) 1-(2-Chloro)phenethyl-5-chloromethyl-1H-imidazole hydrochloride

The title compound was obtained in a yield of 89% according to the same procedure as Preparation 1-2) using the compound prepared in Preparation 18-1). The product thus obtained was directly used in the next reaction without purification.

Preparation 19: Synthesis of 1-(3-chloro)phenethyl-5-chloromethyl-1H-imidazole hydrochloride

19-1) 1-(3-Chloro)phenethyl-5-hydroxymethyl-1H-imidazole

The title compound was obtained in a yield of 72% according to the same procedure as Preparation 1-1) except that dihydroxyacetone dimer and 3-chlorophenethylamine were used as starting materials.

^1H NMR(CDCl_3) δ 2.95(t, 2H), 3.90(br, 1H), 4.10 (t, 2H), 4.37(s, 2H), 6.74(s, 1H), 6.85(m, 1H), 6.98(s, 1H), 7.10(m, 3H)

FAB 237 (M+H), $\text{C}_{12}\text{H}_{13}\text{N}_2\text{OCl}$ (M)

19-2) 1-(3-Chloro)phenethyl-5-chloromethyl-1H-imidazole hydrochloride

The title compound was obtained in a yield of 91% according to the same procedure as Preparation 1-2) using the compound prepared in Preparation 19-1). The product thus obtained was directly used in the next reaction without purification.

Preparation 20: Synthesis of 1-(3-phenyl)propyl-5-chloromethyl-1H-imidazole hydrochloride

20-1) 1-(3-Phenyl)propyl-5-hydroxymethyl-1H-imidazole

The title compound was obtained in a yield of 73% according to the same procedure as Preparation 1-1) except that dihydroxyacetone dimer and 3-phenylpropylamine were used as starting materials.

^1H NMR(CDCl_3) δ 2.11(m, 2H), 2.61(t, 2H), 3.98(t, 2H), 4.25(br, 1H), 4.53(s, 1H), 6.76(s, 1H), 7.10-7.60(m, 6H)

FAB 217 (M+H), $\text{C}_{13}\text{H}_{16}\text{N}_2\text{O}$ (M)

20-2) 1-(3-Phenyl)propyl-5-chloromethyl-1H-imidazole hydrochloride

The title compound was obtained in a yield of 91% according to the same procedure as Preparation 1-2) using the compound prepared in Preparation 20-1). The product thus obtained was directly used in the next reaction without purification.

Preparation 21: Synthesis of 1-(naphthalen-2-yl)methyl-5-chloromethyl-1H-imidazole hydrochloride**21-1) 1-(Naphthalen-2-yl)methyl-5-hydroxymethyl-1H-imidazole**

The title compound was obtained in a yield of 58% according to the same procedure as Preparation 1-1) except that dihydroxyacetone dimer and (naphthalen-2-yl)methylamine were used as starting materials.

^1H NMR(CDCl_3) δ 4.36(s, 2H), 5.28(s, 2H), 6.89(s, 1H), 7.17(d, 1H), 7.35(m, 2H), 7.41(s, 1H), 7.50(s, 1H), 7.65(m, 1H), 7.69(m, 2H)

FAB 239 (M+H), $\text{C}_{15}\text{H}_{14}\text{N}_2\text{O}$ (M)

21-2) 1-(Naphthalen-2-yl)methyl-5-chloromethyl-1H-imidazole hydrochloride

The title compound was obtained in a yield of 87% according to the same procedure as Preparation 1-2) using the compound prepared in Preparation 21-1). The product thus obtained was directly used in the next reaction without purification.

Preparation 22: Synthesis of 1-[2-(naphthalen-1-yl)ethyl]-5-

chloromethyl-1H-imidazole hydrochloride**22-1) 1-[2-(Naphthalen-1-yl)ethyl]-5-hydroxymethyl-1H-imidazole**

The title compound was obtained in a yield of 58% according to the same procedure as Preparation 1-1) except that dihydroxyacetone dimer and (naphthalen-1-yl)ethylamine were used as starting materials.

^1H NMR(CDCl_3) δ 3.44(t, 2H), 4.23(t, 2H), 4.38(s, 2H), 6.79(s, 1H), 7.07(d, 1H), 7.17(s, 1H), 7.24(t, 1H), 7.32-7.48(m, 2H), 7.62(d, 1H), 7.74(d, 1H), 7.92(d, 1H)

FAB 253 (M+H), $\text{C}_{16}\text{H}_{16}\text{N}_2\text{O}$ (M)

22-2) 1-[2-(Naphthalen-1-yl)ethyl]-5-chloromethyl-1H-imidazole hydrochloride

The title compound was obtained in a yield of 87% according to the same procedure as Preparation 1-2) using the compound prepared in Preparation 22-1). The product thus obtained was directly used in the next reaction without purification.

Preparation 23: Synthesis of 1-(4-bromo)phenethyl-5-chloromethyl-1H-imidazole hydrochloride**23-1) 1-(4-Bromo)phenethyl-5-hydroxymethyl-1H-imidazole**

The title compound was obtained in a yield of 72% according to the same procedure as Preparation 1-1) except that dihydroxyacetone dimer and 4-bromophenethylamine were used as starting materials.

^1H NMR(CDCl_3) δ 2.94(t, 2H), 3.76(br, 1H), 4.11 (t, 2H), 4.37(s, 2H), 6.74(s, 1H), 6.85(d, 2H), 6.84(d, 2H), 7.12(s, 1H), 7.29(d, 2H)

FAB 281 (M+H), $\text{C}_{12}\text{H}_{13}\text{N}_2\text{OBr}$ (M)

23-2) 1-(4-Bromo)phenethyl-5-chloromethyl-1H-imidazole hydrochloride

The title compound was obtained in a yield of 91% according to

the same procedure as Preparation 1-2) using the compound prepared in Preparation 23-1). The product thus obtained was directly used in the next reaction without purification.

Preparation 24: Synthesis of 1-(4-fluoro)phenethyl-5-chloromethyl-1H-imidazole hydrochloride

24-1) 1-(4-Fluoro)phenethyl-5-hydroxymethyl-1H-imidazole

The title compound was obtained in a yield of 72% according to the same procedure as Preparation 1-1) except that dihydroxyacetone dimer and 4-fluorophenethylamine were used as starting materials.

^1H NMR(CDCl_3) δ 2.99(t, 2H), 3.76(br, 1H), 4.15(t, 2H), 4.45(s, 2H), 6.80-7.20(m, 5H), 7.26(s, 1H)

FAB 221 (M+H), $\text{C}_{12}\text{H}_{13}\text{N}_2\text{OF}$ (M)

24-2) 1-(4-Fluoro)phenethyl-5-chloromethyl-1H-imidazole hydrochloride

The title compound was obtained in a yield of 91% according to the same procedure as Preparation 1-2) using the compound prepared in Preparation 24-1). The product thus obtained was directly used in the next reaction without purification.

Preparation 25: Synthesis of 1-(4-methyl)phenethyl-5-chloromethyl-1H-imidazole hydrochloride

25-1) 1-(4-Methyl)phenethyl-5-hydroxymethyl-1H-imidazole

The title compound was obtained in a yield of 72% according to the same procedure as Preparation 1-1) except that dihydroxyacetone dimer and 4-methylphenethylamine were used as starting materials.

^1H NMR(CDCl_3) δ 3.02(t, 2H), 2.99(t, 2H), 3.76(br, 1H), 4.19(t, 2H), 4.47(s, 2H), 6.83(s, 1H), 6.94(d, 2H), 7.06(d, 2H), 7.28(s, 1H)

FAB 217 (M+H), $\text{C}_{13}\text{H}_{16}\text{N}_2\text{O}$ (M)

25-2) 1-(4-Methyl)phenethyl-5-chloromethyl-1H-imidazole hydrochloride

The title compound was obtained in a yield of 91% according to the same procedure as Preparation 1-2) using the compound prepared in Preparation 25-1). The product thus obtained was directly used in the next reaction without purification.

Preparation 26: Synthesis of 1-(4-chloro)phenethyl-5-chloromethyl-1H-imidazole hydrochloride**26-1) 1-(4-Chloro)phenethyl-5-hydroxymethyl-1H-imidazole**

The title compound was obtained in a yield of 73% according to the same procedure as Preparation 1-1) except that dihydroxyacetone dimer and 4-chlorophenethylamine were used as starting materials.

^1H NMR(CDCl_3) δ 3.04(t, 2H), 4.18(t, 2H), 4.48(s, 2H), 6.79(s, 1H), 6.96(d, 2H), 7.20-7.40(m, 3H)

FAB 237 (M+H), $\text{C}_{12}\text{H}_{13}\text{N}_2\text{OCl}$ (M)

26-2) 1-(4-Chloro)phenethyl-5-chloromethyl-1H-imidazole hydrochloride

The title compound was obtained in a yield of 91% according to the same procedure as Preparation 1-2) using the compound prepared in Preparation 26-1). The product thus obtained was directly used in the next reaction without purification.

Preparation 27: Synthesis of 1-[2-(naphthalen-2-yl)ethyl]-5-chloromethyl-1H-imidazole hydrochloride**27-1) 1-[2-(Naphthalen-2-yl)ethyl]-5-hydroxymethyl-1H-imidazole**

The title compound was obtained in a yield of 58% according to the same procedure as Preparation 1-1) except that dihydroxyacetone dimer and 2-(naphthalen-2-yl)ethylamine were used as starting materials.

^1H NMR(CDCl_3) δ 3.22(t, 2H), 4.28(t, 2H), 4.48(s, 2H), 6.84(s,

1H), 7.19(d, 1H), 7.24(d, 2H), 7.44(m, 2H), 7.52(s, 1H), 7.76(m, 3H)

FAB 253 (M+H), C₁₆H₁₆N₂O (M)

27-2) 1-[2-(Naphthalen-2-yl)ethyl]-5-chloromethyl-1H-imidazole
hydrochloride

The title compound was obtained in a yield of 88% according to the same procedure as Preparation 1-2) using the compound prepared in Preparation 27-1). The product thus obtained was directly used in the next reaction without purification.

Example 17: Synthesis of 3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-1-[1-(2-methoxy)phenethyl-1H-imidazol-5-yl]methyl-4-(naphthalen-1-yl)-1H-pyrrole(17)

62_{mg}(0.2 mmol) of the compound prepared in Preparation 6 was dissolved in 2_{ml} of dimethylformamide, 26.4_{mg}(0.66 mmol) of sodium hydride(60%) was added thereto at 0°C and then the mixture was stirred for 5 minutes. To the mixture was added 63_{mg}(2.2 mmol) of the compound prepared in Preparation 15 and the whole mixture was stirred at room temperature for 2 hours. The solvent was removed by distillation under reduced pressure and 3_{ml} of water was added to the residue. The mixture was then extracted twice with 10_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated, and subjected to silica gel column chromatography (eluent: dichloromethane/methanol=95/5, v/v) to obtain 78_{mg}(Yield 75%) of the title compound.

¹H NMR(CDCl₃) δ 2.39(s, 2H), 2.71(br, 1H), 2.90(t, 2H), 2.95-3.15(m, 5H), 3.31(br, 1H), 3.52(br, 1H), 3.76(s, 3H), 4.06(t, 2H), 4.83(s, 2H), 6.68(s, 1H), 6.75-6.95(m, 3H), 7.23(s, 1H), 7.25(s, 1H), 7.21(t, 1H), 7.30- 7.48(m, 4H), 7.50(s, 1H), 7.75(d, 1H), 7.81(d, 1H), 8.06(d, 1H)

FAB 523 (M+H), C₃₂H₃₄N₄O₃ (M)

Example 18: Synthesis of 1-[1-(2-methoxy)phenethyl-1H-imidazol-5-yl]methyl-3-[4-methylpiperazin-1-yl]carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(18)

62_{mg}(0.2 mmol) of the compound prepared in Preparation 8 was dissolved in 2_{ml} of dimethylformamide, 26.4_{mg}(0.66 mmol) of sodium hydride(60%) was added thereto at 0°C and then the mixture was stirred for 5 minutes. To the mixture was added 63_{mg}(2.2 mmol) of the compound prepared in Preparation 15 and the whole mixture was stirred at room temperature for 2 hours. The solvent was removed by distillation under reduced pressure and 3_{ml} of water was added to the residue. The mixture was then extracted twice with 10_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated, and subjected to silica gel column chromatography (eluent: dichloromethane/methanol= 95/5, v/v) to obtain 75_{mg}(Yield 70%) of the title compound.

¹H NMR(CDCl₃) δ 1.09(br, 2H), 1.70-2.10(br+s, 5H), 2.85(t, 2H), 2.99 (br, 2H), 3.40(br, 2H), 3.76(s, 3H), 4.04(t, 2H), 4.85(s, 2H), 6.69(d, 1H), 6.80-6.92(m, 3H), 7.04(s, 1H), 7.08(s, 1H), 7.25(t, 1H), 7.30(d, 1H), 7.35- 7.50(m, 4H), 7.77(d, 1H), 7.80(d, 1H), 8.02(d, 1H)

FAB 534 (M+H), C₃₃H₃₄N₅O₂ (M)

Example 19: Synthesis of 3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-1-[1-(4-methoxy)phenethyl-1H-imidazol-5-yl]methyl-4-(naphthalen-1-yl)-1H-pyrrole(19)

62_{mg}(0.2 mmol) of the compound prepared in Preparation 6 was dissolved in 2_{ml} of dimethylformamide, 26.4_{mg}(0.66 mmol) of sodium hydride(60%) was added thereto at 0°C and then the mixture was stirred for 5 minutes. To the mixture was added 63_{mg}(2.2 mmol) of the

compound prepared in Preparation 16 and the whole mixture was stirred at room temperature for 2 hours. The solvent was removed by distillation under reduced pressure and 3_{ml} of water was added to the residue. The mixture was then extracted twice with 10_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated, and subjected to silica gel column chromatography (eluent: dichloromethane/methanol=95/5, v/v) to obtain 83_{mg} (Yield 80%) of the title compound.

¹H NMR(CDCl₃) δ 2.38(br, 2H), 2.72(t, 2H), 2.85-3.15(m, 7H), 3.31(br, 1H), 3.72(s, 3H), 3.97(t, 2H), 4.78(s, 2H), 6.69(d, 1H), 6.77 (d, 2H), 6.85(d, 2H), 7.03(s, 1H), 7.06(s, 1H), 7.24-7.50(m, 5H), 7.73(d, 1H), 7.82(d, 1H), 8.05(d, 1H).

FAB 523 (M+H), C₃₂H₃₄N₄O₃ (M)

Example 20: Synthesis of 1-[1-(4-methoxy)phenethyl-1H-imidazol-5-yl]methyl-3-[4-methylpiperazin-1-yl]carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(20)

62_{mg} (0.2 mmol) of the compound prepared in Preparation 8 was dissolved in 2_{ml} of dimethylformamide, 26.4_{mg} (0.66 mmol) of sodium hydride(60%) was added thereto at 0°C and then the mixture was stirred for 5 minutes. To the mixture was added 63_{mg} (2.2 mmol) of the compound prepared in Preparation 16 and the whole mixture was stirred at room temperature for 2 hours. The solvent was removed by distillation under reduced pressure and 3_{ml} of water was added to the residue. The mixture was then extracted twice with 10_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated, and subjected to silica gel column chromatography (eluent: dichloromethane/methanol=95/5, v/v) to obtain 83_{mg} (Yield 78%) of the title compound.

¹H NMR(CDCl₃) δ 1.05(br, 2H), 1.70-2.10(br+s, 4H), 2.24(br, 1H), 2.72(t, 2H), 2.89(br, 2H), 3.30(br, 1H), 3.73(s, 3H), 3.98(t, 2H),

4.79(s, 2H), 6.69(d, 1H), 6.76(d, 2H), 6.86(d, 2H), 7.08(m, 2H),
7.30-7.50(m, 5H), 7.74(d, 1H), 7.80(d, 1H), 8.00(d, 1H)

FAB 534 (M+H), $C_{33}H_{35}N_5O_2$ (M)

Example 21: Synthesis of 1-[1-(2-fluoro)phenethyl-1H-imidazol-5-yl]methyl-3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole(21)

62_{mg}(0.2 mmol) of the compound prepared in Preparation 6 was dissolved in 2_{ml} of dimethylformamide, 26.4_{mg}(0.66 mmol) of sodium hydride(60%) was added thereto at 0°C and then the mixture was stirred for 5 minutes. To the mixture was added 61_{mg}(2.2 mmol) of the compound prepared in Preparation 17 and the whole mixture was stirred at room temperature for 2 hours. The solvent was removed by distillation under reduced pressure and 3_{ml} of water was added to the residue. The mixture was then extracted twice with 10_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated, and subjected to silica gel column chromatography (eluent: dichloromethane/methanol=95/5, v/v) to obtain 78_{mg}(Yield 77%) of the title compound.

¹H NMR(CDCl₃) δ 2.38(br, 2H), 2.70(br, 1H), 2.81(t, 2H), 2.90-3.38(m, 7H), 4.03(t, 2H), 4.91(s, 2H), 6.71(d, 2H), 6.92(m, 1H), 6.95-7.12(m, 4H), 7.19(m, 1H), 7.30-7.65(m, 4H), 7.73(d, 1H), 7.80(d, 1H), 8.05(d, 1H)

FAB 511 (M+H), $C_{31}H_{31}N_4O_2F$ (M)

Example 22: Synthesis of 1-[1-(2-fluoro)phenethyl-1H-imidazol-5-yl]methyl-3-[4-methylpiperazin-1-yl]carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(22)

62_{mg}(0.2 mmol) of the compound prepared in Preparation 8 was

dissolved in 2_{ml} of dimethylformamide, 26.4_{mg}(0.66 mmol) of sodium hydride(60%) was added thereto at 0°C and then the mixture was stirred for 5 minutes. To the mixture was added 61_{mg}(2.2 mmol) of the compound prepared in Preparation 17 and the whole mixture was stirred at room temperature for 2 hours. The solvent was removed by distillation under reduced pressure and 3_{ml} of water was added to the residue. The mixture was then extracted twice with 10_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated, and subjected to silica gel column chromatography (eluent: dichloromethane/methanol=95/5, v/v) to obtain 78_{mg}(Yield 75%) of the title compound.

¹H NMR(CDCl₃) δ 1.04(br, 2H), 1.70-2.10(br+s, 5H), 2.81(m, 2H), 3.90 (br, 2H), 3.32 (br, 2H), 4.05(t, 2H), 4.93(s, 2H), 6.72(d, 1H), 6.90(t, 1H), 6.95-7.05(m, 2H), 7.10(d, 2H), 7.20(m, 1H), 7.25-7.50(m, 4H), 7.75(d, 1H), 7.82(d, 2H), 8.00(d, 1H)

FAB 522 (M+H), C₃₂H₃₂N₅OF (M)

Example 23: Synthesis of 1-[1-(2-chloro)phenethyl-1H-imidazol-5-yl]methyl-3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole(23)

62_{mg}(0.2 mmol) of the compound prepared in Preparation 6 was dissolved in 2_{ml} of dimethylformamide, 26.4_{mg}(0.66 mmol) of sodium hydride(60%) was added thereto at 0°C and then the mixture was stirred for 5 minutes. To the mixture was added 64_{mg}(2.2 mmol) of the compound prepared in Preparation 18 and the whole mixture was stirred at room temperature for 3 hours. The solvent was removed by distillation under reduced pressure and 3_{ml} of water was added to the residue. The mixture was then extracted twice with 10_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated, and subjected to silica gel column chromatography (eluent: dichloromethane/methanol=

95/5, v/v) to obtain 75_{mg} (Yield 71%) of the title compound.

¹H NMR(CDCl₃) δ 2.39(br, 2H), 2.71(br, 1H), 2.90-3.38(m, 9H), 4.06(t, 2H), 4.87(s, 2H), 6.71(s, 1H), 6.87(m, 1H), 7.00-7.20(m, 4H), 7.30-7.60(m, 6H), 7.73(d, 1H), 7.89(d, 1H), 8.06(d, 1H)

FAB 527 (M+H), C₃₁H₃₁N₄O₂Cl (M)

Example 24: Synthesis of 1-[1-(2-chloro)phenethyl-1H-imidazol-5-yl] methyl-3-[4-methylpiperazin-1-yl]carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(24)

62_{mg} (0.2 mmol) of the compound prepared in Preparation 8 was dissolved in 2_{ml} of dimethylformamide, 26.4_{mg} (0.66 mmol) of sodium hydride (60%) was added thereto at 0°C and then the mixture was stirred for 5 minutes. To the mixture was added 64_{mg} (2.2 mmol) of the compound prepared in Preparation 18 and the whole mixture was stirred at room temperature for 2 hours. The solvent was removed by distillation under reduced pressure and 3_{ml} of water was added to the residue. The mixture was then extracted twice with 10_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated, and subjected to silica gel column chromatography (eluent: dichloromethane/methanol=95/5, v/v) to obtain 84_{mg} (Yield 78%) of the title compound.

¹H NMR(CDCl₃) δ 1.04(br, 1H), 1.70-2.10(br+s, 5H), 2.35(br, 1H), 2.92 (t+br, 4H), 3.32(br, 2H), 4.08(t, 2H), 4.88(s, 2H), 6.71(s, 1H), 6.87(m, 1H), 7.09(m, 3H), 7.18(m, 1H), 7.30-7.55(m, 6H), 7.75(d, 1H), 7.81(d, 1H), 8.01(d, 1H)

FAB 538 (M+H), C₃₂H₃₂N₅OCl (M)

Example 25: Synthesis of 1-[1-(3-chloro)phenethyl-1H-imidazol-5-yl] methyl-3-[N-(2-methoxyethyl)-N-methyl]carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(25)

62_{mg}(0.2 mmol) of the compound prepared in Preparation 6 was dissolved in 2_{ml} of dimethylformamide, 26.4_{mg}(0.66 mmol) of sodium hydride(60%) was added thereto at 0°C and then the mixture was stirred for 5 minutes. To the mixture was added 64_{mg}(2.2 mmol) of the compound prepared in Preparation 19 and the whole mixture was stirred at room temperature for 2 hours. The solvent was removed by distillation under reduced pressure and 3_{ml} of water was added to the residue. The mixture was then extracted twice with 10_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated, and subjected to silica gel column chromatography (eluent: dichloromethane/methanol= 95/5, v/v) to obtain 80_{mg}(Yield 76%) of the title compound.

¹H NMR(CDCl₃) δ 2.37(br, 2H), 2.71(m, 3H), 2.90-3.20(m, 6H), 3.30(br, 1H), 3.99(t, 2H), 4.86(s, 2H), 6.69(d, 1H), 6.81(d, 1H), 7.00(s, 1H), 7.05- 7.20(m, 5H), 7.30-7.50(m, 4H), 7.74(d, 1H), 7.81(d, 1H), 8.04(d, 1H)

FAB 527 (M+H), C₃₁H₃₁N₄O₂Cl (M)

Example 26: Synthesis of 1-[1-(3-chloro)phenethyl-1H-imidazol-5-yl] methyl-3-[4-methylpiperazin-1-yl]carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(26)

62_{mg}(0.2 mmol) of the compound prepared in Preparation 8 was dissolved in 2_{ml} of dimethylformamide, 26.4_{mg}(0.66 mmol) of sodium hydride(60%) was added thereto at 0°C and then the mixture was stirred for 5 minutes. To the mixture was added 64_{mg}(2.2 mmol) of the compound prepared in Preparation 19 and the whole mixture was stirred at room temperature for 2 hours. The solvent was removed by distillation under reduced pressure and 3_{ml} of water was added to the residue. The mixture was then extracted twice with 10_{ml} of ethyl

acetate, dried over anhydrous sodium sulfate, concentrated, and subjected to silica gel column chromatography (eluent: dichloromethane/methanol=95/5, v/v) to obtain 85_{mg} (Yield 79%) of the title compound.

¹H NMR(CDCl₃) δ 1.05(br, 2H), 1.70-2.10(br+s, 5H), 2.69(t, 2H), 2.90 (br, 2H), 3.32(br, 2H), 3.98(t, 2H), 4.87(s, 2H), 6.70(d, 1H), 6.79(d, 1H), 6.98(s, 1H), 7.05-7.21(m, 3H), 7.30-7.50(m, 6H), 7.74(d, 1H), 7.82(d, 1H), 7.99(d, 1H)

FAB 538 (M+H), C₃₂H₃₂N₅OCl (M)

Example 27: Synthesis of 3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)-1-[1-(3-phenyl)propyl-1H-imidazol-5-yl]methyl-1H-pyrrole(27)

62_{mg} (0.2 mmol) of the compound prepared in Preparation 6 was dissolved in 2_{mℓ} of dimethylformamide, 26.4_{mg} (0.66 mmol) of sodium hydride(60%) was added thereto at 0°C and then the mixture was stirred for 5 minutes. To the mixture was added 62_{mg} (2.2 mmol) of the compound prepared in Preparation 20 and the whole mixture was stirred at room temperature for 2 hours. The solvent was removed by distillation under reduced pressure and 3_{mℓ} of water was added to the residue. The mixture was then extracted twice with 10_{mℓ} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated, and subjected to silica gel column chromatography (eluent: dichloromethane/methanol=95/5, v/v) to obtain 76_{mg} (Yield 75%) of the title compound.

¹H NMR(CDCl₃) δ 1.91(m, 2H), 2.24(t, 2H), 2.56(m, 5H), 2.90-3.07(m, 4H), 3.18(br, 1H), 4.03(t, 2H), 5.12(s, 2H), 6.57(s, 1H), 6.90-7.20(m, 8H), 7.21-7.52(m, 3H), 7.66(d, 1H), 7.72(d, 1H), 7.89(d, 1H), 8.06(br, 1H)

FAB 507 (M+H), C₃₂H₃₄N₄O₂ (M)

Example 28: Synthesis of 3-[4-methylpiperazin-1-yl]carbonyl-4-naphthalen-1-yl)-1-[1-(3-phenyl)propyl-1H-imidazol-5-yl]methyl-1H-pyrrole(28)

62_{mg}(0.2 mmol) of the compound prepared in Preparation 8 was dissolved in 2_{ml} of dimethylformamide, 26.4_{mg}(0.66 mmol) of sodium hydride(60%) was added thereto at 0°C and then the mixture was stirred for 5 minutes. To the mixture was added 62_{mg}(2.2 mmol) of the compound prepared in Preparation 20 and the whole mixture was stirred at room temperature for 2 hours. The solvent was removed by distillation under reduced pressure and 3_{ml} of water was added to the residue. The mixture was then extracted twice with 10_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated, and subjected to silica gel column chromatography (eluent: dichloromethane/methanol= 95/5, v/v) to obtain 77_{mg}(Yield 74%) of the title compound.

¹H NMR(CDCl₃) δ 1.01(br, 2H), 2.80-2.01(s+br+m, 6H), 2.30(br, 1H), 2.55 (t, 2H), 2.86(br, 2H), 3.30(br, 2H), 3.79(t, 2H), 5.00(s, 2H), 6.58(s, 1H), 7.00-7.20(m, 8H), 7.36(m, 1H), 7.41(m, 2H), 7.50(s, 1H), 7.74(d, 1H), 7.80(d, 1H), 8.00(d, 1H)

FAB 518(M+H), C₃₃H₃₅N₅O (M)

Example 29: Synthesis of 3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)-1-[1-(naphthalen-2-yl)methyl-1H-imidazol-5-yl]methyl-1H-pyrrole(29)

62_{mg}(0.2 mmol) of the compound prepared in Preparation 6 was dissolved in 2_{ml} of dimethylformamide, 26.4_{mg}(0.66 mmol) of sodium hydride(60%) was added thereto at 0°C and then the mixture was stirred for 5 minutes. To the mixture was added 65_{mg}(2.2 mmol) of the compound prepared in Preparation 21 and the whole mixture was stirred

at room temperature for 2 hours. The solvent was removed by distillation under reduced pressure and 3_{ml} of water was added to the residue. The mixture was then extracted twice with 10_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated, and subjected to silica gel column chromatography (eluent: dichloromethane/methanol=95/5, v/v) to obtain 85_{mg} (Yield 80%) of the title compound.

¹H NMR(CDCl₃) δ 2.36(br, 2H), 2.72(br, 1H), 2.98(br, 3H), 3.02(br, 2H), 3.31(br, 1H), 3.73(br, 1H), 5.10(s, 2H), 5.47(s, 2H), 6.58(s, 1H), 7.03(s, 1H), 7.08(d, 1H), 7.15(d, 1H), 7.21(s, 1H), 7.34-7.53(m, 7H), 7.60(s, 1H), 7.70-7.83(m, 4H), 7.97(d, 1H)

FAB 529 (M+H), C₃₄H₃₀N₄O₂ (M)

Example 30: Synthesis of 3-[4-methylpiperazin-1-yl]carbonyl-4-(naphthalen-1-yl)-1-[1-naphthalen-2-yl)methyl-1H-imidazol-5-yl]methyl-1H-pyrrole(30)

62_{mg} (0.2 mmol) of the compound prepared in Preparation 8 was dissolved in 2_{ml} of dimethylformamide, 26.4_{mg} (0.66 mmol) of sodium hydride(60%) was added thereto at 0°C and then the mixture was stirred for 5 minutes. To the mixture was added 65_{mg} (2.2 mmol) of the compound prepared in Preparation 21 and the whole mixture was stirred at room temperature for 2 hours. The solvent was removed by distillation under reduced pressure and 3_{ml} of water was added to the residue. The mixture was then extracted twice with 10_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated, and subjected to silica gel column chromatography (eluent: dichloromethane/methanol=95/5, v/v) to obtain 74_{mg} (Yield 69%) of the title compound.

¹H NMR(CDCl₃) δ 0.98(br, 2H), 1.70-2.00(s+br, 5H), 2.81(br, 2H), 3.37 (br, 1H), 4.88(s, 2H), 5.10(s, 2H), 6.57(s, 1H), 7.02(s, 1H), 7.08(d, 1H), 7.16(d, 1H), 7.21(s, 1H), 7.34-7.52(m, 7H), 7.60(s, 1H),

7.70-7.83(m, 4H), 7.97(d, 1H)

FAB 540(M+H), C₃₅H₃₃N₅O (M)

Example 31: Synthesis of 3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)-1-{1-[2-(naphthalen-1-yl)ethyl]-1H-imidazol-5-yl}methyl-1H-pyrrole(31)

62_{mg}(0.2 mmol) of the compound prepared in Preparation 6 was dissolved in 2_{ml} of dimethylformamide, 26.4_{mg}(0.66 mmol) of sodium hydride(60%) was added thereto at 0°C and then the mixture was stirred for 5 minutes. To the mixture was added 68_{mg}(2.2 mmol) of the compound prepared in Preparation 22 and the whole mixture was stirred at room temperature for 2 hours. The solvent was removed by distillation under reduced pressure and 3_{ml} of water was added to the residue. The mixture was then extracted twice with 10_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated, and subjected to silica gel column chromatography (eluent: dichloromethane/methanol=95/5, v/v) to obtain 77_{mg}(Yield 71%) of the title compound.

¹H NMR(CDCl₃) δ 2.34(br, 2H), 2.68(br, 1H), 2.80-3.20(m, 5H), 3.23(t, 2H), 3.29(br, 2H), 4.12(t, 2H), 4.45(s, 2H), 6.43(d, 1H), 6.84(d, 1H), 6.97 (m, 2H), 7.21-7.52(m, 10H), 7.72(d, 1H), 7.78-7.85(m, 2H), 8.01(d, 1H)

FAB 543 (M+H), C₃₅H₃₄N₄O₂ (M)

Example 32: Synthesis of 3-[4-methylpiperazin-1-yl]carbonyl-4-(naphthalen-1-yl)-1-{1-[2-(naphthalen-1-yl)ethyl]-1H-imidazol-5-yl}methyl-1H-pyrrole(32)

62_{mg}(0.2 mmol) of the compound prepared in Preparation 8 was dissolved in 2_{ml} of dimethylformamide, 26.4_{mg}(0.66 mmol) of sodium

hydride(60%) was added thereto at 0°C and then the mixture was stirred for 5 minutes. To the mixture was added 68_{mg}(2.2 mmol) of the compound prepared in Preparation 22 and the whole mixture was stirred at room temperature for 2 hours. The solvent was removed by distillation under reduced pressure and 3_{ml} of water was added to the residue. The mixture was then extracted twice with 10_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated, and subjected to silica gel column chromatography (eluent: dichloromethane/methanol=95/5, v/v) to obtain 83_{mg}(Yield 75%) of the title compound.

¹H NMR(CDCl₃) δ 1.01(br, 2H), 1.70-2.00(br+s, 5H), 2.89(br, 2H), 3.27 (t, 2H), 3.40(br, 2H), 4.16(t, 2H), 4.50(s, 2H), 6.45(d, 1H), 6.90(d, 1H), 6.97(d, 1H), 6.99(s, 1H), 7.25-7.55(m, 8H), 7.73-7.95(m, 5H), 8.00(d, 1H)

FAB 554(M+H), C₃₆H₃₅N₅O (M)

Example 33: Synthesis of 1-[1-(4-bromo)phenethyl-1H-imidazol-5-yl]methyl-3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole(33)

62_{mg}(0.2 mmol) of the compound prepared in Preparation 6 was dissolved in 2_{ml} of dimethylformamide, 26.4_{mg}(0.66 mmol) of sodium hydride(60%) was added thereto at 0°C and then the mixture was stirred for 5 minutes. To the mixture was added 74_{mg}(2.2 mmol) of the compound prepared in Preparation 23 and the whole mixture was stirred at room temperature for 2 hours. The solvent was removed by distillation under reduced pressure and 3_{ml} of water was added to the residue. The mixture was then extracted twice with 10_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated, and subjected to silica gel column chromatography (eluent: dichloromethane/methanol=95/5, v/v) to obtain 88_{mg}(Yield 77%) of the title compound.

^1H NMR(CDCl_3) δ 2.38(br, 3H), 2.67(t, 2H), 2.90-3.23(m, 7H), 3.30(br, 1H), 3.97(t, 2H), 4.88(s, 1H), 6.69(d, 1H), 6.82(d, 2H), 7.08(d, 2H), 7.27-7.53(m, 7H), 7.73(d, 1H), 7.80(d, 1H), 8.02(d, 1H)

FAB 571 (M+H), $\text{C}_{31}\text{H}_{31}\text{N}_4\text{O}_2\text{Br}$ (M)

Example 34: Synthesis of 1-[1-(4-bromo)phenethyl-1H-imidazol-5-yl]methyl-3-[4-methylpiperazin-1-yl]carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(34)

62_{mg}(0.2 mmol) of the compound prepared in Preparation 8 was dissolved in 2_{ml} of dimethylformamide, 26.4_{mg}(0.66 mmol) of sodium hydride(60%) was added thereto at 0°C and then the mixture was stirred for 5 minutes. To the mixture was added 74_{mg}(2.2 mmol) of the compound prepared in Preparation 23 and the whole mixture was stirred at room temperature for 2 hours. The solvent was removed by distillation under reduced pressure and 3_{ml} of water was added to the residue. The mixture was then extracted twice with 10_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated, and subjected to silica gel column chromatography (eluent: dichloromethane/methanol=95/5, v/v) to obtain 82_{mg}(Yield 70%) of the title compound.

^1H NMR(CDCl_3) δ 1.04(br, 2H), 1.80-2.00(br+s, 4H), 2.48(br, 1H), 2.66 (t, 2H), 2.90(br, 2H), 3.31(br, 1H), 2.96(t, 2H), 4.88(s, 2H), 6.70(s, 1H), 6.82(d, 2H), 7.10(d, 2H), 7.25-7.60(m, 7H), 7.75(d, 1H), 7.82(d, 1H), 8.01(d, 1H)

FAB 582(M+H), $\text{C}_{32}\text{H}_{32}\text{N}_5\text{OBr}$ (M)

Example 35: Synthesis of 1-[1-(4-fluoro)phenethyl-1H-imidazol-5-yl]methyl-3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole(35)

62_{mg}(0.2 mmol) of the compound prepared in Preparation 6 was dissolved in 2_{ml} of dimethylformamide, 26.4_{mg}(0.66 mmol) of sodium hydride(60%) was added thereto at 0°C and then the mixture was stirred for 5 minutes. To the mixture was added 60_{mg}(2.2 mmol) of the compound prepared in Preparation 24 and the whole mixture was stirred at room temperature for 2 hours. The solvent was removed by distillation under reduced pressure and 3_{ml} of water was added to the residue. The mixture was then extracted twice with 10_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated, and subjected to silica gel column chromatography (eluent: dichloromethane/methanol=95/5, v/v) to obtain 77_{mg}(Yield 76%) of the title compound.

¹H NMR(CDCl₃) δ 2.34(br, 3H), 2.70(t, 2H), 2.90-3.20(br, 6H), 3.30(br, 1H), 3.96(t, 2H), 4.86(s, 1H), 6.68(d, 1H), 6.90(m, 4H), 7.05(s, 1H), 7.09(s, 1H), 7.25-7.52(m, 5H), 7.73(d, 1H), 8.05(d, 1H)

FAB 511 (M+H), C₃₁H₃₁N₄O₂F (M)

Example 36: Synthesis of 1-[1-(4-fluoro)phenethyl-1H-imidazol-5-yl]methyl-3-[4-methylpiperazin-1-yl]carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(36)

62_{mg}(0.2 mmol) of the compound prepared in Preparation 8 was dissolved in 2_{ml} of dimethylformamide, 26.4_{mg}(0.66 mmol) of sodium hydride(60%) was added thereto at 0°C and then the mixture was stirred for 5 minutes. To the mixture was added 60_{mg}(2.2 mmol) of the compound prepared in Preparation 24 and the whole mixture was stirred at room temperature for 2 hours. The solvent was removed by distillation under reduced pressure and 3_{ml} of water was added to the residue. The mixture was then extracted twice with 10_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated, and subjected to silica gel column chromatography (eluent: dichloromethane/methanol=

95/5, v/v) to obtain 78_{mg} (Yield 75%) of the title compound.

¹H NMR(CDCl₃) δ 1.05(br, 2H), 1.70-2.00(br+s, 4H), 2.25(br, 1H), 2.70 (t, 2H), 2.90(br, 2H), 3.30(br, 2H), 3.88(t, 2H), 4.87(s, 2H), 6.69(s, 1H), 6.90(m, 4H), 7.10(m, 2H), 7.29(m, 2H), 7.35-7.50(m, 3H), 7.74(d, 1H), 7.82(d, 1H), 8.00(d, 1H)

FAB 522(M+H), C₃₂H₃₂N₅OF (M)

Example 37: Synthesis of 3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-1-[1-(4-methyl)phenethyl-1H-imidazol-5-yl]methyl-4-(naphthalen-1-yl)-1H-pyrrole(37)

62_{mg} (0.2 mmol) of the compound prepared in Preparation 6 was dissolved in 2_{ml} of dimethylformamide, 26.4_{mg} (0.66 mmol) of sodium hydride (60%) was added thereto at 0°C and then the mixture was stirred for 5 minutes. To the mixture was added 60_{mg} (2.2 mmol) of the compound prepared in Preparation 25 and the whole mixture was stirred at room temperature for 2 hours. The solvent was removed by distillation under reduced pressure and 3_{ml} of water was added to the residue. The mixture was then extracted twice with 10_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated, and subjected to silica gel column chromatography (eluent: dichloromethane/methanol=95/5, v/v) to obtain 78_{mg} (Yield 80%) of the title compound.

¹H NMR(CDCl₃) δ 2.02(br, 1H), 2.28(s, 3H), 2.38(br, 2H), 2.70(br, 1H), 2.75(t, 2H), 2.95-3.20(m, 5H), 3.31(br, 1H), 3.99(t, 2H), 4.77(s, 2H), 6.67(s, 1H), 6.85(d, 2H), 7.06(m, 4H), 7.25-7.50(m, 5H), 7.74(d, 1H), 7.81(d, 1H), 8.07(d, 1H)

FAB 507 (M+H), C₃₂H₃₄N₄O₂ (M)

Example 38: Synthesis of 1-[1-(4-methyl)phenethyl-1H-imidazol-5-yl]methyl-3-[4-methylpiperazin-1-yl]carbonyl-4-(naphthalen-1-yl)-1H-

pyrrole(38)

62_{mg}(0.2 mmol) of the compound prepared in Preparation 8 was dissolved in 2_{ml} of dimethylformamide, 26.4_{mg}(0.66 mmol) of sodium hydride(60%) was added thereto at 0°C and then the mixture was stirred for 5 minutes. To the mixture was added 60_{mg}(2.2 mmol) of the compound prepared in Preparation 25 and the whole mixture was stirred at room temperature for 3 hours. The solvent was removed by distillation under reduced pressure and 3_{ml} of water was added to the residue. The mixture was then extracted twice with 10_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated, and subjected to silica gel column chromatography (eluent: dichloromethane/methanol=95/5, v/v) to obtain 81_{mg}(Yield 78%) of the title compound.

¹H NMR(CDCl₃) δ 1.07(br, 1H), 1.70-2.10(br+s, 6H), 2.28(s, 3H), 2.75(t, 2H), 2.90(br, 2H), 3.33(br, 2H), 4.00(t, 2H), 4.78(s, 2H), 6.72(s, 1H), 6.86(m, 2H), 7.04-7.23(m, 4H), 7.25-7.60(m, 5H), 7.75(d, 1H), 7.82(d, 1H), 8.01(d, 1H)

FAB 518 (M+H), C₃₃H₃₅N₅O (M)

Example 39: Synthesis of 1-[1-(4-chloro)phenethyl-1H-imidazol-5-yl]methyl-3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole(39)

62_{mg}(0.2 mmol) of the compound prepared in Preparation 6 was dissolved in 2_{ml} of dimethylformamide, 26.4_{mg}(0.66 mmol) of sodium hydride(60%) was added thereto at 0°C and then the mixture was stirred for 5 minutes. To the mixture was added 64_{mg}(2.2 mmol) of the compound prepared in Preparation 26 and the whole mixture was stirred at room temperature for 2 hours. The solvent was removed by distillation under reduced pressure and 3_{ml} of water was added to the

residue. The mixture was then extracted twice with 10_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated, and subjected to silica gel column chromatography (eluent: dichloromethane/methanol=95/5, v/v) to obtain 74_{mg} (Yield 70%) of the title compound.

¹H NMR(CDCl₃) δ 2.38(br, 2H), 2.70(t, 2H), 2.90-3.20(m, 7H), 3.30(br, 1H), 3.97(t, 2H), 4.88(s, 2H), 6.69(d, 1H), 6.88(d, 2H), 7.04(s, 1H), 7.09(s, 1H), 7.19(d, 1H), 7.24-7.50(m, 5H), 7.75(d, 1H), 7.81(d, 1H), 8.02(d, 1H)

FAB 527 (M+H), C₃₁H₃₁N₄O₂Cl (M)

Example 40: Synthesis of 1-[1-(4-chloro)phenethyl-1H-imidazol-5-yl]methyl]-3-[4-methylpiperazin-1-yl]carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(40)

62_{mg}(0.2 mmol) of the compound prepared in Preparation 8 was dissolved in 2_{ml} of dimethylformamide, 26.4_{mg}(0.66 mmol) of sodium hydride(60%) was added thereto at 0°C and then the mixture was stirred for 5 minutes. To the mixture was added 64_{mg}(2.2 mmol) of the compound prepared in Preparation 26 and the whole mixture was stirred at room temperature for 2 hours. The solvent was removed by distillation under reduced pressure and 3_{ml} of water was added to the residue. The mixture was then extracted twice with 10_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated, and subjected to silica gel column chromatography (eluent: dichloromethane/methanol=95/5, v/v) to obtain 84_{mg} (Yield 78%) of the title compound.

¹H NMR(CDCl₃) δ 1.08(br, 2H), 1.80(br, 2H), 1.95(s, 3H), 2.73(t, 2H), 2.93(br, 2H), 3.35(br, 2H), 4.00(t, 2H), 4.90(s, 2H), 6.71(d, 1H), 6.91(d, 2H), 7.13-7.60(m, 9H), 7.78(d, 1H), 7.82(d, 1H), 8.01(d, 1H)

FAB 538 (M+H), C₃₂H₃₂N₅OCl (M)

Example 41: Synthesis of 3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)-1-{1-[2-(naphthalen-2-yl)ethyl]-1H-imidazol-5-yl}methyl-1H-pyrrole(41)

62_{mg}(0.2 mmol) of the compound prepared in Preparation 6 was dissolved in 2_{ml} of dimethylformamide, 26.4_{mg}(0.66 mmol) of sodium hydride(60%) was added thereto at 0°C and then the mixture was stirred for 5 minutes. To the mixture was added 67_{mg}(2.2 mmol) of the compound prepared in Preparation 27 and the whole mixture was stirred at room temperature for 2 hours. The solvent was removed by distillation under reduced pressure and 3_{ml} of water was added to the residue. The mixture was then extracted twice with 10_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated, and subjected to silica gel column chromatography (eluent: dichloromethane/methanol=95/5, v/v) to obtain 79_{mg}(Yield 71%) of the title compound.

¹H NMR(CDCl₃) δ 2.96(br, 1H), 2.39(br, 2H), 2.71(br, 1H), 2.80-3.15(m, 7H), 3.32(br, 1H), 4.10(t, 2H), 4.78(s, 1H), 6.66(s, 1H), 7.09(m, 3H), 7.42(m, 8H), 7.63(m, 1H), 7.75(m, 3H), 7.82(d, 1H), 8.06(d, 1H)

FAB 543 (M+H), C₃₅H₃₄N₄O₂ (M)

Example 42: Synthesis of 3-[4-methylpiperazin-1-yl]carbonyl-4-(naphthalen-1-yl)-1-{1-[2-(naphthalen-2-yl)ethyl]-1H-imidazol-5-yl}methyl-1H-pyrrole(42)

62_{mg}(0.2 mmol) of the compound prepared in Preparation 8 was dissolved in 2_{ml} of dimethylformamide, 26.4_{mg}(0.66 mmol) of sodium hydride(60%) was added thereto at 0°C and then the mixture was stirred for 5 minutes. To the mixture was added 67_{mg}(2.2 mmol) of the

compound prepared in Preparation 27 and the whole mixture was stirred at room temperature for 2 hours. The solvent was removed by distillation under reduced pressure and 3_{ml} of water was added to the residue. The mixture was then extracted twice with 10_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated, and subjected to silica gel column chromatography (eluent: dichloromethane/methanol=95/5, v/v) to obtain 82_{mg}(Yield 74%) of the title compound.

¹H NMR(CDCl₃) δ 1.05(br, 2H), 1.70-2.00(s+br, 4H), 2.34(br, 1H), 2.90 (t, 2H), 3.01(br, 2H), 3.32(br, 2H), 4.08(t, 2H), 4.78(s, 2H), 6.65(d, 2H), 7.10(m, 3H), 7.21-7.42(m, 7H), 7.64(m, 1H), 7.75(m, 3H), 7.82(d, 1H), 8.01(d, 1H)

FAB 554(M+H), C₃₆H₃₅N₅O (M)

Example 43: Synthesis of 1-[1-(4-hydroxy)phenethyl-1H-imidazol-5-yl]methyl-3-[4-methylpiperazin-1-yl]carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(43)

53mg(0.1 mmol) of the compound prepared in Example 20 was dissolved in 1_{ml} of dichloromethane, 75_{mg}(0.3 mmol) of borontribromide (BBr₃) was added thereto, and the mixture was stirred for 3 hours. 1_{ml} of methanol was added to stop the reaction and the solvent was removed by distillation under reduced pressure. The residue was subjected to silica gel column chromatography(eluent: dichloromethane/methanol=20/80, v/v) to obtain 26mg(Yield 50%) of the title compound.

¹H NMR(CDCl₃) δ 1.20(br, 2H), 1.80-2.05(br+s, 4H), 2.65(t, 2H), 3.00-3.60(br, 5H), 3.98(t, 2H), 4.88(s, 2H), 6.72(m, 5H), 7.09(s, 1H), 7.14(d, 1H), 7.23(s, 1H), 7.27(s, 1H), 7.33(d, 1H), 7.40-7.53(m, 3H), 7.77(d, 1H), 7.82(d, 1H), 7.93(d, 1H)

FAB 520(M+H), C₃₂H₃₃O₂N₅ (M)

Preparation 28: Synthesis of 4-chloromethyl-1-trityl-1H-imidazole hydrochloride

28-1) 4-Hydroxymethyl-1-trityl-1H-imidazole

3.99g(29.6 mmol) of hydroxymethylimidazole hydrochloride was dissolved in a mixture of 30_{ml} of dimethylformamide and 10_{ml} triethylamine, and then a solution of 9.35g(33.5 mmol) of triphenylmethyl chloride in 110_{ml} of dimethylformamide was added slowly thereto. After 2 hours, 500_{ml} of ice water was added to the reaction mixture to obtain a solid. This solid was recrystallized from dioxane to give 8.82g(Yield 87%) of the title compound.

m.p.: 227-229°C

28-2) 4-Chloromethyl-1-trityl-1H-imidazole hydrochloride

1.50g(4.41 mmol) of the compound prepared in Preparation 28-1) was dissolved in 50_{ml} of chloroform, 0.94_{ml}(13.2 mmol) of thionyl chloride was slowly added thereto at 0°C, and the mixture was stirred at room temperature for 2 hours. The organic solvent was removed under reduced pressure to give 1.66g(4.20 mmol, Yield 95%) of the title compound, which was directly used in the next reaction without purification.

Preparation 29: Synthesis of 4-(5-chloromethyl-1H-imidazol-1-ylmethyl) benzonitrile hydrochloride

29-1) 4-Acetoxymethyl-1-trityl-1H-imidazole

To 100_{ml} of pyridine were added 5.00g(14.7 mmol) of the compound prepared in Preparation 28-1) and 1.65g(16.2 mmol) of acetic anhydride, and the mixture was stirred at room temperature for 24 hours. The reaction solution was distilled under reduced pressure to remove the

pyridine and then the residue was dissolved in 200_{ml} of ethyl acetate and washed with 100_{ml} of aqueous sodium chloride solution. The organic solvent was eliminated by distillation under reduced pressure and the residue was subjected to column chromatography (eluent: dichloromethane/methanol=20/1, v/v) to give 5.22g(13.7 mmol, Yield 93%) of the title compound.

¹H NMR(CDCl₃) δ 2.01(s, 3H), 4.95(s, 2H), 6.88(s, 1H), 7.08(s, 5H), 7.27(s, 10H), 7.45 (s, 1H)

29-2) 4-(4-Acetoxymethyl-1-trityl-1H-imidazol-3-ylmethyl)benzonitrile bromide

5.00g(13.1 mmol) of the compound prepared in Preparation 29-1) was dissolved in 20_{ml} of dichloromethane, 2.82g(14.4 mmol) of 4-cyanobenzyl bromide was added thereto, and the mixture was stirred at room temperature for 60 hours. The organic solvent was removed by distillation under reduced pressure and the residue was subjected to column chromatography(eluent: dichloromethane/methanol=5/1, v/v) to give 5.31g(9.17 mmol, Yield 70%) of the title compound.

¹H NMR(CDCl₃ + CD₃OD) δ 1.95(s, 3H), 4.95(s, 2H), 5.45(s, 2H), 7.11- 7.40(m, 18H), 7.65(d, 2H), 8.21(s, 1H)

29-3) 4-(5-Acetoxymethyl-1H-imidazol-1-ylmethyl)benzonitrile

9.10g(15.7 mmol) of the compound prepared in Preparation 29-2) was dissolved in 500_{ml} of dichloromethane, 6.06_{ml}(78.7 mmol) of trifluoroacetic acid and 12.5_{ml}(78.7 mmol) of triethylsilane were slowly added thereto at 0°C, and the mixture was stirred at room temperature for 1 hour. The organic solvent was removed by distillation under reduced pressure, and then the residue was adjusted to pH 10 with saturated K₂CO₃ aqueous solution and extracted with 300_{ml} of ethyl acetate. The organic solvent was removed by distillation under reduced

pressure and the residue was subjected to column chromatography using ethyl acetate as an eluent to give 3.60 g(14.1 mmol, Yield 90%) of the title compound.

^1H NMR(CDCl_3) δ 1.90(s, 3H), 4.97(s, 2H), 5.25(s, 2H), 7.14(d, 2H), 7.21(d, 1H), 7.67(s, 1H), 7.75(d, 2H)

29-4) 4-(5-Hydroxymethyl-1H-imidazol-1-ylmethyl)benzonitrile

4.20g(16.5 mmol) of the compound prepared in Preparation 29-3) was dissolved in 200 mL of methanol, 4.50g(32.9 mmol) of K_2CO_3 was added thereto, and the mixture was stirred at room temperature for 20 minutes. The organic solvent was removed by distillation under reduced pressure at room temperature. The residue was then extracted with 300 mL of ethyl acetate and the extract was subjected to column chromatography(eluent: dichloromethane/methanol=10/1, v/v) to give 3.19g (15.0 mmol, Yield 91%) of the title compound.

^1H NMR($\text{CDCl}_3 + \text{CD}_3\text{OD}$) δ 4.28(s, 2H), 5.18(s, 2H), 6.84(s, 1H), 7.12(d, 2H), 7.42(s, 1H), 7.55(d, 2H)

29-5) 4-(5-Chloromethyl-1H-imidazol-1-ylmethyl)benzonitrile hydrochloride

3.00g(14.1 mmol) of the compound prepared in Preparation 29-4) was dissolved in 40 mL of chloroform, 5.02 mL (70.5 mmol) of thionyl chloride was added slowly thereto at 0°C , and the mixture was stirred at room temperature for 2 hours. The organic solvent was removed under reduced pressure to obtain 3.50g(13.1 mmol, Yield 93%) of the title compound. This compound was directly used in the next reaction without purification.

Preparation 30: Synthesis of 4-(3-chloro-1-propenyl)-1-trityl-1H-imidazole

30-1) Methyl 3-(1H-imidazol-4-yl)acrylate

500mg(3.62 mmol) of 3-(1H-imidazol-4-yl)acrylic acid was added to 20_{ml} of methanolic HCl and the mixture was stirred at room temperature for 10 hours. The solvent was removed under reduced pressure and then 30_{ml} of dichloromethane was added to the residue. The mixture was washed sequentially with saturated NaHCO₃ solution, aqueous sodium chloride solution and water. The organic layer was dried over anhydrous magnesium sulfate and concentrated to give 510mg(3.35 mmol, Yield 93%) of the title compound. This compound was used directly in the next reaction without purification.

30-2) Methyl 3-(1-trityl-1H-imidazol-4-yl)acrylate

350mg(2.30 mmol) of the compound prepared in Preparation 30-1) and 705mg(2.53 mmol) of triphenylmethylchloride were dissolved in 10_{ml} of dimethylformamide, and 350_{μl}(2.53 mmol) of triethylamine was added thereto. After 2 hours, 100_{ml} of ice water was added to the reaction mixture to obtain a solid. This solid was filtered, washed with diethylether and hexane, and then dried to give 810mg(2.05 mmol, Yield 87%) of the title compound.

¹H NMR(CDCl₃) δ 3.75(s, 3H), 6.35(d, 1H), 7.05-7.50(m, 18H)

30-3) 1-(1-Trityl-1H-imidazol-4-yl)propen-3-ol

800mg(2.03 mmol) of the compound prepared in Preparation 30-2) was added to 20_{ml} of dry dichloromethane. After the mixture was cooled down to -78°C, 6.1_{ml}(1M solution in hexane) of diisobutylaluminum hydride was added thereto. Temperature was raised slowly to room temperature and then 2_{ml} of water was added to the mixture to stop the reaction. 3_{ml} of 1N NaOH was added and then 2_{ml} of water was added, and the mixture was filtered through cellite. The organic layer of the filtrate was separated and combined with the

dichloromethane-extract from the aqueous layer. The mixture was dried over anhydrous magnesium sulfate. The organic solvent was removed under reduced pressure and the residue was subjected to column chromatography(eluent: dichloromethane/methanol=20/1, v/v) to give 671 mg(1.83 mmol, Yield 90%) of the title compound.

^1H NMR(CDCl_3) δ 4.25(s, 2H), 6.45(s, 2H), 6.78(s, 1H), 7.10-7.50(m, 16H)

30-4) 4-(3-Chloropropenyl)-1-trityl-1H-imidazole

650mg(1.77 mmol) of the compound prepared in Preparation 30-3) was added to 10_{ml} of chloroform. 135 μl (1.9 mmol) of thionyl chloride was added thereto at 0°C and the mixture was stirred at room temperature for 2 hours. The organic solvent was removed by distillation under reduced pressure and the residue was dissolved in 10_{ml} of ethyl acetate. The solution was washed with saturated NaHCO_3 aqueous solution and the organic solvent was distilled under reduced pressure to give 647mg(1.68 mmol, Yield 95%) of the title compound.

^1H NMR(CDCl_3) δ 4.22(d, 2H), 6.40-6.55(m, 2H), 6.81(s, 1H), 7.10-7.50 (m, 16H).

Preparation 31: Synthesis of 5-chloromethyl-1-methylimidazole hydrochloride

31-1) 5-Hydroxymethyl-1-methylimidazole

The title compound was obtained in a yield of 32% according to the procedure described in J.M.Dener, L-H Zhang, H. Rapoport, *J. Org. Chem.*, 1993, 58, 1159 using dihydroxyacetone and methylamine hydrochloride as starting materials.

^1H NMR(CDCl_3) δ 3.67(s, 3H), 4.58(s, 2H), 5.37(brs, 1H), 6.76(s, 1H), 7.32(s, 1H)

31-2) 5-Chloromethyl-1-methylimidazole hydrochloride

The title compound was obtained in a yield of 95% according to the same procedure as Preparation 28-2) except that the compound prepared in Preparation 31-1) was used as a starting material.

Preparation 32: Synthesis of 1-(4-bromobenzyl)-5-chloromethyl-1H-imidazole hydrochloride**32-1) 1-(4-Bromobenzyl)-5-hydroxymethyl-1H-imidazole**

The title compound was obtained in a yield of 50% according to the procedure described in J.M.Dener, L-H Zhang, H. Rapoport, *J. Org. Chem.*, 1993, 58, 1159 using dihydroxyacetone dimer and 4-bromobenzylamine hydrochloride as starting materials.

^1H NMR ($\text{CDCl}_3 + \text{CD}_3\text{OD}$) δ 4.46(s, 2H), 5.26(s, 2H), 7.00(s, 1H), 7.07(d, 2H), 7.50(d, 2H), 7.65(s, 1H)

32-2) 1-(4-Bromobenzyl)-5-chloromethyl-1H-imidazole hydrochloride

The title compound was obtained in a yield of 96% according to the same procedure as Preparation 28-2) except that the compound prepared in Preparation 32-1) was used as a starting material. The product thus obtained was directly used in the next reaction without purification.

Preparation 33: Synthesis of 5-chloromethyl-1-isobutylimidazole hydrochloride**33-1) 5-Hydroxymethyl-1-isobutylimidazole**

The title compound was obtained in a yield of 45% according to the same procedure as Preparation 31-1) using dihydroxyacetone and

isobutylamine hydrochloride as starting materials.

^1H NMR(CDCl_3) δ 0.90(d, 6H), 1.76(m, 1H), 3.62(d, 2H), 4.24(brs, 1H), 4.60(s, 2H), 6.85(s, 1H), 7.45(s, 1H)

FAB (M+H): 155

33-2) 5-Chloromethyl-1-isobutylimidazole hydrochloride

The title compound was obtained in a yield of 95% according to the same procedure as Preparation 28-2) except that the compound prepared in Preparation 33-1) was used as a starting material.

Preparation 34: Synthesis of 5-chloromethyl-1-cyclohexylmethylimidazole hydrochloride

34-1) 5-Hydroxymethyl-1-cyclohexylmethylimidazole

The title compound was obtained in a yield of 45% according to the same procedure as Preparation 31-1) using dihydroxyacetone and cyclohexylmethylamine hydrochloride as starting materials.

^1H NMR(CDCl_3) δ 0.94(m, 2H), 1.16(m, 3H), 1.50-1.70(m, 6H), 3.65(d, 2H), 4.24(brs, 1H), 4.60(s, 2H), 6.85(s, 1H), 7.45(s, 1H)

FAB (M+H): 195

34-2) 5-Chloromethyl-1-cyclohexylmethylimidazole hydrochloride

The title compound was obtained in a yield of 95% according to the same procedure as Preparation 28-2) except that the compound prepared in Preparation 34-1) was used as a starting material.

Preparation 35: Synthesis of 5-chloromethyl-1-pentylimidazole hydrochloride

35-1) 5-Hydroxymethyl-1-pentylimidazole

The title compound was obtained in a yield of 50% according to the same procedure as Preparation 31-1) using dihydroxyacetone and pentylamine hydrochloride as starting materials.

^1H NMR(CDCl_3) δ 0.90(t, 3H), 1.08(brs, 2H), 1.30(m, 4H), 1.45(m, 2H), 3.64(t, 2H), 4.24(brs, 1H), 4.60(s, 2H), 6.84(s, 1H), 7.44(s, 1H)

FAB (M+H): 169

35-2) 5-Chloromethyl-1-pentylimidazole hydrochloride

The title compound was obtained in a yield of 90% according to the same procedure as Preparation 28-2) except that the compound prepared in Preparation 35-1) was used as a starting material.

Preparation 36: Synthesis of 5-chloromethyl-1-octylimidazole hydrochloride

36-1) 5-Hydroxymethyl-1-octylimidazole

The title compound was obtained in a yield of 52% according to the same procedure as Preparation 31-1) using dihydroxyacetone and octylamine hydrochloride as starting materials.

^1H NMR(CDCl_3) δ 0.88(t, 3H), 1.18(brs, 2H), 1.30(brs, 10H), 1.42(m, 2H), 3.67(t, 2H), 4.23(brs, 1H), 4.60(s, 2H), 6.84(s, 1H), 7.44(s, 1H)

FAB (M+H): 211

36-2) 5-Chloromethyl-1-octylimidazole hydrochloride

The title compound was obtained in a yield of 93% according to the same procedure as Preparation 28-2) except that the compound prepared in Preparation 36-1) was used as a starting material.

Preparation 37: Synthesis of 5-chloromethyl-1-decylimidazole hydrochloride

37-1) 5-Hydroxymethyl-1-decylimidazole

The title compound was obtained in a yield of 52% according to the same procedure as Preparation 31-1) using dihydroxyacetone and decylamine hydrochloride as starting materials.

^1H NMR(CDCl_3) δ 0.88(t, 3H), 1.04(brs, 2H), 1.30(brs, 14H), 1.42(m, 2H), 3.68(t, 2H), 4.23(brs, 1H), 4.60(s, 2H), 6.84(s, 1H), 7.44(s, 1H)

FAB (M+H): 239

37-2) 5-Chloromethyl-1-decylimidazole hydrochloride

The title compound was obtained in a yield of 93% according to the same procedure as Preparation 28-2) except that the compound prepared in Preparation 37-1) was used as a starting material.

Preparation 38: Synthesis of 5-chloromethyl-1-(3-methyl)butylimidazole hydrochloride

38-1) 5-Hydroxymethyl-1-(3-methyl)butylimidazole

The title compound was obtained in a yield of 52% according to the same procedure as Preparation 31-1) using dihydroxyacetone and isoamylamine hydrochloride as starting materials.

^1H NMR(CDCl_3) δ 0.90(d, 6H), 1.32(m, 2H), 1.65(m, 1H), 3.67(t, 2H), 4.23(brs, 1H), 4.60(s, 2H), 6.84(s, 1H), 7.44(s, 1H)

FAB (M+H): 169

38-2) 5-Chloromethyl-1-(3-methyl)butylimidazole hydrochloride

The title compound was obtained in a yield of 93% according to

the same procedure as Preparation 28-2) except that the compound prepared in Preparation 38-1) was used as a starting material.

Preparation 39: Synthesis of 5-chloromethyl-1-(2-methoxy)ethylimidazole hydrochloride

39-1) 5-Hydroxymethyl-1-(2-methoxy)ethylimidazole

The title compound was obtained in a yield of 60% according to the same procedure as Preparation 31-1) using dihydroxyacetone and 2-methoxyethylamine hydrochloride as starting materials.

^1H NMR(CDCl_3) δ 3.38(s, 3H), 3.42(t, 2H), 3.65(t, 2H), 4.23(brs, 1H), 4.60(s, 2H), 6.84(s, 1H), 7.44(s, 1H)

FAB (M+H): 157

39-2) 5-Chloromethyl-1-(2-methoxy)ethylimidazole hydrochloride

The title compound was obtained in a yield of 93% according to the same procedure as Preparation 28-2) except that the compound prepared in Preparation 39-1) was used as a starting material.

Preparation 40: Synthesis of 5-chloromethyl-1-(3-methoxy)propylimidazole hydrochloride

40-1) 5-Hydroxymethyl-1-(3-methoxy)propylimidazole

The title compound was obtained in a yield of 61% according to the same procedure as Preparation 31-1) using dihydroxyacetone and 3-methoxypropylamine hydrochloride as starting materials.

^1H NMR(CDCl_3) δ 1.72(m, 2H), 3.32(s, 3H), 3.46(t, 2H), 3.63(t, 2H), 4.23(brs, 1H), 4.60(s, 2H), 6.84(s, 1H), 7.44(s, 1H)

FAB (M+H): 171

40-2) 5-Chloromethyl-1-(3-methoxy)propylimidazole hydrochloride

The title compound was obtained in a yield of 90% according to the same procedure as Preparation 28-2) except that the compound prepared in Preparation 40-1) was used as a starting material.

Preparation 41: Synthesis of 5-chloromethyl-1-(3-ethoxy)propylimidazole hydrochloride**41-1) 5-Hydroxymethyl-1-(3-ethoxy)propylimidazole**

The title compound was obtained in a yield of 61% according to the same procedure as Preparation 31-1) using dihydroxyacetone and 3-ethoxypropylamine hydrochloride as starting materials.

^1H NMR(CDCl_3) δ 1.20(t, 3H), 1.72(m, 2H), 3.50(s, 4H), 3.63(t, 2H), 4.23(brs, 1H), 4.60(s, 2H), 6.84(s, 1H), 7.44(s, 1H)

FAB (M+H): 185

41-2) 5-Chloromethyl-1-(3-ethoxy)propylimidazole hydrochloride

The title compound was obtained in a yield of 90% according to the same procedure as Preparation 28-2) except that the compound prepared in Preparation 41-1) was used as a starting material.

Preparation 42: Synthesis of 5-chloromethyl-1-(3-isopropoxy)propylimidazole hydrochloride**42-1) 5-Hydroxymethyl-1-(3-isopropoxy)propylimidazole**

The title compound was obtained in a yield of 61% according to the same procedure as Preparation 31-1) using dihydroxyacetone and 3-isopropoxypropylamine hydrochloride as starting materials.

^1H NMR(CDCl_3) δ 1.15(d, 6H), 1.71(m, 2H), 3.45-3.55(m, 3H), 3.63(t, 2H), 4.23(brs, 1H), 4.60(s, 2H), 6.84(s, 1H), 7.44(s, 1H)

FAB (M+H): 199

42-2) 5-Chloromethyl-1-(3-isopropoxy)propylimidazole hydrochloride

The title compound was obtained in a yield of 90% according to the same procedure as Preparation 28-2) except that the compound prepared in Preparation 42-1) was used as a starting material.

Example 44: Synthesis of 1-(1H-imidazol-4-yl)methyl-4-(naphthalen-1-yl)-3-(thiophen-2-yl)carbonyl-1H-pyrrole(44)

44-1) 3-(Naphthalen-1-yl)-1-(thiophen-2-yl)-prop-2-en-1-one

3.12g(20 mmol) of 1-naphthaldehyde and 2.52g(20 mmol) of 2-acetylthiophene were dissolved in 20_{ml} of methanol and 800mg(20 mmol) of sodium hydroxide was slowly added thereto. The mixture was reacted at room temperature for 8 hours and then the solid thus produced was filtered and dried. The filtrate was adjusted to pH 4-6 using 1N hydrochloric acid solution and extracted with ethyl acetate. The organic solvent was removed under reduced pressure and the residue was subjected to column chromatography(eluent: hexane/ethyl acetate=4/1, v/v) to give 4.23g(16 mmol, Yield 80%) of the title compound together with the filtered solid.

¹H NMR(CDCl₃) δ 7.13-7.31(m, 2H), 7.55-7.70(m, 3H), 7.70(d, 1H), 7.85-7.90(m, 4H), 8.28(d, 1H), 8.70(d, 1H)

44-2) 4-(Naphthalen-1-yl)-3-(thiophen-2-yl)carbonyl-1H-pyrrole

2.64g(9.99 mmol) of the compound prepared in Example 44-1) and 2.35g(12.0 mmol) of tosylmethylisocyanide were dissolved in 30_{ml} of tetrahydrofuran. 1.35g(12.0 mmol) of potassium t-butoxide was slowly added thereto and the mixture was refluxed for 30 minutes. The solvent was removed under reduced pressure and then 15_{ml} of water and

20_{ml} of ethyl acetate was added to the residue. The mixture was shaken thoroughly and filtered to obtain the resulting solid. This solid was washed with diethylether and dried to give 1.97 g(6.48 mmol, Yield 65%) of the title compound.

¹H NMR(CDCl₃) δ 6.90(s, 1H), 7.12(s, 1H), 7.20-7.45(m, 4H), 7.55(s, 1H), 7.61(s, 1H), 7.70-8.00(m, 4H), 11.4(s, 1H)

44-3) 4-(Naphthalen-1-yl)-3-(thiophen-2-yl)carbonyl-1-(1-trityl-1H-imidazol-4-yl)methyl-1H-pyrrole

200mg(0.99 mmol) of the compound prepared in Example 44-2) was dissolved in 5_{ml} of dimethylformamide, 95mg(4.0 mmol) of sodium hydride(50%) was added thereto at 0°C, and the mixture was stirred for 5 minutes. 391mg(0.99 mmol) of the compound prepared in Preparation 28-2) was added to the reaction solution and stirred at room temperature for 5 hours. The solvent was removed by distillation under reduced pressure and the residue was extracted with ethyl acetate. The extract was dried over anhydrous magnesium sulfate, concentrated and subjected to column chromatography(eluent: hexane/ethyl acetate=1/3, v/v) to give 205mg(0.33 mmol, Yield 33%) of the title compound.

¹H NMR(CDCl₃) δ 5.02(s, 2H), 6.75(s, 1H), 6.79(s, 1H), 6.86(t, 1H), 7.10-7.52(m, 23H), 7.71(d, 1H), 7.78(d, 1H), 7.89(d, 1H)

44-4) 1-(1H-Imidazol-4-yl)methyl-4-(naphthalen-1-yl)-3-(thiophen-2-yl)carbonyl-1H-pyrrole

190mg(0.304 mmol) of the compound prepared in Example 44-3) was dissolved in a solvent mixture of trifluoroacetic acid/dichloromethane(0.5_{ml}/0.5_{ml}) and the solution was stirred at room temperature for 2 hours. The organic solvent was removed under reduced pressure. The residue was dissolved in 10_{ml} of ethyl acetate, washed with saturated Na₂CO₃ solution and water, dried over anhydrous magnesium

sulfate, concentrated and subjected to column chromatography(eluent: ethyl acetate) to give 103mg(0.269 mmol, Yield 88%) of the title compound.

^1H NMR(CDCl_3) δ 4.87(s, 2H), 6.55(s, 1H), 6.72(s, 1H), 6.88(t, 1H), 7.11-7.34(m, 7H), 7.50-7.67(m, 3H), 7.83(d, 1H)

FAB MS: 384(M+1)

Examples 45 to 72:

The compounds represented in the following Tables 2-1 to 2-3 were obtained according to the similar procedure as Example 44.

Table 2-1

COM. NO.	^1H NMR(CDCl_3) δ	FAB MS (M+1)
45	4.85(s, 2H), 6.51(s, 1H), 6.67(s, 1H), 7.06(s, 1H), 7.14(s, 1H), 7.21-7.32(m, 7H), 7.61-7.74(m, 3H), 7.82(d, 1H)	384
46	4.95(s, 2H), 6.58(s, 1H), 6.76(s, 1H), 7.13-7.35(m, 9H), 7.61-7.68(m, 4H), 7.91(d, 1H)	378
47	4.92(s, 2H), 6.61(s, 1H), 6.70(s, 1H), 7.02(d, 2H), 7.17-7.35(m, 9H), 7.62(d, 1H), 7.70(d, 1H), 7.95(d, 1H)	456
48	5.03(s, 2H), 6.76(s, 1H), 6.85(s, 1H), 7.07(t, 1H), 7.34-7.54(m, 9H), 7.72-7.79(m, 3H), 7.94(d, 1H)	456
49	5.00(s, 2H), 6.72(s, 1H), 6.77(s, 1H), 7.21-7.38(m, 11H), 7.62(d, 1H), 7.70(d, 1H), 7.78(d, 1H)	456
50	2.23(s, 3H), 5.02(s, 2H), 6.74-7.10(m, 5H), 7.17-7.50(m, 8H), 7.65(d, 1H), 7.71(d, 1H), 7.86(d, 1H)	392
51	($\text{CDCl}_3 + \text{CD}_3\text{OD}$) 2.05(s, 3H), 5.09(s, 2H), 6.84(s, 1H), 6.99-7.05(m, 8H), 7.23-7.36(m, 3H), 7.70(d, 1H), 7.86(d, 1H)	392

Table 2-2

COM. NO.	^1H NMR(CDCl_3) δ	FAB MS (M+1)
52	2.21(s, 3H), 4.92(s, 2H), 6.62(s, 1H), 6.83(s, 1H), 7.14-7.35(m, 8H), 7.61-7.73(m, 5H), 7.88(d, 1H)	392
53	3.66(s, 3H), 5.04(s, 2H), 6.85(s, 1H), 6.82(d, 1H), 6.90(m, 1H), 7.12-7.17(m, 2H), 7.26-7.36(m, 8H), 7.67(t, 1H), 7.74(d, 1H), 7.93(d, 1H)	408
54	3.75(s, 3H), 5.02(s, 2H), 6.71(m, 3H), 6.80(t, 1H), 7.20-7.35(m, 6H), 7.60-7.75(m, 4H), 7.91(d, 1H)	408
55	4.83(s, 2H), 6.51(s, 1H), 6.63(s, 1H), 6.85(m, 1H), 7.03-7.29(m, 10H), 7.61-7.69(m, 2H), 7.83(d, 1H)	412
56	5.01(s, 2H), 6.72(s, 1H), 6.77(s, 1H), 7.22-7.35(m, 11H), 7.61-7.80(m, 3H)	412
57	4.82(s, 2H), 6.63(s, 1H), 6.72(s, 1H), 7.02-7.24(m, 10H), 7.56-7.70(m, 3H)	446
58	4.91(s, 2H), 6.65(s, 1H), 6.77(m, 1H), 7.20-7.31(m, 7H), 7.61(m, 3H), 7.81(d, 1H)	396
59	4.92(s, 2H), 6.45(m, 1H), 6.71(m, 2H), 7.20-7.32(m, 9H), 7.63-7.77(m, 3H)	414
60	5.09(s, 2H), 6.80-7.20(m, 4H), 7.15-7.35(m, 4H), 7.40(d, 1H), 7.45-7.50(m, 3H), 7.60(m, 1H), 7.65(d, 1H), 7.75(d, 1H)	403
61	1.87(s, 3H), 3.55(s, 2H), 5.07(s, 2H), 6.84(s, 2H), 7.08(d, 2H), 7.28-7.48(m, 6H), 7.57(d, 2H), 7.63(t, 1H), 7.71(d, 1H), 7.90(d, 1H)	438
62	2.03(s, 3H), 2.74(m, 2H), 2.91(m, 2H), 5.00(s, 2H), 6.67(s, 1H), 7.02(d, 2H), 7.14-7.43(m, 11H), 7.72-7.89(m, 3H)	452
63	1.98(s, 3H), 2.75(t, 2H), 3.90(t, 2H), 4.85(s, 2H), 6.60-6.72(m, 4H), 7.11-7.45(m, 9H), 7.68-7.82(m, 3H)	468
64	2.01(s, 3H), 3.61(s, 2H), 4.98(s, 2H), 6.61(s, 2H), 6.74(m, 2H), 7.10-7.48(m, 10H), 7.71-7.88(m, 3H)	438

Table 2-3

COM NO.	¹ H NMR(CDCl ₃) δ	FAB MS (M+1)
65	4.92(s, 2H), 6.62(s, 1H), 6.70(s, 1H), 7.12-7.27(m, 14H), 7.53-7.62(m, 4H), 7.81(d, 1H)	454
66	4.97(s, 2H), 6.87(d, 1H), 7.15-7.46(m, 15H), 7.55-7.73(m, 4H), 7.86(m, 1H)	454
67	5.10(s, 2H), 6.70(t, 2H), 6.80-6.95(m, 4H), 7.15(t, 1H), 7.21-7.45(m, 7H), 7.50(t, 1H), 7.60(d, 2H), 7.71(d, 1H), 7.75-7.80(m, 2H), 7.91(m, 1H)	470
68	3.82(s, 2H), 4.95(s, 2H), 6.57(s, 1H), 6.63(s, 1H), 6.92(d, 2H), 7.04(d, 2H), 7.20-7.32(m, 10H), 7.51-7.68(m, 4H), 7.82(d, 1H)	468
69	4.82(s, 2H), 6.41(s, 1H), 6.70(s, 1H), 6.95(s, 1H), 7.16-7.32(m, 9H), 7.51(d, 1H), 7.59(d, 1H), 7.67(m, 3H), 7.90(d, 1H), 8.05(d, 1H)	428
70	2.38(s, 3H), 3.65(s, 3H), 4.91(s, 2H), 6.69(s, 1H), 6.97(d, 1H), 7.00(t, 1H), 7.04(d, 1H), 7.10-7.16(m, 3H), 7.19(d, 1H), 7.34(s, 1H), 7.42(s, 1H), 7.57(d, 1H), 7.67(s, 2H)	395
71	0.61(t, 3H), 1.02(m, 2H), 1.25(m, 2H), 2.31(m, 1H), 2.47(m, 1H), 5.05(s, 2H), 6.57(s, 1H), 6.63(s, 1H), 6.80(d, 1H), 6.87(s, 1H), 7.22-7.35(m, 7H), 7.61-7.72(m, 3H)	502
72	3.71(d, 1H), 3.85(d, 1H), 4.85(s, 2H), 6.61(s, 1H), 6.73(d, 1H), 6.92-7.41(m, 14H), 7.62-7.73(m, 3H)	536

Example 73: Synthesis of 1-[1-(4-cyanobenzyl)-1H-imidazol-5-yl]methyl-4-(naphthalen-1-yl)-3-(thiophen-2-yl)carbonyl-1H-pyrrole(73)

80mg(0.3 mmol) of the compound prepared in Preparation 29-5) and 90mg(0.3 mmol) of the compound prepared in Example 44-2) were dissolved in 2ml of dimethylformamide, 36mg of sodium hydride(60%) was added thereto, and the mixture was stirred for 2 hours. The solvent

was removed by distillation under reduced pressure and the residue was subjected to column chromatography(eluent: dichloromethane/methanol=10/1, v/v) to give 83mg(0.17 mmol, Yield 56%) of the title compound.

^1H NMR(CDCl_3) δ 5.02(s, 2H), 5.08(s, 1H), 6.73(s, 1H), 6.85(s, 1H), 7.03(t, 1H), 7.32-7.45(m, 11H), 7.63(s, 1H), 7.75(d, 1H), 7.82(d, 1H), 8.02 (d, 1H)

FAB MS : 499 (M+1)

Example 74 例 77:

The compounds represented in the following Table 3 were obtained according to the similar procedure as Example 73.

Table 3

COM. NO.	^1H NMR(CDCl_3) δ	FAB MS (M+1)
74	4.82(s, 2H), 5.12(s, 1H), 6.30(s, 1H), 6.41(s, 1H), 6.77-7.08(m, 12H), 7.31-7.46(m, 3H), 7.68(d, 1H)	499
75	5.00(s, 2H), 5.05(s, 2H), 6.76(s, 1H), 6.82(s, 1H), 7.23-7.40(m, 12H), 7.63(d, 2H), 7.72(d, 1H), 7.90(d, 1H)	493
76	5.02(s, 2H), 5.08(s, 2H), 6.65(s, 1H), 6.78(s, 1H), 6.98(t, 1H), 7.23-7.42(m, 12H), 7.65-7.73(m, 3H), 7.82(d, 1H)	571
77	5.03(s, 2H), 5.10(s, 2H), 6.78(s, 1H), 6.87(s, 1H), 7.32-7.45(m, 12H), 7.74(d, 3H), 7.81(d, 1H), 7.88(d, 1H)	571

Example 78: Synthesis of 3-(4-fluorobenzoyl)-1-(1-methyl-1H-imidazol-4-yl)methyl-4-(naphthalen-1-yl)-1H-pyrrole(78)

The title compound was obtained in a yield of 75% according to

the same procedure as Example 44-3) except that 3-(4-fluorobenzoyl)-4-(naphthalen-1-yl)-1H-pyrrole and the compound prepared in Preparation 31-2) were used.

$^1\text{H NMR}(\text{CDCl}_3)$ δ 3.42(s, 3H), 5.01(s, 2H), 6.73(m, 3H), 7.11(s, 1H), 7.24-7.57(m, 8H), 7.67-7.75(m, 2H)

FAB MS (M+1): 410

Example 79: Synthesis of 1-(1-methyl-1H-imidazol-4-yl)methyl-4-(naphthalen-1-yl)-3-(4-phenoxybenzoyl)-1H-pyrrole(79)

The title compound was obtained in a yield of 70% according to the same procedure as Example 44-3) except that 4-(naphthalen-1-yl)-3-(4-phenoxybenzoyl)-1H-pyrrole and the compound prepared in Preparation 31-2) were used.

$^1\text{H NMR}(\text{CDCl}_3)$ δ 3.52(s, 3H), 5.12(s, 2H), 6.63(d, 2H), 6.76(d, 1H), 6.85(d, 2H), 7.12(t, 1H), 7.20(s, 1H), 7.28-7.40(m, 7H), 7.51(d, 2H), 7.68(d, 2H), 7.74(d, 1H), 7.83(d, 1H)

FAB MS (M+1): 484

Example 80: Synthesis of (S)-1-(1H-imidazol-4-yl)methyl-3-[N-(1-methoxycarbonyl-3-methylthio)propyl]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole(80)

80-1) Ethyl 3-(naphthalen-1-yl)acrylate

22.4g(0.10 mol) of triethylphosphonoacetate was dissolved in 500 mL of tetrahydrofuran and 12.4g(1.1 mol) of potassium t-butoxide was slowly added thereto. To this solution was slowly added 15.6g(0.10 mol) of 1-naphthaldehyde dissolved in 20 mL of tetrahydrofuran and the mixture was stirred for 8 hours. The organic solvent was removed by distillation under reduced pressure. The residue was dissolved in ethyl

acetate, washed twice with water, dried over anhydrous magnesium sulfate, concentrated and subjected to column chromatography(eluent: hexane/ethyl acetate=95/5, v/v) to give 20.3g(0.090 mol, Yield 90%) of the title compound.

^1H NMR(CDCl_3) δ 1.42(t, 3H), 4.30(q, 2H), 6.50(d, 1H), 7.40-7.60(m, 3H), 7.73(d, 1H), 7.82(m, 2H), 8.20(d, 1H), 8.50(d, 1H)

80-2) 3-Ethoxycarbonyl-4-(naphthalen-1-yl)-1H-pyrrole

500mg(1.89 mmol) of ethyl 3-(naphthalen-1-yl)acrylate prepared in Example 80-1) and 368mg(1.89 mmol) of tosylmethyisocyanide were dissolved in 10 mL of tetrahydrofuran. 255mg(2.27 mmol) of potassium t-butoxide dissolved in tetrahydrofuran(10 mL) was slowly added thereto and the mixture was refluxed for 30 minutes. 10 mL of water was added to the reaction solution to stop the reaction and the solvent was removed under reduced pressure. The residue was extracted with diethylether, washed with aqueous sodium chloride solution and dried over anhydrous magnesium sulfate. The solvent was removed under reduced pressure and the residue was subjected to column chromatography(eluent: ethyl acetate/hexane=1/3, v/v) to give 385mg(1.45 mmol, Yield 77%) of the title compound.

^1H NMR(CDCl_3) δ 0.86(t, 3H), 4.02(q, 2H), 6.81(s, 1H), 7.48-7.61(m, 5H), 7.90-7.97(m, 3H), 8.92(s, 1H)

80-3) 3-Ethoxycarbonyl-1-(1H-imidazol-4-yl)methyl-4-(naphthalen-1-yl)-1H-pyrrole

The title compound was obtained in a yield of 39% by applying the procedure described in Examples 44-3) and 44-4) from the compounds prepared in Example 80-2) and Preparation 28-2).

^1H NMR(CDCl_3) δ 1.11(t, 3H), 4.20(q, 2H), 5.05(s, 2H), 6.78(s, 1H), 6.89(s, 1H), 7.38-7.49(m, 6H), 7.85-7.97(m, 3H)

80-4) 3-Hydroxycarbonyl-1-(1H-imidazol-4-yl)methyl-4-(naphthalen-1-yl)-1H-pyrrole

220mg(0.64 mmol) of the compound prepared in Example 80-3) was dissolved in 5_{mℓ} of 50% ethanol, 216mg(3.8 mmol) of potassium hydroxide was added dropwise thereto, and the mixture was refluxed for 7 hours. The reaction solution was cooled down to room temperature, adjusted to pH 4-5, extracted with ethyl acetate and dried over anhydrous sodium sulfate. The solvent therein was removed under reduced pressure to give 162mg(0.51 mmol, Yield 80%) of the title compound. This compound was directly used in the next reaction without purification.

¹H NMR(CD₃OD + CDCl₃) δ 5.01(s, 2H), 6.82(s, 1H), 6.87(s, 1H), 7.42- 7.70(m, 7H), 7.82-7.89(m, 3H)

80-5) (S)-1-(1H-Imidazol-4-yl)methyl-3-[N-(1-methoxycarbonyl-3-methylthio)propyl]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole

200mg(0.60 mmol) of the compound prepared in Example 80-4) was dissolved in 2_{mℓ} of dimethylformamide, and then 150_{mg}(0.78 mmol) of EDC and 105_{mg}(0.78 mmol) of HOBT were added thereto. The resulting mixture was stirred at 0°C for 5 minutes. To the reaction solution was added 120_{mg}(0.60 mmol) of L-methionine methylester, which was then stirred at room temperature for 5 hours. The solvent was removed under reduced pressure and then 10_{mℓ} of saturated NaHCO₃ solution was added to the residue. The resulting solution was extracted with ethyl acetate, washed with aqueous sodium chloride solution and water, dried over anhydrous sodium sulfate and concentrated. The residue was subject to column chromatography(eluent: dichloromethane/methanol=20/1, v/v) to give 104_{mg}(0.225 mmol, Yield 37%) of the title compound.

^1H NMR(CDCl_3) δ 1.21(m, 1H), 1.55(m, 3H), 1.80(s, 3H), 3.42(s, 3H), 4.43(m, 1H), 5.05(s, 2H), 5.60(d, 1H), 6.71(s, 1H), 6.95(s, 1H), 7.21-7.45(m, 7H), 7.75-7.87(m, 3H)

FAB MS : 463 (M+1)

Example 81: Synthesis of (S)-3-[N-(1-hydroxycarbonyl-3-methylthio)propyl]carbamoyl-1-(1H-imidazol-4-yl)methyl-4-(naphthalen-1-yl)-1H-pyrrole(81)

70mg(0.15 mmol) of the compound prepared in Example 80-5) was dissolved in 2ml of a solvent mixture of tetrahydrofuran/methanol/water(3/2/1, v/v/v), 10mg(0.18 mmol) of lithium hydroxide was added thereto, and the mixture was stirred at room temperature for 4 hours. The solvent was removed under reduced pressure to give 68mg(0.15 mmol, Yield 99.7%) of the lithium salt of the title compound.

^1H NMR($\text{CD}_3\text{OD} + \text{CDCl}_3$) δ 1.25(m, 1H), 1.49(m, 3H), 1.85(s, 3H), 4.41 (m, 1H), 5.11(s, 2H), 5.58(d, 1H), 6.70(s, 1H), 6.89(s, 1H), 7.15-7.38(m, 7H), 7.76-7.81(m, 3H)

FAB MS : 449 (M+1)

Examples 82 to 98:

The compounds represented in the following Tables 4-1 and 4-2 were obtained according to the similar procedure as Example 80.

Table 4-1

COM. NO.	¹ H NMR(CDCl ₃) δ	FAB MS (M+1)
82	5.02(s, 2H), 6.69(d, 2H), 6.77(s, 1H), 6.92-7.18(m, 5H), 7.40-7.58(m, 6H), 7.75-7.87(m, 4H)	393
83	4.06(d, 2H), 5.01(s, 2H), 5.57(t, 1H), 6.46(d, 1H), 6.71(s, 1H), 6.83(s, 1H), 6.92-7.05(m, 3H), 7.42-7.55(m, 7H), 7.74-7.81(m, 3H)	407
84	0.45(brs, 2H), 1.22(brs, 4H), 2.95(brs, 2H), 3.37(brs, 2H), 5.04(s, 2H), 6.65(s, 1H), 6.92(s, 1H), 7.08(s, 1H), 7.31-7.45(m, 6H), 7.72(d, 1H), 7.82(d, 1H), 8.12 (d, 1H)	385
85	2.32(brs, 2H), 2.22(brs, 2H), 3.23(brs, 2H), 3.65(brs, 2H), 5.06(s, 2H), 6.72(s, 1H), 6.95(s, 1H), 7.12(s, 1H), 7.31-7.48(m, 6H), 7.81(d, 1H), 7.85(d, 1H), 8.11(d, 1H)	387
86	1.41(brs, 2H), 2.86-3.25(m, 6H), 4.97(s, 2H), 6.68(s, 1H), 6.85(s, 1H), 7.06(s, 1H), 7.21-7.35(m, 6H), 7.72(d, 1H), 7.78(d, 1H), 7.95(d, 1H)	403
87	2.04(brs, 4H), 3.62(brs, 4H), 5.03(s, 2H), 6.91(d, 2H), 7.22-7.48(m, 7H), 7.81-7.88(m, 2H), 8.02(m, 1H)	435
88	1.46(brs, 2H), 2.21(brs, 2H), 3.14(brs, 4H), 5.11(s, 2H), 6.88(s, 1H), 7.02(s, 1H), 7.11(s, 1H), 7.32-7.51(m, 5H), 7.62(s, 1H), 7.72-7.80(m, 2H), 8.05(d, 1H)	386
89	(CDCl ₃ + CD ₃ OD) 2.05(s, 3H), 3.33(brs, 8H), 5.13(s, 2H), 6.90(s, 1H), 7.06(s, 1H), 7.21(s, 1H), 7.30-7.55(m, 4H), 7.64(s, 1H), 7.81(s, 1H), 7.88(d, 1H), 8.06(d, 1H)	400
90	2.62(brs, 2H), 3.15(brs, 2H), 3.86(brs, 1H), 4.35(brs, 1H), 5.06(s, 2H), 6.83(s, 1H), 6.90(s, 1H), 7.15-7.60(m, 6H), 7.73(d, 1H), 7.82(d, 1H), 8.06(d, 1H)	389
91	0.22(m, 1H), 0.63(m, 1H), 0.83(m, 1H), 1.24(m, 1H), 2.61(brs, 2H), 3.24(brs, 2H), 3.65(brs, 1H), 4.94(s, 2H), 6.71(s, 1H), 6.84(s, 1H), 6.94(s, 1H), 7.24-7.42(m, 6H), 7.62-7.70(m, 2H), 7.94(d, 1H)	401

Table 4-2

COM. NO.	¹ H NMR(CDCl ₃) δ	FAB MS (M+1)
92	1.37(brs, 2H), 1.96(brs, 2H), 3.52(brs, 4H), 5.21(s, 2H), 7.08(s, 1H), 7.20(s, 1H), 7.37(s, 1H), 7.54(m, 5H), 7.70(s, 1H), 7.94(m, 2H), 8.23(d, 1H)	399
93	2.12(brs, 2H), 3.02(brs, 2H), 4.98(s, 2H), 5.24(m, 1H), 6.82(s, 1H), 7.03(s, 1H), 7.20-7.34(m, 6H), 7.62-7.71(m, 3H), 7.93(d, 1H)	361
94	2.24(brs, 2H), 3.04(brs, 4H), 3.11(s, 3H), 5.03(s, 2H), 6.77(s, 1H), 6.89(s, 1H), 7.14-7.31(m, 6H), 7.56-7.63(m, 3H), 7.87(d, 1H)	375
95	2.51(m, 2H), 3.10(s, 3H), 3.21(m, 2H), 3.47(s, 3H), 5.05(s, 2H), 6.68(s, 1H), 7.05-7.48(m, 7H), 7.74-7.85(m, 2H), 8.09(d, 1H)	389
96	2.58-3.50(brs, 8H), 5.16(s, 2H), 6.98(d, 1H), 7.08(s, 1H), 7.20-7.27(m, 2H), 7.47(t, 1H), 7.67(s, 1H), 7.71(t, 1H), 8.08(d, 1H), 8.15(d, 1H), 8.80(d, 1H)	388
97	3.40(m, 4H), 3.70-4.45(brs, 8H), 3.11(s, 3H), 5.18(s, 2H), 6.98(d, 1H), 7.12(s, 1H), 7.17-7.22(m, 2H), 7.25(d, 1H), 7.30(d, 1H), 7.35(t, 1H), 7.62(d, 1H), 7.90(s, 1H)	413
98	(CD ₃ OD) 3.86(s, 2H), 4.83(s, 2H), 5.58(t, 1H), 6.37(d, 1H), 6.52(s, 2H), 6.81(s, 1H), 7.05-7.35(m, 9H), 7.51(d, 1H), 7.54(d, 1H), 7.58(d, 1H)	432

Example 99: Synthesis of 1-(1-methyl-1H-imidazol-5-yl)methyl-3-(morpholin-4-yl)carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(99)

To the compound prepared in Example 85 was introduced trityl protecting group according to the same procedure as Preparation 28-1), and then the title compound was obtained in a yield of 55% by applying the procedures described in Preparation 29-2) and 29-3) using

methyl iodide.

^1H NMR(CDCl_3) δ 2.80-3.45(m, 8H), 3.58(s, 3H), 5.19(s, 2H), 6.75(d, 1H), 7.18(d, 1H), 7.21(s, 1H), 7.35(d, 1H), 7.40-7.50(m, 3H), 7.72(d, 1H), 8.03(d, 1H)

FAB MS : 401 (M+1)

Example 100: Synthesis of (S)-1-[1-(4-cyanobenzyl)-1H-imidazol-5-yl]methyl-3-[N-(1-methoxycarbonyl-3-methylthio)propyl]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole(100)

100-1) 1-[1-(4-Cyanobenzyl)-1H-imidazol-5-ylmethyl]-3-hydroxycarbonyl-4-(naphthalen-1-yl)-1H-pyrrole

The title compound was obtained in a yield of 75% from the compounds prepared in Example 80-2) and Preparation 29-5) by sequentially applying the procedures of Example 73 and Example 80-4).

^1H NMR($\text{CDCl}_3 + \text{CD}_3\text{OD}$) δ 5.02(s, 2H), 5.10(s, 2H), 6.76(s, 1H), 7.07(m, 2H), 7.25-7.82(m, 12H)

100-2) (S)-1-[1-(4-Cyanobenzyl)-1H-imidazol-5-yl]methyl-3-[N-(1-methoxycarbonyl-3-methylthio)propyl]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole

The title compound was obtained in a yield of 35% according to the same procedure as Example 80-5) except that the compound prepared in Example 100-1) was used.

^1H NMR(CDCl_3) δ 1.85(s, 3H), 2.04(m, 1H), 2.13(m, 1H), 2.42(t, 2H), 3.61(s, 3H), 4.83(m, 1H), 5.02(s, 2H), 5.11(s, 2H), 6.63(s, 1H), 7.01(d, 2H), 7.13(d, 1H), 7.22-7.43(m, 7H), 7.65-7.92(m, 4H)

FAB MS : 578 (M+1)

Example 101: Synthesis of (S)-1-[1-(4-cyanobenzyl)-1H-imidazol-5-yl]methyl-3-[N-(1-hydroxycarbonyl-3-methylthio)propyl]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole(101)

halen-1-yl)-1H-pyrrole(101)

Lithium salt of the title compound was obtained in a yield of 96% according to the similar procedure as Example 81 from the compound prepared in Example 100-2).

^1H NMR($\text{CDCl}_3 + \text{CD}_3\text{OD}$) δ 1.82(s, 3H), 2.00(m, 1H), 2.11(m, 1H), 2.36(t, 2H), 4.82(m, 1H), 4.89(s, 2H), 5.02(s, 2H), 6.49(s, 1H), 6.88(d, 2H), 7.11(d, 1H), 7.17-7.32(m, 7H), 7.62-7.83(m, 4H)

FAB MS : 564(M+1)

Examples 102 and 103:

The compounds represented in the following Table 5 were obtained according to the similar procedure as Examples 100 and 101.

Table 5

COM. NO.	^1H NMR δ	FAB MS (M+1)
102	(CDCl_3) 0.67(d, 3H), 0.78(d, 3H), 0.82(m, 1H), 0.90(m, 1H), 1.10(m, 1H), 3.52(s, 3H), 4.32(m, 1H), 5.02(s, 2H), 5.17(s, 2H), 6.72(s, 1H), 6.83(s, 1H), 7.23-7.34(m, 3H), 7.41-7.92(m, 10H)	560
103	($\text{CDCl}_3 + \text{CD}_3\text{OD}$) 0.62(d, 3H), 0.71(d, 3H), 0.79(m, 1H), 0.88(m, 1H), 0.98(m, 1H), 4.12(m, 1H), 4.97(s, 2H), 5.08(s, 2H), 6.77(s, 1H), 6.82(s, 1H), 7.14-7.30(m, 4H), 7.38-7.84(m, 9H)	546

Examples 104 and 105:

The compounds represented in the following Table 6 were obtained according to the similar procedure as Example 101.

Table 6

COM. NO.	¹ H NMR (CDCl ₃) δ	FAB MS (M+1)
104	1.95(brs, 2H), 2.33(brs, 1H), 2.95(brs, 5H), 4.93(s, 2H), 5.05(s, 2H), 6.62(s, 1H), 7.05(s, 1H), 7.11(d, 2H), 7.28(m, 2H), 7.51(m, 3H), 7.63(m, 3H), 7.81-7.88(m, 2H), 7.95(d, 1H)	502
105	1.12(brs, 2H), 1.88(brs, 2H), 1.90(s, 3H), 2.95(brs, 2H), 3.34(brs, 2H), 4.97(s, 2H), 5.07(s, 2H), 6.60(s, 1H), 7.02(s, 1H), 7.10(d, 2H), 7.29(m, 2H), 7.46(m, 3H), 7.60(m, 3H), 7.80(d, 1H), 7.85(d, 1H), 7.97(d, 1H)	515

Example 106: Synthesis of 1-[2-(1H-imidazol-1-yl)ethyl]-3-(morpholin-4-yl)carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(106)

106-1) 2-(1H-Imidazol-1-yl)ethyl p-tosylate

0.24g(2.41 mmol) of 2-(1H-imidazol-1-yl)ethanol and 0.55g(2.88 mmol) of tosylchloride were dissolved in 20_{ml} of dichloromethane, 0.67 ml of triethylamine was slowly added thereto at 0°C, and the mixture was stirred at room temperature for 4 hours. The organic solvent was removed under reduced pressure. The residue was dissolved in 10_{ml} of ethyl acetate, washed sequentially with 1N hydrochloric acid solution, saturated sodium bicarbonate solution and aqueous sodium chloride solution, dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to column chromatography(eluent: dichloromethane/methanol=20/1, v/v) to give 0.30g(1.13 mmol, Yield 47%) of the title compound.

¹H NMR(CDCl₃) δ 2.42(s, 3H), 4.17-4.28(m, 4H), 6.88(s, 1H), 6.99(s, 1H), 7.29(d, 2H), 7.45(s, 1H), 7.64(d, 2H)

106-2) 3-Hydroxycarbonyl-4-(naphthalen-1-yl)-1H-pyrrole

The title compound was obtained in a yield of 80% by hydrolyzing the compound prepared in Example 80-2) according to the same procedure as Example 80-4).

^1H NMR ($\text{CDCl}_3 + \text{CD}_3\text{OD}$) δ 7.12(m, 3H), 7.20-7.31(m, 3H), 7.50(d, 1H), 7.68(d, 1H), 7.76(d, 1H)

106-3) 3-(Morpholin-4-yl)carbonyl-4-(naphthalen-1-yl)-1H-pyrrole

The title compound was obtained in a yield of 99% according to the same procedure as Example 80-5) from the compound prepared in Example 106-2) and morpholine.

^1H NMR (CDCl_3) δ 2.68-3.62(brs, 8H), 6.88(s, 1H), 7.20(s, 1H), 7.30-7.62(m, 4H), 7.78(d, 1H), 7.85(d, 1H), 8.08(d, 1H), 10.34(s, 1H)

106-4) 1-[2-(1H-Imidazol-1-yl)ethyl]-3-(morpholin-4-yl)carbonyl-4-(naphthalen-1-yl)-1H-pyrrole

The title compound was obtained in a yield of 51% by reacting the compound prepared in Example 106-1) with the compound prepared in Example 106-3) according to the same procedure as Example 44-3).

^1H NMR (CDCl_3) δ 2.20-3.72(brs, 12H), 7.20(s, 1H), 7.40-7.55(m, 8H), 7.82(d, 1H), 7.88(d, 1H), 8.05(d, 1H)

FAB MS : 401 (M+1)

Example 107: Synthesis of (S)-1-[3-(1H-imidazol-4-yl)propyl]-3-[N-(1-methoxycarbonyl-3-methylthio)propyl]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole(107)**107-1) 3-Ethoxycarbonyl-4-(naphthalen-1-yl)-1-[3-(1-trityl-1H-imidazol-4-yl)allyl]-1H-pyrrole**

The title compound was obtained in a yield of 85% by reacting

the compound prepared in Example 80-2) with the compound prepared in Preparation 30-4) according to the same procedure as Example 44-3).

^1H NMR (CDCl_3) δ 0.82(t, 3H), 3.95(q, 2H), 4.67(s, 2H), 6.23(d, 1H), 6.47(m, 1H), 6.63(s, 1H), 7.02(s, 1H), 7.25-7.81(m, 24H)

107-2) 3-Ethoxycarbonyl-4-(naphthalen-1-yl)-1-[3-(1-trityl-1H-imidazol-4-yl)propyl]-1H-pyrrole

300mg(0.49 mmol) of the compound prepared in Example 107-1) was dissolved in 2ml of methanol, catalytic amount of Pd/C was added thereto, and the mixture was stirred for 1 hour under hydrogen atmosphere. The mixture was filtered to remove the catalyst and the solvent therein was removed under reduced pressure. The residue was subject to column chromatography(eluent: dichloromethane/methanol=98/2, v/v) to give 246mg(0.40 mmol, Yield 82%) of the title compound.

^1H NMR (CDCl_3) δ 0.92(t, 3H), 2.22(m, 2H), 2.73(t, 2H), 4.01(m, 4H), 6.70(s, 1H), 6.82(s, 1H), 7.32-7.73(m, 21H), 7.91(m, 3H)

107-3) (S)-1-[3-(1H-Imidazol-4-yl)propyl]-3-[N-(1-methoxycarbonyl-3-methylthio)propyl]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole

The compound prepared in Example 107-2) was treated according to the procedures of Example 44-4) and 80-4) to eliminate the trityl group and hydrolyze. Then, the product thus obtained was reacted with (L)-methionine methylester according to the same procedure as Example 80-5) to give the title compound in a yield of 29%.

^1H NMR (CDCl_3) δ 1.65(m, 2H), 1.90(s, 3H), 2.12(m, 2H), 2.31(m, 2H), 2.73(m, 2H), 3.54(s, 3H), 4.02(m, 2H), 4.56(m, 1H), 5.77(d, 1H), 6.72(s, 1H), 6.90(s, 1H), 7.42-7.67(m, 7H), 7.82-8.01(m, 5H)

FAB MS : 491(M+1)

Example 108: Synthesis of (S)-3-[N-(1-hydroxycarbonyl-3-methylthio)

propyl]carbamoyl-1-[3-(1H-imidazol-4-yl)propyl]-4-(naphthalen-1-yl)-1H-pyrrole(108)

The title compound was obtained in a yield of 95% according to the same procedure as Example 81 except that the compound prepared in Example 107-3) was used.

¹H NMR (CDCl₃) δ 1.57(m, 2H), 1.88(s, 3H), 2.08(m, 2H), 2.29(m, 2H), 2.77(m, 2H), 4.12(m, 2H), 4.49(m, 1H), 5.69(d, 1H), 6.77(s, 1H), 6.92(s, 1H), 7.34-7.58(m, 7H), 7.80-7.89(m, 5H)

FAB MS : 477(M+1)

Example 109: Synthesis of 1-[3-(1H-imidazol-4-yl)propyl]-3-(morpholin-4-yl)carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(109)

The title compound was obtained in a yield of 42% according to the same procedure as Example 107-3) except that morpholine was used to the compound prepared in Example 107-2).

¹H NMR (CDCl₃) δ 2.16(m, 2H), 2.35(brs, 2H), 2.63(m, 2H), 2.80-3.50(brs, 6H), 3.54(s, 3H), 3.96(m, 2H), 6.74(d, 1H), 6.76(s, 1H), 7.07(s, 1H), 7.33(t, 1H), 7.36-7.50(m, 4H), 7.76(d, 1H), 7.84(d, 1H), 8.08(d, 1H)

FAB MS : 415(M+1)

Example 110: Synthesis of 1-[1-(4-cyanobenzyl)-1H-imidazol-5-yl]methyl-3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole(110)

110-1) 3-[N-(2-Methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole

100 mg(0.42 mmol) of the compound prepared in Example 106-2)

and 38_{mg}(0.4 mmol) of N-(2-methoxyethyl)-N-methylamine were reacted according to the similar procedure as Example 80-5) to give 110_{mg}(0.35 mmol, Yield 85%) of the title compound.

¹H NMR (CDCl₃) δ 2.21(s, 3H), 2.64(brs, 1H), 2.75(brs, 1H), 3.02(s, 3H), 3.13(brs, 1H), 3.32(brs, 1H), 6.72(s, 1H), 7.05(m, 2H), 7.21(m, 2H), 7.54(m, 1H), 7.78(m, 2H), 8.04(d, 1H), 8.78(brs, 1H)

110-2) 1-[1-(4-Cyanobenzyl)-1H-imidazol-5-yl]methyl-3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole

98_{mg}(0.32 mmol) of the compound prepared in Example 110-1) and 85_{mg}(0.32 mmol) of the compound prepared in Preparation 29-5) were reacted according to the similar procedure as Example 44-3) to give 115_{mg}(0.23 mmol, Yield 72%) of the title compound.

¹H NMR (CDCl₃) δ 2.41(s, 3H), 2.75(brs, 2H), 3.07(s, 3H), 3.17(brs, 1H), 3.32(brs, 1H), 4.91(s, 2H), 5.11(s, 2H), 6.71(s, 1H), 7.05(s, 1H), 7.17(d, 1H), 7.40-7.68(m, 9H), 7.78(d, 1H), 7.88(d, 1H), 8.06(d, 1H)

FAB MS : 504 (M+1)

Example 111: Synthesis of 1-[1-(4-bromobenzyl)-1H-imidazol-5-yl]methyl-3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole(111)

105_{mg}(0.29 mmol) of the compound prepared in Example 110-1) and 78_{mg}(0.29 mmol) of the compound prepared in Preparation 32-2) were reacted according to the similar procedure as Example 44-3) to give 121_{mg}(0.21 mmol, Yield 75%) of the title compound.

¹H NMR (CDCl₃) δ 2.37(s, 3H), 2.72(brs, 2H), 3.04(s, 3H), 3.15(brs, 1H), 3.31(brs, 1H), 4.95(s, 2H), 5.10(s, 2H), 6.67(s, 1H), 7.11(s, 1H), 7.23-7.65(m, 10H), 7.81(d, 1H), 7.89(d, 1H), 8.02(d, 1H)

FAB MS : 557 (M+1)

Example 112: Synthesis of 1-[1-(4-bromobenzyl)-1H-imidazol-5-yl] methyl-3-(morpholin-4-yl)carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(112)

100_{mg}(0.33 mmol) of the compound prepared in Example 106-3) and 105_{mg}(0.33 mmol) of the compound prepared in Preparation 32-2) were reacted according to the similar procedure as Example 44-3) to give 130_{mg}(0.23 mmol, Yield 71%) of the title compound.

¹H NMR (CDCl₃) δ 2.04(brs, 2H), 2.25(brs, 1H), 3.03(brs, 5H), 4.93(s, 2H), 5.07(s, 2H), 6.62(s, 1H), 7.10(m, 3H), 7.29(m, 2H), 7.41(m, 3H), 7.60(m, 3H), 7.81(d, 1H), 7.89(d, 1H), 8.01(d, 1H)

FAB MS : 555 (M+1)

Example 113: Synthesis of 1-[1-(4-cyanobenzyl)-1H-imidazol-5-yl] methyl-3-(morpholin-4-yl)thiocarbonyl-4-(naphthalen-1-yl)-1H-pyrrole (113)

20_{mg}(0.04 mmol) of the compound prepared in Example 104 and 18_{mg} of 2,4-bis(phenylthio)-1,3-dithia-2,4-diphosphatan-2,4-disulfide were dissolved in 1_{mL} of tetrahydrofuran, and the mixture was stirred at room temperature for 3 hours. To the reaction solution was added 2_{mL} of saturated sodium bicarbonate solution. The resulting mixture was extracted with ethyl acetate and dried over anhydrous sodium sulfate. The organic solvent was removed under reduced pressure and the residue was subjected to column chromatography(eluent: dichloromethane/methanol =9/1, v/v) to give 9_{mg}(0.017 mmol, Yield 43%) of the title compound.

¹H NMR (CDCl₃) δ 1.88(brs, 2H), 2.64(brs, 6H), 4.86(s, 2H), 5.01(s, 2H), 6.67(s, 1H), 7.14(m, 3H), 7.26-7.58(m, 8H), 7.81(m, 2H), 8.03(d, 1H)

FAB MS : 518 (M+1)

Example 114: Synthesis of 3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-1-(1-methyl-1H-imidazol-5-yl)methyl-4-(naphthalen-1-yl)-1H-pyrrole(114)

114-1) 4-(Naphthalen-1-yl)-1H-pyrrole-3-carboxylic acid

2.64g(10 mmol) of the compound prepared in Example 80-2) was dissolved in 50_{ml} of 50% ethanol, and 2.24g(40 mmol) of potassium hydroxide was added thereto. The reaction mixture was refluxed for 7 hours, cooled down to room temperature, adjusted to pH 4-5, extracted with ethyl acetate, dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure to obtain 1.62g(8.1 mmol, Yield 81%) of the title compound. The product thus obtained was directly used in the next reaction without purification.

¹H NMR(CDCl₃) δ 6.60(s, 1H), 7.32-7.49(m, 5H), 7.54(s, 1H), 7.84(m, 2H), 9.92(s, 1H)

FAB (M+H): 236

114-2) 3-[N-(2-Methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole

234_{mg}(1 mmol) of the compound prepared in Example 114-1) was dissolved in 2_{ml} of dimethylformamide, and then 230_{mg}(1.2 mmol) of EDC, 101_{mg}(1 mmol) of triethylamine and 162_{mg}(1.2 mmol) of HOBT were added thereto. The resulting mixture was stirred at 0°C for 5 minutes. To the reaction solution was added 124_{mg}(1 mmol) of N-(2-methoxyethyl)-N-methylamine hydrochloride, which was then stirred at room temperature for 5 hours. The solvent was removed under reduced pressure and then 10_{ml} of saturated potassium carbonate solution was added to the residue. The resulting solution was extracted with 20_{ml} of ethyl acetate, washed with 10_{ml} of 1N aqueous hydrochloric acid

solution, washed with aqueous sodium chloride solution and water, dried over anhydrous sodium sulfate and concentrated to give 246_{mg}(0.8 mmol) of the title compound.

¹H NMR(CDCl₃) δ 2.46(s, 2H), 2.80-3.40(m, 7H), 3.40(s, 1H), 6.80(s, 1H), 7.00(s, 1H), 7.42(m, 4H), 7.73(d, 1H), 7.81(d, 1H), 8.17(d, 1H), 10.66 (s, 1H)

FAB (M+H): 309

114-3) 1-(1-Methyl-1H-imidazol-5-yl)methyl-3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole

618_{mg}(2.0 mmol) of the compound prepared in Example 114-2) was dissolved in 10_{ml} of dimethylformamide, 264_{mg}(6.6 mmol) of sodium hydride(60%) was added thereto at 0°C and then the mixture was stirred for 5 minutes. To the mixture was added 367_{mg}(2.2 mmol) of 5-chloromethyl-1-methylimidazole hydrochloride and the whole mixture was stirred at room temperature for 5 hours. The solvent was removed by distillation under reduced pressure and 10_{ml} of water was added to the residue. The mixture was then extracted twice with 20_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated, and subjected to silica gel column chromatography(eluent: dichloromethane/methanol=90/10, v/v) to obtain 644_{mg}(Yield 80%) of the title compound.

¹H NMR(CDCl₃) δ 2.42(s, 2H), 2.71(m, 1H), 3.10(brs, 6H), 3.30(brs, 1H), 3.50(s, 3H), 5.09(s, 2H), 6.70(s, 1H), 7.05(s, 1H), 7.15(s, 1H), 7.30-7.49 (m, 4H), 7.72(d, 1H), 7.84(d, 2H), 8.08(d, 1H)

FAB (M+H): 403

Example 115: Synthesis of 1-(1-isobutyl-1H-imidazol-5-yl)methyl-3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole (115)

618_{mg}(2.0 mmol) of the compound prepared in Example 114-2) was dissolved in 10_{ml} of dimethylformamide, 264_{mg}(6.6 mmol) of sodium hydride(60%) was added thereto at 0°C, and the mixture was stirred for 5 minutes. 459_{mg}(2.2 mmol) of the compound prepared in Preparation 33-2) was added to the mixture, which was then stirred at room temperature for 5 hours. The solvent was removed by distillation under reduced pressure and 10_{ml} of water was added to the residue. The resulting mixture was extracted twice with 20_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated and subjected to silica gel column chromatography(eluent: dichloromethane/methanol=90/10, v/v) to obtain 667_{mg}(Yield 75%) of the title compound.

¹H NMR(CDCl₃) δ 0.90(d, 6H), 1.75(m, 1H), 2.41(brs, 2H), 2.72(brs, 1H), 3.01(brs, 6H), 3.32(brs, 1H), 3.62(d, 2H), 5.13(s, 2H), 6.72(s, 1H), 7.09(s, 1H), 7.19(s, 1H), 7.30-7.49(m, 4H), 7.78(d, 1H), 7.84(d, 2H), 8.08 d, 1H)

FAB (M+H): 445

Example 116: Synthesis of 1-(1-cyclohexylmethyl-1H-imidazol-5-yl)methyl-3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole (116)

618_{mg}(2.0 mmol) of the compound prepared in Example 114-2) was dissolved in 10_{ml} of dimethylformamide, 264_{mg}(6.6 mmol) of sodium hydride(60%) was added thereto at 0°C, and the mixture was stirred for 5 minutes. 647_{mg}(2.2 mmol) of the compound prepared in Preparation 34-2) was added to the mixture, which was then stirred at room temperature for 5 hours. The solvent was removed by distillation under reduced pressure and 10_{ml} of water was added to the residue. The resulting mixture was extracted twice with 20_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated and subjected to silica

gel column chromatography(eluent: dichloromethane/methanol=90/10, v/v) to obtain 726_{mg}(Yield 75%) of the title compound.

¹H NMR(CDCl₃) δ 0.87(m, 2H), 1.12(m, 3H), 1.30(brs, 1H), 1.40-1.80(m, 5H), 2.41(brs, 2H), 2.72(brs, 1H), 3.01(brs, 6H), 3.32(brs, 1H), 3.63(d, 2H), 5.09(s, 2H), 6.72(s, 1H), 7.09(s, 1H), 7.19(s, 1H), 7.25(s, 1H), 7.30-7.49 (m, 3H), 7.78(d, 1H), 7.83(d, 2H), 8.08(d, 1H)

FAB (M+H): 485

Example 117: Synthesis of 3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)-1-(1-pentyl-1H-imidazol-5-yl)methyl-1H-pyrrole (117)

618_{mg}(2.0 mmol) of the compound prepared in Example 114-2) was dissolved in 10_{ml} of dimethylformamide, 264_{mg}(6.6 mmol) of sodium hydride(60%) was added thereto at 0°C, and the mixture was stirred for 5 minutes. 429_{mg}(2.2 mmol) of the compound prepared in Preparation 35-2) was added to the mixture, which was then stirred at room temperature for 5 hours. The solvent was removed by distillation under reduced pressure and 10_{ml} of water was added to the residue. The resulting mixture was extracted twice with 20_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated and subjected to silica gel column chromatography(eluent: dichloromethane/methanol=90/10, v/v) to obtain 714_{mg}(Yield 78%) of the title compound.

¹H NMR(CDCl₃) δ 0.90(t, 3H), 1.08(brs, 2H), 1.30(m, 2H), 1.45(m, 2H), 2.41(brs, 2H), 2.72(brs, 1H), 3.01(brs, 6H), 3.32(brs, 1H), 3.63(t, 2H), 5.09 (s, 2H), 6.72(s, 1H), 7.09(s, 1H), 7.19(s, 1H), 7.25(s, 1H), 7.30-7.49(m, 3H), 7.78(d, 1H), 7.83(d, 2H), 8.08(d, 1H)

FAB (M+H): 459

Example 118: Synthesis of 3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)-1-(1-octyl-1H-imidazol-5-yl)methyl-1H-pyrrole(118)

618_{mg}(2.0 mmol) of the compound prepared in Example 114-2) was dissolved in 10_{ml} of dimethylformamide, 264_{mg}(6.6 mmol) of sodium hydride(60%) was added thereto at 0°C, and the mixture was stirred for 5 minutes. 508_{mg}(2.2 mmol) of the compound prepared in Preparation 36-2) was added to the mixture, which was then stirred at room temperature for 5 hours. The solvent was removed by distillation under reduced pressure and 10_{ml} of water was added to the residue. The resulting mixture was extracted twice with 20_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated and subjected to silica gel column chromatography(eluent: dichloromethane/methanol=90/10, v/v) to obtain 760_{mg}(Yield 76%) of the title compound.

¹H NMR(CDCl₃) δ 0.87(t, 3H), 1.17(brs, 2H), 1.30(brs, 10H), 1.44(m, 2H), 2.41(brs, 2H), 2.72(brs, 1H), 3.01(brs, 6H), 3.32(brs, 1H), 3.62(t, 2H), 5.09(s, 2H), 6.72(s, 1H), 7.09(s, 1H), 7.19(s, 1H), 7.25(s, 1H), 7.30-7.49 (m, 3H), 7.78(d, 1H), 7.83(d, 2H), 8.08(d, 1H)

FAB (M+H): 501

Example 119: Synthesis of 1-(1-decyl-1H-imidazol-5-yl)methyl-3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole (119)

618_{mg}(2.0 mmol) of the compound prepared in Example 114-2) was dissolved in 10_{ml} of dimethylformamide, 264_{mg}(6.6 mmol) of sodium hydride(60%) was added thereto at 0°C, and the mixture was stirred for 5 minutes. 567_{mg}(2.2 mmol) of the compound prepared in Preparation 37-2) was added to the mixture, which was then stirred at room temperature for 5 hours. The solvent was removed by distillation under reduced pressure and 10_{ml} of water was added to the residue. The resulting mixture was extracted twice with 20_{ml} of ethyl acetate,

dried over anhydrous sodium sulfate, concentrated and subjected to silica gel column chromatography(eluent: dichloromethane/methanol=90/10, v/v) to obtain 667_{mg}(Yield 75%) of the title compound.

¹H NMR(CDCl₃) δ 0.87(t, 3H), 1.17(brs, 2H), 1.30(brs, 14H), 1.44(m, 2H), 2.41(brs, 2H), 2.72(brs, 1H), 3.01(brs, 6H), 3.32(brs, 1H), 3.62(t, 2H), 5.09(s, 2H), 6.72(s, 1H), 7.09(s, 1H), 7.19(s, 1H), 7.25(s, 1H), 7.30-7.49 (m, 3H), 7.78(d, 1H), 7.83(d, 2H), 8.08(d, 1H)

FAB (M+H): 529

Example 120: Synthesis of 3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-1-[1-(3-methylbutyl)-1H-imidazol-5-yl]methyl-4-(naphthalen-1-yl)-1H-pyrrole(120)

618_{mg}(2.0 mmol) of the compound prepared in Example 114-2) was dissolved in 10_{ml} of dimethylformamide, 264_{mg}(6.6 mmol) of sodium hydride(60%) was added thereto at 0°C, and the mixture was stirred for 5 minutes. 429_{mg}(2.2 mmol) of the compound prepared in Preparation 38-2) was added to the mixture, which was then stirred at room temperature for 5 hours. The solvent was removed by distillation under reduced pressure and 10_{ml} of water was added to the residue. The resulting mixture was extracted twice with 20_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated and subjected to silica gel column chromatography(eluent: dichloromethane/methanol=90/10, v/v) to obtain 667_{mg}(Yield 75%) of the title compound.

¹H NMR(CDCl₃) δ 0.91(d, 6H), 1.31(q, 2H), 1.67(m, 1H), 2.41(brs, 2H), 2.72(brs, 1H), 3.01(brs, 6H), 3.32(brs, 1H), 3.62(t, 2H), 5.09(s, 2H), 6.72 (s, 1H), 7.09(s, 1H), 7.19(s, 1H), 7.25(s, 1H), 7.30-7.49(m, 3H), 7.78(d, 1H), 7.83(d, 2H), 8.08(d, 1H)

FAB (M+H): 459

Example 121: Synthesis of 1-[1-(2-methoxyethyl)-1H-imidazol-5-yl]methyl-3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole (121)

618_{mg}(2.0 mmol) of the compound prepared in Example 114-2) was dissolved in 10_{ml} of dimethylformamide, 264_{mg}(6.6 mmol) of sodium hydride(60%) was added thereto at 0°C, and the mixture was stirred for 5 minutes. 429_{mg}(2.2 mmol) of the compound prepared in Preparation 39-2) was added to the mixture, which was then stirred at room temperature for 5 hours. The solvent was removed by distillation under reduced pressure and 10_{ml} of water was added to the residue. The resulting mixture was extracted twice with 20_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated and subjected to silica gel column chromatography(eluent: dichloromethane/methanol=90/10, v/v) to obtain 667_{mg}(Yield 75%) of the title compound.

¹H NMR(CDCl₃) δ 2.41(brs, 2H), 2.72(brs, 1H), 3.01(brs, 6H), 3.32(brs, 1H), 3.37(s, 3H), 3.45(t, 2H), 3.63(t, 2H), 5.09(s, 2H), 6.72(s, 1H), 7.09 (s, 1H), 7.19(s, 1H), 7.25(s, 1H), 7.30-7.49(m, 3H), 7.78(d, 1H), 7.83(d, 2H), 8.08(d, 1H)

FAB (M+H): 447

Example 122: Synthesis of 3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-1-[1-(3-methoxypropyl)-1H-imidazol-5-yl]methyl-4-(naphthalen-1-yl)-1H-pyrrole (122)

618_{mg}(2.0 mmol) of the compound prepared in Example 114-2) was dissolved in 10_{ml} of dimethylformamide, 264_{mg}(6.6 mmol) of sodium hydride(60%) was added thereto at 0°C, and the mixture was stirred for 5 minutes. 459_{mg}(2.2 mmol) of the compound prepared in Preparation 40-2) was added to the mixture, which was then stirred at

room temperature for 5 hours. The solvent was removed by distillation under reduced pressure and 10_{ml} of water was added to the residue. The resulting mixture was extracted twice with 20_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated and subjected to silica gel column chromatography(eluent: dichloromethane/methanol=90/10, v/v) to obtain 683_{mg}(Yield 70%) of the title compound.

¹H NMR(CDCl₃) δ 1.71(m, 2H), 2.41(brs, 2H), 2.72(brs, 1H), 3.01(brs, 6H), 3.31(s, 3H), 3.32(brs, 1H), 3.48(t, 2H), 3.63(t, 2H), 5.09(s, 2H), 6.72 (s, 1H), 7.09(s, 1H), 7.19(s, 1H), 7.25(s, 1H), 7.30-7.49(m, 3H), 7.78(d, 1H), 7.83(d, 2H), 8.08(d, 1H)

FAB (M+H): 461

Example 123: Synthesis of 1-[1-(3-ethoxypropyl)-1H-imidazol-5-yl]methyl-3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole (123)

618_{mg}(2.0 mmol) of the compound prepared in Example 114-2) was dissolved in 10_{ml} of dimethylformamide, 264_{mg}(6.6 mmol) of sodium hydride(60%) was added thereto at 0°C, and the mixture was stirred for 5 minutes. 459_{mg}(2.2 mmol) of the compound prepared in Preparation 41-2) was added to the mixture, which was then stirred at room temperature for 5 hours. The solvent was removed by distillation under reduced pressure and 10_{ml} of water was added to the residue. The resulting mixture was extracted twice with 20_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated and subjected to silica gel column chromatography(eluent: dichloromethane/methanol=90/10, v/v) to obtain 712_{mg}(Yield 71%) of the title compound.

¹H NMR(CDCl₃) δ 1.20(t, 3H), 1.70(m, 2H), 2.41(brs, 2H), 2.72(brs, 1H), 3.01(brs, 6H), 3.32(brs, 1H), 3.50(m, 4H), 3.63(t, 2H), 5.09(s, 2H), 6.72(s, 1H), 7.09(s, 1H), 7.19(s, 1H), 7.25(s, 1H),

7.30-7.49(m, 3H), 7.78(d, 1H), 7.83(d, 2H), 8.08(d, 1H)

FAB (M+H): 475

Example 124: Synthesis of 1-[1-(3-isopropoxypropyl)-1H-imidazol-5-yl]methyl-3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole (124)

618_{mg}(2.0 mmol) of the compound prepared in Example 114-2) was dissolved in 10_{ml} of dimethylformamide, 264_{mg}(6.6 mmol) of sodium hydride(60%) was added thereto at 0°C, and the mixture was stirred for 5 minutes. 459_{mg}(2.2 mmol) of the compound prepared in Preparation 42-2) was added to the mixture, which was then stirred at room temperature for 5 hours. The solvent was removed by distillation under reduced pressure and 10_{ml} of water was added to the residue. The resulting mixture was extracted twice with 20_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated and subjected to silica gel column chromatography(eluent: dichloromethane/methanol=90/10, v/v) to obtain 751_{mg}(Yield 73%) of the title compound.

¹H NMR(CDCl₃) δ 1.16(d, 6H), 1.70(m, 2H), 2.41(brs, 2H), 2.72(brs, 1H), 3.01(brs, 6H), 3.32(brs, 1H), 3.45-3.55(m, 3H), 3.63(t, 2H), 5.09(s, 2H), 6.72(s, 1H), 7.09(s, 1H), 7.19(s, 1H), 7.25(s, 1H), 7.30-7.49(m, 3H), 7.78 (d, 1H), 7.83(d, 2H), 8.08(d, 1H)

FAB (M+H): 489

Example 125: Synthesis of 1-[1-(4-bromobenzyl)-1H-imidazol-5-yl]methyl-3-[4-methylpiperazin-1-yl]carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(125)

125-1) 3-[4-Methylpiperazin-1-yl]carbonyl-4-(naphthalen-1-yl)-1H-pyrrole

The title compound was obtained in a yield of 90% according to

the same procedure as Example 80-5) from the compound prepared in Example 106-2) and 4-methylpiperazine.

^1H NMR (CDCl_3) δ 1.15(br, 2H), 1.87(br, 2H), 1.92(s, 3H), 2.96(br, 2H), 3.41(br, 2H), 6.83(s, 1H), 7.09(s, 1H), 7.36-7.42(m, 4H), 7.73(d, 1H), 7.75 (d, 1H), 8.10(d, 1H), 10.52(s, 1H)

FAB(M+H): 320

125-2) 1-[1-(4-Bromobenzyl)-1H-imidazol-5-yl]methyl-3-[4-methylpiperazin-1-yl]carbonyl-4-(naphthalen-1-yl)-1H-pyrrole

64_{mg}(0.2 mmol) of the compound prepared in Example 125-1) was dissolved in 2_{ml} of dimethylformamide, 26_{mg}(0.66 mmol) of sodium hydride was added thereto at 0°C, and the mixture was stirred for 5 minutes. 66_{mg}(0.22 mmol) of the compound prepared in Preparation 32-2) was added to the mixture, which was then stirred at room temperature for 5 hours. The solvent was removed by distillation under reduced pressure and 10_{ml} of water was added to the residue. The resulting mixture was extracted twice with 10_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated and subjected to silica gel column chromatography(eluent: dichloromethane/methanol=90/10, v/v) to obtain 91_{mg}(Yield 80%) of the title compound.

^1H NMR (CDCl_3) δ 1.15(br, 2H), 1.77(br, 2H), 1.86(s, 3H), 2.82(br, 2H), 3.28(br, 2H), 4.87(s, 2H), 3.88(s, 2H), 6.55(s, 1H), 6.79(d, 2H), 6.97(s, 1H), 7.16(s, 1H), 7.36(d, 1H), 7.36-7.39(m, 5H), 7.50(s, 1H), 7.71(d, 1H), 7.79(d, 1H), 7.93(d, 1H)

FAB(M+H): 568

Example 126: Synthesis of 1-[1-(4-chlorobenzyl)-1H-imidazol-5-yl]methyl-3-[4-methylpiperazin-1-yl]carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(126)

64_{mg}(0.2 mmol) of the compound prepared in Example 125-1) was dissolved in 2_{ml} of dimethylformamide, 26_{mg}(0.66 mmol) of sodium hydride was added thereto at 0°C, and the mixture was stirred for 5 minutes. 55_{mg}(0.22 mmol) of 1-(4-chlorobenzyl)-5-chloromethylimidazole hydrochloride prepared according to the similar procedure as Preparation 32 was added to the mixture, which was then stirred at room temperature for 5 hours. The solvent was removed by distillation under reduced pressure and 10_{ml} of water was added to the residue. The resulting mixture was extracted twice with 10_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated and subjected to silica gel column chromatography(eluent: dichloromethane/methanol=90/10, v/v) to obtain 77_{mg}(Yield 57%) of the title compound.

¹H NMR (CDCl₃) δ 1.15(br, 2H), 1.77(br, 2H), 1.86(s, 3H), 2.82(br, 2H), 3.28(br, 2H), 4.92(s, 2H), 4.95(s, 2H), 6.60(s, 1H), 6.91(d, 2H), 6.01(s, 1H), 7.22(s, 1H), 7.26-7.36(m, 3H), 7.36-7.48(m, 2H), 7.56(s, 1H), 7.77(d, 1H), 7.82(d, 1H), 7.93(d, 1H)

FAB(M+H): 524

Example 127: Synthesis of 1-[1-(4-fluorobenzyl)-1H-imidazol-5-yl]methyl-3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole(127)

62_{mg}(0.2 mmol) of the compound prepared in Example 114-2) was dissolved in 2_{ml} of dimethylformamide, 26_{mg}(0.66 mmol) of sodium hydride was added thereto at 0°C, and the mixture was stirred for 5 minutes. 51_{mg}(0.22 mmol) of 1-(4-fluorobenzyl)-5-chloromethylimidazole hydrochloride prepared according to the similar procedure as Preparation 32 was added to the mixture, which was then stirred at room temperature for 5 hours. The solvent was removed by distillation under reduced pressure and 10_{ml} of water was added to the residue. The

resulting mixture was extracted twice with 10_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated and subjected to silica gel column chromatography(eluent: dichloromethane/methanol=90/10, v/v) to obtain 77_{mg}(Yield 80%) of the title compound.

¹H NMR(CDCl₃) δ 2.12(br, 3H), 2.72(br, 1H), 3.00-3.20(m, 5H), 3.32(s, 1H), 4.97(s, 2H), 3.98(s, 2H), 6.64(s, 1H), 6.95-7.10(m, 5H), 7.21(s, 1H), 7.33(m, 1H), 7.40-7.51(m, 3H), 7.66(s, 1H), 7.74(d, 1H), 7.81(d, 1H), 8.08(d, 1H)

FAB(M+H):497

Example 128: Synthesis of 1-[1-(4-fluorobenzyl)-1H-imidazol-5-yl]methyl-3-[4-methylpiperazin-1-yl]carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(128)

64_{mg}(0.2 mmol) of the compound prepared in Example 125-1) was dissolved in 2_{ml} of dimethylformamide, 26_{mg}(0.66 mmol) of sodium hydride was added thereto at 0°C, and the mixture was stirred for 5 minutes. 51_{mg}(0.22 mmol) of 1-(4-fluorobenzyl)-5-chloromethylimidazole hydrochloride prepared according to the similar procedure as Preparation 32 was added to the mixture, which was then stirred at room temperature for 5 hours. The solvent was removed by distillation under reduced pressure and 10_{ml} of water was added to the residue. The resulting mixture was extracted twice with 10_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated and subjected to silica gel column chromatography(eluent: dichloromethane/methanol=90/10, v/v) to obtain 79_{mg}(Yield 80%) of the title compound.

¹H NMR (CDCl₃) δ 1.15(br, 2H), 1.77(br, 2H), 1.86(s, 3H), 2.82(br, 2H), 3.28(br, 2H), 4.92(s, 2H), 4.97(s, 2H), 6.60(s, 1H), 6.93(d, 2H), 6.01(s, 1H), 7.22(s, 1H), 7.25-7.36(m, 3H), 7.36-7.47(m, 2H), 7.57(s, 1H), 7.78(d, 1H), 7.82(d, 1H), 7.93(d, 1H)

FAB (M+H) 508

Preparation 43: Synthesis of 5-chloromethyl-1-(4-methoxybenzyl)-imidazole hydrochloride

43-1) 5-Hydroxymethyl-1-(4-methoxybenzyl)imidazole

The title compound was obtained in a yield of 30% according to the same procedure as Preparation 31-1) using dihydroxyacetone and 4-methoxybenzylamine hydrochloride as starting materials.

^1H NMR($\text{CDCl}_3+\text{CD}_3\text{OD}$) δ 3.75(s, 3H), 4.50(s, 2H), 5.15(s, 2H), 6.86(m, 3H), 7.08(d, 2H), 7.42(s, 1H)

FAB(M+H):219

43-2) 5-Chloromethyl-1-(4-methoxybenzyl)imidazole hydrochloride

The title compound was obtained in a yield of 95% according to the same procedure as Preparation 28-2) except that the compound prepared in Preparation 43-1) was used as a starting material.

Preparation 44: Synthesis of 5-chloromethyl-1-(3-chlorobenzyl)-imidazole hydrochloride

44-1) 5-Hydroxymethyl-1-(3-chlorobenzyl)imidazole

The title compound was obtained in a yield of 60% according to the same procedure as Preparation 31-1) using dihydroxyacetone and 3-chlorobenzylamine hydrochloride as starting materials.

^1H NMR($\text{CDCl}_3+\text{CD}_3\text{OD}$) δ 3.81(s, 3H), 4.47(s, 2H), 5.25(s, 2H), 6.99(s, 1H), 7.05(m, 1H), 7.14(s, 1H), 7.30(d, 2H), 7.61(s, 1H)

FAB(M+H):239.5

44-2) 5-Chloromethyl-1-(3-chlorobenzyl)imidazole hydrochloride

The title compound was obtained in a yield of 92% according to the same procedure as Preparation 28-2) except that the compound prepared in Preparation 44-1) was used as a starting material.

Preparation 45: Synthesis of 5-chloromethyl-1-(2-chlorobenzyl)imidazole hydrochloride

45-1) 5-Hydroxymethyl-1-(2-chlorobenzyl)imidazole

The title compound was obtained in a yield of 60% according to the same procedure as Preparation 31-1) using dihydroxyacetone and 2-chlorobenzylamine hydrochloride as starting materials.

^1H NMR(CDCl_3) δ 3.24(s, 2H), 4.44(s, 2H), 5.26(s, 2H), 6.78(d, 1H), 6.90(s, 1H), 7.15(m, 1H), 7.21(m, 1H), 7.34(d, 1H), 7.38(s, 1H)

FAB(M+H):239.5

45-2) 5-Chloromethyl-1-(2-chlorobenzyl)imidazole hydrochloride

The title compound was obtained in a yield of 92% according to the same procedure as Preparation 28-2) except that the compound prepared in Preparation 45-1) was used as a starting material.

Preparation 46: Synthesis of 5-chloromethyl-1-(2-fluorobenzyl)imidazole hydrochloride

46-1) 5-Hydroxymethyl-1-(2-fluorobenzyl)imidazole

The title compound was obtained in a yield of 71% according to the same procedure as Preparation 31-1) using dihydroxyacetone and 2-fluorobenzylamine hydrochloride as starting materials.

^1H NMR(CDCl_3) δ 3.25(s, 2H), 4.45(s, 2H), 5.27(s, 2H), 6.79(d, 1H), 7.17(m, 1H), 7.26(m, 1H), 7.35(d, 1H), 7.38(s, 1H)

FAB(M+H): 223

46-2) 5-Chloromethyl-1-(2-fluorobenzyl)imidazole hydrochloride

The title compound was obtained in a yield of 93% according to the same procedure as Preparation 28-2) except that the compound prepared in Preparation 46-1) was used as a starting material.

Preparation 47: Synthesis of 5-chloromethyl-1-(4-methylbenzyl)imidazole hydrochloride**47-1) 5-Hydroxymethyl-1-(4-methylbenzyl)imidazole**

The title compound was obtained in a yield of 65% according to the same procedure as Preparation 31-1) using dihydroxyacetone and 4-methylbenzylamine hydrochloride as starting materials.

^1H NMR(CDCl_3) δ 2.32(s, 3H), 4.50(s, 2H), 5.19(s, 2H), 6.95(s, 1H), 7.05(d, 2H), 7.15(d, 2H), 7.59(s, 1H)

FAB(M+H): 219

47-2) 5-Chloromethyl-1-(4-methylbenzyl)imidazole hydrochloride

The title compound was obtained in a yield of 91% according to the same procedure as Preparation 28-2) except that the compound prepared in Preparation 47-1) was used as a starting material.

Preparation 48: Synthesis of 5-chloromethyl-1-(3-methylbenzyl)imidazole hydrochloride**48-1) 5-Hydroxymethyl-1-(3-methylbenzyl)imidazole**

The title compound was obtained in a yield of 60% according to the same procedure as Preparation 31-1) using dihydroxyacetone and 3-methylbenzylamine hydrochloride as starting materials.

^1H NMR(CDCl_3) δ 2.27(s, 3H), 4.45(s, 2H), 4.52(br, 1H), 5.13(s,

2H), 6.80(d, 1H), 6.90(m, 2H), 7.08(m, 1H), 7.17(m, 1H), 7.34(s, 1H)

FAB(M+H): 219

48-2) 5-Chloromethyl-1-(3-methylbenzyl)imidazole hydrochloride

The title compound was obtained in a yield of 92% according to the same procedure as Preparation 28-2) except that the compound prepared in Preparation 48-1) was used as a starting material.

Example 129: Synthesis of 1-[1-(4-methoxybenzyl)-1H-imidazol-5-yl]methyl-3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole(129)

62_{mg}(0.2 mmol) of the compound prepared in Example 114-2) was dissolved in 2_{mL} of dimethylformamide, 26_{mg}(0.66 mmol) of sodium hydride was added thereto at 0°C, and the mixture was stirred for 5 minutes. 60_{mg}(0.22 mmol) of the compound prepared in Preparation 43-2) was added to the mixture, which was then stirred at room temperature for 5 hours. The solvent was removed by distillation under reduced pressure and 10_{mL} of water was added to the residue. The resulting mixture was extracted twice with 10_{mL} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated and subjected to silica gel column chromatography(eluent: dichloromethane/methanol=95/5, v/v) to obtain 77_{mg}(Yield 76%) of the title compound.

¹HNMR(CDCl₃) δ 2.41(m, 2H), 2.75(m, 1H), 3.03(m, 5H), 3.10(m, 1H), 3.34(m, 1H), 3.76(m, 3H), 4.91(s, 2H), 4.93(s, 2H), 6.62(d, 1H), 6.82(d, 2H), 6.90-7.07(m, 3H), 7.21(s, 1H), 7.32(m, 1H), 7.43(m, 2H), 7.60(s, 1H), 7.74(d, 1H), 7.82(d, 1H), 8.08(d, 1H)

FAB (M+H): 509, C₃₁H₃₂N₄O₃

Example 130: Synthesis of 1-[1-(4-methoxybenzyl)-1H-imidazol-5-yl]

methyl-3-(4-methylpiperazin-1-yl)carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(130)

64_{mg}(0.2 mmol) of the compound prepared in Example 125-1) was dissolved in 2_{ml} of dimethylformamide, 26_{mg}(0.66 mmol) of sodium hydride was added thereto at 0°C, and the mixture was stirred for 5 minutes. 60_{mg}(0.22 mmol) of the compound prepared in Preparation 43-2) was added to the mixture, which was then stirred at room temperature for 5 hours. The solvent was removed by distillation under reduced pressure and 10_{ml} of water was added to the residue. The resulting mixture was extracted twice with 10_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated and subjected to silica gel column chromatography(eluent: dichloromethane/methanol=90/10, v/v) to obtain 79_{mg}(Yield 80%) of the title compound.

¹H NMR(CDCl₃) δ 1.06(br, 2H), 1.72(m, 2H), 1.82(s, 3H), 2.86(br, 2H), 3.28(br, 2H), 3.75(s, 3H), 4.91(s, 2H), 4.93(s, 2H), 6.63(d, 1H), 6.82(d, 2H), 6.90-7.07(m, 3H), 7.23(s, 1H), 7.33(m, 1H), 7.44(m, 2H), 7.61(s, 1H), 7.75(d, 1H), 7.82(d, 1H), 8.08(d, 1H)

FAB (M+H): 520, C₃₂H₃₃N₅O₂

Example 131: Synthesis of 1-[1-(3-chlorobenzyl)-1H-imidazol-5-yl]methyl-3-(4-methylpiperazin-1-yl)carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(131)

64_{mg}(0.2 mmol) of the compound prepared in Example 125-1) was dissolved in 2_{ml} of dimethylformamide, 26_{mg}(0.66 mmol) of sodium hydride was added thereto at 0°C, and the mixture was stirred for 5 minutes. 61_{mg}(0.22 mmol) of the compound prepared in Preparation 44-2) was added to the mixture, which was then stirred at room temperature for 5 hours. The solvent was removed by distillation under

reduced pressure and 10_{ml} of water was added to the residue. The resulting mixture was extracted twice with 10_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated and subjected to silica gel column chromatography(eluent: dichloromethane/methanol=90/10, v/v) to obtain 75_{mg}(Yield 71%) of the title compound.

¹H NMR(CDCl₃) δ 1.02(br, 2H), 1.78(br, 2H), 1.87(s, 3H), 2.84(br, 2H), 3.30(br, 2H), 6.64(m, 2H), 7.01(s, 1H), 7.10-7.30(m, 4H), 7.31-7.47(m, 4H), 7.53(s, 1H), 7.73(d, 1H), 7.81(d, 1H), 7.96(d, 1H)

FAB (M+H): 524, C₃₁H₃₀N₅OCl

Example 132: Synthesis of 1-[1-(3-chlorobenzyl)-1H-imidazol-5-yl]methyl-3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole(132)

62_{mg}(0.2 mmol) of the compound prepared in Example 114-2) was dissolved in 2_{ml} of dimethylformamide, 26_{mg}(0.66 mmol) of sodium hydride was added thereto at 0°C, and the mixture was stirred for 5 minutes. 61_{mg}(0.22 mmol) of the compound prepared in Preparation 44-2) was added to the mixture, which was then stirred at room temperature for 5 hours. The solvent was removed by distillation under reduced pressure and 10_{ml} of water was added to the residue. The resulting mixture was extracted twice with 10_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated and subjected to silica gel column chromatography(eluent: dichloromethane/methanol=95/5, v/v) to obtain 80_{mg}(Yield 78%) of the title compound.

¹H NMR(CDCl₃) δ 2.39(br, 2H), 2.71(m, 1H), 3.02(br, 4H), 3.09(br, 1H), 3.32(br, 1H), 4.09(br, 1H), 4.97(s, 2H), 5.04(s, 2H), 6.64(d, 1H), 6.90(m, 1H), 7.02(d, 2H), 7.20-7.40(m, 4H), 7.40-7.60(m, 3H), 7.74(d, 1H), 7.76(d, 1H), 7.85(s, 1H), 8.04(d, 1H)

FAB (M+H): 513, C₃₀H₂₉N₄O₂Cl

Example 133: Synthesis of 1-[1-(2-chlorobenzyl)-1H-imidazol-5-yl]methyl-3-(4-methylpiperazin-1-yl)carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(133)

64_{mg}(0.2 mmol) of the compound prepared in Example 125-1) was dissolved in 2_{ml} of dimethylformamide, 26_{mg}(0.66 mmol) of sodium hydride was added thereto at 0°C, and the mixture was stirred for 5 minutes. 61_{mg}(0.22 mmol) of the compound prepared in Preparation 45-2) was added to the mixture, which was then stirred at room temperature for 5 hours. The solvent was removed by distillation under reduced pressure and 10_{ml} of water was added to the residue. The resulting mixture was extracted twice with 10_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated and subjected to silica gel column chromatography(eluent: dichloromethane/methanol=90/10, v/v) to obtain 80_{mg}(Yield 76%) of the title compound.

¹H NMR(CDCl₃) δ 1.06(br, 2H), 1.80(br, 2H), 1.86(s, 3H), 2.84(br, 2H), 3.30(br, 2H), 4.98(s, 2H), 5.11(s, 2H), 6.63(m, 2H), 7.01(s, 1H), 7.12-7.30 (m, 4H), 7.32-7.46(m, 4H), 7.53(s, 1H), 7.73(d, 1H), 7.81(d, 1H), 7.97(d, 1H)

FAB (M+H): 524, C₃₁H₃₀N₅OCl

Example 134: Synthesis of 1-[1-(2-chlorobenzyl)-1H-imidazol-5-yl]methyl-3-[N-(2-methoxyethyl)-N-methyl]carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(134)

62_{mg}(0.2 mmol) of the compound prepared in Example 114-2) was dissolved in 2_{ml} of dimethylformamide, 26_{mg}(0.66 mmol) of sodium hydride was added thereto at 0°C, and the mixture was stirred for 5 minutes. 61_{mg}(0.22 mmol) of the compound prepared in Preparation

45-2) was added to the mixture, which was then stirred at room temperature for 5 hours. The solvent was removed by distillation under reduced pressure and 10_{ml} of water was added to the residue. The resulting mixture was extracted twice with 10_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated and subjected to silica gel column chromatography(eluent: dichloromethane/methanol=95/5, v/v) to obtain 77_{mg}(Yield 75%) of the title compound.

¹H NMR(CDCl₃) δ 2.37(br, 2H), 2.72(m, 1H), 3.01(br, 4H), 3.10(br, 1H), 3.32(br, 1H), 4.18(br, 1H), 5.04(s, 2H), 5.17(s, 2H), 6.65(d, 1H), 6.76(d, 2H), 7.04(d, 1H), 7.13-7.35(m, 4H), 7.36-7.50(m, 4H), 7.71(s, 1H), 7.75(d, 1H), 7.82(d, 1H), 8.01(d, 1H)

FAB (M+H): 513, C₃₀H₂₉N₄O₂Cl

Example 135: Synthesis of 1-[1-(2-fluorobenzyl)-1H-imidazol-5-yl] methyl-3-(4-methylpiperazin-1-yl)carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(135)

64_{mg}(0.2 mmol) of the compound prepared in Example 125-1) was dissolved in 2_{ml} of dimethylformamide, 26_{mg}(0.66 mmol) of sodium hydride was added thereto at 0°C, and the mixture was stirred for 5 minutes. 51_{mg}(0.22 mmol) of the compound prepared in Preparation 46-2) was added to the mixture, which was then stirred at room temperature for 5 hours. The solvent was removed by distillation under reduced pressure and 10_{ml} of water was added to the residue. The resulting mixture was extracted twice with 10_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated and subjected to silica gel column chromatography(eluent: dichloromethane/methanol=90/10, v/v) to obtain 79_{mg}(Yield 77%) of the title compound.

¹H NMR(CDCl₃) δ 1.06(br, 2H), 1.80(br, 2H), 1.86(s, 3H), 2.93(br, 2H), 3.35(br, 2H), 5.03(s, 2H), 5.06(s, 2H), 6.66(m, 2H), 6.87(m,

1H), 7.12-7.30 (m, 4H), 7.32-7.46(m, 4H), 7.58(s, 1H), 7.77(d, 1H), 7.82(d, 1H), 7.97(d, 1H)

FAB (M+H): 508, C₃₁H₃₀N₅O₂

Example 136: Synthesis of 1-[1-(4-methylbenzyl)-1H-imidazol-5-yl] methyl-3-(4-methylpiperazin-1-yl)carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(136)

64_{mg}(0.2 mmol) of the compound prepared in Example 125-1) was dissolved in 2_{ml} of dimethylformamide, 26_{mg}(0.66 mmol) of sodium hydride was added thereto at 0°C, and the mixture was stirred for 5 minutes. 57_{mg}(0.22 mmol) of the compound prepared in Preparation 47-2) was added to the mixture, which was then stirred at room temperature for 5 hours. The solvent was removed by distillation under reduced pressure and 10_{ml} of water was added to the residue. The resulting mixture was extracted twice with 10_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated and subjected to silica gel column chromatography(eluent: dichloromethane/methanol=90/10, v/v) to obtain 81_{mg}(Yield 80%) of the title compound.

¹H NMR(CDCl₃) δ 1.09(br, 2H), 1.83(br, 2H), 1.86(s, 3H), 2.24(s, 3H), 2.93(br, 2H), 3.30(br, 2H), 4.86(s, 2H), 4.91(s, 2H), 6.59(d, 1H), 6.87(m, 2H), 7.01(s, 1H), 7.07(d, 2H), 7.15(s, 1H), 7.25(m, 1H), 7.50(m, 3H), 7.53(s, 1H), 7.73(d, 1H), 7.78(d, 1H), 7.97(d, 1H)

FAB (M+H): 504, C₃₂H₃₃N₅O

Example 137: Synthesis of 1-[1-(4-methylbenzyl)-1H-imidazol-5-yl] methyl-3-(morpholin-4-yl)carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(137)

62_{mg}(0.2 mmol) of the compound prepared in Example 106-3) was dissolved in 2_{ml} of dimethylformamide, 26_{mg}(0.66 mmol) of sodium

hydride was added thereto at 0°C, and the mixture was stirred for 5 minutes. 57_{mg}(0.22 mmol) of the compound prepared in Preparation 47-2) was added to the mixture, which was then stirred at room temperature for 5 hours. The solvent was removed by distillation under reduced pressure and 10_{ml} of water was added to the residue. The resulting mixture was extracted twice with 10_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated and subjected to silica gel column chromatography(eluent: dichloromethane/methanol=95/5, v/v) to obtain 80_{mg}(Yield 81%) of the title compound.

¹H NMR(CDCl₃) δ 2.29(s, 3H), 2.30-3.60(br, 8H), 4.94(s, 1H), 4.99(s, 2H), 6.61(d, 1H), 6.91(d, 1H), 7.07(d, 1H), 7.12(d, 2H), 7.21(s, 1H), 7.32(d, 1H), 7.35-7.50(m, 4H), 7.71(s, 1H), 7.77(d, 1H), 7.84(d, 1H), 7.98(d, 1H)

FAB (M+H): 491, C₃₁H₃₀N₄O₂

Example 138: Synthesis of 1-[1-(3-methylbenzyl)-1H-imidazol-5-yl]methyl-3-(4-methylpiperazin-1-yl)carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(138)

64_{mg}(0.2 mmol) of the compound prepared in Example 125-1) was dissolved in 2_{ml} of dimethylformamide, 26_{mg}(0.66 mmol) of sodium hydride was added thereto at 0°C, and the mixture was stirred for 5 minutes. 57_{mg}(0.22 mmol) of the compound prepared in Preparation 48-2) was added to the mixture, which was then stirred at room temperature for 5 hours. The solvent was removed by distillation under reduced pressure and 10_{ml} of water was added to the residue. The resulting mixture was extracted twice with 10_{ml} of ethyl acetate, dried over anhydrous sodium sulfate, concentrated and subjected to silica gel column chromatography(eluent: dichloromethane/methanol=90/10, v/v) to obtain 74_{mg}(Yield 73%) of the title compound.

^1H NMR(CDCl_3) δ 1.06(br, 2H), 1.80(br, 2H), 1.84(s, 3H), 2.91(br, 2H), 3.27(br, 2H), 4.86(s, 2H), 4.89(s, 2H), 6.57(d, 1H), 6.71(d, 1H), 6.77 (s, 1H), 6.97(s, 1H), 7.01(d, 1H), 7.15(m, 2H), 7.25(d, 1H), 7.37(m, 3H), 7.51(s, 1H), 7.70(d, 1H), 7.72(d, 1H), 7.98(d, 1H).

FAB (M+H): 504, $\text{C}_{32}\text{H}_{33}\text{N}_5\text{O}$

Preparation 49: Synthesis of 3-(naphthalen-1-yl)carbonyl-1H-pyrrole

49-1) Methyl N-methyl-1-naphthalen hydroxamate

3.44g(20 mmol) of 1-naphthoic acid was dissolved in 20 mL of dimethylformamide, and then 4.6g(24 mmol) of EDC, 2.02g(20 mmol) of triethylamine and 3.24g(24 mmol) of HOBT were added thereto. The resulting mixture was stirred at 0 $^\circ\text{C}$ for 5 minutes. To the reaction solution was added 1.85g(20 mmol) of N,O-dimethylhydroxylamine hydrochloride, which was then stirred at room temperature for 5 hours. The solvent was removed under reduced pressure and then 100 mL of saturated potassium carbonate solution was added to the residue. The resulting solution was extracted with ethyl acetate. Then, the organic layer was washed sequentially with 1N aqueous hydrochloric acid solution, aqueous sodium chloride solution and water, dried over anhydrous sodium sulfate and concentrated to give 3.04g(1.50 mmol) of the title compound.

^1H NMR(CDCl_3) δ 2.42(s, 3H), 3.24(s, 3H), 7.47(m, 4H), 7.67(d, 1H), 7.74(m, 2H),

FAB 216 (M+H)

49-2) 1-(Naphthalen-1-yl)-prop-2-en-1-one

2.03g(9.4 mmol) of the compound prepared in Preparation 49-1) was dissolved in 20 mL of dry tetrahydrofuran, and then 20 mL of 1N vinylmagnesiumbromide-tetrahydrofuran solution was added slowly thereto

at 0°C. The mixture was stirred at room temperature for 30 minutes and 20_{ml} of 1N hydrochloric acid was added thereto, and then the resulting solution was extracted with 50_{ml} of ethyl acetate. The organic layer was dried over anhydrous magnesium sulfate and the solvent was removed under reduced pressure to give 1.63g(9 mmol; Yield 96%) of the title compound.

¹H NMR(CDCl₃) δ 6.92(m, 1H), 7.51(m, 4H), 7.74(d, 1H), 7.85(m, 2H), 7.98(d, 1H), 8.31(d, 1H)

49-3) 3-(Naphthalen-1-yl)carbonyl-1H-pyrrole

901_{mg}(5 mmol) of the compound prepared in Preparation 49-2) and 1.01g(5.5 mmol) of tosylmethylisocyanide were dissolved in 10_{ml} of tetrahydrofuran. 555_{mg}(5.5 mmol) of potassium t-butoxide dissolved in 10_{ml} of tetrahydrofuran was slowly added thereto and the mixture was stirred for 30 minutes. 10_{ml} of water was added to the reaction solution to stop the reaction and the solvent was removed under reduced pressure. 20_{ml} of water was added to the residue and the resulting mixture was extracted with ethyl acetate, washed with aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The solvent was removed under reduced pressure and the residue was subjected to silica gel column chromatography(eluent: ethyl acetate/hexane=1/3, v/v) to obtain 884_{mg}(4 mmol, Yield 80%) of the title compound.

¹H NMR(CDCl₃) δ 6.57(s, 1H), 6.66(s, 1H), 6.79(s, 1H), 7.36(m, 3H), 7.48(d, 1H), 7.77(d, 1H), 7.82(d, 1H), 8.04(d, 1H), 9.91(s, 1H)

Preparation 50: Synthesis of 4-(naphthalen-1-yl)carbonyl-3-[N-(2-methoxyethyl)-N-methylcarbamoyl]-1H-pyrrole

50-1) 4-(Naphthalen-1-yl)-4-oxo-2-butenic acid

5.88g(60 mmol) of dry maleic acid was dissolved in 100_{ml} of dry tetrahydrofuran and the mixture was cooled down to 78°C. 4.14g(20 mmol) of 1-bromonaphthalene was dissolved in 100_{ml} of dry tetrahydrofuran and 13.8_{ml} of 1.6N n-butyllithium-hexane solution was added thereto at 78°C. This reaction solution was stirred for 5 minutes and then it was added to the dry maleic acid solution prepared in advance using cannula. The resulting mixture was stirred for 10 minutes, and water was added thereto to stop the reaction. The solvent was removed under reduced pressure, and the residue was acidified by 1N aqueous hydrochloric acid solution and extracted with ethyl acetate. The organic layer was washed with water and aqueous sodium chloride solution, dried over anhydrous magnesium sulfate, concentrated under reduced pressure and subjected to column chromatography(eluent: ethyl acetate/hexane=2/1, v/v) to give 1.35g(6.0 mmol; Yield 30%) of the title compound.

¹H NMR(CDCl₃) δ 6.81(d, 1H), 7.52-7.65(m, 3H), 7.85(d, 1H), 7.89(d, 1H), 7.92(d, 1H), 8.06(d, 1H), 8.56(d, 1H)

50-2)N-(2-methoxyethyl)-N-methyl-4-(naphthalen-1-yl)-4-oxo-2-butenamide

1.3g(5.9 mmol) of the compound prepared in Preparation 50-1) was dissolved in 10_{ml} of dimethylformamide, and then 1.7g(8.9 mmol) of EDC and 1.2g(8.9 mmol) of HOBT were added thereto at 0°C. The resulting mixture was stirred for 5 minutes. To the reaction solution were added 530_{mg}(5.9 mmol) of N-(2-methoxyethyl)-N-methylamine and 1.2_{ml}(8.9 mmol) of triethylamine, the mixture of which was then stirred at room temperature for 2 hours. The solvent was removed under reduced pressure and then 50_{ml} of water was added to the residue. The resulting solution was extracted with ethyl acetate. The organic layer was washed with aqueous sodium chloride solution, dried over anhydrous magnesium sulfate and concentrated under reduced pressure. The residue

was subjected to column chromatography(eluent: ethyl acetate/hexane=1/1, v/v) to give 1.4g(4.7 mmol; Yield 80%) of the title compound.

^1H NMR(CDCl_3) δ 3.05(s, 3H), 3.32(s, 3H), 3.54(m, 2H), 3.65(m, 2H), 7.40-7.58(m, 4H), 7.71(t, 1H), 7.89(m, 2H), 8.03(d, 1H), 8.54(d, 1H)

50-3) 3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)carbonyl-1H-pyrrole

1.4g(4.7 mmol) of the compound prepared in Preparation 50-2) and 1.0g(5.1 mmol) of tosylmethylisocyanide were dissolved in 20 ml of tetrahydrofuran. 790mg(7.0 mmol) of potassium t-butoxide was added thereto and the mixture was stirred at room temperature for 3 hours. 2 ml of water was added to the reaction solution to stop the reaction and the solvent was removed under reduced pressure. The residue was extracted with ethyl acetate, washed with aqueous sodium chloride solution and dried over anhydrous magnesium sulfate. The solvent was removed under reduced pressure and the residue was subjected to column chromatography(eluent: ethyl acetate/hexane=2/3, v/v) to give 1.2g(3.6 mmol, Yield 76%) of the title compound.

^1H NMR(CDCl_3) δ 3.04(s, 3H), 3.35(s, 3H), 3.47(m, 2H), 3.64(m, 2H), 6.55(d, 1H), 6.63(m, 1H), 7.21-7.40(m, 4H), 7.74(m, 2H), 8.00(m, 1H), 11.4 (br, 1H)

Example 139: Synthesis of 1-[1-(4-cyanobenzyl)-1H-imidazol-5-yl]methyl-3-(naphthalen-1-yl)carbonyl-1H-pyrrole(139)

The title compound was obtained in a yield of 35% according to the same procedure as Example 1 except that the compound prepared in Preparation 29-5) and the compound prepared in Preparation 49-3) were used.

^1H NMR(CDCl_3) δ 4.86(s, 2H), 4.95(s, 2H), 6.52(s, 1H), 6.61(s, 1H), 6.89(m, 3H), 7.20(s, 1H), 7.49(m, 6H), 7.75(s, 1H), 7.87(d, 1H), 7.95(d, 1H), 8.11(d, 1H)

FAB : 417 (M+1)

Example 140: Synthesis of 1-[1-(4-bromobenzyl)-1H-imidazol-5-yl]methyl-3-(naphthalen-1-yl)carbonyl-1H-pyrrole(140)

The title compound was obtained in a yield of 20% according to the same procedure as Example 1 except that the compound prepared in Preparation 32-2) and the compound prepared in Preparation 49-3) were used.

^1H NMR(CDCl_3) δ 4.84(s, 2H), 4.92(s, 2H), 6.54(s, 1H), 6.67(s, 1H), 6.78(d, 2H), 6.93(s, 1H), 7.22(s, 1H), 7.38(d, 2H), 7.50(m, 3H), 7.58(d, 1H), 7.89(d, 1H), 7.95(d, 1H), 8.13(d, 1H), 8.16(s, 1H)

FAB : 470 (M+1)

Example 141: Synthesis of 1-[1-(4-bromobenzyl)-1H-imidazol-5-yl]methyl-3-[N-(2-methoxyethyl)-N-methyl]carbonyl-4-(naphthalen-1-yl)carbonyl-1H-pyrrole(141)

The title compound was obtained in a yield of 81% according to the same procedure as Example 1 except that the compound prepared in Preparation 32-2) and the compound prepared in Preparation 50-3) were used.

^1H NMR(CDCl_3) δ 2.94(s, 3H), 3.25(s, 3H), 3.42(m, 2H), 3.48(m, 2H), 4.72 (s, 2H), 4.78(s, 2H), 6.64(m, 4H), 7.28-7.48(m, 8H), 7.81(m, 2H), 8.14(m, 1H)

FAB : 585 (M+1)

Preparation 51: Synthesis of ethyl 1-naphthoyleglycinate hydrochloride

51-1) Ethyl N-(diphenylmethylene)glycinate

Glycine ethylester hydrochloride salt and diphenylketimine were reacted according to the procedure described in M. J. O' Donnell, R. L. Polt, *J. Org. Chem* 47, 2663, 1982 to give the title compound in a yield of 90%.

^1H NMR(CDCl_3) δ 1.20(t,3H), 4.12(m,4H), 7.10-7.40(m,8H), 7.59(d,2H)

51-2) Ethyl 1-naphthoylethylglycinate hydrochloride

1-Naphthoylethylchloride and the compound prepared in Preparation 51-1) were reacted according to the procedure described in J. Singh, et. al. *Tetrahedron Lett.*, 34(2), 211, 1993 to give the title compound in a yield of 48%.

^1H NMR($\text{DMSO}-d_6$) δ 1.78(s,3H), 3.65(q,1H), 3.95-4.15(m,2H), 6.33(s, 1H), 7.58-7.85(m,3H), 8.15(d,1H), 8.31(d,1H), 8.38(d,2H), 8.42(d,2H)

Preparation 52: Synthesis of 2-[1-(4-chlorobenzyl)-1H-imidazol-5-yl]thioacetamide**52-1) 1-(4-Chlorobenzyl)-5-hydroxymethyl-1H-imidazole**

The title compound was obtained in a yield of 50% according to the similar procedure described in J.M.Dener, L-H Zhang, H.Rapoport, *J. Org. Chem.*, 1993, 58, 1159 using dihydroxyacetone dimer and 4-chlorobenzylamine hydrochloride as starting materials.

^1H NMR($\text{CDCl}_3+\text{CD}_3\text{OD}$) δ 4.50(s,2H), 5.20(s,2H), 6.94(s,1H), 7.06(d,2H), 7.32(d,2H), 7.46(s,1H)

52-2) 1-(4-Chlorobenzyl)-5-chloromethyl-1H-imidazole hydrochloride

3.00g(13.5 mmol) of the compound prepared in Preparation 52-1) was dissolved in 40_{ml} of chloroform, 2.88_{ml}(40.5 mmol) of thionylchloride was slowly added thereto at 0°C, and the mixture was stirred at room temperature for 2 hours. The organic solvent was removed under reduced pressure to give 3.64g(13.1 mmol, Yield 97%) of the title compound. This compound was used directly in the next reaction without purification.

52-3) [1-(4-Chlorobenzyl)-1H-imidazol-5-yl]acetonitrile

1.2g(4.3 mmol) of the compound prepared in Preparation 52-2) was dissolved in 10_{ml} of dimethylsulfoxide and 1.3g(26 mmol) of sodiumcyanide was added thereto. The mixture was stirred at room temperature for 6 hours. 30_{ml} of water was added thereto and the resulting mixture was extracted with ethyl acetate(20_{ml} × 3). The organic layer was dried over anhydrous sodium sulfate and concentrated to give 0.96g(4.1 mmol, Yield 96%) of the title compound. This compound was used in the next reaction without purification.

¹H NMR(CDCl₃) δ 3.70(s,2H), 5.12(s,2H), 6.88(s,1H), 7.34(d,2H), 7.62(d, 2H), 7.71(s,1H)

52-4) 2-[1-(4-Chlorobenzyl)-1H-imidazol-5-yl]thioacetamide

150mg(0.64 mmol) of the compound prepared in Preparation 52-3) was dissolved in a solvent mixture of 1_{ml} of pyridine and 0.3_{ml} of triethylamine and then saturated by bubbling hydrogen sulfide gas through the solution for 30 minutes. The reaction solution was stirred at room temperature for 12 hours. The solvent was removed under reduced pressure and 10_{ml} of water was added thereto. The mixture was extracted with 10_{ml} of ethyl acetate. The organic layer was dried over anhydrous sodium sulfate, concentrated and subjected to silica gel column chromatography (eluent: methylene chloride/methanol=20/1, v/v) to

give 110mg(0.41 mmol, Yield 64%) of the title compound.

^1H NMR($\text{CDCl}_3+\text{CD}_3\text{OD}$) δ 3.21(s,2H), 5.05(s,2H), 6.76(s,1H), 7.24(d,2H), 7.61(d,2H), 7.67(s,1H)

FAB :266(M+1)

Preparation 53: Synthesis of 2-[1-[1-(benzyloxycarbonyl)piperidin-4-yl]methyl-1H-imidazol-5-yl]thioacetamide

53-1) 4-Aminomethyl-1-(benzyloxycarbonyl)piperidine

22.2g(0.2 mol) of 4-aminomethylpiperidine was dissolved in 250ml of toluene and 21.2g(0.2 mol) of benzaldehyde was added thereto. The reaction mixture was refluxed for 3 hours with Dean-stack to remove water, and then cooled down to 0°C. 34.2g(0.2 mol) of benzylchloroformate was added slowly thereto while stirring. The mixture was stirred at room temperature for 3 hours and 220ml of 1N aqueous KHSO_4 solution was added thereto. The mixture was extracted three times with 200ml of diethylether, and the aqueous layer was basified with 1N aqueous sodium hydroxide solution. The aqueous solution was saturated with sodium chloride. The aqueous layer was extracted three times with 100ml of dichloromethane, dried over anhydrous magnesium sulfate and distilled under reduced pressure to give 38g(Yield 91%, Molecular weight 248) of the title compound.

^1H NMR(CDCl_3) δ 1.11(s,2H), 1.49(s,3H), 1.70(d,2H), 2.57(d,2H), 2.78(s, 2H), 4.20(s,2H), 5.12(s,2H), 7.34-7.35(m, 5H)

53-2) 1-[1-(Benzyloxycarbonyl)piperidin-4-yl]methyl-5-hydroxymethyl-2-mercapto-1H-imidazole

24.8g(0.1 mol) of the compound prepared in Preparation 53-1) and 6.0g(0.1 mol) of acetic acid were dissolved in 50ml of n-butanol, a solution wherein 12.6g(0.13 mol) of potassium thiocyanate, 15.2g(0.1

mol) of 1,3-dihydroxyacetone dimer and 10.0g(0.17 mol) of acetic acid were dissolved in 50_{ml} of n-butanol was added thereto, and the whole mixture was stirred at room temperature. After 48 hours, the solvent was removed by distillation under reduced pressure, and then the residue was dissolved in 200_{ml} of ethyl acetate and washed three times with 100_{ml} of water. The organic layer was dried over anhydrous magnesium sulfate and concentrated to give 27g(75 mmol, Yield 75%) of the title compound.

¹H NMR(CDCl₃) δ 1.22(d,2H), 1.57(d,2H), 2.30(s,1H), 2.72(s,2H), 3.96(s, 2H), 4.15(d,2H), 4.46(s,2H), 5.10(s,2H), 6.62(s,1H), 7.26-7.37(m,5H)

53-3) 1-[1-(Benzyloxycarbonyl)piperidin-4-yl]methyl-5-hydroxymethyl-1H-imidazole

18.05g(50 mmol) of the compound prepared in Preparation 53-2) was added to a mixture of 100_{ml} of 10% nitric acid and 10_{ml} of ethyl acetate, the reaction mixture was cooled with cold ice water and then stirred at room temperature for 3 hours. The mixture was basified using 4N aqueous sodium hydroxide solution and extracted twice with 100_{ml} of ethyl acetate. The extracted organic solution was dried over magnesium sulfate and distilled under reduced pressure to give 12.3g(38 mmol, Yield 75%) of the title compound.

¹H NMR(CDCl₃) δ 1.16(d,2H), 1.56(d,2H), 1.98(s,1H), 2.70(s,2H), 3.88(d, 2H), 4.18(s,2H), 4.49(s,1H), 4.56(s,3H), 5.10(s,2H), 6.82(s,1H), 7.27-7.40(m, 6H)

53-4) 1-[1-(Benzyloxycarbonyl)piperidin-4-yl]methyl-5-chloromethyl-1H-imidazole hydrochloride

9.9g(30 mmol) of the compound prepared in Preparation 53-3) was dissolved in 50_{ml} of chloroform, and 7.1g(60 mmol) of

thionylchloride was slowly added thereto at 0°C. The reaction solution was stirred for 2 hours and the solvent was removed by distillation under reduced pressure to give 9.9g (Yield 95%, Molecular weight 347.5) of hydrochloride salt of the title compound. This compound was used directly in the next reaction without purification.

53-5) {1-[1-(Benzyloxycarbonyl)piperidin-4-yl]methyl-1H-imidazol-5-yl}acetonitrile

The title compound was obtained in a yield of 39% according to the similar procedure as Preparation 52-3) using the compound prepared in Preparation 53-4).

¹H NMR(CDCl₃) δ 1.19(br,2H), 1.60(br,2H), 1.90(m,1H), 2.72(br,2H), 3.71(s,2H), 3.81(d,2H), 4.22(br,2H), 5.11(s,2H), 7.03(s,1H), 7.29-7.36(m, 5H), 7.51(s,1H)

53-6) 2-{1-[1-(Benzyloxycarbonyl)piperidin-4-yl]methyl-1H-imidazol-5-yl}thioacetamide

The title compound was obtained in a yield of 74% according to the similar procedure as Preparation 52-4) using the compound prepared in Preparation 53-5).

¹H NMR(CDCl₃) δ 1.21(br,2H), 1.63(br,2H), 1.87(m,1H), 2.71(br,2H), 3.31 (s,2H), 3.84(d,2H), 4.25(br,2H), 5.12(s,2H), 7.10(s,1H), 7.33-7.41(m,5H), 7.62(s,1H)

FAB : 373 (M+1)

Preparation 54: Synthesis of methyl 3-chloro-3-(naphthalen-1-yl)-2-oxopropionate

7.80g(49.9 mmol) of 1-naphthaldehyde and 7.15g(49.9 mmol) of methyl dichloroacetate were dissolved in 100ml of t-butanol, and 6.15g(54.8 mmol) of potassium t-butoxide was added thereto at 0°C.

The mixture was stirred at room temperature for 24 hours and then 50_{ml} of water was added to stop the reaction. The solvent was removed under reduced pressure and the residue was extracted with ethyl acetate. The organic layer was dried over magnesium sulfate, concentrated and subjected to silica gel column chromatography(eluent: n-hexane/ethyl acetate=90/10, v/v) to give 2.5g(9.52 mmol, Yield 19%) of the title compound.

¹H NMR(CDCl₃) δ 3.78(s,3H), 6.92(s,1H), 7.45-7.73(m,4H), 7.95(m,2H), 8.12(d,1H)

Preparation 55: Synthesis of methyl 2-chloro-3-(naphthalen-1-yl)-3-oxopropionate

55-1) Methyl 3-(naphthalen-1-yl)-3-oxopropionate

10.2g(59.9 mmol) of 1-acetonaphthone and 4.8g(60% in mineral oil, 120 mmol) of sodium hydride were added to 100_{ml} of dimethylcarbonate and the mixture was refluxed for 24 hours. The solvent was removed under reduced pressure, 100_{ml} of 1N aqueous HCl solution was added to the residue, and the resulting mixture was extracted with 100_{ml} of ethyl acetate. The organic layer was washed with water(100_{ml} × 3), dried over anhydrous magnesium sulfate and concentrated. The residue was subjected to silica gel column chromatography(eluent: n-hexane/ethyl acetate=90/10, v/v) to give 10.0g(43.8 mmol, Yield 73%) of the title compound.

¹H NMR(CDCl₃) δ 3.75(s,3H), 4.14(s,2H), 7.45-7.68(m,3H), 7.82-8.08(m, 3H), 8.77(d,1H)

55-2) Methyl 2-chloro-3-(naphthalen-1-yl)-3-oxopropionate

4.56g(20.0 mmol) of the compound prepared in Preparation 55-1) was dissolved in 50_{ml} of 1,2-dichloroethane, and 2.70g(20.0 mmol) of

sulfonyl chloride was slowly added thereto at 0°C. The mixture was stirred at room temperature for 3 hours. The solvent was removed by distillation under reduced pressure to give 4.70g(17.9 mmol, Yield 89%) of the title compound.

¹H NMR(CDCl₃) δ 3.75(s,3H), 5.82(s,2H), 7.50-7.72(m,3H), 7.85-8.15(m, 3H), 8.65(d,1H)

Example 142: Synthesis of 4-ethoxycarbonyl-2-(1H-imidazol-5-ylmethyl)-5-(naphthalen-1-yl)oxazole(142)

142-1) Ethyl 2-[(1H-imidazol-5-yl)acetylamino]-3-(naphthalen-1-yl)-3-oxo-propionate

293mg(0.997 mmol) of the compound prepared in Preparation 51-2), 162mg(0.996 mmol) of 4-imidazoleacetic acid hydrochloride, 135mg(0.999 mmol) of HOBT and 191mg(0.996 mmol) of EDC were added to 10_{ml} of dimethylformamide, and then 202mg(1.99 mmol) of triethylamine was slowly added thereto while stirring. The mixture was stirred at room temperature for 5 hours and then the solvent therein was removed under reduced pressure. To the residue was added 30_{ml} of ethyl acetate, which was then washed with saturated sodium bicarbonate solution and water. The organic layer was dried over anhydrous magnesium sulfate, concentrated and subjected to silica gel column chromatography(eluent: dichloromethane/methanol=95/5, v/v) to give 200mg(0.547 mmol, Yield 55%) of the title compound.

¹H NMR(CDCl₃) δ 0.92(t,3H), 3.70(s,2H), 3.98-4.15(m,2H), 6.20(d,1H), 6.92(s,1H), 7.55(m,4H), 7.65(s,1H), 7.89(d,1H), 8.06(d,1H), 8.12(br,1H), 8.21(d,1H), 8.45(d,1H)

142-2) 4-Ethoxycarbonyl-2-(1H-imidazol-5-ylmethyl)-5-(naphthalen-1-yl)oxazole

100mg(0.27 mmol) of the compound prepared in Example 142-1) was dissolved in 5ml of THF and then refluxed for 6 hours. The solvent was removed by distillation under reduced pressure and the residue was subjected to silica gel column chromatography(eluent: dichloromethane/methanol=95/5, v/v) to give 40mg(0.12 mmol, Yield 44%) of the title compound.

^1H NMR(CDCl_3) δ 0.98(t,3H), 4.13(q,2H), 4.27(s,2H), 6.92(s,1H), 7.45- 7.58(m,4H), 7.65-7.75(m,2H), 7.89(d,1H), 7.97(d,1H)

FAB : 348 (M+1)

Example 143: Synthesis of 2-(1H-imidazol-5-ylmethyl)-4-(morpholin-4-yl)carbonyl-5-(naphthalen-1-yl)oxazole(143)

31mg(0.09 mmol) of the compound prepared in Example 142-2) was dissolved in a solvent mixture of tetrahydrofuran/methanol/water(0.6 ml/0.3ml/1ml), and 6mg(0.13 mmol) of lithium hydroxide was added thereto. The reaction solution was stirred at room temperature for 3 hours, and the solvent was removed under reduced pressure. The residue was adjusted to pH 6 using 0.1N aqueous hydrochloric acid solution and then extracted with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate and concentrated. The concentrate was dissolved in 1ml of dimethylformamide, 18mg(0.13 mmol) of HOBT and 26mg(0.13 mmol) of EDC were added thereto at 0°C, and the mixture was stirred for 10 minutes. 9 μl (0.09 mmol) of morpholine and 18 μl (0.13 mmol) of triethylamine were added thereto and the mixture was stirred at room temperature for 2 hours. The reaction solution was treated according to the same procedure as Example 142-1) to give 14mg(0.04 mmol, Yield 45%) of the title compound.

^1H NMR(CDCl_3) δ 2.97(br,2H), 3.24(br,2H), 3.43(br,2H), 3.57(br,2H), 4.27(s,2H), 6.95(s,1H), 7.52-7.67(m,6H), 7.81-7.95(m,3H)

FAB : 389 (M+1)

Example 144: Synthesis of 4-ethoxycarbonyl-2-(1H-imidazol-5-ylmethyl)-5-(naphthalen-1-yl)thiazole(144)

105mg(0.287 mmol) of the compound prepared in Example 142-1) and 116mg(0.287 mmol) of Lawesson's Reagent were dissolved in 10_{ml} of tetrahydrofuran, and the mixture was refluxed for 6 hours. The solvent was removed under reduced pressure, 10_{ml} of saturated sodium bicarbonate solution was added to the residue, and then the resulting mixture was extracted with ethyl acetate. The organic layer was dried over anhydrous magnesium sulfate, concentrated and subjected to silica gel column chromatography(eluent: dichloromethane/methanol=95/5, v/v) to give 26mg(0.075 mmol, Yield 26%) of the compound of Example 142-2) and 24mg(0.066 mmol, Yield 23%) of the title compound.

¹H NMR(CDCl₃) δ 0.63(t,3H), 3.92(q,2H), 4.42(s,2H), 6.97(s,1H), 7.405- 7.75(m,6H), 7.85-7.95(m,2H)

FAB : 364 (M+1)

Example 145: Synthesis of 2-[1-(4-chlorobenzyl)-1H-imidazol-5-yl methyl]-4-methoxycarbonyl-5-(naphthalen-1-yl)thiazole(145)

130mg(0.49 mmol) of the compound prepared in Preparation 52-4) and 129mg(0.49 mmol) of the compound prepared in Preparation 54 were dissolved in 5_{ml} of ethanol, and the mixture was refluxed for 5 hours. The solvent was removed by distillation under reduced pressure and the residue was subjected to silica gel column chromatography (eluent: dichloromethane/methanol=40/1, v/v) to give 45mg(0.095 mmol, Yield 19%) of the title compound.

¹H NMR(CDCl₃) δ 3.50(s,3H), 4.26(s,2H), 5.11(s,2H), 6.92(d,2H),

7.07(s, 1H), 7.21-7.43(m, 7H), 7.53(s, 1H), 7.83(m, 2H)

FAB : 474 (M+1)

Example 146: Synthesis of 2-[1-(4-chlorobenzyl)-1H-imidazol-5-ylmethyl]-4-(morpholin-4-yl)carbonyl-5-(naphthalen-1-yl)thiazole(146)

The title compound was obtained in a yield of 23% according to the similar procedure as Example 143 using the compound prepared in Example 145.

¹H NMR(CDCl₃) δ 2.63(br, 2H), 3.02(br, 2H), 3.24(br, 2H), 3.42(br, 2H), 4.26(s, 2H), 5.21(s, 2H), 7.02(m, 2H), 7.18(s, 1H), 7.31(m, 2H), 7.43-7.60(m, 5H), 7.78-7.96(m, 3H)

FAB : 529 (M+1)

Example 147: Synthesis of 2-[1-(4-chlorobenzyl)-1H-imidazol-5-ylmethyl]-4-[N-(2-methoxy)ethyl-N-methylcarbamoyl]-5-(naphthalen-1-yl)thiazole (147)

The title compound was obtained in a yield of 41% according to the similar procedure as Example 143 using the compound prepared in Example 145 except that N-(2-methoxyethyl)methylamine was used instead of morpholine

¹H NMR(CDCl₃) δ 2.68(br, 3H), 2.89-3.39(m, 7H), 4.22(s, 2H), 5.17(s, 2H), 7.01(m, 2H), 7.15(s, 1H), 7.33(m, 2H), 7.40-7.61(m, 5H), 7.71-7.82(m, 3H)

FAB : 531 (M+1)

Example 148: Synthesis of 2-[1-(4-chlorobenzyl)-1H-imidazol-5-ylmethyl]-5-methoxycarbonyl-4-(naphthalen-1-yl)thiazole(148)

250mg(0.95 mmol) of the compound prepared in Preparation 52-4) and 249mg(0.95 mmol) of the compound prepared in Preparation 55-2) were dissolved in 10_{ml} of ethanol, and the mixture was refluxed for 24 hours. The solvent was removed by distillation under reduced pressure and the residue was subjected to silica gel column chromatography (eluent: dichloromethane/methanol=40/1, v/v) to give 180mg(0.38 mmol, Yield 40%) of the title compound.

¹H NMR(CDCl₃) δ 3.53(s,3H), 4.22(s,2H), 5.12(s,2H), 6.91(m,2H), 7.11(s, 1H), 7.21-7.54(m,7H), 7.83(m,3H)

FAB : 474 (M+1)

Example 149: Synthesis of 2-[1-(4-chlorobenzyl)-1H-imidazol-5-ylmethyl]-5-(morpholin-4-yl)carbonyl-4-(naphthalen-1-yl)thiazole(149)

The title compound was obtained in a yield of 39% according to the similar procedure as Example 143 using the compound prepared in Example 148.

¹H NMR(CDCl₃) δ 2.38(br,2H), 2.82(br,2H), 3.21(br,2H), 3.42(br,2H), 4.27(s,2H), 5.21(s,2H), 6.98(m,2H), 7.25(m,3H), 7.50-7.61(m,5H), 7.89-7.99 (m,3H)

FAB : 529 (M+1)

Example 150: Synthesis of 2-{1-[1-(benzyloxycarbonyl)piperidin-4-ylmethyl]-1H-imidazol-5-ylmethyl}-5-methoxycarbonyl-4-(naphthalen-1-yl)thiazole(150)

124mg(0.33 mmol) of the compound prepared in Preparation 53-6) and 87mg(0.33 mmol) of the compound prepared in Preparation 55-2) were dissolved in 10_{ml} of ethanol, and the mixture was refluxed for 20 hours. The solvent was removed by distillation under reduced pressure

and the residue was subjected to silica gel column chromatography (eluent: dichloromethane/methanol=95/5, v/v) to give 95mg(0.16 mmol, Yield 48%) of the title compound.

^1H NMR(CDCl_3) δ 1.10(br,2H), 1.53(br,3H), 2.50(br,2H), 3.62(s,3H), 3.81(d,2H), 4.19(br,2H), 4.41(s,2H), 5.14(d,2H), 7.16(s,1H), 7.27-7.61(m, 10H), 7.78(s,1H), 7.91(d,1H), 7.96(d,1H)

FAB : 595 (M+1)

Example 151: Synthesis of 2-{1-[1-(benzyloxycarbonyl)piperidin-4-ylmethyl]-1H-imidazol-5-ylmethyl}-5-[N-(2-methoxy)ethyl-N-methylcarbamoyl]-4-(naphthalen-1-yl)thiazole(151)

The title compound was obtained in a yield of 36% according to the similar procedure as Example 143 using the compound prepared in Example 150 except that N-(2-methoxyethyl)methylamine was used instead of morpholine.

^1H NMR(CDCl_3) δ 2.68(br,3H), 2.89-3.39(m,7H), 4.22(s,2H), 5.17(s,2H), 7.01(m,2H), 7.15(s,1H), 7.33(m,2H), 7.40-7.61(m,5H), 7.71-7.82(m,3H)

FAB : 638 (M+1)

Preparation 56: Synthesis of 4-(5-chloromethyl-1H-imidazol-1-ylmethyl)-piperidine-1-carboxylic acid benzylester

56-1) 4-Aminomethyl-piperidine-1-carboxylic acid benzylester

22.2g(0.2 mol) of 4-aminomethyl-piperidine was dissolved in 250ml of toluene and then 21.2g(0.2 mol) of benzaldehyde was added thereto. The mixture was refluxed for 3 hours with Dean-stack and cooled down to 0°C, and then 34.2g(0.2 mol) of benzylchloroformate was added thereto while stirring. After the mixture was stirred for 3 hours,

1N aqueous potassium hydrosulfate solution(220_{ml}) was added thereto at room temperature. The mixture was extracted three times with 200_{ml} of diethylether, and then the aqueous layer was basified with sodium hydroxide. The aqueous solution was saturated with sodium chloride and extracted three times with 100_{ml} of dichloromethane. The organic solution was dried over magnesium sulfate and distilled under reduced pressure to obtain 38g(Yield 91%, Molecular weight 248) of the title compound.

¹H NMR(CDCl₃) δ 1.11(s,2H), 1.49(s,3H), 1.70(d,2H), 2.57(d,2H), 2.78(s, 2H), 4.20(s,2H), 5.12(s,2H), 7.34-7.35(m,5H)

FAB(M+H): 249

56-2) 4-(5-Hydroxymethyl-2-mercapto-1H-imidazol-1-ylmethyl)-piperidine-1-carboxylic acid benzylester

24.8g(0.1 mol) of the compound prepared in Preparation 56-1) and 6.0g(0.1 mol) of acetic acid were dissolved in 50_{ml} of n-buthanol, and then the resulting solution was added to a solution wherein 12.6g(0.13 mol) of potassium thiocyanate, 15.2g(0.1 mol) of 1,3-dihydroxyacetone dimer and 10.0g(0.17 mol) of acetic acid were dissolved in 50_{ml} of n-butanol. The whole mixture was stirred for 48 hours. The solvent was removed by distillation under reduced pressure, 200_{ml} of ethyl acetate was added thereto, and the mixture was washed three times with 100_{ml} of water. The organic layer was dried over magnesium sulfate, and the solvent was removed by distillation under reduced pressure to obtain 27g(75 mmol, Yield 75%, Molecular weight 361) of the title compound.

¹H NMR(CDCl₃) δ 1.22(d,2H), 1.57(d,2H), 2.30(s,1H), 2.72(s,2H), 3.96 (s,2H), 4.15(d,2H), 4.46(s,2H), 5.10(s,2H), 6.62(s,1H), 7.26-7.37(m,5H)

FAB(M+H): 362

56-3) 4-(5-Hydroxymethyl-1H-imidazol-1-ylmethyl)-piperidine-1-carboxylic acid benzylester

18.05g(50 mmol) of the compound prepared in Preparation 56-2) was added to a mixture of 100_{ml} of nitric acid(10%) and 10_{ml} of ethyl acetate. The whole mixture was soaked in cold ice water for 5 minutes, and stirred at room temperature for 3 hours. The mixture was basified with 4N aqueous sodium hydroxide solution, and then extracted twice with 100_{ml} of ethyl acetate. The organic extract was dried over magnesium sulfate and distilled under reduced pressure to obtain 12.3g (38 mmol, Yield 75%, Molecular weight 329) of the title compound.

¹H NMR(CDCl₃) δ 1.16(d,2H), 1.56(d,2H), 1.98(s,1H), 2.70(s,2H), 3.88 (d,2H), 4.18(s,2H), 4.49(s,1H), 4.56(s,3H), 5.10(s,2H), 6.82(s,1H), 7.27-7.40 (m,5H)

FAB(M+H): 330

56-4) 4-(5-Chloromethyl-1H-imidazol-1-ylmethyl)-piperidine-1-carboxylic acid benzylester

9.9g(30 mmol) of the compound prepared in Preparation 56-3) was dissolved in 50_{ml} of chloroform, and 7.1g(60 mmol) of thionyl chloride was slowly added thereto at 0°C. The mixture was stirred for 2 hours, the solvent was removed by distillation under reduced pressure, and the residual hydrochloric acid was removed under vacuum to obtain 9.9g(Yield 95%, Molecular weight 347.5) of hydrochloric acid salt of the title compound.

¹H NMR(CDCl₃) δ 1.12(d,2H), 1.53(d,2H), 2.65(s,2H), 3.82(d,2H), 4.22 (s,2H), 4.42(s,1H), 4.49(s,3H), 5.12(s,2H), 6.60(s,1H), 7.30-7.41(m,5H)

FAB(M+H): 349

Preparation 57: Synthesis of 1-(4-chlorobenzyl)-5-chloromethyl-1H-imidazole hydrochloride

57-1) 1-(4-Chlorobenzyl)-5-hydroxymethyl-1H-imidazole

The title compound was obtained in a yield of 50% according to the procedure described in J.M.Dener, L-H Zhang, H.Rapoport, *J.Org.Chem.*, 1993, 58, 1159 using dihydroxyacetone dimer and 4-chlorobenzylamine hydrochloride as starting materials.

¹H NMR(CDCl₃+CD₃OD) δ 4.46(s,2H), 5.26(s,2H), 7.00(s,1H), 7.07(d,2H), 7.50(d,2H), 7.65(s,1H)

57-2) 1-(4-Chlorobenzyl)-5-chloromethyl-1H-imidazole hydrochloride

The title compound was obtained in a yield of 96% according to the similar procedure as Preparation 56-4) except that the compound prepared in Preparation 57-1) was used as a starting material. This compound was directly used in the next reaction without purification.

Preparation 58: Synthesis of 4-[N-(2-methoxyethyl)-N-methyl] carbamoyl-3-(naphthalen-1-yl)-1H-pyrazole

58-1) N-t-Butyl-N'-(naphthalen-1-ylmethylene)-hydrazine

5.0g(32 mmol) of 1-naphthaldehyde and 3.99g(32 mmol) of t-butylhydrazine hydrochloride were dissolved in 100_{mL} of methanol, and then the mixture was reacted with 1_{mL} of acetic acid at room temperature for 24 hours. After the solvent was removed by distillation under reduced pressure, 20_{mL} of ethyl acetate was added to the residue. The mixture was washed with saturated sodium hydrogen carbonate solution. Then, the separated organic layer was dried over anhydrous magnesium sulfate and distilled under reduced pressure to remove the solvent to obtain 6.3g(28 mmol, Yield 86%) of the title compound.

^1H NMR(CDCl_3) δ 1.70(s,9H), 7.23(s,1H), 7.32(m,1H),
7.42(m,2H), 7.80 (d,1H), 7.90(d,2H), 8.60(d,1H), 9.91(s,1H), 12.1(br,1H)

FAB(M+H): 227

58-2) 1-(t-Butyl)-3-(naphthalen-1-yl)-1H-pyrazole-4-carboxylic acid ethyl ester

6.3g(28 mmol) of the compound prepared in Preparation 58-1) and 2.44g(30.8 mmol) of ethylpropiolate were dissolved in a solvent mixture of 27 mL of acetic acid and 32 mL of acetonitrile, and the whole mixture was reacted in the air for 3 days. The solvent was removed, and the residue was subjected to silica gel column chromatography (eluent: ethyl acetate/n-hexane=9/1, v/v) to obtain 6.76g(21 mmol, Yield 75%) of the title compound.

^1H NMR(CDCl_3) δ 0.80(t,3H), 1.65(s,9H), 3.98(q,2H),
7.38(m,2H), 7.48 (m,1H), 7.55(m,1H), 7.74(m,1H), 7.85(m,2H),
8.21(s,1H), 11.31(br,1H)

FAB(M+H): 323

58-3) 3-(Naphthalen-1-yl)-1H-pyrazole-4-carboxylic acid ethylester

3.65g(11.3 mmol) of the compound prepared in Preparation 58-2) was dissolved in 50 mL of formic acid, and the resulting solution was boiled for 12 hours under reflux. The solvent therein was removed by distillation under reduced pressure, and ethyl acetate was added thereto. The mixture was washed with saturated aqueous sodium hydrogen carbonate solution and dried over anhydrous magnesium sulfate. The solvent was removed by distillation under reduced pressure, and the residue was subjected to silica gel column chromatography(eluent: ethyl acetate/n-hexane=6/4, v/v) to obtain 1.1g(4.1 mmol, Yield 37%) of the title compound(see, *J.Hetero.Chem.*, 31, 1447, 1944).

^1H NMR(CDCl_3) δ 0.80(t,3H), 3.98(q,2H), 7.35-7.60(m,5H),

7.90(m,2H), 7.94(s,1H)

FAB(M+H): 267

58-4) 3-(Naphthalen-1-yl)-1H-pyrazole-4-carboxylic acid

1.1g(4.1 mmol) of the compound prepared in Preparation 58-3) and 2.1g(12.4 mmol) of potassium hydroxide were dissolved in 50_{ml} of a solvent mixture of methanol/water(1:1, v/v). The mixture was reacted under reflux for 12 hours. The solvent was removed by distillation under reduced pressure. The residue was acidified with 1N aqueous hydrochloric acid solution, extracted with 50_{ml} of ethyl acetate and dried over anhydrous magnesium sulfate. The solvent was removed by distillation under reduced pressure to obtain 910mg(3.8 mmol, Yield 92%) of the title compound.

¹H NMR(CD₃OD+CDCl₃) δ 7.30(m,3H), 7.56(d,1H), 7.80-7.95 (m,3H), 8.07 (s,1H)

FAB(M+H): 239

58-5) 4-[N-(2-Methoxyethyl)-N-methyl]carbamoyl-3-(naphthalen-1-yl)-1H-pyrazole

238mg(1 mmol) of the compound prepared in Preparation 58-4) was dissolved in 10_{ml} of dimethylformamide, and 230mg(1.2 mmol) of EDC, 101mg(1 mmol) of triethylamine and 162mg(1.2 mmol) of HOBT(1-hydroxybenzotriazole) were added thereto, and then the mixture was stirred at 0^oC for 5 minutes. To the mixture was added 124mg (1 mmol) of N-(2-methoxyethyl)-N-methylamine hydrochloride, which was then stirred at room temperature for 5 hours. The solvent was removed under reduced pressure, 10_{ml} of saturated aqueous potassium carbonate solution was added to the residue. The mixture was extracted with 20 _{ml} of ethyl acetate, washed with 10_{ml} of 1N aqueous hydrochloric acid solution, washed with saturated sodium chloride solution and water, dried

over anhydrous sodium sulfate, and concentrated to obtain 247mg(0.8 mmol, Yield 80%) of the title compound.

^1H NMR(CDCl_3) δ 2.40(s,2H), 2.81(s,1H), 2.84(s,1H), 2.96(s,1H), 3.02 (s,4H), 3.15(s,1.5H), 3.34(s,1.5H), 7.24-7.52(m,4H), 7.59(s,1H), 7.77(m,2H), 7.93(d,1H)

FAB(M+H): 310

Preparation 59: Synthesis of 4-(morpholin-4-yl)carbonyl-3-(naphthalen-1-yl)-1H-pyrazole

238_{mg}(1 mmol) of the compound prepared in Preparation 58-4) was dissolved in 10_{ml} of dimethylformamide, and 230_{mg}(1.2 mmol) of EDC and 162_{mg}(1.2 mmol) of HOBt were added thereto, and the mixture was stirred at 0°C for 5 minutes. To the whole mixture was added 87_{mg}(1 mmol) of morpholine, which was then stirred at room temperature for 5 hours. The solvent was removed under reduced pressure and 10_{ml} of saturated aqueous potassium carbonate solution was added to the residue. The mixture was extracted with 20_{ml} of ethyl acetate, washed with 10_{ml} of 1N hydrochloric acid solution, washed with saturated aqueous sodium chloride solution and water, dried over anhydrous sodium sulfate, and concentrated to obtain 240_{mg}(0.8 mmol, Yield 80%) of the title compound.

^1H NMR(CDCl_3) δ 2.5(br,2H), 2.95(br,2H), 3.15(br,2H), 3.40(br,2H), 7.50 (m,4H), 7.95(m,4H), 9.73(br,1H)

FAB(M+H): 308

Example 152: Synthesis of 1-[1-(1-benzyloxycarbonyl-piperidin-4-ylmethyl)-1H-imidazol-5-ylmethyl]-4-[N-(2-methoxyethyl)-N-methyl] carbamoyl-3-(naphthalen-1-yl)-1H-pyrazole(152)

616mg(2.0 mmol) of the compound prepared in Preparation 56-4) was dissolved in 10_{ml} of dimethylformamide, 264mg(6.6 mmol) of sodium hydride(60%) was added thereto at 0°C, and the whole mixture was stirred for 5 minutes. To the mixture was added 765mg(2.2 mmol) of the compound prepared in Preparation 58-5) and the resulting mixture was stirred at room temperature for 5 hours. The solvent was removed by distillation under reduced pressure, and 10_{ml} of water was added to the residue. The mixture was then extracted twice with 20_{ml} of ethyl acetate, dried over magnesium sulfate, concentrated, and subjected to silica gel column chromatography(eluent: dichloromethane/methanol=90/10, v/v) to obtain 930mg(Yield 75%) of the title compound.

¹H NMR(CDCl₃) δ 1.11(m,2H), 1.37(br,1H), 1.50(br,2H), 2.35(br,1H), 2.55 (br,2H), 2.71(br,1H), 2.90-3.21(m,7H), 3.35(br,1H), 3.90(br,2H), 3.98(d,1H), 4.50(d,1H), 5.02(s,2H), 5.10(s,2H), 7.21-7.40(m,6H), 7.41-7.60(m,4H), 7.70 (s,1H), 7.80(s,1H), 7.95(m,2H), 8.13(d,1H)

FAB(M+H): 621

Example 153: Synthesis of 1-[1-(1-methoxycarbonylpiperidin-4-ylmethyl)-1H-imidazole-5-ylmethyl]-4-[N-(2-methoxyethyl)-N-methyl]carbamoyl-3-(naphthalen-1-yl)-1H-pyrazole(153)

153-1) 1-[1-(Piperidin-4-ylmethyl)-1H-imidazole-5-ylmethyl]-4-[N-(2-methoxyethyl)-N-methyl]carbamoyl-3-(naphthalen-1-yl)-1H-pyrazole

227mg(0.36 mmol) of the compound prepared in Example 152 was dissolved in methanol, 20mg of palladium hydroxide carbon was added thereto, and then the mixture was reacted under 1 atm of hydrogen for 2 hours. After the reaction was completed, the mixture was filtered and the solvent was removed. The filtrate was subjected to silica gel column chromatography(eluent: ammonia water/methanol=

15/85, v/v) to obtain 128mg(0.26 mmol, Yield 74%) of the title compound.

¹H NMR(CDCl₃) δ 1.08(s,2H), 1.53(m,4H), 2.33(s,2H), 2.64(br,4H), 3.20(m,6H), 3.31(s,1H), 3.75(d,2H), 4.13(m,2H), 5.10(s,2H), 6.71(s,1H), 7.11(s,1H), 7.30(m,9H), 7.74(d,1H), 7.81(d,1H), 7.90(s,1H), 8.06(d,1H)

FAB(M+H): 486

153-2) 1-[1-(1-methoxycarbonylpiperidin-4-ylmethyl)-1H-imidazol-5-ylmethyl]-4-[N-(2-methoxyethyl)-N-methyl]carbamoyl-3-(naphthalen-1-yl)-1H-pyrazole

30mg(62 μmol) of the compound prepared in Example 153-1) was added to 2ml of dichloromethane, 5.4mg(6.9 μmol) of methylchloroformate was added thereto by an injector, and the mixture was stirred for 2 hours. The solvent was removed under reduced pressure, and the residue was subjected to silica gel column chromatography(eluent: dichloromethane/methanol=80/20, v/v) to obtain 27.8mg(5.3 μmol, Yield 85%) of the title compound.

¹H NMR(CDCl₃) δ 1.11(br,2H), 1.33(br,1H), 1.53(br,2H), 2.39(s,2H), 2.70 (br,4H), 2.90-3.20(br,6H), 3.32(s,1H), 3.62(s,3H), 3.78(d,2H), 4.16(m,2H), 5.16(s,2H), 6.74(s,1H), 7.10(s,1H), 7.21-7.50(m,14H), 7.76(d,1H), 7.84(d,1H), 7.91(s,1H), 8.07(d,1H)

FAB(M+H): 545

Example 154: Synthesis of 1-[1-(4-bromobenzyl)-1H-imidazol-5-ylmethyl]-4-[N-(2-methoxyethyl)-N-methyl]carbamoyl-3-(naphthalen-1-yl)-1H-pyrazole(154)

The title compound was obtained in a yield 81% according to the same procedure as Example 152 except that the compound prepared in

Preparation 32-2) and the compound prepared in Preparation 58-5) were used.

^1H NMR(CDCl_3) δ 2.41(s,2H), 2.82(s,1H), 2.85(s,1H), 2.98(s,1H), 3.04(s,4H), 3.17(s,1.5H), 3.36(s,1.5H), 5.11(s,2H), 5.21(s,2H), 6.95(d,2H), 7.25(d,2H), 7.35-7.60(m,5H), 7.64(s,1H), 7.72(s,1H), 7.81(m,2H), 8.11(d,1H)

FAB(M+H): 558

Example 155: Synthesis of 1-[1-(4-chlorobenzyl)-1H-imidazol-5-yl methyl]-4-[N-(2-methoxyethyl)-N-methyl]carbamoyl-3-(naphthalen-1-yl)-1H-pyrazole(155)

The title compound was obtained in a yield 81% according to the same procedure as Example 152 except that the compound prepared in Preparation 57-2) and the compound prepared in Preparation 58-5) were used.

^1H NMR(CDCl_3) δ 2.41(s,2H), 2.82(s,1H), 2.85(s,1H), 2.98(s,1H), 3.04 (s,4H), 3.17(s,1.5H), 3.36(s,1.5H), 5.20(s,2H), 5.25(s,2H), 6.97(d,2H), 7.26(d,2H), 7.35-7.46(m,5H), 7.47(s,1H), 7.58(s,1H), 7.88(m,2H), 8.11(d,1H)

FAB(M+H): 514

Example 156: Synthesis of 1-[1-(4-cyanobenzyl)-1H-imidazol-5-yl methyl]-4-[N-(2-methoxyethyl)-N-methyl]carbamoyl-3-(naphthalen-1-yl)-1H-pyrazole(156)

The title compound was obtained in a yield 81% according to the same procedure as Example 152 except that the compound prepared in Preparation 29-5) and the compound prepared in Preparation 58-5) were used.

^1H NMR(CDCl_3) δ 2.41(s,2H), 2.82(s,1H), 2.85(s,1H), 2.98(s,1H), 3.04 (s,4H), 3.17(s,1.5H), 3.36(s,1.5H), 5.20(s,2H), 5.31(s,2H), 6.99(d,2H), 7.26 (d,2H), 7.35-7.46(m,5H), 7.48(s,1H), 7.57(s,1H), 7.89(m,2H), 8.12(d,1H)

FAB(M+H): 505

Example 157: Synthesis of 1-[1-methyl-1H-imidazol-5-ylmethyl]-4-[N-(2-methoxyethyl)-N-methyl]carbamoyl-3-(naphthalen-1-yl)-1H-pyrazole (157)

The title compound was obtained in a yield 81% according to the same procedure as Example 152 except that 1-methyl-5-chloromethyl-1H-imidazole hydrochloride and the compound prepared in Preparation 58-5) were used.

^1H NMR(CDCl_3) δ 2.42(br,2H), 2.71(br,1H), 3.10(br,5H), 3.30(br,1H), 3.50(s,3H), 5.17(s,2H), 6.69(s,1H), 7.09(s,1H), 7.41(m,9H), 7.74(d,1H), 7.83 (d,1H), 7.89(s,1H), 8.05(d,1H)

FAB(M+H): 404

Example 158: Synthesis of 1-[1-(1-benzyloxycarbonyl-piperidin-4-ylmethyl)-1H-imidazol-5-ylmethyl]-4-(morpholin-4-yl)carbonyl-3-(naphthalen-1-yl)-1H-pyrazole(158)

612mg(2.0 mmol) of the compound prepared in Preparation 59 was dissolved in 10 ml of dimethylformamide, 264mg(6.6 mmol) of sodium hydride(60%) was added thereto at 0°C, and the whole mixture was stirred for 5 minutes. To the mixture was added 765mg(2.2 mmol) of the compound prepared in Preparation 56-4), which was then stirred at room temperature for 5 hours. The solvent was removed by distillation under reduced pressure and 10 ml of water was added to the

residue. The resulting mixture was then extracted twice with 20 ml of ethyl acetate, dried over magnesium sulfate, concentrated, and subjected to silica gel column chromatography (eluent: dichloromethane/methanol=90/10, v/v) to obtain 930mg (Yield 75%) of the title compound.

¹H NMR(CDCl₃) δ 1.11(m,2H), 1.37(br,1H), 1.50(br,2H), 1.62(br,1H), 2.35(br,1H), 2.55(br,2H), 2.71(br,1H), 3.14(br,2H), 3.35(br,2H), 3.90(br,2H), 4.15(m,4H), 5.02(s,2H), 5.10(s,2H), 7.21-7.40(m,6H), 7.41-7.60(m,4H), 7.70 (s,1H), 7.80(s,1H), 7.95(m,2H), 8.13(d,1H)

FAB(M+H): 619

Example 159: Synthesis of 1-[1-(1-methoxycarbonylpiperidin-4-ylmethyl)-1H-imidazol-5-ylmethyl]-4-(morpholin-4-yl)carbonyl-3-(naphthalen-1-yl)-1H-pyrazole(159)

159-1) 1-[1-(Piperidin-4-ylmethyl)-1H-imidazol-5-ylmethyl]-4-(morpholin-4-yl)carbonyl-3-(naphthalen-1-yl)-1H-pyrazole

227mg(0.36 mmol) of the compound prepared in Example 158 was dissolved in methanol, 20mg of palladium hydroxide carbon was added thereto, and the mixture was reacted under 1 atm of hydrogen for 2 hours. After the reaction was completed, the mixture was filtered and the solvent therein was removed. The residue was subjected to silica gel column chromatography (eluent: ammonia water/methanol=15/85, v/v) to give 120mg(0.26 mmol, Yield 74%) of the title compound.

¹H NMR(CDCl₃) δ 1.06(m,2H), 1.43(m,3H), 2.36(br,5H), 2.41-3.79(br,13H), 3.78(d,2H), 5.22(s,2H), 6.88(s,1H), 7.12(d,2H), 7.26(m,1H), 7.35(m,3H), 7.63(s,1H), 7.75(d,1H), 7.80(d,1H), 7.93(d,1H)

FAB(M+H): 484

159-2) 1-[1-(1-Methoxycarbonylpiperidin-4-ylmethyl)-1H-imidazol-5-yl

methyl]-4-(morpholin-4-yl)carbonyl-3-(naphthalen-1-yl)-1H-pyrazole

30mg(62 μ mol) of the compound prepared in Example 159-1) was dissolved in 2ml of dichloromethane, 5.4mg(6.9 μ mol) of methylchloroformate was added thereto by injector, and the whole mixture was stirred for 2 hours. The solvent was removed under reduced pressure, and the residue was subjected to silica gel column chromatography(eluent: dichloromethane/methanol=80/20, v/v) to give 27.8mg(5.3 μ mol, Yield 85%) of the title compound.

^1H NMR(CDCl_3) δ 1.05(br,2H), 1.32(br,1H), 1.53(br,2H), 2.31-2.72(m,5H), 3.03~3.33(m,7H), 3.62(s,3H), 3.66(m,2H), 4.13(br,2H), 5.12(s,2H), 6.71 (s,1H), 7.03(s,1H), 7.14(s,1H), 7.24~7.43(m,5H), 7.74(d,1H), 7.82(d,1H), 8.10(d,1H)

FAB(M+H): 543

Example 160: Synthesis of 1-[1-(4-bromobenzyl)-1H-imidazol-5-yl methyl]-4-(morpholin-4-yl)carbonyl-3-(naphthalen-1-yl)-1H-pyrazole(160)

The title compound was obtained in a yield 81% according to the same procedure as Example 152 except that the compound prepared in Preparation 32-2) and the compound prepared in Preparation 59 were used.

^1H NMR(CDCl_3) δ 2.35(br,2H), 2.80(br,2H), 3.15(br,2H), 3.35(br,2H), 5.29(s,2H), 5.31(s,2H), 7.00(d,2H), 7.20-7.35(m,3H), 7.40-7.60(m,4H), 7.72 (s,1H), 7.80(s,1H), 7.90(m,2H), 8.01(d,1H)

FAB(M+H): 556

Example 161: Synthesis of 1-[1-(4-chlorobenzyl)-1H-imidazol-5-yl methyl]-4-(morpholin-4-yl)carbonyl-3-(naphthalen-1-yl)-1H-pyrazole(161)

The title compound was obtained in a yield 81% according to the

same procedure as Example 152 except that the compound prepared in Preparation 57-2) and the compound prepared in Preparation 59 were used.

^1H NMR(CDCl_3) δ 2.35(br,2H), 2.80(br,2H), 3.15(br,2H), 3.35(br,2H), 5.29(s,2H), 5.31(s,2H), 7.00(d,2H), 7.20-7.35(m,3H), 7.40-7.60(m,4H), 7.72 (s,1H), 7.80(s,1H), 7.90(m,2H), 8.01(d,1H)

FAB(M+H): 512

Example 162: Synthesis of 1-[1-(4-cyanobenzyl)-1H-imidazol-5-ylmethyl]-4-(morpholin-4-yl)carbonyl-3-(naphthalen-1-yl)-1H-pyrazole(162)

The title compound was obtained in a yield 81% according to the same procedure as Example 152 except that the compound prepared in Preparation 29-5) and the compound prepared in Preparation 59 were used.

^1H NMR(CDCl_3) δ 2.35(br,2H), 2.80(br,2H), 3.15(br,2H), 3.35(br,2H), 5.28(s,2H), 5.34(s,2H), 7.03(d,2H), 7.20-7.35(m,3H), 7.40-7.60(m,4H), 7.72 (s,1H), 7.80(s,1H), 7.90(m,2H), 8.01(d,1H)

FAB(M+H): 503

Example 163: Synthesis of 1-(1-methyl-1H-imidazol-5-ylmethyl)-4-(morpholin-4-yl)carbonyl-3-(naphthalen-1-yl)-1H-pyrazole(163)

The title compound was obtained in a yield 81% according to the same procedure as Example 152 except that 1-methyl-5-chloromethyl-1H-imidazole hydrochloride and the compound prepared in Preparation 59 were used.

^1H NMR(CDCl_3) δ 2.35(br,2H), 2.80(br,2H), 3.15(br,2H), 3.35(br,2H), 3.62(s,3H), 5.29(s,2H), 7.20-7.35(m,3H), 7.40-7.60(m,2H), 7.72(s,1H), 7.80 (s,1H), 7.90(m,2H), 8.01(d,1H)

FAB(M+H): 402

Experimental Example 1**Analysis of in vitro inhibitory activity for Ras farnesyl transferase**

In the present experiment, Ras farnesyl transferase produced by genetic recombination techniques according to the improved Pompliano's method (Pompliano et al., *Biochemistry*, 1992, 31, 3800) was used, and Ras substrate(Ras-CVLS) protein described in Korean Patent Appln. No. 97-14409 was used after it has been purified according to the known method(see, Chung et al., *Biochimica et Biophysica Acta*, 1992, 278, 1129).

The enzyme reaction was performed in 50 μ l of 50mM Sodium HEPES buffer solution containing 25mM of potassium chloride, 25mM of magnesium chloride, 10mM of DTT and 50 μ M of zinc chloride. 1.5 μ M of Ras substrate protein, 0.15 μ M of tritium-farnesylpyrophosphate and 4.5nM of farnesyl transferase were used.

More specifically, in the initial step, farnesyl transferase was added to the above buffer solution, reaction was maintained for 30 minutes at 37°C and then the reaction was stopped by adding 1ml of ethanol solution containing 1M HCl. The formed precipitates were adsorbed to GF/B filter using Hopper harvester(Hopper #FH 225V) for filter-binding, washed with ethanol, and then radioactivity of the dried filter was measured using LKB β counter. Enzyme titer was measured in the unsaturated state of substrate where the concentrations of Ras substrate protein and farnesyl transferase have quantitative relationship. The compound according to the present invention dissolved in dimethyl sulfoxide(DMSO) was added to the reaction solution in an amount of

less than 5% of the total reaction solution, and then the enzyme inhibitory activity thereof was measured. The enzyme inhibitory activity was represented by percentage of the amount of farnesyl incorporated into the Ras substrate protein in the presence of the test compound to that in the absence of the test compound. IC_{50} of the test compound was defined as the concentration at which 50% of the enzyme activity was inhibited.

To evaluate the selective enzyme inhibitory activity of the compound according to the present invention, inhibitory activity on geranylgeranyl transferase was measured. Geranylgeranyl transferase was purified from bovine brain according to the method modified from Schaber's method(Schaber et al., J. Biol. Chem. 1990, 265, 14701), and substantially the same experimental procedure as that for farnesyl transferase was performed on geranylgeranyl pyrophosphate and Ras-CVIL substrate protein.

The test results are represented in the following Table 7.

Experimental Example 2

Analysis of in vivo inhibitory activity for Ras farnesyl transferase

In the present experiment, Rat2 cell line which expresses C-Harvey-Ras protein having transforming activity and Rat2 cell line(Korean patent application No. 97-14409) which is transformed with fused protein of H-Ras substituted with polybasic lysine domain at C-terminus of K-Ras were used. The experiment was performed by the modified Declue's method(Declue. J. E. et al., Cancer Research, 1991, 51, 712). Hereinafter, the experimental method will be described in more detail.

3×10^5 cells of transformed Rat2 fibroblast cell line were sprayed on 60mm cell cultivation dish and cultivated for 48 hours in a cell incubator at 37°C and after 50% or more of density was reached, it was treated with the test compounds. The compound according to the present invention dissolved in dimethylsulfoxide(DMSO) was used. 1% concentration of dimethylsulfoxide was used in both control and test groups. After 4 hours from the treatment with the compound, methionine labeled with 150 μ Ci of radioactive isotope [35 S] per 1ml of medium was added and after cultivating for 20 hours, the cells were washed with physiological saline water. The cells were lysed using 1ml of cold cell lysis buffer solution(50mM of Sodium HEPES buffer solution containing 5mM of magnesium chloride, 1mM of DTT, 1% NP 40, 1mM of EDTA, 1mM of PMSF, 2 μ M of leupeptin, 2 μ M of pepstatin A and 2 μ M of antipain) and the supernatant wherein the cells were lysed was obtained by high-velocity centrifugation of 12,000g $\times$ 5 minutes. The amount of radioisotope in the supernatant was measured and standardized to obtain a quantitative result in immunoprecipitation reaction and then, Y13-259, a monoclonal antibody specifically binding to Ras protein(Furth, M. E. et al., J. Virol, 1982, 43, 294) was added and reacted for 15 hours at 4°C. Protein A(combined with goat anti-murine immunoglobulin antibody)-agarose suspension was added to the solution and reacted for 1 hour at 4°C and then, to remove the unspecific binding product, immunoprecipitates were washed with a buffer solution (50mM Tris chloride buffer solution containing 50mM of sodium chloride, 0.5% of sodium dioxycolate, 0.5% of NP 40 and 0.1% of SDS). The precipitates were added to a buffer solution for electrophoresis and boiled and then, electrophoresis was performed using 13.5% of SDS polyacrylamide gel. After electrophoresis, the gel was fixed and dried. Then, the gel was exposed to X-ray film, developed

and printed. From the result of the experiment, intensities of band of protein combined with or without farnesyl of Ras protein were measured, and the concentration of the test compound inhibiting 50% of farnesyl binding was defined as CIC_{50} , an in vivo Ras farnesyl transferase inhibitory activity. The test results are shown in the following Table 7.

Table 7-1

COM. NO.	H-Ras $IC_{50}(\mu M)$	H-Ras $CIC_{50}(\mu M)$	K-Ras $IC_{50}(\mu M)$	K-Ras $CIC_{50}(\mu M)$
1	0.0011	0.025	0.0035	10
2	0.00085	0.025	0.002	10-50
3	0.001	0.025	0.0024	15
4	0.047	0.1-1	0.75	10-100
5	0.0037	0.025	0.0085	10-50
6	0.001	0.025	0.002	10-50
7	0.0006	0.025	0.0022	10-50
8	0.004	0.025	0.008	10-50
9	0.005	0.025	0.0066	10-50
10	0.00085	0.0125	0.005	10-50
11	0.004	0.025	0.008	10-50
12	0.005	0.025	0.0066	10-50
13	0.00085	0.0125	0.005	10-50
14	0.002	0.0125	0.005	10-50
15	0.005	0.025	0.01	10-50
16	0.0012	0.0125	0.005	10-50
17	0.002	0.025	0.003	10-50
18	0.001	0.025	0.002	10-50

Table 7-2

COM. NO.	H-Ras IC ₅₀ (μ M)	H-Ras CIC ₅₀ (μ M)	K-Ras IC ₅₀ (μ M)	K-Ras CIC ₅₀ (μ M)
19	0.001	0.020	0.003	10-50
20	0.001	0.020	0.002	10-50
21	0.001	0.021	0.001	10-50
22	0.001	0.020	0.002	10-50
23	0.002	0.023	0.002	10-50
24	0.002	0.025	0.003	10-50
25	0.002	0.015	0.005	10-50
26	0.002	0.015	0.003	10-50
27	0.006	0.025	0.005	10-30
28	0.001	0.020	0.002	10-30
29	0.002	0.010	0.004	10-20
30	0.002	0.010	0.004	10-20
31	0.002	0.012	0.005	10-20
32	0.002	0.015	0.003	10-50
33	0.002	0.018	0.003	10-50
34	0.002	0.020	0.003	10-50
35	0.001	0.025	0.002	10-50
36	0.001	0.025	0.002	10-50
37	0.002	0.025	0.003	10-50
38	0.002	0.025	0.004	10-50
39	0.002	0.020	0.003	10-50
40	0.002	0.025	0.003	10-50
41	0.003	0.015	0.004	10-50

Table 7-3

COM. NO.	H-Ras IC ₅₀ (μ M)	H-Ras CIC ₅₀ (μ M)	K-Ras IC ₅₀ (μ M)	K-Ras CIC ₅₀ (μ M)
42	0.004	0.015	0.003	10-50
43	0.001	0.015	0.002	5-10
44	0.25	1-50	0.1-10	10-100
45	0.13	1-50	0.1-10	10-100
46	0.12	1-50	0.1-10	10-100
47	0.09	1-50	0.1-10	10-100
48	0.15	1-50	10	10-100
49	0.03	0.7	0.485	<20
50	0.15	1-50	0.1-10	10-100
51	0.27	1-50	0.1-10	10-100
52	0.07	1-50	0.1-10	10-100
53	0.3	1-50	0.1-10	10-100
54	0.39	1-50	0.1-10	10-100
55	0.06	1-50	0.1-10	10-100
56	0.04	1-50	0.1-10	10-100
57	0.038	1-50	0.1-10	10-100
58	0.025	1-50	0.1-10	10-100
59	0.57	1-50	0.1-10	10-100
60	0.2	1-50	0.1-10	10-100
61	0.74	1-50	0.1-10	10-100
62	0.068	1-50	0.1-10	10-100
63	0.23	1-50	0.1-10	10-100

Table 7-4

COM. NO.	H-Ras IC ₅₀ (μ M)	H-Ras CIC ₅₀ (μ M)	K-Ras IC ₅₀ (μ M)	K-Ras CIC ₅₀ (μ M)
64	0.16	1-50	0.1-10	10-100
65	0.42	1-50	0.1-10	10-100
66	0.12	1-50	0.1-10	10-100
67	0.02	3	0.1-10	> 50
68	0.12	1-50	0.1-10	10-100
69	0.55	1-50	0.1-10	10-100
70	0.21	1-50	0.1-10	10-100
71	0.12	1-50	0.1-10	10-100
72	0.05	1-50	0.1-10	10-100
73	0.002	0.2	0.02	> 10
74	0.01	0.1-1	0.01-0.1	10-100
75	0.005	0.2	0.16	20
76	0.004	0.1-1	0.1-10	10-100
77	0.004	0.1	0.12	20
78	0.0045	0.1	0.2	10-100
79	0.005	0.1	0.1	> 50
80	8.21	1-50	0.1-10	10-100
81	0.68	1-50	0.1-10	10-100
82	0.4	1-50	0.1-10	10-100
83	0.26	1-50	18.5	10-100
84	0.72	1-50	1.83	10-100
85	0.03	4	0.1-10	> 50

Table 7-5

COM. NO.	H-Ras IC ₅₀ (μ M)	H-Ras CIC ₅₀ (μ M)	K-Ras IC ₅₀ (μ M)	K-Ras CIC ₅₀ (μ M)
86	0.03	2	0.1-10	> 50
87	0.06	1-50	0.1-10	10-100
88	0.6	1-50	0.1-10	10-100
89	0.014	1	0.1-10	10-100
90	0.0425	1-50	0.1-10	10-100
91	2.15	1-50	0.1-10	10-100
92	0.07	1-50	0.1-10	10-100
93	0.32	1-50	0.1-10	10-100
94	0.2	1-50	0.1-10	10-100
95	0.0007	0.01-0.1	0.1-1	10-50
96	0.23	1-50	1-10	10-100
97	12	10-100	10-100	10-100
98	0.90-0.9	1-50	0.1-10	10-100
99	0.0030	0.1	0.1	> 50
100	1.8	> 1	0.1-10	> 20
101	0.01	> 5	0.8	> 50
102	0.45	1-50	22	> 50
103	0.064	0.1-10	1.7	10-100
104	0.0005	< 0.05	0.006	< 10
105	0.0004	0.05	0.09	> 50
106	0.9	1-50	10-100	10-100
107	10	1-50	0.1-10	10-100

Table 7-6

COM. NO.	H-Ras IC ₅₀ (μ M)	H-Ras CIC ₅₀ (μ M)	K-Ras IC ₅₀ (μ M)	K-Ras CIC ₅₀ (μ M)
108	0.26	1-50	0.1-10	10-100
109	8.6	1-50	1-50	10-100
110	0.0006	0.008	0.0015	10
111	0.002	0.03	0.002	4
112	0.004	0.015	0.006	10
113	0.004	<0.1	<0.1	10-100
114	0.001	0.015	0.100	<100
115	0.002	0.025	0.035	<50
116	0.004	0.030	0.062	<50
117	-	-	-	<50
118	-	-	-	<40
119	-	-	-	<30
120	-	-	-	<20
121	-	-	-	<40
122	-	-	-	<30
123	-	-	-	<40
124	-	-	-	<20
125	0.002	0.006	0.004	4
126	0.001	0.012	0.004	5
127	0.002	0.015	0.005	5
128	0.002	0.010	0.010	5
129	0.003	0.025	0.004	10-50

Table 7-7

COM. NO.	H-Ras IC ₅₀ (μ M)	H-Ras CIC ₅₀ (μ M)	K-Ras IC ₅₀ (μ M)	K-Ras CIC ₅₀ (μ M)
130	0.002	0.025	0.003	10-50
131	0.001	0.0125	0.0023	10-50
132	0.0035	0.025	0.011	10-50
133	0.00065	0.025	0.002	10-50
134	0.0027	0.025	0.002	10-50
135	0.0024	0.03	0.004	10-50
136	0.0016	0.025	0.0024	10-50
137	0.0017	0.020	0.0021	10-20
138	0.0014	0.025	0.0035	10-50
139	0.005	0.07	37	7
140	0.09	1-10	10-50	10-50
141	0.23	1-10	10-100	10-50
142	12	>50	>50	>50
143	1.2	20	>50	>50
144	0.38	5	50	>50
145	0.007	0.1	0.07	25
146	0.09	1	10	50
147	0.002	0.05	0.03	10
148	1.7	30	>50	>50
149	5	50	>50	>50
150	8	>50	>50	>50
151	4.6	50	>50	>50

Table 7-8


COM. NO.	H-Ras IC ₅₀ (μ M)	H-Ras CIC ₅₀ (μ M)	K-Ras IC ₅₀ (μ M)	K-Ras CIC ₅₀ (μ M)
152	0.023	0.1	0.07	10
153	0.03	0.15	0.1	20
154	0.03	0.15	0.2	10
155	0.02	0.1	0.2	15
156	0.02	0.1	0.2	40
157	0.01	0.1	5	> 50
158	0.25	1	2	30
159	0.3	1.2	4	50
160	0.3	1.5	3	40
161	0.2	1	2	50
162	0.25	1	2	50
163	0.15	1	10	> 50



WHAT IS CLAIMED IS:


1. An imidazole derivative represented by the following formula (1):

[Formula 1]




in which

- n_1 represents an integer of 1 to 4,
 A represents hydrogen; straight-chain or branched C_1 - C_{10} -alkyl which may be optionally substituted by C_3 - C_7 -cycloalkyl or lower alkoxy; or a radical selected from the following group:



wherein

- R_1 and $R_{1'}$ independently of one another represent hydrogen, halogen, cyano, nitro, hydroxycarbonyl, aminocarbonyl, aminothiocarbonyl, lower alkoxy, phenoxy, phenyl, benzyloxy, or lower alkyl which may be optionally substituted by C_3 - C_6 -cycloalkyl,
 R_2 represents hydrogen or lower alkyl, or represents -E-F wherein E is $-CH_2-$, $-C(O)-$ or $-S(O)_2-$ and F is hydrogen; lower alkyl which may be optionally substituted by phenoxy or biphenyl; lower alkoxy which may be optionally substituted by aryl; phenyl; benzyl; benzyloxy; or amino which may be optionally substituted by lower alkyl, benzyl or C_5 - C_6 -cycloalkyl,
 R_3 represents hydrogen, lower alkyl or phenyl,
 R_4 represents a radical selected from the following group:



wherein

n_2 and n_3 independently of one another denote 0, 1, 2, 3 or 4,


R_5 and R_9 independently of one another represent hydrogen, lower alkyl, lower alkoxy, phenoxy, phenyl, hydroxy or halogen,

R_6 and R_8 independently of one another represent hydrogen, lower alkyl, lower alkoxy, phenoxy, phenyl, cyano, hydroxy or halogen,

R_7 represents hydrogen; lower alkyl which may be optionally substituted by C_3 - C_6 -cycloalkyl; lower alkoxy; hydroxy; C_3 - C_6 -cycloalkyl; di(lower alkyl)amino; phenyl; phenoxy; or halogen,

R_{10} represents hydrogen, lower alkyl or lower alkoxy,

Y represents a radical selected from the following group:




wherein

X represents O or S,

B represents hydrogen, or lower alkyl which may be optionally substituted by hydroxy, mercapto, lower alkoxy, lower alkylthio or aryl,

C represents hydrogen, or lower alkyl which may be optionally substituted by aryl; or represents a radical selected from the following group:




wherein

R_{11} and R_{12} independently of one another represent hydrogen, lower alkyl, lower alkoxy, halogen, cyano, hydroxycarbonyl, amino-carbonyl, aminothiocabonyl, hydroxy, phenyl or phenoxy,

R_{13} and R_{14} independently of one another represent hydrogen, lower

alkyl, aryl or $-(CH_2)_{n_4}-X-R_{15}$ wherein X is defined as previously described, n_4 is an integer of 2 to 4 and R_{15} is lower alkyl,

D represents amino acid residue or lower alkyl ester of amino acid residue; or represents a radical selected from the following group:



wherein

R_{10} is defined as previously described,

Q represents O, S, S=O or SO₂,

Z represents O, S, S=O, SO₂, C=O or C=S, or represents CH-R₂₀ or N-R₂₀ (wherein R₂₀ is hydrogen, lower alkyl or hydroxy),




n_5 denotes an integer of 1 to 3,

R_{16} and R_{17} independently of one another represents hydrogen; aryl; lower alkyl which may be optionally substituted by aryl or

cyanoaryl; or $-\frac{1}{2}(\text{CH}_2)_{n_4}-\text{Q}-\text{R}_{10}$ wherein n_4 , Q and R_{10} are defined as previously described,

R_{18} and R_{19} independently of one another represents hydrogen; halogen; hydroxy; cyano; lower alkyl; lower alkoxy; alkoxyalkyl; alkylthio; hydroxycarbonyl; aminocarbonyl; aminothiocarbonyl; alkylsulfonyl; alkylthioalkyl; alkylthioalkyloxy; aryl; or oxy, thio, sulfonyl or lower alkyl substituted by aryl,


G represents a radical selected by the following group:



wherein

R_{11} and R_{12} are defined as previously described,


I represents lower alkoxy, or represents a radical selected from the following group:



wherein


R_{16} , R_{17} and Z are defined as previously described,


L represents a radical selected from the following group:




wherein Z and Q are defined as previously described,

provided that (1) n_2 is other than 0 when R_3 is hydrogen, and





(2) Y is other than , when A is



, or a pharmaceutically acceptable salt or isomer thereof.

2. The compound of claim 1 wherein
 - n_1 represents an integer of 1 to 3,
 - A represents hydrogen; straight-chain or branched C₁-C₁₀-alkyl which may be optionally substituted by C₃-C₇-cycloalkyl or lower alkoxy; or a radical selected from the following group:



wherein

R_1 and R_1' independently of one another represent hydrogen, halogen, cyano, nitro, hydroxycarbonyl, aminocarbonyl, aminothiocarbonyl, lower alkoxy, phenoxy, phenyl, benzyloxy, or lower alkyl which may be optionally substituted by C₃-C₆-cycloalkyl,


R_2 represents hydrogen or lower alkyl, or represents -E-F wherein E is -CH₂-, -C(O)- or -S(O)₂- and F is hydrogen; lower alkyl which may be optionally substituted by phenoxy or biphenyl; lower alkoxy which may be optionally substituted by aryl; phenyl; benzyl; benzyloxy; or amino which may be optionally substituted



by lower alkyl, benzyl or C₅-C₆-cycloalkyl,

R₃ represents hydrogen or lower alkyl,

R₄ represents a radical selected from the following group:



wherein

n₂ and n₃ independently of one another denote 0, 1, 2, 3 or 4,


R₅, R₆, R₈ and R₉ independently of one another represent hydrogen,

lower alkyl, lower alkoxy, hydroxy or halogen,

R₇ represents hydrogen; lower alkyl which may be optionally substituted by C₃-C₆-cycloalkyl; lower alkoxy; hydroxy; C₃-C₆-cycloalkyl; or halogen,

R₁₀ represents hydrogen, methyl or methoxy,

Y represents a radical selected from the following group:




wherein

X represents O or S,

B represents hydrogen, or lower alkyl which may be optionally substituted by lower alkoxy or aryl,

C represents hydrogen, or lower alkyl which may be optionally substituted by aryl; or represents a radical selected from the following group:




wherein

R_{11} and R_{12} independently of one another represent hydrogen, lower alkyl, lower alkoxy, halogen, cyano, aminocarbonyl, phenyl or phenoxy,

R_{13} and R_{14} independently of one another represent hydrogen, lower

alkyl, aryl or $-\frac{1}{2}(\text{CH}_2)_{n_4}-\text{X}-R_{15}$ wherein X is defined as previously described, n_4 is 2 and R_{15} is lower alkyl,

D represents amino acid residue or lower alkyl ester of amino acid residue; or represents a radical selected from the following group:



wherein

R_{10} is defined as previously described,

Q represents O, S, S=O or SO₂,

Z represents O, S, S=O, SO₂ or C=O, or represents CH-R₂₀ or N-R₂₀ (wherein R₂₀ is hydrogen, lower alkyl or hydroxy),




n_5 denotes an integer of 1 to 3,

R_{16} and R_{17} independently of one another represents hydrogen; aryl; lower alkyl which may be optionally substituted by aryl or

cyanoaryl; or $-\frac{1}{2}(\text{CH}_2)_{n_4}-\text{Q}-\text{R}_{10}$ wherein n_4 , Q and R_{10} are defined as previously described,

R_{18} and R_{19} independently of one another represents hydrogen; halogen; hydroxy; cyano; lower alkyl; lower alkoxy; alkoxyalkyl; alkylthio; hydroxycarbonyl; aminocarbonyl; aminothiocarbonyl; alkylsulfonyl; alkylthioalkyl; alkylthioalkyloxy; aryl; or oxy, thio, sulfonyl or lower alkyl substituted by aryl,


G represents a radical selected by the following group:



wherein

R_{11} and R_{12} are defined as previously described,


I represents lower alkoxy, or represents a radical selected from the following group:



wherein

R_{16} , R_{17} and Z are defined as previously described,

L represents a radical selected from the following group:




wherein Z and Q are defined as previously described,





provided that (1) n_2 is other than 0 when R_3 is hydrogen, and


(2) Y is other than



, when A is



3. The compound of claim 1 wherein Y represents



and C represents

4. The compound of claim 1 which is selected from a group consisting of:

3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-1-[1-(3,4-methylenedioxybenzyl)-1H-imidazol-5-ylmethyl]-4-(naphthalen-1-yl)-1H-pyrrole(1),

1-[1-(3,4-methylenedioxybenzyl)-1H-imidazol-5-ylmethyl]-3-(morpholin-4-yl)carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(2),

1-[1-(3,4-methylenedioxybenzyl)-1H-imidazol-5-ylmethyl]-3-(4-methylpiperazin-1-yl)carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(3),

3-{N-[2-(N,N-dimethylamino)ethyl]-N-methyl}carbamoyl-1-[1-(3,4-methylenedioxybenzyl)-1H-imidazol-5-ylmethyl]-4-(naphthalen-1-yl)-1H-pyrrole(4),

- 3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-naphthalen-1-yl)-1-[1-naphthalen-1-ylmethyl]-1H-imidazol-5-ylmethyl]-1H-pyrrole(5),
- 3-(morpholin-4-yl)carbonyl-4-(naphthalen-1-yl)-1-[1-naphthalen-1-ylmethyl]-1H-imidazol-5-ylmethyl]-1H-pyrrole(6),
- 3-(4-methylpiperazin-1-yl)carbonyl-4-(naphthalen-1-yl)-1-[1-(naphthalen-1-ylmethyl)-1H-imidazol-5-ylmethyl]-1H-pyrrole(7),
- 3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-1-[1-((R)- α -methylbenzyl)-1H-imidazol-5-ylmethyl]-4-(naphthalen-1-yl)-1H-pyrrole(8),
- 1-[1-((R)- α -methylbenzyl)-1H-imidazol-5-ylmethyl]-3-(morpholin-4-yl)carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(9),
- 1-[1-((R)- α -methylbenzyl)-1H-imidazol-5-ylmethyl]-3-(4-methylpiperazin-1-yl)carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(10),
- 3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-1-[1-((S)- α -methylbenzyl)-1H-imidazol-5-ylmethyl]-4-(naphthalen-1-yl)-1H-pyrrole(11),
- 1-[1-((S)- α -methylbenzyl)-1H-imidazol-5-ylmethyl]-3-(morpholin-4-yl)carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(12),
- 1-[1-((S)- α -methylbenzyl)-1H-imidazol-5-ylmethyl]-3-(4-methylpiperazin-1-yl)carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(13),
- 3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)-1-[1-(phenethyl)-1H-imidazol-5-ylmethyl]-1H-pyrrole(14),
- 3-(morpholin-4-yl)carbonyl-4-(naphthalen-1-yl)-1-[1-(phenethyl)-1H-imidazol-5-ylmethyl]-1H-pyrrole(15),
- 3-(4-methylpiperazin-1-yl)carbonyl-4-(naphthalen-1-yl)-1-[1-(phenethyl)-1H-imidazol-5-ylmethyl]-1H-pyrrole(16),
- 3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-1-[1-(2-methoxy)phenethyl-1H-imidazol-5-yl]methyl-4-(naphthalen-1-yl)-1H-pyrrole(17),
- 1-[1-(2-methoxy)phenethyl-1H-imidazol-5-yl]methyl-3-[4-methylpiperazin-1-yl]carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(18),
- 3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-1-[1-(4-methoxy)phenethyl-1H-imidazol-5-yl]methyl-4-(naphthalen-1-yl)-1H-pyrrole(19),



- 1-[1-(4-methoxy)phenethyl-1H-imidazol-5-yl]methyl-3-[4-methylpiperazin-1-yl]carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(20),
- 1-[1-(2-fluoro)phenethyl-1H-imidazol-5-yl]methyl-3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole(21),
- 1-[1-(2-fluoro)phenethyl-1H-imidazol-5-yl]methyl-3-[4-methylpiperazin-1-yl]carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(22),
- 1-[1-(2-chloro)phenethyl-1H-imidazol-5-yl]methyl-3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole(23),
- 1-[1-(2-chloro)phenethyl-1H-imidazol-5-yl]methyl-3-[4-methylpiperazin-1-yl]carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(24),
- 1-[1-(3-chloro)phenethyl-1H-imidazol-5-yl]methyl-3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole(25),
- 1-[1-(3-chloro)phenethyl-1H-imidazol-5-yl]methyl-3-[4-methylpiperazin-1-yl]carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(26),
- 3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)-1-[1-(3-phenyl)propyl-1H-imidazol-5-yl]methyl-1H-pyrrole(27),
- 3-[4-methylpiperazin-1-yl]carbonyl-4-naphthalen-1-yl)-1-[1-(3-phenyl)propyl-1H-imidazol-5-yl]methyl-1H-pyrrole(28),
- 3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)-1-[1-(naphthalen-2-yl)methyl-1H-imidazol-5-yl]methyl-1H-pyrrole(29),
- 3-[4-methylpiperazin-1-yl]carbonyl-4-(naphthalen-1-yl)-1-[1-naphthalen-2-yl)methyl-1H-imidazol-5-yl]methyl-1H-pyrrole(30),
- 3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)-1-{1-[2-(naphthalen-1-yl)ethyl]-1H-imidazol-5-yl}methyl-1H-pyrrole(31),
- 3-[4-methylpiperazin-1-yl]carbonyl-4-(naphthalen-1-yl)-1-{1-[2-(naphthalen-1-yl)ethyl]-1H-imidazol-5-yl}methyl-1H-pyrrole(32),
- 1-[1-(4-bromo)phenethyl-1H-imidazol-5-yl]methyl-3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole(33),
- 1-[1-(4-bromo)phenethyl-1H-imidazol-5-yl]methyl-3-[4-methylpiperazin-1-yl]carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(34),



* C L H S *

- 1-[1-(4-fluoro)phenethyl-1H-imidazol-5-yl]methyl-3-[N-(2-methoxyethyl)-N-methyl]carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(35),
- 1-[1-(4-fluoro)phenethyl-1H-imidazol-5-yl]methyl-3-[4-methylpiperazin-1-yl]carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(36),
- 3-[N-(2-methoxyethyl)-N-methyl]carbonyl-1-[1-(4-methyl)phenethyl-1H-imidazol-5-yl]methyl-4-(naphthalen-1-yl)-1H-pyrrole(37),
- 1-[1-(4-methyl)phenethyl-1H-imidazol-5-yl]methyl-3-[4-methylpiperazin-1-yl]carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(38),
- 1-[1-(4-chloro)phenethyl-1H-imidazol-5-yl]methyl-3-[N-(2-methoxyethyl)-N-methyl]carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(39),
- 1-[1-(4-chloro)phenethyl-1H-imidazol-5-yl]methyl-3-[4-methylpiperazin-1-yl]carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(40),
- 3-[N-(2-methoxyethyl)-N-methyl]carbonyl-4-(naphthalen-1-yl)-1-{1-[2-(naphthalen-2-yl)ethyl]-1H-imidazol-5-yl}methyl-1H-pyrrole(41),
- 3-[4-methylpiperazin-1-yl]carbonyl-4-(naphthalen-1-yl)-1-{1-[2-(naphthalen-2-yl)ethyl]-1H-imidazol-5-yl}methyl-1H-pyrrole(42),
- 1-[1-(4-hydroxy)phenethyl-1H-imidazol-5-yl]methyl-3-[4-methylpiperazin-1-yl]carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(43),
- 1-(1H-imidazol-4-yl)methyl-4-(naphthalen-1-yl)-3-(thiophen-2-yl)carbonyl-1H-pyrrole(44),
- 1-(1H-imidazol-4-yl)methyl-4-(naphthalen-1-yl)-3-(thiophen-3-yl)carbonyl-1H-pyrrole(45),
- 3-benzoyl-1-(1H-imidazol-4-yl)methyl-4-(naphthalen-1-yl)-1H-pyrrole(46),
- 3-(2-bromobenzoyl)-1-(1H-imidazol-4-yl)methyl-4-(naphthalen-1-yl)-1H-pyrrole(47),
- 3-(3-bromobenzoyl)-1-(1H-imidazol-4-yl)methyl-4-(naphthalen-1-yl)-1H-pyrrole(48),
- 3-(4-bromobenzoyl)-1-(1H-imidazol-4-yl)methyl-4-(naphthalen-1-yl)-1H-pyrrole(49),
- 1-(1H-imidazol-4-yl)methyl-3-(2-methylbenzoyl)-4-(naphthalen-1-yl)-1H-pyrrole(49),



* C L M S *

ole(50),

1-(1H-imidazol-4-yl)methyl-3-(3-methylbenzoyl)-4-(naphthalen-1-yl)-1H-pyrrole(51),

1-(1H-imidazol-4-yl)methyl-3-(4-methylbenzoyl)-4-(naphthalen-1-yl)-1H-pyrrole(52),

1-(1H-imidazol-4-yl)methyl-3-(3-methoxybenzoyl)-4-(naphthalen-1-yl)-1H-pyrrole(53),

1-(1H-imidazol-4-yl)methyl-3-(4-methoxybenzoyl)-4-(naphthalen-1-yl)-1H-pyrrole(54),

3-(2-chlorobenzoyl)-1-(1H-imidazol-4-yl)methyl-4-(naphthalen-1-yl)-1H-pyrrole(55),

3-(4-chlorobenzoyl)-1-(1H-imidazol-4-yl)methyl-4-(naphthalen-1-yl)-1H-pyrrole(56),

3-(2,4-dichlorobenzoyl)-1-(1H-imidazol-4-yl)methyl-4-(naphthalen-1-yl)-1H-pyrrole(57),

3-(4-fluorobenzoyl)-1-(1H-imidazol-4-yl)methyl-4-(naphthalen-1-yl)-1H-pyrrole(58),

3-(2,4-difluorobenzoyl)-1-(1H-imidazol-4-yl)methyl-4-(naphthalen-1-yl)-1H-pyrrole(59),

3-(4-cyanobenzoyl)-1-(1H-imidazol-4-yl)methyl-4-(naphthalen-1-yl)-1H-pyrrole(60),

1-(1H-imidazol-4-yl)methyl-3-(4-methylthiomethyl-benzoyl)-4-(naphthalen-1-yl)-1H-pyrrole(61),

1-(1H-imidazol-4-yl)methyl-3-[4-(2-methylthioethyl)benzoyl]-4-(naphthalen-1-yl)-1H-pyrrole(62),

1-(1H-imidazol-4-yl)methyl-3-[4-(2-methylthioethoxy)benzoyl]-4-(naphthalen-1-yl)-1H-pyrrole(63),

1-(1H-imidazol-4-yl)methyl-3-(3-methylthiomethyl-benzoyl)-4-(naphthalen-1-yl)-1H-pyrrole(64),

1-(1H-imidazol-4-yl)methyl-4-(naphthalen-1-yl)-3-(3-phenylbenzoyl)-1H-pyrrole(65),



* C L M S *

- ole(65),
1-(1H-imidazol-4-yl)methyl-4-(naphthalen-1-yl)-3-(4-phenylbenzoyl)-1H-pyrrole(66),
1-(1H-imidazol-4-yl)methyl-4-(naphthalen-1-yl)-3-(4-phenoxybenzoyl)-1H-pyrrole(67),
3-(4-benzylbenzoyl)-1-(1H-imidazol-4-yl)methyl-4-(naphthalen-1-yl)-1H-pyrrole(68),
1-(1H-imidazol-4-yl)methyl-4-(naphthalen-1-yl)-3-(naphthalen-1-yl)carbonyl-1H-pyrrole(69),
1-(1H-imidazol-4-yl)methyl-3-(4-methylbenzoyl)-4-(N-methylindol-3-yl)-1H-pyrrole(70),
5-n-butyl-3-(2,4-dichlorobenzoyl)-1-(1H-imidazol-4-yl)methyl-4-(naphthalen-1-yl)-1H-pyrrole(71),
5-benzyl-3-(2,4-dichlorobenzoyl)-1-(1H-imidazol-4-yl)methyl-4-(naphthalen-1-yl)-1H-pyrrole(72),
1-[1-(4-cyanobenzyl)-1H-imidazol-5-yl]methyl-4-(naphthalen-1-yl)-3-(thiophen-2-yl)carbonyl-1H-pyrrole(73),
1-[1-(4-cyanobenzyl)-1H-imidazol-5-yl]methyl-4-(naphthalen-1-yl)-3-(thiophen-3-yl)carbonyl-1H-pyrrole(74),
3-benzoyl-1-[1-(4-cyanobenzyl)-1H-imidazol-5-yl]methyl-4-(naphthalen-1-yl)-1H-pyrrole(75),
3-(3-bromobenzoyl)-1-[1-(4-cyanobenzyl)-1H-imidazol-5-yl]methyl-4-(naphthalen-1-yl)-1H-pyrrole(76),
3-(4-bromobenzoyl)-1-[1-(4-cyanobenzyl)-1H-imidazol-5-yl]methyl-4-(naphthalen-1-yl)-1H-pyrrole(77),
3-(4-fluorobenzoyl)-1-(1-methyl-1H-imidazol-4-yl)methyl-4-(naphthalen-1-yl)-1H-pyrrole(78),
1-(1-methyl-1H-imidazol-4-yl)methyl-4-(naphthalen-1-yl)-3-(4-phenoxybenzoyl)-1H-pyrrole(79),
(S)-1-(1H-imidazol-4-yl)methyl-3-[N-(1-methoxycarbonyl-3-methylthio)propyl



]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole(80),
(S)-3-[N-(1-hydroxycarbonyl-3-methylthio)propyl]carbamoyl-1-(1H-imidazol-4-yl)methyl-4-(naphthalen-1-yl)-1H-pyrrole(81),
1-(1H-imidazol-4-yl)methyl-4-(naphthalen-1-yl)-3-(N-phenylcarbamoyl)-1H-pyrrole(82),
3-(N-benzylcarbamoyl)-1-(1H-imidazol-4-yl)methyl-4-(naphthalen-1-yl)-1H-pyrrole(83),
1-(1H-imidazol-4-yl)methyl-4-(naphthalen-1-yl)-3-(piperidin-1-yl)carbonyl-1H-pyrrole(84),
1-(1H-imidazol-4-yl)methyl-3-(morpholin-4-yl)carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(85),
1-(1H-imidazol-4-yl)methyl-4-(naphthalen-1-yl)-3-(thiomorpholin-4-yl)carbonyl-1H-pyrrole(86),
1-(1H-imidazol-4-yl)methyl-4-(naphthalen-1-yl)-3-(S,S-dioxothiomorpholin-4-yl)carbonyl-1H-pyrrole(87),
1-(1H-imidazol-4-yl)methyl-4-(naphthalen-1-yl)-3-(piperazin-1-yl)carbonyl-1H-pyrrole(88),
1-(1H-imidazol-4-yl)methyl-3-(4-methylpiperazin-1-yl)carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(89),
1-(1H-imidazol-4-yl)methyl-4-(naphthalen-1-yl)-3-(thiazolidin-3-yl)carbonyl-1H-pyrrole(90),
3-(4-hydroxypiperidin-1-yl)carbonyl-1-(1H-imidazol-4-yl)methyl-4-(naphthalen-1-yl)-1H-pyrrole(91),
1-(1H-imidazol-4-yl)methyl-4-(naphthalen-1-yl)-3-(4-oxopiperidin-1-yl)carbonyl-1H-pyrrole(92),
3-N-(2-hydroxyethyl)carbamoyl-1-(1H-imidazol-4-yl)methyl-4-(naphthalen-1-yl)-1H-pyrrole(93),
3-[N-(2-hydroxyethyl)-N-methyl]carbamoyl-1-(1H-imidazol-4-yl)methyl-4-(naphthalen-1-yl)-1H-pyrrole(94),
1-(1H-imidazol-4-yl)methyl-3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-(na



phthalen-1-yl)-1H-pyrrole(95),
1-(1H-imidazol-4-yl)methyl-3-(morpholin-4-yl)carbonyl-4-(quinolin-4-yl)-1H-pyrrole(96),
4-(1,2-dihydroacenaphthylen-5-yl)-1-(1H-imidazol-4-yl)methyl-3-(morpholin-4-yl)carbonyl-1H-pyrrole(97),
3-N-(4-cyanobenzyl)carbamoyl-1-(1H-imidazol-4-yl)methyl-4-(naphthalen-1-yl)-1H-pyrrole(98),
1-(1-methyl-1H-imidazol-5-yl)methyl-3-(morpholin-4-yl)carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(99),
(S)-1-[1-(4-cyanobenzyl)-1H-imidazol-5-yl]methyl-3-[N-(1-methoxycarbonyl-3-methylthio)propyl]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole(100),
(S)-1-[1-(4-cyanobenzyl)-1H-imidazol-5-yl]methyl-3-[N-(1-hydroxycarbonyl-3-methylthio)propyl]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole(101),
(S)-1-[1-(4-cyanobenzyl)-1H-imidazol-5-yl]methyl-3-[N-(1-methoxycarbonyl-3-methyl)butyl]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole(102),
(S)-1-[1-(4-cyanobenzyl)-1H-imidazol-5-yl]methyl-3-[N-(1-hydroxycarbonyl-3-methyl)butyl]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole(103),
1-[1-(4-cyanobenzyl)-1H-imidazol-5-yl]methyl-3-(morpholin-4-yl)carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(104),
1-[1-(4-cyanobenzyl)-1H-imidazol-5-yl]methyl-3-(4-methylpiperazin-1-yl)carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(105),
1-[2-(1H-imidazol-1-yl)ethyl]-3-(morpholin-4-yl)carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(106),
(S)-1-[3-(1H-imidazol-4-yl)propyl]-3-[N-(1-methoxycarbonyl-3-methylthio)propyl]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole(107),
(S)-3-[N-(1-hydroxycarbonyl-3-methylthio)propyl]carbamoyl-1-[3-(1H-imidazol-4-yl)propyl]-4-(naphthalen-1-yl)-1H-pyrrole(108),
1-[3-(1H-imidazol-4-yl)propyl]-3-(morpholin-4-yl)carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(109),
1-[1-(4-cyanobenzyl)-1H-imidazol-5-yl]methyl-3-[N-(2-methoxyethyl)-N-meth



* C L H S *

- yl]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole(110),
1-[1-(4-bromobenzyl)-1H-imidazol-5-yl]methyl-3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole(111),
1-[1-(4-bromobenzyl)-1H-imidazol-5-yl]methyl-3-(morpholin-4-yl)carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(112),
1-[1-(4-cyanobenzyl)-1H-imidazol-5-yl]methyl-3-(morpholin-4-yl)thiocarbonyl-4-(naphthalen-1-yl)-1H-pyrrole(113),
3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-1-(1-methyl-1H-imidazol-5-yl)methyl-4-(naphthalen-1-yl)-1H-pyrrole(114),
1-(1-isobutyl-1H-imidazol-5-yl)methyl-3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole(115),
1-(1-cyclohexylmethyl-1H-imidazol-5-yl)methyl-3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole(116),
3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)-1-(1-pentyl-1H-imidazol-5-yl)methyl-1H-pyrrole(117),
3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)-1-(1-octyl-1H-imidazol-5-yl)methyl-1H-pyrrole(118),
1-(1-decyl-1H-imidazol-5-yl)methyl-3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole(119),
3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-1-[1-(3-methylbutyl)-1H-imidazol-5-yl]methyl-4-(naphthalen-1-yl)-1H-pyrrole(120),
1-[1-(2-methoxyethyl)-1H-imidazol-5-yl]methyl-3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole(121),
3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-1-[1-(3-methoxypropyl)-1H-imidazol-5-yl]methyl-4-(naphthalen-1-yl)-1H-pyrrole(122),
1-[1-(3-ethoxypropyl)-1H-imidazol-5-yl]methyl-3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole(123),
1-[1-(3-isopropoxypropyl)-1H-imidazol-5-yl]methyl-3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole(124),
1-[1-(4-bromobenzyl)-1H-imidazol-5-yl]methyl-3-[4-methylpiperazin-1-yl]carb



* C L H S *

onyl-4-(naphthalen-1-yl)-1H-pyrrole(125),
1-[1-(4-chlorobenzyl)-1H-imidazol-5-yl]methyl-3-[4-methylpiperazin-1-yl]carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(126),
1-[1-(4-fluorobenzyl)-1H-imidazol-5-yl]methyl-3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole(127),
1-[1-(4-fluorobenzyl)-1H-imidazol-5-yl]methyl-3-[4-methylpiperazin-1-yl]carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(128),
1-[1-(4-methoxybenzyl)-1H-imidazol-5-yl]methyl-3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole(129),
1-[1-(4-methoxybenzyl)-1H-imidazol-5-yl]methyl-3-(4-methylpiperazin-1-yl)carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(130),
1-[1-(3-chlorobenzyl)-1H-imidazol-5-yl]methyl-3-(4-methylpiperazin-1-yl)carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(131),
1-[1-(3-chlorobenzyl)-1H-imidazol-5-yl]methyl-3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole(132),
1-[1-(2-chlorobenzyl)-1H-imidazol-5-yl]methyl-3-(4-methylpiperazin-1-yl)carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(133),
1-[1-(2-chlorobenzyl)-1H-imidazol-5-yl]methyl-3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)-1H-pyrrole(134),
1-[1-(2-fluorobenzyl)-1H-imidazol-5-yl]methyl-3-(4-methylpiperazin-1-yl)carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(135),
1-[1-(4-methylbenzyl)-1H-imidazol-5-yl]methyl-3-(4-methylpiperazin-1-yl)carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(136),
1-[1-(4-methylbenzyl)-1H-imidazol-5-yl]methyl-3-(morpholin-4-yl)carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(137),
1-[1-(3-methylbenzyl)-1H-imidazol-5-yl]methyl-3-(4-methylpiperazin-1-yl)carbonyl-4-(naphthalen-1-yl)-1H-pyrrole(138),
1-[1-(4-cyanobenzyl)-1H-imidazol-5-yl]methyl-3-(naphthalen-1-yl)carbonyl-1H-pyrrole(139),
1-[1-(4-bromobenzyl)-1H-imidazol-5-yl]methyl-3-(naphthalen-1-yl)carbonyl-






- 1H-pyrrole(140),
 1-[1-(4-bromobenzyl)-1H-imidazol-5-yl]methyl-3-[N-(2-methoxyethyl)-N-methyl]carbamoyl-4-(naphthalen-1-yl)carbonyl-1H-pyrrole(141),
 4-ethoxycarbonyl-2-(1H-imidazol-5-ylmethyl)-5-(naphthalen-1-yl)oxazole(142),
 2-(1H-imidazol-5-ylmethyl)-4-(morpholin-4-yl)carbonyl-5-(naphthalen-1-yl)oxazole(143),
 4-ethoxycarbonyl-2-(1H-imidazol-5-ylmethyl)-5-(naphthalen-1-yl)thiazole(144),
 2-[1-(4-chlorobenzyl)-1H-imidazol-5-ylmethyl]-4-methoxycarbonyl-5-(naphthalen-1-yl)thiazole(145),
 2-[1-(4-chlorobenzyl)-1H-imidazol-5-ylmethyl]-4-(morpholin-4-yl)carbonyl-5-(naphthalen-1-yl)thiazole(146),
 2-[1-(4-chlorobenzyl)-1H-imidazol-5-ylmethyl]-4-[N-(2-methoxy)ethyl-N-methylcarbamoyl]-5-(naphthalen-1-yl)thiazole(147),
 2-[1-(4-chlorobenzyl)-1H-imidazol-5-ylmethyl]-5-methoxycarbonyl-4-(naphthalen-1-yl)thiazole(148),
 2-[1-(4-chlorobenzyl)-1H-imidazol-5-ylmethyl]-5-(morpholin-4-yl)carbonyl-4-(naphthalen-1-yl)thiazole(149),
 2-{1-[1-(benzyloxycarbonyl)piperidin-4-ylmethyl]-1H-imidazol-5-ylmethyl}-5-methoxycarbonyl-4-(naphthalen-1-yl)thiazole(150),
 2-{1-[1-(benzyloxycarbonyl)piperidin-4-ylmethyl]-1H-imidazol-5-ylmethyl}-5-[N-(2-methoxy)ethyl-N-methylcarbamoyl]-4-(naphthalen-1-yl)thiazole(151),
 1-[1-(1-benzyloxycarbonyl-piperidin-4-ylmethyl)-1H-imidazol-5-ylmethyl]-4-[N-(2-methoxyethyl)-N-methyl]carbamoyl-3-(naphthalen-1-yl)-1H-pyrazole(152),
 1-[1-(1-methoxycarbonylpiperidin-4-ylmethyl)-1H-imidazole-5-ylmethyl]-4-[N-(2-methoxyethyl)-N-methyl]carbamoyl-3-(naphthalen-1-yl)-1H-pyrazole(153),
 1-[1-(4-bromobenzyl)-1H-imidazol-5-ylmethyl]-4-[N-(2-methoxyethyl)-N-methyl]carbamoyl-3-(naphthalen-1-yl)-1H-pyrazole(154),




1-[1-(4-chlorobenzyl)-1H-imidazol-5-ylmethyl]-4-[N-(2-methoxyethyl)-N-methyl]carbamoyl-3-(naphthalen-1-yl)-1H-pyrazole(155),
 1-[1-(4-cyanobenzyl)-1H-imidazol-5-ylmethyl]-4-[N-(2-methoxyethyl)-N-methyl]carbamoyl-3-(naphthalen-1-yl)-1H-pyrazole(156),
 1-[1-methyl-1H-imidazol-5-ylmethyl]-4-[N-(2-methoxyethyl)-N-methyl]carbamoyl-3-(naphthalen-1-yl)-1H-pyrazole(157),
 1-[1-(1-benzylloxycarbonyl-piperidin-4-ylmethyl)-1H-imidazol-5-ylmethyl]-4-(morpholin-4-yl)carbonyl-3-(naphthalen-1-yl)-1H-pyrazole(158),
 1-[1-(1-methoxycarbonylpiperidin-4-ylmethyl)-1H-imidazol-5-ylmethyl]-4-(morpholin-4-yl)carbonyl-3-(naphthalen-1-yl)-1H-pyrazole(159),
 1-[1-(4-bromobenzyl)-1H-imidazol-5-ylmethyl]-4-(morpholin-4-yl)carbonyl-3-(naphthalen-1-yl)-1H-pyrazole(160),
 1-[1-(4-chlorobenzyl)-1H-imidazol-5-ylmethyl]-4-(morpholin-4-yl)carbonyl-3-(naphthalen-1-yl)-1H-pyrazole(161),
 1-[1-(4-cyanobenzyl)-1H-imidazol-5-ylmethyl]-4-(morpholin-4-yl)carbonyl-3-(naphthalen-1-yl)-1H-pyrazole(162), and
 1-(1-methyl-1H-imidazol-5-ylmethyl)-4-(morpholin-4-yl)carbonyl-3-(naphthalen-1-yl)-1H-pyrazole(163).

5. A process for preparing an imidazole derivative of formula (1) as defined in claim 1 characterized in that


(a) a compound represented by the following formula (2):



wherein n_1 is defined as claim 1 and Tr represents trityl, is reacted in a solvent in the presence of a base with a compound represented by the following formula (3):




wherein B, C, D and X are defined as claim 1, then the trityl group in the product thus obtained is eliminated in the presence of trifluoroacetic acid to produce a compound represented by the following formula (1a):




wherein n_1 , B, C, D and X are defined as claim 1; or

(b) a compound represented by the following formula (4):




wherein n_1 and A are defined as claim 1, is reacted in a solvent in the presence of a base with the compound of formula (3) to produce a compound represented by the following formula (1b):



wherein n_1 , A, B, C, D and X are defined as claim 1; or


(c) a compound represented by the following formula (5):



(5)




is reacted in a solvent in the presence of a base with the compound of formula (3), the trityl group in the product thus obtained is eliminated in the presence of trifluoroacetic acid to produce a compound represented by the following formula (6):



(6)


wherein B, C, D and X are defined as claim 1, and then hydrogenation reaction is carried out to produce a compound represented by the following formula (1c):



(1c)


wherein B, C, D and X are defined as claim 1; or

(d) a compound represented by the following formula (7):




(7)


wherein n_1 , A, B and C are defined as claim 1, is hydrolyzed to produce a compound represented by the following formula (8):



wherein n_1 , A, B and C are defined as claim 1, which is then reacted with a compound represented by the following formula (9):




wherein R_{16} and R_{17} are defined as claim 1, in the presence of a coupling agent to produce a compound represented by the following formula (1d):




wherein n_1 , A, B, C, R_{16} and R_{17} are defined as claim 1; or

(e) the carbonyl group in a compound represented by the following formula (1e):




wherein n_1 , A, B, C and D are defined as claim 1, is converted into the thiocarbonyl group in the presence of a sulfurizing agent to produce a compound represented by the following formula (1f):



wherein n_1 , A, B, C and D are defined as claim 1; or


(f) a compound represented by the following formula (1g):



wherein B, C, D and X are defined as claim 1, is coupled in a solvent with a compound represented by the following formula (10):




wherein R_2 is defined as claim 1 and T represents hydroxy or reactive leaving group, to produce a compound represented by the following formula (1h):




wherein R_2 , B, C, D and X are defined as claim 1; or

(g) a compound represented by the following formula (11):




wherein A, G and I are defined as claim 1, is cyclized in an inert solvent to produce a compound represented by the following formula (1i):




wherein A, G and I are defined as claim 1; or

(h) the amide group in the compound of formula (11) is converted into the thioamide group to produce a compound represented by the following formula (12):




wherein A, G and I are defined as claim 1, which is then cyclized in an inert solvent to produce a compound represented by the following formula (1j):




wherein A, G and I are defined as claim 1; or



(i) a compound represented by the following formula (13):




wherein A is defined as claim 1, is reacted in a solvent with a compound represented by the following formula (14a):




wherein G and I are defined as claim 1, to produce the compound of formula (1j); or

(j) the compound of formula (13) is reacted in a solvent with a compound represented by the following formula (14b):




wherein G and I are defined as claim 1, to produce a compound represented by the following formula (1k):



wherein A, G and I are defined as claim 1; or


(k) a compound represented by the following formula (1l):



wherein A and G are defined as claim 1 and I' represents lower alkoxy, is hydrolyzed in the presence of a base and the product thus obtained is reacted in a solvent in the presence of a coupling agent with a compound represented by the following formula (15):




wherein I'' is identical with I except that lower alkoxy is not included, to produce a compound represented by the following formula (1m):




wherein A and G are defined as claim 1 and I'' are defined as above;
or


(l) a compound represented by the following formula (16):



wherein A is defined as claim 1, is reacted in a solvent in the presence of a base with a compound represented by the following formula (17):




wherein G and L are defined as claim 1, to produce a compound represented by the following formula (1n):




wherein A, G and L are defined as claim 1; or

(m) a compound represented by the following formula (18):




wherein Cbz represents benzyloxycarbonyl, is reacted in a solvent in the presence of a base with the compound of formula (17) and deprotected to produce a compound represented by the following formula (1o):




wherein G and L are defined as claim 1, which is then coupled with a



compound represented by the following formula (19):




wherein E and F are defined as claim 1, to produce a compound represented by the following formula (1p):




wherein E, F, G and L are defined as claim 1.

6. A compound represented by the following formula (3):



wherein B, C, D and X are defined as claim 1.

7. A compound represented by the following formula (8):



wherein n_1 , A, B and C are defined as claim 1.

8. A pharmaceutical composition comprising as active ingredient a therapeutically effective amount of a compound of formula (1) as defined

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

in claim 1 or a pharmaceutically acceptable salt or isomer thereof together with a pharmaceutically acceptable carrier.

9. The pharmaceutical composition of claim 8 useful for preventing and treating cancer.

10. The pharmaceutical composition of claim 8 useful for preventing and treating restenosis.

11. The pharmaceutical composition of claim 8 useful for preventing and treating atherosclerosis.

12. The pharmaceutical composition of claim 8 useful for preventing and treating infections from hepatitis delta and related viruses.



This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED