

DESCRIPTION

NITROGEN-CONTAINING HETEROARYL COMPOUND, AND PHARMACEUTICAL USE THEREOF

5 Technical Field

[0001]

The present invention relates to a nitrogen-containing heteroaryl compound having a GLUT9 inhibitory activity, or a pharmaceutically acceptable salt thereof, a pharmaceutical
10 composition containing the same, and a pharmaceutical use thereof.

Background Art

[0002]

Uric acid is a poorly-soluble substance having a
15 molecular weight of 168 and a dissociation constant (pKa value) of 5.75 and is present in the form of uric acid or a conjugate base (urate) thereof, depending on the pH, when it is in the body fluid. In human and many other primates, due to the functional absence of urate oxidase (uricase) in the liver,
20 uric acid is the final metabolite in purine metabolism. About 70% of the uric acid in the body resulted from dietary intake or endogenous production is eliminated through urine via the kidney, and the remaining 30% is eliminated through stools via the intestinal tube.

25 [0003]

GLUT9 belongs to the family of glucose transporters encoded by SLC2A9 (Solute carrier family 2, facilitated glucose transporter member 9) genes, which was cloned as a molecule expressed in kidney, liver, placenta and the like in human
30 (Non-Patent Document 1). According to a series of subsequent reports, the genome-wide association analysis confirmed the correlation between mutations in this molecule and blood uric acid levels, and this molecule functioned as a high-affinity, high-capacity uric acid transporter (Non-Patent Documents 2 and
35 3). It has also been reported that GLUT9 has two isoforms

(GLUT9S and GLUT9L) due to N-terminal intracellular portion splice, and both has the same uric acid transport activity (Non-Patent Documents 4 and 5). Moreover, it has become evident that depressed function of this molecule causes severe hypouricemia from the analysis of GLUT9 genetic variation family (Non-Patent Documents 6 and 7).

Uric acid transport kinetics in the kidney have been studied since early times. Thus, uric acid first passes through glomerular, and then undergoes two-way transport via transporter which is either reabsorption or secretion, and eventually about 90% of the amount of the uric acid which has passed through glomerular is reabsorbed. GLUT9 is, from the information mentioned above, considered to play the main role in the uric acid reabsorption in the kidney, and thus an important molecule that controls blood uric acid levels. Therefore, a GLUT9 inhibitor is expected to reduce blood uric acid levels and be effective for hyperuricemia and pathological conditions associated therewith.

[0004]

In Japan, hyperuricemia is defined as a condition wherein serum uric acid level exceeds 7.0 mg/dL, based on the concentration of uric acid dissolved in blood (Non-Patent Document 8). Persistent hyperuricemia causes gouty arthritis or kidney damage (gouty kidney) resulted from deposition of urate crystals in tissues. A prolonged disease period in gouty arthritis cases results in granuloma formation such as gouty tophus which is primarily due to the urate.

[0005]

Further, in the recent years, hyperuricemia has been recognized as a lifestyle disease, and there has been many reports suggesting that hyperuricemia is associated with various pathological conditions. Remedy of hyperuricemia can be a potential treatment and prophylaxis for those pathological condition. Pathological conditions listed below are generally known to be associated with hyperuricemia and are particularly

suggested to have a relationship with high uric acid.

[0006]

1) Chronic kidney disease (CKD)

Many epidemiological studies have shown that
5 hyperuricemia is a risk factor for development of terminal
kidney failure or CKD onset (Non-Patent Documents 9, 10 and 11),
and there is a report of intervention trial involving use of
uric acid lowering agents which observed a renoprotective
effect (Non-Patent Document 12). It is also reported that gene
10 polymorphism of GLUT9 is responsible for CKD onset (Non-Patent
Document 13).

[0007]

2) Hypertension

Many clinical studies have gradually convinced that
15 hyperuricemia is closely associated with the onset of
hypertension (Non-Patent Documents 14 and 15). Also, there are
clinical results reporting that blood pressure has been
decreased by treating hyperuricemia (Non-Patent Document 16).

[0008]

20 3) Diabetes

In a meta-analysis which puts together multiple
prospective clinical studies, hyperuricemia is reported as an
independent risk factor for type 2 diabetes (Non-Patent
Document 17). Also studies involving use of mice and cultured
25 cells show that high concentration uric acid suppresses insulin
secretion and induces insulin resistance (Non-Patent Documents
18 and 19).

[0009]

4) Cardiac disease (cardiovascular disease, cardiac failure, 30 atrial fibrillation)

The blood uric acid level has been reported as an
independent risk factor for cardiovascular events (Non-Patent
Document 20). Besides the events, a study of correlations
between the characteristics of coronary artery and uric acid
35 using intravascular ultrasound (IVUS) shows that hyperuricemia

is associated with plaque volume and calcified lesion (Non-Patent Document 21). Hyperuricemia is also observed in many patients with chronic cardiac failure. A Japanese epidemiological study conducted under the Japanese Cardiac Registry of Heart Failure in Cardiology (J-CARE-CARD) test demonstrated that cardiac failure patients with hyperuricemia had a significantly higher rate of all-cause death and cardiac death (Non-Patent Document 22). In the recent years, a complication of hyperuricemia and atrial fibrillation has been attracting attention. It has been reported that prevalence of atrial fibrillation increases according to the serum uric acid level, and prevalence of hyperuricemia with 8 mg/dL or greater is significantly higher than those of 6.9 mg/dL or less (Non-Patent Document 23).

[0010]

5) Arteriosclerotic disease

Frequency of hyperuricemia in patients with hypertriglyceridemia is as high as about 30%, which is reported to be closely related with hyperuricemia and hyperlipemia (Non-Patent Documents 24 and 25).

[0011]

6) Non-alcoholic fatty liver disease (NAFLD) and non-alcoholic steatohepatitis (NASH)

NAFLD is associated with fatty liver and is a chronic hepatic disease of unknown cause which is commonly diagnosed in people who do not drink alcohol. Further, a pathological condition with a more progressed inflammation and fibrosis is called "NASH" which may cause hepatic cirrhosis and hepatoma. Many NAFLD patients have a complication with hyperuricemia, where the serum uric acid level is an independent risk factor for NAFLD. There is a meta-analysis result reporting that 1 mg/dL increase in the serum uric acid level increases the risk of NAFLD onset by 21% (Non-Patent Document 26).

[0012]

7) Psoriasis

It has been long known that psoriasis patients generally have a higher uric acid level relative to healthy subjects, as seen in the report that hyperuricemia was observed in about half of psoriasis patients (Non-Patent Document 27). In addition, as it has been reported that risks of cardiovascular disease and kidney damage are significantly higher in psoriasis patients relative to healthy subjects, hyperuricemia may be a factor that plays a role in increasing these risks (Non-Patent Documents 28 and 29).

10 [0013]

As described above, a GLUT9 inhibitor is considered to be an agent for the treatment or prophylaxis of pathological conditions that involve high blood uric acid levels; specifically, hyperuricemia, gout (for example, gouty arthritis, gouty kidney, and gouty tophus) and the like. Further, it is considered to have potential to be useful as an agent for the treatment or prophylaxis of pathological conditions which are generally known to have a complication with hyperuricemia and are particularly suggested to have association with high uric acid; specifically, chronic kidney disease (CKD), hypertension, diabetes, cardiac disease (for example, cardiovascular disease, cardiac failure, and atrial fibrillation), arteriosclerotic disease, non-alcoholic fatty liver disease (NAFLD), non-alcoholic steatohepatitis (NASH), psoriasis and the like.

25 **Document List**

Non-Patent Document

[0014]

[Non-Patent Document 1] Genomics. 2000 Jun 1; 66(2):217-20.

[Non-Patent Document 2] PLoS Genet. 2007 Nov; 3(11):e194.

30 [Non-Patent Document 3] Nat Genet. 2008 Apr; 40(4):437-42.

[Non-Patent Document 4] J Biol Chem. 2008 Oct 3; 283(40):26834-8.

[Non-Patent Document 5] ADMET & DMPK. 2017; 5(2):59-74.

[Non-Patent Document 6] Am J Hum Genet. 2008 Dec; 83(6):744-51.

35 [Non-Patent Document 7] Nephrol Dial Transplant. 2012 Mar;

27(3):1035-41.

[Non-Patent Document 8] Japanese guideline for the management of hyperuricemia and gout; second edition, edited by the Guideline Revision Committee, Japanese Society of Gout and

5 Nucleic Acid Metabolism, Osaka, Medical Review Co., Ltd., 2010

[Non-Patent Document 9] Am J Kidney Dis. 2004 Oct; 44(4):642-50.

[Non-Patent Document 10] J Am Soc Nephrol. 2008 Dec; 19(12):2407-13.

[Non-Patent Document 11] PLoS One. 2014 Jun 24; 9(6):e100801.

10 [Non-Patent Document 12] Am J Kidney Dis. 2015 Dec; 66(6):945-50.

[Non-Patent Document 13] Clin J Am Soc Nephrol. 2014 Jun 6; 9(6):1059-65.

[Non-Patent Document 14] Hypertension. 2006 Dec; 48(6):1031-6.

15 [Non-Patent Document 15] Ann Rheum Dis. 2013 Aug; 72(8):1321-7.

[Non-Patent Document 16] JAMA. 2008 Aug 27; 300(8):924-32.

[Non-Patent Document 17] PLoS One. 2013; 8(2):e56864.

[Non-Patent Document 18] Mol Cell Endocrinol. 2013 Aug 15; 375(1-2):89-96.

20 [Non-Patent Document 19] Biochem Biophys Res Commun. 2014 May 16; 447(4):707-14.

[Non-Patent Document 20] National Health and Nutrition Examination Survey. JAMA. 2000 May 10; 283(18):2404-10.

[Non-Patent Document 21] Coron Artery Dis. 2014 Jun; 25(4):343-
25 8.

[Non-Patent Document 22] Int J Cardiol. 2011 Sep 1; 151(2):143-7.

[Non-Patent Document 23] Hypertens Res. 2014 Aug; 37(8):785-9.

[Non-Patent Document 24] Br J Rheumatol. 1994 Aug; 33(8):731-4.

30 [Non-Patent Document 25] Ther Res 33:1397-1405, 2012

[Non-Patent Document 26] J Clin Endocrinol Metab. 2015 Nov; 100(11):4198-207.

[Non-Patent Document 27] Am J Dermatopathol. 1981 Winter; 3(4):397-404.

35 [Non-Patent Document 28] Arch Dermatol. 2008 Nov; 144(11):1518-

9.

[Non-Patent Document 29] BMJ. 2013 Oct 15; 347:f5961.

Summary of the Invention

[0015]


5 The present invention provides a nitrogen-containing heteroaryl compound having a GLUT9 inhibitory activity, or a pharmaceutically acceptable salt thereof, a pharmaceutical composition containing the same, a pharmaceutical use thereof, and the like. Accordingly, the present invention encompasses
10 the embodiments exemplified below.

[0016]

[Item 1]

A compound of Formula [I], or a pharmaceutically acceptable salt thereof:

15 [0017]



[0018]

wherein

=X- is =C(R⁵)- or =N-;

20 -L-COOH is

- (1) -COOH,
- (2) -C(R⁷¹)(R⁷²)-COOH,
- (3) -C(R⁷³)(R⁷⁴)-C(R⁷⁵)(R⁷⁶)-COOH, or
- (4) -O-C(R⁷⁷)(R⁷⁸)-COOH;

25 n is 0, 1, or 2;

m is 0, 1, 2, or 3;

R¹ is each independently halogen or C₁₋₃ alkyl;

R² is

- (1) halogen,
- (2) hydroxy,
- (3) cyano,
- 5 (4) C₁₋₆ alkyl optionally substituted with 1 to 3 substituents independently selected from the group consisting of cyano and C₁₋₃ alkoxy,
- (5) halo C₁₋₆ alkyl,
- (6) C₁₋₆ alkoxy,
- 10 (7) halo C₁₋₆ alkoxy,
- (8) -COOR²¹ wherein R²¹ is hydrogen or C₁₋₃ alkyl,
- (9) -CON(R²²)(R²³) wherein R²² and R²³ are each independently hydrogen or C₁₋₃ alkyl,
- (10) C₃₋₆ cycloalkyl or
- 15 (11) a 4- to 6-membered saturated heterocyclic group containing 1 or 2 hetero atom as a ring atom in addition to the carbon atoms, wherein the hetero atom is independently selected from the group consisting of oxygen, nitrogen and sulfur atoms, and

R³ is

- 20 (1) hydrogen,
- (2) halogen,
- (3) cyano,
- (4) C₁₋₃ alkyl,
- (5) halo C₁₋₃ alkyl,
- 25 (6) C₁₋₃ alkoxy, or
- (7) -COOR³¹ wherein R³¹ is hydrogen or C₁₋₃ alkyl or

R² and R³, together with the carbon atoms that they are bonded to, form a 4- to 6-membered saturated heterocycle containing 1 or 2 hetero atom as a ring atom in addition to the
30 carbon atoms, wherein the hetero atom is independently selected from the group consisting of oxygen, nitrogen and sulfur atoms;

R⁴ is


- (1) C₁₋₈ alkyl optionally substituted with 1 to 3 substituents independently selected from the following Group A,
- 35 (2) halo C₁₋₆ alkyl,

(3) $-\text{CON}(\text{R}^{41})(\text{R}^{42})$ wherein R^{41} and R^{42} are each independently hydrogen or C_{1-6} alkyl,

(4) C_{3-6} cycloalkyl optionally substituted with 1 to 3 substituents independently selected from C_{1-3} alkoxy, or

5 (5) a 4- to 6-membered saturated heterocyclic group containing 1 or 2 hetero atom as a ring atom in addition to the carbon atoms, wherein the hetero atom is independently selected from the group consisting of oxygen, nitrogen and sulfur atoms, and wherein the ring atom in the heterocyclic group bonded to

10 [0019]



is a carbon atom,

[0020]

and

Group A consists of

15 (a) hydroxy,

(b) C_{1-3} alkoxy optionally substituted with one hydroxy or one C_{1-3} alkoxy,

(c) halo C_{1-3} alkoxy,

(d) C_{3-6} cycloalkyl optionally substituted with one hydroxy, and

20 (e) phenyl, and

R^5 is hydrogen, halogen or C_{1-3} alkyl or

R^4 and R^5 , together with the carbon atoms that they are bonded to, form C_{3-6} cycloalkane;

R^6 are each independently halogen, hydroxy, C_{1-3} alkyl or
25 C_{1-3} alkoxy; and

R^{71} , R^{72} , R^{73} , R^{74} , R^{75} , R^{76} , R^{77} , and R^{78} are each independently hydrogen or C_{1-3} alkyl.

[0021]

[Item 2]

30 The compound according to Item 1 or a pharmaceutically acceptable salt thereof, wherein $=\text{X}-$ is $=\text{C}(\text{R}^5)-$.

[0022]

[Item 3]

The compound according to Item 1 or a pharmaceutically acceptable salt thereof, wherein =X- is =N-.

[0023]

5 [Item 4]

The compound according to any one of Items 1 to 3 or a pharmaceutically acceptable salt thereof, wherein -L-COOH is -COOH.

[0024]

10 [Item 5]

The compound according to any one of Items 1 to 4 or a pharmaceutically acceptable salt thereof, wherein n is 0 or 1.

[0025]

[Item 6]

15 The compound according to any one of Items 1 to 4 or a pharmaceutically acceptable salt thereof, wherein n is 0.

[0026]

[Item 7]

The compound according to any one of Items 1 to 4 or a
20 pharmaceutically acceptable salt thereof, wherein n is 1.

[0027]

[Item 8]

The compound according to any one of Items 1 to 7 or a
pharmaceutically acceptable salt thereof, wherein m is 0 or 1.

25 [0028]

[Item 9]

The compound according to any one of Items 1 to 7 or a pharmaceutically acceptable salt thereof, wherein m is 0.

[0029]

30 [Item 10]

The compound according to any one of Items 1 to 7 or a pharmaceutically acceptable salt thereof, wherein m is 1.

[0030]

[Item 11]

35 The compound according to any one of Items 1 to 10 or a

pharmaceutically acceptable salt thereof, wherein R¹ is halogen.

[0031]

[Item 12]

The compound according to any one of Items 1 to 10 or a
5 pharmaceutically acceptable salt thereof, wherein R¹ is C₁₋₃
alkyl.

[0032]

[Item 13]

The compound according to any one of Items 1 to 12 or a
10 pharmaceutically acceptable salt thereof, wherein R³ is

(1) hydrogen, or

(2) halogen.

[0033]

[Item 14]

15 The compound according to any one of Items 1 to 12 or a
pharmaceutically acceptable salt thereof, wherein R³ is
hydrogen.

[0034]

[Item 15]

20 The compound according to any one of Items 1 to 12 or a
pharmaceutically acceptable salt thereof, wherein R³ is halogen.

[0035]

[Item 16]

The compound according to any one of Items 1 to 15 or a
25 pharmaceutically acceptable salt thereof, wherein
R⁴ is

(1) C₁₋₈ alkyl optionally substituted with 1 to 3 substituents
independently selected from the following Group A,

(2) halo C₁₋₆ alkyl, or

30 (3) -CON(R⁴¹)(R⁴²) wherein R⁴¹ and R⁴² are each independently
hydrogen or C₁₋₆ alkyl, and

Group A consists of

(a) hydroxy,

(b) C₁₋₃ alkoxy optionally substituted with one hydroxy or one

35 C₁₋₃ alkoxy,

- (c) halo C₁₋₃ alkoxy,
- (d) C₃₋₆ cycloalkyl optionally substituted with one hydroxy, and
- (e) phenyl.

[0036]

5 [Item 17]

The compound according to any one of Items 1 to 15 or a pharmaceutically acceptable salt thereof, wherein R⁴ is C₁₋₈ alkyl optionally substituted with 1 to 3 substituents independently selected from the following Group A, and

10 Group A consists of

- (a) hydroxy,
- (b) C₁₋₃ alkoxy optionally substituted with one hydroxy or one C₁₋₃ alkoxy,
- (c) halo C₁₋₃ alkoxy,
- 15 (d) C₃₋₆ cycloalkyl, and
- (e) phenyl.

[0037]

[Item 18]


The compound according to any one of Items 1 to 15 or a
20 pharmaceutically acceptable salt thereof, wherein R⁴ is - CON(R⁴¹)(R⁴²) wherein R⁴¹ and R⁴² are each independently hydrogen or C₁₋₆ alkyl.

[0038]

[Item 19]

25 A compound of the following formula or a pharmaceutically acceptable salt thereof:

[0039]




(Example 21).

[0040]

[Item 20]

A compound of the following formula or a pharmaceutically
5 acceptable salt thereof:

[0041]




(Example 35).

[0042]

[Item 21]

10 A compound of the following formula or a pharmaceutically
acceptable salt thereof:

[0043]




(Example 67).

[0044]

[Item 22]

A compound of the following formula or a pharmaceutically acceptable salt thereof:

5 [0045]




(free form of Example 76).

[0046]

[Item 23]

A compound of the following formula or a pharmaceutically acceptable salt thereof:

10 [0047]




(Example 84).

[0048]

[Item 24]

A compound of the following formula or a pharmaceutically acceptable salt thereof:

15 [0049]




(Example 98).

[0050]

[Item 25]

A compound of the following formula or a pharmaceutically
5 acceptable salt thereof:

[0051]




(Example 107).

[0052]

[Item 26]

10 A compound of the following formula or a pharmaceutically
acceptable salt thereof:

[0053]




(Example 109).

[0054]

[Item 27]

A compound of the following formula or a pharmaceutically acceptable salt thereof:

5 [0055]




(Example 116).

[0056]

[Item 28]

A compound of the following formula or a pharmaceutically acceptable salt thereof:

10 [0057]



(Example 118).


[0058]

[Item 29]

A compound of the following formula or a pharmaceutically acceptable salt thereof:

15

[0059]



(Example 126).

[0060]

[Item 30]

A pharmaceutical composition comprising the compound
 5 according to any one of Items 1 to 29 or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

[0061]

[Item 31]

10 A GLUT9 inhibitor comprising the compound according to any one of Items 1 to 29 or a pharmaceutically acceptable salt thereof.

[0062]

[Item 32]

15 An agent for the treatment or prophylaxis of a disease selected from the group consisting of hyperuricemia and gout, which comprises the compound according to any one of Items 1 to 29 or a pharmaceutically acceptable salt thereof.

[0063]

20 [Item 33]

A method for the inhibition of GLUT9 in a mammal in need of such inhibition, which comprises administering a pharmaceutically effective amount of the compound according to any one of Items 1 to 29 or a pharmaceutically acceptable salt
 25 thereof to the mammal.

[0064]

[Item 34]

A method for the treatment or prophylaxis of a disease

selected from the group consisting of hyperuricemia and gout in a mammal in need of such treatment or prophylaxis, which comprises administering a pharmaceutically effective amount of the compound according to any one of Items 1 to 29 or a
5 pharmaceutically acceptable salt thereof to the mammal.

[0065]

[Item 35]

Use of the compound according to any one of Items 1 to 29 or a pharmaceutically acceptable salt thereof for the
10 manufacture of a GLUT9 inhibitor.

[0066]

[Item 36]

Use of the compound according to any one of Items 1 to 29 or a pharmaceutically acceptable salt thereof for the
15 manufacture of an agent for the treatment or prophylaxis of a disease selected from the group consisting of hyperuricemia and gout.

[0067]

[Item 37]

20 The compound according to any one of Items 1 to 29 or a pharmaceutically acceptable salt thereof, for use in the inhibition of GLUT9.

[0068]

[Item 38]

25 The compound according to any one of Items 1 to 29 or a pharmaceutically acceptable salt thereof, for use in the treatment or prophylaxis of a disease selected from the group consisting of hyperuricemia and gout.

[0069]

30 [Item 39]

A commercial package comprising the composition according to Item 30 and a written matter associated therewith, the written matter stating that the composition can or should be used for the treatment or prophylaxis of a disease selected
35 from the group consisting of hyperuricemia and gout.

[0070]

[Item 40]

A kit comprising the composition according to Item 30 and a written matter associated therewith, the written matter
5 stating that the composition can or should be used for the treatment or prophylaxis of a disease selected from the group consisting of hyperuricemia and gout.

Embodiments of the Invention

[0071]

10 The definitions of the terms used herein are as follows.

[0072]

The following wavy line:

[0073]



15 [0074]

in the partial structure means an abbreviation of a bonding partner.

[0075]

Examples of the "halogen" include fluorine, chlorine,
20 bromine and iodine.

[0076]

The "C₁₋₃ alkyl" means a straight- or branched-chain saturated hydrocarbon group having 1 to 3 carbon atoms. Examples of the "C₁₋₃ alkyl" include methyl, ethyl, n-propyl,
25 and isopropyl.

[0077]

The "C₁₋₆ alkyl" means a straight- or branched-chain saturated hydrocarbon group having 1 to 6 carbon atoms. Examples of the "C₁₋₆ alkyl" include methyl, ethyl, n-propyl,
30 isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, n-pentyl, isopentyl and n-hexyl.

[0078]

The "C₁₋₈ alkyl" means a straight- or branched-chain

saturated hydrocarbon group having 1 to 8 carbon atoms.

Examples of the "C₁₋₈ alkyl" include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, n-pentyl, isopentyl, n-hexyl, n-heptyl and n-octyl.

5 [0079]

The "halo C₁₋₃ alkyl" means the above-mentioned "C₁₋₃ alkyl" substituted with 1 to 5 halogen independently selected from the group consisting of the above-mentioned "halogen".

Examples of the "halo C₁₋₃ alkyl" include monofluoromethyl,
10 difluoromethyl, trifluoromethyl, 2-fluoroethyl, 2-chloroethyl, 2-bromoethyl, 1,1-difluoroethyl, 2,2,2-trifluoroethyl, pentafluoroethyl, 3-fluoropropyl, 3-chloropropyl, 1,1-difluoropropyl and 3,3,3-trifluoropropyl.

[0080]

15 The "halo C₁₋₆ alkyl" means the above-mentioned "C₁₋₆ alkyl" substituted with 1 to 5 halogen independently selected from the group consisting of the above-mentioned "halogen". Examples of the "halo C₁₋₆ alkyl" include monofluoromethyl, difluoromethyl, trifluoromethyl, 2-fluoroethyl, 2-chloroethyl,
20 2-bromoethyl, 1,1-difluoroethyl, 2,2,2-trifluoroethyl, pentafluoroethyl, 3-fluoropropyl, 3-chloropropyl, 1,1-difluoropropyl, 3,3,3-trifluoropropyl, 4,4,4-trifluorobutyl, 5,5,5-trifluoropentyl and 6,6,6-trifluorohexyl.

[0081]

25 The "C₁₋₃ alkoxy" means a group wherein the above-mentioned "C₁₋₃ alkyl" is bonded to an oxygen atom. Examples of the "C₁₋₃ alkoxy" include methoxy, ethoxy, n-propoxy and isopropoxy.

[0082]

30 The "C₁₋₆ alkoxy" means a group wherein the above-mentioned "C₁₋₆ alkyl" is bonded to an oxygen atom. Examples of the "C₁₋₆ alkoxy" include methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, isobutoxy, sec-butoxy, tert-butoxy, n-pentyloxy, isopentyloxy and n-hexyloxy.

35 [0083]

The "halo C₁₋₃ alkoxy" means the above-mentioned "C₁₋₃ alkoxy" substituted with 1 to 5 halogen independently selected from the group consisting of the above-mentioned "halogen". Examples of the "halo C₁₋₃ alkoxy" include monofluoromethoxy, difluoromethoxy, trifluoromethoxy, 2-fluoroethoxy, 2-chloroethoxy, 2-bromoethoxy, 1,1-difluoroethoxy, 2,2,2-trifluoroethoxy, pentafluoroethoxy, 3-fluoropropoxy, 3-chloropropoxy, 1,1-difluoropropoxy and 3,3,3-trifluoropropoxy. [0084]

10 The "halo C₁₋₆ alkoxy" means the above-mentioned "C₁₋₆ alkoxy" substituted with 1 to 5 halogen independently selected from the group consisting of the above-mentioned "halogen". Examples of the "halo C₁₋₆ alkoxy" include monofluoromethoxy, difluoromethoxy, trifluoromethoxy, 2-fluoroethoxy, 2-chloroethoxy, 2-bromoethoxy, 1,1-difluoroethoxy, 2,2,2-trifluoroethoxy, pentafluoroethoxy, 3-fluoropropoxy, 3-chloropropoxy, 1,1-difluoropropoxy, 3,3,3-trifluoropropoxy, 4,4,4-trifluorobutoxy, 5,5,5-trifluoropentyloxy and 6,6,6-trifluorohexyloxy.


20 [0085]

The "C₃₋₆ cycloalkyl" means a monocyclic saturated hydrocarbon group having 3 to 6 carbon atoms. Examples of the "C₃₋₆ cycloalkyl" include cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl.

25 [0086]

The "C₃₋₆ cycloalkane" means a monocyclic saturated hydrocarbon having 3 to 6 carbon atoms. Examples of the "C₃₋₆ cycloalkane" include cyclopropane, cyclobutane, cyclopentane and cyclohexane. Examples of the "C₃₋₆ cycloalkane" formed by R⁴ and R⁵, together with the carbon atoms that they are bonded to, include the following rings:

[0087]




[0088]

Examples of the "4- to 6-membered saturated heterocyclic group containing 1 or 2 hetero atom as a ring atom in addition to the carbon atoms, wherein the hetero atom is independently selected from the group consisting of oxygen, nitrogen and sulfur atoms" include oxetanyl, tetrahydrofuranyl, tetrahydropyranyl, azetidiny, pyrrolidinyl and piperidyl.

[0089]

For example, examples of the "4- to 6-membered saturated heterocyclic group containing 1 or 2 hetero atom as a ring atom in addition to the carbon atoms, wherein the hetero atom is independently selected from the group consisting of oxygen, nitrogen and sulfur atoms, and wherein the ring atom in the heterocyclic group bonded to

[0090]




is a carbon atom"

[0091]

in (5) of R⁴ include the following groups:

[0092]



[0093]


Examples of the "4- to 6-membered saturated heterocycle containing 1 or 2 hetero atom as a ring atom in addition to the carbon atoms, wherein the hetero atom is independently selected from the group consisting of oxygen, nitrogen and sulfur atoms" include oxetane, tetrahydrofuran, tetrahydropyran, 1,4-dioxane,

pyrrolidine and piperidine.

[0094]

For example, examples of the "4- to 6-membered saturated heterocycle containing 1 or 2 hetero atom as a ring atom in addition to the carbon atoms, wherein the hetero atom is independently selected from the group consisting of oxygen, nitrogen and sulfur atoms" formed by R^2 and R^3 , together with the carbon atoms that they are bonded to, include the following rings:

10 [0095]



[0096]

With regard to the term "substituted", for example, C_{1-6} alkyl "optionally substituted with 1 to 3 substituents independently selected from cyano and C_{1-3} alkoxy" in (4) of R^2 means unsubstituted C_{1-6} alkyl, or C_{1-6} alkyl substituted with 1 to 3 substituents independently selected from cyano and C_{1-3} alkoxy at any substitutable position.

[0097]

20 The "compound of Formula [I]" is hereinafter also referred to as "Compound [I]".

[0098]

Specific embodiments of each group of Compound [I] are exemplified below, which should not be construed as limitative. Compound [I] also encompasses combinations of two or more embodiments selected appropriately from the specific embodiments of each group.

[0099]

-L-COOH is preferably

- (1) -COOH,
- (2) -C(R⁷¹)(R⁷²)-COOH, or
- (3) -C(R⁷³)(R⁷⁴)-C(R⁷⁵)(R⁷⁶)-COOH.

5 -L-COOH is more preferably -COOH.

[0100]

n is preferably 0 or 1.

[0101]

m is preferably 0, 1 or 2.

10 m is more preferably 0 or 1.

[0102]

R² is preferably

- (1) halogen,
- (2) hydroxy,
- 15 (3) cyano,
- (4) C₁₋₆ alkyl optionally substituted with 1 to 3 substituents independently selected from the group consisting of cyano and C₁₋₃ alkoxy,
- (5) halo C₁₋₆ alkyl,
- 20 (6) C₁₋₆ alkoxy,
- (7) halo C₁₋₆ alkoxy,
- (8) -COOR²¹ wherein R²¹ is hydrogen or C₁₋₃ alkyl, or
- (9) -CON(R²²)(R²³) wherein R²² and R²³ are each independently hydrogen or C₁₋₃ alkyl.

25 R² is more preferably

- (2) hydroxy,
- (6) C₁₋₆ alkoxy, or
- (7) halo C₁₋₆ alkoxy.

[0103]

30 R³ is preferably

- (1) hydrogen,
- (2) halogen,
- (3) cyano,
- (4) C₁₋₃ alkyl,
- 35 (5) halo C₁₋₃ alkyl,

- (6) C₁₋₃ alkoxy, or
(7) -COOR³¹ wherein R³¹ is hydrogen or C₁₋₃ alkyl.

R³ is more preferably

- (1) hydrogen, or

- 5 (2) halogen.

[0104]

R⁴ is preferably

(1) C₁₋₈ alkyl optionally substituted with 1 to 3 substituents independently selected from the following Group A,

- 10 (2) halo C₁₋₆ alkyl, or

(3) -CON(R⁴¹)(R⁴²) wherein R⁴¹ and R⁴² are each independently hydrogen or C₁₋₆ alkyl.

Group A consists of

- (a) hydroxy,

- 15 (b) C₁₋₃ alkoxy optionally substituted with one hydroxy or one C₁₋₃ alkoxy,

- (c) halo C₁₋₃ alkoxy,

- (d) C₃₋₆ cycloalkyl, and

- (e) phenyl.

- 20 Group A preferably consists of

- (a) hydroxy,

(b) C₁₋₃ alkoxy optionally substituted with one hydroxy or one C₁₋₃ alkoxy, and

- (c) halo C₁₋₃ alkoxy.

- 25 [0105]

R⁵ is preferably hydrogen.

[0106]

A preferable embodiment is Compound [I] wherein

=X- is =C(R⁵)- or =N-;

- 30 -L-COOH is

- (1) -COOH,

- (2) -C(R⁷¹)(R⁷²)-COOH, or

- (3) -C(R⁷³)(R⁷⁴)-C(R⁷⁵)(R⁷⁶)-COOH;

n is 0 or 1;

- 35 m is 0, 1 or 2;

R¹ is halogen or C₁₋₃ alkyl;

R² is

- (1) halogen,
- (2) hydroxy,
- 5 (3) cyano,
- (4) C₁₋₆ alkyl optionally substituted with 1 to 3 substituents independently selected from the group consisting of cyano and C₁₋₃ alkoxy,
- (5) halo C₁₋₆ alkyl,
- 10 (6) C₁₋₆ alkoxy,
- (7) halo C₁₋₆ alkoxy,
- (8) -COOR²¹ wherein R²¹ is hydrogen or C₁₋₃ alkyl, or
- (9) -CON(R²²)(R²³) wherein R²² and R²³ are each independently hydrogen or C₁₋₃ alkyl;

15 R³ is

- (1) hydrogen,
- (2) halogen,
- (3) cyano,
- (4) C₁₋₃ alkyl,
- 20 (5) halo C₁₋₃ alkyl,
- (6) C₁₋₃ alkoxy, or
- (7) -COOR³¹ wherein R³¹ is hydrogen or C₁₋₃ alkyl;

R⁴ is

- (1) C₁₋₈ alkyl optionally substituted with 1 to 3 substituents
- 25 independently selected from the following Group A,
- (2) halo C₁₋₆ alkyl, or
- (3) -CON(R⁴¹)(R⁴²) wherein R⁴¹ and R⁴² are each independently hydrogen or C₁₋₆ alkyl, and

Group A consists of

- 30 (a) hydroxy,
- (b) C₁₋₃ alkoxy optionally substituted with one hydroxy or one C₁₋₃ alkoxy,
- (c) halo C₁₋₃ alkoxy,
- (d) C₃₋₆ cycloalkyl, and
- 35 (e) phenyl;

R⁵ is hydrogen;

R⁶ are each independently halogen, hydroxy, C₁₋₃ alkyl or C₁₋₃ alkoxy; and

R⁷¹, R⁷², R⁷³, R⁷⁴, R⁷⁵, R⁷⁶, R⁷⁷, and R⁷⁸ are each
5 independently hydrogen or C₁₋₃ alkyl.

[0107]

A more preferable embodiment is Compound [I] wherein

=X- is =C(R⁵)- or =N-;

-L-COOH is -COOH;

10 n is 0 or 1;

m is 0 or 1;

R¹ is halogen or C₁₋₃ alkyl;

R² is

(2) hydroxy,

15 (6) C₁₋₆ alkoxy, or

(7) halo C₁₋₆ alkoxy;

R³ is

(1) hydrogen, or

(2) halogen;

20 R⁴ is

(1) C₁₋₈ alkyl optionally substituted with 1 to 3 substituents
independently selected from the following Group A,

(2) halo C₁₋₆ alkyl, or

(3) -CON(R⁴¹)(R⁴²) wherein R⁴¹ and R⁴² are each independently
25 hydrogen or C₁₋₆ alkyl, and

Group A consists of

(a) hydroxy,

(b) C₁₋₃ alkoxy optionally substituted with one hydroxy or one
C₁₋₃ alkoxy,

30 (c) halo C₁₋₃ alkoxy,

(d) C₃₋₆ cycloalkyl, and


(e) phenyl;

R⁵ is hydrogen ; and

R⁶ is halogen, hydroxy, C₁₋₃ alkyl or C₁₋₃ alkoxy.

35 [0108]

Another preferable embodiment is a compound of Formula [II], or a pharmaceutically acceptable salt thereof:
[0109]



5 [0110]

wherein

n is 0 or 1;

m is 0 or 1;

R¹ is halogen or C₁₋₃ alkyl;

10 R² is hydroxy, C₁₋₆ alkoxy or halo C₁₋₆ alkoxy;

R³ is hydrogen or halogen;

R⁴ is

(1) C₁₋₈ alkyl optionally substituted with 1 to 3 substituents independently selected from the following Group A,

15 (2) halo C₁₋₆ alkyl, or

(3) -CON(R⁴¹)(R⁴²) wherein R⁴¹ and R⁴² are each independently hydrogen or C₁₋₆ alkyl, and

Group A consists of

(a) hydroxy,

20 (b) C₁₋₃ alkoxy optionally substituted with one hydroxy or one C₁₋₃ alkoxy,

(c) halo C₁₋₃ alkoxy,

(d) C₃₋₆ cycloalkyl, and

(e) phenyl; and


25 R⁶ is halogen, hydroxy, C₁₋₃ alkyl or C₁₋₃ alkoxy.

[0111]

Another preferable embodiment is a compound of Formula

[III], or a pharmaceutically acceptable salt thereof:

[0112]



[III]

[0113]

5 wherein

n is 0 or 1;

m is 0 or 1;

R¹ is halogen or C₁₋₃ alkyl;

R² is hydroxy, C₁₋₆ alkoxy or halo C₁₋₆ alkoxy;

10 R³ is hydrogen or halogen;

R⁴ is

(1) C₁₋₈ alkyl optionally substituted with 1 to 3 substituents independently selected from the following Group A,

(2) halo C₁₋₆ alkyl, or

15 (3) -CON(R⁴¹)(R⁴²) wherein R⁴¹ and R⁴² are each independently hydrogen or C₁₋₆ alkyl, and

Group A consists of

(a) hydroxy,

(b) C₁₋₃ alkoxy optionally substituted with one hydroxy or one

20 C₁₋₃ alkoxy,

(c) halo C₁₋₃ alkoxy,

(d) C₃₋₆ cycloalkyl, and

(e) phenyl; and

R⁶ is halogen, hydroxy, C₁₋₃ alkyl or C₁₋₃ alkoxy.

25 [0114]

The "pharmaceutically acceptable salt" may be any salt known in the art as long as it is not associated with undue

toxicity. Specific examples thereof include salts with inorganic acid, salts with organic acid, salts with inorganic base, and salts with organic base. Various forms of pharmaceutically acceptable salts are well known in the art and, 5 for example, they are described in the following documents.

- (a) Berge et al., J. Pharm. Sci., 66, p 1-19 (1977),
- (b) Stahl et al., "Handbook of Pharmaceutical Salt: Properties, Selection, and Use" (Wiley-VCH, Weinheim, Germany, 2002),
- (c) Paulekuhn et al., J. Med. Chem., 50, p 6665-6672 (2007)

10 The pharmaceutically acceptable salt of Compound [I] can be obtained by reacting Compound [I] with an inorganic acid, an organic acid, an inorganic base or an organic base, according to a known method.
[0115]

15 Examples of the salt with inorganic acid include salts with hydrofluoric acid, hydrochloric acid, hydrobromic acid, hydriodic acid, nitric acid, phosphoric acid and sulfuric acid.

Examples of the salt with organic acid include salts with acetic acid, adipic acid, alginic acid, 4-aminosalicylic acid, 20 anhydromethylenecitric acid, benzoic acid, benzenesulfonic acid, camphoric acid, camphor-10-sulfonic acid, carbonic acid, citric acid, edetic acid, ethane-1,2-disulfonic acid, dodecylsulfuric acid, ethanesulfonic acid, fumaric acid, glucoheptonic acid, gluconic acid, glucuronic acid, glycollylarsanilic acid, 25 hydroxynaphthoic acid, 2-hydroxy-1-ethanesulfonic acid, lactic acid, lactobionic acid, malic acid, maleic acid, mandelic acid, methanesulfonic acid, methylsulfuric acid, methylnitric acid, methylenebis(salicylic acid), galactaric acid, naphthalene-2-sulfonic acid, 2-naphthoic acid, 1,5-naphthalenedisulfonic acid, 30 oleic acid, oxalic acid, pamoic acid, pantothenic acid, pectic acid, picric acid, propionic acid, polygalacturonic acid, salicylic acid, stearic acid, succinic acid, tannic acid, tartaric acid, teoclic acid, thiocyanic acid, trifluoroacetic acid, p-toluenesulfonic acid, undecanoic acid, aspartic acid 35 and glutamic acid.

[0116]

Examples of the salt with inorganic base include salts with lithium, sodium, potassium, magnesium, calcium, barium, aluminium, zinc, bismass and ammonium.

5 Examples of the salt with organic base include salts with arecoline, betaine, choline, clemizole, ethylene diamine, N-methylglucamine, N-benzylphenethylamine, tris(hydroxymethyl)methylamine, arginine and lysine.

[0117]

10 Preferable embodiments of the "pharmaceutically acceptable salt" are as follows.

Examples of the salt with inorganic acid include salts with hydrochloric acid, nitric acid, sulfuric acid, phosphoric acid and hydrobromic acid.

15 Examples of the salt with organic acid include salts with oxalic acid, maleic acid, citric acid, fumaric acid, lactic acid, malic acid, succinic acid, tartaric acid, acetic acid, trifluoroacetic acid, benzoic acid, glucuronic acid, oleic acid, pamoic acid, methanesulfonic acid, benzenesulfonic acid, p-
20 toluenesulfonic acid and 2-hydroxy-1-ethanesulfonic acid.

Examples of the salt with inorganic base include salts with sodium, potassium, calcium, magnesium and zinc.

Examples of the salt with organic base include salts with tris(hydroxymethyl)methylamine, N-methylglucamine and lysine.

25 [0118]

Compound [I] or a pharmaceutically acceptable salt thereof may be present as a solvate.

The "solvate" is Compound [I] or a pharmaceutically acceptable salt thereof which is coordinated with a solvent
30 molecule, and also encompasses hydrates. The solvate is preferably a pharmaceutically acceptable solvate, and examples thereof include a hydrate, an ethanolate and a dimethyl sulfoxidate of Compound [I] or a pharmaceutically acceptable salt thereof.

35 Specific examples include semihydrate, monohydrate,

dihydrate and monoethanolate of Compound [I], monohydrate of hydrochloride of Compound [I], and 2/3 ethanolate of dihydrochloride of Compound [I]. These solvates can be obtained according to a known method.

5 [0119]

Compound [I] or a pharmaceutically acceptable salt thereof may be present as a tautomer. In this case, Compound [I] or a pharmaceutically acceptable salt thereof can be a single tautomer or a mixture thereof.

10 [0120]

Compound [I] or a pharmaceutically acceptable salt thereof may have a carbon-carbon double bond. In this case, Compound [I] or a pharmaceutically acceptable salt thereof can be present as an E form, a Z form, or a mixture thereof.

15 [0121]

Compound [I] or a pharmaceutically acceptable salt thereof may contain a stereoisomer that should be recognized as a cis/trans isomer. In this case, Compound [I] or a pharmaceutically acceptable salt thereof can be present as a cis form, a trans form, or a mixture thereof.

[0122]

Compound [I] or a pharmaceutically acceptable salt thereof may contain one or more asymmetric carbons. In this case, Compound [I] or a pharmaceutically acceptable salt thereof may be present as a single enantiomer, a single diastereomer, a mixture of enantiomers or a mixture of diastereomers.

[0123]

Compound [I] or a pharmaceutically acceptable salt thereof may be present as an atropisomer. In this case, Compound [I] or a pharmaceutically acceptable salt thereof may be present as a single atropisomer or a mixture thereof.

[0124]

Compound [I] or a pharmaceutically acceptable salt thereof may simultaneously contain plural structural

characteristics derived from the above-mentioned isomers.
Moreover, Compound [I] or a pharmaceutically acceptable salt thereof may contain the above-mentioned isomers at any ratio.
[0125]

5 The formulae, chemical structures and compound names indicated herein without specifying the stereochemistry thereof encompass all the above-mentioned isomers that may be present unless a particular note to the stereochemistry is made herein.
[0126]

10 A diastereomeric mixture can be separated into each diastereomer by conventional methods such as chromatography and crystallization. Alternatively, each diastereomer can also be produced by using a stereochemically single starting material, or by a synthesis method employing a stereoselective reaction.
15 [0127]

An enantiomeric mixture can be separated into each single enantiomer by a method well known in the art.

For example, first, a diastereomeric mixture can be prepared by reacting an enantiomeric mixture with a
20 substantially pure enantiomer compound known as a chiral auxiliary. Next, the obtained diastereomeric mixture can be separated into a single diastereomer having high isomer ratio or a substantially pure single diastereomer by a conventional method such as fractional crystallization and chromatography.
25 Finally, the separated diastereomer can be converted to a desired enantiomer by removing the added chiral auxiliary by cleavage.

Moreover, an enantiomeric mixture can also be directly separated into each enantiomer by a chromatography method using
30 a chiral solid phase well known in the art. Alternatively, one of enantiomers can also be obtained by using a substantially pure optically active starting material or by employing stereoselective synthesis (asymmetric induction) of a prochiral intermediate using a chiral auxiliary and an asymmetric
35 catalyst.

[0128]

The absolute steric configuration can be determined by the X-ray crystal analysis of the crystalline product or intermediate. In this case, a crystalline product or
5 intermediate derivatized with a reagent having an asymmetric center with a known steric configuration may be used if necessary.

[0129]

Compound [I] or a pharmaceutically acceptable salt
10 thereof may be labeled with isotope (e.g., ^2H , ^3H , ^{14}C , and ^{35}S ,).

[0130]

Compound [I] or a pharmaceutically acceptable salt thereof is preferably substantially pure, more preferably has a
15 purity of 80% or more.

[0131]

As used herein, the pharmaceutical composition may be produced according to a method known per se in the art of pharmaceutical preparations, by mixing Compound [I] or a
20 pharmaceutically acceptable salt thereof with a suitable amount of at least one type of pharmaceutically acceptable carrier and the like as appropriate. The content of Compound [I] or a pharmaceutically acceptable salt thereof in the pharmaceutical composition varies depending on the dosage form, dose and the
25 like, and is, for example, 0.1 to 100 wt% of the whole composition.

[0132]

Examples of the dosage form of Compound [I] or a pharmaceutically acceptable salt thereof include oral
30 preparations such as tablet, capsule, granule, powder, troche, syrup, emulsion, and suspension, and parenteral preparations such as external preparation, suppository, injection, eye drop, nasal preparations, and pulmonary preparation.

[0133]

Examples of the "pharmaceutically acceptable carrier" include various organic or inorganic carrier substances conventionally used as preparation materials, and specifically include excipient, disintegrant, binder, glidant, and lubricant for solid preparations; solvent, solubilizing agent, suspending agent, isotonicity agent, buffering agent, and soothing agent for liquid preparations; and base, emulsifier, moistening agent, stabilizer, stabilizing agent, dispersant, plasticizer, pH adjuster, absorption enhancer, gelling agent, preservative, filler, solvent, solubilizing agent, and suspending agent for semi-solid preparations. Where necessary, additives such as preservative, antioxidant, colorant, and sweetening agent may be used.

[0134]

Examples of the "excipient" include lactose, sucrose, D-mannitol, D-sorbitol, corn starch, dextrin, microcrystalline cellulose, crystalline cellulose, carmellose, carmellose calcium, sodium carboxymethyl starch, low-substituted hydroxypropylcellulose and gum arabic.

[0135]

Examples of the "disintegrant" include carmellose, carmellose calcium, carmellose sodium, sodium carboxymethyl starch, croscarmellose sodium, crospovidone, low-substituted hydroxypropylcellulose, hydroxypropylmethylcellulose and crystalline cellulose.

[0136]

Examples of the "binder" include hydroxypropylcellulose, hydroxypropylmethylcellulose, povidone, crystalline cellulose, sucrose, dextrin, starch, gelatin, carmellose sodium and gum arabic.

[0137]

Examples of the "glidant" include light anhydrous silicic acid and magnesium stearate.

[0138]

Examples of the "lubricant" include magnesium stearate,

calcium stearate and talc.

[0139]

Examples of the "solvent" include purified water, ethanol, propylene glycol, macrogol, sesame oil, corn oil and olive oil.

5 [0140]

Examples of the "solubilizing agent" include propylene glycol, D-mannitol, benzyl benzoate, ethanol, triethanolamine, sodium carbonate and sodium citrate.

[0141]

10 Examples of the "suspending agent" include benzalkonium chloride, carmellose, hydroxypropylcellulose, propylene glycol, povidone, methylcellulose and glycerol monostearate.

[0142]

Examples of the "isotonic agent" include glucose, D-
15 sorbitol, sodium chloride and D-mannitol.

[0143]

Examples of the "buffering agent" include sodium hydrogenphosphate, sodium acetate, sodium carbonate and sodium citrate.

20 [0144]

Examples of the "soothing agent" include benzyl alcohol.

[0145]

Examples of the "base" include water, animal and vegetable oils (e.g., olive oil, corn oil, arachis oil, sesame
25 oil, and castor oil), lower alcohols (e.g., ethanol, propanol, propylene glycol, 1,3-butylene glycol, and phenol), higher fatty acids and esters thereof, wax, higher alcohols, polyalcohols, hydrocarbons (e.g., white vaseline, liquid paraffin, and paraffin), hydrophilic vaseline, purified lanolin,
30 absorptive ointment, hydrous lanolin, hydrophilic ointment, starch, pullulan, gum arabic, tragacanth gum, gelatin, dextran, cellulose derivatives (e.g., methyl cellulose, carboxymethyl cellulose, hydroxyethyl cellulose, and hydroxypropyl cellulose), synthetic polymers (e.g., carboxyvinyl polymer, sodium
35 polyacrylate, polyvinyl alcohol, and polyvinyl pyrrolidone),

propylene glycol, Macrogol (e.g., Macrogol 200 to 600), and combinations of two or more types thereof.

[0146]

Examples of the "preservative" include ethyl
5 parahydroxybenzoate, chlorobutanol, benzyl alcohol, sodium dehydroacetate and sorbic acid.

[0147]

Examples of the "antioxidant" include sodium sulfite and ascorbic acid.

10 [0148]

Examples of the "colorant" include food colors (e.g., Food Color Red No. 2 or 3, and Food Color Yellow No. 4 or 5) and β -carotene.

[0149]

15 Examples of the "sweetening agent" include saccharin sodium, dipotassium glycyrrhizinate and aspartame.

[0150]

As used herein, the pharmaceutical composition can be administered orally or parenterally (e.g., topical, rectal,
20 intravenous, intramuscular, and subcutaneous administration) to human as well as mammals other than human (e.g., mouse, rat, hamster, guinea pig, rabbit, cat, dog, swine, bovine, horse, sheep, and monkey). The dose varies depending on the subject of administration, disease, symptom, dosage form, administration
25 route and the like. For example, the daily dose for oral administration to an adult patient is generally within the range of about 0.01 mg to 1 g based on the active ingredient (i.e., Compound [I]). This amount can be administered in one to several portions.

30 [0151]

Since Compound [I] or a pharmaceutically acceptable salt thereof has a GLUT9 inhibitory action, it is useful as a GLUT9 inhibitor.

[0152]

35 The expression "have GLUT9 inhibitory action" or "inhibit

GLUT9" means elimination or attenuation of GLUT9 activity by inhibiting a GLUT9 function, for example, it means specific inhibition of GLUT9 function under the below-mentioned condition of Test Example 1.

5 The "GLUT9 inhibitor" means a substance which inhibits a GLUT9 function.

 The "GLUT9" is preferably "human GLUT9".

[0153]

 In one embodiment, since Compound [I] or a
10 pharmaceutically acceptable salt thereof has a GLUT9 inhibitory action, it is expected to be effective for diseases that involve GLUT9.

[0154]

 That is, Compound [I] or a pharmaceutically acceptable
15 salt thereof is expected to be useful for the treatment or prophylaxis of a disease selected from the group consisting of hyperuricemia and gout.

 Examples of the "gout" include gouty arthritis, gouty kidney and gouty tophus.

20 [0155]

 In another embodiment, Compound [I] or a pharmaceutically acceptable salt thereof is expected to be useful for the treatment or prophylaxis of diseases selected from the group consisting of chronic kidney disease (CKD), hypertension,
25 diabetes, cardiac disease, arteriosclerotic disease, non-alcoholic fatty liver disease (NAFLD), non-alcoholic steatohepatitis (NASH) and psoriasis.

 Examples of the "cardiac disease" include cardiovascular disease, cardiac failure and atrial fibrillation.

30 [0156]

 As used herein, the "treatment" encompasses improving symptoms, preventing the aggravation of symptoms, maintaining the remission of symptoms, preventing the exacerbation of symptoms, and preventing the relapse of symptoms.

35 As used herein, the "prophylaxis" means suppressing the

onset of symptoms.

[0157]

Compound [I] or a pharmaceutically acceptable salt thereof can be used in combination with one or a plurality of other medicaments (hereinafter to be also referred to as a concomitant drug) according to a method generally employed in the medical field (hereinafter to be referred to as combined use).

[0158]

10 The timing of administering Compound [I] or a pharmaceutically acceptable salt thereof and the concomitant drug is not limited, and they may be administered to the subject as a combination preparation, or the both preparations may be administered simultaneously or separately at certain intervals.

15 In addition, the pharmaceutical composition containing Compound [I] or a pharmaceutically acceptable salt thereof and the concomitant drug may be used in the form of a kit. The dose of the concomitant drug is similar to the clinically-employed dose and can be appropriately selected according to the

20 administration subject, disease, symptom, dosage form, administration route, administration time, combination and the like. The administration form of the concomitant drug is not particularly limited as long as it is combined with Compound [I] or a pharmaceutically acceptable salt thereof.

25 [0159]

Examples of the concomitant drug include

(1) an agent for the treatment and/or prophylaxis of hyperuricemia, and

(2) an agent for the treatment and/or prophylaxis of gout,

30 and at least one of these agents can be used in combination with Compound [I] or a pharmaceutically acceptable salt thereof.

[0160]

As long as the embodiment disclosed herein does not contradict other embodiments disclosed herein, any combination

35 of any of two or more such embodiments is intended to be

encompassed by the technical scope of the present invention.

[0161]

The production methods of Compound [I] or a
pharmaceutically acceptable salt thereof are explained in the
5 following, which should not be construed as limitative. Unless
otherwise referred, the salt of each compound in general
production methods can be selected appropriately from the
above-mentioned "pharmaceutically acceptable salt".

The compound obtained in each step can be, if necessary,
10 isolated or purified according to a method known per se such as
distillation, recrystallization and column chromatography, or
directly used in the next step without isolation or
purification.


[0162]

15 [General Production Method]

[0163]

Production Method A1: Production method of Compound [IA] or a
salt thereof

[0164]




[0165]

wherein

R^{4a} is

- 5 (1) C_{1-8} alkenyl optionally substituted with 1 to 3 substituents independently selected from Group A,
- (2) halo C_{1-6} alkenyl,
- (4) C_{3-6} cycloalkenyl optionally substituted with 1 to 3 substituents independently selected from C_{1-3} alkoxy, or
- 10 (5) a 4- to 6-membered unsaturated heterocyclic group containing one carbon-carbon double bond, and containing 1 or 2 hetero atom as a ring atom in addition to the carbon atoms, wherein the hetero atom is independently selected from the group consisting of oxygen, nitrogen and sulfur atoms, and

wherein the ring atom in the heterocyclic group of Compound [A1-Q2] bonded to the boron is a carbon atom, and the ring atom in the heterocyclic group of Compound [A1-4] bonded to [0166]



5 is a carbon atom,

[0167]

R^{4A} is


(1) C₁₋₈ alkyl optionally substituted with 1 to 3 substituents independently selected from Group A,

10 (2) halo C₁₋₆ alkyl,

(4) C₃₋₆ cycloalkyl optionally substituted with 1 to 3 substituents independently selected from C₁₋₃ alkoxy, or

(5) a 4- to 6-membered saturated heterocyclic group containing 1 or 2 hetero atom as a ring atom in addition to the carbon

15 atoms, wherein the hetero atom is independently selected from the group consisting of oxygen, nitrogen and sulfur atoms, and wherein the ring atom in the heterocyclic group bonded to [0168]



is a carbon atom,

20 [0169]

Group A is as defined above,

Q¹¹ and Q²¹ are each independently halogen,

Q³¹ is a leaving group (e.g., halogen and sulfonyloxy (e.g., methanesulfonyloxy, trifluoromethanesulfonyloxy,

25 benzenesulfonyloxy, and toluenesulfonyloxy)),

Q¹², Q¹³, Q²², Q²³, Q³², and Q³³ are hydroxy, or Q¹² and Q¹³, Q²² and Q²³, and Q³² and Q³³, together with the boron atom that they are bonded to, each independently optionally form a borate,

Q⁴¹ is a protecting group for a carboxy group (e.g., methyl,

benzyl, and tert-butyl), and
the other symbols are as defined above.

[0170]

(Step A1-1)

5 Compound [A1-2] or a salt thereof can be obtained by
subjecting Compound [A1-1] or a salt thereof and Compound [A1-
Q3] to Suzuki coupling reaction. For example, Compound [A1-2]
or a salt thereof can be obtained by reacting Compound [A1-1]
or a salt thereof with Compound [A1-Q3] under heating in the
10 presence of a base and a palladium catalyst, in a solvent.

Examples of the solvent include ether solvents such as
1,4-dioxane, tetrahydrofuran, and 1,2-dimethoxyethane; alcohol
solvents such as methanol, and ethanol; hydrocarbon solvents
such as benzene, toluene, and xylene; polar solvents such as
15 N,N-dimethylformamide, dimethyl sulfoxide, and acetonitrile;
mixed solvents thereof, and mixed solvents of the above-
mentioned solvent and water. A preferable solvent is a mixed
solvent of 1,4-dioxane and water, a mixed solvent of
tetrahydrofuran and water, a mixed solvent of 1,2-
20 dimethoxyethane and water, a mixed solvent of toluene and water,
or a mixed solvent of the above-mentioned mixed solvent and an
alcohol solvent such as ethanol.

Examples of the base include potassium phosphate,
potassium carbonate, sodium carbonate, cesium carbonate and
25 cesium fluoride. A preferable base is potassium phosphate,
potassium carbonate or sodium carbonate.

Examples of the palladium catalyst include palladium
complexes such as tetrakis(triphenylphosphine)palladium(0),
bis(triphenylphosphine)palladium(II) dichloride, and [1,1'-
30 bis(diphenylphosphino)-ferrocene]palladium(II) dichloride; and
palladium complexes prepared in reaction system from a
palladium compound (e.g., palladium(II) acetate, and
tris(dibenzylideneacetone)dipalladium(0)), and a phosphine
ligand (e.g., 2-dicyclohexylphosphino-2',6'-dimethoxybiphenyl,
35 and 2-dicyclohexylphosphino-2',4',6'-triisopropylbiphenyl). A

preferable palladium catalyst is
tetrakis(triphenylphosphine)palladium(0) or [1,1'-
bis(diphenylphosphino)-ferrocene]palladium(II) dichloride.
When Q³¹ is chlorine, a method using palladium(II) acetate and
5 2-dicyclohexylphosphino-2',6'-dimethoxybiphenyl is preferably
employed.

The reaction temperature under heating is, for example,
40°C to 140°C, preferably 70°C to 110°C.

Compound [A1-1] or a salt thereof is a commercially
10 available product, or can be obtained by a known method.

Compound [A1-Q3] is a commercially available product, or
can be obtained by a known method.

[0171]

(Step A1-2)

15 Compound [A1-3] or a salt thereof can be obtained by
subjecting Compound [A1-2] or a salt thereof to a halogenation
reaction. For example, Compound [A1-3] or a salt thereof can
be obtained by reacting Compound [A1-2] or a salt thereof with
a halogenating agent in a solvent.

20 Examples of the solvent include polar solvents such as
N,N-dimethylformamide, dimethyl sulfoxide, acetonitrile, and
acetic acid; and halogen solvents such as dichloromethane, and
chloroform. A preferable solvent is N,N-dimethylformamide or
acetonitrile.

25 Examples of the halogenating agent include N-
bromosuccinimide, N-iodosuccinimide, N-chlorosuccinimide and
bromine. A preferable halogenating agent is N-bromosuccinimide
or N-iodosuccinimide.

The reaction temperature is, for example, 0°C to 120°C,
30 preferably 0°C to room temperature (about 25°C).

[0172]

(Step A1-3)

Compound [A1-4] or a salt thereof can be obtained by
reacting Compound [A1-3] or a salt thereof and Compound [A1-Q2]
35 according to Step A1-1.

Compound [A1-Q2] is a commercially available product, or can be obtained by a known method.

[0173]

(Step A1-4)

5 Compound [A1-5] or a salt thereof can be obtained by
subjecting Compound [A1-4] or a salt thereof to a hydrogenation
reaction. For example, Compound [A1-5] or a salt thereof can
be obtained by reacting Compound [A1-4] or a salt thereof under
hydrogen gas atmosphere in the presence of a palladium catalyst,
10 in a solvent.

Examples of the solvent include ether solvents such as
1,4-dioxane, tetrahydrofuran, and 1,2-dimethoxyethane; alcohol
solvents such as methanol, and ethanol; ester solvents such as
ethyl acetate, and acetic acid isobutyl, and mixed solvents
15 thereof. A preferable solvent is methanol, ethyl acetate, or a
mixed solvent of methanol or ethyl acetate and tetrahydrofuran.

Examples of the palladium catalyst include 5% or 10%
palladium on carbon (dry product, wet product). A preferable
palladium catalyst is 10% palladium on carbon (wet product).

20 The reaction temperature is, for example, room
temperature (about 25°C) to 50°C, preferably room temperature
(about 25°C).

The pressure is, for example, 1 atm to 4 atm, preferably
1 atm.

25 [0174]

(Step A1-5)

Compound [A1-6] or a salt thereof can be obtained by
subjecting Compound [A1-5] or a salt thereof to Sandmeyer
reaction. For example, Compound [A1-6] or a salt thereof can
30 be obtained by subjecting Compound [A1-5] or a salt thereof to
a diazotization in a solvent, and then reacting the resulting
compound with a halogenating agent.

Examples of the solvent include dibromomethane,
acetonitrile, tetrahydrofuran, N,N-dimethylformamide, and water.
35 A preferable solvent is dibromomethane.

Examples of the diazotizing agent include nitrites such as isobutyl nitrite, t-butyl nitrite and isopentyl nitrite, and sodium nitrite. A preferable diazotizing agent is isobutyl nitrite, t-butyl nitrite, isopentyl nitrite or sodium nitrite.

5 Examples of the halogenating agent include bromotrimethylsilane, and copper(II) bromide. A preferable halogenating agent is bromotrimethylsilane.

The reaction temperature is, for example, 0°C to 80°C, preferably 0°C to room temperature (about 25°C).

10 [0175]

(Step A1-6)

Compound [A1-7] or a salt thereof can be obtained by reacting Compound [A1-6] or a salt thereof with Compound [A1-Q1] according to Step A1-1.

15 Compound [A1-Q1] is a commercially available product, or can be obtained by a known method.

[0176]

(Step A1-7)

Compound [IA] or a salt thereof can be obtained by
20 subjecting Compound [A1-7] or a salt thereof to a deprotection reaction to remove Q⁴¹. The deprotection reaction can be carried out in a suitable condition depending on the type of Q⁴¹.

For example, when Q⁴¹ is methyl, Compound [IA] or a salt
25 thereof can be obtained by subjecting Compound [A1-7] or a salt thereof to a hydrolysis reaction in the presence of a base, in a solvent.

Examples of the solvent include a mixed solvent of water and an alcohol solvent such as methanol and ethanol, and a
30 mixed solvent of an alcohol solvent, water and tetrahydrofuran. A preferable solvent is a mixed solvent of methanol and water.

Examples of the base include lithium hydroxide, sodium hydroxide, and potassium hydroxide. A preferable base is sodium hydroxide.

35 The reaction temperature is, for example, room


temperature (about 25°C) to 80°C, preferably room temperature (about 25°C) to 50°C.

[0177]

Production Method A2: Alternative step of Step A1-3 and Step

5 A1-4

[0178]



[0179]

wherein

10 $\text{R}^{4\text{AA}}$ is

(1) C_{1-8} alkyl optionally substituted with 1 to 3 substituents independently selected from Group A, or

(2) halo C_{1-6} alkyl,

Group A is as defined above,

15 Q^{24} is halogen, and

the other symbols are as defined above.

[0180]

(Step A2-1)

Compound [A2-1] or a salt thereof can be obtained by
20 subjecting Compound [A1-3] or a salt thereof and Compound [A2-Q2] to Negishi coupling reaction. For example, Compound [A2-1] or a salt thereof can be obtained by reacting Compound [A1-3] or a salt thereof with Compound [A2-Q2] in the presence of a palladium catalyst and a ligand, in a solvent.

25 Examples of the solvent include ether solvents such as 1,4-dioxane, tetrahydrofuran, and 1,2-dimethoxyethane; and hydrocarbon solvents such as benzene, toluene, and xylene. A preferable solvent is tetrahydrofuran or toluene.

Examples of the palladium catalyst include palladium(II)
30 acetate, and tris(dibenzylideneacetone)dipalladium(0). A preferable palladium catalyst is palladium(II) acetate.

Examples of the ligand include 2-dicyclohexylphosphino-

2',6'-dimethoxybiphenyl, 2-dicyclohexylphosphino-2',6'-diisopropoxybiphenyl, and 2-dicyclohexylphosphino-2',6'-bis(N,N-dimethylamino)biphenyl. A preferable ligand is 2-dicyclohexylphosphino-2',6'-bis(N,N-dimethylamino)biphenyl.


5 The reaction temperature is, for example, 0°C to 50°C, preferably 0°C to room temperature (about 25°C).

Compound [A2-Q2] is a commercially available product, or can be obtained by a known method.

[0181]

10 Production Method A3: Alternative production method of Compound [IA] or a salt thereof

[0182]



[0183]

15 wherein each symbol is as defined above, provided that Q^{31} is preferably a group having reactivity equal to or lower than Q^{11} (for example, when Q^{11} is chlorine, Q^{31} is chlorine), and those skilled in the art can easily select such a group.

[0184]

(Step A3-1)

Compound [A3-2] or a salt thereof can be obtained by reacting Compound [A3-1] or a salt thereof with Compound [A1-
5 Q1] according to Step A1-1.

Compound [A3-1] or a salt thereof is a commercially available product, or can be obtained by a known method.

[0185]

(Step A3-2)

10 Compound [A3-3] or a salt thereof can be obtained by reacting Compound [A3-2] or a salt thereof with Compound [A1-Q3] according to Step A1-1.

[0186]

(Step A3-3)

15 Compound [A3-4] or a salt thereof can be obtained by reacting Compound [A3-3] or a salt thereof according to Step A1-5.

[0187]

(Step A3-4)

20 Compound [A3-5] or a salt thereof can be obtained by reacting Compound [A3-4] or a salt thereof with Compound [A1-Q2] according to Step A1-1.

[0188]

(Step A3-5)


25 Compound [A1-7] or a salt thereof can be obtained by reacting Compound [A3-5] or a salt thereof according to Step A1-4.

[0189]

Production Method A4: Alternative step of Step A3-4 and Step

30 A3-5

[0190]



[0191]

wherein each symbol is as defined above.

[0192]


5 (Step A4-1)

Compound [A4-1] or a salt thereof can be obtained by reacting Compound [A3-4] or a salt thereof with Compound [A2-Q2] according to Step A2-1.

[0193]

10 Production Method A5: Production method of Compound [IAA] or a salt thereof

[0194]



[0195]

wherein

$\text{R}^{4\text{X}}$ and $\text{R}^{5\text{A}}$, together with the carbon atoms that they are bonded
 5 to, form C_{3-6} cycloalkane, and
 the other symbols are as defined above.

[0196]

(Step A5-1)

Compound [A5-2] or a salt thereof can be obtained by
 10 reacting Compound [A5-1] or a salt thereof with Compound [A1-

Q3] according to Step A1-1.

Compound [A5-1] or a salt thereof is a commercially available product, or can be obtained by a known method.

[0197]

5 (Step A5-2)

Compound [A5-3] or a salt thereof can be obtained by reacting Compound [A5-2] or a salt thereof according to Step A1-5.

[0198]

10 (Step A5-3)

Compound [A5-4] or a salt thereof can be obtained by reacting Compound [A5-3] or a salt thereof with Compound [A1-Q1] according to Step A1-1.

[0199]


15 (Step A5-4)

Compound [IAA] or a salt thereof can be obtained by reacting Compound [A5-4] or a salt thereof according to Step A1-7.

[0200]

20 Production Method A6: Production method of Compound [IAAA] or a salt thereof

[0201]



[0202]

wherein

p is 1, 2, 3, 4, 5 or 6,

5 Q^{34} is methyl, ethyl or tert-butyl,

Q^{14} is sulfonyloxy (e.g., methanesulfonyloxy, trifluoromethanesulfonyloxy, benzenesulfonyloxy, and toluenesulfonyloxy),

R^{4AAA} is selected from Group A, and

10 the other symbols are as defined above.

[0203]

(Step A6-1)

Compound [A6-2] or a salt thereof can be obtained by reacting Compound [A6-1] or a salt thereof with a metal
15 alkoxide, in a solvent.

Examples of a combination of the solvent and metal alkoxide include a combination of methanol and sodium methoxide, a combination of ethanol and sodium ethoxide, a combination of benzyl alcohol and sodium benzyloxide, a combination of tert-

butanol and potassium tert-butoxide, and a combination of tetrahydrofuran or N,N-dimethylformamide, and sodium methoxide, sodium ethoxide, sodium benzyloxide or potassium tert-butoxide. A preferable combination of the solvent and metal alkoxide is a
5 combination of tetrahydrofuran and potassium tert-butoxide.

The reaction temperature is, for example, -5°C to 30°C, preferably 0°C to 15°C.

Compound [A6-1] or a salt thereof is a commercially available product, or can be obtained by a known method.

10 [0204]

(Step A6-2)

Compound [A6-3] or a salt thereof can be obtained by reacting Compound [A6-2] or a salt thereof with Compound [A1-Q3] according to Step A1-1.

15 [0205]

(Step A6-3)

Compound [A6-4] or a salt thereof can be obtained by subjecting Compound [A6-3] or a salt thereof to a deprotection reaction to remove Q³⁴. The deprotection reaction can be
20 carried out in a suitable condition depending on the type of Q³⁴.

For example, when Q³⁴ is tert-butyl, Compound [A6-4] or a salt thereof can be obtained by reacting Compound [A6-3] or a salt thereof with an acid, in a solvent.

25 Examples of the solvent include methanol, ethanol, 2-propanol, tetrahydrofuran, toluene, and mixed solvent thereof. A preferable solvent is a mixed solvent of ethanol and tetrahydrofuran.

Examples of the acid include hydrochloric acid,
30 hydrobromic acid, sulfuric acid, and trifluoroacetic acid. A preferable acid is hydrochloric acid.

The reaction temperature is, for example, 15°C to 60°C, preferably 30°C to 40°C.

[0206]

35 (Step A6-4)

Compound [A6-5] or a salt thereof can be obtained by subjecting Compound [A6-4] or a salt thereof to a halogenation reaction according to Step A1-2.

[0207]

5 (Step A6-5a)

Compound [A6-6a] or a salt thereof can be obtained by subjecting Compound [A6-5] or a salt thereof and Compound [A1-R4a] to Sonogashira coupling reaction. For example, Compound [A6-6a] or a salt thereof can be obtained by reacting Compound
10 [A6-5] or a salt thereof with Compound [A1-R4a] in the presence of a base, a palladium catalyst and a copper catalyst, in a solvent.

Examples of the solvent include N,N-dimethylformamide, acetonitrile, and tetrahydrofuran. A preferable solvent is
15 acetonitrile.

Examples of the base include triethylamine, diisopropylethylamine, and diisopropylamine. A preferable base is triethylamine.

Examples of the palladium catalyst include
20 tetrakis(triphenylphosphine)palladium(0), and bis(triphenylphosphine)palladium(II) dichloride. A preferable palladium catalyst is bis(triphenylphosphine)palladium(II) dichloride.

Examples of the copper catalyst include copper(I) iodide,
25 and copper(I) bromide. A preferable copper catalyst is copper(I) iodide.

The reaction temperature is, for example, 15°C to 50°C, preferably 25°C to 40°C.

[0208]

30 (Step A6-5b)

Compound [A6-6b] or a salt thereof can be obtained by subjecting Compound [A6-5] or a salt thereof and Compound [A1-R4b] to Heck reaction. For example, Compound [A6-6b] or a salt thereof can be obtained by reacting Compound [A6-5] or a salt
35 thereof with Compound [A1-R4b] in the presence of a base and a

palladium catalyst, in a solvent.

[0209]

(Step A6-6)

Compound [A6-7] or a salt thereof can be obtained by
5 subjecting Compound [A6-6a] or a salt thereof or Compound [A6-6b] or a salt thereof to a hydrogenation reaction according to Step A1-4.

[0210]

(Step A6-7)

10 Compound [A6-8] or a salt thereof can be obtained by subjecting Compound [A6-7] or a salt thereof to a sulfonylation reaction of hydroxy. The sulfonylation reaction can be carried out in a suitable condition depending on the type of Q¹⁴.

For example, when Q¹⁴ is trifluoromethanesulfonyloxy,
15 Compound [A6-8] or a salt thereof can be obtained by reacting Compound [A6-7] or a salt thereof with trifluoromethanesulfonic anhydride in the presence of a base, in a solvent.

Examples of the solvent include toluene, dichloromethane, pyridine, and mixed solvents of the above-mentioned solvent and
20 water. A preferable solvent is a mixed solvent of toluene and water.

Examples of the base include inorganic bases such as dipotassium hydrogenphosphate, tripotassium phosphate, and potassium carbonate, and organic bases such as pyridine, 4-
25 dimethylaminopyridine, 2,6-lutidine, triethylamine, and diisopropylethylamine. A preferable base is dipotassium hydrogenphosphate.

The reaction temperature is, for example, 0°C to 30°C, preferably 5°C to 10°C.

30 [0211]

(Step A6-8)

Compound [A6-9] or a salt thereof can be obtained by reacting Compound [A6-8] or a salt thereof with Compound [A1-Q1] according to Step A1-6.

35 [0212]


(Step A6-9)

Compound [IAAA] or a salt thereof can be obtained by
subjecting Compound [A6-9] or a salt thereof to a hydrolysis
reaction according to Step A1-7.

5 [0213]

Production Method B1: Production method of Compound [IB] or a
salt thereof

[0214]



10 [0215]

wherein each symbol is as defined above.

[0216]

(Step B1-1)

Compound [B1-2] or a salt thereof can be obtained by reacting Compound [B1-1] or a salt thereof with Compound [A1-Q1] according to Step A1-1.

Compound [B1-1] or a salt thereof is a commercially available product, or can be obtained by a known method.

[0217]

(Step B1-2)

Compound [B1-3] or a salt thereof can be obtained by reacting Compound [B1-2] or a salt thereof according to Step A1-5.

[0218]

(Step B1-3)

Compound [B1-4] or a salt thereof can be obtained by reacting Compound [B1-3] or a salt thereof with Compound [A1-Q3] according to Step A1-1.

[0219]

(Step B1-4)

Compound [B1-5] or a salt thereof can be obtained by reacting Compound [B1-4] or a salt thereof at room temperature (about 25°C) according to Step A1-7.

[0220]

(Step B1-5)

Compound [B1-6] or a salt thereof can be obtained by subjecting Compound [B1-5] or a salt thereof and Compound [B1-R4] or a salt thereof to an amidation reaction. For example, Compound [B1-6] or a salt thereof can be obtained by reacting Compound [B1-5] or a salt thereof with Compound [B1-R4] or a salt thereof in the presence of a condensing agent and an optional base, in a solvent.

Examples of the solvent include ether solvents such as 1,4-dioxane, tetrahydrofuran, and 1,2-dimethoxyethane; halogen solvents such as dichloromethane, and chloroform; and polar solvents such as N,N-dimethylformamide, and acetonitrile. A preferable solvent is N,N-dimethylformamide or acetonitrile.

Examples of the condensing agent include N,N'-

dicyclohexylcarbodiimide, 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide, and HATU [O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate] [alias: 2-(7-aza-1H-benzotriazol-1-yl)-1,1,3,3-tetramethyluronium
5 hexafluorophosphate]. A preferable condensing agent is HATU.

Examples of the optional base include triethylamine, and N,N-diisopropylethylamine. A preferable base is triethylamine.

The reaction temperature is, for example, room temperature (about 25°C) to 60°C, preferably room temperature
10 (about 25°C).

Compound [B1-R4] or a salt thereof is a commercially available product, or can be obtained by a known method.

[0221]


(Step B1-6)

15 Compound [IB] or a salt thereof can be obtained by reacting Compound [B1-6] or a salt thereof according to Step A1-7.

[0222]

Production Method C1: Production method of Compound [IC] or a
20 salt thereof

[0223]



[0224]

wherein

L^{4C} is C_{1-8} alkylene or C_{3-6} cycloalkylene,

5 R^{4C} is C_{1-3} alkyl optionally substituted with one hydroxy or one C_{1-3} alkoxy, or halo C_{1-3} alkyl,

Q^{25} is halogen, and

the other symbols are as defined above.

[0225]

10 (Step C1-1)

Compound [C1-2] or a salt thereof can be obtained by reacting Compound [C1-1] or a salt thereof with Compound [A1-Q1] according to Step A1-1.

Compound [C1-1] or a salt thereof is a commercially available product, or can be obtained by a known method.

[0226]

(Step C1-2)

Compound [C1-3] or a salt thereof can be obtained by

subjecting Compound [C1-2] or a salt thereof and Compound [C1-
Q2] to an alkylation reaction. For example, Compound [C1-3] or
a salt thereof can be obtained by reacting Compound [C1-2] or a
salt thereof with Compound [C1-Q2] in the presence of a base,
5 in a solvent.

Examples of the solvent include ether solvents such as
1,4-dioxane, tetrahydrofuran and 1,2-dimethoxyethane, and N,N-
dimethylformamide. A preferable solvent is tetrahydrofuran or
N,N-dimethylformamide.

10 Examples of the base include sodium hydride, potassium t-
butoxide, and sodium t-butoxide. A preferable base is sodium
hydride.

The reaction temperature is, for example, 0°C to 70°C,
preferably 0°C to room temperature (about 25°C).

15 Compound [C1-Q2] is a commercially available product, or
can be obtained by a known method.

[0227]

(Step C1-3)

Compound [C1-4] or a salt thereof can be obtained by
20 reacting Compound [C1-3] or a salt thereof with Compound [A1-
Q3] according to Step A1-1.

[0228]

(Step C1-4)

Compound [IC] or a salt thereof can be obtained by
25 reacting Compound [C1-4] or a salt thereof according to Step
A1-7.

Examples

[0229]

Next, the production method of Compound [I] or a
30 pharmaceutically acceptable salt thereof is concretely
explained by referring to Examples, which should not be
construed as limitative.


In the following Examples, the following abbreviation is
used.

35 HATU: O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium

hexafluorophosphate [alias: 2-(7-aza-1H-benzotriazol-1-yl)-
1,1,3,3-tetramethyluronium hexafluorophosphate]
[0230]


Production Example 1

- 5 Synthesis of 4-(3-(3,5-difluoro-4-methoxyphenyl)-6-(3-methoxypropyl)pyrazin-2-yl)benzoic acid (Example 67)
[0231]



[0232]

- 10 Step 1-1: Methyl 4-(3-aminopyrazin-2-yl)benzoate
[0233]



[0234]

- Under inert gas atmosphere, to a solution of 3-
15 chloropyrazin-2-amine (3.00 g, 23.2 mmol), (4-(methoxycarbonyl)phenyl)boric acid (5.00 g, 27.8 mmol) and [1,1'-bis(diphenylphosphino)-ferrocene]palladium(II) dichloride dichloromethane adduct (0.378 g, 0.463 mmol) in tetrahydrofuran (100 mL) was added 2M-aqueous potassium phosphate solution
20 (23.2 mL, 46.3 mmol), and the mixture was stirred at 70°C for 1 hr. The reaction solution was diluted with water and ethyl acetate, and separated, and the organic layer was washed successively with water and saturated brine, and dried over sodium sulfate. The sodium sulfate was removed by filtration,
25 and the filtrate was concentrated under reduced pressure. The residue was purified by silica gel chromatography (hexane:ethyl


acetate =2:1 to 1:2) to give methyl 4-(3-aminopyrazin-2-yl)benzoate (2.56 g, yield 48%).

¹H-NMR (DMSO-D₆) δ: 3.87 (3H, s), 6.27 (2H, br s), 7.84 (2H, dt, J = 8.4, 1.8 Hz), 7.89 (1H, d, J = 2.5 Hz), 7.97 (1H, d, J = 2.5 Hz), 8.04 (2H, dt, J = 8.6, 1.8 Hz).

[0235]

Step 1-2: Methyl 4-(3-amino-6-bromopyrazin-2-yl)benzoate

[0236]



10 [0237]


Under ice cooling, to a suspension of methyl 4-(3-aminopyrazin-2-yl)benzoate (3.15 g, 13.7 mmol) in acetonitrile (80 mL) was added N-bromosuccinimide (2.56 g, 14.4 mmol), and the mixture was stirred for 30 min. To the reaction solution was added water (160 mL), and the mixture was stirred at room temperature for 1 hr. The precipitated solid was collected by filtration, and dried under reduced pressure to give methyl 4-(3-amino-6-bromopyrazin-2-yl)benzoate (3.93 g, yield 92%).

¹H-NMR (DMSO-D₆) δ: 3.88 (3H, s), 6.57 (2H, br s), 7.82 (2H, dt, J = 8.5, 1.8 Hz), 8.05 (2H, dt, J = 8.6, 1.8 Hz), 8.12 (1H, s).

[0238]

Step 1-3: Methyl (E)-4-(3-amino-6-(3-methoxyprop-1-en-1-yl)pyrazin-2-yl)benzoate

[0239]



25 [0240]

Under inert gas atmosphere, to a suspension of methyl 4-(3-amino-6-bromopyrazin-2-yl)benzoate (2.00 g, 6.49 mmol), (E)-2-(3-methoxyprop-1-en-1-yl)-4,4,5,5-tetramethyl-1,3,2-


dioxaborolane (1.54 g, 7.79 mmol) and [1,1'-bis(diphenylphosphino)-ferrocene]palladium(II) dichloride dichloromethane adduct (0.106 g, 0.130 mmol) in toluene (20 mL) was added 2M-aqueous potassium phosphate solution (4.87 mL, 9.74 mmol), and the mixture was stirred at 100°C for 2 hr. The mixture was allowed to cool to room temperature, and diluted with ethyl acetate (50 mL), and the insoluble substance was removed by filtration through Celite. The organic layer was washed successively with water and saturated brine, and dried over magnesium sulfate. The magnesium sulfate was removed by filtration, and the filtrate was concentrated under reduced pressure. The obtained residue was purified by silica gel chromatography (hexane:ethyl acetate =2:1 to 1:4) to give methyl (E)-4-(3-amino-6-(3-methoxyprop-1-en-1-yl)pyrazin-2-yl)benzoate (1.78 g, yield 91%).

¹H-NMR (DMSO-D₆) δ: 3.27 (3H, s), 3.88 (3H, s), 4.04 (2H, dd, J = 5.4, 1.3 Hz), 6.37 (2H, br s), 6.46-6.53 (1H, m), 6.59 (1H, dt, J = 15.7, 1.2 Hz), 7.86 (2H, dt, J = 8.5, 1.8 Hz), 8.03-8.06 (3H, m).

[0241]

Step 1-4: Methyl 4-(3-amino-6-(3-methoxypropyl)pyrazin-2-yl)benzoate

[0242]



[0243]

To methyl (E)-4-(3-amino-6-(3-methoxyprop-1-en-1-yl)pyrazin-2-yl)benzoate (777 mg, 2.60 mmol) were added methanol (13 mL) and 10% palladium on carbon catalyst (50% wet, 155 mg), and the mixture was stirred under hydrogen gas atmosphere for 24 hr. The palladium on carbon catalyst was removed from the reaction solution by filtration, and the filtrate was concentrated under reduced pressure. The residue


was purified by silica gel chromatography (hexane:ethyl acetate =1:4) to give methyl 4-(3-amino-6-(3-methoxypropyl)pyrazin-2-yl)benzoate (772 mg, yield 98%).

1H-NMR (DMSO-D6) δ : 1.80-1.88 (2H, m), 2.64 (2H, t, J = 7.6 Hz),
5 3.21 (3H, s), 3.34 (2H, t, J = 6.4 Hz), 3.87 (3H, s), 6.02 (2H, br s), 7.84-7.87 (3H, m), 8.03 (2H, dt, J = 8.6, 1.8 Hz).

[0244]

Step 1-5: Methyl 4-(3-bromo-6-(3-methoxypropyl)pyrazin-2-yl)benzoate

10 [0245]



[0246]


A solution of methyl 4-(3-amino-6-(3-methoxypropyl)pyrazin-2-yl)benzoate (5.00 g, 16.6 mmol) in
15 dibromomethane (140 mL) was stirred at room temperature, and isopentyl nitrite (2.40 mL, 18.3 mmol) was added thereto. To the reaction solution was added dropwise a solution of bromotrimethylsilane (2.41 mL, 18.3 mmol) in dibromomethane (20 mL) over 10 min, and the reaction solution was stirred at room
20 temperature for 24 hr. To the reaction solution was added saturated aqueous sodium hydrogencarbonate solution (50 mL), and the mixture was extracted with ethyl acetate (100 mL). The organic layer was washed successively with water and saturated brine, and dried over sodium sulfate. The sodium sulfate was
25 removed by filtration, and the filtrate was concentrated under reduced pressure. The residue was purified by silica gel chromatography (hexane:ethyl acetate =7:1 to 2:1) to give methyl 4-(3-bromo-6-(3-methoxypropyl)pyrazin-2-yl)benzoate (4.31 g, yield 71%).

30 1H-NMR (DMSO-D6) δ : 1.89-1.96 (2H, m), 2.82-2.87 (2H, m), 3.21 (3H, s), 3.36 (2H, t, J = 6.2 Hz), 3.89 (3H, s), 7.85 (2H, dt, J = 8.5, 1.8 Hz), 8.08 (2H, dt, J = 8.4, 1.8 Hz), 8.43 (1H, s).

[0247]

Step 1-6: Methyl 4-(3-(3,5-difluoro-4-methoxyphenyl)-6-(3-methoxypropyl)pyrazin-2-yl)benzoate

[0248]



[0249]


Under inert gas atmosphere, to a solution of methyl 4-(3-bromo-6-(3-methoxypropyl)pyrazin-2-yl)benzoate (4.30 g, 11.8 mmol) and (3,5-difluoro-4-methoxyphenyl)boric acid (2.66 g, 14.1 mmol) in toluene (44 mL) were added [1,1'-bis(diphenylphosphino)-ferrocene]palladium(II) dichloride dichloromethane adduct (0.192 g, 0.235 mmol) and 2M-aqueous potassium phosphate solution (8.83 mL, 17.7 mmol), and the mixture was stirred at 100°C for 4 hr. The mixture was allowed to cool to room temperature, water (100 mL) was added thereto, and the mixture was extracted with ethyl acetate (200 mL). The organic layer was washed successively with water and saturated brine, and dried over sodium sulfate. The sodium sulfate was removed by filtration, and the filtrate was concentrated under reduced pressure. The obtained residue was purified by silica gel chromatography (hexane:ethyl acetate =4:1 to 2:1) to give methyl 4-(3-(3,5-difluoro-4-methoxyphenyl)-6-(3-methoxypropyl)pyrazin-2-yl)benzoate (4.86 g, yield 96%).

¹H-NMR (DMSO-D₆) δ: 1.94-2.01 (2H, m), 2.90-2.94 (2H, m), 3.23 (3H, s), 3.40 (2H, t, J = 6.2 Hz), 3.85 (3H, s), 3.93 (3H, s), 7.05-7.12 (2H, m), 7.56 (2H, dt, J = 8.5, 1.8 Hz), 7.95 (2H, dt, J = 8.4, 1.8 Hz), 8.65 (1H, s).

[0250]

Step 1-7: 4-(3-(3,5-Difluoro-4-methoxyphenyl)-6-(3-methoxypropyl)pyrazin-2-yl)benzoic acid

[0251]



[0252]

To a solution of methyl 4-(3-(3,5-difluoro-4-methoxyphenyl)-6-(3-methoxypropyl)pyrazin-2-yl)benzoate (1.00 g, 2.33 mmol) in methanol (14 mL) was added 4N-aqueous sodium hydroxide solution (3.50 mL, 14.0 mmol), and the mixture was stirred at 50°C for 2 hr. The mixture was allowed to cool to room temperature, 10wt%-aqueous citric acid solution (10.5 mL) and water (15 mL) were added thereto, and the mixture was stirred at room temperature for 2 hr. The precipitated solid was collected by filtration, and dried under reduced pressure to give 4-(3-(3,5-difluoro-4-methoxyphenyl)-6-(3-methoxypropyl)pyrazin-2-yl)benzoic acid (824 mg, yield 85%).

¹H-NMR (DMSO-D₆) δ: 1.92-2.02 (2H, m), 2.91 (2H, t, J = 7.7 Hz), 3.24 (3H, s), 3.40 (2H, t, J = 6.4 Hz), 3.93 (3H, s), 7.09 (2H, d, J = 9.5 Hz), 7.53 (2H, d, J = 8.6 Hz), 7.92 (2H, d, J = 8.6 Hz), 8.64 (1H, s), 13.10 (1H, s).

[0253]

Step 1-8: Crystals of 4-(3-(3,5-difluoro-4-methoxyphenyl)-6-(3-methoxypropyl)pyrazin-2-yl)benzoic acid


To 4-(3-(3,5-difluoro-4-methoxyphenyl)-6-(3-methoxypropyl)pyrazin-2-yl)benzoic acid (50 mg) were added 2-propanol (0.075 mL) and n-heptane (0.025 mL), and the mixture was stirred at 100°C to give a solution. The stirring was stopped, and the mixture was allowed to cool to room temperature. The precipitated solid was collected by filtration, and dried under reduced pressure to give crystals of 4-(3-(3,5-difluoro-4-methoxyphenyl)-6-(3-methoxypropyl)pyrazin-2-yl)benzoic acid (31 mg, yield 62%).

[0254]

Production Example 2

Synthesis of 4-(5-(butylcarbamoyl)-2-(4-methoxyphenyl)pyridin-3-yl)benzoic acid hydrochloride (Example 76)


[0255]



[0256]

Step 2-1: Methyl 5-amino-6-(4-methoxyphenyl)nicotinate

[0257]



10 [0258]

To a mixture of methyl 5-amino-6-bromonicotinate (0.500 g, 2.16 mmol), (4-methoxyphenyl)boric acid (0.660 g, 4.34 mmol) and potassium phosphate (1.43 g, 6.73 mmol) were added 1,2-dimethoxyethane (12 mL) and water (4 mL). Under inert gas atmosphere, [1,1'-bis(diphenylphosphino)-ferrocene]palladium(II) dichloride dichloromethane adduct (0.180 g, 0.220 mmol) was added thereto, and the mixture was stirred at 100°C for 3.5 hr. The reaction solution was diluted with water and ethyl acetate, and the insoluble substance was removed by filtration. The organic layer was washed with saturated brine, and dried over sodium sulfate. The sodium sulfate was removed by filtration, and the filtrate was concentrated under reduced pressure. The residue was dissolved in ethyl acetate (10 mL), and 4N-hydrogen chloride ethyl acetate solution (2 mL) was added thereto. The precipitated solid was collected by filtration, and dissolved in water (20 mL). To this solution was added saturated aqueous sodium hydrogencarbonate solution, and the resulting solid was

15

20

25


collected by filtration, and dried under reduced pressure to give methyl 5-amino-6-(4-methoxyphenyl)nicotinate (0.448 g, yield 80%).

¹H-NMR (DMSO-D₆) δ: 3.80 (3H, s), 3.84 (3H, s), 5.36 (2H, br s),
5 7.02 (2H, dt, J = 9.5, 2.4 Hz), 7.63–7.68 (3H, m), 8.37 (1H, d, J = 1.8 Hz).

[0259]

Step 2-2: Methyl 5-bromo-6-(4-methoxyphenyl)nicotinate

[0260]




[0261]

Under inert gas atmosphere, a solution of methyl 5-amino-6-(4-methoxyphenyl)nicotinate (0.448 g, 1.74 mmol) in dibromomethane was stirred at room temperature, and isopentyl
15 nitrite (0.256 mL, 1.91 mmol) was added thereto. To the reaction solution was added dropwise a solution of bromotrimethylsilane (0.249 mL, 1.91 mmol) in dibromomethane, and the mixture was stirred at room temperature for 2 hr. To the reaction solution was added saturated aqueous sodium
20 hydrogencarbonate solution (10 mL), and the mixture was extracted with chloroform. The organic layer was washed with saturated brine, and dried over sodium sulfate. The sodium sulfate was removed by filtration, and the filtrate was concentrated under reduced pressure. The residue was purified
25 by silica gel chromatography (hexane:ethyl acetate =6:1 to 4:1) to give methyl 5-bromo-6-(4-methoxyphenyl)nicotinate (0.308 g, yield 55%).

[0262]

Step 2-3: Methyl 5-(4-(tert-butoxycarbonyl)phenyl)-6-(4-methoxyphenyl)nicotinate
30

[0263]



[0264]


To a mixture of methyl 5-bromo-6-(4-methoxyphenyl)nicotinate (0.308 g, 0.956 mmol), (4-(tert-butoxycarbonyl)phenyl)boronic acid (0.425 g, 1.91 mmol) and potassium phosphate (0.609 g, 2.87 mmol) were added toluene (4.5 mL) and water (1.5 mL). Under inert gas atmosphere, [1,1'-bis(diphenylphosphino)-ferrocene]palladium(II) dichloride dichloromethane adduct (0.078 g, 0.096 mmol) was added thereto, and the mixture was stirred at 100°C for 4 hr. The reaction solution was diluted with water and ethyl acetate, and the insoluble substance was removed by filtration. The organic layer was washed with saturated brine, and dried over sodium sulfate. The sodium sulfate was removed by filtration, and the filtrate was concentrated under reduced pressure. The residue was purified by silica gel chromatography (hexane:ethyl acetate =5:1 to 3:1), ethyl acetate (1 mL) and hexane (10 mL) were added thereto, and the obtained suspension was stirred. The insoluble substance was collected by filtration, and dried under reduced pressure to give methyl 5-(4-(tert-butoxycarbonyl)phenyl)-6-(4-methoxyphenyl)nicotinate (0.149 mg, yield 37%).

¹H-NMR (DMSO-D₆) δ: 1.53 (9H, s), 3.73 (3H, s), 3.91 (3H, s), 6.85 (2H, dt, J = 9.4, 2.5 Hz), 7.29 (2H, dt, J = 9.4, 2.5 Hz), 7.37 (2H, dt, J = 8.4, 1.8 Hz), 7.86 (2H, dt, J = 8.3, 1.8 Hz), 8.17 (1H, d, J = 2.1 Hz), 9.14 (1H, d, J = 2.1 Hz).

[0265]

Step 2-4: 5-(4-(tert-Butoxycarbonyl)phenyl)-6-(4-methoxyphenyl)nicotinic acid

[0266]




[0267]

Methyl 5-(4-(tert-butoxycarbonyl)phenyl)-6-(4-methoxyphenyl)nicotinate (137 mg, 0.327 mmol) was dissolved in 5 methanol (2 mL) and tetrahydrofuran (2 mL). To this solution was added 4M-aqueous lithium hydroxide solution (0.50 mL, 2.00 mmol), and the mixture was stirred at room temperature for 16 hr. To the reaction solution was added 1M-hydrochloric acid (2.0 mL), the methanol and tetrahydrofuran were evaporated
 10 under reduced pressure, and the residue was diluted with water (1 mL). The precipitated solid was collected by filtration, and dried under reduced pressure to give 5-(4-(tert-butoxycarbonyl)phenyl)-6-(4-methoxyphenyl)nicotinic acid (56 mg).

15 [0268]

Step 2-5: tert-Butyl 4-(5-(butylcarbamoyl)-2-(4-methoxyphenyl)pyridin-3-yl)benzoate

[0269]



20 [0270]


To a solution of 5-(4-(tert-butoxycarbonyl)phenyl)-6-(4-methoxyphenyl)nicotinic acid (56 mg, 0.138 mmol) obtained in the previous step in N,N-dimethylformamide were added successively butan-1-amine (0.050 mL, 0.51 mmol), triethylamine
 25 (0.060 mL, 0.43 mmol) and HATU (80 mg, 0.21 mmol), and the mixture was stirred at room temperature for 23 hr. The reaction solution was diluted with saturated aqueous sodium

hydrogencarbonate solution and water, and extracted with ethyl acetate. The organic layer was washed successively with water and saturated brine, and dried over sodium sulfate. The sodium sulfate was removed by filtration, and the filtrate was concentrated under reduced pressure. The residue was purified by silica gel chromatography (hexane:ethyl acetate =3:1 to 1:1) to give tert-butyl 4-(5-(butylcarbamoyl)-2-(4-methoxyphenyl)pyridin-3-yl)benzoate (19 mg, yield 29%).

[0271]

Step 2-6: 4-(5-(Butylcarbamoyl)-2-(4-methoxyphenyl)pyridin-3-yl)benzoic acid hydrochloride

[0272]



[0273]

tert-Butyl 4-(5-(butylcarbamoyl)-2-(4-methoxyphenyl)pyridin-3-yl)benzoate (18 mg, 0.039 mmol) was dissolved in trifluoroacetic acid (1.0 mL), and the solution was stirred at room temperature for 1.5 hr. The reaction solution was concentrated under reduced pressure, and to the obtained residue was added 4N-hydrogen chloride ethyl acetate solution. The resulting precipitate was collected by filtration, and dried under reduced pressure to give 4-(5-(butylcarbamoyl)-2-(4-methoxyphenyl)pyridin-3-yl)benzoic acid hydrochloride (7.9 mg, yield 45%).


¹H-NMR (DMSO-D₆) δ: 0.91 (3H, t, J = 7.4 Hz), 1.32-1.40 (2H, m), 1.50-1.57 (2H, m), 3.31 (2H, dd, J = 12.8, 6.8 Hz), 3.74 (3H, d, J = 0.7 Hz), 6.86 (2H, dd, J = 8.9, 2.9 Hz), 7.28 (2H, dd, J = 8.8, 1.6 Hz), 7.39 (2H, d, J = 8.1 Hz), 7.91 (2H, d, J = 8.1 Hz), 8.25 (1H, br s), 8.74 (1H, br s), 9.07 (1H, s).

[0274]

Production Example 3

Synthesis of 4-(2-(4-methoxyphenyl)-5-(3-methoxypropyl)pyridin-3-yl)benzoic acid (Example 84)


[0275]



5 [0276]

Step 3-1: 3-(Benzyloxy)-5-bromo-2-chloropyridine

[0277]



[0278]

10 To a solution of 5-bromo-2-chloropyridin-3-ol (50.2 g, 241 mmol) in N,N-dimethylformamide (200 mL) were added successively benzyl bromide (33.0 mL, 278 mmol) and potassium carbonate (48.6 g, 352 mmol), and the mixture was stirred at room temperature for 4 hr. To the reaction solution was added
15 water (600 mL), and the mixture was stirred for 2 hr. The precipitate was collected by filtration, and dried under reduced pressure to give 3-(benzyloxy)-5-bromo-2-chloropyridine (69.7 g, yield 96%).


1H-NMR (DMSO-D6) δ : 5.29 (2H, s), 7.33-7.47 (5H, m), 7.98 (1H, d, J = 2.1 Hz), 8.14 (1H, d, J = 1.8 Hz).

20

[0279]

Step 3-2: (E)-3-(Benzyloxy)-2-chloro-5-(3-methoxyprop-1-en-1-yl)pyridine

[0280]



25

[0281]


To a suspension of 3-(benzyloxy)-5-bromo-2-chloropyridine (25.7 mg, 86.0 mmol) and potassium phosphate (54.9 g, 259 mmol) in tetrahydrofuran (180 mL) were added successively water (130 mL), (E)-2-(3-methoxyprop-1-en-1-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (20.1 mL, 95.0 mmol) and [1,1'-bis(diphenylphosphino)-ferrocene]palladium(II) dichloride dichloromethane adduct (3.54 g, 4.33 mmol), and the mixture was stirred at room temperature for 2 hr. [1,1'-Bis(diphenylphosphino)-ferrocene]palladium(II) dichloride dichloromethane adduct (1.70 g, 2.04 mmol) was added again thereto, and the mixture was stirred at room temperature for 3 hr, warmed to 33°C, and stirred for 1 hr. The reaction solution was diluted with ethyl acetate (180 mL), and the insoluble substance was removed by filtration. The organic layer of the filtrate was washed with water and saturated brine, silica gel (50 g) was added thereto, and the mixture was stirred at room temperature for 1 hr. The silica gel was removed by filtration, and the filtrate was concentrated under reduced pressure. The residue was purified by silica gel chromatography (hexane:ethyl acetate =15:1 to 4:3) to give (E)-3-(benzyloxy)-2-chloro-5-(3-methoxyprop-1-en-1-yl)pyridine (20.9 g, yield 83%).

¹H-NMR (DMSO-D₆) δ: 3.29 (3H, s), 4.06-4.07 (2H, m), 5.29 (2H, s), 6.59-6.60 (2H, m), 7.32-7.49 (5H, m), 7.81 (1H, d, J = 1.8 Hz), 8.02 (1H, d, J = 2.1 Hz).

[0282]

Step 3-3: (E)-3-(Benzyloxy)-2-(4-methoxyphenyl)-5-(3-methoxyprop-1-en-1-yl)pyridine

[0283]



[0284]

To a solution of (E)-3-(benzyloxy)-2-chloro-5-(3-


methoxyprop-1-en-1-yl)pyridine (17.8 g, 61.4 mmol), 4-methoxyphenylboric acid (11.2 g, 73.7 mmol), palladium(II) acetate (0.276 g, 1.13 mmol) and 2-dicyclohexylphosphino-2',6'-dimethoxybiphenyl (1.01 g, 2.46 mmol) in 1,2-dimethoxyethane
 5 (138 mL) was added 2M-aqueous potassium phosphate solution (46.1 mL, 92.2 mmol), and the mixture was stirred at 50°C for 5 hr. The reaction solution was allowed to cool to room temperature, water (100 mL) was added thereto, and the mixture was extracted with ethyl acetate (200 mL). The organic layer
 10 was washed successively with water and saturated brine, and dried over sodium sulfate. The sodium sulfate was removed by filtration, and the filtrate was concentrated under reduced pressure. The residue was purified by silica gel chromatography (hexane:ethyl acetate =3:1 to 1:1) to give (E)-
 15 3-(benzyloxy)-2-(4-methoxyphenyl)-5-(3-methoxyprop-1-en-1-yl)pyridine (23.1 g, yield 104%).

¹H-NMR (DMSO-D₆) δ: 3.30 (3H, s), 3.77 (3H, s), 4.07-4.09 (2H, m), 5.26 (2H, s), 6.52-6.66 (2H, m), 6.95 (2H, dt, J = 9.6, 2.5 Hz), 7.30-7.46 (5H, m), 7.72 (1H, d, J = 1.6 Hz), 7.93 (2H, dt, J = 9.6, 2.5 Hz), 8.27 (1H, d, J = 1.6 Hz).

[0285]

Step 3-4: 2-(4-Methoxyphenyl)-5-(3-methoxypropyl)pyridin-3-ol

[0286]



25 [0287]

To (E)-3-(benzyloxy)-2-(4-methoxyphenyl)-5-(3-methoxyprop-1-en-1-yl)pyridine (23.1 g) obtained in the previous step were added methanol (230 mL) and 10% palladium on carbon catalyst (50% wet, 4.62g), and the mixture was stirred
 30 under hydrogen gas atmosphere for 24 hr. The palladium on carbon catalyst was removed from the reaction solution by filtration, and the filtrate was concentrated under reduced


pressure. To the obtained crude crystals was added ethyl acetate (50 mL), the mixture was stirred at 80°C for 20 min, and hexane (150 mL) was added thereto. The mixture was stirred for additional 2 hr while allowed to cool to room temperature.

5 The precipitate was collected by filtration, and dried under reduced pressure to give 2-(4-methoxyphenyl)-5-(3-methoxypropyl)pyridin-3-ol (13.6 g, yield in two step 81%).
1H-NMR (DMSO-D6) δ : 1.74-1.81 (2H, m), 2.54-2.58 (2H, m), 3.23 (3H, s), 3.32 (2H, t, J = 6.4 Hz), 3.77 (3H, s), 6.95 (2H, dt, J = 9.5, 2.5 Hz), 7.09 (1H, d, J = 1.8 Hz), 7.95-7.99 (3H, m), 9.94 (1H, br s).

[0288]

Step 3-5: 2-(4-Methoxyphenyl)-5-(3-methoxypropyl)pyridin-3-yl trifluoromethanesulfonate

15 [0289]



[0290]

To a solution of 2-(4-methoxyphenyl)-5-(3-methoxypropyl)pyridin-3-ol (13.7 g, 50.0 mmol) in N,N-dimethylformamide (100 mL) was added potassium carbonate (15.2 g, 110 mmol). The reaction solution was ice-cooled, N-(5-chloropyridin-2-yl)-1,1,1-trifluoro-N-((trifluoromethyl)sulfonyl)methanesulfonamide (21.6 g, 55.0 mmol) was added thereto, and the mixture was stirred at room temperature for 2 hr. The reaction solution was diluted with water, and extracted with ethyl acetate. The organic layer was washed successively with water and saturated brine, and dried over sodium sulfate. The sodium sulfate was removed by filtration, and the filtrate was concentrated under reduced pressure. The residue was purified by silica gel chromatography (hexane:ethyl acetate =4:1 to 1:1) to give 2-(4-

20
25
30


methoxyphenyl)-5-(3-methoxypropyl)pyridin-3-yl
trifluoromethanesulfonate (18.9 g, yield 93%).

¹H-NMR (DMSO-D₆) δ: 1.82-1.89 (2H, m), 2.73-2.77 (2H, m), 3.23
(3H, s), 3.34 (2H, t, J = 6.4 Hz), 3.81 (3H, s), 7.06 (2H, dt,
5 J = 9.5, 2.5 Hz), 7.68 (2H, dt, J = 9.6, 2.5 Hz), 7.90 (1H, d,
J = 1.6 Hz), 8.59 (1H, d, J = 1.6 Hz).

[0291]

Step 3-6: Methyl 4-(2-(4-methoxyphenyl)-5-(3-
methoxypropyl)pyridin-3-yl)benzoate

10 [0292]



[0293]

To a solution of 2-(4-methoxyphenyl)-5-(3-
methoxypropyl)pyridin-3-yl trifluoromethanesulfonate (18.9 g,
15 46.7 mmol) and (4-(methoxycarbonyl)phenyl)boric acid (10.1 g,
56.1 mmol) in 1,2-dimethoxyethane (105 mL) were added
successively 2M-aqueous potassium phosphate solution (35.0 mL,
70.0 mmol) and [1,1'-bis(diphenylphosphino)-
ferrocene]palladium(II) dichloride dichloromethane adduct
20 (0.763 g, 0.934 mmol), and the mixture was stirred at 80°C for
1 hr. The reaction solution was allowed to cool to room
temperature, and water was added thereto. The mixture was
extracted with ethyl acetate, and the organic layer was dried
over sodium sulfate. The sodium sulfate was removed by
25 filtration, and the filtrate was concentrated under reduced
pressure. To the obtained residue were added ethyl acetate (50
mL) and hexane (150 mL), the mixture was stirred at room
temperature for 30 min, and the insoluble substance was removed
by filtration. The filtrate was concentrated under reduced
30 pressure, and the obtained residue was purified by silica gel
chromatography (hexane:ethyl acetate =3:1 to 1:1) to give


methyl 4-(2-(4-methoxyphenyl)-5-(3-methoxypropyl)pyridin-3-yl)benzoate (16.9 g, 43.4 mmol).

¹H-NMR (DMSO-D₆) δ: 1.83-1.90 (2H, m), 2.68-2.73 (2H, m), 3.24 (3H, s), 3.36 (2H, t, J = 6.4 Hz), 3.71 (3H, s), 3.83 (3H, s),
5 6.79 (2H, dt, J = 9.4, 2.5 Hz), 7.18 (2H, dt, J = 9.4, 2.5 Hz),
7.33 (2H, dt, J = 8.4, 1.8 Hz), 7.64 (1H, d, J = 2.1 Hz), 7.88 (2H, dt, J = 8.5, 1.9 Hz), 8.51 (1H, d, J = 2.1 Hz).

[0294]

Step 3-7: 4-(2-(4-Methoxyphenyl)-5-(3-methoxypropyl)pyridin-3-yl)benzoic acid
10

[0295]



[0296]

To a solution of methyl 4-(2-(4-methoxyphenyl)-5-(3-methoxypropyl)pyridin-3-yl)benzoate (2.00 g, 5.11 mmol) in
15 methanol (30.6 mL) was added 4N-aqueous sodium hydroxide solution (7.66 mL, 30.6 mmol), and the mixture was stirred at
50°C for 2 hr. The mixture was allowed to cool to room temperature, 10wt%-aqueous citric acid solution (23.0 mL) and
20 water (46 mL) were added thereto, and the mixture was stirred at room temperature for 2 hr. The precipitated solid was
collected by filtration, and dried under reduced pressure to give 4-(2-(4-methoxyphenyl)-5-(3-methoxypropyl)pyridin-3-yl)benzoic acid (1.66 g, yield 86%).

25 ¹H-NMR (DMSO-D₆) δ: 1.84-1.93 (2H, m), 2.72 (2H, t, J = 7.8 Hz),
3.25 (3H, s), 3.38 (2H, t, J = 6.4 Hz), 3.72 (3H, s), 6.81 (2H, d, J = 8.8 Hz), 7.20 (2H, d, J = 8.8 Hz), 7.32 (2H, d, J = 8.3 Hz),
7.65 (1H, d, J = 2.2 Hz), 7.87 (2H, d, J = 8.6 Hz), 8.52 (1H, d, J = 2.2 Hz), 12.99 (1H, br s).

30 [0297]

Step 3-8: Crystals (Form II) of 4-(2-(4-methoxyphenyl)-5-(3-methoxypropyl)pyridin-3-yl)benzoic acid

To 4-(2-(4-methoxyphenyl)-5-(3-methoxypropyl)pyridin-3-yl)benzoic acid (300 mg) was added methyl t-butyl ether (3 mL),
5 and the mixture was stirred at 70°C to give a solution. While stirring, the mixture was allowed to cool to room temperature, and then for 3 days. The precipitated solid was collected by filtration, and dried under reduced pressure to give crystals (Form II) of 4-(2-(4-methoxyphenyl)-5-(3-methoxypropyl)pyridin-
10 3-yl)benzoic acid (191 mg, yield 64%).

[0298]

Step 3-9: Crystals (Form X) of 4-(2-(4-methoxyphenyl)-5-(3-methoxypropyl)pyridin-3-yl)benzoic acid

4-(2-(4-Methoxyphenyl)-5-(3-methoxypropyl)pyridin-3-yl)benzoic acid (20 mg) was suspended in methanol (0.12 mL),
15 and the suspension was stirred at room temperature for 2 weeks. The resulting solid was collected by filtration, and dried under reduced pressure to give crystals (Form X) of 4-(2-(4-methoxyphenyl)-5-(3-methoxypropyl)pyridin-3-yl)benzoic acid
20 (8.3 mg, yield 42%).

[0299]

Step 3-10: Crystals (Form VIII) of 4-(2-(4-methoxyphenyl)-5-(3-methoxypropyl)pyridin-3-yl)benzoic acid

To 4-(2-(4-methoxyphenyl)-5-(3-methoxypropyl)pyridin-3-yl)benzoic acid (600 mg) was added methanol (3.6 mL), and the
25 mixture was stirred at room temperature for 4 days. To this mixture was added a trace amount of Form X, and the mixture was stirred for additional 3 days. The resulting solid was collected by filtration, and dried under reduced pressure to
30 give crystals (Form VIII) of 4-(2-(4-methoxyphenyl)-5-(3-methoxypropyl)pyridin-3-yl)benzoic acid.

[0300]

Step 3-11: Crystals (Form XV) of 4-(2-(4-methoxyphenyl)-5-(3-methoxypropyl)pyridin-3-yl)benzoic acid

35 Form II (15 mg) and Form VIII (15 mg) of 4-(2-(4-


methoxyphenyl)-5-(3-methoxypropyl)pyridin-3-yl)benzoic acid were suspended in a mixed solvent of 1-propanol (0.24 mL) and water (0.24 mL), and the suspension was stirred at room temperature for 11 days. The resulting solid was collected by filtration, and dried under reduced pressure to give crystals (Form XV) of 4-(2-(4-methoxyphenyl)-5-(3-methoxypropyl)pyridin-3-yl)benzoic acid.

[0301]

Production Example 4

10 Synthesis of 4-(2-(4-methoxyphenyl)-5-(3-methoxypropyl)pyridin-3-yl)-2-methylbenzoic acid (Example 116)


[0302]



[0303]

15 Step 4-1: Methyl 4-(2-(4-methoxyphenyl)-5-(3-methoxypropyl)pyridin-3-yl)-2-methylbenzoate

[0304]



[0305]

20 To a solution of 2-(4-methoxyphenyl)-5-(3-methoxypropyl)pyridin-3-yl trifluoromethanesulfonate (2.00 g, 4.93 mmol), which was synthesized by a method similar to that of Step 3-5 in Production Example 3, and (4-(methoxycarbonyl)-3-methylphenyl)boric acid (1.05 g, 5.43 mmol) in 1,2-dimethoxyethane (11.1 mL) were added successively 2M-aqueous potassium phosphate solution (3.70 mL, 7.40 mmol) and [1,1'-bis(diphenylphosphino)-ferrocene]palladium(II) dichloride


dichloromethane adduct (0.081 g, 0.099 mmol), and the mixture was stirred at 80°C for 2 hr. The reaction solution was allowed to cool to room temperature, and water was added thereto. The mixture was extracted with ethyl acetate, and the organic layer was dried over sodium sulfate. The sodium sulfate was removed by filtration, and the filtrate was concentrated under reduced pressure. The residue was purified by silica gel chromatography (hexane:ethyl acetate =3:1 to 1:1) to give methyl 4-(2-(4-methoxyphenyl)-5-(3-methoxypropyl)pyridin-3-yl)-2-methylbenzoate (1.85 g, yield 92%).

¹H-NMR (DMSO-D₆) δ: 1.83-1.90 (2H, m), 2.47 (3H, s), 2.68-2.72 (2H, m), 3.24 (3H, s), 3.36 (2H, t, J = 6.4 Hz), 3.71 (3H, s), 3.80 (3H, s), 6.80 (2H, dt, J = 9.4, 2.5 Hz), 7.01 (1H, dd, J = 8.1, 1.4 Hz), 7.20 (2H, dt, J = 9.4, 2.5 Hz), 7.26 (1H, d, J = 1.4 Hz), 7.63 (1H, d, J = 2.1 Hz), 7.69 (1H, d, J = 8.1 Hz), 8.49 (1H, d, J = 2.1 Hz).

[0306]

Step 4-2: 4-(2-(4-Methoxyphenyl)-5-(3-methoxypropyl)pyridin-3-yl)-2-methylbenzoic acid

[0307]



[0308]

To a solution of methyl 4-(2-(4-methoxyphenyl)-5-(3-methoxypropyl)pyridin-3-yl)-2-methylbenzoate (2.30 g, 5.67 mmol) in methanol (34 mL) was added 4N-aqueous sodium hydroxide solution (8.51 mL, 34.0 mmol), and the mixture was stirred at 50°C for 2 hr. The mixture was allowed to cool to room temperature, 10wt%-aqueous citric acid solution (25.5 mL) and water (55 mL) were added thereto, and the mixture was stirred

at room temperature for 1 hr. The precipitated solid was collected by filtration, and dried under reduced pressure to give 4-(2-(4-methoxyphenyl)-5-(3-methoxypropyl)pyridin-3-yl)-2-methylbenzoic acid (2.05 g, yield 92%).

5 ¹H-NMR (DMSO-D₆) δ: 1.81-1.91 (2H, m), 2.47 (3H, s), 2.70 (2H, t, J = 7.7 Hz), 3.24 (3H, s), 3.36 (2H, t, J = 6.2 Hz), 3.71 (3H, s), 6.81 (2H, d, J = 8.8 Hz), 6.98 (1H, dd, J = 8.1, 1.4 Hz), 7.18-7.24 (3H, m), 7.62 (1H, d, J = 2.3 Hz), 7.69 (1H, d, J = 8.1 Hz), 8.49 (1H, d, J = 2.3 Hz), 12.81 (1H, br s).

10 [0309]

Step 4-3: Crystals of 4-(2-(4-methoxyphenyl)-5-(3-methoxypropyl)pyridin-3-yl)-2-methylbenzoic acid


To 4-(2-(4-methoxyphenyl)-5-(3-methoxypropyl)pyridin-3-yl)-2-methylbenzoic acid (50 mg) was added 2-propanol (0.300
15 mL), and the mixture was stirred at 100°C to give a solution. The stirring was stopped, and the mixture was allowed to cool to room temperature. The precipitated solid was collected by filtration, and dried under reduced pressure to give crystals of 4-(2-(4-methoxyphenyl)-5-(3-methoxypropyl)pyridin-3-yl)-2-
20 methylbenzoic acid (36 mg, yield 72%).

[0310]

Production Example 5

Synthesis of 4-(2-(4-methoxyphenyl)-5-(propoxymethyl)pyridin-3-yl)benzoic acid (Example 118)


25 [0311]



[0312]

Step 5-1: (5-Chloro-6-(4-methoxyphenyl)pyridin-3-yl)methanol

[0313]




[0314]

To a mixture of (5,6-dichloropyridin-3-yl)methanol (200 mg, 1.12 mmol) and (4-methoxyphenyl)boric acid (188 mg, 1.24 mmol) was added toluene (5 mL). Under inert gas atmosphere, [1,1'-bis(diphenylphosphino)-ferrocene]palladium(II) dichloride dichloromethane adduct (45.9 mg, 0.056 mmol) and 2M-aqueous potassium phosphate solution (1.12 mL, 2.24 mmol) were added successively thereto, and the mixture was stirred at 70°C for 1 hr. The mixture was allowed to cool to room temperature, and the reaction solution was diluted with water and ethyl acetate, and separated. The organic layer was washed successively with water and saturated brine, and dried over sodium sulfate. The sodium sulfate was removed by filtration, and the filtrate was concentrated under reduced pressure. The residue was purified by silica gel chromatography (hexane:ethyl acetate =2:1 to 1:2) to give (5-chloro-6-(4-methoxyphenyl)pyridin-3-yl)methanol (250 mg, yield 89%).

[0315]

Step 5-2: 3-Chloro-2-(4-methoxyphenyl)-5-(propoxymethyl)pyridine

[0316]



[0317]


Under inert gas atmosphere, to a suspension of sodium hydride (44.0 mg, 1.10 mmol) in tetrahydrofuran (5 mL) were added successively 1-bromopropane (0.455 mL, 5.01 mmol) and a solution of (5-chloro-6-(4-methoxyphenyl)pyridin-3-yl)methanol (250 mg, 1.00 mmol) in tetrahydrofuran (3 mL), and the mixture was stirred at 100°C for 24 hr. The reaction solution was

allowed to cool to room temperature, diluted with water (20 mL), and extracted with ethyl acetate (20 mL). The organic layer was washed successively with water and saturated brine, and dried over sodium sulfate. The sodium sulfate was removed by filtration, and the filtrate was concentrated under reduced pressure. The residue was purified by silica gel chromatography (hexane:ethyl acetate =3:1 to 1:1) to give 3-chloro-2-(4-methoxyphenyl)-5-(propoxymethyl)pyridine (123 mg, yield 42%).

10 [0318]

Step 5-3: Methyl 4-(2-(4-methoxyphenyl)-5-(propoxymethyl)pyridin-3-yl)benzoate

[0319]



15 [0320]


Under inert gas atmosphere, to a mixture of 3-chloro-2-(4-methoxyphenyl)-5-(propoxymethyl)pyridine (50.0 mg, 0.171 mmol), palladium(II) acetate (3.9 mg, 0.017 mmol), 2-dicyclohexylphosphino-2',6'-dimethoxybiphenyl (14.1 mg, 0.034 mmol) and (4-(methoxycarbonyl)phenyl)boric acid (93.0 mg, 0.514 mmol) were added successively toluene (2 mL) and 2M-aqueous potassium phosphate solution (0.343 mL, 0.685 mmol), and the mixture was stirred at 100°C for 2 hr. The reaction solution was allowed to cool to room temperature, diluted with water (10 mL), and extracted with ethyl acetate (50 mL). The organic layer was washed successively with water and saturated brine, and dried over sodium sulfate. The sodium sulfate was removed by filtration, and the filtrate was concentrated under reduced pressure. The residue was purified by silica gel chromatography (hexane:ethyl acetate =2:1 to 1:2) to give methyl 4-(2-(4-methoxyphenyl)-5-(propoxymethyl)pyridin-3-

yl)benzoate (47.4 mg, yield 70%).

[0321]

Step 5-4: 4-(2-(4-Methoxyphenyl)-5-(propoxymethyl)pyridin-3-yl)benzoic acid

5 [0322]



[0323]

To a solution of methyl 4-(2-(4-methoxyphenyl)-5-(propoxymethyl)pyridin-3-yl)benzoate (47.4 mg, 0.121 mmol) in
10 methanol (0.90 mL) was added 4N-aqueous sodium hydroxide solution (0.182 mL, 0.726 mmol), and the mixture was stirred at 50°C for 2 hr. The reaction solution was allowed to cool to room temperature, 10wt%-aqueous citric acid solution (0.546 mL) and water (2 mL) were added thereto, and the mixture was
15 stirred. The precipitated solid was collected by filtration, and dried under reduced pressure to give 4-(2-(4-methoxyphenyl)-5-(propoxymethyl)pyridin-3-yl)benzoic acid (43.5 mg, yield 95%).


1H-NMR (DMSO-D6) δ : 0.88 (3H, t, J = 7.4 Hz), 1.51-1.62 (2H, m),
20 3.45 (2H, t, J = 6.6 Hz), 3.71 (3H, s), 4.57 (2H, s), 6.81 (2H, d, J = 8.8 Hz), 7.21 (2H, d, J = 8.8 Hz), 7.30 (2H, d, J = 8.3 Hz), 7.71 (1H, d, J = 2.1 Hz), 7.86 (2H, d, J = 8.6 Hz), 8.60 (1H, d, J = 2.1 Hz), 13.01 (1H, br s).

[0324]

25 Production Example 6

Alternative production method of 4-(3-(3,5-difluoro-4-methoxyphenyl)-6-(3-methoxypropyl)pyrazin-2-yl)benzoic acid (Example 67)


[0325]



[0326]

Step 6-1: 2-(tert-Butoxy)-3-chloropyrazine

[0327]



[0328]

Under nitrogen atmosphere, 2,3-dichloropyrazine (150.0 g, 1007 mmol) was dissolved in tetrahydrofuran (450 mL), and a solution of potassium tert-butoxide (129.9 g, 1158 mmol) in tetrahydrofuran (600 mL) was added dropwise thereto under ice-cooling from a dropping funnel. The dropping funnel was washed with tetrahydrofuran (150 mL), and the wash solution was added dropwise to the reaction mixture. The reaction mixture was stirred under ice-cooling for 1 hr, water (450 mL) was added thereto, and the mixture was separated. The organic layer was washed with 10% brine to give a solution of 2-(tert-butoxy)-3-chloropyrazine in tetrahydrofuran. The obtained solution of 2-(tert-butoxy)-3-chloropyrazine in tetrahydrofuran was used in the next step, regarded as yield 100%.


20 The solution of 2-(tert-butoxy)-3-chloropyrazine in tetrahydrofuran was synthesized by the same production method, and concentrated, and the NMR was measured.

¹H-NMR (DMSO-D₆) δ: 1.60 (9H, s), 8.01 (1H, d, J = 2.8 Hz), 8.19 (1H, d, J = 2.8 Hz).

25 [0329]

Step 6-2: Ethyl 4-(3-(tert-butoxy)pyrazin-2-yl)benzoate

[0330]



[0331]

Under nitrogen atmosphere, to 2-(tert-butoxy)-3-chloropyrazine in tetrahydrofuran solution (corresponding to 5 1007 mmol) were added (4-(ethoxycarbonyl)phenyl)boric acid (195.3 g, 1007 mmol) and tetrahydrofuran (150 mL). Then, 2-dicyclohexylphosphino-2',6'-dimethoxybiphenyl (8.27 g, 20.1 mmol) and palladium(II) acetate (2.26 g, 10.1 mmol) were added thereto. The reaction system was degassed under reduced 10 pressure, and replaced with nitrogen. The procedure was repeated three times in total. To this mixture was added dropwise a solution of tripotassium phosphate (363.3 g, 1712 mmol) in water (600 mL) over about 30 min at 40°C. The reaction mixture was stirred at the same temperature for about 15 1 hr, allowed to cool, and separated. The organic layer was washed twice with 10% brine (600 mL). To the organic layer was added activated carbon (15.00 g), and the mixture was stirred at room temperature for 2 hr. The activated carbon was removed by filtration, and washed with tetrahydrofuran (450 mL). The 20 combined filtrate was concentrated under reduced pressure until the volume became 400 mL to give a solution of ethyl 4-(3-(tert-butoxy)pyrazin-2-yl)benzoate in tetrahydrofuran. The obtained solution of ethyl 4-(3-(tert-butoxy)pyrazin-2-yl)benzoate in tetrahydrofuran was used in the next step, 25 regarded as yield 100%.

The solution of ethyl 4-(3-(tert-butoxy)pyrazin-2-yl)benzoate in tetrahydrofuran was synthesized by the same production method, and concentrated to dryness, and the solid was collected by filtration with a mixed solvent of 30 ethanol/water (2/1), and the NMR was measured.


¹H-NMR (DMSO-D₆) δ: 1.35 (3H, t, J = 7.1 Hz), 1.62 (9H, s),

4.35 (2H, q, J = 7.1 Hz), 8.05 (2H, dt, J = 6.8, 2.0 Hz), 8.14 (2H, dt, J = 6.8, 2.0 Hz), 8.22 (1H, d, J = 2.5 Hz), 8.31 (1H, d, J = 2.5 Hz).

[0332]

5 Step 6-3: Ethyl 4-(3-hydroxypyrazin-2-yl)benzoate

[0333]



[0334]


Under nitrogen atmosphere, to ethyl 4-(3-(tert-
10 butoxy)pyrazin-2-yl)benzoate in tetrahydrofuran solution
(corresponding to 1007 mmol) was added ethanol (300 mL), and
then 4N hydrochloric acid (300 mL, 1200 mmol) was added
dropwise thereto at room temperature, and the mixture was
stirred for about 1 hr. To the reaction suspension was added
15 water (750 mL), and the mixture was stirred at room temperature
for 1 hr. Water (750 mL) was added again thereto, and the
mixture was stirred at room temperature for 2 hr. The
precipitated solid was collected by filtration, and the
obtained solid was washed twice with a mixed solvent of
20 water/ethanol (4/1, 300 mL), and dried under reduced pressure
at 60°C to give ethyl 4-(3-hydroxypyrazin-2-yl)benzoate (232.1
g, 950.6 mmol, yield 94.4% from 2,3-dichloropyrazine).

¹H-NMR (DMSO-D₆) δ: 1.34 (3H, t, J = 7.1 Hz), 4.34 (2H, q, J =
7.1 Hz), 7.53 (1H, d, J = 3.7 Hz), 7.55 (1H, d, J = 3.7 Hz),
25 8.02 (2H, dt, J = 8.6, 1.8 Hz), 8.46 (2H, dt, J = 8.6, 1.8 Hz),
12.67 (1H, s).

[0335]

Step 6-4: Ethyl 4-(3-hydroxy-6-iodopyrazin-2-yl)benzoate

[0336]



[0337]


Under nitrogen atmosphere, to ethyl 4-(3-hydroxypyrazin-2-yl)benzoate (100 g, 409mmol) was added acetonitrile (500 mL),
 5 and then, 1,8-diazabicyclo[5.4.0]undec-7-ene (31.2 g, 205 mmol) was added thereto. To this mixture was added dropwise a solution of N-iodosuccinimide (101 g, 450 mmol) in acetonitrile (750 mL) over about 1 hr at room temperature from a dropping funnel. The dropping funnel was washed with acetonitrile (50
 10 mL), the wash solution was added dropwise to the reaction mixture, and the mixture was stirred at room temperature for about 2 hr. To the reaction suspension was added dropwise a solution of sodium sulfite (12.4 g, 123 mmol) in water (600 mL), and the mixture was stirred for 20 min. Then, a solution of
 15 conc. hydrochloric acid (21.3 g, 205 mmol) in water (600 mL) was added dropwise thereto, and the mixture was stirred at 45 to 55°C for 30 min, and then at room temperature for about 30 min. The precipitated solid was collected by filtration, and the obtained solid was washed twice with a mixed solvent of
 20 acetonitrile/water (1/2, 300 mL), and dried under reduced pressure at 50°C to give ethyl 4-(3-hydroxy-6-iodopyrazin-2-yl)benzoate (135 g, yield 89.3%).

¹H-NMR (DMSO-D₆) δ: 1.34 (3H, t, J = 7.1 Hz), 4.35 (2H, q, J = 7.1 Hz), 7.98 (1H, br s), 8.03 (2H, d, J = 8.6 Hz), 8.35 (2H, d, J = 8.6 Hz), 12.85 (1H, s).
 25

[0338]

Step 6-5: Ethyl 4-(3-hydroxy-6-(3-methoxyprop-1-yn-1-yl)pyrazin-2-yl)benzoate

[0339]




[0340]

Under nitrogen atmosphere, to ethyl 4-(3-hydroxy-6-iodopyrazin-2-yl)benzoate (200 g, 540 mmol) was added
 5 acetonitrile (1200 mL), and triethylamine (164 g, 1621 mmol) was added thereto, and then, copper(I) iodide (4.12 g, 21.6 mmol), triphenylphosphine (2.83 g, 10.8mmol) and bis(triphenylphosphine)palladium(II) dichloride (3.79 g, 5.40 mmol) were added thereto. The reaction system was degassed
 10 under reduced pressure, and replaced with nitrogen. The procedure was repeated three times in total. To this mixture was added dropwise a solution of methylpropargyl ether (56.8 g, 810 mmol) in acetonitrile (200 mL) over about 1 hr at 40°C, and the mixture was stirred at the same temperature for about 2 hr.
 15 To the reaction mixture was added acetonitrile (600 mL), and the mixture was concentrated under reduced pressure until the volume became 1000 mL. To the residue was added dropwise acetic acid (64.89 g, 1081 mmol) at 40°C, and the mixture was stirred at the same temperature for 1 hr, and then at room
 20 temperature for an additional 1 hr. The precipitated solid was collected by filtration, and washed with acetonitrile (400 mL). The obtained solid was suspended in acetonitrile (1600 mL), and the suspension was stirred at 70°C for 1 hr, and then at room temperature for 10 hr. The resulting solid was collected by
 25 filtration, washed twice with acetonitrile (400 mL), and dried under reduced pressure at 50°C to give ethyl 4-(3-hydroxy-6-(3-methoxyprop-1-yn-1-yl)pyrazin-2-yl)benzoate (121 g, yield 71.7%).

¹H-NMR (DMSO-D₆) δ: 1.34 (3H, t, J = 7.2 Hz), 3.34 (3H, s),
 30 4.35 (4H, q, J = 7.2 Hz), 7.85 (1H, s), 8.03 (2H, d, J = 8.6 Hz), 8.40 (2H, d, J = 8.6 Hz), 12.96 (1H, s).

[0341]

Step 6-6: Ethyl 4-(3-hydroxy-6-(3-methoxypropyl)pyrazin-2-yl)benzoate
[0342]




5 [0343]

Under nitrogen atmosphere, to ethyl 4-(3-hydroxy-6-(3-methoxyprop-1-yn-1-yl)pyrazin-2-yl)benzoate (30.0 g, 96.1 mmol) was added tetrahydrofuran (360 mL), and 5% palladium on carbon catalyst (50% wet, 1.50 g) was added thereto. The reaction
10 system was replaced with hydrogen, and the mixture was stirred under 0.2 MPa of hydrogen pressure for 4 hr. The 5% palladium on carbon catalyst was removed by filtration, and washed with tetrahydrofuran (120 mL), and the combined filtrate was concentrated under reduced pressure until the volume became 150
15 mL. To the residue was added dropwise heptane (120 mL) at 40°C, and the mixture was stirred at the same temperature for 10 min. Heptane (480 mL) was added dropwise thereto at 45°C, and the mixture was stirred at the same temperature for 30 min, and then at room temperature for 30 min. The precipitated solid
20 was collected by filtration, and the obtained solid was washed with heptane (150 mL), and dried under reduced pressure at 50°C to give ethyl 4-(3-hydroxy-6-(3-methoxypropyl)pyrazin-2-yl)benzoate (28.0 g, yield 92.1%).

¹H-NMR (DMSO-D₆) δ: 1.34 (3H, t, J = 6.9 Hz), 1.84-1.91 (2H, m),
25 2.61 (2H, t, J = 7.6 Hz), 3.24 (3H, s), 3.38 (2H, t, J = 6.4 Hz), 4.34 (2H, q, J = 6.9 Hz), 7.37 (1H, s), 8.02 (2H, d, J = 8.6 Hz), 8.47 (2H, d, J = 8.6 Hz), 12.49 (1H, s).

[0344]

Step 6-7: Ethyl 4-(6-(3-methoxypropyl)-3-((trifluoromethyl)sulfonyl)oxy)pyrazin-2-yl)benzoate
30 [0345]



[0346]

Under nitrogen atmosphere, to a solution of dipotassium hydrogenphosphate (42.1 g, 242 mmol) in water (85 mL) was added
 5 toluene (153 mL), and ethyl 4-(3-hydroxy-6-(3-methoxypropyl)pyrazin-2-yl)benzoate (17.0 g, 53.7 mmol) was added thereto. To this mixture was added dropwise trifluoromethanesulfonic anhydride (22.7 g, 80.5 mmol) over 1 hr at 5 to 10°C, and the mixture was stirred at the same
 10 temperature for 1 hr. The reaction mixture was separated, and the organic layer was washed with 20% brine (68 g) to give a solution of ethyl 4-(6-(3-methoxypropyl)-3-((trifluoromethyl)sulfonyl)oxy)pyrazin-2-yl)benzoate in toluene. The obtained solution of ethyl 4-(6-(3-methoxypropyl)-3-
 15 methoxypropyl)-3-((trifluoromethyl)sulfonyl)oxy)pyrazin-2-yl)benzoate in toluene was used in the next step, regarded as yield 100%.


The solution of ethyl 4-(6-(3-methoxypropyl)-3-((trifluoromethyl)sulfonyl)oxy)pyrazin-2-yl)benzoate in
 20 toluene was synthesized by the same production method, and concentrated to dryness, and the residue was purified by silica gel chromatography (hexane:ethyl acetate =4:1), and the NMR was measured.

¹H-NMR (DMSO-D₆) δ: 1.36 (3H, t, J = 7.2 Hz), 1.97-2.04 (2H, m),
 25 3.00 (2H, t, J = 7.7 Hz), 3.23 (3H, s), 3.41 (2H, t, J = 6.2 Hz), 4.37 (2H, q, J = 7.2 Hz), 8.00 (2H, dt, J = 8.6, 1.8 Hz), 8.15 (2H, dt, J = 8.6, 1.8 Hz), 8.53 (1H, s).

[0347]

Step 6-8: Ethyl 4-(3-(3,5-difluoro-4-methoxyphenyl)-6-(3-methoxypropyl)pyrazin-2-yl)benzoate
 30

[0348]



[0349]

Under nitrogen atmosphere, to a solution of ethyl 4-(6-
5 (3-methoxypropyl)-3-(((trifluoromethyl)sulfonyl)oxy)pyrazin-2-yl)benzoate in toluene (corresponding to 53.7 mmol) was added tetrahydrofuran (34 mL), and (3,5-difluoro-4-methoxyphenyl)boric acid (12.1 g, 64.5 mmol) was added thereto, and the used container was washed with tetrahydrofuran (17 mL).
10 Bis(triphenylphosphine)palladium(II) dichloride (0.377 g, 0.537 mmol) was added thereto, and the reaction system was degassed under reduced pressure, and replaced with nitrogen. To this mixture was added dropwise a solution of tripotassium phosphate (13.7 g, 64.5 mmol) in water (65 mL) over 2 hr at 70°C, and the
15 mixture was stirred at the same temperature for 2 hr. The reaction mixture was separated, and the organic layer was washed twice with 20% brine (68 g), and concentrated under reduced pressure until the volume became 68 mL. To the residue was added toluene until the volume became 136 mL, activated
20 carbon (3.4 g) and metal scavenger (Fuji Silysia Chemical Ltd, SCAVENGER SH SILICA, 1.0 g) were added thereto, and the mixture was stirred at room temperature for 2 hr. The activated carbon and metal scavenger were removed by filtration, and washed with toluene (51 mL). The combined filtrate was concentrated under
25 reduced pressure, to the residue was added 2-propanol (102 mL), and the mixture was concentrated under reduced pressure. The procedure was repeated twice in total. To the residue was added 2-propanol until the volume became 85 mL, and the mixture was stirred at room temperature for 1 hr, and then under ice-
30 cooling for 2 hr. The precipitated solid was collected by filtration, washed with cooled 2-propanol (51 mL), and dried under reduced pressure at 50°C to give ethyl 4-(3-(3,5-


difluoro-4-methoxyphenyl)-6-(3-methoxypropyl)pyrazin-2-yl)benzoate (21.9 g, yield 92.1%).

¹H-NMR (DMSO-D₆) δ: 1.33 (3H, t, J = 7.1 Hz), 1.96-2.03 (2H, m), 2.91-2.95 (2H, m), 3.25 (3H, s), 3.42 (2H, t, J = 6.4 Hz), 3.95
5 (3H, s), 4.33 (2H, q, J = 7.1 Hz), 7.08-7.14 (2H, m), 7.57 (2H, dt, J = 8.5, 1.8 Hz), 7.96 (2H, dt, J = 8.5, 1.8 Hz), 8.66 (1H, s).

[0350]

Step 6-9: 4-(3-(3,5-Difluoro-4-methoxyphenyl)-6-(3-
10 methoxypropyl)pyrazin-2-yl)benzoic acid

[0351]



[0352]

Under nitrogen atmosphere, to ethyl 4-(3-(3,5-difluoro-4-
15 methoxyphenyl)-6-(3-methoxypropyl)pyrazin-2-yl)benzoate (5.0 g, 11.3 mmol) was added ethanol (15 mL), and 2N aqueous sodium hydroxide solution (7.5 mL, 15.0 mmol) was added thereto, and the mixture was stirred at 40°C for 1.5 hr. The reaction mixture was filtered through 0.45 μm membrane filter, and
20 washed with a mixed solvent of ethanol/water (3/1.4, 22 mL). To the combined filtrate was added dropwise 3N hydrochloric acid (5.5 mL, 16.5 mmol) at room temperature, and the mixture was stirred at room temperature for 0.5 hr. To this mixture was added dropwise water (10 mL) at 35°C, and the mixture was
25 stirred at the same temperature for 30 min, and then at room temperature for about 2 hr. The precipitated solid was collected by filtration, and the obtained solid was washed successively with a mixed solvent of ethanol/water (1/2, 22.5 mL) and water (30 mL), and dried under reduced pressure at 50°C
30 to give 4-(3-(3,5-difluoro-4-methoxyphenyl)-6-(3-

methoxypropyl)pyrazin-2-yl)benzoic acid (4.45 g, yield 95.1%).
1H-NMR (DMSO-D6) δ : 1.96-2.03 (2H, m), 2.93 (2H, t, J = 7.7 Hz),
3.25 (3H, s), 3.42 (2H, t, J = 6.4 Hz), 3.95 (3H, s), 7.07-7.14
(2H, m), 7.54 (2H, d, J = 8.3 Hz), 7.94 (2H, d, J = 8.3 Hz),
5 8.66 (1H, s), 13.11 (1H, s).









[0353]

The compounds of the other Examples were obtained
according to the above-mentioned general production methods or
by a method similar to that of the Production Example, using
10 the other known methods as necessary. The structural formulas
and property data of the compounds of Examples 1 to 153 are
shown in the following Table 1-1 to Table 1-20. The MS value
marked with "-COOH" in the tables means a value of fragment
after decarboxylation.

15









[0354]

Table 1-1

Ex.	Structure	NMR	MS (M+H)	MS (M-H)
1		¹ H-NMR (DMSO-D ₆) δ: 0.92 (3H, t, J = 7.4 Hz), 1.38 (2H, td, J = 14.8, 7.5 Hz), 1.68-1.77 (2H, m), 2.85 (2H, t, J = 7.7 Hz), 3.74 (3H, s), 6.87 (2H, dt, J = 9.4, 2.5 Hz), 7.29 (2H, dt, J = 9.4, 2.5 Hz), 7.49 (2H, dt, J = 8.4, 1.8 Hz), 7.88 (2H, dt, J = 8.5, 1.8 Hz), 8.59 (1H, t, J = 4.9 Hz), 13.01 (1H, br s).	363	361
2		¹ H-NMR (DMSO-D ₆) δ: 0.93 (3H, t, J = 7.3 Hz), 1.35-1.44 (2H, m), 1.71-1.79 (2H, m), 1.92 (3H, s), 2.88 (2H, t, J = 7.7 Hz), 3.73 (3H, s), 6.72 (1H, dd, J = 8.3, 2.5 Hz), 6.76 (1H, d, J = 2.5 Hz), 7.04 (1H, d, J = 8.6 Hz), 7.42 (2H, dd, J = 6.7, 1.8 Hz), 7.81 (2H, dd, J = 6.7, 1.8 Hz), 8.61 (1H, s), 13.00 (1H, br s).	377	375
3		¹ H-NMR (DMSO-D ₆) δ: 0.93 (3H, t, J = 7.3 Hz), 1.34-1.43 (2H, m), 1.70-1.78 (2H, m), 2.88 (2H, t, J = 7.7 Hz), 3.76 (3H, s), 6.71 (1H, dd, J = 12.3, 2.5 Hz), 6.87 (1H, dd, J = 8.7, 2.4 Hz), 7.43-7.51 (3H, m), 7.84 (2H, dd, J = 6.7, 1.8 Hz), 8.64 (1H, s), 13.04 (1H, br s).	381	379
4		¹ H-NMR (DMSO-D ₆) δ: 0.90 (3H, t, J = 7.3 Hz), 1.35 (2H, td, J = 14.9, 7.4 Hz), 1.70 (2H, dt, J = 15.8, 7.0 Hz), 1.98 (3H, s), 2.84 (2H, t, J = 7.6 Hz), 3.70 (3H, s), 6.78-6.84 (2H, m), 7.21-7.29 (3H, m), 7.74 (1H, dd, J = 7.9, 1.2 Hz), 7.78 (1H, s), 8.62 (1H, s), 12.97 (1H, br s).	377	375
5		¹ H-NMR (DMSO-D ₆) δ: 0.91 (3H, t, J = 7.4 Hz), 1.31-1.43 (2H, m), 1.65-1.76 (2H, m), 2.85 (2H, t, J = 7.7 Hz), 3.72 (3H, s), 6.85 (2H, d, J = 8.6 Hz), 7.29 (2H, d, J = 8.6 Hz), 7.56 (1H, dd, J = 10.4, 1.4 Hz), 7.68 (1H, t, J = 7.6 Hz), 7.85 (1H, dd, J = 7.9, 1.2 Hz), 8.66 (1H, s), 13.37 (1H, br s).	381	379
6		¹ H-NMR (DMSO-D ₆) δ: 0.93 (3H, t, J = 7.3 Hz), 1.32-1.45 (2H, m), 1.67-1.79 (2H, m), 2.85 (2H, t, J = 7.6 Hz), 3.58 (3H, s), 3.74 (3H, s), 6.90 (2H, dt, J = 9.4, 2.5 Hz), 7.00-7.06 (2H, m), 7.32 (2H, dt, J = 9.3, 2.5 Hz), 7.57 (1H, d, J = 7.9 Hz), 8.58 (1H, s), 12.66 (1H, br s).	393	391
7		¹ H-NMR (DMSO-D ₆) δ: 0.93 (3H, t, J = 7.4 Hz), 1.34-1.45 (2H, m), 1.69-1.80 (2H, m), 2.89 (2H, t, J = 7.6 Hz), 7.45-7.50 (4H, m), 7.84-7.89 (4H, m), 8.67 (1H, s), 13.03 (2H, br s).	377	375
8		¹ H-NMR (DMSO-D ₆) δ: 0.92 (3H, t, J = 7.4 Hz), 1.32-1.43 (2H, m), 1.67-1.77 (2H, m), 2.07 (3H, s), 2.84 (2H, t, J = 7.7 Hz), 3.75 (3H, s), 6.81 (1H, d, J = 8.6 Hz), 6.99-7.07 (1H, m), 7.28 (1H, dd, J = 2.3, 0.7 Hz), 7.49 (2H, dt, J = 8.4, 1.8 Hz), 7.88 (2H, dt, J = 8.4, 1.7 Hz), 8.57 (1H, s), 13.02 (1H, br s).	377	375

[0355]

Table 1-2

Ex.	Structure	NMR	MS (M+H)	MS (M-H)
9		¹ H-NMR (DMSO-D ₆) δ: 0.92 (3H, t, J = 7.4 Hz), 1.33-1.45 (2H, m), 1.69-1.79 (2H, m), 2.89 (2H, t, J = 7.7 Hz), 7.48 (2H, d, J = 8.6 Hz), 7.54 (2H, d, J = 8.1 Hz), 7.80 (2H, d, J = 8.1 Hz), 7.88 (2H, d, J = 8.6 Hz), 8.69 (1H, s), 13.08 (1H, s).	358	356
10		¹ H-NMR (DMSO-D ₆) δ: 0.92 (3H, t, J = 7.3 Hz), 1.33-1.44 (2H, m), 1.68-1.78 (2H, m), 2.87 (2H, t, J = 7.7 Hz), 7.16 (2H, t, J = 8.9 Hz), 7.39 (2H, dd, J = 8.8, 5.5 Hz), 7.48 (2H, d, J = 8.6 Hz), 7.88 (2H, d, J = 8.6 Hz), 8.63 (1H, s), 13.05 (1H, br s).	351	349
11		¹ H-NMR (DMSO-D ₆) δ: 0.92 (3H, t, J = 7.3 Hz), 1.31-1.44 (2H, m), 1.66-1.78 (2H, m), 2.86 (2H, t, J = 7.7 Hz), 3.82 (3H, s), 7.03-7.12 (2H, m), 7.19-7.26 (1H, m), 7.51 (2H, d, J = 8.6 Hz), 7.90 (2H, d, J = 8.6 Hz), 8.61 (1H, s), 13.05 (1H, br s).	381	379
12		¹ H-NMR (DMSO-D ₆) δ: 1.61-1.68 (2H, m), 1.69-1.76 (2H, m), 2.31 (2H, t, J = 6.2 Hz), 2.80 (2H, t, J = 6.1 Hz), 3.66 (3H, s), 6.70 (2H, dt, J = 9.4, 2.5 Hz), 7.09 (2H, dt, J = 9.5, 2.5 Hz), 7.21 (2H, dt, J = 8.2, 1.8 Hz), 7.86 (2H, dt, J = 8.3, 1.8 Hz), 8.36 (1H, s), 12.97 (1H, br s).	360	358
13		¹ H-NMR (DMSO-D ₆) δ: 0.93 (3H, t, J = 7.3 Hz), 1.34-1.45 (2H, m), 1.69-1.80 (2H, m), 2.89 (2H, t, J = 7.7 Hz), 7.49 (2H, d, J = 8.6 Hz), 7.58 (2H, d, J = 8.1 Hz), 7.69 (2H, d, J = 8.8 Hz), 7.89 (2H, d, J = 8.6 Hz), 8.69 (1H, s), 13.09 (1H, br s).	401	399
14		¹ H-NMR (DMSO-D ₆) δ: 0.92 (3H, t, J = 7.3 Hz), 1.32-1.44 (2H, m), 1.67-1.77 (2H, m), 2.85 (2H, t, J = 7.6 Hz), 3.75 (3H, s), 6.90 (2H, d, J = 8.8 Hz), 7.22-7.35 (4H, m), 7.78 (1H, t, J = 7.9 Hz), 8.61 (1H, s), 13.32 (1H, br s).	381	379
15		¹ H-NMR (DMSO-D ₆) δ: 0.92 (3H, t, J = 7.4 Hz), 1.33-1.44 (2H, m), 1.67-1.78 (2H, m), 2.86 (2H, t, J = 7.7 Hz), 3.79 (3H, s), 7.02 (1H, d, J = 9.0 Hz), 7.37 (1H, dd, J = 8.7, 2.4 Hz), 7.51 (2H, d, J = 8.6 Hz), 7.78 (1H, d, J = 2.5 Hz), 7.89 (2H, d, J = 8.6 Hz), 8.61 (1H, s), 12.81 (1H, br s).	407	405
16		¹ H-NMR (DMSO-D ₆) δ: 0.92 (3H, t, J = 7.4 Hz), 1.31-1.45 (2H, m), 1.66-1.78 (2H, m), 2.86 (2H, t, J = 7.7 Hz), 3.83 (3H, s), 7.05 (1H, d, J = 8.8 Hz), 7.19 (1H, dd, J = 8.6, 2.1 Hz), 7.45-7.54 (3H, m), 7.90 (2H, d, J = 8.6 Hz), 8.61 (1H, s), 13.09 (1H, br s).	397	395

[0356]

Table 1-3

Ex.	Structure	NMR	MS (M+H)	MS (M-H)
17		¹ H-NMR (DMSO-D ₆) δ: 0.92 (3H, t, J = 7.4 Hz), 1.33-1.44 (2H, m), 1.68-1.77 (2H, m), 2.87 (2H, t, J = 7.7 Hz), 3.90 (3H, s), 7.18 (1H, d, J = 9.0 Hz), 7.50 (2H, d, J = 8.6 Hz), 7.56 (1H, dd, J = 8.8, 2.3 Hz), 7.73 (1H, d, J = 2.1 Hz), 7.91 (2H, d, J = 6.7 Hz), 8.63 (1H, s), 13.09 (1H, br s).	388	386
18		¹ H-NMR (DMSO-D ₆) δ: 0.91 (3H, t, J = 7.3 Hz), 1.30-1.40 (2H, m), 1.58-1.66 (2H, m), 2.67 (2H, t, J = 7.7 Hz), 3.71 (3H, s), 6.79 (2H, dt, J = 9.4, 2.5 Hz), 7.19 (2H, dt, J = 9.5, 2.5 Hz), 7.30 (2H, dt, J = 8.4, 1.8 Hz), 7.62 (1H, d, J = 2.3 Hz), 7.85 (2H, dt, J = 8.6, 1.8 Hz), 8.50 (1H, d, J = 2.1 Hz), 12.93 (1H, br s).	362	360
19		¹ H-NMR (DMSO-D ₆) δ: 0.92 (3H, t, J = 7.3 Hz), 1.32-1.44 (2H, m), 1.66-1.79 (2H, m), 2.87 (2H, t, J = 7.6 Hz), 3.87 (3H, s), 7.18 (1H, d, J = 8.8 Hz), 7.48-7.53 (3H, m), 7.66 (1H, d, J = 2.3 Hz), 7.91 (2H, d, J = 8.6 Hz), 8.63 (1H, s), 13.06 (1H, br s).	431	429
20		¹ H-NMR (DMSO-D ₆) δ: 1.56-1.63 (2H, m), 1.75-1.83 (2H, m), 2.87 (2H, t, J = 7.6 Hz), 3.22 (3H, s), 3.36 (2H, t, J = 6.3 Hz), 3.75 (3H, s), 6.88 (2H, d, J = 8.7 Hz), 7.31 (2H, d, J = 8.7 Hz), 7.51 (2H, d, J = 7.8 Hz), 7.90 (2H, d, J = 8.1 Hz), 8.60 (1H, s), 13.05 (1H, br s).	393	391
21		¹ H-NMR (DMSO-D ₆) δ: 0.92 (3H, t, J = 7.4 Hz), 1.31-1.44 (2H, m), 1.66-1.78 (2H, m), 2.87 (2H, t, J = 7.7 Hz), 3.93 (3H, s), 7.08 (2H, d, J = 9.2 Hz), 7.51 (2H, d, J = 8.3 Hz), 7.92 (2H, d, J = 8.3 Hz), 8.64 (1H, s), 13.11 (1H, br s).	399	397
22		¹ H-NMR (DMSO-D ₆) δ: 0.93 (3H, t, J = 7.4 Hz), 1.33-1.45 (2H, m), 1.68-1.79 (2H, m), 2.80-3.00 (8H, m), 7.33 (2H, d, J = 8.3 Hz), 7.41 (2H, d, J = 8.3 Hz), 7.49 (2H, d, J = 8.3 Hz), 7.87 (2H, d, J = 8.3 Hz), 8.66 (1H, s), 13.04 (1H, br s).	404	402
23		¹ H-NMR (DMSO-D ₆) δ: 0.93 (3H, t, J = 7.4 Hz), 1.32-1.46 (2H, m), 1.68-1.80 (2H, m), 2.90 (2H, t, J = 7.6 Hz), 3.67 (3H, s), 7.03 (1H, d, J = 8.1 Hz), 7.24 (1H, s), 7.50 (2H, d, J = 8.3 Hz), 7.54 (1H, d, J = 8.1 Hz), 7.90 (2H, d, J = 8.3 Hz), 8.69 (1H, s), 13.11 (1H, br s).	431	429
24		¹ H-NMR (DMSO-D ₆) δ: 0.93 (3H, t, J = 7.4 Hz), 1.33-1.45 (2H, m), 1.68-1.80 (2H, m), 2.90 (2H, t, J = 7.6 Hz), 7.31 (1H, d, J = 8.1 Hz), 7.47-7.57 (3H, m), 7.72 (1H, t, J = 7.9 Hz), 7.91 (2H, d, J = 8.3 Hz), 8.70 (1H, s), 13.08 (1H, br s).	419	417









[0357]

Table 1-4

Ex.	Structure	NMR	MS (M+H)	MS (M-H)
25		¹ H-NMR (DMSO-D ₆) δ: 0.91 (3H, t, J = 7.3 Hz), 1.31-1.39 (2H, m), 1.57-1.64 (2H, m), 2.68 (2H, t, J = 7.6 Hz), 3.69 (3H, s), 6.76-6.79 (2H, m), 7.16 (2H, dt, J = 9.4, 2.5 Hz), 7.32 (2H, d, J = 8.1 Hz), 7.89 (2H, d, J = 8.1 Hz), 8.57 (1H, d, J = 9.9 Hz), 13.03 (1H, br s).	378	380
26		¹ H-NMR (DMSO-D ₆) δ: 0.93 (3H, t, J = 7.3 Hz), 1.33-1.44 (2H, m), 1.69-1.79 (2H, m), 2.38 (3H, s), 2.89 (2H, t, J = 7.7 Hz), 7.23 (1H, d, J = 8.3 Hz), 7.50 (2H, d, J = 8.3 Hz), 7.56 (2H, d, J = 8.3 Hz), 7.89 (2H, d, J = 8.3 Hz), 8.68 (1H, s), 13.07 (1H, br s).	415	413
27		¹ H-NMR (DMSO-D ₆) δ: 0.93 (3H, t, J = 7.4 Hz), 1.33-1.44 (2H, m), 1.69-1.80 (2H, m), 2.90 (2H, t, J = 7.6 Hz), 7.42 (1H, d, J = 8.3 Hz), 7.51 (2H, d, J = 8.3 Hz), 7.75 (1H, s), 7.78 (1H, d, J = 8.3 Hz), 7.91 (2H, d, J = 8.1 Hz), 8.70 (1H, s), 13.13 (1H, br s).	435	433
28		¹ H-NMR (DMSO-D ₆) δ: 0.93 (3H, t, J = 7.4 Hz), 1.34-1.46 (2H, m), 1.69-1.80 (2H, m), 2.92 (2H, t, J = 7.7 Hz), 7.53 (2H, d, J = 8.6 Hz), 7.83 (1H, d, J = 8.6 Hz), 7.92 (2H, d, J = 8.6 Hz), 7.96-8.03 (2H, m), 8.74 (1H, s), 13.09 (1H, br s).	469	467
29		¹ H-NMR (DMSO-D ₆) δ: 1.56-1.63 (2H, m), 1.76-1.84 (2H, m), 2.91 (2H, t, J = 7.6 Hz), 3.20 (3H, s), 3.35 (2H, t, J = 6.4 Hz), 7.50 (2H, dd, J = 6.7, 1.8 Hz), 7.58 (2H, d, J = 8.1 Hz), 7.70 (2H, d, J = 8.1 Hz), 7.89 (2H, dd, J = 6.7, 1.8 Hz), 8.69 (1H, s), 13.07 (1H, br s).	429	431
30		¹ H-NMR (DMSO-D ₆) δ: 0.91 (3H, t, J = 7.4 Hz), 1.30-1.40 (2H, m), 1.58-1.66 (2H, m), 2.67 (2H, t, J = 7.7 Hz), 3.79 (3H, s), 6.93-6.96 (1H, m), 7.01 (1H, t, J = 8.7 Hz), 7.11 (1H, dd, J = 12.7, 2.1 Hz), 7.32 (2H, dt, J = 8.4, 1.8 Hz), 7.65 (1H, d, J = 2.3 Hz), 7.88 (2H, dt, J = 8.5, 1.8 Hz), 8.51 (1H, d, J = 2.1 Hz), 13.00 (1H, br s).	378	380
31		¹ H-NMR (DMSO-D ₆) δ: 0.94 (3H, t, J = 7.4 Hz), 1.35-1.45 (2H, m), 1.79-1.71 (2H, m), 2.89 (2H, t, J = 7.7 Hz), 7.37-7.42 (4H, m), 7.50 (2H, dt, J = 8.4, 1.8 Hz), 7.91 (2H, dt, J = 8.4, 1.8 Hz), 8.66 (1H, s), 13.07 (1H, br s).	367	365
32		¹ H-NMR (DMSO-D ₆) δ: 0.93 (3H, t, J = 7.3 Hz), 1.35-1.44 (2H, m), 1.77-1.70 (2H, m), 2.85 (2H, t, J = 7.7 Hz), 3.13 (2H, t, J = 8.7 Hz), 4.54 (2H, t, J = 8.7 Hz), 6.65 (1H, d, J = 8.3 Hz), 6.99 (1H, dd, J = 8.3, 2.0 Hz), 7.35 (1H, d, J = 1.4 Hz), 7.52 (2H, dt, J = 8.4, 1.8 Hz), 7.90 (2H, dt, J = 8.5, 1.8 Hz), 8.58 (1H, s), 13.02 (1H, s).	375	373

[0358]

Table 1-5

Ex.	Structure	NMR	MS (M+H)	MS (M-H)
33		¹ H-NMR (DMSO-D ₆) δ: 0.93 (3H, t, J = 7.4 Hz), 1.34-1.44 (2H, m), 1.69-1.77 (2H, m), 2.85 (2H, t, J = 7.6 Hz), 6.69 (2H, dt, J = 9.2, 2.4 Hz), 7.19 (2H, dt, J = 9.2, 2.4 Hz), 7.50 (2H, dt, J = 8.4, 1.7 Hz), 7.89 (2H, dt, J = 8.4, 1.7 Hz), 8.57 (1H, s), 9.69 (1H, s), 13.03 (1H, br s).	349	347
34		¹ H-NMR (DMSO-D ₆) δ: 0.66-0.69 (2H, m), 0.92-0.98 (5H, m), 1.44-1.35 (2H, m), 1.70-1.78 (2H, m), 1.87-1.92 (1H, m), 2.87 (2H, t, J = 7.6 Hz), 7.02 (2H, dt, J = 8.5, 1.8 Hz), 7.25 (2H, dt, J = 8.4, 1.8 Hz), 7.50 (2H, dt, J = 8.4, 1.8 Hz), 7.89 (2H, dt, J = 8.5, 1.8 Hz), 8.62 (1H, s), 13.04 (1H, s).	373	371
35		¹ H-NMR (DMSO-D ₆) δ: 1.54-1.63 (2H, m), 1.72-1.83 (2H, m), 2.88 (2H, t, J = 7.6 Hz), 3.20 (3H, s), 3.34 (2H, t, J = 6.5 Hz), 3.93 (3H, s), 7.09 (2H, d, J = 9.5 Hz), 7.52 (2H, d, J = 8.6 Hz), 7.92 (2H, d, J = 8.6 Hz), 8.64 (1H, s), 13.10 (1H, br s).	429	427
36		¹ H-NMR (DMSO-D ₆) δ: 0.98 (3H, t, J = 7.4 Hz), 1.73-1.85 (2H, m), 2.87 (2H, t, J = 7.5 Hz), 7.49 (2H, d, J = 8.6 Hz), 7.58 (2H, d, J = 8.1 Hz), 7.70 (2H, d, J = 8.3 Hz), 7.89 (2H, d, J = 8.6 Hz), 8.69 (1H, s), 13.08 (1H, br s).	387	385
37		¹ H-NMR (DMSO-D ₆) δ: 0.96 (3H, t, J = 7.3 Hz), 1.70-1.82 (2H, m), 2.82 (2H, t, J = 7.6 Hz), 3.74 (3H, s), 6.87 (2H, d, J = 8.8 Hz), 7.30 (2H, d, J = 8.8 Hz), 7.49 (2H, d, J = 8.6 Hz), 7.88 (2H, d, J = 8.8 Hz), 8.58 (1H, s), 13.04 (1H, br s).	349	347
38		¹ H-NMR (DMSO-D ₆) δ: 0.93 (3H, t, J = 7.2 Hz), 1.34-1.44 (2H, m), 1.70-1.77 (2H, m), 2.84 (2H, t, J = 7.5 Hz), 3.58 (2H, s), 3.75 (3H, s), 6.88 (2H, d, J = 8.7 Hz), 7.22 (2H, d, J = 7.8 Hz), 7.31-7.36 (4H, m), 8.54 (1H, s), 12.35 (1H, s).	377	331 (-COOH)
39		¹ H-NMR (DMSO-D ₆) δ: 0.95 (3H, t, J = 7.3 Hz), 1.36-1.46 (2H, m), 1.53-1.61 (2H, m), 2.02 (3H, s), 2.68 (2H, t, J = 7.6 Hz), 3.68 (3H, s), 6.71 (2H, d, J = 8.7 Hz), 7.09 (2H, d, J = 8.7 Hz), 7.23 (2H, d, J = 8.1 Hz), 7.88 (2H, d, J = 8.1 Hz), 8.41 (1H, s), 12.98 (1H, br s).	376	374
40		¹ H-NMR (DMSO-D ₆) δ: 0.94 (3H, t, J = 7.4 Hz), 1.35-1.44 (2H, m), 1.70-1.78 (2H, m), 2.30 (3H, s), 2.87 (2H, t, J = 7.7 Hz), 7.13 (2H, d, J = 8.1 Hz), 7.26 (2H, d, J = 8.1 Hz), 7.49 (2H, d, J = 8.6 Hz), 7.88 (2H, dt, J = 8.6, 1.7 Hz), 8.63 (1H, s), 13.04 (1H, s).	347	345

[0359]

Table 1-6

Ex.	Structure	NMR	MS (M+H)	MS (M-H)
41		¹ H-NMR (DMSO-D ₆) δ: 0.93 (3H, t, J = 7.3 Hz), 1.61-1.71 (2H, m), 2.64 (2H, t, J = 9.7 Hz), 3.71 (3H, s), 6.79 (2H, dd, J = 6.7, 2.1 Hz), 7.19 (2H, dd, J = 6.8, 2.2 Hz), 7.30 (2H, dd, J = 6.7, 1.8 Hz), 7.62 (1H, d, J = 2.3 Hz), 7.85 (2H, dd, J = 6.7, 1.8 Hz), 8.50 (1H, d, J = 2.3 Hz), 12.97 (1H, br s).	346	348
42		¹ H-NMR (DMSO-D ₆) δ: 0.86-1.00 (2H, m), 1.08-1.35 (4H, m), 1.47-1.81 (7H, m), 2.67 (2H, t, J = 8.0 Hz), 3.71 (3H, s), 6.79 (2H, d, J = 9.0 Hz), 7.18 (2H, d, J = 8.8 Hz), 7.29 (2H, d, J = 8.6 Hz), 7.62 (1H, d, J = 2.1 Hz), 7.85 (2H, d, J = 8.6 Hz), 8.49 (1H, d, J = 2.1 Hz), 13.00 (1H, br s).	416	414
43		¹ H-NMR (DMSO-D ₆) δ: 1.81-1.92 (2H, m), 2.72 (2H, t, J = 7.6 Hz), 3.36-3.53 (6H, m), 3.71 (3H, s), 4.56 (1H, t, J = 5.3 Hz), 6.80 (2H, d, J = 9.0 Hz), 7.19 (2H, d, J = 8.8 Hz), 7.30 (2H, d, J = 8.3 Hz), 7.64 (1H, d, J = 2.1 Hz), 7.85 (2H, d, J = 8.6 Hz), 8.51 (1H, d, J = 2.1 Hz), 12.97 (1H, br s).	408	406
44		¹ H-NMR (DMSO-D ₆) δ: 1.32-1.47 (1H, m), 1.49-1.74 (5H, m), 1.89 (4H, dd, J = 8.8, 6.5 Hz), 2.68 (2H, t, J = 7.5 Hz), 3.71 (3H, s), 4.75 (1H, s), 6.80 (2H, d, J = 9.0 Hz), 7.19 (2H, d, J = 8.8 Hz), 7.30 (2H, d, J = 8.6 Hz), 7.63 (1H, d, J = 2.3 Hz), 7.85 (2H, d, J = 8.6 Hz), 8.51 (1H, d, J = 2.1 Hz), 12.96 (1H, br s).	418	416
45		¹ H-NMR (DMSO-D ₆) δ: 0.87 (3H, t, J = 6.9 Hz), 1.28-1.36 (4H, m), 1.60-1.68 (2H, m), 2.66 (2H, t, J = 7.7 Hz), 3.71 (3H, s), 6.79 (2H, dt, J = 9.4, 2.5 Hz), 7.19 (2H, dt, J = 9.3, 2.5 Hz), 7.30 (2H, dd, J = 6.6, 1.7 Hz), 7.62 (1H, d, J = 2.1 Hz), 7.85 (2H, dd, J = 6.7, 1.8 Hz), 8.50 (1H, d, J = 2.1 Hz), 12.97 (1H, br s).	374	376
46		¹ H-NMR (DMSO-D ₆) δ: 1.17-1.58 (5H, m), 1.65-1.90 (5H, m), 2.58-2.70 (1H, m), 3.71 (3H, s), 6.79 (2H, d, J = 8.8 Hz), 7.18 (2H, d, J = 8.8 Hz), 7.30 (2H, d, J = 8.6 Hz), 7.62 (1H, d, J = 2.3 Hz), 7.85 (2H, d, J = 8.6 Hz), 8.53 (1H, d, J = 2.1 Hz), 13.00 (1H, s).	388	386
47		¹ H-NMR (DMSO-D ₆) δ: 0.92-1.22 (5H, m), 1.55-1.67 (6H, m), 2.55 (2H, d, J = 6.9 Hz), 3.71 (3H, s), 6.79 (2H, dt, J = 9.5, 2.4 Hz), 7.19 (2H, dt, J = 9.4, 2.5 Hz), 7.26 (2H, d, J = 8.3 Hz), 7.57 (1H, d, J = 2.1 Hz), 7.84 (2H, d, J = 8.3 Hz), 8.44 (1H, d, J = 2.1 Hz), 13.13 (1H, br s).	400	402
48		¹ H-NMR (DMSO-D ₆) δ: 0.93 (3H, t, J = 7.9 Hz), 1.34-1.44 (2H, m), 1.47 (6H, s), 1.69-1.77 (2H, m), 2.83 (2H, t, J = 7.6 Hz), 3.76 (3H, s), 6.88 (2H, d, J = 9.0 Hz), 7.30-7.38 (6H, m), 8.53 (1H, s), 12.36 (1H, s).	405	359 (-COOH)









[0360]

Table 1-7

Ex.	Structure	NMR	MS (M+H)	MS (M-H)
49		¹ H-NMR (DMSO-D ₆) δ: 0.93 (3H, t, J = 7.5 Hz), 1.34-1.44 (2H, m), 1.69-1.77 (2H, m), 1.86-1.93 (2H, m), 2.66 (2H, t, J = 6.3 Hz), 2.85 (2H, t, J = 7.6 Hz), 4.14 (2H, t, J = 4.9 Hz), 6.60 (1H, d, J = 8.4 Hz), 6.92 (1H, dd, J = 8.5, 1.9 Hz), 7.23 (1H, d, J = 1.9 Hz), 7.52 (2H, d, J = 8.6 Hz), 7.91 (2H, d, J = 8.6 Hz), 8.58 (1H, s), 13.05 (1H, br s).	389	387
50		¹ H-NMR (DMSO-D ₆) δ: 0.93 (3H, t, J = 7.3 Hz), 1.34-1.44 (2H, m), 1.69-1.77 (2H, m), 2.85 (2H, t, J = 7.8 Hz), 4.19-4.26 (4H, m), 6.74-6.79 (2H, m), 6.92 (1H, br s), 7.52 (2H, d, J = 8.1 Hz), 7.91 (2H, d, J = 8.1 Hz), 8.59 (1H, s), 13.07 (1H, br s).	391	389
51		¹ H-NMR (DMSO-D ₆) δ: 1.42-1.53 (2H, m), 1.60-1.72 (2H, m), 2.67 (2H, t, J = 7.6 Hz), 3.38-3.46 (2H, m), 3.71 (3H, s), 4.38 (1H, br s), 6.79 (2H, d, J = 8.8 Hz), 7.19 (2H, d, J = 8.8 Hz), 7.30 (2H, d, J = 8.6 Hz), 7.63 (1H, d, J = 2.1 Hz), 7.85 (2H, d, J = 8.6 Hz), 8.50 (1H, d, J = 2.1 Hz), 12.97 (1H, br s).	378	376
52		¹ H-NMR (DMSO-D ₆) δ: 0.03-0.11 (2H, m), 0.37-0.45 (2H, m), 0.66-0.80 (1H, m), 1.54 (2H, dd, J = 15.4, 7.1 Hz), 2.74 (2H, dd, J = 14.4, 6.6 Hz), 3.71 (3H, s), 6.79 (2H, d, J = 8.8 Hz), 7.18 (2H, d, J = 8.8 Hz), 7.29 (2H, d, J = 8.6 Hz), 7.64 (1H, d, J = 2.1 Hz), 7.85 (2H, d, J = 8.6 Hz), 8.51 (1H, d, J = 2.3 Hz), 12.99 (1H, br s).	374	372
53		¹ H-NMR (DMSO-D ₆) δ: 1.50-1.57 (2H, m), 1.61-1.67 (2H, m), 1.71-1.82 (2H, m), 1.95-2.02 (2H, m), 2.67-2.76 (1H, m), 3.25 (3H, s), 3.49 (1H, br s), 3.72 (3H, s), 6.81 (2H, d, J = 8.4 Hz), 7.20 (2H, d, J = 8.4 Hz), 7.32 (2H, d, J = 8.1 Hz), 7.58 (1H, d, J = 1.7 Hz), 7.87 (2H, d, J = 8.1 Hz), 8.54 (1H, d, J = 1.7 Hz), 13.00 (1H, br s).	418	416
54		¹ H-NMR (DMSO-D ₆) δ: 1.20-1.32 (2H, m), 1.55-1.67 (2H, m), 1.87-1.95 (2H, m), 2.09-2.16 (2H, m), 2.61-2.70 (1H, m), 3.15-3.25 (1H, m), 3.27 (3H, s), 3.72 (3H, s), 6.81 (2H, d, J = 8.4 Hz), 7.20 (2H, d, J = 8.4 Hz), 7.31 (2H, d, J = 8.0 Hz), 7.65 (1H, d, J = 1.5 Hz), 7.86 (2H, d, J = 8.0 Hz), 8.56 (1H, d, J = 1.5 Hz), 13.00 (1H, br s).	418	416
55		¹ H-NMR (DMSO-D ₆) δ: 0.92 (3H, t, J = 7.3 Hz), 1.25 (9H, s), 1.33-1.43 (2H, m), 1.68-1.77 (2H, m), 2.86 (2H, t, J = 7.7 Hz), 7.30 (2H, d, J = 8.8 Hz), 7.33 (2H, d, J = 8.8 Hz), 7.49 (2H, d, J = 8.3 Hz), 7.87 (2H, d, J = 8.3 Hz), 8.62 (1H, s), 13.08 (1H, br s).	389	387
56		¹ H-NMR (DMSO-D ₆) δ: 1.26 (3H, t, J = 7.7 Hz), 2.71 (2H, q, J = 7.7 Hz), 3.72 (3H, s), 6.81 (2H, d, J = 9.0 Hz), 7.20 (2H, d, J = 9.0 Hz), 7.32 (2H, d, J = 8.6 Hz), 7.66 (1H, d, J = 2.1 Hz), 7.87 (2H, d, J = 8.6 Hz), 8.54 (1H, d, J = 2.1 Hz), 12.99 (1H, br s).	334	332

[0361]

Table 1-8

Ex.	Structure	NMR	MS (M+H)	MS (M-H)
57		¹ H-NMR (DMSO-D ₆) δ: 1.30 (6H, d, J = 6.9 Hz), 2.99-3.10 (1H, m), 3.72 (3H, s), 6.81 (2H, d, J = 8.9 Hz), 7.20 (2H, d, J = 8.9 Hz), 7.33 (2H, d, J = 8.1 Hz), 7.67 (1H, d, J = 2.1 Hz), 7.87 (2H, d, J = 8.1 Hz), 8.58 (1H, d, J = 2.1 Hz), 12.99 (1H, br s).	348	346
58		¹ H-NMR (DMSO-D ₆) δ: 0.93 (6H, d, J = 6.6 Hz), 1.87-1.99 (1H, m), 2.56 (2H, d, J = 7.2 Hz), 3.72 (3H, s), 6.81 (2H, d, J = 8.9 Hz), 7.21 (2H, d, J = 8.9 Hz), 7.32 (2H, d, J = 8.7 Hz), 7.61 (1H, d, J = 2.1 Hz), 7.87 (2H, d, J = 8.7 Hz), 8.48 (1H, d, J = 2.1 Hz), 12.99 (1H, br s).	362	360
59		¹ H-NMR (DMSO-D ₆) δ: 1.58-1.87 (6H, m), 2.04-2.15 (2H, m), 3.04-3.15 (1H, m), 3.72 (3H, s), 6.81 (2H, d, J = 9.0 Hz), 7.20 (2H, d, J = 9.0 Hz), 7.32 (2H, d, J = 8.6 Hz), 7.65 (1H, d, J = 2.1 Hz), 7.87 (2H, d, J = 8.6 Hz), 8.57 (1H, d, J = 2.1 Hz), 12.99 (1H, br s).	374	372
60		¹ H-NMR (DMSO-D ₆) δ: 1.74-1.83 (4H, m), 2.88-2.99 (1H, m), 3.41-3.52 (2H, m), 3.72 (3H, s), 3.94-4.01 (2H, m), 6.81 (2H, d, J = 9.0 Hz), 7.20 (2H, d, J = 9.0 Hz), 7.33 (2H, d, J = 8.2 Hz), 7.67 (1H, br s), 7.87 (2H, d, J = 8.2 Hz), 8.59 (1H, br s), 12.91 (1H, br s).	390	388
61		¹ H-NMR (DMSO-D ₆) δ: 1.75-1.84 (2H, m), 2.68-2.75 (2H, m), 3.42-3.49 (2H, m), 3.72 (3H, s), 4.51-4.56 (1H, m), 6.81 (2H, d, J = 9.0 Hz), 7.20 (2H, d, J = 9.0 Hz), 7.32 (2H, d, J = 8.1 Hz), 7.64 (1H, br s), 7.87 (2H, d, J = 8.1 Hz), 8.52 (1H, br s), 12.99 (1H, br s).	364	362
62		¹ H-NMR (DMSO-D ₆) δ: 2.66-2.79 (2H, m), 2.93 (2H, dd, J = 9.8, 6.4 Hz), 3.71 (3H, s), 6.80 (2H, d, J = 8.8 Hz), 7.19 (2H, d, J = 8.8 Hz), 7.31 (2H, d, J = 8.6 Hz), 7.79 (1H, d, J = 2.1 Hz), 7.86 (2H, d, J = 8.3 Hz), 8.59 (1H, d, J = 2.3 Hz), 13.01 (1H, br s).	402	400
63		¹ H-NMR (DMSO-D ₆) δ: 1.06-1.22 (2H, m), 1.40-1.70 (6H, m), 1.70-1.85 (3H, m), 2.67 (2H, t, J = 7.9 Hz), 3.71 (3H, s), 6.79 (2H, d, J = 9.0 Hz), 7.18 (2H, d, J = 9.0 Hz), 7.29 (2H, d, J = 8.6 Hz), 7.63 (1H, d, J = 2.1 Hz), 7.85 (2H, d, J = 8.3 Hz), 8.50 (1H, d, J = 2.1 Hz), 13.01 (1H, br s).	402	400
64		¹ H-NMR (DMSO-D ₆) δ: 0.92 (3H, t, J = 7.3 Hz), 1.32-1.44 (2H, m), 1.68-1.78 (2H, m), 2.87 (2H, t, J = 7.7 Hz), 3.90 (3H, d, J = 1.6 Hz), 7.21 (1H, dd, J = 11.9, 2.0 Hz), 7.29 (1H, t, J = 1.7 Hz), 7.53 (2H, d, J = 8.3 Hz), 7.93 (2H, d, J = 8.6 Hz), 8.65 (1H, s), 13.09 (1H, br s).	415	413

[0362]

Table 1-9

Ex.	Structure	NMR	MS (M+H)	MS (M-H)
65		¹ H-NMR (DMSO-D ₆) δ: 0.92 (3H, t, J = 7.4 Hz), 1.32-1.44 (2H, m), 1.68-1.78 (2H, m), 2.88 (2H, t, J = 7.7 Hz), 3.82 (3H, s), 7.42 (2H, s), 7.53 (2H, d, J = 8.6 Hz), 7.93 (2H, d, J = 8.6 Hz), 8.65 (1H, s), 13.13 (1H, br s).	431	429
66		¹ H-NMR (DMSO-D ₆) δ: 0.96 (3H, t, J = 7.4 Hz), 1.71-1.83 (2H, m), 2.85 (2H, t, J = 7.6 Hz), 3.93 (3H, s), 7.09 (2H, d, J = 9.5 Hz), 7.52 (2H, d, J = 8.6 Hz), 7.92 (2H, d, J = 8.6 Hz), 8.64 (1H, s), 13.11 (1H, br s).	385	383
67		¹ H-NMR (DMSO-D ₆) δ: 1.92-2.02 (2H, m), 2.91 (2H, t, J = 7.7 Hz), 3.24 (3H, s), 3.40 (2H, t, J = 6.4 Hz), 3.93 (3H, s), 7.09 (2H, d, J = 9.5 Hz), 7.53 (2H, d, J = 8.6 Hz), 7.92 (2H, d, J = 8.6 Hz), 8.64 (1H, s), 13.10 (1H, s).	415	413
68		¹ H-NMR (DMSO-D ₆) δ: 0.91 (3H, t, J = 7.4 Hz), 1.29-1.39 (2H, m), 1.58-1.66 (2H, m), 2.65-2.71 (2H, m), 3.90 (3H, s), 6.95 (2H, d, J = 9.7 Hz), 7.34 (2H, dd, J = 6.7, 1.8 Hz), 7.70 (1H, d, J = 2.3 Hz), 7.90 (2H, dd, J = 6.5, 1.8 Hz), 8.54 (1H, d, J = 2.1 Hz), 13.04 (1H, br s).	396	398
69		¹ H-NMR (DMSO-D ₆) δ: 0.94 (3H, t, J = 7.4 Hz), 1.35-1.44 (2H, m), 1.70-1.78 (2H, m), 2.87 (2H, t, J = 7.7 Hz), 3.78 (3H, s), 6.95 (2H, dt, J = 9.4, 2.5 Hz), 7.18 (2H, d, J = 9.0 Hz), 7.36 (2H, dt, J = 9.4, 2.5 Hz), 8.65 (1H, s), 14.03 (1H, s).	399	353 (-COOH)
70		¹ H-NMR (DMSO-D ₆) δ: 0.93 (3H, t, J = 7.4 Hz), 1.34-1.43 (2H, m), 1.69-1.77 (2H, m), 2.82 (2H, t, J = 7.6 Hz), 3.76 (3H, s), 4.69 (2H, s), 6.86-6.91 (4H, m), 7.35-7.31 (4H, m), 8.49 (1H, s), 13.02 (1H, s).	393	391
71		¹ H-NMR (DMSO-D ₆) δ: 0.93 (3H, t, J = 7.3 Hz), 1.33-1.43 (2H, m), 1.68-1.76 (2H, m), 2.23 (2H, t, J = 7.7 Hz), 2.76 (2H, t, J = 7.7 Hz), 2.82 (2H, t, J = 7.7 Hz), 3.75 (3H, s), 6.87 (2H, dt, J = 9.5, 2.5 Hz), 7.15 (2H, d, J = 8.1 Hz), 7.27 (2H, d, J = 8.1 Hz), 7.31 (2H, dt, J = 9.4, 2.5 Hz), 8.50 (1H, s).	391	389
72		¹ H-NMR (DMSO-D ₆) δ: 0.93 (3H, t, J = 7.4 Hz), 1.34-1.42 (2H, m), 1.70-1.77 (2H, m), 2.84 (2H, t, J = 7.6 Hz), 3.22 (2H, br s), 3.76 (3H, s), 6.91 (2H, dt, J = 9.4, 2.5 Hz), 7.03-7.10 (2H, m), 7.20 (1H, t, J = 7.7 Hz), 7.35 (2H, dt, J = 9.4, 2.5 Hz), 8.53 (1H, s).	395	349 (-COOH)

[0363]

Table 1-10

Ex.	Structure	NMR	MS (M+H)	MS (M-H)
73		¹ H-NMR (DMSO-D ₆) δ: 1.10 (3H, t, J = 7.0 Hz), 2.91 (2H, t, J = 6.7 Hz), 3.46 (2H, q, J = 7.0 Hz), 3.66 (2H, t, J = 6.7 Hz), 3.72 (3H, s), 6.81 (2H, d, J = 8.8 Hz), 7.20 (2H, d, J = 8.8 Hz), 7.31 (2H, d, J = 8.5 Hz), 7.70 (1H, d, J = 2.2 Hz), 7.87 (2H, d, J = 8.5 Hz), 8.54 (1H, d, J = 2.2 Hz), 12.99 (1H, br s).	378	376
74		¹ H-NMR (DMSO-D ₆) δ: 0.94 (3H, t, J = 7.3 Hz), 1.17 (3H, t, J = 7.6 Hz), 1.33-1.46 (2H, m), 1.69-1.79 (2H, m), 2.60 (2H, q, J = 7.6 Hz), 2.87 (2H, t, J = 7.8 Hz), 7.17 (2H, d, J = 8.1 Hz), 7.29 (2H, d, J = 8.1 Hz), 7.50 (2H, d, J = 8.1 Hz), 7.88 (2H, d, J = 8.1 Hz), 8.63 (1H, s), 13.09 (1H, br s).	361	359
75		¹ H-NMR (DMSO-D ₆) δ: 0.94 (3H, t, J = 7.3 Hz), 1.34-1.46 (2H, m), 1.69-1.81 (2H, m), 2.90 (2H, t, J = 7.6 Hz), 7.34 (2H, d, J = 8.4 Hz), 7.46-7.53 (4H, m), 7.90 (2H, d, J = 8.1 Hz), 8.67 (1H, s), 13.11 (1H, br s).	417	415
76		¹ H-NMR (DMSO-D ₆) δ: 0.91 (3H, t, J = 7.4 Hz), 1.32-1.40 (2H, m), 1.50-1.57 (2H, m), 3.31 (2H, dd, J = 12.8, 6.8 Hz), 3.74 (3H, d, J = 0.7 Hz), 6.86 (2H, dd, J = 8.9, 2.9 Hz), 7.28 (2H, dd, J = 8.8, 1.6 Hz), 7.39 (2H, d, J = 8.1 Hz), 7.91 (2H, d, J = 8.1 Hz), 8.25 (1H, br s), 8.74 (1H, br s), 9.07 (1H, s).	405	403
77		¹ H-NMR (DMSO-D ₆) δ: 1.11 (3H, t, J = 7.0 Hz), 1.83-1.92 (2H, m), 2.70-2.76 (2H, m), 3.38-3.46 (4H, m), 3.72 (3H, s), 6.81 (2H, d, J = 8.8 Hz), 7.20 (2H, d, J = 8.8 Hz), 7.32 (2H, d, J = 8.6 Hz), 7.65 (1H, d, J = 2.2 Hz), 7.87 (2H, d, J = 8.6 Hz), 8.52 (1H, d, J = 2.2 Hz), 12.98 (1H, br s).	392	390
78		¹ H-NMR (DMSO-D ₆) δ: 3.36 (3H, s), 3.73 (3H, s), 4.55 (2H, s), 6.82 (2H, d, J = 8.8 Hz), 7.23 (2H, d, J = 8.8 Hz), 7.33 (2H, d, J = 8.6 Hz), 7.74 (1H, d, J = 2.1 Hz), 7.88 (2H, d, J = 8.6 Hz), 8.62 (1H, d, J = 2.1 Hz), 13.01 (1H, br s).	350	348
79		¹ H-NMR (DMSO-D ₆) δ: 0.86 (3H, t, J = 7.3 Hz), 0.92 (3H, t, J = 7.3 Hz), 1.33-1.43 (2H, m), 1.50-1.62 (2H, m), 1.68-1.78 (2H, m), 2.53 (2H, t, J = 7.6 Hz), 2.86 (2H, t, J = 7.7 Hz), 7.13 (2H, d, J = 8.3 Hz), 7.26 (2H, d, J = 8.1 Hz), 7.47 (2H, d, J = 8.6 Hz), 7.85 (2H, d, J = 8.3 Hz), 8.61 (1H, s), 13.04 (1H, s).	375	373
80		¹ H-NMR (DMSO-D ₆) δ: 1.16-1.26 (2H, m), 1.45-1.55 (2H, m), 1.58-1.73 (4H, m), 2.10-2.17 (1H, m), 2.67 (2H, d, J = 7.4 Hz), 3.71 (3H, s), 6.79 (2H, dt, J = 9.5, 2.5 Hz), 7.19 (2H, dt, J = 9.4, 2.5 Hz), 7.30 (2H, dt, J = 8.4, 1.8 Hz), 7.62 (1H, d, J = 2.1 Hz), 7.85 (2H, dt, J = 8.5, 1.8 Hz), 8.49 (1H, d, J = 2.1 Hz), 12.97 (1H, br s).	386	388

[0364]

Table 1-11

Ex.	Structure	NMR	MS (M+H)	MS (M-H)
81		¹ H-NMR (DMSO-D ₆) δ: 1.70-1.86 (4H, m), 1.96-2.05 (2H, m), 2.57-2.66 (1H, m), 2.76 (2H, d, J = 7.4 Hz), 3.71 (3H, s), 6.79 (2H, dt, J = 9.5, 2.4 Hz), 7.18 (2H, dt, J = 9.4, 2.5 Hz), 7.29 (2H, dt, J = 8.4, 1.7 Hz), 7.58 (1H, d, J = 2.1 Hz), 7.85 (2H, dt, J = 8.4, 1.8 Hz), 8.47 (1H, d, J = 2.1 Hz), 12.97 (1H, br s).	372	374
82		¹ H-NMR (DMSO-D ₆) δ: 0.92 (3H, t, J = 7.4 Hz), 1.29 (3H, t, J = 7.1 Hz), 1.32-1.43 (2H, m), 1.66-1.77 (2H, m), 2.84 (2H, t, J = 7.6 Hz), 4.00 (2H, q, J = 6.9 Hz), 6.85 (2H, d, J = 8.8 Hz), 7.28 (2H, d, J = 8.8 Hz), 7.47 (2H, d, J = 8.6 Hz), 7.87 (2H, d, J = 8.6 Hz), 8.58 (1H, s), 13.13 (1H, br s).	377	375
83		¹ H-NMR (DMSO-D ₆) δ: 0.94 (3H, t, J = 7.4 Hz), 1.35-1.42 (2H, m), 1.69-1.77 (2H, m), 2.48 (3H, s), 2.85 (2H, t, J = 7.7 Hz), 3.75 (3H, s), 6.89 (2H, dt, J = 9.4, 2.5 Hz), 7.18 (1H, dd, J = 8.1, 1.6 Hz), 7.32 (2H, dt, J = 9.4, 2.5 Hz), 7.41 (1H, d, J = 1.6 Hz), 7.72 (1H, d, J = 8.1 Hz), 8.59 (1H, s), 12.92 (1H, br s).	377	375
84		¹ H-NMR (DMSO-D ₆) δ: 1.84-1.93 (2H, m), 2.72 (2H, t, J = 7.8 Hz), 3.25 (3H, s), 3.38 (2H, t, J = 6.4 Hz), 3.72 (3H, s), 6.81 (2H, d, J = 8.8 Hz), 7.20 (2H, d, J = 8.8 Hz), 7.32 (2H, d, J = 8.3 Hz), 7.65 (1H, d, J = 2.2 Hz), 7.87 (2H, d, J = 8.6 Hz), 8.52 (1H, d, J = 2.2 Hz), 12.99 (1H, br s).	378	376
85		¹ H-NMR (DMSO-D ₆) δ: 0.94 (6H, d, J = 6.4 Hz), 1.50-1.65 (3H, m), 2.68 (2H, t, J = 7.8 Hz), 3.72 (3H, s), 6.81 (2H, d, J = 8.9 Hz), 7.20 (2H, d, J = 8.9 Hz), 7.31 (2H, d, J = 8.3 Hz), 7.65 (1H, d, J = 2.1 Hz), 7.87 (2H, d, J = 8.3 Hz), 8.52 (1H, d, J = 2.1 Hz), 12.99 (1H, br s).	376	374
86		¹ H-NMR (DMSO-D ₆) δ: 0.92 (3H, t, J = 7.3 Hz), 1.32-1.44 (2H, m), 1.68-1.79 (2H, m), 2.87 (2H, t, J = 7.7 Hz), 3.27 (3H, s), 4.39 (2H, s), 7.25 (2H, d, J = 8.6 Hz), 7.33 (2H, d, J = 8.3 Hz), 7.47 (2H, d, J = 8.6 Hz), 7.86 (2H, d, J = 8.6 Hz), 8.63 (1H, s), 13.02 (1H, br s).	377	375
87		¹ H-NMR (DMSO-D ₆) δ: 0.92 (3H, t, J = 7.4 Hz), 1.33-1.45 (2H, m), 1.67-1.79 (2H, m), 2.87 (2H, t, J = 7.7 Hz), 4.04 (2H, s), 7.29 (2H, d, J = 8.6 Hz), 7.38 (2H, d, J = 8.6 Hz), 7.48 (2H, d, J = 8.6 Hz), 7.87 (2H, d, J = 8.6 Hz), 8.64 (1H, s), 13.06 (1H, br s).	372	370
88		¹ H-NMR (DMSO-D ₆) δ: 0.92 (3H, t, J = 7.4 Hz), 1.17 (6H, d, J = 6.9 Hz), 1.32-1.44 (2H, m), 1.67-1.78 (2H, m), 2.81-2.91 (3H, m), 7.18 (2H, d, J = 8.1 Hz), 7.28 (2H, d, J = 8.3 Hz), 7.49 (2H, d, J = 8.6 Hz), 7.87 (2H, d, J = 8.6 Hz), 8.61 (1H, s), 13.04 (1H, br s).	375	373

[0365]

Table 1-12

Ex.	Structure	NMR	MS (M+H)	MS (M-H)
89		¹ H-NMR (DMSO-D ₆) δ: 0.92 (3H, t, J = 7.3 Hz), 1.33-1.44 (2H, m), 1.67-1.80 (2H, m), 2.87 (2H, t, J = 7.6 Hz), 4.19-4.28 (1H, m), 4.59 (2H, t, J = 6.4 Hz), 4.90 (2H, dd, J = 8.4, 5.9 Hz), 7.34 (2H, d, J = 8.8 Hz), 7.37 (2H, d, J = 8.6 Hz), 7.49 (2H, d, J = 8.6 Hz), 7.87 (2H, d, J = 8.6 Hz), 8.63 (1H, s), 13.05 (1H, br s).	389	387
90		¹ H-NMR (DMSO-D ₆) δ: 0.94 (3H, t, J = 7.4 Hz), 1.35-1.45 (2H, m), 1.71-1.78 (2H, m), 2.88 (2H, t, J = 7.7 Hz), 7.10-7.15 (2H, m), 7.29 (1H, t, J = 73.5 Hz), 7.42 (2H, dt, J = 9.2, 2.4 Hz), 7.51 (2H, dt, J = 8.4, 1.8 Hz), 7.90 (2H, dt, J = 8.4, 1.8 Hz), 8.65 (1H, s), 13.07 (1H, s).	399	397
91		¹ H-NMR (DMSO-D ₆) δ: 0.92 (3H, t, J = 7.3 Hz), 1.31-1.44 (2H, m), 1.66-1.78 (2H, m), 2.84 (2H, t, J = 7.6 Hz), 3.74 (3H, d, J = 3.7 Hz), 6.83-6.92 (3H, m), 6.97 (1H, d, J = 1.6 Hz), 7.33 (2H, d, J = 9.0 Hz), 7.70 (1H, d, J = 8.1 Hz), 8.58 (1H, s).	379	377
92		¹ H-NMR (DMSO-D ₆) δ: 0.92 (3H, t, J = 7.4 Hz), 1.33-1.44 (2H, m), 1.67-1.77 (2H, m), 2.85 (2H, t, J = 7.6 Hz), 3.75 (3H, s), 6.91 (2H, d, J = 8.8 Hz), 7.29-7.37 (3H, m), 7.57 (1H, d, J = 1.6 Hz), 7.70 (1H, d, J = 7.9 Hz), 8.61 (1H, s), 13.47 (1H, br s).	397	395
93		¹ H-NMR (DMSO-D ₆) δ: 0.92 (3H, t, J = 7.3 Hz), 1.31-1.44 (2H, m), 1.66-1.77 (2H, m), 2.49-2.54 (2H, m), 2.82 (4H, t, J = 7.6 Hz), 3.75 (3H, s), 6.88 (2H, d, J = 8.8 Hz), 7.09 (1H, dd, J = 7.9, 1.6 Hz), 7.15 (1H, dd, J = 11.1, 1.6 Hz), 7.25 (1H, t, J = 7.9 Hz), 7.31 (2H, d, J = 9.0 Hz), 8.55 (1H, s), 12.23 (1H, br s).	409	407
94		¹ H-NMR (DMSO-D ₆) δ: 1.83-1.90 (2H, m), 2.51-2.55 (2H, m), 2.69 (2H, t, J = 7.6 Hz), 2.81 (2H, t, J = 7.6 Hz), 3.25 (3H, s), 3.37 (2H, t, J = 6.3 Hz), 3.72 (3H, s), 6.79 (2H, d, J = 8.4 Hz), 7.10 (2H, d, J = 7.8 Hz), 7.17-7.20 (4H, m), 7.57 (1H, d, J = 1.8 Hz), 8.45 (1H, d, J = 1.5 Hz), 12.12 (1H, br s).	406	404
95		¹ H-NMR (DMSO-D ₆) δ: 1.83-1.92 (2H, m), 2.50-2.57 (2H, m), 2.72 (2H, t, J = 7.7 Hz), 2.84 (2H, t, J = 7.7 Hz), 3.25 (3H, s), 3.37 (2H, t, J = 6.3 Hz), 3.91 (3H, s), 6.91-6.99 (2H, m), 7.14 (2H, d, J = 8.0 Hz), 7.24 (2H, d, J = 8.0 Hz), 7.65 (1H, d, J = 1.8 Hz), 8.50 (1H, d, J = 1.8 Hz), 12.13 (1H, s).	442	440
96		¹ H-NMR (DMSO-D ₆) δ: 0.24-0.30 (2H, m), 0.47-0.54 (2H, m), 1.00-1.11 (1H, m), 2.60 (2H, d, J = 6.9 Hz), 3.73 (3H, s), 6.81 (2H, d, J = 8.7 Hz), 7.21 (2H, d, J = 8.4 Hz), 7.32 (2H, d, J = 8.1 Hz), 7.71 (1H, d, J = 1.5 Hz), 7.88 (2H, d, J = 8.1 Hz), 8.57 (1H, d, J = 1.8 Hz), 12.99 (1H, br s).	360	358

[0366]

Table 1-13

Ex.	Structure	NMR	MS (M+H)	MS (M-H)
97		¹ H-NMR (DMSO-D ₆) δ: 1.95-2.02 (2H, m), 2.54 (2H, t, J = 7.7 Hz), 2.85 (2H, t, J = 7.6 Hz), 2.90 (2H, t, J = 7.6 Hz), 3.25 (3H, s), 3.41 (2H, t, J = 6.2 Hz), 3.94 (3H, s), 7.09 (2H, d, J = 9.9 Hz), 7.25 (2H, d, J = 8.1 Hz), 7.34 (2H, d, J = 8.1 Hz), 8.58 (1H, s), 12.14 (1H, s).	443	441
98		¹ H-NMR (DMSO-D ₆) δ: 0.94 (6H, d, J = 6.7 Hz), 2.06-2.19 (1H, m), 2.75 (2H, d, J = 7.2 Hz), 3.93 (3H, s), 7.10 (2H, d, J = 9.5 Hz), 7.52 (2H, d, J = 8.3 Hz), 7.92 (2H, d, J = 8.3 Hz), 8.61 (1H, s), 13.07 (1H, br s).	399	397
99		¹ H-NMR (DMSO-D ₆) δ: 0.91 (6H, t, J = 5.4 Hz), 1.86-1.99 (1H, m), 2.57 (2H, d, J = 7.2 Hz), 3.90 (3H, s), 6.96 (2H, d, J = 9.7 Hz), 7.34 (2H, d, J = 8.6 Hz), 7.67 (1H, d, J = 2.3 Hz), 7.90 (2H, d, J = 8.6 Hz), 8.51 (1H, d, J = 2.1 Hz), 12.97 (1H, br s).	398	396
100		¹ H-NMR (DMSO-D ₆) δ: 0.85 (3H, t, J = 7.1 Hz), 1.19-1.41 (6H, m), 1.56-1.69 (2H, m), 2.66 (2H, t, J = 7.7 Hz), 3.71 (3H, s), 6.79 (2H, d, J = 9.0 Hz), 7.19 (2H, d, J = 8.8 Hz), 7.27 (2H, d, J = 8.3 Hz), 7.62 (1H, d, J = 2.1 Hz), 7.84 (2H, d, J = 8.6 Hz), 8.49 (1H, d, J = 2.1 Hz).	390	388
101		¹ H-NMR (DMSO-D ₆) δ: 3.70 (3H, s), 4.04 (2H, s), 6.79 (2H, d, J = 8.8 Hz), 7.14-7.23 (3H, m), 7.23-7.36 (6H, m), 7.65 (1H, d, J = 2.1 Hz), 7.84 (2H, d, J = 8.6 Hz), 8.57 (1H, d, J = 2.3 Hz), 13.01 (1H, br s).	396	394
102		¹ H-NMR (DMSO-D ₆) δ: 2.91 (2H, t, J = 6.6 Hz), 3.27 (3H, s), 3.63 (2H, t, J = 6.6 Hz), 3.72 (3H, s), 6.81 (2H, d, J = 8.8 Hz), 7.20 (2H, d, J = 8.8 Hz), 7.31 (2H, d, J = 8.3 Hz), 7.69 (1H, d, J = 2.2 Hz), 7.87 (2H, d, J = 8.3 Hz), 8.54 (1H, d, J = 2.2 Hz), 12.99 (1H, br s).	364	362
103		¹ H-NMR (DMSO-D ₆) δ: 0.93 (3H, t, J = 7.3 Hz), 1.61-1.71 (2H, m), 2.66 (2H, t, J = 7.5 Hz), 3.90 (3H, s), 6.95 (2H, d, J = 9.7 Hz), 7.34 (2H, dt, J = 8.5, 1.8 Hz), 7.70 (1H, d, J = 2.1 Hz), 7.90 (2H, dt, J = 8.3, 1.8 Hz), 8.54 (1H, d, J = 2.1 Hz), 13.04 (1H, br s).	382	384
104		¹ H-NMR (DMSO-D ₆) δ: 1.88-2.00 (2H, m), 2.73 (2H, t, J = 7.6 Hz), 3.64 (2H, t, J = 6.3 Hz), 3.72 (3H, s), 4.06 (2H, q, J = 9.5 Hz), 6.81 (2H, d, J = 8.7 Hz), 7.20 (2H, d, J = 8.4 Hz), 7.32 (2H, d, J = 8.1 Hz), 7.67 (1H, d, J = 1.8 Hz), 7.87 (2H, d, J = 8.1 Hz), 8.53 (1H, d, J = 1.8 Hz), 13.04 (1H, br s).	446	444

[0367]

Table 1-14

Ex.	Structure	NMR	MS (M+H)	MS (M-H)
105		¹ H-NMR (DMSO-D ₆) δ: 2.98 (4H, s), 3.72 (3H, s), 6.81 (2H, d, J = 8.7 Hz), 7.17-7.23 (3H, m), 7.25-7.34 (6H, m), 7.66 (1H, s), 7.87 (2H, d, J = 8.1 Hz), 8.52 (1H, s), 13.05 (1H, br s).	410	408
106		¹ H-NMR (DMSO-D ₆) δ: 1.83-1.94 (2H, m), 2.73 (2H, t, J = 7.8 Hz), 3.26 (3H, s), 3.38 (2H, t, J = 6.3 Hz), 3.80 (3H, s), 6.93-7.07 (2H, m), 7.13 (1H, d, J = 12.9 Hz), 7.34 (2H, d, J = 8.1 Hz), 7.68 (1H, s), 7.89 (2H, d, J = 8.1 Hz), 8.54 (1H, s), 13.04 (1H, br s).	396	394
107		¹ H-NMR (DMSO-D ₆) δ: 1.83-1.94 (2H, m), 2.74 (2H, t, J = 7.8 Hz), 3.25 (3H, s), 3.38 (2H, t, J = 6.3 Hz), 3.91 (3H, s), 6.96 (2H, d, J = 9.6 Hz), 7.35 (2H, d, J = 8.4 Hz), 7.72 (1H, d, J = 1.8 Hz), 7.91 (2H, d, J = 8.4 Hz), 8.56 (1H, d, J = 1.8 Hz), 13.09 (1H, br s).	414	412
108		¹ H-NMR (DMSO-D ₆) δ: 1.82-1.92 (2H, m), 2.51-2.56 (2H, m), 2.70 (2H, t, J = 7.7 Hz), 2.83 (2H, t, J = 7.6 Hz), 3.25 (3H, s), 3.37 (2H, t, J = 6.2 Hz), 3.81 (3H, s), 7.00-7.14 (5H, m), 7.20 (2H, d, J = 8.1 Hz), 7.60 (1H, d, J = 2.2 Hz), 8.47 (1H, d, J = 2.2 Hz), 12.13 (1H, br s).	424	422
109		¹ H-NMR (DMSO-D ₆) δ: 1.95-2.02 (2H, m), 2.51 (3H, s), 2.92 (2H, t, J = 8.4 Hz), 3.25 (3H, s), 3.42 (2H, t, J = 6.2 Hz), 3.95 (3H, s), 7.12 (2H, d, J = 9.7 Hz), 7.23 (1H, dd, J = 8.1, 1.4 Hz), 7.43 (1H, d, J = 1.4 Hz), 7.77 (1H, d, J = 8.1 Hz), 8.64 (1H, s), 12.96 (1H, s).	429	427
110		¹ H-NMR (DMSO-D ₆) δ: 0.91 (3H, t, J = 7.4 Hz), 1.31-1.42 (2H, m), 1.65-1.76 (2H, m), 2.14 (6H, s), 2.81 (2H, t, J = 7.7 Hz), 3.74 (3H, s), 6.88 (2H, d, J = 9.0 Hz), 7.03 (2H, s), 7.33 (2H, d, J = 8.8 Hz), 8.52 (1H, s).	391	389
111		¹ H-NMR (DMSO-D ₆) δ: 0.92 (3H, t, J = 7.4 Hz), 0.97 (3H, t, J = 7.5 Hz), 1.32-1.44 (2H, m), 1.67-1.77 (2H, m), 2.78 (2H, q, J = 7.5 Hz), 2.84 (2H, t, J = 7.6 Hz), 3.73 (3H, s), 6.88 (2H, d, J = 9.0 Hz), 7.23 (1H, d, J = 1.6 Hz), 7.29 (2H, d, J = 9.0 Hz), 7.35 (1H, dd, J = 8.1, 1.8 Hz), 7.73 (1H, d, J = 8.1 Hz), 8.57 (1H, s), 12.92 (1H, br s).	391	389
112		¹ H-NMR (DMSO-D ₆) δ: 1.93-2.03 (2H, m), 2.92 (2H, t, J = 7.7 Hz), 3.24 (3H, s), 3.41 (2H, t, J = 6.2 Hz), 3.94 (3H, s), 7.13 (2H, d, J = 9.5 Hz), 7.26 (1H, dd, J = 8.1, 1.6 Hz), 7.37 (1H, dd, J = 11.7, 1.5 Hz), 7.82 (1H, t, J = 7.9 Hz), 8.67 (1H, s), 13.38 (1H, br s).	433	431

[0368]

Table 1-15

Ex.	Structure	NMR	MS (M+H)	MS (M-H)
113		¹ H-NMR (DMSO-D ₆) δ: 1.93-2.02 (2H, m), 2.90 (2H, t, J = 7.7 Hz), 3.24 (3H, s), 3.40 (2H, t, J = 6.4 Hz), 3.83 (3H, s), 7.07-7.15 (2H, m), 7.24-7.28 (2H, m), 7.32 (1H, dt, J = 15.0, 5.5 Hz), 7.80 (1H, t, J = 7.9 Hz), 8.63 (1H, s), 13.37 (1H, br s).	415	413
114		¹ H-NMR (DMSO-D ₆) δ: 0.93 (3H, t, J = 7.3 Hz), 1.31-1.42 (2H, m), 1.58-1.68 (2H, m), 2.49 (3H, s), 2.67 (2H, t, J = 7.6 Hz), 3.73 (3H, s), 6.82 (2H, d, J = 9.0 Hz), 7.00 (1H, dd, J = 8.0, 1.5 Hz), 7.21-7.23 (3H, m), 7.62 (1H, d, J = 2.0 Hz), 7.70 (1H, d, J = 8.0 Hz), 8.50 (1H, d, J = 2.0 Hz), 12.84 (1H, br s).	376	374
115		¹ H-NMR (DMSO-D ₆) δ: 1.82-1.91 (2H, m), 2.49 (3H, s), 2.72 (2H, t, J = 7.7 Hz), 3.24 (3H, s), 3.36 (2H, t, J = 6.2 Hz), 3.90 (3H, s), 6.97 (2H, d, J = 9.5 Hz), 7.04 (1H, dd, J = 8.1, 1.6 Hz), 7.23 (1H, s), 7.69 (1H, d, J = 2.1 Hz), 7.74 (1H, d, J = 8.1 Hz), 8.53 (1H, d, J = 2.3 Hz), 12.90 (1H, br s).	428	426
116		¹ H-NMR (DMSO-D ₆) δ: 1.81-1.91 (2H, m), 2.47 (3H, s), 2.70 (2H, t, J = 7.7 Hz), 3.24 (3H, s), 3.36 (2H, t, J = 6.2 Hz), 3.71 (3H, s), 6.81 (2H, d, J = 8.8 Hz), 6.98 (1H, dd, J = 8.1, 1.4 Hz), 7.18-7.24 (3H, m), 7.62 (1H, d, J = 2.3 Hz), 7.69 (1H, d, J = 8.1 Hz), 8.49 (1H, d, J = 2.3 Hz), 12.81 (1H, br s).	392	390
117		¹ H-NMR (DMSO-D ₆) δ: 1.08 (3H, t, J = 7.1 Hz), 3.11 (2H, t, J = 6.5 Hz), 3.46 (2H, q, J = 7.0 Hz), 3.80 (2H, t, J = 6.5 Hz), 3.93 (3H, s), 7.10 (2H, d, J = 9.7 Hz), 7.52 (2H, d, J = 8.6 Hz), 7.92 (2H, d, J = 8.6 Hz), 8.65 (1H, s), 13.10 (1H, s).	415	413
118		¹ H-NMR (DMSO-D ₆) δ: 0.88 (3H, t, J = 7.4 Hz), 1.51-1.62 (2H, m), 3.45 (2H, t, J = 6.6 Hz), 3.71 (3H, s), 4.57 (2H, s), 6.81 (2H, d, J = 8.8 Hz), 7.21 (2H, d, J = 8.8 Hz), 7.30 (2H, d, J = 8.3 Hz), 7.71 (1H, d, J = 2.1 Hz), 7.86 (2H, d, J = 8.6 Hz), 8.60 (1H, d, J = 2.1 Hz), 13.01 (1H, br s).	378	376
119		¹ H-NMR (DMSO-D ₆) δ: 1.86-1.97 (2H, m), 2.83 (2H, t, J = 7.6 Hz), 3.21 (3H, s), 3.39-3.51 (6H, m), 3.75 (3H, s), 6.91 (2H, d, J = 9.0 Hz), 7.26 (2H, d, J = 9.0 Hz), 7.34 (2H, d, J = 8.6 Hz), 7.89 (2H, d, J = 8.6 Hz), 8.15 (1H, br s), 8.68 (1H, d, J = 1.8 Hz), 13.04 (1H, br s).	422	420
120		¹ H-NMR (DMSO-D ₆) δ: 1.82-1.92 (2H, m), 2.70 (2H, t, J = 7.7 Hz), 3.24 (3H, s), 3.36 (2H, t, J = 6.4 Hz), 3.72 (3H, s), 6.84 (2H, d, J = 8.8 Hz), 7.14 (1H, dd, J = 8.1, 1.6 Hz), 7.22 (2H, d, J = 8.8 Hz), 7.42 (1H, d, J = 1.6 Hz), 7.64-7.70 (2H, m), 8.51 (1H, d, J = 2.1 Hz), 13.39 (1H, br s).	412	410

[0369]

Table 1-16

Ex.	Structure	NMR	MS (M+H)	MS (M-H)
121		¹ H-NMR (DMSO-D ₆) δ: 1.96-2.03 (2H, m), 2.93 (2H, t, J = 7.7 Hz), 3.25 (3H, s), 3.42 (2H, t, J = 6.4 Hz), 3.96 (3H, s), 7.17 (2H, d, J = 9.2 Hz), 7.36 (1H, dd, J = 8.1, 1.6 Hz), 7.64 (1H, d, J = 1.6 Hz), 7.75 (1H, d, J = 8.1 Hz), 8.68 (1H, s), 13.52 (1H, s).	449	447
122		¹ H-NMR (DMSO-D ₆) δ: 1.82-1.92 (2H, m), 2.72 (2H, t, J = 7.7 Hz), 3.24 (3H, s), 3.36 (2H, t, J = 6.2 Hz), 3.91 (3H, s), 7.00 (2H, d, J = 9.7 Hz), 7.18 (1H, dd, J = 8.0, 1.7 Hz), 7.47 (1H, d, J = 1.6 Hz), 7.71 (1H, d, J = 7.9 Hz), 7.76 (1H, d, J = 2.1 Hz), 8.55 (1H, d, J = 2.1 Hz), 13.42 (1H, br s).	446	448
123		¹ H-NMR (DMSO-D ₆) δ: 1.51-1.74 (4H, m), 2.69 (2H, t, J = 7.5 Hz), 3.22 (3H, s), 3.35 (2H, t, J = 6.3 Hz), 3.72 (3H, s), 6.81 (2H, d, J = 8.9 Hz), 7.20 (2H, d, J = 8.9 Hz), 7.32 (2H, d, J = 8.1 Hz), 7.64 (1H, d, J = 1.5 Hz), 7.87 (2H, d, J = 8.1 Hz), 8.52 (1H, d, J = 1.5 Hz), 12.99 (1H, br s).	392	390
124		¹ H-NMR (DMSO-D ₆) δ: 1.82-1.91 (2H, m), 2.72 (2H, t, J = 7.7 Hz), 3.24 (3H, s), 3.36 (2H, t, J = 6.2 Hz), 3.91 (3H, s), 6.70 (1H, dd, J = 8.1, 1.6 Hz), 6.87 (1H, d, J = 1.6 Hz), 7.00 (2H, d, J = 9.7 Hz), 7.70 (1H, d, J = 2.1 Hz), 7.71 (1H, d, J = 8.1 Hz), 8.54 (1H, d, J = 2.1 Hz), 11.37 (1H, br s), 14.00 (1H, br s).	430	428
125		¹ H-NMR (DMSO-D ₆) δ: 1.50-1.59 (2H, m), 1.62-1.72 (2H, m), 2.67 (2H, t, J = 7.6 Hz), 3.20 (3H, s), 3.34 (2H, t, J = 6.4 Hz), 3.72 (3H, s), 6.84 (2H, d, J = 9.0 Hz), 7.14 (1H, dd, J = 8.0, 1.7 Hz), 7.22 (2H, d, J = 8.8 Hz), 7.41 (1H, d, J = 1.8 Hz), 7.65-7.69 (2H, m), 8.51 (1H, d, J = 2.1 Hz), 13.39 (1H, br s).	426	424
126		¹ H-NMR (DMSO-D ₆) δ: 1.50-1.59 (2H, m), 1.62-1.71 (2H, m), 2.47 (3H, s), 2.67 (2H, t, J = 7.5 Hz), 3.20 (3H, s), 3.33 (2H, t, J = 6.4 Hz), 3.71 (3H, s), 6.81 (2H, d, J = 9.0 Hz), 6.98 (1H, dd, J = 8.0, 1.5 Hz), 7.19-7.23 (3H, m), 7.61 (1H, d, J = 2.1 Hz), 7.68 (1H, d, J = 8.1 Hz), 8.48 (1H, d, J = 2.1 Hz), 12.81 (1H, br s).	406	404
127		¹ H-NMR (DMSO-D ₆) δ: 1.49-1.58 (2H, m), 1.60-1.71 (2H, m), 2.47-2.53 (2H, m), 2.65 (2H, t, J = 7.5 Hz), 2.80 (2H, t, J = 7.6 Hz), 3.20 (3H, s), 3.33 (2H, t, J = 6.4 Hz), 3.71 (3H, s), 6.77 (2H, d, J = 9.0 Hz), 7.08 (2H, d, J = 8.3 Hz), 7.16 (2H, d, J = 8.3 Hz), 7.19 (2H, d, J = 8.8 Hz), 7.55 (1H, d, J = 2.3 Hz), 8.44 (1H, d, J = 2.1 Hz), 12.15 (1H, br s).	420	418
128		¹ H-NMR (DMSO-D ₆) δ: 1.08 (3H, t, J = 7.1 Hz), 2.49-2.55 (2H, m), 2.82 (2H, t, J = 7.6 Hz), 2.89 (2H, t, J = 6.6 Hz), 3.44 (2H, q, J = 6.9 Hz), 3.64 (2H, t, J = 6.7 Hz), 3.90 (3H, s), 6.94 (2H, d, J = 9.7 Hz), 7.11 (2H, d, J = 8.3 Hz), 7.22 (2H, d, J = 8.3 Hz), 7.67 (1H, d, J = 2.1 Hz), 8.51 (1H, d, J = 2.1 Hz), 12.15 (1H, br s).	442	420









[0370]

Table 1-17

Ex.	Structure	NMR	MS (M+H)	MS (M-H)
129		¹ H-NMR (DMSO-D ₆) δ: 1.52-1.72 (4H, m), 2.71 (2H, t, J = 7.6 Hz), 3.21 (3H, s), 3.35 (2H, t, J = 6.3 Hz), 3.91 (3H, s), 6.94-6.99 (2H, m), 7.36 (2H, d, J = 8.1 Hz), 7.71 (1H, d, J = 1.8 Hz), 7.92 (2H, d, J = 8.1 Hz), 8.56 (1H, d, J = 1.8 Hz), 13.06 (1H, br s).	428	426
130		¹ H-NMR (DMSO-D ₆) δ: 1.50-1.73 (4H, m), 2.51 (3H, s), 2.71 (2H, t, J = 7.3 Hz), 3.22 (3H, s), 3.35 (2H, t, J = 6.4 Hz), 3.91 (3H, s), 6.94-7.01 (2H, m), 7.06 (1H, d, J = 8.1 Hz), 7.25 (1H, br s), 7.70 (1H, br s), 7.76 (1H, d, J = 8.1 Hz), 8.54 (1H, br s), 12.90 (1H, br s).	442	440
131		¹ H-NMR (DMSO-D ₆) δ: 1.50-1.75 (4H, m), 2.71 (2H, t, J = 7.6 Hz), 3.22 (3H, s), 3.32-3.38 (2H, m), 3.93 (3H, s), 6.97-7.06 (2H, m), 7.20 (1H, d, J = 8.1 Hz), 7.48 (1H, br s), 7.71-7.77 (2H, m), 8.56 (1H, br s), 13.46 (1H, br s).	462	460
132		¹ H-NMR (DMSO-D ₆) δ: 1.50-1.73 (4H, m), 2.50-2.56 (2H, m), 2.69 (2H, t, J = 7.5 Hz), 2.84 (2H, t, J = 7.6 Hz), 3.21 (3H, s), 3.31-3.37 (2H, m), 3.91 (3H, s), 6.91-7.00 (2H, m), 7.13 (2H, d, J = 8.0 Hz), 7.24 (2H, d, J = 8.0 Hz), 7.64 (1H, d, J = 1.7 Hz), 8.50 (1H, d, J = 1.7 Hz), 12.13 (1H, br s).	456	454
133		¹ H-NMR (DMSO-D ₆) δ: 1.08 (3H, t, J = 7.1 Hz), 2.49 (3H, s), 3.10 (2H, t, J = 6.5 Hz), 3.46 (2H, q, J = 7.0 Hz), 3.79 (2H, t, J = 6.5 Hz), 3.93 (3H, s), 7.11 (2H, d, J = 9.5 Hz), 7.21 (1H, dd, J = 8.1, 1.4 Hz), 7.41 (1H, d, J = 1.6 Hz), 7.75 (1H, d, J = 8.1 Hz), 8.64 (1H, s), 12.96 (1H, br s).	429	427
134		¹ H-NMR (DMSO-D ₆) δ: 1.08 (3H, t, J = 7.1 Hz), 2.53 (2H, t, J = 7.6 Hz), 2.83 (2H, t, J = 7.5 Hz), 3.08 (2H, t, J = 6.6 Hz), 3.45 (2H, q, J = 7.0 Hz), 3.79 (2H, t, J = 6.5 Hz), 3.93 (3H, s), 7.08 (2H, d, J = 9.7 Hz), 7.24 (2H, d, J = 8.3 Hz), 7.32 (2H, d, J = 8.3 Hz), 8.58 (1H, s), 12.07 (1H, br s).	443	441
135		¹ H-NMR (DMSO-D ₆) δ: 1.56-1.63 (2H, m), 1.75-1.82 (2H, m), 2.51 (3H, s), 2.89 (2H, t, J = 7.6 Hz), 3.22 (3H, s), 3.36 (2H, t, J = 6.5 Hz), 3.94 (3H, s), 7.11 (2H, d, J = 9.2 Hz), 7.23 (1H, dd, J = 8.1, 1.4 Hz), 7.42 (1H, d, J = 1.4 Hz), 7.77 (1H, d, J = 8.1 Hz), 8.64 (1H, s), 12.96 (1H, s).	443	441
136		¹ H-NMR (DMSO-D ₆) δ: 1.56-1.63 (2H, m), 1.75-1.83 (2H, m), 2.89 (2H, t, J = 7.6 Hz), 3.22 (3H, s), 3.36 (2H, t, J = 6.5 Hz), 3.94 (3H, s), 7.11 (2H, d, J = 9.2 Hz), 7.23 (1H, dd, J = 8.1, 1.5 Hz), 7.42 (1H, d, J = 1.5 Hz), 7.77 (1H, d, J = 8.1 Hz), 8.64 (1H, s), 12.96 (1H, s).	463	461

[0371]

Table 1-18

Ex.	Structure	NMR	MS (M+H)	MS (M-H)
137		¹ H-NMR (DMSO-D ₆) δ: 1.11 (3H, t, J = 7.0 Hz), 2.49 (3H, s), 2.90 (2H, t, J = 6.7 Hz), 3.46 (2H, q, J = 7.0 Hz), 3.65 (2H, t, J = 6.7 Hz), 3.73 (3H, s), 6.83 (2H, d, J = 8.7 Hz), 7.00 (1H, d, J = 8.1 Hz), 7.23 (2H, d, J = 8.7 Hz), 7.23 (1H, brs), 7.68 (1H, d, J = 1.8 Hz), 7.71 (1H, d, J = 8.1 Hz), 8.53 (1H, d, J = 1.8 Hz), 12.84 (1H, s).	392	390
138		¹ H-NMR (DMSO-D ₆) δ: 1.11 (3H, t, J = 7.0 Hz), 2.91 (2H, t, J = 6.7 Hz), 3.47 (2H, q, J = 7.0 Hz), 3.66 (2H, t, J = 6.7 Hz), 3.74 (3H, s), 6.86 (2H, d, J = 9.0 Hz), 7.16 (1H, dd, J = 8.1, 1.5 Hz), 7.23 (2H, d, J = 9.0 Hz), 7.43 (1H, d, J = 1.5 Hz), 7.71 (1H, d, J = 8.1 Hz), 7.74 (1H, d, J = 2.1 Hz), 8.55 (1H, d, J = 1.8 Hz), 13.41 (1H, s).	412	410
139		¹ H-NMR (DMSO-D ₆) δ: 1.10 (3H, t, J = 7.0 Hz), 2.51 (2H, t, J = 7.6 Hz), 2.81 (2H, t, J = 7.6 Hz), 2.88 (2H, t, J = 6.7 Hz), 3.46 (2H, q, J = 7.0 Hz), 3.65 (2H, t, J = 6.7 Hz), 3.73 (3H, s), 6.79 (2H, d, J = 9.0 Hz), 7.09 (2H, d, J = 8.1 Hz), 7.18 (2H, d, J = 8.1 Hz), 7.21 (2H, d, J = 9.0 Hz), 7.62 (1H, d, J = 1.8 Hz), 8.48 (1H, d, J = 1.8 Hz), 12.18 (1H, s).	406	404
140		¹ H-NMR (DMSO-D ₆) δ: 1.10 (3H, t, J = 7.0 Hz), 2.93 (2H, t, J = 6.6 Hz), 3.46 (2H, q, J = 7.1 Hz), 3.66 (2H, t, J = 6.6 Hz), 3.91 (3H, s), 6.97 (2H, d, J = 9.3 Hz), 7.35 (2H, d, J = 8.1 Hz), 7.76 (1H, d, J = 1.8 Hz), 7.92 (2H, d, J = 8.1 Hz), 8.59 (1H, d, J = 2.1 Hz), 13.06 (1H, s).	414	412
141		¹ H-NMR (DMSO-D ₆) δ: 1.10 (3H, t, J = 7.0 Hz), 2.51 (3H, s), 2.92 (2H, t, J = 6.4 Hz), 3.43-3.49 (2H, m), 3.66 (2H, t, J = 6.7 Hz), 3.92 (3H, s), 6.96-7.07 (3H, m), 7.22-7.25 (1H, m), 7.73-7.77 (2H, m), 8.57 (1H, d, J = 1.8 Hz), 12.91 (1H, br s).	428	426
142		¹ H-NMR (DMSO-D ₆) δ: 1.10 (3H, t, J = 6.9 Hz), 2.93 (2H, t, J = 6.6 Hz), 3.46 (2H, q, J = 6.9 Hz), 3.66 (2H, t, J = 6.6 Hz), 3.93 (3H, s), 7.02 (2H, d, J = 9.3 Hz), 7.19 (1H, d, J = 7.8 Hz), 7.47 (1H, s), 7.73 (1H, d, J = 7.8 Hz), 7.81 (1H, s), 8.59 (1H, s), 13.47 (1H, s).	448	446
143		¹ H-NMR (DMSO-D ₆) δ: 0.88 (3H, t, J = 7.5 Hz), 1.50-1.61 (2H, m), 2.47 (3H, s), 3.45 (2H, t, J = 6.6 Hz), 3.72 (3H, s), 4.56 (2H, s), 6.82 (2H, d, J = 8.8 Hz), 6.99 (1H, dd, J = 8.0, 1.5 Hz), 7.20-7.26 (3H, m), 7.70 (2H, dd, J = 5.1, 3.0 Hz), 8.59 (1H, d, J = 2.1 Hz), 12.82 (1H, br s).	392	390
144		¹ H-NMR (DMSO-D ₆) δ: 0.88 (3H, t, J = 7.4 Hz), 1.50-1.61 (2H, m), 2.48-2.54 (2H, m), 2.80 (2H, t, J = 7.5 Hz), 3.44 (2H, t, J = 6.6 Hz), 3.71 (3H, s), 4.55 (2H, s), 6.79 (2H, d, J = 9.0 Hz), 7.08 (2H, d, J = 8.1 Hz), 7.17 (2H, d, J = 8.3 Hz), 7.21 (2H, d, J = 9.0 Hz), 7.64 (1H, d, J = 2.3 Hz), 8.54 (1H, d, J = 2.1 Hz), 12.09 (1H, br s).	406	404


[0372]

Table 1-19

Ex.	Structure	NMR	MS (M+H)	MS (M-H)
145		¹ H-NMR (DMSO-D ₆) δ: 0.88 (3H, t, J = 7.4 Hz), 1.51-1.62 (2H, m), 3.46 (2H, t, J = 6.6 Hz), 3.90 (3H, s), 4.59 (2H, s), 6.97 (2H, d, J = 9.7 Hz), 7.35 (2H, d, J = 8.3 Hz), 7.78 (1H, d, J = 2.1 Hz), 7.91 (2H, d, J = 8.6 Hz), 8.65 (1H, d, J = 2.1 Hz), 12.99 (1H, br s).	414	412
146		¹ H-NMR (DMSO-D ₆) δ: 0.88 (3H, t, J = 7.4 Hz), 1.51-1.62 (2H, m), 2.50-2.55 (2H, m), 2.83 (2H, t, J = 7.6 Hz), 3.45 (2H, t, J = 6.6 Hz), 3.90 (3H, s), 4.57 (2H, s), 6.96 (2H, d, J = 9.7 Hz), 7.12 (2H, d, J = 8.3 Hz), 7.23 (2H, d, J = 8.3 Hz), 7.71 (1H, d, J = 2.1 Hz), 8.59 (1H, d, J = 1.8 Hz), 12.12 (1H, br s).	442	440
147		¹ H-NMR (DMSO-D ₆) δ: 0.88 (3H, t, J = 7.4 Hz), 1.51-1.62 (2H, m), 2.49 (3H, s), 3.45 (2H, t, J = 6.6 Hz), 3.91 (3H, s), 4.59 (2H, s), 6.99 (2H, d, J = 9.7 Hz), 7.05 (1H, dd, J = 8.0, 1.5 Hz), 7.23 (1H, d, J = 1.4 Hz), 7.72-7.78 (2H, m), 8.63 (1H, d, J = 2.1 Hz), 12.89 (1H, br s).	428	426
148		¹ H-NMR (DMSO-D ₆) δ: 1.56-1.63 (2H, m), 1.74-1.82 (2H, m), 2.54 (2H, t, J = 7.6 Hz), 2.83-2.89 (4H, m), 3.21 (3H, s), 3.36 (2H, t, J = 6.5 Hz), 3.94 (3H, s), 7.09 (2H, d, J = 9.5 Hz), 7.25 (2H, d, J = 8.3 Hz), 7.33 (2H, d, J = 8.3 Hz), 8.58 (1H, s), 12.13 (1H, s).	457	455
149		¹ H-NMR (DMSO-D ₆) δ: 0.94 (6H, d, J = 6.7 Hz), 2.07-2.18 (1H, m), 2.49 (3H, s), 2.75 (2H, d, J = 7.2 Hz), 3.93 (3H, s), 7.10 (2H, d, J = 9.5 Hz), 7.22 (1H, dd, J = 8.1, 1.4 Hz), 7.40 (1H, d, J = 1.6 Hz), 7.76 (1H, d, J = 8.1 Hz), 8.59 (1H, s), 12.91 (1H, br s).	413	411
150		¹ H-NMR (DMSO-D ₆) δ: 0.94 (6H, d, J = 6.7 Hz), 2.06-2.17 (1H, m), 2.52 (2H, t, J = 7.6 Hz), 2.72 (2H, d, J = 7.2 Hz), 2.83 (2H, t, J = 7.5 Hz), 3.93 (3H, s), 7.08 (2H, d, J = 9.5 Hz), 7.23 (2H, d, J = 8.3 Hz), 7.32 (2H, d, J = 8.3 Hz), 8.53 (1H, s), 12.13 (1H, br s).	427	425
151		¹ H-NMR (DMSO-D ₆) δ: 0.94 (6H, d, J = 6.5 Hz), 2.07-2.19 (1H, m), 2.76 (2H, d, J = 7.2 Hz), 3.94 (3H, s), 7.16 (2H, d, J = 9.5 Hz), 7.35 (1H, dd, J = 8.1, 1.8 Hz), 7.61 (1H, d, J = 1.6 Hz), 7.74 (1H, d, J = 8.1 Hz), 8.62 (1H, s), 13.50 (1H, br s).	433	431
152		¹ H-NMR (DMSO-D ₆) δ: 0.93 (3H, t, J = 7.4 Hz), 1.35-1.44 (2H, m), 1.70-1.77 (2H, m), 2.51 (3H, s), 2.88 (2H, t, J = 7.6 Hz), 3.95 (3H, s), 7.11 (2H, d, J = 9.5 Hz), 7.23 (1H, dd, J = 8.1, 1.5 Hz), 7.42 (1H, d, J = 1.5 Hz), 7.77 (1H, d, J = 8.1 Hz), 8.64 (1H, s), 12.95 (1H, s).	413	411

[0373]

Table 1-20

Ex.	Structure	NMR	MS (M+H)	MS (M-H)
153		¹ H-NMR (DMSO-D ₆) δ: 0.93 (3H, t, J = 7.4 Hz), 1.34-1.43 (2H, m), 1.69-1.77 (2H, m), 2.54 (2H, t, J = 7.6 Hz), 2.83-2.88 (4H, m), 3.94 (3H, s), 7.09 (2H, d, J = 9.2 Hz), 7.25 (2H, d, J = 8.3 Hz), 7.33 (2H, d, J = 8.3 Hz), 8.58 (1H, s), 12.12 (1H, s).	427	425

[0374]

5 The formulation examples of the present invention include the following formulations. However, the present invention is not limited by such formulation examples.

[0375]

Formulation Example 1 (Production of capsule)

- | | | |
|----|-------------------------------|-------|
| 10 | 1) Compound of Example 1 | 30 mg |
| | 2) Microcrystalline cellulose | 10 mg |
| | 3) Lactose | 19 mg |
| | 4) Magnesium stearate | 1 mg |

15 1), 2), 3) and 4) are mixed and filled in a gelatin capsule.

[0376]

Formulation Example 2 (Production of tablet)

- | | | |
|----|--------------------------|------|
| | 1) Compound of Example 1 | 10 g |
| | 2) Lactose | 50 g |
| 20 | 3) Corn starch | 15 g |
| | 4) Carmellose calcium | 44 g |
| | 5) Magnesium stearate | 1 g |

25 The total amount of 1), 2), 3) and 30 g of 4) are kneaded with water, vacuum dried and sieved. The sieved powder is mixed with 14 g of 4) and 1 g of 5), and the mixture is tableted by a tableting machine. In this way, 1000 tablets containing 10 mg of the compound of Example 1 per tablet are obtained.

[0377]

Test Example 1: Evaluation of human GLUT9 inhibitory activity

30 A method for measuring GLUT9 inhibitory activity using human GLUT9 expressing cells transduced with Uricase/Hyper is

shown hereinafter. GLUT9 inhibitory activity (IC_{50} value) of the test compound was calculated based on the intracellular uptake amount of labeled uric acid ($[^{14}C]$ uric acid) transported by GLUT9.

5 [0378]

1. Preparation of human GLUT9 expression plasmid

Human GLUT9S coding region (NM_001001290.1) was amplified by PCR method using PrimeSTAR MAX DNA Polymerase (Takara Bio). The obtained PCR product was subjected to agarose gel
10 electrophoresis, and the desired size of PCR product was purified by QIAquick Gel Extraction Kit (QIAGEN). Next, pEBMulti-Bsd vector (Wako Pure Chemical Industries, Ltd.) was subjected to digestion using restriction enzymes *Sal*I and *Eco*R V, and linked to the PCR product containing GLUT9S by In-Fusion
15 reaction using In-Fusion HD Cloning Kit (Takara Bio). XL10-Gold Competent Cells (Agilent Technologies) were transformed in In-Fusion reaction solution, and the obtained transfectants were cultured with shaking overnight in LB medium to which Blasticidin S HCl (Thermo Fisher Scientific) was added. Human
20 GLUT9 expression plasmid DNA was prepared from the collected transfectants using EndoFree Plasmid Kit (Takara Bio).

[0379]

2. Establish of human GLUT9 stably expressing cell

HEK293T cells (American Type Culture Collection, DS
25 Pharma Biomedical) were seeded in a 6 well plate at 1×10^6 cells/well. Next day, human GLUT9 expression plasmid was transfected by using Lipofectamine 3000 Reagent (Thermo Fisher Scientific) as follows. Lipofectamine 3000 Reagent (5 μ L) was diluted with Opti-MEM (125 μ L, Thermo Fisher Scientific).
30 Separately, plasmid DNA (2.5 μ g) was diluted with Opti-MEM (125 μ L), and P3000 Reagent (5 μ L) was added thereto. The obtained solution was mixed with the previously obtained diluted solution, and the mixture was left stand at room temperature for 5 min, and added to the cells. Next day, the transfected
35 cells were diluted and reseeded in a 6 well plate at 4×10^5

cells/well, and drug selection was performed in the presence of 30 µg/mL of Blasticidin S HCl, and thereby, Blasticidin-resistant human GLUT9 stably expressing cell line was established. Mock cells were prepared by introducing pEBMulti-Bsd vector into HEK293T cells in a similar manner.

[0380]

3. Preparation of Uricase/Hyper expression plasmid

Uricase/Hyper expression plasmid was prepared by linking the coding regions of *Aspergillus Flavus*-derived uricase gene and pHyPer-Cyto vector (Evrogen) HyPer gene to the pEBMulti-Hyg vector (Wako Pure Chemical Industries, Ltd.) and then introducing it. The preparation method was performed by reference to Gout and Nucleic Acid Metabolism 2013, 37(2), 93-101, and seven repeated linker sequence of glycine-glycine-glycine-serine was prepared herein.

[0381]

4. Preparation of human GLUT9-Uricase/Hyper coexpressing cell

Human GLUT9 stably expressing cells were seeded at 3.6×10^7 cells per 225cm² flask. Next day, Uricase/Hyper expression plasmid was transfected by using Lipofectamine 3000 Reagent as follows. Lipofectamine 3000 Reagent (40 µL) was diluted with Opti-MEM (750 µL). Separately, plasmid DNA (20 µg) was diluted with Opti-MEM (750 µL), and P3000 Reagent (40 µL) was added thereto. The obtained solution was mixed with the previously obtained diluted solution, and the mixture was left stand at room temperature for 5 min, and added to the cells. Next day, transfected cells were collected to prepare human GLUT9-Uricase/Hyper coexpressing cells.

[0382]

5. Evaluation of GLUT9 inhibitory activity

Uricase/Hyper transiently-transfected human GLUT9 stably expressing cells or mock cells (blank) were seeded in a 96 well plate (Corning) at 1.6×10^5 cells/well, and cultured overnight at 37°C, 5% CO₂. D-MEM/high glucose (Wako Pure Chemical Industries, Ltd.) containing 10% Fetal Bovine Serum

(Lifetechnology) and 100 units/ml penicillin/100 µg/ml streptomycin (GIBCO) was used as a medium. High K⁺ buffer (129.8 mM KCl, 1.2 mM KH₂PO₄, 1.2 mM MgSO₄·7H₂O, 1.3 mM CaCl₂·2H₂O, 25 mM HEPES, pH 7.4 with 1 M Tris) and the medium were
5 mixed in equal amount to prepare Assay Buffer. The medium in each well was removed, and the test compound solution (final 1% DMSO) diluted with Assay Buffer was added thereto at 50 µl/well, and the mixture was left stand at room temperature for 30 to 60 min. For the solvent control and blank, Assay Buffer
10 containing DMSO alone was added at 50 µl/well, and the mixture was left stand at room temperature for 30 to 60 min. In addition, uric acid solution (containing [¹⁴C]uric acid as a tracer) diluted with Assay Buffer was added to each well at 15 µl/well (final 300 µM uric acid), and the uptake reaction was
15 performed at room temperature for 6 min. After the completion of the reaction, the cells were washed three times with ice-cooled Wash Buffer (Hank's Balanced Salt Solution containing 0.01% Bovine Serum Albumin) at 150 µl/well, and 0.1N aqueous NaOH solution was added thereto at 25 µl/well to dissolve the
20 cell. MicroScint-20 (Perkin-Elmer) was added thereto at 150 µl/well, the plate was shaken, and CPM of [¹⁴C] was measured by TopCount NXT (Perkin-Elmer).

Data was obtained by deducting average of CPM in blank well from average of CPM in each treated well. The inhibitory
25 rate of the test compound in each concentration was calculated from the following formula: $[(A-B)/A] \times 100$, A is data of solvent control, B is data of test compound treatment. IC₅₀ value (50% inhibition concentration) of the test compound was obtained by applying the inhibitory rate of the test compound in each
30 concentration to logistic curve.

[0383]

The results are shown in Table 2-1 to Table 2-6. With regard to Examples 7, 15, 22, 33, 47, 54, 60 and 76 in tables, the GLUT9 inhibitory rate at 10 µM of compound are shown
35 therein.

[0384]

Table 2-1

Example	humans GLUT9 inhibitory activity (IC ₅₀ value (μM))
1	0.0807
2	0.9461
3	0.1341
4	0.6124
5	0.0626
6	0.1409
7	9% inhibition at 10 μM
8	0.1504
9	0.4831
10	1.7769
11	0.0659
12	5.0637
13	0.2188
14	0.0914
15	5% inhibition at 10 μM
16	0.0819
17	0.6922
18	0.0942
19	3.0737
20	0.1198
21	0.0607
22	9% inhibition at 10 μM
23	0.9916
24	0.1342
25	0.0837
26	0.1543
27	0.2548
28	4.2515
29	0.1578
30	0.0892

[0385]

Table 2-2

Example	humans GLUT9 inhibitory activity (IC ₅₀ value (μM))
31	0.1336
32	0.0888
33	28% inhibition at 10 μM
34	0.1241
35	0.1193
36	0.1891
37	0.1106
38	0.8652
39	1.4456
40	0.3405
41	0.0890
42	6.5049
43	7.1322
44	1.6522
45	0.0506
46	2.7530
47	47% inhibition at 10 μM
48	7.1178
49	0.2005
50	0.1333
51	0.9618
52	0.1186
53	0.4600
54	43% inhibition at 10 μM
55	3.9895
56	0.1798
57	0.3443
58	0.0520
59	0.1315
60	45% inhibition at 10 μM

[0386]

Table 2-3

Example	humans GLUT9 inhibitory activity (IC ₅₀ value (μM))
61	5.5905
62	0.1267
63	0.6586
64	0.0987
65	0.0946
66	0.0770
67	0.1024
68	0.0578
69	0.8563
70	1.6718
71	0.1318
72	0.5258
73	0.1044
74	0.1410
75	0.0844
76	10% inhibition at 10 μM
77	0.0585
78	2.2102
79	0.3686
80	0.8421
81	0.1924
82	0.2450
83	0.0662
84	0.1167
85	0.8527
86	1.8180
87	1.7851
88	0.9282
89	1.7982
90	0.1397

[0387]

Table 2-4

Example	humans GLUT9 inhibitory activity (IC ₅₀ value (μM))
91	0.2470
92	0.1810
93	0.1837
94	0.0918
95	0.0576
96	0.0887
97	0.0795
98	0.0605
99	0.0514
100	0.0820
101	0.1642
102	0.7899
103	0.0720
104	0.1388
105	0.4533
106	0.1007
107	0.0627
108	0.0712
109	0.0799
110	0.2649
111	0.0826
112	0.5607
113	1.0001
114	0.0604
115	0.0497
116	0.0684
117	0.5432
118	0.1078
119	0.5012
120	0.5391

[0388]

Table 2-5

Example	humans GLUT9 inhibitory activity (IC ₅₀ value (μM))
121	0.5007
122	0.2579
123	0.0589
124	0.1826
125	0.2287
126	0.0497
127	0.0688
128	0.0732
129	0.0569
130	0.0521
131	0.1552
132	0.0519
133	0.3134
134	0.5248
135	0.0613
136	0.2502
137	0.0917
138	0.4993
139	0.1672
140	0.0713
141	0.0639
142	0.2263
143	0.1253
144	0.3270
145	0.1065
146	0.1387
147	0.0868
148	0.0807
149	0.0574
150	0.1018

[0389]

Table 2-6

Example	humans GLUT9 inhibitory activity (IC ₅₀ value (μM))
151	0.1064
152	0.0590
153	0.1023

Industrial Applicability


5 [0390]

Since Compound [I] or a pharmaceutically acceptable salt thereof has a GLUT9 inhibitory activity, it may be useful for the treatment or prophylaxis of a disease selected from the group consisting of hyperuricemia and gout.

10

CLAIMS

1. A compound of Formula [I], or a pharmaceutically acceptable salt thereof:



wherein

=X- is =C(R⁵)- or =N-;

-L-COOH is

(1) -COOH,

10 (2) -C(R⁷¹)(R⁷²)-COOH,

(3) -C(R⁷³)(R⁷⁴)-C(R⁷⁵)(R⁷⁶)-COOH, or

(4) -O-C(R⁷⁷)(R⁷⁸)-COOH;

n is 0, 1, or 2;

m is 0, 1, 2, or 3;

15 R¹ is each independently halogen or C₁₋₃ alkyl;

R² is

(1) halogen,

(2) hydroxy,

(3) cyano,

20 (4) C₁₋₆ alkyl optionally substituted with 1 to 3 substituents independently selected from the group consisting of cyano and C₁₋₃ alkoxy,

(5) halo C₁₋₆ alkyl,

(6) C₁₋₆ alkoxy,

25 (7) halo C₁₋₆ alkoxy,

(8) -COOR²¹ wherein R²¹ is hydrogen or C₁₋₃ alkyl,

(9) -CON(R²²)(R²³) wherein R²² and R²³ are each independently

hydrogen or C₁₋₃ alkyl,

(10) C₃₋₆ cycloalkyl or

(11) a 4- to 6-membered saturated heterocyclic group containing
 1 or 2 hetero atom as a ring atom in addition to the carbon
 5 atoms, wherein the hetero atom is independently selected from
 the group consisting of oxygen, nitrogen and sulfur atoms, and

R³ is

(1) hydrogen,

(2) halogen,

10 (3) cyano,

(4) C₁₋₃ alkyl,

(5) halo C₁₋₃ alkyl,

(6) C₁₋₃ alkoxy, or

(7) -COOR³¹ wherein R³¹ is hydrogen or C₁₋₃ alkyl or

15 R² and R³, together with the carbon atoms that they are
 bonded to, form a 4- to 6-membered saturated heterocycle
 containing 1 or 2 hetero atom as a ring atom in addition to the
 carbon atoms, wherein the hetero atom is independently selected
 from the group consisting of oxygen, nitrogen and sulfur atoms;

20 R⁴ is


(1) C₁₋₈ alkyl optionally substituted with 1 to 3 substituents
 independently selected from the following Group A,

(2) halo C₁₋₆ alkyl,

(3) -CON(R⁴¹)(R⁴²) wherein R⁴¹ and R⁴² are each independently
 25 hydrogen or C₁₋₆ alkyl,

(4) C₃₋₆ cycloalkyl optionally substituted with 1 to 3
 substituents independently selected from C₁₋₃ alkoxy, or

(5) a 4- to 6-membered saturated heterocyclic group containing
 1 or 2 hetero atom as a ring atom in addition to the carbon
 30 atoms, wherein the hetero atom is independently selected from
 the group consisting of oxygen, nitrogen and sulfur atoms, and
 wherein the ring atom in the heterocyclic group bonded to



is a carbon atom, and

Group A consists of

(a) hydroxy,

(b) C₁₋₃ alkoxy optionally substituted with one hydroxy or one

5 C₁₋₃ alkoxy,

(c) halo C₁₋₃ alkoxy,

(d) C₃₋₆ cycloalkyl optionally substituted with one hydroxy, and

(e) phenyl, and

R⁵ is hydrogen, halogen or C₁₋₃ alkyl or

10 R⁴ and R⁵, together with the carbon atoms that they are bonded to, form C₃₋₆ cycloalkane;

R⁶ are each independently halogen, hydroxy, C₁₋₃ alkyl or C₁₋₃ alkoxy; and

R⁷¹, R⁷², R⁷³, R⁷⁴, R⁷⁵, R⁷⁶, R⁷⁷, and R⁷⁸ are each
15 independently hydrogen or C₁₋₃ alkyl.

2. The compound according to Claim 1 or a pharmaceutically acceptable salt thereof, wherein -L-COOH is -COOH.

20 3. The compound according to Claim 1 or 2 or a pharmaceutically acceptable salt thereof, wherein n is 0 or 1.

4. The compound according to any one of Claims 1 to 3 or a pharmaceutically acceptable salt thereof, wherein m is 0 or 1.

25

5. The compound according to any one of Claims 1 to 4 or a pharmaceutically acceptable salt thereof, wherein R³ is

(1) hydrogen or

(2) halogen.

30

6. The compound according to any one of Claims 1 to 5 or a pharmaceutically acceptable salt thereof, wherein

R⁴ is

(1) C₁₋₈ alkyl optionally substituted with 1 to 3 substituents independently selected from the following Group A,

(2) halo C₁₋₆ alkyl, or

5 (3) -CON(R⁴¹)(R⁴²) wherein R⁴¹ and R⁴² are each independently hydrogen or C₁₋₆ alkyl, and

Group A consists of

(a) hydroxy,

(b) C₁₋₃ alkoxy optionally substituted with one hydroxy or one


10 C₁₋₃ alkoxy,

(c) halo C₁₋₃ alkoxy,

(d) C₃₋₆ cycloalkyl optionally substituted with one hydroxy, and


(e) phenyl.

15 7. A compound of the following formula or a pharmaceutically acceptable salt thereof:




(Example 21).

8. A compound of the following formula or a pharmaceutically
20 acceptable salt thereof:



(Example 35).


9. A compound of the following formula or a pharmaceutically acceptable salt thereof:



5

(Example 67).


10. A compound of the following formula or a pharmaceutically acceptable salt thereof:



(free form of Example 76).


10

11. A compound of the following formula or a pharmaceutically acceptable salt thereof:



(Example 84).


12. A compound of the following formula or a pharmaceutically acceptable salt thereof:



5

(Example 98).


13. A compound of the following formula or a pharmaceutically acceptable salt thereof:



(Example 107).


10

14. A compound of the following formula or a pharmaceutically acceptable salt thereof:



(Example 109).


15. A compound of the following formula or a pharmaceutically acceptable salt thereof:



5

(Example 116).


16. A compound of the following formula or a pharmaceutically acceptable salt thereof:



(Example 118).

10

17. A compound of the following formula or a pharmaceutically acceptable salt thereof:



(Example 126).

18. A pharmaceutical composition comprising the compound according to any one of Claims 1 to 17 or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

19. A GLUT9 inhibitor comprising the compound according to any one of Claims 1 to 17 or a pharmaceutically acceptable salt thereof.

20. An agent for the treatment or prophylaxis of a disease selected from the group consisting of hyperuricemia and gout, which comprises the compound according to any one of Claims 1 to 17 or a pharmaceutically acceptable salt thereof.

21. A method for the inhibition of GLUT9 in a mammal in need of such inhibition, which comprises administering a pharmaceutically effective amount of the compound according to any one of Claims 1 to 17 or a pharmaceutically acceptable salt thereof to the mammal.

22. A method for the treatment or prophylaxis of a disease selected from the group consisting of hyperuricemia and gout in a mammal in need of such treatment or prophylaxis, which comprises administering a pharmaceutically effective amount of the compound according to any one of Claims 1 to 17 or a pharmaceutically acceptable salt thereof to the mammal.

23. Use of the compound according to any one of Claims 1 to 17 or a pharmaceutically acceptable salt thereof for the manufacture of a GLUT9 inhibitor.

5

24. Use of the compound according to any one of Claims 1 to 17 or a pharmaceutically acceptable salt thereof for the manufacture of an agent for the treatment or prophylaxis of a disease selected from the group consisting of hyperuricemia and
10 gout.

25. The compound according to any one of Claims 1 to 17 or a pharmaceutically acceptable salt thereof, for use in the inhibition of GLUT9.

15

26. The compound according to any one of Claims 1 to 17 or a pharmaceutically acceptable salt thereof, for use in the treatment or prophylaxis of a disease selected from the group consisting of hyperuricemia and gout.

20