

NEW ENZYME INHIBITOR COMPOUNDS


FIELD OF THE INVENTION

5 The present invention relates to compounds which are inhibitors of SSAO activity. The invention also relates to pharmaceutical compositions comprising these compounds and to the use of these compounds in the treatment or prevention of medical conditions wherein inhibition of SSAO activity is beneficial, such as inflammatory diseases, immune disorders and the inhibition of tumour growth.

10

BACKGROUND ART

Semicarbazide-sensitive amine oxidase (SSAO) activity is an enzyme activity expressed by Vascular Adhesion Protein-1 (VAP-1) or Amine Oxidase, Copper
15 Containing 3 (AOC3), belongs to the copper-containing amine oxidase family of enzymes (EC.1.4.3.6). Therefore inhibitors of the SSAO enzyme may also modulate the biological functions of the VAP-1 protein. Members of this enzyme family are sensitive to inhibition by semicarbazide and utilize cupric ion and protein-derived topa quinone (TPQ) cofactor in the oxidative deamination of primary amines to
20 aldehydes, hydrogen peroxide, and ammonia according to the following reaction:



Known substrates for human SSAO include endogenous methylamine and
25 aminoacetone as well as some xenobiotic amines such as benzylamine [Lyles, *Int. J. Biochem. Cell Biol.* **1996**, 28, 259-274; Klinman, *Biochim. Biophys. Acta* **2003**, 1647(1-2), 131-137; Mátyus et al., *Curr. Med. Chem.* **2004**, 11(10), 1285-1298; O'Sullivan et al., *Neurotoxicology* **2004**, 25(1-2), 303-315]. In analogy with other copper-containing amine oxidases, DNA-sequence analysis and structure
30 determination suggest that the tissue-bound human SSAO is a homodimeric glycoprotein consisting of two 90–100 kDa subunits anchored to the plasma membrane by a single N-terminal membrane spanning domain [Morris et al., *J. Biol. Chem.* **1997**, 272, 9388-9392; Smith et al., *J. Exp. Med.* **1998**, 188, 17-27; Airene et al., *Protein Science* **2005**, 14, 1964-1974; Jakobsson et al., *Acta Crystallogr. D Biol. Crystallogr.* **2005**, 61(Pt 11), 1550-1562].
35

SSAO activity has been found in a variety of tissues including vascular and non-vascular smooth muscle tissue, endothelium, and adipose tissue [Lewinsohn, *Braz. J. Med. Biol. Res.* **1984**, 17, 223-256; Nakos & Gossrau, *Folia Histochem. Cytobiol.* **1994**, 32, 3-10; Yu et al., *Biochem. Pharmacol.* **1994**, 47, 1055-1059; Castillo et al., *Neurochem. Int.* **1998**, 33, 415-423; Lyles & Pino, *J. Neural. Transm. Suppl.* **1998**, 52, 239-250; Jaakkola et al., *Am. J. Pathol.* **1999**, 155, 1953-1965; Morin et al., *J. Pharmacol. Exp. Ther.* **2001**, 297, 563-572; Salmi & Jalkanen, *Trends Immunol.* **2001**, 22, 211-216]. In addition, SSAO protein is found in blood plasma and this soluble form appears to have similar properties as the tissue-bound form [Yu et al., *Biochem. Pharmacol.* **1994**, 47, 1055-1059; Kurkijärvi et al., *J. Immunol.* **1998**, 161, 1549-1557]. It has recently been shown that circulating human and rodent SSAO originates from the tissue-bound form [Göktürk et al., *Am. J. Pathol.* **2003**, 163(5), 1921-1928; Abella et al., *Diabetologia* **2004**, 47(3), 429-438; Stolen et al., *Circ. Res.* **2004**, 95(1), 50-57], whereas in other mammals the plasma/serum SSAO is also encoded by a separate gene called AOC4 [Schwelberger, *J. Neural. Transm.* **2007**, 114(6), 757-762].

The precise physiological role of this abundant enzyme has yet to be fully determined, but it appears that SSAO and its reaction products may have several functions in cell signalling and regulation. For example, recent findings suggest that SSAO plays a role in both GLUT4-mediated glucose uptake [Enrique-Tarancon et al., *J. Biol. Chem.* **1998**, 273, 8025-8032; Morin et al., *J. Pharmacol. Exp. Ther.* **2001**, 297, 563-572] and adipocyte differentiation [Fontana et al., *Biochem. J.* **2001**, 356, 769-777; Mercier et al., *Biochem. J.* **2001**, 358, 335-342]. In addition, SSAO has been shown to be involved in inflammatory processes where it acts as an adhesion protein for leukocytes [Salmi & Jalkanen, *Trends Immunol.* **2001**, 22, 211-216; Salmi & Jalkanen, in "Adhesion Molecules: Functions and Inhibition" K. Ley (Ed.), **2007**, pp. 237-251], and might also play a role in connective tissue matrix development and maintenance [Langford et al., *Cardiovasc. Toxicol.* **2002**, 2(2), 141-150; Göktürk et al., *Am. J. Pathol.* **2003**, 163(5), 1921-1928]. Moreover, a link between SSAO and angiogenesis has recently been discovered [Noda et al., *FASEB J.* **2008**, 22(8), 2928-2935], and based on this link it is expected that inhibitors of SSAO have an anti-angiogenic effect.

Several studies in humans have demonstrated that SSAO activity in blood plasma is elevated in conditions such as congestive heart failure, diabetes mellitus, Alzheimer's disease, and inflammation [Lewinsohn, *Braz. J. Med. Biol. Res.* **1984**, 17, 223–256; Boomsma et al., *Cardiovasc. Res.* **1997**, 33, 387–391; Ekblom, *Pharmacol. Res.* **1998**, 37, 87–92; Kurkijärvi et al., *J. Immunol.* **1998**, 161, 1549–1557; Boomsma et al., *Diabetologia* **1999**, 42, 233–237; Meszaros et al., *Eur. J. Drug Metab. Pharmacokinet.* **1999**, 24, 299–302; Yu et al., *Biochim. Biophys. Acta* **2003**, 1647(1–2), 193–199; Mátyus et al., *Curr. Med. Chem.* **2004**, 11(10), 1285–1298; O'Sullivan et al., *Neurotoxicology* **2004**, 25(1–2), 303–315; del Mar Hernandez et al., *Neurosci. Lett.* **2005**, 384(1–2), 183–187]. The mechanisms underlying these alterations of enzyme activity are not clear. It has been suggested that reactive aldehydes and hydrogen peroxide produced by endogenous amine oxidases contribute to the progression of cardiovascular diseases, diabetic complications and Alzheimer's disease [Callingham et al., *Prog. Brain Res.* **1995**, 106, 305–321; Ekblom, *Pharmacol. Res.* **1998**, 37, 87–92; Yu et al., *Biochim. Biophys. Acta* **2003**, 1647(1–2), 193–199; Jiang et al., *Neuropathol Appl Neurobiol.* **2008**, 34(2), 194–204]. Furthermore, the enzymatic activity of SSAO is involved in the leukocyte extravasation process at sites of inflammation where SSAO has been shown to be strongly expressed on the vascular endothelium [Salmi et al., *Immunity* **2001**, 14(3), 265–276; Salmi & Jalkanen, in “*Adhesion Molecules: Functions and Inhibition*” K. Ley (Ed.), **2007**, pp. 237–251]. Accordingly, inhibition of SSAO has been suggested to have a therapeutic value in the prevention of diabetic complications and in inflammatory diseases [Ekblom, *Pharmacol. Res.* **1998**, 37, 87–92; Salmi et al., *Immunity* **2001**, 14(3), 265–276; Salter-Cid et al., *J. Pharmacol. Exp. Ther.* **2005**, 315(2), 553–562].

SSAO knockout animals are phenotypically overtly normal but exhibit a marked decrease in the inflammatory responses evoked in response to various inflammatory stimuli [Stolen et al., *Immunity* **2005**, 22(1), 105–115]. In addition, antagonism of its function in wild type animals in multiple animal models of human disease (e.g. carrageenan-induced paw inflammation, oxazolone-induced colitis, lipopolysaccharide-induced lung inflammation, collagen-induced arthritis, endotoxin-induced uveitis) by the use of antibodies and/or small molecules has been shown to be protective in decreasing the leukocyte infiltration, reducing the severity of the disease phenotype and reducing levels of inflammatory cytokines and chemokines

[Kirton et al., *Eur. J. Immunol.* **2005**, 35(11), 3119-3130; Salter-Cid et al., *J. Pharmacol. Exp. Ther.* **2005**, 315(2), 553-562; McDonald et al., *Annual Reports in Medicinal Chemistry* **2007**, 42, 229-243; Salmi & Jalkanen, in "Adhesion Molecules: Functions and Inhibition" K. Ley (Ed.), **2007**, pp. 237-251; Noda et al., *FASEB J.* **2008** 22(4), 1094-1103; Noda et al., *FASEB J.* **2008**, 22(8), 2928-2935]. This anti-inflammatory protection seems to be afforded across a wide range of inflammatory models all with independent causative mechanisms, rather than being restricted to one particular disease or disease model. This would suggest that SSAO may be a key nodal point for the regulation of the inflammatory response, and it is therefore likely that SSAO inhibitors will be effective anti-inflammatory drugs in a wide range of human diseases. VAP-1 has also been implicated in the progression and maintenance of fibrotic diseases including those of the liver and lung. Weston and Adams (J Neural Transm. **2011**, 118(7), 1055-64) have summarised the experimental data implicating VAP-1 in liver fibrosis, and Weston et al (EASL Poster 2010) reported that blockade of VAP-1 accelerated the resolution of carbon tetrachloride induced fibrosis. In addition VAP-1 has been implicated in inflammation of the lung (e.g. Singh et al., **2003**, Virchows Arch 442:491-495) suggesting that VAP-1 blockers would reduce lung inflammation and thus be of benefit to the treatment of cystic fibrosis by treating both the pro-fibrotic and pro-inflammatory aspects of the disease.

SSAO (VAP-1) is up regulated in gastric cancer and has been identified in the tumour vasculature of human melanoma, hepatoma and head and neck tumours (Yoong KF, McNab G, Hubscher SG, Adams DH. (1998), J Immunol **160**, 3978-88.; Irjala H, Salmi M, Alanen K, Grenman R, Jalkanen S (2001), Immunol. **166**, 6937-6943; Forster-Horvath C, Dome B, Paku S, et al. (2004), Melanoma Res. **14**, 135-40.). One report (Marttila-Ichihara F, Castermans K, Auvinen K, Oude Egbrink MG, Jalkanen S, Griffioen AW, Salmi M. (2010), J Immunol. **184**, 3164-3173.) has shown that mice bearing enzymically inactive VAP-1 grow melanomas more slowly, and have reduced tumour blood vessel number and diameter. The reduced growth of these tumours was also reflected in the reduced (by 60-70%) infiltration of myeloid suppressor cells. Encouragingly VAP-1 deficiency had no effect on vessel or lymph formation in normal tissue.


Small molecules of different structural classes have previously been disclosed as SSAO inhibitors, for example in WO 02/38153 (tetrahydroimidazo[4,5-c]pyridine derivatives), in WO 03/006003 (2-indanylhhydrazine derivatives), in WO 2005/014530 (allylhhydrazine and hydroxylamine (aminooxy) compounds) and in WO 2007/120528 (allylamino compounds). Additional SSAO inhibitors are disclosed in PCT/EP2009/062011 and PCT/EP2009/062018.

The invention described here relates to a new class of SSAO inhibitors with biological, pharmacological, and pharmacokinetic characteristics that make them suitable for use as prophylactic or therapeutic agents in a wide range of human inflammatory diseases and immune disorders. This therapeutic capacity is designed to block SSAO enzyme action, reducing the levels of pro-inflammatory enzyme products (aldehydes, hydrogen peroxide and ammonia) whilst also decreasing the adhesive capacity of immune cells and correspondingly their activation and final extra-vasation. Diseases where such an activity is expected to be therapeutically beneficial include all diseases where immune cells play a prominent role in the initiation, maintenance or resolution of the pathology, such as multiple sclerosis, arthritis and vasculitis.

Detailed Description of the Invention

It has surprisingly been found that the compounds of formula (I) below are inhibitors of SSAO. They are therefore useful for the treatment or prevention of diseases in which inhibition of SSAO activity is beneficial, such as inflammation, inflammatory diseases, immune or autoimmune disorders, and inhibition of tumour growth.

According to the invention there is provided a compound of formula (I) or a pharmaceutically acceptable salt, or N-oxide thereof:



(I)

wherein

R^1 is phenyl or 6-membered heteroaryl, optionally substituted with one or more substituents selected from halogen, cyano, C_{1-4} -alkyl, halo- C_{1-4} -alkyl, C_{1-4} alkoxy- C_{1-4} alkyl, hydroxy- C_{1-4} -alkyl, cyano- C_{1-4} -alkyl, amino- C_{1-4} -alkyl, C_{1-4} -alkylamino- C_{1-4} -

alkyl, di(C₁₋₄-alkyl)amino-C₁₋₄-alkyl, -NR^{4A}R^{4B}, -NR⁶C(O)OR⁵, -NR⁶C(O)R⁵, -NR⁶C(O)NR^{4A}R^{4B}, -C(O)NR^{4A}R^{4B}, -C(O)R⁵, -C(O)OR⁵, and -NR⁶S(O)₂R⁵;

R² is -B-Q-[R³]_n or -B-R³;

5

wherein n = 1, 2, 3, or 4

B is a bond, O, NR⁴, -C(O)- or C₁₋₃-alkylene;

10 Q is saturated or partially unsaturated monocyclic 3-7 membered heterocyclic or C₃₋₇-cycloalkyl ring;

when R² is -B-Q-[R³]_n, R³ is independently selected from: 3-7 membered heterocycl-yl-, 3-7 membered heterocycl-yl-C₁₋₄-alkyl-, (3-7 membered heterocycl-yl-C₁₋₄-alkyl)-amino-C₁₋₄-alkyl-, amino-C₁₋₄-alkoxy-C₁₋₄-alkyl-, (amino-C₁₋₄-alkyl)-amino-C₁₋₄-alkyl-, -C₁₋₄-alkyl-NR⁶C(O)OR⁵, -C₁₋₄-alkyl-NR⁶C(O)NR^{4A}R^{4B}, -C₁₋₄-alkyl-C(O)NR^{4A}R^{4B}, (3-7 membered heterocycl-yl-C₁₋₄-alkyl)-C(O)-, -C₁₋₄-alkyl-C(O)OR⁵, -OC(O)R⁵, or

20 -C(O)NR^{9A}R^{9B} wherein R^{9A} and R^{9B} together with the nitrogen to which they are attached form a 3-7 membered cyclic amino group substituted with one or more substituents selected from: C₁₋₄-alkyl, C₁₋₄alkoxy-C₁₋₄alkyl-, C₃₋₇-cycloalkyl, or

-C(O)NR⁶R^{10B} wherein R^{10B} is:

25 (i) 3-7 membered heterocycl-yl- or 3-7 membered heterocycl-yl-C₁₋₄-alkyl-, or -C₁₋₄-alkyl-NR⁶C(O)R⁵; or

(ii) 5 or 6 membered heteroaryl-C₁₋₄-alkyl-, wherein the heteroaryl ring is optionally substituted with one or more substituents selected from halogen, cyano, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, and wherein the C₁₋₄-alkyl part is optionally substituted by one or more C₁₋₄-alkyl- groups, or the C₁₋₄-alkyl part is substituted with two C₁₋₄-alkyl groups which, together with the carbon atom to which they are attached, join together to form a spiro 3-6 membered cycloalkyl ring; and wherein

35 when R² is -B-R³, R³ is -NR⁶R^{11B}, and R^{11B} is 3-7 membered heterocycl-yl-C₁₋₄-alkyl-;

R^{4A} , R^{4B} and R^5 are each independently selected from hydrogen, C_{1-4} -alkyl-, 3-7 membered heterocyclyl- C_{1-4} -alkyl-, amino- C_{1-4} -alkyl-, 3-7 membered heterocyclyl-, - C_{1-4} -alkyl- $NR^6C(O)OR^5$, C_{3-7} -cycloalkyl, or

5

R^{4A} and R^{4B} together with the nitrogen to which they are attached form a 3-7 membered cyclic amino group, optionally substituted by one or more substituents selected from: C_{1-4} -alkyl, $-NR^{4A}R^{4B}$; and wherein


- 10 unless otherwise specified, 3-7 membered heterocyclyl, or the heterocyclyl part of the 3-7 membered heterocyclyl- C_{1-4} -alkyl-, (3-7 membered heterocyclyl- C_{1-4} -alkyl)-amino- C_{1-4} -alkyl-, or (3-7 membered heterocyclyl- C_{1-4} -alkyl)- $C(O)$ - group is optionally substituted with one or more substituents selected from oxo, C_{1-4} -alkyl-, $-C(O)OR^5$, $-C(O)R^5$, $-C(O)NR^{4A}R^{4B}$, $-NR^{4A}R^{4B}$, $-C_{1-4}$ -alkyl- $C(O)NR^{4A}R^{4B}$, or C_{1-4} alkoxy- C_{1-4} alkyl;
- 15 and

where present, the diradical $-C_{1-4}$ -alkyl- group directly attached to Q is optionally substituted with one or more groups independently selected from halogen, amino, methoxy, hydroxyl; and wherein

20

R^4 and R^6 are each independently selected from hydrogen or C_{1-4} -alkyl; and

X is selected from the radicals of formulae (1-16) wherein the bond marked * is attached to R^1 - and the bond marked ** is attached to $-R^2$:



wherein Y is selected from hydrogen, hydroxyl, amino, -NHR^6 , -OCH_3 ;

- 5 Z is selected from hydrogen, fluorine, hydroxyl, C_{1-4} -alkoxy, halo- C_{1-4} -alkyl, CONH_2 , cyano, SO_2NH_2 , amino, -NHR^6 ;

W is selected from H, C_{1-4} -alkyl, halo- C_{1-4} -alkyl,

- 10 **PROVIDED THAT** when R^2 is $\text{-B-Q-[R}^3\text{]}_n$, and R^3 is 3-7 membered heterocyclyl-, the R^3 heterocyclic ring atom directly bonded to Q is not nitrogen.

In a related embodiment, the present invention makes available a compound of formula (I) or a pharmaceutically acceptable salt, or N-oxide thereof:



wherein

R^1 is phenyl or 6-membered heteroaryl, optionally substituted with one or more substituents selected from halogen, cyano, C_{1-4} -alkyl, halo- C_{1-4} -alkyl, C_{1-4} alkoxy- C_{1-}

alkyl, hydroxy-C₁₋₄-alkyl, cyano-C₁₋₄-alkyl, amino-C₁₋₄-alkyl, C₁₋₄-alkylamino-C₁₋₄-alkyl, di(C₁₋₄-alkyl)amino-C₁₋₄-alkyl, -NR^{4A}R^{4B}, -NR⁶C(O)OR⁵, -NR⁶C(O)R⁵, -NR⁶C(O)NR^{4A}R^{4B}, -C(O)NR^{4A}R^{4B}, -C(O)R⁵, -C(O)OR⁵, and -NR⁶S(O)₂R⁵;

5 R² is -B-Q-[R³]_n or -B-R³;

wherein n = 1, 2, 3, or 4

B is a bond, O, NR⁴, -C(O)- or C₁₋₃-alkylene;

10

Q is saturated or partially unsaturated monocyclic 3-7 membered heterocyclic or C₃₋₇-cycloalkyl ring;

when R² is -B-Q-[R³]_n, R³ is independently selected from: 3-7 membered heterocycl-
 15 heterocycl-_{yl}, 3-7 membered heterocycl-_{yl}-C₁₋₄-alkyl-, (3-7 membered heterocycl-_{yl}-C₁₋₄-alkyl)-amino-C₁₋₄-alkyl-, amino-C₁₋₄-alkoxy-C₁₋₄-alkyl-, (amino-C₁₋₄-alkyl)-amino-C₁₋₄-alkyl-, -C₁₋₄-alkyl-NR⁶C(O)OR⁵, -C₁₋₄-alkyl-NR⁶C(O)NR^{4A}R^{4B}, -C₁₋₄-alkyl-C(O)NR^{4A}R^{4B}, (3-7 membered heterocycl-_{yl}-C₁₋₄-alkyl)-C(O)-, -C₁₋₄-alkyl-C(O)OR⁵, -OC(O)R⁵, or

20

-C(O)NR^{9A}R^{9B} wherein R^{9A} and R^{9B} together with the nitrogen to which they are attached form a 3-7 membered cyclic amino group substituted with one or more substituents selected from: C₁₋₄-alkyl, C₁₋₄alkoxy-C₁₋₄alkyl-, C₃₋₇-cycloalkyl, or

25 -C(O)NR⁶R^{10B} wherein R^{10B} is 3-7 membered heterocycl-_{yl} or 3-7 membered heterocycl-_{yl}-C₁₋₄-alkyl-, or -C₁₋₄-alkyl-NR⁶C(O)R⁵; or

when R² is -B-R³, R³ is -NR⁶R^{11B}, wherein R^{11B} is 3-7 membered heterocycl-_{yl}-C₁₋₄-alkyl-;

30

R^{4A}, R^{4B} and R⁵ are each independently selected from hydrogen, C₁₋₄-alkyl-, 3-7 membered heterocycl-_{yl}-C₁₋₄-alkyl-, amino-C₁₋₄-alkyl-, 3-7 membered heterocycl-_{yl}-, -C₁₋₄-alkyl-NR⁶C(O)OR⁵, C₃₋₇-cycloalkyl,

or R^{4A} and R^{4B} together with the nitrogen to which they are attached form a 3-7 membered cyclic amino group, optionally substituted by one or more substituents selected from: C_{1-4} -alkyl, $-NR^{4A}R^{4B}$;


- 5 unless otherwise specified, 3-7 membered heterocyclyl, or the heterocyclyl part of the 3-7 membered heterocyclyl- C_{1-4} -alkyl-, (3-7 membered heterocyclyl- C_{1-4} -alkyl)-amino- C_{1-4} -alkyl-, or (3-7 membered heterocyclyl- C_{1-4} -alkyl)-C(O)- group is optionally substituted with one or more substituents selected from C_{1-4} -alkyl-, $-C(O)OR^5$, $-C(O)R^5$, $-C(O)NR^{4A}R^{4B}$, $-NR^{4A}R^{4B}$, $-C_{1-4}$ -alkyl-C(O) $NR^{4A}R^{4B}$, or C_{1-4} alkoxy- C_{1-4} alkyl;
- 10 and

where present, the diradical $-C_{1-4}$ -alkyl- group directly attached to Q is optionally substituted with one or more groups independently selected from halogen, amino, methoxy, hydroxyl;

15

R^4 and R^6 are each independently selected from hydrogen or C_{1-4} -alkyl; and

X is selected from the radicals of formulae (1-16) wherein the bond marked * is attached to R^1 - and the bond marked ** is attached to $-R^2$:



wherein Y is selected from hydrogen, hydroxyl, amino, -NHR^6 , -OCH_3 ;

- 5 Z is selected from hydrogen, fluorine, hydroxyl, C_{1-4} -alkoxy, halo- C_{1-4} -alkyl, CONH_2 , cyano, SO_2NH_2 , amino, -NHR^6 ;

W is selected from H, C_{1-4} -alkyl, halo- C_{1-4} -alkyl,

- 10 **PROVIDED THAT** when R^2 is $\text{-B-Q-[R}^3\text{]}_n$, and R^3 is 3-7 membered heterocyclyl-, the heterocyclic ring atom directly bonded to Q is not nitrogen.

It is expected that compounds of the invention may be prepared in the form of hydrates, and solvates. Any reference herein, including the claims herein, to

- 15 “compounds with which the invention is concerned” or “compounds of the invention” or “the present compounds”, and the like, includes reference to salts, hydrates, and solvates of such compounds. The term ‘solvate’ is used herein to describe a molecular complex comprising the compound of the invention and a stoichiometric

amount of one or more pharmaceutically acceptable solvent molecules, for example, ethanol. The term 'hydrate' is employed when said solvent is water.

Individual compounds of the invention may exist in an amorphous form and /or
5 several polymorphic forms and may be obtained in different crystal habits. Any reference herein, including the claims herein, to "compounds with which the invention is concerned" or "compounds of the invention" or "the present compounds", and the like, includes reference to the compounds irrespective of amorphous or polymorphic form.

10

Since compounds of the invention have a nitrogen atom in an aromatic ring they may form N-oxides, and the invention includes compounds of the invention in their N-oxide form.

15 DEFINITIONS

The following definitions shall apply throughout the specification and the appended claims, unless otherwise stated or indicated.

20 The term "C₁₋₄-alkyl" denotes a straight or branched alkyl group having from 1 to 4 carbon atoms. For parts of the range C₁₋₄-alkyl all subgroups thereof are contemplated such as C₁₋₃-alkyl, C₁₋₂-alkyl, C₂₋₄-alkyl, C₂₋₃-alkyl and C₃₋₄-alkyl. Examples of said C₁₋₄-alkyl include methyl, ethyl, *n*-propyl, isopropyl, *n*-butyl, isobutyl, *sec*-butyl and *tert*-butyl.

25

Unless otherwise specified, the term "C₃₋₇-cycloalkyl" refers to a monocyclic saturated or partially unsaturated hydrocarbon ring system having from 3 to 7 carbon atoms. Examples of said C₃₋₇-cycloalkyl include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclohexenyl, cycloheptyl, and cycloheptenyl. For parts of
30 the range "C₃₋₇-cycloalkyl" all subgroups thereof are contemplated such as C₃₋₇-cycloalkyl, C₃₋₆-cycloalkyl, C₃₋₅-cycloalkyl, C₃₋₄-cycloalkyl, C₄₋₇-cycloalkyl, C₄₋₆-cycloalkyl, C₄₋₅-cycloalkyl, C₅₋₇-cycloalkyl, C₅₋₆-cycloalkyl, and C₆₋₇-cycloalkyl.

The term "C₁₋₄-alkoxy" refers to a straight or branched C₁₋₄-alkyl group which is
35 attached to the remainder of the molecule through an oxygen atom. For parts of the

range C₁₋₄-alkoxy, all subgroups thereof are contemplated such as C₁₋₃-alkoxy, C₁₋₂-alkoxy, C₂₋₄-alkoxy, C₂₋₃-alkoxy and C₃₋₄-alkoxy. Examples of said C₁₋₄-alkoxy include methoxy, ethoxy, *n*-propoxy, isopropoxy, *n*-butoxy, isobutoxy, *sec*-butoxy and *tert*-butoxy.

5

The term "hydroxy-C₁₋₄-alkyl" denotes a straight or branched C₁₋₄-alkyl group that has one or more hydrogen atoms thereof replaced with OH. Examples of said hydroxy-C₁₋₄-alkyl include hydroxymethyl, 2-hydroxyethyl and 2,3-dihydroxypropyl.

10 The term "halo-C₁₋₄-alkyl" denotes a straight or branched C₁₋₄-alkyl group that has one or more hydrogen atoms thereof replaced with halogen. Examples of said halo-C₁₋₄-alkyl include fluoromethyl, trifluoromethyl, trichloromethyl and 2-fluoroethyl.

The term "cyano-C₁₋₄-alkyl" denotes a straight or branched C₁₋₄-alkyl group that has
15 one or more hydrogen atoms thereof replaced with cyano. Examples of said cyano-C₁₋₄-alkyl include cyanomethyl, 2-cyanoethyl and 3-cyanopropyl.

The term "amino-C₁₋₄-alkyl" denotes a straight or branched C₁₋₄-alkyl group substituted with an amino group. Examples of said amino-C₁₋₄-alkyl group include
20 aminomethyl and 2-aminoethyl.

The term "C₁₋₄-alkylamino-C₁₋₄-alkyl" denotes an amino-C₁₋₄-alkyl group as defined above, wherein the amino group is substituted with a straight or branched C₁₋₄-alkyl group. Examples of said C₁₋₄-alkylamino-C₁₋₄-alkyl include methylaminoethyl and
25 ethylaminopropyl.

The term "di(C₁₋₄-alkyl)amino-C₁₋₄-alkyl" denotes an amino-C₁₋₄-alkyl group as defined above, wherein the amino group is disubstituted with straight or branched C₁₋₄-alkyl groups, which can be the same or different. Examples of said di(C₁₋₄-alkyl)amino-C₁₋₄-alkyl include
30 *N,N*-dimethylaminomethyl, *N*-ethyl-*N*-methylaminoethyl and *N,N*-diethylaminomethyl.

The terms "heteroaryl" and "heteroaromatic ring" denote a monocyclic heteroaromatic ring comprising 5 to 6 ring atoms in which one or more of the ring
35 atoms are other than carbon, such as nitrogen, sulphur or oxygen. Examples of

heteroaryl groups include furyl, pyrrolyl, thienyl, oxazolyl, isoxazolyl, imidazolyl, thiazolyl, isothiazolyl, pyridinyl, pyrimidinyl, tetrazolyl, pyrazolyl, pyridazinyl, pyrazinyl and thiadiazolyl.

- 5 The terms "heterocyclyl" and "heterocyclic ring" denote a non-aromatic, fully saturated or partially unsaturated, preferably fully saturated, monocyclic ring system having from 3 to 7 ring atoms, especially 5 or 6 ring atoms, in which one or more of the ring atoms are other than carbon, such as nitrogen, sulphur or oxygen. Examples of heterocyclic groups include piperidinyl, morpholinyl, homomorpholinyl, azepanyl, piperazinyl, oxo-piperazinyl, diazepinyl, tetrahydropyridinyl, tetrahydropyranyl, pyrrolidinyl, tetrahydrofuranlyl, and dihydropyrrolyl, groups.

The term "heterocyclic-C₁₋₄-alkyl" refers to a heterocyclic ring that is directly linked to a straight or branched C₁₋₄-alkyl group via a carbon or nitrogen atom of said ring.

- 15 Examples of said heterocyclic-C₁₋₄-alkyl include piperidin-4-ylmethyl, piperidin-1-ylmethyl, morpholin-4-yl-methyl and piperazin-4-ylmethyl. The C₁₋₄-alkyl part, which includes methylene, ethylene, propylene or butylene, is optionally substituted by one or more substituents selected from halogen, amino, methoxy, or hydroxyl.

- 20 The term "C₁₋₃-alkylene" denotes a straight or branched divalent saturated hydrocarbon chain having from 1 to 3 carbon atoms. The C₁₋₃-alkylene chain may be attached to the rest of the molecule and to the radical group through one carbon within the chain or through any two carbons within the chain. Examples of C₁₋₃-alkylene radicals include methylene [-CH₂-], 1,2-ethylene [-CH₂-CH₂-], 1,1-ethylene [-CH(CH₃)-], 1,2-propylene [-CH₂-CH(CH₃)-] and 1,3-propylene [-CH₂-CH₂-CH₂-].
- 25 When referring to a "C₁₋₃-alkylene" radical, all subgroups thereof are contemplated, such as C₁₋₂-alkylene and C₂₋₃-alkylene.

- "Halogen" refers to fluorine, chlorine, bromine or iodine, preferably fluorine and chlorine, most preferably fluorine.

"Hydroxy" refers to the -OH radical.

"Cyano" refers to the -CN radical.

“Oxo” refers to the carbonyl group =O.

“Optional” or “optionally” means that the subsequently described event or circumstance may but need not occur, and that the description includes instances
5 where the event or circumstance occurs and instances in which it does not.

“Pharmaceutically acceptable” means being useful in preparing a pharmaceutical composition that is generally safe, non-toxic and neither biologically nor otherwise undesirable and includes being useful for veterinary use as well as human
10 pharmaceutical use.

“Treatment” as used herein includes prophylaxis of the named disorder or condition, or amelioration or elimination of the disorder once it has been established.

15 “An effective amount” refers to an amount of a compound that confers a therapeutic effect on the treated subject. The therapeutic effect may be objective (i.e., measurable by some test or marker) or subjective (i.e., subject gives an indication of or feels an effect).

20 “Prodrugs” refers to compounds that may be converted under physiological conditions or by solvolysis to a biologically active compound of the invention. A prodrug may be inactive when administered to a subject in need thereof, but is converted *in vivo* to an active compound of the invention. Prodrugs are typically rapidly transformed *in vivo* to yield the parent compound of the invention, *e.g.* by
25 hydrolysis in the blood. The prodrug compound usually offers advantages of solubility, tissue compatibility or delayed release in a mammalian organism (see Silverman, R. B., *The Organic Chemistry of Drug Design and Drug Action*, 2nd Ed., Elsevier Academic Press (2004), pp. 498-549). Prodrugs of a compound of the invention may be prepared by modifying functional groups, such as a hydroxy,
30 amino or mercapto groups, present in a compound of the invention in such a way that the modifications are cleaved, either in routine manipulation or *in vivo*, to the parent compound of the invention. Examples of prodrugs include, but are not limited to, acetate, formate and succinate derivatives of hydroxy functional groups or phenyl carbamate derivatives of amino functional groups.

Throughout the specification and the appended claims, a given chemical formula or name shall also encompass all salts, hydrates, solvates, N-oxides and prodrug forms thereof. Further, a given chemical formula or name shall encompass all tautomeric and stereoisomeric forms thereof. Tautomers include enol and keto forms. Stereoisomers include enantiomers and diastereomers. Enantiomers can be present in their pure forms, or as racemic (equal) or unequal mixtures of two enantiomers. Diastereomers can be present in their pure forms, or as mixtures of diastereomers. Diastereomers also include geometrical isomers, which can be present in their pure *cis* or *trans* forms or as mixtures of those.

The compounds of formula (I) may be used as such or, where appropriate, as pharmacologically acceptable salts (acid or base addition salts) thereof. The pharmacologically acceptable addition salts mentioned below are meant to comprise the therapeutically active non-toxic acid and base addition salt forms that the compounds are able to form. Compounds that have basic properties can be converted to their pharmaceutically acceptable acid addition salts by treating the base form with an appropriate acid. Exemplary acids include inorganic acids, such as hydrogen chloride, hydrogen bromide, hydrogen iodide, sulphuric acid, phosphoric acid; and organic acids such as formic acid, acetic acid, propanoic acid, hydroxyacetic acid, lactic acid, pyruvic acid, glycolic acid, maleic acid, malonic acid, oxalic acid, benzenesulphonic acid, toluenesulphonic acid, methanesulphonic acid, trifluoroacetic acid, fumaric acid, succinic acid, malic acid, tartaric acid, citric acid, salicylic acid, *p*-aminosalicylic acid, pamoic acid, benzoic acid, ascorbic acid and the like. Exemplary base addition salt forms are the sodium, potassium, calcium salts, and salts with pharmaceutically acceptable amines such as, for example, ammonia, alkylamines, benzathine, and amino acids, such as, e.g. arginine and lysine. The term addition salt as used herein also comprises solvates which the compounds and salts thereof are able to form, such as, for example, hydrates, alcoholates and the like.

The group X

In the compounds of the invention, X may be selected from any one of the radicals of formula 1-16.

Currently preferred embodiments of the invention include those where X is:

- the formula 1 and R^1 , R^2 , Y, Z and W are as defined above; or
the formula 2 and R^1 , R^2 , Y, Z and W are as defined above; or
the formula 3 and R^1 , R^2 , Y, and Z are as defined above; or
the formula 4 and R^1 , R^2 , Y and W are as defined above; or
5 the formula 5 and R^1 , R^2 , and Y are as defined above; or
the formula 6 and R^1 , R^2 , Y, and Z are as defined above.

The group B

- In an embodiment of the invention, B is a bond, O, NR^4 such as NH, NCH_3 , or
10 NCH_2CH_3 , $-C(O)-$ or C_{1-3} alkylene such as methylene, ethylene or propylene radicals.
In a currently preferred embodiment B is a bond, $-C(O)-$ or methylene. In another
preferred embodiment B is a bond.

The group Y

- 15 In a currently preferred embodiment of the invention Y is selected from hydrogen,
hydroxyl, amino (NH_2), $-NHR^6$ such as $NHCH_3$, $NHCH_2CH_3$, or $-OCH_3$. In another
currently preferred embodiment Y is H, OH, or NH_2 . In an alternative currently
preferred embodiment Y is hydrogen

20 The group Z

Z is selected from hydrogen, fluorine, hydroxyl, C_{1-4} -alkoxy such as methoxy or
ethoxy, halo- C_{1-4} -alkyl such as fluoromethoxy, difluoromethoxy or trimethoxy,
 $CONH_2$, cyano, SO_2NH_2 , amino, $-NHR^6$ such as $NHCH_3$, $NHCH_2CH_3$. In a presently
preferred embodiment of the invention Z is hydrogen or hydroxyl.

25

The group W

- In a currently preferred embodiment of the invention W is selected from H, C_{1-4} -alkyl
such as methyl, ethyl, propyl, isopropyl, or halo- C_{1-4} -alkyl such as fluoromethyl,
difluoromethyl or trifluoromethyl. In another currently preferred embodiment W is
30 hydrogen.

The group R^1

- In one embodiment of the invention R^1 is phenyl or 6-membered heteroaryl such as
pyridine, pyridazine, pyrimidine, pyrazine, optionally substituted with one or more
35 substituents selected from halogen such as chloro or fluoro, C_{1-4} -alkyl such

as methyl, ethyl, propyl or isopropyl, halo-C₁₋₄-alkyl such as fluoromethyl, difluoromethyl or trifluoromethyl, C₁₋₄alkoxy-C₁₋₄alkyl, hydroxy-C₁₋₄-alkyl such as hydroxymethyl or hydroxyethyl, cyano-C₁₋₄-alkyl such as cyanomethyl or cyanoethyl, amino-C₁₋₄-alkyl such as aminomethyl, aminoethyl or aminopropyl, C₁₋₄-alkylamino-C₁₋₄-alkyl, di(C₁₋₄-alkyl)amino-C₁₋₄-alkyl, -NR^{4A}R^{4B}, -NR⁶C(O)OR⁵, -NR⁶C(O)R⁵, -NR⁶C(O)NR^{4A}R^{4B}, -C(O)NR^{4A}R^{4B}, -C(O)R⁵, -C(O)OR⁵, and -NR⁶S(O)₂R⁵.

In a currently preferred embodiment of the invention R¹ is optional substituted with one or more substituents selected from halogen such as fluoro or chloro, cyano, hydroxyl, C₁₋₄-alkyl such as methyl or ethyl, halo-C₁₋₄-alkyl such as fluoromethyl, difluoromethyl or trifluoromethyl, C₁₋₄alkoxy-C₁₋₄alkyl, hydroxy-C₁₋₄-alkyl, cyano-C₁₋₄-alkyl such as cyanomethyl or cyanoethyl, amino-C₁₋₄-alkyl, C₁₋₄-alkylamino-C₁₋₄-alkyl, di(C₁₋₄-alkyl)amino-C₁₋₄-alkyl, -NR^{4A}R^{4B}.

In another currently preferred embodiment R¹ is heteroaryl such as pyridine-2-yl, pyridine-3-yl or pyridine-4-yl optionally substituted with one or more substituents selected from fluoro, chloro, and C₁₋₄-alkyl such as methyl, ethyl, propyl, or isopropyl.

In an alternative embodiment R¹ is phenyl, optionally substituted at one or more of the para-, meta- and ortho- positions by one or more substituents selected from hydrogen, fluoro, chloro, cyano, hydroxyl, C₁₋₄-alkyl such as methyl, ethyl, propyl or isopropyl, or fluoromethyl, difluoromethyl, or trifluoromethyl.

In a currently preferred embodiment R¹ is phenyl substituted at the para position by a substituent selected from, fluoro, chloro, cyano, hydroxyl, C₁₋₄-alkyl such as methyl, ethyl, propyl or isopropyl, or fluoromethyl, difluoromethyl, or trifluoromethyl. In an alternative currently preferred embodiment the para substituent is selected from fluoro, chloro or methyl.

In another currently preferred embodiment R¹ is phenyl substituted at the meta-position by hydrogen.

In a further currently preferred embodiment R^1 is phenyl substituted at the ortho position by a substituent selected from hydrogen, fluoro, methyl, fluoromethyl, difluoromethyl, or trifluoromethyl. In another preferred embodiment R^1 is phenyl substituted at the ortho position by hydrogen, fluoro or methyl.

5

In a currently preferred embodiment of the invention R^1 is a mono, di, or tri substituted phenyl ring wherein the ortho, meta and/or para positions may be any combination of the substituents discussed above.

10 In a preferred embodiment the optional substituents of R^1 have a length of 4 atoms or fewer, preferably of 3 atoms or fewer, more preferably of 2 atoms or fewer.

The group R^2

In one currently preferred embodiment of the invention R^2 is $-B-Q-[R^3]_n$. n can be 1,
15 2, 3, or 4. In another currently preferred embodiment n is 1 or 2.

The ring Q is a saturated or partially unsaturated monocyclic 3-7 membered heterocyclic or C_{3-7} -cycloalkyl ring substituted with R^3 . In a currently preferred embodiment Q is a 7-membered saturated or partially unsaturated 7-membered
20 heterocyclic ring such as a homomorpholine ring, or a bridged homomorpholine ring wherein the bridge is formed by an ethylene or propylene radical, or a 7-membered cycloalkyl ring such as cycloheptane.

In an alternative preferred embodiment Q is a 5- or 6-membered saturated or
25 partially unsaturated 5 or 6 membered heterocyclic such as tetrahydrofuranyl, tetrahydropyranyl, piperidinyl, piperazinyl, morpholinyl, cyclohexyl, or any of the foregoing rings comprising a bridge formed by an ethylene or propylene radical, or a 5 or 6-membered cycloalkyl ring such cyclopentyl or cyclohexyl. In an embodiment Q is piperidinyl, piperazinyl, or morpholinyl.

30

In a currently preferred embodiment R^2 is $-B-Q-[R^3]_n$, wherein R^3 is selected from:

(i) 3-7 membered heterocyclyl- such as 2-, or 4-pyrrolidyl, 2-, 3-, or 4-piperidinyl, 2-, or 3-piperazinyl, or 2- or 3-morpholinyl; 3-7 membered heterocyclyl- C_{1-4} -alkyl- such as piperidin-4-ylmethyl, piperidin-1-ylmethyl, morpholin-4-yl-methyl, morpholin-
35 2-yl-methyl, and morpholin-3-yl-methyl and piperazin-4-ylmethyl, piperazin-2-

ylmethyl or piperazin-3-ylmethyl, or piperidin-4-ylethyl, piperidin-1-ylethyl, morpholin-4-yl-ethyl, morpholin-2-yl-ethyl, and morpholin-3-yl-ethyl and piperazin-4-ylethyl, piperazin-2-ylethyl or piperazin-3-ylethyl, or piperidin-4-ylpropyl, piperidin-1-ylpropyl, morpholin-4-yl-propyl, morpholin-2-yl-propyl, and morpholin-3-yl-propyl and
 5 piperazin-4-ylpropyl, piperazin-2-ylpropyl or piperazin-3-ylpropyl, or piperidin-4-ylbutyl, piperidin-1-ylbutyl, morpholin-4-yl-butyl, morpholin-2-yl-butyl, and morpholin-3-yl-butyl and piperazin-4-ylbutyl, piperazin-2-ylbutyl or piperazin-3-ylbutyl; (3-7 membered heterocyclyl-C₁₋₄-alkyl)-amino-C₁₋₄-alkyl- such as (piperidine-4-ylmethyl)aminomethyl, amino-C₁₋₄-alkoxy-C₁₋₄-alkyl-, (amino-C₁₋₄-alkyl)-amino-C₁₋₄-
 10 alkyl)-, -C₁₋₄-alkyl-NR⁶C(O)OR⁵, -C₁₋₄-alkyl-NR⁶C(O)NR^{4A}R^{4B}, -C₁₋₄-alkyl-C(O)NR^{4A}R^{4B}, (3-7 membered heterocyclyl-C₁₋₄-alkyl)-C(O)-, -C₁₋₄-alkyl-C(O)OR⁵, -OC(O)R⁵, or

(ii) -C(O)NR^{9A}R^{9B} wherein R^{9A} and R^{9B} together with the nitrogen to which they
 15 are attached form a 3-7 membered cyclic amino group substituted with one or more substituents selected from: C₁₋₄-alkyl, C₁₋₄alkoxy-C₁₋₄alkyl-, C₃₋₇-cycloalkyl. In a preferred embodiment the cyclic amino group is pyrrolidyl, piperidinyl, piperazinyl, or morpholinyl each of which is substituted on a ring carbon or nitrogen atom by one or more substituents selected from methyl, ethyl, propyl, iso-propyl, n-butyl, sec-butyl,
 20 tert-butyl, methoxyethyl, cyclopropyl or cyclobutyl. In a currently preferred embodiment the cyclic amino group is piperazinyl substituted on the 4-position by methyl, ethyl, propyl, iso-propyl, sec-butyl, or cyclopropyl, or

(iii) -C(O)NR⁶R^{10B} wherein R^{10B} is 3-7 membered heterocyclyl- such as defined
 25 above, or 3-7 membered heterocyclyl-C₁₋₄-alkyl- such as defined above, or -C₁₋₄-alkyl-NR⁶C(O)R⁵; or

R^{10B} is 5 or 6 membered heteroaryl-C₁₋₄-alkyl- such as tetrazolylmethyl, wherein the heteroaryl ring is optionally substituted with one or more substituents selected from
 30 C₁₋₄-alkyl or halo-C₁₋₄-alkyl, and wherein the C₁₋₄-alkyl part of the heteroaryl-C₁₋₄-alkyl- group is optionally substituted by one or more C₁₋₄-alkyl- groups, or the C₁₋₄-alkyl part is substituted with two C₁₋₄-alkyl groups which, together with the carbon atom to which they are attached, join together to form a spiro 3-6 membered cycloalkyl ring. In a preferred embodiment R^{10B} is tetrazolylmethyl-, wherein the
 35 tetrazole group is optionally substituted with one or more substituents selected from

C₁₋₄-alkyl or halo-C₁₋₄-alkyl, and wherein the methyl of the tetrazolylmethyl is substituted with two C₁₋₄-alkyl groups which, together with the carbon atom to which they are attached, join together to form a spiro cyclopropyl, cyclobutyl or cyclopentyl group.

5

The groups R^{4A}, R^{4B} and R⁵ are each independently selected from hydrogen, C₁₋₄-alkyl- such as methyl, ethyl, propyl, iso-propyl, n-butyl, sec-butyl, tert-butyl, 3-7 membered heterocyclyl-C₁₋₄-alkyl- as defined previously, amino-C₁₋₄-alkyl- such as aminomethyl, amino ethyl, 3-7 membered heterocyclyl- as defined above, -C₁₋₄-alkyl-NR⁶C(O)OR⁵, or C₃₋₇-cycloalkyl such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl,

or R^{4A} and R^{4B} together with the nitrogen to which they are attached form a 3-7 membered cyclic amino group such as pyrrolidyl, piperidinyl, homopiperidinyl, piperazinyl, homopiperizinyl or morpholinyl, optionally substituted by one or more substituents selected from: C₁₋₄-alkyl such as methyl, ethyl, propyl, iso-propyl, n-butyl, sec-butyl, tert-butyl, -NR^{4A}R^{4B} such as -NH₂, -NHCH₃, NHCH₂CH₃, or NH(CH₃)₂.

In the currently preferred embodiments the 3-7 membered heterocyclyl (other than the ring Q), or the heterocyclyl part of the 3-7 membered heterocyclyl-C₁₋₄-alkyl-, (3-7 membered heterocyclyl-C₁₋₄-alkyl)-amino-C₁₋₄-alkyl-, or (3-7 membered heterocyclyl-C₁₋₄-alkyl)-C(O)- group is optionally substituted with one or more substituents selected from oxo, C₁₋₄-alkyl- such as methyl, ethyl, propyl, iso-propyl, n-butyl, sec-butyl, tert-butyl, -C(O)OR⁵, -C(O)R⁵, -C(O)NR^{4A}R^{4B}, -NR^{4A}R^{4B} such as -NH₂, -NHCH₃, NHCH₂CH₃, or N(CH₃)₂, -C₁₋₄-alkyl-C(O)NR^{4A}R^{4B}, or C₁₋₄alkoxy-C₁₋₄alkyl such as methoxyethyl.

In a presently preferred embodiment the R³ group includes a divalent radical -C₁₋₄-alkyl- directly attached to the Q ring, such that R³ may be, for example 3-7 membered heterocyclyl-C₁₋₄-alkyl-, (3-7 membered heterocyclyl-C₁₋₄-alkyl)-amino-C₁₋₄-alkyl-, amino-C₁₋₄-alkoxy-C₁₋₄-alkyl-, (amino-C₁₋₄-alkyl)-amino-C₁₋₄-alkyl-, -C₁₋₄-alkyl-NR⁶C(O)OR⁵, -C₁₋₄-alkyl-NR⁶C(O)NR^{4A}R^{4B}, or -C₁₋₄-alkyl-C(O)NR^{4A}R^{4B}. In a preferred embodiment that -C₁₋₄-alkyl- radical is optionally substituted with one or more groups independently selected from halogen, amino, methoxy, and hydroxyl.

35

In an embodiment the $-C_{1-4}$ -alkyl- radical is selected from methylene, ethylene, propylene or butylene, any of which is optionally substituted by one or more groups independently selected from halogen, amino, methoxy, and hydroxyl. For example, the R^3 group includes $-CH_2-C(O)NR^{4A}R^{4B}$, $-(CH_2)_2-C(O)NR^{4A}R^{4B}$, $-(CH_2)_3-$
 5 $C(O)NR^{4A}R^{4B}$ or 3-7 membered heterocyclyl- CH_2- , 3-7 membered heterocyclyl- $(CH_2)_2-$, or 3-7 membered heterocyclyl- $(CH_2)_3-$.

R^4 and R^6 are each independently selected from hydrogen or C_{1-4} -alkyl such as methyl, ethyl, propyl, iso-propyl, n-butyl, sec-butyl, tert-butyl; and

10


In another embodiment, R^2 is $-B-R^3$ and R^3 is $-NR^6R^{11B}$, wherein R^{11B} is 3-7 membered heterocyclyl- C_{1-4} -alkyl- as defined previously and R^6 is as defined previously;

15 In an alternative currently preferred embodiment R^2 is $-B-Q-[R^3]_n$, and R^3 is:

(i) $-C(O)NR^6R^{10B}$ where R^6 is methyl or hydrogen and R^{10B} is a 3-7 membered heterocyclyl- such as piperidinyl including piperidine-4-yl and 1-methylpiperidine-4-yl or R^{10B} is a 3-7 membered heterocyclyl- C_{1-4} -alkyl- including morpholine-4-ylmethyl, morpholine-4-ylethyl, morpholine-4-ylpropyl, piperidine-4-ylmethyl-, piperidine-4-
 20 ylethyl-, piperidine-4-ylpropyl-, piperazine-1-ylmethyl, or piperazine-1-ylethyl wherein the nitrogen atom in the piperidine 1-position or the piperazine 4-position is substituted with a substituent selected from hydrogen, methyl, ethyl, isopropyl, methoxyethyl-.

25 In a currently preferred embodiment R^3 is $-C_{1-4}$ -alkyl- $C(O)NR^{4A}R^{4B}$ where R^{4A} is hydrogen and R^{4B} is amino ethyl, or R^{4A} and R^{4B} together with the nitrogen to which they are attached form a pyrrolidyl or piperidinyl ring optionally substituted by one or more substituents selected from $-NH_2$, $-NHCH_3$, $NHCH_2CH_3$, or $N(CH_3)_2$.

30 In a preferred embodiment R^2 is:




wherein

T is a trivalent nitrogen atom or a methyne (i.e.CH);

R⁶ is hydrogen or C₁₋₄-alkyl such as methyl

R^{10B} is 3-7 membered heterocycl- group such as morpholine or piperidine, or 3-7 membered heterocycl-C₁₋₄-alkyl- such as morpholinylmethyl, morpholinylethyl, morpholinylpropyl, piperidinylmethyl, piperidinylethyl, piperidinylpropyl, piperazinylmethyl, piperazinylethyl or piperazinylpropyl any of which heterocyclic rings is optionally substituted by one or more substituents selected from C₁₋₄-alkyl- and C₁₋₄alkoxy-C₁₋₄alkyl.

10 In another embodiment R² is:



wherein


T is a trivalent nitrogen atom or a methyne (i.e.CH);

P is a direct bond or a diradical selected from methylene, ethylene, or propylene;

15 R⁶ is hydrogen or C₁₋₄-alkyl;

R¹² is selected from hydrogen, C₁₋₄-alkyl such as methyl, ethyl, propyl, butyl, isopropyl, and C₁₋₄alkoxy-C₁₋₄alkyl such as methoxyethyl-.

In another embodiment R² is:



20

wherein


T is a trivalent nitrogen atom or a methyne (i.e.CH);

P is a diradical selected from methylene, ethylene, or propylene;

R⁶ is hydrogen or C₁₋₄-alkyl;

25 R¹² is selected from hydrogen, C₁₋₄-alkyl such as methyl, ethyl, propyl, butyl, isopropyl, and C₁₋₄alkoxy-C₁₋₄alkyl such as methoxyethyl-.

In another embodiment R² is:



wherein

R^3 is $-C_{1-4}\text{-alkyl-C(O)NR}^{4A}R^{4B}$ such as $-\text{CH}_2\text{-C(O)NR}^{4A}R^{4B}$, $-(\text{CH}_2)_2\text{-C(O)NR}^{4A}R^{4B}$, or $-(\text{CH}_2)_3\text{-C(O)NR}^{4A}R^{4B}$ wherein R^{4A} and R^{4B} are each independently selected from hydrogen, $C_{1-4}\text{-alkyl-}$ such as methyl, ethyl, propyl, and amino- $C_{1-4}\text{-alkyl-}$, or R^{4A} and R^{4B} together with the nitrogen to which they are attached form a 3-7 membered cyclic amino group such as pyrrolidine, piperidine, piperazine or morpholine, any of which is optionally substituted by one or more substituents selected from: $C_{1-4}\text{-alkyl}$, or $-\text{NR}^{4A}R^{4B}$.

In any of the compounds of the invention, the R^1 group may be any one of the specific R^1 groups of the corresponding position of any of the examples described herein.

In any of the compounds of the invention, the R^2 group may be any one of the specific R^2 groups of the corresponding position of any of the examples described herein.

In any of the compounds of the invention, the R^3 group may be any one of the specific R^3 groups of the corresponding position of any of the examples described herein.

Specific currently preferred embodiments of the invention include:

4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]-N-(piperidin-4-ylmethyl)piperidine-1-carboxamide

4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]-N-(1-methylpiperidin-4-yl)piperidine-1-carboxamide

4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]-N-[(1-methylpiperidin-4-yl)methyl]piperidine-1-carboxamide

4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]-N-[(1-ethylpiperidin-4-yl)methyl]piperidine-1-carboxamide

4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]-N-methyl-N-[(1-methylpiperidin-4-yl)methyl]piperidine-1-carboxamide

- 4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]-N-[2-(piperazin-1-yl)ethyl]piperidine-1-carboxamide
- 4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]-N-[2-(1-methylpiperidin-4-yl)ethyl]piperidine-1-carboxamide
- 5 4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]-N-[3-(morpholin-4-yl)propyl]piperidine-1-carboxamide
- 4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]-N-[[1-(propan-2-yl)piperidin-4-yl]methyl]piperidine-1-carboxamide
- 4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]-N-[[1-(2-methoxyethyl)piperidin-10 4-yl]methyl]piperidine-1-carboxamide
- 4-[1-(4-Chlorophenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]-N-[(1-methylpiperidin-4-yl)methyl]piperazine-1-carboxamide
- N-(2-Aminoethyl)-2-{4-[1-(4-chlorophenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl] morpholin-3-yl}acetamide
- 15 2-{4-[1-(4-Chlorophenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]morpholin-3-yl}-1-[(3S)-3-(dimethylamino)pyrrolidin-1-yl]ethan-1-one
- or a pharmaceutically acceptable salt, or N-oxide thereof.

In one aspect, the invention relates to a compound of formula (I) for use in therapy.

20 The compounds as defined above are useful as inhibitors of SSAO activity. As such, they are useful in the treatment or prevention of conditions and diseases in which inhibition of SSAO activity is beneficial. More specifically, they are useful for the treatment or prevention of inflammation, inflammatory diseases, immune or autoimmune disorders, cystic fibrosis, or inhibition of tumour growth.

25

In particular, it is believed that compounds of formula (I) are useful for the treatment or prevention of arthritis (such as rheumatoid arthritis, juvenile rheumatoid arthritis, osteoarthritis and psoriatic arthritis), synovitis, vasculitis, conditions associated with inflammation of the bowel (such as Crohn's disease, ulcerative colitis, inflammatory

30 bowel disease and irritable bowel syndrome), atherosclerosis, multiple sclerosis, Alzheimer's disease, vascular dementia, pulmonary inflammatory diseases (such as asthma, chronic obstructive pulmonary disease and acute respiratory distress syndrome), fibrotic diseases (including idiopathic pulmonary fibrosis, cardiac fibrosis and systemic sclerosis (scleroderma)), inflammatory diseases of the skin (such as

35 contact dermatitis, atopic dermatitis and psoriasis), systemic inflammatory response

syndrome, sepsis, inflammatory and/or autoimmune conditions of the liver (such as autoimmune hepatitis, primary biliary cirrhosis, alcoholic liver disease, sclerosing cholangitis, and autoimmune cholangitis), diabetes (type I or II) and/or the complications thereof, chronic heart failure, congestive heart failure, ischemic
5 diseases (such as stroke and ischemia-reperfusion injury), and myocardial infarction and/or the complications thereof.

It is believed that the compounds of the invention are especially useful for the treatment or prevention of vasculitis, including, but not limited to, giant cell arteritis,
10 Takayasu's arteritis, Polyarteritis nodosa, Kawasaki disease, Wegener's granulomatosis, Churg-Strauss syndrome, microscopic polyangiitis, Henoch-Schönlein purpura, cryoglobulinemia, cutaneous leukocytoclastic angiitis and primary angiitis of the central nervous system.

15 It is also believed that the compounds of the invention are especially useful for the treatment of rheumatoid arthritis, chronic obstructive pulmonary disease or atopic dermatitis.

In view of the evidence cited in the above introduction that VAP-1 is up regulated in
20 several cancers, including gastric cancer, melanoma, hepatoma and head and neck tumours and that mice bearing enzymatically inactive VAP-1 grow melanomas more slowly, and in view of the link between VAP-1 and angiogenesis, it is also expected that the compounds of the invention are anti-angiogenic and therefore have utility in the treatment of cancers by inhibition of tumour growth.

25 The invention thus includes the compounds of formula (I) above for use in the treatment or prevention of the above-mentioned conditions and diseases. The invention also includes the use of said compounds in the manufacture of a medicament for the treatment or prevention of the above-mentioned conditions and
30 diseases. The invention furthermore includes methods for treatment or prevention of such conditions and diseases, comprising administering to a mammal, including man, in need of such treatment an effective amount of a compound as defined above.

Methods delineated herein include those wherein the subject is identified as in need of a particular stated treatment. Identifying a subject in need of such treatment can be in the judgment of a subject or a health care professional and can be subjective (e.g. opinion) or objective (e.g. measurable by a test or diagnostic method).

5

In other aspects, the methods herein include those further comprising monitoring subject response to the treatment administrations. Such monitoring may include periodic sampling of subject tissue, fluids, specimens, cells, proteins, chemical markers, genetic materials, etc. as markers or indicators of the treatment regimen.

10 In other methods, the subject is prescreened or identified as in need of such treatment by assessment for a relevant marker or indicator of suitability for such treatment.

In one embodiment, the invention provides a method of monitoring treatment
15 progress. The method includes the step of determining a level of diagnostic marker (Marker) (e.g., any target or cell type delineated herein modulated by a compound herein) or diagnostic measurement (e.g., screen, assay) in a subject suffering from or susceptible to a disorder or symptoms thereof delineated herein, in which the subject has been administered a therapeutic amount of a compound herein
20 sufficient to treat the disease or symptoms thereof. The level of Marker determined in the method can be compared to known levels of Marker in either healthy normal controls or in other afflicted patients to establish the subject's disease status. In preferred embodiments, a second level of Marker in the subject is determined at a time point later than the determination of the first level, and the two levels are
25 compared to monitor the course of disease or the efficacy of the therapy. In certain preferred embodiments, a pre-treatment level of Marker in the subject is determined prior to beginning treatment according to this invention; this pre-treatment level of Marker can then be compared to the level of Marker in the subject after the treatment commences, to determine the efficacy of the treatment.

30

In certain method embodiments, a level of Marker or Marker activity in a subject is determined at least once. Comparison of Marker levels, e.g., to another measurement of Marker level obtained previously or subsequently from the same patient, another patient, or a normal subject, may be useful in determining whether
35 therapy according to the invention is having the desired effect, and thereby

permitting adjustment of dosage levels as appropriate. Determination of Marker levels may be performed using any suitable sampling/expression assay method known in the art or described herein. Preferably, a tissue or fluid sample is first removed from a subject. Examples of suitable samples include blood, urine, tissue, 5 mouth or cheek cells, and hair samples containing roots. Other suitable samples would be known to the person skilled in the art. Determination of protein levels and/or mRNA levels (e.g., Marker levels) in the sample can be performed using any suitable technique known in the art, including, but not limited to, enzyme immunoassay, ELISA, radiolabeling/assay techniques, blotting/chemiluminescence 10 methods, real-time PCR, and the like.

COMPOSITIONS

A currently preferred embodiment of the invention is a pharmaceutical composition 15 comprising a compound of formula (I), together with one or more pharmaceutically acceptable carriers and/or excipients.

For clinical use, the compounds of the invention are formulated into pharmaceutical formulations for various modes of administration. It will be appreciated that 20 compounds of the invention may be administered together with a physiologically acceptable carrier, excipient, or diluent. The pharmaceutical compositions of the invention may be administered by any suitable route, preferably by oral, rectal, nasal, topical (including buccal and sublingual), sublingual, transdermal, intrathecal, transmucosal or parenteral (including subcutaneous, intramuscular, intravenous and 25 intradermal) administration.

Other formulations may conveniently be presented in unit dosage form, e.g., tablets and sustained release capsules, and in liposomes, and may be prepared by any methods well known in the art of pharmacy. Pharmaceutical formulations are usually 30 prepared by mixing the active substance, or a pharmaceutically acceptable salt thereof, with conventional pharmaceutically acceptable carriers, diluents or excipients. Examples of excipients are water, gelatin, gum arabicum, lactose, microcrystalline cellulose, starch, sodium starch glycolate, calcium hydrogen phosphate, magnesium stearate, talcum, colloidal silicon dioxide, and the like. Such 35 formulations may also contain other pharmacologically active agents, and


conventional additives, such as stabilizers, wetting agents, emulsifiers, flavouring agents, buffers, and the like. Usually, the amount of active compounds is between 0.1-95% by weight of the preparation, preferably between 0.2-20% by weight in preparations for parenteral use and more preferably between 1-50% by weight in preparations for oral administration.

The formulations can be further prepared by known methods such as granulation, compression, microencapsulation, spray coating, etc. The formulations may be prepared by conventional methods in the dosage form of tablets, capsules, granules, powders, syrups, suspensions, suppositories or injections. Liquid formulations may be prepared by dissolving or suspending the active substance in water or other suitable vehicles. Tablets and granules may be coated in a conventional manner. To maintain therapeutically effective plasma concentrations for extended periods of time, compounds of the invention may be incorporated into slow release formulations.

The dose level and frequency of dosage of the specific compound will vary depending on a variety of factors including the potency of the specific compound employed, the metabolic stability and length of action of that compound, the patient's age, body weight, general health, sex, diet, mode and time of administration, rate of excretion, drug combination, the severity of the condition to be treated, and the patient undergoing therapy. The daily dosage may, for example, range from about 0.001 mg to about 100 mg per kilo of body weight, administered singly or multiply in doses, e.g. from about 0.01 mg to about 25 mg each. Normally, such a dosage is given orally but parenteral administration may also be chosen.


PREPARATION OF COMPOUNDS OF THE INVENTION

The compounds of formula (I) above may be prepared by, or in analogy with, conventional methods. The preparation of intermediates and compounds according to the Examples of the present invention may in particular be illuminated by the following Schemes. Definitions of variables in the structures in Schemes herein are commensurate with those of corresponding positions in the formulas delineated herein.

Scheme 1. General synthetic routes for preparation of compounds of formula (Ia)


wherein W, Y, Z, Q, R¹ and R³ are as defined in formula (I);

- 5 Compounds of general formula (Ia) can easily be prepared from 1*H*-pyrrolo[2,3-*c*]pyridines (IIa) by either introduction of the Q ring (or protected Q ring) followed by introduction of R¹ or by reversing these steps to give intermediates of general formula (IIIa). Compounds of general formula (IIIa) can then be converted to compounds of general formula (Ia) by standard synthetic methods. For example,
- 10 condensation of 1*H*-pyrrolo[2,3-*c*]pyridines (IIa) with tert-butyl 4-oxopiperidine-1-carboxylate, reduction, introduction of R¹ by an arylation reaction and Boc deprotection can be used to give compounds of general formula (IVa). Functionalisation of compounds of general formula (IVa) by for example, urea formation, amide coupling or reductive amination gives compounds of general
- 15 formula (Ia).

Scheme 2. General synthetic route for preparation of compounds of formula (Ib)

wherein W, Y, Z, R¹ and R² are as defined in formula (I);

Compounds of general formula (Ib) can easily be prepared from bromoindoles (IIb) by either introduction of R² (for example by nucleophilic substitution) followed by R¹ (for example by a Suzuki reaction), or by reversing these steps (with an appropriate protecting group strategy).


Scheme 3. General synthetic route for preparation of compounds of formula (Ic).

wherein Y, Z, R¹ and R² are as defined in formula (I);

Compounds of general formula (Ic) can easily be prepared by reductive amination of 3-amino-pyridine-4-carbaldehydes of general formula (IIc) to give compounds of

general formula (IIIc) and subsequent cyclisation to give pyrazolo[3,4-c]pyridines of general formula (IVc). Alternatively, pyrazolo[3,4-c]pyridines of general formula (IVc) can be prepared by cyclisation of (3-fluoropyridin-4-yl)carbonyl compounds of general formula (Vc) with hydrazine. Compounds of general formula (Ic) can be prepared from compounds of general formula (IVc) by standard N-arylation reactions.


Scheme 4. General synthetic route for preparation of compounds of formula (Id).



wherein W, Y, R¹ and R² are as defined in formula (I);


Compounds of general formula (Id) can easily be prepared according to standard methods known in the scientific literature, for example, by lithiation of 7-bromo-4-chloro-5H-pyrrolo[3,2-d]pyrimidines (IId) and reaction with an aldehyde, followed by reduction and subsequent introduction of R¹ (for example by an arylation reaction). Such methods are known to those skilled in the art, for example in WO2008070507 and Antilla et al., JOC, 69, 5578, 2004.

Scheme 5. General synthetic route for preparation of compounds of formula (Ie).



wherein Y, R¹ and R² are as defined in formula (I);

Compounds of general formula (Ie) can easily be prepared according to standard methods known in the scientific literature, for example, by condensation of 5-chloropyrimidines (IIe) with a Weinreb amide and subsequent reaction with a hydrazine. Such methods are known to those skilled in the art, for example in WO2003039469 and Verma et al, Tet. Lett., 50, 383, 2009.


Scheme 6. General synthetic route for preparation of compounds of formula (If).

wherein Y, Z, R¹ and R² are as defined in formula (I);

5


Compounds of general formula (If) can easily be prepared by the condensation of pyrazine-2-carbonitriles with a Grignard reagent to give amine intermediates (IIIf). Functionalisation of amines (IIIf) to give amides or ureas of general formula (IIIIf) and cyclisation with phosphorus oxychloride gives compounds of general formula (If).

10

Scheme 7. General synthetic route for preparation of compounds of formula (Ig).


wherein W, Z, R¹ and R² are as defined in formula (I);

15 Compounds of general formula (Ig) can be prepared from 1H-pyrrolo[2,3-d]pyridazines (IIg) by bromination with CuBr (for example, as described in Gallou et al., Syn. Lett., 2, 211-214, 2007) and subsequent introduction of R¹ (for example by an arylation reaction) and R² (for example by a Buchwald-Hartwig reaction).

20 **Scheme 8.** General synthetic route for preparation of compounds of formula (Ih).

wherein Y, Z, R¹ and R² are as defined in formula (I);

25 Compounds of general formula (Ih) can be prepared by sequential alkylations of 1H,2H,3H-imidazo[4,5-c]pyridin-2-ones (IIh), for example as described in WO2008054749.


Scheme 9. General synthetic route for preparation of compounds of formula (li).

wherein W, Y, Z, R^1 and R^2 are as defined in formula (I);

5

Compounds of general formula (li) can be prepared from 1H-pyrrolo[2,3-d]pyridazines (IIIi) by bromination with CuBr (for example, as described in Gallou et al., Syn. Lett., 2, 211-214, 2007) followed by either introduction of R^2 (for example by nucleophilic substitution) followed by R^1 (for example by a Suzuki reaction), or by reversing these steps (with an appropriate protecting group strategy).

10

Scheme 10. General synthetic route for preparation of compounds of formula (lj).


wherein W, Z, R^1 and R^2 are as defined in formula (I);

15

Compounds of general formula (lj) can be prepared from 1H-pyrrolo[2,3-d]pyridazines (IIj) by bromination with CuBr (for example, as described in Gallou et al., Syn. Lett., 2, 211-214, 2007) followed by either introduction of R^2 (for example

by nucleophilic substitution) followed by R^1 (for example by a Suzuki reaction), or by reversing these steps (with an appropriate protecting group strategy).


Scheme 11. General synthetic route for preparation of compounds of formula (Ik).



wherein Y, Z, R^1 and R^2 are as defined in formula (I);

Compounds of general formula (Ik) can be prepared by cyclisation of compounds of general formula (IIk) with hydrazines, for example, as described in Deeb et al., Journal of the Chinese Chemical Society, 37(3), 287-94; 1990.


Scheme 12. General synthetic route for preparation of compounds of formula (II).



wherein Z, R^1 and R^2 are as defined in formula (I);

Compounds of general formula (II) can be prepared by cyclisation of compounds of general formula (III) with hydrazines, for example, as described in Haider et al., Journal of the Chemical Society, Perkin Transactions 1: Organic and Bio-Organic Chemistry, 1, 169-72; 1986.


Scheme 13. General synthetic route for preparation of compounds of formula (Im).



wherein W, Y, R^1 and R^2 are as defined in formula (I);

Compounds of general formula (Im) can be prepared by sequential alkylation / arylation of 7H-pyrrolo[2,3-d]pyrimidines (IIm), for example as described in WO2009080682.


- 5 **Scheme 14.** General synthetic route for preparation of compounds of formula (In).



wherein Y, Z, R¹ and R² are as defined in formula (I);

- 10 Compounds of general formula (In) can be prepared by cyclisation of compounds of general formula (IIn) with hydrazines, for example, as described in Filaok et al., Journal of Organic Chemistry, 73(10), 3900-3906, 2008.

Scheme 15. General synthetic route for preparation of compounds of formula (Io).




- 15 wherein Z, R¹ and R² are as defined in formula (I);

Compounds of general formula (Io) can be prepared according to Scheme 15, for example as described in Haider et al., Journal of the Chemical Society, Perkin Transactions 1: Organic and Bio-Organic Chemistry, 1, 169-72; 1986.

20

Scheme 16. General synthetic route for preparation of compounds of formula (Ip).



wherein R¹ and R² are as defined in formula (I);

Compounds of general formula (Ip) can be prepared according to Scheme 16, for example as described in WO2007134828.

Optionally, a compound of formula (I) can also be transformed into another
5 compound of formula (I) in one or more synthetic steps.

The following abbreviations have been used:

Ac	acetyl
Ac ₂ O	acetic anhydride
AcOH	acetic acid
aq	aqueous
Ar	aryl
Boc	tert-butoxycarbonyl
nBuLi	n-butyllithium
calcd	calculated
CDI	carbonyldiimidazole
conc	concentrated
d	day
DCE	dichloroethane
DCM	dichloromethane
DIBALH	diisobutylaluminium hydride
DIPEA	diisopropylethylamine
DMAP	4-dimethylaminopyridine
DMF	dimethylformamide
EDC	1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride
ES+	electrospray ionization
EtOAc	ethyl acetate
EtOH	ethanol
Ex	Example
h	hour(s)
HBTU	O-benzotriazole-N,N,N',N'-tetramethyl-uronium-hexafluoro-phosphate
HOBt	1-hydroxybenzotriazole hydrate
HPLC	High Performance Liquid Chromatography
HRMS	High-Resolution Mass Spectrometry

Int	Intermediate
LCMS	Liquid Chromatography Mass Spectrometry
LDA	lithium diisopropylamide
M	molar
Me	methyl
mCPBA	meta-chloroperbenzoic acid
MeCN	acetonitrile
MeOH	methanol
min	minute(s)
Ms	methanesulfonate
MS	Mass Spectrometry
NaBH(OAc) ₃	sodium triacetoxyborohydride
NIS	N-iodosuccinimide
NMP	N-methylpyrrolidone
R _f	Retention time
RT	room temperature
sat	saturated
SCX	Strong Cation Exchange
SM	starting material
TFA	trifluoroacetic acid
THF	tetrahydrofuran


EXAMPLES AND INTERMEDIATE COMPOUNDS

Experimental Methods

5 Reactions were conducted at room temperature unless otherwise specified. Microwave reactions were performed with a Biotage microwave reactor using process vials fitted with aluminum caps and septa. Hydrogenations were performed using a Thales H-Cube. Preparative flash chromatography was performed on Merck silica gel 60 (230-400 mesh) or using a Flash Master Personal system equipped
10 with Strata SI-1 silica gigatubes, or using a CombiFlash Companion system equipped with RediSep silica columns. Reverse phase column chromatography was performed on a Gilson system (Gilson 321 pump and Gilson FC204 fraction collector) equipped with Merck LiChroprep® RP-18 (40-63µm) columns. Reverse Phase HPLC was performed on a Gilson system with a UV detector equipped with
15 Phenomenex Synergi Hydro RP 150 x 10 mm, or YMC ODS-A 100/150 x 20 mm columns. The purest fractions were collected, concentrated and dried under vacuum. Compounds were typically dried in a vacuum oven at 40 °C prior to purity analysis. Compound analysis was performed by HPLC/LCMS using an Agilent 1100 HPLC system / Waters ZQ mass spectrometer connected to an Agilent 1100 HPLC
20 system with a Phenomenex Synergi, RP-Hydro column (150 x 4.6 mm, 4 µm, 1.5 mL per min, 30 °C, gradient 5-100% MeCN (+0.085% TFA) in water (+0.1% TFA) over 7 min, 200-300 nm). Accurate masses (HRMS) were measured using a Thermo Scientific LTQ Orbitrap XL equipped with an Advio TriVersa NanoMate electrospray ion source (during the analyses the calibration was checked by three
25 masses. Spectra were acquired in positive electrospray mode. The acquired mass range was m/z 100-2000. Samples were dissolved in DMSO to give 10 mM solutions which were then further diluted with MeOH or 10 mM NH₄OAc in MeOH to ~0.1 M solutions prior to analysis). The values reported correspond to the protonated molecular ions [MH]⁺. The compounds prepared were named using ACD
30 Name 6.0, 7.0 or 10.0.

INTERMEDIATE 1


***tert*-Butyl 4-{1*H*-pyrrolo[2,3-*c*]pyridin-3-yl}-1,2,3,6-tetrahydropyridine-1-carboxylate**



6-Azaindole (4.48 g, 37.9 mmol) was dissolved in MeOH (70 mL) and KOH (4.68 g, 83.4 mmol) and tert-butyl 4-oxopiperidine-1-carboxylate (8.31 g, 41.7 mmol) were added. The reaction mixture was heated at 70 °C for 18 h. The residue was partitioned between water (250 mL) and DCM (250 mL) and the aq phase was extracted with DCM (2 x 250 mL). The combined organic fractions were dried (MgSO₄) and concentrated *in vacuo* to give the title compound as a yellow foam (11.3 g, 99%). LCMS (ES⁺): 300.1 [MH]⁺.

INTERMEDIATE 2


tert-Butyl 4-{1*H*-pyrrolo[2,3-*c*]pyridin-3-yl}piperidine-1-carboxylate



Intermediate 1 (11.3 g, 37.7 mmol) was dissolved in EtOH (200 mL) and hydrogenated over 10% Pd/C in an H-cube at 90 °C and 90 bar. The reaction mixture was concentrated *in vacuo* to give the title compound as a yellow solid (11.1 g, 97%). LCMS (ES⁺): 302.1 [MH]⁺.

INTERMEDIATE 3


tert-Butyl 4-[1-(4-chlorophenyl)-1*H*-pyrrolo[2,3-*c*]pyridin-3-yl]piperidine-1-carboxylate



Intermediate 2 (11.1 g, 36.7 mmol) was dissolved in DMF (60 mL) and 1-chloro-4-iodo-benzene (10.5 g, 44.0 mmol), *N,N*-dimethylethylenediamine (789 µL, 7.33 mmol), K₃PO₄ (16.3 g, 77.0 mmol) and CuI (698 mg, 3.67 mmol) were added under


nitrogen. The reaction mixture was heated in a microwave at 160 °C for 20 min and concentrated *in vacuo*. The residue was partitioned between water (250 mL) and DCM (250 mL) and the aq phase was extracted with DCM (2 x 250 mL). The combined organic fractions were dried (MgSO₄) and concentrated *in vacuo*. The residue was purified by column chromatography to give the title compound as a yellow solid (6.86 g, 45%). LCMS (ES⁺): 411.9 [MH]⁺, HPLC: Rt 5.91 min, 76% purity.

INTERMEDIATE 4

10 **4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]piperidine**

Intermediate 3 (6.86 g, 16.6 mmol) was dissolved in DCM (200 mL) and TFA (50 mL) and stirred for 2 h. The solvents were removed *in vacuo* and the residue was dissolved in 1 M aq Na₂CO₃ (200 mL) and extracted with DCM (3 x 200 mL). The combined organic fractions were dried (MgSO₄) and concentrated *in vacuo*. The residue was purified by column chromatography to give the title compound as a red gum (3.18 g, 61%). LCMS (ES⁺): 312.1 [MH]⁺. HPLC: Rt 3.61 min, 96% purity.

INTERMEDIATE 5


20 **4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]-N-(piperidin-4-ylmethyl)piperidine-1-carboxamide**

CDI (936 mg, 5.77 mmol) was dissolved in DCM (50 mL), a solution of tert-butyl 4-(aminomethyl)piperidine-1-carboxylate (1.24 g, 5.77 mmol) and DIPEA (1.25 mL, 7.22 mmol) in DCM (10 mL) was added and the reaction mixture was stirred for 18 h. A solution of Intermediate 4 (1.50 g, 4.81 mmol) and DIPEA (1.25 mL, 7.22 mmol) in DCM (10 mL) was added and the reaction mixture was stirred for 24 h, diluted with 1 M aq Na₂CO₃ (100 mL) and extracted with DCM (3 x 100 mL). The combined

organic fractions were dried (MgSO_4) and concentrated *in vacuo*. The residue was purified by column chromatography, dissolved in DCM (10 mL) and TFA (2.5 mL) and stirred for 1 h. The reaction mixture was concentrated *in vacuo* and the residue was dissolved in 1 M aq Na_2CO_3 (50 mL) and extracted with DCM (3 x 50 mL). The combined organic fractions were dried (MgSO_4) and concentrated *in vacuo*. The residue was purified by column chromatography to give the title compound as a pale yellow solid (1.41 g, 65%). LCMS (ES^+): 452.0 $[\text{MH}]^+$. HPLC: Rt 3.98 min, 97% purity.

10 INTERMEDIATE 6


N-(3-Aminopropyl)-4-[1-(4-chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]piperidine-1-carboxamide



Intermediate 6 (178 mg, 8%) was prepared similarly to Intermediate 5, using tert-butyl N-(3-aminopropyl)carbamate instead of tert-butyl 4-(aminomethyl)piperidine-1-carboxylate. LCMS (ES^+): 412.3 $[\text{MH}]^+$. HPLC: Rt 3.82 min, 100% purity.

INTERMEDIATE 7

tert-Butyl N-({4-[(3-aminopyridin-4-yl)methyl]morpholin-2-yl}methyl)carbamate




3-Amino-pyridine-4-carbaldehyde (513 mg, 4.20 mmol) was dissolved in DCE (7.3 mL) and tert-butyl (morpholin-2-ylmethyl)carbamate (999 mg, 4.62 mmol) and $\text{NaBH}(\text{OAc})_3$ (1.07 g, 5.04 mmol) were added. The reaction mixture was heated using a microwave at 60 °C for 5 min, diluted with DCM (10 mL) and quenched with sat aq Na_2CO_3 (5 mL).

The aq phase was extracted with DCM (3 x 20 mL) and the combined organic fractions were dried (MgSO_4) and concentrated *in vacuo* to give the crude title compound as a yellow gum (1.37 g, 100%). LCMS (ES^+): 323.1 $[\text{MH}]^+$.

INTERMEDIATE 8


tert-Butyl N-[(4-{1H-pyrazolo [3,4-c]pyridin-3-yl}morpholin-2-yl)methyl]carbamate



5 Intermediate 7 (1.35 g, 4.20 mmol) was dissolved in AcOH (55 mL), and a solution of NaNO₂ (290 mg, 4.20 mmol) in water (438 µL) was added. The reaction mixture was stirred for 5 min and concentrated *in vacuo*. The residue was dissolved in EtOAc (40 mL) and washed with sat aq Na₂CO₃ (2 x 20 mL). The organic fraction
10 was dried (MgSO₄) and concentrated *in vacuo* to give the crude title compound as a yellow gum (999 mg, 71%). LCMS (ES⁺): 334.0 [MH]⁺.

INTERMEDIATE 9


tert-Butyl N-({4-[1-(4-methylphenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]morpholin-2-yl}methyl)carbamate



Intermediate 8 (999 mg, 3.00 mmol) was dissolved in DMF (4 mL) and 1-methyl-4-iodobenzene (784 mg, 3.60 mmol), *N,N*-dimethylethylenediamine (64.5 µL, 0.60 mmol), K₃PO₄ (1.34 g, 6.29 mmol) and CuI (57.1 mg, 0.30 mmol) were added. The
20 reaction mixture was heated using a microwave at 140 °C for 20 min. The solvents were removed *in vacuo* and the residue was purified by column chromatography to give the crude title compound as a yellow gum (438 mg, 35%). LCMS (ES⁺): 424.0 [MH]⁺.

25 INTERMEDIATE 10

{4-[1-(4-Methylphenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]morpholin-2-yl}methanamine dihydrochloride




Intermediate 9 (438 mg, 1.03 mmol) was dissolved in 1.25 M HCl in EtOH (10 mL) and stirred overnight. The solvents were removed *in vacuo* to give the crude title compound as an orange gum (400 mg, 98%). LCMS (ES⁺): 324.0 [MH]⁺.

5

INTERMEDIATE 11

tert-Butyl 2-[(acetyloxy)methyl]morpholine-4-carboxylate




Ac₂O (5.17 mL, 54.7 mmol) was dissolved in DCM (200 mL), DMAP (611 mg, 5.00 mmol), DIPEA (9.52 mL, 54.7 mmol) and *tert*-butyl-2-(hydroxymethyl) morpholine-4-carboxylate (10.0 g, 49.7 mmol) were added and the reaction mixture was stirred for 1 h. The reaction mixture was washed with sat aq NH₄Cl (3 x 100 mL) and the aq fraction was extracted with DCM (2 x 100 mL). The combined organic fractions were dried (MgSO₄) and concentrated *in vacuo* to give the crude title compound as an off white solid (14.1 g). LCMS (ES⁺): 282.1 [MNa]⁺.

15

INTERMEDIATE 12

{4-[(3-Aminopyridin-4-yl)methyl]morpholin-2-yl}methyl acetate




Intermediate 11 (12.8 g, 49.5 mmol) was dissolved in TFA (20 mL) and DCM (80 mL) and the reaction mixture was stirred overnight and concentrated *in vacuo*. The residue was dissolved in DCE (80 mL), cooled to 0 °C and Et₃N (6.90 mL, 49.5 mmol) was added drop-wise. 3-Amino-pyridine-4-carbaldehyde (6.04 g, 49.5 mmol) and MeOH (50 mL) were added and the reaction mixture was stirred for 30 min. NaBH(OAc)₃ (12.6 g, 59.4 mmol) was added portion-wise and the reaction mixture was stirred overnight. Further 3-amino-pyridine-4-carbaldehyde (6.04 g, 49.5 mmol) and NaBH(OAc)₃ (25.2 g, 119 mmol) were added portion-wise over 2 d. The

25

reaction mixture was stirred at 60 °C for 30 h, cooled to 0 °C, quenched with sat aq Na₂CO₃ (50 mL) and diluted with DCM (100 mL). The organic fraction was washed with sat aq Na₂CO₃ (20 mL) and sat aq NH₄Cl (2 x 20 mL), dried (MgSO₄) and concentrated *in vacuo*. The residue was purified by filtration through a pad of silica to give the crude title compound as a yellow gum (5.77 g, 44%). LCMS (ES⁺): 266.1 [MH]⁺.

INTERMEDIATE 13

(4-{1H-Pyrazolo[3,4-c]pyridin-3-yl}morpholin-2-yl)methyl acetate

10


Intermediate 12 (5.77 g, 21.8 mmol) was dissolved in AcOH (282 mL), cooled to 0 °C, and a solution of NaNO₂ (1.50 g, 21.8 mmol) in water (2.29 mL) was added. The reaction mixture was stirred for 5 min and concentrated *in vacuo*. The residue was dissolved in EtOAc (200 mL) and washed with sat aq Na₂CO₃ (2 x 100 mL), dried (MgSO₄) and concentrated *in vacuo* to give the title compound as a dark yellow gum (3.85 g, 64%). LCMS (ES⁺): 277.1 [MH]⁺.

15

INTERMEDIATE 14

{4-[1-(4-Methylphenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]morpholin-2-yl}methyl acetate

20




Intermediate 13 (3.85 g, 13.9 mmol) was dissolved in DMF (20 mL) and 1-methyl-4-iodo-benzene (3.64 g, 16.7 mmol), *N,N*-dimethylethylenediamine (300 µL, 2.78 mmol), K₃PO₄ (6.21 g, 29.2 mmol) and CuI (265 mg, 1.39 mmol) were added. The reaction mixture was heated in a microwave reactor at 140 °C for 1 h and concentrated *in vacuo*. The residue was purified by column chromatography to give the title compound as a yellow gum (260 mg, 5%). LCMS (ES⁺): 367.0 [MH]⁺. HPLC: Rt 5.07 min, 97.4%.

25

INTERMEDIATE 15

{4-[1-(4-Chlorophenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]morpholin-2-yl}methyl acetate




5

Intermediate 13 (200 mg, 0.72 mmol), 4-chlorophenylboronic acid (226 mg, 1.45 mmol), $\text{Cu}(\text{OAc})_2$ (263 mg, 1.45 mmol) and pyridine (292 μL , 3.62 mmol) were suspended in DCM (10 mL) and stirred for 36 h. The reaction mixture was concentrated *in vacuo* and purified by column chromatography to give the crude title compound as a yellow solid (80.0 mg, 29%). LCMS (ES^+): 387.0 $[\text{MH}]^+$.

10

INTERMEDIATE 16

{4-[1-(4-Methylphenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]morpholin-2-yl}methanol




15

Intermediate 14 (200 mg, 0.55 mmol) was dissolved in MeOH (4 mL) and K_2CO_3 (302 mg, 2.18 mmol) was added. The reaction mixture was stirred for 30 min and concentrated *in vacuo*. The residue was dissolved in DCM (20 mL) and water (10 mL) and the aq phase was extracted with DCM (3 x 50 mL). The combined organic fractions were dried (MgSO_4) and concentrated *in vacuo* to yield the title compound as a dark brown gum (168 mg, 95%). LCMS (ES^+): 325.1 $[\text{MH}]^+$.

20

INTERMEDIATE 17

{4-[1-(4-Chlorophenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]morpholin-2-yl}methanol




25

Intermediate 17 (65.0 mg, 91%) was prepared similarly to Intermediate 16, using

Intermediate 15 instead of Intermediate 14. LCMS (ES⁺): 345.0 [MH]⁺.

INTERMEDIATE 18


**{4-[1-(4-Methylphenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]morpholin-2-yl}methyl
methanesulfonate**



Intermediate 16 (292 mg, 0.90 mmol) was dissolved in DCM (7 mL), cooled to 0 °C and Et₃N (138 μL, 0.99 mmol) and methanesulfonyl chloride (76.6 μL, 0.99 mmol) were added. The reaction mixture was stirred for 1 h, diluted with DCM (10 mL) and washed with sat aq NH₄Cl (2 x 5 mL) and sat aq Na₂CO₃ (2 x 5 mL). The organic fraction was dried (MgSO₄) and concentrated *in vacuo* to give the title compound as a brown gum (277 mg, 77%). LCMS (ES⁺): 403.0 [MH]⁺.

INTERMEDIATE 19


**{4-[1-(4-Chlorophenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]morpholin-2-yl}methyl
methanesulfonate**



Intermediate 19 (60.0 mg, 75%) was prepared similarly to Intermediate 18, using Intermediate 17 instead of Intermediate 16. LCMS (ES⁺): 422.9 [MH]⁺.

INTERMEDIATE 20


tert-Butyl 4-({4-[(3-aminopyridin-4-yl)methyl]morpholin-2-yl}methyl)piperazine-1-carboxylate



3-Amino-pyridine-4-carbaldehyde (584 mg, 4.78 mmol) was dissolved in DCM (10 mL) and tert-butyl 4-(morpholin-2-ylmethyl)piperazine-1-carboxylate (1.50 g, 5.26 mmol) and NaBH(OAc)₃ (1.11 g, 5.26 mmol) were added. The reaction mixture was heated in a microwave at 60 °C for 2.5 min, diluted with DCM (20 mL) and quenched with sat aq Na₂CO₃ (10 mL). The organic fraction was washed with sat aq NH₄Cl (10 mL). The combined aq fractions were extracted with DCM (2 x 20 mL) and the combined organic fractions were dried (MgSO₄) and concentrated *in vacuo* to give the crude title compound as a yellow gum (2.46 g). LCMS (ES⁺): 392.1 [MH]⁺.

10 INTERMEDIATE 21

tert-Butyl 4-[(4-{1H-pyrazolo[3,4-c]pyridin-3-yl}morpholin-2-yl)methyl]piperazine-1-carboxylate




Intermediate 20 (1.87 g, 4.76 mmol) was dissolved in AcOH (62 mL), cooled to 0 °C and a solution of NaNO₂ (330 mg, 4.76 mmol) in water (502 µL) was added. The reaction mixture was stirred for 5 min and concentrated *in vacuo*. The residue was dissolved in EtOAc (100 mL) and washed with sat aq Na₂CO₃ (2 x 50 mL). The organic fraction was dried (MgSO₄) and concentrated *in vacuo* to yield the title compound as a brown gum (1.80 g, 93%). LCMS (ES⁺): 403.1 [MH]⁺.

20

INTERMEDIATE 22

tert-Butyl 4-[(4-{1-(4-chlorophenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl}morpholin-2-yl)methyl]piperazine-1-carboxylate




25 Intermediate 21 (1.08 g, 2.69 mmol), 4-chlorobenzeneboronic acid (840 mg, 5.37 mmol), Cu(OAc)₂ (976 mg, 5.37 mmol) and pyridine (1.08 mL, 13.4 mmol) were suspended in DCE (19 mL) and stirred overnight. The solvents were removed *in vacuo* and the residue was purified by column chromatography to give the title

compound as a pale yellow solid (58.0 mg, 4%). LCMS (ES⁺): 513.0 [MH]⁺. HPLC: Rt 4.81 min, 97.1% purity.

INTERMEDIATES 23 to 30




- 5 Intermediates 23-30 were prepared similarly to Intermediate 20, by reductive amination of 3-amino-pyridine-4-carbaldehyde with the appropriate amine; see Table 1 below.

Table 1: Reductive aminations of 3-amino-pyridine-4-carbaldehyde




10

Int	Structure	Crude yield	LCMS (ES ⁺)	Intermediate Name
23		3.16 g 63%	266.1 [MH] ⁺	Methyl 2-{4-[(3-aminopyridin-4-yl)methyl]morpholin-3-yl}acetate
24		11.3 g 100%	264.1 [MH] ⁺	Ethyl 1-[(3-aminopyridin-4-yl)methyl]piperidine-4-carboxylate
25		541 mg 70%	237.1 [MH] ⁺	4-({[2-(Morpholin-4-yl)ethyl]amino}methyl)pyridin-3-amine
26		585 mg 53%	336.1 [MH] ⁺	tert-Butyl 4-(2-({[(3-aminopyridin-4-yl)methyl]amino}ethyl)piperazine-1-carboxylate
27		861 mg 42%	250.1 [MH] ⁺	1-[(3-Aminopyridin-4-yl)methyl]piperidin-4-yl acetate

Int	Structure	Crude yield	LCMS (ES ⁺)	Intermediate Name
28		165 mg 36%	293.3 [MH] ⁺	4-([3-(Morpholin-4-ylmethyl)morpholin-4-yl]methyl)pyridin-3-amine
29		1.67 g 70%	293.1 [MH] ⁺	tert-Butyl 4-[(3-aminopyridin-4-yl)methyl]piperazine-1-carboxylate
30		1.76 g 40%	250.1 [MH] ⁺	Methyl 1-[(3-aminopyridin-4-yl)methyl]piperidine-2-carboxylate

INTERMEDIATE 31

tert-Butyl N-[(3-aminopyridin-4-yl)methyl]-N-[2-(morpholin-4-yl)ethyl]carbamate



5


Intermediate 25 (341 mg, 1.44 mmol) was dissolved in DCM (15 mL), Boc₂O (346 mg, 1.59 mmol) was added and the reaction mixture was stirred for 1.5 h. The reaction mixture was quenched with sat aq Na₂CO₃ (40 mL) and the aq fraction was extracted with DCM (2 x 20 mL). The combined organic fractions were washed with brine (30 mL), dried (MgSO₄) and concentrated *in vacuo* to give the crude title compound as a brown oil (329 mg). LCMS (ES⁺): 337.0 [MH]⁺.

10

INTERMEDIATE 32

tert-Butyl 4-(2-([(3-aminopyridin-4-yl)methyl][(tert-butoxy)carbonyl]amino)ethyl) piperazine-1-carboxylate

15




Intermediate 32 was prepared similarly to Intermediate 31, using Intermediate 26 instead of Intermediate 25, to give the title compound as a brown oil (409 mg, 54%).
LCMS (ES⁺): 436.1 [MH]⁺. HPLC: Rt 4.18 min, 93% purity.

5

INTERMEDIATES 33 to 40

Intermediates 33-30 were prepared similarly to Intermediate 21, by cyclisation of 3-aminopyridines 23-24 and 27-32 with NaNO₂; see Table 2 below.

10 **Table 2: Cyclisation of 3-aminopyridines**




Int	Structure	SM / Crude yield	LCMS (ES ⁺)	Intermediate Name
33		Int 23 2.00 g 100%	277.1 [MH] ⁺	2-(4-{1H-Pyrazolo[3,4-c]pyridin-3-yl}morpholin-3-yl)acetate
34		Int 24 12.2 g 100%	275.1 [MH] ⁺	Ethyl 1-{1H-pyrazolo[3,4-c]pyridin-3-yl}piperidine-4-carboxylate

Int	Structure	SM / Crude yield	LCMS (ES ⁺)	Intermediate Name
35		Int 31 243 mg 72%	348.1 [MH] ⁺	<i>tert</i> -Butyl N-[2-(morpholin-4-yl)ethyl]-N-{1H-pyrazolo[3,4-c]pyridin-3-yl} carbamate
36		Int 32 399 mg 94%	447.1 [MH] ⁺	<i>tert</i> -Butyl 4-(2-([(tert-butoxy)carbonyl](1H-pyrazolo[3,4-c]pyridin-3-yl)amino)ethyl)piperazine-1-carboxylate
37		Int 27 618 mg 69%	261.1 [MH] ⁺	1-{1H-Pyrazolo[3,4-c]pyridin-3-yl}piperidin-4-yl acetate
38		Int 28 92.0 mg crude	304.1 [MH] ⁺	3-(Morpholin-4-ylmethyl)-4-{1H-pyrazolo[3,4-c]pyridin-3-yl}morpholine
39		Int 29 933 mg 54%	304.2 [MH] ⁺	<i>tert</i> -Butyl 4-{1H-pyrazolo[3,4-c]pyridin-3-yl}piperazine-1-carboxylate
40		Int 30 650 g 89%	261.1 [MH] ⁺	Methyl 1-{1H-pyrazolo[3,4-c]pyridin-3-yl}piperidine-2-carboxylate


INTERMEDIATES 41 to 43

Intermediates 41-43 were prepared similarly to Intermediate 22, by N-arylation of 1H-pyrazolo[3,4-c]pyridines; see Table 3 below.

Table 3: N-Arylation of 1H-pyrazolo[3,4-c]pyridines

Int	Structure	Int / Crude yield	LCMS (ES ⁺)	Intermediate Name
41		Int 34 619 mg 13%	384.9 [MH] ⁺	Ethyl 1-[1-(4-chlorophenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]piperidine-4-carboxylate
42		Int 39 80.0 mg 75%	414.0 [MH] ⁺	tert-Butyl 4-[1-(4-chlorophenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]piperazine-1-carboxylate
43		Int 40 260 mg 28%	370.9 [MH] ⁺	Methyl 1-[1-(4-chlorophenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]piperidine-2-carboxylate


5 INTERMEDIATE 44

1-[1-(4-Chlorophenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]piperidine-4-carboxylic acid hydrochloride

Intermediate 41 (834 mg, 2.17 mmol) was dissolved in 1:1 THF/water (16 mL),
 10 LiOH.H₂O (200 mg, 4.77 mmol) was added and the reaction mixture was stirred for
 3 h. The THF was removed *in vacuo* and the reaction mixture was acidified to pH 1
 with 1 M aq HCl (5 mL). The precipitate was collected by filtration and washed with
 water to give the title compound as an orange solid (450 mg, 53%). LCMS (ES⁺):
 357.0 [MH]⁺. HPLC: Rt 4.92 min, 99.6% purity.

INTERMEDIATE 45

4-({1-[1-(4-Chlorophenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]piperidin-4-yl}carbonyl) morpholine




5

Intermediate 44 (200 mg, 0.51 mmol) was dissolved in DMF (2 mL), HBTU (231 mg, 0.61 mmol) was added and the reaction mixture was stirred for 30 min. Morpholine (53.4 μ L, 0.61 mmol) and DIPEA (266 μ L, 1.53 mmol) were added and the reaction mixture was stirred overnight. The solvents were removed *in vacuo* and the residue was diluted with EtOAc (25 mL), washed with sat aq NH_4Cl (4 x 25 mL), dried (MgSO₄) and concentrated *in vacuo*. The residue was purified by column chromatography to give the title compound as a light yellow solid (78.7 mg, 36%). HRMS (ESI⁺) calcd for C₂₂H₂₄ClN₅O₂ 426.1691, found 426.1691. HPLC: Rt 4.96 min, 100% purity.

15

INTERMEDIATE 46

tert-Butyl 4-({1-[1-(4-chlorophenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]piperidin-4-yl}carbonyl)piperazine-1-carboxylate




Intermediate 46 was prepared similarly to Intermediate 45, using *tert*-butyl 1-piperazinecarboxylate instead of morpholine to give the title compound as yellow gum (260 mg, 97%). LCMS (ES⁺): 525.1 [MH]⁺. HPLC: Rt 6.14 min, 100% purity.

INTERMEDIATE 47

1-[1-(4-Chlorophenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]piperidine-2-carboxylic acid hydrochloride

25




Intermediate 47 was prepared similarly to Intermediate 44, using Intermediate 43 instead of Intermediate 41, to give the crude title compound as a brown solid (332 mg). LCMS (ES⁺): 357.0 [MH]⁺.

5

INTERMEDIATE 48

2-{4-[1-(4-Chlorophenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]morpholin-3-yl}ethan-1-ol




10 Example 40 (50.0 mg, 0.13 mmol) was dissolved in DCM (1 mL), cooled to 0 °C and DIBALH (0.78 mL, 1.0 M in heptane, 0.78 mmol) was added portion-wise over 6 days. The reaction mixture was stirred for 1 week, cooled to 0 °C and quenched with water (1 mL). The reaction mixture was filtered and concentrated *in vacuo* to give the crude title compound as a yellow gum (51.0 mg). LCMS (ES⁺): 359.0 [MH]⁺.

15

INTERMEDIATE 49

2-{4-[1-(4-chlorophenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]morpholin-3-yl}ethyl methanesulfonate




20 Intermediate 48 (50.0 mg, 0.14 mmol) was dissolved in DCM (1.5 mL), cooled to 0 °C and Et₃N (42.7 μL, 0.31 mmol) and methanesulfonyl chloride (11.9 μL, 0.15 mmol) were added. The reaction mixture was stirred for 20 h, diluted with DCM (5 mL), washed with sat aq NH₄Cl (3 x 5 mL) and sat aq Na₂CO₃ (5 mL), dried

(MgSO₄) and concentrated *in vacuo* to give the title compound as a dark yellow gum (60.0 mg, 99%). LCMS (ES⁺): 437.0 [MH]⁺.

INTERMEDIATE 50


5 **1-[1-(4-Chlorophenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]piperazine dihydrochloride**



Intermediate 42 (80.0 mg, 0.19 mmol) was dissolved in HCl (1.25 M in EtOH, 10 mL) and the reaction mixture was stirred for 18 h. The solvents were removed *in vacuo* to give the title compound as an orange solid (76.0 mg, 100%). LCMS (ES⁺): 314.0 [MH]⁺. HPLC: Rt 3.83 min, 90.0% purity.

INTERMEDIATE 51


15 **tert-Butyl N-({4-[1-(4-chlorophenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]morpholin-2-yl}methyl)carbamate**



Intermediate 8 (290 mg, 0.87 mmol), 4-chlorophenylboronic acid (272 mg, 1.74 mmol), anhydrous copper (II) acetate (316 mg, 1.74 mmol) and pyridine (350 μ L, 4.35 mmol) were suspended in DCM (12 mL) and stirred for 24 h. The residue was dissolved in MeOH (15 mL) and purified using an SCX-2 cartridge and by column chromatography to give the title compound as a light yellow solid (100 mg, 26%). LCMS (ES⁺): 444.1 [MH]⁺. HPLC: Rt 6.01 min, 84% purity.

INTERMEDIATE 52

25 **{4-[1-(4-Chlorophenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]morpholin-2-yl}methanamine dihydrochloride**




Intermediate 52 was prepared similarly to Intermediate 10, using Intermediate 51 instead of Intermediate 9, to give the title compound as an orange solid (62.0 mg). LCMS (ES⁺): 344.1 [MH]⁺.

5

INTERMEDIATE 53

tert-Butyl N-[(3-aminopyridin-4-yl)methyl]-N-[2-(2-oxoimidazolidin-1-yl)ethyl] carbamate




10 3-Amino-pyridine-4-carbaldehyde (0.86 g, 7.04 mmol), 1-(2-aminoethyl)imidazolidin-2-one (1.00 g, 7.74 mmol) and AcOH (0.44 mL, 7.75 mmol) were dissolved in DCM (20 mL) and stirred for 1 h. NaBH(OAc)₃ (2.24 g, 10.6 mmol) was added and the reaction mixture was stirred for 3 h and diluted with DCM (10mL) and water (20mL). Na₂CO₃ (2.24 g, 21.1 mmol) and di-tert-butyl-dicarbonate (1.84 g, 8.45 mmol) were
15 added and the reaction mixture was stirred for 20 h. The aqueous fraction was extracted with DCM (50mL) and the combined organic fractions were washed with sat aq NaHCO₃ (40 mL), dried (MgSO₄) and concentrated *in vacuo*. The residue was purified by column chromatography to give the title compound (539 mg, 23%) as a yellow gum. LCMS (ES⁺): 336.2 [MH]⁺.

20

INTERMEDIATE 54

tert-Butyl N-[2-(2-oxoimidazolidin-1-yl)ethyl]-N-{1H-pyrazolo[3,4-c]pyridin-3-yl}carbamate




25

Intermediate 54 was prepared similarly to Intermediate 21, using Intermediate 53 instead of Intermediate 20, to give the title compound as a light brown solid (432 mg, 78%). LCMS (ES⁺): 347.2 [MH]⁺.

INTERMEDIATE 55


tert-Butyl N-[1-(4-chlorophenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]-N-[2-(2-oxoimidazolidin-1-yl)ethyl]carbamate



Intermediate 55 was prepared similarly to Intermediate 22, using Intermediate 54 instead of Intermediate 21, to give the title compound as a yellow solid (74.0 mg, 13%). LCMS (ES⁺): 457.0 [MH]⁺. HPLC: Rt 5.33min, 100% purity.

EXAMPLE 1


4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]-1-(pyrrolidin-3-yl)piperidine



Intermediate 4 (205 mg, 0.66 mmol) and tert-butyl 3-oxopyrrolidine-1-carboxylate (229 μ L, 243 mg) were dissolved in DCM (5 mL) and NaBH(OAc)₃ (348 mg, 1.64 mmol) was added. The reaction mixture was stirred for 18 h, diluted with 1 M aq Na₂CO₃ (50 mL) and extracted with DCM (2 x 50 mL). The combined organic fractions were dried (MgSO₄) and concentrated *in vacuo*. The residue was purified by column chromatography, dissolved in DCM (10 mL) and TFA (2.5 mL) and stirred for 2 h. The reaction mixture was concentrated *in vacuo* and the residue was dissolved in 1 M aq Na₂CO₃ (50 mL) and extracted with DCM (2 x 50 mL). The combined organic fractions were dried (MgSO₄) and concentrated *in vacuo*. The residue was purified by reverse phase HPLC to give the title compound as a colourless gum (67.0 mg, 27%). HRMS (ESI⁺) calcd for C₂₂H₂₅CIN₄ 381.184, found 381.1846. HPLC: Rt 3.36 min, 98% purity.

EXAMPLE 2


4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]-1-(piperidin-4-yl)piperidine



Example 2 (72.0 mg, 19%) was prepared similarly to Example 1, using tert-butyl 4-oxopiperidine-1-carboxylate instead of tert-butyl 3-oxopyrrolidine-1-carboxylate. HRMS (ESI+) calcd for C₂₃H₂₇ClN₄ 395.1997, found 395.1998. HPLC: Rt 3.52 min, 99% purity.

EXAMPLE 3


4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]-1-(piperidin-4-ylmethyl)piperidine



Example 3 (39.0 mg, 20%) was prepared similarly to Example 1, using tert-butyl 4-formylpiperidine-1-carboxylate instead of tert-butyl 3-oxopyrrolidine-1-carboxylate. HRMS (ESI+) calcd for C₂₄H₂₉ClN₄ 409.2153, found 409.2155. HPLC: Rt 3.55 min, 99% purity.

EXAMPLE 4


1-{4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]piperidin-1-yl}-2-(piperidin-4-yl)ethan-1-one



Intermediate 4 (200 mg, 0.64 mmol), 2-{1-[(tert-butoxy)carbonyl]piperidin-4-yl}acetic acid (203 mg, 0.83 mmol), HOBt (113 mg, 0.83 mmol) and DIPEA (290 μ L, 1.67 mmol) were dissolved in DMF (5 mL) and EDC (160 mg, 0.83 mmol) was added. The reaction mixture was stirred for 18 h and concentrated *in vacuo*. The residue


was dissolved in EtOAc (25 mL) and washed with 10% aq citric acid (25 mL), 1 M aq Na₂CO₃ (25 mL) and water (25 mL), dried (MgSO₄) and concentrated *in vacuo*. The residue was purified by column chromatography, dissolved in DCM (10 mL) and TFA (2 mL) and stirred for 2 h. The reaction mixture was concentrated *in vacuo*, dissolved in 1 M aq Na₂CO₃ (25 mL) and extracted with DCM (3 x 25 mL). The combined organic fractions were dried (MgSO₄) and concentrated *in vacuo*. The residue was purified by column chromatography to give the title compound as a colourless gum (20.5 mg, 7%). HRMS (ESI+) calcd for C₂₅H₂₉CIN₄O 437.2103, found 437.21. HPLC: Rt 3.92 min, 96% purity.

EXAMPLE 5

1-({4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]piperidin-1-yl}carbonyl)-4-methylpiperazine

Intermediate 4 (200 mg, 0.64 mmol), DIPEA (245 μ L, 1.41 mmol) and DMAP (7.80 mg, 0.06 mmol) were dissolved in DCM (10 mL) and 4-methylpiperazine-1-carbonyl chloride hydrochloride (140 mg, 0.70 mmol) was added. The reaction mixture was stirred for 18 h, diluted with 1 M aq Na₂CO₃ (50 mL) and extracted with DCM (3 x 50 mL). The combined organic fractions were dried (MgSO₄) and concentrated *in vacuo*. The residue was purified by column chromatography and reverse phase HPLC to give the title compound as a white solid (84.0 mg, 30%). HRMS (ESI+) calcd for C₂₄H₂₈CIN₅O 438.2055, found 438.2057. HPLC: Rt 3.92 min, 100% purity.

EXAMPLE 6

4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]-N-(piperidin-4-ylmethyl)piperidine-1-carboxamide


CDI (187 mg, 1.15 mmol) was dissolved in DCM (10 mL), a solution of tert-butyl 4-(aminomethyl)piperidine-1-carboxylate (247 mg, 1.15 mmol) and DIPEA (251 μ L, 1.15 mmol) in DCM (2 mL) was added and the reaction mixture was stirred for 18 h. A solution of Intermediate 4 (300 mg, 0.96 mmol) and DIPEA (251 μ L, 1.15 mmol) in DCM (2 mL) was added and the reaction mixture was stirred for 24 h, mixed with 1 M aq Na₂CO₃ (50 mL) and extracted with DCM (3 x 50 mL). The combined organic fractions were dried (MgSO₄) and concentrated *in vacuo*. The residue was purified by column chromatography, dissolved in DCM (10 mL) and TFA (2.5 mL) and stirred for 1 h. The reaction mixture was concentrated *in vacuo*, dissolved in 1 M aq Na₂CO₃ (50 mL) and extracted with DCM (3 x 50 mL). The combined organic fractions were dried (MgSO₄) and concentrated *in vacuo*. The residue was purified by reverse phase HPLC to give the title compound as a colourless gum (52.0 mg, 12%). HRMS (ESI+) calcd for C₂₅H₃₀ClN₅O 452.2212, found 452.2213. HPLC: Rt 3.92 min, 100% purity.

15





EXAMPLES 7-26





Examples 7-26 were prepared similarly to Example 6, by CDI (or triphosgene) coupling of Intermediate 4 with the appropriate amine, and subsequent Boc deprotection (where required); see Table 4 below.






Table 4: Urea couplings with Intermediate 4 and subsequent Boc deprotection (where required)







Ex	Structure	Name	Yield	LCMS, HPLC
7		4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]-N-(piperidin-4-yl)piperidine-1-carboxamide	80.0 mg 19%	HRMS (ESI+) calcd for C ₂₄ H ₂₈ ClN ₅ O 438.2055, found 438.2057. HPLC: Rt 3.91 min, 100% purity.
8		4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]-N-(1-methylpiperidin-4-yl)piperidine-1-carboxamide	91.0 mg 21%	HRMS (ESI+) calcd for C ₂₅ H ₃₀ ClN ₅ O 452.2212, found 452.2214. HPLC: Rt 4.01 min, 99% purity.
9		4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]-N-[1-(propan-2-yl)piperidin-4-yl]piperidine-1-carboxamide	125 mg 27%	HRMS (ESI+) calcd for C ₂₇ H ₃₄ ClN ₅ O 480.2525, found 480.2521. HPLC: Rt 4.15 min, 99% purity.

10		N-(1-Acetylpiperidin-4-yl)-4-[1-(4-chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]piperidine-1-carboxamide	184 mg 40%	HRMS (ESI+) calcd for C ₂₆ H ₃₀ CIN ₅ O ₂ 480.2161, found 480.2159. HPLC: Rt 4.50 min, 100% purity.
11		4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]-N-[(1-methylpiperidin-4-yl)methyl]piperidine-1-carboxamide	96.0 mg 21%	HRMS (ESI+) calcd for C ₂₆ H ₃₂ CIN ₅ O 466.2368, found 466.2379. HPLC: Rt 4.02 min, 99% purity.
12		4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]-N-[(1-ethylpiperidin-4-yl)methyl]piperidine-1-carboxamide	122 mg 26%	HRMS (ESI+) calcd for C ₂₇ H ₃₄ CIN ₅ O 480.2525, found 480.2521. HPLC: Rt 4.10 min, 99% purity.
13		4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]-N-methyl-N-[(1-methylpiperidin-4-yl)methyl]piperidine-1-carboxamide; formic acid	4.57 mg 1%	LCMS (ES ⁺): 480.1 [MH] ⁺ . HPLC: Rt 4.22 min, 96% purity.


14		N-([1-(Carbamoylmethyl)piperidin-4-yl]methyl)-4-[1-(4-chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]piperidine-1-carboxamide; formic acid	32.8 mg 11%	HRMS (ESI+) calcd for C ₂₇ H ₃₃ ClN ₆ O ₂ 509.2426, found 509.2418. HPLC: Rt 3.93 min, 100% purity.
15		4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]-N-methyl-N-([1-(propan-2-yl)piperidin-4-yl]methyl)piperidine-1-carboxamide	28.2 mg 6%	HRMS (ESI+) calcd for C ₂₉ H ₃₈ ClN ₅ O 508.2838, found 508.2835. HPLC: Rt 4.39 min, 100% purity.
16		1-({4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]piperidin-1-yl}carbonyl)-4-cyclopropylpiperazine	49.2 mg 11%	HRMS (ESI+) calcd for C ₂₆ H ₃₀ ClN ₅ O 464.2212, found 464.2208. HPLC: Rt 4.11 min, 100% purity.
17		1-({4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]piperidin-1-yl}carbonyl)-4-(propan-2-yl)piperazine	38.0 mg 11%	HRMS (ESI+) calcd for C ₂₆ H ₃₂ ClN ₅ O 466.2368, found 466.2359. HPLC: Rt 4.19 min, 100% purity.

18		1-({4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]piperidin-1-yl}carbonyl)-4-(2-methoxyethyl)piperazine	69.5 mg 15%	HRMS (ESI+) calcd for C ₂₆ H ₃₂ ClN ₅ O ₂ 482.2317, found 482.2314. HPLC: Rt 4.09 min, 99% purity.
19		(3S)-1-({4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]piperidin-1-yl}carbonyl)-3-(propan-2-yl)piperazine	58.0 mg 16%	HRMS (ESI+) calcd for C ₂₆ H ₃₂ ClN ₅ O 466.2368, found 466.2363. HPLC: Rt 4.17 min, 100% purity.
20		4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]-N-(morpholin-2-ylmethyl)piperidine-1-carboxamide	120 mg 27%	HRMS (ESI+) calcd for C ₂₄ H ₂₈ ClN ₅ O ₂ 454.2004, found 454.2004. HPLC: Rt 3.93 min, 100% purity.
21		4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]-N-[(1,4-dimethylpiperazin-2-yl)methyl]piperidine-1-carboxamide	123 mg 29%	HRMS (ESI+) calcd for C ₂₆ H ₃₃ ClN ₆ O 481.2477, found 481.2475. HPLC: Rt 3.79 min, 100% purity.
22		4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]-N-[2-(morpholin-4-yl)ethyl]piperidine-1-carboxamide	91.0 mg 20%	HRMS (ESI+) calcd for C ₂₅ H ₃₀ ClN ₅ O ₂ 468.2161, found 468.2158. HPLC: Rt 4.34 min, 100% purity.

23		4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]-N-[2-(piperazin-1-yl)ethyl]piperidine-1-carboxamide	110 mg 24%	LCMS (ES ⁺): 467.0 [MH] ⁺ . HPLC: Rt 3.66 min, 99 % purity.
24		4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]-N-[2-(1-methylpiperidin-4-yl)ethyl]piperidine-1-carboxamide	83.2 mg 18%	HRMS (ESI ⁺) calcd for C ₂₇ H ₃₄ CIN ₅ O 480.2525, found 480.2525. HPLC: Rt 4.13 min, 99% purity.
25		4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]-N-[2-(4-methylpiperazin-1-yl)ethyl]piperidine-1-carboxamide	72.4 mg 16%	HRMS (ESI ⁺) calcd for C ₂₆ H ₃₃ CIN ₆ O 481.2477, found 481.2474. HPLC: Rt 3.76 min, 100% purity.
26		4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]-N-[3-(morpholin-4-yl)propyl]piperidine-1-carboxamide	91.0 mg 20%	LCMS (ES ⁺): 482.0 [MH] ⁺ . HPLC: Rt 4.01 min, 100% purity.

EXAMPLE 27


4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]-N-[[1-(propan-2-yl)piperidin-4-yl]methyl]piperidine-1-carboxamide; formic acid



Intermediate 5 (250 mg, 0.55 mmol) and acetone (81.1 μ L, 1.11 mmol) were dissolved in DCM (200 mL) and stirred for 1 h. NaBH(OAc)₃ (293 mg, 1.38 mmol) was added and the reaction mixture was stirred for 18 h, diluted with 1 M aq Na₂CO₃ (50 mL) and extracted with DCM (3 x 50 mL). The combined organic fractions were dried (MgSO₄) and concentrated *in vacuo*. The residue was purified by reverse phase HPLC (formic acid buffered) to give the title compound as a white solid (20.5 mg, 7%). LCMS (ES⁺): 494.1 [MH]⁺. HPLC: Rt 4.16 min, 98% purity.

EXAMPLE 28


4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]-N-[[1-(2-methoxyethyl)piperidin-4-yl]methyl]piperidine-1-carboxamide; formic acid



Intermediate 5 (250 mg, 0.55 mmol) was dissolved in MeCN (3 mL) and K₂CO₃ (229 mg, 1.66 mmol) and 1-bromo-2-methoxyethane (52.0 μ L, 0.55 mmol) were added. The reaction mixture was heated in a microwave reactor at 100 °C for 30 min, diluted with water (50 mL) and extracted with DCM (3 x 50 mL). The combined organic fractions were dried (MgSO₄) and concentrated *in vacuo*. The residue was purified by reverse phase HPLC (formic acid buffered) to give the title compound as a white solid (11.3 mg, 4%). LCMS (ES⁺): 510.0 [MH]⁺. HPLC: Rt 4.08 min, 94% purity.

EXAMPLE 29


N-[3-({4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]piperidin-1-yl}carbonylamino)propyl]acetamide



Intermediate 6 (64.0 mg, 0.16 mmol) was dissolved in DCM (5 mL) and Et₃N (22.8 μ L, 0.16 mmol) and Ac₂O (15.4 μ L, 0.16 mmol) were added. The reaction mixture was stirred for 18 h then diluted with sat aq Na₂CO₃ (25 mL) and extracted with DCM (3 x 25 mL). The combined organic fractions were dried (MgSO₄) and concentrated *in vacuo*. The residue was purified by reverse phase HPLC to give the title compound as a white solid (63.0 mg, 89%). HRMS (ESI+) calcd for C₂₄H₂₈ClN₅O₂ 454.2004, found 454.2004. HPLC: Rt 4.34 min, 98% purity.

EXAMPLE 30


Propan-2-yl N-({4-[1-(4-methylphenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]morpholin-2-yl}methyl)carbamate



Intermediate 10 (100 mg, 0.25 mmol) and isopropyl chloroformate (278 μ L, 0.28 mmol) were added to a mixture of DCM (2 mL) and sat aq K₂CO₃ (3.5 mL) and the reaction mixture was stirred for 1 h. The aq fraction was extracted with DCM (2 x 50 mL) and the combined organic fractions were dried (MgSO₄) and concentrated *in vacuo*. The residue was purified by column chromatography to give the title compound as a white solid (18.9 mg, 18%). HRMS (ESI+) calcd for C₂₂H₂₇N₅O₃ 410.2187, found 410.2188. HPLC: Rt 5.31 min, 98% purity.

EXAMPLE 31


3-Cyclopropyl-1-({4-[1-(4-methylphenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]morpholin-2-yl}methyl)urea



Cyclopropylamine (17.5 μ L, 0.25 mmol) and CDI (40.9 mg, 0.25 mmol) were dissolved in DMF (1 mL) and stirred for 6 h. A solution of Intermediate 10 (100 mg, 0.25 mmol) in DMF (1 mL) and DIPEA (92.6 μ L, 0.56 mmol) were added and the reaction mixture was stirred at 50 $^{\circ}$ C overnight. The solvents were removed *in vacuo* and the residue was dissolved in DCM (10 mL) and washed with sat aq Na_2CO_3 (5 mL). The aq fraction was extracted with DCM (10 mL) and the combined organic fractions were dried (MgSO_4) and concentrated *in vacuo*. The residue was purified by column chromatography to give the title compound as a light yellow solid (32.7 mg, 32%). HRMS (ESI+) calcd for $\text{C}_{22}\text{H}_{26}\text{N}_6\text{O}_2$ 407.2190, found 407.2192. HPLC: Rt 4.59 min, 98% purity.

EXAMPLE 32


2-({4-[1-(4-Methylphenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]morpholin-2-yl}methoxy)ethan-1-amine




tert-Butyl N-(2-hydroxyethyl)carbamate (159 μ L, 1.03 mmol) was dissolved in DMF (0.50 mL), NaH (49.2 mg, 60% dispersion in mineral oil, 1.03 mmol) was added and the reaction mixture was stirred at 50 $^{\circ}$ C for 30 min. A solution of Intermediate 18 (92.0 mg, 0.23 mmol) in DMF (0.5 mL) was added drop-wise and the reaction mixture was stirred at 65 $^{\circ}$ C for 18 h and at 80 $^{\circ}$ C for 4 h. The reaction mixture was cooled to 0 $^{\circ}$ C and quenched with water (1 mL). The reaction mixture was concentrated *in vacuo* and the residue was purified by column chromatography to give the title compound as an orange gum (3.05 mg, 4%). HRMS (ESI+) calcd for $\text{C}_{20}\text{H}_{25}\text{N}_5\text{O}_2$ 368.2081, found 368.2084. HPLC: Rt 3.96 min, 97.4% purity.

EXAMPLE 33

(2-Aminoethyl)({4-[1-(4-methylphenyl)-1H-pyrazolo[3,4-c]pyridin-3-


yl)morpholin-2-yl)methyl)amine trihydrochloride

Intermediate 18 (92.0 mg, 0.23 mmol), *tert*-butyl N-(2-aminoethyl)carbamate (110 mg, 0.69 mmol), K₂CO₃ (126 mg, 0.91 mmol) and Cs₂CO₃ (100 mg, 0.31 mmol) were suspended in MeCN (2 mL) and DMF (1 mL) and the reaction mixture was heated at 90 °C for 20 h. The reaction mixture was concentrated *in vacuo* and the residue was purified by column chromatography, dissolved in 1.25 M HCl in EtOH (2.5 mL) and stirred for 18 h. The reaction mixture was concentrated *in vacuo* to give the title compound as a dark yellow solid (1.52 mg, 30%). LCMS (ES⁺): 367.0 [MH]⁺. HPLC: Rt 3.57 min, 98.2% purity.

EXAMPLE 34**4-[1-(4-Methylphenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]-2-(morpholin-4-ylmethyl)morpholine**

Intermediate 18 (92.0 mg, 0.23 mmol), morpholine (60.0 µL, 0.69 mmol), K₂CO₃ (126 mg, 0.91 mmol) and Cs₂CO₃ (100 mg, 0.31 mmol) were suspended in MeCN (2 mL) and DMF (1 mL) and the reaction mixture was heated at 90 °C for 20 h. The reaction mixture was concentrated *in vacuo* and the residue was purified by column chromatography to give the title compound as a yellow solid (7.46 mg, 8%). HRMS (ESI⁺) calcd for C₂₂H₂₇N₅O₂ 394.2238, found 394.2239. HPLC: Rt 3.94 min, 94.5% purity.


EXAMPLE 35**4-[1-(4-Chlorophenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]-2-[(4-methylpiperazin-1-yl)methyl)morpholine**



Example 35 (1.23 mg, 2%) was prepared similarly to Example 34, using Intermediate 19 instead of Intermediate 18 and 1-methylpiperazine instead of morpholine. LCMS (ES+): 427.0 [MH]⁺. HPLC: Rt 3.85 min, 97.1% purity.

EXAMPLE 36


4-[1-(4-Chlorophenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]-2-(piperazin-1-ylmethyl) morpholine trihydrochloride



Intermediate 22 (75.0 mg, 0.15 mmol) was dissolved in 1.25 M HCl in EtOH (15 mL) and the reaction mixture was stirred for 18 h and concentrated *in vacuo* to give the title compound as an orange solid (74.6 mg, 97%). HRMS (ESI+) calcd for C₂₁H₂₅ClN₆O 413.1851, found 413.1853. HPLC: Rt 3.76 min, 97.8% purity.

EXAMPLE 37

3-Aminopropyl 4-({4-[1-(4-chlorophenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]morpholin-2-yl}methyl)piperazine-1-carboxylate trihydrochloride




Triphosgene (14.2 mg, 0.05 mmol) was dissolved in DCM (1 mL) and a solution of tert-butyl N-(3-hydroxypropyl)carbamate (25.2 mg, 0.14 mmol) and DIPEA (25.0 μL, 0.14 mmol) in DCM (1 mL) was added. The reaction mixture was stirred for 1 h, a solution of Example 36 (50.0 mg, 0.10 mmol) and DIPEA (25.0 μL, 0.14 mmol) in DCM (1 mL) was added and the reaction mixture was stirred for 4 d. The reaction mixture was diluted with DCM (10 mL) and washed with sat aq NH₄Cl (5 x 10 mL). The organic fraction was dried (MgSO₄) and concentrated *in vacuo*. The residue was purified by reverse phase chromatography, dissolved in 1.25 M HCl in EtOH (5

mL) and stirred for 16 h. The solvents were removed *in vacuo* to give the title compound as an orange solid (19.5 mg, 35%). HRMS (ESI+) calcd for C₂₅H₃₂ClN₇O₃ 514.2328, found 514.2326. HPLC: Rt 3.79 min, 98.3% purity.

EXAMPLE 38


N-(3-Aminopropyl)-4-({4-[1-(4-chlorophenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]morpholin-2-yl}methyl)piperazine-1-carboxamide trihydrochloride



Example 38 (24.6 mg, 41%) was prepared similarly to Example 37, using tert-butyl N-(3-aminopropyl)carbamate instead of tert-butyl N-(3-hydroxypropyl)carbamate. HRMS (ESI+) calcd for C₂₅H₃₃ClN₈O₂ 513.2488, found 513.2486. HPLC: Rt 3.73 min, 100% purity.

EXAMPLE 39

4-({4-[1-(4-Chlorophenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]morpholin-2-yl}methyl)-N-ethylpiperazine-1-carboxamide




Triphosgene (14.2 mg, 0.05 mmol) was dissolved in DCM (1 mL) and a solution of ethylamine (9.52 μ L, 0.14 mmol) and DIPEA (25.0 μ L, 0.14 mmol) in DCM (1 mL) was added. The reaction mixture was stirred for 1 h and a solution of Example 36 (50.0 mg, 0.10 mmol) and DIPEA (25.0 μ L, 0.14 mmol) in DCM (1 mL) was added. The reaction mixture was stirred for 4 d, diluted with DCM (10 mL) and washed with sat aq NH₄Cl (5 x 10 mL). The organic fraction was dried (MgSO₄) concentrated *in vacuo*. The residue was purified by reverse phase chromatography to give the title compound as a pale yellow solid (18.6 mg, 40%). HRMS (ESI+) calcd for C₂₄H₃₀ClN₇O₂ 484.2222, found 484.2219. HPLC: Rt 4.15 min, 99.3% purity.

EXAMPLE 40

Methyl 2-{4-[1-(4-chlorophenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]morpholin-3-


yl}acetate



Intermediate 33 (1.64 g, 5.94 mmol), 4-chlorophenylboronic acid (1.86 g, 11.9 mmol), $\text{Cu}(\text{OAc})_2$ (2.16 g, 11.9 mmol) and pyridine (2.39 mL, 29.7 mmol) were suspended in DCE (41 mL) and stirred overnight. The reaction mixture was purified by column chromatography to give the title compound as a yellow gum (866 mg, 38%). HRMS (ESI+) calcd for $\text{C}_{19}\text{H}_{19}\text{ClN}_4\text{O}_3$ 387.1218, found 387.1218. HPLC: Rt 5.32 min, 100% purity.

EXAMPLE 41


4-[1-(4-Chlorophenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]-3-(morpholin-4-ylmethyl)morpholine



Example 41 was prepared similarly to Example 40, by N-arylation of Intermediate 38 to give the title compound as a yellow gum (6.00 mg, 5%). HRMS (ESI+) calcd for $\text{C}_{21}\text{H}_{24}\text{ClN}_5\text{O}_2$ 414.1691, found 414.1693. HPLC: Rt 4.04 min, 97.6% purity.

EXAMPLE 42

4-[1-(4-Chlorophenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]-3-[2-(4-methylpiperazin-1-yl)ethyl]morpholine




Intermediate 49 (60.0 mg, 0.14 mmol), 1-methyl-piperazine (45.7 μL , 0.41 mmol) and K_2CO_3 (75.9 mg, 0.55 mmol) were suspended in MeCN (1 mL) and the reaction mixture was heated at 50 °C for 4 h and at 75 °C for 4 h. The reaction

mixture was filtered, concentrated *in vacuo* and the residue was purified by column chromatography to give the title compound as a dark yellow gum (3.47 mg, 6%). HRMS (ESI+) calcd for C₂₃H₂₉ClN₆O 441.2164, found 441.2164. HPLC: Rt 3.89 min, 99.1% purity.

EXAMPLE 43


1-[1-(4-Chlorophenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]-N-[(1-methylpiperidin-4-yl)methyl]piperidine-2-carboxamide



Intermediate 47 (66.0 mg, 0.17 mmol) was dissolved in DMF (2 mL), cooled to 0 °C and HBTU (63.3 mg, 0.17 mmol), (1-methyl-4-piperidiny)methanamine (25.7 mg, 0.20 mmol) and DIPEA (58.2 µL, 0.33 mmol) were added. The reaction mixture was stirred at 0 °C for 1 h and at RT for 18 h, diluted with DCM (10 mL) and washed with sat aq NH₄Cl (3 x 5 mL). The organic fraction was dried (MgSO₄) and concentrated *in vacuo*. The residue was purified by column chromatography to give the title compound as a yellow solid (8.62 mg, 11%). LCMS (ES⁺): 467.0 [MH]⁺. HPLC: Rt 4.19 min, 100% purity.


EXAMPLE 44

1-(4-Chlorophenyl)-N-[2-(morpholin-4-yl)ethyl]-1H-pyrazolo[3,4-c]pyridin-3-amine




Example 44 was prepared similarly to Example 40, by N-arylation of Intermediate 35 and subsequent Boc deprotection (HCl in EtOH) to give the title compound as a pale brown gum (2.27 mg, 1%). HRMS (ESI+) calcd for C₁₈H₂₀ClN₅O 358.1429, found 358.1434. HPLC: Rt 4.00 min, 97% purity.

EXAMPLE 45

1-(4-Chlorophenyl)-N-[2-(piperazin-1-yl)ethyl]-1H-pyrazolo[3,4-c]pyridin-3-amine


Example 45 was prepared similarly to Example 40, by N-arylation of Intermediate 36 and subsequent Boc deprotection (HCl in Et₂O) to give the title compound as a pale green gum (9.98 mg, 10%). HRMS (ESI+) calcd for C₁₈H₂₁ClN₆ 357.1589, found 357.1592. HPLC: Rt 3.57 min, 99.5% purity.

EXAMPLE 46

1-(4-Chlorophenyl)-N-[2-(4-methylpiperazin-1-yl)ethyl]-1H-pyrazolo[3,4-c]pyridin-3-amine

Example 45 (24.6 mg, 0.07 mmol) was dissolved in MeOH (2 mL), formaldehyde (55.9 mg, 37 % in water, 0.69 mmol) was added and the reaction mixture was stirred for 30 min. NaBH(OAc)₃ (17.5 mg, 0.08 mmol) was added and the reaction mixture was stirred overnight and concentrated *in vacuo*. The residue was purified by reverse phase HPLC to give the title compound as a pale green gum (16.0 mg, 63%). HRMS (ESI+) calcd for C₁₉H₂₃ClN₆ 371.1745, found 371.1751. HPLC: Rt 3.54 min, 100% purity.

EXAMPLE 47


1-[1-(4-Chlorophenyl)-1H-pyrazolo [3,4-c]pyridin-3-yl]-N-(piperidin-4-ylmethyl)piperidine-4-carboxamide dihydrochloride

Intermediate 44 (200 mg, 0.51 mmol) was dissolved in DMF (2 mL) and HBTU (231

mg, 0.61 mmol) was added. The reaction mixture was stirred for 30 min, *tert*-butyl 4-(aminomethyl)piperidine-1-carboxylate (131 mg, 0.61 mmol) and DIPEA (266 μ L, 1.53 mmol) were added and the reaction mixture was stirred overnight. The solvents were removed *in vacuo* and the residue was diluted with EtOAc (25 mL), washed with sat aq NH_4Cl (4 x 25 mL), dried (MgSO_4) and concentrated *in vacuo*. The residue was purified by column chromatography and half of the product was dissolved in 1.25 M HCl in EtOH (10 mL) and stirred for 18 h. The reaction mixture was concentrated *in vacuo* to give the title compound as an orange solid (77.9 mg, 58%). HRMS (ESI+) calcd for $\text{C}_{24}\text{H}_{29}\text{ClN}_6\text{O}$ 453.2164, found 453.2163. HPLC: Rt 4.07 min, 98.1% purity.

EXAMPLE 48


4-({1-[1-(4-Chlorophenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]piperidin-4-yl)methyl}morpholine dihydrochloride



Intermediate 45 (100 mg, 0.23 mmol) was dissolved in THF (1 mL) and 1.0 M BH_3 in THF (1.88 mL, 1.88 mmol) was added portion-wise with heating at 67 $^{\circ}\text{C}$ for 2 d. The reaction mixture was cooled to 0 $^{\circ}\text{C}$, quenched with cold water (2 mL) and concentrated *in vacuo*. The residue was purified by column chromatography, dissolved in 1.25 M HCl in EtOH (5 mL), stirred for 4 h and concentrated *in vacuo* to give the title compound as an orange solid (5.68 mg, 5%). HRMS (ESI+) calcd for $\text{C}_{22}\text{H}_{26}\text{ClN}_5\text{O}$ 412.1899, found 412.1896. HPLC: Rt 4.13 min, 98.1% purity.


EXAMPLES 49-50

Examples 49-50 were prepared similarly to Example 48, by borane reduction of Intermediate 46 and Boc protected Example 47, and subsequent Boc deprotection; see Table 5 below.

Table 5: Borane reduction and subsequent Boc deprotection


Ex	Structure	Name	Int / Yield	LCMS, HPLC
49		1-({1-[1-(4-Chlorophenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]piperidin-4-yl}methyl)piperazine	Int 46 3.35 mg 3%	HRMS (ESI+) calcd for C ₂₂ H ₂₇ ClN ₆ 411.2058, found 411.2061. HPLC: Rt 3.75 min, 100% purity.
50		({1-[1-(4-Chlorophenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]piperidin-4-yl}methyl)(piperidin-4-ylmethyl)amine	Ex 47(Boc) 9.13 mg 9 %	HRMS (ESI+) calcd for C ₂₄ H ₃₁ ClN ₆ 439.2371, found 439.2372. HPLC: Rt 3.74 min, 99.6% purity.

EXAMPLE 51

4-[1-(4-Chlorophenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]-N-[(1-methylpiperidin-4-yl)methyl]piperazine-1-carboxamide

Triphosgene (14.2 mg, 0.05 mmol) was dissolved in DCM (1 mL) and a solution of (1-methyl-4-piperidiny)methanamine (18.4 mg, 0.14 mmol) and DIPEA (25.0 μ L, 0.14 mmol) in DCM (1 mL) was added. The reaction mixture was stirred for 1 h and a solution of Intermediate 50 (37.0 mg, 0.10 mmol) and DIPEA (25.0 μ L, 0.14 mmol) in DCM (1 mL) was added. The reaction mixture was stirred for 18 h, diluted with DCM (10 mL), washed with sat aq NH_4Cl (5 x 10 mL), dried (MgSO_4) and concentrated *in vacuo*. The residue was purified by reverse phase chromatography to give the title compound as a yellow solid (17.0 mg, 38%). LCMS (ES^+): 468.0 $[\text{MH}]^+$. HPLC: Rt 4.13 min, 98.7% purity.


EXAMPLE 52

1-[1-(4-Methylphenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]piperidin-4-yl acetate


Intermediate 37 (618 mg, 2.37 mmol) was dissolved in DMF (3.5 mL) and 1-methyl-4-iodo-benzene (621 mg, 2.85 mmol), *N,N*-dimethylethylenediamine (51.1 μ L, 0.42 mmol), K_3PO_4 (1.06 g, 4.99 mmol) and CuI (45.2 mg, 0.24 mmol) were added. The reaction mixture was heated using a microwave reactor at 60 $^{\circ}\text{C}$ for 10 min. The reaction mixture was concentrated *in vacuo* and the residue was purified by column chromatography to give the title compound as a yellow gum (248 mg, 30%). HRMS (ESI^+) calcd for $\text{C}_{20}\text{H}_{22}\text{N}_4\text{O}_2$ 351.1816, found 351.1819. HPLC: Rt 5.44 min, 100% purity.

EXAMPLE 53

2-[4-[1-(4-Chlorophenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]morpholin-3-yl]acetic

acid hydrochloride


Example 40 (1.00 g, 2.84 mmol) was dissolved in 1:1 THF/water (16 mL), LiOH.H₂O (262 mg, 6.24 mmol) was added and the reaction mixture was stirred for 3 h. The THF was removed *in vacuo* and the reaction mixture was acidified to pH 1 with 1 M aq HCl (5 mL). The precipitate was collected by filtration and washed with water to give the title compound as an orange solid (28.3 mg, 3%). HRMS (ESI+) calcd for C₁₈H₁₇ClN₄O₃ 373.1062, found 373.1062. HPLC: Rt 4.40 min, 97% purity.

EXAMPLE 54**N-(2-Aminoethyl)-2-{4-[1-(4-chlorophenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]morpholin-3-yl}acetamide dihydrochloride**



Example 53 (180 mg, 0.44 mmol) was dissolved in DMF (2.1 mL) and cooled to 0 °C, and HBTU (167 mg, 0.44 mmol), *tert*-butyl N-(2-aminoethyl)carbamate (84.6 mg, 0.53 mmol) and DIPEA (76.6 µL, 0.44 mmol) were added. The reaction mixture was stirred at 0 °C for 2.5 h and purified by column chromatography. The residue was dissolved in 1.25 M HCl in EtOH (2.5 mL) and stirred for 2 h. The reaction mixture was concentrated *in vacuo* to give the title compound as an orange solid (46.4 mg, 22%). HRMS (ESI+) calcd for C₂₀H₂₃ClN₆O₂ 415.1644, found 415.1638. HPLC: Rt 3.97 min, 99% purity.

EXAMPLES 55-58

Examples 55-58 were prepared similarly to Example 54, by amide coupling to Example 53 (no HCl salt formation step); see Table 6 below.


Table 6: Amide couplings to Example 53

Ex	Structure	Name	Yield	LCMS, HPLC
55		2-{4-[1-(4-Chlorophenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]morpholin-3-yl}-1-(4-methylpiperazin-1-yl)ethan-1-one	7.95 mg 12%	LCMS (ES+): 454.9 $[\text{MH}]^+$. HPLC: Rt 4.09 min, 99.8% purity.
56		2-{4-[1-(4-Chlorophenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]morpholin-3-yl}-1-[(3S)-3-(dimethylamino)pyrrolidin-1-yl]ethan-1-one	30.9 mg 22%	LCMS (ES+): 468.9 $[\text{MH}]^+$. HPLC: Rt 4.11min, 98.6% purity.

57		2-{4-[1-(4-Chlorophenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]morpholin-3-yl}-N-(1-methylpiperidin-4-yl)acetamide	31.8 mg 46%	LCMS (ES ⁺): 468.9 [MH] ⁺ . HPLC: Rt 4.06 min, 98.8% purity.
58		2-{4-[1-(4-Chlorophenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]morpholin-3-yl}-N-[(1-methylpiperidin-4-yl)methyl]acetamide	17.9 mg 25%	LCMS (ES ⁺): 482.9 [MH] ⁺ . HPLC: Rt 4.10 min, 98.1% purity.

EXAMPLE 59


({4-[1-(4-Chlorophenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]morpholin-2-yl}methyl)urea



- Intermediate 52 (50.0 mg, 0.12 mmol) and DIPEA (41.8 μ L, 0.24 mmol) were dissolved in DMF (2mL). Trimethylsilyl isocyanate (29.2 μ L, 0.22 mmol) was added and the reaction mixture was stirred for 48 h. The reaction mixture was concentrated *in vacuo* and the residue was dissolved in EtOAc (10 mL) and washed with water (2 x 5 mL). The organic fraction was concentrated *in vacuo* and purified by reverse phase HPLC to give the title compound as a white solid (16.5 mg, 35%). LCMS (ES⁺): 387.2 [MH]⁺. HPLC: Rt 4.40 min, 96.8% purity.

EXAMPLE 60


1-({4-[1-(4-Chlorophenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]morpholin-2-yl}methyl)-3-methylurea



- Intermediate 52 (50.0 mg, 0.12 mmol) and DIPEA (62.7 μ L, 0.36 mmol) were dissolved in THF (2mL). N-methylcarbamoyl chloride (12.3mg, 0.132mmol) was added and the reaction mixture was stirred overnight. The reaction mixture was concentrated *in vacuo* and the residue was dissolved in EtOAc (10 mL) and washed with water (2 x 5 mL). The organic fraction was concentrated *in vacuo* and purified by reverse phase HPLC to give the title compound as an off white solid (11.1 mg, 23%). LCMS (ES⁺): 401.1 [MH]⁺. HPLC: Rt 4.55min, 99.7% purity.

EXAMPLE 61


4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]-N-(2H-1,2,3,4-tetrazol-5-yl)methyl)piperidine-1-carboxamide; trifluoroacetic acid



CDI (172 mg, 1.06 mmol) was dissolved in DCM (10 mL) and a suspension of 1H-1,2,3,4-tetrazol-5-ylmethanamine hydrochloride (143 mg, 1.06 mmol) and DIPEA (368 μ L, 2.12 mmol) in DCM was added. The reaction mixture was stirred for 6 h and a solution of Intermediate 4 (300 mg, 0.96 mmol) and DIPEA (368 μ L, 2.12 mmol) in DCM (2 mL) was added. The reaction mixture was stirred for 3 d and concentrated *in vacuo*. The residue was purified by reverse phase HPLC to give the title compound as a white solid (134 mg, 25%). LCMS (ES⁺): 437.0 [MH]⁺. HPLC: Rt 4.29 min, 100% purity.

EXAMPLE 62


4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]-N-[cyclopropyl(2H-1,2,3,4-tetrazol-5-yl)methyl]piperidine-1-carboxamide; trifluoroacetic acid



Example 62 (252 mg, 53%) was prepared similarly to Example 61, using cyclopropyl(1H-1,2,3,4-tetrazol-5-yl)methanamine instead of 1H-1,2,3,4-tetrazol-5-ylmethanamine hydrochloride. LCMS (ES⁺): 477.1 [MH]⁺. HPLC: Rt 4.69 min, 99% purity.

EXAMPLE 63

4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]-N-[1-(2H-1,2,3,4-tetrazol-5-yl)cyclobutyl]piperidine-1-carboxamide; trifluoroacetic acid




Example 63 (92.0 mg, 19%) was prepared similarly to Example 61, using 1-(1H-1,2,3,4-tetrazol-5-yl)cyclobutan-1-amine instead of 1H-1,2,3,4-tetrazol-5-ylmethanamine hydrochloride. LCMS (ES⁺): 477.1 [MH]⁺. HPLC: Rt 4.57 min, 99%

purity.

EXAMPLE 64

1-(2-[[1-(4-Chlorophenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]amino]ethyl)imidazolidin-2-one



Intermediate 55 (70.0 mg, 0.15 mmol) was dissolved in HCl in EtOH (1.25 M, 20 mL) and stirred for 5 d. The reaction mixture was concentrated *in vacuo*, partitioned between DCM (15 mL) and sat aq NaHCO₃ (10 mL) and the organic fraction was dried (MgSO₄) and concentrated *in vacuo* to give the title compound (47.5 mg, 87%) as a white solid. LCMS (ES⁺): 357.1 [MH]⁺. HPLC: Rt 4.89min, 99.6% purity.

BIOLOGICAL TESTS

Biological Assays of the SSAO Enzyme Inhibitors

All primary assays were performed at RT. with purified recombinantly expressed human SSAO. Enzyme was prepared essentially as described in Öhman et al. (Protein Expression and Purification **46** (2006) 321–331). In addition, secondary- and selectivity assays were performed using SSAO prepared from various tissues or purified rat recombinant SSAO. The enzyme activity was assayed with benzylamine as substrate by measuring either benzaldehyde production, using ¹⁴C-labeled substrate, or by utilizing the production of hydrogen peroxide in a horseradish peroxidase (HRP) coupled reaction. Briefly, test compounds were dissolved in dimethyl sulfoxide (DMSO) to a concentration of 10 mM. Dose-response measurements were assayed by either creating 1:10 serial dilutions in DMSO to produce a 7 point curve or by making 1:3 serial dilutions in DMSO to produce 11 point curves. The top concentrations were adjusted depending on the potency of the compounds and subsequent dilution in reaction buffer yielded a final DMSO concentration ≤ 2%.

Hydrogen peroxide detection:

In a horseradish peroxidase (HRP) coupled reaction, hydrogen peroxide oxidation of 10-acetyl-3,7-dihydroxyphenoxazine produced resorufin, which is a highly fluorescent compound (Zhout and Panchuk-Voloshina. Analytical Biochemistry **253** (1997) 169-174; Amplex[®] Red Hydrogen Peroxide/peroxidase Assay kit, Invitrogen
5 A22188). Enzyme and compounds in 50 mM sodium phosphate, pH 7.4 were set to pre-incubate in flat-bottomed microtiter plates for approximately 15 minutes before initiating the reaction by addition of a mixture of HRP, benzylamine and Amplex reagent. Benzylamine concentration was fixed at a concentration corresponding to the Michaelis constant, determined using standard procedures. Fluorescence
10 intensity was then measured at several time points during 1 – 2 hours, exciting at 544 nm and reading the emission at 590 nm. For the human SSAO assay final concentrations of the reagents in the assay wells were: SSAO enzyme 1 µg/ml, benzylamine 100 µM, Amplex reagent 20 µM, HRP 0.1 U/mL and varying concentrations of test compound. The inhibition was measured as % decrease of
15 the signal compared to a control without inhibitor (only diluted DMSO). The background signal from a sample containing no SSAO enzyme was subtracted from all data points. Data was fitted to a four parameter logistic model and IC₅₀ values were calculated using the GraphPad Prism 4 or XLfit 4 programs.

20 Aldehyde detection:

SSAO activity was assayed using 14C-labeled benzylamine and analysed by measuring radioactive benzaldehyde. In a white 96-well optiplate (Packard), 20 µL of diluted test compound was pre-incubated at RT. with 20 µL SSAO enzyme for approximately 15 minutes with continuous agitation. All dilutions were made with
25 PBS. The reaction was initiated by adding 20 µL of the benzylamine substrate solution containing [7-14C] Benzylamine hydrochloride (CFA589, GE Healthcare). The plate was incubated for 1 hour as above after which the reaction was stopped by acidification (10 µL 1 M HCl). Then 90 µL Micro Scint-E solution (Perkin-Elmer) was added to each well and the plate was continuously mixed for 15 minutes. Phase
30 separation occurred instantly and activity was read in a Topcount scintillation counter (Perkin-Elmer). In the final reaction well, the human recombinant SSAO concentration was 10 µg/ml. In order to optimize sensitivity, the substrate concentration was decreased as compared to the HRP coupled assay in order to get a higher fraction of radioactive product. In the human SSAO assay, benzylamine
35 concentration was 40 µM (0.2 µCi/mL). Data was analysed as above.

All of the exemplified compounds of the invention had an IC₅₀ value of 1-2500 nM at SSAO (See Table 7).

5 **Table 7: SSAO inhibitory activity**

(A: <100nM, B: 100-500nM, C: 500-2500nM)

Compound	IC ₅₀ (nM)	Compound	IC ₅₀ (nM)	Compound	IC ₅₀ (nM)
1	A	23	A	45	B
2	B	24	A	46	B
3	B	25	A	47	B
4	A	26	A	48	A
5	A	27	A	49	B
6	A	28	A	50	B
7	A	29	A	51	A
8	A	30	A	52	A
9	A	31	A	53	C
10	A	32	B	54	A
11	A	33	B	55	A
12	A	34	B	56	A
13	A	35	B	57	B
14	A	36	A	58	B
15	A	37	B	59	A
16	A	38	B	60	A
17	A	39	B	61	A
18	A	40	A	62	A
19	A	41	B	63	A
20	A	42	A	64	A
21	A	43	B		
22	A	44	C		

CLAIMS

1. A compound of formula (I) or a pharmaceutically acceptable salt, or N-oxide thereof:



wherein

R¹ is phenyl or 6-membered heteroaryl, optionally substituted with one or more substituents selected from halogen, cyano, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, C₁₋₄alkoxy-C₁₋₄alkyl, hydroxy-C₁₋₄-alkyl, cyano-C₁₋₄-alkyl, amino-C₁₋₄-alkyl, C₁₋₄-alkylamino-C₁₋₄-alkyl, di(C₁₋₄-alkyl)amino-C₁₋₄-alkyl, -NR^{4A}R^{4B}, -NR⁶C(O)OR⁵, -NR⁶C(O)R⁵, -NR⁶C(O)NR^{4A}R^{4B}, -C(O)NR^{4A}R^{4B}, -C(O)R⁵, -C(O)OR⁵, and -NR⁶S(O)₂R⁵;

R² is -B-Q-[R³]_n or -B-R³;

wherein n = 1, 2, 3, or 4

B is a bond, O, NR⁴, -C(O)- or C₁₋₃-alkylene;

Q is saturated or partially unsaturated monocyclic 3-7 membered heterocyclic or C₃₋₇-cycloalkyl ring;

when R² is -B-Q-[R³]_n, R³ is independently selected from: 3-7 membered heterocyclyl-, 3-7 membered heterocyclyl-C₁₋₄-alkyl-, (3-7 membered heterocyclyl-C₁₋₄-alkyl)-amino-C₁₋₄-alkyl-, amino-C₁₋₄-alkoxy-C₁₋₄-alkyl-, (amino-C₁₋₄-alkyl)-amino-C₁₋₄-alkyl-, -C₁₋₄-alkyl-NR⁶C(O)OR⁵, -C₁₋₄-alkyl-NR⁶C(O)NR^{4A}R^{4B}, -C₁₋₄-alkyl-C(O)NR^{4A}R^{4B}, (3-7 membered heterocyclyl-C₁₋₄-alkyl)-C(O)-, -C₁₋₄-alkyl-C(O)OR⁵, -OC(O)R⁵, or

-C(O)NR^{9A}R^{9B} wherein R^{9A} and R^{9B} together with the nitrogen to which they are attached form a 3-7 membered cyclic amino group substituted with one or more substituents selected from: C₁₋₄-alkyl, C₁₋₄alkoxy-C₁₋₄alkyl-, C₃₋₇-cycloalkyl, or

-C(O)NR⁶R^{10B} wherein R^{10B} is:

(i) 3-7 membered heterocyclyl- or 3-7 membered heterocyclyl-C₁₋₄-alkyl-, or -C₁₋₄-alkyl-NR⁶C(O)R⁵; or

(ii) 5 or 6 membered heteroaryl-C₁₋₄-alkyl-, wherein the heteroaryl ring is optionally substituted with one or more substituents selected from halogen, cyano, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, and wherein the C₁₋₄-alkyl part is optionally substituted by one or more C₁₋₄-alkyl- groups, or the C₁₋₄-alkyl part is substituted with two C₁₋₄-alkyl groups which, together with the carbon atom to which they are attached, join together to form a spiro 3-6 membered cycloalkyl ring; and wherein

10

when R² is -B-R³, R³ is -NR⁶R^{11B}, and R^{11B} is 3-7 membered heterocyclyl-C₁₋₄-alkyl-;

R^{4A}, R^{4B} and R⁵ are each independently selected from hydrogen, C₁₋₄-alkyl-, 3-7 membered heterocyclyl-C₁₋₄-alkyl-, amino-C₁₋₄-alkyl-, 3-7 membered heterocyclyl-, -C₁₋₄-alkyl-NR⁶C(O)OR⁵, C₃₋₇-cycloalkyl, or

15

R^{4A} and R^{4B} together with the nitrogen to which they are attached form a 3-7 membered cyclic amino group, optionally substituted by one or more substituents selected from: C₁₋₄-alkyl, -NR^{4A}R^{4B}; and wherein

20

unless otherwise specified, 3-7 membered heterocyclyl, or the heterocyclyl part of the 3-7 membered heterocyclyl-C₁₋₄-alkyl-, (3-7 membered heterocyclyl-C₁₋₄-alkyl)-amino-C₁₋₄-alkyl-, or (3-7 membered heterocyclyl-C₁₋₄-alkyl)-C(O)- group is optionally substituted with one or more substituents selected from oxo, C₁₋₄-alkyl-, -C(O)OR⁵, -C(O)R⁵, -C(O)NR^{4A}R^{4B}, -NR^{4A}R^{4B}, -C₁₋₄-alkyl-C(O)NR^{4A}R^{4B}, or C₁₋₄alkoxy-C₁₋₄alkyl; and

25


where present, the diradical -C₁₋₄-alkyl- group directly attached to Q is optionally substituted with one or more groups independently selected from halogen, amino, methoxy, hydroxyl; and wherein

30

R⁴ and R⁶ are each independently selected from hydrogen or C₁₋₄-alkyl; and

X is selected from the radicals of formulae (1-16) wherein the bond marked * is attached to R¹- and the bond marked ** is attached to -R²:

35



wherein Y is selected from hydrogen, hydroxyl, amino, -NHR^6 , -OCH_3 ;


- 5 Z is selected from hydrogen, fluorine, hydroxyl, C_{1-4} -alkoxy, halo- C_{1-4} -alkyl, CONH_2 , cyano, SO_2NH_2 , amino, -NHR^6 ;

W is selected from H, C_{1-4} -alkyl, halo- C_{1-4} -alkyl,

- 10 **PROVIDED THAT** when R^2 is $\text{-B-Q-[R}^3\text{]}_n$, and R^3 is 3-7 membered heterocyclyl-, the R^3 heterocyclic ring atom directly bonded to Q is not nitrogen.

2. A compound of formula (I) or a pharmaceutically acceptable salt, or N-oxide thereof:

15



wherein

R^1 is phenyl or 6-membered heteroaryl, optionally substituted with one or more substituents selected from halogen, cyano, C_{1-4} -alkyl, halo- C_{1-4} -alkyl, C_{1-4} alkoxy- C_1 .

alkyl, hydroxy-C₁₋₄-alkyl, cyano-C₁₋₄-alkyl, amino-C₁₋₄-alkyl, C₁₋₄-alkylamino-C₁₋₄-alkyl, di(C₁₋₄-alkyl)amino-C₁₋₄-alkyl, -NR^{4A}R^{4B}, -NR⁶C(O)OR⁵, -NR⁶C(O)R⁵, -NR⁶C(O)NR^{4A}R^{4B}, -C(O)NR^{4A}R^{4B}, -C(O)R⁵, -C(O)OR⁵, and -NR⁶S(O)₂R⁵;

5 R² is -B-Q-[R³]_n or -B-R³;

wherein n = 1, 2, 3, or 4

B is a bond, O, NR⁴, -C(O)- or C₁₋₃-alkylene;

10

Q is saturated or partially unsaturated monocyclic 3-7 membered heterocyclic or C₃₋₇-cycloalkyl ring;

when R² is -B-Q-[R³]_n, R³ is independently selected from: 3-7 membered heterocycl-
 15 heterocycl-alkyl-, 3-7 membered heterocycl-alkyl-C₁₋₄-alkyl-, (3-7 membered heterocycl-alkyl-C₁₋₄-alkyl)-amino-C₁₋₄-alkyl-, amino-C₁₋₄-alkoxy-C₁₋₄-alkyl-, (amino-C₁₋₄-alkyl)-amino-C₁₋₄-alkyl-, -C₁₋₄-alkyl-NR⁶C(O)OR⁵, -C₁₋₄-alkyl-NR⁶C(O)NR^{4A}R^{4B}, -C₁₋₄-alkyl-C(O)NR^{4A}R^{4B}, (3-7 membered heterocycl-alkyl-C₁₋₄-alkyl)-C(O)-, -C₁₋₄-alkyl-C(O)OR⁵, -OC(O)R⁵, or

20

-C(O)NR^{9A}R^{9B} wherein R^{9A} and R^{9B} together with the nitrogen to which they are attached form a 3-7 membered cyclic amino group substituted with one or more substituents selected from: C₁₋₄-alkyl, C₁₋₄alkoxy-C₁₋₄alkyl-, C₃₋₇-cycloalkyl, or

25 -C(O)NR⁶R^{10B} wherein R^{10B} is 3-7 membered heterocycl- or 3-7 membered heterocycl-alkyl-, or -C₁₋₄-alkyl-NR⁶C(O)R⁵; or

when R² is -B-R³, R³ is -NR⁶R^{11B}, wherein R^{11B} is 3-7 membered heterocycl-alkyl-;

30

R^{4A}, R^{4B} and R⁵ are each independently selected from hydrogen, C₁₋₄-alkyl-, 3-7 membered heterocycl-alkyl-, amino-C₁₋₄-alkyl-, 3-7 membered heterocycl-, -C₁₋₄-alkyl-NR⁶C(O)OR⁵, C₃₋₇-cycloalkyl,

or R^{4A} and R^{4B} together with the nitrogen to which they are attached form a 3-7 membered cyclic amino group, optionally substituted by one or more substituents selected from: C_{1-4} -alkyl, $-NR^{4A}R^{4B}$;


- 5 unless otherwise specified, 3-7 membered heterocyclyl, or the heterocyclyl part of the 3-7 membered heterocyclyl- C_{1-4} -alkyl-, (3-7 membered heterocyclyl- C_{1-4} -alkyl)-amino- C_{1-4} -alkyl-, or (3-7 membered heterocyclyl- C_{1-4} -alkyl)-C(O)- group is optionally substituted with one or more substituents selected from C_{1-4} -alkyl-, $-C(O)OR^5$, $-C(O)R^5$, $-C(O)NR^{4A}R^{4B}$, $-NR^{4A}R^{4B}$, $-C_{1-4}$ -alkyl-C(O) $NR^{4A}R^{4B}$, or C_{1-4} alkoxy- C_{1-4} alkyl;
- 10 and

where present, the diradical $-C_{1-4}$ -alkyl- group directly attached to Q is optionally substituted with one or more groups independently selected from halogen, amino, methoxy, hydroxyl;

15

R^4 and R^6 are each independently selected from hydrogen or C_{1-4} -alkyl; and

X is selected from the radicals of formulae (1-16) wherein the bond marked * is attached to R^1 - and the bond marked ** is attached to $-R^2$:



wherein Y is selected from hydrogen, hydroxyl, amino, -NHR^6 , -OCH_3 ;

- 5 Z is selected from hydrogen, fluorine, hydroxyl, C_{1-4} -alkoxy, halo- C_{1-4} -alkyl, CONH_2 , cyano, SO_2NH_2 , amino, -NHR^6 ;

W is selected from H, C_{1-4} -alkyl, halo- C_{1-4} -alkyl,

- 10 **PROVIDED THAT** when R^2 is $\text{-B-Q-[R}^3\text{]}_n$, and R^3 is 3-7 membered heterocyclyl-, the heterocyclic ring atom directly bonded to Q is not nitrogen.

3. A compound according to claim 1 or 2 wherein X is selected from the radicals of formulae 1 or 3.

15

4. A compound according to any one of claims 1 to 3 wherein R^1 is phenyl optionally substituted with one or more substituents as defined in claim 1.


5. A compound according to any one of claims 1 to 4 wherein R^1 is optionally substituted by halogen, cyano, C_{1-4} -alkyl, halo- C_{1-4} -alkyl.

6. A compound according to any one of claims 1 to 5 wherein B is a bond.

7. A compound as claimed in any of claims 1 to 6 wherein R^2 is $-B-Q-[R^3]_n$, and Q is a saturated or partially unsaturated 5 or 6 membered heterocyclic or cycloalkyl ring.

8. A compound as claimed in claim 7 wherein Q is selected from tetrahydrofuranyl, tetrahydropyranyl, piperidiny, piperaziny, morpholinyl, cyclohexyl, or any of the foregoing rings comprising a bridge formed by an ethylene or propylene radical.

9. A compound according to any one of claims 1 to 7 wherein R^2 is:




wherein

T is N or CH

R^6 is hydrogen or C_{1-4} -alkyl

R^{10B} is 3-7 membered heterocyclyl-, or 3-7 membered heterocyclyl- C_{1-4} -alkyl-, either of which heterocyclic rings is optionally substituted by one or more substituents selected from C_{1-4} -alkyl- and C_{1-4} alkoxy- C_{1-4} alkyl.

10. A compound according to claim 9 wherein R^2 is:



wherein:


T is N or CH;

P is a direct bond or a diradical selected from methylene, ethylene, or propylene;

R^6 is hydrogen or C_{1-4} -alkyl;

R¹² is selected from hydrogen, C₁₋₄-alkyl- and C₁₋₄alkoxy-C₁₋₄alkyl-.

11. A compound according to claim 9 wherein R² is:



5 wherein

T is N or CH;


P is a diradical selected from methylene, ethylene, or propylene;

R⁶ is hydrogen or C₁₋₄-alkyl;

R¹² is selected from hydrogen, C₁₋₄-alkyl, and C₁₋₄alkoxy-C₁₋₄alkyl-.

10

12. A compound according to any one of claims 1 to 7 wherein R² is:



wherein

15 R³ is -C₁₋₄-alkylC(O)NR^{4A}R^{4B} wherein

R^{4A} and R^{4B} are each independently selected from hydrogen, C₁₋₄-alkyl-, and amino-C₁₋₄-alkyl-, or

R^{4A} and R^{4B} together with the nitrogen to which they are attached form a 3-7 membered cyclic amino group, optionally substituted by one or more substituents

20 selected from: C₁₋₄-alkyl, or -NR^{4A}R^{4B}; or

13. A compound as claimed in claim 1 which is:

4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]-N-(piperidin-4-ylmethyl)piperidine-1-carboxamide

25 4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]-N-(1-methylpiperidin-4-yl)piperidine-1-carboxamide

4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]-N-[(1-methylpiperidin-4-yl)methyl]piperidine-1-carboxamide

30 **4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]-N-[(1-ethylpiperidin-4-yl)methyl]piperidine-1-carboxamide**

4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]-N-methyl-N-[(1-methylpiperidin-4-yl)methyl]piperidine-1-carboxamide

4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]-N-[2-(piperazin-1-yl)ethyl]piperidine-1-carboxamide

5 **4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]-N-[2-(1-methylpiperidin-4-yl)ethyl]piperidine-1-carboxamide**

4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]-N-[3-(morpholin-4-yl)propyl]piperidine-1-carboxamide

4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]-N-[[1-(propan-2-yl)piperidin-4-yl)methyl]piperidine-1-carboxamide

4-[1-(4-Chlorophenyl)-1H-pyrrolo[2,3-c]pyridin-3-yl]-N-[[1-(2-methoxyethyl)piperidin-4-yl)methyl]piperidine-1-carboxamide

4-[1-(4-Chlorophenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]-N-[(1-methylpiperidin-4-yl)methyl]piperazine-1-carboxamide

15 N-(2-Aminoethyl)-2-{4-[1-(4-chlorophenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl] morpholin-3-yl}acetamide

2-{4-[1-(4-Chlorophenyl)-1H-pyrazolo[3,4-c]pyridin-3-yl]morpholin-3-yl}-1-[(3S)-3-(dimethylamino)pyrrolidin-1-yl]ethan-1-one

20 or a pharmaceutically acceptable salt, or N-oxide thereof.

14. A pharmaceutical composition comprising a compound as claimed in any of the preceding claims, together with one or more pharmaceutically acceptable carriers and/or excipients

15. A compound as claimed in any of claims 1 to 13 for use in the treatment of inflammation, an inflammatory disease, an immune or an autoimmune disorder, or inhibition of tumour growth.

16. The use of a compound as claimed in any of claims 1 to 13 in the manufacture of a composition for treatment of inflammation, an inflammatory disease, an immune or an autoimmune disorder, or inhibition of tumour growth.

17. A method for the treatment of inflammation, an inflammatory disease, an immune or an autoimmune disorder, or inhibition of tumour growth, which comprises

administering to a subject suffering such disease an effective amount of a compound of formula (I) as claimed in any of claims 1 to 13.

18. A compound as claimed in any one of claims 1 to 13, or use as claimed in claim 15, or the method as claimed in claim 17 wherein the inflammation or inflammatory disease or immune or autoimmune disorder is arthritis (including rheumatoid arthritis, juvenile rheumatoid arthritis, osteoarthritis and psoriatic arthritis), synovitis, vasculitis, a condition associated with inflammation of the bowel (including Crohn's disease, ulcerative colitis, inflammatory bowel disease and irritable bowel syndrome), atherosclerosis, multiple sclerosis, Alzheimer's disease, vascular dementia, a pulmonary inflammatory disease (including asthma, chronic obstructive pulmonary disease and acute respiratory distress syndrome), a fibrotic disease (including cystic fibrosis, idiopathic pulmonary fibrosis, cardiac fibrosis and systemic sclerosis (scleroderma)), an inflammatory disease of the skin (including contact dermatitis, atopic dermatitis and psoriasis), systemic inflammatory response syndrome, sepsis, an inflammatory and/or autoimmune condition of the liver (including autoimmune hepatitis, primary biliary cirrhosis, alcoholic liver disease, sclerosing cholangitis, and autoimmune cholangitis), diabetes (type I or II) and/or the complications thereof, chronic heart failure, congestive heart failure, an ischemic disease (including stroke and ischemia-reperfusion injury) or myocardial infarction and/or the complications thereof.

19. A compound as claimed in any one of claims 1 to 13, or use as claimed in claim 16, or method as claimed in claim 17, wherein the inflammatory disease is rheumatoid arthritis, chronic obstructive pulmonary disease or atopic dermatitis.

20. A compound as claimed in any one of claims 1 to 13, or use as claimed in claim 15, or the method as claimed in claim 17 for inhibition of tumour growth.