

IMPROVED PROCESSES FOR PREPARING PEPTIDE CONJUGATES AND LINKERS

The present invention relates to novel processes, compounds and intermediates useful in the preparation of certain antibody-linker-peptide conjugates.

BACKGROUND


Covalent conjugation of biochemical molecules can be employed to bring together two or more molecules to form a bioconjugate that displays the combined properties of each of the individual components. This technique has been used to increase plasma half-life and decrease immunogenicity of therapeutic agents, such as peptides. Typically, the therapeutic agent is conjugated to a macromolecular carrier directly or via a linker. Common macromolecular carriers include antibodies, albumin and synthetic polymers.

US7521425 and US8288349 describe processes for preparing compounds useful as linkers.

The reference to any art in this specification is not, and should not be taken as, an acknowledgement of any form or suggestion that the referenced art forms part of the common general knowledge.


Background of conjugation process

Antibody-drug conjugate **5** has been described in US8288349 (whose contents are hereby incorporated entirely), the production of which involves several stages. Initially, the peptide **2** and the linker **1** are prepared separately. The peptide **2** is then conjugated to the linker **1** to form the linker-peptide complex (**3**). After purification, the conjugated linker-peptide complex **3** is combined with the antibody (**4**) so as to allow the azetidinone moiety of **3** form a covalent bond with the antibody **4**, thereby resulting in an assembled peptide-linker-antibody complex; the antibody-drug conjugate **5** (scheme I). The linker peptide complex is prepared by a lengthy multi-step process requiring generation of the linker **1**, **8** and conjugation to the peptide **2** (scheme II).



5 Peptide-linker-antibody bioconjugate

Scheme (I) Preparation of peptide-linker-antibody bioconjugate 5. The peptide is shown schematically, with the available lysine side-chain indicated. The lysine residue forming the covalent bond with the linker is preferably the only unmodified lysine in the peptide, to avoid multiple species forming. The lysine may be located as the N-terminal residue, the C-terminal residue, or anywhere within the peptide chain (for example SEQ ID NOs:1 and 2). The antibody is also shown schematically, with the reactive side chain indicated. Typically, the antibody is a catalytic antibody such as an aldolase catalytic antibody comprising a reactive lysine in the antibody combining site (antigen recognition site, or CDR), as further described herein and also in US7521425, US8288349, and US8252902.



Scheme (II) Preparation of peptide-linker 3. (a) Cl_2SO , reflux. (b) $n\text{-BuLi}$, -70°C , THF. (c) 0°C , 91%. (d) H_2 , 10% Pd/C, MeOH, HCl, 40°C , 68%. (e) DIPEA, CH_2Cl_2 , rt, 83%. (f) DIC, THF, 0°C to rt, 18 h, 100%. (g) NMM, DMF, $0\text{--}5^\circ\text{C}$, 10 min.

Synthesis of acid linker 6

The original synthesis of acid 6 is shown in scheme II. Acid 9 is treated with Cl_2SO at reflux to provide crude acid chloride 10. In a separate flask, 2-azetidinone 11 is deprotonated at cryogenic temperature with $n\text{-BuLi}$ in THF to generate Li anion 12 which, without isolation, is reacted with acid chloride 10 to give intermediate 13 in 91% yield. The nitro group on 13 is then reduced using catalytic hydrogenation with 10% Pd/C in MeOH to give aniline HCl salt 14b in 68% yield. The last step involves the reaction between 14b and diglycolic anhydride (15) in CH_2Cl_2 in the presence of DIPEA to provide acid 6 in 83% yield.

During preliminary experiments to determine the scalability of scheme II, it was found that the yield for the coupling between acid chloride 10 and azetidinone 11 was not reproducible and dropped to about 40-55% when the reaction was run on about 200-g scale. There therefore exists a need to find an

alternative mechanism to generate acid **6** with the additional goal of avoiding running the process at cryogenic temperature.

Synthesis of Peptide-Linker **3**

Acid **6** and *N*-hydroxysuccinimide (**7**) undergo reaction with *N,N'*-diisopropylcarbodiimide (DIC) as coupling reagent to afford *N*-hydroxysuccinimido ester **8** in quantitative yield after urea byproduct filtration and trituration in petroleum ether (scheme 2, reaction f). Crude **8** is immediately used in the subsequent reaction with peptide **2** with *N*-methyilmorpholine (NMM) as base in DMF to provide conjugate **3** in a process that takes about 2 days. The isolation of crude **3** from the reaction mixture involves neutralization to pH = 6.0 with acetic acid, removal of DMF under vacuum, and dissolution of the resulting residue in 0.1 M ammonium acetate buffer.

Crude **3** is then subjected to chromatographic purification (0.1 M NaClO₄/MeCN buffer). The fractions with low purity (<60%) are discarded and the fractions in the 60-95% purity range re-chromatographed (0.1% TFA water/MeCN buffer). The fractions in the 80-95% purity range are re-chromatographed under the same conditions and the fractions with purity below 80% discarded. The fractions with >96% purity and no single impurity above 1.5% are pooled and lyophilized to give clean **3** in 40.4% yield (molar basis). This chromatographic purification of **3** takes about 2 days. The lyophilized fractions are then reconstituted (in other words, a full redissolution of the solid in an appropriate solvent), in a 1:1 CH₃CN/H₂O mixture to generate a homogeneous lot of intermediate **3**, in a process that takes about 2 days.


After filtration of some insoluble material, the filtrates are subjected to a second lyophilization to generate **3** in about 11.9% yield, in a process that takes about 2 days. While this lengthy and energy- and solvent-intensive process is quite suitable for the generation of small batches of material (~50 g), its implementation in the manufacture of larger quantities of **3** is relatively impractical due to the very low throughput, high scale-up costs, and lengthy process time of about 10 days. There therefore exists a need to develop a process to generate large amounts of peptide-linker conjugate **3**, capable of

producing multi-hundred gram quantities under cGMP conditions in a time- and cost-effective manner.

SUMMARY OF THE INVENTION


The present invention provides an improved process for the preparation of

- 5 peptide-linker antibody conjugate according to formula **5a**:




comprising

- (i) reacting **9a** and **11** together in the presence of 1-propanephosphonic acid anhydride (T3P) to create compound **13a**




- (ii) catalytic hydrogenation with Pd/C of compound **13a** in a THF:H₂O solution of at least about 50% THF to produce a compound of formula **14a**




- 15 (iii) combining a solution of **14a** in THF with a compound according to formula **15** in a reaction substantially free of base to produce compound **6a**


6



- (iv) reacting a compound according to the formula **6a** with a compound according to formula **19a** in THF in the presence of DCC to produce compound **20a**




- (v) combining **20a** with a ϵ -amino containing peptide **2** dissolved in an aprotic polar ^{15th} solvent to produce bioconjugate **3a**




- (vi) dissolving compound **3a** in DMSO:

- (vii) adding histidine buffer at pH between about 5.5 to about 7.5 to the solution of DMSO and **3a** of step (vi);
- (viii) Adding an antibody comprising comprises a variable light region comprising SEQ ID NO:5 and a variable heavy region comprising SEQ ID NO:6 to the solution of step (vii), so as to have a peptide:antibody molar ratio of between about 1.8:1 to about 3:1;
- (ix) Agitating the mixture formed in step (viii) at a medium speed so as to avoid foaming the reaction mixture for at least about 1hr at between about pH 5.5 and about pH7.5 and at a temperature of between about 5°C and 35°C;
- (x) Filtration of the solution from (ix) to extract the resultant peptide-linker antibody conjugate **5a**.




wherein q = 1, 2, 3, 4, or 5, X= F or Cl, m=3, 4, or 5.

- The present invention further provides an improved process for the preparation of peptide-linker according to formula **3a**




comprising


- (i) reacting **9a** and **11** together in the presence of 1-propanephosphonic acid anhydride (T3P) to create compound **13a**:




- (ii) catalytic hydrogenation with Pd/C of compound **13a** in a THF:H₂O solution of at least about 50% THF to produce a compound of formula **14a**




- (iii) combining a solution of **14a** in THF with a compound according to formula **15** in a reaction substantially free of base to produce compound **6a**



- (iv) reacting compound **6a** with a compound according to formula **19a** in THF in the presence of DCC to produce compound **20a**



- (v) optionally crystallizing compound **20a**;
- (vi) combining **20a** with a ϵ -amino containing peptide **2** dissolved in an aprotic polar 15th solvent to produce peptide-linker **3a**



(vii) optionally precipitating peptide-linker **3a** by adding a DMF solution of **3a** to MeCN;


5 wherein $q = 1, 2, 3, 4$, or 5 , $X = F$ or Cl , and $m = 3, 4$, or 5 .

In some aspects of the invention, the improved process provides a reduction in cost based on the improved yield, with an increase in the yield of peptide-linker conjugate **3a** from starting peptide **2** from about 30% to about 85%. There are other components such as energy savings and solvent savings, which are also significant, but likely have a smaller economic impact. In particular, the elimination of 2 chromatographies in the improved process results in a solvent savings estimated at 1500-2000 liters/kg product. The reduction in solvent also provides energy savings since elution solvent does not have to be removed via distillation. Significant energy saving are also associated with elimination of 2 lyophilization steps in the improved process.


In some aspects of the invention, the improved process provides a reduction in cycle time by avoiding chromatography and lyophilization after the synthesis of the peptide-linker conjugate **3a**. For the cycle time, the new process delivers material in about 2 days, compared with the existing cycle time of about 10 days. Therefore, the new process delivers a reduction in cycle time of 80% or greater.

In some aspects of the invention, the peptide **2** comprises the specific sequence of SEQ ID NO:1, wherein AcK is acyl lysine, to form peptide **2a**.

SEQ ID NO:1: C(O)CH₃-Q(AcK)YQPLDE(AcK)DKTLYDQFMLQQG-NH₂ **2a**



In some aspects of the invention, the peptide-linker conjugate **3b** comprises the formula:



It will be appreciated that other peptides and non-peptidic agents may usefully be conjugated to the linkers of the invention.

The present invention provides an optimized and scalable process to manufacture peptide-linker conjugate **3a** that avoids the chromatographic purification and lyophilization that is typically required for the isolation of this type of compound and results in significantly higher yields.

The invention provides an operationally simple protocol that couples the peptide to the linker in DMF, optionally followed by precipitation with MeCN as anti-solvent and filtration to give material that meets specifications for use in clinical batches. A scalable synthesis of the linker is also described which features the *N*-acylation of 2-azetidinone **11** promoted by 1-propanephosphonic acid anhydride (T3P) under mild conditions in the first step. This new protocol provides reproducible yields, does not resort to chromatographic purification, and avoids both the need for performing the reaction at cryogenic temperatures and the use of *n*-BuLi as was required in the original route.

The number of operations during the second step of the synthesis (**13a** reduction to **14a**) has been simplified by telescoping compound **14a** into the next

step (reaction with diglycolic anhydride to form acid **6a**), thus avoiding an additional isolation.


Furthermore, an efficient activation method for acid **6a** has been developed by means of the corresponding ester of formula **20a**, which display an
5 excellent balance between stability for isolation purposes and reactivity during the conjugation with peptide **2**.

The present invention further provides an improved and efficient conjugation protocol for the coupling of a linker and a peptide as part of a program to manufacture a peptide-linker-antibody bioconjugate for the treatment
10 of cancer. This novel approach allows for the isolation of peptide-linker conjugate **3a** via direct precipitation from a MeCN/DMF mixture followed by filtration. The resource-intensive isolation protocol performed originally, and common to this class of compounds, involving extensive chromatographic purification and lyophilization has thus been avoided. One of the major improvements is the
15 dramatic increase in yield for this step, which has gone up from 12% to 83%. Further advantages are the reduction in solvent and energy consumption, which substantially lowers the cost per gram and turns the process into a greener alternative. An extensive screen of all the key parameters resulted in reaction conditions that provided material of satisfactory quality to meet stringent
20 specifications in terms of purity and residual solvent content. This method represents an innovative approach toward the isolation of this type of material that departs from the traditional chromatographic purification.


The present invention also provides an optimized route to the activated linker that eliminates cryogenic conditions and the use of *n*-BuLi, avoids the
25 isolation of one of the intermediates through telescoping, and further provides a novel pentafluorophenol ester as a suitable substrate with the desired balance between stability for isolation purposes and reactivity in the final conjugation step with the peptide therapeutic agent.

All these improvements have translated into the manufacture of significant
30 quantities of peptide-linker conjugate **3a** under cGMP conditions.

In some aspects, the invention provides for a new process for preparing compound **13a**, comprising reacting **9a** and **11** together in the presence of 1-propanephosphonic acid anhydride (T3P) to create compound **13a**



- 5 wherein q is 1, 2, 3, 4, or 5. Where q is 2, the reaction may comprise **9** and **11** in the presence of T3P to produce **13**:



- The improved process of the invention provides significant advantages over the original process, as 2-azetidinone **11** is typically quite unreactive toward amide bond formation. In addition, this improved process is reproducible at least on several hundred gram-scale. In some aspects, the invention provides a compound prepared according to such a process.

- The reaction may be carried out in a 1st solvent selected from the group consisting of tetrahydrofuran (THF), 2-methyltetrahydrofuran, NMP, N,N-dimethylformamide (DMF), N,N-dimethylacetamide (DMAc), 1-methyl-2-pyrrolidinone, ethyl acetate (EtOAc), and acetonitrile (MeCN). Particularly suitable 1st solvents include DMF, EtOAc, and MeCN. In some aspects, the 1st solvent is selected between EtOAc and MeCN. In some aspects, the 1st solvent is DMF. In some aspects, the 1st solvent may be MeCN.

- In some aspects, the reaction is carried out in the presence of a 1st base. The 1st base may be selected from the group consisting of trimethylamine, triethylamine, tributylamine, DIPEA, pyridine, DBU, DABCO, 2,3-lutidine, 2,4-lutidine, 2,5-lutidine, 2,6-lutidine, 3,4-lutidine, 3,5-lutidine. Particularly suitable 1st bases include DIPEA, triethylamine and pyridine. In some aspects, the 1st base is DIPEA.

The 1st base may be present in an amount relative to acid **9a** in a range whose lower limit is selected from the group consisting of about 0.5, 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, and 5; and whose upper limit is selected from the group consisting of about 2.5, 3, 3.5, 4, 4.5, 5, 6, 7, 8, 9 and 10. In some aspects, the 1st base is present in an amount between about 1 and about 10 equivalents of acid **9**. In some aspects, the 1st base is present in an amount between about 2 and about 5 equivalents of acid **9a**. In some aspects, the 1st base is present in an amount between about 2 and about 4 equivalents of acid **9a**. In some aspects, the 1st base is present in an amount between about 2.5 and about 3.5 equivalents of acid **9a**. In some aspects, the 1st base is present in an amount of about 3 equivalents of acid **9a**. In some aspects, the 1st base is DIPEA and is present in an amount of about 3 equivalents of acid **9a**.

The 2-azetidinone **11** may be present in an amount relative to acid **9a** in a range whose lower limit is selected from the group consisting of about 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1, 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9, 2, 2.5, 3, 3.5, 4, 4.5, and 5; and whose upper limit is selected from the group consisting of about 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9, 2, 2.5, 3, 3.5, 4, 4.5, 5, 6, 7, 8, 9 and 10. In some aspects, the 2-azetidinone **11** is present in an amount between about 0.1 and about 10 equivalents of acid **9a**. In some aspects, the 2-azetidinone **11** is present in an amount between about 0.5 and about 3 equivalents of acid **9a**. In some aspects, the 2-azetidinone **11** is present in an amount between about 0.5 and about 5 equivalents of acid **9a**. In some aspects, the 2-azetidinone **11** is present in an amount between about 0.5 and about 1.5 equivalents of acid **9a**. In some aspects, the 2-azetidinone **11** is present in an amount between about 1.0 and about 1.5 equivalents of acid **9a**. In some aspects, the 2-azetidinone **11** is present in an amount of about 1.2 equivalents of acid **9a**.

The T3P may be present in an amount relative to acid **9a** in a range whose lower limit is selected from the group consisting of about 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1, 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9, 2, 2.5, 3, 3.5, 4, 4.5, and 5; and whose upper limit is selected from the group consisting of about 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9, 2, 2.5, 3, 3.5, 4, 4.5, 5, 6, 7, 8, 9 and 10. In some

aspects, the T3P is present in an amount between about 0.1 and about 10 equivalents of acid **9a**. In some aspects, the T3P is present in an amount between about 0.5 and about 3 equivalents of acid **9a**. In some aspects, the T3P is present in an amount between about 0.5 and about 5 equivalents of acid **9a**. In some aspects, the T3P is present in an amount between about 0.5 and about 1.5 equivalents of acid **9a**. In some aspects, the T3P is present in an amount between about 1.0 and about 1.5 equivalents of acid **9a**. In some aspects, the T3P is present in an amount of about 1.3 equivalents of acid **9a**.

In some aspects, the T3P may be provided in a 2nd solvent selected from the group consisting of EtOAc, N,N-dimethylformamide and butyl acetate. In some aspects, the T3P may be present in an amount between 1% and 99% in the solution, provided that molar ratio between T3P and substrate is kept between about 0.1:1 and about 10:1. The T3P may be provided in the second solvent in a range whose lower limit is selected from the group consisting of about 1%, about 5%, about 10%, about 15%, about 20%, about 25%, about 30%, about 35%, about 40%, about 45% and about 50%; and whose upper limit is selected from the group consisting of about 50%, about 55%, about 60%, about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, about 95% and about 99%. In some aspects, the T3P is provided in an about 50% solvent solution. The T3P may be provided in an about 50% EtOAc solution.

Accordingly, in some aspects, the addition of T3P to a mixture of compounds **9a** and **11** in DMF and pyridine as base gave desired product in about 20% to about 45% yield. In some aspects, the ratio of reaction constituents is about 3 equiv of DIPEA, about 1.2 equiv of **11** and about 1.2 equiv of T3P (in about 50% EtOAc) in MeCN to about 1 equivalent of acid **9a**. The advantage of these proportions is to provide consistent yields of at least about 20%. In some aspects, this combination yields at least about 25%. In some aspects, this combination yields at least about 30%. In some aspects, this combination yields at least about 35%. In some aspects, this combination yields at least about 40%. In some aspects, these proportions provide about 40% yield.

In some aspects, about 5 equivalents of DIPEA, about 1.5 equiv of **11** and about 1.5 equiv of T3P are used, in MeCN. The advantage of these proportions is to provide consistent yields of about 55-60%, in particular when the reaction was performed at RT (i.e. about 15°C to about 25°C, preferably 18°C to about 22°C, and most preferably about 20°C) for about 2 h.

Some of the advantages of this method are the operational simplicity (slow addition of a 50% T3P solution in EtOAc to a mixture containing the **2** coupling partners and base in MeCN), mild reaction conditions, and yield reproducibility.

In some aspects, the reaction may be performed at RT (e.g. about 20°C) for about 18 hr. In some aspects, the temperature of the reaction may be operated at a range whose lowest value is selected from the group consisting of about 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 12, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, and 30°C, and whose upper value is selected from the group consisting of about 25, 26, 27, 28, 29, 30, 32, 35, 36, 37, 40, 45, and 50°C. In some aspects, the temperature range is between about 0°C and about 50°C. In some aspects, the temperature range is between about 10°C and about 30°C. In some aspects, the temperature is room temperature. In some aspects, the temperature is 20°C.


Upon reaction completion the MeCN may be removed under vacuum and the residue redissolved in a 3rd solvent comprising a C₁-C₄ alkyl acetate. The C₁-C₄ alkyl acetate may be selected from the group consisting of *i*-propyl acetate, methyl acetate, ethyl acetate, *n*-propyl acetate, and *n*-butyl acetate. In some aspects, the residue may be redissolved in *i*-propyl acetate

The organic phase may then be washed with aqueous citric acid (other acids, such as acetic acid may also be used), and, after a solvent switch from the 3rd solvent to a 4th solvent selected from the group consisting of 2-propanol, 1-propanol, 1-butanol, 2-butanol, and tert-butanol, intermediate **13a** precipitates from solution in high purity (>98%). In some aspects, the 3rd to 4th solvent switch may be from *i*-propyl acetate to 2-propanol.

In some aspects, it can be advantageous to treat the intermediate **13a** with activated carbon to remove some color and trace impurities present before using

this material in the next nitro-reduction step (i.e. the conversion of intermediate **13a** to salt **14a**). The activated carbon treatment may be carried out in a 5th solvent, such as EtOAc. Other suitable 5th solvents are tetrahydrofuran, 2-methyltetrahydrofuran, methanol, ethanol, methyl acetate, and i-propyl acetate.


- 5 The activated carbon may be selected from the group consisting of, SX-Plus, Darco® S-51HF, E Supra USP, SX-Ultra, CASP, Darco® G-60 and CGSP (all available from Norit®). In some aspects of the invention, the activated carbon is Darco® G-60, which provides the lowest amount of the trace impurity **16a** (where $q = 1, 2, 3, 4, \text{ or } 5$), or **16** where $q = 2$.




In some aspects, the purity of acid **9a** is at least about 85%. In some aspects, the purity of acid **9a** is at least about 90%. In some aspects, the purity of acid **9a** is at least about 95%. In some aspects, the purity of acid **9a** is at least about 96%. In some aspects, the purity of acid **9a** is at least about 97%. In some aspects, the purity of acid **9a** is at least about 98%. In some aspects, the purity of acid **9a** is at least about 99%. The purity of the acid **9a** can have a great impact on the reproducibility of the reaction. When **9a** of purity less than 85% is used, 2 carbon treatments may be necessary to remove highly colored impurities, which causes the yield of coupling product **13a** to drop to 48%.

- 20 A further benefit of carbon treatment was that the level of impurity **16a**, resulting from the opening of the azetidinone ring by the newly formed amino group in the subsequent nitro reduction step, may be kept below acceptable levels. When the carbon treatment used Darco® G-60 in EtOAc, the level of impurity of **16a** can be about 0.3%.

- 25 In some aspects, the invention provides for a new process for preparing compound **14a**, comprising catalytic hydrogenation with Pd/C of compound **13a** in a THF:H₂O solution of at least about 50% THF to produce a compound of formula **14a**:




wherein $q = 1, 2, 3, 4$, or 5 . Where $q=2$, **13a** may be **13** and **14a** may be **14**:



In some aspects, the THF:H₂O solution comprises at least about 60% THF. In some aspects, the THF:H₂O solution comprises at least about 70% THF. In some aspects, the THF:H₂O solution comprises at least about 80% THF. In some aspects, the THF:H₂O solution comprises at least about 90% THF. In some aspects, the THF:H₂O solution comprises about 90% THF. In some aspects, the THF:H₂O solution comprises up to about 99% THF. In some aspects, the THF:H₂O solution comprises up to about 95% THF. In some aspects, the ratio of THF:H₂O is between about 8-10:1. In some aspects, the ratio of THF:H₂O is about 9:1.

In some aspects, the 5% or 10% Pd/C is employed and used in 1-100% weight/weight ratio with respect to **13a**. In some aspects, the catalytic hydrogenation takes place at a pressure of between about 1-50 psig. In some aspects, the catalytic hydrogenation takes place at about 15 psig. In some aspects, **13a** is subjected to charcoal treatment substantially as described herein before the reaction. In some aspects, the catalyst may be filtered off and the filtrates treated with about 1 equiv of a strong acid (such as HCl (e.g. 12M), or similar, as discussed above) to generate and isolate the corresponding (HCl) salt.

Previously, the nitro-reduction step of compound **13a** to compound **14a** was carried out in methanol (MeOH) and provided approximately 68% yield of **14a**. However, it has been found that using methanol results in a considerable amount of the impurity **17a** (**17**, where $q=2$) (between about 1 and about 5%):



There therefore exists a need to improve the process of converting compound **13a** to compound **14a**. One potential solution to avoid the generation of **17a** would be to use dioxane to prevent its formation. However, dioxane is a known carcinogen, and there therefore exists a need to find a safer alternative.

A proposed solution comprises the use of THF in conjunction with catalytic hydrogenation at a pressure range of between 1-50 psig (pound-force per square inch gauge). 15 psig with 10% Pd/C (10 wt%) in THF gave a fast reaction (<1 h) but the formation of variable levels of **16a** were observed (1–5%) (other amounts of Pd metal by weight of the C support may also be useful; from 1-100%). Several additives were tested to prevent or reduce the formation of **16a** (HCl, HOAc, H₂O, all at 10%) but compound **16a** was still detected (1.8%, 1.9% and 1.6% respectively), as well as slower conversion.


The presence of water was a concern due to the potential opening of the azetidinone ring, but running the reaction in THF/H₂O 9:1 (vol/vol) only caused a modest increase in impurity level.

The best results were obtained when starting material **13a** was subjected to a charcoal treatment as described above, which kept the amount of impurity **16a** at low levels (<0.3%) in the absence of additives.


Once full conversion of **13a** to **14a** (in some aspects, **13** to aniline **14**) was attained, the catalyst (10% Pd/C) may be removed by filtration and the filtrates treated with 1 equiv of 12 M HCl to generate and isolate the corresponding HCl salt (which is the preferred form for long-term storage, as anilines have a tendency to undergo oxidation, giving rise to highly colored products). Other strong acids are suitable, such as sulphuric and nitric. The selection of the concentration and acid is largely determined so as to minimize volume and material loss.

In some aspects, the invention provides a process for preparing compound **6a**, comprising combining a solution of **14a** in a 6th solvent with a

compound according to formula **15** in a reaction substantially free of base to produce compound **6a**




wherein $q = 1, 2, 3, 4$ or 5 ;



5 from a solution of compound **14a** in THF,
Compound **15** may be added as a solution or as a solid. The 6th solvent may be selected from the group consisting of THF, C1-C4 alkyl acetate, toluene, chloroform, methyl THF. In some aspects, the 6th solvent is THF. In some aspects, the reaction is conducted in the absence of H₂O.

10 Where $q=2$, compound **6** may be prepared as described above using compound **14**:



In part, some aspects of the invention are based on the surprising discovery that treating the THF filtrates containing **14a** with about 1 equiv of diglycolic anhydride (**15**) in the absence of base led to the efficient formation of **6a**. Compounds of the formula **14a**, (such as aniline, **14**), are only poorly nucleophilic, and require typically deprotonation with a base to increase their reactivity. Moreover, the HCL salt of compound **14**, compound **14b**, is solid, and can be stored for long periods of time, whereas a solution of **14** or **14a** generally requires immediate usage. However, these disadvantages were outweighed by the surprising advantage of the present process of reduction in overall cycle time and cost of the reaction and purification.

In some aspects, compound **14a** is reacted with about 1 equivalent of compound **15**. In some aspects, compound **14a** is reacted with at least about 1 equivalent of compound **15**. In some aspects, compound **14a** is reacted with between about 1 and about 10 equivalents of compound **15**.

5 As a result, after reaction completion to form **6a**, the present invention also provides for an additional process step; wherein the solution of compound **6a** is distilled under vacuum to remove THF, following which, a 7th solvent comprising C₁-C₄ alkyl acetate is added to attain a solvent composition of between about 25%THF/75% C₁-C₄ alkyl acetate and about 100% C₁-C₄ alkyl acetate. This may
10 represent a total solvent volume of between about 5 and about 15 mls of solvent per gram of **6a**. In some aspects, the 7th solvent may comprise one of methyl acetate, ethyl acetate, *i*-propyl acetate, *n*-propyl acetate, *n*-butyl acetate. In some aspects, the 7th solvent is either methyl acetate or *i*-propyl acetate. In some aspects, the 7th solvent is methyl acetate. In some aspects, the 7th solvent is *i*-
15 propyl acetate.

Isolation in this manner results in a yield of 85–90% and typical residual solvent levels of 0.2 and 0.5 wt% for THF and *i*-PrOAc, respectively.

No change in yield or quality is seen when a final solvent ratio of between 25:75 6th solvent: 7th solvent (THF:*i*-PrOAc) and 100% 7th solvent (*i*-PrOAc) is
20 employed. The telescoping of the THF solution of free base allows for a much simpler process and saves a considerable amount of time.

The reaction of **14a** with **15** to form **6a** is very fast (for example, <15 min in 60 volumes of solvent) and, after concentration of the THF solution, the addition of the 7th solvent (e.g. *i*-propyl acetate) and cooling causes acid **6a** to
25 precipitate from solution in excellent yield (85–90%) and purity (>98%). This isolation method, while high yielding and fast, can result in relatively high residual solvent levels (e.g. 0.5–1% THF, 1-2% *i*-propyl acetate). Consequently, it was desirable to develop an alternative solvent exchange method to better control the final crystallization.

30 Crystallization of **6a** from the 7th solvent comprising a C₁-C₄ alkyl acetate (e.g. *i*-PrOAc) at a high temperature (e.g. 70°C) causes product degradation

(high temperature was needed to dissolve the material in a reasonable volume of solvent), whereas dissolution in acetone followed by solvent displacement with *i*-PrOAc and crystallization provided material that contained 0.5% residual *i*-PrOAc. Therefore, there exists a need to develop an optimized crystallization protocol, in order to further reduce the residual amount of solvent, (<0.25 wt%).

Accordingly, in some aspects, the invention provides for a further process step to crystallize acid **6a**, comprising:

- (i) dissolving acid **6a** in an 8th solvent comprising THF;
- (ii) treating with activated carbon and then filtering off said activated carbon;
- 10 (iii) concentrating the acid **6a** in THF solution to between about 2 and about 20 vol;
- (iv) adding between about 1 and about 50 vol of a 1st alcohol consisting of C₁-C₆ alkyl alcohol;
- (v) concentrating the solution of acid **6a** in THF and 2-propanol to between
15 about 2 and about 50 vol;
- (vi) cooling the concentrated solution of acid **6** to between about -25°C and about 10°C.

In some aspects, the 8th solvent is selected from the group consisting of THF, C₁-C₄ alkyl acetate, toluene, and acetonitrile. In some aspects the 8th
20 solvent is THF. In some aspects, **6a** may be dissolved in the 8th solvent at a temperature of between about 10°C and about 67°C. The 8th solvent may be present in an amount of between about 10 to about 50 volumes, and may be about 35 volumes.

In some aspects of the invention, such as where **6a** is provided in a
25 solution containing C₁-C₄ alkyl acetate, the solute may be removed, and **6a** redissolved in the 8th solvent, which may be THF. In some aspects of the invention, acid **6a** may be dissolved in between about 20 to about 50 volumes of 8th solvent, and in some aspects about 35 vol. In some aspects, acid **6a** may be dissolved in the 8th solvent at between about 15°C and about 60°C. In some
30 aspects, the temperature is between 20°C and about 40°C. In some aspects, the temperature is about 30°C.

After being dissolved in the 8th solvent at step (i), the acid **6a** may be subjected to an additional step; (ii): wherein the acid **6a** is treated with activated carbon. The carbon treatment provides the advantage that removal of colour appears to lead to more predictable crystallization behavior

5 The activated carbon may be selected from the group consisting of CGSP, SX-Plus, Darco S-51HF, E Supra USP, SX-Ultra, CASP, Darco G-60 and Darco-KBB. In some aspects of the invention, the activated carbon is Darco-KBB. In some aspects, the activated carbon is added to a final amount of at least about 5% by weight of **6a**. In some aspects, the activated carbon is added to a final
10 amount of at least about 10% by weight of **6a**. In some aspects, the activated carbon is added to a final amount of at least about 15% by weight of **6a**. In some aspects, the activated carbon is added to a final amount of about 20% by weight of **6a**.

 The acid **6a** / 8th solvent solution may be treated with activated carbon for
15 at least about 15 mins, and in some aspects, at least about 1 hr, and may be between about 1 and about 24 hrs.

 In some aspects, the activated carbon is filtered off (for example, using a filter aid, such as celite). In some aspects, the solution (filtrates) may then be concentrated (for example, under reduced pressure, such as between about 0
20 and about 1 atmospheres pressure) at step (iii) to between about 2 and about 20 volumes. In some aspects, the filtrates may then be concentrated to between about 5 and about 15 volumes. In some aspects, the filtrates may then be concentrated to about 10 volumes.

 In some aspects, between about 1 and about 50 volumes of 1st alcohol
25 may be added to the solution at step (iv). In some aspects, excess 1st alcohol may be added to initiate precipitation of **6a** from the 8th solvent. In some aspects, between about 1 and about 30 volumes of 1st alcohol may be added. In some aspects, between about 1 and about 20 volumes of 1st alcohol may be added. In some aspects, between about 5 and about 50 volumes of 1st alcohol may be
30 added. In some aspects, between about 5 and about 30 volumes of 1st alcohol may be added. In some aspects, between about 12 and about 16 volumes of 1st

alcohol may be added. In some aspects, about 14 volumes of 1st alcohol may be added.

The 1st alcohol may be selected from the group consisting of C₁-C₆ alkyl alcohols, including methanol, ethanol, 1-propanol, 2-propanol, 1-butanol, 2-butanol, 1-pentanol, 2-pentanol, 3-pentanol, 1-hexanol, 2-hexanol, 3-hexanol, 3-methyl-1-butanol, 2-methyl-1-butanol, 2,2-dimethyl-1-propanol, 3-methyl-2-butanol, and 2-methyl-2-butanol. In some aspects, the 1st alcohol may be a primary or secondary C₁-C₆ alkyl alcohol. In some aspects, the 1st alcohol may be a secondary C₁-C₆ alkyl alcohol. In some aspects, the 1st alcohol may be 2-propanol.

In some aspects, 1st alcohol may be added to the concentrated solution of acid **6** at step (iv) at a ratio of between about 0.5 to about 10: about 1. In some aspects, the ratio is between about 0.5:1 and about 5:1. In some aspects, excess 1st alcohol is added. In some aspects, the ratio is between about 1:1 and about 3:1. In some aspects, the ratio is between about 1:1 and about 2:1. In some aspects, the ratio is between about 1.2:1 and about 6:1. In some aspects, the ratio is between about 1:3 and about 1.5:1. In some aspects, the ratio is about 1.4:1.

In some aspects, the solution of acid **6a** in 8th solvent and 1st alcohol from step (iv) may then be concentrated to between about 1 and about 50 vol. In some aspects, the solution of acid **6a** in 8th solvent and 1st alcohol may be concentrated to between about 1 and about 30 vol. In some aspects, the solution of acid **6a** in 8th solvent and 1st alcohol may be concentrated to between about 5 and about 20 vol. In some aspects, the solution of acid **6a** in 8th solvent and 1st alcohol may be concentrated to between about 5 and about 20 vol. In some aspects, the solution of acid **6a** in 8th solvent and 1st alcohol may be concentrated to between about 5 and about 15 vol. In some aspects, the solution of acid **6a** in 8th solvent and 1st alcohol may be concentrated to about 10 vol.

In some aspects, the solution of acid **6a** in 8th solvent and 1st alcohol from step (iv) may be concentrated, ideally under reduced pressure. In some aspects,

atmospheric pressure may be used. In some aspects, between a full vacuum and 1 atmosphere may be used to concentrate **6a**.

In some aspects, the solution from step (v) may then be cooled so as to enable acid **6a** to crystallize. In some aspects, the solution from step (v) may be cooled to between about -25 °C to about 10°C. In some aspects, the solution from step (v) may be cooled to between about -20 °C to about 10°C. In some aspects, the solution from step (v) may be cooled to between about -20 °C to about 5°C. In some aspects, the solution from step (v) may be cooled to between about -5 °C to about 10°C. In some aspects, the solution from step (v) may be cooled to between about -5 °C to about 5°C. In some aspects, the solution from step (v) may be cooled to between about -5 °C to about 0°C. In some aspects, the solution from step (v) may be cooled to between about -1 °C to about 4°C. In some aspects, the solution from step (v) may be cooled to between about -0 °C to about 4°C. In some aspects, the solution from step (v) may be cooled to between about 0 °C to about 5°C.


In some aspects, after filtering the carbon off, the mixture may be concentrated (for example, by distillation at reduced pressure (0-1 atm) to 10 vol, to which about 14 vol of 1st alcohol may be added. The solution may then be further concentrated to about 10 vol, and then cooled to between about -20°C to about 10°C, and preferably between about 0 and about 5°C. Such a protocol affords acid **6a** in 80% recovery and >99% chemical purity. This especially advantageous embodiment of the invention delivers significantly smaller particles (~30 µ) than the original THF/*i*-PrOAc crystallization (100–200 µ) and the smaller particle size may explain why less solvent was trapped in the crystals.

In some aspects, the invention is based on the successful identification of activated esters that may be easily prepared in high yield and stored over prolonged periods of time before being conjugated to peptides and proteins and the like, and methods of making said activated esters.

A need was identified to develop a process for the conjugation of peptide **2** to acid **6a** that allowed for the isolation of peptide-linker conjugate **3a** in high yield and purity without resorting to the time and resource expensive chromatographic

purifications and lyophilizations present in the original synthesis. It is further desirable that the improved process provides a chemical purity of $\geq 95\%$ with no single impurity above 2% and residual solvent content below 0.25% (wt/wt) for each individual solvent.

5 Accordingly, in some aspects, the invention provides for compounds and intermediates of the formulae:




10 wherein q is 1, 2, 3, 4, or 5, n = 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, and X is any halogen, and m = 3, 4 or 5. X may be F or Cl, and preferably is F. M may be 4 or 5, and preferably is 5. In some aspects, n = 1-10. In some aspects, n = 1, 2, 3, 4, 5, or 6. In some aspects, q = 1, 2, or 3. In some aspects, q=2.


15 In some aspects, the invention provides a method of generating activated esters of formula **20** and **20a**, **20b**, **20c**, **20d**, and **20e**. In some aspects, the invention provides a method of generating activated esters of formula **20**, **20a**, **20c**, and **20d**.

The preparation of the acid chloride derivative of **6a** with either Cl₂SO or (COCl)₂ or activation with reagents such as CDI or chloroformate was ruled out due to their high reactivity, as the resulting intermediates would readily cyclize to give compounds such as **18a**, for example, morpholine-3,5-dione **18**, where q = 2.


26



In part, the invention is based on the discovery that when acid **6a** (such as **6**) and compounds such as tri-, tetra-, and penta-halo substituted phenol compounds of formula **19a** (such as pentafluorophenol **19**) undergo reaction in the presence of DCC in a 10th solvent such as THF, esters **20a** are obtained in 80–85% yield (such as **20**). Other suitable 10th solvents in place of THF include dichloromethane, 2-methyltetrahydrofuran, N,N-dimethylformamide, and N,N-dimethylacetamide.



Accordingly, in some aspects the invention provides a process where a compound according to formula **6b** is reacted with a compound according to formula **19a** in a 10th solvent in the presence of DCC to form a compound according to formula **20b**.



wherein q = 1, 2, 3, 4, or 5, n = 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29 or 30, X = F or Cl, and m = 3, 4, or 5.

In some aspects, the ratio of **19a** relative to **6b** is at least about 1. In some aspects, the ratio of **19a:6b** is about 1:1. In some aspects, excess **19a** is added to **6b**. In some aspects, the ratio of **19a:6b** is about 2:1.

In some aspects, the reaction takes place in a 10th solvent selected from the group consisting of THF, dichloromethane, 2-methyltetrahydrofuran, N,N-dimethylformamide, and N,N-dimethylacetamide. In some aspects, the 10th solvent is THF.

5 In some aspects, the DCC is added to the solution of **6b** and **19a** at between about -10 to about 10°C, preferably between about 0 to about 10°C, and more preferably between about 3 and about 5°C. After a mixing time of at least about 1 min and preferably at least about 10 mins, the reaction solution may be heated to about RT (for example, between about 15 and about 25°C, preferably
10 between about 18 and about 22°C, most preferably about 20°C), and stirred for at least 4, preferably at least 6, more preferably at least 12 hrs, most preferably for about 18 hrs.

After reaction, the solution may be filtered (to remove byproducts, such as dicyclohexylurea), and washed with an 11th solvent selected from the group
15 consisting of THF, dichloromethane, 2-methyltetrahydrofuran, N,N-dimethylformamide, and N,N-dimethylacetamide. In some aspects, the 11th solvent is THF. Following this, the solids may be resuspended in the 11th solvent, and mixed with acetone (in some aspects, the acetone may be substitute with C₁-C₄ alkylacetate, toluene, MTBE, or acetonitrile). The ratio of 11th solvent to
20 acetone may be between about 2:1 to about 1:2, and is preferably about 1:1. IN some aspects, it is desirable to use excess acetone.

The solution may be cooled to at least 15°C, and preferably at least about 12°C, and most preferably at least about 10°C, and preferably up to about 0°C. Optionally, the solution may then be stirred for at least 10 mins, and preferably at
25 least 30, at least 60 and at least 80 mins. The solution may then be filtered and the solid washed with acetone (or similar, as discussed above).

The filtrates may be resuspended in a 12th solvent selected from the group consisting of THF, dichloromethane, 2-methyltetrahydrofuran, N,N-dimethylformamide, and N,N-dimethylacetamide. In some aspects, the 12th
30 solvent is THF. The 12th solvent may be mixed with a 2nd alcohol (which may be selected from the same group as the 1st alcohol, excepting methanol, and may

be 2-propanol) to a ratio of between about 2:1 to about 1:2, and preferably about 1:1 to create a slurry. In some aspects, the ration is at least about 1:1 so as to provide an excess of 2nd alcohol.

5 The slurry may be stirred for at least 1 hr, at least 6 hr, at least 12 hr. The slurry may be stirred at between about 4 and about 30°C, or between about 12 and about 25°C or between about 18 and about 22°C, or about 20, or at RT.

The slurry may be filtered, washed with a 3rd alcohol (which may be selected form the same group as the 2nd alcohol, and may be 2-propanol), and dried. The obtained solid may be dried under vacuum. The obtained solid may be
10 dried at between RT and about 50°C, and preferably about 40°C. The solid may be dried for at least about 1, at least about 4, at least about 12 and preferably at least about 16 hrs.

Compound **20** may be similarly prepared by the advantageous processes of the invention, using compounds such as **6** and **19** as starting material.
15 Compound **20a** may be similarly prepared by the advantageous processes of the invention, using compounds such as **6a** and **19a** as starting material. Compound **20c** may be similarly prepared by the advantageous processes of the invention, using compounds such as **6** and **19a** as starting material. Compound **20d** may be similarly prepared by the advantageous processes of the invention, using
20 compounds such as **6a** and **19** as starting material. Compound **20e** may be similarly prepared by the advantageous processes of the invention, using compounds such as **6b** and **19** as starting material.

In some aspects, the addition reaction compound **6** and compound **19** in the presence of DCC in THF produces compound **20** at a yield of at least 80%,
25 and in some cases, at least 85%.

In some aspects acid **6b** and **19a** are reacted together at a temperature of between about -25°C and about 50°C. In some aspects acid **6b** and **19a** are reacted together at a temperature range whose lower limit is selected from the group consisting of about -25, -20, -15, -10, -5, -1, 0, 1, 4, 5, 10, 15, 16, 17, 18,
30 19, 20, 21, and 22°C and whose upper limit is selected from the group consisting of about -1, 0, 1, 4, 5, 10, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 30, 35, 40,

45, and 50°C. In some aspects acid **6b** and **19a** are reacted together at a temperature of between about -0°C and about 20°C. In some aspects, the reaction is at about -15°C.

In some aspects acid **6b** and **19a** are reacted together for at least 1 hr. In
5 some aspects acid **6b** and **19a** are reacted together for at least 2 hrs. In some aspects acid **6b** and **19a** are reacted together for at least 3 hrs. In some aspects acid **6b** and **19a** are reacted together for at least 4 hrs. In some aspects acid **6b** and **19a** are reacted together for at least 6 hrs. In some aspects acid **6b** and **19a** are reacted together for at least 8 hrs. In some aspects acid **6b** and **19a** are
10 reacted together for at least 12 hrs. In some aspects linker **6b** and **19a** are reacted together for at least 18 hrs. In some aspects linker **6b** and **19a** are reacted together for about 24 hrs.

In some aspects, acid **6b** and **19a** are reacted together at between about 1:1 and about 1:10 ratio.


15 In some aspects, a coupling agent may be added. The role of this coupling agent is the activation of acid **6b** to produce a more reactive intermediate that is capable of reacting with phenol **19a** to afford the desired ester. In the absence of coupling agent, no ester bond formation occurs. The coupling agent may be selected from the group consisting of DCC, CDI, CDMT, DCMT, DIC, DPPA,
20 EDC, HATU, HBTU, PyBOP, PyBroP, PyCloP, TBTU, and T3P. In some aspects, the coupling agent is DCC.

Esters **20**, **20a**, **20b**, **20c**, **20d** and **20e** may be isolated via chromatographic purification. In a further aspect of the invention, esters **20**, **20a**, **20b**, **20c**, **20d** and **20e** may be precipitated from a 14th solvent such as
25 isopropanol (as exemplified in Example 11). Other suitable 14th solvents include ethanol and butanol.

Ester **20** is a fine, white solid that is stable for months at room temperature in contact with air, without the need for any special storage conditions, and that shows no detectable amounts of cyclic morpholine-3,5-dione **18**. In addition, it
30 displays excellent reactivity in the final conjugation step to form **1** (vide infra).

In some aspects, the invention provides an improved process for generating compound **3b** comprising

- (i) mixing 1 equiv of peptide **2** in an aprotic, polar 15th solvent, and
- (ii) combining excess **20b** with peptide **2** to produce **3b**



wherein $n = 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29$ or 30 , $q = 1, 2, 3, 4$, or 5 , $X = F$ or Cl , and $m = 3, 4$, or 5 .

10 In some aspects, the aprotic, polar 15th solvent is selected from the group consisting of DMF, DMAc, DMSO or NMP. In some aspects, the aprotic polar solvent is DMF.

15 In some aspects, peptide **2** can be either a salt such as trifluoroacetate salt of **2** or a free base form. Where peptide **2** is a salt, then step (i) is carried out in a 2nd base. Where peptide **2** is provided as a free base form, then the 2nd base is not required.

20 In some aspects, peptide **2** is dissolved in between about 5 and about 50 volumes of the aprotic polar 15th solvent. In some aspects, peptide **2** is dissolved in between about 10 and about 20 volumes of the aprotic polar 15th solvent. In some aspects, peptide **2** is dissolved in about 15 volumes of the aprotic polar 15th solvent.

In some aspects of the invention, the 2nd base is selected from the group consisting of NMM, TEA, DIPEA, pyridine, DABCO, DBU, and lutidine. In some aspects, the 2nd base may be TEA, DIPEA or pyridine.

The 2nd base may be present in an amount of between about 1 and about 5 equiv. In some aspects, the 2nd base is present in an amount of about 1 to about 5 equiv. In some aspects, the 2nd base is present in an amount of about 1 to about 3 equiv. In some aspects, the 2nd base is present in an amount of about 1 to about 2 equiv. In some aspects, the 2nd base is present in an amount of about 1 to about 1.5 equiv. In some aspects, the 2nd base is present in an amount of about 1.2 equiv.

However, both peptide **2** and peptide-linker conjugate **3** tend to form thick gels in aprotic, polar 15th solvents such as DMF, DMAc, and DMSO after a few hours, either in the presence or absence of water. The solution to this problem is to react peptide **2** with **20b** within about 18 hrs of dissolving peptide **2** in the aprotic polar 15th solvent; preferably within about 12 hrs, more preferably within about 8 hrs, more preferably within about 6 hrs more preferably within about 4 hrs, more preferably within about 2 hrs, more preferably within about 1 hr. In addition, the problem of gelification is made less acute by use of at least, 10, and favourably, at least 15 volumes of aprotic polar 15th solvent.

In some aspects, between about 1 and about 30 equiv of **20b** may be added to about 1 equiv of peptide **2**. In some aspects, between about 1 and about 20 equiv of **20b** may be added to about 1 equiv of peptide **2**. In some aspects, between about 1 and about 10 equiv of **20b** may be added to about 1 equiv of peptide **2**. In some aspects, between about 2 and about 30 equiv of **20b** may be added to about 1 equiv of peptide **2**. In some aspects, between about 3 and about 30 equiv of **20b** may be added to about 1 equiv of peptide **2**. In some aspects, between about 3 and about 20 equiv of **20b** may be added to about 1 equiv of peptide **2**. In some aspects, between about 3 and about 10 equiv of **20b** may be added to about 1 equiv of peptide **2**. In some aspects, at least 3 equiv of **20b** is added to about 1 equiv of peptide **2**. In some aspects, about 3 equiv of **20b** may be added to about 1 equiv of peptide **2**.

In some aspects, the **20b** and **2** are combined in between about 1 to about 10 equiv of 3rd base, compared to peptide **2**. In some aspects, the **20b** and **2** are combined in between about 1 to about 5 equiv of 3rd base, compared to peptide **2**. In some aspects, the **20b** and **2** are combined in between about 1 to about 2 equiv of 3rd base, compared to peptide **2**. In some aspects, the **20b** and **2** are combined in between about 1 to about 1.5 equiv of 3rd base, compared to peptide **2**. In some aspects, the **20b** and **2** are combined in about 1.3 equiv of 3rd base, compared to peptide **2**. The 3rd base may be selected from the group consisting of NMM, TEA, DIPEA and pyridine. The 3rd base may be NMM.

Due to the low solubility of peptide **2**, no conversion to peptide-linker conjugate **1** was observed in solvents such as MeCN or MeOH. The addition of water to these solvents to help bring peptide **2** into solution led to low conversions due to activated linker decomposition and to the formation of thick gels after a few hours.

In accordance with the present invention, aprotic, polar solvents (such as DMF), may fully dissolve peptide **2** and allow full consumption of the peptide **2** at RT (e.g. about 20°C) with about 3 equiv of compound **20b**. The cleanest reaction was obtained in DMF in combination with NMM as base. Lower purities were obtained in DMSO, NMP and DMAc with bases such as TEA, DIPEA or pyridine.

In some aspects, the reaction comprises mixing about 20 volumes of DMF, between about 2 to about 3 equiv of **20b**, and about 20 equiv of NMM. The reaction may be at RT. The reaction may be carried out for at least 10 minutes.

Conditions for the isolation of peptide-linker conjugate **3** were also investigated. The simplest approach was the identification of a suitable anti-solvent that would precipitate peptide-linker conjugate **3** from solution followed by filtration. A number of organic solvents were tested for this purpose. Whereas toluene, EtOAc, THF, and MTBE gave sticky, gel-like solids that were difficult to filter, some aspects of the invention are based on the surprising discovery that MeCN produced fine, free-flowing solids that are much more easily handled.

Accordingly, in some aspects of the invention, **3** may be subsequently isolated by precipitation with MeCN, followed by filtration.

In some aspects, the proportion of anti-solvent:aprotic, polar 15th solvent is between about 5:1 to about 20:1, and is preferably about 9:1.

Filtration in the presence of an inert gas (N₂, Ar, CO₂ etc) at this point was desirable to prevent moisture adsorption on the cake, otherwise the product may
5 turn into a gummy solid. The workup procedure involved the slow transfer of the DMF solution of peptide-linker conjugate **3** after reaction completion into MeCN to give a final 9:1 MeCN/DMF mixture. After the resulting precipitate was aged (where the solid is afforded sufficient time to precipitate out of solution), the solid was filtered and dried at between about 0 to about 50°C under vacuum. In some
10 aspects, the precipitate was aged for up to 24 hrs. In some aspects, the precipitate was aged for up to 12 hrs. In some aspects, the precipitate was aged for up to 6 hrs. In some aspects, the precipitate was aged for up to 3 hrs. In some aspects, the precipitate was aged for up to 2 hrs. In some aspects, further stirring after about 2 hrs makes the solid sticky and difficult to handle. In some aspects,
15 the solid was dried at about 0 to about 40°C. In some aspects, the solid was dried at about 20 to about 50°C. Higher temperatures are likely to have a detrimental effect.

Accordingly, in some aspects of the invention, the filtration may be conducted in the absence of water vapour or atmosphere. In some aspects, the
20 filtration may be conducted under inert gas. The inert gas may be selected from the group consisting of N₂, Argon, CO₂, etc.

In some aspects of the invention, the peptide **2** and compound **20b** may be conjugated together at a temperature of between about -30°C and about 30°C. In some aspects of the invention, the peptide **3** and the compound **20b**
25 may be conjugated together at a temperature range whose lower limit is selected from the group consisting of about -30, -25, -20, -18, -17, -16, -15, -14, -13, -12, -10, -5, -1, 0, 1, 2, 3, 4, 5, 10, 15 and 18 °C, and whose upper limit is selected from the group consisting of about -14, -13, -12, -10, -5, -1, 0, 1, 2, 3, 4, 5, 10, 15, 20, 25 and 30 °C. In some aspects of the invention, the peptide **2** and compound
30 **20b** may be conjugated together at a temperature of between about -30°C and

about 20°C. In some aspects of the invention, the peptide **2** and compound **20b** may be conjugated together at a temperature of about -15°C.

In some aspects conjugation between **2** and **20b** proceeds even at -30 °C but can require between about 18 to about 20 hrs, and may necessitate up to
5 about 5 equiv of pentafluorophenol ester linker to fully consume the peptide. In addition, the longer reaction times can give more byproduct formation as well as hazy mixtures due to partial product precipitation. After further experimentation, a satisfactory compromise was surprisingly found between the kinetics of the process and a satisfactory impurity profile. Accordingly, in some aspects, the
10 invention provides for a process that runs the reaction in about 15 volumes of DMF, preferably between about 15 to about 50 volumes of DMF, (enough to easily dissolve peptide **2** and prevent gelification) in the presence of about 3 equiv of compound **20b** and between about 1 and about 5, and preferably about 1.2 equiv NMM at between about -15 to about -18 °C. After about 7hr, less than
15 about 1% of unreacted **2** remained and the total byproduct of ester **20b** was kept at or below about 0.3%.

After 7 hrs, less than 1% of unreacted **2** remained and the total pentafluorophenol ester byproduct level was kept at 0.3%. The mixture may then be filtered through a 0.45 micron in-line filter and slowly added into the anti-
20 solvent, such as MeCN, while using low agitation to promote larger particle size. The amount of MeCN may be advantageously chosen such that the final desired ratio of MeCN: DMF is achieved. In some aspects, the amount of MeCN is about 5 to about 20 times the amount of aprotic polar 15th solvent, and may be arranged so as to give a final ratio of MeCN:DMF of 9:1.

25 Solid precipitation typically occurs immediately and, after an aging period (for example, about 1-2 hrs), the solids may be filtered, washed with fresh MeCN, and dried under vacuum.

In some aspects, a DMF content of 4.5% (wt/wt) may still be present in the solid, most likely due to the intentional slow agitation during the quick
30 precipitation of the product from solution and the resulting entrapment of DMF. Screening the solid through a # 20 hand-sieve followed by a MeCN reslurry



(wherein the solid was suspended in a solvent and stirred without fully dissolving the solid to remove impurities), and subsequent drying at between about 20 and about 50°C, and preferably at about 40°C provides material in 83% yield.

In some aspects, a high agitation speed or subsurface addition of the product solution into MeCN during the precipitation can minimize the amount of trapped DMF.


Accordingly, in some aspects the invention provides a means to minimize the entrapment of DMF in the isolated product solids, comprising the addition of the product solution into MeCN with vigorous mixing. This can be accomplished using high speed agitation in the isolation vessel, or preferably, by also adding the product solution through an addition tube whose outlet is below the surface of the MeCN and near a region of high shear (close to the impeller blades). The vigorous agitation allows the product solution to be more thoroughly dispersed before precipitation and reduces the concentration of DMF in the region of precipitation.

Room temperature (RT) may be between about 15°C and about 25°C. In some aspects, RT may be between about 18°C and about 22°C. In some aspects, RT may be between about 18°C and about 20°C. In some aspects, RT may be between about 20°C.

In some aspects, the invention provides for use of PNP esters rather than tri-, tetra- and penta- halo substituted phenyls. Accordingly, in some aspects the invention provides for compounds and intermediates of the formulae:



, and




, wherein q is 1, 2, 3, 4, or 5, n = 1, 2, 3, 4,


5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27,

28, 29, 30, and X is any halogen. In some aspects, $n = 1-10$. In some aspects, $n = 1, 2, 3, 4, 5$, or 6 . In some aspects, $q = 1, 2$, or 3 . In some aspects, $q = 2$.

In some aspects, the invention provides a method of generating activated esters of formula **22**, **22a**, and **22b**, comprising reacting compound **21** with
 5 compound **6**, **6a**, or **6b** respectively (using the same chemistry as described for reaction of **6b** and **19b**). Compounds **22**, **22a**, and **22b** can be conjugated with ϵ -amino group-bearing side chains (e.g. a lysine side chain) of peptides **2** to form peptide-linker conjugates **3**, **3a**, and **3b** respectively.



10 The surprising part of this aspect of the invention is that use of PNP esters of the formula **22**, **22a** and **22b** for the coupling with the peptide **2** afford products **3**, **3a**, **3b** respectively that do not contain corresponding PNP ester impurities of **3**, **3a**, **3b**. Possible chemical structures of these impurities are shown below.



$R_1, R_2, R_3, R_4 = \text{H or p-nitrophenyl}$

15 These types of impurities are typically formed when penta halophenol esters are used as activating groups during coupling reaction of **20b** with **2** resulting in the formation of corresponding pentahalo ester impurities (see example 16). These impurities are very reactive and can potentially react with a monoclonal antibody used in subsequent conjugation steps, such as h38C2,
 20 resulting in the formation undesired impurities in the drug substance (see discussion in example 16). In some aspects, the invention describes a process for preparation of peptide-linker-antibody **5** of the formula shown below. In some aspects, the antibody is h38C2 or variants thereof. In some aspects, the peptide-linker is **3**, **3a** or **3b**.

ABBREVIATIONS

	<i>n</i> -BuLi:	<i>n</i> -butyl lithium
	CDI:	1,1'-carbonyldiimidazole
	CDMT:	2-chloro-4,6-dimethoxy-1,3,5-triazine
5	DBU:	1,8-diazabicyclo[5.4.0]undec-7-ene
	DABCO:	1,4-diazabicyclo[2.2.2]octane
	DCC:	<i>N,N'</i> -dicyclohexylcarbodiimide
	DCMT:	2,4-dichloro-6-methoxy-1,3,5-triazine
	DIC:	<i>N,N'</i> -diisopropylcarbodiimide
10	DIPEA:	diisopropylethylamine
	DMAc:	<i>N,N</i> -dimethylacetamide
	DMF:	<i>N,N</i> -dimethylformamide
	DMSO:	dimethylsulfoxide
	DPPA	diphenylphosphoryl azide
15	EDC:	1-[3-(dimethylaminopropyl)]-3-ethylcarbodiimide hydrochloride
	EtOAc:	ethyl acetate
	HATU:	<i>O</i> -(7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate
20	HBTU:	<i>O</i> -(benzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate
	HOAc:	acetic acid
	MeCN:	acetonitrile
	MeOH:	methanol
25	MTBE:	tert-butyl methyl ether
	NH ₄ OAc:	ammonium acetate

	NMM:	N-methylmorpholine
	NMP:	1-methyl-2-pyrrolidinone
	PFP	pentafluorophenyl
	PNP	para nitrophenyl
5	<i>i</i> -PrOAc:	<i>i</i> -propyl acetate
	PyBOP:	benzotriazol-1-yloxytri(pyrrolidino)phosphonium hexafluorophosphate
	PyBroP:	bromotri(pyrrolidino)phosphonium hexafluorophosphate
	PyCloP:	chlorotri(pyrrolidino)phosphonium hexafluorophosphate
10	T3P:	1-propanephosphonic acid anhydride
	TBTU:	O-benzotriazol-1-yl-1,1,3,3-tetramethyluronium tetrafluoroborate
	TEA :	triethylamine
	THF:	tetrahydrofuran
15	HIC	Hydrophobic interaction chromatography
	SEC	Size Exclusion Chromatography

SEQUENCE LIST

SEQ ID NO:	Description	Sequence
1	Ang2 binding peptide X2 is Acyl Lysine X9 is acyl lysine	QxYQPLDExD KTLYDQFMLQ QG
2	Ang2 binding peptide X1 is C(O)CH ₃ X3 is Acyl Lysine X10 is acyl lysine X24 is NH ₂	xQxYQPLDEx DKTLYDQFML QQGx
3	h38C2 light chain	ELQMTQSPSS LSASVGDRVT ITCRSSQSLL HTYGSPYLNW YLQKFGQSPK LLIYKVSNRF SGVPSRFSGS GSGTDFTLTI SSLQPEDFAV YFCSQGTHLP YTFGGGTKVE IKRTVAAPSV FIFPPSDEQL KSGTASVVCL LNNFYPREAK VQWKVDNALQ SGNSQESVTE QDSKDSTYSL SSTLTLSKAD YEKHKVYACE VTHQGLSSPV TKSFNRGEC

4	h38C2 heavy chain	EVQLVESGGG LVQPGGSLRL SCAASGFTFS NYWMSWVRQS PEKGLEWVSE IRLRSDNYAT HYAESVKGGRF TISRDN SKNT LYLQMNSLRA EDTGIYYCKT YFYFSYWGQ GTLVTVSSAS TKGPSVFPLA PSSKSTSGGT AALGCLVKDY FPEPVTVSWN SGALTSGVHT FPAVLQSSGL YSLSSVVTVP SSSLGTQTYI CNVNHKPSNT KVDKRVEPKS CDKTHTCPPC PAPELLGGPS VFLFPPKPKD TLMISRTPEV TCVVVDVSHE DPEVKFNWYV DGVEVHNAKT KPREEQYNST YRVVSVLTVL HQDWLNGKEY KCKVSNKALP APIEKTISKA KGQPREPQVY TLPPSREEMT KNQVSLTCLV KGFYPSTIAV EWESNGQPEN NYKTTTPVLD SDGSFFLYSK LTVDKSRWQQ GNVFSCSVMH EALHNHYTQK SLSPGK
5	VL h38C2	ELQMTQSPSS LSASVGRVT ITCRSSQSL HTYGSPYLNW YLQKPGQSPK LLIYKVSNRFG SGVPDRFSGS GSGTDFTLT I SSLQPEDFAV YFCSQGTSLP YTFGGGTKVE IK
6	VH h38C2	EVQLVESGGG LVQPGGSLRL SCAASGFTFS NYWMSWVRQS PEKGLEWVSE IRLRSDNYAT HYAESVKGGRF TISRDN SKNT LYLQMNSLRA EDTGIYYCKT YFYFSYWGQ GTLVTVSS
7	VL m38C2	DVVMTQTPLS LPVRLGQAS ISCRSSQSL HTYGSPYLNW YLQKPGQSPK LLIYKVSNRFG SGVPDRFSGS GSGTDFTLT I SRVEAEDLG V YFCSQGTSLP YTFGGGTKLE IK
8	VH m38C2	EVKLVESGGG LVQPGGTMKL SCEISGLTFR NYWMSWVRQS PEKGLEWVAE IRLRSDNYAT HYAESVKGGRF TISRDDSKSR LYLQMNSLRT EDTGIYYCKY YFYFSYWGQ GTLVTVSA
9	h38C2-IgG1 LC constant region genus x46=V or A, x84=V or L	RTVAAPSVFI FPPSDEQLKS GTASVVCLLN NFYPREAKVQ WKVDNQLQSG NSQESVTEQD SKDSTYSLSS TLTLKADYE KHKYACEVT HQGLSSPVTK SFNRGEC
10	h38C2-IgG1 LC constant region Km(1)	RTVAAPSVFI FPPSDEQLKS GTASVVCLLN NFYPREAKVQ WKVDNQLQSG NSQESVTEQD SKDSTYSLSS TLTLKADYE KHKLYACEVT HQGLSSPVTK SFNRGEC
11	h38C2-IgG1 LC constant region Km(1,2)	RTVAAPSVFI FPPSDEQLKS GTASVVCLLN NFYPREAKVQ WKVDNQLQSG NSQESVTEQD SKDSTYSLSS TLTLKADYE KHKLYACEVT HQGLSSPVTK SFNRGEC
12	h38C2-IgG1 LC constant region Km(3)	RTVAAPSVFI FPPSDEQLKS GTASVVCLLN NFYPREAKVQ WKVDNQLQSG NSQESVTEQD SKDSTYSLSS TLTLKADYE KHKVYACEVT HQGLSSPVTK SFNRGEC
13	H38C2 IgG1 HC constant region	AS TKGPSVFPLA PSSKSTSGGT AALGCLVKDY FPEPVTVSWN SGALTSGVHT FPAVLQSSGL YSLSSVVTVP SSSLGTQTYI CNVNHKPSNT KVDKRVEPKS CDKTHTCPPC PAPELLGGPS VFLFPPKPKD TLMISRTPEV TCVVVDVSHE DPEVKFNWYV DGVEVHNAKT KPREEQYNST YRVVSVLTVL HQDWLNGKEY KCKVSNKALP APIEKTISKA KGQPREPQVY TLPPSREEMT KNQVSLTCLV KGFYPSTIAV EWESNGQPEN NYKTTTPVLD SDGSFFLYSK LTVDKSRWQQ GNVFSCSVMH EALHNHYTQK SLSPGK
14	h38C2-IgG2 HC:	AS TKGPSVFPLA PCSRSTSEST AALGCLVKDY FPEPVTVSWN SGALTSGVHT FPAVLQSSGL YSLSSVVTVP SSNFGTQTYT CNVDHKPSNT KVDKTVKRC CVECPCPAP PVAGPSVFLF PPKPKDTLMI SRTPEVTCVV VDVSHEDPEV QFNWYVDGVE VHNAKTKPRE EQFNSTFRV SVLTVVHQDW LNGKEYKCKV SNKGLPSSIE KTISKTKGQP REPQVYTLPP SREEMTKNQV SLTCLVKGFY PSDIAVEWES NGQPENNYKT TPPLDSDGS FFLYSLKLTVD KSRWQQGNVF SCSVMHEALH NHYTQKSLSL SPGK

DEFINITIONS

- "About" or "approximately," when used in connection with a measurable numerical variable, refers to the indicated value of the variable and to all values of the variable that are within the experimental error of the indicated value (e.g.,
- 5 within the 95% confidence interval for the mean) or within 10 percent of the indicated value, whichever is greater.

An "antibody" is an immunoglobulin molecule capable of specific binding to a target, such as a carbohydrate, polynucleotide, lipid, polypeptide, etc., through at least one antigen recognition site, located in the variable region of the immunoglobulin molecule. As used herein, the term "antibody" encompasses not only intact polyclonal or monoclonal antibodies, but also any antigen binding fragment (i.e., "antigen-binding portion") or single chain thereof, fusion proteins comprising an antibody, and any other modified configuration of the immunoglobulin molecule that comprises an antigen recognition site including, for example without limitation, scFv, single domain antibodies (e.g., shark and camelid antibodies), maxibodies, minibodies, intrabodies, diabodies, triabodies, tetrabodies, v-NAR and bis-scFv (see, e.g., Hollinger and Hudson, 2005, Nature Biotechnology 23(9): 1126-1136). An antibody includes an antibody of any class, such as IgG, IgA, or IgM (or sub-class thereof), and the antibody need not be of any particular class. Depending on the antibody amino acid sequence of the constant region of its heavy chains, immunoglobulins can be assigned to different classes. There are five major classes of immunoglobulins: IgA, IgD, IgE, IgG, and IgM, and several of these may be further divided into subclasses (isotypes), e.g., IgG1, IgG2, IgG3, IgG4, IgA1 and IgA2. The heavy-chain constant regions that correspond to the different classes of immunoglobulins are called alpha, delta, epsilon, gamma, and mu, respectively. The subunit structures and three-dimensional configurations of different classes of immunoglobulins are well known.

Two types of human light chain constant regions are known: lambda (CL- λ) and kappa (CL- κ). There are three known CL- κ variants, based on the polymorphisms V/A at position 46 and A/L at position 84 (numbering according to SEQ ID NO:9). The 3 identified CL- κ polymorphisms are Km(1):V⁴⁶/L⁸⁴, Km(1,2):A⁴⁶/L⁸⁴, and Km(3) A⁴⁶/V⁸⁴. Antibodies of the present invention may therefore comprise a constant kappa domain according to any one of SEQ ID NOs:9, 10, 11 or 12, or variants thereof that comprise no more than 5, 4, 3, 2, or 1 amino acid insertions, substitutions or deletions. It is understood by the skilled person that residue R¹ of SEQ ID NOs:9, 10, 11 and 12 by some counting methods may

be included in the variable domain, and that the constant domains may therefore also be considered as beginning from residue T² of said sequences.

The term "antigen binding portion" of an antibody, as used herein, refers to one or more fragments of an intact antibody that retain the ability to specifically
5 bind to a given antigen (e.g., target X). Antigen binding functions of an antibody can be performed by fragments of an intact antibody. Examples of binding fragments encompassed within the term "antigen binding portion" of an antibody include Fab; Fab'; F(ab')₂; an Fd fragment consisting of the VH and CH1 domains; an Fv fragment consisting of the VL and VH domains of a single arm of
10 an antibody; a single domain antibody (dAb) fragment (Ward et al., 1989 Nature 341:544-546), and an isolated complementarity determining region (CDR).

A "variable region" of an antibody refers to the variable region of the antibody light chain or the variable region of the antibody heavy chain, either alone or in combination. As known in the art, the variable regions of the heavy
15 and light chain each consist of four framework regions (FRs) connected by three complementarity determining regions (CDRs) also known as hypervariable regions, contribute to the formation of the antigen binding site of antibodies. If variants of a subject variable region are desired, particularly with substitution in amino acid residues outside of a CDR region (i.e., in the framework region),
20 appropriate amino acid substitution, preferably, conservative amino acid substitution, can be identified by comparing the subject variable region to the variable regions of other antibodies which contain CDR1 and CDR2 sequences in the same canonical class as the subject variable region (Chothia and Lesk, J Mol Biol 196(4): 901-917, 1987). When choosing FR to flank subject CDRs, e.g.,
25 when humanizing or optimizing an antibody, FRs from antibodies which contain CDR1 and CDR2 sequences in the same canonical class are preferred.


A "CDR" of a variable domain are amino acid residues within the variable region that are identified in accordance with the definitions of the Kabat, Chothia, the accumulation of both Kabat and Chothia, AbM, contact, and/or conformational
30 definitions or any method of CDR determination well known in the art. Antibody CDRs may be identified as the hypervariable regions originally defined by Kabat

et al. See, e.g., Kabat et al., 1992, Sequences of Proteins of Immunological Interest, 5th ed., Public Health Service, NIH, Washington D.C. The positions of the CDRs may also be identified as the structural loop structures originally described by Chothia and others. See, e.g., Chothia et al., 1989, Nature 342:877-883. Other approaches to CDR identification include the "AbM definition," which is a compromise between Kabat and Chothia and is derived using Oxford Molecular's AbM antibody modelling software (now Accelrys®), or the "contact definition" of CDRs based on observed antigen contacts, set forth in MacCallum et al., 1996, J. Mol. Biol., 262:732-745. In another approach, referred to herein as the "conformational definition" of CDRs, the positions of the CDRs may be identified as the residues that make enthalpic contributions to antigen binding. See, e.g., Makabe et al., 2008, Journal of Biological Chemistry, 283:1156-1166. Still other CDR boundary definitions may not strictly follow one of the above approaches, but will nonetheless overlap with at least a portion of the Kabat CDRs, although they may be shortened or lengthened in light of prediction or experimental findings that particular residues or groups of residues or even entire CDRs do not significantly impact antigen binding. As used herein, a CDR may refer to CDRs defined by any approach known in the art, including combinations of approaches. The methods used herein may utilize CDRs defined according to any of these approaches. For any given embodiment containing more than one CDR, the CDRs may be defined in accordance with any of Kabat, Chothia, extended, AbM, contact, and/or conformational definitions.

The contents of US2006205670 (US7521425) are incorporated herein by reference. US2006205670 (US7521425) describes a number of compositions and techniques directly applicable to the present application, in particular at paragraphs [0153]-[0233], describing antibodies, useful fragments and variants and modifications thereof, combining sites and CDRs, antibody preparation, expression, humanization, amino acid modification, glycosylation, ADCC, CDC, increasing serum half life of antibodies, expression vectors, mammalian host systems, and folding, amongst other elements of antibody technology.

“Combining site”, as used herein, (also known as the antibody binding site) refers to the region of the immunoglobulin or Ig domains that combine (or can combine) with the determinant of an appropriate antigen (or a structurally similar protein). The term generally includes the CDRs and the adjacent framework residues that are involved in antigen binding.

“Aldolase antibodies” as used herein, refers to antibodies containing combining site portions that, when unencumbered (for example by conjugation), catalyze an aldol addition reaction between an aliphatic ketone donor and an aldehyde acceptor. Aldolase antibodies are capable of being generated by immunization of an immune-responsive animal with an immunogen that includes a 1,3 diketone hapten of the formula:



coupled to a carrier protein, and further characterized by having a lysine with a reactive ϵ -amino group in the combining site of the antibody. Aldolase antibodies are further characterized by their catalytic activity being subject to inhibition with the 1,3-diketone hapten by formation of a complex between the 1,3-diketone hapten and the ϵ -amino group of the lysine of the catalytic antibody.

As discussed, in certain embodiments, certain antibodies that can be used in conjunction with compounds of the invention may require a reactive side chain in the antibody combining site. A reactive side chain may be present naturally or may be placed in an antibody by mutation. The reactive residue of the antibody combining site may be associated with the antibody, such as when the residue is encoded by nucleic acid present in the lymphoid cell first identified to make the antibody. Alternatively, the amino acid residue may arise by purposely mutating the DNA so as to encode the particular residue (e.g. WO 01/22922). The reactive residue may be a non-natural residue arising, for example, by biosynthetic incorporation using a unique codon, tRNA, and aminoacyl-tRNA as discussed herein. In another approach, the amino acid residue or its reactive functional groups (e.g., a nucleophilic amino group or sulfhydryl group) may be attached to an amino acid residue in the antibody combining site. Thus, covalent linkage with

the antibody occurring "through an amino acid residue in a combining site of an antibody" as used herein means that linkage can be directly to an amino acid residue of an antibody combining site or through a chemical moiety that is linked to a side chain of an amino acid residue of an antibody combining site. In some
5 embodiments, the amino acid is cysteine, and the reactive group of the side chain is a sulfhydryl group. In other embodiments, the amino acid residue is lysine, and the reactive group of the side chain is the ϵ -amino group. In some embodiments, the amino acid is Lys93 on the heavy chain according to Kabat numbering. In some embodiments, the amino acid is Lys-99 on HC h38C2 (SEQ
10 ID NO:4).

Catalytic antibodies are one source of antibodies with suitable combining sites that comprise one or more reactive amino acid side chains. Such antibodies include aldolase antibodies, beta lactamase antibodies, esterase antibodies, and amidase antibodies.

15 One embodiment comprises an aldolase antibody such as the mouse monoclonal antibodies mAb 33F12 and mAb 38C2, as well as suitably chimeric and humanized versions of such antibodies (e.g. h38C2, SEQ ID NOs:3 and 4). Mouse mAb 38C2 (and h38C2) has a reactive lysine near to but outside HCDR3, and is the prototype of a new class of catalytic antibodies that were generated by
20 reactive immunization and mechanistically mimic natural aldolase enzymes. See C.F. Barbas 3rd *et al.*, Science 278:2085-2092 (1997). Other aldolase catalytic antibodies that may be used include the antibodies produced by the hybridoma 85A2, having ATCC accession number PTA-1015; hybridoma 85C7, having ATCC accession number PTA-1014; hybridoma 92F9, having ATCC accession
25 number PTA-1017; hybridoma 93F3, having ATCC accession number PTA-823; hybridoma 84G3, having ATCC accession number PTA-824; hybridoma 84G11, having ATCC accession number PTA-1018; hybridoma 84H9, having ATCC accession number PTA-1019; hybridoma 85H6, having ATCC accession number PTA-825; hybridoma 90G8, having ATCC accession number PTA-1016. Through
30 a reactive lysine, these antibodies catalyze aldol and retro-aldol reactions using the enamine mechanism of natural aldolases. Aldolase antibodies and methods

of generating aldolase antibodies are disclosed in U.S. Patents Nos. 6,210,938, 6,368,839, 6,326,176, 6,589,766, 5,985,626, and 5,733,75, which are incorporated herein by reference.

Compounds of the invention may also be formed by linking a compound of
5 the invention to a reactive cysteine, such as those found in the combining sites of thioesterase and esterase catalytic antibodies. Suitable thioesterase catalytic antibodies are described by K.D. Janda *et al.*, Proc. Natl. Acad. Sci. U.S.A. 91:2532-2536 (1994). Suitable esterase antibodies are described by P. Wirsching *et al.*, Science 270:1775-1782 (1995). Reactive amino acid-containing antibodies
10 may be prepared by means well known in the art, including mutating an antibody combining site residue to encode for the reactive amino acid or chemically derivatizing an amino acid side chain in an antibody combining site with a linker that contains the reactive group.

The antibody may be a humanized antibody. Where compounds of the
15 invention are covalently linked to the combining site of an antibody, and such antibodies are humanized, it is important that such antibodies be humanized with retention of high linking affinity for the Z group. Various forms of humanized murine aldolase antibodies are contemplated. One embodiment uses the humanized aldolase catalytic antibody h38c2 IgG1 or h38c2 Fab with human
20 constant domains C_κ and C_{γ1}1. C. Rader *et al.*, J. Mol. Bio. 332:889-899 (2003) discloses the gene sequences and vectors that may be used to produce h38c2 Fab and h38c2 IgG1. Human germline V_κ gene DPK-9 and human J_κ gene JK4 were used as frameworks for the humanization of the kappa light chain variable domain of m38c2, and human germline gene DP-47 and human J_H gene JH4
25 were used as frameworks for the humanization of the heavy chain variable domain of m38c2. Figure 7A of US2006205670 (herein incorporated by reference) illustrates a sequence alignment between the variable light and heavy chains in m38c2, h38c2, and human germlines. h38c2 may utilize IgG1, IgG2, IgG3, or IgG4 constant domains, including any of the allotypes thereof. In certain
30 embodiments of compounds of the invention wherein the antibody is h38c2 IgG1 with the G1m(f) allotype, Z binds to the side chain of the lysine residue at position

99 of the heavy chain. Another embodiment uses a chimeric antibody comprising the variable domains (V_L and V_H) of h38c2 (SEQ ID NOs:5 and 6) and the constant domains from an IgG1, IgG2, IgG3, or IgG4. The antibody may be a full-length antibody, Fab, Fab', $F(ab')_2$, F_v , dsF_v , scF_v , V_H , V_L , diabody, or minibody.

- 5 The antibody may be a full length antibody, and may be selected from the group consisting of IgG1, IgG2, IgG_{2Δa}, IgG3, IgG4, IgG_{4Δb}, IgG_{4Δc}, IgG₄ S228P, IgG_{4Δb} S228P and IgG_{4Δc} S228P. The antibody or antigen binding portion thereof may comprise the V_H and V_L domains from h38c2. The antibody may be an antibody comprising the V_L and V_H domains from h38c2 and a constant domain selected
- 10 from the group consisting of IgG1, IgG2, IgG_{2Δa}, IgG3, IgG4, IgG_{4Δb}, IgG_{4Δc}, IgG₄ S228P, IgG_{4Δb} S228P and IgG_{4Δc} S228P. The antibody may be h38C2 IgG1 (SEQ ID NOs:3 and 4). The antibody may be h38C2 IgG2 (SEQ ID NOs:3 and 14). The antibody may be a humanized version of a murine aldolase antibody comprising a constant region from a human IgG, IgA, IgM, IgD, or IgE antibody.
- 15 In another embodiment, the antibody is a chimeric antibody comprising the V_L and V_H region from a murine aldolase antibody and a constant region from a human IgG, IgA, IgM, IgD, or IgE antibody. In some embodiments, the antibody comprises the V_L and V_H regions from m38C2 (SEQ ID NOs:7 and 8). In further embodiments, the antibody is a fully human version of a murine aldolase
- 20 antibody comprising a polypeptide sequence from natural or native human IgG, IgA, IgM, IgD, or IgE antibody. In some aspects, the antibody may comprise a light chain variable region (V_L) comprising a V_L CDR1, V_L CDR2, and V_L CDR3 of the V_L sequence shown in SEQ ID NO:5; and a heavy chain variable region (V_H) comprising a V_H CDR1, V_H CDR2, and V_H CDR3 of the V_H sequence shown in
- 25 SEQ ID NO:6. As outlined above, the CDRs may be determined by a number of known methods of the art.

In some aspects, the antibody the antibody comprises a light chain at least 95% identical to SEQ ID NO:3 and a heavy chain at least 95% identical to SEQ ID NO:4. The light chain may be at least 96% identical to SEQ ID NO:3. The light

30 chain may be at least 96% identical to SEQ ID NO:3. The light chain may be at least 97% identical to SEQ ID NO:3. The light chain may be at least 98%

identical to SEQ ID NO:3. The light chain may be at least 99% identical to SEQ ID NO:3. The heavy chain may be at least 96% identical to SEQ ID NO:4. The heavy chain may be at least 97% identical to SEQ ID NO:4. The heavy chain may be at least 98% identical to SEQ ID NO:4. The heavy chain may be at least 99% identical to SEQ ID NO:4. In some aspects, the light chain may differ from SEQ ID NO:3 by one amino acid. In some aspects, the heavy chain may differ from SEQ ID NO:4 by one amino acid. In some aspects, the differences between the light chain and SEQ ID NO:3 may be located in the constant region only. In some aspects, the differences between the heavy chain and SEQ ID NO:4 may be located in the constant region only.

In some aspects, the antibodies of the present invention comprise a light chain comprising a light chain constant region comprising a sequence selected from the group consisting of SEQ ID NOs:9, 10, 11, and 12, or variants thereof comprising between 1 and 5 amino acid substitutions, deletions or insertions, and a light chain variable region comprising SEQ ID NO:3.

In some aspects, the antibodies of the present invention comprise a heavy chain comprising a heavy chain constant region comprising a sequence selected from the group consisting of SEQ ID NOs:13, and 14, or variants thereof comprising between 1 and 5 amino acid substitutions, deletions or insertions, and a heavy chain variable region comprising SEQ ID NO:4.

In some aspects, the antibodies of the present invention comprise a light chain constant region comprising a sequence selected from the group consisting of SEQ ID NOs:9, 10, 11, and 12, or variants thereof comprising between 1 and 5 amino acid substitutions, deletions or insertions, and a heavy chain constant region comprising a sequence selected from the group consisting of SEQ ID NOs:13, and 14, or variants thereof comprising between 1 and 5 amino acid substitutions, deletions or insertions. In some aspects, the heavy chain is SEQ ID NO:13, or variants thereof comprising between 1 and 5 amino acid substitutions, deletions or insertions.

Amino acid sequence modification(s) of the antibodies described herein are contemplated. For example, it may be desirable to improve the binding

affinity and/or other biological properties of the antibody. Amino acid sequence variants of an antibody are prepared by introducing appropriate nucleotide changes into the antibody nucleic acid, or by peptide synthesis. Such modifications include, for example, deletions from, insertions into, and/or
5 substitutions of residues within the amino acid sequences of the antibody. Any combination of deletion, insertion, and substitution is made to arrive at the final construct, provided that the final construct possesses the desired characteristics. The amino acid changes also may alter post-translational processes of the antibody, such as changing the number or position of glycosylation sites.

10 An antibody or antibody portion of the invention can be derivatized or linked to another molecule (e.g. another peptide or protein). In general, the antibodies or portion thereof is derivatized such that the ability of the linker to covalently conjugate to the antibody combining is not affected adversely by the derivatization or labelling. Accordingly, the antibodies and antibody portions of
15 the invention are intended to include both intact and modified forms of the antibodies described herein. E.g. an antibody or antibody portion of the invention can be functionally linked (by chemical coupling, genetic fusion, noncovalent association or otherwise) to one or more other molecular entities, such as another antibody (e.g. a bispecific antibody or a diabody), a detection agent, a
20 cytotoxic agent, a pharmaceutical agent, and/or a protein or peptide that can mediate association of the antibody or antibody portion with another molecule (such as a streptavidin core region or a polyhistidine tag).

In other embodiments, the antibody or antigen binding portion thereof of the invention may be a fusion antibody or an antibody linked to another
25 polypeptide. In some aspects, only the variable regions of the antibody are linked to the polypeptide. In some aspects, the antibody is covalently conjugated to a peptide in such a way so as to not interfere with the binding ability of the combining site.

The polypeptide may be a therapeutic agent, such as a targeting agent,
30 peptide, protein agonist, protein antagonist, metabolic regulator, hormone, toxin, growth factor or other regulatory protein, or may be a diagnostic agent, such as

an enzyme that may be easily visualized, such as horseradish peroxidase. In addition, fusion antibodies can be created in which two (or more) single-chain antibodies are linked to one another. This is useful if one wants to create a divalent or polyvalent antibody on a single polypeptide chain, or if one wants to
5 create a bispecific antibody. By "peptide", it is understood that the term encompasses chains of amino acids commonly referred to as peptides, polypeptides and proteins. In some aspects, the peptide is at least three amino acids in length. In some aspects, the peptide is less than about 500 amino acids in length. In some aspects, the peptide is less than about 300 amino acids in
10 length. In some aspects, the peptide is less than about 200 amino acids in length. In some aspects, the peptide is less than about 150 amino acids in length. In some aspects, the peptide is less than about 100 amino acids in length. In some aspects, the peptide is less than about 80 amino acids in length. In some aspects, the peptide is less than about 70 amino acids in length. In some
15 aspects, the peptide is less than about 60 amino acids in length. In some aspects, the peptide is less than about 50 amino acids in length. In some aspects, the peptide is less than about 40 amino acids in length.

One type of derivatized antibody is produced by crosslinking two or more antibodies (of the same type or of different types, e.g. to create bispecific
20 antibodies). Suitable crosslinkers include those that are heterobifunctional, having two distinctly reactive groups separated by an appropriate spacer (e.g. m-maleimidobenzoyl-N-hydroxysuccinimide ester) or homobifunctional (e. g. disuccinimidyl suberate).

Another type of derivatized antibody is a labelled antibody. Useful
25 detection agents with which an antibody or antibody portion of the invention may be derivatized include fluorescent compounds, including fluorescein, fluorescein isothiocyanate, rhodamine, 5-dimethylamine-1-naphthalenesulfonyl chloride, phycoerythrin, lanthanide phosphors and the like. An antibody may also be labelled with enzymes that are useful for detection, such as horseradish
30 peroxidase, galactosidase, luciferase, alkaline phosphatase, glucose oxidase and the like. When an antibody is labelled with a detectable enzyme, it is detected by

adding additional reagents that the enzyme uses to produce a reaction product that can be discerned. For example, when the agent horseradish peroxidase is present, the addition of hydrogen peroxide and diaminobenzidine leads to a colored reaction product, which is detectable. An antibody may also be labelled
5 with biotin, and detected through indirect measurement of avidin or streptavidin binding. An antibody may be labelled with a magnetic agent, such as gadolinium. An antibody may also be labelled with a predetermined polypeptide epitopes recognized by a secondary reporter (e. g., leucine zipper pair sequences, binding sites for secondary antibodies, metal binding domains, epitope tags). In some
10 embodiments, labels are attached by spacer arms of various lengths to reduce potential steric hindrance.

The antibody may also be derivatized with a chemical group such as polyethylene glycol (PEG), a methyl or ethyl group, or a carbohydrate group. These groups may be useful to improve the biological characteristics of the
15 antibody, e.g. to increase serum half-life or to increase tissue binding. In some aspects, antibodies of the invention relate to h38C2. In some aspects, h38C2 comprises SEQ ID NO:3 and 4 and variants thereof. In this context, "variants thereof" relates to antibodies that comprise a light chain variable region (V_L) comprising a V_L CDR1, V_L CDR2, and V_L CDR3 of the V_L sequence shown
20 in SEQ ID NO:5; and a heavy chain variable region (V_H) comprising a V_H CDR1, V_H CDR2, and V_H CDR3 of the V_H sequence shown in SEQ ID NO:6. Preferably, the h38C2 is an IgG1 antibody. Preferably the h38C2 variant comprises the V_L as set forth in SEQ ID NO:5 and the V_H as set forth in SEQ ID NO:6, and further comprises a light chain constant region at least 95% identical to one or more of
25 SEQ ID NOs:9, 10, 11 and 12, and a heavy chain constant region at least 95% identical to SEQ ID NO:13. In some aspects, the identity of each constant chain independently to one of SEQ ID NO:12 or 13 may be at least 96%, at least 97%, at least 98% or at least 99%. In some aspects, the light chain constant region differs by no more than 5 amino acid residues from one or more of SEQ ID
30 NOs:9, 10, 11 and 12. In some aspects, the light chain constant region comprises

SEQ ID NO:12. In some aspects, the light chain constant region differs by no more than 5, 4, 3, 2, or 1 amino acid from SEQ ID NO:12.

EXAMPLES

The versatility of the invention is illustrated by the following Examples, which illustrate typical embodiments of the invention and are not limiting of the claims or specification in any way.

Example 1 Summary of experimental conditions for the preparation of compound 13

Several routes to generate **13** were explored. In the past, KF on alumina has been employed in several types of transformations to promote a slow deprotonation of the azetidinone, but low conversion to **13** was observed over a 3-day period when this approach was tried (data not shown).

Experimental conditions

1. A first route to generate **13** employed Cl_2SO to make the acid chloride **10** and *n*-BuLi to deprotonate azetidinone **11**. In a first flask, acid **9** was treated with Cl_2SO to prepare acid chloride **10**. Upon reaction completion, excess Cl_2SO was removed by distillation and the acid chloride was dissolved in THF. In a second flask azetidinone **11** was treated with *n*-BuLi in THF at the specified temperature. The acid chloride solution prepared in the first flask was added to the second flask and the mixture warmed to RT. After aqueous workup, the product was purified by chromatography.
2. The second route to generate **13** employed $(\text{COCl})_2$ to make the acid chloride **10** and LiHMDS to deprotonate azetidinone **11**. A similar protocol was employed as for the first route, above.
3. The third route explored used CDI/DIPEA. CDI was added to a solution of acid **9** in CH_2Cl_2 . After 1 hr at RT, 2-azetidinone (**11**) was added and the mixture is stirred at RT for the desired amount of time.
4. The fourth route explored was a combination of T3P and DIPEA. To a solution of acid **9**, 2-azetidinone **11**, and DIPEA in a suitable solvent was added a 50% solution of T3P in EtOAc. The mixture was stirred at the desired temperature for the specified amount of time. Upon reaction completion, an

aqueous workup was performed and compound **13** was purified by recrystallization from a suitable solvent.

	Equiv 11	Base (Equiv)	Other reagents (Equiv)	Temp. (°C)	Solvent	Reaction time (h)	Yield (%)
1	1.2	BuLi (1.3)	Cl ₂ SO (5),	-70 to 0	THF	2	2
1	0.85	BuLi (1.3)	Cl ₂ SO (5)	-70 to 0	THF	2	55
2	1.2	LiHMDS (1.1)	(COCl) ₂ (2)	-30 to 0	THF	18	17
3	1.2	DIPEA (1)	CDI (1.05 equiv)	RT	CH ₂ Cl ₂	1	0
4	1.2	TEA (5)	T3P (1.5 equiv)	RT	THF	1	0
4	1.2	pyridine (7)	T3P (2 equiv)	0	DMF	1	0
4	1.2	DIPEA (5)	T3P (1 equiv)	0	DMF	18	59 (IPA recryst)
4	1.2	DIPEA (5)	T3P (1 equiv)	0	MeCN	18	31 (IPA recryst)
4	1.5	DIPEA (5)	T3P (1 equiv)	2 h @ 0-5, then 18 h RT	MeCN	20	40 (hexanes/IPA recryst)
4	1.35	DIPEA (3)	T3P (1 equiv)	RT	MeCN	18	46 (IPA recryst)
4	1.2	DIPEA (5)	T3P (1 equiv)	RT	MeCN	18	39 (IPA recryst)
4	1.5	DIPEA (5)	T3P (1.5 equiv)	RT	MeCN	3.5	55 (IPA recryst)
4	1.5	DIPEA (5)	T3P (1.5 equiv)	RT	MeCN	2	59-65 (IPA recryst)

Table 1 Summary of experimental conditions for the coupling of acid **9** with 2-azetidinone (**11**).


5 It was found that the preparation of acid chloride **10** using Cl₂SO or oxalyl chloride and the subsequent reaction with the anion of 2-azetidinone prepared via deprotonation with BuLi or LiHMDS led to the desired product **13** but generally gave low yield and purity.

10 The activation of acid **9** with CDI was readily achieved, but the subsequent coupling with 2-azetidinone or its anion failed in solvents such as CH₂Cl₂, THF, and EtOAc, most likely due to the low nucleophilicity of this substrate.

The best results were obtained from the fourth route, using DIPEA/T3P.

Example 2 Synthesis of **13**

53




To a solution of 3-(4-nitrophenyl)propanoic acid (**9**, 440 g, 2.25 mol), 2-azetidinone (**11**, 240 g, 3.37 mol), and DIPEA (1.46 kg, 11.3 mol) in MeCN (4.4 L) was added a 50% (wt/wt) solution of 1-propanephosphonic acid anhydride (T3P) in EtOAc (2.15 kg, 3.38 mol) over 1 hr while the internal temperature was held at 20–25 °C. After the resulting mixture had been stirred at 22°C for 20 hrs, HPLC analysis showed 0.9% of unreacted **9**.

The reaction was concentrated under reduced pressure to about 2 L and the residue was taken up in *i*-PrOAc (6.6 L) at 30–35°C. The organic layer was washed with 10% aqueous citric acid (4 L). The aqueous layer was back-extracted with *i*-PrOAc (2.7 L) and to the combined organic extracts was added Darco G-60 (90 g). The mixture was stirred at 25°C for 4 hrs and filtered through celite. The celite filter was washed with *i*-PrOAc (0.5 L) and the filtrates were concentrated under reduced pressure to about 1 L. 2-Propanol (2.2 L) was added and the distillation was resumed to a final volume of about 1 L. 2-Propanol (2.2 L) was added and the mixture was cooled to 3°C and held at this temperature for 1 hr. The solid was filtered and washed with 2-propanol (0.55 L). The wet cake was transferred back to the reactor and dissolved in EtOAc (3.24 L). Darco G-60 (90 g) was added and the mixture was stirred at 25°C for 2 hrs. The suspension was filtered through celite and the celite pad was washed with EtOAc (0.5 L). The filtrates were concentrated under reduced pressure to about 1 L and 2-propanol (1.4 L) was added. The solution was concentrated to about 1 L and additional 2-propanol (1.4 L) was added. The suspension was cooled to 3°C and held at this temperature for 1 hr. The solid was filtered, washed with 2-propanol (0.5 L), and dried at 30–35°C under vacuum for 8 hrs to give 273 g (49%) of **13**.

HPLC retention time: 2.68 min. HPLC purity: 98.0% (a/a). Mp: 104–106 °C. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 2.88–3.05 (m, 6 H) 3.42 (t, *J* = 5.27 Hz, 2 H) 7.44–7.53 (m, 2 H) 8.07–8.16 (m, 2 H). ¹³C NMR (100 MHz, DMSO-*d*₆) δ

ppm 29.37, 36.05, 36.89, 37.00, 123.83, 130.06, 146.38, 149.49, 166.20, 169.38.
MS (ES⁺): 249 (M + H)⁺.

Example 3 Synthesis of 14b




5 One hypothesis explored was that it was necessary to isolate compound
14 as a solid, in order facilitate long term storage. Accordingly, an improved
process was devised for the generation of a salt form of compound 14;
compound 14b. To a 1-L Atlantis reactor was added 10% palladium on carbon
(Pd/C) (50.00 mg), THF (11.00 mL), and compound 13 (0.5 g). The reactor was
10 sealed and hydrogenated at 10 psi and 25°C for 1hr. HPLC indicates complete
reaction. The catalyst was removed by filtration. Compound 14 was isolated as
a solution. HPLC purity >99%.

Formation and Isolation of HCl salt:

A 190 mL solution from the nitro reduction carried out by containing the
15 free base compound 14 (5.85g) was cooled to 0-5°C. HCl (12 M, 1.5 mL) was
added to give a yellowish suspension. The slurry was concentrated under
vacuum to ~60 mL. Isopropyl acetate (117 mL) was added over 30 min. The
slurry was reconcentrated under vacuum to a final volume of 60 mL. The slurry
was granulated for 30 min at 0-5°C and then filtered. The solids were dried in
20 vacuum oven at 40°C overnight to give 6.6 g of HCl salt 14b.

While an improvement over the previous process, the above method has
the disadvantages of still being relatively time and resource consuming.
Accordingly, the counterintuitive decision was made to isolate compound 14 as a
free base, despite the potential disadvantages of stability of the compound in
25 solution, and the likely concomitant requirement to use the solution immediately.

Example 4 Summary of conditions for preparation of compound 14



Compound **13** was dissolved in THF and 10% Pd/C added. The reaction was then hydrogenated at 10 psig for 16 hrs at about 25°C. Upon reaction completion, the catalyst was filtered off and the filtrates containing aniline **14**

used in the next step (coupling with diglycolic anhydride **15**). In assays, altering the volume of THF (e.g. from about 20 to about 60 volumes) or the reaction time (e.g. from about 1 to about 16 hours) did not alter the yield or purity of the reaction; accordingly these variables were determined to be discretionary.

Example 5 Activated Carbon Screen

10% wt equivalent of 10% palladium and carbon were charged dry under nitrogen to a reactor followed by a solution of starting material in 10 volumes of tetrahydrofuran. A portion of additive was added to each and hydrogenated at 50 psi at RT overnight. The complete reaction was filtered to remove the catalyst and the resulting filtrate analyzed for undesired side products. The results from the carbon screen are shown in Table 2.

Carbon	Recovery	% of 16 by HPLC
CGSP	82%	0.2438
SX-Plus	91%	0.6257
Darco S-51HF	93%	0.44
E Supra USP	91%	0.3332
SX-Ultra	91%	0.3585
CASP	92%	0.5264
Darco G-60	90%	0.2633

Table 2 Levels of impurity **16** after carbon treatment screen.

Example 6 Summary of Reaction conditions for preparing compound 6

To a solution of compound **14** in the corresponding solvent is added diglycolic anhydride (**15**). A base such as DIPEA can be added but it is not necessary. The reaction is stirred for the specified amount of time at the desired temperature. After an aqueous workup, the crude material is purified via slurry in a suitable solvent.

Base	Solvent	Reaction time (h)	Yield (%)
DIPEA (1.1)	CH ₂ Cl ₂	2	68% (hexanes trituration)


equiv)			
DIPEA (1.1 equiv)	CH ₂ Cl ₂	2	50% (IPA trituration)
No base	THF	0.5	57-64% (IPA trituration)
No base	THF	0.5	74-94% (MeOAc trituration)

Table 3 Summary of experimental conditions for the coupling of aniline **14** with diglycolic anhydride (**15**) to produce **6**, using 1 equiv of **15**, at RT.

Example 7 Selection of 7th Solvent

An equal amount of compounds **14** (free base) and **15** were dissolved in 20 volumes of THF (20 mL/g of **14**). The mixture was stirred at RT until reaction completion (about 30 min). The solvent was removed under reduced pressure and the solid residue was triturated in 5 volumes of 7th solvent (one of hexane, 2-propanol, methyl acetate or *i*-propyl acetate). The solids were filtered and dried, and assessed for yield and purity (Table 5). Use of hexane gave a 68% yield of 97% purity. Use of 2-propanol gave only a 50% yield. Use of methyl acetate gave an 80% yield of 96.8% purity. Use of *i*-propyl acetate gave an 80% yield of approximately 95.5% purity.

Example 8 Synthesis of compound **6**




Scheme V

A 30-L hydrogenation reactor was charged with nitro compound **13** (142 g, 0.57 mol), 10% Pd/C (14 g), and THF (8.52 L). Stirring was started and the mixture was hydrogenated at 10 psig and 25°C for 1 hr. After HPLC analysis showed complete consumption of the starting material, the mixture was filtered through a celite-precoated 0.5-micron cartridge filter and the filter was washed with THF (3 L). The filtrates containing aniline **14** were transferred to a 20-L jacketed reactor and diglycolic anhydride (**15**, 75 g, 0.65 mol) was added. After 15 min at 22°C, HPLC analysis of the mixture showed complete consumption of **14** and the reaction was concentrated to 2 L under reduced pressure. The residue was seeded with a small amount of acid **6** crystals (1.06 g) and, after 15 min, *i*-PrOAc (1.78 L) was added over a 1-h period. The suspension was

concentrated under reduced pressure to 1.5 L and the residue was cooled to 3°C. After 1 hr, the solid was filtered, washed with cold *i*-PrOAc (0.5 L) and dried under vacuum at 40°C to give 160 g (84%) of acid **6**. Analysis showed residual THF (0.60% wt/wt) and *i*-PrOAc (0.62% wt/wt).

- 5 This material was reworked as follows to improve color and reduce the amount of residual solvent. Acid **6** (150 g, 0.45 mol) was dissolved in THF (5.25 L) at 30°C. Activated carbon Darco KBB (30 g) was added and the mixture was stirred for 1 hr at this temperature. The suspension was filtered through celite and the celite pad was washed with THF (0.5 L). The filtrates were concentrated
- 10 under reduced pressure to 1.5 L and to the residue was added *i*-PrOAc (3 L) over 30 min. The mixture was held at RT (e.g. about 20°C) for 2 hrs and it was then concentrated under reduced pressure to 1.5 L. The suspension was cooled to 3°C, stirred for 1 hr, and filtered. The solid was washed with *i*-PrOAc (80 mL) and dried under vacuum at 40°C for 12 hrs to give 120 g (80% yield) of **6**.
- 15 Analysis showed residual THF (0.13% wt/wt) and *i*-PrOAc (0.27% wt/wt). HPLC retention time: 2.18 min. HPLC purity: 99.3% (a/a). Mp: 139–140°C. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 2.72–2.90 (m, 4 H) 3.00 (t, *J* = 5.27 Hz, 2 H) 3.38–3.42 (m, 2 H) 3.43 (s, 1 H) 4.12 (s, 2 H) 4.16 (s, 2 H) 7.10–7.16 (m, 2 H) 7.47–7.53 (m, 2 H) 9.77 (s, 1 H) 12.84 (br. s., 1 H). ¹³C NMR (100 MHz, DMSO-
- 20 *d*₆) δ ppm 29.16, 35.98, 36.84, 37.89, 68.54, 70.89, 120.09, 128.95, 136.24, 136.84, 166.20, 168.10. MS (ES⁺): 335 (M + H)⁺.

Example 9 Preparation of compound 6 from compounds 9 and 11




Scheme X

Scheme X: (a) T3P (50% in EtOAc), DIPEA, MeCN, 20 °C, 20 h, 60-68%. (b) H₂ (15 psig), 10% Pd/C (10 wt%), THF, 25°C, 1 h. (c) THF, 22 °C, 15 min, 80% (2 steps)

5 Example 10 Summary of conditions for preparation of compound 20


Acid **6** and pentafluorophenol (**19**) were dissolved in the corresponding solvent. The coupling agent was added and the reaction stirred for the specified amount of time at the desired temperature. After an aqueous workup, crude **20** was purified by either chromatography or slurry in a suitable solvent.

 19 Equivalents	Coupling reagent	Temp. (°C)	Solvent	Reaction time (h)	Yield (%)
1.1	EDC·HCl (1.1 equiv)	0 to rt	CH ₂ Cl ₂	18	77% (IPA slurry)
1	EDC·HCl (1.2 equiv)	0 to rt	DMF	18	62% (IPA slurry)
1.1	DCC (1.05 equiv)	0 to rt	THF	18	65% (after chromatography)
1.1	DCC (1.05 equiv)	0 to rt	THF	18	79% (IPA slurry)

10 **Table 4** Summary of experimental conditions for the coupling of acid **6** with pentafluorophenol (**19**).

Example 11 Synthesis of pentafluorophenol ester **20**

59



Scheme VI

Scheme VI Conditions: DCC, THF, 0°C to RT, 18 hr, 80-85% yield.

To a cold (3–5°C) solution of acid **6** (80 g, 239 mmol) and pentafluorophenol (**19**, 49 g, 266 mmol) in THF (1.1 L) was added a solution of *N,N'*-dicyclohexylcarbodiimide (52 g, 252 mmol) in THF (400 mL) over 10 min. After 15 min, the mixture was warmed to 22°C and stirred for 18 hr. HPLC analysis showed complete reaction. The suspension was filtered to remove the dicyclohexylurea byproduct and the solid was washed with THF (110 mL). The filtrates were concentrated to about 1.8 L and acetone (1.58 L) was added. The suspension was cooled to 10°C and stirred for 1.5 hr. The remaining dicyclohexylurea was filtered off and the solid was washed with acetone (25 mL). The filtrates were concentrated to about 1.8 L and 2-propanol (2.75 L) was added. The slurry was stirred at RT (e.g. about 20°C) for 16 hr and the solid was filtered, washed with 2-propanol (110 mL) and dried under vacuum at 40 °C for 16 hr to give 95 g (83%) of pentafluorophenol ester **20** as a white solid.

HPLC retention time: 3.25 min. HPLC purity: 96.6%. Mp: 94–99°C. ¹H NMR (400 MHz, CDCl₃) δ ppm 2.85–3.07 (m, 7 H) 3.53 (t, *J* = 5.27 Hz, 2 H) 4.26 (s, 2 H) 4.62 (s, 2 H) 7.10–7.22 (m, 2 H) 7.40–7.53 (m, 2 H) 8.47 (s, 1 H). ¹³C NMR (100 MHz, CDCl₃) δ ppm 29.37, 35.83, 36.50, 37.98, 67.89, 69.03, 71.46, 71.83, 120.08, 120.16, 128.97, 128.08, 135.07, 136.87, 165.03, 166.13, 166.24, 170.10. MS (ES⁺): 501 (M + H)⁺.

Example 12 experimental conditions for preparation of compound 3

The peptide **2a** (SEQ ID NO:1) was dissolved in a suitable solvent and the temperature of the resulting solution was adjusted to the desired value.

- Pentafluoro ester **20** and base were added and the reaction stirred for the specified amount of time at the desired temperature. If an anti-solvent was employed to precipitate product **3** from solution, the anti-solvent was added at the same temperature that the coupling between **2** and **20** is run. The mixture was
- 5 then allowed to warm to RT and stirred for 1 hr. Product **3** was isolated by filtration.

20 (equiv)	Base (equiv.)	Solvent (volumes)	Anti-solvent (Vol)	Temp. (°C)	Reaction time (h)	Yield (%)	Comments
3	NMM (20)	DMF (20)	MTBE	20	0.3	Not isolated	High level of impurities
3	NMM (20)	NMP (20)	MTBE	20	0.3	Not isolated	High level of impurities
3	DIPEA (20)	DMF (20)	MTBE	20	0.3	Not isolated	High level of impurities
3	DIPEA(20)	NMP (20)	MTBE	20	0.3	Not isolated	High level of impurities
1	NMM (20)	DMF (20)	MTBE	20	0.3	Not isolated	Product oiled out
1.2	NMM (20)	DMF (20)	EtOAc	20	0.3	Not isolated	Product oiled out
1.5	NMM (20)	DMF (20)	THF	20	0.3	Not isolated	Product oiled out
1	NMM (20)	NMP (20)	MTBE	20	0.3	Not isolated	Product oiled out
1.2	NMM (20)	NMP (20)	EtOAc	20	0.3	Not isolated	Product oiled out
1.5	NMM (20)	NMP (20)	THF	20	0.3	Not isolated	Product oiled out
2	NMM (20)	DMF (20)	MTBE	20	0.3	Not isolated	HPLC purity: 78%
3	NMM (20)	DMF (20)	MTBE	20	0.3	Not isolated	HPLC purity: 84%
3	NMM (20)	MeOH (40)		20			No desired product detected
3	NMM (20)	MeCN (40)		20			No desired product detected
4	NMM (20)	DMSO	2-MeTHF (80)	20	1	Not isolated	HPLC purity: 88%
4	NMM (20)	DMF	2-MeTHF (80)	20	1	Not isolated	HPLC purity: 94%
4	NMM (20)	DMAc	2-MeTHF (80)	20	1	Not isolated	HPLC purity: 90%
4	NMM (1)	DMF (13)/MeCN (13)		20	2		Incomplete reaction
4	NMM (1.3)	DMF (13)/MeCN		20	2		Incomplete reaction


		(13)					
4	NMM (1.5)	DMF (13)/MeCN (13)		20	2		Incomplete reaction
5	NMM (1)	DMF (13)/MeCN (13)		20	2		Incomplete reaction
5	NMM (1.3)	DMF (13)/MeCN (13)		20	2		Incomplete reaction
5	NMM (1.5)	DMF (13)/MeCN (13)		20	2		Incomplete reaction
4	NMM (1.5)	DMF (26)	MeCN (9)	20	0.5	80%	HPLC purity: 99.2%
4	NMM (1.5)	DMF (15)	MeCN (9)	20	0.5	81%	HPLC purity: 92% due to presence of pentafluoroester byproducts not detected with previous HPLC method
2	NMM (1.5)	DMF (15)		0	0.5	Not isolated	Pentafluoroester impurities level below 3%
3	NMM (1.5)	DMF (15)		16	0.5	Not isolated	Pentafluoroester impurities level: 6%
1.2	NMM (1.5)	DMF (15)		-20	2	Not isolated	Pentafluoroester impurities level <1% but incomplete reaction
1.5	NMM (1.5)	DMF (15)		-20	2	Not isolated	Pentafluoroester impurities level <1% but incomplete reaction
2	NMM (1.5)	DMF (15)		-20	2	Not isolated	Pentafluoroester impurities level <1% but incomplete reaction
1.5	NMM (1.5)	DMF (15)		-20	8	Not isolated	Incomplete reaction
3	NMM (1.5)	DMF (15)		-20	4	Not isolated	Complete reaction. Pentafluoroester impurities level: 2.2%
4	NMM (1.5)	DMF (15)		-20	5	Not isolated	Complete reaction. Pentafluoroester impurities level: 3.7%
3	NMM (1.5)	DMF (15)	MeCN (144)	-15	4.5	79	Final conditions. Unreacted starting material: 1.96%. HPLC purity: 97%. Pentafluoroester

							impurities level: 1.19%
--	--	--	--	--	--	--	----------------------------

Table 5 Summary of experimental conditions for the coupling of peptide **2** with pentafluoro ester **20**.

Example 13 Synthesis of peptide-linker conjugate 3

To a cold (-15°C) solution of peptide **2** (100 g, 35.2 mmol) in DMF (0.95 L) was added NMM (4.8 mL, 43.6 mmol) followed by pentafluorophenol ester **20** (50 g, 100 mmol) in small portions over 5 min. The mixture was stirred at -15°C for 7 hrs at which point HPLC analysis showed less than 1% of unreacted **2**. The mixture was filtered through a 0.45 micron in-line filter and added to a second reactor containing MeCN (12.8 L) at RT (e.g. about 20°C) over 5 min. A white precipitate formed immediately. The first reactor was washed with DMF (100 mL) and this wash was added to the second reactor through the in-line filter. The slurry was stirred at RT (e.g. about 20°C) for about 1 hr and the solid was filtered, washed with MeCN (3×1 L), and dried under vacuum at RT (e.g. about 20°C) for 6 hr to give 94 g (86%) of **3** as a white solid. GC headspace showed residual DMF (4.5% wt/wt) and MeCN (0.95% wt/wt). With the goal of reducing solvent content to meet specifications, the solid was passed through a #20 hand-sieve and then slurried in MeCN (10 L) at about RT for 1 hr with good agitation. The solid was filtered, washed with MeCN (2×1 L) and dried at 30°C for 24 hrs and at 40°C for an additional 24 hrs to give 91 g of **3** (97% recovery, 83% overall yield). Residual DMF (0.05 % wt/wt) and MeCN (0.19% wt/wt). HPLC purity: 96.5% (a/a). HPLC retention time: 17.99 min. MS (ES⁺): 3156.6 Da ($M + H$)⁺.



Scheme VII: Final conjugation and isolation conditions for peptide-linker conjugate **3**. Conditions: NMM (1.3 equiv), DMF (15 vol), -15 to -18°C , 7 h. Filtration to remove insolubles (0.45 micron). Drip in filtrates into MeCN (~ 144 vol) to precipitate product. Final composition: MeCN/DMF 9/1. Wash filter with

DMF (1 vol). Stir at RT for 1-2 hrs. Filter under N₂. Wash cake with 100% MeCN (3 X 10 vol). Dry at RT (e.g. about 20°C). Reslurry in MeCN (100 vol) to remove residual DMF. Dry 1 at 40°C under vacuum Yield: 83%.


Example 14 Conjugation of acid 6 with peptide 2 to form conjugate 3

5 Acid 6 was originally activated as its *N*-hydroxysuccinimido ester prior to coupling with peptide 2. This approach was explored by treating acid 6 with *N*-hydroxysuccinimide (7) using *N,N'*-dicyclohexylcarbodiimide (DCC) as coupling reagent to generate intermediate 8. It proved difficult to isolate intermediate 8 by crystallization.

10 In addition, efforts to isolate intermediate 8 by chromatography on silica demonstrated signs of instability of intermediate 8 when in contact with silica were observed by TLC analysis.

Even though 8 was isolated as a foamy solid in an excellent 92% yield after chromatography, both HPLC and ¹H NMR analyses showed a mixture of
15 two major components. One of them was identified as desired 8 but increasing amounts of the second component were noticed after the solid had been set aside for only a few hours.

Further investigation resulted in the identification of morpholine-3,5-dione
18 as the second component by LC-MS, which formed via intramolecular
20 displacement of the *N*-hydroxysuccinimido group by the anilide nitrogen on the molecule.



This cyclic byproduct became the major constituent of the isolated solid after only a few days at RT. Since this side-reaction was envisioned to become even
25 more problematic on scale due to the longer timelines for processing and isolation, a search for an alternative activating group for acid 6 was undertaken.



Based on the information obtained from experimentation related to the *N*-hydroxysuccinimido ester, it became clear that a fine balance between reactivity

toward the amino group on the lysine residue of peptide **2** and stability to prevent morpholine-3,5-dione **18** formation was necessary.

Example 15 Improving the impurity profile

The reaction between peptide **2** and compound **20** was very fast (<30 min) when run at RT (e.g. about 20°C) and the purities of the isolated material were in the 90–95% range. Up to 4 major impurities were detected that caused a drop in purity compared to the purity of the peptide **2** starting material.

These 4 byproducts were characterized as pentafluorophenol ester derivatives of peptide-linker conjugate **3** based on MS data (all of them displayed the same mass) and, when combined together, they amounted to 7–8%. Interestingly, each impurity is a distinct monoester corresponding to the four different carboxylic acid groups on the peptide backbone and no impurities with multiple esters were detected. Therefore, it appears that once a pentafluorophenol monoester derivative forms, further esterification on that same molecule is extremely slow. The structures of these impurities are shown below.



This fact is difficult to rationalize based on the existing physical separation between the acid groups. It was therefore desirable to minimize their presence in order to obtain $\geq 95\%$ purity specification for peptide-linker conjugate **3**. A major driving force for this was that any attempts to upgrade the chemical purity of peptide-linker conjugate **3** by trituration or recrystallization had failed due to the extremely low solubility of peptide-linker conjugate **3** and related byproducts.

A final study was carried out to determine the stability of the pentafluorophenol ester impurities that were identified during the peptide-linker conjugation reaction. It was reasoned that these byproducts would also couple to the monoclonal antibody during the final process to prepare the drug substance


and might complicate the final purification of **5**. As a result, a sample of peptide-linker conjugate **3** with high pentafluorophenol ester impurities levels was placed in the same 50:50 20 mM histidine/propylene glycol buffer at pH 6.5 that is employed to conjugate peptide-linker **3** with monoclonal antibody **4**. The results are shown in Table 6. After only 30 min, one of the three impurities was not detectable and the levels of the other two had decreased considerably. At the same time, the overall purity had increased by about the same amount. After 60 min, the remaining 2 had almost completely vanished and the purity of **3** stayed unchanged. This result seems to indicate that these impurities are short-lived in this aqueous medium and revert back to the desired product **3**.

	Time point (Minutes)				
	0	30	60	90	120
Purity (1, %)	95.6	96.5	96.5	96.6	96.7
Impurity 1 (%)	0.75	0.26	0.09	n/d	n/d
Impurity 2 (%)	0.53	n/d	n/d	n/d	n/d
Impurity 3 (%)	0.68	0.14	0.05	n/d	n/d

Table 6 Pentafluorophenol ester impurities time-dependent stability study in 50:50 20 mM histidine/propylene glycol buffer at pH 6.5. n/d: not detected.


Example 16 Improved process for preparing 3

66




Scheme XI: (a) T3P (50% in EtOAc), DIPEA, MeCN, 20 °C, 20 h, 60-68%. (b) H₂ (15 psig), 10% Pd/C (10 wt%), THF, 25°C, 1 h. (c) THF, 22 °C, 15 min, 80% (2 steps). (d) DCC, THF, 0 °C to rt, 18 h, 80-85%. (e) NMM (1.3 equiv), DMF (15 vol), -15 to -18 °C, 7

5 Example 17 Preparation of conjugate 3 using PNP esters



Scheme XII



Scheme XIII

To a solution of compound **6** (0.5 g, 0.0015 moles, 1 equiv.) in THF (10 mL) compound **21** (p-nitrophenol) (0.23 g, 0.00165 moles, 1.1 equiv.) was added. The mixture was stirred at 0-5°C. The mixture was slightly cloudy. To this mixture DCC (0.325 g, 0.0016 moles, 1.05 equiv.) was added. The mixture was stirred at RT overnight. Next day the slurry was filtered through buchner funnel to remove precipitated DCU. The filtrate was evaporated to dryness on a rotary evaporator. The residue was slurried in 8 mL of isopropanol and stirred at 0-5°C for 1 hr. The

product slurry was filtered, washed with 5 mL of isopropanol and dried under high vacuum overnight, to provide a yield of 0.531 g (68 %) of compound **22**.

Compound **22** (0.3 g, 0.00066 moles, 3 equiv.) was placed in 35 mL round bottom flask. Solids were dissolved in 5 mL of DMF. To the solution of peptide **2a** (SEQ ID NO:1) free-base (0.624 g, 0.000225 moles, 1 equiv.) was added as a solid. The reaction mixture was stirred at 0-5°C for 8 hr. At the end of reaction period, HPLC analysis of the reaction mixture indicated that all of peptide **2a** had disappeared. The mixture was added slowly in a dropwise manner to 150 mL of acetonitrile. The precipitated product slurry was stirred at room temperature for 3 hrs. The slurry was filtered through Buchner funnel using Whatman # 2 filter paper. The slurry was filtered within 10 min. The product cake was washed with 15 mL of acetonitrile. The product compound **3d** was dried under high vacuum for 16 hrs to afford a white product; compound **3d**, in a yield of 0.612 g (86 %).

Example 18 Summary of reaction completions and product purities

1. Reaction completion and product purity for compounds **6**, **13**, **14**, and **20** were evaluated by HPLC using the following conditions: Column: Zorbax SB-CN 3.5 μ m, 3 $\times$ 75 mm. Column temperature: 45°C. Detection: UV @ 210 nm. Mobile phase: A: water (0.05% TFA); B: MeCN. Gradient: 0 min: 95/5; 3.7 min: 5/95; 4.3 min: 5/95; 4.4 min: 95/5. Flow: 1.2 mL/min.
2. Reaction completion for peptide-linker conjugate **3** was evaluated by HPLC using the following conditions: Column: XBridge BEH130 C18 3.5 μ m, 4.6 $\times$ 150 mm. Column temperature: 30°C. Detection: UV @ 210 nm. Mobile phase: A: 60/40 water/MeOH (0.1% formic acid); B: 60/40 MeCN/MeOH (0.085% formic acid). Gradient: 0 min: 95/5; 3 min: 95/5; 20 min: 60/40; 23 min: 0/100; 25 min: 0/100; 25.1 min: 95/5; 30 min: 95/5. Flow: 1 mL/min.
3. Product purity for compound **3** was evaluated by HPLC using the following conditions: Column: YMC-Pack ODS-A, 250 $\times$ 4.6 mm I.D., S-5 μ m 12 nm, P/N AA12S05-2546WT. Column temperature: 60°C. Autosampler temperature: 5°C. Injection volume: 25 μ L. Detection: UV @ 220 nm. Mobile phase: A: 0.1 M NaClO₄, pH adjusted to 3.1 with H₃PO₄; B: 0.1% TFA in ACN. Dissolving solvent: 1:1 Water / Dimethylformamide. Sample concentration: 1

mg/mL. Gradient: 0 min: 73/27; 2 min: 73/27; 32 min: 70/30; 42 min: 50/50; 42.1 min: 73/27; 50 min: 73/27 min. Flow: 1.5 mL/min.

4. ^1H NMR and ^{13}C NMR spectra were recorded on a 400 MHz spectrometer in either CDCl_3 or $\text{DMSO}-d_6$ as both solvent and internal standard.

- 5 Mass data for compounds **6**, **13**, **14**, and **20** was obtained on a Agilent 1100 Series LC/MSD SL spectrometer (ESI). Mass data for **3** was obtained using a MicroMass Q-ToF Global mass spectrometer (ESI).

Example 19 Conjugation of 3b with h38C2 antibody

- Conjugation of **3b** was performed at a temperature range of between
10 about 5°C to about 35°C. The solution of **3b** in a mixture of cosolvent and histidine buffer was added to a solution of h38C2 antibody (SEQ ID NO:3 and SEQ ID NO:4) in the buffer mixture containing histidine, glycine and sucrose. The conjugation was performed for about 2-about 24hr. At the end of conjugation
15 reaction the mixture was filtered through a 0.2 micron filter. The solution was then passed through Q-membrane filter to remove remaining residual peptide **3b**. The solution was then concentrated and diafiltered through UF-DF membrane using histidine/glycine buffer solution at the desired pH; preferably pH 6.5. The concentrated solution of peptide-linker-antibody **5** was then diluted with
20 histidine/glycine buffer containing polysorbate 20 and sucrose to the desired concentration; preferably to about 20 mg/mL.

Example 20 Change in the cosolvent of the Conjugation Process

- The process for the formation of the drug substance **5** was originally described in US8288349 whose contents are hereby incorporated. The original process involved use of propylene glycol as the cosolvent in the conjugation
25 process between peptide-linker **3** and the antibody h38C2. Peptide-linker **3b** has very low solubility in the solvent propylene glycol. Also, compounds of the type **3**, in particular **3b**, exist in two different solid forms, one being completely amorphous and the other solid form exhibits partial partial crystalline morphology. Both solid forms exhibit different solubility characteristics in different solvents and
30 buffers. The more amorphous solid form is usually more soluble in different solvents and buffer solutions. However, it is difficult to control conditions of the

process for preparation of **3b** that would consistently result in the formation and isolation of only the solid form with higher solubility in the conjugation mixture. The manufacturing process of **3b** typically affords the product as a mixture of two different solid forms. This renders the conjugation mixture during preparation of **5** heterogeneous, making kinetics of the conjugation reaction variable. During preparation of bioconjugate **5**, it is desirable to have a conjugation mixture as a homogeneous solution at the beginning of the reaction in order to afford consistent reaction kinetics and more robust manufacturing process.

Surprisingly it was found that with DMSO as solvent, both the solid forms of **3b** were readily soluble at ambient temperature at the desired concentration of 12 mg/mL and up to 100 mg/mL. In solvents other than DMSO such as acetonitrile and methanol, the more crystalline solid form had limited solubility, as shown in Table 7.

Solvent	Solubility (mg/ml)
Acetonitrile	<0.01
Methanol	<0.01
Acetone	<0.01
DMSO	>100
THF	0.01
Acetonitrile/Water (2:1 v/v)	0.39
Acetonitrile/Water (1:1 v/v)	0.73
THF/Water (2:1 v/v)	0.12
Acetone/Water (2:1 v/v)	0.43
Methanol/Water (2:1 v/v)	

Table 7 Comparison of solubility of **3b** in different solvents.

In the original process, propylene glycol was used as a cosolvent primarily to assist in the dispersion of the hydrophobic peptide-linker **3b** in the aqueous histidine buffer solution at pH 6.5. In the propylene glycol/histidine buffer mixture, both solid forms of **3b** dissolve slowly over period of 2-3 hours albeit at different rates. This renders the conjugation mixture at the beginning of the conjugation reaction heterogeneous, making kinetics of the conjugation reaction variable. During preparation of bioconjugate **5**, it was found to be advantageous to have a conjugation mixture as a homogeneous solution at the start of the reaction in

order to afford predictable reaction rate. It was discovered that replacing propylene glycol with DMSO (up to 10% v/v) as cosolvent in the conjugation process of **5** resulted the reaction mixture that was homogeneous. This in turn afforded more consistent and predictable reaction rate of the conjugation process and a more robust manufacturing process for formation of **5**.

The h338C2 used in the process consists of a mixture of mAb species that contain fully conjugatable mAb, partially conjugatable mAb, and non-conjugatable mAb. The conjugation of h38C2 with the peptide **3b** affords **5** as a mixture of conjugate containing fully conjugated two loaded (+2) species, partially loaded species (+1) and unconjugatable mAb (0 loaded species).

In order to investigate the impact of different conjugation reaction variables on the reaction rate, product quality and yield, a set of experiments were executed using design of experiments (DOE) approach. Three parameters were chosen for evaluation: pH of the conjugation mixture, temperature of the conjugation and the ratio of peptide to starting antibody. Other parameters such as composition of the buffer solution (20 mM histidine buffer), antibody concentration (16.7 mg/mL) and concentration of DMSO in solution (7 %) were kept constant. The centre point for each parameter was pH6.5, 20°C and a peptide:antibody ratio of 2.4:1. The low levels tested were pH5.5, 5°C and a ratio of 1.8, and the high levels were pH7.5, 35°C and a ratio of 3.

Several experiments whose variable values were at the center-point of all the parameter ranges were selected to provide information on possible non-linear effects. The protocol for these experiments was briefly as follows: Peptide **3b** (25 mg) was placed in a 20mL vial equipped with a magnetic stir bar. The solid was dissolved in DMSO (2.1 mL). Histidine buffer (20 mM, pH 6.5, 2.1 mL) was added to the solution. The mixture was stirred for 90 mins at 20°C. The solution of antibody h38C2 (25.13 mL, 19.9 mg/mL) was added to a separate jacketed reactor. To this solution was added the solution of peptide **3b** in DMSO / histidine buffer with a syringe over 2-3 min period. The mixture was stirred at 20°C. The reaction was monitored for completion by SEC and HIC chromatography methods. The protocol for the remaining tests as identical, except in the

respective pH, temperature and ratio of peptide:antibody. The in-situ yield of 5 and time of reaction completion (hours) were evaluated by HIC and SEC chromatographic methods. Results of the DOE experiment are shown in Table 8.

pH	Temp.	Peptide Ratio	% +2 by HIC	% +1 by HIC	% 0 by HIC	% Peptide by Org. SEC	Reaction Compl. (hr)
5.5	35	3	81.7	17.3	0.9	10.1	1
			83.6	15.7	0.7	-	3
			83.9	15.4	0.7	10.2	5
5.5	5	3	20.3	48.5	31.2	15	1
			37.4	47.2	15.3	13	3
			40.5	47	12.5	9.4	5
			73	25	2	8.9	16
5.5	35	1.8	57.8	38.3	3.9	0.6	1
			58.4	37.6	4	0.4	2
			57.8	38.1	4.1	0.2	3
			62.4	35.3	2.3	0	16
5.5	5	1.8	25.5	49.8	24.7	9.2	1
			37.1	53.4	9.5	3.2	3
			53.3	39.6	7.1	1.2	5
			58	39.4	5.5	0.8	7
			61.8	33.8	4.4	0.6	11
6.5	20	2.4	82	17	1	5.7	2
			83.4	16.5	1	4.9	4
			89.6	10.4	0	3.4	24
6.5	20	2.4	77.9	20.7	1.3	4.6	1
			82.9	16.3	0.8	3.7	3
			83.5	15.7	0.8	3.4	5
7.5	35	3	83	16.5	0.5	10.3	0.75

			85.5	14	0.5	9.6	3
7.5	5	3	47.3	43.3	9.4	14.7	1
			75.6	22	1.6	9.9	3
			80.8	18.2	1	9	5
			81.9	17.2	0.9	8.8	6
7.5	35	1.8	69.1	28.2	2.7	0	1
			69.5	28	2.5	0	3
			69.5	28.1	2.4	0	5
7.5	5	1.8	64.7	32.1	3.2	0.1	2
			68.3	28	3.6	0.1	4
			69.6	27.2	3.2	0.1	24

Table 8 DOE for conjugation parameters.**Conclusions from the DOE studies are as follows:**

Within the given design space for the conjugation:

- 5 - At a given pH, increased yield of +2 is obtained at higher temperature and higher peptide: mAb ratio.
- At a given peptide ratio, higher yield of +2 is obtained at higher pH and higher temperature.
- At given temperature, higher pH and higher peptide ratio affords higher
- 10 yield of +2.
- pH of the conjugation mixture has maximum effect on the reaction rate. At low pH the conjugation reaction is slower.

Within the reaction design space as defined by parameters and their range limits, the optimum yield of +2 of the conjugate is obtained within the standard

15 reaction time of 3 hrs when the peptide ratio is between 2.1–3 equiv., pH is 6.5–7.5 and temperature is between 20–30°C. However the desired product can still be obtained within the broader range limits for pH (5.5–7.5) and temperature (5–35°C) as long as peptide ratio is between 1.8–3 equiv. and the conjugation

 reaction is allowed to proceed to its completion. To minimize cost of goods and

20 maximize productivity, the conjugation reaction is normally performed with peptide/antibody ratio of 2.1 at RT, (for example, 22 °C) and pH of 6.5. At higher


temperatures (e.g. 30–35°C) and at higher pH (e.g. about pH7.5) the degradation of peptide **3b** occurs at a faster rate. Addition of significant excess of peptide **3b** to the conjugation mixture can potentially lead to formation of more process related impurities, particularly at high pH conditions. Therefore, while the process can operate within the defined parameters, it is preferable to operate within the optimized ranges.

In some aspects, the invention provides for a method of conjugation of peptide-linker to an antibody as herein described, comprising


- (i) dissolving compound **3b** in DMSO; preferably at between about 5 and about 100 mg/ml, more preferably at between about 10 and about 15mg/ml, most preferably at about 12mg/ml; preferably for a duration of at least about 1min, at least about 5 mins, and preferably at least about 15 mins (as the reaction is stable, there need not be an upper limit to the duration);
- (ii) adding histidine buffer (as described herein and below) (pH between about 5.5 to about 7.5), preferably about pH 6.5 to the solution of compound **3b** made in (i); preferably about an about equal volume, preferably to between about 2 and about 10mg/ml, more preferably at between about 5 and about 8mg/ml, most preferably at about 6mg/ml;
- (iii) Adding h38C2 antibody;
- (iv) Agitating at a medium speed so as to avoid foaming the reaction mixture for at least about 1hr, preferably at about between about 15°C to about 35°C, more preferably at RT, and most preferably at about 22°C;
- (v) Filtration of the solution from (iv).

Accordingly, in some aspects, the invention provides for a process for conjugating a peptide of formula **3b** to an antibody comprising the steps:

- (i) dissolving compound **3b** in DMSO:



- (ii) adding histidine buffer at pH between about 5.5 to about 7.5 to the solution of DMSO and **3b** of step (i);
- (iii) Adding an antibody comprising a variable light region comprising SEQ ID NO:5 and a variable heavy region comprising SEQ ID NO:6 to the solution of step (ii), so as to have a peptide:antibody molar ratio of between about 1.8:1 to about 3:1;
- (iv) Agitating the mixture formed in step (vi) at a medium speed so as to avoid foaming the reaction mixture for at least about 1hr at between about pH 5.5 and about pH7.5 and at a temperature of between about 5°C and 35°C;
- (v) Filtration of the solution from (iv) to extract the resultant peptide-linker antibody conjugate **5b**.



wherein $n = 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29$ or 30 , $q = 1, 2, 3, 4$, or 5 .


- In some aspects, the antibody may be in a solution comprising between about 1mM to about 100mM histidine, preferably about 5 to about 20mM, more preferably about 10 mM histidine; between about 1 and about 100 mM glycine, preferably between about 1 and about 20mM glycine, more preferably about 10mM glycine buffer; and further comprising between about 0.1 and about 10% (w/w) sucrose, preferably between about 0.5% and about 5% (w/w) sucrose, more preferably between about 1% and about 3% (w/w) sucrose, and most preferably about about 2% (w/w) sucrose. Preferably the final concentration of the antibody in the conjugation mixture may be between about 10 and about 30 mg/ml, more preferably between about about 14 and about 20 ml, most preferably about 16.7 mg/mL.

In some aspects, the agitation may be done in a stainless steel reactor with bottom mounted magnetic coupled agitator (6 bladed with 6" diameter blades) at 70 rpm or in a glass reactor using overhead mechanical agitator with



Teflon paddle at 120 rpm speed. The reaction mixture may be agitated for at least about 1hr, more preferably at least about 2hr, more preferably between about 1hr and overnight, further preferably between about 2hr and about 6hr, and most preferably about 3hr.

- 5 In some aspects, the filtration is by a 0.2 μ m filter; following which, the filtrate may be subjected to Q-filtration. Following this, the filtrate may be subjected to a UF/DF filtration, such as through a Hydrosart ultrafiltration membrane with 30 Kilodalton Molecular Weight Cut off with the buffer containing 20 mM histidine and 200 mM glycine at pH 6.5, and formulation with polysorbate 20 (0.1 % w/w) and sucrose (2 % w/w) followed by further 0.2 μ m filtration.
- 10 20 (0.1 % w/w) and sucrose (2 % w/w) followed by further 0.2 μ m filtration.

In some aspect, the above reaction relates to compound **3b** and compound **5b**. In some aspects of the above, $n=1$, and the process relates to compound **3a** and compound **5a**. In some aspects, $n=1$ and $q=1$, and the process relates to compound **3** and compound **5**.




76



5


Example 21 Improved process for preparing 5 from 9 and 11

77



Scheme XV: (a) T3P (50% in EtOAc), DIPEA, MeCN, 20°C, 20 h, 60-68%. (b) H₂ (15 psig), 10% Pd/C (10 wt%), THF, 25°C, 1 h. (c) THF, 22 °C, 15 min, 80% (2 steps). (d) DCC, THF, 0°C to rt, 18 h, 80-85%. (e) NMM (1.3 equiv), DMF (15 vol), -15 to -18°C, 7. (f) agitation at 22°C, pH6.5,

Example 22 Improved process for preparing 5 from 3 and 4



Scheme XVI

Scheme XVI: (f) agitation at 22°C, pH6.5,

The invention thus has been disclosed broadly and illustrated in reference to representative embodiments described above. Those skilled in the art will recognize that various modifications can be made to the present invention without departing from the spirit and scope thereof. All publications, patent
5 applications, and issued patents, are herein incorporated by reference to the same extent as if each individual publication, patent application or issued patent were specifically and individually indicated to be incorporated by reference in its entirety. Definitions that are contained in text incorporated by reference are excluded to the extent that they contradict definitions in this disclosure.

10 It is appreciated that certain features of the invention, which are, for clarity, described in the context of separate embodiments, may also be provided in combination in a single embodiment. Conversely, various features of the invention which are, for brevity, described in the context of a single embodiment, may also be provided separately or in any suitable sub-combination.

15 It is specifically contemplated that any limitation discussed with respect to one embodiment of the invention may apply to any other embodiment of the invention. Furthermore, any composition of the invention may be used in any method of the invention, and any method of the invention may be used to produce or to utilize any composition of the invention. In particular, any aspect of
20 the invention described in the claims, alone or in combination with one or more additional claims and/or aspects of the description, is to be understood as being combinable with other aspects of the invention set out elsewhere in the claims and/or description.

The use of the term "or" in the claims is used to mean "and/or" unless
25 explicitly indicated to refer to alternatives only or the alternative are mutually exclusive, although the disclosure supports a definition that refers to only alternatives and "and/or."


As used herein the specification, "a" or "an" may mean one or more, unless clearly indicated otherwise. As used herein in the claim (s), when used in
30 conjunction with the word "comprising," the words "a" or "an" may mean one or more than one. As used herein "another" may mean at least a second or more.

The words “comprises/comprising” and the words “having/including” when used herein with reference to the present invention are used to specify the presence of stated features, integers, steps or components but does not preclude the presence or addition of one or more other features, integers, steps,

5 components or groups thereof.


CLAIMS

1. A process for preparing a compound according to formula 5a




comprising


- (i) reacting 9a and 11 together in the presence of 1-propanephosphonic acid anhydride (T3P) to create compound 13a




- (ii) catalytic hydrogenation with Pd/C of compound 13a in a THF:H₂O solution of at least about 50% THF to produce a compound of formula 14a




- (iii) combining a solution of 14a in THF with a compound according to formula 15 in a reaction substantially free of base to produce compound 6a



- (iv) reacting a compound according to the formula 6a with a compound according to formula 19a in THF in the presence of DCC to produce compound 20a




- (v) combining **20a** with a ϵ -amino containing peptide **2** dissolved in an aprotic polar ^{15th} solvent to produce compound **3a**



- (vi) dissolving compound **3a** in DMSO;
- (vii) adding histidine buffer at pH between about 5.5 to about 7.5 to the solution of DMSO and **3a** of step (vi);
- (viii) Adding an antibody comprising a variable light region comprising SEQ ID NO:5 and a variable heavy region comprising SEQ ID NO:6 to the solution of step (vii), so as to have a peptide:antibody molar ratio of between about 1.8:1 to about 3:1;
- (ix) Agitating the mixture formed in step (viii) at a medium speed so as to avoid foaming the reaction mixture for at least about 1hr at between about pH 5.5 and about pH7.5 and at a temperature of between about 5°C and 35°C; and
- (x) Filtration of the solution from (ix) to extract the resultant peptide-linker antibody conjugate **5a**


wherein $q = 1, 2, 3, 4$, or 5 , $X = F$ or Cl , and $m = 3, 4$, or 5 .

2. The process as claimed in claim 1, further comprising a process for preparing a compound according to Formula **3a**




comprising


- (i) reacting **9a** and **11** together in the presence of 1-propanephosphonic acid anhydride (T3P) to create compound **13a**




- (ii) catalytic hydrogenation with Pd/C of compound **13a** in a THF:H₂O solution of at least about 50% THF to produce a compound of formula **14a**




- (iii) combining a solution of **14a** in THF with a compound according to formula **15** in a reaction substantially free of base to produce compound **6a**



- (iv) reacting a compound according to the formula **6a** with a compound according to formula **19a** in THF in the presence of DCC to produce compound **20a**




- (v) combining 20a with a ϵ -amino containing peptide 2 dissolved in an aprotic polar ^{15th} solvent to produce bioconjugate 3b




wherein $q = 1, 2, 3, 4$, or 5 , $X = F$ or Cl , and $m = 3, 4$, or 5 .

3. The process as claimed in claim 1, further comprising a process for preparing a compound of formula 13a




wherein q is $1, 2, 3, 4$, or 5 , characterized in that




are reacted together in the presence of 1-propanephosphonic acid anhydride (T3P).

4. The process as claimed in claim 1, wherein the reaction between 9a and 11 is carried out in the presence of a 1st solvent selected from the group consisting of tetrahydrofuran (THF), 2-methyltetrahydrofuran, 1-methyl-2-pyrrolidinone (NMP), N,N-dimethylformamide (DMF), N,N-dimethylacetamide (DMAc), 1-methyl-2-pyrrolidinone, ethyl acetate (EtOAc), and acetonitrile (MeCN), and preferably MeCN.
5. The process as claimed in claim 4, wherein the reaction is carried out in the presence of a 1st base selected from the group consisting of trimethylamine, triethylamine, tributylamine, DIPEA, pyridine, DBU, DABCO, 2,3-lutidine, 2,4-lutidine, 2,5-lutidine, 2,6-lutidine, 3,4-lutidine, 3,5-lutidine, and preferably DIPEA.

6. The process as claimed in claim 1, further comprising a process for preparing a compound of formula **4a**,




wherein q is 1, 2, 3, 4, or 5, comprising catalytic hydrogenation with Pd/C of compound **13a**,




and characterized in that the reaction is conducted in a THF:H₂O solution of at least about 50% THF.

7. The process as claimed in claim 1, wherein the THF:H₂O solution comprises THF in an amount selected from the group consisting of at least about 60%, at least about 70%, at least about 80%, and at least about 90%.
8. The process as claimed in claim 1, wherein prior to hydrogenation in THF/H₂O, compound **13a** is subjected to treatment with activated charcoal.
9. (Original) The process as claimed in claim 8, wherein the activated charcoal is selected from the group consisting of SX-Plus, Darco® S-51HF, E Supra USP, SX-Ultra, CASP, Darco® G-60 and CGSP.
10. The process as claimed in claim 1, further comprising a process for preparing a compound of formula **6a**



wherein q = 1, 2, 3, 4 or 5; comprising reacting a solution of compound **14a**




in a 6th solvent comprising THF,



and characterized in that the reaction is carried out substantially free of a base.


11. The process as claimed in claim 1, wherein the compound **14a** is reacted with between about 1 and about 10 equivalents of compound **15**.
12. The process as claimed in claim 1, wherein the solution of compound **6a** is distilled under vacuum, following which a 7th solvent comprising C₁-C₄ alkyl acetate is added to attain a solvent composition of between about 25% THF/75% C₁-C₄ alkyl acetate and about 100% C₁-C₄ alkyl acetate.
13. The process as claimed in claim 12, wherein the 7th solvent is selected from the group consisting of methyl acetate, ethyl acetate, *i*-propyl acetate, *n*-propyl acetate, *n*-butyl acetate, and preferably is *i*-propyl acetate.
14. The process as claimed in claim 12, wherein the ratio between 6th and 7th solvents is between 25:75 and 0:100.
15. The process as claimed in claim 1, further comprising a process for crystallization of a compound according to the formula **6a**, comprising
 - (i) dissolving acid **6a** in an 8th solvent comprising THF;
 - (ii) optionally treating with activated carbon and then filtering off said activated carbon;
 - (iii) concentrating the acid **6a** in THF solution to between about 2 and about 20 vol;
 - (iv) adding between about 1 and about 50 vol of a 9th solvent consisting of 2-propanol;
 - (v) concentrating the solution of acid **6a** in THF and 2-propanol to between about 2 and about 50 vol;
 - (vi) cooling the concentrated solution of acid **6** to between about -25°C and about 10°C.

16. The process as claimed in claim 1, further comprising a process for preparing a compound according to the formula 20b comprising reacting a compound according to the formula 6b with a compound according to formula 19a in a 10th solvent in the presence of DCC



wherein $q = 1, 2, 3, 4$, or 5 , $n = 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29$ or 30 , $X = F$ or Cl , and $m = 3, 4$, or 5 .

17. The process of claim 1, further comprising a process for conjugating an ϵ -amino containing peptide 2 to a linker of the formula 6b comprising:
- dissolving peptide 2 in an aprotic, polar 15th solvent,
 - reacting 6b with between about 1 and about 10 equivalents of 19a in THF to produce compound 20b;
 - combining 20b with peptide 2 to produce bioconjugate 3b



wherein n = 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29 or 30, q = 1, 2, 3, 4, or 5, X= F or Cl, and m=3, 4, or 5.

18. The process as claimed in claim 1, wherein at least about 1 equivalent of compound **19a** is added to compound **6a** or **6b**.
19. The process as claimed in claim 1, wherein the DCC is added to the reaction at a temperature of between -10°C and +10°C, and preferably between about +3°C and +5°C.
20. The process as claimed in claims 1, wherein the reaction between compound **19a** and either **6a** or **6b** takes place in a 10th solvent selected from the group consisting of THF, dichloromethane, 2-methyltetrahydrofuran, N,N-dimethylformamide, and N,N-dimethylacetamide, and preferably THF.
21. The process as claimed in claim 1, wherein X is F, and m is 5.
22. The process as claimed in claim 1 wherein compound **3a** or compound **3b** is dissolved in DMSO at between about 5 and about 100 mg/ml, and preferably at between about 10 and about 15 mg/ml.
23. The process as claimed in claim 1, wherein the histidine buffer is added to the DMSO solution of compound **3a** or **3b** at an equal volume, plus or minus 10%.
24. The process as claimed in claim 1, wherein the concentration of compound **3a** or compound **3b** following addition of the histidine buffer is between about 2 and about 10mg/ml, and is preferably about 6mg/ml.
25. The process as claimed in claim 1, wherein the antibody comprises a light chain constant region comprising a sequence selected from the group consisting of SEQ ID NOs:9, 10, 11, and 12, or variants thereof comprising between 1 and 5 amino acid substitutions, deletions or insertions, and a heavy chain constant region comprising a sequence selected from the group consisting of SEQ ID NOs:13, and 14, or variants thereof comprising between 1 and 5 amino acid substitutions, deletions or insertions.
26. The process as claimed in claim 25, wherein the antibody comprises a light chain comprising SEQ ID NO:3, or variants thereof comprising between 1 and 5 amino acid substitutions, deletions or insertions, and a heavy chain comprising SEQ ID NO:4, or variants thereof comprising between 1 and 5 amino acid substitutions, deletions or insertions.

27. The process as claimed in claimed 1, wherein $q=2$, n when present =1, m when present = 5, and x when present =F.