PCT WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 7 : C12N 15/82, 9/10

A1

(11) International Publication Number: **WO 00/34490**

(43) International Publication Date: 15 June 2000 (15.06.00)

(21) International Application Number: PCT/JP99/06881

(22) International Filing Date: 8 December 1999 (08.12.99)

(30) Priority Data:
10/350584 9 December 1998 (09.12.98) JP

(71)(72) Applicants and Inventors: SEKI, Tatsuji [JP/JP]; 5–13–53–209, Mino, Mino–shi, Osaka 562–0001 (JP). FUJIYAMA, Kazuhito [JP/JP]; A18–308, 1–28, Yamadanishi, Suita–shi, Osaka 565–0824 (JP).

(72) Inventor; and
(75) Inventor/Applicant *(for US only)*: YOSHIDA, Toshiomi [JP/JP]; A1–505, 2–4, Yamadanishi, Suita–shi, Osaka 565–0824 (JP).

(74) Agent: YAMAMOTO, Shusaku; Crystal Tower, 15th Floor, 2–27, Shiromi 1–chome, Chuo–ku, Osaka–shi, Osaka 540–6015 (JP).

(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published
With international search report.
Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(54) Title: A METHOD FOR MANUFACTURING GLYCOPROTEINS HAVING HUMAN–TYPE GLYCOSYLATION

(57) Abstract

The present invention provides a method for manufacturing a glycoprotein having a human–type sugar chain comprising a step in which a transformed plant cell is obtained by introducing to a plant cell the gene of glycosyltransfetase and the gene of an exogenous glycoprotein, and a step in which the obtained transformed plant cell is cultivated.