

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
25 October 2001 (25.10.2001)

PCT

(10) International Publication Number
WO 01/79388 A3

- (51) International Patent Classification⁷: **C10G 21/16**, 21/06
- (21) International Application Number: PCT/US01/08895
- (22) International Filing Date: 20 March 2001 (20.03.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
09/551,659 18 April 2000 (18.04.2000) US
- (71) Applicant: **EXXONMOBIL RESEARCH AND ENGINEERING COMPANY** [US/US]; P.O. Box 900, 1545 Route 22 East, Annandale, NJ 08801 (US).
- (72) Inventors: **GREANEY, Mark, Alan**; 1265 River Road, Upper Black Eddy, PA 18972 (US). **BEGASSE, John, Nicholas**; 30 Amelia Avenue, Livingston, NJ 07039 (US).
- (74) Agents: **SCUORZO, Linda, M.** et al.; ExxonMobil Research and Engineering Company, P.O. Box 900, Annandale, NJ 08801 (US).
- (81) Designated States (*national*): AU, CA, ID, JP, SG.
- (84) Designated States (*regional*): European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR).
- Published:
— with international search report
- (88) Date of publication of the international search report:
18 April 2002
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: METHOD FOR ISOLATING ENRICHED SOURCE OF CONDUCTING POLYMERS PRECURSORS

(57) Abstract: An embodiment of the present invention is a method for isolating conducting polymer precursors by contacting a non-basic heterocyclic nitrogen containing hydrocarbon stream having a boiling point of from 232 °C /450 °F) to 566 °C (1050 °F) with an effective amount of a treating agent selected from the group consisting of alkylene and polyalkylene glycols and glycol ethers and mixtures thereof, having a molecular weight of less than 1000 and 1200, respectively, at conditions effective to maintain the reactants in a liquid phase to produce a first stream enriched in non-basic heterocyclic nitrogen containing hydrocarbon compounds and a second treated stream having a decreased non-basis heterocyclic nitrogen content. Optionally, an effective amount of a mineral acid may be added to the treating agent to enhance the process.

WO 01/79388 A3