



CALCITONIN RECEPTOR-BINDING PEPTIDES

5 This invention relates to calcitonin receptor binding reagents which are capable of complexing with metal ions, including radioactive metal ions, and to labeled embodiments of such reagents for use in imaging sites in a mammalian body or for use in therapy, particularly for use in cancer therapy.

10 BACKGROUND OF THE INVENTION

Great strides have been made over the past 60 years in reducing long-term mortality trends for some types of cancer, such as stomach cancer. However, during the same period, mortality trends for other cancers have remained stable or increased. For example, lung cancer is the most frequent cancer worldwide, representing the leading cause of cancer mortality
15 among men and women. Breast cancer is the commonest cancer among women and the second leading cause of cancer mortality in women, and ovarian cancer mortality rates are increasing in some countries. Childhood and adult lymphatic cancers, such as leukemias and non-Hodgkin's lymphomas, also continue to represent significant causes of cancer mortality. Early diagnosis and effective treatment remains a goal for all of these cancers.

20 Several years ago, site directed diagnosis and therapy were proposed, to allow *in vivo* targeting of particular sites of disease within an animal's body. In general, site-directed diagnosis or therapy employs a targeting moiety, such as an antibody specific for the disease site or for the organism which caused the disease, coupled to a label in the case of a diagnostic agent or to a cytotoxic agent in the case of a therapeutic agent. A very large body of literature
25 exists relating to radiolabeling antibodies or antibody fragments for diagnostic imaging purposes. Similarly, a number of site directed therapeutic agents employing monoclonal antibodies and a variety of radioisotopes have been proposed over the years, *e.g.*, as set forth in U.S.Pat.Nos. 4,454,106; 4,472,509; 4,828,991; 5,246,691; 5,355,394; and 5,641,471; in EP 429624; EP 585986; WO 90/15625, and the like. Such antibody-based agents produce side
30 effects related to the immune responses of the treated animal to the antibody, even if antibody fragments or humanized antibodies are employed as the targeting moiety.

The disadvantages of antibody-based site-directed diagnostic and therapeutic agents can be avoided when targeting moieties having lower molecular weights, such as receptor-specific peptides or small molecules are employed. However, coupling of a peptide or small molecule to a label or cytotoxic agent, while retaining the compound's receptor specificity, can be technically difficult. Methods for radiolabeling peptides and other small molecules which preserve the ability of the compound to bind specifically to a receptor are disclosed in commonly owned U.S. Patent Nos. 5,225,180; 5,405,597; 5,443,815; 5,508,020; 5,552,525; 5,561,220; 5,620,675; 5,645,815; 5,654,272; 5,711,931; 5,716,596; 5,720,934; and 5,736,122; in abandoned U.S. Patent Application Serial No. 07/955,466; and in WO92/13572, WO93/10747, WO93/17719, WO93/21962, WO93/23085, WO93/25244, WO94/00489, WO94/07918, and WO94/28942. The methods disclosed in these patents and publications are particularly suitable for manufacture of site-directed diagnostic imaging agents. Commonly assigned U.S. Pat. Nos. 5,620,675; 5,716,596; WO 94/00489; WO 95/03330; WO 95/00553; WO 95/31221; and WO 96/04308 disclose somatostatin peptide analogs which may be used for site-directed radiotherapy. Commonly assigned WO 95/33497 discloses somatostatin analogs, gpIIb/IIIa receptor-binding peptides, and leukocyte-binding peptides which may be used for site-directed radiodiagnosis or radiotherapy. Commonly assigned WO 96/30055 discloses vasoactive intestinal peptide (VIP) receptor-binding peptides which may be used for site-directed radiodiagnosis or radiotherapy.

Tumor cells often occasionally express or overexpress a particular receptor or receptor subtype, as indicated by receptor binding studies. In some types of cancer, tumor cell markers can change as the disease progresses, possibly reflecting the stage of the disease and thus the patient's prognosis. The kind of receptor that a tumor cell expresses can be characteristic of the tumor's etiology and can thus provide a relatively specific marker for the tumor. For example, radiolabeled somatostatin analogs have been shown to bind specifically to neuroendocrine tumors, melanomas, lung cancer, and certain breast cancers. One such analog, ¹¹¹In-OCTREOSCAN, has received marketing approval for use in imaging neuroendocrine tumors. A second radiolabeled somatostatin analog, ^{99m}Tc-Depreotide, has completed Phase III clinical trials for use in imaging lung cancers. ¹²³I-vasoactive intestinal peptide has been shown to target adenocarcinomas of the colon and stomach.

Calcitonin (CT) is a 32 amino acid peptide secreted from the thyroid in response to elevated serum calcium levels. Calcitonin has a number of biological effects, which are mediated by calcitonin receptors present on the surfaces of cells in the target organ. High affinity receptors for CT have been identified in bone, kidney, lung, and central nervous system. In bone, CT inhibits bone resorption by osteoclasts; in kidney, CT increases excretion of calcium ions; and in the central nervous system, the peptide induces analgesia, gastric acid secretion, and appetite inhibition. Small amounts of CT have been administered to animals and humans without toxic effects, and salmon CT is used clinically to treat such bone disorders as Paget's disease, hypercalcemia of malignancy, and osteoporosis. Intravenously-

administered calcitonin clears the blood rapidly and is excreted primarily in urine. The major sites of localization for administered CT are kidney, liver and the epiphyses of the long bones.

Circulating CT levels are considered to be a marker for some types or stages of cancer, for example, medullary thyroid carcinoma, small-cell lung cancer, carcinoids, breast cancer, and gastrointestinal cancer. High affinity CT receptors have been identified in lymphoid cells, human lung cancer cell lines, human breast cancer cell lines, and in primary breast cancer tissue. Findlay *et al.* (1981) *Biochem. J.* 196: 513-520 reports that CT receptors are overexpressed in certain breast, lung, ovarian and lymphoma cancer cell lines.

The amino acid sequences of CT from several species (human, salmon and eel) are set forth below:

hCT CGNLSTCMLGTYTQDFNKFHTFPQTAIGVG.AP.amide (SEQ ID NO. 1)
 sCT CSNLSTCVLGKLSQELHKLQTYPRNTGSG.TP.amide (SEQ ID NO. 2)
 eCT CSNLSTCVLGKLSQELHKLQTYPRTDVGAGTP.amide (SEQ ID NO. 3)

(where single-letter abbreviations for amino acids can be found in Zubay, *Biochemistry* 2d ed., 1988, MacMillan Publishing: New York, p. 33, and where the underlined amino acids between the two cysteine residues in the amino terminal portion of the peptide represent a disulfide bond). Among species, nine residues, including the carboxyl-terminal proline amide and the disulfide-bonded cysteine residues at positions 1 and 7 are conserved. The salmon and eel CTs are more potent than human CT both *in vitro* and *in vivo*.

CT peptide analogs have been developed in which the chemically-labile disulfide is replaced with stable carbon-carbon linkages formed between 2-aminosuberic acid, as described

in U.S.PAT.No. 4,086,221. CT analogs in which the amino terminal, midregion, or carboxyl terminal portion of the molecule are deleted demonstrate only weak binding to CT receptors. Many amino acid substitutions may be made between residues 8 and 22 of the CT molecule to generate biologically active CT analogs. Some CT analogs with only minimal sequence
5 homology to any natural form of CT have biological activity similar to that of salmon CT. Truncated CT peptide derivatives (such as Cbz-LHKLQY-OMe) also retain substantial receptor binding activity.

Since tumors may express or overexpress different receptors, a variety of radiodiagnostic and radiotherapeutic agents are needed to afford optimal diagnostic and
10 therapeutic modalities against cancer.

SUMMARY OF THE INVENTION

CT receptors on cell surfaces of lung and ovarian adenocarcinoma, breast cancers and
15 lymphomas can be exploited as markers to locate, identify, and treat such tumors *in vivo*. The present inventors have for the first time developed small, synthetic compounds, including CT-derived peptides, possessing the capacity for high-affinity binding to CT receptors and favorable pharmacokinetics, thereby permitting efficient *in vivo* localization of diagnostic and therapeutic agents at tumor sites. The reagents of the invention provide the basis of rapid, cost-effective,
20 non-invasive diagnostic imaging procedures useful for tumor detection, disease staging and evaluation of metastatic spread of tumors characterized by expression or overexpression of CT receptors. The reagents of the invention also provide the basis for assessment of therapeutic effectiveness of other treatment modalities, for example, by localization of CT receptor-expressing tumor cells following surgery, radiation therapy or chemotherapy. The reagents of
25 the invention may also be used as targeting moieties for site-directed radiotherapy.

The invention provides CT receptor binding reagents comprising CT receptor binding compounds, preferably CT peptides, CT derivatives, or CT analogues, which are covalently linked to a radiometal chelator. The CT receptor binding compounds employed in the reagents of the invention have a molecular weight of less than about 10,000 daltons. In some
30 embodiments, the reagents of the invention are characterized as peptides, by virtue of the presence of a peptide linkage either in the CT receptor binding portion of the reagent or by

virtue of the presence of a peptide linkage in the radiometal chelator. The reagents of the invention have a CT receptor binding affinity that is not less than about one-tenth the affinity of radioiodinated native CT for said receptor, when compared in a standardized assay such as the assays described in Example 4 below. In preferred embodiments, the reagents of the invention
5 have a CT receptor binding affinity equal to or greater than native CT or radioiodinated species of native CT for said receptor, when compared in said standardized assay.

The radiopharmaceuticals of the invention may be employed as site specific diagnostic or therapeutic agents. When labeled with technetium-99m, iodine-123, and iodine-131, the reagents of the invention may be employed as scintigraphic imaging agents. When labeled with
10 a magnetic, paramagnetic, supermagnetic, or superparamagnetic metal, the reagents of the invention may be employed as magnetic resonance contrast agents. When labeled with a cytotoxic radionuclides, the reagents of the invention may be used for site-directed radiotherapy. The invention also provides pharmaceutical compositions comprising the radiolabeled CT receptor-binding compounds of the invention and a pharmaceutically acceptable carrier.
15 Methods for making the CT receptor binding reagents of the invention and radiolabeled embodiments thereof are also provided.

The invention also provides kits for preparing radiolabeled CT receptor binding compounds from the reagents of the invention. The kits of the invention comprise a sealed vial containing a predetermined quantity of a reagent of the invention and optionally a sufficient
20 amount of a reducing agent to radiolabel the reagent.

This invention provides methods for using the radiolabeled CT receptor-binding reagents of the invention diagnostically and therapeutically. In one embodiment, methods are provided for using reagents of the invention, in labeled form, for imaging sites within a mammalian body by obtaining *in vivo* images. These methods comprise the steps of
25 administering an effective diagnostic amount of labeled reagents of the invention and detecting the label localized at the site within the mammalian body.

The invention also provides methods for alleviating diseases characterized by expression or overexpression of CT receptors, comprising the step of administering a therapeutically effective amount of a radiolabeled CT receptor-binding reagents of the invention to the animal.

30 Other aspects and advantages of the present invention are apparent in the following more detailed description of preferred embodiments set forth below.

DETAILED DESCRIPTION OF THE INVENTION

The present invention provides CT receptor binding reagents useful in the preparation of CT receptor binding pharmaceutical agents for diagnosis and therapy. For the purposes of this invention, the term "CT receptor binding compound" is intended to encompass naturally-occurring CT, fragments of CT, analogues of CT, and derivatives of CT that specifically bind to the CT receptor expressed in a variety of cell types recognized by those with skill in the art. Compounds designed to mimic the receptor-binding properties of CT are also included in this definition and encompassed by the invention.

For the purposes of this invention, the term "CT receptor binding affinity" is intended to mean binding affinity as measured by any methods known to those of skill in the art, including, *inter alia*, those methods which measure binding affinity by a dissociation constant, an inhibition constant or an IC_{50} value.

The term "having a CT receptor binding affinity of at least one-tenth the affinity of radioiodinated CT for said receptor" is intended to mean that the dissociation constant (K_d) of the reagent is not less than ten times less than the K_d of radioiodinated CT as measured in a CT receptor direct binding or competitive inhibition assay, or that the or inhibition constant (K_i) or IC_{50} of the reagent is not more than 10 times that of radioiodinated CT, as measured in a CT receptor competitive inhibition assay.

The term "having a CT receptor binding affinity equal to or greater than native CT or radioiodinated species of CT for said CT receptor" is intended to encompass reagents having a dissociation constant (K_d) equal to or greater than that of unlabeled CT or radioiodinated CT, as measured in a CT receptor direct binding or competitive inhibition assay. Alternatively, this term may be interpreted as encompassing reagents having an inhibition constant (K_i) or IC_{50} equal to or less than that of native CT or radioiodinated CT as measured in a CT receptor competitive inhibition assay.


In accordance with the invention, the reagents of the invention have a CT receptor binding affinity of at least ten times greater than the affinity of radioiodinated native CT for said receptor, when compared in a standardized assay as described below. In preferred embodiments, the reagents of the invention have a CT receptor binding affinity equal to or

greater than native CT or radioiodinated species of native CT for said receptor, when compared in said standardized assay.

Dissociation constants or binding inhibition constants of the radiopharmaceuticals of the invention in relation to the CT receptor, as well as comparison of the affinity or avidity of such binding with binding of ^{125}I -labeled CT itself, may be determined using known methods such as those set forth in *Receptors, A Quantitative Approach*, A. Levitzki, The Benjamin/Cummings Publishing Company (California, 1984). Preferably a standardized CT receptor binding assay, such as the assays set forth in Example 4 below, is employed to measure CT receptor binding affinity of the reagents of the invention.


In addition to the CT receptor binding compound, the reagents provided by this invention comprise a radiometal chelator which is covalently linked to the compound in such a way that the binding specificity of the compound for the CT receptor is not substantially altered.

Any radiometal chelator may be covalently linked to a CT receptor binding compound to provide a reagent of the invention. For example, a CT receptor-binding compound, preferably a peptide, may be covalently linked to a chelator having the formula:



where $(\text{pgp})^s$ is hydrogen or a thiol protecting group and (aa) is any α - or β -amino acid not comprising a thiol group. In a preferred embodiment, the amino acid is glycine.


As another example, a radiometal chelator useful in the reagents of the invention comprises a single thiol-containing group of formula:




wherein A is H, HOOC, H_2NOC , (peptide)-NHOC, (peptide)-OOC, R^cNCO , or R^d ; B is H, SH or -NHR^c , $\text{-N(R}^c\text{)-(peptide)}$ or R^d ; Z is H or R^d ; X is SH or -NHR^c , $\text{-N(R}^c\text{)-(peptide)}$ or R^d ; R^a , R^b , R^c and R^d are independently H or straight or branched chain or cyclic lower alkyl; n is 0, 1 or 2; R^c is $\text{C}_1\text{-C}_4$ alkyl, an amino acid or a peptide comprising 2 to about 10 amino acids; and: (1) where B is -NHR^c or $\text{-N(R}^c\text{)-(peptide)}$, X is SH and n is 1 or 2; (2) where X is -NHR^c or $\text{-N(R}^c\text{)-(peptide)}$, B is SH and n is 1 or 2; (3) where B is H or R^d , A is HOOC, H_2NOC , (peptide)-NHOC, or (peptide)-OOC, X is SH and n is 0 or 1; (4) where A is H or R^d , then where B is SH, X is -NHR^c or $\text{-N(R}^c\text{)-(peptide)}$ and where X is SH, B is -NHR^c or $\text{-N(R}^c\text{)-(peptide)}$ and n is 1 or 2; (5) where X is H or R^d , A is HOOC, H_2NOC , (peptide)-NHOC, or (peptide)-OOC and B is SH; (6) where Z is methyl, X is methyl, A is HOOC, H_2NOC , (peptide)-NHOC, or (peptide)-

OOC and B is SH and n is 0; and (7) where B is SH, X is not SH and where X is SH, B is not SH.

Preferred embodiments of this radiometal chelator have the chemical formula:






5 wherein (amino acid)¹ and (amino acid)² are each independently any primary α - or β -amino acid that does not comprise a thiol group, Z is a thiol-containing moiety that is cysteine, homocysteine, isocysteine, penicillamine, 2-mercaptoethylamine or 3-mercaptoethylamine, and R¹ is lower (C¹-C⁴) alkyl, an amino acid or a peptide comprising 2 to 10 amino acids. When Z is cysteine, homocysteine, isocysteine or penicillamine, the carbonyl group of said moiety is covalently linked to a hydroxyl group, a NR³R⁴ group, wherein each of R³ and R⁴ are
10 independently H or lower (C¹-C⁴) alkyl, an amino acid or a peptide comprising 2 to 10 amino acids; or




wherein Y is a thiol-containing moiety that is cysteine, homocysteine, isocysteine, penicillamine, 2-mercaptoacetate or 3-mercaptoethylamine, (amino acid)¹ and (amino acid)² are
15 each independently any primary α - or β -amino acid that does not comprise a thiol group, and R² is H or lower (C¹-C⁴) alkyl, an amino acid or a peptide comprising 2 to 10 amino acids. When Y is cysteine, homocysteine, isocysteine or penicillamine, the amino group of said moiety is covalently linked to -H, an amino acid or a peptide comprising 2 to 10 amino acids.

20 In particular embodiments of this aspect of the invention, the formula of this radiometal chelator is selected from the group consisting of:



25 and




wherein (amino acid)¹ and (amino acid)² are each independently any naturally-occurring, modified, substituted or altered α - or β -amino acid not containing a thiol group; A is H, HOOC, H₂NOC, (amino acid or peptide)-NHOC, (amino acid or peptide)-OOC or R⁴; B is H,
30 SH or -NHR³, -N(R³)-(amino acid or peptide) or R⁴; Z is H or R⁴; X is SH or -NHR³, -N(R³)-

(amino acid or peptide) or R^4 ; R^1 , R^2 , R^3 and R^4 are independently H or straight or branched chain or cyclic lower alkyl; n is an integer that is either 0, 1 or 2; (peptide) is a peptide of 2 to about 10 amino acids; and: (1) where B is $-NHR^3$ or $-N(R^3)$ -(amino acid or peptide), X is SH and n is 1 or 2; (2) where X is $-NHR^3$ or

- 5 $-N(R^3)$ -(amino acid or peptide), B is SH and n is 1 or 2; (3) where B is H or R^4 , A is $HOOC$, H_2NOC , (amino acid or peptide)- $NHOC$, (amino acid or peptide)- OOC , X is SH and n is 0 or 1; (4) where A is H or R^4 , then where B is SH, X is $-NHR^3$ or $-N(R^3)$ -(amino acid or peptide) and where X is SH, B is $-NHR^3$ or $-N(R^3)$ -(amino acid or peptide) and n is 1 or 2; (5) where X is H or R^4 , A is $HOOC$, H_2NOC , (amino acid or peptide)- $NHOC$, (amino acid or peptide)- OOC and
10 B is SH; (6) where Z is methyl, X is methyl, A is $HOOC$, H_2NOC , (amino acid or peptide)- $NHOC$, (amino acid or peptide)- OOC and B is SH and n is 0; and (7) where B is SH, X is not SH and where X is SH, B is not SH.

Specific preferred embodiments of this aspect of the invention include radiometal chelators having a formula selected from the group consisting of: -Gly-Gly-Cys-, Cys-Gly-Gly-,
15 Gly-Gly-Cys-, $-(\epsilon\text{-Lys})\text{-Gly-Cys-}$, $(\delta\text{-Orn})\text{-Gly-Cys-}$, $-(\gamma\text{-Dab})\text{-Gly-Cys-}$, $-(\beta\text{-Dap})\text{-Lys-Cys-}$, and $-(\beta\text{-Dap})\text{-Gly-Cys-}$. (In these formulae, it will be understood that $\epsilon\text{-Lys}$ represents a lysine residue in which the ϵ -amino group, rather than the typical α -amino group, is covalently linked to the carboxyl group of the adjacent amino acid to form a peptide bond; $\delta\text{-Orn}$ represents an ornithine residue in which the δ -amino group, rather than the typical α -amino group, is
20 covalently linked to the carboxyl group of the adjacent amino acid to form a peptide bond; $\gamma\text{-Dab}$ represents a 2,4-diaminobutyric acid residue in which the γ -amino group is covalently linked to the carboxyl group of the adjacent amino acid to form a peptide bond; and $\beta\text{-Dap}$ represents a 1,3-diaminopropionic acid residue in which the β -amino group is covalently linked to the carboxyl group of the adjacent amino acid to form a peptide bond.)

25 In another embodiment, the radiometal chelator of the reagent of the invention is a bisamino-bisthiol chelator having the formula:




30

wherein each R can be independently H, CH₃ or C₂H₅; each (pgp)^s can be independently a thiol protecting group or H; m, n and p are independently 2 or 3; A is linear or cyclic lower alkyl, aryl, heterocyclyl, combinations or substituted derivatives thereof; and X is peptide.

Alternatively, the bisamino bithiol chelator in this embodiment of the invention has the

5 formula:



10 wherein each R is independently H, CH₃ or C₂H₅; m, n and p are independently 2 or 3; A is linear or cyclic lower alkyl, aryl, heterocyclyl, combinations or substituted derivatives thereof; V is H or CO-peptide; R' is H or peptide; provided that when V is H, R' is peptide and when R' is H, V is CO-peptide. For purposes of this invention, chelating moieties having these structures will be referred to as "BAT" moieties.

15 Alternatively, the radiometal chelator used in the reagent of the invention may have a formula selected from the group consisting of:
diethylenetriaminepentaacetic acid (DTPA)



where each R is independently H, C₁ to C₄ alkyl, or aryl and one R is covalently linked to a


20 bivalent linker;

ethylenediaminetetraacetic acid (EDTA): $(\text{HOOCCH}_2)_2\text{N}(\text{CR}_2)(\text{CR}_2)\text{N}(\text{CH}_2\text{COOH})$;

where each R is independently H, C₁ to C₄ alkyl, or aryl and one R is covalently linked to a bivalent linker;

1,4,7,10-tetraazadodecanetetraacetic acid;

25




where n is an integer that is 2 or 3 and where each R is independently H, C₁ to C₂ alkyl, or aryl and one R is covalently linked to the CT receptor binding compound, and desferrioxamine.

Most radiometals may be chelated to reagents of the invention comprising the above-mentioned radiometal chelators.


The reagents of the invention may also comprise a radiometal chelator selected from the group consisting of:

- (i) a group having the formula:



I.

- (ii) a group having the formula:




II.

wherein n, m and p are each integers that are independently 0 or 1; each R' is independently H, lower alkyl, C₂-C₄ hydroxyalkyl, or C₂-C₄ alkoxyalkyl, and each R is independently H or R'', where R'' is substituted or unsubstituted lower alkyl or phenyl not comprising a thiol group, and one R or R' is L, where L is a bivalent linker moiety linking the metal chelator to the targeting moiety and wherein when one R' is L, NR' is an amine.

In preferred embodiments, L is a C_1 - C_6 linear, branched chain or cyclic alkyl group, a carboxylic ester, a carboxamide, a sulfonamide, an ether, a thioether, an amine, an alkene, an alkyne, a 1,2-, 1,3- or 1,4-linked, optionally substituted, benzene ring, or an amino acid or peptide of 2 to about 10 amino acids, or combinations thereof.

- 5 In preferred embodiments, R'' is a C_1 - C_6 linear, branched or cyclic alkyl group; a $-C_qOC_r-$, $-C_qNHC_r-$ or $-C_qSC_r-$ group, where q and r are integers each independently 1 to 5 wherein the sum of q + r is not greater than 6; (C_1-C_6) alkyl-X, where X is a hydroxyl group, a substituted amine, a guanidine, an amidine, a substituted thiol group, or a carboxylic acid, ester, phosphate, or sulfate group; a phenyl group or a phenyl group substituted with a halogen,
- 10 hydroxyl, substituted amine, guanidine, amidine, substituted thiol, ether, phosphate, or sulfate group; an indole group; a C_1 - C_6 heterocyclic group containing 1 to 3 nitrogen, oxygen or sulfur atoms or combinations thereof.


In accordance with the invention, the radiometal chelator of the CT receptor-binding reagent may have the formula:



III.

- wherein R^1 and R^2 are each independently H, lower alkyl, C_2 - C_4 hydroxyalkyl, or C_2 - C_4 alkoxyalkyl; R^3 , R^4 , R^5 and R^6 are independently H, substituted or unsubstituted lower alkyl or phenyl not comprising a thiol group; R^7 and R^8 are each independently H, lower alkyl, lower
- 20 hydroxyalkyl or lower alkoxyalkyl; L is a bivalent linker group and Z is a CT peptide.


Additional preferred metal chelators of the invention include chelators of formula:



IV.

wherein R^1 and R^2 are each independently H, lower alkyl, C_2-C_4 hydroxyalkyl, or C_2-C_4 alkoxyalkyl; R^3 , R^4 , R^5 and R^6 are independently H, substituted or unsubstituted lower alkyl or phenyl not comprising a thiol group, and one of R^3 , R^4 , R^5 or R^6 is $Z-L-HN(CH_2)_n-$, where L is a bivalent linker group, Z is a targeting moiety, and n is an integer from 1 to 6; R^7 and R^8 are each independently H, lower alkyl, lower hydroxyalkyl or lower alkoxyalkyl; and X is an amino group, a substituted amino group or $-NR^1-Y$, where Y is an amino acid, an amino acid amide, or a peptide comprising from 2 to 10 amino acids.


More preferred metal chelators of the invention include chelators having the formula:



V.

wherein R^1 and R^2 are each independently H, lower alkyl, lower hydroxyalkyl, or lower alkenylalkyl; R^3 and R^4 are independently H, substituted or unsubstituted lower alkyl or phenyl not comprising a thiol group; n is an integer from 1 to 6; L is a bivalent linker group; and Z is a CT peptide moiety.

Additional more preferred chelating moieties include chelators of formula:



VI.

wherein L is a bivalent linker group and Z is a CT peptide moiety.

Most preferred chelating moieties of the invention include chelators having the following formulae:

(amino acid)¹-(amino acid)²-cysteine-,

(amino acid)¹-(amino acid)²-isocysteine-,
 (amino acid)¹-(amino acid)²-homocysteine-,
 (amino acid)¹-(amino acid)²-penicillamine-,
 (amino acid)¹-(amino acid)²-2-mercaptoethylamine-,
 5 (amino acid)¹-(amino acid)²-2-mercaptopropylamine-,
 (amino acid)¹-(amino acid)²-2-mercapto-2-methylpropylamine-,
 (amino acid)¹-(amino acid)²-3-mercaptopropylamine-,

wherein (amino acid) in a primary α - or β -amino acid not comprising a thiol group and wherein the chelator is attached to either a targeting moiety or a linker group *via* a covalent bond with
 10 the carboxyl terminus of the chelator or a side chain on one of the amino acid groups.

Most preferred chelators also include chelators of the above formula wherein (amino acid)¹ is either an α,ω - or β,ω -amino acid wherein the α - or β -amino group is a free amine and the α,ω - or β,ω -amino acid is covalently linked *via* the ω amino group.

Other most preferred chelators include those selected from the group consisting of:


15 -cysteine-(amino acid)-(α,β - or β,γ -diamino acid);
 -isocysteine-(amino acid)-(α,β - or β,γ -diamino acid);
 -homocysteine-(amino acid)-(α,β - or β,γ -diamino acid);
 -penicillamine-(amino acid)-(α,β - or β,γ -diamino acid);
 2-mercaptoacetic acid-(amino acid)-(α,β - or β,γ -diamino acid);
 20 2- or 3-mercaptopropionic acid-(amino acid)-(α,β - or β,γ -diamino acid);
 2-mercapto-2-methylpropionic acid-(amino acid)-(α,β - or β,γ -diamino acid);

wherein (amino acid) in a primary α - or β -amino acid not comprising a thiol group and wherein the chelator is attached to either a targeting moiety or a linker group *via* a covalent bond with the amino terminus of the chelator or a side chain on one of the amino acid groups.

25 Particularly preferred metal chelators are selected from the group consisting of:


Gly-Gly-Cys-, Arg-Gly-Cys-, -(ϵ -Lys)-Gly-Cys-, -(δ -Orn)-Gly-Cys-, -(γ -Dab)-Gly-Cys-, -(β -Dap)-Lys-Cys-. and -(β -Dap)-Gly-Cys-. (In these formulae, the amino acid designations have the same meaning as is set forth above.)

An example of a radiometal chelator having structure III above is Gly-Gly-Cys-,
 30 wherein the chelating moiety has the structure:




VII.

- 5 Chelating ligands having structure type VII form oxotechnetium complexes having the structure:




VIII.

- An example of radiometal chelators having structure type V as shown above is Lys-(ω-peptide)-Gly-Cys.amide which forms a chelator of structure:
- 10



IX.


- 15 Chelating ligands having structure type IX form oxotechnetium complexes having the structure:



X.

5


An example of a reagent of the invention comprising a radiometal chelator having structure II as shown above is (targeting moiety)-Cys-Gly- α,β -diaminopropionamide which forms a chelator of structure:



XI.

10

Radiodiagnostic agents having structure type XI form oxotechnetium complexes having the structure:



XII.

15

In the radiometal chelators and CT receptor binding reagents provided by the invention that contain a thiol covalently linked to a thiol protecting group $\{(pgp)^S\}$, the thiol-protecting groups may be the same or different and may be but are not limited to:

-CH₂-aryl (aryl is phenyl or alkyl or alkyloxy substituted phenyl);

-CH-(aryl)₂, (aryl is phenyl or alkyl or alkyloxy substituted phenyl);

-C-(aryl)₂, (aryl is phenyl or alkyl or alkyloxy substituted phenyl);

-CH₂-(4-methoxyphenyl);

-CH-(4-pyridyl)(phenyl)₂;

5 -C(CH₃)₃

-9-phenylfluorenyl;

-CH₂NHCOR (R is unsubstituted or substituted alkyl or aryl);

-CH₂-NHCOOR (R is unsubstituted or substituted alkyl or aryl);

-CONHR (R is unsubstituted or substituted alkyl or aryl);

10 -CH₂-S-CH₂-phenyl

Preferred protecting groups have the formula -CH₂-NHCOR wherein R is a lower alkyl having between 1 and 8 carbon atoms, phenyl or phenyl-substituted with lower alkyl, hydroxyl, lower alkoxy, carboxy, or lower alkoxy carbonyl. The most preferred protecting group is an acetamidomethyl group.

15 When the reagent of the invention comprises a CT receptor binding compound which is a peptide, the peptide preferably comprises the amino acid sequence:

CH₃CO.SNLSTX—

SEQ ID NO.: 10

wherein X is selected from the group consisting of a cysteine residue, a homocysteine residue, and a homohomocysteine residue. Alternatively, the peptide may comprise the amino acid
20 sequence:

CH₃CO.X¹NLSTX²—

SEQ ID NO.: 11

wherein X¹ is selected from the group consisting of an alanine residue, a glycine residue, and a serine residue; and X² is selected from the group consisting of a cysteine residue, a homocysteine residue, and a homohomocysteine residue. Such peptides include naturally-
25 occurring human CT and CT peptide analogs such as those which are specifically embodied in the amino acid sequences set forth below:

CH₃CO.SNLST.Hhc.VLGKLSCELHKLQTYPRNTNTGSGTP.amide

(SEQ ID No. 5),

CH₃CO.SNLST.Hcy.VLGKLSCELHKLQTYPRNTNTGSGTP.amide

(SEQ ID No. 6),

CH₃CO.SNLST.Cys.VLGKLSCELHKLQTYPRNTNTGSGTP.amide

(SEQ ID No. 7),

30 and

SNLST.Asu.VLGKLSCELHKLQTYPRNTNTGSGTP.amide

(SEQ ID No. 8).

Particularly preferred embodiments of the reagents of the invention include:

CH₂CO.SNLST.Hhc.VLGKLSC(BAT)ELHKLQTYPRTNTGSGTP.amide (SEQID No.4),
CH₂CO.SNLST.Hhc.VLGKLSQELHKLQTYPRTNTGSGTP(ε-K)GC.amide,
CH₂CO.SNLST.Hhc.VLGKLSC(CH₂CO.GGCK.amide)ELHKLQTYPRTNTGSGTP.amide,
CH₂CO.SNLST.Hhc.VLGKLSC(CH₂CO.(β-Dap)KCK.amide)ELHKLQTYPRTNTGSGTP.amide,
CH₂CO.SNLST.Hhc.VLGKLSC(CH₂CO.(ε-K)GCE.amide)ELHKLQTYPRTNTGSGTP.amide,
CH₂CO.SNLST.Hcy.VLGKLSC(CH₂CO.GGCK.amide)ELHKLQTYPRTNTGSGTP.amide,
CH₂CO.SNLST.Hcy.VLGKLSC(CH₂CO.(β-Dap)KCK.amide)ELHKLQTYPRTNTGSGTP.amide,
CH₂CO.SNLST.Hcy.VLGKLSC(CH₂CO.(ε-K)GCE.amide)ELHKLQTYPRTNTGSGTP.amide,
CH₂CO.SNLST.Cys.VLGKLSC(CH₂CO.GGCK.amide)ELHKLQTYPRTNTGSGTP.amide,
CH₂CO.SNLST.Cys.VLGKLSC(CH₂CO.(β-Dap)KCK.amide)ELHKLQTYPRTNTGSGTP.amide,
CH₂CO.SNLST.Cys.VLGKLSC(CH₂CO.(ε-K)GCE.amide)ELHKLQTYPRTNTGSGTP.amide,
SNLST.Asu.VLGKLSC(CH₂CO.(β-Dap)KCK.amide)ELHKLQTYPRTNTGSGTP.amide, and
SNLST.Asu.VLGKLSC(CH₂CO.(β-Dap)KCK.amide)ELHKLQTYPRTDVGAGTP.amide.

All naturally-occurring amino acids are abbreviated using standard abbreviations (which
5 can be found in G. Zubay, *Biochemistry* (2d. ed.), 1988 (MacMillen Publishing: New York)
p.33). For the purposes of this invention, the naturally-occurring amino acids are characterized
as lipophilic (alanine, isoleucine, leucine, methionine, phenylalanine, tyrosine, proline,
tryptophan and valine, as well as *S*-alkylated derivatives of cysteine), hydrophilic (asparagine,
glutamine, threonine, serine), acidic (glutamic acid and aspartic acid), basic (arginine, histidine
10 and lysine). ε-K, δ-Orn, γ-Dab and β-Dap have the meanings set forth above. (BAT) represents
N⁶,N⁹-bis(2-mercapto-2-methyl-propyl)-6,9-diazanonanoic acid; K.(BAT) and Lys.(BAT)
represent the amino acid lysine, acylated at the ε-amino group on the amino acid sidechain to
(BAT); C(BAT) and Cys(BAT) represent *S-(N⁶,N⁹-bis(2-mercapto-2methylpropyl)-6,9--*
diazanonan-1-yl)cysteine; (BAM) is *(N¹,N²-bis(2-mercapto-2-methylpropyl)-1,4,10-*
15 *triazadecane*; (BAT-BM) is *N-{2-(N¹,N²-bis(2-maleimidoethyl)aminoethyl)-N⁹-(*t*-*
butoxycarbonyl)-N⁶,N⁹-bis(2-methyl-2-triphenyl-methylthiopropyl)-6,9-diazanonanamide;
(BAT-BS) is *N-{2-(N¹,N²-bis(2-succinimidoethyl)aminoethyl)-N⁶,N⁹-bis(2-mercapto-2-*
methylpropyl)-6,9-diazanonanamide; (BMH) is *bis-maleimido*hexane; (BSH) is *bis-*
*succinimido*hexane; (BMME) is *bis-maleimidomethylether*; (BSEE) is *bis-*
20 *succinimidoethylether*; (BMEE) is *bis-maleimidoethylether*; and (BSME) is *bis-*

succinimidomethylether. As used herein, the following amino acids and amino acid analogues are intended to be represented by the following abbreviations: Ac_m is the sulfhydryl protecting group acetamidomethyl; Pen is penicillamine; Aca is 6-aminocaproic acid; Hly is homolysine; Apc is L-{S-(3-aminopropyl)cysteine; F_ε is D-phenylalanine; W_δ is D-tryptophan; Y_δ is D-tyrosine; Cpa is L-(4-chlorophenyl)alanine; Thp is 4-amino-tetrahydrothiopyran-4-carboxylic acid; D-Nal is D-2-naphthylalanine; Dpg is dipropylglycine; Nle is norleucine; Hcy is homocysteine; Hhc is homohomocysteine; Aib is aminoisobutyric acid; Nal is 2-naphthylalanine; D-Nal is D-2-naphthylalanine; Ain is 2-aminoindan-2-carboxylic acid; Achxa is 4-amino-cyclohexylalanine; Amf is 4-aminomethyl-phenylalanine; Aec is S-(2-aminoethyl)cysteine; Apc is S-(3-aminopropyl)cysteine; Aes is O-(2-aminoethyl)serine; Aps is O-(3-aminopropyl)serine; Abu is 2-aminobutyric acid; Nva is norvaline; and Asu is 2-amino suberic acid, wherein the amino terminal amino acids of peptides containing an Asu residue are cyclized *via* an amide bond between the amino terminal amino group and the side chain carboxylic acid moiety of the Asu residue.

In accordance with the invention, CT receptor binding peptides may comprise one or more amino acid derivatives having a radiometal chelator linked to an amino acid sidechain. Preferably, the radiometal chelator is incorporated into the peptide at the carbocyl terminus of the CT receptor binding peptide. More preferably, the radiometal chelator is incorporated into the synthetic CT receptor binding peptide at the sidechain sulfur atom of a cysteine corresponding to position 14 of the native peptide. Most preferably, the radiometal chelator is incorporated into a sidechain of an amino acid of the CT receptor binding peptide having the sequence

CH₃CO.SNLS**T.Hhc**.VLGKLSCELHKLQTYPRNTGSGTP.amide (SEQ ID No. 9).

Most preferably, the radiometal chelator is incorporated into the CT receptor binding peptide at the sidechain sulfur atom of the cysteine at position 13 of the peptide of SEQ ID NO:9 (distinguished by the bold highlighting in SEQ ID NO:9).

Additional embodiments of the reagents of the invention comprise at least two synthetic CT receptor binding compounds, each compound being covalently linked to a radiometal chelator, and a polyvalent linker forming a covalent linkage selected from the group consisting of a linkage to each compound, a linkage to each chelator, and a linkage to one compound and to the chelator of the other compound. Additional permutations of this embodiment may also

occur, in accordance with the invention. Polyvalent linkers suitable for use in this embodiment of the invention comprise at least two identical functional groups capable of covalently bonding to CT analogues, CT receptor binding compounds, CT peptides or radiometal chelators, or capable of binding both to a CT receptor binding compound and to a radiometal chelator.

- 5 Preferred functional groups include, without limitation, primary amines, secondary amines, hydroxyl groups, carboxylic acid groups or thiol-reactive groups. In preferred embodiments, the polyvalent linkers comprise *bis*-succinimidylmethylether (BSME), *bis*-succinimidylethylether (BSEE), 4-(2,2-dimethylacetyl)benzoic acid (DMBA), *N*-(2-(*N'*,*N'*-*bis*(2-succinimido-ethyl)aminoethyl))-*N''*,*N''*-*bis*(2-methyl-2-mercapto-propyl)-6,9-diazanonanamide (BAT-BS), *tris*(succinimidylethyl)amine (TSEA), *bis*-succinimidohexane (BSH), 4-(O-CH₂CO-Gly-Gly-Cys.amide)-2-methylpropiophenone (ETAC), *tris*(acetamidoethyl)amine, *bis*-acetamidomethyl ether, *bis*-acetamidoethyl ether, α,ϵ -*bis*-acetyllysine, lysine and 1,8-*bis*-acetamido-3,6-dioxa-octane, or derivatives thereof.
- 10

- CT receptor binding compounds provided by the present invention can be chemically synthesized *in vitro* using any suitable synthetic method. Preferably, CT peptides may be synthesized in accordance with the invention using recombinant methods. More preferably, CT peptides, CT peptide derivatives, and CT peptide analogues can generally advantageously be prepared in accordance with the present invention using a peptide synthesizer. The CT peptides of this invention are preferably synthesized by covalently linking the radiometal chelator to the peptide during chemical synthesis *in vitro*, using techniques well known to those with skill in the art such as solid phase peptide synthesis. In this manner, radiometal chelators may be incorporated into the peptide in a site-selective fashion at virtually any position in the peptide thereby avoiding a decrease in the affinity and specificity of the peptide for the CT receptor.
- 15
- 20

- In accordance with the invention, CT receptor binding peptides are prepared having a protected thiol-containing amino acid, typically a cysteine residue, incorporated into the peptide. Following cleavage of the peptide from the synthetic resin and cyclization of the amino terminal residues, the protected thiol group is deprotected and elaborated with a prosthetic group containing a radiometal chelator and a thiol-reactive group.
- 25

- As set forth above, the CT receptor binding reagents of the invention are capable of being radiolabeled to provide radiodiagnostic or radiotherapeutic agents. An exemplary radiodiagnostic application using the radiolabeled reagents of the invention is scintigraphic
- 30

imaging, wherein the location and extent of CT receptor-bearing tumors may be determined. The term "scintigraphic imaging agent" as used herein is meant to encompass a radiolabeled reagent capable of being detected with a radioactivity detecting means (including but not limited to a gamma-camera or a scintillation detector probe). In the radiotherapeutic embodiments of the invention, CT receptor binding reagents are labeled with a cytotoxic radionuclides and are useful in the treatment of diseases or other ailments in animals, preferably humans, such diseases or ailments being characterized by expression or overexpression of CT receptors. CT-related diseases or ailments include but are not limited to breast cancer, ovarian cancer, lung cancer, lymphoma and other diseases characterized by the growth of malignant or benign tumors capable of binding CT receptor binding compounds, CT or derivatives or analogues thereof *via* the expression of CT receptors on the cell surface of cells comprising such tumors.

Any radiometal may be complexed with the reagents of the invention to provide a radiodiagnostic or radiotherapeutic agent. For example, the reagents of the invention may be radiolabeled with technetium-99m, iodine-125, or iodine-123 to provide a scintigraphic imaging agent. This invention also provides CT receptor-binding reagents capable of forming a complex with a magnetic, paramagnetic, supermagnetic, or superparamagnetic metal atom, ion or particle. The CT receptor binding reagents of the invention can also advantageously be radiolabeled with a cytotoxic radioisotope selected from the group consisting of scandium-47, copper-67, gallium-72, yttrium-90, tin-117m, iodine-125, iodine-131, samarium-153, gadolinium-159, dysprosium-165, holmium-166, ytterbium-175, lutetium-177, rhenium-186, rhenium-188, astatine-211, bismuth-212, and bismuth-213 to provide a radiotherapeutic agent.

When the reagents of the invention are used to form a complex of radioactive technetium or rhenium, the technetium complex, preferably a salt of technetium-99m pertechnetate, or rhenium in the form of perrhenate, is reacted with the reagent in the presence of a reducing agent. Preferred reducing agents are dithionite, stannous and ferrous ions; the most preferred reducing agent is stannous chloride. Alternatively, the complex may be formed by reacting a reagent of this invention with a pre-formed labile complex of technetium or rhenium and another compound known as a transfer ligand. This process is known as ligand exchange and is well known to those skilled in the art. The labile complex may be formed using such transfer ligands as tartrate, citrate, gluconate or mannitol, for example. Among the

technetium-99m pertechnetate and rhenium salts useful with the present invention are included the alkali metal salts such as the sodium salt, or ammonium salts or lower alkyl ammonium salts.

The invention is also embodied in a kit for preparing radiometal-labeled reagents for use as radiopharmaceuticals. The kit of the invention comprises a sealed vial containing a predetermined quantity of the CT receptor binding reagent, and optionally, when the radiometal is technetium-99m, rhenium-186, or rhenium-188, a reducing agent. For example, an appropriate quantity of a reagent of the invention is introduced into a vial containing a reducing agent, such as stannous chloride, in an amount sufficient to label the reagent with technetium-99m, rhenium-186 or rhenium-188. An appropriate amount of a transfer ligand as described (such as tartrate, citrate, gluconate, glucoheptanate or mannitol, for example) can also be included in the kit. The kit may also contain conventional pharmaceutical adjunct materials such as, for example, pharmaceutically acceptable salts to adjust the osmotic pressure, buffers, preservatives and the like. The components of the kit may be in liquid, frozen or dry form. In a preferred embodiment, kit components are provided in lyophilized form. Technetium-99m, rhenium-186 and rhenium-188 labeled radiopharmaceuticals according to the present invention may be prepared by the addition of an appropriate amount of technetium-99m, rhenium-186 or rhenium-188, or radionuclide complexes thereof, into the vials and reaction under conditions described in the Examples below.

The kit of the invention may be embodied in a form suitable for diagnostic imaging or as a therapeutic agent using a radioisotope of iodine, including iodine-123 and iodine-131, and preferably iodine-123. In this embodiment, the kit comprises a sealed vial containing a predetermined quantity of a CT receptor binding reagent capable of being radiolabeled with an iodine isotope. CT receptor binding reagents suitable for use in this embodiment include CT itself, a CT derivative, a CT analogue, CT mimetics and CT peptidomimetics that specifically bind to the CT receptor. When peptide and peptidomimetic CT receptor binding reagents are employed in this embodiment, a tyrosine residue in the reagent may be radioiodinated. Such a tyrosine residue may occur naturally in the peptide or peptidomimetic, or the tyrosine residue may be added at a position in the peptide or peptidomimetic that does not disrupt binding of the reagent to CT receptors. Dose, sites and routes of administration, formulations and administered

specific radioactivity using the kit of this embodiment are as described herein for technetium and rhenium-labeled reagents for scintigraphic and therapeutic uses.

The imaging agents provided by the invention have utility for tumor imaging, particularly for imaging primary and metastatic neoplastic sites characterized by neoplastic cells that express or overexpress CT receptors, and in particular such primary and especially metastatic breast, lung and ovarian tumor-derived cells that have been clinically recalcitrant to detection using conventional methodologies. The imaging reagents provided by the present invention can also be used for visualizing organs such as the kidney or bone for diagnosing disorders in these organs.

For diagnostic purposes, an effective diagnostic amount of the diagnostic or radiodiagnostic agent of the invention is administered, preferably intravenously. In radiodiagnostic embodiments, localization of the radiolabel is detected using conventional methodologies such as gamma scintigraphy. In non-radioactive diagnostic embodiments, localization of sites of accumulation of the paramagnetic metal-labeled diagnostic agents of the invention is achieved using magnetic resonance imaging methodologies.

In accordance with this invention, for scintigraphic imaging the technetium-99m labeled reagents of the invention are administered in a single unit injectable dose. The technetium-99m labeled reagents provided by the invention may be administered intravenously in any conventional medium for intravenous injection such as an aqueous saline medium, or in blood plasma medium. Generally, the unit dose to be administered has a radioactivity of about 0.01 mCi to about 100 mCi, preferably 1 mCi to 20 mCi. The solution to be injected at unit dosage is from about 0.01 mL to about 10 mL. After intravenous administration, imaging *in vivo* can take place in a matter of a few minutes. However, imaging can take place, if desired, in hours or even longer, after the radiolabeled reagent is injected into a patient. In most instances, a sufficient amount of the administered dose will accumulate in the area to be imaged within about 0.1 of an hour to permit the taking of scintiphotos. Any conventional method of scintigraphic imaging for diagnostic purposes can be utilized in accordance with this invention.

For the purposes of this invention, radiotherapy encompasses any therapeutic effect ranging from pain palliation to tumor ablation or remission of symptoms associated with the particular cancer being treated. When the reagents of the invention are used for therapeutic

purposes, they are radiolabeled with an effective amount of a cytotoxic radioisotope. For this purpose, an amount of cytotoxic radioisotope from about 10mCi to about 200mCi may be administered *via* any suitable clinical route, preferably by intravenous injection.

In accordance with this invention, effective radiodiagnostic and radiotherapeutic agents may be identified as follows. Reagents of the invention comprising CT receptor binding compounds, including CT fragments, CT peptide analogues and CT derivatives, are synthesized using the methods of the invention and a radiometal chelator is covalently linked to the compound. The reagents are then complexed with a radiometal or a non-radioactive isotope having chelation characteristic similar to the desired radiometal, and CT receptor binding is then evaluated in *in vitro* competition binding assays as described herein using radioiodinated CT. As an example of this methodology, ReO is employed to evaluate the suitability of CT receptor-binding peptides for use as 99m-technetium radiolabeled scintigraphic imaging agents, as disclosed in Example 4 below

The methods for making and labeling these compounds are more fully illustrated in the following Examples, which illustrate certain aspects of the above-described invention and advantageous results and are shown by way of illustration and not limitation.

EXAMPLE 1

Synthesis of BAT Chelators

BAT chelators, in particular S-cysteine derived and ϵ -amino Lysine derived BAT chelators, are prepared according to the methods of co-owned and co-pending U.S. Serial No. 08/414,424, incorporated by reference herein.

EXAMPLE 2

Solid Phase Peptide Synthesis

Solid phase peptide synthesis (SPPS) was carried out on a 0.25 millimole (mmole) scale using an Applied Biosystems Model 431A Peptide Synthesizer and using 9-fluorenylmethyloxycarbonyl (Fmoc) amino-terminus protection, coupling with dicyclohexylcarbodiimide/hydroxybenzotriazole or 2-(1H-benzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate/ hydroxybenzotriazole (HBTU/HOBT), and using *p*-

hydroxymethyl henoxymethyl-polystyrene (HMP) resin or Sasrin™ or chlorotriyl resin for carboxyl-terminus acids or Rink amide resin for carboxyl-terminus amides.

Where appropriate, Fmoc-Cys(BAT) and *N*α-Fmoc-*N*ε-(BAT)Lys were synthesized as described in co-owned and co-pending U.S. Serial No. 08/414,424, incorporated by reference
5 herein.

Where appropriate, 2-chloroacetyl, 2-bromoacetyl and 2-bromo-3-phenylpropionyl groups are introduced either by using the appropriate 2-halo acid as the last residue coupled during SPPS, or by treating the N-terminus free amino acid peptide bound to the resin with either 2-halo acid/ diisopropylcarbodiimide/*N*-hydroxysuccinimide/NMP or 2-halo acid
10 anhydride/ diisopropylethylamine/NMP.

Where appropriate, HPLC-purified 2-haloacylated peptides are cyclized by stirring an 0.1-1.0 mg/mL solution in phosphate or bicarbonate buffer or dilute ammonium hydroxide (pH 8.0), optionally containing 0.5-1.0 mM EDTA, or acetonitrile or THF for 1-48 h followed optionally by acidification with acetic acid, lyophilization and HPLC purification.

Where appropriate, thiol-containing peptides are reacted with chloroacetyl-containing, thiol-protected Technetium-99m complexing moieties at pH 10 for 0.5-4 hours at room temperature, followed by acetic acid acidification and evaporation of the solution to give the corresponding peptide-sulfide adduct. Deprotection and purification are routinely performed as described to yield the chelator-peptide conjugate.
15

Where appropriate, BSME, BSEE and BSH adducts are prepared by reacting single thiol-containing peptides (5 to 50 mg/mL in DMF buffered to pH 7 with *N*-methylmorpholine or *N*-ethyl-morpholine, or 50mM sodium phosphate buffer, pH 7-8, optionally containing 0.5mM EDTA or DMF or THF or acetonitrile) with 0.5 molar equivalents of BMME (*bis*-maleimidomethylether), BMEE (*bis*-maleimidoethylether) or BMH (*bis*-maleimidohexane),
20 respectively, pre-dissolved in acetonitrile at room temperature for approximately 1-18 hours. The solution was concentrated and the product was purified by HPLC.

Where appropriate, TSEA adducts are prepared by reacting single thiol-containing peptide (at concentrations of 10 to 100 mg/mL peptide in DMF buffered to pH 7 with *N*-methylmorpholine or *N*-ethylmorpholine, or 5 to 50 mg/mL peptide in 50mM sodium
30 phosphate, pH 7-8, optionally containing 0.5mM EDTA or DMF or THF or acetonitrile) with 0.33 molar equivalents of TMEA (*tris*(2-maleimidoethyl)amine) pre-dissolved in acetonitrile or

DMF, with or without 1 molar equivalent of triethanolamine, at room temperature for approximately 1-18h. Such reaction mixtures containing adducts are concentrated and the adducts are then purified using HPLC.

Where appropriate, (BAM) (*N',N'*-bis(2-mercapto-2-methylpropyl)-1,4,10-triazadecane) is conjugated to the peptide by first activating the peptide carboxylate with a mixture of diisopropylcarbodiimide/ N-hydroxysuccinimide or HBTU/HOBt in DMF, NMP or methylene chloride, followed by coupling in the presence of diisopropylethylamine. After coupling, the conjugates are deprotected as described above.

Where appropriate, (BAT) (*N',N'*-bis(2-mercapto-2-methylpropyl)-6,9-diazanonanoic acid) is incorporated into peptides as protected amino acid derivatives, such as (*N*α(Fmoc)-*N*ε(*N*-Boc)-*S,S'*-bistrityl-BAT)lysine (prepared from *N*α(Fmoc)-lysine and *N*ε(*N*-Boc)-*S,S'*-bistrityl-BAT as described in Example 2 of co-owned and co-pending U.S. Patent Application Serial No. 08/044,825, incorporated by reference), or as (*N*(Fmoc)-*S,S'*-bistrityl-BAT)cysteine (prepared as described in Example 1F of co-owned and copending U.S. Serial No.8/414,424, incorporated by reference) during peptide synthesis and then deprotected after cleavage of the completed peptide from the synthetic resin.

Where appropriate, BAT-BS (*N*-{2-(*N',N'*-bis(2-succinimidoethyl) aminoethyl)}-*N',N'*-bis(2-methyl-2-mercaptoethyl)-6,9-diazanonanamide) adducts are prepared by reacting single thiol-containing peptide (at concentrations of 2 to 50 mg/mL peptide in DMF buffered to pH 7 with *N*-methylmorpholine or *N*-ethylmorpholine, or in 50mM sodium phosphate (pH 7-8), optionally containing 0.5mM EDTA or DMF or THF or acetonitrile) with 0.5 molar equivalents of BAT-BM (*N*-{2-(*N',N'*-bis(2-maleimidoethyl)aminoethyl)}-*N'*-(*t*-butoxycarbonyl)-*N',N'*-bis(2-methyl-2-triphenylmethylthiopropyl)-6,9-diazanonanamide) pre-dissolved in acetonitrile or THF, at room temperature for approximately 1-18h. The solution is then evaporated to dryness and (BAT-BS)-peptide conjugates deprotected by treatment with 10mL TFA and 0.2mL triethylsilane for 1h. The solution is concentrated, the product adducts precipitated with ether, and then purified by HPLC.

Where appropriate, peptide precursors are cyclized (between the amino- and carboxyl-termini) by reaction of the sidechain-protected, N-terminal free amine and C-terminal free acid with diphenylphosphoryl azide.

SasrinTM resin-bound peptides are cleaved using a solution of 1% TFA in dichloromethane to yield the protected peptide. Where appropriate, protected peptide precursors are cyclized between the amino- and carboxyl-termini by reaction of sidechain-protected, amino-terminal free amine and carboxyl-terminal free acid using diphenylphosphorylazide.

HMP or Rink amide resin-bound products are routinely cleaved and protected cyclized peptides deprotected using a solution comprised of trifluoroacetic acid (TFA), or TFA and methylene chloride, optionally comprising water, thioanisole, ethanedithiol, and triethylsilane or triisopropylsilane in ratios of 100 : 5 : 5 : 2.5 : 2, for 0.5 - 3 hours at room temperature. Where appropriate, products were re-*S*-tritylated in triphenolmethanol/ TFA, and *N*-Boc groups re-introduced into the peptide using (Boc)₂O.

Where appropriate, thiol functionalities within the peptide or peptidomimetic sequence designed for further elaboration with a prosthetic group were protected using compound such as *S*-*t*-butyl (to produce mixed *t*-butyl disulfides) or *p*-methoxybenzyl. *S*-*t*-butyl groups are removed by treatment with a solution of dithiothreitol or mercaptoethanol, while *p*-methoxybenzyl groups are removed using boron trifluoride etherate in trifluoroacetic acid in the presence of a free radical scavenger such as *m*-cresol. Prosthetic peptides containing radiometal binding moieties are prepared by SPPS ending with an N-terminal 2-haloacetyl group. The prosthetic group are removed from the resin and any thiol groups are-protected, *for example*, with a trityl group. The haloacetyl sequence is then coupled with the thiol-containing peptide under essentially the same conditions as described above for preparing cyclic thioethers. Removal of remaining protecting groups is then achieved using the methods described herein to yield the final product.

Crude peptides are purified by preparative high pressure liquid chromatography (HPLC) using a Waters Delta Pak C18 column and gradient elution using 0.1% trifluoroacetic acid (TFA) in water modified with acetonitrile. Acetonitrile is evaporated from the eluted fractions which are then lyophilized. The identity of each product is confirmed by fast atom bombardment mass spectroscopy (FABMS) or by electrospray mass spectroscopy (ESMS).

CT receptor binding peptides, derivatives and analogues synthesized as provided herein, as well as the products of such synthesis identified by ESMS or FABMS, are shown in Table I below.

TABLE I

	Peptide	Peptide	ReO Complex
1.	<u>CH₂CO.SNLS.T.Hhc.</u> VLGKLS(C(BA ¹))ELHKLQTYPRNTGSGTP.amide	3679*	3877*
2.	<u>CH₂CO.SNLS.T.Hhc.</u> VLGKLSQELHKLQTYPRNTGSGTP(ε-K)GC.amide	3687*	3886
3.	<u>CH₂CO.SNLS.T.Hhc.</u> VLGKLSCELHKLQTYPRNTGSGTP.amide	3373*	NA
4.	<u>CH₂CO.SNLS.T.Hhc.</u> VLGKLS(C(CH ₂ CO.GGCK.amide)ELHKLQTYPRNTGSGTP.amide	3873*	4078
5.	<u>CH₂CO.SNLS.T.Hhc.</u> VLGKLS(C(CH ₂ CO.(β-Dap)KCK.amide)ELHKLQTYPRNTGSGTP.amide	3777*	3975
6.	<u>CH₂CO.SNLS.T.Hhc.</u> VLGKLS(C(CH ₂ CO.(ε-K)GCE.amide)ELHKLQTYPRNTGSGTP.amide	3849*	4047
7.	<u>CH₂CO.SNLS.T.Hey.</u> VLGKLSCELHKLQTYPRNTGSGTP.amide	3360*	NA
8.	<u>CH₂CO.SNLS.T.Hey.</u> VLGKLS(C(CH ₂ CO.GGCK.amide)ELHKLQTYPRNTGSGTP.amide	3762*	3962
9.	<u>CH₂CO.SNLS.T.Hey.</u> VLGKLS(C(CH ₂ CO.(β-Dap)KCK.amide)ELHKLQTYPRNTGSGTP.amide	3862*	4063
10.	<u>CH₂CO.SNLS.T.Hey.</u> VLGKLS(C(CH ₂ CO.(ε-K)GCE.amide)ELHKLQTYPRNTGSGTP.amide	3835	4033
11.	<u>CH₂CO.SNLS.T.Cys.</u> VLGKLSCELHKLQTYPRNTGSGTP.amide	3346*	NA
12.	<u>CH₂CO.SNLS.T.Cys.</u> VLGKLS(C(CH ₂ CO.GGCK.amide)ELHKLQTYPRNTGSGTP.amide	3748*	3947
13.	<u>CH₂CO.SNLS.T.Cys.</u> VLGKLS(C(CH ₂ CO.(β-Dap)KCK.amide)ELHKLQTYPRNTGSGTP.amide	3848*	4048
14.	<u>CH₂CO.SNLS.T.Cys.</u> VLGKLS(C(CH ₂ CO.(ε-K)GCE.amide)ELHKLQTYPRNTGSGTP.amide	3820	4019
15.	<u>SNLS.T.Asu.</u> VLGKLSCELHKLQTYPRNTGSGTP.amide	3356	NA
16.	<u>SNLS.T.Asu.</u> VLGKLS(C(CH ₂ CO.(β-Dap)KCK.amide)ELHKLQTYPRNTGSGTP.amide	3858	4057

Peptide	Peptide	ReO Complex
17. <u>SNLST</u> .Asu.VLGKLSCELHKLQTYPRTDVGAGTP.amide	3338	NA
18. <u>SNLST</u> .Asu.VLGKLSCT(ch.co.(β-Dap)KCK.amide)ELHKLQTYPRTDVGAGTP.amide	3841	4040

* = M^{*} determined by electrospray mass spectrometry

M^{*} determined by fast atom bombardment mass spectrometry for all other peptides.

EXAMPLE 3

A General Method for Radiolabeling with Technetium-99m

5 0.1 mg of a peptide prepared as in Example 2 was dissolved in 0.1 mL or 0.2 mL of water or 0.9% saline. Technetium-99m gluceptate was prepared by reconstituting a Glucoscan vial (E.I. DuPont de Nemours, Inc., Wilmington, DE) with 0.25 mL of Technetium-99m sodium pertechnetate containing up to 200 mCi and allowed to stand for 15 minutes at room temperature. 25 μ L of Technetium-99m gluceptate was then added to the peptide and the
10 reaction allowed to proceed at room temperature for 15 to 60 min or at 100°C for 10 to 30 min, and then filtered through a 0.2 μ m filter.

 The Technetium-99m labeled peptide purity was determined by reverse-phase HPLC using the following conditions: a Waters Delta Pak C-18, 5 μ , 3.9mm x 150mm analytical column was loaded with each radiolabeled peptide, and the peptides eluted at a solvent flow rate
15 equal to 1 mL/min (Delta-Pak). Gradient elution was performed using a gradient of 20-50% Solvent B/Solvent A (Solvent A is 0.1% CF₃COOH in water and Solvent B is 0.1% CF₃COOH in 90/10 CH₃CN/H₂O) for 20 min., followed by 100% B/A for 3 min.

 Radioactive components were detected using an in-line radiometric detector linked to an integrating recorder. Technetium-99m gluceptate and Technetium-99m sodium pertechnetate
20 elute between 1 and 4 minutes under these conditions, whereas the Technetium-99m labeled peptides eluted after a much greater amount of time. Peptides were detected by in-line spectrophotometric detection at 220nm.

 Non-radioactive rhenium complexes were prepared by co-dissolving each of the peptide reagents of the invention with about one molar equivalent of tetrabutylammonium
25 oxotetrabromorhenate (+5), prepared as described by Cotton *et al.* (1966, *Inorg. Chem.* 5: 9-16) in dimethylformamide or acetonitrile/water and stirred for 0.5-5 days. The rhenium complexes were isolated by reverse phase HPLC as described above for Technetium-99m labeled peptides and were characterized by FABMS or ESMS. Non-radioactive peptides were detected as peptides by in-line spectrophotometric detection at 220nm.

30 Radioactive rhenium complexes, using for example Re-186 or Re-188, are prepared from the appropriate perrhenate salts using the same protocol as for Technetium-99m labeling,

or by adding a reducing agent to a solution of the peptide and perrhenate, or optionally using a ligand transfer agent such as citrate and incubating the reaction at a temperature between room temperature and 100°C for between 5 and 60 min.

Results of HPLC purification of peptides, Technetium-99m labeled peptides and ReO-
5 complexed peptides are shown in Table II.

TABLE II

	Peptide	Peptide	ReO Complex
1.	CH ₂ CO.SNLST.Hhc.VLGKLS(BAT)ELHKLQTYPRNTGSGTP.amide	13.7	15.6
2.	CH ₂ CO.SNLST.Hhc.VLGKLSQELHKLQTYPRNTGSGTP(e-K)GC.amide	15.0	13.5
3.	CH ₂ CO.SNLST.Hhc.VLGKLSCEI.HKI.QTYPRNTGSGTP.amide	19.2	NA
4.	CH ₂ CO.SNLST.Hhc.VLGKLS(CCH ₂ CO.GGCK.amide)ELHKLQTYPRNTGSGTP.amide	21.2	17.8
5.	CH ₂ CO.SNLST.Hhc.VLGKLS(CCH ₂ CO.(β-Dap)KCK.amide)ELHKLQTYPRNTGSGTP.amide	12.1	12.9
6.	CH ₂ CO.SNLST.Hhc.VLGKLS(CCH ₂ CO.(ε-K)GCE.amide)ELHKLQTYPRNTGSGTP.amide	11.9	12.4
7.	CH ₂ CO.SNLST.Hcy.VLGKLSCEI.HKLQTYPRNTGSGTP.amide	13.1	NA
8.	CH ₂ CO.SNLST.Hcy.VLGKLS(CCH ₂ CO.GGCK.amide)ELHKLQTYPRNTGSGTP.amide	12.4	13.5
9.	CH ₂ CO.SNLST.Hcy.VLGKLS(CCH ₂ CO.(β-Dap)KCK.amide)ELHKLQTYPRNTGSGTP.amide	11.0	11.8
10.	CH ₂ CO.SNLST.Hcy.VLGKLS(CCH ₂ CO.(ε-K)GCE.amide)ELHKLQTYPRNTGSGTP.amide	12.5	13.1
11.	CH ₂ CO.SNLST.Cys.VLGKLSCEI.HKLQTYPRNTGSGTP.amide	10.7	NA
12.	CH ₂ CO.SNLST.Cys.VLGKLS(CCH ₂ CO.GGCK.amide)ELHKLQTYPRNTGSGTP.amide	9.98	11.1
13.	CH ₂ CO.SNLST.Cys.VLGKLS(CCH ₂ CO.(β-Dap)KCK.amide)ELHKLQTYPRNTGSGTP.amide	8.23	9.2
14.	CH ₂ CO.SNLST.Cys.VLGKLS(CCH ₂ CO.(ε-K)GCE.amide)ELHKLQTYPRNTGSGTP.amide	9.98	10.7
15.	SNLST.Asu.VLGKLSCEI.HKLQTYPRNTGSGTP.amide	12.7	NA
16.	SNLST.Asu.VLGKLS(CCH ₂ CO.(β-Dap)KCK.amide)ELHKLQTYPRNTGSGTP.amide	10.7	11.4

Peptide	Peptide	ReO Complex
17. <u>SNLST</u> <u>Asu</u> .VLGKLSCELHKLQTYPRTDVGAGTP.amide	13.2	NA
18. <u>SNLST</u> <u>Asu</u> .VLGKLSCE(ch ₃ co.(β-Dap)KCK.amide)ELHKLQTYPRTDVGAGTP.amide	11.1	11.7

Data represents HPLC retention times in minutes.

EXAMPLE 4

Biological Assays

A. Binding Inhibition Assays

Peptide reagents within the scope of the invention, or ReO complexed embodiments thereof, were tested using *in vitro* assays that measure their ability to inhibit the specific binding of ^{125}I -CT to the CT receptor, using membranes from rat brain or from a breast tumor cell line, and on whole cells from breast tumor cell lines, as described in detail below.

Assays using microsomal membrane fractions of rat brain and T-47D cells (obtained from the American Type Culture Collection, Rockville, MD, ATCC Accession No. HTB-133) were used to identify analogs with high affinity for the CT receptor according to the method of Fisher *et al.* (1977, *British J. Cancer* 35: 777-784). Briefly, tissue was minced and homogenized. The membranes were washed several times, assayed for protein content and used in the binding assay. Membrane protein was incubated with 0.1 μCi of ^{125}I -salmon CT (Amersham, Cleveland, OH) in the presence or absence of varying concentrations of reagents to be tested. Both uncomplexed and ReO complexed reagents were tested. One hundred percent specific binding of ^{125}I -CT to the CT receptor was defined as the difference between total ^{125}I -CT binding and nonspecific binding of ^{125}I -CT measured in the presence of a receptor-saturating concentration (1 μM) of excess unlabeled salmon CT (Sigma, St. Louis, MO). The concentration at which the tested reagents inhibited specific binding of ^{125}I -CT by 50% was defined as the IC_{50} .

Results of binding to membrane preparations of CT receptors for each of the tested reagents are shown in Table III. These results indicate that peptide reagents within the scope of the invention and ReO complexes of such reagents bind with high affinity to CT receptor-expressing tumor cell and brain membranes.

Similar experiments were performed with additional peptide reagents within the scope of the invention, using whole T-47D cells and MCF-7 cells (ATCC Accession No. HTB-22). The cell binding assays were done essentially by the method of Findlay *et al.* (1990, *J. Endocrinol.* 130: 321-326). Briefly, cells were washed in saline and resuspended in Hank's balanced salt solution. One to two million cells were incubated with 0.1 μCi ^{125}I -salmon CT in the absence and presence of 1 μM unlabeled salmon CT to determine a 100% specific binding

value. Cells were incubated with 0.1 μCi of ^{125}I -CT in the presence or absence of varying concentrations of reagents to be tested, in uncomplexed and/or ReO-complexed form. IC_{50} values were determined as described above. Results for each of the tested reagents are shown in Table IV.

TABLE III
Displacement of 125 I-CT from CTR in T-47D and Rat Brain Membranes
by CT-Mimetic Peptides

	Peptide Structure	T-47D			Rat brain	
		IC ₅₀	K _i	K _i	IC ₅₀	K _i
1.	<u>(CH₂CO.SNLST.Hhc).VLGKLSC(BAT)ELHKLQTYPRNTGSGTP.amide</u>	0.53	0.41	2.6	2.6	ND
2.	<u>(CH₂CO.SNLST.Hhc).VLGKLSQELHKLQTYPRNTGSGTP(ε-K)GC.amide</u>	16	12	18	18	ND
1.	<u>(CH₂CO.SNLST.Hhc).VLGKLSC(BAT)ELHKLQTYPRNTGSGTP.amide(ReO)</u>	1.5	1.2	3.0	3.0	ND
2.	<u>(CH₂CO.SNLST.Hhc).VLGKLSQELHKLQTYPRNTGSGTP(ε-K)GC.amide(ReO)</u>	28	22	22	22	ND

TABLE IV
Displacement of 125 I-CT from T47D (Cells and Membranes) and MCF-7 Cells

Sequence	IC ₅₀ (nM)	IC ₅₀ (nM)	IC ₅₀ (nM)
3. <u>CH₂CO.SNLST.Hhc.VLGKLSCELHKLQTYPRNTGSGTP.amide</u>	1.6	nd	nd
4. <u>CH₂CO.SNLST.Hhc.VLGKLSCH(CH₂CO.GGCK.amide)ELHKLQTYPRNTGSGTP.amide</u>	2.6	nd	nd
5. <u>CH₂CO.SNLST.Hhc.VLGKLSCH(CH₂CO.(β-Dap)KCK.amide)ELHKLQTYPRNTGSGTP.amide•ReO</u>	nd	0.42	5.8
6. <u>CH₂CO.SNLST.Hhc.VLGKLSCH(CH₂CO.(ε-K)GCE.amide)ELHKLQTYPRNTGSGTP.amide•ReO</u>	nd	0.33	3.3
8. <u>CH₂CO.SNLST.Hcy.VLGKLSCH(CH₂CO.GGCK.amide)ELHKLQTYPRNTGSGTP.amide•ReO</u>	nd	0.83	4.8
9. <u>CH₂CO.SNLST.Hcy.VLGKLSCH(CH₂CO.(β-Dap)KCK.amide)ELHKLQTYPRNTGSGTP.amide•ReO</u>	nd	0.84	nd
10. <u>CH₂CO.SNLST.Hcy.VLGKLSCH(CH₂CO.(ε-K)GCE.amide)ELHKLQTYPRNTGSGTP.amide•ReO</u>	nd	0.89	nd
12. <u>CH₂CO.SNLST.Cys.VLGKLSCH(CH₂CO.GGCK.amide)ELHKLQTYPRNTGSGTP.amide•ReO</u>	nd	1.0	5.2
13. <u>CH₂CO.SNLST.Cys.VLGKLSCH(CH₂CO.(β-Dap)KCK.amide)ELHKLQTYPRNTGSGTP.amide•ReO</u>	nd	2.9	nd
14. <u>CH₂CO.SNLST.Cys.VLGKLSCH(CH₂CO.(ε-K)GCE.amide)ELHKLQTYPRNTGSGTP.amide•ReO</u>	nd	2.6	1.9
16. <u>SNLST.Asu.VLGKLSCH(CH₂CO.(β-Dap)KCK.amide)ELHKLQTYPRNTGSGTP.amide•ReO</u>	nd	2.7	8.0
18. <u>SNLST.Asu.VLGKLSCH(CH₂CO.(β-Dap)KCK.amide)ELHKLQTYPRTDVGAGTP.amide•ReO</u>	nd	0.55	7.0
<u>CSNLSTCVLGKLSQELHKLQTYPRNTGSGTP-NH₂ (salmon calcitonin)</u>	nd	nd	12.3

nd = not done

wherein:
x T-47D cell membranes
y Whole T-47D cells..
z Whole MCF-7 cells

These results indicated that peptide reagents within the scope of the invention and ReO complexes of such peptides were capable of specifically binding to CT receptors on whole cells and that the reagents were potent inhibitors of CT binding in two different CT receptor-expressing breast tumor cell lines.

B. CT Receptor Expression in Breast Cancer Cell Lines

The binding of ^{125}I -CT to whole cells was used to assess the CT receptor density in seven different human breast cancer cell lines. The site density per cell was determined for MCF-7 cells in the presence of different concentrations of CT to achieve saturation of the receptors. The data was then linearized by the method of Scatchard *et al.* (1949, N.Y. Acad. Sci. 51: 600-672) to estimate receptor density. Briefly, the saturation curves were linearized and the K_d was calculated as $(-1/\text{slope})$, and the B_{max} was equal to the x-axis intercept of the curve. Other cell lines (each of which was obtained from the ATCC, Rockville, MD) were compared to MCF-7 cells at a single concentration of ^{125}I -CT and their CT receptor density estimated thereby. The data are summarized in Table V.

Table V

Cell Line	CT Receptors/Cell x 10^3
Du4475	21.5
ZR75	39.5
MDA-MB-157	51.0
MCF-7	1,000
Bt474	166
HS578T	0
T-47D	360

These data show that 86% (6 of 7) of the breast cancer cell lines assayed were positive for CT receptors. Thus, the target receptor for the reagents of the invention is present in most breast cancer cell lines tested. Recently, the target receptor for the reagents of the invention was found to be overexpressed in primary breast cancers (Gillespie, *et al.* (1997) *Int. J. Cancer* **73**,

812-815). These results indicate that the CT peptides of the invention are appropriate for preparing useful site specific reagents for diagnosis and treatment of tumors in humans.

C. *In vivo* Assay

Breast cancer cell lines were screened for receptor expression and the best cell lines chosen for xenograft implantation in immune deficient mice. These tumor models were used to evaluate the tumor targeting potential *in vivo* of the high affinity peptides identified in the *in vitro* assays above.

CT receptor expressing breast cancer cells (T47D and MCF-7) were implanted into immune deficient mice or Sprague-Dawley rats and allowed to grow tumors. For testing new ^{99m}Tc -CT peptides, tumor-bearing mice or rats were injected with approximately 0.025 mCi at approximately 6 mCi/10 nmol peptide. The animals were cervically dislocated and imaged statically for 5 minutes using a gamma camera. The biodistribution of the ^{99m}Tc -CT peptide was then determined by counting blood, tumor, target organs and muscle in a gamma counter along with standard aliquots of the injected dose. The time points for biodistribution were chosen to represent early, middle and late phases of ^{99m}Tc -CT analog clearance. Biodistribution studies indicated that optimal signal to noise ratios occurred at 90 minutes post-injection. To assess the tumor imaging potential of selected analogs, the ratio of radioactivity in blood and various tissues were compared to that in tumor. Representative biodistribution data after 90 minutes are shown in Table VI below.

TABLE VI

Peptide	Species	Tumor			Biodistribution					PK (min)	
		(%ID/g)	Tum:Bld	Tum:Con	Blood (% ID/g)	Lung (% ID)	GI tract (% ID)	Liver (% ID)	Kidney s (% ID)	t1/2α	t1/2β
8	mouse ^b	0.42	2.8	2.8	0.16	1.23 %ID/g					
8	rat				0.026	0.10 %ID/g	4.8	3.7	45.6	1.6?	20?
2	mouse ^a	1.6	3.3	10		0.24	5.4	5.5	18		
4	rat				0.027	0.53	2.0	4.0	35	1.4	19
5	rat				0.048	0.56	4.5	5.8	31	1.6	17
6	rat				0.044	0.10	1.8	2.5	32.2	1.6	21
6	rabbit				0.012	0.20	5.3	2.3	14.1		
6 ^c	mouse ^b	2.0	9.9	25	0.23	2.5 %ID/g				3.1	25.9
6 ^c	mouse ^b	3.9	4.2	2.5	0.94	6.5 %ID/g		2.7	53		
6	mouse ^a	1.5	1.6	2.1	0.91	2.3 %ID/g		1.5	27		

Peptide numbering corresponds to that of Tables I through IV.

^aT-47D tumor in nude mouse

^bMCF-7 breast cancer xenograft in nude mouse

^cDifferent preparations

It should be understood that the foregoing disclosure emphasizes certain specific embodiments of the invention and that all modifications or alternatives equivalent thereto are within the spirit and scope of the invention as set forth in the appended claims.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: Dean, Richard T.
Bush, Larry R.
Pearson, Daniel A.
Lister-James, John
- (ii) TITLE OF INVENTION: CALCITONIN RECEPTOR BINDING REAGENTS
- (iii) NUMBER OF SEQUENCES: 11
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: Diatide, Inc.
 - (B) STREET: 9 Delta Drive
 - (C) CITY: Londonderry
 - (D) STATE: NH
 - (E) COUNTRY: USA
 - (F) ZIP: 03053
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Floppy disk
 - (B) COMPUTER: IBM PC compatible
 - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 - (D) SOFTWARE: PatentIn Release #1.0, Version #1.30
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER:
 - (B) FILING DATE:
 - (C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER: US 08/847,007
 - (B) FILING DATE: 01-MAY-1997
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: McDaniels, Patricia A.
 - (B) REGISTRATION NUMBER: 33,194
 - (C) REFERENCE/DOCKET NUMBER: DITI 125.1PCT
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: (603) 437-8970
 - (B) TELEFAX: (603) 437-8977

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 32 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS:
 - (D) TOPOLOGY: both
- (ii) MOLECULE TYPE: peptide
- (vi) ORIGINAL SOURCE:
 - (A) ORGANISM: CALCITONIN--HUMAN
- (ix) FEATURE:
 - (A) NAME/KEY: Disulfide-bond

(B) LOCATION: 1..7

(ix) FEATURE:

(A) NAME/KEY: Modified-site

(B) LOCATION: 32

(D) OTHER INFORMATION: /product= "PROLINE 32"

/label= amide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

Cys	Gly	Asn	Leu	Ser	Thr	Cys	Met	Leu	Gly	Thr	Tyr	Thr	Gln	Asp	Phe
1				5					10					15	

Asn	Lys	Phe	His	Thr	Phe	Pro	Gln	Thr	Ala	Ile	Gly	Val	Gly	Ala	Pro
			20				25						30		

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 32 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS:

(D) TOPOLOGY: both

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:

(A) ORGANISM: CALCITONIN--SALMON

(ix) FEATURE:

(A) NAME/KEY: Disulfide-bond

(B) LOCATION: 1..7

(ix) FEATURE:

(A) NAME/KEY: Modified-site

(B) LOCATION: 32

(D) OTHER INFORMATION: /product= "PROLINE 32"

/label= amide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Cys	Ser	Asn	Leu	Ser	Thr	Cys	Val	Leu	Gly	Lys	Leu	Ser	Gln	Glu	Leu
1				5					10					15	

His	Lys	Leu	Gln	Thr	Tyr	Pro	Arg	Thr	Asn	Thr	Gly	Ser	Gly	Thr	Pro
			20				25					30			

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 32 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS:

(D) TOPOLOGY: both

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:

(A) ORGANISM: CALCITONIN--EEL

(ix) FEATURE:

(A) NAME/KEY: Disulfide-bond
(B) LOCATION: 1..7

(ix) FEATURE:

(A) NAME/KEY: Modified-site
(B) LOCATION: 32
(D) OTHER INFORMATION: /product= "PROLINE 32"

/label= amide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

Cys	Ser	Asn	Leu	Ser	Thr	Cys	Val	Leu	Gly	Lys	Leu	Ser	Gln	Glu	Leu
1				5					10					15	
His	Lys	Leu	Gln	Thr	Tyr	Pro	Arg	Thr	Asp	Val	Gly	Ala	Gly	Thr	Pro
			20					25					30		

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 31 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS:
(D) TOPOLOGY: both

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

(A) NAME/KEY: Region
(B) LOCATION: 1..6
(D) OTHER INFORMATION: /product= "S1 is acetylated and X 6

is Hhc"

/label= cyclized

/note= "The peptide is cyclized by a thioether bond between the N-terminal acetyl group of serine 1 and the thiol of homohomocysteine 6"

(ix) FEATURE:

(A) NAME/KEY: Modified-site
(B) LOCATION: 13
(D) OTHER INFORMATION: /product= "C 13"

/label= BAT

/note= "The thiol group of cysteine 13 is attached to a bisamino bithiol chelator "

(ix) FEATURE:

(A) NAME/KEY: Modified-site
(B) LOCATION: 31
(D) OTHER INFORMATION: /product= "Proline 31"

/label= amide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Ser	Asn	Leu	Ser	Thr	Xaa	Val	Leu	Gly	Lys	Leu	Ser	Cys	Glu	Leu	His
1				5					10					15	

Lys Leu Gln Thr Tyr Pro Arg Thr Asn Thr Gly Ser Gly Thr Pro
 20 25 30

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 31 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: both

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

- (A) NAME/KEY: Region
- (B) LOCATION: 1..6
- (D) OTHER INFORMATION: /product= "S1 is acetylated and X6

is Hhc"

/label= cyclized

/note= "The peptide is cyclized by a thioether linkage between
 the N-terminal acetyl group of serine 1 and the thiol of
 homohomocysteine 6"

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 31
- (D) OTHER INFORMATION: /product= "PROLINE 31"

/label= amide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Ser Asn Leu Ser Thr Xaa Val Leu Gly Lys Leu Ser Cys Glu Leu His
 1 5 10 15

Lys Leu Gln Thr Tyr Pro Arg Thr Asn Thr Gly Ser Gly Thr Pro
 20 25 30

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 31 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: both

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

- (A) NAME/KEY: Region
- (B) LOCATION: 1..6
- (D) OTHER INFORMATION: /product= "S1 is acetylated and X6

is Hcy"

/label= cyclized

/note= "The peptide is cyclized by a thioether linkage between
 the N-terminal acetyl group of serine 1 and the thiol of
 homocysteine 6."

(ix) FEATURE:

(A) NAME/KEY: Modified-site
 (B) LOCATION: 31
 (D) OTHER INFORMATION: /product= "Proline 31"
 /label= amide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Ser	Asn	Leu	Ser	Thr	Xaa	Val	Leu	Gly	Lys	Leu	Ser	Cys	Glu	Leu	His
1				5					10					15	
Lys	Leu	Gln	Thr	Tyr	Pro	Arg	Thr	Asn	Thr	Gly	Ser	Gly	Thr	Pro	
			20					25					30		

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 31 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: both

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

(A) NAME/KEY: Region
 (B) LOCATION: 1..6
 (D) OTHER INFORMATION: /product= "S1 is acetylated"

/label= cyclized

/note= "The peptide is cyclized by a thioether bond between the N-terminal acetyl group of serine 1 and the thiol of cysteine 6."

(ix) FEATURE:

(A) NAME/KEY: Modified-site
 (B) LOCATION: 31
 (D) OTHER INFORMATION: /product= "Proline 31"

/label= amide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

Ser	Asn	Leu	Ser	Thr	Cys	Val	Leu	Gly	Lys	Leu	Ser	Cys	Glu	Leu	His
1				5					10					15	
Lys	Leu	Gln	Thr	Tyr	Pro	Arg	Thr	Asn	Thr	Gly	Ser	Gly	Thr	Pro	
			20					25					30		

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 31 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: both

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

(A) NAME/KEY: Region

(B) LOCATION: 1..6
 (D) OTHER INFORMATION: /product= "X6 is Asu"
 /label= cyclized
 /note= "The peptide is cyclized by a bond between serine 1 and
 2-amino suberic acid 6. "

(ix) FEATURE:
 (A) NAME/KEY: Modified-site
 (B) LOCATION: 31
 (D) OTHER INFORMATION: /product= "Proline 31"
 /label= amide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Ser	Asn	Leu	Ser	Thr	Xaa	Val	Leu	Gly	Lys	Leu	Ser	Cys	Glu	Leu	His
1				5					10					15	
Lys	Leu	Gln	Thr	Tyr	Pro	Arg	Thr	Asn	Thr	Gly	Ser	Gly	Thr	Pro	
			20					25					30		

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 31 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: both

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:
 (A) NAME/KEY: Region
 (B) LOCATION: 1..6
 (D) OTHER INFORMATION: /product= "S1 is acetylated and X6
 is Hhc"
 /label= cyclized
 /note= "The peptide is cyclized by a thioether bond between the
 N-terminal acetyl group of serine 1 and the thiol of
 homohomocysteine 6."

(ix) FEATURE:
 (A) NAME/KEY: Modified-site
 (B) LOCATION: 31
 (D) OTHER INFORMATION: /product= "Proline 31"
 /label= amide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

Ser	Asn	Leu	Ser	Thr	Xaa	Val	Leu	Gly	Lys	Leu	Ser	Cys	Glu	Leu	His
1				5					10					15	
Lys	Leu	Gln	Thr	Tyr	Pro	Arg	Thr	Asn	Thr	Gly	Ser	Gly	Thr	Pro	
			20					25					30		

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 6 amino acids
 (B) TYPE: amino acid

- (C) STRANDEDNESS:
- (D) TOPOLOGY: circular

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

- (A) NAME/KEY: Region
- (B) LOCATION: 1..6
- (D) OTHER INFORMATION: /product= "S1 is acetylated and X6 is C, Hcy, or Hhc"
/label= cyclized
/note= "The peptide is cyclized by a thioether bond between the N-terminal acetyl group of serine 1 and the thiol of X6"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Ser Asn Leu Ser Thr Xaa
1 5

(2) INFORMATION FOR SEQ ID NO:11:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 6 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS:
 - (D) TOPOLOGY: circular

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

- (A) NAME/KEY: Region
- (B) LOCATION: 1..6
- (D) OTHER INFORMATION: /product= "X1 is A,G, or S; X6 is C, Hcy, or Hhc"
/label= cyclized
/note= "The peptide is cyclized by a thioether bond between the N-terminal methylene and the thiol of X6."

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

Xaa Asn Leu Ser Thr Xaa
1 5

We claim:

1. A synthetic, calcitonin receptor-binding compound having a molecular weight of less than about 10,000 daltons and being covalently linked to a radiometal chelator to form a reagent characterized in that:

said reagent has a binding affinity for a calcitonin receptor equal to or greater than the binding affinity of radioiodinated native calcitonin for said receptor.

2. A synthetic, calcitonin receptor-binding compound having a molecular weight of less than about 10,000 daltons and being covalently linked to a radiometal chelator to form a reagent characterized in that:


said reagent has a binding affinity for a calcitonin receptor of not less than about one-tenth the binding affinity of radioiodinated native calcitonin for said receptor.

3. The reagent of any of claims 1 or 2, wherein the chelator is selected from the group consisting of:

a) $C(pgp)^5-(aa)-C(pgp)^5$

wherein $(pgp)^5$ is H or a thiol protecting group, and (aa) is any primary α - or β -amino acid;

b) a chelator comprising a single thiol-containing moiety of formula:



wherein A is H, HOOC, H_2NOC , (amino acid or peptide)-NHOC, (amino acid or peptide)-OOC or R^4 ;

B is H, SH, $-NHR^3$, $-N(R^3)$ -(amino acid or peptide), or R^4 ;

X is H, SH, $-NHR^3$, $-N(R^3)$ -(amino acid or peptide) or R^4 ;

Z is H or R⁴;

R¹, R², R³ and R⁴ are independently H or lower straight or branched chain or cyclic alkyl;

n is 0, 1 or 2;

(amino acid) is any primary α - or β -amino acid not containing a thiol group;

(peptide) is a peptide of 2 to about 10 amino acids;

and

where B is -NHR³ or -N(R³)-(amino acid or peptide), X is SH, and n is 1 or 2;

where X is -NHR³ or -N(R³)-(amino acid or peptide), B is SH, and n is 1 or 2;

where B is H or R⁴, A is HOOC, H₂NOC, (amino acid or peptide)-NHOC, or (amino acid or peptide)-OOC; X is SH; and n is 0 or 1;

where A is H or R⁴, then where B is SH, X is -NHR³ or -N(R³)-(amino acid or peptide)

and where X is SH, B is -NHR³ or -N(R³)-(amino acid or peptide) and n is 1 or 2;


where X is H or R⁴, A is HOOC, H₂NOC, (amino acid or peptide)-NHOC, or (amino acid or peptide)-OOC; and B is SH;

where Z is methyl, X is methyl; A is HOOC, H₂NOC, (amino acid or peptide)-NHOC, or (amino acid or peptide)-OOC; B is SH and n is 0;

where B is SH, X is not SH and where X is SH, B is not SH;

and wherein the thiol moiety is in the reduced form;

c)




wherein each R is independently H, CH₃ or C₂H₅;

each (pgp)^s is independently a thiol protecting group or H;

m, n and p are independently 2 or 3;

A = linear or cyclic lower alkyl, aryl, heterocyclyl, a combination thereof, or a substituted derivative thereof;

d)



wherein

each R is independently H, CH₃ or C₂H₅;

m, n and p are independently 2 or 3;


A = linear or cyclic lower alkyl, aryl, heterocyclyl, a combination thereof or a substituted derivative thereof;

V = H or -CO-peptide;

R' = H or peptide;

and wherein when V = H, R' = peptide and when R' = H, V = -CO-peptide;

e)



wherein:

n, m and p are each independently 0 or 1,


each R' is independently H, lower alkyl, hydroxyalkyl (C₂-C₄), or alkoxyalkyl (C₂-C₄);

each R is independently H or R'', where R'' is substituted or unsubstituted lower alkyl or phenyl not comprising a thiol group;

one R or R' is L, wherein when an R' is L, -NR'₂ is an amine; and

L is a bivalent group linking the chelator to the compound;

f)



wherein: n, m and p are each independently 0 or 1,
 each R' is independently H, lower alkyl, hydroxyalkyl (C₂-C₄), or alkoxyalkyl (C₂-C₄);
 each R is independently H or R'', where R'' is substituted or unsubstituted lower alkyl or phenyl not comprising a thiol group;
 one R or R' is L, wherein when an R' is L, -NR'₂ is an amine; and
 L is a bivalent group linking the chelator to the compound;

g) (HOOCCH₂)₂N(CR₂)(CR₂)N(CH₂COOH)(CR₂)(CR₂)N(CH₂COOH);


where each R is independently H, C₁ to C₄ alkyl, or aryl and one R is covalently linked to a bivalent linker;

h) (HOOCCH₂)₂N(CR₂)(CR₂)N(CH₂COOH);

where each R is independently H, C₁ to C₄ alkyl, or aryl and one R is covalently linked to a bivalent linker;

i) 1,4,7,10-tetraazadodecanetetraacetic acid;

j)



where n is an integer that is 2 or 3 and where each R is independently H, C₁ to C₄ alkyl, or aryl and one R is covalently linked to the calcitonin receptor binding compound; and

k) desferrioxamine.

4. The reagent of claim 3, wherein the chelator is selected from the group consisting of:

- (amino acid)¹-(amino acid)²-{A-CZ(B)-{C(R¹R²)_n-X};
- {A-CZ(B)-{C(R¹R²)_n-X}-(amino acid)¹-(amino acid)²;
- (a primary α,β- or β,γ-diamino acid)-(amino acid)¹-{A-CZ(B)-{C(R¹R²)_n-X}; and
- {A-CZ(B)-{C(R¹R²)_n-X}-(amino acid)¹-(a primary α,β- or α,γ-diamino acid);

wherein (amino acid)¹ and (amino acid)² are each independently any naturally-occurring, modified, substituted or altered α- or β-amino acid not containing a thiol group.

5. The reagent of claim 4, wherein the chelator is selected from the group consisting of:

- (amino acid)¹-(amino acid)²-cysteine;
- (amino acid)¹-(amino acid)²-isocysteine-;
- (amino acid)¹-(amino acid)²-homocysteine-;
- (amino acid)¹-(amino acid)²-penicillamine-;
- (amino acid)¹-(amino acid)²-2-mercaptoethylamine-;
- (amino acid)¹-(amino acid)²-2-mercaptopropylamine-;
- (amino acid)¹-(amino acid)²-2-mercapto-2-methylpropylamine-; and
- (amino acid)¹-(amino acid)²-3-mercaptopropylamine-;

wherein a carboxyl terminus or a side chain of said chelator is covalently bonded to the compound.

6. The reagent of claim 5, wherein (amino acid)¹ is selected from the group consisting of an α,β-diamino acid having a free α-amine and a β,γ-diamino acid having a free β- amine.

7. The reagent of claim 4, wherein the chelator is selected from the group consisting of:


- cysteine-(amino acid)-(α,β- or β,γ-diamino acid);
- isocysteine-(amino acid)-(α,β- or β,γ-diamino acid);
- homocysteine-(amino acid)-(α,β- or β,γ-diamino acid);
- penicillamine-(amino acid)-(α,β- or β,γ-diamino acid);
- 2-mercaptoacetic acid-(amino acid)-(α,β- or β,γ-diamino acid);
- 2- or 3-mercaptopropionic acid-(amino acid)-(α,β- or β,γ-diamino acid); and
- 2-mercapto-2-methylpropionic acid-(amino acid)-(α,β- or β,γ-diamino acid);

wherein an amino terminus or a side chain of said chelator is covalently bonded to the compound.

8. The reagent of claim 1 or claim 2, wherein the chelator is selected from the group consisting of:

- Gly-Gly-Cys-;
- Ala-Gly-Cys-;
- (ε-Lys)-Gly-Cys-;
- (δ-Orn)-Gly-Cys-;
- (γ-Dab)-Gly-Cys-;
- (β-Dap)-Lys-Cys-;
- (β-Dap)-Gly-Cys-; and
- Cys(BAT).

9. The reagent of claim 3, wherein the chelator has a formula:



wherein:

R¹ and R² are each independently H, lower alkyl, hydroxyalkyl (C₂-C₄) or alkoxyalkyl (C₂-C₄);


R^3 , R^4 , R^5 , and R^6 are independently H, substituted or unsubstituted lower alkyl or phenyl not comprising a thiol group;

R^7 and R^8 are each independently H, lower alkyl, lower hydroxyalkyl or lower alkoxyalkyl;

L is a bivalent linking moiety; and

Z is a peptide.

10. The reagent of claim 3, wherein the chelator has a formula:



wherein:

R^1 and R^2 are each independently H, lower alkyl, hydroxyalkyl (C_2-C_4), or alkoxyalkyl (C_2-C_4);

R^3 , R^4 , R^5 , and R^6 are independently H, substituted or unsubstituted lower alkyl or phenyl not comprising a thiol group, and one of R^3 , R^4 , R^5 , and R^6 is $Z-L-(CR_2)_n$;


R^7 and R^8 are each independently H, lower alkyl, lower hydroxyalkyl or lower alkoxyalkyl;

L is a bivalent linking moiety;

Z is a peptide; and


X is $-NH_2$, $-NR^1R^2$, or $-NR^1-Y$, where Y is an amino acid, an amino acid amide, or a peptide having from 2 to about 20 amino acids.

11. The reagent of claim 3, wherein the chelator has a formula:



wherein: R^1 and R^2 are each independently H, lower alkyl, hydroxyalkyl (C_2-C_4) or alkoxyalkyl (C_2-C_4);
 R^3 , R^4 , R^5 , and R^6 are independently H, substituted or unsubstituted lower alkyl or phenyl not comprising a thiol group;
 n is an integer from 1 to 6;
 L is a bivalent linking moiety; and
 Z is a peptide.

12. The reagent of claim 3, wherein the chelator has a formula:



wherein:

L is a linker group; and
 Z is a peptide.

13. The reagent of any of claims 1 or 2, wherein the compound is a peptide.

14. The reagent of claim 13, wherein the peptide has an amino acid sequence
CH₃CO.SNLST.Hhc.VLGKLSCELHKLQTYPRNTGSGTP.amide
 (SEQ ID No. 9)

and the chelator is incorporated into the peptide at a sidechain of an amino acid of said sequence.

15. The reagent of claim 13, wherein the peptide comprises a calcitonin receptor-binding domain which is cyclized by a thioether.

16. The reagent of claim 15, wherein the peptide comprises an amino acid sequence:
CH₃CO.SNLSTX— SEQ ID. NO.:10

wherein X is selected from the group consisting of a cysteine residue, a homocysteine residue, and a homohomocysteine residue.

17. The reagent of claim 15, wherein the peptide comprises an amino acid sequence:

CH₃CO.X¹NLSTX²— SEQ ID NO.: 11

wherein X¹ is selected from the group consisting of an alanine residue, a glycine residue, and a serine residue; and

X² is selected from the group consisting of a cysteine residue, a homocysteine residue, and a homohomocysteine residue.

18. A reagent having a formula selected from the group consisting of:

CH₃CO.SNLST.Hhc.VLGKLSC(BAT)ELHKLQTYPRTNTGSGTP.amide (SEQ ID No. 4);
CH₃CO.SNLST.Hhc.VLGKLSQELHKLQTYPRTNTGSGTP(ε-K)GC.amide;
CH₃CO.SNLST.Hhc.VLGKLSC(CH₃CO.GGCK.amide)ELHKLQTYPRTNTGSGTP.amide;
CH₃CO.SNLST.Hhc.VLGKLSC(CH₃CO.(β-Dap)KCK.amide)ELHKLQTYPRTNTGSGTP.amide;
CH₃CO.SNLST.Hhc.VLGKLSC(CH₃CO.(ε-K)GCE.amide)ELHKLQTYPRTNTGSGTP.amide;
CH₃CO.SNLST.Hcy.VLGKLSC(CH₃CO.GGCK.amide)ELHKLQTYPRTNTGSGTP.amide;
CH₃CO.SNLST.Hcy.VLGKLSC(CH₃CO.(β-Dap)KCK.amide)ELHKLQTYPRTNTGSGTP.amide;
CH₃CO.SNLST.Hcy.VLGKLSC(CH₃CO.(ε-K)GCE.amide)ELHKLQTYPRTNTGSGTP.amide;
CH₃CO.SNLST.Cys.VLGKLSC(CH₃CO.GGCK.amide)ELHKLQTYPRTNTGSGTP.amide;
CH₃CO.SNLST.Cys.VLGKLSC(CH₃CO.(β-Dap)KCK.amide)ELHKLQTYPRTNTGSGTP.amide;
CH₃CO.SNLST.Cys.VLGKLSC(CH₃CO.(ε-K)GCE.amide)ELHKLQTYPRTNTGSGTP.amide;
SNLST.Asu.VLGKLSC(CH₃CO.(β-Dap)KCK.amide)ELHKLQTYPRTNTGSGTP.amide;
SNLST.Asu.VLGKLSC(CH₃CO.(β-Dap)KCK.amide)ELHKLQTYPRTDVGAGTP.amide.;
CH₃CO.SNLST.Hhc.VLGKLSCELHKLQTYPRTNTGSGTP.amide (SEQ ID No. 5);
CH₃CO.SNLST.Hcy.VLGKLSCELHKLQTYPRTNTGSGTP.amide (SEQ ID No. 6);
CH₃CO.SNLST.Cys.VLGKLSCELHKLQTYPRTNTGSGTP.amide (SEQ ID No. 7); and
SNLST.Asu.VLGKLSCELHKLQTYPRTNTGSGTP.amide (SEQ ID No. 8).

19. The reagent of claim 18, having the formula:

CH₃CO.SNLST.Hhc.VLGKLSC(CH₃CO.(ε-K)GCE.amide)ELHKLQTYPRTNTGSGTP.amide.

20. A reagent comprising:

a) at least two synthetic, calcitonin receptor-binding compounds, each compound being covalently linked to a radiometal chelator; and

b) a polyvalent linker forming a covalent linkage selected from the group consisting of a linkage to each compound, a linkage to each chelator, and a linkage to one compound and to the chelator of the other compound;

said reagent being characterized in that:

the reagent has a molecular weight of less than about 10,000 daltons; and

when said chelator is chelated to a radiometal, said reagent has a binding affinity for a calcitonin receptor of not less than about one-tenth the binding affinity of radioiodinated native calcitonin for said receptor.

21. A scintigraphic imaging agent comprising the reagent of any of claims 1 through 20 and technetium-99m.

22. A composition comprising the reagent of any of claims 1 through 20 and a stannous ion.

23. A radiotherapeutic agent comprising the reagent of any of claims 1 through 20 and a cytotoxic radioisotope.

24. The agent of claim 23, wherein the radioisotope is selected from the group consisting of scandium-47, copper-67, gallium-72, yttrium-90, tin-117m, iodine-125, iodine-131, samarium-153, gadolinium-159, dysprosium-165, holmium-166, ytterbium-175, lutetium-177, rhenium-186, rhenium-188, astatine-211, bismuth-212, and bismuth-213.

25. A complex formed by reacting the reagent of any of claims 1 through 20 with technetium-99m, rhenium-186 or rhenium-188 in the presence of a reducing agent.

26. The complex of claim 25, wherein the reducing agent is a stannous ion.

27. A complex formed by labeling the reagent of any of claims 1 through 20 with technetium-99m by ligand exchange of a prereduced technetium-99m complex.

28. A complex formed by labeling the reagent of any of claims 1 through 20 with rhenium-186 or rhenium-188 by ligand exchange of a prereduced rhenium complex.

29. A kit for preparing a radiopharmaceutical preparation, said kit comprising a sealed vial containing a predetermined quantity of the reagent of any of claims 1 through 20 and a sufficient amount of a reducing agent to label the reagent with technetium-99m, rhenium-186 or rhenium-188.

30. A kit for preparing a radiopharmaceutical preparation, said kit comprising a sealed vial containing a predetermined quantity of the reagent of any of claims 1 through 20.

31. A method of labeling the reagent of any of claims 1 through 20, comprising the step of reacting the reagent with technetium-99m, rhenium-186 or rhenium-188 in the presence of a reducing agent.

32. The method of Claim 30, wherein the reducing agent is a stannous ion.

33. A method of making the reagent of any of claims 1 through 20 by chemical *in vitro* synthesis.

34. The method of claim 33, wherein the synthesis is solid phase peptide synthesis.

35. The method of any of claims 33 or 34, wherein the chelator is covalently linked to the compound during synthesis.

36. Use of the reagent of any of claims 1 through 20 for manufacture of a medicament for imaging a site within a mammalian body.

37. Use of the reagent of any of claims 1 through 20 for manufacture of a medicament for treating a disease characterized by the presence of calcitonin receptors.