CLAIMS

1. A cannabidiol-containing hemostatic composition, comprising, in parts by mass, 0.05 part to 1 part of cannabidiol, 5 parts to 25 parts of *Bletilla striata* extract, 8 parts to 20 parts of *Corydalis yanhusuo* extract, 7 parts to 30 parts of *Panax notoginseng* extract, 25 parts to 60 parts of polyvinylpyrrolidone, 2 parts to 8 parts of poloxamer, 0.05 part to 4 parts of an antioxidant, 0.5 part to 4 parts of polyethylene glycol and 35 parts to 80 parts of absolute ethanol.

2. The composition according to claim 1, wherein the composition comprises, in parts by mass, 0.1 part to 1 part of cannabidiol, 7 parts to 20 parts of *Bletilla striata* extract, 7.5 parts to 20 parts of *Corydalis yanhusuo* extract, 7 parts to 10 parts of *Panax notoginseng* extract, 25 parts to 60 parts of polyvinylpyrrolidone, 2 parts to 4 parts of poloxamer, 0.05 part to 4 parts of an antioxidant, 0.5 part to 4 parts of polyethylene glycol and 35 parts to 80 parts of absolute ethanol.

3. The composition according to claim 1, wherein the polyvinylpyrrolidone is polyvinylpyrrolidone K60;

the poloxamer is poloxamer 407;

the polyethylene glycol is polyethylene glycol 6000;

the antioxidant is selected from any one or a combination of at least two of butylhydroxyanisole, dibutylhydroxytoluene, propyl gallate, tea polyphenol, phytic acid, sodium phytate, tert-butyl hydroquinone, licorice root antioxidant, phospholipid, dilauryl thiodipropionate, 4-hexylresorcinol, rosemary extract, ascorbic acid, vitamin E and bamboo leaf antioxidant;

the polysaccharide content of the *Bletilla striata* extract is higher than 60%;

the purity of the cannabidiol is not less than 90%.

4. The composition according to claim 1, wherein the antioxidant is dibutylhydroxytoluene.

5. The composition according to claim 1, wherein the *Bletilla striata* extract has a polysaccharide content higher than 70%.

6. The composition according to claim 1, wherein the *Bletilla striata* extract has a polysaccharide content of 99%.

7. The composition according to claim 1, wherein the cannabidiol has a purity not less than 99%.

8. The composition according to any one of claims 3-7, wherein the composition

comprises, in parts by mass, 25 parts to 30 parts of polyvinylpyrrolidone, 2 parts to 4 parts of poloxamer, 0.05 part to 0.2 part of dibutylhydroxytoluene, 0.5 part to 2 parts of polyethylene glycol, 7 parts to 10 parts of *Bletilla striata* extract, 8 parts to 12 parts of *Corydalis yanhusuo* extract, 9 parts to 10 parts of *Panax notoginseng* extract, 36.5 parts to 40 parts of absolute ethanol, and 0.05 part to 0.4 part of cannabidiol.

9. A preparation method for the composition according to claim 1, comprising the following steps:

step one: weighing out a proper amount of an antioxidant and cannabidiol in proportion and dissolving in absolute ethanol, subsequently adding polyvinylpyrrolidone, poloxamer and polyethylene glycol in proportion, adding absolute ethanol, dissolving, stirring for mixing well, and allowing to stand at room temperature for 6-12 h to obtain gel matrix (a);

step two: adding *Bletilla striata* extract, *Panax notoginseng* extract and *Corydalis yanhusuo* extract in sequence for mixing with the gel matrix (a), stirring at room temperature for mixing well, allowing to stand at room temperature for 24-48 h to obtain gel (b), and subpackaging to finish the preparation.

10. A liquid bandage comprising the composition according to any one of claims 1-8.

11. Use of the composition according to any one of claims 1-8 in the preparation of a liquid bandage for hemostasis, analgesia, and anti-inflammation.