Patent Claims

1. Use of 3-(1-{3-[5-(1-Methyl-piperidin-4-ylmethoxy)-pyrimidin-2-yl]-benzyl}-6-oxo-1,6-dihydro-pyridazin-3-yl)-benzonitrile or a pharmaceutically acceptable salt and/or solvate thereof for the manufacture of a medicament for the treatment of non-small cell lung cancer (NSCLC), characterized by deletion in the EGFR exon 19, wherein the medicament is to be used in combination with N-[3-[[2-[4-(4-acetylpiperazin-1-yl)-2-methoxy-anilino]-5-(trifluoromethyl)pyrimidin-4-yl]amino]phenyl]prop-2-enamide.

2. Use according to claim 1 of 3-(1-{3-[5-(1-Methyl-piperidin-4-ylmethoxy)-pyrimidin-2-yl]-benzyl}-6-oxo-1,6-dihydro-pyridazin-3-yl)-benzonitrile hydrochloride hydrate for the manufacture of a medicament for the treatment of non-small cell lung cancer (NSCLC), characterized by deletion in the EGFR exon 19, wherein the medicament is to be used in combination with N-[3-[[2-[4-(4-acetylpiperazin-1-yl)-2-methoxy-anilino]-5-(trifluoromethyl)pyrimidin-4-yl]amino]phenyl]prop-2-enamide.

3. Use according to claim 1 or 2, wherein 3-(1-{3-[5-(1-Methyl-piperidin-4-ylmethoxy)-pyrimidin-2-yl]-benzyl}-6-oxo-1,6-dihydro-pyridazin-3-yl)-benzonitrile or a pharmaceutically acceptable salt and/or solvate thereof or
3-(1-{3-[5-(1-Methyl-piperidin-4-ylmethoxy)-pyrimidin-2-yl]-benzyl}-6-oxo-1,6-dihydro-pyridazin-3-yl)-benzonitrile hydrochloride hydrate
is to be used in an amount of 250 mg to 12500 mg per week.