substitutions; and the LCDR3 includes the amino acid sequence of SEQ ID NO: 45 or SEQ ID NO: 45 with one or two single amino acid substitutions; or (b) an amino acid sequence at least 80%, at least 85%, at least 90%, at least 95% or 100% identical to SEQ ID NO: 42. In certain examples the nucleic acid sequence that encodes the light chain polypeptide subunit can encode the amino acid sequence SEQ ID NO: 58.

[0020] In certain examples of the provided polynucleotide composition, the nucleic acid sequence encoding the heavy chain polypeptide subunit and the nucleic acid sequence encoding the light chain polypeptide subunit can be on separate vectors, or they can be situated on a single vector.

[0021] The provided polynucleotide composition can further include, e.g., a nucleic acid sequence that encodes a J-chain, or functional fragment thereof, or a functional variant thereof. In certain examples the J-chain or fragment thereof can include the amino acid sequence SEQ ID NO: 49 or a functional fragment thereof. Moreover, the J-chain or fragment thereof can be a modified J-chain that further includes a heterologous polypeptide. The heterologous polypeptide can, in certain examples, be directly or indirectly fused to the J-chain or fragment thereof. In certain examples, the heterologous polypeptide can include an antibody or antigen-binding fragment thereof. In certain examples, the heterologous polypeptide can be, e.g., a scFv that can specifically bind to CD3ε. In certain examples the modified J-chain can include the amino acid sequence SEQ ID NO: 64 (V15J) or SEQ ID NO: 66 (J15V). In those examples where the modified J-chain further comprises a signal peptide, the modified J-chain can include the amino acid sequence SEQ ID NO: 63 (V15J) or SEQ ID NO: 65 (J15V). In certain examples of the provided polynucleotide composition, the nucleic acid sequence that encodes a J-chain, or functional fragment thereof, or a functional variant thereof can include SEQ ID NO: 68 or SEQ ID NO: 69.

[0022] In certain examples of the provided polynucleotide composition, the nucleic acid sequence encoding the heavy chain polypeptide subunit, the nucleic acid sequence encoding the light chain polypeptide subunit, and the nucleic acid sequence encoding the J-chain can be situated on a single vector, or they can be situated on two or three separate vectors. In one aspect, there is provided a vector or combination of vectors comprising the provided nucleic acid sequences. In another aspect, there is provided a host cell comprising the provided polynucleotide, the provided polynucleotide composition, or the provided vector or vectors, wherein the host cell can express the binding molecule as provided herein, or a subunit thereof. In yet another aspect, there is provided a method of producing the binding molecule as provided herein, comprising culturing the provided host cell, and recovering the binding molecule.

[0023] In certain examples, the disclosure provides a method for directing complement-mediated, T-cell-mediated, or both complement-mediated and T-cell-mediated killing of a CD20-expressing cell where the method includes contacting a CD20-expressing cell with a multimeric binding molecule as provided herein or a composition that includes the provided binding molecule. According to these examples, the binding molecule can direct complement-mediated, T-cell-mediated, or both complement-mediated and T-cell-mediated killing of a CD-20-expressing cell at higher potency than an equivalent amount of a monospecific, bivalent IgG1 antibody or fragment thereof that specifically binds to the same CD20 epitope as the CD20 antigen binding domain. In certain examples, the monospecific, bivalent IgG1 antibody is 1.5.3, which includes a VH having the amino acid sequence SEQ ID NO: 38 and a VL having the amino acid sequence SEQ ID NO: 42. In one aspect, there is provided an *in vitro* method for directing complement-mediated killing of a CD20-expressing cell comprising contacting a CD20-expressing cell with the binding molecule provided herein or the composition provided herein, wherein the binding molecule can direct complement-mediated killing of a CD-20-expressing cell at higher potency than an equivalent amount of a monospecific, bivalent IgG1 antibody or fragment thereof that specifically binds to CD20, wherein the monospecific, bivalent IgG1 antibody is 1.5.3, which comprises a VH having the amino acid sequence SEQ ID NO: 38 and a VL having the amino acid sequence SEQ ID NO: 42. In certain examples, the CD-20-expressing cell is a lymphoma cell line, *e.g.*, a Ramos cell line, a Raji cell line, a Daudi cell line, a Namalwa cell line, a Granta cell line, a Z138 cell line, a DoHH2 cell line, or a DB cell line. In certain examples, the CD20-expressing cell is a Raji cell line, and the binding molecule directs complement-mediated killing with an IC_{50} at least four-fold, at least ten-fold, at least 50-fold, or at least 100-fold lower than the IC_{50} of an equivalent amount of the monospecific bivalent IgG1 antibody, as measured, *e.g.*, in μg/ml. In certain examples, where the CD20-expressing cell is a Ramos cell line, the binding molecule can direct complement-mediated killing with an IC_{50} at least ten-fold, at least 20-fold, at least 30-fold, at least 40-fold, at least 50-fold, at least 60-fold, at least 70-fold, at least 80-fold, at least 90-fold, or at least 100-fold lower than the IC_{50} of an equivalent amount of the monospecific bivalent IgG1 antibody, as measured, *e.g.*, as molar equivalents. In certain examples, the CD-20-expressing cell is a malignant B cell in a subject with cancer, *e.g.*, a CD20-positive leukemia, lymphoma, or myeloma. In one aspect, there is provided use of the binding molecule provided herein or the composition provided herein in the preparation of a medicament for treating cancer in as subject in need thereof, wherein the cancer is a CD20-positive leukemia, lymphoma, or myeloma. In certain examples, the cancer is minimally responsive or non-responsive to rituximab therapy. In certain examples the subject is a human.

[0024] In other examples, the disclosure provides a method for directing complement-mediated, T-cell-mediated, or both complement-mediated and T-cell-mediated killing of a CD20-expressing cell, where the method includes contacting a CD20-expressing cell with a dimeric, pentameric, or hexameric binding molecule including two, five, or six bivalent binding units, respectively, where each binding unit includes two IgA or IgM heavy chain constant regions or fragments thereof and two antigen binding domains, where at least one antigen binding domain of the binding molecule is a CD20 antigen binding domain, and where the binding molecule can direct complement-mediated, T-cell-mediated, or both complement-mediated and T-cell-mediated killing of a CD-20-expressing cell at higher potency than an equivalent amount of a

WHAT IS CLAIMED IS:

1. A pentameric binding molecule comprising five bivalent binding units and a J-chain or functional fragment or variant thereof, wherein each binding unit comprises two IgM heavy chain constant regions, each associated with an antigen-binding domain, wherein at least one antigen binding domain of the binding molecule is a CD20 antigen binding domain comprising:

an antibody heavy chain variable region (VH) comprising immunoglobulin complementarity determining regions HCDR1, HCDR2, HCDR3, and an antibody light chain variable region (VL) comprising immunoglobulin complementarity determining regions LCDR1, LCDR2, and LCDR3, wherein the HCDR1 comprises the amino acid sequence of SEQ ID NO: 39; the HCDR2 comprises the amino acid sequence of SEQ ID NO: 40; the HCDR3 comprises the amino acid sequence of SEQ ID NO: 41; the LCDR1 comprises the amino acid sequence of SEQ ID NO: 43; the LCDR2 comprises the amino acid sequence of SEQ ID NO: 44; and the LCDR3 comprises the amino acid sequence of SEQ ID NO: 45;

wherein the binding molecule can direct complement-mediated killing of a CD-20-expressing cell at higher potency than an equivalent amount of a monospecific, bivalent reference IgG1 antibody that specifically binds to CD20; and

wherein the monospecific, bivalent reference IgG1 antibody is 1.5.3, which comprises a VH having the amino acid sequence SEQ ID NO: 38 and a VL having the amino acid sequence SEQ ID NO: 42.

2. The binding molecule of claim 1, wherein the VH comprises an amino acid sequence at least 80%, at least 85%, at least 90%, at least 95% or 100% identical to SEQ ID NO: 38, and the VL comprises an amino acid sequence at least 80%, at least 85%, at least 90%, at least 95% or 100% identical to SEQ ID NO: 42.

3. The binding molecule of claim 1 or claim 2, wherein each binding unit comprises two IgM heavy chains each comprising a VH situated amino terminal to the IgM constant region, and two immunoglobulin light chains each comprising a VL situated amino terminal to an immunoglobulin light chain constant region.

4. The binding molecule of any one of claims 1 to 3, wherein the IgM heavy chain constant regions each comprise a Cμ1 domain, a Cμ2 domain, a Cμ3 domain and a Cμ4-tp domain.

5. The binding molecule of claim 4, wherein the IgM heavy chain constant regions are human IgM constant regions.

6. The binding molecule of claim 5, wherein the binding units are identical, wherein the IgM heavy chains in each binding unit comprise the amino acid sequence SEQ ID NO: 56, and wherein the light chains in each binding unit comprise the amino acid sequence SEQ ID NO: 58.

7. The binding molecule of any one of claims 1 to 6, wherein the J-chain or functional fragment or variant thereof comprises the amino acid sequence SEQ ID NO: 49 or a functional fragment or variant thereof.

8. The binding molecule of any one of claims 1 to 7, wherein the J-chain or functional fragment or variant thereof is a modified J-chain further comprising a heterologous polypeptide, wherein the heterologous polypeptide is directly or indirectly fused to the J-chain or functional fragment or variant thereof.

9. The binding molecule of claim 8, wherein the heterologous polypeptide is fused to the J-chain or functional fragment or variant thereof via a peptide linker.

10. The binding molecule of claim 8 or claim 9, wherein a heterologous polypeptide is fused to the N-terminus of the J-chain or functional fragment or variant thereof, the C-terminus of the J-chain or functional fragment or variant thereof, or to both the N-terminus and C-terminus of the J-chain or functional fragment or variant thereof.

11. The binding molecule of any one of claims 8 to 10, wherein the heterologous polypeptide comprises an antibody or antigen-binding fragment thereof.

12. The binding molecule of claim 11, wherein the antigen-binding fragment comprises an Fab fragment, an Fab' fragment, an $F(ab')_2$ fragment, an Fd fragment, an Fv fragment, a single-chain Fv (scFv) fragment, a disulfide-linked Fv (sdFv) fragment, or any combination thereof.

13. The binding molecule of claim 11 or claim 12, wherein the heterologous polypeptide comprises a scFv antibody fragment that can specifically bind to CD3ε.

14. The binding molecule of claim 13, wherein the modified J-chain comprises the amino acid sequence SEQ ID NO: 64 (V15J) or SEQ ID NO: 66 (J15V).

15. The binding molecule of any one of claims 1 to 14, wherein the CD20-expressing cell is a Raji cell line, and wherein the binding molecule can direct complement-mediated killing with an IC_{50} at least four-fold, at least ten-fold, at least 50-fold, or at least 100-fold lower than the IC_{50} of an equivalent amount of the monospecific bivalent IgG1 antibody, as measured in µg/ml, or wherein the CD20-expressing cell is a Ramos cell line, and wherein the binding molecule can direct complement-mediated killing with an IC_{50} at least ten-fold, at least 20-fold, at least 30-fold, at least 40-fold, at least 50-fold, at least 60-fold, at least 70-fold, at least 80-fold, at least 90-fold, or at least 100-fold lower than the IC_{50} of an equivalent amount of the monospecific bivalent IgG1 antibody, as measured as molar equivalents,

wherein the monospecific bivalent IgG1 antibody is 1.5.3, which comprises a VH having the amino acid sequence SEQ ID NO: 38 and a VL having the amino acid sequence SEQ ID NO: 42.

16. The binding molecule of any one of claims 1 to 14, wherein the CD-20-expressing cell is a malignant B cell isolated from a subject with cancer.

17. The binding molecule of claim 16, wherein the cancer is a CD20-positive leukemia, lymphoma, or myeloma.

18. The binding molecule of any one of claim 16 or claim 17, wherein the cancer is minimally responsive or non-responsive to rituximab therapy.

19. The binding molecule of any one of claims 16 to 18, wherein the subject is human.

20. A composition comprising the binding molecule of any one of claims 1 to 19, and a carrier.

21. A polynucleotide comprising a nucleic acid sequence that encodes a heavy chain polypeptide subunit of the binding molecule of any one of claims 1 to 19, wherein the heavy chain polypeptide subunit comprises an IgM heavy chain constant region, and at least the antibody VH portion of the CD20 antigen binding domain.

22. A composition comprising the polynucleotide of claim 21.

23. The composition of claim 22, further comprising a nucleic acid sequence that encodes a light chain polypeptide subunit of the binding molecule of any one claims 1 to 19.

24. The composition of claim 23, wherein the nucleic acid sequence encoding the heavy chain polypeptide subunit and the nucleic acid sequence encoding the light chain polypeptide subunit are on separate vectors.

25. The composition of claim 23, wherein the nucleic acid sequence encoding the heavy chain polypeptide subunit and the nucleic acid sequence encoding the light chain polypeptide subunit are on a single vector.

26. The composition of any one of claims 22 to 25, further comprising a nucleic acid sequence that encodes the J-chain, or functional fragment or variant thereof of the binding molecule of any one of claims 1 to 19, situated on a separate vector or on the same vector as the nucleic acid sequence encoding the heavy chain polypeptide subunit and/or the nucleic acid sequence encoding the light chain polypeptide subunit.

27. A vector or combination of vectors comprising the nucleic acid sequences of any one of claims 21 to 26.

28. A host cell comprising the polynucleotide of claim 21, the composition of any one of claims 22 to 26, or the vector or vectors of claim 27, wherein the host cell can express the binding molecule of any one of claims 1 to 19, or a subunit thereof.

29. A method of producing the binding molecule of any one of claims 1 to 19, comprising culturing the host cell of claim 28, and recovering the binding molecule.

30. An *in vitro* method for directing complement-mediated killing of a CD20-expressing cell comprising contacting a CD20-expressing cell with the binding molecule of any one of claims 1 to 19 or the composition of claim 20, wherein the binding molecule can direct complement-mediated killing of a CD-20-expressing cell at higher potency than an equivalent amount of a monospecific, bivalent IgG1 antibody or fragment thereof that specifically binds to CD20, wherein the monospecific, bivalent IgG1 antibody is 1.5.3, which comprises a VH having the amino acid sequence SEQ ID NO: 38 and a VL having the amino acid sequence SEQ ID NO: 42.

31. Use of the binding molecule of any one of claims 1 to 19 or the composition of claim 20 in the preparation of a medicament for treating cancer in a subject in need thereof, wherein the cancer is a CD20-positive leukemia, lymphoma, or myeloma.

32. The use of claim 31, wherein the cancer is minimally responsive or non-responsive to rituximab therapy.

33. The use of claim 31 or 32, wherein the subject is human.