

9705783-0
IPOS copy



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07H 15/248, C07C 233/32, 323/41, A61K 31/70, 31/645		A1	(11) International Publication Number: WO 97/01570
			(43) International Publication Date: 16 January 1997 (16.01.97)
(21) International Application Number: PCT/EP96/02718		(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 21 June 1996 (21.06.96)		Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(30) Priority Data: MI95A001367 27 June 1995 (27.06.95) IT			
(71) Applicant (for all designated States except US): INDENA S.P.A. [IT/IT]; Viale Ortles, 12, I-20139 Milano (IT).			
(72) Inventors; and (75) Inventors/Applicants (for US only): BOMBARDELLI, Ezio [IT/IT]; Via Ortles, 12, I-20139 Milano (IT). GABETTA, Bruno [IT/IT]; Viale Ortles, 12, I-20139 Milano (IT).			
(74) Agent: MINOJA, Fabrizio; Studio Consulenza Brevettuale, Via Rossini, 8, I-20122 Milano (IT).			
(54) Title: COLCHICINE DERIVATIVES, THE USE THEREOF AND FORMULATIONS CONTAINING THEM			
(57) Abstract			
The present invention relates to novel colchicine derivatives having antiproliferative, antineoplastic, anti-inflammatory and muscle relaxant activities; said derivatives include novel colchine nitrogen amides for use either as such or after derivatization of the hydroxyl at C₃ of the aromatic ring and at C₁₀ of the tropolone ring. These novel compounds have a cytotoxicity on human tumoral cell lines comparable with colchicine but, in comparison with the latter, they are much more active on cells resistant to the usual antitumorals. The compounds can be included in pharmaceutical formulations useful for the intravenous, oral and topical administrations.			