

DESCRIPTION

NITROGEN-CONTAINING HETEROCYCLIC COMPOUND AND AGRICULTURAL
FUNGICIDE

5

TECHNICAL FIELD

[0001]

The present invention relates to novel nitrogen-containing heterocyclic compounds and an agricultural fungicide which contains at least one selected from the group consisting of the nitrogen-containing heterocyclic compounds as an active ingredient.

10

Priority is claimed on Japanese Patent Application No. 2010-000194, filed January 4, 2010, the content of which is incorporated herein by reference.

15

BACKGROUND ART

[0002]

20

In agricultural crop cultivation, a variety of control agents are used to deal with crop diseases. However, there are very few agents which fully satisfy the requirements for use as a control agent due to reasons such as an insufficient control effect, limited use due to the emergence of agent-resistant pathogens, phytotoxic or contaminating effects on plants, or toxicity to humans, domestic animals, and fish, and an adverse effect on the environment. Therefore, the need for agents having fewer disadvantages of this kind and which are safe to use is required.

[0003]

25

Related to the present invention, in the following Patent Document 1 or 2,

quinoline derivatives having similar chemical structures to the compounds according to the present invention, and agricultural fungicides containing the quinoline derivatives as active ingredients, are disclosed.

5

DOCUMENTS OF RELATED ART

PATENT DOCUMENTS

[0004]

[Patent Document 1] Pamphlet of WO2005/070917

[Patent Document 2] Pamphlet of WO2007/011022

10

SUMMARY OF THE INVENTION

PROBLEMS TO BE SOLVED BY THE INVENTION

[0005]

An object of the present invention is to provide a novel nitrogen-containing
15 heterocyclic compound, and a salt thereof or N-oxide compound thereof, and an
agricultural fungicide, which has reliable effects, is safe to use and contains at least one
selected from the group consisting of the nitrogen-containing heterocyclic compound, the
salt and N-oxide compound thereof as an active ingredient.

20 MEANS TO SOLVE THE PROBLEMS

[0006]

The inventors have studied intensively in order to solve the above problems.
As a result, a nitrogen-containing heterocyclic compound represented by Formula (I), a
salt thereof or an N-oxide compound thereof were obtained. It was also discovered that
25 the nitrogen-containing heterocyclic compound, the salt thereof or the N-oxide


compound thereof are useful as an active ingredient of an agricultural fungicide, which has reliable effects and is safe to use. The present invention was completed with further studies based on these findings.

[0007]

5 That is, the present invention includes the following aspects.

<1> A nitrogen-containing heterocyclic compound represented by Formula (I), a salt thereof, or an N-oxide compound thereof.

[0008]



Formula (I)

10 [0009]

In Formula (I), R represents a group represented by $CR^1R^2R^3$, an unsubstituted or substituted C_{6-10} aryl group, or a cyano group;

R^1 to R^3 each independently represents a hydrogen atom, an unsubstituted or substituted C_{1-8} alkyl group, an unsubstituted or substituted C_{2-8} alkenyl group, an unsubstituted or substituted C_{2-8} alkynyl group, an unsubstituted or substituted C_{3-8} cycloalkyl group, an unsubstituted or substituted C_{4-8} cycloalkenyl group, an unsubstituted or substituted C_{6-10} aryl group, an unsubstituted or substituted heterocyclic group, an unsubstituted or substituted C_{1-8} acyl group, an unsubstituted or substituted (1-imino) C_{1-8} alkyl group, an unsubstituted or substituted carboxyl group, an

unsubstituted or substituted carbamoyl group, an unsubstituted or substituted hydroxyl group, an unsubstituted or substituted amino group, an unsubstituted or substituted mercapto group, a substituted sulfonyl group, a halogeno group, a cyano group, or a nitro group;

- 5 excepting in which: R^1 to R^3 are all hydrogen atoms; R^1 to R^3 are all unsubstituted C_{1-8} alkyl groups; any one of R^1 to R^3 is a hydrogen atom and the remaining two are both unsubstituted C_{1-8} alkyl groups; and, any one of R^1 to R^3 is an unsubstituted C_{1-8} alkyl group and the remaining two are both hydrogen atoms;

- R^1 and R^2 may be joined to form an unsubstituted or substituted 5- to
10 8-membered ring, or to form $O=$, $R^a R^b C=$, or $R'-N=$;

R^a represents a hydrogen atom or an unsubstituted or substituted C_{1-8} alkyl group;

R^b represents a hydrogen atom or an unsubstituted or substituted C_{1-8} alkyl group;

- 15 R' represents an unsubstituted or substituted hydroxyl group, or an unsubstituted or substituted C_{1-8} alkyl group;

- R^4 each independently represents an unsubstituted or substituted C_{1-8} alkyl group, an unsubstituted or substituted C_{2-8} alkenyl group, an unsubstituted or substituted C_{2-8} alkynyl group, an unsubstituted or substituted C_{3-8} cycloalkyl group, an unsubstituted or substituted C_{4-8} cycloalkenyl group, an unsubstituted or substituted C_{6-10} aryl group, an unsubstituted or substituted heterocyclic group, an unsubstituted or substituted C_{1-8} acyl group, an unsubstituted or substituted (1-imino) C_{1-8} alkyl group, an unsubstituted or substituted carboxyl group, an unsubstituted or substituted carbamoyl group, an unsubstituted or substituted hydroxyl group, an unsubstituted or substituted amino group,
25 an unsubstituted or substituted mercapto group, a substituted sulfonyl group, a halogeno

group, a cyano group, or a nitro group;

m represents a number of R^4 and is an integer of 0 to 6;

R^5 each independently represents an unsubstituted or substituted C_{1-8} alkyl group, an unsubstituted or substituted C_{2-8} alkenyl group, an unsubstituted or substituted C_{2-8} alkynyl group, an unsubstituted or substituted C_{3-8} cycloalkyl group, an unsubstituted or substituted C_{4-8} cycloalkenyl group, an unsubstituted or substituted C_{6-10} aryl group, an unsubstituted or substituted heterocyclic group, an unsubstituted or substituted C_{1-8} acyl group, an unsubstituted or substituted (1-imino) C_{1-8} alkyl group, an unsubstituted or substituted carboxyl group, an unsubstituted or substituted carbamoyl group, an unsubstituted or substituted hydroxyl group, an unsubstituted or substituted amino group, an unsubstituted or substituted mercapto group, a substituted sulfonyl group, a halogeno group, a cyano group, or a nitro group;

n represents a number of R^5 and is an integer of 0 to 5;

any one of R^1 to R^3 and any one of R^5 may be joined to form an unsubstituted or substituted 5- to 8-membered ring;

A represents: a 5- to 7-membered hydrocarbon ring or a 5- to 7-membered heterocyclic ring when R is a group represented by $CR^1R^2R^3$; or a benzene ring when R is an unsubstituted or substituted C_{6-10} aryl group or a cyano group;

D represents a 5- to 7-membered hydrocarbon ring or a 5- to 7-membered heterocyclic ring;

X represents an oxygen atom, a sulfur atom, a sulfenyl group, a sulfonyl group, an unsubstituted or substituted carbon atom, or an unsubstituted or substituted nitrogen atom;


Y represents a carbon atom or a nitrogen atom; and

Z represents a carbon atom or a nitrogen atom.

[0010]

<2> A nitrogen-containing heterocyclic compound represented by Formula (II), a salt thereof, or an N-oxide compound thereof.

[0011]



Formula (II)

5


In Formula (II), each of R, R⁴, R⁵, n, D, X, Y, and Z represents the same meaning as those in Formula (I) described above in <1>; and

m₁ represents a number of R⁴ and is an integer of 0 to 6.

[0012]

10 <3> A nitrogen-containing heterocyclic compound represented by Formula (III), a salt thereof, or an N-oxide compound thereof.

[0013]



Formula (III)

In Formula (III), each of R, R⁴, R⁵, m₁ and X represents the same meaning as

15 those in Formula (II) described above in <2>; and

n_1 represents a number of R^5 and is an integer of 0 to 4.

[0014]

<4> An agricultural fungicide comprising, as an active ingredient, at least one selected from the group consisting of the nitrogen-containing heterocyclic compound, the salt thereof, and the N-oxide compound thereof, of any one <1> to <3>.

[0015]

<5> An intermediate product of the nitrogen-containing heterocyclic compound represented by Formula (I), the salt thereof, or the N-oxide compound thereof, described above in <1>, the intermediate product being selected from the group consisting of 8-fluoro-3-hydroxyquinoline, 7,8-difluoro-3-hydroxyquinoline, 7,8-difluoro-3-iodoquinoline, 8-fluoro-3-hydroxy-2-methylquinoline, and 7,8-difluoro-3-hydroxy-2-methylquinoline.

EFFECT OF THE INVENTION

[0016]

The nitrogen-containing heterocyclic compound, the salt thereof, and N-oxide compound thereof according to the present invention are novel compounds useful as an active ingredient of an agricultural fungicide which has reliable effects and is safe to use.

The agricultural fungicide according to the present invention is an agent which has excellent control effects, does not cause phytotoxicity in plants, and has little toxicity to humans, domestic animals, and fish or on the environment.

MODES FOR CARRYING OUT THE INVENTION

[0017]

Hereinafter, the present invention will be categorized into 1) a

nitrogen-containing heterocyclic compound represented by Formula (I), and a salt thereof or an N-oxide compound thereof, and 2) an agricultural fungicide and will be described in detail.

[0018]

- 5 1) Nitrogen-containing heterocyclic compound represented by Formula (I), and salt thereof, or N-oxide compound thereof.

The nitrogen-containing heterocyclic compound according to the present invention is represented by Formula (I) (hereinafter, sometimes referred to as “compound (I)”), is preferably represented by Formula (II) (hereinafter, sometimes referred to as “compound (II)”), and is more preferably represented by Formula (III) (hereinafter, sometimes referred to as “compound (III)”).

10

[0019]

The nitrogen-containing heterocyclic compound, the salt thereof, or the N-oxide compound thereof according to the present invention may be a hydrate, a solvate, a crystal polymorphism, or the like. Also, an asymmetric carbon atom, stereoisomers based on a double bond or the like, or a mixture thereof, may be present in the nitrogen-containing heterocyclic compound, the salt thereof, or the N-oxide compound thereof according to the present invention.

15

[0020]

20 First of all, the meaning of “unsubstituted” and “unsubstituted” in Formulae (I), (II), and (III) will be explained. The term “unsubstituted” in the present specification means that the specified group is solely formed of a group serving as a mother nucleus. When only the name of the group serving as the mother nucleus is mentioned without mention of “substituted”, it means “unsubstituted” unless otherwise stated.

25 On the other hand, the term “substituted” means that a hydrogen atom of a group

serving as a mother nucleus has been substituted with a substituent having the same structure as or a different structure to the mother nucleus. The “substituent” is a different group which is bonded to the group serving as a mother nucleus. The “substituent” may be one or more. At least two substituents may be the same or
 5 different.

Terms such as “C₁₋₆” or the like indicate that the number of carbon atoms in a group serving as a mother nucleus is 1 to 6 or the like. The number of carbon atoms does not include carbon atoms in a substituent. For example, a butyl group having an epoxy group as a substituent is categorized as a C₂ alkoxy C₄ alkyl group.

10 [0021]

The “substituent” is not particularly limited as long as it is chemically acceptable and has the effects of the present invention.

Examples of the “substituent” include: a halogen atom such as a fluorine atom, a chlorine atom, a bromine atom, or an iodine atom; a C₁₋₆ alkyl group such as a methyl
 15 group, an ethyl group, an n-propyl group, an i-propyl group, an n-butyl group, an s-butyl group, an i-butyl group, a t-butyl group, an n-pentyl group, or an n-hexyl group; a C₃₋₆ cycloalkyl group such as a cyclopropyl group, a cyclobutyl group, a cyclopentyl group, or a cyclohexyl group; a C₂₋₆ alkenyl group such as a vinyl group, a 1-propenyl group, a 2-propenyl group, a 1-butenyl group, a 2-butenyl group, a 3-butenyl group, a
 20 1-methyl-2-propenyl group, a 2-methyl-2-propenyl group, a 1-pentenyl group, a 2-pentenyl group, a 3-pentenyl group, a 4-pentenyl group, a 1-methyl-2-butenyl group, a 2-methyl-2-butenyl group, a 1-hexenyl group, a 2-hexenyl group, a 3-hexenyl group, a 4-hexenyl group, and a 5-hexenyl group; a C₃₋₆ cycloalkenyl group such as a 2-cyclopropenyl group, a 2-cyclopentenyl group, or a 3-cyclohexenyl group; a C₂₋₆
 25 alkynyl group such as an ethynyl group, a 1-propynyl group, 2-propynyl group, a

1-butynyl group, a 2-butynyl group, a 3-butynyl group, a 1-methyl-2-propynyl group, a 2-methyl-3-butynyl group, a 1-pentynyl group, a 2-pentynyl group, a 3-pentynyl group, a 4-pentynyl group, a 1-methyl-2-butynyl group, a 2-methyl-3-pentynyl group, a 1-hexynyl group, or a 1,1-dimethyl-2-butynyl group;

5 [0022]

a C₁₋₆ alkoxy group such as a methoxy group, an ethoxy group, an n-propoxy group, an i-propoxy group, an n-butoxy group, an s-butoxy group, an i-butoxy group, or a t-butoxy group; a C₂₋₆ alkenyloxy group such as a vinyloxy group, an allyloxy group, a propenyloxy group, or a butenyloxy group; a C₂₋₆ alkynyloxy group such as an ethynyloxy group or a propargyloxy group; a C₆₋₁₀ aryl group such as a phenyl group or a naphthyl group; a C₆₋₁₀ aryloxy group such as a phenoxy group or a 1-naphthoxy group; a C₇₋₁₁ aralkyl group such as a benzyl group or a phenethyl group; a C₇₋₁₁ aralkyloxy group such as a benzyloxy group or a phenethyloxy group; a C₁₋₇ acyl group such as a formyl group, an acetyl group, a propionyl group, a benzoyl group, or a cyclohexylcarbonyl group; a C₁₋₇ acyloxy group such as a formyloxy group, an acetyloxy group, a propionyloxy group, a benzoyloxy group, or a cyclohexylcarbonyloxy group; a C₁₋₆ alkoxy carbonyl group such as a methoxycarbonyl group, an ethoxycarbonyl group, an n-propoxycarbonyl group, an i-propoxycarbonyl group, an n-butoxycarbonyl group, and a t-butoxycarbonyl group; carboxyl group;

20 [0023]

a hydroxyl group; an oxo group; a C₁₋₆ haloalkyl group such as a chloromethyl group, a chloroethyl group, a trifluoromethyl group, a 1,2-dichloro-n-propyl group, a 1-fluoro-n-butyl group, or a perfluoro-n-pentyl group; a C₂₋₆ haloalkenyl group such as a 2-chloro-1-propenyl group or a 2-fluoro-1-butenyl group; a C₂₋₆ haloalkynyl group such as a 4,4-dichloro-1-butynyl group, a 4-fluoro-1-pentynyl group or a 5-bromo-2-pentynyl

group; a C₁₋₆ haloalkoxy group such as a 2-chloro-n-propoxy group or a 2,3-dichlorobutoxy group; a C₂₋₆ haloalkenyloxy group such as a 2-chloropropenyloxy group or a 3-bromobutenyloxy group; a C₆₋₁₀ haloaryl group such as a 4-chlorophenyl group, a 4-fluorophenyl group, or a 2,4-dichlorophenyl group; a C₆₋₁₀ haloaryloxy group
 5 such as a 4-fluorophenyloxy group, or a 4-chloro-1-naphthoxy group; a C₁₋₇ haloacyl group such as a chloroacetyl group, a trifluoroacetyl group, a trichloroacetyl group, or a 4-chlorobenzoyl group;

[0024]

a cyano group; an isocyano group; a nitro group; an isocyanate group; a cyanate
 10 group; an azide group; an amino group; a C₁₋₆ alkylamino group such as a methylamino group, a dimethylamino group, or a diethylamino group; a C₆₋₁₀ arylamino group such as an anilino group or a naphthylamino group; a C₇₋₁₁ aralkylamino group such as a benzylamino group or a phenylethylamino group; a C₁₋₇ acylamino group such as a formylamino group, an acetylamino group, a propanoylamino group, a butyrylamino
 15 group, an i-propylcarbonylamino group, or a benzoylamino group; a C₁₋₆ alkoxycarbonylamino group such as a methoxycarbonylamino group, an ethoxycarbonylamino group, an n-propoxycarbonylamino group, or an i-propoxycarbonylamino group; a carbamoyl group; a substituted carbamoyl group such as a dimethylcarbamoyl group, a phenylcarbamoyl group, or an
 20 N-phenyl-N-methylcarbamoyl group; an imino C₁₋₆ alkyl group such as an iminomethyl group, a (1-imino)ethyl group, or a (1-imino)-n-propyl group; a hydroxyimino C₁₋₆ alkyl group such as a hydroxyiminomethyl group, a (1-hydroxyimino)ethyl group, or a (1-hydroxyimino)propyl group; a C₁₋₆ alkoxyimino C₁₋₆ alkyl group such as a methoxyiminomethyl group or a (1-methoxyimino)ethyl group;

25 [0025]

a mercapto group; an isothiocyanate group; a thiocyanate group; a C₁₋₆ alkylthio group such as a methylthio group, an ethylthio group, an n-propylthio group, an i-propylthio group, an n-butylthio group, an i-butylthio group, an s-butylthio group, or a t-butylthio group; a C₂₋₆ alkenylthio group such as a vinylthio group or an allylthio group; a C₂₋₆ alkynylthio group such as an ethynylthio group or a propargylthio group; a C₆₋₁₀ arylthio group such as a phenylthio group or a naphthylthio group; a heteroarylthio group such as a thiazolylthio group or a pyridylthio group; a C₇₋₁₁ aralkylthio group such as a benzylthio group or a phenethylthio group; a (C₁₋₆ alkylthio)carbonyl group such as a (methylthio)carbonyl group, an (ethylthio)carbonyl group, a (n-propylthio)carbonyl group, a (i-propylthio)carbonyl group, a (n-butylthio)carbonyl group, a (i-butylthio)carbonyl group, a (s-butylthio)carbonyl group, or a (t-butylthio)carbonyl group;

[0026]

a C₁₋₆ alkylsulfinyl group such as a methylsulfinyl group, an ethylsulfinyl group, or a t-butylsulfinyl group; a C₂₋₆ alkenylsulfinyl group such as an allylsulfinyl group; a C₂₋₆ alkynylsulfinyl group such as a propargylsulfinyl group; a C₆₋₁₀ arylsulfinyl group such as a phenylsulfinyl group; a heteroarylsulfinyl group such as a thiazolylsulfinyl group or a pyridylsulfinyl group; a C₇₋₁₁ aralkylsulfinyl group such as a benzylsulfinyl group or a phenethylsulfinyl group; a C₁₋₆ alkylsulfonyl group such as a methylsulfonyl group, an ethylsulfonyl group, or a t-butylsulfonyl group; a C₂₋₆ alkenylsulfonyl group such as an allylsulfonyl group; a C₂₋₆ alkynylsulfonyl group such as a propargylsulfonyl group; a C₆₋₁₀ arylsulfonyl group such as a phenylsulfonyl group; a heteroarylsulfonyl group such as a thiazolylsulfonyl group or a pyridylsulfonyl group; a C₇₋₁₁ aralkylsulfonyl group such as a benzylsulfonyl group or a phenethylsulfonyl group;

25 [0027]

a 5-membered heteroaryl group such as a pyrrolyl group, a furyl group, a thienyl group, an imidazolyl group, a pyrazolyl group, an oxazolyl group, an isoxazolyl group, a thiazolyl group, an isothiazolyl group, a triazolyl group, an oxadiazolyl group, a thiadiazolyl group, or a tetrazolyl group; a 6-membered heteroaryl group such as a pyridyl group, pyrazinyl group, pyrimidinyl group, a pyridazinyl group, or a triazinyl group; a saturated heterocyclic group such as an aziridinyl group, an epoxy group, a pyrrolidinyl group, a tetrahydrofuranyl group, a piperidyl group, a piperazinyl group, or a morpholinyl group; a tri C₁₋₆ alkylsilyl group such as a trimethylsilyl group, a triethylsilyl group, or a t-butyl dimethylsilyl group; a triphenylsilyl group; and the like.

10 [0028]

The “substituent” may have a different “substituent”.

[0029]

[R]

R represents a group represented by CR¹R²R³, an unsubstituted or substituted C₆₋₁₀ aryl group, or a cyano group.

R¹ to R³ each independently represents a hydrogen atom, an unsubstituted or substituted C₁₋₈ alkyl group, an unsubstituted or substituted C₂₋₈ alkenyl group, an unsubstituted or substituted C₂₋₈ alkynyl group, an unsubstituted or substituted C₃₋₈ cycloalkyl group, an unsubstituted or substituted C₄₋₈ cycloalkenyl group, an unsubstituted or substituted C₆₋₁₀ aryl group, an unsubstituted or substituted heterocyclic group, an unsubstituted or substituted C₁₋₈ acyl group, an unsubstituted or substituted (1-imino) C₁₋₈ alkyl group, an unsubstituted or substituted carboxyl group, an unsubstituted or substituted carbamoyl group, an unsubstituted or substituted hydroxyl group, an unsubstituted or substituted amino group, an unsubstituted or substituted mercapto group, a substituted sulfonyl group, a halogeno group, a cyano group, or a nitro

group.

However, there are no cases in which R^1 to R^3 are all hydrogen atoms. Also, there are no cases in which R^1 to R^3 are all unsubstituted C_{1-8} alkyl groups. Also, when any one of R^1 to R^3 is a hydrogen atom, there are no cases in which the remaining two are both unsubstituted C_{1-8} alkyl groups. Also, when any one of R^1 to R^3 is an unsubstituted C_{1-8} alkyl group, there are no cases in which the remaining two are both hydrogen atoms.

[0030]

A " C_{1-8} alkyl group" is a saturated hydrocarbon having 1 to 8 carbon atoms.

- 10 The C_{1-8} alkyl group may be a straight chain or a branched chain. Examples of the C_{1-8} alkyl group include a methyl group, an ethyl group, an n-propyl group, an n-butyl group, an n-pentyl group, an n-hexyl group, an n-heptyl group, an n-octyl group, an i-propyl group, an i-butyl group, an s-butyl group, a t-butyl group, an i-pentyl group, a neopentyl group, a 2-methylbutyl group, a 2,2-dimethylpropyl group, an i-hexyl group and the like.
- 15 Among these, a C_{1-6} alkyl group is preferable.

[0031]

Examples of "substituted C_{1-8} alkyl group" include:

- a cycloalkylalkyl group such as a cyclopropylmethyl group, a 2-cyclopropylethyl group, a cyclopentylmethyl group, or a 2-cyclohexylethyl group,
- 20 preferably a C_{3-6} cycloalkyl C_{1-6} alkyl group;
- a cycloalkenylalkyl group such as a cyclopentenylmethyl group, a 3-cyclopentenylmethyl group, a 3-cyclohexenylmethyl group, or a 2-(3-cyclohexenyl)ethyl group, preferably a C_{4-6} cycloalkenyl C_{1-6} alkyl group;
- a haloalkyl group such as a fluoromethyl group, a chloromethyl group, a
- 25 bromomethyl group, a difluoromethyl group, a dichloromethyl group, a dibromomethyl

group, a trifluoromethyl group, a trichloromethyl group, a tribromomethyl group, a 2,2,2-trifluoroethyl group, a 2,2,2-trichloroethyl group, a pentafluoroethyl group, a 4-fluorobutyl group, a 4-chlorobutyl group, a 3,3,3-trifluoropropyl group, a 2,2,2-trifluoro-1-trifluoromethylethyl group, a perfluorohexyl group, a perchlorohexyl group, a perfluorooctyl group, a perchlorooctyl group, a 2,4,6-trichlorohexyl group, a perfluorodecyl group, or a 2,2,4,4,6,6-hexachlorooctyl group, preferably a C₁₋₆ haloalkyl group;

an arylalkyl group (aralkyl group) such as a benzyl group, a phenethyl group, a 3-phenylpropyl group, a 1-naphthylmethyl group, or a 2-naphthylmethyl group, preferably a C₆₋₁₀ aryl C₁₋₆ alkyl group;

[0032]

a heteroarylalkyl group such as a 2-pyridylmethyl group, a 3-pyridylmethyl group, 4-pyridylmethyl group, a 2-(2-pyridyl)ethyl group, a 2-(3-pyridyl)ethyl group, a 2-(4-pyridyl)ethyl group, a 3-(2-pyridyl)propyl group, a 3-(3-pyridyl)propyl group, a 3-(4-pyridyl)propyl group, a 2-pyrazinylmethyl group, a 3-pyrazinylmethyl group, a 2-(2-pyrazinyl)ethyl group, a 2-(3-pyrazinyl)ethyl group, a 3-(2-pyrazinyl)propyl group, a 3-(3-pyrazinyl)propyl group, a 2-pyrimidylmethyl group, a 4-pyrimidylmethyl group, a 2-(2-pyrimidyl)ethyl group, a 2-(4-pyrimidyl)ethyl group, a 3-(2-pyrimidyl)propyl group, a 3-(4-pyrimidyl)propyl group, a 2-furylmethyl group, a 3-furylmethyl group, a 2-(2-furyl)ethyl group, a 2-(3-furyl)ethyl group, a 3-(2-furyl)propyl group, or a 3-(3-furyl)propyl group, preferably a 5- to 6-membered heteroaryl C₁₋₆ alkyl group;

a hydroxyalkyl group such as a hydroxymethyl group, a 1-hydroxyethyl group, a 2-hydroxyethyl group, a 1-hydroxypropyl group, a 3-hydroxypropyl group, a 1-hydroxy-1-methylethyl group, a 2-hydroxy-1,1-dimethylethyl group, a 2-hydroxy-1,1-dimethylpropyl group, or a 2-hydroxy-2-methylpropyl group, preferably a

hydroxyl C₁₋₆ alkyl group;

an alkoxyalkyl group such as a methoxymethyl group, an ethoxymethyl group, a 2-methoxyethyl group, a 2-ethoxyethyl group, a methoxy-n-propyl group, an n-propoxymethyl group, an i-propoxyethyl group, an s-butoxymethyl group, a t-butoxyethyl group, a 2,2-dimethoxyethyl group, or a 2,2-dimethoxy-1,1-dimethylethyl group, preferably a C₁₋₆ alkoxy C₁₋₆ alkyl group;

an acyloxyalkyl group such as a formyloxymethyl group, an acetoxymethyl group, a 2-acetoxyethyl group, a propionyloxymethyl group, or a propionyloxyethyl group, preferably C₁₋₇ acyloxy C₁₋₆ alkyl group;

a trialkylsilyloxyalkyl group such as a trimethylsilyloxymethyl group or a t-butyltrimethylsilyloxymethyl group, preferably a tri C₁₋₆ alkylsilyloxy C₁₋₆ alkyl group;

an arylsulfonyloxyalkyl group such as a tosyloxymethyl group or a 2-tosyloxy-1,1-dimethylethyl group, preferably a C₁₋₆ alkyl-substituted C₆₋₁₀ arylsulfonyloxy C₁₋₆ alkyl group;

a cyanoalkyl group such as a cyanomethyl group, a 2-cyanoethyl group, or a 1-cyano-1-methylethyl group, preferably a cyano C₁₋₆ alkyl group;

an acylalkyl group such as a formylmethyl group, a 2-formylethyl group, a 3-formylpropyl group, a 1-formyl-1-methylethyl group, a 2-formyl-1,1-dimethylethyl group, an acetylmethyl group, a 2-acetyethyl group, a 3-acetylpropyl group, 1-acetyl-1-methylethyl group, or a 2-acetyl-1,1-dimethylethyl group, preferably a C₁₋₆ acyl C₁₋₆ alkyl group;

a 2-hydroxyiminoalkyl group such as a 2-hydroxyiminoethyl group, a 2-hydroxyimino-1-methylethyl group, a 2-hydroxy-1,1-dimethylethyl group, or a 2-hydroxyiminopropyl group, preferably 2-hydroxyimino C₂₋₆ alkyl group;

an acylalkyl group such as an acetylmethyl group, a 2-acetyethyl group, a

3-acetylpropyl group, a 1-acetyl-1-methylethyl group, or a 2-acetyl-1,1-dimethylethyl group, preferably a formyl C₁₋₆ alkyl group;

a carboxyalkyl group such as a carboxymethyl group, a 2-carboxyethyl group, a 3-carboxypropyl group, a 1-carboxy-1-methylethyl group, or a

5 2-carboxy-1,1-dimethylethyl group, preferably a carboxy C₁₋₆ alkyl group;

an alkoxycarbonylalkyl group such as a methoxycarbonylmethyl group, a 2-methoxycarbonylethyl group, a 3-methoxycarbonylpropyl group, a 1-methoxycarbonyl-1-methylethyl group, or a 2-methoxycarbonyl-1,1-dimethylethyl group, preferably a C₁₋₆ alkoxycarbonyl C₁₋₆ alkyl group;

10 an azidoalkyl group such as an azidomethyl group, a 2-azidoethyl group, or a 1-azido-1-methylethyl group, preferably an azido C₁₋₆ alkyl group; and the like.

[0033]

A “C₂₋₈ alkenyl group” is an unsaturated hydrocarbon group including 2 to 8 carbon atoms having at least one carbon-carbon double bond. The C₂₋₈ alkenyl group

15 may be a straight chain or a branched chain. Examples of the C₂₋₈ alkenyl group include a vinyl group, a 1-propenyl group, an isopropenyl group, an allyl group, a 1-butenyl group, a 2-butenyl group, a 3-butenyl group, a 1-pentenyl group, a 2-pentenyl group, a 3-pentenyl group, a 4-pentenyl group, a 1-hexenyl group, a 2-hexenyl group, a 3-hexenyl group, a 4-hexenyl group, a 5-hexenyl group, a 1-heptenyl group, a 6-heptenyl group, a 20 1-octenyl group, a 7-octenyl group, a 1-methyl-allyl group, a 2-methyl-allyl group, a 1-methyl-2-butenyl group, a 2-methyl-2-butenyl group, and the like. Among these, a C₂₋₆ alkenyl group is preferable.

[0034]

Examples of a “substituted C₂₋₈ alkenyl group” include: a haloalkenyl group

25 such as a 3-chloro-2-propenyl group, 4-chloro-2-butenyl group, a 4,4-dichloro-3-butenyl

group, 4,4-difluoro-3-butenyl group, a 3,3-dichloro-2-propenyl group, a 2,3-dichloro-2-propenyl group, a 3,3-difluoro-2-propenyl group, or a 2,4,6-trichloro-2-hexenyl group, preferably C₂₋₆ haloalkenyl group;

a hydroxyalkenyl group such as a 3-hydroxy-1-propenyl group, a 4-hydroxy-1-butenyl group, a 1-hydroxyallyl group, or a 1-hydroxy-2-methylallyl group, preferably a hydroxy C₂₋₆ alkenyl group; and the like.

[0035]

A “C₂₋₈ alkynyl group” is an unsaturated hydrocarbon group including 2 to 8 carbon atoms having at least one carbon-carbon triple bond. The C₂₋₈ alkynyl group may be a straight chain or a branched chain. Examples of the C₂₋₈ alkynyl group include an ethynyl group, a 1-propynyl group, a propargyl group, a 1-butylnyl group, a 2-butylnyl group, a 3-butylnyl group, a 1-pentylnyl group, a 2-pentylnyl group, a 3-pentylnyl group, a 4-pentylnyl group, a 1-hexynyl group, a 1-methyl-2-propynyl group, a 2-methyl-3-butylnyl group, a 1-methyl-2-butylnyl group, a 2-methyl-3-pentylnyl group, a 1,1-dimethyl-2-butylnyl group, and the like. Among these, a C₂₋₆ alkynyl group is preferable.

[0036]

Examples of a “substituted C₂₋₈ alkynyl group” include a haloalkynyl group such as a 3-chloro-1-propynyl group, a 3-chloro-1-butylnyl group, a 3-bromo-1-butylnyl group, a 3-bromo-2-propynyl group, a 3-iodo-2-propynyl group, a 3-bromo-1-hexynyl group, a 4,4,6,6-tetrafluoro-1-dodecynyl group, a 5,5-dichloro-2-methyl-3-pentylnyl group, or a 4-chloro-1,1-dimethyl-2-butylnyl group, preferably a C₂₋₆ haloalkynyl group, and the like.

[0037]

A “C₃₋₈ cycloalkyl group” is an alkyl group including 3 to 8 carbon atoms having a cyclic moiety. Examples of the C₃₋₈ cycloalkyl group include a cyclopropyl

group, a cyclobutyl group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, and the like. Among these, a C₃₋₆ cycloalkyl group is preferable.

[0038]

Examples of a “substituted C₃₋₈ cycloalkyl group” include an alkyl-substituted
5 cycloalkyl group such as a 2,3,3-trimethylcyclobutyl group, a
4,4,6,6-tetramethylcyclohexyl group, or a 1,3-dibutylcyclohexyl group, preferably a C₃₋₆
cycloalkyl group in which 1 to 3 C₁₋₆ alkyl group(s) is(are) substituted, and the like.

[0039]

A “C₄₋₈ cycloalkenyl group” is an alkenyl group including 4 to 8 carbon atoms
10 having a cyclic moiety. Examples of the C₄₋₈ cycloalkenyl group include a
1-cyclobutenyl group, a 1-cyclopentenyl group, a 3-cyclopentenyl group, a
1-cyclohexenyl group, a 3-cyclohexenyl group, a 3-cycloheptenyl group, a
4-cyclooctenyl group, and the like.

[0040]

15 Examples of a “substituted C₄₋₈ cycloalkenyl group” include an alkyl-substituted
cycloalkenyl group such as a 2-methyl-3-cyclohexenyl group or a
3,4-dimethyl-3-cyclohexenyl group, preferably a C₄₋₆ cycloalkenyl group in which 1 to 3
C₁₋₆ alkyl group(s) is(are) substituted, and the like.

[0041]

20 A “C₆₋₁₀ aryl group” is a monocyclic or polycyclic aryl group having 6 to 10
carbon atoms. In the polycyclic aryl group, if at least one ring is an aromatic ring, the
remaining ring(s) may be any of a saturated alicyclic ring, an unsaturated alicyclic ring,
and an aromatic ring. Examples of the C₆₋₁₀ aryl group include a phenyl group, a
naphthyl group, an azulenyl group, an indenyl group, an indanyl group, a tetralinyl group,
25 and the like. Among these, a phenyl group is preferable.

Examples of a “substituted C₆₋₁₀ aryl group” include an alkyl-substituted aryl group such as a 2-chlorophenyl group, a 3,5-dichlorophenyl group, a 4-fluorophenyl group, a 3,5-difluorophenyl group, a 4-trifluoromethylphenyl group, or a 2-methoxy-1-naphthyl group, a halogeno-substituted aryl group, and an

5 alkoxy-substituted aryl group, preferably a C₁₋₆ alkyl-substituted C₆₋₁₀ aryl group, a halogeno-substituted C₆₋₁₀ aryl group, and a C₁₋₆ alkoxy substituted aryl group.

[0042]

A “heterocyclic group” includes 1 to 4 heteroatom(s) selected from the group consisting of a nitrogen atom, an oxygen atom, and a sulfur atom, as a constituent atom

10 of a ring. The heterocyclic group may be monocyclic or polycyclic.

Examples of the heterocyclic group include a 5-membered heteroaryl group, a 6-membered heteroaryl group, a condensed heteroaryl group, a saturated heterocyclic group, a partially unsaturated heterocyclic group and the like.

[0043]

15 Examples of the 5-membered heteroaryl group include: a pyrrolyl group such as a pyrrol-1-yl group, a pyrrol-2-yl group, or a pyrrol-3-yl group; a furyl group such as a furan-2-yl group or a furan-3-yl group; a thienyl group such as a thiophen-2-yl group, or a thiophen-3-yl group; an imidazolyl group such as an imidazol-1-yl group, an imidazol-2-yl group, an imidazol-4-yl group, or an imidazol-5-yl group; a pyrazolyl

20 group such as a pyrazol-1-yl group, a pyrazol-3-yl group, a pyrazol-4-yl group, or a pyrazol-5-yl group; an oxazolyl group such as an oxazol-2-yl group, an oxazol-4-yl group, or an oxazol-5-yl group; an isoxazolyl group such as an isoxazol-3-yl group, an isoxazol-4-yl group, or an isoxazol-5-yl group; a thiazolyl group such as a thiazol-2-yl group, a thiazol-4-yl group, or a thiazol-5-yl group; an isothiazolyl group such as an

25 isothiazol-3-yl group, an isothiazol-4-yl group, or a thiazol-5-yl group; a triazolyl group

such as a 1,2,3-triazol-1-yl group, a 1,2,3-triazol-4-yl group, a 1,2,3-triazol-5-yl group, a 1,2,4-triazol-1-yl group, a 1,2,4-triazol-3-yl group, or a 1,2,4-triazol-5-yl group; an oxadiazolyl group such as a 1,2,4-oxadiazol-3-yl group, a 1,2,4-oxadiazol-5-yl group, or a 1,3,4-oxadiazol-2-yl group; a thiadiazolyl group such as a 1,2,4-thiadiazol-3-yl group, 1,2,4-thiadiazol-5-yl group, or a 1,3,4-thiadiazol-2-yl group; a tetrazolyl group such as a tetrazol-1-yl group or a tetrazol-2-yl group; and the like.

[0044]

Examples of the 6-membered heteroaryl group include: a pyridyl group such as a pyridin-2-yl group, a pyridin-3-yl group, or a pyridin-4-yl group; a pyrazinyl group such as a pyrazin-2-yl group or a pyrazin-3-yl group; a pyrimidinyl group such as a pyrimidin-2-yl group, a pyrimidin-4-yl group, or a pyrimidin-5-yl group; a pyridazinyl group such a pyridazin-3-yl group or a pyridazin-4-yl group; a triazinyl group; and the like.

[0045]

Examples of the condensed heteroaryl group include: indol-1-yl group, an indol-2-yl group, an indol-3-yl group, an indol-4-yl group, an indol-5-yl group, an indol-6-yl group, an indol-7-yl group; a benzofuran-2-yl group, a benzofuran-3-yl group, a benzofuran-4-yl group, a benzofuran-5-yl group, a benzofuran-6-yl group, a benzofuran-7-yl group; a benzothiophen-2-yl group, a benzothiophen-3-yl group, a benzothiophen-4-yl group, a benzothiophen-5-yl group, a benzothiophen-6-yl group, a benzothiophen-7-yl group; a benzoimidazol-1-yl group, a benzoimidazol-2-yl group, a benzoimidazol-4-yl group, a benzoimidazol-5-yl group, a benzoxazol-2-yl group, a benzoxazol-4-yl group, a benzoxazol-5-yl group, a benzothiazol-2-yl group, a benzothiazol-4-yl group, a benzothiazol-5-yl group; a quinolin-2-yl group, a quinolin-3-yl group, a quinolin-4-yl group, a quinolin-5-yl group, a quinolin-6-yl group,

a quinolin-7-yl group, a quinolin-8-yl group; and the like.

[0046]

Examples of the other heterocyclic group include: a 3-membered saturated heterocyclic ring such as an aziridin-1-yl group, an aziridin-2-yl group, or an oxiranyl group; a 5-membered saturated heterocyclic ring such as a pyrrolidin-1-yl group, a pyrrolidin-2-yl group, a pyrrolidin-3-yl group, a tetrahydrofuran-2-yl group, a tetrahydrofuran-3-yl group, or a [1,3]dioxiran-2-yl group; a 6-membered saturated heterocyclic ring such as a piperidin-1-yl group, a piperidin-2-yl group, a piperidin-3-yl group, a piperidin-4-yl group, a piperazin-1-yl group, a piperazin-2-yl group, a morpholin-2-yl group, a morpholin-3-yl group, or a morpholin-4-yl group; a 1,3-benzodioxol-4-yl group, a 1,3-benzodioxol-5-yl group, a 1,4-benzodioxan-5-yl group, a 1,4-benzodioxan-6-yl group, a 3,4-dihydro-2H-1,5-benzodioxepin-6-yl group, a 3,4-dihydro-2H-1,5-benzodioxepin-7-yl group, a 2,3-dihydrobenzofuran-4-yl group, a 2,3-dihydrobenzofuran-5-yl group, a 2,3-dihydrobenzofuran-6-yl group, or a 2,3-dihydrobenzofuran-7-yl group; and the like.

[0047]

Examples of a “substituted heterocyclic group” include: a 4-chloro-2-pyridinyl group, a 3-chloro-2-pyrazinyl group, a 4-methyl-2-pyridinyl group, a 5-trifluoromethyl-2-pyrimidinyl group, a 3-methyl-2-quinolyl group, and the like.

[0048]

A “C₁₋₈ acyl group” is a group in which a hydrogen atom, a C₁₋₇ alkyl group, a C₂₋₇ alkenyl group, a C₂₋₇ alkynyl group, a C₆₋₇ aryl group, or a 5- to 7-membered heterocyclic group is bonded to a carbonyl group.

Examples of the C₁₋₈ acyl group include: a formyl group; an alkylcarbonyl group such as an acetyl group, a propionyl group, an n-propylcarbonyl group, an

n-butylcarbonyl group, a pentanoyl group, a valeryl group, an octanoyl group, an i-propylcarbonyl group, an i-butylcarbonyl group, a pivaloyl group, or an isovaleryl group, preferably a C₁₋₆ alkylcarbonyl group; an alkenylcarbonyl group such as an acryloyl group, or a metacryloyl group, preferably a C₂₋₆ alkenylcarbonyl group; an alkynylcarbonyl group such as a propioloyl group, preferably a C₂₋₆ alkynylcarbonyl group; an arylcarbonyl group such as a benzoyl group; a heterocyclic carbonyl group such as a 2-pyridylcarbonyl group or a thienylcarbonyl group; and the like.

[0049]

Examples of a “substituted C₁₋₈ acyl group” include: a haloacyl group such as a monofluoroacetyl group, a monochloroacetyl group, a monobromoacetyl group, a difluoroacetyl group, a dichloroacetyl group, a dibromoacetyl group, a trifluoroacetyl group, a trichloroacetyl group, a tribromoacetyl group, a 3,3,3-trifluoropropionyl group, a 3,3,3-trichloropropionyl group, or a 2,2,3,3,3-pentafluoropropionyl group, preferably a C₁₋₇ haloacyl group; and the like.

[0050]

A “(1-imino)C₁₋₈ alkyl group” is an iminomethyl group or a group in which a C₁₋₇ alkyl group is bonded to an iminomethyl group. Examples of the (1-imino)C₁₋₈ alkyl group include an iminomethyl group, a (1-imino)ethyl group, a (1-imino)propyl group, a (1-imino)butyl group, a (1-imino)pentyl group, a (1-imino)hexyl group, a (1-imino)heptyl group, and the like. Among these, a (1-imino) C₁₋₆ alkyl group is preferable.

Examples of a “substituted (1-imino)C₁₋₈ alkyl group” include: a (1-hydroxyimino)alkyl group such as a hydroxyiminomethyl group, a (1-hydroxyimino)ethyl group, a (1-hydroxyimino)propyl group, or a (1-hydroxyimino)butyl group, preferably a (1-hydroxyimino)C₁₋₆ alkyl group; a

(1-alkoxyimino)alkyl group such as a methoxyiminomethyl group, a (1-ethoxyimino)methyl group, a (1-methoxyimino)ethyl group, a (1-t-butoxyimino)ethyl group, or a (1-ethoxyimino)ethyl group, preferably a (1-(C₁₋₆ alkoxy)imino)C₁₋₆ alkyl group; and the like.

5 [0051]

A “substituted carboxyl group” is a group in which a C₁₋₆ alkyl group, a C₂₋₆ alkenyl group, a C₂₋₆ alkynyl group, a C₆₋₁₀ aryl group, a C₆₋₁₀ aryl C₁₋₆ alkyl group, or a 5 to 6-membered heterocyclic group is bonded to a carbonyl group.

[0052]

10 Examples of the “substituted carboxyl group” include: an alkoxycarbonyl group such as a methoxycarbonyl group, an ethoxycarbonyl group, an n-propoxycarbonyl group, an i-propoxycarbonyl group, an n-butoxycarbonyl group, an i-butoxycarbonyl group, a t-butoxycarbonyl group, an n-pentyloxycarbonyl group, or an n-hexyloxycarbonyl group, preferably a C₁₋₆ alkoxycarbonyl group;

15 an alkenyloxycarbonyl group such as a vinyloxycarbonyl group or an allyloxycarbonyl group, preferably a C₂₋₆ alkenyloxycarbonyl group;

an alkynyloxycarbonyl group such as an ethynyloxycarbonyl group or a propargyloxycarbonyl group, preferably a C₂₋₆ alkynyloxycarbonyl group;

20 an aryloxycarbonyl group such as a phenoxycarbonyl group or a naphthoxycarbonyl group, preferably a C₆₋₁₀ aryloxycarbonyl group;

an aralkyloxycarbonyl group such as a benzyloxycarbonyl group, preferably a C₆₋₁₀ aryl C₁₋₆ alkoxycarbonyl group; and the like.

[0053]

25 A “substituted carbamoyl group” is a group in which a C₁₋₆ alkyl group, a C₂₋₆ alkenyl group, a C₂₋₆ alkynyl group, a C₆₋₁₀ aryl group, a C₆₋₁₀ aryl C₁₋₆ alkyl group, or a 5

to 6-membered heterocyclic group is bonded to a carbamoyl group.

[0054]

Examples of the “substituted carbamoyl group” include: a monoalkylcarbamoyl group or a dialkylcarbamoyl group, such as a methylcarbamoyl group, an ethylcarbamoyl group, a dimethylcarbamoyl group, or a diethylcarbamoyl group, preferably a mono C₁₋₆ alkylcarbamoyl group or a di C₁₋₆ alkylcarbamoyl group; a monoarylcarbamoyl group such as a phenylcarbamoyl group or a 4-methylphenylcarbamoyl group, preferably a mono C₆₋₁₀ arylcarbamoyl group; and the like.

[0055]

Examples of a “substituted hydroxyl group” include: an alkoxy group such as a methoxy group, an ethoxy group, an n-propoxy group, an n-butoxy group, an n-pentyloxy group, an n-hexyloxy group, a decyloxy group, a dodecyloxy group, a lauryloxy group, an i-propoxy group, an i-butoxy group, an s-butoxy group, a t-butoxy group, a 1-ethylpropoxy group, an i-hexyloxy group, a 4-methylpentoxy group, a 3-methylpentoxy group, a 2-methylpentoxy group, a 1-methylpentoxy group, a 3,3-dimethylbutoxy group, a 2,2-dimethylbutoxy group, a 1,1-dimethylbutoxy group, a 1,2-dimethylbutoxy group, a 1,3-dimethylbutoxy group, a 2,3-dimethylbutoxy group, a 1-ethylbutoxy group, or a 2-ethylbutoxy group, preferably a C₁₋₆ alkoxy group;

[0056]

a cycloalkylalkoxy group such as a cyclopropylmethyloxy group or a 2-cyclopentylethyloxy group, preferably a C₃₋₈ cycloalkyl C₁₋₆ alkoxy group; an aralkyloxy group such as a benzyloxy group, preferably a C₆₋₁₀ aryl C₁₋₆ alkoxy group; a haloalkoxy group such as a chloromethoxy group, a dichloromethoxy group, a trichloromethoxy group, a trifluoromethoxy group, a 1-fluoroethoxy group, a 1,1-difluoroethoxy group, a 2,2,2-trifluoroethoxy group, or a pentafluoroethoxy group,

preferably a C₁₋₆ haloalkoxy group; an alkenyloxy group such as a vinyloxy group, a 1-propenyloxy group, an allyloxy group, a 1-butenyloxy group, a 2-butenyloxy group, a 3-butenyloxy group, a 1-pentenyloxy group, a 2-pentenyloxy group, a 3-pentenyloxy group, a 4-pentenyloxy group, a 1-hexenyloxy group, a 2-hexenyloxy group, a 3-hexenyloxy group, a 4-hexenyloxy group, a 5-hexenyloxy group, a 1-methyl-2-propenyloxy group, a 2-methyl-2-propenyloxy group, a 1-methyl-2-butenyloxy group, or a 2-methyl-2-butenyloxy group, preferably a C₂₋₆ alkenyloxy group;

[0057]

an alkynyloxy group such as an ethynyloxy group, a propynyloxy group, a propargyloxy group, a 1-butyloxy group, a 2-butyloxy group, a 3-butyloxy group, a 1-pentyloxy group, a 2-pentyloxy group, a 3-pentyloxy group, a 4-pentyloxy group, a 1-hexyloxy group, a 1-methyl-2-propynyloxy group, a 2-methyl-3-butyloxy group, a 1-methyl-2-butyloxy group, a 2-methyl-3-pentyloxy group, or a 1,1-dimethyl-2-butyloxy group, preferably a C₂₋₆ alkynyloxy group; a cycloalkyloxy group such as a cyclopropyloxy group, a cyclobutyloxy group, a cyclopentyloxy group, a cyclohexyloxy group, a cycloheptyloxy group, a cyclooctyloxy group, a 2-methylcyclopropyloxy group, a 2-ethylcyclopropyloxy group, a 2,3,3-trimethylcyclobutyloxy group, a 2-methylcyclopentyloxy group, a 2-ethylcyclohexyloxy group, a 2-ethylcyclooctyloxy group, a 4,4,6,6-tetramethylcyclohexyloxy group, or a 1,3-dibutylcyclohexyloxy group, preferably a C₃₋₆ cycloalkyloxy group; an aryloxy group such as a phenyloxy group, a naphthyloxy group, an azulenyloxy group, an indenyloxy group, an indanyloxy group, or a tetralinyloxy group, preferably a C₆₋₁₀ aryloxy group;

an arylalkyloxy group (aralkyloxy group) such as a benzyloxy group, a

phenethyloxy group, or a 2-naphthylmethyloxy group, preferably a C₆₋₁₀ aryl C₁₋₆ alkyloxy group;

an acyloxy group such as an acetyloxy group, a propionyloxy group, an n-propylcarbonyloxy group, an i-propylcarbonyloxy group, an n-butylcarbonyloxy group, an i-butylcarbonyloxy group, a pentanoyloxy group, or a pivaloyloxy group, preferably a C₁₋₇ acyloxy group;

an alkoxycarbonylalkyloxy group such as a methoxycarbonylmethyloxy group or a 1-methoxycarbonyl-1-methylethyloxy group, preferably a C₁₋₆ alkoxycarbonyl C₁₋₆ alkoxy group;

a trialkylsilyloxy group such as a trimethylsilyloxy group or a t-butyltrimethylsilyloxy group, preferably a tri C₁₋₆ alkylsilyloxy group;

and the like.

[0058]

Examples of a “substituted amino group” include: an alkylamino group such as a methylamino group, an ethylamino group, an n-propylamino group, an n-butylamino group, a dimethylamino group, or a diethylamino group, preferably a mono C₁₋₆ alkylamino group or a di C₁₋₆ alkylamino group; a mono C₁₋₆ alkylideneamino group such as a methylideneamino group, or an ethylideneamino group; a monoarylamino group such as a phenylamino group or a 4-methylphenylamino group, preferably a mono C₆₋₁₀ arylamino group; a diarylamino group such as a di-1-naphthylamino group, preferably a di C₆₋₁₀ arylamino group; an aralkylamino group such as a benzylamino group, preferably a C₆₋₁₀ aryl C₁₋₆ alkylamino group; an acylamino group such as an acetylamino group, a trifluoroacetylamino group, or a benzoylamino group, preferably a C₁₋₆ acylamino group; an alkoxycarbonylamino group such as a methoxycarbonylamino group or a t-butoxycarbonylamino group, preferably a C₁₋₆ alkoxycarbonylamino group;

and the like.

[0059]

Examples of a “substituted mercapto group” include: an alkylthio group such as a methylthio group or an ethylthio group, preferably a C₁₋₆ alkylthio group; an arylthio
 5 group such as a phenylthio group or a 4-methylphenylthio group, preferably a C₆₋₁₀ arylthio group; an acylthio group such as an acetylthio group or a benzoylthio group, preferably a C₁₋₆ acylthio group; and the like.

[0060]

Examples of a “substituted sulfonyl group” include: an alkylsulfonyl group such
 10 as a methylsulfonyl group, an ethylsulfonyl group, an n-propylsulfonyl group, an i-propylsulfonyl group, an n-butylsulfonyl group, an i-butylsulfonyl group, an s-butylsulfonyl group, a t-butylsulfonyl group, an n-pentylsulfonyl group, an i-pentylsulfonyl group, a neopentylsulfonyl group, a 1-ethylpropylsulfonyl group, an n-hexylsulfonyl group, or an i-hexylsulfonyl group, preferably a C₁₋₆ alkylsulfonyl group;
 15 a haloalkylsulfonyl group such as a trifluoromethylsulfonyl group, preferably a C₁₋₆ haloalkylsulfonyl group; an arylsulfonyl group such as a phenylsulfonyl group or a 4-methylphenylsulfonyl group, preferably a C₆₋₁₀ arylsulfonyl group; a sulfo group; an alkoxysulfonyl group such as a methoxysulfonyl group or an ethoxysulfonyl group, preferably a C₁₋₆ alkoxysulfonyl group; a sulfamoyl group; a sulfamoyl group such as an
 20 N-methylsulfamoyl group, an N-ethylsulfamoyl group, a N,N-dimethylsulfamoyl group, or an N,N-diethylsulfamoyl group, preferably a mono C₁₋₆ alkylsulfamoyl group or a di C₁₋₆ alkylsulfamoyl group; a monoarylsulfamoyl group such as a phenylsulfamoyl group or a 4-methylsulfamoyl group, preferably a mono C₆₋₁₀ arylsulfamoyl group; and the like.

[0061]

25 Examples of a “halogeno group” include a fluorine atom, a chlorine atom, a

bromine atom, an iodine atom, and the like.

[0062]

As a preferred combination of a group represented by $CR^1R^2R^3$, the following combination may be included.

5 R^1 is an unsubstituted or substituted hydroxyl group;

R^2 is a hydrogen atom or an unsubstituted or substituted C_{1-8} alkyl group; and

R^3 is a hydrogen atom, an unsubstituted or substituted C_{1-8} alkyl group, an unsubstituted or substituted C_{2-8} alkenyl group, an unsubstituted or substituted C_{2-8} alkynyl group, an unsubstituted or substituted C_{3-8} cycloalkyl group, an unsubstituted or substituted C_{6-10} aryl group, an unsubstituted or substituted heterocyclic group, an unsubstituted or substituted C_{1-8} acyl group, an unsubstituted or substituted (1-imino) C_{1-8} alkyl group, an unsubstituted or substituted C_{1-8} alkoxy carbonyl group, an unsubstituted or substituted hydroxyl group, or a cyano group.

[0063]

15 The following combination may also be included as a preferred example.

R^1 is an unsubstituted or substituted C_{1-8} alkyl group;

R^2 is an unsubstituted or substituted C_{1-8} alkyl group; and

R^3 is a substituted C_{1-8} alkyl group, an unsubstituted or substituted C_{2-8} alkenyl group, an unsubstituted or substituted C_{2-8} alkynyl group, an unsubstituted or substituted C_{3-8} cycloalkyl group, an unsubstituted or substituted C_{6-10} aryl group, an unsubstituted or substituted heterocyclic group, an unsubstituted or substituted C_{1-8} acyl group, an unsubstituted or substituted (1-imino) C_{1-8} alkyl group, an unsubstituted or substituted C_{1-8} alkoxy carbonyl group, an unsubstituted or substituted carbamoyl group, an unsubstituted or substituted sulfonyl group, or a cyano group.

25 [0064]

R^1 and R^2 may be joined to form an unsubstituted or substituted 5- to 8-membered ring, or to form $O=$, $R^a R^b C=$, or $R' N=$.

Here, R^a represents a hydrogen atom or an unsubstituted or substituted C_{1-8} alkyl group. R^b represents a hydrogen atom or an unsubstituted or substituted C_{1-8} alkyl group. R' represents an unsubstituted or substituted hydroxyl group, or an unsubstituted or substituted C_{1-8} alkyl group.

The unsubstituted or substituted C_{1-8} alkyl group in R^a , R^b , and R' may be the same as the " C_{1-8} alkyl group" given as the examples in R^1 to R^3 above.

The substituted hydroxyl group in R' may be the same as the "substituted hydroxyl group" given as the examples in R^1 to R^3 above.

[0065]

Examples of the unsubstituted or substituted 5- to 8-membered ring, which may be formed from R^1 and R^2 being joined, include: an aliphatic hydrocarbon ring such as a cyclopropane ring, a cyclobutane ring, a cyclopentane ring, a cyclohexane ring, a cycloheptane ring, or a cyclooctane ring, preferably a C_{3-8} cycloalkane ring; an unsaturated heterocyclic ring such as an oxirane ring, a [1,3]dioxirane ring, a dihydro-2H-pyran ring, a dihydro-2H-thiopyran ring, and a tetrahydropyridine ring, preferably an oxygen-containing 3- to 5-membered unsaturated heterocyclic ring.

[0066]

The unsubstituted or substituted C_{6-10} aryl group in R may be the same as the " C_{6-10} aryl group" given as the examples in R^1 to R^3 above. It is preferably a phenyl group.

[0067]

$[R^4]$

R^4 each independently represents an unsubstituted or substituted C_{1-8} alkyl group,

an unsubstituted or substituted C₂₋₈ alkenyl group, an unsubstituted or substituted C₂₋₈ alkynyl group, an unsubstituted or substituted C₃₋₈ cycloalkyl group, an unsubstituted or substituted C₄₋₈ cycloalkenyl group, an unsubstituted or substituted C₆₋₁₀ aryl group, an unsubstituted or substituted heterocyclic group, an unsubstituted or substituted C₁₋₈ acyl group, an unsubstituted or substituted (1-imino)C₁₋₈ alkyl group, an unsubstituted or substituted carboxyl group, an unsubstituted or substituted carbamoyl group, an unsubstituted or substituted hydroxyl group, an unsubstituted or substituted amino group, an unsubstituted or substituted mercapto group, a substituted sulfonyl group, a halogeno group, a cyano group, or a nitro group.

10 The groups represented by R⁴ may be the same as the examples given in the groups represented by R¹ to R³.

In Formulae (I), (II) and (III), m and m1 represent the number of R⁴ and are an integer of 0 to 6.

15 R⁴ preferably represents a C₁₋₆ alkyl group, a C₁₋₆ haloalkyl group, a C₂₋₆ alkenyl group, a C₃₋₈ cycloalkyl group, a hydroxyl group, a C₁₋₆ alkoxy group, or a halogeno group.

[0068]

[R⁵]

20 R⁵ each independently represents an unsubstituted or substituted C₁₋₈ alkyl group, an unsubstituted or substituted C₂₋₈ alkenyl group, an unsubstituted or substituted C₂₋₈ alkynyl group, an unsubstituted or substituted C₃₋₈ cycloalkyl group, an unsubstituted or substituted C₄₋₈ cycloalkenyl group, an unsubstituted or substituted C₆₋₁₀ aryl group, an unsubstituted or substituted heterocyclic group, an unsubstituted or substituted C₁₋₈ acyl group, an unsubstituted or substituted (1-imino)C₁₋₈ alkyl group, an unsubstituted or substituted carboxyl group, an unsubstituted or substituted carbamoyl group, an

25

unsubstituted or substituted hydroxyl group, an unsubstituted or substituted amino group, an unsubstituted or substituted mercapto group, a substituted sulfonyl, a halogeno group, a cyano group, or a nitro group.

The groups represented by R^5 may be the same as the examples given in the groups represented by R^1 to R^3 .

In Formula (I) and (II), n represents the number of R^5 and is an integer of 0 to 5.

In Formula (III), n_1 represents the number of R^5 and is an integer of 0 to 4.

R^5 preferably represents a C_{1-6} alkyl group, a C_{1-6} haloalkyl group, a C_{6-10} aryl C_{1-6} alkyl group, a C_{3-8} cycloalkyl group, a C_{6-10} aryl group, a C_{1-7} acyl group, a C_{1-6} alkoxycarbonyl group, a C_{1-6} alkoxy group, an amino group, a mono C_{1-6} alkylamino group, di C_{1-6} alkylamino group, a C_{1-6} alkoxycarbonylamino group, a C_{1-6} alkylthio group, a C_{1-6} alkylsulfonyl group, a halogeno group, a cyano group or a nitro group.

[0069]

Any one of R^1 to R^3 and any one of R^5 , by being joined, may form an unsubstituted or substituted 5- to 8-membered ring.

Examples of the 5- to 8-membered ring include: an aromatic hydrocarbon ring such as a benzene ring; a C_{5-8} cycloalkene ring such as a cyclopentene ring, a cyclopentadiene ring, a cyclohexene ring, a cycloheptene ring, or a cyclooctene ring; and the like.

[0070]

[A, D]

When R is a group represented by $CR^1R^2R^3$, A represents a 5- to 7-membered hydrocarbon ring or a 5- to 7-membered heterocyclic ring.

Examples of the 5- to 7-membered hydrocarbon ring include: an aromatic hydrocarbon ring such as a benzene ring; a C_{5-7} cycloalkene ring such as a cyclopentene

ring, a cyclohexene ring, a cycloheptene ring; an aromatic 5- to 7-membered heterocyclic ring such as a furan ring, a thiophene ring, a pyrrole ring, an imidazole ring, a pyrazole ring, a thiazole ring, an oxazole ring, an isoxazole ring, a pyridine ring, a pyrazine ring, a pyrimidine ring, a pyridazine ring, an azepine ring, or a diazepine ring; an unsaturated 5-
 5 to 7-membered heterocyclic ring such as a dihydro-2H-pyran ring, a dihydro-2H-thiopyran ring, or a tetrahydropyridine ring; and the like. Among these, an aromatic hydrocarbon ring is preferable and a benzene ring is more preferable.

When R is an unsubstituted or substituted C₆₋₁₀ aryl group or a cyano group, A represents a benzene ring.

10 That is, it is preferable that the compound according to the present invention be a compound (II) or (III).

[0071]

D represents a 5- to 7-membered hydrocarbon ring or a 5- to 7-membered heterocyclic ring. Examples of the 5- to 7-membered hydrocarbon ring and the 5- to
 15 7-membered heterocyclic ring include the same as the examples given above as A, and among these, an aromatic hydrocarbon ring is preferable and a benzene ring is more preferable.

That is, it is more preferable that the compound according to the present invention be a compound (III).

20 [0072]

[X, Y, Z]

X represents an oxygen atom, a sulfur atom, a sulfenyl group, a sulfonyl group, an unsubstituted or substituted carbon atom, or an unsubstituted or substituted nitrogen atom.

25 Examples of a substituent of the carbon atom include an oxo group, a C₂₋₆

alkenylamino group, and a hydroxyl group.

[0073]

Y represents a carbon atom or a nitrogen atom. Z represents a carbon atom or a nitrogen atom. It is preferable that Y and Z be both carbon atoms, and D be an aromatic
5 ring which includes both Y and Z.

[0074]

A salt or an N-oxide compound according to the present invention is not particularly limited as long as it is an agriculturally acceptable salt or N-oxide compound. Examples of the salt include: a salt of an inorganic acid such as hydrochloric acid or
10 sulfuric acid; a salt of an organic acid such as acetic acid or lactic acid; a salt of an alkali metal such as lithium, sodium, or potassium; a salt of an alkali earth metal such as calcium or magnesium; a salt of a transition metal such as iron or copper; a salt of an organic base such as ammonia, triethylamine, tributylamine, pyridine, or hydrazine; and the like.


15 [0075]

(Production of the compound according to the present invention)

The compound according to the present invention may be produced in accordance with the following synthesis methods.

(Synthesis method 1)

20 [0076]



(In the formulae, R, R⁴, R⁵, A, D, X, Y, Z, m and n represent the same meaning as those described above. Q represents a halogen atom.)


The compound represented by Formula (I) may be produced by reacting a
 5 compound represented by Formula (1) and a compound represented by Formula (2) by conventional methods.

According to the present invention, 7,8-difluoro-3-iodo-quinoline is a useful intermediate product.

[0077]

10 (Synthesis method 2)

[0078]



(In the formulae, R, R⁴, R⁵, A, D, Q, X, Y, Z, m and n represent the same meaning as those described above.)

15 The compound represented by Formula (I) may be produced by reacting a compound represented by Formula (3) and a compound represented by Formula (4) by


conventional methods.

According to the present invention, 8- fluoro-3-hydroxyquinoline, 7,8-difluoro-3-hydroxyquinoline, 8-fluoro-3-hydroxy-2-methylquinoline, or 7,8-difluoro-3-hydroxy-2-methylquinoline is a useful intermediate product.

5 [0079]

(Synthesis method 3)

[0080]



(In the formulae, R, R⁴, R⁵, A, D, X, Y, Z, m and n represent the same meaning
 10 as those described above. M represents lithium or magnesium.)

A compound represented by Formula (5) is obtained by lithiation or magnesium
 complexation using an alkyllithium reagent or a Grignard reagent or a manufactured
 complex prepared from an alkyllithium reagent or a Grignard reagent, and then a
 compound represented by Formula (6) is added thereto, resulting in the production of a
 15 compound represented by Formula (7).

Examples of the alkyllithium reagent used in the lithiation include methyllithium,
 n-butyllithium, s- butyllithium, t-butyllithium, and the like.

Examples of the Grignard reagent used in the magnesium complexation include
 methylmagnesium chloride, ethylmagnesium chloride, n-butyllmagnesium chloride,
 20 i-propylmagnesium chloride, and the like. Also, for example, a manufactured complex


prepared from n-butyilmagnesium chloride and n-butyllithium may be used.

A solvent used in the lithiation or magnesium complexation is not particularly limited as long as it forms an anhydrous reaction system without dissolving the compound to react therewith or exhibit any particular interaction therewith. Suitable examples thereof include: an alkane-based solvent such as pentane, hexane, heptane, ISOPAR (registered trademark) E, or ISOPAR (registered trademark) G; an aromatic-based solvent such as benzene, toluene, or ortho-xylene; an ether-based solvent such as diethyl ether or tetrahydrofuran; and a mixture thereof. Among these, an ether-based solvent such as diethyl ether or tetrahydrofuran is preferable. The reaction may be performed in a nitrogen atmosphere and an anhydrous system, and prepared at a temperature of -10°C to -78°C .

[0081]

(Synthesis method 4)

[0082]



(In the formulae, R , R^4 , R^5 , A , D , Y , Z , m and n represent the same meaning as those described above.)


A compound represented by Formula (8) is produced by oxidation by reacting an oxidizing agent with a compound represented by Formula (7). The above oxidation reaction may be performed without particular limitation as long as the reaction oxidizes a

secondary hydroxyl group. For example, an oxidation method such as a Jones oxidation, an ozone oxidation, or a Swern oxidation, or using an oxidizing reagent, such as manganese dioxide, or a Dess-Martin reagent, may be adopted.

[0083]

5 (Synthesis method 5)

[0084]



(In the formulae, R^4 , R^5 , A, D, X, Y, Z, m and n represent the same meaning as those described above. $R^{1'}$ and $R^{2'}$, among the above R^1 to R^3 , represent an


10 unsubstituted or substituted alkyl group, an unsubstituted or substituted alkenyl group, an unsubstituted or substituted alkynyl group, an unsubstituted or substituted aryl group, or an unsubstituted or substituted heterocyclic ring. Hal represents a halogen atom.)

A compound represented by Formula (10) may be produced by reacting an equivalent of a Grignard reagent with a compound represented by Formula (9). Also, a
 15 compound represented by Formula (11) may be produced by reacting at least two equivalents of a Grignard reagent with the compound represented by Formula (9).

[0085]

(Synthesis method 6)

[0086]



[0087]


(In the formulae, R^4 , R^5 , $R^{2'}$, A, D, X, Y, Z, m, n and Hal represent the same meaning as those described above. G represents a leaving group, such as an alkoxy group, or a halogen atom.)

A compound represented by Formula (13) may be produced by reacting one equivalent of a Grignard reagent with a compound represented by Formula (12). Also, a compound represented by Formula (14) may be produced by reacting at least two equivalents of a Grignard reagent with the compound represented by Formula (12).

10 [0088]

(Synthesis method 7)

[0089]




(In the formulae, R^4 , R^5 , A, D, X, Y, Z, m, and n represent the same meaning as those described above. J represents an alkoxycarbonyl group or a cyano group. K^1 and K^2 represent an alkyl group.)

A compound represented by Formula (16) may be produced by hydrolysis using conventional methods. Also, a compound represented by Formula (17) may be produced by reacting an alkylating agent in the presence of a base.

[0090]

5 (Synthesis method 8)

[0091]



(In the formulae, R, R^4 , R^5 , A, D, X, Y, Z, m, n and Hal represent the same meaning as those described above. $R^{4'}$ represents an unsubstituted or substituted alkoxy group, an unsubstituted or substituted alkyl group, an unsubstituted or substituted alkenyl group, an unsubstituted or substituted alkynyl group, an unsubstituted or substituted aryl group, or an unsubstituted or substituted heterocyclic ring, an unsubstituted or substituted amino group, or an unsubstituted or substituted thioalkyl group.)

An N-oxide compound represented by Formula (19) may be produced by


15 oxidizing a compound represented by Formula (18) using conventional methods such as

the use of an oxidizing agent. A compound represented by Formula (20) may be produced by reacting a conventional halogenating agent such as phosphorus oxychloride with the compound represented by Formula (19). And then, the compound represented by Formula (20) is subjected to a nucleophilic substitution reaction or a coupling reaction using an organic metal catalyst to synthesize a compound represented by Formula (21).

[0092]

(Synthesis method 9)

[0093]



(In the formulae, R , R^4 , R^5 , A , D , X , Y , Z , m , and n represent the same meaning as those described above. R' represents a C_{1-8} alkyl group. M represents an alkali metal such as lithium, sodium, or potassium. L represents a halogen atom.

The compound represented by Formula (I) may be prepared by a Suzuki coupling reaction of a boronic acid derivative represented by Formula (22) or Formula (23) and a halide derivative represented by Formula (24).

[0094]

The N-oxide compound may be prepared by a conventional oxidation reaction. For example, in solvents or in the absence of solvents, it may be prepared by causing

contact between the compound represented by Formula (I) and peroxide such as hydrogen peroxide.

[0095]

In any reaction, after the reaction is completed, the resultant is subjected to a normal post reaction operation in an organic synthetic chemistry, followed by subjecting to a conventional separation and purification, if needed, to efficiently isolate a desired product.

[0096]

The structure of the desired product may be identified and confirmed using ^1H -NMR spectrum, IR spectrum, or mass spectrum measurements, or by elemental analysis or the like.

[0097]

2) Agricultural fungicide

An agricultural fungicide according to the present invention contains, as an active ingredient thereof, at least one selected from the group consisting of the nitrogen-containing heterocyclic compound, the salt thereof, and the N- oxide compound thereof.

[0098]

The fungicide according to the present invention has an excellent fungicidal capacity over a wide variety of filamentous fungi, such as bacteria which belong to Oomycetes, Ascomycetes, Deuteromycetes, or Basidiomycetes.

[0099]

The fungicide according to the present invention may be used to prevent various diseases that occur in the cultivation of agricultural crops including flowers, turf, and grass through seed treatment, foliar spraying, soil application or submerged application.

[0100]

For example, it may be used to prevent:

sugar beet: Leaf spot (*Cercospora beticola*), Black root rot (*Aphanomyces cochlioides*), Root rot (*Thanatephorus cucumeris*), Leaf rot (*Thanatephorus cucumeris*);

5 peanut: Brown leaf spot (*Mycosphaerella arachidis*), Black rust (*Mycosphaerella berkeleyi*);

cucumber: Powdery mildew (*Sphaerotheca fuliginea*), Downy mildew (*Pseudoperonospora cubensis*), Gummy stem blight (*Mycosphaerella melonis*), Wilt (*Fusarium oxysporum*), Sclerotinia rot (*Sclerotinia sclerotiorum*), Gray mold (*Botrytis cinerea*), Anthracnose (*Colletotrichum orbiculare*), Scab (*Cladosporium cucumerinum*), Target leaf spot (*Corynespora cassicola*), Damping-off (*Pythium debaryanum*, *Rhizoctonia solani* Kuhn), Bacterial spot (*Pseudomonas syringae* pv. *Lecrymans*);

tomato: Gray mold (*Botrytis cinerea*), Leaf mold (*Cladosporium fulvum*), Phytophthora root rot (*Phytophthora infestans*);

15 eggplant: Gray mold (*Botrytis cinerea*), Black rot (*Corynespora melongenae*), Powdery mildew (*Erysiphe cichoracearum*), Sooty mold (*Mycovellosiella nattrassii*);

strawberry: Gray mold (*Botrytis cinerea*), Powdery mildew (*Sphaerotheca humuli*), Anthracnose (*Colletotrichum acutatum*, *Colletotrichum fragariae*), Phytophthora root rot (*Phytophthora cactorum*);

20 onion: Gray mold neck rot (*Botrytis allii*), Gray mold (*Botrytis cinerea*), Leaf blight (*Botrytis squamosa*), Downy mildew (*Peronospora destructor*);

cabbage: Clubroot (*Plasmodiophora brassicae*), Soft rot (*Erwinia carotovora*), Downy mildew (*Peronospora parasitica*);

kidney bean: Sclerotinia rot (*Sclerotinia sclerotiorum*), Gray mold (*Botrytis cinerea*);

25

apple: Powdery mildew (*Podosphaera leucotricha*), Scab (*Venturia inaequalis*),
 Monilia disease (*Monilinia mali*), Brooks fruit spot (*Mycosphaerella pomi*), Brown
 canker (*Valsa mali*), Alternaria blotch (*Alternaria mali*), Cedar apple rust
 (5 *Gymnosporangium yamadae*), White rot (*Botryosphaeria berengeriana*), Anthracnose
 (*Glomerella cingulata*, *Collectotrichum acutatum*), Marssonina blotch (*Diplocarpon mali*),
 Fly speck (*Zygophiala jamaicensis*), Sooty blotch (*Gloeodes pomigena*);

persimmon: Powdery mildew (*Phyllactinia kakicola*), Anthracnose
 (*Gloeosporium kaki*), Angular leaf spot (*Cercospora kaki*);

[0101]

10 peach: Brown rot (*Monilinia fructicola*), Scab (*Cladosporium carpophilum*),
 Phomopsis bacterial rot (*Phomopsis* sp.);

cherry: Brown rot (*Monilinia fructicola*);

grape: Gray mold (*Botrytis cinerea*), Powdery mildew (*Uncinula necator*), Ripe
 rot (*Glomerella cingulata*, *Colletotrichum acutatum*), Downy mildew (*Plasmopara*
 15 *viticola*), Bird's eye rot (*Elsinoe ampelina*), Leaf spot (*Pseudocercospora vitis*), Black rot
 (*Guignardia bidwellii*);

pear: Scab (*Venturia nashicola*), Cedar apple rust (*Gymnosporangium asiaticum*),
 Leaf spot (*Alternaria kikuchiana*), Black rot (*Botryosphaeria berenengeriana*), Powdery
 mildew (*Phyllactinia mali*);

20 tea: Collar rot (*Pestalotia theae*), Anthracnose (*Colletotrichum theae-sinensis*);

citrus: Scab (*Elsinoe fawcetti*), Blue mold (*Penicillium italicum*), Green mold
 (*Penicillium digitatum*), Gray mold (*Botrytis cinerea*), Melanose (*Diaporthe citri*),
 Bacterial canker (*Xanthomonas campestris* pv. *Citri*);

wheat: Powdery mildew (*Erysiphe graminis* f.sp. *tritici*), Petch (*Gibberella zeae*),
 25 Leaf rust (*Puccinia recondita*), Brown snow mold (*Pythium iwayamai*), Red snow mold

(*Monographella nivalis*), Black eye spot (*Pseudocercospora herpotrichoides*), Leaf blight (*Septoria tritici*), Glume blotch (*Leptosphaeria nodorum*), Typhula snow blight (*Typhula incarnata*), Sclerotinia snow blight (*Myriosclerotinia borealis*), Take-all (*Gaeumanomyces graminis*);

5 [0102]

barley: Leaf stripe (*Pyrenophora graminea*), Rhynchosporium scald (*Rhynchosporium secalis*), Loose smut (*Ustilago tritici*, *U.nuda*);

rice: Neck rot (*Pyricularia oryzae*), Sheath blight (*Rhizoctonia solani*), Bakanae disease (*Gibberella fujikuroi*), Helminthosporium blight (*Cochliobolus niyabeanus*),

10 Damping-off (*Pythium graminicolum*), Bacterial blight (*Xanthomonas oryzae*), Bacterial damping-off (*Burkholderia plantarii*), Brown stripe (*Acidovorax avenae*), Bacterial grain rot (*Burkholderia glumae*);

tobacco: Sclerotinia rot (*Sclerotinia sclerotiorum*), Powdery mildew (*Erysiphe cichoracearum*);

15 tulip: Gray mold (*Botrytis cinerea*);

bent grass: Sclerotinia snow blight (*Sclerotinia borealis*), Brown blight (*Pythium asphanidermatum*);

orchard grass: Powdery mildew (*Erysiphe graminis*);

soybean: Purple blotch (*Cercospora kikuchii*), Downy mildew (*Peronospora*

20 *Manshurica*), Stem rot (*Phytophthora sojae*);

potato · tomato: *Phytophthora* root rot (*phytophthora infestans*);

or the like.

[0103]

Also, the fungicide according to the present invention exhibits an excellent

25 fungicidal effect on resistant bacteria as well. Examples of the resistant bacteria

include: gray mold germs (*Botrytis cinerea*), sugar beet brown spot germs (*Cercospora beticola*), apple scab germs (*Venturia inaequalis*), pear scab germs (*Venturia nashicola*), resistant to a benzimidazole-based fungicide such as thiophanate-methyl, benomyl, or carbendazim; Gray mold germs (*Botrytis cinerea*) resistant to a dicarboximide-based
5 fungicide (such as vinclozolin, procymidone, or iprodione), and the like.

[0104]

Examples of diseases in which application of the fungicide according to the present invention is more preferable include Scab of apple, Gray mold of cucumber, Powdery mildew of wheat, Late blight of tomato, Leaf rust of wheat, Neck rot of rice,
10 Wilt disease of cucumber, and the like.

[0105]

Also, the fungicide according to the present invention is an agent with low phytotoxicity, low toxicity to fish or warm-blooded animals, and high safety.

The fungicide according to the present invention may be used in an
15 agrochemically acceptable form, that is, in a form of an agrochemical formulation, such as a water-dispersible powder, granules, a powder, an emulsion, a water-soluble powder, a suspension, or water-dispersible granules.

[0106]

Examples of an additive and a carrier used in the formulation of a solid include:
20 vegetable powder such as soybean flour or wheat flour; mineral fine powder such as diatomite, apatite, plaster, talc, bentonite, pyrophyllite, or clay; an organic or inorganic compound such as sodium benzoate, urea, or sulfate of soda; and the like.

[0107]

Examples of a solvent used in the formulation of a liquid include: an aromatic
25 hydrocarbon such as kerosene, xylene, or a petroleum-based aromatic hydrocarbon;

cyclohexane, cyclohexanone, dimethyl formamide, dimethyl sulfoxide, an alcohol, acetone, trichloroethylene, methylisobutylketone, mineral oil, vegetable oil, water, and the like.

[0108]

5 Also, in the formulations, in order to attain a uniform and stable form, a surfactant may be added as necessary.

 The surfactant which may be added is not particularly limited. Examples of the surfactant include: a non-ionic surfactant such as polyoxyethylene-added alkylphenylether, polyoxyethylene-added alkylether, polyoxyethylene-added higher fatty acid ester, polyoxyethylene-added sorbitan higher fatty acid ester, polyoxyethylene-added
10 tristyrylphenylether; a sulfuric ester salt of polyoxyethylene-added alkylphenylether, an alkylbenzene sulfonate, a sulfuric ester salt of higher alcohol, an alkylnaphthalene sulfonate, a polycarboxylate, a lignin sulfonate, a formaldehyde condensate of alkylnaphthalene sulfonate, an isobutylene-anhydrous maleic acid copolymer, and the
15 like.

[0109]

 The thus obtained water-dispersible powder, the emulsion, a flowable agent, the water-soluble powder, or the water-dispersible granules is diluted with water to a predetermined concentration and used as a solution, a suspension, or an emulsion to be
20 sprayed onto plants. Also, the powder and the granules are used sprayed onto plants as they are.

[0110]

 The amount of the fungicide according to the present invention is, normally, with respect to the total amount of the formulation, preferably 0.01 to 90% by weight,
25 and more preferably 0.05 to 85% by weight.

[0111]

The application rate of the fungicide according to the present invention, although it differs depending on weather conditions, formulation form, time of application, method of application, area of application, control target disease, or target
5 crop, is, normally as an amount of active ingredient compound per 1 hectare, 1 to 1,000 g, and preferably 10 to 100 g.

[0112]

When the water-dispersible powder, the emulsion, the suspension, the water-soluble powder, or the water-dispersible granules is diluted with water, the applied
10 concentration is 1 to 1,000 ppm, and preferably 10 to 250 ppm.

[0113]

The fungicide according to the present invention may be mixed with other fungicides or insecticides/acaricides, or synergists.

Representative examples of other fungicides, insecticides, acaricides, and plant
15 growth regulators that may be mixed and used with the fungicide according to the present invention are shown below.

Fungicide:

benzimidazole series such as benomyl, carbendazim, fuberidazole, thiabendazole, or thiophanate-methyl;
20 dicarboximide series such as chlozolate, iprodione, procymidone, or vinclozoline;

DMI-fungicide such as imazalil, oxpoconazole, perfluoate, prochloraz, triflumizole, trifenox, fenarimol, nuarimol, azaconazole, bitertanol, bromuconazole, cyproconazole, difenoconazole, diniconazole, epoxyconazole,
25 fenbuconazole, fluquinconazole, fulsilazole, flutriafol, hexaconazole, imibenconazole,

ipconazole, metconazole, myclobutanil, penconazole, propiconazole, prothioconazole, simeconazole, tebuconazole, tetraconazole, triadimefon, triadimenol, triticonazole, etaconazole, furconazole-cis, ipconazole, or imibenconazole;

phenylamide series such as benalaxyl, furalaxyl, metalaxyl, metalaxyl-M,

5 oxadixyl, or ofurace;

amine series such as aldimorph, dodemorph, fenpropimorph, tridemorph,

fenpropidine, piperalin, or spiroxamine;

phosphorothiolate series such as EDDP, iprobenfos, pyrazophos;

dithiolane series such as isoprothiolane;

10 carboxamide series such as benodanil, boscalid, carboxin, fenfuran, flutolanil, furametpyr, mepronil, oxycarboxin, penthiopyrad, or thifluzamide;

hydroxy-(2-amino)pyrimidine such as bupurimate, dimethirimol, or ethirimol;

[0114]

AP fungicide (anilinopyrimidine) such as cyprodinil, mepanipyrim, or

15 pyrimethanil;

N-phenylcarbamate such as diethofencarb; QoI-fungicide (Qo inhibitor) such as

azoxystrobin, picoxystrobin, pyraclostrobin, kresoxim-methyl, trifloxystrobin,

dimoxystrobin, metominostrobin, orysastrobin, famoxadone, fluoxastrobin, fenamidone,

metominofen, pyribencarb;

20 PP fungicide (phenylpyrrole) such as fenpiconyl, fludioxonil;

quinoline series such as quinoxifen;

AH fungicide (aromatic hydrocarbon) such as biphenyl, chloroneb, dicloran,

quintozene, tecnazene, or tolktfos-methyl;

MBI-R such as phthalide, pyroquilon, or tricyclazole;

25 MBI-D such as carpropamid, diclocymet, or fenoxanil;

- SBI agent such as fenhexamid, pyributicarb, or terbinafin;
 phenylurea such as pencycuron;
 QII-fungicide (QI inhibitor) such as cyazofamid;
 benzamide such as zoxamide;
 5 enopyranuron such as blasticidin, mildiomyacin;
 hexopyranosyl such as kasugamycin;
 glucopyranocyl such as streptomycin, validamycin;
 cyanoacetamide such as cymoxanil;
 carbamate such as propamocarb, prothiocarb, or polycarbamate;
 10 uncoupling agent such as binapacryl, dinocap, ferimzone, or fluazinam;
 organotin compound such as triphenyltin acetate, triphenyltin chloride, or
 triphenyltin hydroxide;
 [0115]
 organophosphate such as phosphoric acid, tolclofos-methyl, or fosetyl;
 15 phthalamic acid such as techlofthalam;
 benzotriazine such as triazoxide;
 benzenesulfonamide such as flusulfamide;
 pyridazinone such as diclomezin;
 CAA fungicide (carboxylic acid amid) such as dimethomorph, flumorph,
 20 benthiavalicarb, iprovalicarb, or mandipropamid;
 tetracycline such as oxytetracycline;
 thiocarbamate such as methasulfocarb; or
 other compounds such as etridiazole, polyoxin, oxolinic acid, hydroxyisoxazol,
 octhinoline, silthiofam, diflumetorim, acibenzolar-S-methyl, probenazole, tiadinil,
 25 ethaboxam, cyflufenamid, proquinazid, metrafenone, fluopicolide, copper(II) hydroxide,

organocopper, sulfur, ferbam, manzeb, maneb, metiram, propineb, thiuram, zineb, ziram, captan, captafol, folpet, chlorothalonil, dichlofluanid, tolylfluanid, dodine, guazatine, iminoctadine, anilazine, dithianon, chloropicrin, dazomet, metam-sodium, chinomethionat, cyprofuram, silthiofam, agro-bacterium, or fluoroimide.

5 [0116]

Insecticides · acaricides:

Organophosphorous and carbamate series insecticides:

fenthion, fenitrothion, diazinon, chlorpyrifos, ESP, vamidothion, phenthoate, dimethoate, formothion, malathon, trichlorhon, thiometon, phosmet, dichlorvos, acephate, 10 EPBP, methyl parathion, oxydemeton-methyl, ethion, salithion, cyanophos, isoxathion, pyridaphenthion, phosalone, methidathion, sulprofos, chlorfenvinphos, tetrachlorvinphos, dimethylvinhos, propaphos, isofenphos, ethylthiometon, profenofos, pyraclofos, monocrotophos, Azinphos-methyl, aldicarb, methomyl, thiodicarb, carbofuran, carbosulfan, benfuracarb, furathiocarb, propoxur, BPMC, MTMC, MIPC, carbaryl, 15 pirimicarb, ethiofencarb, fenoxycarb, EDDP or the like.

Pyrethroid series insecticides:

permethrin, cypermethrin, Deltamethrine, fenvalerate, fenpropathrin, pyrethrin, allethrin, tetramethrine, resmethrin, dimethrin, propathrin, phenothrin, prothrin, fluvalinate, cyfluthrin, cyhalothrin, flucythrinate, ethofenprox, cycroprothrin, 20 tralomethrin, silafluofen, brofenprox, acrinathrin or the like.

Other insecticides of benzoyl urea series:

microbial agrochemicals such as diflubenzuron, chlorfluazuron, hexaflumuron, triflumuron, tetrabenzuron, flufenoxuron, flucycloxuron, buprofezin, pyriproxyfen, methoprene, benzoepin, diafenthiuron, acetamiprid, imidacloprid, nitenpyram, fipronil, 25 cartap, thiocyclam, bensultap, nicotine sulfate, rotenone, metaldehyde, machine oil, BT,

insect pathogenic viruses, or the like.

[0117]

Nematocides:

Fenamiphos, fosthiazate or the like.

5 Acaricides:

chlorobenzilate, phenisobromolate, dicofol, amitraz, BPPS, benzomate, hexythiazox, fenbutatin oxide, polynactins, chinomethionat, CPCBS, tetradifon, avermectin, milbemectin, clofentezine, cyhexatin, pyridaben, fenpyroximate, tebufenpyrad, pyrimidifen, phenothiocarb, dienochlor or the like.

10 Plant growth regulator:

abscisic acid, indolebutyric acid, uniconazole, ethylchlozate, ethephon, cloxyfonac, chlormequat, chlorella extract, calcium peroxide, cyanamide, dichlorprop, gibberellin, daminozide, decyl alcohol, trinexapac-ethyl, mepiquat chloride, paclobutrazol, paraffin, wax, piperonyl butoxide, pyraflufen-ethyl, flurprimidol, prohydrojasmon, prohexadione-calcium, benzylaminopurine, pendimethalin, forchlorfenuron, maleic hydrazide potassium, 1-naphthylacetamide, 4-CPA, MCPB, corrin, oxyquinoline sulfate, ethychlozate, butralin, 1-methylcyclopropene, aviglycine hydrochloride.

[Examples]

[0118]


20 Hereinafter, the present invention is described in more detail with reference to examples, however, the present invention is not in the least limited by the following examples.

[0119]

Example 1

25 Synthesis of 1-[2-(quinolin-3-yloxy)-phenyl]-ethanone

[0120]



[0121]

0.73 g of 3-quinolinol, 1.8 g of cesium carbonate, 0.18 g of dipivaloylmethane,
 5 2.0 g of 2-bromoacetophenone, and 0.50 g of copper(I) chloride were dissolved in 10 ml
 of N-methylpyrrolidone, and the mixture was stirred for 3 hours at 130°C. The resultant
 was purified by silica gel column chromatography, and 0.82 g of
 1-[2-(quinolin-3-yloxy)-phenyl]-ethanone (Compound Number 1) was obtained.

[0122]


10 Compounds represented by Compound Numbers 2 to 13 were synthesized in the
 same manner as that of Example 1.

[0123]

Example 2

Synthesis of 3-(2-t-butyl-phenoxy)-quinoline

15 [0124]



[0125]

0.60 g of 2-tert-butylphenol, 1.3 g of cesium carbonate, 0.07 g of
 dipivaloylmethane, 0.42 g of 3-bromoquinoline, and 0.20 g of copper(I) chloride were

dissolved in 4 ml of N-methylpyrrolidone, and the mixture was stirred for one day at 130°C. The resultant was purified by silica gel column chromatography, and 0.04 g of 3-(2-t-butyl-phenoxy)-quinoline (Compound Number 14) was obtained.

[0126]


5 Compounds represented by Compound Numbers 15 to 19 were synthesized in the same manner as that of Example 2.

[0127]

Example 3

Synthesis of 2-chloro-6-(quinolin-3-yloxy)-benzamide

10 [0128]



[0129]

1.18 g of 2-chloro-6-(quinolin-3-yloxy)-benzonitrile was dissolved in 5 ml of 80% sulfuric acid and the mixture was stirred for 3 hours at 100°C. After being
15 neutralized with an aqueous solution of sodium hydroxide, the liquid was separated with ethyl acetate, the organic layer was concentrated and 1.03 g of 2-chloro-6-(quinolin-3-yloxy)-benzamide (Compound Number 20) was obtained.

[0130]


20 A compound represented by Compound Number 21 was synthesized in the same manner as that of Example 3.

[0131]

Example 4

Synthesis of 2-bromo-N,N-dimethyl-6-(quinolin-3-yloxy)-benzamide

[0132]



[0133]


- 5 After 0.34 g of 2-bromo-6-(quinolin-3-yloxy)-benzamide was dissolved in 2 ml of N-methylpyrrolidone, 47 mg of sodium hydride was added thereto at room temperature. After 0.21 g of methyl iodide was added to the reaction solution, the mixture was heated to 100°C and stirred for 4 hours. After the liquid was separated with ethyl acetate, the solvent was evaporated and the resultant was purified by silica gel
- 10 column chromatography to obtain 0.11 g of 2-bromo-N,N-dimethyl-6-(quinolin-3-yloxy)-benzamide (Compound Number 22).

[0134]

Example 5

Synthesis of 1-[2-(quinolin-3-yloxy)-phenyl]-ethanone oxime

- 15 [0135]



[0136]

- 0.46 g of 1-[2-(quinolin-3-yloxy)-phenyl]-ethanone and 0.15 g of hydroxylamine hydrochloride were dissolved in 3 ml of pyridine and then the mixture

was stirred for 24 hours at room temperature. After the reaction mixture was treated with dilute hydrochloric acid and separated with ethyl acetate, the organic layer was concentrated to obtain 0.54 g of 1-[2-(quinolin-3-yloxy)-phenyl]-ethanone oxime (Compound Number 23).

5 [0137]


Compounds represented by Compound Numbers 24 to 26 were synthesized in the same manner as that of Example 5.

[0138]

Example 6

10 Synthesis of 3-[2-(1,1-dimethoxy-ethyl)-phenoxy]-quinoline

[0139]



[0140]

After 0.13 g of 1-[2-(quinolin-3-yloxy)-phenyl]-ethanone and 0.06 g of trimethyl orthoformate were dissolved in 1.5 ml of methanol, 0.02 g of tetrabutylammonium tribromide was added and the mixture was stirred for 4 days at room temperature. The reaction mixture was concentrated and then purified by silica gel column chromatography to obtain 0.08 g of 3-[2-(1,1-dimethoxy-ethyl)-phenoxy]-quinoline (Compound Number 27).


20 [0141]

Example 7

Synthesis of

3-[2-(2,2,2-trifluoro-1-methyl-1-trimethylsilanyloxy-ethyl)-phenoxy]-quinoline

[0142]



[0143]

5 After 0.26 g of 1-[2-(quinolin-3-yloxy)-phenyl]-ethanone and 0.16 g of (trifluoromethyl)trimethylsilane were dissolved in 3 ml of tetrahydrofuran, 2 drops of tetrabutylammonium fluoride (1.0 M tetrahydrofuran solution) was added under ice-cooling and then stirred for 4 hours. After the reaction mixture was concentrated and purified by silica gel column chromatography to obtain 0.34 g of


10 3-[2-(2,2,2-trifluoro-1-methyl-1-trimethylsilanyloxy-ethyl)-phenoxy]-quinoline (Compound Number 28).

[0144]

Example 8

Synthesis of 1,1,1-trifluoro-2-[2-(quinolin-3-yloxy)-phenyl]-propan-2-ol

15 [0145]



[0146]

After 0.17 g of 3-[2-(2,2,2-trifluoro-1-methyl-1-trimethylsilanyloxy-ethyl)-phenoxy]-quinoline

was dissolved in 1.5 ml of tetrahydrofuran, 1 ml of tetrabutylammonium fluoride (1.0 M tetrahydrofuran solution) was added and the mixture was stirred for 16 hours. The reaction mixture was concentrated and then purified by silica gel column chromatography to obtain 0.11 g of

5 1,1,1-trifluoro-2-[2-(quinolin-3-yloxy)-phenyl]-propan-2-ol (Compound Number 29).


[0147]

Example 9

Synthesis of

2-[2-fluoro-6-(8-fluoro-quinolin-3-yloxy)-phenyl]-3,3-dimethyl-butan-2-ol

10 [0148]



[0149]

After 0.15 g of 1-[2-fluoro-6-(8-fluoro-quinolin-3-yloxy)-phenyl]-ethanone was dissolved in 1.5 ml of tetrahydrofuran, and then the solution was cooled to -78°C,

15 0.3 ml of t-butylmagnesium chloride (2.0 M diethyl ether solution) was added dropwise. After the reaction temperature was increased to room temperature, the reaction solution was treated with dilute hydrochloric acid, and then the liquid was separated with ethyl acetate. The organic layer was concentrated and then purified by silica gel column chromatography to obtain 0.04 g of

20 2-[2-fluoro-6-(8-fluoro-quinolin-3-yloxy)-phenyl]-3,3-dimethylbutan-2-ol (Compound Number 30).

[0150]


Compounds represented by Compound Numbers 31 to 35 were synthesized in the same manner as that of Example 9.

[0151]

Example 10

Synthesis of 2-[2-fluoro-6-(8-fluoroquinolin-3-yloxy)phenyl]propan-2-ol and 2-[2-fluoro-6-(8-fluoro-2-methylquinolin-3-yloxy)phenyl]propan-2-ol

[0152]



10

[0153]

After 0.3 g of 2-fluoro-6-(8-fluoroquinolin-3-yloxy)-benzoic acid ethyl ester was dissolved in 4.5 ml of tetrahydrofuran, the solution was cooled to -78°C , and 0.77 ml of methylmagnesium chloride (3.0 M diethyl ether solution) was added dropwise. After the reaction temperature was increased to room temperature, the reaction solution was treated with dilute hydrochloric acid and the liquid was separated with ethyl acetate.

15

The organic layer was concentrated and then purified by silica gel column chromatography to obtain 0.12 g of

2-[2-fluoro-6-(8-fluoroquinolin-3-yloxy)phenyl]propan-2-ol (Compound Number 36)

20

and 0.05 g of 2-[2-fluoro-6-(8-fluoro-2-methylquinolin-3-yloxy)phenyl]propan-2-ol (Compound Number 37).

[0154]


Compounds represented by Compound Numbers 38 to 49 were synthesized in the same manner as that of Example 10.

[0155]

Example 11

5 Synthesis of 7-(quinolin-3-ylamino)-indan-1-one

[0156]



[0157]

0.43 g of 3-bromoquinoline, 0.29 g of 7-aminoindan-1-one, 0.08 g of copper(I)
10 iodide, 0.41 g of potassium carbonate, and 4 ml of N-methylpyrrolidone were mixed, and then the mixture was stirred for 2 hours at 200°C. The resultant mixture was purified by silica gel column chromatography to obtain 0.15 g of 7-(quinolin-3-ylamino)-indan-1-one (Compound Number 50).

[0158]


15 A compound represented by Compound Number 51 was synthesized in the same manner as that of Example 11.

[0159]

Example 12

Synthesis of 3-(t-butyl-benzyl)-8-fluoroquinoline

20 [0160]



[0161]

After 0.69 g of triol salt derived from 8-fluoroquinoline-3-boric acid and 1,1,1-tris(hydroxymethyl)ethane by a method reported in the literature (Angew. Chem. Int. Ed., 2008, 47, 928-931) and 0.45 g of 1-bromomethyl-2-*t*-butylbenzene were dissolved in 10 ml of toluene, 0.46 g of tetrakis triphenylphosphine palladium was added and the mixture was stirred for 3 hours. After the solvent of the reaction mixture was evaporated, the resultant was purified by silica gel column chromatography to obtain 0.14 g of 3-(*t*-butyl-benzyl)-8-fluoroquinoline (Compound Number 52).

10 [0162]


A compound represented by Compound Number 53 was synthesized in the same manner as that of Example 12.

[0163]

Example 13

15 Synthesis of (2-*t*-butyl-phenyl)-(8-fluoro-quinolin-3-yl)-methanol

[0164]



[0165]

After 3 ml of tetrahydrofuran was cooled to -10°C, 0.91 ml of n-butyllithium (2.6 M hexane solution) and 1.35 ml of n-butylmagnesium chloride (0.91 M tetrahydrofuran solution) were added. After 30 minutes of stirring, 0.98 g of 8-fluoro-3-iodoquinoline was added and the mixture was stirred for 15 minutes. 0.49 g of 2-t-butylbenzaldehyde was added to the reaction solution and the mixture was stirred for 3 hours. After dilute hydrochloric acid was added to the reaction mixture, the liquid was separated with ethyl acetate. After the organic layer was concentrated and purified by silica gel column chromatography, 0.55 g of (2-t-butyl-phenyl)-(8-fluoro-quinolin-3-yl)-methanol (Compound Number 54) was obtained.

[0166]


Compounds represented by Compound Numbers 55 to 56 were synthesized in the same manner as that of Example 13.

[0167]

Example 14

Synthesis of (2-t-butyl-phenyl)-(8-fluoro-quinolin-3-yl)-methanone

[0168]



[0169]

After 0.32 g of (2-t-butyl-phenyl)-(8-fluoro-quinolin-3-yl)-methanol was dissolved in 3 ml of dichloromethane, 0.51 g of Dess-Martin reagent was added and the

mixture was stirred for 1 hour at room temperature. After water was added and then the liquid was separated with ethyl acetate, the organic layer was concentrated and purified by silica gel column chromatography to obtain 0.21 g of (2-*t*-butyl-phenyl)-(8-fluoro-quinolin-3-yl)-methanone (Compound Number 57).

5 [0170]

Compounds represented by Compound Numbers 58 to 59 were synthesized in the same manner as that of Example 14.

[0171]

Example 15

10 Synthesis of 2-chloro-3-(1,1-dimethylindan-7-yloxy) quinoline

0.2 g of 3-(1,1-dimethylindan-7-yloxy) quinolone was dissolved in 5 mL of chloroform, and 0.2 g of *m*-CPBA (70%) was added and stirred overnight at room temperature. After the reaction mixture was diluted with chloroform and washed with saturated sodium bicarbonate water, magnesium sulfate was added to dry the resultant.

15 The solvent was evaporated under reduced pressure to obtain 0.2 g of 3-(1,1-dimethylindan-7-yloxy) quinoline-N-oxide.

The resultant was dissolved in phosphorus oxychloride and heated under reflux for 4 hours. Phosphorus oxychloride was evaporated under reduced pressure and the residue was dissolved in ethyl acetate and was washed with saturated sodium bicarbonate water. The organic layer was dried with magnesium sulfate and evaporated under reduced pressure. The crude product obtained was purified by silica gel column chromatography, and 0.13 g of 2-chloro-3-(1,1-dimethylindan-7-yloxy) quinoline (Compound Number 60) was obtained.

[0172]

25 A compound represented by Compound Number 61 was synthesized in the same

manner as that of Example 15.

[0173]

Example 16

Synthesis of 3-(2-cyanophenoxy)quinoline

5 After 0.27 g of 3-hydroxyquinoline was dissolved in N-methyl-2-pyrrolidone, 0.087 g of sodium hydride (60% oil dispersion) was added under ice-cooling, and the mixture was stirred for 30 minutes. At the same temperature, 0.27 g of 2-fluorobenzonitrile was added and then the mixture was stirred for 6 hours at room temperature. After the reaction mixture was poured into ice water, extracted with ethyl
10 acetate, and washed with saturated saline solution, the resultant was dried with magnesium sulfate. After the solvent was evaporated under reduced pressure, the crude product was purified by silica gel column chromatography to obtain 0.37 g of 3-(2-cyanophenoxy)quinoline (Compound Number 62).

[0174]

15 Compounds represented by Compound Numbers 63 to 66 were synthesized in the same manner as that of Example 16.

[0175]


Example 17

Synthesis of difluoro-[2-fluoro-6-(8-fluoroquinolin-3-yloxy)-ethyl acetate

20 [0176]

Step 1) Synthesis of 8-fluoro-3-(3-fluoro-2-iodophenoxy)-quinoline

[0177]



[0178]


16 ml of N-methylpyrrolidone was added to 1.06 g of 8-fluoro-3-hydroxyquinoline, 1.94 g of 2,6-difluoroiodobenzene, and 1.7 g of potassium carbonate, and the mixture was stirred for 24 hours at 130°C. The reaction solution was purified by silica gel column chromatography, and 0.55 g of 8-fluoro-3-(3-fluoro-2-iodophenoxy)-quinoline was obtained.

[0179]

Step 2) Synthesis of difluoro-[2-fluoro-6-(8-fluoroquinolin-3-yloxy)-ethyl acetate

10 acetate

[0180]



[0181]

After 0.4 g of 8-fluoro-3-(3-fluoro-2-iodophenoxy)-quinoline was dissolved in 4 ml of dimethylsulfoxide, 0.25 g of copper (powder) and 0.41 g of bromodifluoroethyl acetate were added thereto, and then the mixture was stirred for 24 hours at 40°C. The reaction solution was purified by silica gel column chromatography, and 0.27 g of difluoro-[2-fluoro-6-(8-fluoroquinolin-3-yloxy)-ethyl acetate (Compound Number 292)

15

was obtained.


[0182]

Example 18

Synthesis of 1,1-difluoro-1-[2-fluoro-6-(8-fluoroquinolin-3-yloxy)-

5 phenyl]-2-methyl-propan-2-ol

[0183]



[0184]

0.18 g of difluoro-[2-fluoro-6-(8-fluoroquinolin-3-yloxy)-ethyl acetate was
 10 dissolved in 3 ml of tetrahydrofuran and 0.4 ml of methylmagnesium chloride (3.0 M
 tetrahydrofuran solution) was added dropwise at 0°C. After the mixture was stirred for
 2 hours at 0°C, dilute hydrochloric acid was added, and the liquid was separated with
 ethyl acetate. The organic layer was concentrated and purified by silica gel column
 chromatography to obtain 0.10 g of
 15 1,1-difluoro-1-[2-fluoro-6-(8-fluoroquinolin-3-yloxy)-phenyl]-2-methyl-propan-2-ol
 (Compound Number 293).


[0185]

Example 19

Synthesis of 1-[2-(7,8-difluoroquinolin-3-yloxy)-6-fluoro-phenyl]ethanone

20

[0186]



[0187]


After 0.91 g of 7,8-difluoro-3-hydroxyquinoline and 0.94 g of 2,6-difluoroacetophenone were dissolved in 10 ml of dimethylformamide, 0.9 g of potassium carbonate was added. The reaction solution was heated to 100°C and the mixture was stirred for 4.5 hours. Dilute hydrochloric acid was added to the reaction solution and the solution was separated by ethyl acetate. The organic layer was concentrated and purified by silica gel column chromatography to obtain 0.64 g of 1-[2-(7,8-difluoroquinolin-3-yloxy)-6-fluoro-phenyl]-ethanone (Compound Number 391).

[0188]

Example 20

Synthesis of 2-[2-(7,8-difluoroquinolin-3-yloxy)-6-fluoro-phenyl]-propan-2-ol

[0189]



15

[0190]

After 0.64 g of 1-[2-(7,8-difluoroquinolin-3-yloxy)-6-fluoro-phenyl]-ethanone was dissolved in 10 ml of tetrahydrofuran, the resultant was cooled to 0°C and 1.5 ml of

methylmagnesium chloride (3.0 M tetrahydrofuran solution) was added dropwise.

Dilute hydrochloric acid was added to the reaction solution, and the liquid was separated with ethyl acetate. The organic layer was concentrated and purified by silica gel column chromatography to obtain 0.49 g of

- 5 2-[2-(7,8-difluoroquinolin-3-yloxy)-6-fluoro-phenyl]-propan-2-ol (Compound Number 124).


[0191]

Example 21

Synthesis of

- 10 3-[2-(7,8-difluoroquinolin-3-yloxy)-6-fluoro-phenyl]-3-methyl-butan-2-one

[0192]



[0193]

3 ml of N-methylpyrrolidone was added to 0.78 g of

- 15 7,8-difluoro-3-iodoquinoline, 0.26 g of 3-(2-fluoro-6-hydroxyphenyl)-3-methyl-butan-2-one, 1.05 g of cesium carbonate, 53 mg of dipivaloylmethane, and 0.27 g of copper(I) chloride. After 48 hours of stirring at 130°C, the reaction solution was filtered with CELITE and the liquid was separated with ethyl acetate. The organic layer was concentrated and purified by silica gel column chromatography to obtain 0.05 g of
- 20 3-[2-(7,8-difluoroquinolin-3-yloxy)-6-fluoro-phenyl]-3-methyl-butan-2-one


(Compound Number 267).

[0194]

Example 22

Synthesis of 2-fluoro-6-(8-fluoroquinolin-3-yloxy)-benzaldehyde

5 [0195]



[0196]

After 30 ml of acetonitrile was added to 3.0 g of 8-fluoro-3-hydroxyquinoline, 3.1 g of potassium carbonate and 3.5 g of 2,6-difluoroacetophenone were added. After
 10 the reaction solution was heated under reflux for 3 hours, the reaction solution was filtered with CELITE. After the filtrate was extracted with ethyl acetate, the organic layer was concentrated and purified by silica gel column chromatography to obtain 3.0 g of 2-fluoro-6-(8-fluoroquinolin-3-yloxy)-benzaldehyde (Compound Number 341).

[0197]


15 Example 23

Synthesis of 1-[2-fluoro-6-(8-fluoroquinolin-3-yloxy)-phenyl]-propane-1,2-dione

[0198]

Step 1) Synthesis of [2-fluoro-6-(8-fluoroquinolin-3-yloxy)-phenyl]-
 20 trimethylsilyloxy-acetonitrile

[0199]




[0200]

After 3.2 g of 2-fluoro-6-(8-fluoroquinolin-3-yloxy)-benzaldehyde was dissolved in 70 ml of dichloromethane and the resultant was cooled to 0°C, 1.3 g of titanium tetraisopropoxide was added. After the reaction solution was warmed to room temperature, 4.5 g of trimethylsilyl cyanide was added thereto, and the mixture was stirred for 2 hours. Dilute hydrochloric acid was added to the reaction solution and the solution was extracted with dichloromethane. The organic layer was concentrated and purified by silica gel column chromatography to obtain 3.1 g of [2-fluoro-6-(8-fluoroquinolin-3-yloxy)-phenyl]-trimethylsilyloxy-acetonitrile (approximately 90% purity).

[0201]

Step 2) Synthesis of 1-[2-fluoro-6-(8-fluoroquinolin-3-yloxy)-phenyl]-1-hydroxy-propan-2-one

15 [0202]



[0203]


3.3 g of [2-fluoro-6-(8-fluoroquinolin-3-yloxy)-phenyl]-trimethylsilyloxy-

acetonitrile (approximately 90% purity) was dissolved in 50 ml of diethyl ether and 5.7 ml of methylmagnesium bromide (3.0 M diethyl ether solution) was added dropwise at room temperature. After the mixture was stirred for 1 hour at room temperature, the resultant was heated under reflux for 3 hours, and then treated with dilute hydrochloric acid. The resultant was subjected to extraction with ethyl acetate, and then the organic layer was concentrated to obtain 3.9 g of crude product. 20 ml of 2N hydrochloric acid and 10 ml of tetrahydrofuran were added to 1.9 g of the crude product and the mixture was stirred for 4 hours at room temperature. The reaction solution was extracted with ethyl acetate and then purified by silica gel column chromatography to obtain 0.4 g of 1-[2-fluoro-6-(8-fluoroquinolin-3-yloxy)-phenyl]-1-hydroxy-propan-2-one.

[0204]

Step 3) Synthesis of 1-[2-fluoro-6-(8-fluoroquinolin-3-yloxy)-phenyl]-propane-1,2-dione

[0205]



[0206]

After 0.14 g of 1-[2-fluoro-6-(8-fluoroquinolin-3-yloxy)-phenyl]-1-hydroxy-propan-2-one was dissolved in 10 ml of dichloromethane, 0.91 g of Dess-Martin reagent was added thereto at 0°C. After 3 hours of stirring, the reaction solution was concentrated and purified by silica gel column chromatography to obtain 0.08 g of 1-[2-fluoro-6-(8-fluoroquinolin-3-yloxy)-phenyl]-propane-1,2-dione (Compound

Number 396).

[0207]

Example 24

Synthesis of 1-[2-fluoro-6-(8-fluoroquinolin-3-yloxy)-phenyl]-2-hydroxy-


5 2-methyl-propan-1-one

[0208]

Step 1) Synthesis of 1-[2-fluoro-6-(8-fluoroquinolin-3-yloxy)-phenyl]-2-

hydroxy-2-methyl-propane-1,2-diol

[0209]



10

[0210]

3.3 g of [2-fluoro-6-(8-fluoroquinolin-3-yloxy)-phenyl]-trimethylsilyloxy-

acetonitrile was dissolved in 50 ml of diethyl ether and 5.7 ml of methylmagnesium

bromide (3.0 M diethyl ether solution) was added dropwise at room temperature. After

15 the mixture was stirred for 1 hour at room temperature, the resultant was heated under

reflux for 3 hours, and then treated with dilute hydrochloric acid. The resultant was

subjected to extraction with ethyl acetate and then the organic layer was concentrated to

obtain 3.9 g of crude product. 1.9 g of the crude product was dissolved in 20 ml of

tetrahydrofuran and 3.2 ml of methylmagnesium bromide (3.0 M diethyl ether solution)

20 was added dropwise at 0°C. After 3 hours of stirring at 0°C, 20 ml of 2N hydrochloric


acid was added and the mixture was stirred for one day at room temperature. The

reaction solution was extracted with ethyl acetate and purified by silica gel column chromatography to obtain 0.25 g of 1-[2-fluoro-6-(8-fluoroquinolin-3-yloxy)-phenyl]-2-methyl-propane-1,2-diol as a crude product.

[0211]

- 5 Step 2) Synthesis of 1-[2-fluoro-6-(8-fluoroquinolin-3-yloxy)-phenyl]-2-hydroxy-2-methyl-propan-1-one

[0212]



[0213]


- 10 0.15 g of 1-[2-fluoro-6-(8-fluoroquinolin-3-yloxy)-phenyl]-2-methyl-propane-1,2-diol was dissolved in 10 ml of dichloromethane, and 0.20 g of Dess-Martin reagent was added thereto at 0°C. After 2 hours of stirring, the reaction solution was concentrated and purified by silica gel column chromatography to obtain 0.03 g of 1-[2-fluoro-6-(8-fluoroquinolin-3-yloxy)-phenyl]-2-hydroxy-2-methyl-propan-1-one (Compound Number 398).

[0214]

Example 25

Synthesis of 2-[2-fluoro-6-(7,8-difluoro-2-methylquinolin-3-yloxy)phenyl]propan-2-ol

- 20 [0215]



[0216]

26.3 g of 7,8-difluoro-3-hydroxy-2-methylquinoline was dissolved in 200 mL of dimethylformamide. 27.2 g of 2,6-difluoroacetophenone and 24.0 g of potassium carbonate were added to the solution and the mixture was stirred for 4 hours at 100°C. After the thus obtained reaction solution was poured into ice water, and neutralized with dilute hydrochloric acid, the liquid was separated with ethyl acetate. The organic layer was concentrated and purified by silica gel column chromatography to obtain 17.0 g of 1-[2-fluoro-6-(7,8-difluoro-2-methylquinolin-3-yloxy)-phenyl]-ethanone.

After 11.0 g of 1-[2-fluoro-6-(7,8-difluoro-2-methylquinolin-3-yloxy)-phenyl]-ethanone obtained was dissolved in 80 ml of tetrahydrofuran and cooled to 0°C, 16.6 ml of methylmagnesium chloride (3.0 M tetrahydrofuran solution) was added dropwise. After the resultant was warmed to room temperature and stirred for 2 hours, the reaction solution was treated with dilute hydrochloric acid and the liquid was separated with ethyl acetate. The organic layer was concentrated and purified by silica gel column chromatography to obtain 10.0 g of 2-[2-fluoro-6-(7,8-difluoro-2-methylquinolin-3-yloxy)phenyl]propan-2-ol


(Compound Number 125) was obtained.

[0217]

Example 26

Synthesis of 2-[2-fluoro-6-(7,8-difluoroquinolin-3-yloxy)phenyl]propan-2-ol

[0218]



[0219]

0.5 g of 7,8-difluoro-3-hydroxyquinoline was dissolved in 10 mL of
 5 dimethylformamide. 0.52 g of 2,6-difluoroacetophenone and 0.46 g of potassium
 carbonate were added to the solution, and the mixture was stirred for 2.5 hours at 100°C.
 The thus obtained reaction solution was poured into ice water and then neutralized with
 dilute hydrochloric acid, the liquid was separated with ethyl acetate. The organic layer
 was concentrated and purified by silica gel column chromatography to obtain 0.38 g of
 10 1-[2-fluoro-6-(7,8-difluoroquinolin-3-yloxy)-phenyl]-ethanone (Compound Number
 391).

After 0.38 g of 1-[2-fluoro-6-(7,8-difluoroquinolin-3-yloxy)-phenyl]-ethanone
 obtained was dissolved in 5 ml of tetrahydrofuran and cooled to 0°C, 1.4 ml of
 methylmagnesium bromide (3.0 M diethyl ether solution) was added dropwise. After
 15 the resultant was warmed to room temperature and stirred for 2 hours, the reaction
 solution was treated with dilute hydrochloric acid and the liquid was separated with ethyl
 acetate. The organic layer was concentrated and purified by silica gel column
 chromatography to obtain 0.33 g of 2-[2-fluoro-6-(7,8-difluoroquinolin-3-yloxy)phenyl]
 propan-2-ol (Compound Number 124).

20 [0220]

Hereinafter, the intermediate product according to the present invention is
 described in more detail with reference to examples, however, the intermediate product

according to the present invention is not in the least limited by the following examples.

[0221]

Example 27


Synthesis of 7,8-difluoro-3-hydroxy-2-methylquinoline

5

[0222]

Step 1) Synthesis of 6,7-difluoroisatin

[0223]



[0224]

10

After 15.7 g of 2,3-difluoroaniline was added to 825 mL of water, and then 24.2 g of trichloroacetaldehyde 1-hydrate, 30.8 g of hydroxylamine hydrochloride, 138.6 g of anhydrous sodium sulfate were added thereto, the mixture was stirred for 10 hours at 50°C. After the resultant was cooled to room temperature, 44 mL of 2N hydrochloric acid was added, and then stirred for 30 minutes, followed by collecting the crystal by filtration. After the thus obtained crystals were dried, and then added to concentrated sulfuric acid heated to 70°C, the mixture was stirred for 1 hour at 80 to 90°C. The reaction solution was poured into ice, followed by subjecting to extraction with ethyl acetate, and then washing with saturated saline solution. After the organic layer was dried with magnesium sulfate, the solvent was evaporated under reduced pressure to obtain 26 g of a crude product of 6,7-difluoroisatin.


15

20

[0225]

Step 2) Synthesis of 7,8-difluoro-3-hydroxy-2-methylquinoline

[0226]



[0227]

After 41 g of the crude product of 6,7-difluoroisatin was added to 200 mL of water, 75.3 g of potassium hydroxide (6 equivalents) was added thereto under ice-cooling and the mixture was stirred for 30 minutes. 42 g of bromoacetophenone (1.4 equivalents) was added dropwise to the suspension while maintaining the internal temperature of the reaction solution at 20 to 25°C, and the mixture was further stirred overnight at room temperature. After the resultant was neutralized with concentrated hydrochloric acid, the precipitated crystals were collected by filtration and washed with a small amount of water. The thus obtained crystals were sufficiently dried, and then added little by little to 100 mL of nitrobenzene heated to 130 to 140°C, and the mixture was stirred further for 1 hour at 150°C. After the reaction solution was cooled to room temperature, the precipitated crystals were washed with chloroform to obtain 26.3 g of 7,8-difluoro-3-hydroxy-2-methylquinoline.

¹H NMR (300 MHz, DMSO-d₆) δ 2.57 (s, 3H), 7.4 to 7.7 (m, 3H), 10.60 (bs, 1H).

[0228]

The following compound was produced using the same method.

8-fluoro-3-hydroxy-2-methylquinoline

¹H NMR (300 MHz, DMSO-d₆) δ 2.56 (s, 3H), 7.2 to 7.6 (m, 4H), 10.53 (bs, 1H).

[0229]


Example 28

Synthesis of 7,8-difluoro-3-hydroxyquinoline

[0230]

Step 1) Synthesis of 7,8-difluoroquinoline

5 [0231]




[0232]

In a 3 L eggplant-shaped flask containing a stir bar, 80% sulfuric acid (607.7 g, 49.57 mol) was introduced and cooled to 0°C, and then 2,3-difluoroaniline (160.0 g, 1.24 mol) was slowly added. After the addition, the mixture was stirred for 1 hour at room temperature, sodium iodide (1.85 g, 12.3 mmol) was further added thereto, and then the mixture was heated in an oil bath until the temperature thereof reached 150°C (temperature of the bath). When it reached this temperature, glycerin (125.5 g, 1.36 mol) was added dropwise over a period of one hour, and the mixture was stirred for 1 hour at 150°C. The temperature of the bath was further elevated to 180°C, and the water was evaporated over a period of 2 hours using a distillation apparatus. After confirming that the starting material had disappeared, the mixture was neutralized using 10N aqueous sodium hydroxide while cooling with an ice-water bath (the inner temperature is 60 to 70°C). After the neutralization, before the inner temperature returned to room temperature, the resultant was subjected to extraction with ethyl acetate, and the extract was dried with magnesium sulfate, filtered, and concentrated. The thus obtained crude product was purified by silica gel column chromatography (normal

hexane: ethyl acetate) to obtain 185.5 g of 7,8-difluoroquinoline (91%) as a hazel solid.

[0233]

Step 2) Synthesis of 7,8-difluoro-3-iodoquinoline



[0234]

In a 3 L eggplant-shaped flask containing a stir bar, 7,8-difluoroquinoline (185.5 g, 1.12 mol), N-iodosuccinimide (505.4 g, 2.25 mol), and acetic acid (927 mL) were introduced and the mixture was stirred for 30 hours at 90°C. After cooling, the precipitated crystals were filtered and dried. On the other hand, the filtrate was concentrated under reduced pressure, the residual acetic acid was neutralized with sodium hydrogen carbonate, and the resultant was subjected to extraction with ethyl acetate. In addition, after the filtrate was dried with magnesium sulfate, filtered and concentrated, the thus obtained crude product was purified by silica gel column chromatography (normal hexane: ethyl acetate) to obtain 227.2 g (70%) of 7,8-difluoro-3-iodoquinoline as a hazel solid combined with the previously obtained crystals.

¹H-NMR (300 MHz, CDCl₃) δ 7.39 to 7.51 (m, 2H), 8.55 (m, 1H), 9.08 (d, 1H, J = 2.1 Hz).

[0235]

Step 3) Synthesis of 7,8-difluoro-3-hydroxyquinoline

[0236]



4

5

15

[0238]

The following compound was produced using the same method.

8-fluoro-3-hydroxyquinoline

20

[0239]


The nitrogen-containing heterocyclic compounds obtained from the above

examples are shown in Tables 1 to 19. In addition, compounds synthesized in the same manner as that of any of the above examples are further shown in Table 20 to Table 28.

Also, Et represents an ethyl group, ⁿPr represents an n-propyl group, ⁱPr represents an i-propyl group, ^cPr represents a cyclopropyl group, ⁿBu represents an n-butyl group, ^tBu represents a t-butyl group, Ph represents a phenyl group, Bn represents a benzyl group, and Tos represents toluene sulfonyl group. ¹H-NMR of the compounds in each table is also shown in Table 29 to Table 41.

[0240]

Table 1



Compound No.	(R ⁴) _m	R ¹	R ²	R ³	R ⁶	R ⁶	R ^d	E	-X-	Physical property
1	-	=O		-CH ₃	H	H	H	CH	-O-	75-76
2	-	=O		-H	H	H	H	CH	-O-	76-79
3	-	=O		-OEt	H	H	H	CH	-O-	amorphous
4	-	=O		-OEt	H	H	H	CF	-O-	72-74
5	-	=O		-OEt	H	H	F	CH	-O-	amorphous
6	-	=O		-OEt	H	F	H	CH	-O-	93-95
7	-	=O		-OEt	F	H	H	CH	-O-	amorphous
8	8-F	=O		-OEt	F	H	H	CH	-O-	amorphous
9	-	-F	-F	-F	H	H	H	CH	-O-	amorphous
10	8-F	=O		-CH ₃	F	H	H	CH	-O-	amorphous
11	-	=O		- ^t Bu	H	H	H	CH	-O-	amorphous
12	8-F	=O		- ^t Bu	F	H	H	CH	-O-	amorphous
13	-	-F	-F	-F	H	H	H	N	-O-	amorphous
14	-	-CH ₃	-CH ₃	-CH ₃	H	H	H	CH	-O-	amorphous
15	8-F	-CH ₃	-CH ₃	-CH ₃	H	H	H	CH	-O-	77-79
16	-	-CH ₃	-CH ₃	-(CH ₂) ₂ -			H	CH	-O-	90-94

[0241]

Table 2

Compound No.	(R ⁴) _m	R ¹	R ²	R ³	R ⁵	R ⁶	R ^d	E	-X-	Physical property
17	-	-CH ₃	-CH ₃	-Et	H	H	H	CH	-O-	amorphous
18	8-F	-CH ₃	-CH ₃	-Et	H	H	H	CH	-O-	amorphous
19	8-F	-CH ₃	-CH ₃	-(CH ₂) ₂ -		H	H	CH	-O-	104-106
20	-	=O		-NH ₂	Cl	H	H	CH	-O-	168-188
21	-	=O		-NH ₂	Br	H	H	CH	-O-	204-206
22	-	=O		-N(CH ₃) ₂	Br	H	H	CH	-O-	amorphous
23	-	=N-OH		-CH ₃	H	H	H	CH	-O-	125-126
24	-	=N-OCH ₃		-CH ₃	H	H	H	CH	-O-	amorphous
25	-	=N-O ^t Bu		-CH ₃	H	H	H	CH	-O-	amorphous
26	8-F	=N-OH		-CH ₃	F	H	H	CH	-O-	171-173
27	-	-OCH ₃	-CH ₃	-OCH ₃	H	H	H	CH	-O-	amorphous
28	-	-OSi(CH ₃) ₃	-CH ₃	-CF ₃	H	H	H	CH	-O-	amorphous
29	-	-OH	-CH ₃	-CF ₃	H	H	H	CH	-O-	158-160
30	8-F	-OH	-CH ₃	- ^t Bu	F	H	H	CH	-O-	amorphous
31	-	-OH	-CH ₃	- ^t Bu	H	H	H	CH	-O-	amorphous
32	-	-OH	-CH ₃	-Et	H	H	H	CH	-O-	129-131
33	8-F	-OH	-CH ₃	-Et	H	H	H	CH	-O-	122-124
34	8-F	-OH	-CH ₃	-(CH ₂) ₂ -		H	H	CH	-O-	amorphous
35	8-F	-OH	- ^t Bu	-(CH ₂) ₂ -		H	H	CH	-O-	amorphous
36	8-F	-OH	-CH ₃	-CH ₃	F	H	H	CH	-O-	amorphous
37	2-CH ₃ , 8-F	-OH	-CH ₃	-CH ₃	F	H	H	CH	-O-	amorphous
38	-	-OH	-CH ₃	-CH ₃	H	H	H	CH	-O-	amorphous
39	2- ^t Bu	=O		-H	H	H	H	CH	-O-	amorphous
40	-	-OH	-CH ₃	-CH ₃	H	H	H	CF	-O-	amorphous
41	-	-OH	-CH ₃	-CH ₃	H	H	F	CH	-O-	amorphous
42	-	-OH	-CH ₃	-CH ₃	H	F	H	CH	-O-	amorphous

[0242]

Table 3

Compound No.	(R ⁴) _m	R ¹	R ²	R ³	R ^b	R ^c	R ^d	E	X-	Physical property
43	-	-OH	-CH ₃	-CH ₃	F	H	H	OH	-O-	amorphous
44	8-F	-OH	-CH ₃	-CH ₃	H	H	H	OH	-O-	128-130
45	-	-OH	-CH ₃	-CH ₃	H	H	H	N	-O-	146-147
46	2-Et	-OH	-CH ₃	-CH ₃	H	H	H	OH	-O-	amorphous
47	2-nBu	-OH	-CH ₃	-CH ₃	H	H	H	OH	-O-	amorphous
48	8-F	-OH	-CH ₃	-Et	F	H	H	OH	-O-	amorphous
49	8-Cl	-OH	-CH ₃	-CH ₃	F	H	H	OH	-O-	amorphous
50	-	=O		-CH ₂ CH ₂ -		H	H	OH	-NH-	130-132
51	-	=O		-Ph	H	H	H	OH	-NH-	amorphous
52	8-F	-CH ₃	-CH ₃	-CH ₃	H	H	H	OH	-CH ₂ -	108-111
53	8-F	=O		-OEt	F	H	H	OH	-CH ₂ -	amorphous
54	8-F	-CH ₃	-CH ₃	-CH ₃	H	H	H	OH	-CH(OH)-	173-175
55	-	-CH ₃	-CH ₃	-CH ₂ -OS(CH ₃) ₂ -tBu	H	H	H	OH	-CH(OH)-	140-143
56	-	-CH ₃	-CH ₃	-C(CH ₃) ₂ -CH ₂ OH	H	H	H	OH	-CH(OH)-	136-138
57	8-F	-CH ₃	-CH ₃	-CH ₃	H	H	H	OH	-CO-	182-183
58	-	-CH ₃	-CH ₃	-CH ₂ -OS(CH ₃) ₂ -tBu	H	H	H	OH	-CO-	105-108
59	-	-CH ₃	-CH ₃	-CH ₂ O-Tos	H	H	H	OH	-CO-	amorphous
60	2-Cl	-CH ₃	-CH ₃	-CH ₂ CH ₂ -		H	H	OH	-O-	amorphous
61	1-Oxide, 8-F	-CH ₃	-CH ₃	-Et	H	H	H	OH	-O-	amorphous
62	-		≡ N		H	H	H	OH	-O-	92-95
63	-		≡ N		F	H	H	OH	-O-	149-151
64	-		≡ N		Br	H	H	OH	-O-	167-169
65	-		≡ N		Cl	H	H	OH	-O-	150-151
66	-		≡ N		OMe	H	H	OH	-O-	120-122

[0243]

Table 4

Compound No.	(R ⁴) _m	R ¹	R ²	R ³	R ^b	R ^c	R ^d	E	-X-	Physical property
67	2-CH ₃	-OH	-CH ₃	-CH ₃	H	H	H	CH	-O-	amorphous
68	2- ⁱ Pr	-OH	-CH ₃	-CH ₃	H	H	H	CH	-O-	amorphous
68	2-OCH ₃	-OH	-CH ₃	-CH ₃	H	H	H	CH	-O-	amorphous
70	2-OEt	-OH	-CH ₃	-CH ₃	H	H	H	CH	-O-	amorphous
71	1-Oxide	-OH	-CH ₃	-CH ₃	H	H	H	CH	-O-	154-155
72	-	-OH	-CH ₃	-C(CH ₃) ₂ OH	H	H	H	CH	-O-	amorphous
73	-	-OH	-CH ₃	-CO ₂ Et	H	H	H	CH	-O-	amorphous
74	8-F	-OH	-CH ₃	-Ph	F	H	H	CH	-O-	amorphous
75	8-F	-OH	-CH ₃	-CH ₂ Ph	F	H	H	CH	-O-	amorphous
76	8-F	-OH	-CH ₃	-CH=CH ₂	F	H	H	CH	-O-	amorphous
77	8-F	-OH	-CH ₃	-CH ₂ -CH=CH ₂	F	H	H	CH	-O-	amorphous
78	2-CH=CH ₂ , 8-F	-OH	-CH ₃	-CH=CH ₂	F	H	H	CH	-O-	amorphous
79	2-CH ₂ CH=CH ₂ , 8-F	-OH	-CH ₃	-CH ₂ -CH=CH ₂	F	H	H	CH	-O-	amorphous
80	8-F	-OH	-CH ₃	- ⁱ Pr	F	H	H	CH	-O-	amorphous
81	8-F	-OH	-CH ₃	-C(CH ₃)=CH ₂	F	H	H	CH	-O-	amorphous
82	8-F	-OH	-CH ₃	-CH ₂ -(4-FPh)	F	H	H	CH	-O-	amorphous
83	8-F	-OH	-CH ₃	-CH(CH ₃)-CH=CH ₂	F	H	H	CH	-O-	amorphous
84	8-F	-OH	-CH ₃	-CH ₃	F	H	H	CF	-O-	149-150
85	2-CH ₃ , 8-F	-OH	-CH ₃	-CH ₃	F	H	H	CF	-O-	111-113
86	8-F	-OH	-CH ₃	- ^s Pr	F	H	H	CH	-O-	amorphous
87	2- ^s Pr, 8-F	-OH	-CH ₃	- ^s Pr	F	H	H	CH	-O-	amorphous
88	2-CH ₃ , 8-F	-OH	-CH ₃	-Et	F	H	H	CH	-O-	amorphous
89	2-CH ₃ , 8-F	-OH	-CH ₃	-CH=CH ₂	F	H	H	CH	-O-	amorphous
90	2-CH ₃ , 8-F	-OH	-CH ₃	- ^s Pr	F	H	H	CH	-O-	amorphous
91	2-CH ₃ , 8-F	-OH	-CH ₃	-CH ₂ -(4-CH ₃ OPh)	F	H	H	CH	-O-	amorphous
92	2-CH ₃ , 8-F	-OH	-CH ₃	-CH ₂ -(4-FPh)	F	H	H	CH	-O-	amorphous

[0244]

Table 5

Compound No.	(R ⁴) _m	R ¹	R ²	R ³	R ^b	R ^c	R ^d	E	X-	Physical property
93	2-CH ₃ , 8-Cl	-OH	-CH ₃	-CH ₃	F	H	H	CH	-O-	96-98
94	2,8-(CH ₃) ₂	-OH	-CH ₃	-CH ₃	F	H	H	CH	-O-	amorphous
95	8-F	-OH	-CH ₃	-CH ₃	Cl	H	H	CH	-O-	120-123
96	8-F	-OH	-CH ₃	-Et	Cl	H	H	CH	-O-	39-42
97	2-CH ₃ , 8-F	-OH	-CH ₃	-CH ₃	Cl	H	H	CH	-O-	amorphous
98	8-F	-OH	-CH ₃	-CH ₃	Br	H	H	CH	-O-	107-110
99	8-F	-OH	-CH ₃	-CH ₃	H	Cl	H	CH	-O-	124-126
100	2-CH ₃ , 8-F	-OH	-CH ₃	-CH ₃	Br	H	H	CH	-O-	93-94
101	8-F	-OH	-CH ₃	-CH ₃	H	CH ₃	H	CH	-O-	amorphous
102	8-F	-OH	-CH ₃	-CH ₃	Cl	F	H	CH	-O-	135-137
103	-	-OH	-CH ₃	-CH ₃	Cl	H	H	CH	-O-	amorphous
104	8-F	-OH	-CH ₃	-CH ₃	H	F	H	CH	-O-	122-124
105	2-CH ₃ , 6-F	-OH	-CH ₃	-CH ₃	F	H	H	CH	-O-	amorphous
106	2-CH ₃ , 7-F	-OH	-CH ₃	-CH ₃	F	H	H	CH	-O-	98-100
107	2-CH ₃	-OH	-CH ₃	-CH ₃	F	H	H	CH	-O-	amorphous
108	2-CH ₃	-OH	-CH ₃	-CH ₃	Cl	H	H	CH	-O-	amorphous
109	-	-OCH ₃	-CH ₃	-CH ₃	F	H	H	CH	-O-	93-94
110	2-CH ₃ , 8-F	-OH	-CH ₃	-CH ₃	Cl	F	H	CH	-O-	138-140
111	-	-OH	-CH ₃	-CF ₃	F	H	H	CH	-O-	amorphous
112	-	-OH	-CF ₃	-CF ₃	H	H	H	CH	-O-	170-172
113	-	-OSi(CH ₃) ₃	-CF ₃	-CF ₃	F	H	H	CH	-O-	amorphous
114	-	-OH	-CF ₃	-CF ₃	F	H	H	CH	-O-	130-132
115	-	-OH	-CH ₃	-CH ₃	Cl	F	H	CH	-O-	115-116
116	5,8-F ₂	-OH	-CH ₃	-CH ₃	F	H	H	CH	-O-	154-155
117	2-CH ₃ , 5,8-F ₂	-OH	-CH ₃	-CH ₃	F	H	H	CH	-O-	124-126
118	2-CH ₃ , 7-Cl	-OH	-CH ₃	-CH ₃	F	H	H	CH	-O-	amorphous

[0245]

[Table 6]

Table 6

Compound No.	(R ⁴) _m	R ¹	R ²	R ³	R ^b	R ^c	R ^d	E	-X-	Physical property
119	-	-OH	-CH ₃	-CH ₃	F	F	H	CH	-O-	amorphous
120	5-F	-OH	-CH ₃	-CH ₃	F	H	H	CH	-O-	90-92
121	2-CH ₃ , 5-F	-OH	-CH ₃	-CH ₃	F	H	H	CH	-O-	95-97
122	8-F	-OH	-CF ₃	-CF ₃	F	H	H	CH	-O-	151-153
123	8-F	-OH	-CH ₃	-CH ₃	F	CH ₃	H	CH	-O-	118-120
124	7,8-F ₂	-OH	-CH ₃	-CH ₃	F	H	H	CH	-O-	78-80
125	2-CH ₃ , 7,8-F ₂	-OH	-CH ₃	-CH ₃	F	H	H	CH	-O-	108-110
126	7-F	-OH	-CH ₃	-CH ₃	F	H	H	CH	-O-	amorphous
127	2-CH ₃ , 4,8-F ₂	-OH	-CH ₃	-CH ₃	F	H	H	CH	-O-	130-132
128	2-CH ₃ , 7-F	-OH	-CH ₃	-Et	F	H	H	CH	-O-	307, 1,5580
129	8-F	-OSi(CH ₃) ₃	-CH ₃	-CN	F	H	H	CH	-O-	
130	8-F	-OH	-CH ₃	-COCH ₃	F	H	H	CH	-O-	
131	-	-OOH ₃	-H	-(CH ₂) ₂ -	F	H	H	CH	-O-	
132	-	-OOH ₃	-H	-CH ₃	H	H	H	CH	-O-	amorphous
133	-	-OEt	-H	-(CH ₂) ₂ -		H	H	CH	-O-	amorphous
134	-	-O ⁱ Pr	-H	-(CH ₂) ₂ -		H	H	CH	-O-	amorphous
135	-	-O ⁱ Bu	-H	-CH ₃	H	H	H	CH	-O-	amorphous
136	-	-OH	-H	- ⁱ Bu	H	H	H	CH	-O-	amorphous
137	-	-OH	-H	-(CH ₂) ₂ -		H	H	CH	-O-	amorphous
138	8-F	-OH	-H	-CH ₃	F	H	H	CH	-O-	149-151
139	2-CH ₃ , 8-F	-OH	-H	- ⁱ Bu	F	H	H	CH	-O-	104-106
140	8-F	-OCH ₃	-H	-CH ₃	F	H	H	CH	-O-	amorphous
141	8-F	-OH	-H	-CH ₃	Cl	F	H	CH	-O-	139-141
142	8-F	-OCH ₃	-H	-CH ₃	Cl	F	H	CH	-O-	amorphous
143	8-F	-OH	-H	- ⁱ Pr	Cl	H	H	CH	-O-	amorphous
144	8-F	-OH	-H	-C(CH ₃)=CH ₃	Cl	H	H	CH	-O-	103-104

[0246]

Table 7

Compound No.	(R ⁴) _m	R ¹	R ²	R ³	R ^b	R ^c	R ^d	E	-X-	Physical property
145	8-F	-CH ₃	-H	-t-Bu	Cl	H	H	CH	-O-	amorphous
146	8-F	-CH ₃	-H	-CH ₃	Cl	H	H	CH	-O-	amorphous
147	8-F	-OCH ₃	-H	-CH ₃	Cl	H	H	CH	-O-	amorphous
148	8-F	-CH ₃	-H	-CH ₃	Br	H	H	CH	-O-	amorphous
149	8-F	-OCH ₃	-H	-CH ₃	Br	H	H	CH	-O-	amorphous
150	8-F	-OCH ₃	-H	-CH ₃	CH ₃	H	H	CH	-O-	amorphous
151	2-CH ₃ , 8-F	-CH ₃	-H	-CH ₃	Cl	H	H	CH	-O-	112-114
152	2-CH ₃ , 8-F	-OCH ₃	-H	-CH ₃	Cl	H	H	CH	-O-	amorphous
153	8-F	-OCH ₃	-H	-CH ₃	CN	H	H	CH	-O-	amorphous
154	8-OCH ₃	-OCH ₃	-H	-CH ₃	Br	H	H	CH	-O-	amorphous
155	8-OCH ₃	-OCH ₃	-H	-CH ₃	OCH ₃	H	H	CH	-O-	amorphous
156	8-F	-OCH ₂ CH=CH ₂	-H	-CH ₃	Cl	H	H	CH	-O-	amorphous
157	8-F	-CH ₃	-H	-Et	Cl	H	H	CH	-O-	amorphous
158	8-F	-OSi(CH ₃) ₃	-H	-CF ₃	Cl	H	H	CH	-O-	amorphous
159	8-F	-CH ₃	-H	-CF ₃	Cl	H	H	CH	-O-	115-117
160	8-F	-OCH ₂ Ph	-H	-CH ₃	Cl	H	H	CH	-O-	174-176
161	8-F	-CH ₃	-H	-Ph	Cl	H	H	CH	-O-	amorphous
162	8-F	-CH ₃	-H	-CH=CH ₂	Cl	H	H	CH	-O-	98-100
163	8-F	-CH ₃	-H	-t-Bu	Cl	H	H	CH	-O-	amorphous
164	8-F	-CH ₃	-H	-C≡CH	Cl	H	H	CH	-O-	amorphous
165	8-F	-OSi(CH ₃) ₃	-H	-CN	Cl	H	H	CH	-O-	amorphous
166	8-F	-CH ₃	-H	-CH ₃	NO ₂	H	H	CH	-O-	112-114
167	8-F	-OCH ₃	-H	-CH ₃	NO ₂	H	H	CH	-O-	amorphous
168	8-F	-OCH ₃	-H	-CO ₂ CH ₃	Cl	H	H	CH	-O-	amorphous
169	8-F	-CH ₃	-H	-CN	Cl	H	H	CH	-O-	174-175
170	8-F	-OSi(CH ₃) ₃	-H	-COOCH ₃	Cl	H	H	CH	-O-	amorphous

[0247]

Table 8

Compound No.	$\langle R^4 \rangle_m$	R^1	R^2	R^3	R^b	R^c	R^d	E	X-	Physical property
171	8-F	-OH	-H	-OCH ₃	-OCH ₃	Cl	H	H	-O-	amorphous
172	8-F	-OCH ₃	-H	-OCH ₃	-OCH ₃	Cl	H	H	-O-	amorphous
173	8-F	-OH	-H	-OCH ₃	-O(CH ₂) ₂ OH	Cl	H	H	-O-	amorphous
174	8-F	-OH	-H	-OH	-CH ₂ Ph	Cl	H	H	-O-	amorphous
175	8-F	-OCH ₃	-H	-OCH ₃	-OCH ₂ Ph	Cl	H	H	-O-	amorphous
176	8-F	-OH	-H	-OH	-CH ₃	Br	H	H	-O-	148-150
177	8-F	-OCH ₃	-H	-OCH ₃	-CH ₃	Br	H	H	-O-	amorphous
178	2-CH ₃ , 8-F	-OH	-H	-OH	-CH ₃	Cl	F	H	-O-	130-131
179	2-CH ₃ , 8-F	-OH	-H	-OH	-CH ₃	Br	F	H	-O-	119-121
180	8-F	-OCH ₃	-H	-OCH ₃	-O(CH ₂) ₂ OH	Cl	H	H	-O-	amorphous
181	8-F	-OH	-H	-OH	-CO ₂ CH ₃	Cl	H	H	-O-	amorphous
182	8-F	-OH	-H	-OH	-CH ₂ ON	Cl	H	H	-O-	123-125
183	2-CH ₃ , 8-F	-OCH ₃	-H	-OH	-CH ₃	Cl	F	H	-O-	124-126
184	2-CH ₃ , 8-F	-OCH ₃	-H	-OH	-CH ₃	Br	F	H	-O-	108-110
185	8-F	-OH	-H	-OH	-CH ₃	CF ₃	H	H	-O-	amorphous
186	8-F	-OCH ₃	-H	-OH	-CH ₃	CF ₃	H	H	-O-	amorphous
187	-	-OH	-H	-OH	-CH ₃	Cl	H	H	-O-	77-80
188	-	-OCH ₃	-H	-OH	-CH ₃	Cl	H	H	-O-	amorphous
189	-	-OH	-H	-OH	-CH ₃	H	H	H	-O-	amorphous
190	-	-OH	-H	-OH	-CH ₃	Cl	H	H	-O-	amorphous
191	-	-OH	-H	-OH	-OCH ₃	H	H	H	-O-	157-159
192	-	-OH	-H	-OH	-CH ₃	Cl	F	H	-O-	133-135
193	-	-OH	-H	-OH	-OCH ₃	F	H	H	-O-	129-131
194	5,8-F ₂	-OH	-H	-OH	-CH ₃	F	H	H	-O-	115-116
195	5-F	-OH	-H	-OH	-CH ₃	F	H	H	-O-	114-115
196	8-F	-OH	-H	-OH	-OCH ₃	F	H	H	-O-	100-101

[0248]

Table 9

Compound No.	(R ⁴) _m	R ¹	R ²	R ³	R ⁵	R ⁶	R ^c	R ^d	E	X-	Physical property
197	7,8-F ₂	-OH	-H	-CH ₃		Cl	H	H	CH	-O-	99-102
198	7,8-F ₂	-OCH ₃	-H	-CH ₃		Cl	H	H	CH	-O-	amorphous
199	2-CH ₃ , 7,8-F ₂	-OH	-H	-CH ₃		Cl	H	H	CH	-O-	135-136
200	2-CH ₃ , 7,8-F ₂	-OCH ₃	-H	-CH ₃		Cl	H	H	CH	-O-	127-129
201	7-F	-OH	-H	-CH ₃		Cl	H	H	CH	-O-	92-94
202	2-CH ₃ , 7-F	-OCH ₃	-H	-CH ₃		Cl	H	H	CH	-O-	98-100
203	7-F	-OCH ₃	-H	-CH ₃		Cl	H	H	CH	-O-	amorphous
204	8-F	-OH	-H	-C(=N-OCH ₃)-CH ₃		Cl	H	H	CH	-O-	amorphous
205	8-F	-OH	-H	-CH ₃		I	H	H	CH	-O-	116-118
206	2-CH ₃ , 4,8-F ₂	-OH	-H	-CH ₃		F	H	H	CH	-O-	amorphous
207	8-F	-N(CH ₃) ₂	-H	-CH ₃		F	H	H	CH	-O-	n _D ²⁰ , 1.5936
208	8-F	-OCH ₃	-H	-CH ₃		I	H	H	CH	-O-	61-63
209	8-F	-OCH ₃	-H	-CH ₃		CO ₂ CH ₃	H	H	CH	-O-	amorphous
210	8-F	-OCH ₃	-H	-CH ₃		COCH ₃	H	H	CH	-O-	amorphous
211	8-F	-OCH ₃	-H	-CH ₃		SOCH ₃	H	H	CH	-O-	amorphous
212	8-F	-OCH ₃	-H	-CH ₃		SO ₂ CH ₃	H	H	CH	-O-	amorphous
213	8-F	-OCH ₃	-H	-CH ₃		COCH ₃	H	H	CH	-O-	
214	8-F	-OCH ₃	-H	-CH ₃		ⁿ Pr	H	H	CH	-O-	
215	8-F	-OCH ₃	-H	-CH ₃		Ph	H	H	CH	-O-	amorphous
216	8-F	-OCH ₃	-H	-CH ₃		N(CH ₃) ₂	H	H	CH	-O-	
217	8-F	-OCH ₃	-H	-OCH ₃		F	H	H	CH	-O-	amorphous
218	8-F	-OEt	-H	-OEt		F	H	H	CH	-O-	amorphous
219	8-F	-OH	-H	-H		Cl	H	H	CH	-O-	141-143
220	8-F	-OCH ₃	-H	-H		Cl	H	H	CH	-O-	amorphous
221	8-F	-OCH ₂ CH=CH ₂	-H	-H		Cl	H	H	CH	-O-	amorphous
222	8-F	-OH	-H	-H		Br	H	H	CH	-O-	142-144

[0249]

Table 10

Compound No.	(R ⁴) _m	R ¹	R ²	R ³	R ^p	R ^c	R ^d	E	X-	Physical property
223	B-F	-OCH ₃	-H	-H	Br	H	H	CH	-O-	amorphous
224	B-F	-NHt ⁿ Bu	-H	-H	Cl	H	H	CH	-O-	amorphous
225	-	-CH ₃	-CH ₃	-O(CH ₂) ₂ -CH ₂ OH	H	H	H	CH	-O-	amorphous
226	-	-CH ₃	-CH ₃	-O(CH ₂) ₂ -CH ₂ O-Tos	H	H	H	CH	-O-	amorphous
227	-	-CH ₃	-CH ₃	-O(CH ₂) ₂ OH	H	H	H	CH	-O-	amorphous
228	-	-CH ₃	-CH ₃	-CH ₂ OH	H	H	H	CH	-O-	105-108
229	-	-CH ₃	-CH ₃	-O(CH ₂) ₂ -CHO	H	H	H	CH	-O-	121-123
230	-	-CH ₃	-CH ₃	-O(CH ₂) ₂ -CH(CH ₂) ₂ OH	H	H	H	CH	-O-	amorphous
231	-	-CH ₃	-CH ₃	-O(CH ₂) ₂ -CH=CH-OH	H	H	H	CH	-O-	158-169
232	-	-CH ₃	-CH ₃	-O(CH ₂) ₂ -CN	H	H	H	CH	-O-	amorphous
233	-	-CH ₃	-CH ₃	-O(CH ₂) ₂ -COOCH ₃	H	H	H	CH	-O-	107-109
234	-	-CH ₃	-CH ₃	-CHO	H	H	H	CH	-O-	amorphous
235	-	-CH ₃	-CH ₃	-O(CH ₂) ₂ -CH ₂ OH	F	H	H	CH	-O-	amorphous
236	B-F	-CH ₃	-CH ₃	-O(CH ₂) ₂ -CH ₂ OH	H	H	H	CH	-O-	130-131
237	B-F	-CH ₃	-CH ₃	-O(CH ₂) ₂ -CH ₂ OH	F	H	H	CH	-O-	135-138
238	B-F	-CH ₃	-CH ₃	-O(CH ₂) ₂ -CHO	H	H	H	CH	-O-	143-149
239	B-F	-CH ₃	-CH ₃	-O(CH ₂) ₂ -CHO	F	H	H	CH	-O-	155-158
240	B-F	-CH ₃	-CH ₃	-O(CH ₂) ₂ -COOCH ₃	F	H	H	CH	-O-	147-148
241	B-F	-CH ₃	-CH ₃	-CH ₂ OH	H	H	H	CH	-O-	amorphous
242	B-F	-CH ₃	-CH ₃	-O(CH ₂) ₂ OH	H	H	H	CH	-O-	120-122
243	-	-CH ₃	-CH ₃	-CH ₂ -O(CH ₂) ₂ OH	H	H	H	CH	-O-	amorphous
244	-	-CH ₃	-CH ₃	-CH ₂ CH ₂ OH	H	H	H	CH	-O-	amorphous
245	-	-CH ₃	-CH ₃	-CH ₂ CHO	H	H	H	CH	-O-	93-96
246	-	-CH ₃	-CH ₃	-CH ₂ OH(CH ₂) ₂ OH	H	H	H	CH	-O-	amorphous
247	-	-CH ₃	-CH ₃	-CH ₂ CH ₂ COOCH ₃	H	H	H	CH	-O-	amorphous
248	-	-CH ₃	-CH ₃	-O(CH ₂) ₂ -CH ₂ COOCH ₃	H	H	H	CH	-O-	amorphous

[0250]

Table 11

Compound No.	(R ⁴) _m	R ¹	R ²	R ³	R ^b	R ^c	R ^d	E	-X-	Physical property
249	-	-CH ₃	-CH ₃	-C(CH ₃) ₂ -CO ₂ H	H	H	H	CH	-O-	181-184
250	-	-CH ₃	-CH ₃	-C(CH ₃) ₂ -CO ₂ CH ₃	H	H	H	CH	-O-	123-125
251	2-CH ₃	-CH ₃	-CH ₃	-C(CH ₃) ₂ OH	H	H	H	CH	-O-	amorphous
252	2-CH ₃	-CH ₃	-CH ₃	-CH ₂ CH ₂ OH	H	H	H	CH	-O-	amorphous
253	-	-CH ₃	-CH ₃	-CH(CH ₃)OH	H	H	H	CH	-O-	amorphous
254	-	-CH ₃	-CH ₃	-CH(OH)Et	H	H	H	CH	-O-	amorphous
255	-	-CH ₃	-CH ₃	-CH(CH ₃)-C(CH ₃) ₂ =CH ₂	H	H	H	CH	-O-	amorphous
256	2-CH ₃	-CH ₃	-CH ₃	-O(CH ₃) ₂ -CH(CH ₃)OH	H	H	H	CH	-O-	amorphous
257	2-CH ₃	-CH ₃	-CH ₃	-O(CH ₃) ₂ -COOCH ₃	H	H	H	CH	-O-	amorphous
258	8-F	-CH ₃	-CH ₃	-CH ₂ CH ₂ OH	H	H	H	CH	-O-	amorphous
259	2-CH ₃ , 8-F	-CH ₃	-CH ₃	-CH ₂ CH ₂ OH	H	H	H	CH	-O-	amorphous
260	-	-CH ₃	-CH ₃	-COOCH ₃	H	H	H	CH	-O-	amorphous
261	2-CH ₃ , 8-F	-CH ₃	-CH ₃	-CH ₂ CH ₂ COOCH ₃	H	H	H	CH	-O-	amorphous
262	8-F	-CH ₃	-CH ₃	-COOCH ₃	F	H	H	CH	-O-	amorphous
263	-	-CH ₃	-CH ₃	-C(CH ₃) ₂ OH	F	H	H	CH	-O-	amorphous
264	8-F	-CH ₃	-CH ₃	-C(CH ₃) ₂ OH	F	H	H	CH	-O-	139-141
265	-	-CH ₃	-CH ₃	-COOCH ₃	F	H	H	CH	-O-	amorphous
266	7-F	-CH ₃	-CH ₃	-COOCH ₃	F	H	H	CH	-O-	viscous oil
267	7,8-F ₂	-CH ₃	-CH ₃	-COOCH ₃	F	H	H	CH	-O-	viscous oil
268	8-F	-CH ₃	-CH ₃	-CH(CH ₃)OH	H	H	H	CH	-O-	viscous oil
269	8-F	-CH ₃	-CH ₃	-COOCH ₃	H	H	H	CH	-O-	viscous oil
270	2-CH ₃ , 8-F	-CH ₃	-CH ₃	-CN	H	H	H	CH	-O-	viscous oil
271	-	-CH ₃	-CH ₃	-CN	H	H	H	CH	-O-	amorphous
272	8-F	-CH ₃	-CH ₃	-CN	H	H	H	CH	-O-	126-128
273	8-F	-CH ₃	-CH ₃	-CONH ₂	H	H	H	CH	-O-	172-175
274	8-F	-CH ₃	-CH ₃	-CON(CH ₃) ₂	H	H	H	CH	-O-	164-166

[0251]

Table 12

Table 12

Compound No.	(R ⁴) _m	R ¹	R ²	R ³	R ^b	R ^c	R ^d	E	X-	Physical property	
275	8-F	-CH ₃	-CH ₃	-SO ₂ CH ₃	F	H	H	CH	-O-	175-177	
276	-	-H	-CH ₃	-CH ₃	H	H	H	CH	-O-	96-98	
277	-	-H	-CH ₃	-Et	H	H	H	CH	-O-	amorphous	
278	-	-H	-CH ₃	-Pr	H	H	H	CH	-O-	amorphous	
279	-	-H	-CH ₃	- ⁿ Bu	H	H	H	CH	-O-	²⁰⁴ _D 1.6215	
280	2-CH ₃ , 8-F	-CH	-H	-COCH ₃	F	H	H	CH	-O-	amorphous	
281	-	-H	-H	-C(CH ₃) ₂ OH	H	H	H	CH	-O-	131-132	
282	-	-H	-H	-H	H	H	H	CH	-O-	46-48	
283	-	-H	-H	-H	F	H	H	CH	-O-	74-75	
284	8-F	-H	-H	-H	F	H	H	CH	-O-	82-83	
285	-	-CH=CH-CH=CH-			-	H	H	CH	-O-	105-106	
286	-	-(CH ₂) ₄ -				H	H	CH	-O-	amorphous	
287	-	-(CH ₂) ₃ -				H	H	CH	-O-	68-70	
288	-	-CH=CH-CH ₂ CH ₂ -			-	H	H	CH	-O-	67-59	
289	-	-F	-F	-CO ₂ Et		F	H	H	CH	-O-	106-109
290	-	-F	-F	-CH ₂ OH		F	H	H	CH	-O-	129-131
291	-	-F	-F	-O(CH ₂) ₂ OH		F	H	H	CH	-O-	99-101
292	8-F	-F	-F	-CO ₂ Et		F	H	H	CH	-O-	²⁰⁴ _D 1.5400
293	8-F	-F	-F	-C(CH ₃) ₂ OH		F	H	H	CH	-O-	114-116
294	8-F	-F	-F	-COCH ₃		F	H	H	CH	-O-	60-63
295	8-F	-F	-F	-CON(CH ₃)OCH ₃		F	H	H	CH	-O-	amorphous
296	2-CH ₃ , 8-F	-F	-F	-COCH ₃		F	H	H	CH	-O-	104-106
297	8-F	-(CH ₂) ₂ -		-COCH ₂ Ph		F	H	H	CH	-O-	amorphous
298	8-F	-(CH ₂) ₂ -		-OH		F	H	H	CH	-O-	128-130
299	-	-(CH ₂) ₂ -		-COCH ₂ Ph		F	H	H	CH	-O-	91-93
300	-	-(CH ₂) ₂ -		-OH		F	H	H	CH	-O-	123-124

[0252]

Table 13

Compound No.	(R ⁴) _m	R ¹	R ²	R ³	R ^b	R ^c	R ^d	E	X-	Physical property
301	8-F	-(CH ₂) ₂ -		-OH	Cl	H	H	CH	-O-	
302	2-CH ₃ , 8-F	-(CH ₂) ₂ -		-OH	Cl	H	H	CH	-O-	
303	8-F	-(CH ₂) ₂ -		-COCH ₃ Ph	Cl	H	H	CH	-O-	
304	2-CH ₃ , 8-F	-(CH ₂) ₂ -		-COCH ₃ Ph	Cl	H	H	CH	-O-	
305	-	=CH ₂		-CH ₃	H	H	H	CH	-O-	72-74
306	2-Cl	=CH ₂		-CH ₃	H	H	H	CH	-O-	amorphous
307	2-OH	=CH ₂		-CH ₃	H	H	H	CH	-O-	135-139
308	8-F	-O-CH ₃ -		-CH ₃	F	H	H	CH	-O-	amorphous
309	-	-(CH ₂) ₄ -		-O(CH ₂) ₂ OH	H	H	H	CH	-O-	85-88
310	8-F	-(CH ₂) ₂ -		-CN	F	H	H	CH	-O-	142-144
311	2-CH ₃ , 8-F	-(CH ₂) ₂ -		-CN	F	H	H	CH	-O-	132-134
312	8-F	-(CH ₂) ₂ -		-COCH ₃	F	H	H	CH	-O-	
313	2-CH ₃ , 8-F	-(CH ₂) ₂ -		-COCH ₃	F	H	H	CH	-O-	
314	2-CH ₃ , 8-F	=N-OH		-CH ₃	F	H	H	CH	-O-	168-170
315	8-F	=N-OH		-CH ₃	Cl	H	H	CH	-O-	155-158
316	8-F	=N-OH		-CH ₃	Br	H	H	CH	-O-	145-147
317	8-F	=N-OH		-CH ₃	Cl	F	H	CH	-O-	199-202
318	2-CH ₃ , 7-F	=N-OCH ₃		-CH ₃	F	H	H	CH	-O-	n _D ²⁰ 1.5712
319	2-CH ₃ , 7-F	=N-OEt		-CH ₃	F	H	H	CH	-O-	n _D ³⁰ 1.5629
320	-	=N-O ⁱ Bu		-H	F	H	H	CH	-O-	105-107
321	8-F		≡N		F	H	H	CH	-O-	184-185
322	-		Ph		H	H	H	CH	-O-	85-86
323	-	-OH	-CH ₃	-CH ₃	F	H	H	CH	-CH ₂ -	148-149
324	8-F	-OH	-CH ₃	-CH ₃	H	H	H	CH	-CH ₂ -	114-116
325	2-CH ₃ , 8-F	-OH	-CH ₃	-CH ₃	H	H	H	CH	-CH ₂ -	159-161
326	-	-CH ₃	-CH ₃	-O(CH ₂) ₂ -CH ₂ OH	H	H	H	CH	-CH ₂ -	115-117

[0253]

Table 14

Compound No.	(R ⁴) _m	R ¹	R ²	R ³	R ^b	R ^c	R ^d	E	-X-	Physical property
327	8-F	-OH	-H	-CH ₃	Cl	H	H	OH	-CH ₂ -	151-153
328	8-F	-OCH ₃	-H	-COCH ₃	Cl	H	H	OH	-CH ₂ -	amorphous
329	8-F	=O		-OEt	H	H	H	OH	-CH ₂ -	amorphous
330	-	=O		-OEt	F	H	H	OH	-CH ₂ -	amorphous
331	7-F	=O		-OEt	F	H	H	OH	-CH ₂ -	viscous oil
332	8-F	=O		-H	Cl	H	H	OH	-CH ₂ -	111-112
333	8-F	-OH	-CH ₃	-CH ₃	F	H	H	OH	-CO-	182-193
334	-	-OH	-H	-CH=CH ₂	H	H	H	OH	-CO-	amorphous
335	8-F	=O(CH ₂) ₂		-H	F	H	H	OH	-CO-	43-45
336	-	-OCH ₂ CH ₂ O-		-H	H	H	H	OH	-CO-	134-136
337	-	=CH ₂		-H	H	H	H	OH	-CO-	92-95
338	-	=CH ₂		-CH ₃	H	H	H	OH	-CO-	125-128
339	-	=O(CH ₂) ₂		-H	H	H	H	OH	-CO-	amorphous
340	-			-CH=CH-CH=CH-		H	H	OH	-CO-	viscous oil
341	8-F	=O		-H	F	H	H	OH	-CO-	179-181
342	-	=O		-H	H	H	H	OH	-CO-	120-122
343	-	=O		-CH=CH ₂	H	H	H	OH	-CO-	amorphous
344	8-F	-H	-H	-O(CH ₂) ₂ CH ₂	H	H	H	OH	-CO-	amorphous
345	-	-H	-H	-CH ₂ -N ₃	H	H	H	OH	-CO-	amorphous
346	-	-H	-H	-Ph	Cl	H	H	OH	-CO-	115-117
347	-	-H	-H	-O(CH ₂) ₂ CH ₂	F	H	H	OH	-CO-	amorphous
348	8-F	-H	-H	-O(CH ₂) ₂ CH ₂	F	H	H	OH	-CO-	89-92
349	8-F	-OCH ₂ CH ₂ O-		-H	F	H	H	OH	-CH(OH)-	159-161
350	8-F	-OH	-CH ₃	-CH ₃	F	H	H	OH	-CH(OH)-	amorphous
351	-	=CH ₂		-H	H	H	H	OH	*4	amorphous
352	-	=O		-CO ₂ Et	H	H	H	OH	-O-	amorphous

[0254]

Table 15

Compound No.	(R ¹) _m	R ¹	R ²	R ³	R ^b	R ^c	R ^d	E	-X-	Physical property
353	8-F	=O		-CH ₃	Cl	H	H	CH	-O-	amorphous
354	8-F	=O		-CH ₃	*2	H	H	CH	-O-	146-148
355	8-F	=O		-CH ₃	Br	H	H	CH	-O-	amorphous
356	8-F	=O		-CH ₃	F	H	H	CF	-O-	103-104
357	8-F	=O		*3	F	H	H	CH	-O-	amorphous
358	2-CH ₃ , 8-F	=O		-CH ₃	F	H	H	CH	-O-	96-100
359	2-CH ₃ , 8-F	=O		-CH ₃	Cl	H	H	CH	-O-	amorphous
360	2-CH ₃ , 8-F	=O		-CH ₃	Br	H	H	CH	-O-	amorphous
361	8-F	=O		-CH ₃	Cl	F	H	CH	-O-	amorphous
362	2,4-(CH ₃) ₂ , 8-F	=O		-CH ₃	F	H	H	CH	-O-	164-65
363	8-F	=O		- ^o Pr	Cl	H	H	CH	-O-	amorphous
364	8-F	=O		-C(CH ₃)=CH ₂	Cl	H	H	CH	-O-	amorphous
365	8-F	=O		- ^t Bu	Cl	H	H	CH	-O-	amorphous
366	2-CH ₃ , 8-Cl	=O		-CH ₃	F	H	H	CH	-O-	100-102
367	2,6-(CH ₃) ₂	=O		-CH ₃	F	H	H	CH	-O-	amorphous
368	8-F	=O		-Ph	Cl	H	H	CH	-O-	amorphous
369	8-F	=O		-CH=CH ₂	Cl	H	H	CH	-O-	77-79
370	8-F	=O		-C≡CH	Cl	H	H	CH	-O-	121-123
371	8-F	=O		- ⁿ Bu	Cl	H	H	CH	-O-	amorphous
372	8-F	=O		-CH ₂ Cl	F	H	H	CH	-O-	113-116
373	8-F	=O		-CHCl ₂	F	H	H	CH	-O-	amorphous
374	8-F	=O		-CCl ₃	F	H	H	CH	-O-	amorphous
375	8-F	=O		-COCH ₃	Cl	H	H	CH	-O-	amorphous
376	8-F	=O		-C(CH ₃) ₂ OH	Cl	H	H	CH	-O-	amorphous
377	8-F	=O		- ⁱ Pr	Cl	H	H	CH	-O-	amorphous
378	8-F	=O		-CH ₃	Br	F	H	CH	-O-	132-135

[0255]

Table 16

Compound No.	$\langle R^3 \rangle_m$	R ¹	R ²	R ³	R ^b	R ^c	R ^d	E	-X-	Physical property
379	2-CH ₃ , 8-F	=O		-CH ₃	Cl	F	H	CH	-O-	126-128
380	2-CH ₃ , 8-F	=O		-CH ₃	Br	F	H	CH	-O-	126-128
381	-	=O		-CH ₃	Cl	H	H	CH	-O-	38-40
382	2-CH ₃ , 6-F	=O		-CH ₃	F	H	H	CH	-O-	amorphous
383	2-CH ₃ , 7-F	=O		-CH ₃	F	H	H	CH	-O-	amorphous
384	8-F	=O		-CH ₃ O-COCH ₃	F	H	H	CH	-O-	amorphous
385	-	=O		-CF ₃	F	H	H	CH	-O-	100-104
386	-	=O		-CH ₃	Cl	F	H	CH	-O-	112-124
387	5,8-F ₂	=O		-CH ₃	F	H	H	CH	-O-	45-48
388	5-F	=O		-CH ₃	F	H	H	CH	-O-	amorphous
389	2-CH ₃ , 7-Cl	=O		-CH ₃	F	H	H	CH	-O-	amorphous
390	8-F	=O		-CF ₃	F	H	H	CH	-O-	amorphous
391	7,8-F ₂	=O		-CH ₃	F	H	H	CH	-O-	128-130
392	7,8-F ₂	=O		-CH ₃	Cl	H	H	CH	-O-	105-107
393	2-CH ₃ , 8-F	=O		-COCH ₃	Cl	H	H	CH	-O-	113-114
394	7-F	=O		-CH ₃	F	H	H	CH	-O-	97-99
395	2-CH ₃ , 8-F	=O		-COCH ₃	F	H	H	CH	-O-	169-170
396	8-F	=O		-COCH ₃	F	H	H	CH	-O-	96-98
397	2-CH ₃ , 8-F	=O		-C(CH ₃) ₂ OH	F	H	H	CH	-O-	amorphous
398	8-F	=O		-C(CH ₃) ₂ OH	F	H	H	CH	-O-	amorphous
399	8-F	=O		-C(=N-OCH ₃)-CH ₃	Cl	H	H	CH	-O-	118-119
400	2-CH ₃ , 4,8-F ₂	=O		-CH ₃	F	H	H	CH	-O-	155-156
401	-	=O		-NHPh	H	H	H	CH	-O-	amorphous
402	-	=O		-OCH ₃	Cl	H	H	CH	-O-	amorphous
403	-	=O		-OCH ₃	Br	H	H	CH	-O-	amorphous
404	8-F	=O		-OEt	H	Cl	H	CH	-O-	88-90

Table 17

Compound No.	(R ⁴) _n	R ¹	R ²	R ³	R ^b	R ^c	R ^d	E	X-	Physical property
406	8-F	=O		-OEt	H	CH ₃	H	CH	-O-	88-90
406	8-F	=O		-OCH ₃	Cl	H	H	CH	-O-	²⁰⁶ _D 1.6008
407	8-F	=O		-OPr	Cl	H	H	CH	-O-	²⁰⁷ _D 1.5809
408	8-F	=O		-O-CH ₂ Ph	Cl	H	H	CH	-O-	114-116
408	8-F	=O		-OEt	F	CH ₃	H	CH	-O-	amorphous
410	8-F	=O		-O ⁱ Bu	Cl	H	H	CH	-O-	amorphous
411	8-F	=O		-O-C(CH ₃) ₂ -CO ₂ Et	Cl	H	H	CH	-O-	²²⁴ _D 1.5411
412	8-F	=O		-H	Cl	F	H	CH	-O-	153-155
413	8-F	=O		-H	Cl	H	H	CH	-O-	133-135
414	8-F	=O		-H	Br	H	H	CH	-O-	111-114
415	8-F	=O		-H	Bn	H	H	CH	-O-	92-94
416	8-F	=O		-H	NO ₂	H	H	CH	-O-	106-108
417	8-F	=O		-H	Br	F	H	CH	-O-	140-142
418	8-F	=O		-H	CF ₃	H	H	CH	-O-	125-127
419	-	=O		-H	Cl	H	H	CH	-O-	82-85
420	-	=O		-H	Cl	F	H	CH	-O-	136-138
421	5,8-F ₂	=O		-H	F	H	H	CH	-O-	118-121
422	5-F	=O		-H	F	H	H	CH	-O-	92-94
423	7,8-F ₂	=O		-H	Cl	H	H	CH	-O-	142-144
424	7-F	=O		-H	Cl	H	H	CH	-O-	122-123
425	2-CH ₃ , 8-F	-F	-F	-O(CH ₂) ₂ OH	F	H	H	CH	-O-	110-112
426	2-CH ₃ , 8-F	-F	-F	-CH(CH ₃)OH	F	H	H	CH	-O-	163-165
427	2-CH ₃ , 8-F	-F	-F	-CH(CH ₃)OCH ₃	F	H	H	CH	-O-	amorphous
428	2-CH ₃ , 8-F	-F	-F	-C(CH ₃) ₂ OH	F	H	H	CH	-O-	amorphous
429	2-CH ₃ , 8-F	-F	-F	-CH(CH ₃)OOH ₂ COCH ₃	F	H	H	CH	-O-	141-143
430	8-F	-F	-F	-O ₂ Et	F	H	H	CH	-O-	amorphous

[0257]

Table 18

Compound No.	(R ⁴) _m	R ¹	R ²	R ³	R ^b	R ^o	R ^d	E	-X-	Physical property
431	8-F	-(CH ₂) ₂ -		-C(CH ₃) ₂ OH	F	H	H	CH	-O-	amorphous
432	8-F	-CH ₃	-CH ₃	-CN	F	H	H	CH	-O-	110-112
433	8-F	-NHCO ₂ ^t Bu	-CH ₃	-CH ₃	F	H	H	CH	-O-	127-129
434	8-F	-(CH ₂) ₂ -		-NHCO ₂ ^t Bu	F	H	H	CH	-O-	132-134
435	8-F	-NH ₂	-CH ₃	-CH ₃	F	H	H	CH	-O-	amorphous
436	8-F	-NHCOCH ₃	-CH ₃	-CH ₃	F	H	H	CH	-O-	144-146
437	8-F	-NHCOCF ₃	-CH ₃	-CH ₃	F	H	H	CH	-O-	amorphous
438	-	-COCH ₂ CH ₂ O-		-H	H	H	H	CH	-CH(OH)-	80-83
439	2-CH ₃ , 7,8-F ₂	-COCH ₃	-CH ₃	-CH ₃	F	H	H	CH	-O-	97-98
440	2-CH ₃ , 8-F	-COCH ₃	-CH ₃	-CH ₃	F	H	H	CH	-O-	94-95
441	2-CH ₃ , 7-F	-COCH ₃	-CH ₃	-CH ₃	F	H	H	CH	-O-	104-106
442	8-F	-N(CH ₃) ₂	-H	-CN	Cl	H	H	CH	-O-	amorphous
443	8-F	-NHCOCH ₃	-H	-CN	Cl	H	H	CH	-O-	amorphous
444	8-F	-N(Bn)COCH ₃	-CH ₃	-CH ₃	F	H	H	CH	-O-	195-197
445	8-F	Pyrazolidin-1-yl	-H	-CH ₃	F	H	H	CH	-O-	205-206, 15593
446	8-F	-COCH ₃	-H	-CH ₃	NHCO ₂ ^t Bu	H	H	CH	-O-	amorphous
447	8-F	-COCH ₃	-H	-CH ₃	NH ₂	H	H	CH	-O-	amorphous
448	8-F	-COCH ₃	-H	-CH ₃	NHCH ₃	H	H	CH	-O-	amorphous
449	2-CH ₃ , 8-F	-CH ₃	-CH ₃	-SO ₂ CH ₃	F	H	H	CH	-O-	138-141
450	7,8-F ₂	-CH ₃	-CH ₃	-SO ₂ CH ₃	F	H	H	CH	-O-	157-160
451	2-CH ₃ , 7,8-F ₂	-CH ₃	-CH ₃	-SO ₂ CH ₃	F	H	H	CH	-O-	172-175

[0258]


Table 19

Compound No.	(R ¹) _m	R ¹	R ²	R ³	R ^b	R ^c	R ^d	E	-X-	Physical property
452	8-F	-CH ₃	-CH ₃	-SO ₂ Et	F	H	H	OH	-O-	150-152
453	8-F	-CH ₃	-H	-SO ₂ CH ₃	Cl	H	H	CH	-O-	amorphous

*1="-(N-CH₂CH₂CH₂)₂-"


*3="

*2="8-Fluoro-quinolin-3-yloxy"



[0259]


Table 20



Compound No.	(R ⁴) _m	R ¹	R ²	R ³	R ⁰	R ^d	E	-X-	Physical property
a-1	-		≡ N		H	H	CH	-O-	149-150
a-2	-	-OH	-CH ₃	-CH ₃	H	H	CH	-O-	amorphous
a-3	8-F		=O	-H	H	H	CH	-O-	138-140
a-4	2- ^t Bu, 8-F		=O	-H	H	H	CH	-O-	150-151
a-5	8-F	-OH	-CH ₃	- ^t Bu	H	H	CH	-O-	110-112
a-6	8-F		=O	- ^t Bu	H	H	CH	-O-	90-91
a-7	2-CH ₃ , 8-F	-OH	-CH ₃	-CH ₃	H	H	CH	-O-	amorphous
a-8	8-F	-OH	-CH ₃	-CH ₃	H	H	CH	-O-	amorphous
a-9	2-CH ₃ , 8-F		=O	-CH ₃	H	H	CH	-O-	133-134
a-10	2-CH ₃ , 8-F	-OH	-CH ₃	- ^t Bu	H	H	CH	-O-	133-136
a-11	8-F	-CH ₃	-CH ₃	-CN	H	H	CH	-O-	
a-12	8-F	-CH ₃	-CH ₃	-CO ₂ Et	H	H	CH	-O-	
a-13	8-F	-F	-F	-O(CH ₂) ₂ CH	H	H	CH	-O-	
a-14	8-F	-CH ₃	-CH ₃	-O(CH ₂) ₂ CH	H	H	CH	-O-	
a-15	8-F	-CH ₃	-CH ₃	-COCH ₃	H	H	CH	-O-	

[0260]




Table 21



Compound No.	Q	R ¹	R ²	R ³	R ^b	R ^c	R ^d	E	-X-	Physical property
b-1		-CH ₃	-CH ₃	-C(CH ₃) ₂ OH	H	H	H	CH	-O-	viscous oil
b-2		-CH ₃	-CH ₃	-C(CH ₃) ₂ OH	H	H	H	CH	-O-	133-135
b-3		-CH ₃	-CH ₃	-C(CH ₃) ₂ OH	F	H	H	CH	-O-	viscous oil
b-4		-CH ₃	-CH ₃	-C(CH ₃) ₂ OH	H	H	H	CH	-O-	viscous oil
b-5		-CH ₃	-CH ₃	-C(CH ₃) ₂ OH	H	H	H	CH	-O-	viscous oil
b-6		=O	-CH ₃	-CH ₃	F	H	H	CH	-O-	viscous oil


[0261]

Table 22

Compound No.	Q	R ¹	R ²	R ³	R ^b	R ^c	R ^d	E	-X-	Physical property
b-7		-OH	-CH ₃	-CH ₃	F	H	H	CH	-O-	viscous oil
b-8		-OH	-CH ₃	- ^t Bu	F	H	H	CH	-O-	viscous oil
b-9		-OH	-H	-CH ₃	F	H	H	CH	-O-	viscous oil
b-10		-OCH ₃	-H	-CH ₃	F	H	H	CH	-O-	viscous oil
b-11		=O		-CH ₃	F	H	H	CH	-O-	viscous oil
b-12		-OH	-CH ₃	-CH ₃	F	H	H	CH	-O-	viscous oil
b-13		-CH ₃	-CH ₃	-C(CH ₃) ₂ OH	H	H	H	CH	-O-	viscous oil
b-14		-OH	-H	-CH ₃	F	H	H	CH	-O-	viscous oil


[0262]

Table 23

Compound No.	Q	R ¹	R ²	R ³	R ^b	R ^c	R ^d	E	-X-	Physical property
b-15		-OCH ₃	-H	-CH ₃	F	H	H	OH	-O-	viscous oil
b-16		-OH	-CH ₃	-tBu	F	H	H	OH	-O-	amorphous
b-17		=O		-CH ₃	F	H	H	OH	-O-	78-80
b-18		-OH	-CH ₃	-CH ₃	F	H	H	OH	-O-	101-102
b-19		=C(CH ₃) ₂		-H	F	H	H	OH	-CO-	viscous oil
b-20		=C(CH ₃) ₂		-H	F	H	H	OH	-CO-	amorphous
b-21		=O		-CH ₃	F	H	H	OH	-O-	viscous oil
b-22		-OH	-CH ₃	-CH ₃	F	H	H	OH	-O-	viscous oil


[0263]

Table 24

Compound No.	Q	R ¹	R ²	R ³	R ^b	R ^c	R ^d	E	-X-	Physical property
b-23		=O	-CH ₃	-CH ₃	F	H	H	CH	-O-	viscous oil
b-24				-CH ₃	F	H	H	CH	-O-	87-89
b-25		-OH	-CH ₃	-CH ₃	F	H	H	CH	-O-	
b-26		-OH	-CH ₃	-CH ₃	F	H	H	CH	-O-	

[0264]


Table 25



Compound No.	$(R^4)_m$	R^1	R^2	R^3	-X-	Physical property
C-1	8-F	=O		-OCH ₃	-O-	
C-2	8-F	=O		-H	-O-	135-137
C-3	8-F	=O		-CH ₃	-O-	138-139
C-4	8-F	-OH	-CH ₃	-CH ₃	-O-	129-131
C-5	8-F	-OH	-H	-CH ₃	-O-	96-98

[0265]

Table 26



Compound No.	(R ⁴) _m	R ¹	R ²	R ³	(R ⁵) _n	X	Physical property
d-1	8-F	=O		-OCH ₃	2-CH ₃ 5-OF ₃	-O-	
d-2	8-F	=O		-H	2-CH ₃ 5-OF ₃	-O-	
d-3	8-F	=O		-CH ₃	2-CH ₃ 5-OF ₃	-O-	
d-4	8-F	-OH	-H	-CH ₃	2-CH ₃ 5-OF ₃	-O-	160-162
d-5	8-F	-OCH ₃	-H	-CH ₃	2-CH ₃ 5-OF ₃	-O-	206, 15056
d-6	8-F	-OH	-H	-Pr	2-CH ₃ 5-OF ₃	-O-	
d-7	8-F	-OH	-H	- ^s Pr	2-CH ₃ 5-OF ₃	-O-	
d-8	8-F	-OH	-H	-OF ₃	2-CH ₃ 5-OF ₃	-O-	
d-9	8-F	-OH	-H	-C(CH ₃) ₂ OH	2-CH ₃ 5-OF ₃	-O-	
d-10	8-F	-OH	-H	-C(CH ₃) ₂ -COOCH ₃	2-CH ₃ 5-OF ₃	-O-	
d-11	8-F	-OH	-H	-C(CH ₃) ₂ -CO ₂ CH ₃	2-CH ₃ 5-OF ₃	-O-	

[0266]

Table 27

Compound No.	(R ⁴) _m	R ¹	R ²	R ³	(R ⁵) _n	X-	Physical property
d-12	8-F	-OH	-H	-CH ₂ Ph	2-CH ₃ 5-OF ₃	-O-	
d-13	8-F	-OH	-H	-CH=CH ₂	2-CH ₃ 5-OF ₃	-O-	
d-14	8-F	-OH	-H	-C≡CH	2-CH ₃ 5-OF ₃	-O-	
d-15	8-F	-OH	-H	-OOCCH ₃	2-CH ₃ 5-OF ₃	-O-	
d-16	2-CH ₃ 8-F	-OH	-H	-CH ₃	2-CH ₃ 5-OF ₃	-O-	
d-17	2-CH ₃ 8-F	-OH	-H	-Pr	2-CH ₃ 5-OF ₃	-O-	
d-18	2-CH ₃ 8-F	-OH	-H	- ^s Pr	2-CH ₃ 5-OF ₃	-O-	
d-19	2-CH ₃ 8-F	-OH	-H	-OF ₃	2-CH ₃ 5-OF ₃	-O-	
d-20	2-CH ₃ 8-F	-OH	-H	-O(CH ₂) ₂ OH	2-CH ₃ 5-OF ₃	-O-	
d-21	2-CH ₃ 8-F	-OH	-H	-CH=CH ₂	2-CH ₃ 5-OF ₃	-O-	
d-22	2-CH ₃ 8-F	-OH	-H	-C≡CH	2-CH ₃ 5-OF ₃	-O-	
d-23	8-F	-OH	-CH ₃	-CH ₃	2-CH ₃ 5-Cl	-O-	
d-24	8-F	-OH	-CH ₃	-Pr	2-CH ₃ 5-Cl	-O-	
d-25	8-F	-OH	-CH ₃	- ^s Pr	2-CH ₃ 5-Cl	-O-	
d-26	8-F	-OH	-CH ₃	-OF ₃	2-CH ₃ 5-Cl	-O-	
d-27	8-F	-OH	-CH ₃	-O(CH ₂) ₂ OH	2-CH ₃ 5-Cl	-O-	
d-28	8-F	-OH	-CH ₃	-O(CH ₂) ₂ -COCH ₃	2-CH ₃ 5-Cl	-O-	
d-29	8-F	-OH	-CH ₃	-O(CH ₂) ₂ -CO ₂ CH ₃	2-CH ₃ 5-Cl	-O-	
d-30	8-F	-OH	-CH ₃	-CH ₂ Ph	2-CH ₃ 5-Cl	-O-	
d-31	8-F	-OH	-CH ₃	-CH=CH ₂	2-CH ₃ 5-Cl	-O-	
d-32	8-F	-OH	-CH ₃	-C≡CH	2-CH ₃ 5-Cl	-O-	
d-33	8-F	-OH	-CH ₃	-OOCCH ₃	2-CH ₃ 5-Cl	-O-	
d-34	2-CH ₃ 8-F	-OH	-CH ₃	-CH ₃	2-CH ₃ 5-Cl	-O-	
d-35	2-CH ₃ 8-F	-OH	-CH ₃	-Pr	2-CH ₃ 5-Cl	-O-	

[0267]

Table 28

Compound No.	(R ⁴) _m	R ¹	R ²	R ³	(R ⁵) _n	-X-	Physical property
d-36	2-CH ₃ , 8-F	-OH	-CH ₃	- ⁱ Pr	2-CH ₃ , 5-Cl	-O-	
d-37	2-CH ₃ , 8-F	-OH	-CH ₃	-CF ₃	2-CH ₃ , 5-Cl	-O-	
d-38	2-CH ₃ , 8-F	-OH	-CH ₃	-C(CH ₃) ₂ OH	2-CH ₃ , 5-Cl	-O-	
d-39	2-CH ₃ , 8-F	-OH	-CH ₃	-CH=CH ₂	2-CH ₃ , 5-Cl	-O-	
d-40	2-CH ₃ , 8-F	-OH	-CH ₃	-C≡CH	2-CH ₃ , 5-Cl	-O-	

[0268]

Table 29

Compound No	¹ H-NMR
3	1.16(t, 3H), 4.23(q, 2H), 7.13(d, 1H), 7.29–7.64(m, 6H), 8.02(m, 1H), 8.08(d, 1H), 8.83(d, 1H)
5	1.19(t, 3H), 4.24(q, 2H), 6.80(m, 1H), 7.01(m, 1H), 7.43(d, 1H), 7.50–7.71(m, 3H), 8.04–8.16(m, 2H), 8.83(br, 1H)
7	1.22(t, 3H), 4.31(q, 2H), 6.79(d, 1H), 6.99(t, 1H), 7.38(m, 1H), 7.51–7.72(m, 4H), 8.11(d, 1H), 8.81(d, 1H)
8	1.20(t, 3H), 4.32(q, 2H), 6.85(d, 1H), 7.03(t, 1H), 7.30–7.54(m, 5H), 8.85(d, 1H)
9	7.03(d, 1H), 7.27(d, 1H), 7.30–7.77(m, 6H), 8.12(d, 1H), 8.81(d, 1H)
10	2.61(s, 3H), 6.79(m, 1H), 7.01(t, 1H), 7.31–7.43(m, 2H), 7.47–7.50(m, 2H), 7.58(m, 1H), 8.83(d, 1H)
11	1.28(s, 9H), 6.97(d, 1H), 7.17–7.26(m, 2H), 7.26(m, 1H), 7.52(t, 1H), 7.56(d, 1H), 7.63(m, 1H), 7.69(d, 1H), 8.09(d, 1H), 8.77(d, 1H)
12	1.28(s, 9H), 6.76(d, 1H), 6.98(t, 1H), 7.27–7.50(m, 4H), 7.59(m, 1H), 8.87(d, 1H)
13	7.13(m, 1H), 7.54(m, 1H), 7.69(m, 1H), 7.79(m, 1H), 7.96–8.04(m, 2H), 8.14(d, 1H), 8.25(br, 1H), 8.83(d, 1H)
14	1.35(d, 9H), 6.82(d, 1H), 7.03–7.14(m, 2H), 7.38–7.56(m, 5H), 8.02(d, 1H), 8.73(s, 1H)
17	0.71(t, 3H), 1.39(s, 6H), 1.84(q, 2H), 6.90(m, 1H), 7.12–7.23(m, 2H), 7.41(d, 1H), 7.43–7.68(m, 4H), 8.10(d, 1H), 8.80(br, 1H)
18	0.66(t, 3H), 1.37(s, 6H), 1.85(q, 2H), 6.64(d, 1H), 6.84–6.93(m, 2H), 7.07(m, 1H), 7.15–7.26(m, 2H), 7.41–7.47(m, 2H), 8.82(d, 1H)
22	2.98(s, 3H), 3.12(s, 3H), 6.88(d, 1H), 7.20(t, 1H), 7.41(d, 1H), 7.54(t, 1H), 7.63–7.73(m, 3H), 8.10(d, 1H), 8.79(d, 1H)
24	2.19(s, 3H), 3.89(s, 3H), 7.03(d, 1H), 7.26(m, 1H), 7.36–7.43(m, 2H), 7.48–7.55(m, 2H), 7.59–7.67(m, 2H), 8.09(d, 1H), 8.81(d, 1H)
25	1.18(s, 9H), 2.18(s, 3H), 7.05(d, 1H), 7.26(m, 1H), 7.35–7.64(m, 6H), 8.08(d, 1H), 8.80(d, 1H)
27	1.73(s, 3H), 3.17(s, 6H), 6.98(d, 1H), 7.19(t, 1H), 7.29(m, 1H), 7.48–7.68(m, 4H), 7.82(dd, 1H), 8.08(d, 1H), 8.79(d, 1H)
28	0.20(s, 9H), 1.76(d, 1H), 4.57(br, 1H), 6.62(d, 1H), 6.92(m, 1H), 7.15(m, 1H), 7.33(m, 1H), 7.33(m, 1H), 7.46–7.51(m, 2H), 7.70(s, 1H), 8.83(br, 1H)
30	1.06(s, 9H), 1.76(d, 1H), 4.57(br, 1H), 6.62(d, 1H), 6.92(m, 1H), 7.15(m, 1H), 7.33(m, 1H), 7.46–7.51(m, 2H), 7.70(s, 1H), 8.83(br, 1H)
31	1.03(s, 9H), 1.60(s, 1H), 1.68(s, 3H), 6.86(m, 1H), 7.13–7.24(m, 2H), 7.51–7.71(m, 5H), 8.11(d, 1H), 8.79(d, 1H)
34	1.70(s, 3H), 2.22–2.37(m, 2H), 2.43(br, 1H), 2.91(m, 1H), 3.08(m, 1H), 6.72(d, 1H), 7.07(d, 1H), 7.24(t, 1H), 7.32(m, 1H), 7.45–7.49(m, 2H), 7.59(m, 1H), 8.88(d, 1H)
35	1.07(s, 9H), 2.13(m, 1H), 2.65(m, 1H), 2.93–2.99(m, 3H), 6.69(d, 1H), 7.03(d, 1H), 7.19(t, 1H), 7.33(m, 1H), 7.45–7.52(m, 2H), 7.67(m, 1H), 8.85(d, 1H)

[0269]

Table 30

Compound No	¹ H-NMR
36	1.75(d, 6H), 3.79(d, 1H), 6.69(m, 1H), 6.95(m, 1H), 7.20(m, 1H), 7.35(m, 1H), 7.45-7.51(m, 2H), 7.59(d, 1H), 8.83(d, 1H).
37	1.76(d, 6H), 2.80(s, 3H), 3.87(br, 1H), 6.58(d, 1H), 6.93(m, 1H), 7.19(m, 1H), 7.29-7.45(m, 4H).
38	1.70(s, 6H), 3.01(br, 1H), 6.87(d, 1H), 7.15-7.25(m, 2H), 7.54(t, 1H), 7.60-7.71(m, 4H), 8.12(d, 1H), 8.81(d, 1H).
39	1.28(s, 9H), 7.03(s, 1H), 7.36(d, 1H), 7.52-7.72(m, 4H), 7.95(d, 1H), 8.13(d, 1H), 8.87(d, 1H), 10.42(s, 1H).
40	1.64(s, 6H), 2.53(s, 1H), 7.12(m, 1H), 7.21-7.32(m, 2H), 7.50-7.67(m, 4H), 8.11(d, 1H), 8.85(d, 1H).
41	1.70(s, 6H), 2.77(s, 1H), 6.56(m, 1H), 6.86(m, 1H), 7.54-7.75(m, 5H), 8.13(d, 1H), 8.80(d, 1H).
42	1.68(s, 6H), 2.88(s, 1H), 6.84-6.97(m, 2H), 7.44(m, 1H), 7.52-7.57(m, 2H), 7.62-7.71(m, 2H), 8.12(d, 1H), 8.79(d, 1H).
43	1.77(d, 6H), 4.00(d, 1H), 6.66(m, 1H), 6.90(m, 1H), 8.18(m, 1H), 8.53-7.73(m, 4H), 8.12(d, 1H), 8.79(d, 1H).
46	1.43(t, 3H), 1.72(s, 6H), 2.95(s, 1H), 3.14(q, 2H), 6.80(d, 1H), 7.16-7.25(m, 2H), 7.43-7.48(m, 2H), 7.60-7.67(m, 3H), 8.07(d, 1H).
47	0.94(t, 3H), 1.16(m, 2H), 1.72(s, 6H), 1.87(m, 2H), 2.92(s, 1H), 3.10(t, 2H), 6.81(d, 1H), 7.14-7.23(m, 2H), 7.43-7.48(m, 2H), 7.59-7.67(m, 3H), 8.07(d, 1H).
48	0.93(t, 3H), 1.72(d, 3H), 1.95(m, 1H), 2.08(m, 1H), 3.87(d, 1H), 6.68(d, 1H), 6.95(m, 1H), 7.21(m, 1H), 7.36(m, 1H), 7.44-7.51(m, 2H), 7.59(m, 1H), 8.82(d, 1H).
49	1.75(d, 6H), 3.76(d, 1H), 6.69(m, 1H), 6.94(m, 1H), 7.20(m, 1H), 7.47(m, 1H), 7.59-7.66(m, 2H), 7.78(dd, 1H), 8.90(d, 1H).
51	6.81-6.84(m, 1H), 7.39-7.64(m, 8H), 7.72-7.76(m, 3H), 7.99(d, 1H), 8.07(d, 1H), 8.89(d, 1H), 10.29(s, 1H).
53	1.19(t, 3H), 4.25(q, 2H), 4.31(s, 2H), 7.04(d, 1H), 7.32-7.53(m, 5H), 7.86(s, 1H), 8.84(d, 1H).
59	0.94(s, 6H), 1.26(s, 6H), 2.45(s, 3H), 3.91(s, 2H), 7.03(m, 1H), 7.28-7.37(m, 3H), 7.40(m, 1H), 7.53(d, 1H), 7.61(t, 1H), 7.80-7.88(m, 4H), 8.16(d, 1H), 8.40(d, 1H), 9.27(d, 1H).
60	1.37(s, 6H), 1.98(t, 1H), 2.98(t, 1H), 6.72(d, 1H), 7.08(d, 1H), 7.17(t, 1H), 7.47-7.65(m, 3H), 8.02(dd, 1H).
61	6.96(m, 1H), 7.16-7.32(m, 2H), 7.39-7.49(m, 3H), 7.57(m, 1H), 7.99(d, 1H), 8.45(br, 1H).
67	1.72(s, 6H), 2.77(s, 3H), 3.01(br, 1H), 6.79(d, 1H), 7.13-7.25(m, 2H), 7.43-7.49(m, 2H), 7.59-7.67(m, 3H), 8.05(d, 1H).
68	1.43(d, 6H), 1.73(s, 6H), 3.01(s, 1H), 3.64(m, 1H), 6.78(d, 1H), 7.12-7.25(m, 2H), 7.41-7.46(m, 2H), 7.58-7.65(m, 3H), 8.08(d, 1H).
69	1.70(s, 3H), 3.57(s, 1H), 4.11(s, 3H), 6.83(d, 1H), 7.11-7.23(m, 2H), 7.38(t, 1H), 7.54-7.62(m, 4H), 7.88(d, 1H).

[0270]

Table 31

Compound No.	¹ H-NMR
70	1.35 (t, 3H), 1.71 (s, 6H), 3.61 (s, 1H), 4.56 (q, 2H), 6.84 (d, 1H), 7.11–7.20 (m, 2H), 7.39 (t, 1H), 7.53–7.63 (m, 4H), 7.85 (d, 1H).
72	1.26 (s, 3H), 1.36 (s, 3H), 1.76 (s, 3H), 2.17 (s, 1H), 4.01 (s, 1H), 6.89 (d, 1H), 7.17–7.28 (m, 2H), 7.54 (t, 1H), 7.60–7.71 (m, 4H), 8.11 (d, 1H), 8.78 (d, 1H).
73	1.15 (t, 3H), 1.86 (s, 3H), 3.80 (s, 1H), 3.94–4.08 (m, 2H), 6.95 (d, 1H), 7.23–7.37 (m, 2H), 7.49–7.55 (m, 2H), 7.61–7.71 (m, 3H), 8.10 (d, 1H), 8.72 (d, 1H).
74	2.04 (s, 3H), 4.27 (s, 1H), 6.60 (d, 1H), 6.94–7.04 (m, 3H), 7.10–7.46 (m, 8H), 8.31 (d, 1H).
75	1.84 (d, 3H), 3.12 (d, 1H), 3.36 (d, 1H), 4.17 (d, 1H), 6.50 (d, 1H), 6.89–7.50 (m, 11H), 8.51 (d, 1H).
76	1.83 (d, 3H), 3.92 (d, 1H), 5.01 (d, 1H), 5.16 (d, 1H), 6.25 (dd, 1H), 6.69 (d, 1H), 6.95 (dd, 1H), 7.22 (m, 1H), 7.33 (m, 1H), 7.47–7.50 (m, 2H), 7.56 (d, 1H), 8.76 (d, 1H).
77	1.74 (d, 3H), 2.62 (m, 1H), 2.89 (m, 1H), 3.87 (d, 1H), 5.10 (d, 1H), 5.78 (m, 1H), 6.69 (d, 1H), 6.94 (dd, 1H), 7.19 (m, 1H), 7.33 (m, 1H), 7.48–4.50 (m, 2H), 7.58 (d, 1H), 8.82 (d, 1H).
78	1.84 (d, 3H), 3.97 (d, 1H), 5.03 (d, 1H), 5.17 (d, 1H), 5.71 (dd, 1H), 6.27 (dd, 1H), 6.58 (d, 1H), 6.76 (dd, 1H), 6.93 (dd, 1H), 7.16–7.42 (m, 5H), 7.46 (d, 1H).
79	1.73 (d, 3H), 2.62 (m, 1H), 2.86 (m, 1H), 3.89–3.95 (m, 3H), 5.07–5.20 (m, 4H), 5.82 (m, 1H), 6.18 (m, 1H), 6.64 (d, 1H), 6.94 (dd, 1H), 7.16–7.41 (m, 5H).
80	0.95 (d, 3H), 0.99 (d, 3H), 1.67 (d, 3H), 2.30 (m, 1H), 3.81 (d, 1H), 6.67 (d, 1H), 6.94 (m, 1H), 7.17 (m, 1H), 7.33 (m, 1H), 7.45–7.51 (m, 2H), 7.61 (d, 1H), 8.81 (d, 1H).
81	1.75 (s, 3H), 1.85 (d, 3H), 4.00 (d, 1H), 4.77 (s, 1H), 4.95 (s, 1H), 6.68 (d, 1H), 6.95 (m, 1H), 7.22 (m, 1H), 7.35 (m, 1H), 7.48–7.51 (m, 2H), 7.57 (d, 1H), 8.72 (d, 1H).
82	1.81 (d, 3H), 3.09 (d, 1H), 3.35 (d, 1H), 4.13 (s, 1H), 6.57 (d, 1H), 6.80–7.04 (m, 6H), 7.10–7.51 (m, 4H), 8.65 (d, 1H).
83	1.09 (d, 3H), 1.67–1.70 (m, 3H), 2.84 (t, 2/3H), 2.95 (t, 1/3H), 3.72 (d, 1/3H), 3.94 (d, 2/3H), 4.97–5.10 (m, 2H), 5.78–5.95 (m, 1H), 6.68 (d, 1H), 6.90–6.96 (m, 1H), 7.16–7.25 (m, 1H), 7.31–7.38 (m, 1H), 7.47–7.50 (m, 2H), 7.59–7.62 (m, 1H), 8.81–8.83 (m, 1H).
86	0.39–0.45 (m, 3H), 0.55 (m, 1H), 1.51 (dd, 1H), 1.73 (d, 3H), 3.74 (d, 1H), 6.68 (m, 1H), 6.95 (m, 1H), 7.22 (m, 1H), 7.35 (m, 1H), 7.47–7.50 (m, 2H), 7.59 (d, 1H), 8.82 (d, 1H).
87	0.40–0.51 (m, 3H), 0.59 (m, 1H), 1.09 (m, 2H), 1.38 (m, 2H), 1.55 (m, 1H), 1.76 (d, 3H), 2.54 (m, 1H), 4.08 (d, 1H), 6.62 (d, 1H), 6.91 (dd, 1H), 7.13–7.40 (m, 4H), 7.45 (s, 1H).
88	0.85 (t, 3H), 1.73 (d, 3H), 1.98 (m, 1H), 2.08 (m, 1H), 2.79 (s, 3H), 3.96 (d, 1H), 6.58 (d, 1H), 6.92 (m, 1H), 7.19 (m, 1H), 7.29–7.46 (m, 4H).
89	1.74 (d, 3H), 2.76 (s, 3H), 3.99 (d, 1H), 5.02 (d, 1H), 5.18 (d, 1H), 6.28 (m, 1H), 6.58 (d, 1H), 6.94 (m, 1H), 7.21 (m, 1H), 7.29–7.42 (m, 4H).
90	0.39–0.47 (m, 3H), 0.58 (m, 1H), 1.50 (m, 1H), 1.74 (d, 3H), 2.80 (s, 3H), 3.85 (d, 1H), 6.59 (d, 1H), 6.95 (m, 1H), 7.18 (m, 1H), 7.28–7.46 (m, 4H).

[0271]

Table 32

Compound No.	¹ H-NMR
91	1.83 (d, 3H), 2.63 (s, 3H), 3.06 (d, 1H), 3.30 (d, 1H), 3.67 (s, 3H), 4.21 (br, 1H), 6.43 (d, 1H), 6.68 (d, 2H), 6.89–6.96 (m, 3H), 7.16 (m, 1H), 7.27–7.41 (m, 4H).
92	1.82 (d, 3H), 2.64 (s, 3H), 3.10 (d, 1H), 3.34 (d, 1H), 4.25 (br, 1H), 6.45 (d, 1H), 6.81–7.02 (m, 6H), 7.17 (m, 1H), 7.26–7.42 (m, 3H).
94	1.78 (d, 6H), 2.74 (s, 3H), 2.81 (s, 3H), 4.23 (d, 1H), 6.52 (d, 1H), 6.86 (dd, 1H), 7.10 (m, 1H), 7.37 (t, 1H), 7.48–7.51 (m, 3H).
97	1.81 (s, 6H), 2.82 (s, 3H), 4.02 (s, 1H), 6.80 (dd, 1H), 7.17 (m, 1H), 7.26–7.32 (m, 3H), 7.37–7.40 (m, 2H).
101	1.68 (s, 6H), 2.40 (s, 3H), 6.82 (d, 1H), 7.07 (m, 1H), 7.2–7.6 (m, 5H), 8.86 (d, 1H).
103	1.82 (s, 6H), 4.12 (s, 1H), 6.86 (dd, 1H), 7.15 (m, 1H), 7.28 (m, 1H), 7.47–7.56 (m, 2H), 7.62–7.70 (m, 2H), 8.10 (d, 1H), 8.77 (d, 1H).
105	1.76 (d, 3H), 2.76 (s, 3H), 3.93 (d, 1H), 6.58 (d, 1H), 6.92 (m, 1H), 7.13–7.27 (m, 2H), 7.35–7.42 (m, 2H), 8.02 (m, 1H).
107	1.78 (d, 6H), 2.75 (s, 3H), 6.55 (m, 1H), 6.89 (m, 1H), 7.14 (m, 1H), 7.4–7.7 (m, 4H), 8.05 (dd, 1H).
108	1.82 (s, 6H), 2.77 (s, 3H), 4.18 (s, 1H), 6.77 (dd, 1H), 7.13 (m, 1H), 7.25–7.35 (m, 2H), 7.46 (m, 1H), 7.59–7.64 (m, 2H), 8.04 (d, 1H).
111	2.62 (s, 3H), 6.75 (d, 1H), 6.96 (t, 1H), 7.36 (m, 1H), 7.55 (t, 1H), 7.61 (d, 1H), 7.65–7.76 (m, 2H), 8.12 (d, 1H), 8.79 (d, 1H).
113	0.20 (s, 9H), 6.73 (d, 1H), 6.97 (m, 1H), 7.36 (m, 1H), 7.54–7.77 (m, 4H), 8.12 (d, 1H), 8.71 (d, 1H).
118	1.76 (d, 3H), 2.74 (s, 3H), 3.92 (d, 1H), 6.55 (m, 1H), 6.91 (m, 1H), 7.17 (m, 1H), 7.41–7.51 (m, 2H), 7.57 (d, 1H), 8.04 (d, 1H).
119	1.77 (d, 6H), 3.88 (d, 1H), 6.63 (m, 1H), 7.04 (m, 1H), 7.53–7.59 (m, 2H), 7.65–7.73 (m, 2H), 8.12 (d, 1H), 8.79 (d, 1H).
126	1.77 (d, 6H), 3.97 (d, 1H), 6.64 (m, 1H), 6.91 (m, 1H), 7.17 (dt, 1H), 7.36 (dt, 1H), 7.63 (d, 1H), 7.68–7.77 (m, 2H), 8.80 (d, 1H).
131	2.17–2.25 (m, 2H), 2.83–2.93 (m, 1H), 3.16–3.27 (m, 1H), 3.33 (s, 3H), 4.88 (t, 1H), 6.84 (d, 1H), 7.14 (d, 1H), 7.30 (d, 1H), 7.48–7.54 (m, 2H), 7.59–7.68 (m, 2H), 8.10 (d, 1H), 8.85 (d, 1H).
132	1.44 (d, 3H), 3.25 (s, 3H), 4.73 (q, 1H), 6.96–6.99 (m, 1H), 7.26–7.33 (m, 2H), 7.41 (d, 1H), 7.51 (t, 1H), 7.58–7.66 (m, 3H), 8.10 (d, 1H), 8.82 (d, 1H).
133	0.99 (t, 3H), 2.16–2.25 (m, 1H), 2.88–2.92 (m, 1H), 3.16–3.25 (m, 1H), 3.35–3.43 (m, 1H), 3.48–3.56 (m, 1H), 4.97 (dd, 1H), 6.87 (d, 1H), 7.13 (d, 1H), 7.29 (t, 1H), 7.46–7.52 (m, 2H), 7.58–7.67 (m, 2H), 8.09 (d, 1H), 8.84 (d, 1H).
134	0.74 (t, 3H), 1.35–1.43 (m, 2H), 2.17–2.25 (m, 2H), 2.89 (m, 1H), 3.22 (m, 1H), 3.29 (m, 1H), 3.44 (m, 1H), 4.98 (m, 1H), 6.85 (d, 1H), 7.13 (d, 1H), 7.29 (t, 1H), 7.46–7.52 (m, 2H), 7.58–7.67 (m, 2H), 8.10 (d, 1H), 8.84 (d, 1H).

[0272]

Table 33

Compound No.	¹ H-NMR
135	0.87 (t, 3H), 1.43 (d, 3H), 1.54 (m, 2H), 3.28 (m, 2H), 4.81 (m, 1H), 6.95 (m, 1H), 7.26–7.30 (m, 2H), 7.39 (d, 1H), 7.51 (m, 1H), 7.59–7.66 (m, 3H), 8.10 (d, 1H), 8.82 (d, 1H).
136	1.00 (s, 9H), 1.99 (br, 1H), 4.89 (d, 1H), 6.91 (d, 1H), 7.23 (m, 1H), 7.48 (d, 1H), 7.53 (d, 1H), 7.60–7.68 (m, 4H), 8.10 (d, 1H), 8.80 (d, 1H).
137	2.11 (m, 1H), 2.49 (m, 1H), 2.85 (br, 1H), 2.91 (m, 1H), 3.20 (m, 1H), 5.50 (m, 1H), 6.76 (d, 1H), 7.11 (d, 1H), 7.26 (t, 1H), 7.52 (t, 1H), 7.55–7.71 (m, 3H), 8.13 (d, 1H), 8.82 (s, 1H).
140	1.60 (d, 3H), 3.24 (s, 3H), 4.86 (q, 1H), 6.78 (d, 1H), 7.01 (t, 1H), 7.25–7.35 (m, 2H), 7.42–7.50 (m, 3H), 8.83 (d, 1H).
142	1.61 (d, 3H), 3.18 (s, 3H), 5.03 (q, 1H), 6.94 (dd, 1H), 7.15 (t, 1H), 7.27–7.37 (m, 2H), 7.41–7.49 (m, 2H), 8.81 (d, 1H).
143	0.44–0.67 (m, 4H), 1.62 (m, 1H), 2.86 (d, 1H), 4.57 (t, 1H), 6.84 (dd, 1H), 7.20 (t, 1H), 7.26 (m, 1H), 7.34 (m, 1H), 7.47–7.50 (m, 2H), 7.61 (d, 1H), 8.86 (d, 1H).
145	1.10 (s, 9H), 3.32 (d, 1H), 5.18 (d, 1H), 6.76 (dd, 1H), 7.14 (t, 1H), 7.24 (m, 1H), 7.37 (m, 1H), 7.47–7.53 (m, 2H), 7.71 (d, 1H), 8.83 (d, 1H).
146	1.65 (d, 3H), 2.99 (d, 1H), 5.50 (m, 1H), 6.81 (d, 1H), 7.15–7.27 (m, 2H), 7.36 (m, 1H), 7.48–7.50 (m, 2H), 7.60 (m, 1H), 8.86 (d, 1H).
147	1.62 (d, 3H), 3.20 (s, 3H), 5.07 (q, 1H), 6.91 (dd, 1H), 7.22–7.35 (m, 3H), 7.41–7.49 (m, 3H), 8.81 (d, 1H).
148	1.65 (d, 3H), 3.00 (d, 1H), 5.47 (m, 1H), 6.84 (d, 1H), 7.10 (t, 1H), 7.30–7.51 (m, 4H), 7.61 (d, 1H), 8.86 (d, 1H).
149	1.61 (d, 3H), 3.19 (s, 3H), 5.06 (q, 1H), 6.96 (d, 1H), 7.17 (t, 1H), 7.28 (m, 1H), 7.43–7.51 (m, 4H), 8.81 (d, 1H).
150	1.50 (d, 3H), 2.54 (s, 3H), 3.16 (s, 3H), 4.92 (q, 1H), 6.84 (d, 1H), 7.08 (d, 1H), 7.21 (t, 1H), 7.26–7.34 (m, 2H), 7.38–7.46 (m, 2H), 8.84 (d, 1H).
152	1.61 (d, 3H), 2.83 (s, 3H), 3.16 (s, 3H), 5.05 (q, 1H), 6.85 (dd, 1H), 7.21–7.32 (m, 4H), 7.34–7.40 (m, 2H).
153	1.63 (d, 3H), 3.29 (s, 3H), 4.97 (q, 1H), 7.18 (d, 1H), 7.32–7.51 (m, 5H), 7.61 (d, 1H), 8.83 (d, 1H).
154	1.62 (d, 3H), 3.22 (s, 3H), 4.09 (s, 3H), 5.08 (q, 1H), 6.91 (d, 1H), 6.97 (m, 1H), 7.13 (t, 1H), 7.25 (m, 1H), 7.41–7.47 (m, 3H), 8.76 (d, 1H).
155	1.58 (d, 3H), 3.20 (s, 3H), 3.90 (s, 3H), 4.09 (s, 3H), 5.00 (q, 1H), 6.58 (d, 1H), 6.78 (d, 1H), 6.95 (d, 1H), 7.21–7.28 (m, 2H), 7.39–7.4 (m, 2H), 8.78 (br, 1H).
156	1.63 (d, 3H), 3.82 (d, 2H), 4.98 (dd, 1H), 5.15 (dd, 1H), 5.21 (q, 1H), 5.70 (m, 1H), 6.92 (dd, 1H), 7.24–7.33 (m, 3H), 7.39–7.49 (m, 3H), 8.82 (d, 1H).
157	1.02 (t, 3H), 1.92–2.08 (m, 2H), 2.84 (d, 1H), 5.23 (m, 1H), 6.80 (dd, 1H), 7.18 (t, 1H), 7.24 (m, 1H), 7.35 (m, 1H), 7.45–7.51 (m, 2H), 7.60 (m, 1H), 8.84 (d, 1H).

[0273]

Table 34

Compound No.	¹ H-NMR
160	1.65 (d, 3H), 4.33 (d, 1H), 4.38 (d, 1H), 5.25 (d, 1H), 6.93 (dd, 1H), 7.13 (m, 5H), 7.25–7.42 (m, 6H), 8.79 (d, 1H).
161	6.56 (d, 1H), 6.78 (m, 1H), 7.2–7.5 (m, 11H), 8.50 (d, 1H).
163	0.85 (t, 3H), 1.2–1.4 (m, 4H), 1.9–2.1 (m, 2H), 2.90 (d, 1H), 5.31 (m, 1H), 6.80 (dd, 1H), 7.1–7.6 (m, 6H), 8.83 (d, 1H).
164	2.25 (m, 1H), 3.53 (d, 1H), 6.07 (dd, 1H), 6.86 (m, 1H), 7.2–7.7 (m, 6H), 8.85 (d, 1H).
165	0.13 (s, 9H), 6.45 (s, 1H), 6.94 (d, 1H), 7.29–7.38 (m, 3H), 7.48–7.50 (m, 2H), 7.59 (m, 1H), 8.86 (d, 1H).
167	1.62 (d, 3H), 3.20 (s, 3H), 5.08 (q, 1H), 6.91 (dd, 1H), 7.21–7.34 (m, 3H), 7.42–7.47 (m, 3H), 8.81 (d, 1H).
168	3.51 (s, 3H), 3.65 (s, 3H), 5.44 (s, 1H), 6.89 (m, 1H), 7.2–7.6 (m, 6H), 8.78 (d, 1H).
170	0.02 (s, 9H), 2.26 (s, 3H), 5.52 (s, 1H), 6.86 (dd, 1H), 7.2–7.6 (m, 6H), 8.75 (d, 1H).
171	2.22 (s, 3H), 4.13 (d, 1H), 5.66 (d, 1H), 6.91 (m, 1H), 7.3–7.6 (m, 6H), 8.76 (d, 1H).
172	2.34 (s, 3H), 3.46 (s, 3H), 5.28 (s, 1H), 6.9 (m, 1H), 7.2–7.26 (m, 6H), 8.81 (d, 1H).
173	1.26 (s, 3H), 1.45 (s, 3H), 3.67 (d, 1H), 5.27 (d, 1H), 6.83 (d, 1H), 7.1–7.5 (m, 6H), 7.66 (bs, 1H), 8.82 (d, 1H).
174	2.97 (d, 1H), 3.31 (m, 2H), 5.59 (m, 1H), 6.75 (d, 1H), 7.13–7.49 (m, 11H), 8.77 (br, 1H).
175	3.19 (s, 3H), 3.28–3.44 (m, 2H), 5.18 (t, 1H), 6.82 (d, 1H), 7.11–7.49 (m, 11H), 8.67 (br, 1H).
177	1.60 (d, 3H), 3.17 (s, 3H), 5.04 (q, 1H), 6.97 (dd, 1H), 7.13 (t, 1H), 7.26–7.36 (m, 2H), 7.42–7.49 (m, 2H), 8.81 (d, 1H).
180	1.23 (s, 3H), 1.40 (s, 3H), 3.20 (s, 3H), 4.87 (s, 1H), 6.94 (d, 1H), 7.26–7.35 (m, 3H), 7.45–7.48 (m, 3H), 8.75 (d, 1H).
181	3.59 (d, 1H), 3.70 (s, 3H), 5.79 (d, 1H), 6.8–6.95 (m, 1H), 7.2–7.6 (m, 6H), 8.77 (d, 1H).
185	1.71 (t, 3H), 3.06 (m, 1H), 5.34 (m, 1H), 7.08 (d, 1H), 7.32–7.41 (m, 2H), 7.49–7.52 (m, 3H), 7.67 (dd, 1H), 8.86 (d, 1H).
186	1.67 (d, 3H), 3.16 (s, 3H), 4.85 (q, 1H), 7.21 (d, 1H), 7.32 (m, 1H), 7.41–7.47 (m, 4H), 7.58 (d, 1H), 8.80 (d, 1H).
188	1.63 (d, 3H), 3.23 (s, 3H), 5.10 (q, 1H), 6.89 (dd, 1H), 7.18–7.29 (m, 2H), 7.45 (d, 1H), 7.52 (t, 1H), 7.60–7.69 (m, 2H), 8.10 (d, 1H), 8.78 (d, 1H).
189	1.48 (d, 3H), 1.97 (d, 1H), 5.17 (m, 1H), 7.15 (m, 1H), 7.24–7.35 (m, 2H), 7.45–7.52 (m, 2H), 7.58–7.63 (m, 2H), 8.09 (d, 1H), 8.87 (d, 1H).
190	1.61 (d, 3H), 2.69 (m, 1H), 5.43 (m, 1H), 7.09 (t, 1H), 7.27–7.34 (m, 2H), 7.52 (t, 1H), 7.60–7.67 (m, 2H), 8.10 (d, 1H), 8.87 (d, 1H).
198	1.61 (d, 3H), 2.83 (s, 3H), 3.16 (s, 3H), 5.04 (q, 1H), 6.84 (dd, 1H), 7.19–7.36 (m, 5H).

[0274]

Table 35

Compound No.	¹ H-NMR
203	1.62 (d, 3H), 3.22 (s, 3H), 5.09 (q, 1H), 6.87 (dd, 1H), 7.1–7.4 (m, 3H), 7.46 (d, 1H), 7.6–7.8 (m, 2H), 8.78 (d, 1H).
204	1.78 (s, 3H), 3.74 (s, 3H), 4.14 (d, 1H), 5.85 (d, 1H), 6.93 (m, 1H), 7.2–7.5 (m, 6H), 8.78 (d, 1H).
206	1.77 (d, 1H), 2.71 (s, 3H), 5.52 (m, 1H), 6.24 (dd, 1H), 6.83 (m, 1H), 7.07 (m, 1H), 7.42–7.77 (m, 2H), 7.80 (m, 1H).
207	1.52 (dd, 3H), 2.29 (s, 6H), 3.95 (q, 1H), 6.78 (d, 1H), 6.97 (m, 1H), 7.23–7.34 (m, 2H), 7.43–7.47 (m, 3H), 8.84 (d, 1H).
209	1.60 (d, 3H), 3.19 (s, 3H), 3.92 (s, 3H), 4.82 (q, 1H), 7.0–7.1 (m, 1H), 7.2–7.5 (m, 6H), 8.84 (d, 1H).
210	1.59 (d, 3H), 2.59 (s, 3H), 3.23 (s, 3H), 4.76 (q, 1H), 6.99 (dd, 1H), 7.11 (dd, 1H), 7.2–7.5 (m, 5H), 8.85 (d, 1H).
211	1.60 (d, 3H), 2.51 (s, 3H), 3.23 (s, 3H), 5.04 (q, 1H), 6.78 (dd, 1H), 7.12 (d, 1H), 7.2–7.5 (m, 5H), 8.83 (d, 1H).
212	1.74 (d, 3H), 3.24 (s, 3H), 3.26 (s, 3H), 5.48 (q, 1H), 7.2–7.5 (m, 5H), 8.03 (d, 1H), 8.81 (d, 1H).
215	1.60 (d, 3H), 3.02 (s, 3H), 4.48 (q, 1H), 7.02 (dd, 1H), 7.12 (dd, 1H), 7.2–7.55 (m, 10H), 8.85 (d, 1H).
217	3.42 (s, 6H), 5.65 (s, 1H), 6.79 (d, 1H), 7.01 (t, 1H), 7.29–7.37 (m, 2H), 7.45–7.48 (m, 3H), 8.86 (d, 1H).
218	1.11 (t, 6H), 3.49 (m, 2H), 3.74 (m, 2H), 5.78 (s, 1H), 6.80 (d, 1H), 7.01 (t, 1H), 7.27–7.36 (m, 2H), 7.43–7.49 (m, 3H), 8.86 (d, 1H).
220	3.35 (s, 3H), 4.66 (s, 2H), 6.92 (dd, 1H), 7.29–7.35 (m, 3H), 7.44–7.46 (m, 3H), 8.88 (d, 1H).
221	3.99 (m, 2H), 4.70 (s, 2H), 5.08 (d, 1H), 5.17 (d, 1H), 5.78 (m, 1H), 6.93 (dd, 1H), 7.28–7.35 (m, 3H), 7.44–7.49 (m, 3H), 8.88 (d, 1H).
223	3.36 (s, 3H), 4.66 (s, 2H), 6.96 (d, 1H), 7.22 (t, 1H), 7.33 (m, 1H), 7.45–7.52 (m, 3H), 8.88 (d, 1H).
224	0.83 (t, 3H), 1.30 (m, 2H), 1.40 (m, 2H), 2.60 (t, 2H), 3.99 (s, 2H), 6.90 (d, 1H), 7.20–7.37 (m, 3H), 7.44–7.50 (m, 3H), 8.87 (d, 1H).
225	0.99 (s, 6H), 1.48 (s, 6H), 1.63 (d, 1H), 3.57 (d, 1H), 6.93 (d, 1H), 7.14–7.26 (m, 2H), 7.44–7.71 (m, 5H), 8.09 (d, 1H), 8.77 (br, 1H).
226	0.95 (s, 6H), 1.42 (s, 6H), 2.43 (s, 3H), 4.01 (s, 2H), 6.87 (d, 1H), 7.11 (m, 1H), 7.20 (m, 1H), 7.28 (d, 2H), 7.40–7.54 (m, 3H), 7.59–7.74 (m, 4H), 8.10 (d, 1H), 8.68 (d, 1H).
227	1.23 (s, 6H), 1.52 (s, 6H), 2.49 (s, 1H), 6.94 (d, 1H), 7.18–7.26 (m, 2H), 7.49–7.68 (m, 5H), 8.10 (d, 1H), 8.77 (d, 1H).
230	0.87 (s, 3H), 1.04 (s, 3H), 1.22 (d, 3H), 1.52 (d, 6H), 1.68 (d, 1H), 4.08 (m, 1H), 6.94 (m, 1H), 7.15–7.26 (m, 2H), 7.46–7.54 (m, 2H), 7.59–7.68 (m, 3H), 8.10 (d, 1H), 8.78 (d, 1H).
232	1.38 (s, 6H), 1.65 (s, 6H), 6.95 (m, 1H), 7.19–7.31 (m, 2H), 7.47 (d, 1H), 7.52 (m, 1H), 7.60–7.69 (m, 3H), 8.10 (d, 1H), 8.75 (d, 1H).

[0275]

Table 36

Compound No.	¹ H-NMR
234	1.49 (s, 6H), 6.92 (m, 1H), 7.19–7.33 (m, 2H), 7.48–7.71 (m, 5H), 8.02 (d, 1H), 8.72 (d, 1H), 9.72 (s, 1H)
235	1.05 (s, 6H), 1.57 (s, 6H), 3.64 (d, 1H), 6.73 (d, 1H), 6.87 (m, 1H), 7.15 (m, 1H), 7.81–7.55 (m, 2H), 7.61–7.73 (m, 2H), 8.10 (d, 1H), 8.75 (d, 1H)
243	1.08 (s, 6H), 1.52 (s, 6H), 2.26 (s, 2H), 6.88 (m, 1H), 7.17 (m, 1H), 7.36 (s, 2H), 7.50–7.70 (m, 4H), 8.81 (d, 1H), 8.91 (d, 1H)
244	1.02 (br, 1H), 1.46 (s, 6H), 2.19 (t, 2H), 3.56 (m, 2H), 6.90 (d, 1H), 7.13–7.24 (m, 2H), 7.43 (d, 1H), 7.49–7.54 (m, 2H), 7.60–7.70 (m, 2H), 8.10 (d, 1H), 8.81 (d, 1H)
246	1.06 (d, 3H), 1.50 (d, 6H), 1.96 (dd, 1H), 2.15 (dd, 1H), 3.83 (m, 1H), 6.89 (dd, 1H), 7.13–7.26 (m, 2H), 7.46–7.55 (m, 2H), 7.59–7.70 (m, 3H), 8.11 (d, 1H), 8.82 (d, 1H)
247	1.47 (s, 6H), 2.21 (d, 2H), 3.16 (s, 6H), 4.19 (t, 1H), 6.87 (dd, 1H), 7.13–7.20 (m, 2H), 7.44 (m, 1H), 7.49 (t, 1H), 7.54–7.70 (m, 3H), 8.10 (d, 1H), 8.81 (d, 1H)
248	0.89 (s, 6H), 1.49 (s, 6H), 3.43 (s, 6H), 3.95 (s, 1H), 6.93 (d, 1H), 7.12–7.24 (m, 2H), 7.42 (d, 1H), 7.48–7.67 (m, 4H), 8.09 (d, 1H), 8.78 (d, 1H)
251	1.26 (s, 6H), 1.56 (s, 6H), 2.48 (s, 1H), 2.79 (s, 1H), 6.88 (m, 1H), 7.19–7.30 (m, 3H), 7.43 (t, 1H), 7.56–7.61 (m, 3H), 8.02 (d, 1H)
252	0.97 (t, 1H), 1.47 (s, 6H), 2.19 (t, 2H), 2.84 (s, 3H), 3.57 (m, 1H), 6.83 (dd, 1H), 7.11–7.23 (m, 2H), 7.37 (s, 1H), 7.41–7.46 (m, 2H), 7.56–7.62 (m, 2H), 8.02 (d, 1H)
253	1.09 (d, 3H), 1.40 (s, 3H), 1.45 (s, 3H), 1.72 (bs, 1H), 4.54 (m, 1H), 6.89 (dd, 1H), 7.1–7.3 (m, 2H), 7.4–7.7 (m, 5H), 8.09 (d, 1H), 8.77 (d, 1H)
254	0.91 (t, 3H), 1.2–1.5 (m, 8H), 1.6–1.7 (m, 1H), 4.1–4.2 (m, 1H), 6.90 (dd, 1H), 7.1–7.3 (m, 2H), 7.4–7.7 (m, 5H), 8.10 (d, 1H), 8.77 (d, 1H)
255	1.43 (s, 3H), 1.49 (s, 3H), 1.57 (s, 3H), 1.75 (d, 1H), 4.8–5.0 (m, 3H), 6.91 (dd, 1H), 7.1–7.3 (m, 2H), 7.4–7.7 (m, 5H), 8.10 (d, 1H), 8.81 (d, 1H)
256	0.89 (s, 3H), 1.05–1.10 (m, 6H), 1.23 (br, 6H), 2.81 (s, 3H), 4.09 (m, 1H), 6.88 (d, 1H), 7.18–7.26 (m, 3H), 7.42 (t, 1H), 7.55–7.64 (m, 3H), 8.01 (d, 1H)
257	1.24 (s, 6H), 1.53 (br, 6H), 1.88 (s, 3H), 2.83 (s, 3H), 6.87 (d, 1H), 7.13–7.26 (m, 3H), 7.41–7.46 (m, 2H), 7.56–7.60 (m, 2H), 8.03 (d, 1H)
258	1.11 (br, 1H), 1.45 (s, 6H), 2.37 (t, 2H), 3.55 (m, 2H), 6.90 (m, 1H), 7.15–7.34 (m, 3H), 7.42–7.46 (m, 3H), 7.53 (d, 1H), 8.84 (d, 1H)
259	0.98 (br, 1H), 1.46 (s, 6H), 2.17 (t, 2H), 2.86 (s, 3H), 3.57 (m, 2H), 6.86 (d, 1H), 7.15–7.30 (m, 3H), 7.36–7.37 (m, 3H), 7.45 (m, 1H)
260	1.51 (s, 6H), 2.11 (s, 3H), 6.91 (dd, 1H), 7.2–7.3 (m, 2H), 7.5–7.7 (m, 5H), 8.11 (d, 1H), 8.71 (d, 1H)
261	1.47 (s, 6H), 2.28 (t, 2H), 2.84 (s, 3H), 3.66 (s, 3H), 4.04 (t, 2H), 6.84 (m, 1H), 7.12–7.28 (m, 3H), 7.30–7.45 (m, 4H)

[0276]

Table 37

Compound No.	¹ H-NMR
262	1.59 (s, 6H), 2.18 (s, 3H), 6.72 (m, 1H), 6.96 (m, 1H), 7.2–7.6 (m, 5H), 8.73 (d, 1H).
263	1.3 (s, 6H), 1.63 (s, 6H), 6.74 (d, 1H), 6.91 (m, 1H), 7.18 (m, 1H), 7.5–7.8 (m, 4H), 8.10 (d, 1H), 8.75 (d, 1H).
265	1.60 (s, 6H), 2.18 (s, 3H), 6.70 (d, 1H), 6.92 (m, 1H), 7.20 (m, 1H), 7.5–7.8 (m, 2H), 8.10 (d, 1H), 8.69 (d, 1H).
266	1.59 (d, 6H), 2.18 (s, 3H), 6.68 (d, 2H), 6.92 (dd, 1H), 6.96–7.24 (m, 1H), 7.31–7.37 (m, 1H), 7.55 (d, 1H), 7.67–7.75 (m, 2H), 8.69 (d, 1H).
267	1.59 (d, 6H), 2.18 (s, 3H), 6.70 (d, 1H), 6.96 (dd, 1H), 7.19–7.25 (m, 1H), 7.40–7.49 (m, 2H), 7.53 (d, 1H), 8.75 (d, 1H).
268	0.87 (d, 3H), 1.39 (s, 3H), 1.44 (s, 3H), 1.85 (s, 1H), 4.50 (m, 1H), 6.91 (d, 1H), 7.16–7.33 (m, 3H), 7.43–7.52 (m, 3H), 8.81 (d, 1H).
269	1.51 (s, 6H), 2.14 (s, 3H), 6.92 (d, 1H), 7.22–7.36 (m, 3H), 7.44–7.48 (m, 2H), 7.50–7.58 (m, 2H), 8.73 (d, 1H).
270	1.88 (s, 6H), 2.86 (s, 3H), 6.88 (d, 1H), 7.18–7.43 (m, 5H), 7.50–7.56 (m, 2H).
271	1.89 (s, 6H), 6.93 (d, 1H), 7.20 (m, 1H), 7.31 (m, 1H), 7.54 (t, 1H), 7.63–7.73 (m, 2H), 8.12 (d, 1H), 8.83 (d, 1H).
277	0.82 (t, 3H), 1.20 (t, 3H), 1.64 (m, 1H), 3.04 (m, 1H), 6.98 (m, 1H), 7.19–7.26 (m, 2H), 7.32–7.39 (m, 2H), 7.49 (t, 1H), 7.52–7.65 (m, 2H), 8.10 (d, 1H), 8.83 (br, 1H).
278	0.79 (d, 3H), 0.94 (d, 3H), 1.21 (d, 3H), 1.78–1.90 (m, 1H), 2.80–2.90 (m, 1H), 6.94–6.99 (m, 1H), 7.18–7.4 (m, 2H), 7.35 (d, 1H), 7.49 (t, 1H), 7.58–7.65 (m, 3H), 8.09 (d, 1H), 8.93 (br, 1H).
280	2.24 (s, 3H), 2.72 (s, 3H), 4.13 (d, 1H), 5.51 (d, 1H), 6.69 (d, 1H), 7.02 (t, 1H), 7.2–7.6 (m, 5H).
286	1.73–1.82 (m, 4H), 2.66–2.70 (m, 2H), 2.82–2.84 (m, 2H), 6.82 (d, 1H), 6.98 (d, 1H), 7.13 (t, 1H), 7.31 (d, 1H), 7.48–7.76 (m, 4H), 8.07–8.14 (m, 2H), 8.83 (d, 1H).
289	1.26 (t, 3H), 4.25 (d, 2H), 6.73 (d, 1H), 7.01 (t, 1H), 7.41 (m, 1H), 7.56 (t, 1H), 7.63 (d, 1H), 7.66–7.75 (m, 2H), 8.12 (d, 1H), 8.74 (d, 1H).
295	3.13 (s, 3H), 3.63 (s, 3H), 6.76 (d, 1H), 7.03 (m, 1H), 7.31–7.52 (m, 4H), 7.60 (dd, 1H), 8.78 (d, 1H).
297	1.01 (m, 2H), 1.31 (m, 2H), 4.53 (s, 2H), 6.79 (d, 1H), 6.98 (t, 1H), 7.08–7.11 (m, 2H), 7.16–7.18 (m, 3H), 7.28–7.49 (m, 5H), 8.87 (d, 1H).
308	1.66 (s, 3H), 2.93 (d, 1H), 2.99 (d, 1H), 6.74 (d, 1H), 6.95 (t, 1H), 7.26–7.36 (m, 2H), 7.46–7.48 (m, 2H), 7.54 (d, 1H), 8.85 (d, 1H).
328	3.45 (s, 6H), 4.54 (s, 2H), 5.93 (s, 1H), 7.04 (d, 1H), 7.17 (t, 1H), 7.27 (m, 1H), 7.31 (m, 1H), 7.43 (m, 1H), 7.50 (m, 1H), 7.85 (s, 1H), 8.86 (d, 1H).
329	1.27 (t, 3H), 4.26 (d, 2H), 4.60 (s, 2H), 7.27–7.52 (m, 6H), 7.81 (s, 1H), 7.99 (d, 1H), 8.88 (d, 1H).
330	1.18 (t, 3H), 4.24 (d, 2H), 4.30 (s, 2H), 7.01–7.07 (m, 2H), 7.35 (m, 1H), 7.51 (t, 1H), 7.67 (t, 1H), 7.72 (d, 1H), 7.85 (s, 1H), 8.07 (d, 1H), 8.78 (d, 1H).

[0277]

Table 38

Compound No.	¹ H-NMR
331	1.15 (t, 3H), 4.09 (q, 2H), 4.28 (s, 2H), 7.02–7.07 (m, 2H), 7.27–7.40 (m, 2H), 7.68–7.83 (m, 2H), 8.16 (d, 1H), 8.78 (d, 1H).
334	5.28–5.61 (m, 2H), 5.83–6.12 (m, 2H), 7.18–7.42 (m, 4H), 7.54–8.10 (m, 4H), 8.47 (dd, 1H), 9.02 (d, 1H).
339	1.55 (d, 3H), 1.68 (d, 3H), 6.14 (s, 1H), 7.32–7.41 (m, 2H), 7.51–7.63 (m, 3H), 7.81–7.88 (m, 2H), 8.16 (d, 1H), 8.44 (d, 1H), 9.24 (d, 1H).
340	7.52–7.67 (m, 5H), 7.83–7.88 (m, 2H), 7.96 (m, 1H), 8.08 (d, 1H), 8.18–8.21 (m, 2H), 8.53 (d, 1H), 9.42 (d, 1H).
343	5.94 (dd, 1H), 6.22 (dd, 1H), 6.89 (dd, 1H), 7.54–7.87 (m, 7H), 8.14 (m, 1H), 8.47 (d, 1H), 9.28 (d, 1H).
344	1.66 (s, 3H), 3.02 (s, 2H), 4.50 (m, 1H), 4.70 (m, 1H), 7.31–7.71 (m, 7H), 8.50 (m, 1H), 9.33 (d, 1H).
345	3.07 (t, 2H), 3.58 (t, 2H), 7.35–7.71 (m, 5H), 7.84–7.91 (m, 2H), 8.19 (d, 1H), 8.50 (d, 1H), 9.36 (d, 1H).
347	1.64 (s, 3H), 3.53 (s, 2H), 4.40 (m, 1H), 4.64 (m, 1H), 7.17 (dd, 1H), 7.26–7.42 (m, 2H), 7.63 (m, 1H), 7.82–7.90 (m, 2H), 8.18 (m, 1H), 8.46 (d, 1H), 9.30 (d, 1H).
350	1.67 (d, 3H), 1.75 (d, 3H), 2.60 (bs, 1H), 5.04 (bs, 1H), 6.54 (s, 1H), 7.02–7.10 (m, 2H), 7.18–7.55 (m, 5H), 8.04 (s, 1H), 8.85 (d, 1H).
351	3.94–4.00 (m, 2H), 5.12–5.26 (m, 3H), 5.75 (d, 1H), 6.09 (m, 1H), 6.49 (dd, 1H), 7.10 (m, 1H), 7.38–7.53 (m, 3H), 7.69–7.76 (m, 3H), 8.00 (d, 1H), 8.12 (d, 1H), 9.48 (d, 1H).
352	1.17 (t, 3H), 4.20 (q, 2H), 6.94 (d, 1H), 7.31 (t, 1H), 7.55–7.61 (m, 2H), 7.67–7.77 (m, 3H), 8.04 (d, 1H), 8.14 (d, 1H), 8.80 (br, 1H).
353	2.61 (s, 3H), 6.90 (d, 1H), 7.28–7.38 (m, 3H), 7.45–7.51 (m, 2H), 7.58 (m, 1H), 8.86 (d, 1H).
355	2.61 (s, 3H), 6.93 (d, 1H), 7.23–7.37 (m, 2H), 7.43–7.49 (m, 3H), 7.57 (s, 1H), 8.82 (br, 1H).
359	2.61 (s, 3H), 2.74 (s, 3H), 6.78 (dd, 1H), 7.24–7.46 (m, 5H), 7.46 (s, 1H).
360	1.24 (d, 6H), 3.16 (sep, 1H), 6.88 (d, 1H), 7.2–7.6 (m, 6H), 8.81 (d, 1H).
361	2.60 (s, 3H), 6.93 (dd, 1H), 7.21 (t, 1H), 7.35 (m, 1H), 7.47–7.51 (m, 3H), 8.83 (d, 1H).
363	1.06 (m, 2H), 1.25 (m, 2H), 2.29 (m, 1H), 6.96 (dd, 1H), 7.29–7.38 (m, 3H), 7.43–7.53 (m, 3H), 8.83 (d, 1H).
364	2.00 (s, 3H), 5.79 (s, 1H), 6.06 (s, 1H), 6.90 (d, 1H), 7.26–7.37 (m, 3H), 7.45–7.51 (m, 2H), 7.57 (m, 1H), 8.78 (br, 1H).
365	1.33 (s, 9H), 6.90 (dd, 1H), 7.26–7.37 (m, 3H), 7.46–7.51 (m, 2H), 7.54 (m, 1H), 8.81 (d, 1H).
367	2.62 (s, 3H), 2.65 (s, 3H), 2.80 (s, 3H), 6.59 (d, 1H), 6.92 (t, 1H), 7.26–7.39 (m, 2H), 7.48–7.51 (m, 3H).
368	6.95 (d, 1H), 7.2–7.7 (m, 10H), 7.87 (m, 1H), 8.64 (d, 1H).
371	0.91 (s, 3H), 1.38 (m, 2H), 1.70 (m, 2H), 2.89 (t, 2H), 6.88 (d, 1H), 7.2–7.6 (m, 6H), 8.82 (d, 1H).
373	6.66 (s, 1H), 6.76 (d, 1H), 7.04 (t, 1H), 7.36–7.56 (m, 4H), 7.73 (d, 1H), 8.84 (d, 1H).

[0278]

Table 39

Compound No.	¹ H-NMR
374	6.79 (d, 1H), 7.04 (t, 1H), 7.35-7.56 (m, 4H), 7.74 (s, 1H), 8.82 (d, 1H).
375	2.46 (s, 3H), 6.86 (d, 1H), 7.2-7.7 (m, 6H), 8.80 (d, 1H).
376	1.57 (s, 6H), 2.7 (bs, 1H), 6.89 (dd, 1H), 7.2-7.7 (m, 6H), 8.82 (d, 1H).
377	1.24 (d, 6H), 3.16 (sep, 1H), 6.88 (d, 1H), 7.2-7.6 (m, 6H), 8.81 (d, 1H).
382	2.59 (s, 3H), 2.69 (s, 3H), 6.69 (d, 1H), 6.99 (t, 1H), 7.25 (m, 1H), 7.33-7.41 (m, 3H), 8.00 (m, 1H).
383	2.61 (s, 3H), 2.69 (s, 3H), 6.62 (d, 1H), 6.97 (t, 1H), 7.24-7.37 (m, 2H), 7.48 (s, 1H), 7.61-7.69 (m, 2H).
384	2.15 (s, 3H), 5.11 (s, 2H), 6.78 (d, 1H), 7.03 (t, 1H), 7.33-7.53 (m, 4H), 7.64 (d, 1H), 8.84 (d, 1H).
388	2.61 (s, 3H), 6.80 (d, 1H), 7.02 (m, 1H), 7.22 (m, 1H), 7.39 (m, 1H), 7.58 (m, 1H), 7.79 (d, 1H), 7.92 (d, 1H), 8.81 (d, 1H).
389	2.60 (s, 3H), 2.69 (s, 3H), 6.67 (d, 1H), 6.98 (t, 1H), 7.32-7.45 (m, 3H), 7.58 (d, 1H), 8.03 (s, 1H).
390	6.76 (d, 1H), 7.04 (t, 1H), 7.40 (m, 1H), 7.47-7.57 (m, 3H), 7.75 (s, 1H), 8.82 (d, 1H).
397	1.53 (bs, 6H), 2.73 (s, 3H), 6.67 (d, 1H), 6.99 (t, 1H), 7.2-7.7 (m, 5H).
398	1.53 (d, 6H), 6.98 (t, 1H), 7.3-7.7 (m, 5H), 8.81 (d, 1H).
401	6.97 (d, 1H), 7.10 (t, 1H), 7.26-7.38 (m, 3H), 7.46-7.75 (m, 7H), 8.14 (d, 1H), 8.35 (d, 1H), 8.91 (d, 1H), 9.36 (s, 1H).
402	3.94 (s, 3H), 6.87 (d, 1H), 7.24-7.35 (m, 2H), 7.56 (t, 1H), 7.61-7.73 (m, 3H), 8.11 (d, 1H), 8.79 (d, 1H).
403	3.89 (s, 3H), 6.92 (d, 1H), 7.26 (t, 1H), 7.43 (d, 1H), 7.54 (t, 1H), 7.61-7.74 (m, 3H), 8.11 (d, 1H), 8.79 (d, 1H).
409	1.19 (t, 3H), 2.33 (d, 3H), 4.27 (q, 2H), 6.78 (d, 1H), 7.29-7.35 (m, 2H), 7.44-7.48 (m, 3H), 8.84 (d, 1H).
410	1.41 (s, 9H), 6.97 (d, 1H), 7.28-7.38 (m, 3H), 7.45-7.49 (m, 3H), 8.85 (d, 1H).
427	1.33 (d, 3H), 2.80 (s, 3H), 3.38 (s, 3H), 4.00 (m, 1H), 6.77 (d, 1H), 7.03 (m, 1H), 7.25-7.45 (m, 5H).
428	1.02 (t, 3H), 1.36 (s, 3H), 1.79 (q, 2H), 2.18 (s, 1H), 2.79 (s, 3H), 6.75 (d, 1H), 7.03 (m, 1H), 7.28-7.45 (m, 5H).
430	1.03 (t, 3H), 1.28 (m, 2H), 1.68 (m, 2H), 3.90 (q, 2H), 6.79 (d, 1H), 6.97 (t, 1H), 7.26-7.35 (m, 2H), 7.40-7.50 (m, 3H), 8.83 (d, 1H).
431	0.66-0.87 (m, 2H), 1.12-1.19 (m, 2H), 1.26-1.29 (m, 8H), 1.99 (s, 1H), 6.67 (d, 1H), 6.89 (m, 1H), 7.18 (m, 1H), 7.34 (m, 1H), 7.47-7.50 (m, 2H), 7.62 (m, 1H), 8.82 (d, 1H).
435	1.82 (d, 6H), 5.28 (br, 2H), 6.62 (d, 1H), 6.87 (dd, 1H), 7.19 (m, 1H), 7.33 (m, 1H), 7.43-7.51 (m, 2H), 7.68 (m, 1H), 8.79 (d, 1H).
437	1.93 (d, 6H), 6.72 (m, 1H), 6.91-6.98 (m, 2H), 7.19-7.38 (m, 2H), 7.43-7.52 (m, 2H), 7.58 (m, 1H), 8.83 (d, 1H).
442	2.46 (s, 6H), 5.22 (s, 1H), 6.8-7.0 (m, 1H), 7.2-7.5 (m, 5H), 7.67 (dd, 1H), 8.84 (d, 1H).

[0279]

Table 40

Compound No.	¹ H-NMR
443	2.07 (s, 3H), 6.78 (d, 1H), 6.80 (bs, 1H), 7.03 (d, 1H), 7.2–7.6 (m, 5H), 7.78 (m, 1H), 8.88 (d, 1H).
445	1.54 (dd, 3H), 1.61–1.69 (m, 4H), 2.34–2.39 (m, 2H), 2.54–2.59 (m, 2H), 3.92 (q, 1H), 6.79 (m, 1H), 6.97 (m, 1H), 7.22–7.33 (m, 2H), 7.39 (m, 1H), 7.43–7.47 (m, 2H), 8.83 (d, 1H).
446	1.47 (d, 3H), 1.54 (s, 9H), 3.29 (s, 3H), 4.97 (q, 1H), 6.67 (d, 1H), 7.2–7.5 (m, 5H), 8.04 (d, 1H), 8.66 (bs, 1H), 8.85 (d, 1H).
447	1.50 (d, 3H), 3.26 (s, 3H), 4.88 (bs, 2H), 4.93 (q, 2H), 6.33 (dd, 1H), 6.52 (dd, 1H), 7.07 (t, 1H), 7.2–7.5 (m, 4H), 8.84 (d, 1H).
448	1.47 (d, 3H), 2.89 (d, 3H), 3.23 (s, 3H), 4.94 (q, 2H), 5.78 (bs, 1H), 6.30 (dd, 1H), 6.51 (d, 1H), 7.19 (t, 1H), 7.24–7.43 (m, 5H), 8.84 (d, 1H).
449	2.11 (br, 6H), 2.84 (s, 3H), 2.92 (s, 3H), 6.68 (m, 1H), 6.96 (m, 1H), 7.27–7.44 (m, 4H), 7.49 (d, 1H).
450	2.11 (d, 6H), 2.91 (s, 3H), 6.74 (m, 1H), 6.97 (m, 1H), 7.29–7.51 (m, 3H), 7.67 (m, 1H), 8.82 (d, 1H).
451	2.16 (br, 6H), 2.81 (s, 3H), 2.92 (s, 3H), 6.65 (d, 1H), 6.98 (m, 1H), 7.26–7.42 (m, 3H), 7.50 (s, 1H).
452	1.38 (t, 3H), 2.11 (s, 6H), 3.08 (q, 2H), 6.76 (d, 1H), 6.98 (m, 1H), 7.30–7.38 (m, 2H), 7.47–7.54 (m, 2H), 7.70 (d, 1H), 8.84 (d, 1H).
453	1.99 (d, 3H), 2.95 (s, 3H), 5.20 (q, 1H), 6.84 (m, 1H), 7.23–7.55 (m, 5H), 7.78 (m, 1H), 8.83 (d, 1H).
a-2	1.67 (s, 6H), 6.00 (s, 1H), 7.54–7.61 (m, 2H), 7.66–7.74 (m, 2H), 8.14 (d, 1H), 8.38 (t, 1H), 8.80 (d, 1H).
a-7	1.66 (s, 6H), 2.81 (s, 3H), 6.02 (s, 1H), 7.19–7.43 (m, 6H), 8.40 (d, 1H).
a-8	1.65 (s, 6H), 5.96 (s, 1H), 7.25–7.39 (m, 3H), 7.48–7.58 (m, 3H), 8.41 (m, 1H), 8.85 (d, 1H).
b-1	1.18 (s, 6H), 1.50 (s, 6H), 1.75–1.93 (m, 4H), 2.73 (t, 2H), 2.84 (s, 1H), 2.89 (t, 2H), 6.82 (dd, 1H), 6.96 (d, 1H), 7.08–7.20 (m, 2H), 7.49 (dd, 1H), 8.10 (d, 1H).
b-3	1.27 (s, 6H), 1.60 (d, 6H), 1.76–1.94 (m, 4H), 2.49 (s, 1H), 2.74 (t, 2H), 2.90 (t, 2H), 6.62 (d, 1H), 6.79–6.86 (m, 1H), 6.97 (d, 1H), 7.08–7.15 (m, 1H), 8.10 (d, 1H).
b-4	1.19 (s, 6H), 1.50 (s, 6H), 2.11–2.21 (m, 2H), 2.83 (s, 1H), 2.90 (t, 2H), 2.95 (t, 2H), 6.81 (dd, 1H), 7.08–7.21 (m, 3H), 7.50 (dd, 1H), 8.08 (d, 1H).
b-5	1.19 (s, 6H), 1.50 (s, 6H), 1.62–1.73 (m, 4H), 1.83–1.90 (m, 2H), 2.70–2.73 (m, 2H), 2.92 (s, 1H), 3.00–3.04 (m, 2H), 6.84 (dd, 1H), 7.01 (d, 2H), 7.07–7.20 (m, 2H), 7.48 (dd, 1H), 8.00 (d, 1H).
b-6	1.76–1.95 (m, 4H), 2.59 (s, 3H), 2.75 (t, 2H), 2.91 (t, 2H), 6.63 (d, 1H), 6.88 (t, 1H), 7.03 (d, 1H), 7.24–7.32 (m, 1H), 8.16 (d, 1H).
b-7	1.74 (d, 6H), 1.74–1.95 (m, 4H), 2.76 (t, 2H), 2.92 (t, 2H), 4.26 (d, 1H), 6.55 (dd, 1H), 6.79–6.86 (m, 1H), 7.05 (d, 1H), 7.07–7.15 (m, 1H), 8.18 (d, 1H).
b-8	1.03 (s, 9H), 1.71 (d, 3H), 1.74–1.95 (m, 4H), 2.75 (t, 2H), 2.91 (t, 2H), 5.00 (s, 1H), 6.53 (d, 1H), 6.77–6.84 (m, 1H), 7.06 (s, 1H), 7.08–7.26 (m, 1H), 8.20 (d, 1H).
b-9	1.63 (d, 3H), 1.76–1.95 (m, 4H), 2.75 (t, 2H), 2.77–2.93 (m, 2H), 3.48 (s, 1H), 5.29–5.34 (m, 1H), 6.55 (d, 1H), 6.81 (dd, 1H), 7.03 (d, 1H), 7.13 (q, 1H), 8.16 (d, 1H).

[0280]

Table 41

Compound No.	¹ H-NMR
b-10	1.59 (d, 3H), 1.65-1.94 (m, 4H), 2.74 (t, 2H), 2.90 (t, 2H), 3.27 (s, 3H), 4.90 (q, 1H), 6.62 (d, 1H), 6.84 (dd, 1H), 6.89 (d, 1H), 7.14-7.22 (m, 1H), 8.11 (d, 1H).
b-11	2.65 (s, 3H), 6.65 (d, 1H), 6.92 (t, 1H), 7.21 (d, 1H), 7.27-7.35 (m, 1H), 7.61 (d, 1H), 7.69 (d, 1H), 8.43 (d, 1H).
b-12	1.77 (d, 6H), 4.16 (d, 1H), 6.57 (d, 1H), 6.86 (dd, 1H), 7.09-7.18 (m, 1H), 7.20 (d, 1H), 7.61 (d, 1H), 7.69 (d, 1H), 8.44 (d, 1H).
b-13	1.17 (s, 6H), 1.53 (s, 6H), 2.92 (s, 3H), 2.92 (t, 2H), 3.41 (s, 1H), 3.48 (t, 2H), 6.78 (d, 1H), 6.91 (d, 1H), 6.92-7.15 (m, 2H), 7.42 (dd, 1H), 7.65 (d, 1H).
b-14	1.67 (d, 3H), 2.75 (dd, 1H), 5.31-5.41 (m, 1H), 6.58 (d, 1H), 6.87 (m, 1H), 7.12-7.21 (m, 2H), 7.61 (d, 1H), 7.67 (d, 1H), 8.44 (d, 1H).
b-15	1.59 (d, 3H), 3.29 (s, 3H), 4.93 (q, 1H), 6.65 (d, 1H), 6.90 (t, 1H), 7.07-7.25 (m, 2H), 7.58 (d, 1H), 7.64 (s, 1H), 8.41 (d, 1H).
b-16	1.06 (s, 9H), 1.75 (d, 3H), 4.93 (s, 1H), 6.54 (d, 1H), 6.84 (dd, 1H), 7.07-7.12 (m, 1H), 7.13 (d, 1H), 7.61 (d, 1H), 7.75 (d, 1H), 8.45 (d, 1H).
b-19	1.50 (s, 3H), 1.67 (s, 3H), 1.81-1.97 (m, 4H), 2.80 (m, 2H), 2.97 (m, 2H), 5.90 (s, 1H), 7.19-7.37 (m, 3H), 7.72 (m, 1H), 8.55 (d, 1H).
b-20	1.52 (s, 3H), 1.58 (s, 3H), 5.88 (s, 1H), 7.23-7.42 (m, 4H), 7.62 (d, 1H), 8.38 (d, 1H), 8.83 (d, 1H).
b-21	2.59 (s, 3H), 2.80 (t, 2H), 4.07 (t, 2H), 4.73 (s, 2H), 6.65 (d, 1H), 6.91 (t, 1H), 6.97 (d, 1H), 7.28-7.35 (m, 1H), 8.24 (d, 1H).
b-22	1.73 (d, 6H), 2.02 (s, 1H), 3.02 (t, 2H), 4.07 (t, 2H), 4.73 (s, 2H), 6.57 (d, 1H), 6.82-6.89 (m, 1H), 6.98 (d, 1H), 7.10-7.15 (m, 1H), 8.25 (d, 1H).
b-23	2.63 (s, 3H), 6.60 (d, 1H), 6.77 (d, 2H), 6.89 (t, 1H), 7.25-7.32 (m, 1H), 7.62 (d, 1H), 7.76 (d, 1H), 8.16 (d, 1H).

[0281]

(Formulation)

Examples of the fungicide according to the present invention are shown below,

5 however, adding agents and the addition ratio are not limited to these examples and it is possible to vary them extensively. Also, parts in formulation examples represent parts by weight.

[0282]

Formulation example 1 Water-dispersible powder

10	The compound according to the present invention	40 parts
	Clay	48 parts
	Sodium dioctylsulfo succinate	4 parts
	Sodium lignin sulfonate	8 parts

The above were uniformly mixed and finely pulverized to obtain

15 water-dispersible powder with 40% active ingredient.

[0283]

Formulation example 2 Emulsion

The compound according to the present invention	10 parts
---	----------

SORVESSO 200	53 parts
--------------	----------

5 Cyclohexanone	26 parts
----------------------	----------

Potassium dodecylbenzene sulfonate	1 part
------------------------------------	--------

Polyoxyethylene alkylallylether	10 parts
---------------------------------	----------

The above were mixed and dissolved to obtain an emulsion with 10% active ingredient.

10 [0284]

Formulation example 3 Powder

The compound according to the present invention	10 parts
---	----------

Clay	90 parts
------	----------

The above were uniformly mixed and finely pulverized to obtain powder with

15 10% active ingredient.

[0285]

Formulation example 4 Granule

The compound according to the present invention	5 parts
---	---------

Clay	73 parts
------	----------

20 Bentonite	20 parts
-------------------	----------

Sodium dioctylsulfo succinate	1 part
-------------------------------	--------

Potassium phosphate	1 part
---------------------	--------

The above were well pulverized, mixed, and well kneaded with an addition of water, followed by granulating and drying to obtain granule with 5% active ingredient.

25 [0286]

Formulation example 5 Suspension

	The compound according to the present invention	10 parts
	Polyoxyethylene alkyallylether	4 parts
	Sodium polycarboxylate	2 parts
5	Glycerine	10 parts
	Xanthane gum	0.2 parts
	Water	73.8 parts

The above were mixed and wet pulverized until the particle diameter becomes equal to or less than 3 micron to obtain a suspension with 10% active ingredient.

10 [0287]

Formulation example 6 Water-dispersible granule

	The compound according to the present invention	40 parts
	Clay	36 parts
	Potassium chloride	10 parts
15	Sodium alkylbenzene sulfonate	1 part
	Sodium lignin sulfonate	8 parts
	Formaldehyde condensation product of sodium alkylbenzene sulfonate	5 parts

The above were mixed and finely pulverized, followed by adding a suitable amount of water and kneading therewith to be in clay shape. The clay shaped material was granulated and dried to obtain water-dispersible granule with 40% active ingredient.

[0288]

(Biological test example 1) Control test of apple scab

To apple seedlings grown in clay pots (Varieties "Ralls Janet", 3 to 4 leaf base), the emulsion of the compound according to the present invention with active ingredient of concentration of 100 ppm was sprayed. After being air dried at room temperature, a

conidium of apple scab bacteria (*Venturia inaequalis*) was inoculated, and was held for 2 weeks in a room at 20°C under a high humidity, with light and darkness being repeated every 12 hours. Controlling effect was determined by comparing the state of the appearance of lesion on the leaves with that of untreated.

5 To compounds represented by Compound Numbers 3, 6, 7, 8, 10, 11, 12, 14, 15, 16, 17, 18, 19, 20, 21, 22, 24, 25, 26, 28, 29, 30, 31, 33, 34, 35, 36, 37, 38, 40, 41, 43, 44, 45, 46, 48, 49, 52, 53, 54, 57, 59, 60, 61, 62, 63, 64, 65 and 66, control test of an apple scab was carried out. As a result, all of the compounds showed at least 75% protective value.

10 Also, the same test was carried out to compounds represented by Compound Numbers 67, 68, 70, 73, 74, 75, 76, 77, 78, 79, 80, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 15 152, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 190, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 215, 217, 218, 219, 220, 221, 222, 223, 224, 225, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 20 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 276, 277, 278, 279, 280, 281, 283, 284, 285, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 306, 308, 309, 310, 311, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 327, 328, 329, 330, 331, 332, 333, 334, 335, 344, 347, 348, 349, 350, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 25 368, 369, 370, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380, 381, 382, 383, 384, 385,

386, 387, 388, 389, 390, 391, 392, 393, 394, 395, 396, 397, 398, 399, 400, 402, 403, 404, 406, 407, 408, 409, 410, 411, 412, 413, 414, 416, 417, 418, 419, 423, 424, 425, 426, 427, 428, 429, 430, 431, 432, 433, 434, 435, 437, 439, 440, 441, 442, 443, 445, a-2, a-3, a-4, a-5, a-6, a-7, a-8, a-9, a-10, b-1, b-2, b-3, b-4, b-5, b-6, b-7, b-8, b-9, b-10, b-11, b-12, b-13, b-14, b-15, b-16, b-17, b-20, b-22, b-23, b-24, c-2, c-3, c-4, c-5, d-5, and d-6. As a result, all of the compounds showed at least 75% protective value.

[0289]

(Biological test example 2) Control test of cucumber gray mold

To cucumber seedlings grown in clay pots (Varieties "Cucumis sativus L.", cotyledon base), the emulsion of the compound according to the present invention with active ingredient of concentration of 100 ppm was sprayed. After being air dried at room temperature, a conidium suspension of cucumber gray mold bacteria (*Botrytis cinerea*) was dropwise inoculated, and was held for 4 days in a dark room at 20°C under a high humidity. Controlling effect was determined by comparing the state of the appearance of lesion on the leaves with that of untreated.

To compounds represented by Compound Numbers 10, 11, 12, 15, 18, 19, 22, 26, 29, 30, 31, 33, 34, 35, 36, 37, 38, 42, 43, 44, 48, 49, 54, 57, and 59, control test of a cucumber gray mold was carried out. As a result, all of the compounds showed at least 75% protective value.

Also, the same test was carried out to compounds represented by Compound Numbers 67, 73, 75, 76, 77, 80, 81, 82, 83, 86, 88, 89, 90, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 104, 106, 107, 108, 109, 110, 111, 113, 114, 115, 116, 117, 118, 119, 122, 123, 124, 125, 126, 127, 128, 133, 134, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 156, 157, 158, 159, 160, 162, 163, 164, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 183, 184, 185, 186, 187, 188,

192, 197, 198, 199, 200, 201, 202, 203, 205, 208, 209, 210, 219, 220, 221, 222, 223, 225,
 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244,
 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262,
 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 274, 275, 278, 281, 284, 285, 291, 292,
 5 293, 295, 296, 297, 298, 299, 308, 309, 310, 311, 314, 315, 316, 317, 318, 319, 323, 324,
 325, 326, 327, 328, 332, 333, 338, 340, 344, 348, 353, 355, 359, 360, 361, 363, 364, 365,
 371, 375, 376, 377, 378, 379, 380, 381, 382, 386, 391, 392, 393, 395, 397, 398, 399, 402,
 403, 405, 406, 412, 414, 416, 417, 418, 423, 424, 425, 426, 427, 428, 429, 431, 432, 433,
 434, 435, 436, 437, 439, 440, 441, 442, 444, a-2, a-5, a-6, a-7, a-8, a-10, b-1, b-2, b-3,
 10 b-7, b-8, b-12, b-16, c-3, and c-4. As a result, all of the compounds showed at least
 75% protective value.

[0290]

(Biological test example 3) Submerged application test of rice neck rot

Rice seedlings were grown in pots where commercially available hilling is in a
 15 submerged condition (Varieties “koshihikari”, 1 leaf base), and the emulsion of the
 compound according to the present invention with active ingredient of concentration of
 400 ppm was drip treated to the surface of the water. After 2 days, a conidium
 suspension of rice neck rot bacteria (*Magnaporthe grisea*) was spray inoculated, was held
 for 2 days in a dark room at 25°C under a high humidity, and then was held for 8 days in
 20 a room at 25°C, with light and darkness being repeated in every 12 hours. Controlling
 effect was decided by the following grades after comparing the state of the appearance of
 lesion on the leaves with that of untreated.

A (at least 60% protective value)

B (at least 40% but less than 60% protective value)

25 As a result of the present test, the controlling effect of the following compounds

was evaluated as A.

Compound Number : 5, 20, 21, 23, 25, 28, 32, 33, 34, 36, 37, 38, 43, 44, 48, 58, 66, 70, 71, 73, 78, 84, 85, 88, 89, 90, 106, 107, 227, 228, 234, 236, 260, 262, 348, 353, 356, 358, 375, 376, 401, 402, a-6

5 Also, the controlling effect of the following compounds was evaluated as B.

Compound Number: 1, 3, 7, 12, 13, 22, 46, 54, 55, 75, 95, 98, 139, 151, 261, 359, 360, b-1

[0291]

(Biological test example 4) Submerged application test of rice neck rot

10 Rice seedlings were grown in pots where commercially available hilling is in a submerged condition (Varieties “koshihikari”, 1 leaf base), and the emulsion of the compound according to the present invention with active ingredient of concentration of 400 ppm was drip treated to the surface of the water. After 14 days, a conidium suspension of rice neck rot bacteria (*Magnaporthe grisea*) was spray inoculated, was held
15 for 2 days in a dark room at 25°C under a high humidity, and then was held for 8 days in a room at 25°C, with light and darkness being repeated in every 12 hours. Controlling effect was decided by the following grades after comparing the state of the appearance of lesion on the leaves with that of untreated.

A (at least 60% protective value)

20 B (at least 40% but less than 60% protective value)

As a result of the present test, the controlling effect of the following compounds was evaluated as A.

Compound Number: 21, 36, 37, 38, 43, 44, 48, 84, 88, 89, 95, 106, 109, 125, 262, 266, 267, 292, 350, 375, a-6

25 Also, the controlling effect of the following compound was evaluated as B.

Compound Number: 236

[0292]

(Biological test example 5) Seed treatment test of cucumber wilt

Cucumber seeds (Varieties "Cucumis sativus L.") contaminated by cucumber
 5 wilt bacteria (*Fusarium oxysporum*) were treated with the emulsion of the compound
 according to the present invention to obtain the seeds containing 1 g/kg active ingredient.
 The seeds were sowed and after 3 weeks, controlling effect was determined by
 comparing the degree of disease occurrence with that of untreated.

As a result, the following compounds showed an excellent protective value of at
 10 least 75%.

Compound Number: 18, 36, 37, 43, 44, 48, 77, 81, 83, 88, 92, 94, 100, 103, 109,
 126, 225, 227, 228, 234, 235, 241, 242, 243, 244, 252, 258, 266, 268, 298, 395, 397, 425,
 435, a-6, a-7, a-8, b-8, b-12, c-4

[0293]

15 (Biological test example 6) Seed treatment test of cucumber wilt


Cucumber seeds (Varieties "Cucumis sativus L.") were treated with the emulsion
 of the compound according to the present invention to obtain the seeds containing 1 g/kg
 active ingredient. The seeds were sowed in soil contaminated by cucumber wilt
 bacteria (*Fusarium oxysporum*) and, after 3 weeks, controlling effect was determined by
 20 comparing the degree of disease occurrence with that of untreated.

As a result, the following compounds showed an excellent protective value of at
 least 75%.

Compound Number: 17, 18, 31, 33, 34, 35, 36, 37, 43, 86, 88, 96, 97, 100, 109,
 227, 232, 239, 242, 243, 244, 262, 265, 268, 295, 296, 375, 428, a-7, b-1

CLAIMS

1. A nitrogen-containing heterocyclic compound represented by Formula (I):



- 5 (wherein, in Formula (I), R represents a group represented by $CR^1R^2R^3$, an unsubstituted or substituted C_{6-10} aryl group, or a cyano group;
- R^1 to R^3 each independently represents a hydrogen atom, an unsubstituted or substituted C_{1-8} alkyl group, an unsubstituted or substituted C_{2-8} alkenyl group, an unsubstituted or substituted C_{2-8} alkynyl group, an unsubstituted or substituted C_{3-8}
- 10 cycloalkyl group, an unsubstituted or substituted C_{4-8} cycloalkenyl group, an unsubstituted or substituted C_{6-10} aryl group, an unsubstituted or substituted heterocyclic group, an unsubstituted or substituted C_{1-8} acyl group, an unsubstituted or substituted (1-imino) C_{1-8} alkyl group, an unsubstituted or substituted carboxyl group, an unsubstituted or substituted carbamoyl group, an unsubstituted or substituted hydroxyl
- 15 group, an unsubstituted or substituted amino group, an unsubstituted or substituted mercapto group, a substituted sulfonyl group, a halogeno group, a cyano group, or a nitro group;

excepting in which: R^1 to R^3 are all hydrogen atoms; R^1 to R^3 are all unsubstituted C_{1-8} alkyl groups; any one of R^1 to R^3 is a hydrogen atom and the

remaining two are both unsubstituted C₁₋₈ alkyl groups; and, any one of R¹ to R³ is an unsubstituted C₁₋₈ alkyl group and the remaining two are both hydrogen atoms;

R¹ and R² may be joined to form an unsubstituted or substituted 5- to 8-membered ring, or to form O=, R^aR^bC=, or R'-N=;

5 R^a represents a hydrogen atom or an unsubstituted or substituted C₁₋₈ alkyl group;

R^b represents a hydrogen atom or an unsubstituted or substituted C₁₋₈ alkyl group;

R' represents an unsubstituted or substituted hydroxyl group, or an unsubstituted
10 or substituted C₁₋₈ alkyl group;

R⁴ each independently represents an unsubstituted or substituted C₁₋₈ alkyl group, an unsubstituted or substituted C₂₋₈ alkenyl group, an unsubstituted or substituted C₂₋₈ alkynyl group, an unsubstituted or substituted C₃₋₈ cycloalkyl group, an unsubstituted or substituted C₄₋₈ cycloalkenyl group, an unsubstituted or substituted C₆₋₁₀ aryl group, an
15 unsubstituted or substituted heterocyclic group, an unsubstituted or substituted C₁₋₈ acyl group, an unsubstituted or substituted (1-imino)C₁₋₈ alkyl group, an unsubstituted or substituted carboxyl group, an unsubstituted or substituted carbamoyl group, an unsubstituted or substituted hydroxyl group, an unsubstituted or substituted amino group, an unsubstituted or substituted mercapto group, a substituted sulfonyl group, a halogeno
20 group, a cyano group, or a nitro group;

m represents a number of R⁴ and is an integer of 0 to 6;

R⁵ each independently represents an unsubstituted or substituted C₁₋₈ alkyl group, an unsubstituted or substituted C₂₋₈ alkenyl group, an unsubstituted or substituted C₂₋₈ alkynyl group, an unsubstituted or substituted C₃₋₈ cycloalkyl group, an unsubstituted or
25 substituted C₄₋₈ cycloalkenyl group, an unsubstituted or substituted C₆₋₁₀ aryl group, an

unsubstituted or substituted heterocyclic group, an unsubstituted or substituted C₁₋₈ acyl group, an unsubstituted or substituted (1-imino)C₁₋₈ alkyl group, an unsubstituted or substituted carboxyl group, an unsubstituted or substituted carbamoyl group, an unsubstituted or substituted hydroxyl group, an unsubstituted or substituted amino group, an unsubstituted or substituted mercapto group, a substituted sulfonyl group, a halogeno group, a cyano group, or a nitro group;

n represents a number of R⁵ and is an integer of 0 to 5;

any one of R¹ to R³ and any one of R⁵ may be joined to form an unsubstituted or substituted 5- to 8-membered ring;

A represents: a 5- to 7-membered hydrocarbon ring or a 5- to 7-membered heterocyclic ring when R is a group represented by CR¹R²R³; or a benzene ring when R is an unsubstituted or substituted C₆₋₁₀ aryl group or a cyano group;

D represents a 5- to 7-membered hydrocarbon ring or a 5- to 7-membered heterocyclic ring;


X represents an oxygen atom, a sulfur atom, a sulfenyl group, a sulfonyl group, an unsubstituted or substituted carbon atom, or an unsubstituted or substituted nitrogen atom;

Y represents a carbon atom or a nitrogen atom; and

Z represents a carbon atom or a nitrogen atom),

or a salt thereof, or an N-oxide compound thereof.

2. A nitrogen-containing heterocyclic compound represented by Formula (II):




Formula (II)

(wherein, in Formula (II), each of R, R⁴, R⁵, n, D, X, Y, and Z represents the same meaning as those in Formula (I) of Claim 1; and

m1 represents a number of R⁴ and is an integer of 0 to 6),

5 or a salt thereof, or an N-oxide compound thereof.

3. A nitrogen-containing heterocyclic compound represented by Formula (III):



Formula (III)

(wherein, in Formula (III), each of R, R⁴, R⁵, m1 and X represents the same meaning as those in Formula (II) of Claim 2; and

10 n1 represents a number of R⁵ and is an integer of 0 to 4),

or a salt thereof, or N-oxide compound thereof.

4. An agricultural fungicide comprising, as an active ingredient, at least one

15 selected from the group consisting of the nitrogen-containing heterocyclic compound, the

salt thereof, and the N-oxide compound thereof, of any one of Claims 1 to 3.

5. An intermediate product of the nitrogen-containing heterocyclic compound represented by Formula (I), the salt thereof, or the N-oxide compound thereof, of Claim 1,
- 5 the intermediate product being selected from the group consisting of
- 8-fluoro-3-hydroxyquinoline, 7,8-difluoro-3-hydroxyquinoline,
- 7,8-difluoro-3-iodoquinoline, 8-fluoro-3-hydroxy-2-methylquinoline, and
- 7,8-difluoro-3-hydroxy-2-methylquinoline.