

12. The composition of claim 11 wherein at least one of the additional compounds is an interferon or a ribavirin.

13. The composition of claim 12 wherein the interferon is selected from interferon alpha 2B, pegylated interferon alpha, consensus interferon, interferon alpha 2A, interferon lambda, and lymphoblastoid interferon tau.

14. The composition of claim 11 wherein at least one of the additional compounds is selected from interleukin 2, interleukin 6, interleukin 12, a compound that enhances the development of a type 1 helper T cell response, interfering RNA, anti-sense RNA, Imiquimod, ribavirin, an inosine 5'-monophosphate dehydrogenase inhibitor, amantadine, and rimantadine.

15. The composition of claim 11 wherein at least one of the additional compounds is effective to inhibit the function of a target selected from HCV metalloprotease, HCV serine protease, HCV polymerase, HCV helicase, HCV NS4B protein, HCV entry, HCV assembly, HCV egress, HCV NS5A protein, and IMPDH for the treatment of an HCV infection.

16. ~~A method of treating an HCV infection in a patient, comprising administering to the patient a therapeutically effective amount of~~ Use of a compound of claim 1, or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for treating an HCV infection in a patient.

17. ~~The method-use~~ of claim 16 further comprising administering one, two, or three additional compounds having anti-HCV activity prior to, after or simultaneously with the compound of claim 1, or a pharmaceutically acceptable salt thereof.

18. ~~The method-use~~ of claim 17 wherein at least one of the additional compounds is an interferon or a ribavirin.

19. ~~The method-use~~ of claim 18 wherein the interferon is selected from interferon alpha 2B, pegylated interferon alpha, consensus interferon, interferon alpha 2A, interferon lambda, and lymphoblastoid interferon tau.

20. The ~~method~~use of claim 17 wherein at least one of the additional compounds is selected from interleukin 2, interleukin 6, interleukin 12, a compound that enhances the development of a type 1 helper T cell response, interfering RNA, anti-sense RNA, Imiquimod, ribavirin, an inosine 5'-monophosphate dehydrogenase inhibitor, amantadine, and rimantadine.

21. The ~~method~~use of claim 17 wherein at least one of the additional compounds is effective to inhibit the function of a target selected from HCV metalloprotease, HCV serine protease, HCV polymerase, HCV helicase, HCV NS4B protein, HCV entry, HCV assembly, HCV egress, HCV NS5A protein, and IMPDH for the treatment of an HCV infection.