

BIOMARKER WITH THERAPEUTIC IMPLICATIONS FOR CARCINOMATOSIS

CROSS-REFERENCE TO RELATED APPLICATIONS

5 [001] This application claims the benefit of priority of Singapore provisional application no. 10201902763U, filed 27 March 2019, the contents of it being hereby incorporated by reference in its entirety for all purposes.

FIELD OF THE INVENTION

[002] The present invention relates generally to the field of molecular biology. In particular, the present
10 invention relates to the use of biomarkers for the detection, diagnosis and subsequent treatment of cancer.

BACKGROUND OF THE INVENTION

[003] Colorectal cancer is the third most common cancer and the fourth most common cause of cancer death globally, accounting for 1.4 million new cases and 600 000 deaths per year. Deaths from colorectal
15 cancer are largely due to metastasis with peritoneal carcinomatosis (PC) occurring in 15% of all patients and accounting for up to 30% of all metastases. Compared to other forms of metastatic colorectal cancer without peritoneal involvement, colorectal peritoneal carcinomatosis has consistently demonstrated to have significantly shorter overall survival despite palliative systemic chemotherapy.

[004] Cytoreductive surgery (CRS) and hyperthermic intraperitoneal chemotherapy (HIPEC) have
20 revolutionised the treatment of peritoneal carcinomatosis. Cytoreductive surgery refers to a series of visceral resections and peritonectomy procedures that remove all macroscopic disease. Remaining viable microscopic disease is then eradicated with the instillation of hyperthermic intraperitoneal chemotherapy. The combined treatment modalities of cytoreductive surgery and hyperthermic intraperitoneal chemotherapy have greatly improved survival in patients with peritoneal carcinomatosis of colorectal
25 origin. The median survival of patients treated with cytoreductive surgery and hyperthermic intraperitoneal chemotherapy was 33 months, compared to 6 to 12 months in patients treated with systemic chemotherapy alone. However, despite this dramatic improvement, much more needs to be done to further improve the outcome of treatment for patients with colorectal peritoneal carcinomatosis by improving the hyperthermic intraperitoneal chemotherapy regimen, as surgery is unlikely to improve
30 patient outcome further.

[005] Hence, there is a need for improved patient stratification in order to improve treatment.

SUMMARY

[006] In one aspect, the present disclosure refers to a use of a “plasminogen activator inhibitor 1” (PAI-
35 1) inhibitor in the manufacture of a medicament for treating a subject suffering from peritoneal carcinomatosis, wherein the subject is to be identified by determining the concentration of PAI-1 and determining the level of phosphorylation of “signal transducer and activator of transcript 3” (STAT3) in a

sample obtained from the subject; wherein if the subject shows (a) an increase in PAI-1 concentration and an increase in STAT3 phosphorylation, or (b) a decrease in PAI 1 concentration and an increase in STAT3 phosphorylation, the subject is to be administered the medicament; wherein the increase and/or decrease of the concentration of PAI-1 and STAT3 phosphorylation is compared to a reference value; wherein the sample is selected from the group consisting of ascites, tumour samples, and biopsy samples; wherein the reference values are obtained from a reference group comprising subjects suffering from peritoneal carcinomatosis. In one example, there is described a method of treating a subject suffering from peritoneal carcinomatosis with a “plasminogen activator inhibitor 1” (PAI-1) inhibitor, the method comprising determining the concentration of PAI-1 and determining the level of phosphorylation of “signal transducer and activator of transcript 3” (STAT3) in a sample obtained from the subject; administering the PAI-1 inhibitor to the subject showing (a) an increase in PAI-1 concentration and an increase in STAT3 phosphorylation, or (b) a decrease in PAI 1 concentration and an increase in STAT3 phosphorylation; wherein the increase and/or decrease of the concentration of PAI-1 and STAT3 phosphorylation is compared to a reference value.

[007] In another aspect, the present disclosure refers to a method of detecting or determining susceptibility of a subject suffering from peritoneal carcinomatosis to a treatment with a “plasminogen activator inhibitor 1” (PAI-1) inhibitor, the method comprising determining the concentration of PAI-1 and determining the level of phosphorylation of “signal transducer and activator of transcript 3” (STAT3) in a sample obtained from a subject; wherein the subject is susceptible to the treatment if the subject shows (a) an increase in PAI 1 concentration and an increase in STAT3 phosphorylation, or (b) a decrease in PAI 1 concentration and an increase in STAT3 phosphorylation; wherein the subject is not considered to be susceptible to treatment if the subject shows (c) a decrease in PAI-1 concentration and a decrease in STAT3 phosphorylation; wherein the increase and/or decrease of the concentration of PAI-1 and the level of STAT3 phosphorylation is compared to a reference value; wherein the sample is selected from ascites, tumour samples, and biopsy samples; wherein the reference values are obtained from a reference group comprising subjects suffering from peritoneal carcinomatosis.

[008] In one example, the present disclosure refers to a panel of markers for treating a patient suffering from peritoneal carcinomatosis with a “plasminogen activator inhibitor 1” (PAI-1) inhibitor, or for detecting or determining susceptibility of a subject suffering from peritoneal carcinomatosis to a treatment with a “plasminogen activator inhibitor 1” (PAI-1) inhibitor, wherein the panel of markers comprises PAI-1, and one or more surrogate markers of STAT3 phosphorylation, or p-STAT3.

[009] In another aspect, the present disclosure refers to the use of a panel of markers in the use or the method disclosed herein, wherein the panel comprises PAI-1 and two or more surrogate markers of STAT3 phosphorylation, wherein at least one of the surrogate markers of STAT3 phosphorylation is MMP9, or PAI-1 and p-STAT3; wherein the at two or more surrogate markers are selected from the group consisting of IL6, IL10, CCL2, MMP9, ANGPT1, TGFB1, POSTN, VSIG4, CD44, and CXCL10, wherein one of two or more surrogate markers is MMP9.

BRIEF DESCRIPTION OF THE DRAWINGS

[0010] The invention will be better understood with reference to the detailed description when considered in conjunction with the non-limiting examples and the accompanying drawings, in which:

[0011] **Fig. 1** shows data indicating that the presence of ascites leads to a poorer prognosis in patients regardless of histological subtype. (A) Kaplan-Meier survival curve of all patients with peritoneal carcinomatosis ($P = 0.002$). (B) Kaplan-Meier survival curve of colorectal patients with peritoneal carcinomatosis ($P = 0.001$). (C) Kaplan-Meier survival curve of ovarian patients with peritoneal carcinomatosis ($P = 0.077$).

[0012] **Fig. 2** (A) Addition of cell-free ascites increased cancer cells proliferation in a dose-dependent manner. 0.1% of cell-free ascitic fluid was sufficient to maintain cell viability without proliferation. (B) Treatment with cell-free ascites significantly increased cancer cells migration. (C) Treatment with cell-free ascites dramatically increased the cell settlement of cancer cells *in vitro* independently of serum supplemented media (As: Ascites, SFM: Serum-free media, FBS: Foetal bovine serum).

[0013] **Fig. 3** shows pathways significantly upregulated upon treatment with cell-free ascites. (A) Pathways upregulated in peritoneal carcinomatosis (PC) cell lines treated with 5% versus 0.1% cell-free ascites. (B) Specific pathways upregulated in cell-free ascites treated cells that are not shared in pathways activated by survival signals. These figures show that upon treating cancer cell lines with cell-free ascites, several signalling pathways were found to be upregulated, including the IL6-JAK-STAT3 signalling pathway. This indicates that the activation of STAT3 plays an important role in the disease. The term “treatment”, as referred to in the present figure and context, differs from the definition of “treatment” as provided in the definition section below. In the context of Figure 3, treatment refers to the exposure of cell line models to cell-free ascites collected from patients in an *in vitro* setting. For example, cancer cell lines are exposed to 5% cell-free ascites in an *in vitro* setting and the cells are observed for a change in physical phenotype (e.g. proliferation or migration) or molecular phenotype (e.g. gene expression changes).

[0014] **Fig. 4** (A) Treatment with 5% cell-free ascites activated STAT3 via phosphorylation at Tyr705. (B) Treatment of 5% cell-free ascites of various histological peritoneal carcinomatosis (PC) subtypes resulted in activation of STAT3, where cell-free ascites of colorectal origin showed the most activation.

[0015] **Fig. 5** (A) Representative immunohistochemical staining of p-STAT3 in colorectal primary tumour and its matched metastases. (B) STAT3 activation is more prevalent in the metastases compared to the primary tumour (P: Primary tumour, M: Metastases). In the context of the present invention as a whole, this data highlights that STAT3 signalling pathway is more upregulated in metastases than the primary tumour, and by inference, more reliant on STAT3 signalling. This in turn suggests that metastases are more susceptible to STAT3 inhibition than the primary tumour. In other words, targeting STAT3 signalling in metastases can be more efficacious than targeting the primary tumour.

[0016] **Fig. 6** (A) Bar chart illustrating most differentially expressed epithelial-mesenchymal transition (EMT) genes in established cell line models of colorectal peritoneal carcinomatosis treated with cell-free ascites. (B) Proteins involved in the coagulation pathway were most prevalent in cell-free ascites of colorectal cancer origin. (C) Cytokine array performed on peritoneal carcinomatosis cell-free ascites from various histological subtypes identified abundant PAI-1 levels in cell-free ascites from colorectal peritoneal carcinomatosis.

[0017] **Fig. 7** shows interrogation of PAI-1, STAT3 and EMT expressions and survival analysis in The Cancer Genome Atlas Colorectal Adenocarcinoma (TCGA COADREAD) cohort ($n = 345$). (A) Correlation between PAI-1 and STAT3 expressions. (B) Correlation between PAI-1 expression and EMT signature. Correlations in (A-B) were determined by Pearson correlation coefficient test. Linear regression lines are shown. (C) Kaplan-Meier survival analysis illustrating poorest survival in colorectal cancers with high levels of PAI-1, activated STAT3 signalling, and with enrichment of the epithelial-mesenchymal transition (EMT) signature. (P: PAI-1, S: STAT3 signalling, E: EMT signature)

[0018] **Fig. 8** (A) Receptor Tyrosine Kinase (RTK) phosphorylation array performed on colorectal peritoneal carcinomatosis (PC) cell lines treated with cell-free ascites revealed no activation of JAKs,

suggesting non-canonical mechanism of STAT3 activation. (B) Western blot validation showing JAKs are inactive in cell-free ascites-treated cells.

[0019] **Fig. 9** shows results of a screen which was conducted on (A) cell-free ascites from colorectal peritoneal carcinomatosis (PC) ($n = 55$) and (B) cell-free ascites from various histological peritoneal carcinomatosis (PC) subtypes ($n = 156$) using ELISA. Levels of PAI-1 in ascites and cancer cells p-STAT3 (Y705) levels upon treatment with cell-free ascites were measured and plotted to determine their association. PAI-1 concentrations are plotted on a $\log_2$ scale. p-STAT3 (Y705) levels are shown as optical density reading at 450 nm (OD450). Correlation analyses were determined by Pearson correlation coefficient test.

[0020] **Fig. 10** Untransformed values of PAI-1 and p-STAT3 (Y705) levels were used for gating strategy to identify patient subpopulations which might benefit from PAI-1 inhibition. Three distinct groups of patients can be observed – patients who had high PAI-1 levels and high STAT3 activation, termed PAI-1 paracrine addicted or PPA (right upper quadrant), patients with low PAI-1 levels but high STAT3 activation, termed co-activators predominant or CAP (left upper quadrant), and patients with low PAI-1 levels and low STAT3 activation, termed alternative pathways activation or APA (left lower quadrant). (A) PAI-1 and p-STAT3 gating of colorectal peritoneal carcinomatosis (PC) cell-free ascites. (B) PAI-1 and p-STAT3 gating of various histological peritoneal carcinomatosis (PC) subtypes cell-free ascites. (C) PAI-1 and p-STAT3 cut-off values used to stratify patients into the three distinct groups. (D) Cohort of cell-free ascites used in this analysis.

[0021] **Fig. 11** shows the effect of TM5441 (PAI-1 inhibitor) on the three distinct groups of cell-free ascites-treated Colo-205 cells. (A) Representative inhibitor dose-response curves of PAI-1 paracrine addicted (PPA) group (black solid line), co-activators predominant (CAP) group (black dotted line), alternative pathways activation (APA) group (grey solid line), and foetal bovine serum (FBS; control, grey dotted line) demonstrated a left shift in dose-response curve, indicating responsiveness to PAI-1 inhibition. (B) Differential sensitivity of cell-free ascites to TM5441 corresponding to the three distinct group, with PAI-1 paracrine addicted (PPA) ($n = 18$) being the most sensitive to PAI-1 inhibition, followed by co-activators predominant (CAP) ($n = 59$) and alternative pathways activation (APA) ($n = 17$). (C) Cohort of cell-free ascites used in this analysis.

[0022] **Fig. 12** shows the effect of various pharmacological inhibitions on the three distinct groups of cell-free ascites-treated Colo-205 cells. Representative inhibitor dose-response curves of PAI-1 paracrine addicted (PPA) group (black solid line), co-activators predominant (CAP) group (black dotted line), alternative pathways activation (APA) group (grey solid line), and foetal bovine serum (FBS; control, grey dotted line). Corresponding IC_{50} values are shown in inset, mean $\pm$ s.d. (A) Tiplaxtinin (PAI-1 inhibitor) dose-response curve, (B) Napabucasin (STAT3 inhibitor) dose-response curve, (C) BEZ235 (dual PI3K/mTOR inhibitor) dose-response curve, and (D) Mitomycin C (conventional chemotherapeutic agent used in hyperthermic intraperitoneal chemotherapy (HIPEC) – DNA crosslinker) dose-response curve. Targeting PAI-1, a dominant paracrine factor in cell-free ascites, was shown to be more effective

than targeting downstream signalling pathway activated by cell-free ascites, proliferation pathway, or DNA synthesis.

[0023] **Fig. 13** (A) Signalling pathways affected by PAI-1 inhibition were identified by RNA microarray analysis of cancer cells treated with cell-free ascites representative of PAI-1 paracrine addicted (PPA) (PC085), co-activators predominant (CAP) (PC249) or foetal bovine serum (FBS; control) in the presence of TM5441 or DMSO vehicle. IL6-JAK-STAT3 signalling pathway was significantly downregulated in PAI-1 paracrine addicted (PPA)-treated cells upon PAI-1 inhibition. Normalised enrichment scores less than 0 indicate pathway suppression and scores greater than 0 indicate pathway activation. (B) Treatment of cancer cells with PAI-1 paracrine addicted (PPA) cell-free ascites (PC085 and PC383), co-activators predominant (CAP) cell-free ascites (PC249) and alternative pathways activation (APA) cell-free ascites (PC010) in the presence of various concentrations of TM5441 or DMSO vehicle measured by ELISA confirmed that cells exposed to PAI-1 paracrine addicted (PPA) cell-free ascites relied on PAI-1 to activate STAT3 as they required a lower concentration of TM5441 to suppress STAT3 activation.

[0024] **Fig. 14** (A) Schematic of modified peritoneal cancer index (PCI), used to assess tumour burden in peritoneal carcinomatosis (PC) cell line mouse model. This scoring system is a modification of peritoneal carcinomatosis index (PCI) scoring from Klaver et al. (Klaver Y.L.B., Hendriks T., Lomme R.M.L.M., Rutten H.J.T., Bleichrodt R.P., de Hingh I.H.J.T. (2010) Intraoperative hyperthermic intraperitoneal chemotherapy after cytoreductive surgery for peritoneal carcinomatosis in an experimental model. *British Journal of Surgery*. **97**: 1874-80) and Sugarbaker (Sugarbaker P.H. (1998) Intraperitoneal chemotherapy and cytoreductive surgery for the prevention and treatment of peritoneal carcinomatosis and sarcomatosis. *Seminars in Surgical Oncology*. **14**: 254-61).

[0025] (B) *In vivo* validation of differential sensitivity to PAI-1 inhibition in peritoneal carcinomatosis (PC) cell line mouse model treated with PAI-1 paracrine addicted (PPA) cell-free ascites (PC085), co-activators predominant (CAP) cell-free ascites (PC249) and foetal bovine serum (FBS; control). Images shown are representative of peritoneal metastases formed in response to PAI-1 inhibition or vehicle. Arrows indicate visible tumours. (C) Tumour burden was assessed by modified peritoneal carcinomatosis index (PCI) score. Mice treated with PAI-1 paracrine addicted (PPA) cell-free ascites had significant reduction in tumour burden in response to TM5441 compared to mice treated with co-activators predominant (CAP) cell-free ascites (PC249) and foetal bovine serum ($n = 5$ mice/group).

[0026] **Fig. 15** shows formation of peritoneal tumours in peritoneal carcinomatosis (PC) cell line mouse model exposed to PAI-1 paracrine addicted (PPA) cell-free ascites (PC085) effectively inhibited by intraperitoneal (i.p.) instillation of TM5441, but not when taken orally ($n = 4$ mice/group).

[0027] **Fig. 16** Matched patient's cell-free ascites and its cellular components were used to generate patient-derived ascites-dependent xenograft (PDADX). (A) Representative images of intraperitoneal tumours formed in PC383 patient-derived ascites-dependent xenograft (PDADX) and PC249 patient-derived ascites-dependent xenograft (PDADX) models. Arrows indicate visible tumours. (B) Representative haematoxylin and eosin (H&E) staining and immunohistochemical analyses reveal

patient-derived ascites-dependent xenograft (PDADX) tumours with similar histological features as corresponding patients' tumour tissues, and that these patient-derived ascites-dependent xenograft (PDADX) tumours are of colonic origin (CK20+ CK7- CDX2+). Scale bar, 50 μ M.

[0028] **Fig. 17** shows PAI-1 inhibition is highly efficacious in *in vivo* mouse models that are addicted to PAI-1 paracrine addicted (PPA) cell-free ascites. (A) PAI-1 paracrine addicted (PPA) patient-derived ascites-dependent xenografts (PDADX) (PC383) and co-activators predominant (CAP) patient-derived ascites-dependent xenograft (PDADX) (PC249) were treated with its matched cell-free ascites or foetal bovine serum (FBS) in the presence of DMSO vehicle or 2 mM TM5441 ($n = 4$ mice/group). Tumour burden was quantified by weighing all visible tumours after mice were sacrificed. Only PAI-1 paracrine addicted (PPA) patient-derived ascites-dependent xenograft (PDADX) treated with matched PAI-1 paracrine addicted (PPA) cell-free ascites was susceptible to PAI-1 inhibition and demonstrated significant reduction in tumour burden. (B) Co-activators predominant (CAP) patient-derived ascites-dependent xenograft (PDADX) (PC249) were treated with its matched cell-free ascites or PAI-1 paracrine addicted (PPA) cell-free ascites (PC383) in the presence of DMSO vehicle or 2 mM TM5441 ($n = 4$ mice/group, except in group treated with PC249 cell-free ascites and DMSO ($n = 3$). Co-activators predominant (CAP) patient-derived ascites-dependent xenograft (PDADX) exposed to PAI-1 paracrine addicted (PPA) cell-free ascites became susceptible to PAI-1 inhibition despite not being susceptible in the presence of its matched ascites.

[0029] **Fig. 18** shows a proposed model of paracrine perturbation that can be harnessed for novel therapeutic strategy in peritoneal carcinomatosis (PC).

[0030] **Fig. 19** (A) Workflow to select p-STAT3 surrogate biomarker candidates. (B) Targets prioritisation based on systematic paired correlation analysis. Genes that were chosen for validation with ELISA are shown in bold. Others represent genes that are not in top 25% positively correlated with STAT3 in TCGA COADREAD database and genes that are not in top 25% downregulated/upregulated in TM5441 microarray database.

[0031] **Fig. 20** shows validated surrogate biomarker panel of p-STAT3 ($n = 70$). (A) Correlation between p-STAT3 and selected p-STAT3 surrogate biomarker candidates IL6, IL10, CCL2, MMP9, and ANGPT1. Concentrations of surrogate biomarkers in each patient's cell-free ascites were measured by ELISA and plotted against the degree of STAT3 phosphorylation ($n = 70$ samples/surrogate marker). Correlation analyses were determined by Spearman correlation coefficient test. (B) Receiver operating characteristic (ROC) curve representing ability of individual p-STAT3 surrogate biomarkers to correctly classify PAI-1 paracrine addicted (PPA)/co-activators predominant (CAP) group or alternative pathways activation (APA) group. (C) Cut-off values of p-STAT3 surrogate biomarkers used to classify samples into PAI-1 paracrine addicted (PPA)/co-activators predominant (CAP) group or alternative pathways activation (APA) group. (D) Summary of classification accuracy of individual biomarkers and combinations of composite biomarkers. Biomarker is considered as positive (+) if the concentration of

sample is above the cut-off value. (E) Summary of cut-off values that can be used to identify patients who might be susceptible to PAI-1 inhibition.

[0032] **Fig. 21** shows an alternative surrogate biomarker panel of p-STAT3 ($n = 40$), (A) Correlation between p-STAT3 and selected p-STAT3 surrogate biomarker candidates TGFB1, POSTN, VSIG4, CD44, and CXCL10. Concentrations of surrogate biomarkers in each patient's cell-free ascites were measured by ELISA and plotted against the degree of STAT3 phosphorylation ($n = 40$ samples/surrogate marker). Correlation analyses were determined by Spearman correlation coefficient test. (B) Receiver operating characteristic (ROC) curve representing ability of individual p-STAT3 surrogate biomarkers to correctly classify PAI-1 paracrine addicted (PPA)/co-activators predominant (CAP) group or alternative pathways activation (APA) group. (C) Receiver operating characteristic (ROC) curve of composite biomarker panel comprising of TGFB1, POSTN, VSIG4, CD44, and CXCL10. (D) Receiver operating characteristic (ROC) curve of IL6 using matched samples used in TGFB1, POSTN, VSIG4, CD44, and CXCL10 analysis (left) and receiver operating characteristic (ROC) curve of composite biomarker panel comprising of IL6, TGFB1, POSTN, VSIG4, CD44, and CXCL10 (right). (E) Summary of area under the curve (AUC) of individual biomarkers and composite biomarker panels.

DEFINITIONS

[0033] As used herein, the terms "level" and "concentration" are used synonymously.

[0034] As used herein, the term "biomarker", or "marker", refers to molecular indicators of a specific biological property, a biochemical feature or facet that can be used to determine the presence or absence and/or severity of a particular disease or condition. In other words, "biomarker" is defined as a laboratory measurement that reflects the activity of a disease process. Examples of biomarkers are, but are not limited to, proteins, metabolites, genes, DNA and RNA. Biomarkers, as disclosed herein, refers to isolated biomarkers. Evaluation of such biomarkers and their correlation to a pathological condition or disease can be done by, for example, determining the absence or presence of a marker, differences in expression levels of the same marker in different clinical settings, and/or comparative analysis between diseased and disease-free samples.

[0035] As used herein, the term "surrogate marker" refers to the measurement of biomarker levels in bodily fluid that are indicative of an active biological process or signalling pathway, or clinicopathological grade of disease. For example, surrogate markers described herein refer to one or more biomarkers which can be used as a substitute for or as a proxy for the intended target. Thus, as used herein, a surrogate marker can also refer to a biomarker panel that serves as a substitute parameter for, for example, the level of STAT3 activation in cells via the analysis of patient ascites. For example, as shown herein, biomarkers listed herein can be used as surrogate markers for STAT3 phosphorylation.

[0036] As used herein, the term "PAI-1" refers to plasminogen activator inhibitor-1 (PAI-1), also known as endothelial plasminogen activator inhibitor or serpin E1. PAI-1 is a protein encoded by the SERPINE1 gene in humans. PAI-1's main function is the inhibition of urokinase-type plasminogen activator (uPA) and tissue-type plasminogen activator (tPA), enzymes responsible for the cleavage of plasminogen to

form plasmin. Plasmin mediates the degradation of the extracellular matrix, either by itself or in conjunction with matrix metalloproteinases. In this scenario, PAI-1 inhibits urokinase-type plasminogen activator via active site binding, preventing the formation of plasmin. Additional inhibition is mediated by PAI-1 binding to the urokinase-type plasminogen activator (uPA)/urokinase-type plasminogen activator receptor (uPAR) complex, resulting in the latter's degradation. Thus, PAI-1 can be said to inhibit the serine proteases tissue-type plasminogen activator (tPA) and urokinase-type plasminogen activator (uPA)/urokinase, and hence is an inhibitor of fibrinolysis, the physiological process that degrades blood clots. In addition, PAI-1 inhibits the activity of matrix metalloproteinases, which play a crucial role in invasion of malignant cells through the basal lamina. In humans, PAI-1 is mainly produced by the endothelium (cells lining blood vessels), but is also secreted by other tissue types, such as adipose tissue and stromal tissue.

[0037] As used herein, the term "PAI-1 inhibitor" refers to compounds that are capable of inhibiting or blocking the activity of Plasminogen activator inhibitor-1 (PAI-1). Various compounds and drugs are not limited to a single effect and can therefore be considered to be PAI-1 inhibitors, even if they are structurally different. That is to say, the inhibition of PAI-1 is the combining characteristic of these compounds.

[0038] As used herein, the term "ascites" refers to an abnormal accumulation of fluid within the abdomen. There are many causes of ascites, including but not limited to, cirrhosis of the liver, cancer within the abdomen, congestive heart failure, and tuberculosis. The term "ascites" can also refer to free fluid in the peritoneal cavity. As used herein, when referring to ascites used in the context of treatment, for example, when exposing cells in cell culture to cell-free ascites, this refers to contacting cells *in vitro* to cell-free ascites fluid obtained from a subject, in order to elucidate changes in biomarker levels and observe the overall change in the molecular or physical phenotype of cells.

[0039] As used herein, the term "cell-free ascites" refers to the supernatant component of ascites derived from, for example, patients. The cell-free ascites as referred to herein was collected from the peritoneal cavity at the beginning of the cytoreductive surgery (CRS) or during routine ascitic tap (paracentesis) and was subjected, for example, to centrifugation at 2000 g for 10 minutes to separate the cellular component from the fluid component. Filter-sterilisation using 0.22 µm filter was performed on the fluid component to render it suitable for downstream experiments. A person skilled in the art would appreciate that other sterilisation methods known in the art can be used in order to obtain cell-free ascites suitable for downstream applications.

[0040] As used herein, the term "phosphorylation" refers to a process whereby a protein kinase transfers a phosphate group from an adenosine triphosphate (ATP) or guanosine triphosphate (GTP) to one or more free hydroxyl groups of amino acids. Generally speaking, phosphorylation is one of the on-off switches used in signalling cascades and pathways. Depending on the context of the pathway in question, phosphorylation can be used as an "on" or "off" switch. By way of an example of STAT3 phosphorylation (also termed "p-STAT3" in the present disclosure), in STAT3 signalling, the

phosphorylation of critical amino acid residue (such as Tyrosine 705) on STAT3 induces the formation of STAT3 dimers, which then translocate into the nucleus to regulate specific gene expression and trigger downstream signalling cascades in the cell.

[0041] As used herein, the term “STAT3” refers to the signal transducer and activator of transcription 3, a transcription factor encoded by the STAT3 gene. In response to growth factors, hormones and cytokines, STAT3 is phosphorylated by upstream receptor kinase, thus undergoing dimerization prior to translocation into the nucleus, where the STAT3 dimer acts as a transcription activator. However, a STAT3 pathway can also be activated via a non-canonical pathway, independent of the upstream receptor kinase (see, for example, Interferon Independent Non-Canonical STAT Activation and Virus Induced Inflammation (Viruses. 2018 Apr; 10(4): 196)).

[0042] As used herein, the term “p-STAT3 activation level” is used interchangeably with the term “STAT3 phosphorylation”, “STAT3 activation”, or “level of STAT3 phosphorylation”.

[0043] As used herein, the term “sample” refers to a biological sample, which includes, but is not limited to, any quantity of a substance from a living thing or formerly living thing. Such living things include, but are not limited to, humans, mice, monkeys, rats, rabbits, and other animals. Such substances include, but are not limited to bodily fluids, such as blood, plasma, ascites, serum, urine, cells, organs, tumour samples, biopsy samples, tissues, bone, bone marrow, lymph, lymph nodes, and skin. Such samples can be obtained from subjects known to suffer from the disease, subjects thought to suffer from the disease, and disease-free subjects. A person skilled in the art will appreciate that each type of sample could require different (pre-) processing steps before being able to be used in the claimed methods. By way of various examples, for samples that are in liquid form, centrifugation would need to be performed to separate the cellular and soluble components. For samples in solid form, tissue dissociation would need to be performed using a combination of mechanical dissociation and enzymatic treatment to create single-cell suspensions which can be centrifuged to separate the supernatant and cellular component. Both supernatant/soluble component and cellular component can then be evaluated via our *in vitro* and *in vivo* experiments. A person skilled in the art would be aware of the methods required in order to obtain samples suitable for use in the methods disclosed herein.

[0044] As used herein, the term “peritoneal carcinomatosis” refers to the intra-abdominal spread of cancer, whereby the origin of the carcinomatosis can be a malignancy arising from an intra-abdominal organ, or from the peritoneum (a thin layer of tissue that lines most of the abdominal organs) itself.

[0045] As used herein, the term “cytoreductive surgery (CRS)” refers to the complete removal of macroscopic tumour found in the abdominal cavity, via a series of peritonectomy and visceral resections.

[0046] As used herein, the term “hyperthermic intraperitoneal chemotherapy” refers to a therapy that is used in the eradication of microscopic disease left behind following cytoreductive surgery, which involves the addition of a heated solution of chemotherapeutic drug(s) into the abdominal cavity for 60 to 90 minutes.

[0047] As used herein, the term “paracrine factors” refers to diffusible and soluble proteins secreted by cells to modulate cellular responses in adjacent cells or the cell of origin via paracrine or autocrine interaction. Examples of such paracrine factors are, but are not limited to, interleukin 6 (IL6), transforming growth factor beta (TGF- β), Wnt proteins, Sonic Hedgehog (SHH), vascular endothelial growth factor (VEGF) and epidermal growth factor (EGF).

[0048] As used herein, the term “oncogenic addiction”, refers to a phenomenon whereby cells, when exposed to a certain paracrine factor, are led to the activation of a cellular signalling cascade. For example, STAT3 activation leads to the production and secretion of more of the same paracrine factor, leading to a positive feedback loop (see, for example, Fig. 18). These cells harness the positive feedback biological cycle for growth and activate pathway activation, and is hence addicted to this process. In the same logic, the term “oncogenic addiction to PAI-1” refers to the situation in which PAI-1 activation leads to production of more PAI-1, therefore leading to a positive, PAI-1-based, feedback loop. If the generation of such a positive feedback loop is prevented, the cells, devoid of a critical stimulus which they become accustomed to (that is addicted to), will die.

DETAILED DESCRIPTION

[0049] Presence of ascites in colorectal peritoneal carcinomatosis portends a poor prognosis. It is hypothesised that ascites are biologically relevant, and can be exploited for novel therapy. Exploiting tumour biology to identify novel therapeutic strategies in this disease is shown to have tremendous clinical impact. As shown herein, small molecule inhibitors targeting major signalling pathways in colorectal peritoneal carcinomatosis can be used in the clinical setting.

[0050] Thus, disclosed herein are methods which enable the targeted treatment of patients with peritoneal carcinomatosis. Also shown herein is that, for example, small molecule inhibitors targeting major signalling pathways can be used in the treatment of colorectal peritoneal carcinomatosis, or that these inhibitors can be used in the following clinical settings: in a neoadjuvant setting, to decrease tumour burden in patients who are not candidates for cytoreductive surgery (CRS) and hyperthermic intraperitoneal chemotherapy (HIPEC, also known as IPHC - Intraperitoneal hyperthermic chemoperfusion) to convert them into candidates for cytoreductive surgery and hyperthermic intraperitoneal chemotherapy; in an adjuvant setting, by adding small molecule inhibitors to a hyperthermic intraperitoneal chemotherapy regimen to improve the efficacy in eradicating residual microscopic disease after cytoreductive surgery; in a palliative setting, to decrease debilitating symptoms from peritoneal disease; and in a prophylactic setting, in patients with colorectal cancer who are at high risk of developing peritoneal carcinomatosis.

[0051] Currently, the only form of cure or standard of care for patients having or suffering from peritoneal carcinomatosis is to perform cytoreductive surgery and instil hyperthermic intraperitoneal chemotherapy at the end of the operation. However, current hyperthermic intraperitoneal chemotherapy regimens do not harness knowledge of tumour biology for therapy and merely uses conventional

chemotherapy in the form of a cytotoxic drug. Moreover, patients who qualify for cytoreductive surgery and hyperthermic intraperitoneal chemotherapy only constitute 10% of all peritoneal carcinomatosis patients.

[0052] In the scope of the present disclosure, biomarkers have been identified that can predict response of treatment with intraperitoneal (IP) instillation of a PAI-1 inhibitor (for example, but not limited to, TM5441). Various groups of patients have been identified who are thought to respond to this therapy using the methods disclosed herein. One such group comprises patients with a high concentration of PAI-1 (≥ 20 ng/mL) and concurrently high STAT3 activation (≥ 0.2 OD450), which, without being bound by theory, are thought to be highly susceptible to PAI-1 inhibition. Another group comprises patients with a lower concentration of PAI-1 (< 20 ng/mL) compared to the first group, but high STAT3 activation (≥ 0.2 OD450). Patients in this group are also considered to be susceptible to PAI-1 inhibition. This can be seen, for example, in ascites having high PAI-1 levels and concurrently activate STAT3 signalling in, for example, cancer cells. In the clinical context, intraperitoneal instillation of PAI-1 inhibitor can be used in the neoadjuvant, at the time of hyperthermic intraperitoneal chemotherapy or even in the palliative setting, hence providing a therapeutic option to much more patients than those who qualify for cytoreductive surgery and hyperthermic intraperitoneal chemotherapy. It is also thought that PAI-1 inhibitor can be aerosolized to be used in the palliative setting, for example in pressurized intraperitoneal aerosol chemotherapy (PIPAC).

[0053] In addition, the data generated shows that this strategy applies to patients with colorectal peritoneal carcinomatosis, and that it can also be applied to patients with other histological subtypes of peritoneal carcinomatosis.

[0054] Thus, in one example, subtypes of the peritoneal carcinomatosis can be but are not limited to, colorectal peritoneal carcinomatosis, small bowel peritoneal carcinomatosis, mesothelioma, endometrial peritoneal carcinomatosis, gastric peritoneal carcinomatosis, ovarian peritoneal carcinomatosis, appendiceal peritoneal carcinomatosis, pancreatic peritoneal carcinomatosis, urothelial peritoneal carcinomatosis and Pseudomyxoma peritonei (PMP). In another example, the peritoneal carcinomatosis is of unknown origin. In one example, the subtype of peritoneal carcinomatosis is colorectal peritoneal carcinomatosis.

[0055] As used herein, the term “unknown primary” when used in conjunction with tumours, tumour samples or subtypes, refers to the presence of peritoneal carcinomatosis where the primary tumour is undetermined by clinical, radiological and pathological assessment. Such indetermination can be due to reasons such as, but not limited to, the (small) insufficient size of primary tumour for pathological assessment, or that the primary tumour is encased by extrinsic peritoneal carcinomatosis such that clinical detection is not possible, or the lack of tumour markers with high specificity.

[0056] In the context of the experiments disclosed in the present disclosure, the term “other peritoneal carcinomatosis (PC) histologies” refers to histological subtypes of peritoneal carcinomatosis that originate from the following sites, namely, lung, breast, peritoneum, synchronous gastric and ovary, small bowel,

urothelial, and palate. These samples are grouped under “other PC histologies” due to the small number of samples collected in each subgroup.

[0057] In one example, the peritoneal carcinomatosis is malignant. In another example, the peritoneal carcinomatosis is a primary tumour. In yet another example, the peritoneal carcinomatosis is a metastasis, or secondary tumour.

[0058] In another example, there is disclosed a method of predicting, determining or detecting susceptibility of a subject suffering from peritoneal carcinomatosis to a treatment with an anti-cancer drug or an anti-cancer treatment instilled within the peritoneal cavity based on the levels of PAI-1 within the ascitic fluid in the peritoneal cavity. Without being bound by theory, it is thought that STAT3 activation in cancer cells exposed to ascitic fluid allows the identification of a subgroup of patients who would benefit from inhibition of PAI-1.

[0059] As used herein, the term “susceptibility” refers to the propensity of something, for example a disease, to be likely affected by something else, for example, a treatment for said disease. This effect can be either positive or negative, depending on what is being referenced. For example, if a disease is sensitive to a particular treatment, then the susceptibility of said disease to a particular treatment is a positive effect. It can then be said that the disease is susceptible (or sensitive) to the treatment. On the other hand, if a disease is not susceptible to a given treatment, the disease is then considered to be unresponsive or resistant to said treatment.

[0060] As defined above, the term “predicting susceptibility” refers to the propensity of something, for example a disease, to be likely affected by something else, for example, a treatment for said disease. In other words, to predict susceptibility of a cancer to a particular treatment is to determine whether the cancer would react to a treatment with a certain medicament, or anti-cancer drug, or anti-cancer treatment. It is of note that the term “determining susceptibility” is not synonymous with, for example, “making a prognosis”. The former term only looks at the possible reaction of a disease to a specific drug or therapy, while the latter describes the clinical outcome of the patient defined by parameters such as, but not limited to, the length of time of stable disease (once such a status is acquired), the length of time of overall survival and/or disease-free survival. While in some cases, it may be possible to correlate the effect of one term on the other, that is to say that a disease reacting well to a given treatment (that is, the disease is susceptible to the treatment) may increase the likelihood of said patient receiving a positive prognosis in regards to the overall disease progression, this is not to be taken as a rule. As a person skilled in the art would appreciate, a positive prognosis depends on many patient-specific factors in addition to the disease’s susceptibility for treatment, for example, overall well-being of the patient prior to treatment, metabolism, diet, aggressiveness of the (primary) disease, secondary diseases and/or infections and the like.

[0061] Also disclosed herein is a method of predicting, determining or detecting susceptibility of a subject suffering from peritoneal carcinomatosis to a treatment with an anti-cancer drug or anti-cancer treatment.

[0062] Firstly, it was identified from clinical data in peritoneal carcinomatosis patients operated on that the presence of clinically apparent ascites within the peritoneal cavity leads to a poorer prognosis compared to patients who did not have clinically apparent ascites. This prognostic significance is relevant in peritoneal carcinomatosis of colorectal origin, although it is not limited to this histological subtype (Fig. 1). It is noted that the basis of comparison for the term “poorer prognosis” is the following: A group of patients with poorer prognosis are those patients with clinically apparent ascites during surgery as compared to the ones who do not have clinically apparent ascites during surgery. As used herein, the term “clinically apparent ascites” refers to ascites present in a volume of, for example, 50ml or more during surgery.

[0063] *In vitro* treatment of established cell line models of peritoneal carcinomatosis (Colo-205, HM3-TERT and LP9-TERT) with cell-free ascites collected from patients showed increased proliferation, migration, and establishment of colonies on stromal cells in co-culture models (Fig. 2), indicating that the cell line models of peritoneal carcinomatosis closely mimic the tumour in its *in vivo* setting.

[0064] Gene expression analysis of peritoneal carcinomatosis cell lines treated with cell-free ascites showed that the STAT3 pathway was activated (Fig. 3), indicating that the activation of STAT3 plays an important role in the disease.

[0065] Western blot analysis of STAT3 phosphorylation in cells treated with cell-free ascites revealed that STAT3 signalling pathway was activated through phosphorylation at Tyr705 (Fig. 4a). Screening of cell-free ascites collected from different histological subtypes showed that STAT3 activation was the most prevalent when cells were treated with cell-free ascites collected from colorectal peritoneal carcinomatosis compared to peritoneal carcinomatosis from other anatomical origins (Fig. 4b). Thus, this data shows that cell-free ascites from colorectal peritoneal carcinomatosis induces a higher rate of STAT3 activation compared to cell-free ascites from peritoneal carcinomatosis of other anatomical origins. It is further of note that cell-free ascites collected from peritoneal carcinomatosis of other anatomical origins can also activate STAT3 signalling.

[0066] Immunohistochemistry of primary colorectal cancers matched with peritoneal metastases collected from the same patients showed that STAT3 activation was more prevalent in the metastases compared to the primary tumour (Fig. 5). Thus, this data shows that treatment of depriving metastatic cells of STAT3 signalling works well compared to the strategy of depriving primary tumours of STAT3 signalling. This is because STAT3 activation is more prevalent in metastases than primary tumours. As shown in the data presented herein, the subset of patients whose tumours have high STAT3 signalling due to high PAI-1 levels are more susceptible to PAI-1 inhibition compared to tumours whose STAT3 signalling is not activated by PAI-1. This is even less so in the tumours that do not show any activation in STAT3 signalling.

[0067] Treatment of established cell line models of colorectal peritoneal carcinomatosis with cell-free ascites also led to the enrichment of the epithelial-mesenchymal transition (EMT) signature (Fig. 6a). An unbiased mass spectrometry screen of cell-free ascites along with a cytokine array performed on cell-free

ascites of different histological origins identified the importance of coagulation/thrombolytic factors in cell-free ascites from colorectal peritoneal carcinomatosis (Fig. 6b). It is also shown that PAI-1 (which is involved in prevention of coagulation) is highly enriched in cell-free ascites from colorectal peritoneal carcinomatosis (Fig. 6c). This means that by analysing the proteomics of cell-free ascites, either via mass spectrometry or cytokine array, for example, it was found that the coagulation pathway was enriched in colorectal peritoneal carcinomatosis. From a marker candidate perspective, PAI-1, which is involved in the coagulation cascade, was also shown to be highly abundant. Thus, a phenomenon is being described whereby the presence of an active coagulation pathway is hijacked by cancer cells for oncogenic activation. In other words, and without being bound by theory, it is thought that the presence of activation of the coagulation pathway within the peritoneal cavity leads to oncogenic activation of signalling pathways in cancer cells initiated by coagulation factors or factors involved in prevention of coagulation. It is not intended to describe coagulation in the physical sense of having blood clots in the abdomen.

[0068] Interrogation of The Cancer Genome Atlas (TCGA) database by the inventors showed that colorectal cancers with high levels of PAI-1 activated STAT3 signalling (Fig. 7a) and displayed enrichment of the epithelial-mesenchymal transition signature (Fig. 7b), and had the poorest prognosis (Fig. 7c). Taken together, it was shown that PAI-1 within the ascites can lead to STAT3 activation in cancer cells when these cancer cells are exposed to ascites, culminating in an epithelial-mesenchymal transition (EMT) phenotype that is responsible for the clinical manifestation of a biological aggressive tumour, leading to poor prognosis in these patients. Of note, activation of JAKs was not found in cell lines treated with cell-free ascites, suggesting that ascites activates STAT3 signalling in a non-canonical fashion (Fig. 8). This means that in this situation, STAT3 is activated in a non-canonical fashion, instead of the canonical fashion. Hence, without being bound by theory, it is thought that STAT3 can be activated by other activators (for example, PAI-1), instead of the canonical activator of STAT3 such as, for example, IL6.

[0069] Thus, in one example, the sample is, but is not limited to, ascites, blood, serum, urine, drain fluid, surgical drain fluid, liquid bodily fluids, supernatant obtained from cells, supernatant obtained from organs, supernatant obtained from tissues, lymph, supernatant obtained from lymph nodes, liquid biopsy samples, and supernatant obtained from biopsy samples.

[0070] Supernatant obtained from organs, tissues and the like can refer to liquid obtained from, for example, an organ sample which is macerated, minced, ground or crushed after extraction. Alternatively, for samples that contain little to no fluid, the sample can be placed in a clinical compatible buffer prior to or after mincing. The resulting liquid is termed a supernatant, which can then be used downstream for further analysis.

[0071] In another example, the sample is a liquid sample. In yet another example, the methods disclosed herein can be performed on one or more samples. For example, a method disclosed herein can be performed on two samples. In another example, the determining or measuring of the concentration of

PAI-1 can be performed on one sample, and the determining or measuring of the level of STAT3 activation (for example, by way of phosphorylation) can be performed on another sample. These samples can be of the same or different origins. In one example, the first sample can be a cell-free sample, and the second sample can be a sample containing cells. In another example, the first sample can be ascites, and the second sample can be a biopsy sample. In one example, the concentration of PAI-1 and STAT3 activation (for example, by way of phosphorylation, or by way of surrogate markers) can be measured in a single sample. In other words, the determination of the concentration of PAI-1 and STAT3 activation can be performed on a single sample.

[0072] Having understood that PAI-1 is upstream of STAT3 activation in cancer cells via paracrine signalling, the levels of PAI-1 were systematically elucidated in cell-free ascites collected from patients with peritoneal carcinomatosis. An established cell line model of peritoneal carcinomatosis, Colo-205, was also treated with these cell-free ascites and levels of p-STAT3 elucidated using enzyme-linked immunosorbent assay (ELISA) to establish the magnitude of STAT3 activation. Plotting the levels of PAI-1 in ascites ($\log_2$; x-axis) with the degree of STAT3 phosphorylation (leading to STAT3 activation) (y-axis), it was identified that the PAI-1 levels in ascites were positively correlated with the levels of STAT3 activation in cell-free ascites treated cells, both in the context of colorectal peritoneal carcinomatosis (PC) cell-free ascites, as well as cell-free ascites collected from peritoneal carcinomatosis (PC) of other histological subtypes (Fig. 9a, b).

[0073] The untransformed PAI-1 levels in cell-free ascites were then analysed with the corresponding degree of STAT3 phosphorylation in peritoneal carcinomatosis (PC) cells exposed to these ascites. Setting the phosphorylation of STAT3 (Tyr705) above 0.2 (OD 450) as a definition of activated STAT3 signalling, it was noted that all samples with PAI-1 level above 20ng/mL showed activated STAT3 signalling. This observation prompted the definition of three (sub-)groups as shown in the following section.

[0074] Firstly, ascites with high PAI-1 levels (that is, PAI-1 levels of more than 20ng/ml) were shown to rely heavily on PAI-1 to activate STAT3 signalling. These samples were termed PAI-1 paracrine addicted (PPA). Treatment of cell lines with cell-free ascites collected from these patients led to high levels of STAT3 phosphorylation (high STAT3 activity). Without being bound by theory, it is thought that STAT3 activation in cancer cells is likely exclusively dependent on PAI-1 levels within cell-free ascites collected from this group of patients, highlighting the phenomenon of oncogenic addiction to upstream activator of this pathway. Secondly, cell-free ascites with low PAI-1 levels (that is to say, PAI-1 levels of less than 20ng/ml) and with activated STAT3 signalling in cells exposed to these cell-free ascites were termed co-activators predominant (CAP). Despite having low PAI-1 levels, treatment of cell lines with cell-free ascites collected from these patients still led to high levels of STAT3 phosphorylation (high STAT3 activity). Without being bound by theory, it is thought that in this group, STAT3 signalling was likely to be activated by PAI-1 and a combination of other ligands. Finally, cell-free ascites with low PAI-1 levels (that is to say, PAI-1 levels of less than 20ng/ml) and which failed to activate STAT3 signalling likely

had ligands that activated other signalling pathways. These samples were termed alternative pathways activation (APA). Treatment of cell lines with cell-free ascites collected from these patients did not lead to a significant level of STAT3 phosphorylation (low STAT3 activity). Figures 10a and 10b highlight that the classification of the different forms of cell-free ascites was applicable to both cell-free ascites of colorectal peritoneal carcinomatosis origin, as well as those from other histological subtypes.

[0075] To confirm this theory, Colo-205 cell line were treated in the presence of cell-free ascites from the 3 different sub-groups with TM5441 (PAI-1 inhibitor), with the expectation that cells exposed to PAI-1 paracrine addicted (PPA) cell-free ascites will be highly sensitive towards PAI-1 inhibition. As predicted, differential sensitivity to PAI-1 inhibition according to the PAI-1 and p-STAT3 gating (Fig. 11) was observed. Treatment with another PAI-1 inhibitor (Tiplaxtinin) also showed the same trend of differential response, highlighting that the PAI-1 inhibition is specific, and that the results observed are not due to cytotoxic effects of the inhibitors (Fig. 12a). Subsequently, a STAT3 inhibitor (Napabucasin), a dual PI3K/mTOR inhibitor (BEZ235) and a conventional chemotherapeutic agent used in hyperthermic intraperitoneal chemotherapy (HIPEC) for treatment of colorectal peritoneal carcinomatosis (PC) (Mitomycin C) were tested to compare the efficacy in inhibition of oncogenic addiction to PAI-1 versus inhibition of downstream signalling pathway activated by cell-free ascites and cellular proliferation. It was found that direct targeting of cancer cells in the presence of cell-free ascites is ineffective because cell-free ascites promote chemoresistance in these tumour cells (Fig. 12b-d).

[0076] To further investigate the downstream signalling pathways involved in determining the sensitivity to PAI-1 inhibition, RNA microarray of Colo-205 cells treated with cell-free ascites representative of PAI-1 paracrine addicted (PPA) (PC085), co-activators predominant (CAP) (PC249), or foetal bovine serum (FBS; control) in the presence of TM5441 or DMSO vehicle was performed. Gene set enrichment analysis (GSEA) identified IL6-JAK-STAT3 signalling pathway to be significantly down-regulated in PAI-1 paracrine addicted (PPA) group upon PAI-1 inhibition (Fig. 13a). This finding is consistent with the initial hypothesis that the highly sensitive PAI-1 inhibition in PAI-1 paracrine addicted (PPA) group is attributed to PAI-1-STAT3 signalling pathway. Similarly, measurement of p-STAT3 in cells treated with these cell-free ascites and TM5441 demonstrated differential concentration needed to abrogate STAT3 activation (Fig. 13b).

[0077] As proof of concept that this is not exclusively a biological observation *in vitro*, Colo-205 cells were co-injected with cell-free ascites or foetal bovine serum (FBS) intraperitoneally in BALB/c nude mice to create a peritoneal carcinomatosis (PC) model. These mice were treated with intraperitoneal (i.p.) injection of TM5441. Consistent with the *in vitro* results, significant reduction in tumour burden was observed in PAI-1 paracrine addicted (PPA) cell-free ascites-treated mice (Fig. 14). In one example, the optimal drug delivery route was then assessed by comparing i.p. injection and oral administration of TM5441. I.p. instillation of TM5441 greatly outperformed oral administration in reducing tumour burden in peritoneal carcinomatosis (PC) mouse model (Fig. 15). This finding is in line with what has been observed in peritoneal carcinomatosis (PC) patients, where systemic administration of drugs has generally

been considered to be ineffective due to the peritoneal-plasma barrier, leading to diminished penetration of cytotoxic agents from plasma into peritoneal tumours and ascites.

[0078] Subsequently, two patient-derived ascites-dependent xenografts (PDADXs) were developed, one from PAI-1 paracrine addicted (PPA) group (PC383 patient-derived ascites-dependent xenografts (PDADX)) and one from co-activators predominant (CAP) group (PC249 patient-derived ascites-dependent xenografts (PDADX)), to better recapitulate the PAI-1 addiction theory as an avatar of peritoneal carcinomatosis (PC) patients. Morphological evaluation of the patient-derived ascites-dependent xenograft (PDADX) tumours showed signet ring cell morphology, resembling the histology of the original patients' tumour. Immunohistochemical staining also confirmed that the patient-derived ascites-dependent xenograft (PDADX) tumours are of colonic origin (Fig. 16).

[0079] When treated with TM5441, PC383 patient-derived ascites-dependent xenograft (PDADX) mice exposed to matched cell-free ascites from the same patient elicited a significantly superior inhibition of tumour growth compared to vehicle control and to foetal bovine serum (FBS) group. In contrast, PC249 patient-derived ascites-dependent xenograft (PDADX) mice, which had been exposed to its matched patient's cell-free ascites and treated with TM5441, showed no reduction in tumour burden compared to vehicle control, similar to that of foetal bovine serum (FBS) group (Fig. 17a). When PC249 patient-derived ascites-dependent xenograft (PDADX) mice were exposed to cell-free ascites from PAI-1 paracrine addicted (PPA) group (PC383 ascites), these tumour cells became susceptible to PAI-1 inhibition, despite not being susceptible to PAI-1 inhibition in the presence of its own matched cell-free ascites (Fig. 17b). Taken together, this information describes a previously unknown phenomenon of oncogenic addiction in the context of a closed biological system, where tumours, along with their microenvironment, are segregated from the systemic circulation. Paracrine factors in this context provide the key stimulus for pathway activation; paracrine inhibition provides the critical stop point (Fig. 18).

[0080] The patient-derived ascites-dependent xenograft (PDADX) model has two components. The first component is the solid tumour that is formed by allowing the cellular components from ascites to form nodules in the host (usually in mice). The second component is cell-free ascites collected from the same patient from which the solid tumours had been obtained. The cell-free ascites is co-injected with the cellular component that is being propagated in the mice. This is a model considers the intrinsic phenotype of the cells, as well as the paracrine environment of the tumours within the peritoneal cavity.

[0081] As proof of concept, that the method disclosed herein is able to subclassify patient ascites into PAI-1 paracrine addicted (PPA), co-activators predominant (CAP) and alternative pathways activation (APA) groups, it was sought to identify surrogate markers of STAT3 activation in cells by analysing the cell-free ascites of patients. Briefly, STAT3-related genes were identified from Kyoto Encyclopedia of Genes and Genomes (KEGG) database by compiling all genes that are involved in known STAT3 pathways. Secreted STAT3-related proteins were selected based on extracellular genes listed in NCBI's Biosystems database and proteins that were identified by mass spectrometry analysis of cell-free ascites. Transcriptomics comparisons were performed using two databases to prioritize putative STAT3 surrogate

markers, and to identify genes that are down-regulated and up-regulated in PAI-1 paracrine addicted (PPA) cell-free ascites-treated cells in response to TM5441 (PAI-1 inhibition). Genes were ranked from most down-regulated to most up-regulated, and systematic paired correlation analysis of candidate genes was subsequently performed. The paired analysis for each group was prioritized, as shown in Fig. 19b, and representative genes were chosen from each group based on literature review to streamline the selection to 35 genes. Targets were selected for further evaluation with enzyme-linked immunosorbent assay (ELISA) (Fig. 19) based on rank prioritisation, potential good correlation with p-STAT3 from Luminex assay data, and the importance of the candidate genes in cancer pathogenesis from literature review. Validating this in a cohort of 40 to 70 patients, a 4-biomarker panel was identified that can identify levels of STAT3 activation in cells via patient cell-free ascites analysis (Fig. 20 and Fig. 21). This finding also serves as the basis of, for example, a point of care stratification kit for patients who would benefit from PAI-1 therapy.

[0082] As described above, the present disclosure highlights exemplary cut-off levels of PAI-1 within the cell-free ascitic fluid in patients with peritoneal carcinomatosis and, when coupled with the levels of STAT3 activation in cancer cells exposed to these cell-free ascitic fluids, identifies a subgroup of patients who would benefit from inhibition of PAI-1.

[0083] In one example, the concentration of PAI-1 is between 0 to 450 ng/ml, between 10 to 20 ng/ml, between 15 to 25 ng/ml, or between 19 to 29 ng/ml. In one example, the concentration of PAI-1 is between 0 to 17 ng/ml. In another example, the concentration of PAI-1 is between 0 to 20 ng/ml. In another example, the concentration of PAI-1 in the context of the present invention is either less than 20 ng/ml, or more than or equals to 20 ng/ml.

[0084] In another example, the level of STAT activation, as measured by phosphorylation, is between 0 to 1.7, as measured at an optical density of 450 nm (OD450). In one example, the level of STAT3 activation, as measured by phosphorylation, is between 0.01 to 1, between 0.1 to 0.5, between 0.05 to 0.19, between 0.18 to 0.26, between 0.24 to 0.48, between 0.35 to 0.5, about 0.08, between 0.4 to 0.6, between 0.5 to 0.75, between 0.65 to 0.8, between 0.79 to 0.90, between 0.88 to 0.95, between 0.9 to 1, about 0.1, about 0.15, about 0.17, about 0.18, about 0.19, about 0.2, about 0.21, about 0.22, about 0.23, about 0.24, about 0.25, or about 0.3, as measured at an optical density of 450 nm (OD450). In another example, the level of STAT3 activation, as measured by phosphorylation, in the context of the present invention is either less than 0.2 ng/ml (OD450), or more than 0.2 (OD450).

[0085] The recitation of the term “0” (zero) is included as a value based on the understanding that “0” is also used when the presence of a marker is, for example, below the detection limit of a kit or a detection method, and is therefore immeasurable. In such cases, the observed value is often denoted as “n.a.” or “nil”.

[0086] In order to include most of the PAI-1 paracrine addicted (PPA)/co-activators predominant (CAP) samples, the surrogate biomarker values of the 60 PAI-1 paracrine addicted (PPA)/co-activators predominant (CAP) samples at the lower 5% percentile were selected as the initial cut-offs (extended

data, not shown). For MMP9, the biomarker value with the highest value in the alternative pathways activation (APA) samples, 13 ng/ml was selected to exclude all alternative pathways activation (APA) samples. Based on a panel of 3 or 4 initial cut-offs, 52 PAI-1 paracrine addicted (PPA)/co-activators predominant (CAP) samples and two alternative pathways activation (APA) samples were filtered to be the PAI-1 paracrine addicted (PPA)/co-activators predominant (CAP) samples (extended data, not shown), corresponding to 86.67% and 80.0% accuracy, respectively.

[0087] In order to enhance the accuracy, more stringent cut-off values of IL6 and IL10 were chosen in order to exclude the false-positive alternative pathways activation (APA) samples, and less stringent cut-off values of CCL2 and MMP9 were chosen in order to include the false-negative PAI-1 paracrine addicted (PPA)/co-activators predominant (CAP) samples. The values of the final cut-offs are shown in Fig 20c. Based on a panel of 3, or 4, final cut-offs, 56 PAI-1 paracrine addicted (PPA)/co-activators predominant (CAP) samples and one alternative pathways activation (APA) sample were filtered to be the PAI-1 paracrine addicted (PPA)/co-activators predominant (CAP) samples (extended data, not shown), corresponding to 93.33% and 90.0% accuracy, respectively. The overall accuracy of this composite biomarker panel (passing ≥ 3 biomarker cut-offs) is 92.86% (Fig. 20b).

[0088] Based on the information provided herein, it has been shown that IL6 alone is the statistically most robust predictor of the degree of STAT3 phosphorylation (p-STAT3), based on Stepwise Method and Best Subset Methods with Akaike Information Criterion or Bayesian Information Criterion. Generally speaking, regression analysis is a statistical approach to assess whether a set of independent variables significantly influences the dependent variable. Stepwise Regression is a method of fitting regression models by automatically adding or removing individual predictors and selecting a single model based on statistical significance. Best Subsets Regression is a method of comparing all possible models, using a specified set of predictors, and displays the best-fitting models that contain one predictor, two predictors, and so on. Akaike information criterion is an estimator of out-of-sample prediction error and relative quality of each model, thus providing a means for model selection. Bayesian Information Criterion is a criterion for model selection, among a finite set of models, based on likelihood function, solving the problem of potential overfitting by adding more parameters.

[0089] For example, a cut-off of IL6 at 997 pg/ml can define PAI-1 paracrine addicted (PPA)/co-activators predominant (CAP) samples with 95% accuracy and exclude an alternative pathways activation (APA) samples with 80% accuracy, corresponding to 92.86%. It is further shown that overall prediction accuracy can be increased with an increase in the number of biomarkers used in a composite biomarker panel. For example, a panel of IL6, IL10, CCL2 and MMP9 with a criterion of 3 or 4 positive biomarkers, is capable of identifying a PAI-1 paracrine addicted (PPA)/co-activators predominant (CAP) subject with 93.33% accuracy, and exclude an alternative pathways activation (APA) subject with 90% accuracy, corresponding to 92.86% overall accuracy. Other exemplary panels can be found throughout the present specification. The above means that the minimum number of markers required to obtain a statistically robust outcome is one biomarker (for example, IL6). These biomarkers can be, but are not limited to, the

biomarker listed herein. In one example, the panel or group of biomarkers includes IL6. In one example, a single biomarker is used in the method disclosed herein, wherein the biomarker is, but is not limited to, IL6 (Interleukin 6), IL10 (Interleukin 10), CCL2 (chemokine (C-C motif) ligand 2; also referred to as monocyte chemo-attractant protein 1 (MCP1) or small inducible cytokine A2), MMP9 (matrix metalloproteinase 9, also known as 92 kDa type IV collagenase, 92 kDa gelatinase or gelatinase B (GELB)), TGFB1 (transforming growth factor beta 1), POSTN (Periostin, PN, or osteoblast-specific factor OSF-2), VSIG4 (V-set and immunoglobulin domain containing 4), CD44, and CXCL10 (C-X-C motif chemokine 10, also known as Interferon gamma-induced protein 10 (IP-10) or small-inducible cytokine B10). In another example, two biomarkers are used in the method disclosed herein, wherein the two biomarkers are, but are not limited to, the following combinations: IL6 and IL10; IL6 and CCL2; IL10 and CCL2; IL6 and MMP9; IL10 and MMP9; and CCL2 and MMP9. In yet another example, three biomarkers are used in the method disclosed herein, wherein the three biomarkers are, but are not limited to, the following combinations: IL6, IL10, and CCL2; IL6, IL10, and MMP9; IL6, CCL2, and MMP9; IL10, CCL2, and MMP9. In another example, four biomarkers are used in the method disclosed herein, wherein the four biomarkers are IL6, IL10, CCL2 and MMP9. In yet another example, five biomarkers are used in the method disclosed herein, wherein the five biomarkers are TGFB1, POSTN, VSIG4, CD44 and CXCL10. In another example, six biomarkers are used in the method disclosed herein, wherein the six biomarkers are IL6, TGFB1, POSTN, VSIG4, CD44 and CXCL10. In a further example, the panel disclosed herein comprises the biomarkers, which are but are not limited to, IL6; IL10; CCL2; MMP9; IL6 and IL10; IL6 and CCL2; IL10 and CCL2; IL6 and MMP9; IL10 and MMP9; CCL2 and MMP9; IL6, IL10, and CCL2; IL6, IL10, and MMP9; IL6, CCL2, and MMP9; IL10, CCL2 and IL6; and IL10, CCL2, and MMP9.

[0090] The cut-off values of, for example, four surrogate biomarkers were determined by the screening of 70 patient cell-free ascites. Taking into consideration the flexibility of patient samples, a range of $\pm 5\%$ cut-off value for each biomarker was included. Thus, in one example, the present invention, when referring to cut-off values, refers to the specific cut-off value with a buffer of $\pm 5\%$ or a buffer of $\pm 2\%$.

[0091] In one example, one subgroup or subset of patients is defined as having a PAI-1 level of between 0 to 20 ng/ml, and a p-STAT3 activation level of less than 0.2 (OD450). In another example, one subgroup or subset of patients is defined as having a PAI-1 level of between 0 to 20 ng/ml, and a p-STAT3 activation level of equal to or more than 0.2 (≥ 0.2 ; OD450). In a further example, one subgroup or subset of patients is defined as having a PAI-1 level of equal to or more than 20 ng/ml (≥ 20 ng/ml), and a p-STAT3 activation level of equal to or more than 0.2 (≥ 0.2 ; OD450).

[0092] In yet another example, one subgroup or subset of patients is defined as having a PAI-1 level of between 0 to 17 ng/ml, and a p-STAT3 activation level of less than 0.2 (OD450). In another example, one subgroup or subset of patients is defined as having a PAI-1 level of between 0 to 17 ng/ml, and a p-STAT3 activation level of equal to or more than 0.2 (≥ 0.2 ; OD450). In a further example, one subgroup

or subset of patients is defined as having a PAI-1 level of equal to or more than 17 ng/ml (≥ 17), and a p-STAT3 activation level of equal to or more than 0.2 (≥ 0.2 ; OD450).

[0093] The concentrations of five exemplary surrogate markers (in this case, IL6, IL10, CCL2, MMP9 and ANGPT1), identified using the methods disclosed herein, were plotted against the degree of STAT3 phosphorylation. The resulting graph is shown in Fig 20. Based on Spearman Correlation analysis (R in Fig. 20), IL6, IL10 and CCL2 were selected to be surrogate biomarkers of STAT3 phosphorylation. Although MMP9 shows a weak correlation with phosphorylated STAT3, the concentration of MMP9 in the PAI-1 paracrine addicted (PPA)/co-activators predominant (CAP) samples is significantly higher than that in the alternative pathways activation (APA) samples (unpaired t test, $P < 0.05$). The inclusion of MMP9 as a surrogate biomarker helps to exclude the alternative pathways activation (APA) samples from the PAI-1 paracrine addicted (PPA)/co-activators predominant (CAP) samples.

[0094] Thus, in one example, one subgroup or subset of patients is defined based on the concentration of PAI-1 and p-STAT3. In the event that p-STAT3 is used to determine which subgroup or subset of patients the subject to be tested belongs to (for example, in combination with PAI-1), the measurement of further surrogate markers in addition to p-STAT3 is optional. In another example, no further measurement of surrogate markers is undertaken, if p-STAT3 is measured directly. In another example, if p-STAT3 is not used to determine which subgroup or subset of patients the subject to be tested belongs to, the surrogate markers listed herein are used in place of direct measurements of STAT3 phosphorylation.

[0095] In one example, a concentration of PAI-1 of less than 20 ng/ml indicates that a patient belongs to either the co-activators predominant (CAP) or the alternative pathways activation (APA) subgroup. In another example, a PAI-1 concentration of more than or equals to 20 ng/ml indicates that the patient belongs to the PAI-1 paracrine addicted (PPA) subgroup.

[0096] In one example, a concentration of p-STAT3 of less than 0.2 OD450 indicates that a patient belongs to the alternative pathways activation (APA) subgroup. In another example, a concentration of p-STAT3 of at least 0.2 OD450 or more indicates that a patient belongs to either the PAI-1 paracrine addicted (PPA) or the co-activators predominant (CAP) subgroup.

[0097] In one example, the subgroup or subset of patients is defined as being of the PAI-1 paracrine addicted (PPA) group, if the patient is shown to have a PAI-1 concentration of at least 20 ng/ml or more, and a p-STAT3 concentration of at least 0.2 OD450 or more.

[0098] In one example, the subgroup or subset of patients is defined as being of the co-activators predominant (CAP) group, if the patient is shown to have a PAI-1 concentration of less than 20 ng/ml, and a p-STAT3 concentration of at least 0.2 OD450 or more.

[0099] In one example, the subgroup or subset of patients is defined as being of the alternative pathways activation (APA) group, if the patient is shown to have a PAI-1 concentration of less than 20 ng/ml, and a p-STAT3 concentration of less than 0.2 OD450.

[00100] In one example, the subgroup or subset of patients is defined as being of the PAI-1 paracrine addicted (PPA) group, if the patient is shown to have a PAI-1 concentration of at least 20 ng/ml or more, and an increased p-STAT3 concentration.

[00101] In one example, the subgroup or subset of patients is defined as being of the co-activators predominant (CAP) group, if the patient is shown to have a PAI-1 concentration of less than 20 ng/ml, and an increased p-STAT3 concentration.

[00102] In one example, the subgroup or subset of patients is defined as being of the alternative pathways activation (APA) group, if the patient is shown to have a PAI-1 concentration of less than 20 ng/ml, and a decreased p-STAT3 concentration.

[00103] In one example, the subgroup or subset of patients is defined as being of the PAI-1 paracrine addicted (PPA) group, if the patient is shown to have an increased PAI-1 concentration, and a p-STAT3 concentration of at least 0.2 OD450 or more.

[00104] In one example, the subgroup or subset of patients is defined as being of the co-activators predominant (CAP) group, if the patient is shown to have a decreased PAI-1 concentration, and a p-STAT3 concentration of at least 0.2 OD450 or more.

[00105] In one example, the subgroup or subset of patients is defined as being of the alternative pathways activation (APA) group, if the patient is shown to have a decreased PAI-1 concentration, and a p-STAT3 concentration of less than 0.2 OD450.

[00106] In one example, the concentration of p-STAT3 is measured using one or more surrogate markers, whereby the surrogate markers are, but are not limited to IL6, CCL2, IL10, MMP9, TGFB1, POSTN, VISG4, CD44, CXCL10, and combinations thereof.

[00107] In one example, one subgroup or subset of patients is defined using a panel of markers comprising IL6, CCL2, IL10, MMP9, TGFB1, POSTN, VISG4, CD44, and CXCL10. In another example, one subgroup or subset of patients is defined using a panel of markers comprising IL6, CCL2, IL10, and MMP9. In yet another example, one subgroup or subset of patients is defined using a panel of markers comprising PAI-1 and pSTAT3.

[00108] In one example, the cut-off value for IL6 is a concentration of 997pg/ml. In another example, the cut-off value for IL10 is a concentration of 15 pg/ml. In another example, the cut-off value for CCL2 is a concentration of 450 pg/ml. In yet another example, the cut-off value for MMP9 is a concentration of 3ng/ml. In the above examples, a concentration equal to, or more than, each of the marker-specific cut-off values indicates that the patient belongs to the PAI-1 paracrine addicted (PPA) and co-activators predominant (CAP) subgroup. In other words, the values shown herein can also be termed the cut-off values or “(+)”, for the respective marker. Conversely, if a measured concentration is below the above referenced cut-off value for the same marker, it can be indicated as “(-)” for the respective marker.

[00109] In one example, one subgroup or subset of patients is defined using a panel of markers comprising IL6, CCL2, IL10, and MMP9, whereby a combination of any 2 markers shown to have a

concentration below the cut-off value indicates that the patient belongs to the alternative pathways activation (APA) subgroup.

[00110] In one example, one subgroup or subset of patients is defined using a panel of markers comprising IL6, CCL2, IL10, and MMP9, whereby a combination of any 3 markers shown to have a concentration above the cut-off value indicates that the patient belongs to the PAI-1 paracrine addicted (PPA) and co-activators predominant (CAP) subgroup.

[00111] In another example, one subgroup or subset of patients is defined using a panel of markers comprising IL6, CCL2, IL10, and MMP9, whereby all 4 markers shown to have a concentration above the cut-off value indicates that the patient belongs to the PAI-1 paracrine addicted (PPA) and co-activators predominant (CAP) subgroup.

[00112] In another example, one subgroup or subset of patients is defined using a panel of markers comprising TGFB1, POSTN, VSIG4, CCD44 and CXCL10, whereby all 5 markers shown to have a concentration above the cut-off value indicates that the patient belongs to the PAI-1 paracrine addicted (PPA) and co-activators predominant (CAP) subgroup.

[00113] In yet another example, one subgroup or subset of patients is defined using a panel of markers comprising IL6, TGFB1, POSTN, VSIG4, CCD44 and CXCL10, whereby all 6 markers shown to have a concentration above the cut-off value indicates that the patient belongs to the PAI-1 paracrine addicted (PPA) and co-activators predominant (CAP) subgroup.

[00114] In another example, the concentration of p-STAT3 is determined first, followed by the determination of the concentration of PAI-1.

[00115] In one example, if the patient is shown to belong to the PAI-1 paracrine addicted (PPA) and co-activators predominant (CAP) subgroup based on the concentration measurements for p-STAT3, then a PAI-1 concentration of less than 20 ng/ml indicates that the subject belongs to the co-activators predominant (CAP) subgroup. If the patient is shown to belong to the PAI-1 paracrine addicted (PPA) and co-activators predominant (CAP) subgroup based on the concentration measurements for p-STAT3, then a PAI-1 concentration of at least 20 ng/ml or more indicates that the subject belongs to the PAI-1 paracrine addicted (PPA) subgroup. If the patient is shown to belong to the alternative pathways activation (APA) subgroup based on the concentration measurements for p-STAT3, then a PAI-1 concentration of less than 20 ng/ml indicates that the subject belongs to the alternative pathways activation (APA) subgroup. If the patient is shown to belong to the alternative pathways activation (APA) subgroup based on the concentration measurements for p-STAT3, then a PAI-1 concentration of at least 20 ng/ml or more indicates that the subject belongs to an undetermined subgroup.

[00116] In another example, one subgroup or subset of patients is defined as having an IL6 concentration of less than 997 pg/ml, a CCL2 concentration of less than 450 pg/ml, an IL10 concentration of less than 15 pg/ml, and an MMP9 concentration of less than 3 ng/ml. This group refers to the alternative pathways activation (APA) group, as defined herein.

[00117] In another example, one subgroup or subset of patients is defined as having an IL6 concentration equal to or more than 997 pg/ml (≥ 997 pg/ml), a CCL2 concentration equal to or more than 450 pg/ml (≥ 450 pg/ml), an IL10 concentration equal to or more than 15 pg/ml (≥ 15 pg/ml), and a MMP9 concentration equal to or more than 3 ng/ml (≥ 3 ng/ml). This group collectively refers to the PAI-1 paracrine addicted (PPA) and co-activators predominant (CAP) groups, as defined herein.

[00118] In one example, the methods disclosed herein can be performed in a treatment setting, which is, but is not limited to, neoadjuvant setting, adjuvant setting, palliative setting and prophylactic setting. In another example, the methods disclosed herein can be performed on the same subject in one or more settings.

[00119] As used herein, the term “setting” refers to the timing when the biomarkers are assessed and timing of treatment. For example, the term “neoadjuvant setting” means that ascites fluid has been extracted before the patient has undergone surgery, and that the ascites fluid is extracted via a percutaneous drainage procedure. Upon determining susceptibility of patient, appropriate treatment (depending on which group patients falls into, i.e. PAI-1 paracrine addicted (PPA), co-activators predominant (CAP), alternative pathways activation (APA)) would be provided. In the adjuvant setting, a drain has been inserted prior to surgery to extract ascitic fluid from the intraabdominal cavity. Upon determining the susceptibility of patient, appropriate treatment (depending on which group patient falls into, i.e. PAI-1 paracrine addicted (PPA), co-activators predominant (CAP), alternative pathways activation (APA)) would be provided during hyperthermic intraperitoneal chemotherapy (HIPEC). In the palliative setting, patients do not undergo any surgery and the ascites fluid is extracted for analysis. The patient is treated accordingly (depending on which group patient falls into, i.e. PAI-1 paracrine addicted (PPA), co-activators predominant (CAP), alternative pathways activation (APA)) with a palliative intent.

[00120] In another example, the determination or measurement of the level of STAT3 activation (or phosphorylation) can be performed using surrogate markers. In another example, the level of STAT3 phosphorylation is determined by measuring the concentration of one or more surrogate markers. Alternatively, the level of STAT3 phosphorylation can also be determined by directly measuring the concentration of phosphorylated STAT3 directly. It will be appreciated by a person skilled in the art that STAT3 phosphorylation cannot be determined directly in, for example, a liquid sample, as phosphorylation takes place within cells. Therefore, when measuring levels of STAT3 phosphorylation in liquid form, cell lysis of cells that have been exposed to cell-free ascites *in vitro*, *in vivo*, and in the clinical setting must have taken place. The levels of STAT3 phosphorylation in the resulting sample determined using, for example, enzyme-linked immunosorbent assay (ELISA), or any other method capable of determining said levels. Cell lysis can be performed using methods known to a person skilled in the art, who would be able to ascertain which method is best suited for the sample in hand.

[00121] In another example, the level of STAT3 phosphorylation is determined by measuring the concentration of one or more surrogate markers present in the cell-free ascites. In yet another example, the level of STAT3 phosphorylation (p-STAT3) can also be determined by measuring the p-STAT3 level

of cellular components present in ascites or tumour biopsy. In another example, the level of STAT3 phosphorylation can be determined by directly measuring the concentration of phosphorylated STAT3 directly and by measuring the concentration of one or more surrogate markers.

[00122] As used herein, the term “surrogate marker” refers to one or more (bio-)markers which can be used in substitute or a proxy of the intended target. The term “biomarker” can and is used interchangeable with the term “surrogate marker” in the present disclosure. For example, as disclosed herein, the level of STAT3 activation can be measured by determining the level of IL6. In one example, the relationship between a surrogate marker and the intended target can be proportional, meaning that an increase or decrease in the level or concentration of the surrogate marker is understood to have the same increase or decrease in the level or concentration of the intended target. This relationship can also be linear. However, it is also possible to have a surrogate marker with an anti-proportional relationship to the intended target.

[00123] In one example, the surrogate marker used for the determination of the level STAT3 activation (or phosphorylation) can be, but is not limited to, one or more of the markers as listed in Table 1.

[00124] In one example, the level of STAT3 phosphorylation is determined by measuring the concentration of at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more surrogate markers. In one example, the level of STAT3 phosphorylation is determined by measuring the concentration of at least 3 surrogate markers. In one example, these 3 markers can be, but are not limited, to, IL6, IL10 and CCL2. In one example, the level of STAT3 phosphorylation is determined by measuring the concentration of at least 4 surrogate markers. In one example, these 4 markers can be, but are not limited to, IL6, IL10, CCL2 and MMP9. In one example, the level of STAT3 phosphorylation is determined by measuring the concentration of at least 5 surrogate markers. In one example, these 5 markers can be, but are not limited to, TGFB1, POSTN, VSIG4, CD44 and CXCL10. In one example, the level of STAT3 phosphorylation is determined by measuring the concentration of at least 6 surrogate markers. In one example, the 6 markers can be, but are not limited to, IL6, TGFB1, POSTN, VSIG4, CD44 and CXCL10. In one example, the method disclosed herein is performed using one surrogate marker. In another example, the method disclosed herein is performed using 2 surrogate markers. In another example, the method disclosed herein is performed using 3 surrogate markers. In another example, the method disclosed herein is performed using 4 surrogate markers. In another example, the method disclosed herein is performed using 5 surrogate markers. In another example, the method disclosed herein is performed using 6 surrogate markers. By way of an example, a biomarker panel, for example, will measure the concentration of a defined number biomarkers. In one example, the panel comprises or consists of IL6, IL10, CCL2 and MMP9. In the method disclosed herein, in order for a patient sample to be defined as PAI-1 paracrine addicted (PPA)/co-activators predominant (CAP), the values of the surrogate markers detected in the sample must pass the respective cut-off values defined for each of the surrogate markers. For example, in a panel of 4 markers, at least 3 for the 4 surrogate markers must pass their respective cut-off values. In a panel of 2 markers, for example, depending on the markers chosen, at least one biomarker or both biomarkers must

pass their respective cut-off values. In a panel of 3 markers, for example, depending on the markers chosen, at least two biomarkers or all biomarkers must pass their respective cut-off values. In a panel of 4 markers, for example, depending on the markers chosen, at least 3 biomarkers or all biomarkers must pass their respective cut-off values. Exemplary panels can be found in Fig. 20D.

5 [00125] Table 1: Non-exhaustive list of putative surrogate markers for the determination of the level of STAT3 activation

A1BG	CSF2	IDE	CEACA M6	SPARC	PDIA6	IGKV2 D-30	LACRT
A2M	CSF3	CFI	NBL1	SPINK1	NAMPT	IGKV2 D-29	FRMD7
SERPINA3	CSN2	IFNA1	NDP	SPINK2	CELA3A	IGKV2 D-28	UCN2
ABCA1	CSN3	IFNA2	NID1	SPINT1	EBI3	IGKV2 D-26	ERVH48-1
ABCA3	VCAN	IFNA4	NODAL	SPN	USPL1	IGKV1 D-43	IL33
AOC1	CST1	IFNA5	NPY	SPOCK1	GDF11	IGKV1 D-33	SCGB3A1
ACHE	CST2	IFNA6	NOV	SPP1	MSLN	IGKV1 D-17	BPIFB1
ACPP	CST3	IFNA7	NPPA	SPTBN2	LRRC17	IGKV1 D-13	PKHD1L1
ACTA1	CST4	IFNA8	NPPB	SST	RAMP1	IGKV1 D-12	CPA5
ACTA2	CSTA	IFNA1 0	NPPC	ST14	FSTL3	IGKV1 D-8	MUC16
ACTB	CSTB	IFNA1 3	NUCB1	STC1	LILRB2	IGKV6- 21	TGS1
ACTC1	CTBS	IFNA1 4	NUCB2	STX4	RTN3	IGKV4- 1	CMTM7
ACTG1	CTF1	IFNA1 6	OAS3	XCL2	CRISP3	IGKV3- 20	IL17F
ACTG2	CTGF	IFNA1 7	OMD	TAC1	AKR1A1	IGKV3- 15	SAAL1
ACTN4	CTRB1	IFNA2 1	OGN	TAC3	CCL26	IGKV2- 40	CMTM1
ACTN1	CTRL	IFNAR 2	OMG	SERPINA7	SEMA3A	IGKV2- 30	UCN3
ADA	CTSB	IFNB1	TNFRS F11B	ELOA	CPQ	IGKV2- 28	CGB1
ADAM10	CTSD	IFNG	ORM1	TCN1	WFDC2	IGKV2- 24	CGB2
ADCYAP1	CTSG	IFNW1	ORM2	TCN2	SPON2	IGKV1- 39	ELFN2
ADM	CTSH	IGF1	OSM	TDGF1	SPON1	IGKV1- 27	C1QTNF1
AEBP1	CTSK	IGF2	OXT	TF	OLFM1	IGKV1- 17	C1QTNF2
CRISP1	CTSL	IGF2R	PCSK6	TFF1	LRRN2	IGKV1- 12	C1QTNF3
AFM	CTSV	IGFAL	PRDX1	TFF2	FAM3C	IGKV1-	C1QTNF4

		S				9	
AFP	CTSO	IGFBP 1	SERPIN E1	TFF3	MERTK	IGKV1- 6	C1QTNF5
AGA	CTSS	IGFBP 2	SERPIN B2	TFPI	PIBF1	MRPL1 8	C1QTNF6
AGRP	CTSW	IGFBP 3	PAM	TFRC	CAP1	VPREB 3	VASN
AGT	CTSZ	IGFBP 4	REG3A	TGFA	CRTAP	HILPD A	CTHRC1
AHSG	DAG1	IGFBP 5	PAPPA	TGFB1	ENOX2	NENF	CMTM5
ALAD	DBH	IGFBP 6	SERPIN A5	TGFB2	SEMA4D	IL19	ACSM1
ALB	DCN	IGFBP 7	PCOLC E	TGFB3	SEMA3C	LMCD1	IL22RA2
ALDH3A1	ACE	CYR61	PCSK1	LEFTY2	FBLN5	KLK14	MRGPRD
ALDOA	DDB1	IGHA1	PCSK5	TGFB1	CIB2	KLK12	APOA5
AKR1B1	DEFA1	IGHA2	PCSK2	TGFBR3	PRDX4	PLA2G 3	LRG1
ALOX5	DEFA3	IGHD	PDE4C	THBD	AGR2	IL20	OLFM3
ALPL	DEFA4	IGHE	PDGFA	THBS1	OLFM4	IL22	CPXM2
AMBP	DEFA5	IGHG1	PDGFB	THBS2	CXCL13	DHH	LRRK2
AMH	DEFA6	IGHG2	ENPP1	THBS4	NPC2	SOST	LRIG3
AMY1A	DEFB1	IGHG3	ENPP2	THPO	GNLY	CELA2 B	GPHB5
AMY1B	DEFB4A	IGHG4	PECAM 1	TIMP1	POSTN	GAL	NRN1L
AMY1C	DLG3	IGHM	SERPIN F1	TIMP2	LEFTY1	UTP11	CMTM3
AMY2A	DMBT1	JCHAIN	PF4	TIMP3	SCGB1D2	ANGPT L4	ZG16B
ANG	DPEP1	IGKC	PF4V1	TIMP4	SCGB1D1	IRAK4	SEZ6
ANGPT1	DPT	IGLC1	CFP	TLE2	TNFSF13 B	SERPI NA10	TMIGD2
ANGPT2	DPYSL3	IGLC2	PFN1	TMSB4X	STAG3	EGFL7	LRRC38
ANPEP	EPYC	IGLC3	PGC	CLEC3B	MASP2	CKLF	PODN
ANXA1	HBEGF	IGLC6	PGF	TNF	MTHFD2	CPA4	NAXE
ANXA2	ECM2	IGLL1	PGK1	TNFAIP2	CCL27	C1RL	CPO
ANXA5	ECM1	IHH	SERPIN A1	TNFAIP6	FGL2	GOLM 1	LYZL4
ANXA13	EDN1	IK	SERPIN A4	TNFRSF1A	EDDM3A	BPIFA1	OTOP1
APOF	EDN2	IL1A	SERPIN B5	TNXB	CFHR3	PLA1A	CD109
APCS	EDN3	IL1B	SERPIN B6	TPI1	SMR3B	ANGPT 4	PXDNL
APOA1	EEF1A1	IL1RAP	SERPIN E2	TPO	UTS2	WNT16	CBLN4
APOA2	EGF	IL1RN	SERPIN B8	TPSAB1	SMPDL3A	PRKAG 2	SOGA1
APOA4	EGFR	IL2	SERPIN B9	TPT1	POP1	PCYOX 1	RBBP8NL
APOB	CELA1	IL3	SERPIN B10	CRISP2	LMAN2	IL23A	OVOS2

APOC1	ELANE	IL4	SERPIN I1	TSHB	IL24	ESF1	A2ML1
APOC2	SERPINB1	IL4R	SERPIN B13	TST	KLK11	LSR	SERPINA1 2
APOC3	ENG	IL5	SERPIN I2	TTR	KERA	OAZ3	LRFN5
APOC4	ENO1	IL5RA	PIGR	TNFSF4	ADAMTS 13	GPRC5 B	HAPLN3
APOD	ENO2	IL6	PIP	UBA52	ADAMTS 5	GHRL	TTBK2
APOE	ENO3	IL7	PLA2G 1B	UBB	PRSS23	ERAP1	CMTM4
APOH	STOM	CXCL 8	PLA2G 2A	UBC	EMILIN1	ADA2	CMTM2
APP	STX2	IL9	PLAT	SCGB1A1	FSTL1	IL17D	IL34
KLK3	EPO	IL9R	PLAU	COL14A1	ADAMTS 7	PRKAG 3	TMC8
FASLG	ERBB3	IL10	PLG	VCAM1	KLK8	FAM3B	CCBE1
AREG	EREG	IL11	SERPIN F2	VEGFA	PRR4	TLR9	CBLN2
ARG1	F2	IL12A	PLTP	VEGFB	SCRG1	H2BFS	HFE2
ASAH1	F3	IL12B	PNLIPR P2	VEGFC	NID2	CYTL1	PM20D1
ASIP	F5	IL13	PODXL	VGF	VASH1	WNT4	ERFE
SERPINC1	F7	IL13R A2	POMC	EZR	CLSTN1	SIAE	GDF7
ATP4A	F8	IL15	PON1	VLDLR	CEP164	ADAM TSL4	CPNE9
AVP	F9	IL15R A	PON3	VPREB1	ARSG	LRRN3	CCDC80
AXL	F11	IL16	PPBP	VTN	DKK1	EPDR1	CMTM8
AZGP1	F12	TNFR SF9	PPIA	WNT1	CNOT1	FAM20 A	SPINK13
AZU1	F13A1	IL17A	PPP1R1 A	WNT2	SIPA1L3	BIVM	PRSS3P2
B2M	FABP3	IL18	PPT1	WNT3	SSPO	SEMA4 C	LINGO2
BCHE	FAP	INHA	PPY	WNT5A	FRMD4B	CMTM 6	EEF1A1P5
CFB	FBLN1	INHB A	PRELP	WNT6	PMPCA	LIME1	CFAP58
BGLAP	FBLN2	INHB B	SRGN	WNT7A	SULF1	ODAM	LGI4
BGN	FBN1	INHB C	PRH1	WNT7B	DNAJC9	INTS11	TMPRSS6
BMP1	EFEMP1	CXCL 10	PRH2	WNT8A	KIAA0556	ENOX1	FREM3
BMP2	FKTN	INS	PROC	WNT8B	MTCL1	WDR60	BMPER
BMP3	FCN2	INSL4	PROS1	WNT10B	MCF2L	LGI2	QSOX2
BMP4	GPC4	ISLR	PRSS1	WNT11	DMXL2	THNSL 2	SUPT20H L2
BMP5	FGA	ITGA2 B	PRSS2	WNT2B	ZCCHC11	RNLS	ADAMTS 15
BMP6	FGB	ITGA	PRSS3	WNT9A	MAN2B2	NDUF	KRT78

		M				AF7	
BMP7	FGF1	ITIH1	MASP1	WNT9B	ADNP	ZNF446	ZFC3H1
BMP8B	FGF2	ITIH2	RELN	XDH	CELA3B	KDM4D	DAND5
BMPR2	FGF5	ITIH4	KLK7	YWHAZ	ANGPTL2	SLF2	GKN2
BPI	FGF6	ANOS1	PRSS8	ZNF177	CLCF1	IL26	LIPH
BTC	FGF9	KARS	KLK6	ZP3	DNPEP	SELENOS	C9orf72
BTD	FGF10	KCNK3	HTRA1	PXDN	PYY2	DEFB103B	C3orf58
BTN1A1	FGF12	KISS1	PRTN3	SCG2	LY96	APOBR	LRRC55
C1QBP	GPC5	KIT	PSAP	MANF	PLA2G15	APOM	UCMA
SERPING1	FGG	KLKB1	PYY	PLA2G7	FLRT3	SULF2	GPC2
C1QA	FGL1	KNG1	PSMC5	ADAM12	FLRT2	MYDGF	SCUBE3
C1QB	VEGFD	KRT1	PTGDS	FGF23	FLRT1	PDGFC	SEMA3D
C1QC	FLT1	KRT2	PTGIS	MFAP5	FJX1	CPXM1	IL27
C1R	FLT3LG	KRT9	PTH	MIA	ATXN10	GKN1	ZBTB38
C1S	FMOD	KRT10	PTHLH	GDF5	KLK5	IL36G	UBN2
C2	FN1	KRT31	PTN	EPX	PRDX5	CCL28	BPIFC
C3	FRZB	KRT33A	QSOX1	COLQ	CHRD2	MUC13	EPGN
C4A	FSHB	KRT33B	PTPRG	HIST1H2BG	ABI3BP	RETN	PCSK9
C4B	FUCA2	KRT34	PTPRR	HIST1H2BF	PAMR1	IFNK	NPNT
C4BPA	GAST	KRT35	PTX3	HIST1H2BE	SOSTDC1	GRIPAP1	SERPINA11
C4BPB	KDSR	KRT81	PVR	HIST1H2BI	EGFL6	FAM20C	KLHL34
C5	GAS6	KRT83	PZP	HIST1H2BC	TSKU	ADAMTS9	MDS2
C8A	GBA	KRT85	RARRES2	HIST2H2BE	MOXD1	TWSG1	PRSS33
C8B	GC	KRT86	RBP3	PLA2G6	KLK13	CPA6	MDGA1
C8G	GCG	LALBA	RBP4	SPARCL1	FGF21	EPPIN	IFNL2
C9	BLOC1S1	LAMA2	RDX	LTBP4	GNL3	SLURP1	IFNL3
CA2	KAT2A	LAMA5	REN	ATRNL	TIMM8B	RALGAPA2	IFNL1
CA6	GCNT1	LAMB1	RNASE3	CILP	IL36RN	DSCAML1	PRTG
DDR1	GDF1	LAMB2	RNPEP	PPFIBP2	GREM1	LRFN2	BRICD5
CALCA	GDF2	LAMC1	RPL39	CPZ	FETUB	MTUS1	METRNL
CALR	MSTN	LAMC2	RPS27A	APOL1	FGF22	LRFN1	LAMA1
CAMP	GDF9	LAMP2	RS1	FCN3	LYPD3	LRRN1	HMSD
CAT	GDF10	LBP	S100A4	YARS	DKKL1	COL20	SSC5D

						A1	
SERPINA6	GGT1	LCAT	S100A8	TNFSF11	DKK3	ZSWIM5	CXCL17
CBR3	B4GALT1	LCN1	S100A9	STC2	DKK2	LRRC4C	VSTM1
CCK	GH1	LCN2	S100A11	NPFF	CPAMD8	NCOA5	FAM19A3
KRIT1	GHR	LCP1	S100A13	CDK13	CHIA	SCUBE2	C3orf33
CD5L	GHRH	LDLR	S100B	RNASET2	IL36B	HAMP	LCN1P1
CD9	GIF	LECT2	SAA1	CHRD	IL37	WFDC1	SERPINA9
CD14	GIP	LEP	SAA2	SERPINB7	IL36A	CXCL16	GPIHBP1
MS4A1	GPC3	LGALS1	SAA4	CTSF	IL17C	OPRPN	IFNE
TNFSF8	GLB1	LGALS3	SERPINB3	TNFSF14	IL17B	IL21	C1QL4
CD36	GLE1	LGALS3BP	SERPINB4	TNFSF13	TINAG	ACE2	KLHL17
ENTPD6	GNB2	LGALS4	CLEC11A	TNFSF12	SRPX2	CELA2A	VWA2
CD40	GNL1	LGALS8	SCT	TNFSF10	SMPDL3B	GFRA4	OTOG
CD40LG	GNRH1	LGALS9	CCL1	TNFSF9	BMP10	TINAGL1	GLDN
CD59	SFN	LHB	CCL2	ADAM15	RBMX	IRF2BPL	OSTN
CD63	GP5	LIF	CCL3	ADAM9	ANGPTL3	SIL1	ACTBL2
CD70	GPC1	LIFR	CCL3L1	TNFRSF6B	PCSK1N	GREM2	CLEC18A
ADGRE5	GPI	LIPC	CCL4	DLK1	IGKV1-5	IL25	C6orf58
CDH13	GPLD1	LOX	CCL5	CREG1	IGHV6-1	VWA1	BMP8A
CEL	GPT	LOXL1	CCL7	FGF18	IGHV5-51	ZNF649	IGFL1
CETP	GPX3	LOXL2	CCL8	FGF17	IGHV4-61	CHID1	MROH7
CFL1	GPX5	LPA	CCL11	FGF16	IGHV4-28	LRFN4	VWC2
CFL2	GRN	LPL	CCL13	NRP1	IGHV4-4	METRN	KCP
CTSC	CXCL1	LPO	CCL14	GGH	IGHV3-74	APOO	IL31
CEACAM8	CXCL2	LTA	CCL15	WISP3	IGHV3-73	FKRP	SOGA3
CHEK1	CXCL3	LTB	CCL16	WISP2	IGHV3-72	CRELD2	C10orf99
CHGA	GRP	LTBP1	CCL17	WISP1	IGHV3-66	GLB1L	CCL4L1
CHI3L1	PDIA3	LTBP2	CCL18	PROM1	IGHV3-64	LRFN3	C3P1
CHI3L2	GSN	LTF	CCL19	PROZ	IGHV3-49	TCTN1	CEACAM16
CHIT1	GSTP1	LUM	CCL20	APLN	IGHV3-43	FAM184A	C1QTNF12
CKB	GUSB	LYZ	CCL21	ENDOU	IGHV3-30	GSDMD	SERPINA2
CLCA1	HABP2	TACSTD2	CCL22	HIST1H2BJ	IGHV3-23	MMRN2	TRIM75P
CLU	HBA1	MAN2A1	CCL23	SELENBP1	IGHV3-21	PLBD1	GDF6

CLIC1	HBA2	MAN2 B1	CCL24	TNFSF18	IGHV3-15	PDZD7	PATE2
CNP	HBB	MATN 2	CCL25	ARTN	IGHV3-13	SVEP1	PATE4
CNTF	HBD	MBL2	CXCL6	ANGPTL1	IGHV3-7	PLEKH H3	LOC40057 6
CNTFR	HBE1	MCA M	CXCL1 1	MTMR4	IGHV2-26	ADAM TS20	PRSS57
COL1A1	HBG2	MDH1	CXCL5	INA	IGHV1-58	SCUBE 1	VWC2L
COL1A2	SERPIND1	MECP 2	XCL1	BMP15	IGHV1-45	PDGFD	OVOS
COL2A1	HDGF	MEP1 A	CX3CL 1	LGI1	IGHV1-24	WNT10 A	SPINK14
COL3A1	HDLBP	MEP1 B	SDCBP	IL32	IGHV1-18	ULBP2	CCL3L3
COL4A1	HEXB	MFAP 4	CXCL1 2	NOG	IGHV1-3	BPIFB2	DEFB103 A
COL4A2	CFH	MFGE 8	SDF2	CRLF1	TRDC	SPX	LOC43995 1
COL5A1	HFE	MELT F	SECTM 1	AIMP1	TRBC2	COL18 A1	CTRB2
COL5A2	CFHR1	MFNG	SELE	MMP20	TRBC1	APOL4	CDNF
COL6A1	HGF	SCGB 2A1	SELP	CER1	IGLV10- 54	WNT5 B	IGFL4
COL6A2	HGFAC	KITLG	SEMA3 F	SLIT2	IGLV9-49	AMN	POTEE
COL6A3	HMGB1	MIF	SEMG1	ITGBL1	IGLV8-61	JAM3	SPINK9
COL7A1	HMGB2	CXCL 9	SEMG2	KL	IGLV7-46	INHBE	CBLN3
COL8A1	HMOX1	MMP2	SELEN OP	ADIPOQ	IGLV5-52	FGFBP 2	PYY3
COL11A1	HP	MMP3	SFRP1	LIPG	IGLV5-45	EIF2A	LINGO3
COL12A1	HPGD	MMP7	SFRP2	ITM2B	IGLV5-39	TMPRS S13	SPINK8
COL15A1	HPR	MMP8	SFRP4	LY86	IGLV5-37	EMILI N2	LYPD8
COMP	HPX	MMP9	SFRP5	NAPSA	IGLV4-69	LOXL4	SERPINE3
COPA	HRG	MMP1 0	SFTPB	CABP1	IGLV4-60	C2orf40	POTEI
CORT	HSPA1A	MMP1 2	SFTPC	ADAMTS4	IGLV4-3	RAB11 FIP4	LGALS7B
CP	HSPA1B	MMP1 3	SFTPD	ADAMTS3	IGLV3-25	COL25 A1	SFTPA1
CPA1	HSPA1L	MMP1 4	SH3BG RL	ADAMTS2	IGLV3-22	IL1F10	POTEJ
CPA2	HSPA2	MMP1 9	SHH	GDF15	IGLV3-21	SPINK7	MSMP
CPA3	HSPA6	MOV1 0	SLC2A1	CXCL14	IGLV3-16	RETNL B	MUC5B
CPB1	HSPA7	MPO	SLC4A1	GDF3	IGLV3-12	LOXL3	DEFA1B
CPB2	HSPA8	MSMB	SLIT1	PRDX6	IGLV3-10	AIFM2	POTEF
CPD	HSPB1	MSN	SLIT3	PDIA4	IGLV2-18	LINGO 1	SFTPA2

CPE	HSPD1	MSR1	SLPI	CARTPT	IGLV1-47	HIST1 H2BK	PSAPL1
CPM	HSPG2	MST1	SMARC A4	CLCA3P	IGLV1-36	KRT87 P	LRRC70
CPN1	TNC	MT3	SMPD1	SEMA3E	IGLC7	SSH2	GKN3P
CPN2	HYAL1	NUDT 1	SNCA	FAM20B	IGKV6D- 21	TSLP	MOXD2P
CRH	IAPP	MUC1	SOD1	TNFSF15	IGKV3D- 20	UMOD L1	IGLL5
CRHBP	IBSP	MUC4	SOD3	FGFBP1	IGKV3D- 15	SERPI NB12	APELA
CRP	ICAM1	MUC5 AC	SORD	IL18BP	IGKV3D- 11	SERPI NB11	MICA
CSF1	ICAM4	MYOC	SORL1	GPC6	IGKV3D-7	WNT3 A	ZNF559- ZNF177
CEACAM5	TUBB2A	IGLV3 -19	KRT5	HUS1B	KRT80	IFNL4	LOC10106 0157
KRT6B	IGLV1-40	LDHA	TUBA4 A	KRT6C	YWHAG	PROCR	TRAP1
KRT76	HIST1H1A	CFHR 2	REG1B	IGHV1OR2 1-1	PPIAL4A	KRT24	CSF1R
KRT12	PGAM4	VSIG4	HSP90A B3P	PC	POTEKP	LOC64 2131	C6
F13B	REG1A	TUBA 1A	TUBB4 A	PLXDC2	HSPH1	KRT17	IGLV3-27
LDHB	SLC4A4	KRT7	KRT16	KRT6A	TUBA3E	TUBB6	KRT36
SUCLG1	TUBB	VIM	GTPBP 8	TNXA	ITLN2	CEACA M1	CENPQ
KRT71	TKT	ANTX R1	SRSF9	GFAP	LYVE1	LAMA 3	PPIAL4G
GAPDHS	NLRC5	PGAM 1	ATP5A 1	EPHB4	ITLN1	KRT84	LEKR1
KRT74	ICAM2	TUBA 1C	HIST1H 1B	BCAM	BRSK1	USO1	HNRNPA1
MAN1A1	KRT38	ABCB 9	KRT25	FGFR1	FCGBP	C2ORF 72	HBG1
HIST1H1D	CDH5	KRT37	DSG2	PLS1	ITIH3	TAGLN	HNRNPA1 L2
HSP90AB1	VCL	YWH AB	GDI1	KRT79	KRT27	EXOC3 L4	IGLV7-43
G6PD	SLC4A5	OSBP L11	EPCAM	ESD	KRT13	CD163	KRT3
KRT19	CRTAC1	PGLY RP2	CD44	ETFB	IGLV1-44	USH1C	IFT74
ARHGDIA	VWF	TUBB 4B	GAPDH	KIAA0232	DOCK10	GPX6	HBZ
PRG4	IGHV4-59	KRT15	ANXA2 P2	TUBB2B	NUP214	CA1	KRT73
HSPA9	TGM4	TUBA 8	GDI2	IDH1	YWHAH	CST7	MXRA5
TUBA1B	IGLV2-11	LMNA	S100A6	PRDX2	CRIP1	RXRβ	PI16
PLS3	STX17	YWH AQ	TUBB8	CD248	MCM5	EEF1A 2	SYNE2
C7	MYO19	KRT4	GSTO1	HIST1H1T	KRT77	KRT8	CFHR5

CPS1	MMP11	KRT75	DCD	HIST1H1C	ADH5	KRT14	ALDOC
KPNA2	KRT82	KRT28	KRT32	SHBG	FCGR3A	CNDP1	RARRES1
TUBA3C	YWHAE	IGK@	HNRNP A2B1	FYCO1	KRT18	FNDC3 A	THBS3
PGK2	NEFH	UBA1	HIST1H 1E	METTL18	FCGR3B	PGAM2	ALDOB
IGLV6-57	KRT72	CAD	TLN1	HSPA5	CFD	NEB	PLGLA
CD24	CD26	CD147	FGF7	FGF19	TNFRSF8	RLN2	BDNF
RAGE	TIM3	HSP90 AA1	GP1BA				

[00126] Figure 19a provides an overview of the p-STAT3 surrogate markers selection workflow. Briefly, and as previously mentioned, STAT3-related genes were identified from Kyoto Encyclopedia of Genes and Genomes (KEGG) database by compiling all genes that are involved in known STAT3 pathways.

5 Secreted STAT3-related proteins were selected based on extracellular genes listed in NCBI's Biosystems database and proteins identified in mass spectrometry analysis of cell-free ascites. Transcriptomics comparison was performed using two databases to prioritize putative STAT3 surrogate markers. Database 1 was used to determine genes that are positively correlated with STAT3 in The Cancer Genome Database (TCGA) colorectal cancer (COADREAD) data set. Genes were ranked from the most positively
10 correlated to least correlated with STAT3. Database 2 was derived from microarray analysis of PAI-1 paracrine addicted (PPA) cell-free ascites-treated cells exposed to TM5441 to determine genes that are downregulated and upregulated in PAI-1 paracrine addicted (PPA) cell-free ascites-treated cells in response to TM5441 (PAI-1 inhibition). Upregulated genes were also of interest as these were thought to represent genes that are involved in rescue mechanisms in response to PAI-1 inhibition. Similarly, genes
15 were ranked from most downregulated to most upregulated. Systematic paired correlation analysis of candidate genes was subsequently performed by focusing on top 1% and 25% of genes positively correlated with STAT3 in database 1, and top 1% and 25% of most downregulated and upregulated genes in database 2. The paired analysis for each group were prioritised as shown in Fig. 19b, and representative genes were chosen from each group based on literature review to reduce the list of potential targets to 35
20 genes. Ten targets were selected based on rank prioritisation, potential good correlation with p-STAT3 from Luminex assay data, and the importance of the candidate genes in cancer pathogenesis from literature review for further evaluation with enzyme-linked immunosorbent assay (ELISA). The concentrations of each surrogate marker in cell-free ascites were correlated with p-STAT3 levels in cell-free ascites-treated cells using Spearman correlation analysis.

25 [00127] Thus, the surrogate markers disclosed herein can be selected based on their correlation to STAT3. The surrogate markers disclosed herein can also be selected based on their up- or down-regulation compared to the same markers in cell-free ascites-treated samples or in negative controls. Once ranked, for example by prevalence, priority, or any other criteria, these markers can be, but are not limited to, the top 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%,
30 20%, 21%, 22%, 23%, 24%, or 25% of all the markers listed based on the above criteria. For example,

markers can be selected for being in the top 1% of markers which positively correlate with STAT3. In another example, markers can be selected for being in the top 1% of markers which negatively correlate with STAT3 phosphorylation. As used herein, a positive correlation refers to a proportional relationship between a surrogate marker and its target. A negative correlation therefore refers to an anti-proportional (or inverse) relationship between a surrogate marker and its target. For example, a positive correlation means that an increase in target concentration results in an increase in surrogate marker concentration. A positive correlation can also mean that a decrease in target concentration results in a decrease in surrogate marker concentration. Conversely, a negative correlation means that an increase in target concentration results in a decrease in surrogate marker concentration. Markers can also be selected for being in the top 1% or 25% of markers which are up-regulated or down-regulated compared to a control or any other benchmark.

[00128] In another example, surrogate markers are selected based on their up- and/or down- regulation. Such an up- or down-regulation can be determined based on the level of such markers in, for example, samples which have been treated with a PAI-1 inhibitor.

[00129] A person skilled in the art will readily appreciate that markers can be chosen for reasons and criteria other than listed herein, for example, markers that do not show any apparent correlation to STAT3 but were shown to have a significant effect on for example, a multivariate analysis. It is also appreciated that multiple criteria can be applied to the initial marker pool in order to narrow down and obtain a final list of, for example, surrogate markers.

[00130] Out of the 10 targets selected, five exemplary candidate surrogate markers (IL6, IL10, CCL2, MMP9, and ANGPT1) were validated on 70 patient samples and successfully identified an exemplary composite biomarker panel. As an example, such a composite biomarker panel can consist of four targets (IL6, CCL2, IL10, and MMP9) as surrogate biomarkers of STAT3. This exemplary panel has an overall accuracy of 92.86%.

[00131] In addition, five candidate surrogate markers (TGFB1, POSTN, VSIG4, CD44, and CXCL10) were validated in 40 patient samples, which resulted in the results as shown in Fig 21. This exemplary composite biomarker panel resulted in an area under the curve (AUC) value of 0.83 ($P = 0.001$). Combining IL6 with this composite biomarker panel, an area under the curve (AUC) of 0.98 ($P < 0.0001$) was obtained.

[00132] Thus, in one example, the surrogate markers can be, but are not limited to, one or more of the following: LUM, ANGPT1, IL1B, POSTN, TNC, MMP9, MMP2, TIMP3, DCN, VSIG4, CXCL5, CD36, ANGPT2, SERPINB5, IL6, CCL2, LEP, VCAM1, CCL8, ITGAM, THBS1, FN1, COL5A1, MXRA5, C3, CXCL10, TGFB1, CD44, TIM3, TNFSF13B, CEACAM1, LAMB1, IL10, IL5, IL22. In yet another example, the surrogate markers can be, but are not limited to, one or more of the following: IL6, IL10, CCL2, MMP9, ANGPT1, TGFB1, POSTN, VSIG4, CD44, and CXCL10. In a further example, the surrogate markers can be, but are not limited to, one, or more, or all, of the following: IL6, IL10, CCL2, MMP9 and ANGPT1. In another example, the surrogate markers can be, but are not limited

to, one, or more, or all, of the following: IL6, IL10, CCL2, and MMP9. In another example, one of the surrogate markers is IL6. In yet another example, the combination, or group, or panel of surrogate markers used comprises IL6.

[00133] Thus, in one example, the surrogate markers are, but are not limited to IL6, IL10, CCL2, MMP9, ANGPT1, TGFB1, POSTN, VSIG4, CD44, and CXCL10. In another example, the surrogate markers are, but are not limited to, IL6, IL10, CCL2, MMP9 and ANGPT1. In yet another example, the surrogate markers are, but are not limited to, IL6, IL10, CCL2, and MMP9. In a further example, the surrogate markers comprise IL6, IL10, CCL2, MMP9 and ANGPT1. In another example, the surrogate markers comprise at least IL6, IL10, CCL2, and MMP9. In a further example, the surrogate markers are, but are not limited to, IL6, TGFB1, POSTN, VSIG4, CD44, and CXCL10. In yet another example, the surrogate markers are, but are not limited to, TGFB1, POSTN, VSIG4, CD44, and CXCL10.

[00134] The conclusion drawn from this group of patients is that the cell-free ascites (collected from a subgroup of patients) activates other signalling pathways thereby sustaining cancer cells. It was further noted that there appeared to be an absence of any cell-free ascites with high PAI-1 levels that is associated with low levels of STAT3 phosphorylation, when established cell line models are exposed to these cell-free ascites. This further shows that PAI-1 activates STAT3. In the unlikely event that this statement is not true, one would observe ascitic samples with high PAI-1 levels that do not activate STAT3 signalling when cells are exposed to these cell-free ascites. However, the method disclosed herein is supported and underlined by the fact that ascitic samples with high PAI-1 levels do activate STAT3 signalling when cells are exposed to these cell-free ascites. Taken together, a subset or subgroup of patients was identified, who had cell-free ascites with high PAI-1 levels that had been shown to drive STAT3 activation of cancer cells. Without being bound by theory, it was thought that if paracrine STAT3 activation of cancer cells is dependent on PAI-1 levels within these ascites, this would lead to a phenomenon of oncogenic addiction to a single upstream ligand. That is to say that patients whose cell-free ascites have high PAI-1 and activated STAT3 signalling when cells are considered to be highly susceptible to ligand inhibition of PAI-1 in ascites. This ligand inhibition of PAI-1 can be performed by, for example, intraperitoneal instillation of a PAI-1 inhibitor.


[00135] Next, Colo-205 (an established cell line model of colorectal peritoneal carcinomatosis) was systematically exposed to cell-free ascites collected from patients before subjecting these cells to treatment with TM5441 (PAI-1 inhibitor). As shown in Fig. 11, paracrine activation of Colo-205 led to a differential sensitivity to TM5441. Ascites collected from patients, of which it was thought that the STAT3 activation in cancer cells is dependent on PAI-1 levels within cell-free ascites, were most susceptible to inhibition with TM5441, confirming the finding that cancer cells exposed to these cell-free ascites were oncogenically addicted to PAI-1. In other words, the data shown here indicates that when cells are exposed to cell-free ascites belonging to, for example, the PAI-1 paracrine addicted (PPA) group, these cells are dependent on the ascites to activate STAT3 signalling within them. When PAI-1 is blocked (ligand inhibition) within the cell-free ascites, STAT3 signalling within the cells is inhibited and

the cells die. Cells exposed to co-activators predominant (CAP) group cell-free ascites, for example, are less reliant on PAI-1 for STAT3 activation, however a response is still possible. Concurrently, cells exposed to alternative pathways activation (APA) group cell-free ascites are not reliant on PAI-1 and do not activate STAT3, and are therefore not susceptible to PAI-1 inhibition.


5 [00136] Thus, in one example, there is disclosed a method of detecting or detecting susceptibility of a subject suffering from peritoneal carcinomatosis to treatment with a PAI-1 inhibitor, the method comprising determining the concentration of plasminogen activator inhibitor 1 (PAI-1) and determining the level of phosphorylation of “signal transducer and activator of transcript 3” (STAT3) in a sample obtained from a subject; wherein the subject is susceptible to treatment if the subject shows (a) an
10 increase in PAI-1 concentration and an increase in STAT3 phosphorylation, or (b) a decrease in PAI-1 concentration and an increase in STAT3 phosphorylation; wherein the increase and/or decrease is compared to the concentration of PAI-1 and the level of STAT3 phosphorylation measured in a sample obtained from a reference group.

[00137] Validation in *in vivo* mouse models using cell-free ascites that activates paracrine STAT3
15 signalling via PAI-1 demonstrated sensitivity to intra-peritoneal instillation of TM5441 (Fig. 14, Fig. 15 and Fig. 17). Some examples of PAI-1 inhibitors have been shown to bind to the s4A in PAI-1. The precursors of the PAI-1 inhibitor were identified by *in silico* virtual screening based on the 3D conformation of s4A position (Izuhara et al., *Arterioscler Thromb Vasc Biol.* 2008;28:672-677). The docking models of these precursors to s4A position of PAI-1 have been previously reported (Izuhara et al., *Journal of Cerebral Blood Flow & Metabolism* (2010) 30, 904–912). In one example, the PAI-1
20 inhibitor binds to the s4A position in PAI-1. In another example, the PAI-1 inhibitor is an anti-cancer treatment or anti-cancer drug. In another example, administration of the PAI-1 inhibitor, as disclosed herein, leads to inhibition of PAI-1 activity compared to patients suffering from the same disease.


[00138] In yet another example, the anti-cancer treatment or anti-cancer drug is, but is not limited to, a
25 small molecule, a chemotherapeutic agent, a peptide, an antibody, combinations thereof, and combination therapy. In another example, the anti-cancer drug is, but is not limited to, TM5441 (5-Chloro-2-[[2-[2-[[3-(3-furanyl)phenyl]amino]-2-oxoethoxy]acetyl]amino]benzoic acid sodium salt; CAS 1190221-43-2), TM5007 (N, N-bis [3,3'-carboxy-4,4'-(2,2'-thienyl)-2,2'- thienyl] hexanedicarboxamide; CAS 342595-05-5), TM5275 (5-Chloro-2-[[2-[2-[4-(diphenylmethyl)-1-piperazinyl]-2-oxoethoxy]acetyl]amino]-
30 benzoic acid sodium salt; CAS 1103926-82-4), Tiplaxtinin (2-(1-Benzyl-5-(4-(trifluoromethoxy)phenyl)-1H-indol-3-yl)oxoacetic acid; CAS 393105-53-8), ZK4044, and derivatives thereof. Example structures of the various anti-cancer drugs are shown below:




TM5441



TM5007



TM5275



[00139] In another example, the PAI-1 inhibitor is administered intraperitoneally.

- 5 [00140] In another example, there is disclosed a panel of markers for treating a patient suffering from peritoneal carcinomatosis with a “plasminogen activator inhibitor 1” (PAI-1) inhibitor, or for detecting or determining susceptibility of a subject suffering from peritoneal carcinomatosis to a treatment with a “plasminogen activator inhibitor 1” (PAI-1) inhibitor, wherein the panel of markers comprises PAI-1, and one or more surrogate markers of STAT3 phosphorylation or p-STAT3. In one example, the use of a panel
- 10 of markers in the method as referred to herein is disclosed, wherein the panel comprises PAI-1 and one or more surrogate markers of STAT3 phosphorylation, or PAI-1 and p-STAT3. In one example, the panel comprises PAI-1 and one or more or all of IL6, IL10, CCL2, and MMP9. In another example, the panel comprises PAI-1, and one or more or all of IL6, IL10, CCL2, MMP9 and ANGPT1. In yet another example, the panel comprises PAI-1, and one or more or all of TGFB1, POSTN, VSIG4, CD44, and
- 15 CXCL10. In a further example, the panel comprises PAI-1, and one or more or all of IL6, TGFB1, POSTN, VSIG4, CD44, and CXCL10.

[00141] In another example, there is disclosed use of a PAI-1 inhibitor in the manufacture of a medicament for treating peritoneal carcinomatosis, wherein the medicament is to be administered to a subject determined to belong to a patient group determined to be susceptible to PAI-1 inhibitor treatment.

- 20 In another example, the susceptibility of a subject is determined by measuring the concentration of PAI-1 and STAT3 phosphorylation (p-STAT3), as disclosed herein, and comparing the measured values to cut-off values as disclosed herein.

[00142] In the context of this invention the term “administering” and variations of that term including “administer” and “administration”, includes contacting, applying, delivering or providing a compound or composition of the invention to an organism, or any relevant surface by any appropriate means.

[00143] In As used herein, the term “treatment” refers to any and all uses which remedy a disease state or symptoms, prevent the establishment of disease, or otherwise prevent, hinder, retard, or reverse the progression of disease or other undesirable symptoms.

[00144] In the context of this specification, the terms “therapeutically effective amount” and “diagnostically effective amount”, include within their meaning a sufficient but non-toxic amount of a compound or composition of the invention to provide the desired therapeutic or diagnostic effect. The exact amount required will vary from subject to subject depending on factors such as the species being treated, the age and general condition of the subject, the severity of the condition being treated, the particular agent being administered, the mode of administration, and so forth. Thus, it is not possible to specify an exact “effective amount”. However, for any given case, an appropriate “effective amount” may be determined by one of ordinary skill in the art using only routine experimentation.

[00145] *In vitro* validation with Tiplaxtinin (PAI-1 inhibitor) highlights that inhibition of PAI-1 Michaelis complex is the potential mechanism of how cells are oncogenically addicted to PAI-1. Treatment with Napabucasin (STAT3 inhibitor) highlights that STAT3 inhibition alone is not useful as the Michaelis complex likely activates other signalling cascade in addition to STAT3 signalling. Treatment with dual PI3K/mTOR inhibitor or Mitomycin C (induces DNA damage) highlights the absence of utility of these drugs when cancer cells are exposed to paracrine activation driven by ascites (Fig. 12). Thus, in one example, the concentration of PAI-1 is determined by measuring the concentration of PAI-1 to urokinase-type plasminogen activator (uPA)/tissue-type plasminogen activator (tPA) complex. In another example, the concentration of PAI-1 is determined by measuring the concentration of PAI-1 in cell-free ascites. In another example, the concentration of PAI-1 is determined by measuring PAI-1 in its active and/or latent forms and/or complexes with, including but not limited to, urokinase-type plasminogen activator (uPA), tissue-type plasminogen activator (tPA), vitronectin and combinations thereof. In another example, the concentration of PAI-1 is determined by measuring the concentration of PAI-1 directly, or in one or more complexes. That is to say that PAI-1 need not be in a complex with, for example urokinase-type plasminogen activator (uPA)/tissue-type plasminogen activator (tPA) or other proteins to result in downstream effects. In yet another example, the level of STAT3 phosphorylation is determined by measuring the p-STAT3 level in established cell line models of colorectal peritoneal carcinomatosis. In one example, the cell line models of colorectal peritoneal carcinomatosis are treated with cell-free ascites. In a further example, the level of STAT3 phosphorylation is determined by measuring the p-STAT3 level in established cell line models of colorectal peritoneal carcinomatosis treated with cell-free ascites.

[00146] In another example, there is disclosed a method of treating a subject suffering from peritoneal carcinomatosis with a PAI-1 inhibitor, the method comprising determining the concentration of plasminogen activator inhibitor 1 (PAI-1) and determining the level of phosphorylation of “signal

transducer and activator of transcript 3" (STAT3) in a sample obtained from the subject; administering the PAI-1 inhibitor to the subject showing (a) an increase in PAI-1 concentration and an increase in STAT3 phosphorylation, or (b) a decrease in PAI-1 concentration and an increase in STAT3 phosphorylation; wherein the increase and/or decrease is compared to the levels of PAI-1 and STAT3 phosphorylation measured in a sample obtained from a reference group.

[00147] In one example, the PAI-1 inhibitor is an anti-cancer drug.

[00148] In the methods disclosed herein, the reference group refers to a group of subjects suffering from peritoneal carcinomatosis. In another example the reference group is a group of patients who are not suffering from peritoneal carcinomatosis, but who present with benign tumours.

[00149] Comparison of values obtained in the experiments disclosed herein result in the definition of reference values as disclosed herein (also referred to as cut-off values), which have been determined for each of the markers to be measured. The comparison between the measured values and the cut-off values can be done in a relative, qualitative manner (for example, that the concentration of one marker is more or less than the concentration of other marker) or in a quantitative manner (for example, value X is compared to value Y). Due to the nature of measurements, reference values or cut-off values can also include a buffer around the specific values. For example, a cut-off value with a 2% buffer means that if the cut-off value is 10, the buffer would result in a range of 9.8 to 10.2 being allowable for measurements. Depending on the context of the cut-off value, a buffer can also be applied in only one direction. For example, if the cut-off value is at least 10, then a buffer of 2% would result in a value of 9.8 also being acceptable. If the cut-off value is no more than 10, then a buffer of 2% would result in a value of 10.2 also being acceptable. In another example, the buffer can be 3%, 4% or 5% of the cut-off value in question. In another example, the buffer is 5% of the cut-off value in question. In another example, the buffer is 2% of the cut-off value in question.

[00150] Also envisioned in the scope of the present application is a system for detecting the markers, surrogate or otherwise, disclosed herein. For example, such a detection system is to be capable of diagnosing or detecting or predicting the likelihood of a patient or subject having peritoneal carcinomatosis. Accordingly, the biomarkers as described herein can be incorporated in diagnostic tools, detection systems, methods of diagnosis, methods of predicting or methods of determining the likelihood of a patient having peritoneal carcinomatosis. Exemplary detection system can comprise, for example, a receiving section to receive a sample from a patient suspected to suffer from peritoneal carcinomatosis, wherein the sample is suspected to comprise one or more biomarkers of the present disclosure, and a detection section comprising a substance or substances capable of detecting one or more biomarkers of the present disclosure. The samples used in this system can be, but are not limited to, the sample types disclosed here.

[00151] To assist in detecting the biomarkers of the present disclosure, the detection system can comprise a substance capable of binding or specifically binding to any of the biomarkers disclosed herein. For example, such substances can be biospecific capture reagents such as antibodies (or antigen-binding

fragments thereof), interacting fusion proteins, aptamers or affibodies (which are non-immunoglobulin-derived affinity proteins based on a three-helical bundle protein domain) that recognize the biomarker and/or variants thereof. In use, the substance can, for example, be bound to a solid phase, wherein the biomarkers can be detected methods known in the art, for example, mass spectrometry, or by eluting the biomarkers from the biospecific capture reagents and detecting the eluted biomarkers using methods known in the art, for example, a traditional matrix-assisted laser desorption/ionization (MALDI) or by surface-enhanced laser desorption/ionization (SELDI). For example, the detection system comprised on a biochip, test strip, or microtiter plate.

[00152] A companion biomarker that dictates therapy based on the concept of oncogenic addiction in peritoneal carcinomatosis patients has been identified. The biomarker that has been identified that activates STAT3 and other signalling pathways is part of the coagulation cascade. In other words, activation of the coagulation cascade after surgery can stimulate growth of cancer cells. It is further thought that hyper-activation of coagulation or the fibrinolytic cascade is oncogenic and inhibition of these two processes has shown potential therapeutic relevance. In addition, gene expression profiling of 2 colorectal peritoneal carcinomatosis cell lines treated with cell-free ascites revealed activation of STAT3 signalling. Furthermore, validation experiments on cell-free ascites ($n = 13$) demonstrated STAT3 to be most relevant in colorectal peritoneal carcinomatosis. Clinically, colorectal cancer patients in the TCGA database ($n = 345$) with STAT3 and epithelial-mesenchymal transition (EMT) activation had poorer prognosis. Interestingly, receptor tyrosine kinase arrays showed no phosphorylation of JAK kinase, suggesting non-canonical activation of STAT3 signalling. Cytokine array and mass spectrometry identified potential STAT3 activating ligands independent of JAK kinase including POSTN, CD24, and CD44. Treatment of cell lines exposed to cell-free ascites demonstrated sensitivity to inhibitors of upstream non-canonical STAT3 activator in *in vitro* and *in vivo* settings.

[00153] The invention illustratively described herein may suitably be practiced in the absence of any element or elements, limitation or limitations, not specifically disclosed herein. Thus, for example, the terms "comprising", "including", "containing", etc. shall be read expansively and without limitation. Additionally, the terms and expressions employed herein have been used as terms of description and not of limitation, and there is no intention in the use of such terms and expressions of excluding any equivalents of the features shown and described or portions thereof, but it is recognized that various modifications are possible within the scope of the invention claimed. Thus, it should be understood that although the present invention has been specifically disclosed by preferred embodiments and optional features, modification and variation of the inventions embodied therein herein disclosed may be resorted to by those skilled in the art, and that such modifications and variations are considered to be within the scope of this invention.

[00154] As used in this application, the singular form "a," "an," and "the" include plural references unless the context clearly dictates otherwise. For example, the term "a genetic marker" includes a plurality of genetic markers, including mixtures and combinations thereof.

[00155] As used herein, the term “about”, in the context of concentrations of components of the formulations, typically means +/- 5% of the stated value, more typically +/- 4% of the stated value, more typically +/- 3% of the stated value, more typically, +/- 2% of the stated value, even more typically +/- 1% of the stated value, and even more typically +/- 0.5% of the stated value.

5 [00156] Throughout this disclosure, certain embodiments may be disclosed in a range format. It should be understood that the description in range format is merely for convenience and brevity and should not be construed as an inflexible limitation on the scope of the disclosed ranges. Accordingly, the description of a range should be considered to have specifically disclosed all the possible sub-ranges as well as individual numerical values within that range. For example, description of a range such as from 1 to 6
10 should be considered to have specifically disclosed sub-ranges such as from 1 to 3, from 1 to 4, from 1 to 5, from 2 to 4, from 2 to 6, from 3 to 6 etc., as well as individual numbers within that range, for example, 1, 2, 3, 4, 5, and 6. This applies regardless of the breadth of the range.

[00157] Certain embodiments may also be described broadly and generically herein. Each of the narrower species and sub-generic groupings falling within the generic disclosure also form part of the
15 disclosure. This includes the generic description of the embodiments with a proviso or negative limitation removing any subject matter from the genus, regardless of whether or not the excised material is specifically recited herein.

[00158] The invention has been described broadly and generically herein. Each of the narrower species and sub-generic groupings falling within the generic disclosure also form part of the invention. This
20 includes the generic description of the invention with a proviso or negative limitation removing any subject matter from the genus, regardless of whether or not the excised material is specifically recited herein.

[00159] Other embodiments are within the following claims and non-limiting examples. In addition, where features or aspects of the invention are described in terms of Markush groups, those skilled in the
25 art will recognize that the invention is also thereby described in terms of any individual member or subgroup of members of the Markush group.

EXPERIMENTAL SECTION

Materials and methods

30 Patient recruitment, biospecimen collection and processing

[00160] Patients who were undergoing treatment for peritoneal carcinomatosis at the National Cancer Centre Singapore were identified and recruited. Informed consent was obtained from all patients in accordance to study protocol approved by the SingHealth Centralized Institutional Review Board (CIRB Ref: 2015/2479/F). All experiments were performed in accordance with the relevant guidelines and
35 regulations. Tumour specimens harvested post-operatively were systematically divided into multiple pieces. One portion was snap-frozen in liquid nitrogen immediately and stored in -80°C freezer while another was fixed in formalin to construct formalin-fixed paraffin embedded (FFPE) block. Remaining

tissues were processed in the laboratory to establish primary cell lines and patient derived xenografts. Ascites collected from the peritoneal cavity at the beginning of the cytoreductive surgery (CRS) or during routine ascites tap (paracentesis) was subjected to centrifugation at 2000 g for 10 minutes to separate the cellular and fluid component. Filter-sterilization using 0.22 µm filter was performed on the fluid component to render it suitable for downstream experiments. The cellular component of ascites was used for downstream assays and the generation of patient-derived ascites-dependent xenograft (PDADX) models.

Cell lines

[00161] Human metastatic colon cancer cell line Colo-205 and SNU-C1 were purchased from American Type Culture Collection and cultured in RPMI medium with 10% foetal bovine serum (FBS), 1% penicillin-streptomycin and 1% antimycotic. Human normal peritoneal mesothelial cell lines LP9/TERT and HM3/TERT were purchased from Brigham and Women's Hospital Cell Culture Core and cultured in M199/M106 with 15% iron-supplemented new-born calf serum, 0.4 µg/mL hydrocortisone, 10 ng/mL epidermal growth factor, 1% penicillin-streptomycin and 1% antimycotic. All cells were grown in serum-free medium for overnight prior to experiments.

Primary clinical endpoint

[00162] The primary clinical endpoint was overall survival (OS). OS is defined as the time from surgery to death, regardless of cause. Kaplan-Meier curves were plotted to compare 5-year overall survival (OS) by the presence or absence of ascites. Presence of ascites is defined by accumulation of more than 50 mL fluid in the abdominal cavity. Log-rank test was used to determine statistical significance for curve comparison.

Proliferation assay

[00163] A total of 5000 cells/wells were seeded in 96-well plates and were grown in serum-free RPMI medium supplemented with varying concentration of cell-free ascites. Cell proliferation was assessed using CellTitreGlo assay (Promega, Madison, US) at day 0 and day 5. These experiments were performed in triplicates and repeated 3 times.

Cell migration assay

[00164] Colo-205 or SNU-C1 cells were serum starved for 24 hours and subsequently treated with 3 different cell-culture media: serum-free RPMI, RPMI supplemented with 10% foetal bovine serum (FBS), or serum-free media supplemented with 5% cell-free ascites for 24 hours. Pre-treated cells were then seeded into 6-well transwell migration assay at a density of 600,000 cells/well. The inner chamber of the transwell plate was filled with serum-free media and the outer chamber was filled with 10% foetal bovine serum (FBS) media. Cells were allowed to migrate for 24 hours. These experiments were performed in triplicates and repeated 3 times.

Cell settlement assay

[00165] A total of 70,000 cells/well of LP9/TERT or HM3/TERT were seeded in 12-well plates and were grown to confluency in complete media to form the feeder layer. Subsequently, the mesothelial feeder

layer was serum starved prior to co-culture with cancer cells. 35,000 cells/well of Colo-205 or SNU-C1 were seeded into each well in 3 different medium: serum-free RPMI, RPMI supplemented with 10% foetal bovine serum (FBS), or serum-free RPMI supplemented with 5% cell-free ascites, and incubated for 24 hours. Non-attached cancer cells were removed by gentle washing with complete media for 5 times. The average number of cells settled in three fields per well was counted. The final number of cells settled was determined by the mean of triplicate assays.

Gene expression profiling

[00166] To assess signalling pathways upregulated upon treatment with cell-free ascites, Colo-205 and SNU-C1 cells were treated with 5% and 0.1% cell-free ascites for 24 hours. To assess signalling pathways affected by PAI-1 inhibition, Colo-205 cells were treated with cell-free ascites representative of PAI-1 paracrine addicted (PPA) group, co-activators predominant (CAP) group or foetal bovine serum (FBS; control) in the presence of DMSO vehicle or 27.25 μ M TM5441 for 24 hours. Total RNA was isolated using Qiagen Mini Kit (Qiagen, CA, USA), following the manufacturer's instructions. Gene expression profiling was performed using Affymetrix GeneChip Genome U133 Plus 2.0 microarray platform (Affymetrix, Santa Clara, CA) in accordance to manufacturer's protocols. Microarray data was uploaded into the free programming software R (R Foundation for Statistical Computing, Vienna, Austria) for processing and normalization. Gene Set Enrichment Analysis (GSEA) was used to assess enrichment of genes showing up- and down-regulation using GSEA graphical user interface (GUI) software (<http://www.broadinstitute.org/gsea/>).

Protein immunoblotting

[00167] Colo-205 or SNU-C1 cells were starved in serum-free media overnight before treating with 5% of patients' cell-free ascites for 24 hours. On the next day, cells were harvested and lysed in M-PER (Mammalian Protein Extraction Reagent, Thermo Scientific Inc.) supplemented with Pierce Protease and Phosphatase Inhibitor (Thermo Scientific Inc.) for 1 hour on ice. The lysates were centrifuged at 14,000 g for 20 minutes at 4°C to obtain clear supernatants. Protein concentrations were determined using the Bradford protein assay reagent (Bio-Rad). Specific amount of proteins were calculated (5 μ g for STAT3 and actin; 25 μ g for phospho-STAT3 (Tyr705) and phospho-STAT3 (Ser727); 10 μ g for JAK1, JAK2, phospho-JAK1 (Tyr1022/Tyr1023) and phospho-JAK2 (Tyr1007/Tyr1008)) and aliquoted into 0.2 mL thin wall PCR tubes. Lysates were denatured at 97°C for 5 minutes and resolved in 10% polyacrylamide gels in Tris/glycine/SDS running buffer (24.76 mM Tris, 191.83 mM glycine and 0.1% SDS) and then transferred to 0.45 μ m nitrocellulose membrane (Bio-Rad) in Tris/glycine/methanol transfer buffer (24.76 mM Tris, 191.83 mM glycine and 20% methanol). The membranes were blocked with 5% non-fat milk in 1x PBS containing 0.1% Tween 20 (PBST) for 1 hour at room temperature before blotting with primary antibodies for 1.5 hours. Dilutions of the primary antibodies were: 1:2,000 STAT3 (Cell Signaling Technology; #4904); 1:1,000 phospho-STAT3 (Tyr705) (Cell Signaling Technology; #9145); 1:1,000 phospho-STAT3 (Ser727) (Cell Signaling Technology; #94994); 1:1,000 JAK1 (Santa Cruz Biotechnology; sc-277); 1:1,000 phospho-JAK1 (Tyr1022/Tyr1023) (Santa Cruz Biotechnology; sc-

16773); 1:1,000 JAK2 (Santa Cruz Biotechnology; sc-294); 1:1,000 phospho-JAK2 (Tyr1007/Tyr1008) (Santa Cruz Biotechnology; sc-16566) and 1:100,000 β -actin (Sigma Aldrich; A1978). After 4 washes (5 min per wash) in PBST, the blots were incubated with anti-rabbit or anti-mouse horseradish peroxidase (HRP)-linked secondary antibody (GE Healthcare Life Sciences; NA934 or NA931) for 30 minutes at room temperature. After another 4 washes in PBST, Pierce SuperSignal West Dura Extended Duration Substrate (Thermo Scientific Inc.) was added to the blots and incubated for 5 minutes at room temperature. Excess liquid was dripped off and the blots were wrapped in polyethylene for exposure to UltraCruz® Autoradiography Film (Santa Cruz Biotechnology, CA). Images were scanned using GS-800 TM calibrated densitometer (Bio-Rad).

10 Immunohistochemistry (IHC)

[00168] Formalin-fixed paraffin-embedded (FFPE) specimens from peritoneal carcinomatosis (PC) cases with matched primary tumour and metastases were identified and interrogated using chromogen-based immunohistochemical (IHC) staining. All immunohistochemical (IHC) staining was carried out using the Bond Max Autostainer (Leica Microsystems, Ltd, Milton Keynes, UK) in accordance to the manufacturer's recommendations. Formalin-fixed paraffin-embedded (FFPE) blocks were sectioned into 4 μ m thick sections and mounted on slides. Rabbit monoclonal anti phospho-STAT3 (Tyr705) (#9145L, Cell Signalling Technology, Massachusetts, US, 1:50, pH9, 30 minutes) was optimized and used. Slides were evaluated by two independent scorers, who had no prior knowledge of the clinical data, and staining results were determined based on the percentage of positive staining within the tumour epithelial component in each slide. Formalin-fixed paraffin-embedded (FFPE) samples were also collected from patient-derived ascites-dependent xenograft (PDADX) tumours and probed with antibodies against CK7, CK20 and CDX2 to confirm histology and origin of patient-derived ascites-dependent xenograft (PDADX) tumours formed. Rabbit monoclonal anti-CK7 (#31-1167-00, RevMab Biosciences, California, US, 1:200, pH9, 20 minutes), rabbit polyclonal anti-CK20 (HPA024309, Sigma Aldrich, Missouri, US, 1:200, pH9, 20 minutes) and rabbit monoclonal anti-CDX2 (#12306, Cell Signaling Technology, Massachusetts, US, 1:100, pH 9, 20 minutes) were optimized and used in the immunohistochemical (IHC) staining.

Mass spectrometry

[00169] Mass spectrometry was performed on proteins isolated from the soluble and exosomal components of cell-free ascites from patients with benign serous cystadenofibroma ($n = 1$) and malignant cell-free ascites from patients with colorectal peritoneal carcinomatosis ($n = 3$). Briefly, liquid chromatography coupled with tandem mass spectrometry (LC-MS/MS) analysis of digested peptides was carried out using a Thermo Scientific Inc. Orbitrap Elite and QExactive mass spectrometers (Bremen, Germany) coupled with a Dionex UltiMate 3000 UHPLC system from Thermo Scientific Inc.

35 Cytokine profiling

[00170] Proteome Profiler Human Cytokine array (ARY022B, R&D Systems, Minneapolis, US) consisting of 105 cytokines were used to profile plasma ($n = 1$), benign cell-free ascites from patients

with benign serous cystadenofibroma ($n = 1$) and malignant cell-free ascites from patients with colorectal peritoneal carcinomatosis ($n = 4$), gastric peritoneal carcinomatosis ($n = 1$), and ovarian peritoneal carcinomatosis ($n = 1$). Concentration of proteins in all fluids was quantified with Bradford protein assay (Biorad, Hercules, US) and equal amount of proteins were incubated with the membrane array following the manufacturer's instructions.

Epithelial-Mesenchymal Transition (EMT) gene profiling

[00171] Epithelial-Mesenchymal transition (EMT) gene profiling was performed using RT2 Profiler PCR Arrays (Qiagen, CA, USA) comprising 84 EMT-related genes. RNA was extracted from Colo-205 and SNU-C1 cells grown in complete media and 5% cell-free ascites for 24 hours using RNeasy extraction kit (Qiagen). cDNA was synthesized using RT2 First Strand kit (Qiagen) and reverse-transcription polymerase chain reaction was performed using RT2 SYBR Green Mastermixes (Qiagen). The results were analysed using Qiagen's Gene Globe Data Analysis Centre tool.

Phospho-Receptor Tyrosine Kinases (RTKs) profiling

[00172] Profiling of phosphorylation of receptor tyrosine kinases (RTKs) was performed using human RTK phosphorylation antibody array (RayBiotech, GA, USA) which allows simultaneous detection of relative phosphorylation levels of 71 different human RTKs in cell lysate. Proteins were extracted from cell lysates of Colo-205 and SNU-C1 that were treated with 5% cell-free ascites or 10% foetal bovine serum (FBS) for 24 hours. Total protein concentration was determined with Bradford protein assay (Biorad, Hercules, US) and 40 μ g of protein was used for phospho-RTKs profiling following the manufacturer's instructions.

The Cancer Genome Database (TCGA) survival analysis

[00173] Kaplan-Meier overall survival (OS) curve analysis was used to determine prognostic significance of PAI-1, STAT3 and epithelial-mesenchymal transition (EMT) regulation in The Cancer Genome Database (TCGA) colorectal adenocarcinoma (COADREAD) data set ($n = 345$). Patients were stratified high ($P+$, ≥ 3.071) or low ($P-$, < 3.071) PAI-1 expression, high ($S+$, ≥ 0.074) or low ($S-$, < 0.074) STAT3 expression, and high ($E+$, ≥ 0.096) or low ($E-$, < 0.096) epithelial-mesenchymal transition (EMT) expression based on cut-offs determined by recursive partitioning. Four subtypes ($P-S-E+$, $P+S-E-$, $P+S-E+$, $P+S+E-$) were excluded from analysis due to small sample size ($n < 20$). Log-rank test was used to test for statistical significance.

Enzyme-linked Immunosorbent Assay (ELISA)

[00174] Concentrations of PAI-1 (DSE100), IL6 (D6050), IL10 (D1000B), CCL2 (DCP00), MMP9 (DMP900), ANGPT1 (DANG10), TGFB1 (DB100B), and CXCL10 (DIP100) in cell-free ascites were quantified using human Quantikine ELISA kits from R&D Systems. Concentrations of POSTN (DY3548B) and CD44 (DY7045-05) in cell-free ascites were quantified using human DuoSet ELISA kits from R&D Systems. VSIG4 (ELH-VSIG4-1) concentration in cell-free ascites was quantified using ELISA from RayBiotech. All samples were performed with 2 technical replicates according to the manufacturers' instructions. Total STAT3 and phospho-STAT3 (Tyr705) were detected with ELISA

(7305C and 7300C, Cell Signalling Technology, Massachusetts, US). Proteins were isolated from cell lysates of Colo-205 and SNU-C1 that were treated with 5% cell-free ascites for 24 hours. In all experiments, 25 µg of protein was used for total STAT3 and p-STAT3(Y705) ELISA following the manufacturer's instructions.

5 ***In vitro* drug treatment**

[00175] A total of 5,000 cells/wells were seeded in 96-well plates and were grown for 24 hours in serum-free RPMI medium supplemented with 5% cell-free ascites or complete media, and then treated with various concentrations of TM5441 (PAI-1 inhibitor), Tiplaxtinin (PAI-1 inhibitor), Napabucasin (STAT3 inhibitor), BEZ235 (dual PI3K/mTOR inhibitor) and Mitomycin C (chemotherapeutic agent used in hyperthermic intraperitoneal chemotherapy (HIPEC)) for 72 hours. Cell proliferation was assessed using CellTitreGlo assay (Promega, Madison, US). These experiments were performed in triplicates and were repeated at least 3 times.

Patient-derived ascites-dependent xenografts (PDADXs) generation

[00176] All mice experiments were performed according to protocols approved by SingHealth Institutional Animal Care and Use Committee (IACUC Ref: 2017/SHS/1295). Ascites collected from patients with peritoneal carcinomatosis were centrifuged at 2000 g for 10 minutes to concentrate the cellular components and to separate the fluid component. 1 mL of cell pellet was resuspended with 1 mL of ascitic fluid and 400 µL of the mixture was implanted intraperitoneally into 6-week-old BALB/c nude mice ($n = 5$ mice) to generate patient-derived ascites-dependent xenograft (PDADX) passage 0 (P0). For subsequent passages, patient-derived ascites-dependent xenograft (PDADX) tumours were diced into small pieces using scalpel blades and passed through an 18-G syringe needle. Diced tumours were resuspended with matched patient's ascites at 1:1 ratio and implanted intraperitoneally into 6-week-old BALB/c nude mice ($n = 10$ mice).

***In vivo* PC cell line mouse model drug treatment**

[00177] To determine the efficacy of PAI-1 inhibition in different susceptibility ascites groups *in vivo*, 5×10^6 of Colo-205 cells were co-injected with cell-free ascites representative of PAI-1 paracrine addicted (PPA) group, co-activators predominant (CAP) group or foetal bovine serum (FBS) into abdominal cavity of 6- to 8-week-old BALB/c nude mice (female, $n = 5$ mice/group) and treated with 1.75 mM TM5441 administered intraperitoneally. Ascites and drug treatment were performed by injecting 400 µL of 5% cell-free ascites or 10% foetal bovine serum (FBS) with TM5441 intraperitoneally every 3 days for a duration of 21 days. After 3 weeks, the mice were sacrificed and tumour burden was quantified based on a modified peritoneal carcinomatosis index (PCI) score and presented as total peritoneal carcinomatosis index (PCI) score. Total peritoneal carcinomatosis index (PCI) score was calculated based on the sum of score for each region and ranges from 0 to 39.

[00178] To determine the optimal drug concentration and drug delivery method, a total of 16 female BALB/c nude mice with the age of 6–8 weeks old were selected for the experiment. Each mouse was injected with 5×10^6 of Colo-205 cells intraperitoneally. The mice were divided into 4 groups and given

the following treatments: (i) 5% cell-free ascites with 1% DMSO, (ii) 5% cell-free ascites with 1 mM TM5441, (iii and iv) 5% cell-free ascites with 2 mM TM5441. Treatment was performed by injecting 400 µL of 5% cell-free ascites with DMSO vehicle/drug intraperitoneally (i – iii) and orally (iv) every 3 days, up to 21 days. After 3 weeks, the mice were sacrificed and tumour burden was quantified based on a modified peritoneal carcinomatosis index (PCI) score and presented as total peritoneal carcinomatosis index (PCI) score. Total peritoneal carcinomatosis index (PCI) score was calculated based on the sum of score for each region and ranges from 0 to 39.

Patient-derived ascites-dependent xenografts (PDADXs) drug treatment

[00179] Matched patient's cell-free ascites and its cellular components were used to generate patient-derived ascites-dependent xenografts (PDADXs) to better recapitulate peritoneal carcinomatosis patients. PAI-1 paracrine addicted (PPA) patient-derived ascites-dependent xenograft (PDADX) tumours (100 mg) were implanted into 16 female BALB/c nude mice intraperitoneally and co-activators predominant (CAP) patient-derived ascites-dependent xenograft (PDADX) tumours (100 mg) were implanted into 16 female BALB/c nude mice intraperitoneally. PAI-1 paracrine addicted (PPA) patient-derived ascites-dependent xenograft (PDADX) and co-activators predominant (CAP) patient-derived ascites-dependent xenograft (PDADX) were then divided into 4 groups and given the following treatments: (i) 5% PAI-1 paracrine addicted (PPA) or co-activators predominant (CAP) cell-free ascites with 1% DMSO, (ii) 5% PAI-1 paracrine addicted (PPA) or co-activators predominant (CAP) cell-free ascites with 2 mM TM5441, (iii) 10% foetal bovine serum (FBS) with 1% DMSO and (iv) 10% foetal bovine serum (FBS) with 2 mM TM5441. Treatment was performed via intraperitoneal administration every 3 days for 21 days. Tumour burden was quantified by weighing all visible tumours after mice were sacrificed.

[00180] To determine if susceptibility to PAI-1 inhibition is reliant on cell-free ascites and not tumours, co-activators predominant (CAP) patient-derived ascites-dependent xenograft (PDADX) whose patient's cell-free ascites are not responsive to PAI-1 inhibition was treated with PAI-1 paracrine addicted (PPA) cell-free ascites. Briefly, co-activators predominant (CAP) patient-derived ascites-dependent xenograft (PDADX) tumours (100 mg) were implanted into 16 female BALB/c nude mice intraperitoneally. The mice were divided into 4 groups and given the following treatment: (i) 5% co-activators predominant (CAP) cell-free ascites with 1% DMSO, (ii) 5% co-activators predominant (CAP) cell-free ascites with 2 mM TM5441, (iii) 5% PAI-1 paracrine addicted (PPA) cell-free ascites with 1% DMSO, and (iv) 5% PAI-1 paracrine addicted (PPA) cell-free ascites with 2 mM TM5441. Treatment was performed via intraperitoneal administration every 3 days for 21 days. Tumour burden was quantified by weighing all visible tumours after mice were sacrificed.

p-STAT3 surrogate marker selection

[00181] STAT3-related genes were identified from Kyoto Encyclopedia of Genes and Genomes (KEGG) database by compiling all genes that are involved in known STAT3 pathways. Secreted STAT3-related proteins were selected based on extracellular genes listed in NCBI's Biosystems database and proteins identified in mass spectrometry analysis of cell-free ascites. Transcriptomics comparison was performed

using 2 databases to prioritize putative STAT3 surrogate markers. First database was used to determine genes that are positively correlated with STAT3 in TCGA COADREAD data set. Genes were ranked from most positively correlated to least correlated with STAT3. Second database was derived from microarray analysis of PAI-1 paracrine addicted (PPA) cell-free ascites-treated cells exposed to TM5441 to determine genes that are downregulated and upregulated in PPA cell-free ascites-treated cells in response to PAI-1 inhibition. Upregulated genes were also of interest as these might represent genes that are involved in rescue mechanisms in response to PAI-1 inhibition. Similarly, genes were ranked from most downregulated to most upregulated. Systematic paired correlation analysis of candidate genes was subsequently performed by focusing on top 1% and top 25% of genes positively correlated with STAT3 in database 1, and top 1% and top 25% of most downregulated and upregulated genes in database 2. The paired analysis for each group was prioritised and representative genes were chosen from each group based on literature review to streamline to 35 genes. 10 targets were selected based on rank prioritisation, potential good correlation with p-STAT3 from Luminex assay data, and the importance of the candidate genes in cancer pathogenesis from literature review for further evaluation with ELISA. The concentrations of each surrogate marker in cell-free ascites were correlated with ascites-treated cells p-STAT3 levels using Spearman correlation analysis.

CLAIMS

1. Use of a “plasminogen activator inhibitor 1” (PAI-1) inhibitor in the manufacture of a medicament for treating a subject suffering from peritoneal carcinomatosis, wherein the subject is to be identified by

determining the concentration of PAI-1 and determining the level of phosphorylation of “signal transducer and activator of transcript 3” (STAT3) in a sample obtained from the subject;

wherein if the subject shows (a) an increase in PAI-1 concentration and an increase in STAT3 phosphorylation, or (b) a decrease in PAI-1 concentration and an increase in STAT3 phosphorylation, the subject is to be administered the medicament;

wherein the increase and/or decrease of the concentration of PAI-1 and STAT3 phosphorylation is compared to a reference value;

wherein the sample is selected from the group consisting of ascites, tumour samples, and biopsy samples;

wherein the reference values are obtained from a reference group comprising subjects suffering from peritoneal carcinomatosis.

2. A method of detecting or determining susceptibility of a subject suffering from peritoneal carcinomatosis to a treatment with a “plasminogen activator inhibitor 1” (PAI-1) inhibitor, the method comprising

determining the concentration of PAI-1 and determining the level of phosphorylation of “signal transducer and activator of transcript 3” (STAT3) in a sample obtained from a subject;

wherein the subject is susceptible to the treatment if the subject shows (a) an increase in PAI-1 concentration and an increase in STAT3 phosphorylation, or (b) a decrease in PAI-1 concentration and an increase in STAT3 phosphorylation;

wherein the subject is not considered to be susceptible to treatment if the subject shows (c) a decrease in PAI-1 concentration and a decrease in STAT3 phosphorylation;

wherein the increase and/or decrease of the concentration of PAI-1 and the level of STAT3 phosphorylation is compared to a reference value;

wherein the sample is selected from ascites, tumour samples, and biopsy samples;

wherein the reference values are obtained from a reference group comprising subjects suffering from peritoneal carcinomatosis.

3. The use of claim 1 or the method of claim 2, wherein the concentration of PAI-1 is determined by measuring the concentration of PAI-1 directly, and/or in one or more complexes.

4. The use of claim 1 or 3, or the method of claims 2 to 3, wherein the level of STAT3 phosphorylation is determined by measuring the concentration of two or more surrogate markers, and/or

by measuring the concentration of STAT3 phosphorylation directly, wherein the two or more surrogate markers are selected from the group consisting of IL6, IL10, CCL2, MMP9, ANGPT1, TGFB1, POSTN, VSIG4, CD44, and CXCL10, wherein one of the two or more surrogate markers is MMP9.

5 5. The use or the method of claim 4, wherein the two or more surrogate markers are selected from the group consisting of IL6, IL10, CCL2, MMP9 and ANGPT1, wherein one of the two or more surrogate markers is MMP9.

6. The use or the method of any one of claims 4 to 5, wherein the two or more surrogate markers are selected from the group consisting of IL6, IL10, CCL2, and MMP9, wherein one of the two or more surrogate markers is MMP9.

7. The use or the method of any one of claims 4 to 6, wherein the two or more surrogate markers comprise IL6, IL10, CCL2, MMP9 and ANGPT1, wherein one of the two or more surrogate markers is MMP9; or wherein the surrogate markers comprise at least IL6, IL10, CCL2, and MMP9, wherein one of the two or more surrogate markers is MMP9.

8. The use of claim 1 or 3 to 7, or the method of any one of claims 2 to 7, wherein the PAI-1 inhibitor is an anti-cancer drug or an anti-cancer treatment.

9. The use or the method of claim 8, wherein the anti-cancer drug or the anti-cancer therapy is selected from the group consisting of a small molecule, a chemotherapeutic agent, a peptide, an antibody, combinations thereof, and combination therapy.

10. The use or the method of any one of claims 8 to 9, wherein the anti-cancer drug is selected from the group consisting of TM5441 (5-Chloro-2-[[2-[2-[[3-(3-furanyl)phenyl]amino]-2-oxoethoxy]acetyl]amino]benzoic acid sodium salt; CAS 1190221-43-2), TM5007 (N, N-bis [3,3'-carboxy-4,4'-(2,2'-thienyl)-2,2'- thienyl] hexanedicarboxamide; CAS 342595-05-5), TM5275 (5-Chloro-2-[[2-[2-[4-(diphenylmethyl)-1-piperazinyl]-2-oxoethoxy]acetyl]amino]-benzoic acid sodium salt; CAS 1103926-82-4), tiplaxtinin (2-(1-Benzyl-5-(4-(trifluoromethoxy)phenyl)-1H-indol-3-yl)oxoacetic acid; CAS 393105-53-8), and ZK4044.

11. The use of claim 1 or claims 3 to 10, wherein the PAI-1 inhibitor is to be administered intraperitoneally.

12. The use of claim 1 or claims 3 to 11, or the method according to any one of claims 2 to 10, wherein the peritoneal carcinomatosis is selected from the group consisting of colorectal peritoneal carcinomatosis, small bowel peritoneal carcinomatosis, mesothelioma, endometrial peritoneal

carcinomatosis, gastric peritoneal carcinomatosis, ovarian peritoneal carcinomatosis, appendiceal peritoneal carcinomatosis, pancreatic peritoneal carcinomatosis, urothelial carcinomatosis and Pseudomyxoma peritonei (PMP).

- 5 13. The use of claim 1 or claims 3 to 12, or the method according to any one of claims 2 to 10 and 12, wherein the treatment is to be performed in a treatment setting selected from the groups consisting of neoadjuvant setting, adjuvant setting, palliative setting and prophylactic setting.
14. The use of claim 1 or claims 3 to 13, or the method of any of claims 2 to 10 and 12 to 13, wherein
10 administration of the PAI-1 inhibition leads to inhibition of PAI-1 activity compared to patients suffering from the same disease.
15. Use of a panel of markers in the use of claim 1 or claims 3 to 14, or the method of any one of claims 2 to 10 and 12 to 14, wherein the panel comprises PAI-1 and two or more surrogate markers of
15 STAT3 phosphorylation, wherein at least one of the surrogate markers of STAT3 phosphorylation is MMP9, or PAI-1 and p-STAT3; wherein the two or more surrogate markers are selected from the group consisting of IL6, IL10, CCL2, MMP9, ANGPT1, TGFB1, POSTN, VSIG4, CD44, and CXCL10, wherein one of two or more surrogate markers is MMP9.
- 20 16. The use of claim 15, wherein the panel comprises PAI-1, and two or more or all of IL6, IL10, CCL2, and MMP9, wherein one of the surrogate markers is MMP9.
17. The use of claims 15 to 16, wherein the panel comprises PAI-1, and one or more or all of IL6, IL10, CCL2, MMP9 and ANGPT1, wherein one of the surrogate markers is MMP9.