
In one aspect, the present disclosure provides use of a compound of formula 1 in the manufacture of a medicament for treating aging-associated neuronal loss, memory impairment, or loss of motor coordination in a subject diagnosed with the aging-associated neuronal loss, memory impairment, or loss of motor coordination, wherein the compound of formula 1 is to be administered in the subject and has the following formula:



1

wherein

A is CH_2, O or N-C_{1-6}-alkyl;

R^1 is selected from

- $NHR^{1.1}$, $NMeR^{1.1}$;
- $NHR^{1.2}$, $NMeR^{1.2}$;
- $NHCH_2$-$R^{1.3}$;
- NH-C_{3-6}-cycloalkyl, whereas optionally one carbon atom is replaced by a nitrogen atom, whereas the ring is optionally substituted with one or two residues selected from the group consisting of C_{1-6}-alkyl, O-C_{1-6}-alkyl, $NHSO_2$-phenyl, NHCONH-phenyl, halogen, CN, SO_2-C_{1-6}-alkyl, COO-C_{1-6}-alkyl;
- a $C_{9\text{ or }10}$-bicyclic-ring, whereas one or two carbon atoms are replaced by nitrogen atoms and the ring system is bound via a nitrogen atom to the basic structure of formula **1** and whereas the ring system is optionally substituted with one or two residues selected from the group consisting of C_{1-6}-alkyl, COO-C_{1-6}-alkyl, C_{1-6}-haloalkyl, O-C_{1-6}-alkyl, NO_2, halogen, CN, $NHSO_2$-C_{1-6}-alkyl, methoxy-phenyl;
- a group selected from NHCH(pyridinyl)CH_2COO-C_{1-6}-alkyl, NHCH(CH_2O-C_{1-6}-alkyl)-benzoimidazolyl, optionally substituted with halogen or CN;

- piperidinyl, optionally substituted with one or two residues selected from the group consisting of $NHSO_2$-C_{1-4}-alkyl, m-methoxyphenyl;
- or 1-aminocyclopentyl, optionally substituted with methyl-oxadiazole;

$R^{1.1}$ is phenyl, optionally substituted with one or two residues selected from the group consisting of C_{1-6}-alkyl, C_{2-6}-alkenyl, C_{2-6}-alkynyl, C_{1-6}-haloalkyl, C_{1-6}-alkylene-OH, C_{2-6}-alkenylene-OH, C_{2-6}-alkynylene-OH, $CH_2CON(C_{1-6}\text{-alkyl})_2$, $CH_2NHCONH$-C_{3-6}-cycloalkyl, CN, CO-pyridinyl, $CONR^{1.1.1}R^{1.1.2}$, COO-C_{1-6}-alkyl, $N(SO_2\text{-}C_{1-6}\text{-alkyl})(CH_2CON(C_{1-4}\text{-alkyl})_2)$ O-C_{1-6}-alkyl, O-pyridinyl, SO_2-C_{1-6}-alkyl, SO_2-C_{1-6}-alkylen-OH, SO_2-C_{3-6}-cycloalkyl, SO_2-piperidinyl, SO_2NH-C_{1-6}-alkyl, $SO_2N(C_{1-6}\text{-alkyl})_2$, halogen, CN, CO-morpholinyl, CH_2-pyridinyl or a heterocyclic ring optionally substituted with one or two residues selected from the group consisting of C_{1-6}-alkyl, NHC_{1-6}-alkyl and =O;

$R^{1.1.1}$ H, C_{1-6}-alkyl, C_{3-6}-cycloalkyl, C_{1-6}-haloalkyl, $CH_2CON(C_{1-6}\text{-alkyl},)_2$, CH_2CO-azetindinyl, C_{1-6}-alkylen-C_{3-6}-cycloalkyl, CH_2-pyranyl, CH_2-tetrahydrofuranyl, CH_2-furanyl, C_{1-6}-alkylen-OH or thiadiazolyl, optionally substituted with C_{1-6}-alkyl;

$R^{1.1.2}$ H, C_{1-6}-alkyl, SO_2C_{1-6}-alkyl;

or $R^{1.1.1}$ and $R^{1.1.2}$ together with the nitrogen atom of $CONR^{1.1.1}R^{1.1.2}$ are forming a four-, five- or six-membered carbocyclic ring, optionally containing one N or O, replacing a carbon atom of the ring, optionally substituted with one or two residues selected from the group consisting of C_{1-6}-alkyl, C_{1-4}-alkylene-OH, OH, =O;

or

$R^{1.1}$ is phenyl, wherein two adjacent residues are together forming a five- or six-membered carbocyclic aromatic or non-aromatic ring, optionally containing

independently from each other one or two N, S, or SO_2, replacing a carbon atom of the ring, wherein the ring is optionally substituted with C_{1-4}-alkyl or =O;

$R^{1.2}$ is selected from

- heteroaryl, optionally substituted with one or two residues selected from the group consisting of C_{1-6}-alkyl, C_{2-6}-alkenyl, C_{2-6}-alkynyl, C_{3-6}-cycloalkyl, CH_2COO-C_{1-6}-alkyl, $CONR^{1.2.1}R^{1.2.2}$, $COR^{1.2.3}$, COO-C_{1-6}-alkyl, $CONH_2$, O-C_{1-6}-alkyl, halogen, CN, $SO_2N(C_{1-6}$-alkyl$)_2$ or heteroaryl optionally substituted with one or two residues selected from the group consisting of C_{1-6}-alkyl;
- heteroaryl, optionally substituted with a five- or six-membered carbocyclic non-aromatic ring containing independently from each other two N, O, S, or SO_2, replacing a carbon atom of the ring;
- an aromatic or non-aromatic $C_{9\text{ or }10}$-bicyclic-ring, whereas one or two carbon atoms are replaced by N, O or S each optionally substituted with one or two residues selected from the group consisting of $N(C_{1-6}$-alkyl$)_2$, CONH-C_{1-6}-alkyl, =O;
- a heterocyclic non-aromatic ring, optionally substituted with pyridinyl;
- 4,5-dihydro-naphtho[2,1-d]thiazole, optionally substituted with NHCO-C_{1-6}-alkyl,

$R^{1.2.1}$ H, C_{1-6}-alkyl, C_{1-6}-alkylene-C_{3-6}-cycloalkyl, C_{1-4}-alkylene-phenyl, C_{1-4}-alkylene-furanyl, C_{3-6}-cycloalkyl, C_{1-4}-alkylene-O-C_{1-4}-alkyl, C_{1-6}-haloalkyl or a five- or six-membered carbocyclic non-aromatic ring, optionally containing independently from each other one or two N, O, S, or SO_2, replacing a carbon atom of the ring, optionally substituted with 4-cyclopropylmethyl-piperazinyl

$R^{1.2.2}$ H, C_{1-6}-alkyl;

$R^{1.2.3}$ a five- or six-membered carbocyclic non-aromatic ring, optionally containing independently from each other one or two N, O, S, or SO_2, replacing a carbon atom of the ring;

$R^{1.3}$ is selected from phenyl, heteroaryl or indolyl, each optionally substituted with one or two residues selected from the group consisting of C_{1-6}-alkyl, C_{3-6}-cycloalkyl, O-C_{1-6}-alkyl, O-C_{1-6}-haloalkyl, phenyl, heteroaryl;


R^2 is selected from the group consisting of C_{1-6}-alkylene-phenyl, C_{1-6}-alkylene-naphthyl, and C_{1-6}-alkylene-heteroaryl; each optionally substituted with one, two or three residues selected from the group consisting of C_{1-6}-alkyl, C_{1-6}-haloalkyl, O-C_{1-6}-alkyl, O-C_{1-6}-haloalkyl, halogen;

R^3 is H, C_{1-6}-alkyl;

R^4 is H, C_{1-6}-alkyl;

or **R^3** and **R^4** together are forming a CH_2-CH_2 group.


In another aspect, the present disclosure provides use of a crystalline salt of a compound of the formula below in the manufacture of a medicament for treating aging-associated neuronal loss, memory impairment, or loss of motor coordination in a subject diagnosed with the aging-associated neuronal loss, memory impairment, or loss of motor coordination, wherein the medicament is to be administered in the subject



In another aspect, the present disclosure provides use of a pharmaceutical composition comprising the compound as defined herein in the manufacture of a medicament for treating

aging-associated neuronal loss, memory impairment, or loss of motor coordination in a subject, wherein the pharmaceutical composition is to be administered in the subject and comprises at least one co-crystal of the compound of the formula as defined herein and a pharmaceutically acceptable carrier.

In another aspect, the present disclosure provides use of a pharmaceutical composition comprising a compound of formula 3 in the manufacture of a medicmament for treating aging-associated neuronal loss, memory impairment, or loss of motor coordination in a subject, wherein the compound of formula 3 is to be administered in the subject and has the following formula:



3

wherein

R^1 is H, C_{1-6}-alkyl, C_{0-4}-alkyl-C_{3-6}-cycloalkyl, C_{1-6}-haloalkyl;

R^2 is H, C_{1-6}-alkyl;

X is an anion selected from the group consisting of chloride or ½ dibenzoyltartrate

j is 1 or 2,


and the pharmaceutical composition further comprises a first diluent, a second diluent, a binder, a disintegrant and a lubricant.

IV. Incorporation by Reference

Claims

What is claimed:

1. Use of a compound of formula 1 in the manufacture of a medicament for treating aging-associated neuronal loss, memory impairment, or loss of motor coordination in a subject diagnosed with the aging-associated neuronal loss, memory impairment, or loss of motor coordination, wherein the compound of formula 1 is to be administered in the subject and has the following formula:



1

wherein

A is CH_2, O or N-C_{1-6}-alkyl;

R^1 is selected from

- $NHR^{1.1}$, $NMeR^{1.1}$;
- $NHR^{1.2}$, $NMeR^{1.2}$;
- $NHCH_2$-$R^{1.3}$;
- NH-C_{3-6}-cycloalkyl, whereas optionally one carbon atom is replaced by a nitrogen atom, whereas the ring is optionally substituted with one or two residues selected from the group consisting of C_{1-6}-alkyl, O-C_{1-6}-alkyl, $NHSO_2$-phenyl, NHCONH-phenyl, halogen, CN, SO_2-C_{1-6}-alkyl, COO-C_{1-6}-alkyl;
- a $C_{9 \text{ or } 10}$-bicyclic-ring, whereas one or two carbon atoms are replaced by nitrogen atoms and the ring system is bound via a nitrogen atom to the basic structure of formula 1 and whereas the ring system is optionally substituted with one or two residues selected from the group consisting of C_{1-6}-alkyl, COO-C_{1-6}-alkyl, C_{1-6}-haloalkyl, O-C_{1-6}-alkyl, NO_2, halogen, CN, $NHSO_2$-C_{1-6}-alkyl, methoxy-phenyl;

- a group selected from NHCH(pyridinyl)CH_2COO-C_{1-6}-alkyl, NHCH(CH_2O-C_{1-6}-alkyl)-benzoimidazolyl, optionally substituted with halogen or CN;
- piperidinyl, optionally substituted with one or two residues selected from the group consisting of $NHSO_2$-C_{1-4}-alkyl, m-methoxyphenyl;
- or 1-aminocyclopentyl, optionally substituted with methyl-oxadiazole;

$R^{1.1}$ is phenyl, optionally substituted with one or two residues selected from the group consisting of C_{1-6}-alkyl, C_{2-6}-alkenyl, C_{2-6}-alkynyl, C_{1-6}-haloalkyl, C_{1-6}-alkylene-OH, C_{2-6}-alkenylene-OH, C_{2-6}-alkynylene-OH, $CH_2CON(C_{1-6}\text{-alkyl})_2$, $CH_2NHCONH$-C_{3-6}-cycloalkyl, CN, CO-pyridinyl, $CONR^{1.1.1}R^{1.1.2}$, COO-C_{1-6}-alkyl, $N(SO_2\text{-}C_{1-6}\text{-alkyl})(CH_2CON(C_{1-4}\text{-alkyl})_2)$ O-C_{1-6}-alkyl, O-pyridinyl, SO_2-C_{1-6}-alkyl, SO_2-C_{1-6}-alkylen-OH, SO_2-C_{3-6}-cycloalkyl, SO_2-piperidinyl, SO_2NH-C_{1-6}-alkyl, $SO_2N(C_{1-6}\text{-alkyl})_2$, halogen, CN, CO-morpholinyl, CH_2-pyridinyl or a heterocyclic ring optionally substituted with one or two residues selected from the group consisting of C_{1-6}-alkyl, NHC_{1-6}-alkyl and =O;

$R^{1.1.1}$ H, C_{1-6}-alkyl, C_{3-6}-cycloalkyl, C_{1-6}-haloalkyl, $CH_2CON(C_{1-6}\text{-alkyl},)_2$, CH_2CO-azetindinyl, C_{1-6}-alkylen-C_{3-6}-cycloalkyl, CH_2-pyranyl, CH_2-tetrahydrofuranyl, CH_2-furanyl, C_{1-6}-alkylen-OH or thiadiazolyl, optionally substituted with C_{1-6}-alkyl;

$R^{1.1.2}$ H, C_{1-6}-alkyl, SO_2C_{1-6}-alkyl;

or $R^{1.1.1}$ and $R^{1.1.2}$ together with the nitrogen atom of $CONR^{1.1.1}R^{1.1.2}$ are forming a four-, five- or six-membered carbocyclic ring, optionally containing one N or O, replacing a carbon atom of the ring, optionally substituted with one or two residues selected from the group consisting of C_{1-6}-alkyl, C_{1-4}-alkylene-OH, OH, =O;

or

$R^{1.1}$ is phenyl, wherein two adjacent residues are together forming a five- or six-membered carbocyclic aromatic or non-aromatic ring, optionally containing independently from each other one or two N, S, or SO_2, replacing a carbon atom of the ring, wherein the ring is optionally substituted with C_{1-4}-alkyl or =O;

$R^{1.2}$ is selected from

- heteroaryl, optionally substituted with one or two residues selected from the group consisting of C_{1-6}-alkyl, C_{2-6}-alkenyl, C_{2-6}-alkynyl, C_{3-6}-cycloalkyl, CH_2COO-C_{1-6}-alkyl, $CONR^{1.2.1}R^{1.2.2}$, $COR^{1.2.3}$, COO-C_{1-6}-alkyl, $CONH_2$, O-C_{1-6}-alkyl, halogen, CN, $SO_2N(C_{1-6}$-alkyl$)_2$ or heteroaryl optionally substituted with one or two residues selected from the group consisting of C_{1-6}-alkyl;
- heteroaryl, optionally substituted with a five- or six-membered carbocyclic non-aromatic ring containing independently from each other two N, O, S, or SO_2, replacing a carbon atom of the ring;
- an aromatic or non-aromatic $C_{9\text{ or }10}$-bicyclic-ring, whereas one or two carbon atoms are replaced by N, O or S each optionally substituted with one or two residues selected from the group consisting of $N(C_{1-6}$-alkyl$)_2$, CONH-C_{1-6}-alkyl, =O;
- a heterocyclic non-aromatic ring, optionally substituted with pyridinyl;
- 4,5-dihydro-naphtho[2,1-d]thiazole, optionally substituted with NHCO-C_{1-6}-alkyl,

$R^{1.2.1}$ H, C_{1-6}-alkyl, C_{1-6}-alkylene-C_{3-6}-cycloalkyl, C_{1-4}-alkylene-phenyl, C_{1-4}-alkylene-furanyl, C_{3-6}-cycloalkyl, C_{1-4}-alkylene-O-C_{1-4}-alkyl, C_{1-6}-haloalkyl or a five- or six-membered carbocyclic non-aromatic ring, optionally containing independently from each other one or two N, O, S, or SO_2, replacing a carbon atom of the ring, optionally substituted with 4-cyclopropylmethyl-piperazinyl

$R^{1.2.2}$ H, C_{1-6}-alkyl;

$R^{1.2.3}$ a five- or six-membered carbocyclic non-aromatic ring, optionally containing independently from each other one or two N, O, S, or SO_2, replacing a carbon atom of the ring;

$R^{1.3}$ is selected from phenyl, heteroaryl or indolyl, each optionally substituted with one or two residues selected from the group consisting of C_{1-6}-alkyl, C_{3-6}-cycloalkyl, O-C_{1-6}-alkyl, O-C_{1-6}-haloalkyl, phenyl, heteroaryl;

R^2 is selected from the group consisting of C_{1-6}-alkylene-phenyl, C_{1-6}-alkylene-naphthyl, and C_{1-6}-alkylene-heteroaryl; each optionally substituted with one, two or three residues selected from the group consisting of C_{1-6}-alkyl, C_{1-6}-haloalkyl, O-C_{1-6}-alkyl, O-C_{1-6}-haloalkyl, halogen;

R^3 is H, C_{1-6}-alkyl;

R^4 is H, C_{1-6}-alkyl;

or R^3 and R^4 together are forming a CH_2-CH_2 group.

2. The use of claim 1 wherein the compound of formula 1,

A is CH_2, O or N-C_{1-4}-alkyl;

R^1 is selected from

- $NHR^{1.1}$, $NMeR^{1.1}$;
- $NHR^{1.2}$, $NMeR^{1.2}$;
- $NHCH_2$-$R^{1.3}$;
- NH-C_{3-6}-cycloalkyl, whereas optionally one carbon atom is replaced by a nitrogen atom, whereas the ring is optionally substituted with one or two residues selected from the group consisting of C_{1-6}-alkyl, O-C_{1-6}-alkyl, $NHSO_2$-phenyl, NHCONH-phenyl, halogen, CN, SO_2-C_{1-6}-alkyl, COO-C_{1-6}-alkyl;

- a $C_{9 \text{ or } 10}$-bicyclic-ring, whereas one or two carbon atoms are replaced by nitrogen atoms and the ring system is bound via a nitrogen atom to the basic structure of formula 1 and whereas the ring system is optionally substituted with one or two residues selected from the group consisting of C_{1-6}-alkyl, COO-C_{1-6}-alkyl, C_{1-6}-haloalkyl, O-C_{1-6}-alkyl, NO_2, halogen, CN, $NHSO_2$-C_{1-6}-alkyl, m-methoxyphenyl;
- a group selected from NHCH(pyridinyl)CH_2COO-C_{1-6}-alkyl, NHCH(CH_2O-C_{1-6}-alkyl)-benzoimidazolyl, optionally substituted with Cl;
- or 1-aminocyclopentyl, optionally substituted with methyl-oxadiazolyl;

$R^{1.1}$ is phenyl, optionally substituted with one or two residues selected from the group consisting of C_{1-6}-alkyl, C_{1-6}-haloalkyl, $CH_2CON(C_{1-6}\text{-alkyl})_2$, $CH_2NHCONH$-C_{3-6}-cycloalkyl, CN, $CONR^{1.1.1}R^{1.1.2}$, COO-C_{1-6}-alkyl, O-C_{1-6}-alkyl, SO_2-C_{1-6}-alkyl, SO_2-C_{1-6}-alkylen-OH, SO_2-C_{3-6}-cycloalkyl, SO_2-piperidinyl, SO_2NH-C_{1-6}-alkyl, $SO_2N(C_{1-6}\text{-alkyl})_2$, halogen, CN, CO-morpholinyl, CH_2-pyridinyl or a heterocyclic ring optionally substituted with one or two residues selected from the group consisting of C_{1-6}-alkyl, NHC_{1-6}-alkyl,, =O;

$R^{1.1.1}$ H, C_{1-6}-alkyl, C_{3-6}-cycloalkyl, C_{1-6}-haloalkyl, $CH_2CON(C_{1-6}\text{-alkyl},)_2$, CH_2CO-azetindinyl, C_{1-6}-alkylen-C_{3-6}-cycloalkyl, CH_2-pyranyl, CH_2-tetrahydrofuranyl, CH_2-furanyl, C_{1-6}-alkylen-OH or thiadiazolyl, optionally substituted with C_{1-6}-alkyl;

$R^{1.1.2}$ H, C_{1-6}-alkyl, SO_2C_{1-6}-alkyl;

or $R^{1.1.1}$ and $R^{1.1.2}$ together with the nitrogen atom of $CONR^{1.1.1}R^{1.1.2}$ are forming a four-, five- or six-membered carbocyclic ring, optionally containing one O, replacing a carbon atom of the ring, optionally substituted with one or two residues selected from the group consisting of CH_2OH

$R^{1.2}$ is selected from

- heteroaryl, optionally substituted with one or two residues selected from the group consisting of C_{1-6}-alkyl, C_{3-6}-cycloalkyl, CH_2COO-C_{1-6}-alkyl, $CONR^{1.2.1}R^{1.2.2}$, COO-C_{1-6}-alkyl, $CONH_2$, O-C_{1-6}-alkyl, halogen, CN, CO-pyrrolidinyl, CO-morpholinyl or heteroaryl optionally substituted with one or two residues selected from the group consisting of C_{1-6}-alkyl;
- benzothiazolyl, indazolyl, dihydro-indolyl, indanyl, tetrahydro-quinolinyl, each optionally substituted with one or two residues selected from the group consisting of $N(C_{1-6}\text{-alkyl})_2$, $CONH$-C_{1-6}-alkyl, =O;
- piperidinyl, optionally substituted with pyridinyl;
- 4,5-dihydro-naphtho[2,1-d]thiazole, optionally substituted with $NHCO$-C_{1-6}-alkyl,

$R^{1.2.1}$ H, C_{1-6}-alkyl;

$R^{1.2.2}$ H, C_{1-6}-alkyl;

$R^{1.3}$ is selected from phenyl, pyrazolyl, isoxazolyl, pyrimidinyl, indolyl or oxadiazolyl, each optionally substituted with one or two residues selected from the group consisting of C_{1-6}-alkyl, C_{3-6}-cycloalkyl, O-C_{1-6}-alkyl, O-C_{1-6}-haloalkyl;

R^2 is selected from CH_2-phenyl or CH_2-naphthyl, both optionally substituted with one or two residues selected from the group consisting of C_{1-6}-alkyl, C_{1-6}-haloalkyl, O-C_{1-6}-alkyl, O-C_{1-6}-haloalkyl, halogen; or CH_2-thiophenyl, optionally substituted with one or two residues selected from the group consisting of halogen;

R^3 is H, C_{1-4}-alkyl;

R^4 is H, C_{1-4}-alkyl;

or R^3 and R^4 together are forming a CH_2-CH_2 group.

3. The use of claim 1 wherein the compound of formula 1 is

A is CH_2, O or NMe;

R^1 is selected from

- $NHR^{1.1}$, $NMeR^{1.1}$;
- $NHR^{1.2}$, $NMeR^{1.2}$;
- $NHCH_2$-$R^{1.3}$;
- NH-cyclohexyl, optionally substituted with one or two residues selected from the group consisting of C_{1-4}-alkyl, $NHSO_2$-phenyl, NHCONH-phenyl, halogen;
- NH-pyrrolidinyl, optionally substituted with one or two residues selected from the group consisting of SO_2-C_{1-4}-alkyl, COO-C_{1-4}-alkyl;
- piperidinyl, optionally substituted with one or two residues selected from the group consisting of $NHSO_2$-C_{1-4}-alkyl, m-methoxyphenyl;
- dihydro-indolyl, dihydro-isoindolyl, tetrahydro-quinolinyl or tetrahydro-isoquinolinyl, optionally substituted with one or two residues selected from the group consisting of C_{1-4}-alkyl, COO-C_{1-4}-alkyl, C_{1-4}-haloalkyl, O-C_{1-4}-alkyl, NO_2, halogen;
- a group selected from NHCH(pyridinyl)CH_2COO-C_{1-4}-alkyl, NHCH(CH_2O-C_{1-4}-alkyl)-benzoimidazolyl, optionally substituted with Cl;
- or 1-aminocyclopentyl, optionally substituted with methyl-oxadiazolyl;

$R^{1.1}$ is phenyl, optionally substituted with one or two residues selected from the group consisting of C_{1-4}-alkyl, C_{1-4}-haloalkyl, $CH_2CON(C_{1-4}\text{-alkyl})_2$,

$CH_2NHCONH$-C_{3-6}-cycloalkyl, CN, $CONR^{1.1.1}R^{1.1.2}$, COO-C_{1-4}-alkyl, O-C_{1-4}-alkyl, SO_2-C_{1-4}-alkyl, SO_2-C_{1-4}-alkylen-OH, SO_2-C_{3-6}-cycloalkyl, SO_2-piperidinyl, SO_2NH-C_{1-4}-alkyl, $SO_2N(C_{1-4}$-alkyl$)_2$, halogen, CO-morpholinyl, CH_2-pyridinyl, or imidazolidinyl, piperidinyl, oxazinanyl, pyrazolyl, triazolyl, tetrazolyl, oxazolyl, oxadiazolyl, thiazolyl, pyridinyl, pyrimidinyl, each optionally substituted with one or two residues selected from the group consisting of C_{1-4}-alkyl, NHC_{1-4}-alkyl,, =O;

$R^{1.1.1}$ H, C_{1-6}-alkyl, C_{3-6}-cycloalkyl, C_{1-4}-haloalkyl, $CH_2CON(C_{1-4}$-alkyl,$)_2$, CH_2CO-azetindinyl, C_{1-4}-alkylen-C_{3-6}-cycloalkyl, CH_2-pyranyl, CH_2-tetrahydrofuranyl, CH_2-furanyl, C_{1-4}-alkylen-OH or thiadiazolyl, optionally substituted with C_{1-4}-alkyl;

$R^{1.1.2}$ H, C_{1-4}-alkyl, SO_2C_{1-4}-alkyl;

or $R^{1.1.1}$ and $R^{1.1.2}$ together with the nitrogen atom of $CONR^{1.1.1}R^{1.1.2}$ are forming a four-, five- or six-membered carbocyclic ring, optionally containing one O, replacing a carbon atom of the ring, optionally substituted with one or two residues selected from the group consisting of CH_2OH

$R^{1.2}$ is selected from

- pyridinyl, pyridazinyl, pyrrolyl, pyrazolyl, isoxazolyl, thiazolyl, thiadiazolyl, optionally substituted with one or two residues selected from the group consisting of C_{1-4}-alkyl, C_{3-6}-cycloalkyl, CH_2COO-C_{1-4}-alkyl, $CONR^{1.2.1}R^{1.2.2}$, COO-C_{1-4}-alkyl, $CONH_2$, O-C_{1-4}-alkyl, halogen, CO-pyrrolidinyl, CO-morpholinyl or pyrazolyl, triazolyl, tetrazolyl, isoxazolyl, oxadiazolyl, each optionally substituted with one or two residues selected from the group consisting of C_{1-4}-alkyl;

- benzothiazolyl, indazolyl, dihydro-indolyl, indanyl, tetrahydro-quinolinyl, each optionally substituted with one or two residues selected from the group consisting of $N(C_{1-4}\text{-alkyl})_2$, $CONH\text{-}C_{1-4}\text{-alkyl}$, =O;
- piperidinyl, optionally substituted with pyridinyl;
- 4,5-dihydro-naphtho[2,1-d]thiazole, optionally substituted with $NHCO\text{-}C_{1-4}\text{-alkyl}$,

$R^{1.2.1}$ H, C_{1-4}-alkyl;

$R^{1.2.2}$ H, C_{1-4}-alkyl;

$R^{1.3}$ is selected from phenyl, pyrazolyl, isoxazolyl, pyrimidinyl, indolyl or oxadiazolyl, each optionally substituted with one or two residues selected from the group consisting of C_{1-4}-alkyl, C_{3-6}-cycloalkyl, $O\text{-}C_{1-4}$-alkyl, $O\text{-}C_{1-4}$-haloalkyl;

R^2 is selected from CH_2-phenyl or CH_2-naphthyl, both optionally substituted with one or two residues selected from the group consisting of C_{1-4}-alkyl, C_{1-4}-haloalkyl, $O\text{-}C_{1-4}$-haloalkyl, halogen; or CH_2-thiophenyl, optionally substituted with one or two residues selected from the group consisting of halogen;

R^3 is H;

R^4 is H;

or R^3 and R^4 together are forming a $CH_2\text{-}CH_2$ group.

4. The use of claim 1 wherein formula 1 is

A is CH_2, O or NMe;

R^1 is selected from

- $NHR^{1.1}$, $NMeR^{1.1}$;
- $NHR^{1.2}$, $NMeR^{1.2}$;
- $NHCH_2$-$R^{1.3}$;
- NH-piperidinyl, optionally substituted with pyridinyl;
- NH-cyclohexyl, optionally substituted with one or two residues selected from the group consisting of t-Bu, $NHSO_2$-phenyl, NHCONH-phenyl, F;
- NH-pyrrolidinyl, optionally substituted with one or two residues selected from the group consisting of SO_2Me, COO-t-Bu;
- piperidinyl, optionally substituted with one or two residues selected from the group consisting of $NHSO_2$-n-Bu, m-methoxyphenyl;
- dihydro-indolyl, dihydro-isoindolyl, tetrahydro-quinolinyl or tetrahydro-isoquinolinyl, optionally substituted with one or two residues selected from the group consisting of Me, COOMe, CF_3, OMe, NO_2, F, Br;
- a group selected from NHCH(pyridinyl)CH_2COOMe, NHCH(CH_2OMe)-benzoimidazolyl, optionally substituted with Cl;
- or 1-aminocyclopentyl, optionally substituted with methyl-oxadiazolyl;

$R^{1.1}$ is phenyl, optionally substituted with one or two residues selected from the group consisting of Me, Et, t-Bu, CF_3, CH_2CONMe_2, $CH_2NHCONH$-cyclohexyl, CN, $CONR^{1.1.1}R^{1.1.2}$, COOMe, COOEt, OMe, SO_2Me, $SO_2CH_2CH_2OH$, SO_2Et, SO_2-cyclopropyl, SO_2-piperidinyl, SO_2NHEt, SO_2NMeEt, F, Cl, CO-morpholinyl, CH_2-pyridinyl, or imidazolidinyl, piperidinyl, oxazinanyl, pyrazolyl, triazolyl, tetrazolyl, oxazolyl, oxadiazolyl, thiazolyl, pyridinyl, pyrimidinyl, each optionally substituted with one or two residues selected from the group consisting of Me, NHMe, =O;

$R^{1.1.1}$ H, Me, Et, t-Bu, i-Pr, cyclopropyl, CH_2-i-Pr, CH_2-t-Bu, $CH(CH_3)CH_2CH_3$, CH_2CHF_2, CH_2CONMe_2, CH_2CO-azetindinyl, CH_2-cyclopropyl, CH_2-cyclobutyl, CH_2-pyranyl, CH_2-tetrahydrofuranyl, CH_2-furanyl, CH_2CH_2OH or thiadiazolyl, optionally substituted with Me;

$R^{1.1.2}$ H, Me, Et, SO_2Me, SO_2Et

or $R^{1.1.1}$ and $R^{1.1.2}$ together with the nitrogen atom of $CONR^{1.1.1}R^{1.1.2}$ are forming a four-, five- or six-membered carbocyclic ring, optionally containing one O, replacing a carbon atom of the ring, optionally substituted with one or two residues selected from the group consisting of CH_2OH

$R^{1.2}$ is selected from

- pyridinyl, pyrrolyl, pyrazolyl, isoxazolyl, thiazolyl, thiadiazolyl, optionally substituted with one or two residues selected from the group consisting of Me, Et, Pr, Bu, cyclopropyl, CH_2COOEt, $CONR^{1.2.1}R^{1.2.2}$, COOMe, COOEt, $CONH_2$, OMe, Cl, Br CO-pyrrolidinyl, CO-morpholinyl or pyrazolyl, triazolyl, tetrazolyl, isoxazolyl, oxadiazolyl, each optionally substituted Me;
- benzothiazolyl, indazolyl, dihydro-indolyl, indanyl, tetrahydro-quinolinyl, each optionally substituted with one or two residues selected from the group consisting of NMe_2, CONHMe, =O;
- 4,5-dihydro-naphtho[2,1-d]thiazole, optionally substituted with NHCOMe,

$R^{1.2.1}$ H, Me;

$R^{1.2.2}$ H, Me;

$R^{1.3}$ is selected from phenyl, pyrazolyl, isoxazolyl, pyrimidinyl, indolyl or oxadiazolyl, each optionally substituted with one or two residues selected from the group consisting of Me, Et, Pr, cyclopentyl, OMe, $OCHF_2$;

R^2 is selected from CH_2-phenyl or CH_2-naphthyl, both optionally substituted with one or two residues selected from the group consisting of CH_3, CF_3, OCF_3, F, Cl, Br, Et; or CH_2-thiophenyl, optionally substituted with one or two residues selected from the group consisting of Cl, Br;

R^3 is H;

R^4 is H;

or R^3 and R^4 together are forming a CH_2-CH_2 group.

5. The use of claim 1 wherein formula 1 is

A is CH_2, O or NMe;

R^1 is selected from

- $NHR^{1.1}$
- $NHR^{1.2}$,

$R^{1.1}$ is phenyl, optionally substituted with one or two residues selected from the group consisting of Me, Et, Bu, CF_3, CH_2CONMe_2, $CH_2NHCONH$-cyclohexyl, CN, $CONR^{1.1.1}R^{1.1.2}$, COOMe, COOEt, OMe, SO_2Me, $SO_2CH_2CH_2OH$, SO_2Et, SO_2-cyclopropyl, SO_2-piperidinyl, SO_2NHEt, SO_2NMeEt, F, Cl, CO-morpholinyl, CH_2-pyridinyl, or imidazolidinyl, piperidinyl, oxazinanyl, pyrazolyl, triazolyl, tetrazolyl, oxazolyl, oxadiazolyl, thiazolyl, pyridinyl, pyrimidinyl, each optionally

substituted with one or two residues selected from the group consisting of Me, NHMe, =O;

$R^{1.1.1}$ H, Me, Et, t-Bu, i-Pr, cyclopropyl, CH_2-i-Pr, CH_2-t-Bu, $CH(CH_3)CH_2CH_3$, CH_2CHF_2, CH_2CONMe_2, CH_2CO-azetindinyl, CH_2-cyclopropyl, CH_2-cyclobutyl, CH_2-pyranyl, CH_2-tetrahydrofuranyl, CH_2-furanyl, CH_2CH_2OH or thiadiazolyl, optionally substituted with Me;

$R^{1.1.2}$ H, Me, Et, SO_2Me, SO_2Et

or $R^{1.1.1}$ and $R^{1.1.2}$ together with the nitrogen atom of $CONR^{1.1.1}R^{1.1.2}$ are forming a four-, five- or six-membered carbocyclic ring, optionally containing one O, replacing a carbon atom of the ring, optionally substituted with one or two residues selected from the group consisting of CH_2OH

$R^{1.2}$ is selected from

- pyridinyl, pyrrolyl, pyrazolyl, isoxazolyl, thiazolyl, thiadiazolyl, optionally substituted with one or two residues selected from the group consisting of Me, Et, Pr, Bu, cyclopropyl, CH_2COOEt, $CONR^{1.2.1}R^{1.2.2}$, COOMe, COOEt, $CONH_2$, OMe, Cl, Br CO-pyrrolidinyl, CO-morpholinyl or pyrazolyl, triazolyl, tetrazolyl, isoxazolyl, oxadiazolyl, each optionally substituted Me;
- benzothiazolyl, indazolyl, dihydro-indolyl, indanyl, tetrahydro-quinolinyl, each optionally substituted with one or two residues selected from the group consisting of NMe_2, CONHMe, =O;
- 4,5-dihydro-naphtho[2,1-d]thiazole, optionally substituted with NHCOMe,

$R^{1.2.1}$ H, Me;

$R^{1.2.2}$ H, Me;

R^2 is selected from CH_2-phenyl or CH_2-naphthyl, both optionally substituted with one or two residues selected from the group consisting of CH_3, CF_3, OCF_3, F, Cl, Br, Et













R^3 is H;


R^4 is H.


6. The use of claim 1 wherein formula 1 is


A is CH_2, O or NMe;

























R^1 is selected from


,

,

,

,

,

,

,

,

,

,

,

,









,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,




























,


,

,

,






















,








,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,


,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,
























,




















,


,

,

,


,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,


,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,







,

,


,

,
















,


,

,

,

,

,

;;


wherein X_1 is:


, wherein R^1 is attached to the vacant single bond;

R^2 is selected from


,

,

,

,

,

,

,

,

,

,

,

,

,

,

;

wherein X_2 is:




, wherein R^2 is attached to the vacant single bond;

R^3 is H;

R^4 is H;

or R^3 and R^4 together are forming a CH_2-CH_2 group.

7. Use of a crystalline salt of a compound of the formula below in the manufacture of a medicament for treating aging-associated neuronal loss, memory impairment, or loss of motor coordination in a subject diagnosed with the aging-associated neuronal loss, memory impairment, or loss of motor coordination, wherein the medicament is to be administered in the subject




*2HCl

.

8. Use of a pharmaceutical composition comprising the compound of claim 7 in the manufacture of a medicament for treating aging-associated neuronal loss, memory impairment, or loss of motor coordination in a subject, wherein the pharmaceutical composition is to be administered in the subject and comprises at least one co-crystal of the compound of the formula according to claim 7 and a pharmaceutically acceptable carrier.

9. The use of claim 1 wherein the compound of formula 1 is to be administered in the form of the individual optical isomers, a mixture of the individual enantiomers, a racemate or in the form of the enantiomerically pure compounds.

10. Use of a pharmaceutical composition comprising a compound of formula 3 in the manufacture of a medicmament for treating aging-associated neuronal loss, memory impairment, or loss of motor coordination in a subject, wherein the compound of formula 3 is to be administered in the subject and has the following formula:



3

wherein

R^1 is H, C_{1-6}-alkyl, C_{0-4}-alkyl-C_{3-6}-cycloalkyl, C_{1-6}-haloalkyl;

R^2 is H, C_{1-6}-alkyl;

X is an anion selected from the group consisting of chloride or ½ dibenzoyltartrate

j is 1 or 2,

and the pharmaceutical composition further comprises a first diluent, a second diluent, a binder, a disintegrant and a lubricant.

11. The use of claim 10 wherein the pharmaceutical composition further comprises an additional disintegrant.

12. The use of claim 10 or 11 wherein the pharmaceutical composition further comprises a glidant.

13. The use of claim 10, 11 or 12 wherein the diluent of the pharmaceutical composition further comprises cellulose powder, dibasic calciumphosphatae anhydrous, dibasic calciumphosphate dehydrate, erythritol, low substituted hydroxypropyl cellulose, mannitol, pregelantinized starch, or xylitol.

14. The use of claim 1, wherein the medicament is for treating aging-associated neuronal loss.

15. The use of claim 1, wherein the medicament is for treating aging-associated memory impairment.

16. The use of claim 1, wherein the medicament is for treating aging-associated motor coordination.