NEISSERIA MENINGITIDIS ANTIGENS

This invention relates to antigens from the bacterium *Neisseria meningitidis*.

BACKGROUND

Neisseria meningitidis is a non-motile, gram negative diplococcus human pathogen. It colonises the pharynx, causing meningitis and, occasionally, septicaemia in the absence of meningitis. It is closely related to *N.gonorrhoeae*, although one feature that clearly differentiates meningococcus from gonococcus is the presence of a polysaccharide capsule that is present in all pathogenic meningococci.

N.meningitidis causes both endemic and epidemic disease. In the United States the attack rate is 0.6-1 per 100,000 persons per year, and it can be much greater during outbreaks (see Lieberman *et al.* (1996) Safety and Immunogenicity of a Serogroups A/C *Neisseria meningitidis* Oligosaccharide-Protein Conjugate Vaccine in Young Children. *JAMA* 275(19):1499-1503; Schuchat *et al* (1997) Bacterial Meningitis in the United States in 1995. *N Engl J Med* 337(14):970-976). In developing countries, endemic disease rates are much higher and during epidemics incidence rates can reach 500 cases per 100,000 persons per year. Mortality is extremely high, at 10-20% in the United States, and much higher in developing countries. Following the introduction of the conjugate vaccine against *Haemophilus influenzae*, *N. meningitidis* is the major cause of bacterial meningitis at all ages in the United States (Schuchat *et al* (1997) *supra*).

Based on the organism's capsular polysaccharide, 12 serogroups of *N.meningitidis* have been identified. Group A is the pathogen most often implicated in epidemic disease in sub-Saharan Africa. Serogroups B and C are responsible for the vast majority of cases in the United States and in most developed countries. Serogroups W135 and Y are responsible for the rest of the cases in the United States and developed countries. The meningococcal vaccine currently in use is a tetravalent polysaccharide vaccine composed of serogroups A, C, Y and W135. Although efficacious in adolescents and adults, it induces a poor immune response and short duration of protection, and cannot be used in infants [*eg.* Morbidity and Mortality weekly report, Vol.46, No. RR-5 (1997)]. This is because polysaccharides are T-cell independent antigens that induce a weak immune response that cannot be boosted by repeated immunization. Following the success of the

vaccination against *H.influenzae*, conjugate vaccines against serogroups A and C have been developed and are at the final stage of clinical testing (Zollinger WD "New and Improved Vaccines Against Meningococcal Disease" in: *New Generation Vaccines, supra*, pp. 469-488; Lieberman *et al* (1996) *supra*; Costantino *et al* (1992) Development and phase I clinical testing of a conjugate vaccine against meningococcus A and C. *Vaccine* 10:691-698).

Meningococcus B remains a problem, however. This serotype currently is responsible for approximately 50% of total meningitis in the United States, Europe, and South America. The polysaccharide approach cannot be used because the menB capsular polysaccharide is a polymer of α(2-8)-linked *N*-acetyl neuraminic acid that is also present in mammalian tissue. This results in tolerance to the antigen; indeed, if an immune response were elicited, it would be anti-self, and therefore undesirable. In order to avoid induction of autoimmunity and to induce a protective immune response, the capsular polysaccharide has, for instance, been chemically modified substituting the *N*-acetyl groups with *N*-propionyl groups, leaving the specific antigenicity unaltered (Romero & Outschoorn (1994) Current status of Meningococcal group B vaccine candidates: capsular or non-capsular? *Clin Microbiol Rev* 7(4):559-575).

Alternative approaches to menB vaccines have used complex mixtures of outer membrane proteins (OMPs), containing either the OMPs alone, or OMPs enriched in porins, or deleted of the class 4 OMPs that are believed to induce antibodies that block bactericidal activity. This approach produces vaccines that are not well characterized. They are able to protect against the homologous strain, but are not effective at large where there are many antigenic variants of the outer membrane proteins. To overcome the antigenic variability, multivalent vaccines containing up to nine different porins have been constructed (*eg.* Poolman JT (1992) Development of a meningococcal vaccine. *Infect. Agents Dis.* 4:13-28). Additional proteins to be used in outer membrane vaccines have been the opa and opc proteins, but none of these approaches have been able to overcome the antigenic variability (*eg.* Ala'Aldeen & Borriello (1996) The meningococcal transferrin-binding proteins 1 and 2 are both surface exposed and generate bactericidal antibodies capable of killing homologous and heterologous strains. *Vaccine* 14(1):49-53).

A certain amount of sequence data is available for meningococcal and gonococcal genes and proteins (*eg.* EP-A-0467714, WO96/29412), but this is by no means complete. The provision of further sequences could provide an opportunity to identify secreted or surface-exposed proteins that

are presumed targets for the immune system and which are not antigenically variable. For instance, some of the identified proteins could be components of efficacious vaccines against meningococcus B, some could be components of vaccines against all meningococcal serotypes, and others could be components of vaccines against all pathogenic *Neisseriae*.

THE INVENTION

The invention provides proteins comprising the *N.meningitidis* amino acid sequences disclosed in the examples.

It also provides proteins comprising sequences homologous (*ie.* having sequence identity) to the *N.meningitidis* amino acid sequences disclosed in the examples. Depending on the particular sequence, the degree of sequence identity is preferably greater than 50% (*eg.* 60%, 70%, 80%, 90%, 95%, 99% or more). These homologous proteins include mutants and allelic variants of the sequences disclosed in the examples. Typically, 50% identity or more between two proteins is considered to be an indication of functional equivalence. Identity between the proteins is preferably determined by the Smith-Waterman homology search algorithm as implemented in the MPSRCH program (Oxford Molecular), using an affine gap search with parameters *gap open penalty=12* and *gap extension penalty=1*.

The invention further provides proteins comprising fragments of the *N.meningitidis* amino acid sequences disclosed in the examples. The fragments should comprise at least *n* consecutive amino acids from the sequences and, depending on the particular sequence, *n* is 7 or more (*eg.* 8, 10, 12, 14, 16, 18, 20 or more). Preferably the fragments comprise an epitope from the sequence.

The proteins of the invention can, of course, be prepared by various means (*eg.* recombinant expression, purification from cell culture, chemical synthesis *etc.*) and in various forms (*eg.* native, fusions *etc.*). They are preferably prepared in substantially pure form (*ie.* substantially free from other *N.meningitidis* or host cell proteins)

According to a further aspect, the invention provides antibodies which bind to these proteins. These may be polyclonal or monoclonal and may be produced by any suitable means.

According to a further aspect, the invention provides nucleic acid comprising the *N.meningitidis* nucleotide sequences disclosed in the examples. In addition, the invention provides nucleic acid comprising sequences homologous (*ie.* having sequence identity) to the *N.meningitidis* nucleotide sequences disclosed in the examples.

Furthermore, the invention provides nucleic acid which can hybridise to the *N.meningitidis* nucleic acid disclosed in the examples, preferably under "high stringency" conditions (*eg.* 65°C in a 0.1xSSC, 0.5% SDS solution).

Nucleic acid comprising fragments of these sequences are also provided. These should comprise at least *n* consecutive nucleotides from the *N.meningitidis* sequences and, depending on the particular sequence, *n* is 10 or more (*eg* 12, 14, 15, 18, 20, 25, 30, 35, 40 or more).

According to a further aspect, the invention provides nucleic acid encoding the proteins and protein fragments of the invention.

It should also be appreciated that the invention provides nucleic acid comprising sequences complementary to those described above (*eg.* for antisense or probing purposes).

Nucleic acid according to the invention can, of course, be prepared in many ways (*eg.* by chemical synthesis, from genomic or cDNA libraries, from the organism itself *etc.*) and can take various forms (*eg.* single stranded, double stranded, vectors, probes *etc.*).

In addition, the term "nucleic acid" includes DNA and RNA, and also their analogues, such as those containing modified backbones, and also peptide nucleic acids (PNA) *etc.*

According to a further aspect, the invention provides vectors comprising nucleotide sequences of the invention (*eg.* expression vectors) and host cells transformed with such vectors.

According to a further aspect, the invention provides compositions comprising protein, antibody, and/or nucleic acid according to the invention. These compositions may be suitable as vaccines, for instance, or as diagnostic reagents, or as immunogenic compositions.

The invention also provides nucleic acid, protein, or antibody according to the invention for use as medicaments (*eg.* as vaccines) or as diagnostic reagents. It also provides the use of nucleic acid, protein, or antibody according to the invention in the manufacture of: (i) a medicament for treating or preventing infection due to Neisserial bacteria; (ii) a diagnostic reagent for detecting the presence of Neisserial bacteria or of antibodies raised against Neisserial bacteria; and/or (iii) a reagent which can raise antibodies against Neisserial bacteria. Said Neisserial bacteria may be any species or strain (such as *N.gonorrhoeae*) but are preferably *N.meningitidis*, especially strain A, strain B or strain C.

The invention also provides a method of treating a patient, comprising administering to the patient a therapeutically effective amount of nucleic acid, protein, and/or antibody according to the invention.

According to further aspects, the invention provides various processes.

A process for producing proteins of the invention is provided, comprising the step of culturing a host cell according to the invention under conditions which induce protein expression.

A process for producing protein or nucleic acid of the invention is provided, wherein the protein or nucleic acid is synthesised in part or in whole using chemical means.

A process for detecting polynucleotides of the invention is provided, comprising the steps of: (a) contacting a nucleic probe according to the invention with a biological sample under hybridizing conditions to form duplexes; and (b) detecting said duplexes.

A process for detecting proteins of the invention is provided, comprising the steps of: (a) contacting an antibody according to the invention with a biological sample under conditions suitable for the formation of an antibody-antigen complexes; and (b) detecting said complexes.

Unlike the sequences disclosed in PCT/IB98/01665, the sequences disclosed in the present application are believed not to have any significant homologs in *N.gonorrhoeae*. Accordingly, the sequences of the present invention also find use in the preparation of reagents for distinguishing between *N.meningitidis* and *N.gonorrhoeae*

A summary of standard techniques and procedures which may be employed in order to perform the invention (*eg.* to utilise the disclosed sequences for vaccination or diagnostic purposes) follows. This summary is not a limitation on the invention but, rather, gives examples that may be used, but are not required.

General

The practice of the present invention will employ, unless otherwise indicated, conventional techniques of molecular biology, microbiology, recombinant DNA, and immunology, which are within the skill of the art. Such techniques are explained fully in the literature *eg.* Sambrook *Molecular Cloning; A Laboratory Manual, Second Edition* (1989); *DNA Cloning, Volumes I and ii* (D.N Glover ed. 1985); *Oligonucleotide Synthesis* (M.J. Gait ed, 1984); *Nucleic Acid Hybridization* (B.D. Hames & S.J. Higgins eds. 1984); *Transcription and Translation* (B.D. Hames & S.J. Higgins eds. 1984); *Animal Cell Culture* (R.I. Freshney ed. 1986); *Immobilized Cells and Enzymes* (IRL Press, 1986); B. Perbal, *A Practical Guide to Molecular Cloning* (1984); the *Methods in Enzymology* series (Academic Press, Inc.), especially volumes 154 & 155; *Gene Transfer Vectors for Mammalian Cells* (J.H. Miller and M.P. Calos eds. 1987, Cold Spring Harbor Laboratory); Mayer and Walker, eds. (1987), *Immunochemical Methods in Cell and Molecular Biology* (Academic Press, London); Scopes, (1987) *Protein Purification: Principles and Practice*, Second Edition (Springer-Verlag, N.Y.), and *Handbook of Experimental Immunology, Volumes I-IV* (D.M. Weir and C. C. Blackwell eds 1986).

Standard abbreviations for nucleotides and amino acids are used in this specification.

All publications, patents, and patent applications cited herein are incorporated in full by reference. In particular, the contents of UK patent applications 9800760.2, 9819015.0 and 9822143.5 are incorporated herein.

Definitions

A composition containing X is "substantially free of" Y when at least 85% by weight of the total X+Y in the composition is X. Preferably, X comprises at least about 90% by weight of the total of X+Y in the composition, more preferably at least about 95% or even 99% by weight.

The term "comprising" means "including" as well as "consisting" *eg.* a composition "comprising" X may consist exclusively of X or may include something additional to X, such as X+Y.

The term "heterologous" refers to two biological components that are not found together in nature. The components may be host cells, genes, or regulatory regions, such as promoters. Although the heterologous components are not found together in nature, they can function together, as when a promoter heterologous to a gene is operably linked to the gene. Another example is where a Neisserial sequence is heterologous to a mouse host cell. A further examples would be two epitopes from the same or different proteins which have been assembled in a single protein in an arrangement not found in nature.

An "origin of replication" is a polynucleotide sequence that initiates and regulates replication of polynucleotides, such as an expression vector. The origin of replication behaves as an autonomous unit of polynucleotide replication within a cell, capable of replication under its own control. An origin of replication may be needed for a vector to replicate in a particular host cell. With certain origins of replication, an expression vector can be reproduced at a high copy number in the presence of the appropriate proteins within the cell. Examples of origins are the autonomously replicating sequences, which are effective in yeast; and the viral T-antigen, effective in COS-7 cells.

A "mutant" sequence is defined as DNA, RNA or amino acid sequence differing from but having sequence identity with the native or disclosed sequence. Depending on the particular sequence, the degree of sequence identity between the native or disclosed sequence and the mutant sequence is preferably greater than 50% (*eg.* 60%, 70%, 80%, 90%, 95%, 99% or more, calculated using the Smith-Waterman algorithm as described above). As used herein, an "allelic variant" of a nucleic acid molecule, or region, for which nucleic acid sequence is provided herein is a nucleic acid molecule, or region, that occurs essentially at the same locus in the genome of another or second isolate, and that, due to natural variation caused by, for example, mutation or recombination, has a similar but not identical nucleic acid sequence. A coding region allelic variant typically encodes a protein having similar activity to that of the protein encoded by the gene to which it is being compared. An allelic variant can also comprise an alteration in the 5' or 3' untranslated regions of the gene, such as in regulatory control regions (*eg.* see US patent 5,753,235).

Expression systems

The Neisserial nucleotide sequences can be expressed in a variety of different expression systems; for example those used with mammalian cells, baculoviruses, plants, bacteria, and yeast.

i. Mammalian Systems

Mammalian expression systems are known in the art. A mammalian promoter is any DNA sequence capable of binding mammalian RNA polymerase and initiating the downstream (3') transcription of a coding sequence (*eg.* structural gene) into mRNA. A promoter will have a transcription initiating region, which is usually placed proximal to the 5' end of the coding sequence, and a TATA box, usually located 25-30 base pairs (bp) upstream of the transcription initiation site. The TATA box is thought to direct RNA polymerase II to begin RNA synthesis at the correct site. A mammalian promoter will also contain an upstream promoter element, usually located within 100 to 200 bp upstream of the TATA box. An upstream promoter element determines the rate at which transcription is initiated and can act in either orientation [Sambrook et al. (1989) "Expression of Cloned Genes in Mammalian Cells." In *Molecular Cloning: A Laboratory Manual, 2nd ed.]*.

Mammalian viral genes are often highly expressed and have a broad host range; therefore sequences encoding mammalian viral genes provide particularly useful promoter sequences. Examples include the SV40 early promoter, mouse mammary tumor virus LTR promoter, adenovirus major late promoter (Ad MLP), and herpes simplex virus promoter. In addition, sequences derived from non-viral genes, such as the murine metallotheionein gene, also provide useful promoter sequences. Expression may be either constitutive or regulated (inducible), depending on the promoter can be induced with glucocorticoid in hormone-responsive cells.

The presence of an enhancer element (enhancer), combined with the promoter elements described above, will usually increase expression levels. An enhancer is a regulatory DNA sequence that can stimulate transcription up to 1000-fold when linked to homologous or heterologous promoters, with synthesis beginning at the normal RNA start site. Enhancers are also active when they are placed upstream or downstream from the transcription initiation site, in either normal or flipped orientation, or at a distance of more than 1000 nucleotides from the promoter [Maniatis et al. (1987) *Science 236:*1237; Alberts et al. (1989) *Molecular Biology of the Cell*, 2nd ed.]. Enhancer elements derived from viruses may be particularly useful, because they usually have a broader host range.

Examples include the SV40 early gene enhancer [Dijkema et al (1985) *EMBO J. 4*:761] and the enhancer/promoters derived from the long terminal repeat (LTR) of the Rous Sarcoma Virus [Gorman et al. (1982b) *Proc. Natl. Acad. Sci 79*:6777] and from human cytomegalovirus [Boshart et al. (1985) *Cell 41*:521]. Additionally, some enhancers are regulatable and become active only in the presence of an inducer, such as a hormone or metal ion [Sassone-Corsi and Borelli (1986) *Trends Genet. 2*:215; Maniatis et al. (1987) *Science* 236:1237].

A DNA molecule may be expressed intracellularly in mammalian cells. A promoter sequence may be directly linked with the DNA molecule, in which case the first amino acid at the N-terminus of the recombinant protein will always be a methionine, which is encoded by the ATG start codon. If desired, the N-terminus may be cleaved from the protein by *in vitro* incubation with cyanogen bromide.

Alternatively, foreign proteins can also be secreted from the cell into the growth media by creating chimeric DNA molecules that encode a fusion protein comprised of a leader sequence fragment that provides for secretion of the foreign protein in mammalian cells. Preferably, there are processing sites encoded between the leader fragment and the foreign gene that can be cleaved either *in vivo* or *in vitro*. The leader sequence fragment usually encodes a signal peptide comprised of hydrophobic amino acids which direct the secretion of the protein from the cell. The adenovirus triparite leader is an example of a leader sequence that provides for secretion of a foreign protein in mammalian cells.

Usually, transcription termination and polyadenylation sequences recognized by mammalian cells are regulatory regions located 3' to the translation stop codon and thus, together with the promoter elements, flank the coding sequence. The 3' terminus of the mature mRNA is formed by site-specific post-transcriptional cleavage and polyadenylation [Birnstiel et al. (1985) *Cell 41*:349; Proudfoot and Whitelaw (1988) "Termination and 3' end processing of eukaryotic RNA. In *Transcription and splicing* (ed. B.D. Hames and D.M. Glover); Proudfoot (1989) *Trends Biochem. Sci. 14*:105]. These sequences direct the transcription of an mRNA which can be translated into the polypeptide encoded by the DNA. Examples of transcription terminater/polyadenylation signals include those derived from SV40 [Sambrook et al (1989) "Expression of cloned genes in cultured mammalian cells." In *Molecular Cloning: A Laboratory Manual*].

Usually, the above described components, comprising a promoter, polyadenylation signal, and transcription termination sequence are put together into expression constructs. Enhancers, introns with functional splice donor and acceptor sites, and leader sequences may also be included in an expression construct, if desired. Expression constructs are often maintained in a replicon, such as an extrachromosomal element (*eg.* plasmids) capable of stable maintenance in a host, such as mammalian cells or bacteria. Mammalian replication systems include those derived from animal viruses, which require trans-acting factors to replicate. For example, plasmids containing the replication systems of papovaviruses, such as SV40 [Gluzman (1981) *Cell 23*:175] or polyomavirus, replicate to extremely high copy number in the presence of the appropriate viral T antigen. Additional examples of mammalian replicons include those derived from bovine papillomavirus and Epstein-Barr virus. Additionally, the replicon may have two replicaton systems, thus allowing it to be maintained, for example, in mammalian cells for expression and in a prokaryotic host for cloning and amplification. Examples of such mammalian-bacteria shuttle vectors include pMT2 [Kaufman et al. (1989) *Mol. Cell. Biol. 9*:946] and pHEBO [Shimizu et al. (1986) *Mol. Cell. Biol. 6*:1074].

The transformation procedure used depends upon the host to be transformed. Methods for introduction of heterologous polynucleotides into mammalian cells are known in the art and include dextran-mediated transfection, calcium phosphate precipitation, polybrene mediated transfection, protoplast fusion, electroporation, encapsulation of the polynucleotide(s) in liposomes, and direct microinjection of the DNA into nuclei.

Mammalian cell lines available as hosts for expression are known in the art and include many immortalized cell lines available from the American Type Culture Collection (ATCC), including but not limited to, Chinese hamster ovary (CHO) cells, HeLa cells, baby hamster kidney (BHK) cells, monkey kidney cells (COS), human hepatocellular carcinoma cells (*eg.* Hep G2), and a number of other cell lines.

ii. Baculovirus Systems

The polynucleotide encoding the protein can also be inserted into a suitable insect expression vector, and is operably linked to the control elements within that vector. Vector construction employs techniques which are known in the art. Generally, the components of the expression system include a transfer vector, usually a bacterial plasmid, which contains both a fragment of the baculovirus

genome, and a convenient restriction site for insertion of the heterologous gene or genes to be expressed; a wild type baculovirus with a sequence homologous to the baculovirus-specific fragment in the transfer vector (this allows for the homologous recombination of the heterologous gene in to the baculovirus genome); and appropriate insect host cells and growth media.

After inserting the DNA sequence encoding the protein into the transfer vector, the vector and the wild type viral genome are transfected into an insect host cell where the vector and viral genome are allowed to recombine. The packaged recombinant virus is expressed and recombinant plaques are identified and purified. Materials and methods for baculovirus/insect cell expression systems are commercially available in kit form from, *inter alia*, Invitrogen, San Diego CA ("MaxBac" kit). These techniques are generally known to those skilled in the art and fully described in Summers and Smith, *Texas Agricultural Experiment Station Bulletin No. 1555* (1987) (hereinafter "Summers and Smith").

Prior to inserting the DNA sequence encoding the protein into the baculovirus genome, the above described components, comprising a promoter, leader (if desired), coding sequence of interest, and transcription termination sequence, are usually assembled into an intermediate transplacement construct (transfer vector). This construct may contain a single gene and operably linked regulatory elements; multiple genes, each with its owned set of operably linked regulatory elements; or multiple genes, regulated by the same set of regulatory elements. Intermediate transplacement constructs are often maintained in a replicon, such as an extrachromosomal element (*eg.* plasmids) capable of stable maintenance in a host, such as a bacterium. The replicon will have a replication system, thus allowing it to be maintained in a suitable host for cloning and amplification.

Currently, the most commonly used transfer vector for introducing foreign genes into AcNPV is pAc373. Many other vectors, known to those of skill in the art, have also been designed. These include, for example, pVL985 (which alters the polyhedrin start codon from ATG to ATT, and which introduces a BamHI cloning site 32 basepairs downstream from the ATT; see Luckow and Summers, *Virology* (1989) *17*:31.

The plasmid usually also contains the polyhedrin polyadenylation signal (Miller et al. (1988) *Ann. Rev. Microbiol.*, *42*:177) and a prokaryotic ampicillin-resistance (*amp*) gene and origin of replication for selection and propagation in *E. coli*.

Baculovirus transfer vectors usually contain a baculovirus promoter. A baculovirus promoter is any DNA sequence capable of binding a baculovirus RNA polymerase and initiating the downstream (5' to 3') transcription of a coding sequence (*eg.* structural gene) into mRNA. A promoter will have a transcription initiation region which is usually placed proximal to the 5' end of the coding sequence. This transcription initiation region usually includes an RNA polymerase binding site and a transcription initiation site. A baculovirus transfer vector may also have a second domain called an enhancer, which, if present, is usually distal to the structural gene. Expression may be either regulated or constitutive.

Structural genes, abundantly transcribed at late times in a viral infection cycle, provide particularly useful promoter sequences. Examples include sequences derived from the gene encoding the viral polyhedron protein, Friesen et al., (1986) "The Regulation of Baculovirus Gene Expression," in: *The Molecular Biology of Baculoviruses* (ed. Walter Doerfler); EPO Publ. Nos. 127 839 and 155 476; and the gene encoding the p10 protein, Vlak et al., (1988), *J. Gen. Virol. 69*:765.

DNA encoding suitable signal sequences can be derived from genes for secreted insect or baculovirus proteins, such as the baculovirus polyhedrin gene (Carbonell et al. (1988) *Gene, 73*:409). Alternatively, since the signals for mammalian cell posttranslational modifications (such as signal peptide cleavage, proteolytic cleavage, and phosphorylation) appear to be recognized by insect cells, and the signals required for secretion and nuclear accumulation also appear to be conserved between the invertebrate cells and vertebrate cells, leaders of non-insect origin, such as those derived from genes encoding human α-interferon, Maeda et al., (1985), *Nature 315*:592; human gastrin-releasing peptide, Lebacq-Verheyden et al., (1988), *Molec. Cell. Biol. 8*:3129; human IL-2, Smith et al., (1985) *Proc. Nat'l Acad. Sci. USA, 82*:8404; mouse IL-3, (Miyajima et al., (1987) *Gene 58*:273; and human glucocerebrosidase, Martin et al. (1988) *DNA, 7*:99, can also be used to provide for secretion in insects.

A recombinant polypeptide or polyprotein may be expressed intracellularly or, if it is expressed with the proper regulatory sequences, it can be secreted. Good intracellular expression of nonfused foreign proteins usually requires heterologous genes that ideally have a short leader sequence containing suitable translation initiation signals preceding an ATG start signal. If desired, methionine at the N-terminus may be cleaved from the mature protein by *in vitro* incubation with cyanogen bromide.

Alternatively, recombinant polyproteins or proteins which are not naturally secreted can be secreted from the insect cell by creating chimeric DNA molecules that encode a fusion protein comprised of a leader sequence fragment that provides for secretion of the foreign protein in insects. The leader sequence fragment usually encodes a signal peptide comprised of hydrophobic amino acids which direct the translocation of the protein into the endoplasmic reticulum.

After insertion of the DNA sequence and/or the gene encoding the expression product precursor of the protein, an insect cell host is co-transformed with the heterologous DNA of the transfer vector and the genomic DNA of wild type baculovirus -- usually by co-transfection. The promoter and transcription termination sequence of the construct will usually comprise a 2-5kb section of the baculovirus genome. Methods for introducing heterologous DNA into the desired site in the baculovirus virus are known in the art. (See Summers and Smith *supra*; Ju et al. (1987); Smith et al., *Mol. Cell. Biol.* (1983) *3*:2156; and Luckow and Summers (1989)). For example, the insertion can be into a gene such as the polyhedrin gene, by homologous double crossover recombination; insertion can also be into a restriction enzyme site engineered into the desired baculovirus gene. Miller et al., (1989), *Bioessays 4*:91.The DNA sequence, when cloned in place of the polyhedrin gene in the expression vector, is flanked both 5' and 3' by polyhedrin-specific sequences and is positioned downstream of the polyhedrin promoter.

The newly formed baculovirus expression vector is subsequently packaged into an infectious recombinant baculovirus. Homologous recombination occurs at low frequency (between about 1% and about 5%); thus, the majority of the virus produced after cotransfection is still wild-type virus. Therefore, a method is necessary to identify recombinant viruses. An advantage of the expression system is a visual screen allowing recombinant viruses to be distinguished. The polyhedrin protein, which is produced by the native virus, is produced at very high levels in the nuclei of infected cells at late times after viral infection. Accumulated polyhedrin protein forms occlusion bodies that also contain embedded particles. These occlusion bodies, up to 15 μm in size, are highly refractile, giving them a bright shiny appearance that is readily visualized under the light microscope. Cells infected with recombinant viruses lack occlusion bodies. To distinguish recombinant virus from wild-type virus, the transfection supernatant is plaqued onto a monolayer of insect cells by techniques known to those skilled in the art. Namely, the plaques are screened under the light microscope for the presence (indicative of wild-type virus) or absence (indicative of recombinant

virus) of occlusion bodies. "Current Protocols in Microbiology" Vol. 2 (Ausubel et al. eds) at 16.8 (Supp. 10, 1990); Summers and Smith, *supra*; Miller et al. (1989).

Recombinant baculovirus expression vectors have been developed for infection into several insect cells. For example, recombinant baculoviruses have been developed for, *inter alia*: *Aedes aegypti*, *Autographa californica*, *Bombyx mori*, *Drosophila melanogaster*, *Spodoptera frugiperda*, and *Trichoplusia ni* (WO 89/046699; Carbonell et al., (1985) *J. Virol. 56*:153; Wright (1986) *Nature 321*:718; Smith et al., (1983) *Mol. Cell. Biol. 3*:2156; and see generally, Fraser, *et al.* (1989) *In Vitro Cell. Dev. Biol. 25*:225).

Cells and cell culture media are commercially available for both direct and fusion expression of heterologous polypeptides in a baculovirus/expression system; cell culture technology is generally known to those skilled in the art. *See, eg.* Summers and Smith *supra*.

The modified insect cells may then be grown in an appropriate nutrient medium, which allows for stable maintenance of the plasmid(s) present in the modified insect host. Where the expression product gene is under inducible control, the host may be grown to high density, and expression induced. Alternatively, where expression is constitutive, the product will be continuously expressed into the medium and the nutrient medium must be continuously circulated, while removing the product of interest and augmenting depleted nutrients. The product may be purified by such techniques as chromatography, eg. HPLC, affinity chromatography, ion exchange chromatography, etc.; electrophoresis; density gradient centrifugation; solvent extraction, or the like. As appropriate, the product may be further purified, as required, so as to remove substantially any insect proteins which are also secreted in the medium or result from lysis of insect cells, so as to provide a product which is at least substantially free of host debris, eg. proteins, lipids and polysaccharides.

In order to obtain protein expression, recombinant host cells derived from the transformants are incubated under conditions which allow expression of the recombinant protein encoding sequence. These conditions will vary, dependent upon the host cell selected. However, the conditions are readily ascertainable to those of ordinary skill in the art, based upon what is known in the art.

iii. Plant Systems

There are many plant cell culture and whole plant genetic expression systems known in the art. Exemplary plant cellular genetic expression systems include those described in patents, such as:

US 5,693,506; US 5,659,122; and US 5,608,143. Additional examples of genetic expression in plant cell culture has been described by Zenk, *Phytochemistry* 30:3861-3863 (1991). Descriptions of plant protein signal peptides may be found in addition to the references described above in Vaulcombe et al., *Mol. Gen Genet.* 209:33-40 (1987); Chandler et al., *Plant Molecular Biology* 3:407-418 (1984); Rogers, *J. Biol Chem.* 260:3731-3738 (1985); Rothstein et al., *Gene* 55:353-356 (1987); Whittier et al., Nucleic Acids Research 15:2515-2535 (1987); Wirsel et al., *Molecular Microbiology* 3:3-14 (1989); Yu et al., *Gene* 122:247-253 (1992). A description of the regulation of plant gene expression by the phytohormone, gibberellic acid and secreted enzymes induced by gibberellic acid can be found in R.L. Jones and J. MacMillin, Gibberellins: in: *Advanced Plant Physiology*,. Malcolm B. Wilkins, ed., 1984 Pitman Publishing Limited, London, pp. 21-52. References that describe other metabolically-regulated genes: Sheen, *Plant Cell*, 2:1027-1038(1990); Maas et al., *EMBO J.* 9:3447-3452 (1990); Benkel and Hickey, *Proc. Natl. Acad. Sci.* 84:1337-1339 (1987)

Typically, using techniques known in the art, a desired polynucleotide sequence is inserted into an expression cassette comprising genetic regulatory elements designed for operation in plants. The expression cassette is inserted into a desired expression vector with companion sequences upstream and downstream from the expression cassette suitable for expression in a plant host. The companion sequences will be of plasmid or viral origin and provide necessary characteristics to the vector to permit the vectors to move DNA from an original cloning host, such as bacteria, to the desired plant host. The basic bacterial/plant vector construct will preferably provide a broad host range prokaryote replication origin; a prokaryote selectable marker; and, for Agrobacterium transformations, T DNA sequences for Agrobacterium-mediated transfer to plant chromosomes. Where the heterologous gene is not readily amenable to detection, the construct will preferably also have a selectable marker gene suitable for determining if a plant cell has been transformed. A general review of suitable markers, for example for the members of the grass family, is found in Wilmink and Dons, 1993, *Plant Mol. Biol. Reptr*, 11(2):165-185.

Sequences suitable for permitting integration of the heterologous sequence into the plant genome are also recommended. These might include transposon sequences and the like for homologous recombination as well as Ti sequences which permit random insertion of a heterologous expression cassette into a plant genome. Suitable prokaryote selectable markers include resistance toward

antibiotics such as ampicillin or tetracycline. Other DNA sequences encoding additional functions may also be present in the vector, as is known in the art.

The nucleic acid molecules of the subject invention may be included into an expression cassette for expression of the protein(s) of interest. Usually, there will be only one expression cassette, although two or more are feasible. The recombinant expression cassette will contain in addition to the heterologous protein encoding sequence the following elements, a promoter region, plant 5' untranslated sequences, initiation codon depending upon whether or not the structural gene comes equipped with one, and a transcription and translation termination sequence. Unique restriction enzyme sites at the 5' and 3' ends of the cassette allow for easy insertion into a pre-existing vector.

A heterologous coding sequence may be for any protein relating to the present invention. The sequence encoding the protein of interest will encode a signal peptide which allows processing and translocation of the protein, as appropriate, and will usually lack any sequence which might result in the binding of the desired protein of the invention to a membrane. Since, for the most part, the transcriptional initiation region will be for a gene which is expressed and translocated during germination, by employing the signal peptide which provides for translocation, one may also provide for translocation of the protein of interest. In this way, the protein(s) of interest will be translocated from the cells in which they are expressed and may be efficiently harvested. Typically secretion in seeds are across the aleurone or scutellar epithelium layer into the endosperm of the seed. While it is not required that the protein be secreted from the cells in which the protein is produced, this facilitates the isolation and purification of the recombinant protein.

Since the ultimate expression of the desired gene product will be in a eucaryotic cell it is desirable to determine whether any portion of the cloned gene contains sequences which will be processed out as introns by the host's splicosome machinery. If so, site-directed mutagenesis of the "intron" region may be conducted to prevent losing a portion of the genetic message as a false intron code, Reed and Maniatis, *Cell* 41:95-105, 1985.

The vector can be microinjected directly into plant cells by use of micropipettes to mechanically transfer the recombinant DNA. Crossway, *Mol. Gen. Genet*, 202:179-185, 1985. The genetic material may also be transferred into the plant cell by using polyethylene glycol, Krens, et al., *Nature*, 296, 72-74, 1982. Another method of introduction of nucleic acid segments is high

velocity ballistic penetration by small particles with the nucleic acid either within the matrix of small beads or particles, or on the surface, Klein, et al., *Nature*, 327, 70-73, 1987 and Knudsen and Muller, 1991, *Planta*, 185:330-336 teaching particle bombardment of barley endosperm to create transgenic barley. Yet another method of introduction would be fusion of protoplasts with other entities, either minicells, cells, lysosomes or other fusible lipid-surfaced bodies, Fraley, et al., *Proc. Natl Acad. Sci. USA*, 79, 1859-1863, 1982.

The vector may also be introduced into the plant cells by electroporation. (Fromm et al., *Proc. Natl Acad. Sci. USA* 82:5824, 1985). In this technique, plant protoplasts are electroporated in the presence of plasmids containing the gene construct. Electrical impulses of high field strength reversibly permeabilize biomembranes allowing the introduction of the plasmids. Electroporated plant protoplasts reform the cell wall, divide, and form plant callus.

All plants from which protoplasts can be isolated and cultured to give whole regenerated plants can be transformed by the present invention so that whole plants are recovered which contain the transferred gene. It is known that practically all plants can be regenerated from cultured cells or tissues, including but not limited to all major species of sugarcane, sugar beet, cotton, fruit and other trees, legumes and vegetables. Some suitable plants include, for example, species from the genera *Fragaria*, *Lotus*, *Medicago*, *Onobrychis*, *Trifolium*, *Trigonella*, *Vigna*, *Citrus*, *Linum*, *Geranium*, *Manihot*, *Daucus*, *Arabidopsis*, *Brassica*, *Raphanus*, *Sinapis*, *Atropa*, *Capsicum*, *Datura*, *Hyoscyamus*, *Lycopersion*, *Nicotiana*, *Solanum*, *Petunia*, *Digitalis*, *Majorana*, *Cichorium*, *Helianthus*, *Lactuca*, *Bromus*, *Asparagus*, *Antirrhinum*, *Hererocallis*, *Nemesia*, *Pelargonium*, *Panicum*, *Pennisetum*, *Ranunculus*, *Senecio*, *Salpiglossis*, *Cucumis*, *Browaalia*, *Glycine*, *Lolium*, *Zea*, *Triticum*, *Sorghum*, and *Datura*.

Means for regeneration vary from species to species of plants, but generally a suspension of transformed protoplasts containing copies of the heterologous gene is first provided. Callus tissue is formed and shoots may be induced from callus and subsequently rooted. Alternatively, embryo formation can be induced from the protoplast suspension. These embryos germinate as natural embryos to form plants. The culture media will generally contain various amino acids and hormones, such as auxin and cytokinins. It is also advantageous to add glutamic acid and proline to the medium, especially for such species as corn and alfalfa. Shoots and roots normally develop simultaneously. Efficient regeneration will depend on the medium, on the genotype, and on the

history of the culture. If these three variables are controlled, then regeneration is fully reproducible and repeatable.

In some plant cell culture systems, the desired protein of the invention may be excreted or alternatively, the protein may be extracted from the whole plant. Where the desired protein of the invention is secreted into the medium, it may be collected. Alternatively, the embryos and embryoless-half seeds or other plant tissue may be mechanically disrupted to release any secreted protein between cells and tissues. The mixture may be suspended in a buffer solution to retrieve soluble proteins. Conventional protein isolation and purification methods will be then used to purify the recombinant protein. Parameters of time, temperature pH, oxygen, and volumes will be adjusted through routine methods to optimize expression and recovery of heterologous protein.

iv. Bacterial Systems

Bacterial expression techniques are known in the art. A bacterial promoter is any DNA sequence capable of binding bacterial RNA polymerase and initiating the downstream (3') transcription of a coding sequence (*eg.* structural gene) into mRNA. A promoter will have a transcription initiation region which is usually placed proximal to the 5' end of the coding sequence. This transcription initiation region usually includes an RNA polymerase binding site and a transcription initiation site. A bacterial promoter may also have a second domain called an operator, that may overlap an adjacent RNA polymerase binding site at which RNA synthesis begins. The operator permits negative regulated (inducible) transcription, as a gene repressor protein may bind the operator and thereby inhibit transcription of a specific gene. Constitutive expression may occur in the absence of negative regulatory elements, such as the operator. In addition, positive regulation may be achieved by a gene activator protein binding sequence, which, if present is usually proximal (5') to the RNA polymerase binding sequence. An example of a gene activator protein is the catabolite activator protein (CAP), which helps initiate transcription of the lac operon in Escherichia coli (E. coli) [Raibaud *et al.* (1984) *Annu. Rev. Genet. 18*:173]. Regulated expression may therefore be either positive or negative, thereby either enhancing or reducing transcription.

Sequences encoding metabolic pathway enzymes provide particularly useful promoter sequences. Examples include promoter sequences derived from sugar metabolizing enzymes, such as galactose, lactose (*lac*) [Chang *et al.* (1977) *Nature 198*:1056], and maltose. Additional examples include promoter sequences derived from biosynthetic enzymes such as tryptophan (*trp*) [Goeddel *et al.*

(1980) *Nuc Acids Res. 8*:4057; Yelverton *et al.* (1981) *Nucl. Acids Res. 9*:731; US patent 4,738,921; EP-A-0036776 and EP-A-0121775]. The g-laotamase (*bla*) promoter system [Weissmann (1981) "The cloning of interferon and other mistakes." In *Interferon 3* (ed. I. Gresser)], bacteriophage lambda PL [Shimatake *et al.* (1981) *Nature 292*:128] and T5 [US patent 4,689,406] promoter systems also provide useful promoter sequences.

In addition, synthetic promoters which do not occur in nature also function as bacterial promoters. For example, transcription activation sequences of one bacterial or bacteriophage promoter may be joined with the operon sequences of another bacterial or bacteriophage promoter, creating a synthetic hybrid promoter [US patent 4,551,433]. For example, the *tac* promoter is a hybrid *trp-lac* promoter comprised of both *trp* promoter and *lac* operon sequences that is regulated by the *lac* repressor [Amann *et al.* (1983) *Gene 25*:167; de Boer *et al.* (1983) *Proc. Natl. Acad. Sci. 80*:21]. Furthermore, a bacterial promoter can include naturally occurring promoters of non-bacterial origin that have the ability to bind bacterial RNA polymerase and initiate transcription. A naturally occurring promoter of non-bacterial origin can also be coupled with a compatible RNA polymerase to produce high levels of expression of some genes in prokaryotes. The bacteriophage T7 RNA polymerase/promoter system is an example of a coupled promoter system [Studier *et al.* (1986) *J. Mol. Biol. 189*:113; Tabor *et al.* (1985) *Proc Natl. Acad. Sci. 82*:1074]. In addition, a hybrid promoter can also be comprised of a bacteriophage promoter and an *E. coli* operator region (EPO-A-0 267 851).

In addition to a functioning promoter sequence, an efficient ribosome binding site is also useful for the expression of foreign genes in prokaryotes. In *E. coli*, the ribosome binding site is called the Shine-Dalgarno (SD) sequence and includes an initiation codon (ATG) and a sequence 3-9 nucleotides in length located 3-11 nucleotides upstream of the initiation codon [Shine *et al.* (1975) *Nature 254*:34]. The SD sequence is thought to promote binding of mRNA to the ribosome by the pairing of bases between the SD sequence and the 3' and of *E. coli* 16S rRNA [Steitz *et al.* (1979) "Genetic signals and nucleotide sequences in messenger RNA." In *Biological Regulation and Development: Gene Expression* (ed. R.F. Goldberger)]. To express eukaryotic genes and prokaryotic genes with weak ribosome-binding site [Sambrook *et al.* (1989) "Expression of cloned genes in Escherichia coli." In *Molecular Cloning: A Laboratory Manual*].

A DNA molecule may be expressed intracellularly. A promoter sequence may be directly linked with the DNA molecule, in which case the first amino acid at the N-terminus will always be a methionine, which is encoded by the ATG start codon. If desired, methionine at the N-terminus may be cleaved from the protein by *in vitro* incubation with cyanogen bromide or by either *in vivo* on *in vitro* incubation with a bacterial methionine N-terminal peptidase (EPO-A-0 219 237).

Fusion proteins provide an alternative to direct expression. Usually, a DNA sequence encoding the N-terminal portion of an endogenous bacterial protein, or other stable protein, is fused to the 5' end of heterologous coding sequences. Upon expression, this construct will provide a fusion of the two amino acid sequences. For example, the bacteriophage lambda cell gene can be linked at the 5' terminus of a foreign gene and expressed in bacteria. The resulting fusion protein preferably retains a site for a processing enzyme (factor Xa) to cleave the bacteriophage protein from the foreign gene [Nagai *et al.* (1984) *Nature 309*:810]. Fusion proteins can also be made with sequences from the *lacZ* [Jia *et al.* (1987) *Gene 60*:197], *trpE* [Allen et al. (1987) J. Biotechnol. 5:93; Makoff *et al.* (1989) *J. Gen. Microbiol. 135*:11], and *Chey* [EP-A-0 324 647] genes. The DNA sequence at the junction of the two amino acid sequences may or may not encode a cleavable site. Another example is a ubiquitin fusion protein. Such a fusion protein is made with the ubiquitin region that preferably retains a site for a processing enzyme (*eg.* ubiquitin specific processing-protease) to cleave the ubiquitin from the foreign protein. Through this method, native foreign protein can be isolated [Miller *et al.* (1989) *Bio/Technology 7*:698].

Alternatively, foreign proteins can also be secreted from the cell by creating chimeric DNA molecules that encode a fusion protein comprised of a signal peptide sequence fragment that provides for secretion of the foreign protein in bacteria [US patent 4,336,336]. The signal sequence fragment usually encodes a signal peptide comprised of hydrophobic amino acids which direct the secretion of the protein from the cell. The protein is either secreted into the growth media (gram-positive bacteria) or into the periplasmic space, located between the inner and outer membrane of the cell (gram-negative bacteria). Preferably there are processing sites, which can be cleaved either *in vivo* or *in vitro* encoded between the signal peptide fragment and the foreign gene.

DNA encoding suitable signal sequences can be derived from genes for secreted bacterial proteins, such as the *E. coli* outer membrane protein gene (*ompA*) [Masui *et al.* (1983), in: *Experimental Manipulation of Gene Expression*; Ghrayeb *et al.* (1984) *EMBO J. 3*:2437] and the *E. coli* alkaline

phosphatase signal sequence (*phoA*) [Oka *et al.* (1985) *Proc. Natl. Acad. Sci. 82*:7212]. As an additional example, the signal sequence of the alpha-amylase gene from various Bacillus strains can be used to secrete heterologous proteins from *B. subtilis* [Palva *et al.* (1982) *Proc. Natl. Acad. Sci. USA 79*:5582; EP-A-0 244 042].

Usually, transcription termination sequences recognized by bacteria are regulatory regions located 3' to the translation stop codon, and thus together with the promoter flank the coding sequence. These sequences direct the transcription of an mRNA which can be translated into the polypeptide encoded by the DNA. Transcription termination sequences frequently include DNA sequences of about 50 nucleotides capable of forming stem loop structures that aid in terminating transcription. Examples include transcription termination sequences derived from genes with strong promoters, such as the *trp* gene in *E. coli* as well as other biosynthetic genes.

Usually, the above described components, comprising a promoter, signal sequence (if desired), coding sequence of interest, and transcription termination sequence, are put together into expression constructs. Expression constructs are often maintained in a replicon, such as an extrachromosomal element (*eg.* plasmids) capable of stable maintenance in a host, such as bacteria. The replicon will have a replication system, thus allowing it to be maintained in a prokaryotic host either for expression or for cloning and amplification. In addition, a replicon may be either a high or low copy number plasmid. A high copy number plasmid will generally have a copy number ranging from about 5 to about 200, and usually about 10 to about 150. A host containing a high copy number plasmid will preferably contain at least about 10, and more preferably at least about 20 plasmids. Either a high or low copy number vector may be selected, depending upon the effect of the vector and the foreign protein on the host.

Alternatively, the expression constructs can be integrated into the bacterial genome with an integrating vector. Integrating vectors usually contain at least one sequence homologous to the bacterial chromosome that allows the vector to integrate. Integrations appear to result from recombinations between homologous DNA in the vector and the bacterial chromosome. For example, integrating vectors constructed with DNA from various Bacillus strains integrate into the Bacillus chromosome (EP-A- 0 127 328). Integrating vectors may also be comprised of bacteriophage or transposon sequences.

Usually, extrachromosomal and integrating expression constructs may contain selectable markers to allow for the selection of bacterial strains that have been transformed. Selectable markers can be expressed in the bacterial host and may include genes which render bacteria resistant to drugs such as ampicillin, chloramphenicol, erythromycin, kanamycin (neomycin), and tetracycline [Davies *et al.* (1978) *Annu. Rev Microbiol. 32*:469]. Selectable markers may also include biosynthetic genes, such as those in the histidine, tryptophan, and leucine biosynthetic pathways.

Alternatively, some of the above described components can be put together in transformation vectors. Transformation vectors are usually comprised of a selectable market that is either maintained in a replicon or developed into an integrating vector, as described above.

Expression and transformation vectors, either extra-chromosomal replicons or integrating vectors, have been developed for transformation into many bacteria. For example, expression vectors have been developed for, *inter alia*, the following bacteria: Bacillus subtilis [Palva *et al.* (1982) *Proc. Natl. Acad. Sci. USA 79*:5582; EP-A-0 036 259 and EP-A-0 063 953; WO 84/04541], Escherichia coli [Shimatake *et al.* (1981) *Nature 292*:128; Amann *et al.* (1985) *Gene 40*:183; Studier *et al.* (1986) *J. Mol. Biol. 189*:113; EP-A-0 036 776,EP-A-0 136 829 and EP-A-0 136 907], Streptococcus cremoris [Powell *et al.* (1988) *Appl. Environ. Microbiol. 54*:655]; Streptococcus lividans [Powell *et al.* (1988) *Appl. Environ. Microbiol. 54*:655], Streptomyces lividans [US patent 4,745,056].

Methods of introducing exogenous DNA into bacterial hosts are well-known in the art, and usually include either the transformation of bacteria treated with $CaCl_2$ or other agents, such as divalent cations and DMSO. DNA can also be introduced into bacterial cells by electroporation. Transformation procedures usually vary with the bacterial species to be transformed. See *eg.* [Masson *et al.* (1989) *FEMS Microbiol. Lett. 60*:273; Palva *et al.* (1982) *Proc. Natl. Acad. Sci. USA 79*:5582; EP-A-0 036 259 and EP-A-0 063 953; WO 84/04541, Bacillus], [Miller *et al.* (1988) *Proc. Natl. Acad. Sci. 85*:856; Wang *et al.* (1990) *J. Bacteriol 172*:949, Campylobacter], [Cohen *et al.* (1973) *Proc. Natl. Acad. Sci. 69*:2110; Dower *et al.* (1988) *Nucleic Acids Res. 16*:6127; Kushner (1978) "An improved method for transformation of Escherichia coli with ColE1-derived plasmids. In *Genetic Engineering: Proceedings of the International Symposium on Genetic Engineering* (eds. H.W. Boyer and S. Nicosia); Mandel *et al.* (1970) *J. Mol. Biol. 53*:159; Taketo (1988) *Biochim. Biophys. Acta 949*:318; Escherichia], [Chassy *et al.* (1987) *FEMS Microbiol. Lett.*

44:173 Lactobacillus]; [Fiedler *et al.* (1988) *Anal. Biochem 170*:38, Pseudomonas]; [Augustin *et al.* (1990) *FEMS Microbiol Lett 66*:203, Staphylococcus], [Barany *et al.* (1980) *J. Bacteriol. 144*:698; Harlander (1987) "Transformation of Streptococcus lactis by electroporation, in: *Streptococcal Genetics* (ed. J. Ferretti and R. Curtiss III); Perry *et al.* (1981) *Infect. Immun 32*:1295; Powell *et al.* (1988) *Appl. Environ. Microbiol. 54*:655; Somkuti *et al.* (1987) *Proc. 4th Evr. Cong. Biotechnology 1*:412, Streptococcus].

v. Yeast Expression

Yeast expression systems are also known to one of ordinary skill in the art. A yeast promoter is any DNA sequence capable of binding yeast RNA polymerase and initiating the downstream (3') transcription of a coding sequence (*eg.* structural gene) into mRNA. A promoter will have a transcription initiation region which is usually placed proximal to the 5' end of the coding sequence. This transcription initiation region usually includes an RNA polymerase binding site (the "TATA Box") and a transcription initiation site. A yeast promoter may also have a second domain called an upstream activator sequence (UAS), which, if present, is usually distal to the structural gene. The UAS permits regulated (inducible) expression. Constitutive expression occurs in the absence of a UAS. Regulated expression may be either positive or negative, thereby either enhancing or reducing transcription.

Yeast is a fermenting organism with an active metabolic pathway, therefore sequences encoding enzymes in the metabolic pathway provide particularly useful promoter sequences. Examples include alcohol dehydrogenase (ADH) (EP-A-0 284 044), enolase, glucokinase, glucose-6-phosphate isomerase, glyceraldehyde-3-phosphate-dehydrogenase (GAP or GAPDH), hexokinase, phosphofructokinase, 3-phosphoglycerate mutase, and pyruvate kinase (PyK) (EPO-A-0 329 203). The yeast *PHO5* gene, encoding acid phosphatase, also provides useful promoter sequences [Myanohara *et al.* (1983) *Proc. Natl. Acad. Sci. USA 80*:1].

In addition, synthetic promoters which do not occur in nature also function as yeast promoters. For example, UAS sequences of one yeast promoter may be joined with the transcription activation region of another yeast promoter, creating a synthetic hybrid promoter. Examples of such hybrid promoters include the ADH regulatory sequence linked to the GAP transcription activation region (US Patent Nos. 4,876,197 and 4,880,734). Other examples of hybrid promoters include promoters which consist of the regulatory sequences of either the *ADH2*, *GAL4*, *GAL10*, OR *PHO5* genes,

combined with the transcriptional activation region of a glycolytic enzyme gene such as GAP or PyK (EP-A-0 164 556). Furthermore, a yeast promoter can include naturally occurring promoters of non-yeast origin that have the ability to bind yeast RNA polymerase and initiate transcription. Examples of such promoters include, *inter alia*, [Cohen *et al.* (1980) *Proc. Natl. Acad. Sci. USA* 77:1078; Henikoff *et al.* (1981) *Nature 283*:835; Hollenberg *et al.* (1981) *Curr. Topics Microbiol Immunol. 96*:119; Hollenberg *et al.* (1979) "The Expression of Bacterial Antibiotic Resistance Genes in the Yeast Saccharomyces cerevisiae," in: *Plasmids of Medical, Environmental and Commercial Importance* (eds. K.N. Timmis and A. Puhler); Mercerau-Puigalon *et al.* (1980) *Gene 11*:163; Panthier *et al.* (1980) *Curr. Genet. 2*:109;].

A DNA molecule may be expressed intracellularly in yeast. A promoter sequence may be directly linked with the DNA molecule, in which case the first amino acid at the N-terminus of the recombinant protein will always be a methionine, which is encoded by the ATG start codon. If desired, methionine at the N-terminus may be cleaved from the protein by *in vitro* incubation with cyanogen bromide.

Fusion proteins provide an alternative for yeast expression systems, as well as in mammalian, baculovirus, and bacterial expression systems. Usually, a DNA sequence encoding the N-terminal portion of an endogenous yeast protein, or other stable protein, is fused to the 5' end of heterologous coding sequences. Upon expression, this construct will provide a fusion of the two amino acid sequences. For example, the yeast or human superoxide dismutase (SOD) gene, can be linked at the 5' terminus of a foreign gene and expressed in yeast. The DNA sequence at the junction of the two amino acid sequences may or may not encode a cleavable site. See *eg.* EP-A-0 196 056. Another example is a ubiquitin fusion protein. Such a fusion protein is made with the ubiquitin region that preferably retains a site for a processing enzyme (*eg.* ubiquitin-specific processing protease) to cleave the ubiquitin from the foreign protein. Through this method, therefore, native foreign protein can be isolated (*eg.* WO88/024066).

Alternatively, foreign proteins can also be secreted from the cell into the growth media by creating chimeric DNA molecules that encode a fusion protein comprised of a leader sequence fragment that provide for secretion in yeast of the foreign protein. Preferably, there are processing sites encoded between the leader fragment and the foreign gene that can be cleaved either *in vivo* or *in vitro*. The

leader sequence fragment usually encodes a signal peptide comprised of hydrophobic amino acids which direct the secretion of the protein from the cell.

DNA encoding suitable signal sequences can be derived from genes for secreted yeast proteins, such as the yeast invertase gene (EP-A-0 012 873; JPO. 62,096,086) and the A-factor gene (US patent 4,588,684). Alternatively, leaders of non-yeast origin, such as an interferon leader, exist that also provide for secretion in yeast (EP-A-0 060 057).

A preferred class of secretion leaders are those that employ a fragment of the yeast alpha-factor gene, which contains both a "pre" signal sequence, and a "pro" region. The types of alpha-factor fragments that can be employed include the full-length pre-pro alpha factor leader (about 83 amino acid residues) as well as truncated alpha-factor leaders (usually about 25 to about 50 amino acid residues) (US Patents 4,546,083 and 4,870,008; EP-A-0 324 274). Additional leaders employing an alpha-factor leader fragment that provides for secretion include hybrid alpha-factor leaders made with a presequence of a first yeast, but a pro-region from a second yeast alphafactor. (*eg.* see WO 89/02463.)

Usually, transcription termination sequences recognized by yeast are regulatory regions located 3' to the translation stop codon, and thus together with the promoter flank the coding sequence. These sequences direct the transcription of an mRNA which can be translated into the polypeptide encoded by the DNA. Examples of transcription terminator sequence and other yeast-recognized termination sequences, such as those coding for glycolytic enzymes.

Usually, the above described components, comprising a promoter, leader (if desired), coding sequence of interest, and transcription termination sequence, are put together into expression constructs. Expression constructs are often maintained in a replicon, such as an extrachromosomal element (*eg.* plasmids) capable of stable maintenance in a host, such as yeast or bacteria. The replicon may have two replication systems, thus allowing it to be maintained, for example, in yeast for expression and in a prokaryotic host for cloning and amplification. Examples of such yeast-bacteria shuttle vectors include YEp24 [Botstein *et al.* (1979) *Gene 8*:17-24], pCl/1 [Brake *et al.* (1984) *Proc. Natl. Acad. Sci USA 81*:4642-4646], and YRp17 [Stinchcomb *et al.* (1982) *J. Mol. Biol. 158*:157]. In addition, a replicon may be either a high or low copy number plasmid. A high copy number plasmid will generally have a copy number ranging from about 5 to about 200, and

usually about 10 to about 150. A host containing a high copy number plasmid will preferably have at least about 10, and more preferably at least about 20. Enter a high or low copy number vector may be selected, depending upon the effect of the vector and the foreign protein on the host. See *eg.* Brake *et al.*, *supra.*

Alternatively, the expression constructs can be integrated into the yeast genome with an integrating vector. Integrating vectors usually contain at least one sequence homologous to a yeast chromosome that allows the vector to integrate, and preferably contain two homologous sequences flanking the expression construct. Integrations appear to result from recombinations between homologous DNA in the vector and the yeast chromosome [Orr-Weaver *et al.* (1983) *Methods in Enzymol. 101*:228-245]. An integrating vector may be directed to a specific locus in yeast by selecting the appropriate homologous sequence for inclusion in the vector. See Orr-Weaver *et al.*, *supra.* One or more expression construct may integrate, possibly affecting levels of recombinant protein produced [Rine *et al.* (1983) *Proc. Natl. Acad. Sci. USA 80*:6750]. The chromosomal sequences included in the vector can occur either as a single segment in the vector, which results in the integration of the entire vector, or two segments homologous to adjacent segments in the chromosome and flanking the expression construct in the vector, which can result in the stable integration of only the expression construct.

Usually, extrachromosomal and integrating expression constructs may contain selectable markers to allow for the selection of yeast strains that have been transformed. Selectable markers may include biosynthetic genes that can be expressed in the yeast host, such as *ADE2*, *HIS4*, *LEU2*, *TRP1*, and *ALG7*, and the G418 resistance gene, which confer resistance in yeast cells to tunicamycin and G418, respectively. In addition, a suitable selectable marker may also provide yeast with the ability to grow in the presence of toxic compounds, such as metal. For example, the presence of *CUP1* allows yeast to grow in the presence of copper ions [Butt *et al.* (1987) *Microbiol, Rev. 51*:351].

Alternatively, some of the above described components can be put together into transformation vectors. Transformation vectors are usually comprised of a selectable marker that is either maintained in a replicon or developed into an integrating vector, as described above.

Expression and transformation vectors, either extrachromosomal replicons or integrating vectors, have been developed for transformation into many yeasts. For example, expression vectors have been developed for, *inter alia*, the following yeasts:Candida albicans [Kurtz, *et al.* (1986) *Mol. Cell. Biol. 6*:142], Candida maltosa [Kunze, *et al.* (1985) *J. Basic Microbiol. 25*:141]. Hansenula polymorpha [Gleeson, *et al.* (1986) *J. Gen. Microbiol. 132*:3459; Roggenkamp *et al.* (1986) *Mol. Gen. Genet. 202*:302], Kluyveromyces fragilis [Das, *et al.* (1984) *J Bacteriol. 158*:1165], Kluyveromyces lactis [De Louvencourt *et al.* (1983) *J. Bacteriol. 154*:737; Van den Berg *et al.* (1990) *Bio/Technology 8*:135], Pichia guillerimondii [Kunze *et al.* (1985) *J. Basic Microbiol. 25*:141], Pichia pastoris [Cregg, *et al.* (1985) *Mol. Cell. Biol. 5*:3376; US Patent Nos. 4,837,148 and 4,929,555], Saccharomyces cerevisiae [Hinnen *et al.* (1978) *Proc. Natl. Acad. Sci. USA 75*:1929; Ito *et al.* (1983) *J. Bacteriol. 153*:163], Schizosaccharomyces pombe [Beach and Nurse (1981) *Nature 300*:706], and Yarrowia lipolytica [Davidow, *et al.* (1985) *Curr. Genet. 10*:380471 Gaillardin, *et al.* (1985) *Curr. Genet. 10*:49].

Methods of introducing exogenous DNA into yeast hosts are well-known in the art, and usually include either the transformation of spheroplasts or of intact yeast cells treated with alkali cations. Transformation procedures usually vary with the yeast species to be transformed. See *eg.* [Kurtz *et al.* (1986) *Mol. Cell. Biol. 6*:142; Kunze *et al.* (1985) *J. Basic Microbiol. 25*:141; Candida]; [Gleeson *et al.* (1986) *J. Gen. Microbiol. 132*:3459; Roggenkamp *et al.* (1986) *Mol. Gen. Genet. 202*:302; Hansenula]; [Das *et al.* (1984) *J. Bacteriol. 158*:1165; De Louvencourt *et al.* (1983) *J. Bacteriol. 154*:1165; Van den Berg *et al.* (1990) *Bio/Technology 8*:135; Kluyveromyces]; [Cregg *et al.* (1985) *Mol. Cell. Biol. 5*:3376; Kunze *et al.* (1985) *J. Basic Microbiol. 25*:141; US Patent Nos. 4,837,148 and 4,929,555; Pichia]; [Hinnen *et al.* (1978) *Proc. Natl. Acad. Sci. USA 75*;1929; Ito *et al.* (1983) *J. Bacteriol. 153*:163 Saccharomyces]; [Beach and Nurse (1981) *Nature 300*:706; Schizosaccharomyces]; [Davidow *et al.* (1985) *Curr Genet. 10*:39; Gaillardin *et al.* (1985) *Curr. Genet. 10*:49; Yarrowia].

Antibodies

As used herein, the term "antibody" refers to a polypeptide or group of polypeptides composed of at least one antibody combining site. An "antibody combining site" is the three-dimensional binding space with an internal surface shape and charge distribution complementary to the features of an epitope of an antigen, which allows a binding of the antibody with the antigen. "Antibody"

includes, for example, vertebrate antibodies, hybrid antibodies, chimeric antibodies, humanised antibodies, altered antibodies, univalent antibodies, Fab proteins, and single domain antibodies.

Antibodies against the proteins of the invention are useful for affinity chromatography, immunoassays, and distinguishing/identifying Neisserial proteins.

Antibodies to the proteins of the invention, both polyclonal and monoclonal, may be prepared by conventional methods. In general, the protein is first used to immunize a suitable animal, preferably a mouse, rat, rabbit or goat. Rabbits and goats are preferred for the preparation of polyclonal sera due to the volume of serum obtainable, and the availability of labeled anti-rabbit and anti-goat antibodies. Immunization is generally performed by mixing or emulsifying the protein in saline, preferably in an adjuvant such as Freund's complete adjuvant, and injecting the mixture or emulsion parenterally (generally subcutaneously or intramuscularly). A dose of 50-200 μg/injection is typically sufficient. Immunization is generally boosted 2-6 weeks later with one or more injections of the protein in saline, preferably using Freund's incomplete adjuvant. One may alternatively generate antibodies by in vitro immunization using methods known in the art, which for the purposes of this invention is considered equivalent to *in vivo* immunization. Polyclonal antisera is obtained by bleeding the immunized animal into a glass or plastic container, incubating the blood at 25°C for one hour, followed by incubating at 4°C for 2-18 hours. The serum is recovered by centrifugation (*eg.* 1,000*g* for 10 minutes). About 20-50 ml per bleed may be obtained from rabbits.

Monoclonal antibodies are prepared using the standard method of Kohler & Milstein [*Nature* (1975) 256:495-96], or a modification thereof. Typically, a mouse or rat is immunized as described above. However, rather than bleeding the animal to extract serum, the spleen (and optionally several large lymph nodes) is removed and dissociated into single cells. If desired, the spleen cells may be screened (after removal of nonspecifically adherent cells) by applying a cell suspension to a plate or well coated with the protein antigen. B-cells expressing membrane-bound immunoglobulin specific for the antigen bind to the plate, and are not rinsed away with the rest of the suspension. Resulting B-cells, or all dissociated spleen cells, are then induced to fuse with myeloma cells to form hybridomas, and are cultured in a selective medium (*eg.* hypoxanthine, aminopterin, thymidine medium, "HAT"). The resulting hybridomas are plated by limiting dilution, and are assayed for the production of antibodies which bind specifically to the immunizing antigen

(and which do not bind to unrelated antigens). The selected MAb-secreting hybridomas are then cultured either *in vitro* (*eg.* in tissue culture bottles or hollow fiber reactors), or *in vivo* (as ascites in mice).

If desired, the antibodies (whether polyclonal or monoclonal) may be labeled using conventional techniques. Suitable labels include fluorophores, chromophores, radioactive atoms (particularly ^{32}P and ^{125}I), electron-dense reagents, enzymes, and ligands having specific binding partners. Enzymes are typically detected by their activity. For example, horseradish peroxidase is usually detected by its ability to convert 3,3',5,5'-tetramethylbenzidine (TMB) to a blue pigment, quantifiable with a spectrophotometer. "Specific binding partner" refers to a protein capable of binding a ligand molecule with high specificity, as for example in the case of an antigen and a monoclonal antibody specific therefor. Other specific binding partners include biotin and avidin or streptavidin, IgG and protein A, and the numerous receptor-ligand couples known in the art. It should be understood that the above description is not meant to categorize the various labels into distinct classes, as the same label may serve in several different modes. For example, ^{125}I may serve as a radioactive label or as an electron-dense reagent. HRP may serve as enzyme or as antigen for a MAb. Further, one may combine various labels for desired effect. For example, MAbs and avidin also require labels in the practice of this invention: thus, one might label a MAb with biotin, and detect its presence with avidin labeled with ^{125}I, or with an anti-biotin MAb labeled with HRP. Other permutations and possibilities will be readily apparent to those of ordinary skill in the art, and are considered as equivalents within the scope of the instant invention.

Pharmaceutical Compositions

Pharmaceutical compositions can comprise either polypeptides, antibodies, or nucleic acid of the invention. The pharmaceutical compositions will comprise a therapeutically effective amount of either polypeptides, antibodies, or polynucleotides of the claimed invention.

The term "therapeutically effective amount" as used herein refers to an amount of a therapeutic agent to treat, ameliorate, or prevent a desired disease or condition, or to exhibit a detectable therapeutic or preventative effect. The effect can be detected by, for example, chemical markers or antigen levels. Therapeutic effects also include reduction in physical symptoms, such as decreased body temperature. The precise effective amount for a subject will depend upon the subject's size and health, the nature and extent of the condition, and the therapeutics or combination of

therapeutics selected for administration. Thus, it is not useful to specify an exact effective amount in advance. However, the effective amount for a given situation can be determined by routine experimentation and is within the judgement of the clinician.

For purposes of the present invention, an effective dose will be from about 0.01 mg/ kg to 50 mg/kg or 0.05 mg/kg to about 10 mg/kg of the DNA constructs in the individual to which it is administered.

A pharmaceutical composition can also contain a pharmaceutically acceptable carrier. The term "pharmaceutically acceptable carrier" refers to a carrier for administration of a therapeutic agent, such as antibodies or a polypeptide, genes, and other therapeutic agents. The term refers to any pharmaceutical carrier that does not itself induce the production of antibodies harmful to the individual receiving the composition, and which may be administered without undue toxicity. Suitable carriers may be large, slowly metabolized macromolecules such as proteins, polysaccharides, polylactic acids, polyglycolic acids, polymeric amino acids, amino acid copolymers, and inactive virus particles. Such carriers are well known to those of ordinary skill in the art.

Pharmaceutically acceptable salts can be used therein, for example, mineral acid salts such as hydrochlorides, hydrobromides, phosphates, sulfates, and the like; and the salts of organic acids such as acetates, propionates, malonates, benzoates, and the like. A thorough discussion of pharmaceutically acceptable excipients is available in Remington's Pharmaceutical Sciences (Mack Pub. Co., N.J. 1991).

Pharmaceutically acceptable carriers in therapeutic compositions may contain liquids such as water, saline, glycerol and ethanol. Additionally, auxiliary substances, such as wetting or emulsifying agents, pH buffering substances, and the like, may be present in such vehicles. Typically, the therapeutic compositions are prepared as injectables, either as liquid solutions or suspensions; solid forms suitable for solution in, or suspension in, liquid vehicles prior to injection may also be prepared. Liposomes are included within the definition of a pharmaceutically acceptable carrier.

Delivery Methods

Once formulated, the compositions of the invention can be administered directly to the subject. The subjects to be treated can be animals; in particular, human subjects can be treated.

Direct delivery of the compositions will generally be accomplished by injection, either subcutaneously, intraperitoneally, intravenously or intramuscularly or delivered to the interstitial space of a tissue. The compositions can also be administered into a lesion. Other modes of administration include oral and pulmonary administration, suppositories, and transdermal or transcutaneous applications (*eg.* see WO98/20734), needles, and gene guns or hyposprays. Dosage treatment may be a single dose schedule or a multiple dose schedule.

Vaccines

Vaccines according to the invention may either be prophylactic (*ie.* to prevent infection) or therapeutic (*ie.* to treat disease after infection).

Such vaccines comprise immunising antigen(s), immunogen(s), polypeptide(s), protein(s) or nucleic acid, usually in combination with "pharmaceutically acceptable carriers," which include any carrier that does not itself induce the production of antibodies harmful to the individual receiving the composition. Suitable carriers are typically large, slowly metabolized macromolecules such as proteins, polysaccharides, polylactic acids, polyglycolic acids, polymeric amino acids, amino acid copolymers, lipid aggregates (such as oil droplets or liposomes), and inactive virus particles. Such carriers are well known to those of ordinary skill in the art. Additionally, these carriers may function as immunostimulating agents ("adjuvants"). Furthermore, the antigen or immunogen may be conjugated to a bacterial toxoid, such as a toxoid from diphtheria, tetanus, cholera, *H. pylori*, *etc.* pathogens.

Preferred adjuvants to enhance effectiveness of the composition include, but are not limited to: (1) aluminum salts (alum), such as aluminum hydroxide, aluminum phosphate, aluminum sulfate, etc; (2) oil-in-water emulsion formulations (with or without other specific immunostimulating agents such as muramyl peptides (see below) or bacterial cell wall components), such as for example (a) MF59™ (WO 90/14837; Chapter 10 in *Vaccine design: the subunit and adjuvant approach*, eds. Powell & Newman, Plenum Press 1995), containing 5% Squalene, 0.5% Tween 80, and 0.5% Span 85 (optionally containing various amounts of MTP-PE (see below), although not required) formulated into submicron particles using a microfluidizer such as Model 110Y microfluidizer (Microfluidics, Newton, MA), (b) SAF, containing 10% Squalane, 0.4% Tween 80, 5% pluronic-blocked polymer L121, and thr-MDP (see below) either microfluidized into a submicron emulsion or vortexed to generate a larger particle size emulsion, and (c) Ribi™ adjuvant system (RAS), (Ribi Immunochem, Hamilton, MT) containing 2% Squalene, 0.2% Tween 80, and one or more bacterial

cell wall components from the group consisting of monophosphorylipid A (MPL), trehalose dimycolate (TDM), and cell wall skeleton (CWS), preferably MPL + CWS (Detox™); (3) saponin adjuvants, such as Stimulon™ (Cambridge Bioscience, Worcester, MA) may be used or particles generated therefrom such as ISCOMs (immunostimulating complexes); (4) Complete Freund's Adjuvant (CFA) and Incomplete Freund's Adjuvant (IFA); (5) cytokines, such as interleukins (*eg* IL-1, IL-2, IL-4, IL-5, IL-6, IL-7, IL-12, *etc.*), interferons (*eg.* gamma interferon), macrophage colony stimulating factor (M-CSF), tumor necrosis factor (TNF), etc; and (6) other substances that act as immunostimulating agents to enhance the effectiveness of the composition. Alum and MF59™ are preferred.

As mentioned above, muramyl peptides include, but are not limited to, N-acetyl-muramyl-L-threonyl-D-isoglutamine (thr-MDP), N-acetyl-normuramyl-L-alanyl-D-isoglutamine (nor-MDP), N-acetylmuramyl-L-alanyl-D-isoglutaminyl-L-alanine-2-(1'-2'-dipalmitoyl-*sn*-glycero-3-hydroxyphosphoryloxy)-ethylamine (MTP-PE), *etc.*

The immunogenic compositions (*eg.* the immunising antigen/immunogen/polypeptide/protein/ nucleic acid, pharmaceutically acceptable carrier, and adjuvant) typically will contain diluents, such as water, saline, glycerol, ethanol, etc. Additionally, auxiliary substances, such as wetting or emulsifying agents, pH buffering substances, and the like, may be present in such vehicles.

Typically, the immunogenic compositions are prepared as injectables, either as liquid solutions or suspensions; solid forms suitable for solution in, or suspension in, liquid vehicles prior to injection may also be prepared. The preparation also may be emulsified or encapsulated in liposomes for enhanced adjuvant effect, as discussed above under pharmaceutically acceptable carriers.

Immunogenic compositions used as vaccines comprise an immunologically effective amount of the antigenic or immunogenic polypeptides, as well as any other of the above-mentioned components, as needed. By "immunologically effective amount", it is meant that the administration of that amount to an individual, either in a single dose or as part of a series, is effective for treatment or prevention. This amount varies depending upon the health and physical condition of the individual to be treated, the taxonomic group of individual to be treated (*eg.* nonhuman primate, primate, *etc.*), the capacity of the individual's immune system to synthesize antibodies, the degree of protection desired, the formulation of the vaccine, the treating doctor's assessment of the medical situation,

and other relevant factors. It is expected that the amount will fall in a relatively broad range that can be determined through routine trials.

The immunogenic compositions are conventionally administered parenterally, *eg.* by injection, either subcutaneously, intramuscularly, or transdermally/transcutaneously (*eg.* WO98/20734). Additional formulations suitable for other modes of administration include oral and pulmonary formulations, suppositories, and transdermal applications. Dosage treatment may be a single dose schedule or a multiple dose schedule. The vaccine may be administered in conjunction with other immunoregulatory agents.

As an alternative to protein-based vaccines, DNA vaccination may be employed [*eg.* Robinson & Torres (1997) *Seminars in Immunology* 9:271-283; Donnelly *et al.* (1997) *Annu Rev Immunol* 15:617-648; see later herein].

Gene Delivery Vehicles

Gene therapy vehicles for delivery of constructs including a coding sequence of a therapeutic of the invention, to be delivered to the mammal for expression in the mammal, can be administered either locally or systemically. These constructs can utilize viral or non-viral vector approaches in *in vivo* or *ex vivo* modality. Expression of such coding sequence can be induced using endogenous mammalian or heterologous promoters. Expression of the coding sequence in vivo can be either constitutive or regulated.

The invention includes gene delivery vehicles capable of expressing the contemplated nucleic acid sequences. The gene delivery vehicle is preferably a viral vector and, more preferably, a retroviral, adenoviral, adeno-associated viral (AAV), herpes viral, or alphavirus vector. The viral vector can also be an astrovirus, coronavirus, orthomyxovirus, papovavirus, paramyxovirus, parvovirus, picornavirus, poxvirus, or togavirus viral vector. See generally, Jolly (1994) *Cancer Gene Therapy* 1:51-64; Kimura (1994) *Human Gene Therapy* 5:845-852; Connelly (1995) *Human Gene Therapy* 6:185-193; and Kaplitt (1994) *Nature Genetics* 6:148-153.

Retroviral vectors are well known in the art and we contemplate that any retroviral gene therapy vector is employable in the invention, including B, C and D type retroviruses, xenotropic retroviruses (for example, NZB-X1, NZB-X2 and NZB9-1 (see O'Neill (1985) *J. Virol.* 53:160) polytropic retroviruses

eg. MCF and MCF-MLV (see Kelly (1983) *J. Virol.* 45:291), spumaviruses and lentiviruses. See RNA Tumor Viruses, Second Edition, Cold Spring Harbor Laboratory, 1985.

Portions of the retroviral gene therapy vector may be derived from different retroviruses. For example, retrovector LTRs may be derived from a Murine Sarcoma Virus, a tRNA binding site from a Rous Sarcoma Virus, a packaging signal from a Murine Leukemia Virus, and an origin of second strand synthesis from an Avian Leukosis Virus.

These recombinant retroviral vectors may be used to generate transduction competent retroviral vector particles by introducing them into appropriate packaging cell lines (see US patent 5,591,624). Retrovirus vectors can be constructed for site-specific integration into host cell DNA by incorporation of a chimeric integrase enzyme into the retroviral particle (see WO96/37626). It is preferable that the recombinant viral vector is a replication defective recombinant virus.

Packaging cell lines suitable for use with the above-described retrovirus vectors are well known in the art, are readily prepared (see WO95/30763 and WO92/05266), and can be used to create producer cell lines (also termed vector cell lines or "VCLs") for the production of recombinant vector particles. Preferably, the packaging cell lines are made from human parent cells (*eg.* HT1080 cells) or mink parent cell lines, which eliminates inactivation in human serum.

Preferred retroviruses for the construction of retroviral gene therapy vectors include Avian Leukosis Virus, Bovine Leukemia, Virus, Murine Leukemia Virus, Mink-Cell Focus-Inducing Virus, Murine Sarcoma Virus, Reticuloendotheliosis Virus and Rous Sarcoma Virus. Particularly preferred Murine Leukemia Viruses include 4070A and 1504A (Hartley and Rowe (1976) *J Virol* 19:19-25), Abelson (ATCC No. VR-999), Friend (ATCC No. VR-245), Graffi, Gross (ATCC Nol VR-590), Kirsten, Harvey Sarcoma Virus and Rauscher (ATCC No. VR-998) and Moloney Murine Leukemia Virus (ATCC No. VR-190). Such retroviruses may be obtained from depositories or collections such as the American Type Culture Collection ("ATCC") in Rockville, Maryland or isolated from known sources using commonly available techniques.

Exemplary known retroviral gene therapy vectors employable in this invention include those described in patent applications GB2200651, EP0415731, EP0345242, EP0334301, WO89/02468; WO89/05349, WO89/09271, WO90/02806, WO90/07936, WO94/03622, WO93/25698,

WO93/25234, WO93/11230, WO93/10218, WO91/02805, WO91/02825, WO95/07994, US 5,219,740, US 4,405,712, US 4,861,719, US 4,980,289, US 4,777,127, US 5,591,624. See also Vile (1993) *Cancer Res* 53:3860-3864; Vile (1993) *Cancer Res* 53:962-967; Ram (1993) *Cancer Res* 53 (1993) 83-88; Takamiya (1992) *J Neurosci Res* 33:493-503; Baba (1993) *J Neurosurg* 79:729-735; Mann (1983) *Cell* 33:153; Cane (1984) *Proc Natl Acad Sci* 81:6349; and Miller (1990) *Human Gene Therapy* 1.

Human adenoviral gene therapy vectors are also known in the art and employable in this invention. See, for example, Berkner (1988) *Biotechniques* 6:616 and Rosenfeld (1991) *Science* 252:431, and WO93/07283, WO93/06223, and WO93/07282. Exemplary known adenoviral gene therapy vectors employable in this invention include those described in the above referenced documents and in WO94/12649, WO93/03769, WO93/19191, WO94/28938, WO95/11984, WO95/00655, WO95/27071, WO95/29993, WO95/34671, WO96/05320, WO94/08026, WO94/11506, WO93/06223, WO94/24299, WO95/14102, WO95/24297, WO95/02697, WO94/28152, WO94/24299, WO95/09241, WO95/25807, WO95/05835, WO94/18922 and WO95/09654. Alternatively, administration of DNA linked to killed adenovirus as described in Curiel (1992) *Hum. Gene Ther.* 3:147-154 may be employed. The gene delivery vehicles of the invention also include adenovirus associated virus (AAV) vectors. Leading and preferred examples of such vectors for use in this invention are the AAV-2 based vectors disclosed in Srivastava, WO93/09239. Most preferred AAV vectors comprise the two AAV inverted terminal repeats in which the native D-sequences are modified by substitution of nucleotides, such that at least 5 native nucleotides and up to 18 native nucleotides, preferably at least 10 native nucleotides up to 18 native nucleotides, most preferably 10 native nucleotides are retained and the remaining nucleotides of the D-sequence are deleted or replaced with non-native nucleotides. The native D-sequences of the AAV inverted terminal repeats are sequences of 20 consecutive nucleotides in each AAV inverted terminal repeat (*ie.* there is one sequence at each end) which are not involved in HP formation. The non-native replacement nucleotide may be any nucleotide other than the nucleotide found in the native D-sequence in the same position. Other employable exemplary AAV vectors are pWP-19, pWN-1, both of which are disclosed in Nahreini (1993) *Gene* 124:257-262. Another example of such an AAV vector is psub201 (see Samulski (1987) *J. Virol.* 61:3096). Another exemplary AAV vector is the Double-D ITR vector. Construction of the Double-D ITR vector is disclosed in US Patent 5,478,745. Still other vectors are those disclosed in Carter US Patent 4,797,368 and

Muzyczka US Patent 5,139,941, Chartejee US Patent 5,474,935, and Kotin WO94/288157. Yet a further example of an AAV vector employable in this invention is SSV9AFABTKneo, which contains the AFP enhancer and albumin promoter and directs expression predominantly in the liver. Its structure and construction are disclosed in Su (1996) *Human Gene Therapy* 7:463-470. Additional AAV gene therapy vectors are described in US 5,354,678, US 5,173,414, US 5,139,941, and US 5,252,479.

The gene therapy vectors of the invention also include herpes vectors. Leading and preferred examples are herpes simplex virus vectors containing a sequence encoding a thymidine kinase polypeptide such as those disclosed in US 5,288,641 and EP0176170 (Roizman). Additional exemplary herpes simplex virus vectors include HFEM/ICP6-LacZ disclosed in WO95/04139 (Wistar Institute), pHSVlac described in Geller (1988) *Science* 241:1667-1669 and in WO90/09441 and WO92/07945, HSV Us3::pgC-lacZ described in Fink (1992) *Human Gene Therapy* 3:11-19 and HSV 7134, 2 RH 105 and GAL4 described in EP 0453242 (Breakefield), and those deposited with the ATCC as accession numbers ATCC VR-977 and ATCC VR-260.

Also contemplated are alpha virus gene therapy vectors that can be employed in this invention. Preferred alpha virus vectors are Sindbis viruses vectors. Togaviruses, Semliki Forest virus (ATCC VR-67; ATCC VR-1247), Middleberg virus (ATCC VR-370), Ross River virus (ATCC VR-373; ATCC VR-1246), Venezuelan equine encephalitis virus (ATCC VR923; ATCC VR-1250; ATCC VR-1249; ATCC VR-532), and those described in US patents 5,091,309, 5,217,879, and WO92/10578. More particularly, those alpha virus vectors described in US Serial No. 08/405,627, filed March 15, 1995,WO94/21792, WO92/10578, WO95/07994, US 5,091,309 and US 5,217,879 are employable. Such alpha viruses may be obtained from depositories or collections such as the ATCC in Rockville, Maryland or isolated from known sources using commonly available techniques. Preferably, alphavirus vectors with reduced cytotoxicity are used (see USSN 08/679640).

DNA vector systems such as eukaryotic layered expression systems are also useful for expressing the nucleic acids of the invention. See WO95/07994 for a detailed description of eukaryotic layered expression systems. Preferably, the eukaryotic layered expression systems of the invention are derived from alphavirus vectors and most preferably from Sindbis viral vectors.

Other viral vectors suitable for use in the present invention include those derived from poliovirus, for example ATCC VR-58 and those described in Evans, Nature 339 (1989) 385 and Sabin (1973) *J. Biol. Standardization* 1:115; rhinovirus, for example ATCC VR-1110 and those described in Arnold (1990) *J Cell Biochem* L401; pox viruses such as canary pox virus or vaccinia virus, for example ATCC VR-111 and ATCC VR-2010 and those described in Fisher-Hoch (1989) *Proc Natl Acad Sci* 86:317; Flexner (1989) *Ann NY Acad Sci* 569:86, Flexner (1990) *Vaccine* 8:17; in US 4,603,112 and US 4,769,330 and WO89/01973; SV40 virus, for example ATCC VR-305 and those described in Mulligan (1979) *Nature* 277:108 and Madzak (1992) *J Gen Virol* 73:1533; influenza virus, for example ATCC VR-797 and recombinant influenza viruses made employing reverse genetics techniques as described in US 5,166,057 and in Enami (1990) *Proc Natl Acad Sci* 87:3802-3805; Enami & Palese (1991) *J Virol* 65:2711-2713 and Luytjes (1989) *Cell* 59:110, (see also McMichael (1983) *NEJ Med* 309:13, and Yap (1978) *Nature* 273:238 and *Nature* (1979) 277:108); human immunodeficiency virus as described in EP-0386882 and in Buchschacher (1992) *J. Virol.* 66:2731; measles virus, for example ATCC VR-67 and VR-1247 and those described in EP-0440219; Aura virus, for example ATCC VR-368; Bebaru virus, for example ATCC VR-600 and ATCC VR-1240; Cabassou virus, for example ATCC VR-922; Chikungunya virus, for example ATCC VR-64 and ATCC VR-1241; Fort Morgan Virus, for example ATCC VR-924; Getah virus, for example ATCC VR-369 and ATCC VR-1243; Kyzylagach virus, for example ATCC VR-927; Mayaro virus, for example ATCC VR-66; Mucambo virus, for example ATCC VR-580 and ATCC VR-1244; Ndumu virus, for example ATCC VR-371; Pixuna virus, for example ATCC VR-372 and ATCC VR-1245; Tonate virus, for example ATCC VR-925; Triniti virus, for example ATCC VR-469; Una virus, for example ATCC VR-374; Whataroa virus, for example ATCC VR-926; Y-62-33 virus, for example ATCC VR-375; O'Nyong virus, Eastern encephalitis virus, for example ATCC VR-65 and ATCC VR-1242; Western encephalitis virus, for example ATCC VR-70, ATCC VR-1251, ATCC VR-622 and ATCC VR-1252; and coronavirus, for example ATCC VR-740 and those described in Hamre (1966) *Proc Soc Exp Biol Med* 121:190.

Delivery of the compositions of this invention into cells is not limited to the above mentioned viral vectors. Other delivery methods and media may be employed such as, for example, nucleic acid expression vectors, polycationic condensed DNA linked or unlinked to killed adenovirus alone, for example see US Serial No. 08/366,787, filed December 30, 1994 and Curiel (1992) *Hum Gene Ther* 3:147-154 ligand linked DNA, for example see Wu (1989) *J Biol Chem* 264:16985-16987, eucaryotic cell delivery vehicles cells, for example see US Serial No.08/240,030, filed May 9,

1994, and US Serial No. 08/404,796, deposition of photopolymerized hydrogel materials, hand-held gene transfer particle gun, as described in US Patent 5,149,655, ionizing radiation as described in US5,206,152 and in WO92/11033, nucleic charge neutralization or fusion with cell membranes. Additional approaches are described in Philip (1994) *Mol Cell Biol* 14:2411-2418 and in Woffendin (1994) *Proc Natl Acad Sci* 91:1581-1585.

Particle mediated gene transfer may be employed, for example see US Serial No. 60/023,867. Briefly, the sequence can be inserted into conventional vectors that contain conventional control sequences for high level expression, and then incubated with synthetic gene transfer molecules such as polymeric DNA-binding cations like polylysine, protamine, and albumin, linked to cell targeting ligands such as asialoorosomucoid, as described in Wu & Wu (1987) *J. Biol. Chem.* 262:4429-4432, insulin as described in Hucked (1990) *Biochem Pharmacol* 40:253-263, galactose as described in Plank (1992) *Bioconjugate Chem* 3:533-539, lactose or transferrin.

Naked DNA may also be employed. Exemplary naked DNA introduction methods are described in WO 90/11092 and US 5,580,859. Uptake efficiency may be improved using biodegradable latex beads. DNA coated latex beads are efficiently transported into cells after endocytosis initiation by the beads. The method may be improved further by treatment of the beads to increase hydrophobicity and thereby facilitate disruption of the endosome and release of the DNA into the cytoplasm.

Liposomes that can act as gene delivery vehicles are described in US 5,422,120, WO95/13796, WO94/23697, WO91/14445 and EP-524,968. As described in USSN. 60/023,867, on non-viral delivery, the nucleic acid sequences encoding a polypeptide can be inserted into conventional vectors that contain conventional control sequences for high level expression, and then be incubated with synthetic gene transfer molecules such as polymeric DNA-binding cations like polylysine, protamine, and albumin, linked to cell targeting ligands such as asialoorosomucoid, insulin, galactose, lactose, or transferrin. Other delivery systems include the use of liposomes to encapsulate DNA comprising the gene under the control of a variety of tissue-specific or ubiquitously-active promoters. Further non-viral delivery suitable for use includes mechanical delivery systems such as the approach described in Woffendin *et al* (1994) *Proc. Natl. Acad. Sci. USA* 91(24):11581-11585. Moreover, the coding sequence and the product of expression of such can be delivered through deposition of photopolymerized hydrogel materials. Other conventional methods for gene delivery that can be used for delivery of the coding sequence include, for example, use of

hand-held gene transfer particle gun, as described in US 5,149,655; use of ionizing radiation for activating transferred gene, as described in US 5,206,152 and WO92/11033

Exemplary liposome and polycationic gene delivery vehicles are those described in US 5,422,120 and 4,762,915; in WO 95/13796; WO94/23697; and WO91/14445; in EP-0524968; and in Stryer, Biochemistry, pages 236-240 (1975) W.H. Freeman, San Francisco; Szoka (1980) *Biochem Biophys Acta* 600:1; Bayer (1979) *Biochem Biophys Acta* 550:464; Rivnay (1987) *Meth Enzymol* 149:119; Wang (1987) *Proc Natl Acad Sci* 84:7851; Plant (1989) *Anal Biochem* 176:420.

A polynucleotide composition can comprises therapeutically effective amount of a gene therapy vehicle, as the term is defined above. For purposes of the present invention, an effective dose will be from about 0.01 mg/ kg to 50 mg/kg or 0.05 mg/kg to about 10 mg/kg of the DNA constructs in the individual to which it is administered.

Delivery Methods

Once formulated, the polynucleotide compositions of the invention can be administered (1) directly to the subject; (2) delivered *ex vivo*, to cells derived from the subject; or (3) *in vitro* for expression of recombinant proteins. The subjects to be treated can be mammals or birds. Also, human subjects can be treated.

Direct delivery of the compositions will generally be accomplished by injection, either subcutaneously, intraperitoneally, intravenously or intramuscularly or delivered to the interstitial space of a tissue. The compositions can also be administered into a lesion. Other modes of administration include oral and pulmonary administration, suppositories, and transdermal or transcutaneous applications (*eg.* see WO98/20734), needles, and gene guns or hyposprays. Dosage treatment may be a single dose schedule or a multiple dose schedule.

Methods for the *ex vivo* delivery and reimplantation of transformed cells into a subject are known in the art and described in *eg.* WO93/14778. Examples of cells useful in ex vivo applications include, for example, stem cells, particularly hematopoetic, lymph cells, macrophages, dendritic cells, or tumor cells.

Generally, delivery of nucleic acids for both *ex vivo* and *in vitro* applications can be accomplished by the following procedures, for example, dextran-mediated transfection, calcium phosphate precipitation, polybrene mediated transfection, protoplast fusion, electroporation, encapsulation of the polynucleotide(s) in liposomes, and direct microinjection of the DNA into nuclei, all well known in the art.

Polynucleotide and polypeptide pharmaceutical compositions

In addition to the pharmaceutically acceptable carriers and salts described above, the following additional agents can be used with polynucleotide and/or polypeptide compositions.

A.Polypeptides

One example are polypeptides which include, without limitation: asioloorosomucoid (ASOR); transferrin; asialoglycoproteins; antibodies; antibody fragments; ferritin; interleukins; interferons, granulocyte, macrophage colony stimulating factor (GM-CSF), granulocyte colony stimulating factor (G-CSF), macrophage colony stimulating factor (M-CSF), stem cell factor and erythropoietin. Viral antigens, such as envelope proteins, can also be used. Also, proteins from other invasive organisms, such as the 17 amino acid peptide from the circumsporozoite protein of plasmodium falciparum known as RII.

B.Hormones, Vitamins, *etc.*

Other groups that can be included are, for example: hormones, steroids, androgens, estrogens, thyroid hormone, or vitamins, folic acid.

C.Polyalkylenes, Polysaccharides, *etc.*

Also, polyalkylene glycol can be included with the desired polynucleotides/polypeptides. In a preferred embodiment, the polyalkylene glycol is polyethlylene glycol. In addition, mono-, di-, or polysaccharides can be included. In a preferred embodiment of this aspect, the polysaccharide is dextran or DEAE-dextran. Also, chitosan and poly(lactide-co-glycolide)

D.Lipids, and Liposomes

The desired polynucleotide/polypeptide can also be encapsulated in lipids or packaged in liposomes prior to delivery to the subject or to cells derived therefrom.

Lipid encapsulation is generally accomplished using liposomes which are able to stably bind or entrap and retain nucleic acid. The ratio of condensed polynucleotide to lipid preparation can vary but will generally be around 1:1 (mg DNA:micromoles lipid), or more of lipid. For a review of the use of liposomes as carriers for delivery of nucleic acids, see, Hug and Sleight (1991) *Biochim. Biophys Acta.* 1097:1-17; Straubinger (1983) *Meth. Enzymol.* 101:512-527.

Liposomal preparations for use in the present invention include cationic (positively charged), anionic (negatively charged) and neutral preparations. Cationic liposomes have been shown to mediate intracellular delivery of plasmid DNA (Felgner (1987) *Proc. Natl. Acad. Sci. USA* 84:7413-7416); mRNA (Malone (1989) *Proc. Natl. Acad. Sci. USA* 86:6077-6081); and purified transcription factors (Debs (1990) *J. Biol. Chem.* 265:10189-10192), in functional form.

Cationic liposomes are readily available. For example, N[1-2,3-dioleyloxy)propyl]-N,N,N-triethylammonium (DOTMA) liposomes are available under the trademark Lipofectin, from GIBCO BRL, Grand Island, NY. (See, also, Felgner *supra*). Other commercially available liposomes include transfectace (DDAB/DOPE) and DOTAP/DOPE (Boerhinger). Other cationic liposomes can be prepared from readily available materials using techniques well known in the art. See, *eg.* Szoka (1978) *Proc. Natl. Acad. Sci. USA* 75:4194-4198; WO90/11092 for a description of the synthesis of DOTAP (1,2-bis(oleoyloxy)-3-(trimethylammonio)propane) liposomes.

Similarly, anionic and neutral liposomes are readily available, such as from Avanti Polar Lipids (Birmingham, AL), or can be easily prepared using readily available materials. Such materials include phosphatidyl choline, cholesterol, phosphatidyl ethanolamine, dioleoylphosphatidyl choline (DOPC), dioleoylphosphatidyl glycerol (DOPG), dioleoylphoshatidyl ethanolamine (DOPE), among others. These materials can also be mixed with the DOTMA and DOTAP starting materials in appropriate ratios. Methods for making liposomes using these materials are well known in the art.

The liposomes can comprise multilammelar vesicles (MLVs), small unilamellar vesicles (SUVs), or large unilamellar vesicles (LUVs). The various liposome-nucleic acid complexes are prepared using methods known in the art. See *eg.* Straubinger (1983) *Meth. Immunol.* 101:512-527; Szoka (1978) *Proc. Natl. Acad. Sci. USA* 75:4194-4198; Papahadjopoulos (1975) *Biochim. Biophys. Acta* 394:483; Wilson (1979) *Cell* 17:77); Deamer & Bangham (1976) *Biochim Biophys. Acta* 443:629; Ostro (1977) *Biochem. Biophys Res. Commun.* 76:836; Fraley (1979) *Proc. Natl. Acad. Sci. USA*

76:3348); Enoch & Strittmatter (1979) *Proc Natl Acad. Sci USA* 76:145; Fraley (1980) *J. Biol Chem* (1980) 255:10431; Szoka & Papahadjopoulos (1978) *Proc Natl Acad. Sci. USA* 75:145; and Schaefer-Ridder (1982) *Science* 215:166.

E.Lipoproteins

In addition, lipoproteins can be included with the polynucleotide/polypeptide to be delivered. Examples of lipoproteins to be utilized include: chylomicrons, HDL, IDL, LDL, and VLDL. Mutants, fragments, or fusions of these proteins can also be used. Also, modifications of naturally occurring lipoproteins can be used, such as acetylated LDL. These lipoproteins can target the delivery of polynucleotides to cells expressing lipoprotein receptors. Preferably, if lipoproteins are including with the polynucleotide to be delivered, no other targeting ligand is included in the composition.

Naturally occurring lipoproteins comprise a lipid and a protein portion. The protein portion are known as apoproteins. At the present, apoproteins A, B, C, D, and E have been isolated and identified. At least two of these contain several proteins, designated by Roman numerals, AI, AII, AIV; CI, CII, CIII.

A lipoprotein can comprise more than one apoprotein. For example, naturally occurring chylomicrons comprises of A, B, C, and E, over time these lipoproteins lose A and acquire C and E apoproteins. VLDL comprises A, B, C, and E apoproteins, LDL comprises apoprotein B; and HDL comprises apoproteins A, C, and E.

The amino acid of these apoproteins are known and are described in, for example, Breslow (1985) Annu Rev. Biochem 54:699; Law (1986) Adv. Exp Med. Biol. 151:162; Chen (1986) J Biol Chem 261:12918; Kane (1980) Proc Natl Acad Sci USA 77:2465; and Utermann (1984) Hum Genet 65:232.

Lipoproteins contain a variety of lipids including, triglycerides, cholesterol (free and esters), and phospholipids. The composition of the lipids varies in naturally occurring lipoproteins. For example, chylomicrons comprise mainly triglycerides. A more detailed description of the lipid content of naturally occurring lipoproteins can be found, for example, in *Meth. Enzymol.* 128 (1986). The composition of the lipids are chosen to aid in conformation of the apoprotein for receptor binding activity. The composition of lipids can also be chosen to facilitate hydrophobic interaction and association with the polynucleotide binding molecule.

Naturally occurring lipoproteins can be isolated from serum by ultracentrifugation, for instance. Such methods are described in *Meth. Enzymol. (supra)*; Pitas (1980) *J. Biochem.* 255:5454-5460 and Mahey (1979) *J Clin Invest* 64:743-750. Lipoproteins can also be produced by *in vitro* or recombinant methods by expression of the apoprotein genes in a desired host cell. See, for example, Atkinson (1986) *Annu Rev Biophys Chem* 15:403 and Radding (1958) *Biochim Biophys Acta* 30: 443. Lipoproteins can also be purchased from commercial suppliers, such as Biomedical Techniologies, Inc., Stoughton, Massachusetts, USA. Further description of lipoproteins can be found in Zuckermann *et al.* PCT/US97/14465.

F.Polycationic Agents

Polycationic agents can be included, with or without lipoprotein, in a composition with the desired polynucleotide/polypeptide to be delivered.

Polycationic agents, typically, exhibit a net positive charge at physiological relevant pH and are capable of neutralizing the electrical charge of nucleic acids to facilitate delivery to a desired location. These agents have both in vitro, ex vivo, and in vivo applications. Polycationic agents can be used to deliver nucleic acids to a living subject either intramuscularly, subcutaneously, etc.

The following are examples of useful polypeptides as polycationic agents: polylysine, polyarginine, polyornithine, and protamine. Other examples include histones, protamines, human serum albumin, DNA binding proteins, non-histone chromosomal proteins, coat proteins from DNA viruses, such as (X174, transcriptional factors also contain domains that bind DNA and therefore may be useful as nucleic aid condensing agents. Briefly, transcriptional factors such as C/CEBP, c-jun, c-fos, AP-1, AP-2, AP-3, CPF, Prot-1, Sp-1, Oct-1, Oct-2, CREP, and TFIID contain basic domains that bind DNA sequences.

Organic polycationic agents include: spermine, spermidine, and purtrescine.

The dimensions and of the physical properties of a polycationic agent can be extrapolated from the list above, to construct other polypeptide polycationic agents or to produce synthetic polycationic agents.

Synthetic polycationic agents which are useful include, for example, DEAE-dextran, polybrene. Lipofectin™, and lipofectAMINE™ are monomers that form polycationic complexes when combined with polynucleotides/polypeptides.

Immunodiagnostic Assays

Neisserial antigens of the invention can be used in immunoassays to detect antibody levels (or, conversely, anti-Neisserial antibodies can be used to detect antigen levels). Immunoassays based on well defined, recombinant antigens can be developed to replace invasive diagnostics methods. Antibodies to Neisserial proteins within biological samples, including for example, blood or serum samples, can be detected. Design of the immunoassays is subject to a great deal of variation, and a variety of these are known in the art. Protocols for the immunoassay may be based, for example, upon competition, or direct reaction, or sandwich type assays. Protocols may also, for example, use solid supports, or may be by immunoprecipitation. Most assays involve the use of labeled antibody or polypeptide; the labels may be, for example, fluorescent, chemiluminescent, radioactive, or dye molecules. Assays which amplify the signals from the probe are also known; examples of which are assays which utilize biotin and avidin, and enzyme-labeled and mediated immunoassays, such as ELISA assays.

Kits suitable for immunodiagnosis and containing the appropriate labeled reagents are constructed by packaging the appropriate materials, including the compositions of the invention, in suitable containers, along with the remaining reagents and materials (for example, suitable buffers, salt solutions, *etc.*) required for the conduct of the assay, as well as suitable set of assay instructions.

Nucleic Acid Hybridisation

"Hybridization" refers to the association of two nucleic acid sequences to one another by hydrogen bonding. Typically, one sequence will be fixed to a solid support and the other will be free in solution. Then, the two sequences will be placed in contact with one another under conditions that favor hydrogen bonding. Factors that affect this bonding include: the type and volume of solvent; reaction temperature; time of hybridization; agitation; agents to block the non-specific attachment of the liquid phase sequence to the solid support (Denhardt's reagent or BLOTTO); concentration of the sequences; use of compounds to increase the rate of association of sequences (dextran sulfate or polyethylene glycol); and the stringency of the washing conditions following hybridization. See Sambrook *et al.* [*supra*] Volume 2, chapter 9, pages 9.47 to 9.57.

"Stringency" refers to conditions in a hybridization reaction that favor association of very similar sequences over sequences that differ. For example, the combination of temperature and salt concentration should be chosen that is approximately 120 to 200°C below the calculated Tm of the hybrid under study. The temperature and salt conditions can often be determined empirically in preliminary experiments in which samples of genomic DNA immobilized on filters are hybridized to the sequence of interest and then washed under conditions of different stringencies. See Sambrook *et al.* at page 9.50.

Variables to consider when performing, for example, a Southern blot are (1) the complexity of the DNA being blotted and (2) the homology between the probe and the sequences being detected. The total amount of the fragment(s) to be studied can vary a magnitude of 10, from 0.1 to 1μg for a plasmid or phage digest to 10^{-9} to 10^{-8} g for a single copy gene in a highly complex eukaryotic genome. For lower complexity polynucleotides, substantially shorter blotting, hybridization, and exposure times, a smaller amount of starting polynucleotides, and lower specific activity of probes can be used. For example, a single-copy yeast gene can be detected with an exposure time of only 1 hour starting with 1 μg of yeast DNA, blotting for two hours, and hybridizing for 4-8 hours with a probe of 10^8 cpm/μg. For a single-copy mammalian gene a conservative approach would start with 10 μg of DNA, blot overnight, and hybridize overnight in the presence of 10% dextran sulfate using a probe of greater than 10^8 cpm/μg, resulting in an exposure time of ~24 hours.

Several factors can affect the melting temperature (Tm) of a DNA-DNA hybrid between the probe and the fragment of interest, and consequently, the appropriate conditions for hybridization and washing. In many cases the probe is not 100% homologous to the fragment. Other commonly encountered variables include the length and total G+C content of the hybridizing sequences and the ionic strength and formamide content of the hybridization buffer. The effects of all of these factors can be approximated by a single equation:

$$Tm = 81 + 16.6(\log_{10}Ci) + 0.4[\%(G + C)] - 0.6(\%formamide) - 600/n - 1.5(\%mismatch).$$

where Ci is the salt concentration (monovalent ions) and *n* is the length of the hybrid in base pairs (slightly modified from Meinkoth & Wahl (1984) *Anal. Biochem.* 138: 267-284).

In designing a hybridization experiment, some factors affecting nucleic acid hybridization can be conveniently altered. The temperature of the hybridization and washes and the salt concentration during the washes are the simplest to adjust. As the temperature of the hybridization increases (*ie.* stringency), it becomes less likely for hybridization to occur between strands that are nonhomologous, and as a result, background decreases. If the radiolabeled probe is not completely homologous with the immobilized fragment (as is frequently the case in gene family and interspecies hybridization experiments), the hybridization temperature must be reduced, and background will increase. The temperature of the washes affects the intensity of the hybridizing band and the degree of background in a similar manner. The stringency of the washes is also increased with decreasing salt concentrations.

In general, convenient hybridization temperatures in the presence of 50% formamide are 42°C for a probe with is 95% to 100% homologous to the target fragment, 37°C for 90% to 95% homology, and 32°C for 85% to 90% homology. For lower homologies, formamide content should be lowered and temperature adjusted accordingly, using the equation above. If the homology between the probe and the target fragment are not known, the simplest approach is to start with both hybridization and wash conditions which are nonstringent. If non-specific bands or high background are observed after autoradiography, the filter can be washed at high stringency and reexposed. If the time required for exposure makes this approach impractical, several hybridization and/or washing stringencies should be tested in parallel.

Nucleic Acid Probe Assays

Methods such as PCR, branched DNA probe assays, or blotting techniques utilizing nucleic acid probes according to the invention can determine the presence of cDNA or mRNA. A probe is said to "hybridize" with a sequence of the invention if it can form a duplex or double stranded complex, which is stable enough to be detected.

The nucleic acid probes will hybridize to the Neisserial nucleotide sequences of the invention (including both sense and antisense strands). Though many different nucleotide sequences will encode the amino acid sequence, the native Neisserial sequence is preferred because it is the actual sequence present in cells. mRNA represents a coding sequence and so a probe should be complementary to the coding sequence; single-stranded cDNA is complementary to mRNA, and so a cDNA probe should be complementary to the non-coding sequence.

The probe sequence need not be identical to the Neisserial sequence (or its complement) — some variation in the sequence and length can lead to increased assay sensitivity if the nucleic acid probe can form a duplex with target nucleotides, which can be detected. Also, the nucleic acid probe can include additional nucleotides to stabilize the formed duplex. Additional Neisserial sequence may also be helpful as a label to detect the formed duplex. For example, a non-complementary nucleotide sequence may be attached to the 5' end of the probe, with the remainder of the probe sequence being complementary to a Neisserial sequence. Alternatively, non-complementary bases or longer sequences can be interspersed into the probe, provided that the probe sequence has sufficient complementarity with the a Neisserial sequence in order to hybridize therewith and thereby form a duplex which can be detected.

The exact length and sequence of the probe will depend on the hybridization conditions, such as temperature, salt condition and the like. For example, for diagnostic applications, depending on the complexity of the analyte sequence, the nucleic acid probe typically contains at least 10-20 nucleotides, preferably 15-25, and more preferably at least 30 nucleotides, although it may be shorter than this. Short primers generally require cooler temperatures to form sufficiently stable hybrid complexes with the template.

Probes may be produced by synthetic procedures, such as the triester method of Matteucci *et al.* [*J. Am. Chem. Soc.* (1981) 103:3185], or according to Urdea *et al.* [*Proc. Natl. Acad. Sci. USA* (1983) 80: 7461], or using commercially available automated oligonucleotide synthesizers.

The chemical nature of the probe can be selected according to preference. For certain applications, DNA or RNA are appropriate. For other applications, modifications may be incorporated *eg.* backbone modifications, such as phosphorothioates or methylphosphonates, can be used to increase *in vivo* half-life, alter RNA affinity, increase nuclease resistance *etc.* [*eg.* see Agrawal & Iyer (1995) *Curr Opin Biotechnol* 6:12-19; Agrawal (1996) *TIBTECH* 14:376-387]; analogues such as peptide nucleic acids may also be used [*eg.* see Corey (1997) *TIBTECH* 15:224-229; Buchardt *et al.* (1993) *TIBTECH* 11:384-386].

Alternatively, the polymerase chain reaction (PCR) is another well-known means for detecting small amounts of target nucleic acids. The assay is described in: Mullis *et al.* [*Meth. Enzymol* (1987) 155: 335-350]; US patents 4,683,195 and 4,683,202. Two "primer" nucleotides hybridize

with the target nucleic acids and are used to prime the reaction. The primers can comprise sequence that does not hybridize to the sequence of the amplification target (or its complement) to aid with duplex stability or, for example, to incorporate a convenient restriction site. Typically, such sequence will flank the desired Neisserial sequence.

A thermostable polymerase creates copies of target nucleic acids from the primers using the original target nucleic acids as a template. After a threshold amount of target nucleic acids are generated by the polymerase, they can be detected by more traditional methods, such as Southern blots. When using the Southern blot method, the labelled probe will hybridize to the Neisserial sequence (or its complement).

Also, mRNA or cDNA can be detected by traditional blotting techniques described in Sambrook *et al* [*supra*]. mRNA, or cDNA generated from mRNA using a polymerase enzyme, can be purified and separated using gel electrophoresis. The nucleic acids on the gel are then blotted onto a solid support, such as nitrocellulose. The solid support is exposed to a labelled probe and then washed to remove any unhybridized probe. Next, the duplexes containing the labeled probe are detected. Typically, the probe is labelled with a radioactive moiety.

BRIEF DESCRIPTION OF THE DRAWINGS

Figures 1-7 show biochemical data and sequence analysis pertaining to Examples 1, 2, 3, 7, 13, 16 and 19, respectively, with ORFs 40, 38, 44, 52, 114, 41 and 124.. M1 and M2 are molecular weight markers. Arrows indicate the position of the main recombinant product or, in Western blots, the position of the main *N.meningitidis* immunoreactive band. TP indicates *N.meningitidis* total protein extract; OMV indicates *N.meningitidis* outer membrane vesicle preparation. In bactericidal assay results: a diamond (♦) shows preimmune data; a triangle (▲) shows GST control data; a circle (●) shows data with recombinant *N.meningitidis* protein. Computer analyses show a hydrophilicity plot (upper), an antigenic index plot (middle), and an AMPHI analysis (lower). The AMPHI program has been used to predict T-cell epitopes [Gao *et al.* (1989) *J. Immunol.* 143:3007; Roberts *et al.* (1996) *AIDS Res Hum Retrovir* 12:593; Quakyi *et al.* (1992) *Scand J Immunol* suppl.11:9) and is available in the Protean package of DNASTAR, Inc. (1228 South Park Street, Madison, Wisconsin 53715 USA).

the nucleotide sequences in the examples are derived from combining the results of two or more experiments).

Nucleotide sequences were scanned in all six reading frames to predict the presence of hydrophobic domains using an algorithm based on the statistical studies of Esposti *et al.* [Critical evaluation of the hydropathy of membrane proteins (1990) *Eur J Biochem* 190:207-219]. These domains represent potential transmembrane regions or hydrophobic leader sequences.

Open reading frames were predicted from fragmented nucleotide sequences using the program ORFFINDER (NCBI).

Underlined amino acid sequences indicate possible transmembrane domains or leader sequences in the ORFs, as predicted by the PSORT algorithm (http://www.psort.nibb.ac.jp). Functional domains were also predicted using the MOTIFS program (GCG Wisconsin & PROSITE).

Various tests can be used to assess the *in vivo* immunogenicity of the proteins identified in the examples. For example, the proteins can be expressed recombinantly and used to screen patient sera by immunoblot. A positive reaction between the protein and patient serum indicates that the patient has previously mounted an immune response to the protein in question *ie.* the protein is an immunogen. This method can also be used to identify immunodominant proteins.

The recombinant protein can also be conveniently used to prepare antibodies *eg.* in a mouse. These can be used for direct confirmation that a protein is located on the cell-surface. Labelled antibody (*eg.* fluorescent labelling for FACS) can be incubated with intact bacteria and the presence of label on the bacterial surface confirms the location of the protein.

In particular, the following methods (A) to (S) were used to express, purify and biochemically characterise the proteins of the invention:

A) Chromosomal DNA preparation

N.meningitidis strain 2996 was grown to exponential phase in 100ml of GC medium, harvested by centrifugation, and resuspended in 5ml buffer (20% Sucrose, 50mM Tris-HCl, 50mM EDTA, pH8). After 10 minutes incubation on ice, the bacteria were lysed by adding 10ml lysis solution (50mM NaCl, 1% Na-Sarkosyl, 50μg/ml Proteinase K), and the suspension was incubated at 37°C for 2

hours. Two phenol extractions (equilibrated to pH 8) and one $ChCl_3$/isoamylalcohol (24:1) extraction were performed. DNA was precipitated by addition of 0.3M sodium acetate and 2 volumes ethanol, and was collected by centrifugation. The pellet was washed once with 70% ethanol and redissolved in 4ml buffer (10mM Tris-HCl, 1mM EDTA, pH 8). The DNA concentration was measured by reading the OD at 260 nm.

B) Oligonucleotide design

Synthetic oligonucleotide primers were designed on the basis of the coding sequence of each ORF, using (a) the meningococcus B sequence when available, or (b) the gonococcus/meningococcus A sequence, adapted to the codon preference usage of meningococcus as necessary. Any predicted signal peptides were omitted, by deducing the 5'-end amplification primer sequence immediately downstream from the predicted leader sequence.

The 5' primers included two restriction enzyme recognition sites (*BamHI-NdeI*, *BamHI-NheI*, or *EcoRI-NheI*, depending on the gene's own restriction pattern); the 3' primers included a *XhoI* restriction site. This procedure was established in order to direct the cloning of each amplification product (corresponding to each ORF) into two different expression systems: pGEX-KG (using either *BamHI-XhoI* or *EcoRI-XhoI*), and pET21b+ (using either *NdeI-XhoI* or *NheI-XhoI*).

5'-end primer tail:	CGCGGATCCCATATG	(*BamHI-NdeI*)
	CGCGGATCCGCTAGC	(*BamHI-NheI*)
	CCGGAATTCTAGCTAGC	(*EcoRI-NheI*)
3'-end primer tail:	CCCGCTCGAG	(*XhoI*)

As well as containing the restriction enzyme recognition sequences, the primers included nucleotides which hybridised to the sequence to be amplified. The number of hybridizing nucleotides depended on the melting temperature of the whole primer, and was determined for each primer using the formulae:

$T_m = 4\ (G+C) + 2\ (A+T)$ (tail excluded)

$T_m = 64.9 + 0.41\ (\%\ GC) - 600/N$ (whole primer)

The average melting temperature of the selected oligos were 65-70°C for the whole oligo and 50-55°C for the hybridising region alone.

Table I shows the forward and reverse primers used for each amplification Oligos were synthesized by a Perkin Elmer 394 DNA/RNA Synthesizer, eluted from the columns in 2ml NH_4OH, and deprotected by 5 hours incubation at 56°C. The oligos were precipitated by addition of 0.3M Na-Acetate and 2 volumes ethanol. The samples were then centrifuged and the pellets resuspended in either 100µl or 1ml of water. OD_{260} was determined using a Perkin Elmer Lambda Bio spectrophotometer and the concentration was determined and adjusted to 2-10pmol/µl.

C) Amplification

The standard PCR protocol was as follows: 50-200ng of genomic DNA were used as a template in the presence of 20-40µM of each oligo, 400-800µM dNTPs solution, 1x PCR buffer (including 1.5mM $MgCl_2$), 2.5 units *TaqI* DNA polymerase (using Perkin-Elmer AmpliTaQ, GIBCO Platinum, Pwo DNA polymerase, or Tahara Shuzo Taq polymerase).

In some cases, PCR was optimised by the addition of 10µl DMSO or 50µl 2M betaine.

After a hot start (adding the polymerase during a preliminary 3 minute incubation of the whole mix at 95°C), each sample underwent a double-step amplification: the first 5 cycles were performed using as the hybridization temperature the one of the oligos excluding the restriction enzymes tail, followed by 30 cycles performed according to the hybridization temperature of the whole length oligos. The cycles were followed by a final 10 minute extension step at 72°C.

The standard cycles were as follows:

	Denaturation	**Hybridisation**	**Elongation**
First 5 cycles	30 seconds 95°C	30 seconds 50-55°C	30-60 seconds 72°C
Last 30 cycles	30 seconds 95°C	30 seconds 65-70°C	30-60 seconds 72°C

The elongation time varied according to the length of the ORF to be amplified.

The amplifications were performed using either a 9600 or a 2400 Perkin Elmer GeneAmp PCR System. To check the results, 1/10 of the amplification volume was loaded onto a 1-1.5% agarose gel and the size of each amplified fragment compared with a DNA molecular weight marker.

The amplified DNA was either loaded directly on a 1% agarose gel or first precipitated with ethanol and resuspended in a suitable volume to be loaded on a 1% agarose gel. The DNA fragment corresponding to the right size band was then eluted and purified from gel, using the Qiagen Gel Extraction Kit, following the instructions of the manufacturer. The final volume of the DNA fragment was 30µl or 50µl of either water or 10mM Tris, pH 8.5.

D) Digestion of PCR fragments

The purified DNA corresponding to the amplified fragment was split into 2 aliquots and double-digested with:

- *NdeI/XhoI* or *NheI/XhoI* for cloning into pET-21b+ and further expression of the protein as a C-terminus His-tag fusion

- *BamHI/XhoI* or *EcoRI/XhoI* for cloning into pGEX-KG and further expression of the protein as N-terminus GST fusion.

- *EcoRI/PstI, EcoRI/SalI, SalI/PstI* for cloning into pGex-His and further expression of the protein as N-terminus His-tag fusion

Each purified DNA fragment was incubated (37°C for 3 hours to overnight) with 20 units of each restriction enzyme (New England Biolabs) in a either 30 or 40µl final volume in the presence of the appropriate buffer. The digestion product was then purified using the QIAquick PCR purification kit, following the manufacturer's instructions, and eluted in a final volume of 30 or 50µl of either water or 10mM Tris-HCl, pH 8.5. The final DNA concentration was determined by 1% agarose gel electrophoresis in the presence of titrated molecular weight marker.

E) Digestion of the cloning vectors (pET22B, pGEX-KG, pTRC-His A, and pGex-His)

10µg plasmid was double-digested with 50 units of each restriction enzyme in 200µl reaction volume in the presence of appropriate buffer by overnight incubation at 37°C. After loading the

whole digestion on a 1% agarose gel, the band corresponding to the digested vector was purified from the gel using the Qiagen QIAquick Gel Extraction Kit and the DNA was eluted in 50µl of 10mM Tris-HCl, pH 8.5. The DNA concentration was evaluated by measuring OD_{260} of the sample, and adjusted to 50µg/µl. 1µl of plasmid was used for each cloning procedure.

The vector pGEX-His is a modified pGEX-2T vector carrying a region encoding six histidine residues upstream to the thrombin cleavage site and containing the multiple cloning site of the vector pTRC99 (Pharmacia).

F) Cloning

The fragments corresponding to each ORF, previously digested and purified, were ligated in both pET22b and pGEX-KG. In a final volume of 20µl, a molar ratio of 3:1 fragment/vector was ligated using 0.5µl of NEB T4 DNA ligase (400 units/µl), in the presence of the buffer supplied by the manufacturer. The reaction was incubated at room temperature for 3 hours. In some experiments, ligation was performed using the Boehringer "Rapid Ligation Kit", following the manufacturer's instructions.

In order to introduce the recombinant plasmid in a suitable strain, 100µl *E. coli* DH5 competent cells were incubated with the ligase reaction solution for 40 minutes on ice, then at 37°C for 3 minutes, then, after adding 800µl LB broth, again at 37°C for 20 minutes. The cells were then centrifuged at maximum speed in an Eppendorf microfuge and resuspended in approximately 200µl of the supernatant. The suspension was then plated on LB ampicillin (100mg/ml).

The screening of the recombinant clones was performed by growing 5 randomly-chosen colonies overnight at 37°C in either 2ml (pGEX or pTC clones) or 5ml (pET clones) LB broth + 100µg/ml ampicillin. The cells were then pelletted and the DNA extracted using the Qiagen QIAprep Spin Miniprep Kit, following the manufacturer's instructions, to a final volume of 30µl. 5µl of each individual miniprep (approximately 1g) were digested with either *NdeI/XhoI* or *BamHI/XhoI* and the whole digestion loaded onto a 1-1.5% agarose gel (depending on the expected insert size), in parallel with the molecular weight marker (1Kb DNA Ladder, GIBCO). The screening of the positive clones was made on the base of the correct insert size.

G) Expression

Each ORF cloned into the expression vector was transformed into the strain suitable for expression of the recombinant protein product. 1µl of each construct was used to transform 30µl of *E.coli* BL21 (pGEX vector), *E.coli* TOP 10 (pTRC vector) or *E coli* BL21-DE3 (pET vector), as described above. In the case of the pGEX-His vector, the same *E.coli* strain (W3110) was used for initial cloning and expression. Single recombinant colonies were inoculated into 2ml LB+Amp (100µg/ml), incubated at 37°C overnight, then diluted 1:30 in 20ml of LB+Amp (100µg/ml) in 100ml flasks, making sure that the OD_{600} ranged between 0.1 and 0.15. The flasks were incubated at 30°C into gyratory water bath shakers until OD indicated exponential growth suitable for induction of expression (0.4-0.8 OD for pET and pTRC vectors; 0.8-1 OD for pGEX and pGEX-His vectors). For the pET, pTRC and pGEX-His vectors, the protein expression was induced by addition of 1mM IPTG, whereas in the case of pGEX system the final concentration of IPTG was 0.2mM. After 3 hours incubation at 30°C, the final concentration of the sample was checked by OD. In order to check expression, 1ml of each sample was removed, centrifuged in a microfuge, the pellet resuspended in PBS, and analysed by 12% SDS-PAGE with Coomassie Blue staining. The whole sample was centrifuged at 6000*g* and the pellet resuspended in PBS for further use.

H) GST-fusion proteins large-scale purification.

A single colony was grown overnight at 37°C on LB+Amp agar plate. The bacteria were inoculated into 20ml of LB+Amp liquid culture in a water bath shaker and grown overnight. Bacteria were diluted 1:30 into 600ml of fresh medium and allowed to grow at the optimal temperature (20-37°C) to OD_{550} 0.8-1. Protein expression was induced with 0.2mM IPTG followed by three hours incubation. The culture was centrifuged at 8000rpm at 4°C. The supernatant was discarded and the bacterial pellet was resuspended in 7.5ml cold PBS. The cells were disrupted by sonication on ice for 30 sec at 40W using a Branson sonifier B-15, frozen and thawed twice and centrifuged again. The supernatant was collected and mixed with 150µl Glutatione-Sepharose 4B resin (Pharmacia) (previously washed with PBS) and incubated at room temperature for 30 minutes. The sample was centrifuged at 700*g* for 5 minutes at 4°C. The resin was washed twice with 10ml cold PBS for 10 minutes, resuspended in 1ml cold PBS, and loaded on a disposable column. The resin was washed twice with 2ml cold PBS until the flow-through reached OD_{280} of 0.02-0.06. The GST-fusion protein was eluted by addition of 700µl cold Glutathione elution buffer (10mM reduced

glutathione, 50mM Tris-HCl) and fractions collected until the OD_{280} was 0.1. 21µl of each fraction were loaded on a 12% SDS gel using either Biorad SDS-PAGE Molecular weight standard broad range (M1) (200, 116.25, 97.4, 66.2, 45, 31, 21.5, 14.4, 6.5 kDa) or Amersham Rainbow Marker (M2) (220, 66, 46, 30, 21.5, 14.3 kDa) as standards As the MW of GST is 26kDa, this value must be added to the MW of each GST-fusion protein.

I) His-fusion solubility analysis

To analyse the solubility of the His-fusion expression products, pellets of 3ml cultures were resuspended in buffer M1 [500µl PBS pH 7.2]. 25µl lysozyme (10mg/ml) was added and the bacteria were incubated for 15 min at 4°C. The pellets were sonicated for 30 sec at 40W using a Branson sonifier B-15, frozen and thawed twice and then separated again into pellet and supernatant by a centrifugation step. The supernatant was collected and the pellet was resuspended in buffer M2 [8M urea, 0.5M NaCl, 20mM imidazole and 0.1M NaH_2PO_4] and incubated for 3 to 4 hours at 4°C. After centrifugation, the supernatant was collected and the pellet was resuspended in buffer M3 [6M guanidinium-HCl, 0.5M NaCl, 20mM imidazole and 0.1M NaH_2PO_4] overnight at 4°C. The supernatants from all steps were analysed by SDS-PAGE.

J) His-fusion large-scale purification.

A single colony was grown overnight at 37°C on a LB + Amp agar plate. The bacteria were inoculated into 20ml of LB+Amp liquid culture and incubated overnight in a water bath shaker. Bacteria were diluted 1:30 into 600ml fresh medium and allowed to grow at the optimal temperature (20-37°C) to OD_{550} 0.6-0.8. Protein expression was induced by addition of 1mM IPTG and the culture further incubated for three hours. The culture was centrifuged at 8000rpm at 4°C, the supernatant was discarded and the bacterial pellet was resuspended in 7.5ml of either (i) cold buffer A (300mM NaCl, 50mM phosphate buffer, 10mM imidazole, pH 8) for soluble proteins or (ii) buffer B (urea 8M, 10mM Tris-HCl, 100mM phosphate buffer, pH 8.8) for insoluble proteins.

The cells were disrupted by sonication on ice for 30 sec at 40W using a Branson sonifier B-15, frozen and thawed two times and centrifuged again.

For insoluble proteins, the supernatant was stored at -20°C, while the pellets were resuspended in 2ml buffer C (6M guanidine hydrochloride, 100mM phosphate buffer, 10mM Tris-HCl, pH 7.5) and treated in a homogenizer for 10 cycles. The product was centrifuged at 13000rpm for 40 minutes.

Supernatants were collected and mixed with 150µl Ni^{2+}-resin (Pharmacia) (previously washed with either buffer A or buffer B, as appropriate) and incubated at room temperature with gentle agitation for 30 minutes. The sample was centrifuged at 700*g* for 5 minutes at 4°C. The resin was washed twice with 10ml buffer A or B for 10 minutes, resuspended in 1ml buffer A or B and loaded on a disposable column. The resin was washed at either (i) 4°C with 2ml cold buffer A or (ii) room temperature with 2ml buffer B, until the flow-through reached OD_{280} of 0.02-0.06.

The resin was washed with either (i) 2ml cold 20mM imidazole buffer (300mM NaCl, 50mM phosphate buffer, 20mM imidazole, pH 8) or (ii) buffer D (urea 8M, 10mM Tris-HCl, 100mM phosphate buffer, pH 6.3) until the flow-through reached the $O.D_{280}$ of 0.02-0.06. The His-fusion protein was eluted by addition of 700µl of either (i) cold elution buffer A (300mM NaCl, 50mM phosphate buffer, 250mM imidazole, pH 8) or (ii) elution buffer B (urea 8M, 10mM Tris-HCl, 100mM phosphate buffer, pH 4.5) and fractions collected until the $O.D_{280}$ was 0.1. 21µl of each fraction were loaded on a 12% SDS gel.

K) His-fusion proteins renaturation

10% glycerol was added to the denatured proteins. The proteins were then diluted to 20µg/ml using dialysis buffer I (10% glycerol, 0.5M arginine, 50mM phosphate buffer, 5mM reduced glutathione, 0.5mM oxidised glutathione, 2M urea, pH 8.8) and dialysed against the same buffer at 4°C for 12-14 hours. The protein was further dialysed against dialysis buffer II (10% glycerol, 0.5M arginine, 50mM phosphate buffer, 5mM reduced glutathione, 0.5mM oxidised glutathione, pH 8.8) for 12-14 hours at 4°C. Protein concentration was evaluated using the formula:

$$\text{Protein (mg/ml)} = (1.55 \times OD_{280}) - (0.76 \times OD_{260})$$

L) His-fusion large-scale purification

500ml of bacterial cultures were induced and the fusion proteins were obtained soluble in buffer M1, M2 or M3 using the procedure described above. The crude extract of the bacteria was loaded

onto a Ni-NTA superflow column (Qiagen) equilibrated with buffer M1, M2 or M3 depending on the solubilization buffer of the fusion proteins Unbound material was eluted by washing the column with the same buffer. The specific protein was eluted with the corresponding buffer containing 500mM imidazole and dialysed against the corresponding buffer without imidazole. After each run the columns were sanitized by washing with at least two column volumes of 0.5 M sodium hydroxide and reequilibrated before the next use.

M) Mice immunisations

20μg of each purified protein were used to immunise mice intraperitoneally. In the case of ORF 44, CD1 mice were immunised with $Al(OH)_3$ as adjuvant on days 1, 21 and 42, and immune response was monitored in samples taken on day 56. For ORF 40, CD1 mice were immunised using Freund's adjuvant, rather than $Al(OH)_3$, and the same immunisation protocol was used, except that the immune response was measured on day 42, rather than 56. Similarly, for ORF 38, CD1 mice were immunised with Freund's adjuvant, but the immune response was measured on day 49.

N) ELISA assay (sera analysis)

The acapsulated MenB M7 strain was plated on chocolate agar plates and incubated overnight at 37°C. Bacterial colonies were collected from the agar plates using a sterile dracon swab and inoculated into 7ml of Mueller-Hinton Broth (Difco) containing 0.25% Glucose. Bacterial growth was monitored every 30 minutes by following OD_{620}. The bacteria were let to grow until the OD reached the value of 0.3-0.4. The culture was centrifuged for 10 minutes at 10000rpm. The supernatant was discarded and bacteria were washed once with PBS, resuspended in PBS containing 0.025% formaldehyde, and incubated for 2 hours at room temperature and then overnight at 4°C with stirring. 100μl bacterial cells were added to each well of a 96 well Greiner plate and incubated overnight at 4°C. The wells were then washed three times with PBT washing buffer (0.1% Tween-20 in PBS). 200μl of saturation buffer (2.7% Polyvinylpyrrolidone 10 in water) was added to each well and the plates incubated for 2 hours at 37°C. Wells were washed three times with PBT. 200μl of diluted sera (Dilution buffer: 1% BSA, 0.1% Tween-20, 0.1% NaN_3 in PBS) were added to each well and the plates incubated for 90 minutes at 37°C. Wells were washed three times with PBT. 100μl of HRP-conjugated rabbit anti-mouse (Dako) serum diluted 1:2000 in dilution buffer were added to each well and the plates were incubated for 90 minutes at

37°C. Wells were washed three times with PBT buffer. 100µl of substrate buffer for HRP (25ml of citrate buffer pH5, 10mg of O-phenildiamine and 10µl of H_2O) were added to each well and the plates were left at room temperature for 20 minutes. 100µl H_2SO_4 was added to each well and OD_{490} was followed. The ELISA was considered positive when OD_{490} was 2.5 times the respective pre-immune sera.

O) FACScan bacteria Binding Assay procedure.

The acapsulated MenB M7 strain was plated on chocolate agar plates and incubated overnight at 37°C. Bacterial colonies were collected from the agar plates using a sterile dracon swab and inoculated into 4 tubes containing 8ml each Mueller-Hinton Broth (Difco) containing 0.25% glucose. Bacterial growth was monitored every 30 minutes by following OD_{620}. The bacteria were let to grow until the OD reached the value of 0.35-0.5. The culture was centrifuged for 10 minutes at 4000rpm. The supernatant was discarded and the pellet was resuspended in blocking buffer (1% BSA, 0.4% NaN_3) and centrifuged for 5 minutes at 4000rpm. Cells were resuspended in blocking buffer to reach OD_{620} of 0.07. 100µl bacterial cells were added to each well of a Costar 96 well plate. 100µl of diluted (1:200) sera (in blocking buffer) were added to each well and plates incubated for 2 hours at 4°C. Cells were centrifuged for 5 minutes at 4000rpm, the supernatant aspirated and cells washed by addition of 200µl/well of blocking buffer in each well. 100µl of R-Phicoerytrin conjugated $F(ab)_2$ goat anti-mouse, diluted 1:100, was added to each well and plates incubated for 1 hour at 4°C. Cells were spun down by centrifugation at 4000rpm for 5 minutes and washed by addition of 200µl/well of blocking buffer. The supernatant was aspirated and cells resuspended in 200µl/well of PBS, 0.25% formaldehyde. Samples were transferred to FACScan tubes and read. The condition for FACScan setting were: FL1 on, FL2 and FL3 off; FSC-H threshold:92; FSC PMT Voltage: E 02; SSC PMT: 474; Amp. Gains 7.1; FL-2 PMT: 539; compensation values: 0.

P) OMV preparations

Bacteria were grown overnight on 5 GC plates, harvested with a loop and resuspended in 10 ml 20mM Tris-HCl. Heat inactivation was performed at 56°C for 30 minutes and the bacteria disrupted by sonication for 10 minutes on ice (50% duty cycle, 50% output). Unbroken cells were removed by centrifugation at 5000*g* for 10 minutes and the total cell envelope fraction recovered by centrifugation

at 50000*g* at 4°C for 75 minutes. To extract cytoplasmic membrane proteins from the crude outer membranes, the whole fraction was resuspended in 2% sarkosyl (Sigma) and incubated at room temperature for 20 minutes. The suspension was centrifuged at 10000*g* for 10 minutes to remove aggregates, and the supernatant further ultracentrifuged at 50000*g* for 75 minutes to pellet the outer membranes. The outer membranes were resuspended in 10mM Tris-HCl, pH8 and the protein concentration measured by the Bio-Rad Protein assay, using BSA as a standard.

Q) Whole Extracts preparation

Bacteria were grown overnight on a GC plate, harvested with a loop and resuspended in 1ml of 20mM Tris-HCl. Heat inactivation was performed at 56°C for 30 minutes.

R) Western blotting

Purified proteins (500ng/lane), outer membrane vesicles (5µg) and total cell extracts (25µg) derived from MenB strain 2996 were loaded on 15% SDS-PAGE and transferred to a nitrocellulose membrane. The transfer was performed for 2 hours at 150mA at 4°C, in transferring buffer (0.3 % Tris base, 1.44 % glycine, 20% methanol). The membrane was saturated by overnight incubation at 4°C in saturation buffer (10% skimmed milk, 0.1% Triton X100 in PBS). The membrane was washed twice with washing buffer (3% skimmed milk, 0.1% Triton X100 in PBS) and incubated for 2 hours at 37°C with mice sera diluted 1:200 in washing buffer. The membrane was washed twice and incubated for 90 minutes with a 1:2000 dilution of horseradish peroxidase labelled anti-mouse Ig. The membrane was washed twice with 0.1% Triton X100 in PBS and developed with the Opti-4CN Substrate Kit (Bio-Rad). The reaction was stopped by adding water.

S) Bactericidal assay

MC58 strain was grown overnight at 37°C on chocolate agar plates. 5-7 colonies were collected and used to inoculate 7ml Mueller-Hinton broth. The suspension was incubated at 37°C on a nutator and let to grow until OD_{620} was 0.5-0.8. The culture was aliquoted into sterile 1.5ml Eppendorf tubes and centrifuged for 20 minutes at maximum speed in a microfuge. The pellet was washed once in Gey's buffer (Gibco) and resuspended in the same buffer to an OD_{620} of 0.5, diluted 1:20000 in Gey's buffer and stored at 25°C.

50µl of Gey's buffer/1% BSA was added to each well of a 96-well tissue culture plate. 25µl of diluted mice sera (1:100 in Gey's buffer/0.2% BSA) were added to each well and the plate incubated at 4°C. 25µl of the previously described bacterial suspension were added to each well. 25µl of either heat-inactivated (56°C waterbath for 30 minutes) or normal baby rabbit complement were added to each well. Immediately after the addition of the baby rabbit complement, 22µl of each sample/well were plated on Mueller-Hinton agar plates (time 0). The 96-well plate was incubated for 1 hour at 37°C with rotation and then 22µl of each sample/well were plated on Mueller-Hinton agar plates (time 1). After overnight incubation the colonies corresponding to time 0 and time 1 hour were counted.

Table II gives a summary of the cloning, expression and purification results.

Example 1

The following partial DNA sequence was identified in *N.meningitidis* <SEQ ID 1>:

```
   1  ..ACACTGTTGT TTGCAACGGT TCAGGCAAGT GCTAACCAAT GAAGAGCAAG
  51    AAGAAGATTT ATATTTAGAC CCCGTACAAC GCACTGTTGC CGTGTTGATA
 101    GTCAATTCCG ATAAAGAAGG CACGGGAGAA AAAGAAAAAG TAGAAGAAAA
 151    TTCAGATTGG GCAGTATATT TCAACGAGAA AGGAGTACTA ACAGCCAGAG
 201    AAATCACCyT CAAAGCCGGC GACAACCTGA AAATCAAACA AAACGGCACA
 251    AACTTCACCT ACTCGCTGAA AAAAGACCTC AcAGATCTGA CCAGTGTTGG
 301    AACTGAAAAA TTATCGTTTA GCGCAAACGG CAATAAAGTC AACATcACAA
 351    GCGACACCAA AGGCTTGAAT TTTGCGAAAG AAACGGCTGG sACGAACGgC
 401    GACACCACGG TTCATCTGAA CGGTATTGGT TCGACTTTGA CCGATACGCT
 451    GCTGAATACC GGAGCGACCA CAAACGTAAC CAACGACAAC GTTACCGATG
 501    ACGAGAAAAA ACGTGCGGCA AGCGTTAAAG ACGTATTAAA CGCTGGCTGG
 551    AACATTAAAG GCGTTAAACC CGGTACAACA GCTTCCGATA ACGTTGATTT
 601    CGTCCGCACT TACGACACAG TCGAGTTCTT GAGCGCAGAT ACGAAAACAA
 651    CGACTGTTAA TGTGGAAAGC AAAGACAACG GCAAGAAAAC CGAAGTTAAA
 701    ATCGGTGCGA AGACTTCTGT TATTAAAGAA AAAGAC...
```

This corresponds to the amino acid sequence <SEQ ID 2; ORF40>:

```
   1  ..TLLFATVQAS ANQEEQEEDL YLDPVQRTVA VLIVNSDKEG TGEKEKVEEN
  51    SDWAVYFNEK GVLTAREITX KAGDNLKIKQ NGTNFTYSLK KDLTDLTSVG
 101    TEKLSFSANG NKVNITSDTK GLNFAKETAG TNGDTTVHLN GIGSTLTDTL
 151    LNTGATTNVT NDNVTDDEKK RAASVKDVLN AGWNIKGVKP GTTASDNVDF
 201    VRTYDTVEFL SADTKTTTVN VESKDNGKKT EVKIGAKTSV IKEKD...
```

Further work revealed the complete DNA sequence <SEQ ID 3>:

```
   1  ATGAACAAAA TATACCGCAT CATTTGGAAT AGTGCCCTCA ATGCCTGGGT
  51  CGTCGTATCC GAGCTCACAC GCAACCACAC CAAACGCGCC TCCGCAACCG
 101  TGAAGACCGC CGTATTGGCG ACACTGTTGT TTGCAACGGT TCAGGCAAGT
 151  GCTAACAATG AAGAGCAAGA AGAAGATTTA TATTTAGACC CCGTACAACG
 201  CACTGTTGCC GTGTTGATAG TCAATTCCGA TAAAGAAGGC ACGGGAGAAA
 251  AAGAAAAAGT AGAAGAAAAT TCAGATTGGG CAGTATATTT CAACGAGAAA
 301  GGAGTACTAA CAGCCAGAGA AATCACCCTC AAAGCCGGCG ACAACCTGAA
 351  AATCAAACAA AACGGCACAA ACTTCACCTA CTCGCTGAAA AAAGACCTCA
 401  CAGATCTGAC CAGTGTTGGA ACTGAAAAAT TATCGTTTAG CGCAAACGGC
 451  AATAAAGTCA ACATCACAAG CGACACCAAA GGCTTGAATT TTGCGAAAGA
 501  AACGGCTGGG ACGAACGGCG ACACCACGGT TCATCTGAAC GGTATTGGTT
 551  CGACTTTGAC CGATACGCTG CTGAATACCG GAGCGACCAC AAACGTAACC
 601  AACGACAACG TTACCGATGA CGAGAAAAAA CGTGCGGCAA GCGTTAAAGA
```

```
651  CGTATTAAAC GCTGGCTGGA ACATTAAAGG CGTTAAACCC GGTACAACAG
701  CTTCCGATAA CGTTGATTTC GTCCGCACTT ACGACACAGT CGAGTTCTTG
751  AGCGCAGATA CGAAAACAAC GACTGTTAAT GTGGAAAGCA AAGACAACGG
801  CAAGAAAACC GAAGTTAAAA TCGGTGCGAA GACTTCTGTT ATTAAAGAAA
851  AAGACGGTAA GTTGGTTACT GGTAAAGACA AAGGCGAGAA TGGTTCTTCT
901  ACAGACGAAG GCGAAGGCTT AGTGACTGCA AAAGAAGTGA TTGATGCAGT
951  AAACAAGGCT GGTTGGAGAA TGAAAACAAC AACCGCTAAT GGTCAAACAG
1001 GTCAAGCTGA CAAGTTTGAA ACCGTTACAT CAGGCACAAA TGTAACCTTT
1051 GCTAGTGGTA AAGGTACAAC TGCGACTGTA AGTAAAGATG ATCAAGGCAA
1101 CATCACTGTT ATGTATGATG TAAATGTCGG CGATGCCCTA AACGTCAATC
1151 AGCTGCAAAA CAGCGGTTGG AATTTGGATT CCAAAGCGGT TGCAGGTTCT
1201 TCGGGCAAAG TCATCAGCGG CAATGTTTCG CCGAGCAAGG GAAAGATGGA
1251 TGAAACCGTC AACATTAATG CCGGCAACAA CATCGAGATT ACCCGCAACG
1301 GTAAAAATAT CGACATCGCC ACTTCGATGA CCCCGCAGTT TTCCAGCGTT
1351 TCGCTCGGCG CGGGGGCGGA TGCGCCCACT TTGAGCGTGG ATGGGGACGC
1401 ATTGAATGTC GGCAGCAAGA AGGACAACAA ACCCGTCCGC ATTACCAATG
1451 TCGCCCCGGG CGTTAAAGAG GGGGATGTTA CAAACGTCGC ACAACTTAAA
1501 GGCGTGGCGC AAAACTTGAA CAACCGCATC GACAATGTGG ACGGCAACGC
1551 GCGTGCGGGC ATCGCCCAAG CGATTGCAAC CGCAGGTCTG GTTCAGGCGT
1601 ATTTGCCCGG CAAGAGTATG ATGGCGATCG GCGGCGGCAC TTATCGCGGC
1651 GAAGCCGGTT ACGCCATCGG CTACTCCAGT ATTTCCGACG GCGGAAATTG
1701 GATTATCAAA GGCACGGCTT CCGGCAATTC GCGCGGCCAT TTCGGTGCTT
1751 CCGCATCTGT CGGTTATCAG TGGTAA
```

This corresponds to the amino acid sequence <SEQ ID 4; ORF40-1>:

```
1    MNKIYRIIWN SALNAWVVVS ELTRNHTKRA SATVKTAVLA TLLFATVQAS
51   ANNEEQEEDL YLDPVQRTVA VLIVNSDKEG TGEKEKVEEN SDWAVYFNEK
101  GVLTAREITL KAGDNLKIKQ NGTNFTYSLK KDLTDLTSVG TEKLSFSANG
151  NKVNITSDTK GLNFAKETAG TNGDTTVHLN GIGSTLTDTL LNTGATTNVT
201  NDNVTDDEKK RAASVKDVLN AGWNIKGVKP GTTASDNVDF VRTYDTVEFL
251  SADTKTTTVN VESKDNGKKT EVKIGAKTSV IKEKDGKLVT GKDKGENGSS
301  TDEGEGLVTA KEVIDAVNKA GWRMKTTTAN GQTGQADKFE TVTSGTNVTF
351  ASGKGTTATV SKDDQGNITV MYDVNVGDAL NVNQLQNSGW NLDSKAVAGS
401  SGKVISGNVS PSKGKMDETV NINAGNNIEI TRNGKNIDIA TSMTPQFSSV
451  SLGAGADAPT LSVDGDALNV GSKKDNKPVR ITNVAPGVKE GDVTNVAQLK
501  GVAQNLNNRI DNVDGNARAG IAQAIATAGL VQAYLPGKSM MAIGGGTYRG
551  EAGYAIGYSS ISDGGNWIIK GTASGNSRGH FGASASVGYQ W*
```

Further work identified the corresponding gene in strain A of *N.meningitidis* <SEQ ID 5 >:

```
1    ATGAACAAAA TATACCGCAT CATTTGGAAT AGTGCCCTCA ATGCCTGNGT
51   CGCCGTATCC GAGCTCACAC GCAACCACAC CAAACGCGCC TCCGCAACCG
101  TGAAGACCGC CGTATTGGCG ACACTGTTGT TTGCAACGGT TCAGGCGAAT
151  GCTACCGATG AAGATGAAGA AGAAGAGTTA GAATCCGTAC AACGCTCTGT
201  CGTAGGGAGC ATTCAAGCCA GTATGGAAGG CAGCGGCGAA TTGGAAACGA
251  TATCATTATC AATGACTAAC GACAGCAAGG AATTTGTAGA CCCATACATA
301  GTAGTTACCC TCAAAGCCGG CGACAACCTG AAAATCAAAC AAAACACCAA
351  TGAAAACACC AATGCCAGTA GCTTCACCTA CTCGCTGAAA AAAGACCTCA
401  CAGGCCTGAT CAATGTTGAN ACTGAAAAAT TATCGTTTGG CGCAAACGGC
451  AAGAAAGTCA ACATCATAAG CGACACCAAA GGCTTGAATT TCGCGAAAGA
501  AACGGCTGGG ACGAACGGCG ACACCACGGT TCATCTGAAC GGTATCGGTT
551  CGACTTTGAC CGATACGCTT GCGGGTTCTT CTGCTTCTCA CGTTGATGCG
601  GGTAACCNAA GTACACATTA CACTCGTGCA GCAAGTATTA AGGATGTGTT
651  GAATGCGGGT TGGAATATTA AGGGTGTTAA ANNNGGCTCA ACAACTGGTC
701  AATCAGAAAA TGTCGATTTC GTCCGCACTT ACGACACAGT CGAGTTCTTG
751  AGCGCAGATA CGNAAACAAC GACNGTTAAT GTGGAAAGCA AAGACAACGG
801  CAAGAGAACC GAAGTTAAAA TCGGTGCGAA GACTTCTGTT ATTAAAGAAA
851  AAGACGGTAA GTTGGTTACT GGTAAAGGCA AAGGCGAGAA TGGTTCTTCT
901  ACAGACGAAG GCGAAGGCTT AGTGACTGCA AAAGAAGTGA TTGATGCAGT
951  AAACAAGGCT GGTTGGAGAA TGAAAACAAC AACCGCTAAT GGTCAAACAG
1001 GTCAAGCTGA CAAGTTTGAA ACCGTTACAT CAGGCACAAA TGTAACCTTT
1051 GCTAGTGGTA AAGGTACAAC TGCGACTGTA AGTAAAGATG ATCAAGGCAA
1101 CATCACTGTT ATGTATGATG TAAATGTCGG CGATGCCCTA AACGTCAATC
1151 AGCTGCAAAA CAGCGGTTGG AATTTGGATT CCAAAGCGGT TGCAGGTTCT
1201 TCGGGCAAAG TCATCAGCGG CAATGTTTCG CCGAGCAAGG GAAAGATGGA
1251 TGAAACCGTC AACATTAATG CCGGCAACAA CATCGAGATT AGCCGCAACG
1301 GTAAAAATAT CGACATCGCC ACTTCGATGG CGCCGCAGTT TTCCAGCGTT
1351 TCGCTCGGCG CGGGGGCAGA TGCGCCCACT TTAAGCGTGG ATGACGAGGG
1401 CGCGTTGAAT GTCGGCAGCA AGGATGCCAA CAAACCCGTC CGCATTACCA
```

```
1451  ATGTCGCCCC GGGCGTTAAA GANGGGGATG TTACAAACGT CNCACAACTT
1501  AAAGGCGTGG CGCAAAACTT GAACAACCGC ATCGACAATG TGGACGGCAA
1551  CGCGCGTGCN GGCATCGCCC AAGCGATTGC AACCGCAGGT CTGGTTCAGG
1601  CGTATCTGCC CGGCAAGAGT ATGATGGCGA TCGGCGGCGG CACTTATCGC
1651  GGCGAAGCCG GTTACGCCAT CGGCTACTCC AGTATTTCCG ACGGCGGAAA
1701  TTGGATTATC AAAGGCACGG CTTCCGGCAA TTCGCGCGGC CATTTCGGTG
1751  CTTCCGCATC TGTCGGTTAT CAGTGGTAA
```

This encodes a protein having amino acid sequence <SEQ ID 6; ORF40a>:

```
   1  MNKIYRIIWN SALNAXVAVS ELTRNHTKRA SATVKTAVLA TLLFATVQAN
  51  ATDEDEEEEL ESVQRSVVGS IQASMEGSGE LETISLSMTN DSKEFVDPYI
 101  VVTLKAGDNL KIKQNTNENT NASSFTYSLK KDLTGLINVX TEKLSFGANG
 151  KKVNIISDTK GLNFAKETAG TNGDTTVHLN GIGSTLTDTL AGSSASHVDA
 201  GNXSTHYTRA ASIKDVLNAG WNIKGVKXGS TTGQSENVDF VRTYDTVEFL
 251  SADTXTTTVN VESKDNGKRT EVKIGAKTSV IKEKDGKLVT GKGKGENGSS
 301  TDEGEGLVTA KEVIDAVNKA GWRMKTTTAN GQTGQADKFE TVTSGTNVTF
 351  ASGKGTTATV SKDDQGNITV MYDVNVGDAL NVNQLQNSGW NLDSKAVAGS
 401  SGKVISGNVS PSKGKMDETV NINAGNNIEI SRNGKNIDIA TSMAPQFSSV
 451  SLGAGADAPT LSVDDEGALN VGSKDANKPV RITNVAPGVK XGDVTNVXQL
 501  KGVAQNLNNR IDNVDGNARA GIAQAIATAG LVQAYLPGKS MMAIGGGTYR
 551  GEAGYAIGYS SISDGGNWII KGTASGNSRG HFGASASVGY QW*
```

The originally-identified partial strain B sequence (ORF40) shows 65.7% identity over a 254aa overlap with ORF40a:

```
                                                  10        20        30
orf40.pep                                        TLLFATVQASANQEEQEEDLYLDPVQRTVA
                                                 ||||||||||:|::|::||:|  : |||:|
orf40a       SALNAXVAVSELTRNHTKRASATVKTAVLATLLFATVQANATDEDEEEEL--ESVQRSV-
                    20        30        40        50        60

                   40        50        60        70        80
orf40.pep    VLIVNSDKEGTGEKEKVEEN-SDWAVYFNEKGVLTAREITXKAGDNLKIKQN------GT
             |  :::: ||:|| | :  : :: :  | :  ::    :| ||||||||||       ::
orf40a       VGSIQASMEGSGELETISLSMTNDSKEFVDPYIV----VTLKAGDNLKIKQNTNENTNAS
              70        80        90        100           110       120

                  90        100       110       120       130       140
orf40.pep    NFTYSLKKDLTDLTSVGTEKLSFSANGNKVNITSDTKGLNFAKETAGTNGDTTVHLNGIG
             :|||||||||| | :| ||||||:|||:|||| ||||||||||||||||||||||||||
orf40a       SFTYSLKKDLTGLINVXTEKLSFGANGKKVNIISDTKGLNFAKETAGTNGDTTVHLNGIG
                 130       140       150       160       170       180

                 150       160       170       180       190       200
orf40.pep    STLTDTLLNTGATTNVTNDNVTDDEKKRAASVKDVLNAGWNIKGVKPGTTA--SDNVDFV
             ||||||| :::|: :|   | :  :  ||||:|||||||||||||| |:|:  |:|||||
orf40a       STLTDTLAGSSAS-HVDAGNXST-HYTRAASIKDVLNAGWNIKGVKXGSTTGQSENVDFV
                 190       200       210       220       230       240

                   210       220       230       240
orf40.pep    RTYDTVEFLSADTKTTTVNVESKDNGKKTEVKIGAKTSVIKEKD
             |||||||||||| ||||||||||||:||||||||||||||||
orf40a       RTYDTVEFLSADTXTTTVNVESKDNGKRTEVKIGAKTSVIKEKDGKLVTGKGKGENGSST
                   250       260       270       280       290       300
```

The complete strain B sequence (ORF40-1) and ORF40a show 83.7% identity in 601 aa overlap:

```
                  10        20        30        40        50        60
orf40-1.pep   MNKIYRIIWNSALNAWVVVSELTRNHTKRASATVKTAVLATLLFATVQASANNEEQEEDL
              |||||||||||||||| |:|||||||||||||||||||||||||||||||:|::|::||:|
orf40a        MNKIYRIIWNSALNAXVAVSELTRNHTKRASATVKTAVLATLLFATVQANATDEDEEEEL
                  10        20        30        40        50        60

                  70        80        90        100       110       119
orf40-1.pep   YLDPVQRTVAVLIVNSDKEGTGEKEKVEEN-SDWAVYFNEKGVLTAREITLKAGDNLKIK
               : |||:| |  :::: ||:|| | :  : :: :  | :  ::    :|||||||||||
```

```
orf40a          --ESVQRSV-VGSIQASMEGSGELETISLSMTNDSKEFVDPYIV----VTLKAGDNLKIK
                              70        80        90       100       110

               120       130       140       150       160       170
orf40-1.pep     QN------GTNFTYSLKKDLTDLTSVGTEKLSFSANGNKVNITSDTKGLNFAKETAGTNG
                ||      :::||||||||||| | :| ||||||:|||:|||| ||||||||||||||||
orf40a          QNTNENTNASSFTYSLKKDLTGLINVXTEKLSFGANGKKVNIISDTKGLNFAKETAGTNG
                    120       130       140       150       160       170

                    180       190       200       210       220       230
orf40-1.pep     DTTVHLNGIGSTLTDTLLNTGATTNVTNDNVTDDEKKRAASVKDVLNAGWNIKGVKPGTT
                |||||||||||||||||| :::|: :|   | .  :  ||||:|||||||||||||| |:|
orf40a          DTTVHLNGIGSTLTDTLAGSSAS-HVDAGNXST-HYTRAASIKDVLNAGWNIKGVKXGST
                    180       190       200       210       220       230

                    240       250       260       270       280       290
orf40-1.pep     A--SDNVDFVRTYDTVEFLSADTKTTTVNVESKDNGKKTEVKIGAKTSVIKEKDGKLVTG
                :  |:|||||||||||||||||| ||||||||||||:||||||||||||||||||||||
orf40a          TGQSENVDFVRTYDTVEFLSADTXTTTVNVESKDNGKRTEVKIGAKTSVIKEKDGKLVTG
                    240       250       260       270       280       290

                    300       310       320       330       340       350
orf40-1.pep     KDKGENGSSTDEGEGLVTAKEVIDAVNKAGWRMKTTTANGQTGQADKFETVTSGTNVTFA
                | ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
orf40a          KGKGENGSSTDEGEGLVTAKEVIDAVNKAGWRMKTTTANGQTGQADKFETVTSGTNVTFA
                    300       310       320       330       340       350

                    360       370       380       390       400       410
orf40-1.pep     SGKGTTATVSKDDQGNITVMYDVNVGDALNVNQLQNSGWNLDSKAVAGSSGKVISGNVSP
                ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
orf40a          SGKGTTATVSKDDQGNITVMYDVNVGDALNVNQLQNSGWNLDSKAVAGSSGKVISGNVSP
                    360       370       380       390       400       410

                    420       430       440       450       460       470
orf40-1.pep     SKGKMDETVNINAGNNIEITRNGKNIDIATSMTPQFSSVSLGAGADAPTLSVDGD-ALNV
                ||||||||||||||||||:||||||||||||:|||||||||||||||||||| : ||||
orf40a          SKGKMDETVNINAGNNIEISRNGKNIDIATSMAPQFSSVSLGAGADAPTLSVDDEGALNV
                    420       430       440       450       460       470

                     480       490       500       510       520       530
orf40-1.pep     GSKKDNKPVRITNVAPGVKEGDVTNVAQLKGVAQNLNNRIDNVDGNARAGIAQAIATAGL
                |||  |||||||||||||| |||||| ||||||||||||||||||||||||||||||||||
orf40a          GSKDANKPVRITNVAPGVKXGDVTNVXQLKGVAQNLNNRIDNVDGNARAGIAQAIATAGL
                    480       490       500       510       520       530

                    540       550       560       570       580       590
orf40-1.pep     VQAYLPGKSMMAIGGGTYRGEAGYAIGYSSISDGGNWIIKGTASGNSRGHFGASASVGYQ
                ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
orf40a          VQAYLPGKSMMAIGGGTYRGEAGYAIGYSSISDGGNWIIKGTASGNSRGHFGASASVGYQ
                    540       550       560       570       580       590

orf40-1.pep     WX
                ||
orf40a          WX
```

Computer analysis of these amino acid sequences gave the following results:

Homology with Hsf protein encoded by the type b surface fibrils locus of *H.influenzae* (accession number U41852)

ORF40 and Hsf protein show 54% aa identity in 251 aa overlap:

```
Orf40   1   TLLFATVQASANQEEQEEDLYLDPVQRTVAVLIVNSDXXXXXXXXXXXXXNSDWAVYFNEK 60
            TLLFATVQA+A  E++E    LDPV RT  VL  +SD            NS+W +YF+ K
Hsf     41  TLLFATVQANATDEDEE----LDPVVRTAPVLSFHSDKEGTGEKEVTE-NSNWGIYFDNK 95

Orf40   61  GVLTAREITXKAGDNLKIKQN------GTNFTYSLKKDLTDLTSVGTEKLSFSANGNKVN 114
            GVL A  IT KAGDNLKIKQN       ++FTYSLKKDLTDLTSV TEKLSF ANG+KV+
Hsf     96  GVLKAGAITLKAGDNLKIKQNTDESTNASSFTYSLKKDLTDLTSVATEKLSFGANGDKVD 155
```

```
Orf40   115 ITSDTKGLNFAKETAGTNGDTTVHLNGIGSTLTDTLLNTGAXXXXXXXXXXXXXEKKRAAS 174
            ITSD  GL  AK      G+  VHLNG+ STL D + NTG             EK RAA+
Hsf     156 ITSDANGLKLAK-----TGNGNVHLNGLDSTLPDAVTNTGVLSSSSFTPNDV-EKTRAAT 209

Orf40   175 VKDVLNAGWNIKGVKPGTTASDNVDFVRTYDTVEFLSADTKTTTVNVESKDNGKKTEVKI 234
            VKDVLNAGWNIKG K      ++VD V  Y+ VEF++ D  T  V + +K+NGK TEVK
Hsf     210 VKDVLNAGWNIKGAKTAGGNVESVDLVSAYNNVEFITGDKNTLDVVLTAKENGKTTEVKF 269

Orf40   235 GAKTSVIKEKD 245
              KTSVIKEKD
Hsf     270 TPKTSVIKEKD 280
```

ORF40a also shows homology to Hsf:

```
gi|1666683 (U41852) hsf gene product [Haemophilus influenzae] Length = 2353
 Score = 153 (67.7 bits), Expect = 1.5e-116, Sum P(11) = 1.5e-116
 Identities = 33/36 (91%), Positives = 34/36 (94%)

Query:    16 VAVSELTRNHTKRASATVKTAVLATLLFATVQANAT 51
             V VSELTR HTKRASATV+TAVLATLLFATVQANAT
Sbjct:    17 VVVSELTRTHTKRASATVETAVLATLLFATVQANAT 52

 Score = 161 (71.2 bits), Expect = 1.5e-116, Sum P(11) = 1.5e-116
 Identities = 32/38 (84%), Positives = 36/38 (94%)

Query:   101 VTLKAGDNLKIKQNTNENTNASSFTYSLKKDLTGLINV 138
             +TLKAGDNLKIKQNT+E+TNASSFTYSLKKDLT L +V
Sbjct:   103 ITLKAGDNLKIKQNTDESTNASSFTYSLKKDLTDLTSV 140

 Score = 110 (48.7 bits), Expect = 1.5e-116, Sum P(11) = 1.5e-116
 Identities = 21/29 (72%), Positives = 25/29 (86%)

Query:   138 VTEKLSFGANGKKVNIISDTKGLNFAKET 166
             V++KLS G NG KVNI SDTKGLNFAK++
Sbjct:  1439 VSDKLSLGTNGNKVNITSDTKGLNFAKDS 1467

 Score = 85 (37.6 bits), Expect = 1.5e-116, Sum P(11) = 1.5e-116
 Identities = 18/32 (56%), Positives = 20/32 (62%)

Query:   169 TNGDTTVHLNGIGSTLTDTLAGSSASHVDAGN 200
             T  D  +HLNGI STLTDTL  S A+    GN
Sbjct:  1469 TGDDANIHLNGIASTLTDTLLNSGATTNLGGN 1500

 Score = 92 (40.7 bits), Expect = 1.5e-116, Sum P(11) = 1.5e-116
 Identities = 16/19 (84%), Positives = 19/19 (100%)

Query:   206 RAASIKDVLNAGWNIKGVK 224
             RAAS+KDVLNAGWN++GVK
Sbjct:  1509 RAASVKDVLNAGWNVRGVK 1527

 Score = 90 (39.8 bits), Expect = 1.5e-116, Sum P(11) = 1.5e-116
 Identities = 17/28 (60%), Positives = 20/28 (71%)

Query:   226 STTGQSENVDFVRTYDTVEFLSADTTTT 253
             S   Q EN+DFV TYDTV+F+S D  TT
Sbjct:  1530 SANNQVENIDFVATYDTVDFVSGDKDTT 1557
```

Based on homology with Hsf, it was predicted that this protein from *N.meningitidis*, and its epitopes, could be useful antigens for vaccines or diagnostics.

ORF40-1 (61kDa) was cloned in pET and pGex vectors and expressed in *E.coli*, as described above. The products of protein expression and purification were analyzed by SDS-PAGE. Figure 1A shows the results of affinity purification of the His-fusion protein, and Figure 1B shows the

results of expression of the GST-fusion in *E coli*. Purified His-fusion protein was used to immunise mice, whose sera were used for FACS analysis (Figure 1C), a bactericidal assay (Figure 1D), and ELISA (positive result) These experiments confirm that ORF40-1 is a surface-exposed protein, and that it is a useful immunogen.

Figure 1E shows plots of hydrophilicity, antigenic index, and AMPHI regions for ORF40-1.

Example 2

The following partial DNA sequence was identified in *N.meningitidis* <SEQ ID 7>

```
   1  ATGTTACGTt TGACTGCtTT AGCCGTATGC ACCGCCCTCG CTTTGGGCGC
  51  GTGTTCGCCG CAAAATTCCG ACTCTGCCCC ACAAGCCAAA GaACAGGCGG
 101  TTTCCGCCGC ACAAACCGAA GgCGCGTCCG TTACCGTCAA AACCGCGCGC
 151  GGCGACGTTC AAATACCGCA AAACCCCGAA CGCATCGCCG TTTACGATTT
 201  GGGTATGCTC GACACCTTGA GCAAACTGGG CGTGAAAACC GGTTTGTCCG
 251  TCGATAAAAA CCGCCTGCCG TATTTAGAGG AATATTTCAA AACGACAAAA
 301  CCTGCCGGCA CTTTGTTCGA GCCGGATTAC GAAACGCTCA ACGCTTACAA
 351  ACCGCAGCTC ATCATCATCG GCAGCCGCGC CgCCAAGGCG TTTGACAAAT
 401  TGAAcGAAAT CGCGCCGACC ATCGrmwTGA CCGCCGATAC CGCCAACCTC
 451  AAAGAAAGTG CCAArGAGGC ATCGACGCTG GCGCAAATCT TC..
```

This corresponds to the amino acid sequence <SEQ ID 8; ORF38>:

```
   1  MLRLTALAVC TALALGACSP QNSDSAPQAK EQAVSAAQTE GASVTVKTAR
  51  GDVQIPQNPE RIAVYDLGML DTLSKLGVKT GLSVDKNRLP YLEEYFKTTK
 101  PAGTLFEPDY ETLNAYKPQL IIIGSRAAKA FDKLNEIAPT IXXTADTANL
 151  KESAKEASTL AQIF..
```

Further work revealed the complete nucleotide sequence <SEQ ID 9>:

```
   1  ATGTTACGTT TGACTGCTTT AGCCGTATGC ACCGCCCTCG CTTTGGGCGC
  51  GTGTTCGCCG CAAAATTCCG ACTCTGCCCC ACAAGCCAAA GAACAGGCGG
 101  TTTCCGCCGC ACAAACCGAA GGCGCGTCCG TTACCGTCAA AACCGCGCGC
 151  GGCGACGTTC AAATACCGCA AAACCCCGAA CGCATCGCCG TTTACGATTT
 201  GGGTATGCTC GACACCTTGA GCAAACTGGG CGTGAAAACC GGTTTGTCCG
 251  TCGATAAAAA CCGCCTGCCG TATTTAGAGG AATATTTCAA AACGACAAAA
 301  CCTGCCGGCA CTTTGTTCGA GCCGGATTAC GAAACGCTCA ACGCTTACAA
 351  ACCGCAGCTC ATCATCATCG GCAGCCGCGC CGCCAAGGCG TTTGACAAAT
 401  TGAACGAAAT CGCGCCGACC ATCGAAATGA CCGCCGATAC CGCCAACCTC
 451  AAAGAAAGTG CCAAAGAGCG CATCGACGCG CTGGCGCAAA TCTTCGGCAA
 501  ACAGGCGGAA GCCGACAAGC TGAAGGCGGA AATCGACGCG TCTTTTGAAG
 551  CCGCGAAAAC TGCCGCACAA GGTAAGGGCA AAGGTTTGGT GATTTTGGTC
 601  AACGGCGGCA AGATGTCGGC TTTCGGCCCG TCTTCACGCT TGGGCGGCTG
 651  GCTGCACAAA GACATCGGCG TTCCCGCTGT CGATGAATCA ATTAAAGAAG
 701  GCAGCCACGG TCAGCCTATC AGCTTTGAAT ACCTGAAAGA GAAAAATCCC
 751  GACTGGCTGT TTGTCCTTGA CCGAAGCGCG GCCATCGGCG AAGAGGGTCA
 801  GGCGGCGAAA GACGTGTTGG ATAATCCGCT GGTTGCCGAA ACAACCGCTT
 851  GGAAAAAAGG ACAGGTCGTG TACCTCGTTC CTGAAACTTA TTTGGCAGCC
 901  GGTGGCGCGC AAGAGCTGCT GAATGCAAGC AAACAGGTTG CCGACGCTTT
 951  TAACGCGGCA AAATAA
```

This corresponds to the amino acid sequence <SEQ ID 10; ORF38-1>:

```
   1  MLRLTALAVC TALALGACSP QNSDSAPQAK EQAVSAAQTE GASVTVKTAR
  51  GDVQIPQNPE RIAVYDLGML DTLSKLGVKT GLSVDKNRLP YLEEYFKTTK
 101  PAGTLFEPDY ETLNAYKPQL IIIGSRAAKA FDKLNEIAPT IEMTADTANL
 151  KESAKERIDA LAQIFGKQAE ADKLKAEIDA SFEAAKTAAQ GKGKGLVILV
 201  NGGKMSAFGP SSRLGGWLHK DIGVPAVDES IKEGSHGQPI SFEYLKEKNP
 251  DWLFVLDRSA AIGEEGQAAK DVLDNPLVAE TTAWKKGQVV YLVPETYLAA
```

```
301  GGAQELLNAS KQVADAFNAA K*
```

Computer analysis of this amino acid sequence reveals a putative prokaryotic membrane lipoprotein lipid attachment site (underlined).

Further work identified the corresponding gene in strain A of *N.meningitidis* <SEQ ID 11>:

```
  1  ATGTTACGTT TGACTGCTTT AGCCGTATGC ACCGCCCTCG CTTTGGGCGC
 51  GTGTTCGCCG CAAAATTCCG ACTCTGCCCC ACAAGCCAAA GAACAGGCGG
101  TTTCCGCCGC ACAATCCGAA GGCGTGTCCG TTACCGTCAA AACGGCGCGC
151  GGCGATGTTC AAATACCGCA AAACCCCGAA CGTATCGCCG TTTACGATTT
201  GGGTATGCTC GACACCTTGA GCAAACTGGG CGTGAAAACC GGTTTGTCCG
251  TCGATAAAAA CCGCCTGCCG TATTTAGAGG AATATTTCAA AACGACAAAA
301  CCTGCCGGAA CTTTGTTCGA GCCGGATTAC GAAACGCTCA ACGCTTACAA
351  ACCGCAGCTC ATCATCATCG GCAGCCGCGC AGCCAAAGCG TTTGACAAAT
401  TGAACGAAAT CGCGCCGACC ATCGAAATGA CCGCCGATAC CGCCAACCTC
451  AAAGAAAGTG CCAAAGAGCG TATCGACGCG CTGGCGCAAA TCTTCGGCAA
501  AAAGGCGGAA GCCGACAAGC TGAAGGCGGA AATCGACGCG TCTTTTGAAG
551  CCGCGAAAAC TGCCGCGCAA GGCAAAGGCA AGGGTTTGGT GATTTTGGTC
601  AACGGCGGCA AGATGTCCGC CTTCGGCCCG TCTTCACGAC TGGGCGGCTG
651  GCTGCACAAA GACATCGGCG TTCCCGCTGT TGACGAAGCC ATCAAAGAAG
701  GCAGCCACGG TCAGCCTATC AGCTTTGAAT ACCTGAAAGA GAAAAATCCC
751  GACTGGCTGT TTGTCCTTGA CCGCAGCGCG GCCATCGGCG AAGAGGGTCA
801  GGCGGCGAAA GACGTGTTGA ACAATCCGCT GGTTGCCGAA ACAACCGCTT
851  GGAAAAAAGG ACAAGTCGTT TACCTTGTTC CTGAAACTTA TTTGGCAGCC
901  GGTGGCGCGC AAGAGCTACT GAATGCAAGC AAACAGGTTG CCGACGCTTT
951  TAACGCGGCA AAATAA
```

This encodes a protein having amino acid sequence <SEQ ID 12; ORF38a>:

```
  1  MLRLTALAVC TALALGACSP QNSDSAPQAK EQAVSAAQSE GVSVTVKTAR
 51  GDVQIPQNPE RIAVYDLGML DTLSKLGVKT GLSVDKNRLP YLEEYFKTTK
101  PAGTLFEPDY ETLNAYKPQL IIIGSRAAKA FDKLNEIAPT IEMTADTANL
151  KESAKERIDA LAQIFGKKAE ADKLKAEIDA SFEAAKTAAQ GKGKGLVILV
201  NGGKMSAFGP SSRLGGWLHK DIGVPAVDEA IKEGSHGQPI SFEYLKEKNP
251  DWLFVLDRSA AIGEEGQAAK DVLNNPLVAE TTAWKKGQVV YLVPETYLAA
301  GGAQELLNAS KQVADAFNAA K*
```

The originally-identified partial strain B sequence (ORF38) shows 95.2% identity over a 165aa overlap with ORF38a:

```
                10        20        30        40        50        60
orf38.pep   MLRLTALAVCTALALGACSPQNSDSAPQAKEQAVSAAQTEGASVTVKTARGDVQIPQNPE
            ||||||||||||||||||||||||||||||||||||||||||:||:||||||||||||||||||
orf38a      MLRLTALAVCTALALGACSPQNSDSAPQAKEQAVSAAQSEGVSVTVKTARGDVQIPQNPE
                10        20        30        40        50        60

                70        80        90       100       110       120
orf38.pep   RIAVYDLGMLDTLSKLGVKTGLSVDKNRLPYLEEYFKTTKPAGTLFEPDYETLNAYKPQL
            ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
orf38a      RIAVYDLGMLDTLSKLGVKTGLSVDKNRLPYLEEYFKTTKPAGTLFEPDYETLNAYKPQL
                70        80        90       100       110       120

               130       140       150       160
orf38.pep   IIIGSRAAKAFDKLNEIAPTIXXTADTANLKESAKE-ASTLAQIF
            ||||||||||||||||||||  ||||||||||||  ::||||||
orf38a      IIIGSRAAKAFDKLNEIAPTIEMTADTANLKESAKERIDALAQIFGKKAEADKLKAEIDA
               130       140       150       160       170       180

orf38a      SFEAAKTAAQGKGKGLVILVNGGKMSAFGPSSRLGGWLHKDIGVPAVDEAIKEGSHGQPI
               190       200       210       220       230       240
```

The complete strain B sequence (ORF38-1) and ORF38a show 98.4% identity in 321 aa overlap:

```
orf38a.pep      MLRLTALAVCTALALGACSPQNSDSAPQAKEQAVSAAQSEGVSVTVKTARGDVQIPQNPE
                ||||||||||||||||||||||||||||||||||||||:||:||||||||||||||||||
orf38-1         MLRLTALAVCTALALGACSPQNSDSAPQAKEQAVSAAQTEGASVTVKTARGDVQIPQNPE

orf38a.pep      RIAVYDLGMLDTLSKLGVKTGLSVDKNRLPYLEEYFKTTKPAGTLFEPDYETLNAYKPQL
                ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
orf38-1         RIAVYDLGMLDTLSKLGVKTGLSVDKNRLPYLEEYFKTTKPAGTLFEPDYETLNAYKPQL

orf38a.pep      IIIGSRAAKAFDKLNEIAPTIEMTADTANLKESAKERIDALAQIFGKKAEADKLKAEIDA
                |||||||||||||||||||||||||||||||||||||||||||||||:|||||||||||||
orf38-1         IIIGSRAAKAFDKLNEIAPTIEMTADTANLKESAKERIDALAQIFGKQAEADKLKAEIDA

orf38a.pep      SFEAAKTAAQGKGKGLVILVNGGKMSAFGPSSRLGGWLHKDIGVPAVDEAIKEGSHGQPI
                |||||||||||||||||||||||||||||||||||||||||||||||||:||||||||||
orf38-1         SFEAAKTAAQGKGKGLVILVNGGKMSAFGPSSRLGGWLHKDIGVPAVDESIKEGSHGQPI

orf38a.pep      SFEYLKEKNPDWLFVLDRSAAIGEEGQAAKDVLNNPLVAETTAWKKGQVVYLVPETYLAA
                |||||||||||||||||||||||||||||||||||:|||||||||||||||||||||||||
orf38-1         SFEYLKEKNPDWLFVLDRSAAIGEEGQAAKDVLDNPLVAETTAWKKGQVVYLVPETYLAA

orf38a.pep      GGAQELLNASKQVADAFNAAK
                |||||||||||||||||||||
orf38-1         GGAQELLNASKQVADAFNAAK
```

Computer analysis of these sequences revealed the following:

Homology with a lipoprotein (lipo) of *C.jejuni* (accession number X82427)

ORF38 and lipo show 38% aa identity in 96 aa overlap:

```
Orf38: 40  EGASVTVKTARGDVQIPQNPERIAVYDLGMLDTLSKLGVKTGLS-VDKNRLPYLEEYFKT 98
           EG S  VK + G+ + P+NP ++ + DLG+LDT   L +   ++ V    LP   + FK
Lipo:  51  EGDSFLVKDSLGENKTPKNPSKVVILDLGILDTFDALKLNDKVAGVPAKNLPKYLQQFKN 110

Orf38: 99  TKPAGTLFEPDYETLNAYKPQLIIIGSRAAKAFDKL 134
               G + + D+E +NA KP LIII  R +K +DKL
Lipo:  111 KPSVGGVQQVDFEAINALKPDLIIISGRQSKFYDKL 146
```

Based on this analysis, it was predicted that this protein from *N.meningitidis*, and its epitopes, could be useful antigens for vaccines or diagnostics.

ORF38-1 (32kDa) was cloned in pET and pGex vectors and expressed in *E.coli*, as described above. The products of protein expression and purification were analyzed by SDS-PAGE. Figure 2A shows the results of affinity purification of the His-fusion protein, and Figure 2B shows the results of expression of the GST-fusion in *E.coli*. Purified His-fusion protein was used to immunise mice, whose sera were used for Western blot analysis (Figure 2C) and FACS analysis (Figure 2D). These experiments confirm that ORF38-1 is a surface-exposed protein, and that it is a useful immunogen.

Figure 2E shows plots of hydrophilicity, antigenic index, and AMPHI regions for ORF38-1.

Example 3

The following *N.meningitidis* DNA sequence was identified <SEQ ID 13>:

```
   1  ATGAAACTTC TGACCACCGC AATCCTGTCT TCCGCAATCG CGCTCAGCAG
  51  TATGGCTGCC GCCGCTGGCA CGGACAACCC CACTGTTGCA AAAAAAACCG
 101  TCAGCTACGT CTGCCAGCAA GGTAAAAAAG TCAAAGTAAC CTACGGCTTC
 151  AACAAACAGG GTCTGACCAC ATACGCTTCC GCCGTCATCA ACGGCAAACG
 201  CGTGCAAATG CCTGTCAATT TGGACAAATC CGACAATGTG GAAACATTCT
 251  ACGGCAAAGA AGGCGGTTAT GTTTTGGGTA CCGGCGTGAT GGATGGCAAA
 301  TCCTACCGCA AACAGCCCAT TATGATTACC GCACCTGACA ACCAAATCGT
 351  CTTCAAAGAC TGTTCCCCAC GTTAA
```

This corresponds to the amino acid sequence <SEQ ID 14; ORF44>:

```
   1  MKLLTTAILS SAIALSSMAA AAGTDNPTVA KKTVSYVCQQ GKKVKVTYGF
  51  NKQGLTTYAS AVINGKRVQM PVNLDKSDNV ETFYGKEGGY VLGTGVMDGK
 101  SYRKQPIMIT APDNQIVFKD CSPR*
```

Computer analysis of this amino acid sequence predicted the leader peptide shown underlined.

Further work identified the corresponding gene in strain A of *N.meningitidis* <SEQ ID 15>:

```
   1  ATGAAACTTC TGACCACCGC AATCCTGTCT TCCGCAATCG CGCTCAGCAG
  51  TATGGCTGCT GCTGCCGGCA CGAACAACCC CACCGTTGCC AAAAAAACCG
 101  TCAGCTACGT CTGCCAGCAA GGTAAAAAAG TCAAAGTAAC CTACGGCTTT
 151  AACAAACAGG GCCTGACCAC ATACGCTTCC GCCGTCATCA ACGGCAAACG
 201  TGTGCAAATG CCTGTCAATT TGGACAAATC CGACAATGTG GAAACATTCT
 251  ACGGCAAAGA AGGCGGTTAT GTTTTGGGTA CCGGCGTGAT GGATGGCAAA
 301  TCCTATCGCA AACAGCCTAT TATGATTACC GCACCTGACA ACCAAATCGT
 351  CTTCAAAGAC TGTTCCCCAC GTTAA
```

This encodes a protein having amino acid sequence <SEQ ID 16; ORF44a>:

```
   1  MKLLTTAILS SAIALSSMAA AAGTNNPTVA KKTVSYVCQQ GKKVKVTYGF
  51  NKQGLTTYAS AVINGKRVQM PVNLDKSDNV ETFYGKEGGY VLGTGVMDGK
 101  SYRKQPIMIT APDNQIVFKD CSPR*
```

The strain B sequence (ORF44) shows 99.2% identity over a 124aa overlap with ORF44a:

```
                10        20        30        40        50        60
orf44.pep       MKLLTTAILSSAIALSSMAAAAGTDNPTVAKKTVSYVCQQGKKVKVTYGFNKQGLTTYAS
                ||||||||||||||||||||||||:|||||||||||||||||||||||||||||||||||
orf44a          MKLLTTAILSSAIALSSMAAAAGTNNPTVAKKTVSYVCQQGKKVKVTYGFNKQGLTTYAS
                10        20        30        40        50        60

                70        80        90       100       110       120
orf44.pep       AVINGKRVQMPVNLDKSDNVETFYGKEGGYVLGTGVMDGKSYRKQPIMITAPDNQIVFKD
                ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
orf44a          AVINGKRVQMPVNLDKSDNVETFYGKEGGYVLGTGVMDGKSYRKQPIMITAPDNQIVFKD
                70        80        90       100       110       120

orf44.pep       CSPRX
                |||||
orf44a          CSPRX
```

Computer analysis gave the following results:

Homology with the LecA adhesin of *Eikenella corrodens* (accession number D78153)

ORF44 and LecA protein show 45% aa identity in 91 aa overlap:

```
Orf44   33  TVSYVCQQGKKVKVTYGFNKQGLTTYASAVINGKRVQMPVNLDKSDNVETFYGKEGGYVL 92
            +V+YVCQQG+++ V Y FN  G+ T A   +N + +++P NL  SDNV+T +    GY L
LecA   135  SVAYVCQQGRRLNVNYRFNSAGVPTSAELRVNNRNLRLPYNLSASDNVDTVF-SANGYRL 193

Orf44   93  GTGVMDGKSYRKQPIMITAPDNQIVFKDCSP 123
             T  MD  +YR Q I+++AP+ Q+++KDCSP
```

```
LecA   194 TTNAMDSANYRSQDIIVSAPNGQMLYKDCSP 224
```

Based on homology with the adhesin, it was predicted that this protein from *N.meningitidis*, and its epitopes, could be useful antigens for vaccines or diagnostics.

ORF44-1 (11.2kDa) was cloned in pET and pGex vectors and expressed in *E.coli*, as described above. The products of protein expression and purification were analyzed by SDS-PAGE. Figure 3A shows the results of affinity purification of the His-fusion protein, and Figure 3B shows the results of expression of the GST-fusion in *E.coli*. Purified His-fusion protein was used to immunise mice, whose sera were used for ELISA, which gave positive results, and for a bactericidal assay (Figure 3C). These experiments confirm that ORF44-1 is a surface-exposed protein, and that it is a useful immunogen.

Figure 3D shows plots of hydrophilicity, antigenic index, and AMPHI regions for ORF44-1.

Example 4

The following partial DNA sequence was identified in *N.meningitidis* <SEQ ID 17>

```
   1  ..GGCACCGAAT TCAAAACCAC CCTTTCCGGA GCCGACATAC AGGCAGGGGT
  51    GGGTGAAAAA GCCCGAGCCG ATGCGAAAAT TATCCTAAAA GGCATCGTTA
 101    ACCGCATCCA AACCGAAGAA AAGCTGGAAT CCAACTCGAC CGTATGGCAA
 151    AAGCAGGCCG GAAGCGGCAG CACGGTTGAA ACGCTGAAGC TACCGAGCTT
 201    TGAAGGGCCG GCACTGCCTA AGCTGACCGC TCCCGGCGGC TATATCGCCG
 251    ACATCCCCAA AGGCAACCTC AAAACCGAAA TCGAAAAGCT GGCCAAACAG
 301    CCCGAATATG CCTATCTGAA ACAGCTTCAG ACGGTCAAGG ACGTGAACTG
 351    GAACCAAGTA CAGCTCGCTT ACGACAAATG GGACTATAAA CAGGAAGGCC
 401    TAACCGGAGC CGGAGCCGCA ATTANCGCAC TGGCCGTTAC CGTGGTCACC
 451    TCAGGCGCAG GAACCGGAGC CGTATTGGGA TTAANACGNG TGGCCGCCGC
 501    CGCAACCGAT GCAGCATTT...
```

This corresponds to the amino acid sequence <SEQ ID 18; ORF49>:

```
   1  ..GTEFKTTLSG ADIQAGVGEK ARADAKIILK GIVNRIQTEE KLESNSTVWQ
  51    KQAGSGSTVE TLKLPSFEGP ALPKLTAPGG YIADIPKGNL KTEIEKLAKQ
 101    PEYAYLKQLQ TVKDVNWNQV QLAYDKWDYK QEGLTGAGAA IXALAVTVVT
 151    SGAGTGAVLG LXRVAAAATD AAF..
```

Further work revealed the complete nucleotide sequence <SEQ ID 19>:

```
   1  ATGCAACTGC TGGCAGCCGA AGGCATTCAC CAACACCAAT TGAATGTTCA
  51  GAAAAGTACC CGTTTCATCG GCATCAAAGT GGGTAAAAGC AATTACAGCA
 101  AAAACGAGCT GAACGAAACC AAACTGCCCG TACGCGTTAT CGCCCAAACA
 151  GCCAAAACCC GTTCCGGCTG GGATACCGTA CTCGAAGGCA CCGAATTCAA
 201  AACCACCCTT TCCGGAGCCG ACATACAGGC AGGGGTGGGT GAAAAAGCCC
 251  GAGCCGATGC GAAAATTATC CTAAAAGGCA TCGTTAACCG CATCCAAACC
 301  GAAGAAAAGC TGGAATCCAA CTCGACCGTA TGGCAAAAGC AGGCCGGAAG
 351  CGGCAGCACG GTTGAAACGC TGAAGCTACC GAGCTTTGAA GGGCCGGCAC
 401  TGCCTAAGCT GACCGCTCCC GGCGGCTATA TCGCCGACAT CCCCAAAGGC
 451  AACCTCAAAA CCGAAATCGA AAAGCTGGCC AAACAGCCCG AATATGCCTA
 501  TCTGAAACAG CTTCAGACGG TCAAGGACGT GAACTGGAAC CAAGTACAGC
 551  TCGCTTACGA CAAATGGGAC TATAAACAGG AAGGCCTAAC CGGAGCCGGA
 601  GCCGCAATTA TCGCACTGGC CGTTACCGTG GTCACCTCAG GCGCAGGAAC
 651  CGGAGCCGTA TTGGGATTAA ACGGTGCGGC CGCCGCCGCA ACCGATGCAG
```

```
 701  CATTTGCCTC TTTGGCCAGC CAGGCTTCCG TATCGTTCAT CAACAACAAA
 751  GGCAATATCG GTAACACCCT GAAAGAGCTG GGCAGAAGCA GCACGGTGAA
 801  AAATCTGATG GTTGCCGTCG CTACCGCAGG CGTAGCCGAC AAAATCGGTG
 851  CTTCGGCACT GAACAATGTC AGCGATAAGC AGTGGATCAA CAACCTGACC
 901  GTCAACCTGG CCAATGCGGG CAGTGCCGCA CTGATTAATA CCGCTGTCAA
 951  CGGCGGCAGC CTGAAAGACA ATCTGGAAGC GAATATCCTT GCGGCTTTGG
1001  TGAATACTGC GCATGGAGAG GCAGCAAGTA AAATCAAACA GTTGGATCAG
1051  CACTACATTG CCCATAAGAT TGCCCATGCC ATAGCGGGCT GTGCGGCAGC
1101  GGCGGCGAAT AAGGGCAAGT GTCAAGATGG TGCGATCGGT GCGGCGGTCG
1151  GTGAAATCCT TGGCGAAACC CTACTGGACG GCAGAGACCC TGGCAGCCTG
1201  AATGTGAAGG ACAGGGCAAA AATCATTGCT AAGGCGAAGC TGGCAGCAGG
1251  GGCGGTTGCG GCGTTGAGTA AGGGGGATGT GAGTACGGCG GCGAATGCGG
1301  CTGCTGTGGC GGTAGAGAAT AATTCTTTAA ATGATATACA GGATCGTTTG
1351  TTGAGTGGAA ATTATGCTTT ATGTATGAGT GCAGGAGGAG CAGAAAGCTT
1401  TTGTGAGTCT TATCGACCAC TGGGCTTGCC ACACTTTGTA AGTGTTTCAG
1451  GAGAAATGAA ATTACCTAAT AAATTCGGGA ATCGTATGGT TAATGGAAAA
1501  TTAATTATTA ACACTAGAAA TGGCAATGTA TATTTCTCTG TAGGTAAAAT
1551  ATGGAGTACT GTAAAATCAA CAAAATCAAA TATAAGTGGG GTATCTGTCG
1601  GTTGGGTTTT AAATGTTTCC CCTAATGATT ATTTAAAAGA AGCATCTATG
1651  AATGATTTCA GAAATAGTAA TCAAAATAAA GCCTATGCAG AAATGATTTC
1701  CCAGACTTTG GTAGGTGAGA GTGTTGGTGG TAGTCTTTGT CTGACAAGAG
1751  CCTGCTTTTC GGTAAGTTCA ACAATATCTA AATCTAAATC TCCTTTTAAA
1801  GATTCAAAAA TTATTGGGGA AATCGGTTTG GGAAGTGGTG TTGCTGCAGG
1851  AGTAGAAAAA ACAATATACA TAGGTAACAT AAAAGATATT GATAAATTTA
1901  TTAGTGCAAA CATAAAAAAA TAG
```

This corresponds to the amino acid sequence <SEQ ID 20; ORF49-1>:

```
   1  MQLLAAEGIH QHQLNVQKST RFIGIKVGKS NYSKNELNET KLPVRVIAQT
  51  AKTRSGWDTV LEGTEFKTTL SGADIQAGVG EKARADAKII LKGIVNRIQT
 101  EEKLESNSTV WQKQAGSGST VETLKLPSFE GPALPKLTAP GGYIADIPKG
 151  NLKTEIEKLA KQPEYAYLKQ LQTVKDVNWN QVQLAYDKWD YKQEGLTGAG
 201  AAIIALAVTV VTSGAGTGAV LGLNGAAAAA TDAAFASLAS QASVSFINNK
 251  GNIGNTLKEL GRSSTVKNLM VAVATAGVAD KIGASALNNV SDKQWINNLT
 301  VNLANAGSAA LINTAVNGGS LKDNLEANIL AALVNTAHGE AASKIKQLDQ
 351  HYIAHKIAHA IAGCAAAAAN KGKCQDGAIG AAVGEILGET LLDGRDPGSL
 401  NVKDRAKIIA KAKLAAGAVA ALSKGDVSTA ANAAAVAVEN NSLNDIQDRL
 451  LSGNYALCMS AGGAESFCES YRPLGLPHFV SVSGEMKLPN KFGNRMVNGK
 501  LIINTRNGNV YFSVGKIWST VKSTKSNISG VSVGWVLNVS PNDYLKEASM
 551  NDFRNSNQNK AYAEMISQTL VGESVGGSLC LTRACFSVSS TISKSKSPFK
 601  DSKIIGEIGL GSGVAAGVEK TIYIGNIKDI DKFISANIKK *
```

Computer analysis predicts a transmembrane domain and also indicates that ORF49 has no significant amino acid homology with known proteins. A corresponding ORF from *N.meningitidis* strain A was, however, identified:

ORF49 shows 86.1% identity over a 173aa overlap with an ORF (ORF49a) from strain A of *N. meningitidis:*

```
                                                  10        20        30
orf49.pep                                        GTEFKTTLSGADIQAGVGEKARADAKIILK
                                                 ||||||||:|||||||| ||||:|||||||
orf49a       SKNELNETKLPVRVVAQXAATRSGWDTVLEGTEFKTTLAGADIQAGVXEKARVDAKIILK
                    40        50        60        70        80        90

                     40        50        60        70        80        90
orf49.pep    GIVNRIQTEEKLESNSTVWQKQAGSGSTVETLKLPSFEGPALPKLTAPGGYIADIPKGNL
             |||||||:|||||:|||||||||| |||:|||||||||:|: |||:|||||||:|||||||
orf49a       GIVNRIQSEEKLETNSTVWQKQAGRGSTIETLKLPSFESPTPPKLSAPGGYIVDIPKGNL
                   100       110       120       130       140       150

                    100       110       120       130       140       150
orf49.pep    KTEIEKLAKQPEYAYLKQLQTVKDVNWNQVQLAYDKWDYKQEGLTGAGAAIXALAVTVVT
             |||||||:||||||||||||::|::||||||||||:|||||||| ||||| ||||||||
```

```
orf49a      KTEIEKLSKQPEYAYLKQLQVAKNINWNQVQLAYDRWDYKQEGLTEAGAAIIALAVTVVT
                 160       170       180       190       200       210

                   160       170
orf49.pep   SGAGTGAVLGLXRVAAAATDAAF
            ||||||||||| : ||||||||
orf49a      SGAGTGAVLGLNGAXAAATDAAFASLASQASVSFINNKGDVGKTLKELGRSSTVKNLVVA
                 220       230       240       250       260       270
```

ORF49-1 and ORF49a show 83.2% identity in 457 aa overlap:

```
orf49a.pep  XQLLAEEGIHKHELDVQKSRRFIGIKVGXSNYSKNELNETKLPVRVVAQXAATRSGWDTV
            |||| ||||:|:|:|||| |||||||| ||||||||||||||||:||:| ||||||||
orf49-1     MQLLAAEGIHQHQLNVQKSTRFIGIKVGKSNYSKNELNETKLPVRVIAQTAKTRSGWDTV

orf49a.pep  LEGTEFKTTLAGADIQAGVXEKARVDAKIILKGIVNRIQSEEKLETNSTVWQKQAGRGST
            ||||||||||:|||||||| ||||:|||||||||||||:|||||:|||||||||| |||
orf49-1     LEGTEFKTTLSGADIQAGVGEKARADAKIILKGIVNRIQTEEKLESNSTVWQKQAGSGST

orf49a.pep  IETLKLPSFESPTPPKLSAPGGYIVDIPKGNLKTEIEKLSKQPEYAYLKQLQVAKNINWN
            :|||||||||:|: |||:|||||:|||||||||||||:||||||||||||::|:::|||
orf49-1     VETLKLPSFEGPALPKLTAPGGYIADIPKGNLKTEIEKLAKQPEYAYLKQLQTVKDVNWN

orf49a.pep  QVQLAYDRWDYKQEGLTEAGAAIIALAVTVVTSGAGTGAVLGLNGAXAAATDAAFASLAS
            |||||||:|||||||| ||||||||||||||||||||||||| ||||||||||||||
orf49-1     QVQLAYDKWDYKQEGLTGAGAAIIALAVTVVTSGAGTGAVLGLNGAAAAATDAAFASLAS

orf49a.pep  QASVSFINNKGDVGKTLKELGRSSTVKNLVVAAATAGVADKIGASALXNVSDKQWINNLT
            ||||||||||||::|:|||||||||||||:||:|||||||||||| ||||||||||||
orf49-1     QASVSFINNKGNIGNTLKELGRSSTVKNLMVAVATAGVADKIGASALNNVSDKQWINNLT

orf49a.pep  VNLANAGSAALINTAVNGGSLKDXLEANILAALVNTAHGEAASKIKQLDQHYIVHKIAHA
            |||||||||||||||||||||| ||||||||||||||||||||||||||:|||||||
orf49-1     VNLANAGSAALINTAVNGGSLKDNLEANILAALVNTAHGEAASKIKQLDQHYIAHKIAHA

orf49a.pep  IAGCAAAAAANKGKCQDGAIGAAVGEIVGEALTNGKNPDTLTAKEREQILAYSKLVAGTVS
            |||||||||||||||||||||||||||:||:| :|::| :|::|:| :|:| :||:||:|:
orf49-1     IAGCAAAAAANKGKCQDGAIGAAVGEILGETLLDGRDPGSLNVKDRAKIIAKAKLAAGAVA

orf49a.pep  GVVGGDVNAAANAAEVAVKNNQLSDXEGREFDNEMTACAKQNXPQLCRKNTVKKYQNVAD
            ::  |||::||||| |||:||:|:| : | :::::: |
orf49-1     ALSKGDVSTAANAAAVAVENNSLNDIQDRLLSGNYALCMSAGGAESFCESYRPLGLPHFV

orf49a.pep  KRLAASIAICTDISRSTECRTIRKQHLIDSRSLHSSWEAGLIGKDDEWYKLFSKSYTQAD

orf49-1     SVSGEMKLPNKFGNRMVNGKLIINTRNGNVYFSVGKIWSTVKSTKSNISGVSVGWVLNVS
```

The complete length ORF49a nucleotide sequence <SEQ ID 21> is:

```
   1  NTGCAACTGC TGGCAGAAGA AGGCATCCAC AAGCACGAGT TGGATGTCCA
  51  AAAAAGCCGC CGCTTTATCG GCATCAAGGT AGGTNAGAGC AATTACAGTA
 101  AAAACGAACT GAACGAAACC AAATTGCCTG TCCGCGTCGT CGCCCAAANT
 151  GCAGCCACCC GTTCAGGCTG GGATACCGTG CTCGAAGGTA CCGAATTCAA
 201  AACCACGCTG GCCGGTGCCG ACATTCAGGC AGGTGTANGC GAAAAAGCCC
 251  GTGTCGATGC GAAAATTATC CTCAAAGGCA TTGTGAACCG TATCCAGTCG
 301  GAAGAAAAAT TAGAAACCAA CTCAACCGTA TGGCAGAAAC AGGCCGGACG
 351  CGGCAGCACT ATCGAAACGC TAAAACTGCC CAGCTTCGAA AGCCCTACTC
 401  CGCCCAAATT GTCCGCACCC GGCGGNTATA TCGTCGACAT TCCGAAAGGC
 451  AATCTGAAAA CCGAAATCGA AAAGCTGTCC AAACAGCCCG AGTATGCCTA
 501  TCTGAAACAG CTCCAAGTAG CGAAAAACAT CAACTGGAAT CAGGTGCAGC
 551  TTGCTTACGA CAGATGGGAC TACAAACAGG AGGGCTTAAC CGAAGCAGGT
 601  GCGGCGATTA TCGCACTGGC CGTTACCGTG GTCACCTCAG GCGCAGGAAC
 651  CGGAGCCGTA TTGGGATTAA ACGGTGCGNC CGCCGCCGCA ACCGATGCAG
 701  CATTCGCCTC TTTGGCCAGC CAGGCTTCCG TATCGTTCAT CAACAACAAA
 751  GGCGATGTCG GCAAAACCCT GAAAGAGCTG GGCAGAAGCA GCACGGTGAA
 801  AAATCTGGTG GTTGCCGCCG CTACCGCAGG CGTAGCCGAC AAAATCGGCG
 851  CTTCGGCACT GANCAATGTC AGCGATAAGC AGTGGATCAA CAACCTGACC
 901  GTCAACCTAG CCAATGCGGG CAGTGCCGCA CTGATTAATA CCGCTGTCAA
 951  CGGCGGCAGC CTGAAAGACA NTCTGGAAGC GAATATCCTT GCGGCTTTGG
1001  TCAATACCGC GCATGGAGAA GCAGCCAGTA AAATCAAACA GTTGGATCAG
```

```
1051  CACTACATAG TCCACAAGAT TGCCCATGCC ATAGCGGGCT GTGCGGCAGC
1101  GGCGGCGAAT AAGGGCAAGT GTCAGGATGG TGCGATAGGT GCGGCTGTGG
1151  GCGAGATAGT CGGGGAGGCT TTGACAAACG GCAAAAATCC TGACACTTTG
1201  ACAGCTAAAG AACGCGAACA GATTTTGGCA TACAGCAAAC TGGTTGCCGG
1251  TACGGTAAGC GGTGTGGTCG GCGGCGATGT AAATGCGGCG GCGAATGCGG
1301  CTGAGGTAGC GGTGAAAAAT AATCAGCTTA GCGACNAAGA GGGTAGAGAA
1351  TTTGATAACG AAATGACTGC ATGCGCCAAA CAGAATANTC CTCAACTGTG
1401  CAGAAAAAAT ACTGTAAAAA AGTATCAAAA TGTTGCTGAT AAAAGACTTG
1451  CTGCTTCGAT TGCAATATGT ACGGATATAT CCCGTAGTAC TGAATGTAGA
1501  ACAATCAGAA AACAACATTT GATCGATAGT AGAAGCCTTC ATTCATCTTG
1551  GGAAGCAGGT CTAATTGGTA AAGATGATGA ATGGTATAAA TTATTCAGCA
1601  AATCTTACAC CCAAGCAGAT TTGGCTTTAC AGTCTTATCA TTTGAATACT
1651  GCTGCTAAAT CTTGGCTTCA ATCGGGCAAT ACAAAGCCTT TATCCGAATG
1701  GATGTCCGAC CAAGGTTATA CACTTATTTC AGGAGTTAAT CCTAGATTCA
1751  TTCCAATACC AAGAGGGTTT GTAAAACAAA ATACACCTAT TACTAATGTC
1801  AAATACCCGG AAGGCATCAG TTTCGATACA AACCTANAAA GACATCTGGC
1851  AAATGCTGAT GGTTTTAGTC AAGAACAGGG CATTAAAGGA GCCCATAACC
1901  GCACCAATNT TATGGCAGAA CTAAATTCAC GAGGAGGANG NGTAAAATCT
1951  GAAACCCANA CTGATATTGA AGGCATTACC CGAATTAAAT ATGAGATTCC
2001  TACACTAGAC AGGACAGGTA AACCTGATGG TGGATTTAAG GAAATTTCAA
2051  GTATAAAAAC TGTTTATAAT CCTAAAAANT TTTNNGATGA TAAAATACTT
2101  CAAATGGCTC AANATGCTGN TTCACAAGGA TATTCAAAAG CCTCTAAAAT
2151  TGCTCAAAAT GAAAGAACTA AATCAATATC GGAAAGAAAA AATGTCATTC
2201  AATTCTCAGA AACCTTTGAC GGAATCAAAT TTAGANNNTA TNTNGATGTA
2251  AATACAGGAA GAATTACAAA CATTCACCCA GAATAATTTA A
```

This encodes a protein having amino acid sequence <SEQ ID 22>:

```
   1  XQLLAEEGIH KHELDVQKSR RFIGIKVGXS NYSKNELNET KLPVRVVAQX
  51  AATRSGWDTV LEGTEFKTTL AGADIQAGVX EKARVDAKII LKGIVNRIQS
 101  EEKLETNSTV WQKQAGRGST IETLKLPSFE SPTPPKLSAP GGYIVDIPKG
 151  NLKTEIEKLS KQPEYAYLKQ LQVAKNINWN QVQLAYDRWD YKQEGLTEAG
 201  AAIIALAVTV VTSGAGTGAV LGLNGAXAAA TDAAFASLAS QASVSFINNK
 251  GDVGKTLKEL GRSSTVKNLV VAAATAGVAD KIGASALXNV SDKQWINNLT
 301  VNLANAGSAA LINTAVNGGS LKDXLEANIL AALVNTAHGE AASKIKQLDQ
 351  HYIVHKIAHA IAGCAAAAAN KGKCQDGAIG AAVGEIVGEA LTNGKNPDTL
 401  TAKEREQILA YSKLVAGTVS GVVGGDVNAA ANAAEVAVKN NQLSDXEGRE
 451  FDNEMTACAK QNXPQLCRKN TVKKYQNVAD KRLAASIAIC TDISRSTECR
 501  TIRKQHLIDS RSLHSSWEAG LIGKDDEWYK LFSKSYTQAD LALQSYHLNT
 551  AAKSWLQSGN TKPLSEWMSD QGYTLISGVN PRFIPIPRGF VKQNTPITNV
 601  KYPEGISFDT NLXRHLANAD GFSQEQGIKG AHNRTNXMAE LNSRGGXVKS
 651  ETXTDIEGIT RIKYEIPTLD RTGKPDGGFK EISSIKTVYN PKXFXDDKIL
 701  QMAQXAXSQG YSKASKIAQN ERTKSISERK NVIQFSETFD GIKFRXYXDV
 751  NTGRITNIHP E*
```

Based on the presence of a putative transmembrane domain, it is predicted that these proteins from *N.meningitidis*, and their epitopes, could be useful antigens for vaccines or diagnostics.

Example 5

The following partial DNA sequence was identified in *N.meningitidis* <SEQ ID 23>

```
   1  ..CGGATCGTTG TAGGTTTGCG GATTTCTTGC GCCGTAGTCA CCGTAGTCCC
  51    AAGTATAACC CAAGGCTTTG TCTTCGCCTT TCATTCCGAT AAGGGATATG
 101    ACGCTTTGGT CGGTATAGCC GTCTTGGGAA CCTTTGTCCA CCCAACGCAT
 151    ATCTGCCTGC GGATTCTCAT TGCCGCTTCT TGGCTGCTGA TTTTTCTGCC
 201    TTCGCGTTTT TCAACTTCGC GCTTGAGGGC TTCGGCATAT TTGTCGGCCA
 251    ACGCCATTTC TTTCGGATGC AGCTGCCTAT TGTTCCAATC TACATTCGCA
 301    CCCACCACAG CACCACCACT ACCACCAGTT GCATAG
```

This corresponds to the amino acid sequence <SEQ ID 24; ORF50>:

```
   1  ..RIVVGLRISC AVVTVVPSIT QGFVFAFHSD KGYDALVGIA VLGTFVHPTH
  51    ICLRILIAAS WLLIFLPSRF STSRLRASAY LSANAISFGC SCLLFQSTFA
 101    PTTAPPLPPV A*
```

Computer analysis predicts two transmembrane domains and also indicates that ORF50 has no significant amino acid homology with known proteins.

Based on the presence of a putative transmembrane domain, it is predicted that this protein from *N.meningitidis*, and its epitopes, could be useful antigens for vaccines or diagnostics.

Example 6

The following partial DNA sequence was identified in *N.meningitidis* <SEQ ID 25>

```
   1   .AAGTTTGACT TTACCTGGTT TATTCCGGCG GTAATCAAAT ACCGCCGGTT
  51    GTTTTTTGAA GTATTGGTGG TGTCGGTGGT GTTGCAGCTG TTTGCGCTGA
 101    TTACGCCTCT GTTTTTCCAA GTGGTGATGG ACAAGGTGCT GGTACATCGG
 151    GGATTCTCTA CTTTGGATGT GGTGTCGGTG GCTTTGTTGG TGGTGTCGCT
 201    GTTTGAGATT GTGTTGGGCG GTTTGCGGAC GTATCTGTTT GCACATACGA
 251    CTTCACGTAT TGATGTGGAA TTGGGCGCGC GTTTGTTCCG GCATCTGCTT
 301    TCCCTGCCTT TATCCTATTT CGAGCACAGA CGAGTGGGTG ATACGGTGGC
 351    TCGGGTGCGG GAATTGGAGC AGATTCGCAA TTTCTTGACC GGTCAGGCGC
 401    TGACTTCGGT GTTGGATTTG GCGTTTTCGT TTATCTTTCT GGCGGTGATG
 451    TGGTATTACA GCTCCACTCT GACTTGGGTG GTATTGGCTT CGTTG.....
                                        //
1451    ..........  ..........  ..........  ..........  ..........
1501    ..........  ..........  ..........  ..........  ..ATTTGCGC
1551    CAACCGGACG GTGCTGATTA TCGCCCACCG TCTGTCCACT GTTAAAACGG
1601    CACACCGGAT CATTGCCATG GATAAAGGCA GGATTGTGGA AGCGGGAACA
1651    CAGCAGGAAT TGCTGGCGAA CG..AACGGA TATTACCGCT ATCTGTATGA
1701    TTTACAGAAC GGGTAG
```

This corresponds to the amino acid sequence <SEQ ID 26; ORF39>:

```
   1   ..KFDFTWFIPA VIKYRRLFFE VLVVSVVLQL FALITPLFFQ VVMDKVLVHR
  51     GFSTLDVVSV ALLVVSLFEI VLGGLRTYLF AHTTSRIDVE LGARLFRHLL
 101     SLPLSYFEHR RVGDTVARVR ELEQIRNFLT GQALTSVLDL AFSFIFLAVM
 151     WYYSSTLTWV VLASL..... ..........  ..........  ..........
                                        //
 501     ..........  ....ICANRT VLIIAHRLST VKTAHRIIAM DKGRIVEAGT
 551     QQELLANXNG YYRYLYDLQN G*
```

Further work revealed the complete nucleotide sequence <SEQ ID 27>:

```
   1  ATGTCTATCG TATCCGCACC GCTCCCCGCC CTTTCCGCCC TCATCATCCT
  51  CGCCCATTAC CACGGCATTG CCGCCAATCC TGCCGATATA CAGCATGAAT
 101  TTTGTACTTC CGCACAGAGC GATTTAAATG AAACGCAATG GCTGTTAGCC
 151  GCCAAATCTT TGGGATTGAA GGCAAAGGTA GTCCGCCAGC CTATTAAACG
 201  TTTGGCTATG GCGACTTTAC CCGCATTGGT ATGGTGTGAT GACGGCAACC
 251  ATTTCATTTT GGCCAAAACA GACGGTGAGG GTGAGCATGC CCAATTTTTG
 301  ATACAGGATT TGGTTACGAA TAAGTCTGCG GTATTGTCTT TTGCCGAATT
 351  TTCTAACAGA TATTCGGGCA AACTGATATT GGTTGCTTCC CGCGCTTCGG
 401  TATTGGGCAG TTTGGCAAAG TTTGACTTTA CCTGGTTTAT TCCGGCGGTA
 451  ATCAAATACC GCCGGTTGTT TTTTGAAGTA TTGGTGGTGT CGGTGGTGTT
 501  GCAGCTGTTT GCGCTGATTA CGCCTCTGTT TTTCCAAGTG GTGATGGACA
 551  AGGTGCTGGT ACATCGGGGA TTCTCTACTT TGGATGTGGT GTCGGTGGCT
 601  TTGTTGGTGG TGTCGCTGTT TGAGATTGTG TTGGGCGGTT TGCGGACGTA
 651  TCTGTTTGCA CATACGACTT CACGTATTGA TGTGGAATTG GGCGCGCGTT
 701  TGTTCCGGCA TCTGCTTTCC CTGCCTTTAT CCTATTTCGA GCACAGACGA
 751  GTGGGTGATA CGGTGGCTCG GGTGCGGGAA TTGGAGCAGA TTCGCAATTT
 801  CTTGACCGGT CAGGCGCTGA CTTCGGTGTT GGATTTGGCG TTTTCGTTTA
 851  TCTTTCTGGC GGTGATGTGG TATTACAGCT CCACTCTGAC TTGGGTGGTA
 901  TTGGCTTCGT TGCCTGCCTA TGCGTTTTGG TCGGCATTTA TCAGTCCGAT
 951  ACTGCGGACG CGTCTGAACG ATAAGTTCGC GCGCAATGCA GACAACCAGT
1001  CGTTTTTAGT AGAAAGCATC ACTGCGGTGG GTACGGTAAA GGCGATGGCG
1051  GTGGAGCCGC AGATGACGCA GCGTTGGGAC AATCAGTTGG CGGCTTATGT
```

```
1101  GGCTTCGGGA TTTCGGGTAA CGAAGTTGGC GGTGGTCGGC CAGCAGGGGG
1151  TGCAGCTGAT TCAGAAGCTG GTGACGGTGG CGACGTTGTG GATTGGCGCA
1201  CGGCTGGTAA TTGAGAGCAA GCTGACGGTG GGGCAGCTGA TTGCGTTTAA
1251  TATGCTCTCG GGACAGGTGG CGGCGCCTGT TATCCGTTTG GCGCAGTTGT
1301  GGCAGGATTT CCAGCAGGTG GGGATTTCGG TGGCGCGTTT GGGGGATATT
1351  CTGAATGCGC CGACCGAGAA TGCGTCTTCG CATTTGGCTT TGCCCGATAT
1401  CCGGGGGGAG ATTACGTTCG AACATGTCGA TTTCCGCTAT AAGGCGGACG
1451  GCAGGCTGAT TTTGCAGGAT TTGAACCTGC GGATTCGGGC GGGGGAAGTG
1501  CTGGGGATTG TGGGACGTTC GGGGTCGGGC AAATCCACAC TCACCAAATT
1551  GGTGCAGCGT CTGTATGTAC CGGAGCAGGG ACGGGTGTTG GTGGACGGCA
1601  ACGATTTGGC TTTGGCCGCT CCTGCCTGGC TGCGGCGGCA GGTCGGCGTG
1651  GTCTTGCAGG AGAATGTGCT GCTCAACCGC AGCATACGCG ACAATATCGC
1701  GCTGACGGAT ACGGGTATGC CGCTGGAACG CATTATCGAA GCAGCCAAAC
1751  TGGCGGGCGC ACACGAGTTT ATTATGGAGC TGCCGGAAGG CTACGGCACC
1801  GTGGTGGGCG AACAAGGGGC CGGCTTGTCG GGCGGACAGC GGCAGCGTAT
1851  TGCGATTGCC CGCGCGTTAA TCACCAATCC GCGCATTCTG ATTTTTGATG
1901  AAGCCACCAG CGCGCTGGAT TATGAAAGTG AACGAGCGAT TATGCAGAAC
1951  ATGCAGGCCA TTTGCGCCAA CCGGACGGTG CTGATTATCG CCCACCGTCT
2001  GTCCACTGTT AAAACGGCAC ACCGGATCAT TGCCATGGAT AAAGGCAGGA
2051  TTGTGGAAGC GGGAACACAG CAGGAATTGC TGGCGAAGCC GAACGGATAT
2101  TACCGCTATC TGTATGATTT ACAGAACGGG TAG
```

This corresponds to the amino acid sequence <SEQ ID 28; ORF39-1>:

```
   1  MSIVSAPLPA LSALIILAHY HGIAANPADI QHEFCTSAQS DLNETQWLLA
  51  AKSLGLKAKV VRQPIKRLAM ATLPALVWCD DGNHFILAKT DGEGEHAQFL
 101  IQDLVTNKSA VLSFAEFSNR YSGKLILVAS RASVLGSLAK FDFTWFIPAV
 151  IKYRRLFFEV LVVSVVLQLF ALITPLFFQV VMDKVLVHRG FSTLDVVSVA
 201  LLVVSLFEIV LGGLRTYLFA HTTSRIDVEL GARLFRHLLS LPLSYFEHRR
 251  VGDTVARVRE LEQIRNFLTG QALTSVLDLA FSFIFLAVMW YYSSTLTWVV
 301  LASLPAYAFW SAFISPILRT RLNDKFARNA DNQSFLVESI TAVGTVKAMA
 351  VEPQMTQRWD NQLAAYVASG FRVTKLAVVG QQGVQLIQKL VTVATLWIGA
 401  RLVIESKLTV GQLIAFNMLS GQVAAPVIRL AQLWQDFQQV GISVARLGDI
 451  LNAPTENASS HLALPDIRGE ITFEHVDFRY KADGRLILQD LNLRIRAGEV
 501  LGIVGRSGSG KSTLTKLVQR LYVPEQGRVL VDGNDLALAA PAWLRRQVGV
 551  VLQENVLLNR SIRDNIALTD TGMPLERIIE AAKLAGAHEF IMELPEGYGT
 601  VVGEQGAGLS GGQRQRIAIA RALITNPRIL IFDEATSALD YESERAIMQN
 651  MQAICAMRTV LIIAHRLSTV KTAHRIIAMD KGRIVEAGTQ QELLAKPNGY
 701  YRYLYDLQNG *
```

Computer analysis of this amino acid sequence gave the following results:

Homology with a predicted ORF from *N.meningitidis* (strain A)

ORF39 shows 100% identity over a 165aa overlap with an ORF (ORF39a) from strain A of *N. meningitidis:*

```
                                                    10        20        30
orf39.pep                                          KFDFTWFIPAVIKYRRLFFEVLVVSVVLQL
                                                   ||||||||||||||||||||||||||||||
orf39a       AVLSFAEFSNRYSGKLILVASRASVLGSLAKFDFTWFIPAVIKYRRLFFEVLVVSVVLQL
           110       120       130       140       150       160

                    40        50        60        70        80        90
orf39.pep    FALITPLFFQVVMDKVLVHRGFSTLDVVSVALLVVSLFEIVLGGLRTYLFAHTTSRIDVE
             ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
orf39a       FALITPLFFQVVMDKVLVHRGFSTLDVVSVALLVVSLFEIVLGGLRTYLFAHTTSRIDVE
           170       180       190       200       210       220

                   100       110       120       130       140       150
orf39.pep    LGARLFRHLLSLPLSYFEHRRVGDTVARVRELEQIRNFLTGQALTSVLDLAFSFIFLAVM
             ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
orf39a       LGARLFRHLLSLPLSYFEHRRVGDTVARVRELEQIRNFLTGQALTSVLDLAFSFIFLAVM
           230       240       250       260       270       280

                   160       170       180       190       200       210
orf39.pep    WYYSSTLTWVVLASLXXXXXXXXXXXXXXXXXXXXXXXXXXXXXICANRTVLIIAHRLSTV
```

```
                  ||||||||||||||||
orf39a       WYYSSTLTWVVLASLPAYAFWSAFISPILRTRLNDKFARNADNQSFLVESITAVGTVKAM
          290        300        310        320        330        340
```

ORF39-1 and ORF39a show 99.4% identity in 710 aa overlap:

```
orf39-1.pep   MSIVSAPLPALSALIILAHYHGIAANPADIQHEFCTSAQSDLNETQWLLAAKSLGLKAKV
              ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
orf39a        MSIVSAPLPALSALIILAHYHGIAANPADIQHEFCTSAQSDLNETQWLLAAKSLGLKAKV

orf39-1.pep   VRQPIKRLAMATLPALVWCDDGNHFILAKTDGEGEHAQFLIQDLVTNKSAVLSFAEFSNR
              ||||||||||||||||||||||||||||||| ||||:|||||:|||||||||||||||||
orf39a        VRQPIKRLAMATLPALVWCDDGNHFILAKTDGGGEHAQYLIQDLTTNKSAVLSFAEFSNR

orf39-1.pep   YSGKLILVASRASVLGSLAKFDFTWFIPAVIKYRRLFFEVLVVSVVLQLFALITPLFFQV
              ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
orf39a        YSGKLILVASRASVLGSLAKFDFTWFIPAVIKYRRLFFEVLVVSVVLQLFALITPLFFQV

orf39-1.pep   VMDKVLVHRGFSTLDVVSVALLVVSLFEIVLGGLRTYLFAHTTSRIDVELGARLFRHLLS
              ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
orf39a        VMDKVLVHRGFSTLDVVSVALLVVSLFEIVLGGLRTYLFAHTTSRIDVELGARLFRHLLS

orf39-1.pep   LPLSYFEHRRVGDTVARVRELEQIRNFLTGQALTSVLDLAFSFIFLAVMWYYSSTLTWVV
              ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
orf39a        LPLSYFEHRRVGDTVARVRELEQIRNFLTGQALTSVLDLAFSFIFLAVMWYYSSTLTWVV

orf39-1.pep   LASLPAYAFWSAFISPILRTRLNDKFARNADNQSFLVESITAVGTVKAMAVEPQMTQRWD
              ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
orf39a        LASLPAYAFWSAFISPILRTRLNDKFARNADNQSFLVESITAVGTVKAMAVEPQMTQRWD

orf39-1.pep   NQLAAYVASGFRVTKLAVVGQQGVQLIQKLVTVATLWIGARLVIESKLTVGQLIAFNMLS
              ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
orf39a        NQLAAYVASGFRVTKLAVVGQQGVQLIQKLVTVATLWIGARLVIESKLTVGQLIAFNMLS

orf39-1.pep   GQVAAPVIRLAQLWQDFQQVGISVARLGDILNAPTENASSHLALPDIRGEITFEHVDFRY
              ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
orf39a        GQVAAPVIRLAQLWQDFQQVGISVARLGDILNAPTENASSHLALPDIRGEITFEHVDFRY

orf39-1.pep   KADGRLILQDLNLRIRAGEVLGIVGRSGSGKSTLTKLVQRLYVPEQGRVLVDGNDLALAA
              |||||||||||||||||||||||||||||||||||||||||| ||||||||||||||||
orf39a        KADGRLILQDLNLRIRAGEVLGIVGRSGSGKSTLTKLVQRLYVPAQGRVLVDGNDLALAA

orf39-1.pep   PAWLRRQVGVVLQENVLLNRSIRDNIALTDTGMPLERIIEAAKLAGAHEFIMELPEGYGT
              ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
orf39a        PAWLRRQVGVVLQENVLLNRSIRDNIALTDTGMPLERIIEAAKLAGAHEFIMELPEGYGT

orf39-1.pep   VVGEQGAGLSGGQRQRIAIARALITNPRILIFDEATSALDYESERAIMQNMQAICANRTV
              ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
orf39a        VVGEQGAGLSGGQRQRIAIARALITNPRILIFDEATSALDYESERAIMQNMQAICANRTV

orf39-1.pep   LIIAHRLSTVKTAHRIIAMDKGRIVEAGTQQELLAKPNGYYRYLYDLQNGX
              ||||||||||||||||||||||||||||||||||||||||||||||||||
orf39a        LIIAHRLSTVKTAHRIIAMDKGRIVEAGTQQELLAKPNGYYRYLYDLQNGX
```

The complete length ORF39a nucleotide sequence <SEQ ID 29> is:

```
   1  ATGTCTATCG TATCCGCACC GCTCCCCGCC CTTTCCGCCC TCATCATCCT
  51  CGCCCATTAC CACGGCATTG CCGCCAATCC TGCCGATATA CAGCATGAAT
 101  TTTGTACTTC CGCACAGAGC GATTTAAATG AAACGCAATG GCTGTTAGCC
 151  GCCAAATCTT TGGGATTGAA GGCAAAGGTA GTCCGCCAGC CTATTAAACG
 201  TTTGGCTATG GCGACTTTAC CCGCATTGGT ATGGTGTGAT GACGGCAACC
 251  ATTTTATTTT GGCTAAAACA GACGGTGGGG GTGAGCATGC CCAATATCTA
 301  ATACAGGATT TAACTACGAA TAAGTCTGCG GTATTGTCTT TTGCCGAATT
 351  TTCTAACAGA TATTCGGGCA AACTGATATT GGTTGCTTCC CGCGCTTCGG
 401  TATTGGGCAG TTTGGCAAAG TTTGACTTTA CCTGGTTTAT TCCGGCGGTA
 451  ATCAAATACC GCCGGTTGTT TTTTGAAGTA TTGGTGGTGT CGGTGGTGTT
 501  GCAGCTGTTT GCGCTGATTA CGCCTCTGTT TTTCCAAGTG GTGATGGACA
 551  AGGTGCTGGT ACATCGGGGA TTCTCTACTT TGGATGTGGT GTCGGTGGCT
 601  TTGTTGGTGG TGTCGCTGTT TGAGATTGTG TTGGGCGGTT TGCGGACGTA
 651  TCTGTTTGCA CATACGACTT CACGTATTGA TGTGGAATTG GGCGCGCGTT
```

```
 701  TGTTCCGGCA TCTGCTTTCC CTGCCTTTAT CCTATTTCGA GCACAGACGA
 751  GTGGGTGATA CGGTGGCTCG GGTGCGGGAA TTGGAGCAGA TTCGCAATTT
 801  CTTGACCGGT CAGGCGCTGA CTTCGGTGTT GGATTTGGCG TTTTCGTTTA
 851  TCTTTCTGGC GGTGATGTGG TATTACAGCT CCACTCTGAC TTGGGTGGTA
 901  TTGGCTTCGT TGCCTGCCTA TGCGTTTTGG TCGGCATTTA TCAGTCCGAT
 951  ACTGCGGACG CGTCTGAACG ATAAGTTCGC GCGCAATGCA GACAACCAGT
1001  CGTTTTTAGT AGAAAGCATC ACTGCGGTGG GTACGGTAAA GGCGATGGCG
1051  GTGGAGCCGC AGATGACGCA GCGTTGGGAC AATCAGTTGG CGGCTTATGT
1101  GGCTTCGGGA TTTCGGGTAA CGAAGTTGGC GGTGGTCGGC CAGCAGGGGG
1151  TGCAGCTGAT TCAGAAGCTG GTGACGGTGG CGACGTTGTG GATTGGCGCA
1201  CGGCTGGTAA TTGAGAGCAA GCTGACGGTG GGGCAGCTGA TTGCGTTTAA
1251  TATGCTCTCG GGACAGGTGG CGGCGCCTGT TATCCGTTTG GCGCAGTTGT
1301  GGCAGGATTT CCAGCAGGTG GGGATTTCGG TGGCGCGTTT GGGGGATATT
1351  CTGAATGCGC CGACCGAGAA TGCGTCTTCG CATTTGGCTT TGCCCGATAT
1401  CCGGGGGGAG ATTACGTTCG AACATGTCGA TTTCCGCTAT AAGGCGGACG
1451  GCAGGCTGAT TTTGCAGGAT TTGAACCTGC GGATTCGGGC GGGGGAAGTG
1501  CTGGGGATTG TGGGACGTTC GGGGTCGGGC AAATCCACAC TCACCAAATT
1551  GGTGCAGCGT CTGTATGTAC CGGCGCAGGG ACGGGTGTTG GTGGACGGCA
1601  ACGATTTGGC TTTGGCCGCT CCTGCTTGGC TGCGGCGGCA GGTCGGCGTG
1651  GTCTTGCAGG AGAATGTGCT GCTCAACCGC AGCATACGCG ACAATATCGC
1701  GCTGACGGAT ACGGGTATGC CGCTGGAACG CATTATCGAA GCAGCCAAAC
1751  TGGCGGGCGC ACACGAGTTT ATTATGGAGC TGCCGGAAGG CTACGGCACC
1801  GTGGTGGGCG AACAAGGGGC CGGCTTGTCG GGCGGACAGC GGCAGCGTAT
1851  TGCGATTGCC CGCGCGTTAA TCACCAATCC GCGCATTCTG ATTTTTGATG
1901  AAGCCACCAG CGCGCTGGAT TATGAAAGTG AACGAGCGAT TATGCAGAAC
1951  ATGCAGGCCA TTTGCGCCAA CCGGACGGTG CTGATTATCG CCCACCGTCT
2001  GTCCACTGTT AAAACGGCAC ACCGGATCAT TGCCATGGAT AAAGGCAGGA
2051  TTGTGGAAGC GGGAACACAG CAGGAATTGC TGGCGAAGCC GAACGGATAT
2101  TACCGCTATC TGTATGATTT ACAGAACGGG TAG
```

This encodes a protein having amino acid sequence <SEQ ID 30>:

```
   1  MSIVSAPLPA LSALIILAHY HGIAANPADI QHEFCTSAQS DLNETQWLLA
  51  AKSLGLKAKV VRQPIKRLAM ATLPALVWCD DGNHFILAKT DGGGEHAQYL
 101  IQDLTTNKSA VLSFAEFSNR YSGKLILVAS RASVLGSLAK FDFTWFIPAV
 151  IKYRRLFFEV LVVSVVLQLF ALITPLFFQV VMDKVLVHRG FSTLDVVSVA
 201  LLVVSLFEIV LGGLRTYLFA HTTSRIDVEL GARLFRHLLS LPLSYFEHRR
 251  VGDTVARVRE LEQIRNFLTG QALTSVLDLA FSFIFLAVMW YYSSTLTWVV
 301  LASLPAYAFW SAFISPILRT RLNDKFARNA DNQSFLVESI TAVGTVKAMA
 351  VEPQMTQRWD NQLAAYVASG FRVTKLAVVG QQGVQLIQKL VTVATLWIGA
 401  RLVIESKLTV GQLIAFNMLS GQVAAPVIRL AQLWQDFQQV GISVARLGDI
 451  LNAPTENASS HLALPDIRGE ITFEHVDFRY KADGRLILQD LNLRIRAGEV
 501  LGIVGRSGSG KSTLTKLVQR LYVPAQGRVL VDGNDLALAA PAWLRRQVGV
 551  VLQENVLLNR SIRDNIALTD TGMPLERIIE AAKLAGAHEF IMELPEGYGT
 601  VVGEQGAGLS GGQRQRIAIA RALITNPRIL IFDEATSALD YESERAIMQN
 651  MQAICANRTV LIIAHRLSTV KTAHRIIAMD KGRIVEAGTQ QELLAKPNGY
 701  YRYLYDLQNG *
```

ORF39a is homologous to a cytolysin from *A.pleuropneumoniae*:

```
sp|P26760|RT1B_ACTPL RTX-I TOXIN DETERMINANT B (TOXIN RTX-I SECRETION ATP-
BINDING PROTEIN) (APX-IB) (HLY-IB) (CYTOLYSIN IB) (CLY-IB)
>gi|97137|pir||D43599 cytolysin IB - Actinobacillus pleuropneumoniae (serotype 9)
>gi|38944 (X61112) ClyI-B protein [Actinobacillus pleuropneumoniae] Length = 707
 Score =  931 bits (2379), Expect = 0.0
 Identities = 472/690 (68%), Positives = 540/690 (77%), Gaps = 3/690 (0%)

Query: 20  YHGIAANPADIQHEFCTSAQSDLNETQWXXXXXXXXXXXXVVRQPIKRLAMATLPALVWC 79
           YH IA NP +++H+F    +  L+ T W            V++ I RLA   LPALVW
Sbjct: 20  YHNIAVNPEELKHKFDLEGKG-LDLTAWLLAAKSLELKAKQVKKAIDRLAFIALPALVWR 78

Query: 80  DDGNHFILAKTDGGGEHAQYLIQDLTTNKSAVLSFAEFSNRYSGKLILVASRASVLGSLA 139
           +DG HFIL K D   E  +YLI DL T+   +L  AEF + Y GKLILVASRAS++G LA
Sbjct: 79  EDGKHFILTKIDN--EAKKYLIFDLETHNPRILEQAEFESLYQGKLILVASRASIVGKLA 136

Query: 140 KFDFTWFIPAVIKYRRXXXXXXXXXXXXXXXXXITPLFFQVVMDKVLVHRGFXXXXXXXX 199
           KFDFTWFIPAVIKYR+                  ITPLFFQVVMDKVLVHRGF
Sbjct: 137 KFDFTWFIPAVIKYRKIFIETLIVSIFLQIFALITPLFFQVVMDKVLVHRGFSTLNVITV 196

Query: 200 XXXXXXXFEIVLGGLRTYLFAHTTSRIDVELGARLFRHLLSLPLSYFEHRRVGDTVARVR 259
```

```
                FEIVL GLRTY+FAH+TSRIDVELGARLFRHLL+LP+SYFE+RRVGDTVARVR
Sbjct: 197 ALAIVVLFEIVLNGLRTYIFAHSTSRIDVELGARLFRHLLALPISYFENRRVGDTVARVR 256

Query: 260 ELEQIRNFLTGQALTSVLDLAFSFIFLAVMWYYSSTLTWVVLASLPAYAFWSAFISPILR 319
           EL+QIRNFLTGQALTSVLDL FSFIF AVMWYYS  LT V+L SLP Y  WS FISPILR
Sbjct: 257 ELDQIRNFLTGQALTSVLDLMFSFIFFAVMWYYSPKLTLVILGSLPFYMGWSIFISPILR 316

Query: 320 TRLNDKFARNADNQSFLVESITAVGTVKAMAVEPQMTQRWDNQLAAYVASGFRVTKLAVV 379
            RL++KFAR ADNQSFLVES+TA+ T+KA+AV PQMT  WD QLA+YV++GFRVT LA +
Sbjct: 317 RRLDEKFARGADNQSFLVESVTAINTIKALAVTPQMTNTWDKQLASYVSAGFRVTTLATI 376

Query: 380 GQQGVQLIQKLVTVATLWIGARLVIESKLTVGQLIAFNMLSGQVAAPVIRLAQLWQDFQQ 439
           GQQGVQ IQK+V V TLW+GA LVI   L++GQLIAFNMLSGQV APVIRLAQLWQDFQQ
Sbjct: 377 GQQGVQFIQKVVMVITLWLGAHLVISGDLSIGQLIAFNMLSGQVIAPVIRLAQLWQDFQQ 436

Query: 440 VGISVARLGDILNAPTENASSHLALPDIRGEITFEHVDFRYKADGRLILQDLNLRIRAGE 499
           VGISV RLGD+LN+PTE+    LALP+I+G+ITF ++ FRYK D  +IL D+NL I+ GE
Sbjct: 437 VGISVTRLGDVLNSPTESYQGKLALPEIKGDITFRNIRFRYKPDAPVILNDVNLSIQQGE 496

Query: 500 VLGIVGRSGSGKSTLTKLVQRLYVPAQGRVLVDGNDLALAAPAWLRRQVGVVLQENVLLN 559
           V+GIVGRSGSGKSTLTKL+QR Y+P  G+VL+DG+DLALA P WLRRQVGVVLQ+NVLLN
Sbjct: 497 VIGIVGRSGSGKSTLTKLIQRFYIPENGQVLIDGHDLALADPNWLRRQVGVVLQDNVLLN 556

Query: 560 RSIRDNIALTDTGMPLERIIEAAKLAGAHEFIMELPEGYGTVVGEQGAGLSGGQRQRIAI 619
           RSIRDNIAL D GMP+E+I+ AAKLAGAHEFI EL EGY T+VGEQGAGLSGGQRQRIAI
Sbjct: 557 RSIRDNIALADPGMPMEKIVHAAKLAGAHEFISELREGYNTIVGEQGAGLSGGQRQRIAI 616

Query: 620 ARALITNPRILIFDEATSALDYESERAIMQNMQAICANRTVLIIAHRLSTVKTAHRIIAM 679
           ARAL+ NP+ILIFDEATSALDYESE  IM+NM  IC  RTV+IIAHRLSTVK A RII M
Sbjct: 617 ARALVNNPKILIFDEATSALDYESEHIIMRNMHQICKGRTVIIIAHRLSTVKNADRIIVM 676

Query: 680 DKGRIVEAGTQQELLAKPNGYYRYLYDLQN 709
           +KG+IVE G  +ELLA PNG Y YL+ LQ+
Sbjct: 677 EKGQIVEQGKHKELLADPNGLYHYLHQLQS 706
```

Homology with the HlyB leucotoxin secretion ATP-binding protein of *Haemophilus actinomycetemcomitans* (accession number X53955)

ORF39 and HlyB protein show 71% and 69% amino acid identity in 167 and 55 overlap at the N- and C-terminal regions, respectively:

```
Orf39   1   KFDFTWFIPAVIKYRRXXXXXXXXXXXXXXXXXITPLFFQVVMDKVLVHRGFXXXXXXXX 60
            KFDFTWFIPAVIKYR+                    ITPLFFQVVMDKVLVHRGF
HlyB    137 KFDFTWFIPAVIKYRKIFIETLIVSIFLQIFALITPLFFQVVMDKVLVHRGFSTLNVITV 196

Orf39   61  XXXXXXXFEIVLGGLRTYLFAHTTSRIDVELGARLFRHLLSLPLSYFEHRRVGDTVARVR 120
                FEI+LGGLRTY+FAH+TSRIDVELGARLFRHLL+LP+SYFE RRVGDTVARVR
HlyB    197 ALAIVVLFEIILGGLRTYVFAHSTSRIDVELGARLFRHLLALPISYFEARRVGDTVARVR 256

Orf39   121 ELEQIRNFLTGQALTSVLDLAFSFIFLAVMWYYSSTLTWVVLASLIC 167
            EL+QIRNFLTGQALTS+LDL FSFIF AVMWYYS  LT VVL SL C
HlyB    257 ELDQIRNFLTGQALTSILDLLFSFIFFAVMWYYSPKLTLVVLGSLPC 303

                                    //

Orf39   166 ICANRTVLIIAHRLSTVKTAHRIIAMDKGRIVEAGTQQELLANXNGYYRYLYDLQ 220
            IC NRTVLIIAHRLSTVK A RII MDKG I+E G  QELL +  G Y YL+ LQ
HlyB    651 ICQNRTVLIIAHRLSTVKNADRIIVMDKGEIIEQGKHQELLKDEKGLYSYLHQLQ 705
```

Based on this analysis, it is predicted that this protein from *N.meningitidis*, and its epitopes, could be useful antigens for vaccines or diagnostics.

Example 7

The following partial DNA sequence was identified in *N.meningitidis* <SEQ ID 31>

```
   1  ATGAAATACT TGATCCGCAC CGCCTTACTC GCAGTCGCAG CCGCCGGCAT
  51  CTACGCCTGC CAACCGCAAT CCGAAGCCGC AGTGCAAGTC AAGGCTGAAA
 101  ACAGCCTGAC CGCTATGCGC TTAGCCGTCG CCGACAAACA GGCAGAGATT
 151  GACGGGTTGA ACGCCCAAAk sGACGCCGAA ATCAGA...
```

This corresponds to the amino acid sequence <SEQ ID 32; ORF52>:

```
   1  MKYLIRTALL AVAAAGIYAC QPQSEAAVQV KAENSLTAMR LAVADKQAEI
  51  DGLNAQXDAE IR..
```

Further work revealed the complete nucleotide sequence <SEQ ID 33>:

```
   1  ATGAAATACT TGATCCGCAC CGCCTTACTC GCAGTCGCAG CCGCCGGCAT
  51  CTACGCCTGC CAACCGCAAT CCGAAGCCGC AGTGCAAGTC AAGGCTGAAA
 101  ACAGCCTGAC CGCTATGCGC TTAGCCGTCG CCGACAAACA GGCAGAGATT
 151  GACGGGTTGA ACGCCCAAAT CGACGCCGAA ATCAGACAAC GCGAAGCCGA
 201  AGAATTGAAA GACTACCGAT GGATACACGG CGACGCGGAA GTGCCGGAGC
 251  TGGAAAAATG A
```

This corresponds to the amino acid sequence <SEQ ID 34; ORF52-1>:

```
   1  MKYLIRTALL AVAAAGIYAC QPQSEAAVQV KAENSLTAMR LAVADKQAEI
  51  DGLNAQIDAE IRQREAEELK DYRWIHGDAE VPELEK*
```

Computer analysis of this amino acid sequence predicts a prokaryotic membrane lipoprotein lipid attachment site (underlined).

ORF52-1 (7kDa) was cloned in the pGex vectors and expressed in *E.coli*, as described above. The products of protein expression and purification were analyzed by SDS-PAGE. Figure 4A shows the results of affinity purification of the GST-fusion. Figure 4B shows plots of hydrophilicity, antigenic index, and AMPHI regions for ORF52-1.

Based on this analysis, it is predicted that this protein from *N.meningitidis*, and its epitopes, could be useful antigens for vaccines or diagnostics.

Example 8

The following DNA sequence was identified in *N.meningitidis* <SEQ ID 35>

```
   1  ATGGTTATCG GAATATTACT CGCATCAAGC AAGCATGCTC TTGTCATTAC
  51  TCTATTGTTA AATCCCGTCT TCCATGCATC CAGTTGCGTA TCGCGTTsGG
 101  CAATACGGAA TAAAAtCTGC TGTTCTGCTT TGGCTAAATT TGCCAAATTG
 151  TTTATTGTTT CTTTAGGaGC AGCTTGCTTA GCCGCCTTCG CTTTCGACAA
 201  CGCCCCCACA GGCGCTTCCC AAGCgTTGCC TACCGTTACC GCACCCGTGG
 251  CGATTCCCGC GCCCGCTTCG GCAGCCTGA
```

This corresponds to the amino acid sequence <SEQ ID 36; ORF56>:

```
   1  MVIGILLASS KHALVITLLL NPVFHASSCV SRXAIRNKIC CSALAKFAKL
  51  FIVSLGAACL AAFAFDNAPT GASQALPTVT APVAIPAPAS AA*
```

Further work revealed the complete nucleotide sequence <SEQ ID 37>:

```
   1  ATGGCTTGTA CAGGTTTGAT GGTTTTTCCG TTAATGGTTA TCGGAATATT
```

```
  51  ACTTGCATCA AGCAAGCCTG CTCCTTTCCT TACTCTATTG TTAAATCCCG
 101  TCTTCCATGC ATCCAGTTGC GTATCGCGTT GGGCAATACG GAATAAAATC
 151  TGCTGTTCTG CTTTGGCTAA ATTTGCCAAA TTGTTTATTG TTTCTTTAGG
 201  AGCAGCTTGC TTAGCCGCCT TCGCTTTCGA CAACGCCCCC ACAGGCGCTT
 251  CCCAAGCGTT GCCTACCGTT ACCGCACCCG TGGCGATTCC CGCGCCCGCT
 301  TCGGCAGCCT GA
```

This corresponds to the amino acid sequence <SEQ ID 38; ORF56-1>:

```
   1  MACTGLMVFP LMVIGILLAS SKPAPFLTLL LNPVFHASSC VSRWAIRNKI
  51  CCSALAKFAK LFIVSLGAAC LAAFAFDNAP TGASQALPTV TAPVAIPAPA
 101  SAA*
```

Computer analysis of this amino acid sequence predicts a leader peptide (underlined) and suggests that ORF56 might be a membrane or periplasmic protein.

Based on this analysis, it is predicted that this protein from *N.meningitidis*, and its epitopes, could be useful antigens for vaccines or diagnostics.

Example 9

The following partial DNA sequence was identified in *N.meningitidis* <SEQ ID 39>

```
   1  ATGTTCAGTA TTTTAAATGT GTTTCTTCAT TGTATTCTGG CTTGTGTAGT
  51  CTCTGGTGAG ACGCCTACTA TATTTGGTAT CCTTGCTCTT TTTTACTTAT
 101  TGTATCTTTC TTATCTTGCT GTTTTTAAGA TTTTCTTTTC TTTTTTCTTA
 151  GACAGAGTTT CACTCCGGTC TCCCAGGCTG GAGTGCAAAT GGCATGACCC
 201  TTTGGCTCAC TGGCTCACGG CCACTTCTGC TATTCTGCCG CCTCAGCCTC
 251  CAGGG...
```

This corresponds to the amino acid sequence <SEQ ID 40; ORF63>:

```
   1  MFSILNVFLH CILACVVSGE TPTIFGILAL FYLLYLSYLA VFKIFFSFFL
  51  DRVSLRSPRL ECKWHDPLAH WLTATSAILP PQPPG...
```

Computer analysis of this amino acid sequence predicts a transmembrane region.

Based on this analysis, it is predicted that this protein from *N.meningitidis*, and its epitopes, could be useful antigens for vaccines or diagnostics.

Example 10

The following partial DNA sequence was identified in *N.meningitidis* <SEQ ID 41>

```
   1  ..GTGCGGACGT GGTTGGTTTT TTGGTTGCAG CGTTTGAAAT ACCCGTTGTT
  51    GCTTTGGATT GCGGATATGT TGCTGTACCG GTTGTTGGGC GGCGCGGAAA
 101    TCGAATGCGG CCGTTGCCCT GTGCCGCCGA TGACGGATTG GCAGCATTTT
 151    TTGCCGGCGA TGGGAACGGT GTCGGCTTGG GTGGCGGTGA TTTGGGCATA
 201    CCTGATGATT GAAAGTGAAA AAAACGGAAG ATATTGA
```

This corresponds to the amino acid sequence <SEQ ID 42; ORF69>:

```
   1  ..VRTWLVFWLQ RLKYPLLLWI ADMLLYRLLG GAEIECGRCP VPPMTDWQHF
  51    LPAMGTVSAW VAVIWAYLMI ESEKNGRY*
```

Computer analysis of this amino acid sequence predicts a transmembrane region.

A corresponding ORF from strain A of *N.meningitidis* was also identified.

Homology with a predicted ORF from *N meningitidis* (strain A)

ORF69 shows 96.2% identity over a 78aa overlap with an ORF (ORF69a) from strain A of *N. meningitidis*.

```
                10        20        30        40        50        60
orf69.pep       VRTWLVFWLQRLKYPLLLWIADMLLYRLLGGAEIECGRCPVPPMTDWQHFLPAMGTVSAW
                ||||||||||||||||||| ||||||||||||||||||||||||||||||:||||:||
orf69a          VRTWLVFWLQRLKYPLLLCIADMLLYRLLGGAEIECGRCPVPPMTDWQHFLPTMGTVAAW
                10        20        30        40        50        60

                70       79
orf69.pep       VAVIWAYLMIESEKNGRYX
                |||||||||||||||||||
orf69a          VAVIWAYLMIESEKNGRYX
                70
```

The ORF69a nucleotide sequence <SEQ ID 43> is:

```
   1  GTGCGGACGT GGTTGGTTTT TTGGTTGCAG CGTTTGAAAT ACCCGTTGTT
  51  GCTTTGTATT GCGGATATGC TGCTGTACCG GTTGTTGGGC GGCGCGGAAA
 101  TCGAATGCGG CCGTTGCCCT GTACCGCCGA TGACGGATTG GCAGCATTTT
 151  TTGCCGACGA TGGGAACGGT GGCGGCTTGG GTGGCGGTGA TTTGGGCATA
 201  CCTGATGATT GAAAGTGAAA AAAACGGAAG ATATTGA
```

This encodes a protein having amino acid sequence <SEQ ID 44>:

```
   1  VRTWLVFWLQ RLKYPLLLCI ADMLLYRLLG GAEIECGRCP VPPMTDWQHF
  51  LPTMGTVAAW VAVIWAYLMI ESEKNGRY*
```

Based on this analysis, it is predicted that this protein from *N.meningitidis*, and its epitopes, could be useful antigens for vaccines or diagnostics.

Example 11

The following DNA sequence was identified in *N.meningitidis* <SEQ ID 45>

```
   1  ATGTTTCAAA ATTTTGATTT GGGCGTGTTC CTGCTTGCCG TCCTCCCCGT
  51  GCTGCCCTCC ATTACCGTCT CGCACGTGGC GCGCGGCTAT ACGGCGCGCT
 101  ACTGGGGAGA CAACACTGCC GAACAATACG GCAGGCTGAC ACTGAACCCC
 151  CTGCCCCATA TCGATTTGGT CGGCACAATC ATCgTACCGC TGCTTACTTT
 201  GATGTTCACG CCCTTCCTGT TCGGCTGGGC GCGTCCGATT CCTATCGATT
 251  CGCGCAACTT CCGCAACCCG cGCCTTGCCT GGCGTTGCGT TGCCGCGTCC
 301  GGCCCGCTGT CGAATCTAGC GATGGCTGTw CTGTGGGGCG TGGTTTTGGT
 351  GCTGACTCCG TATGTCGGCG GGGCGTATCA GATGCCGTTG GCTCAAATGG
 401  CAAACTACGG TATTCTGATC AATGCGATTC TGTTCGCGCT CAACATCATC
 451  CCCATCCTGC CTTGGGACGG CGGCATTTTC ATCGACACCT TCCTGTCGGC
 501  GAAATATTCG CAAGCGTTCC GCAAAATCGA ACCTTATGGG ACGTGGATTA
 551  TCCTACTGCT GATGCTGACC sGGGTTTTGG GTGCGTTTAT wGCACCGATT
 601  sTGCGGmTGc GTGATTGCrT TTGTGCAGAT GTwCGTCTGA CTGGCTTTCA
 651  GACGGCATAA
```

This corresponds to the amino acid sequence <SEQ ID 46; ORF77>:

```
   1  MFQNFDLGVF LLAVLPVLPS ITVSHVARGY TARYWGDNTA EQYGRLTLNP
  51  LPHIDLVGTI IVPLLTLMFT PFLFGWARPI PIDSRNFRNP RLAWRCVAAS
 101  GPLSNLAMAV LWGVVLVLTP YVGGAYQMPL AQMANYGILI NAILFALNII
 151  PILPWDGGIF IDTFLSAKYS QAFRKIEPYG TWIILLLMLT XVLGAFIAPI
 201  XRXRDCXCAD VRLTGFQTA*
```

Further work revealed the complete nucleotide sequence <SEQ ID 47>:

```
   1  ATGTTTCAAA ATTTTGATTT GGGCGTGTTT CTGCTTGCCG TCCTGCCCGT
  51  GCTGCTCTCC ATTACCGTCA GGGAGGTGGC GCGCGGCTAT ACGGCGCGCT
 101  ACTGGGGAGA CAACACTGCC GAACAATACG GCAGGCTGAC ACTGAACCCC
 151  CTGCCCCATA TCGATTTGGT CGGCACAATC ATCGTACCGC TGCTTACTTT
 201  GATGTTCACG CCCTTCCTGT TCGGCTGGGC GCGTCCGATT CCTATCGATT
 251  CGCGCAACTT CCGCAACCCG CGCCTTGCCT GGCGTTGCGT TGCCGCGTCC
 301  GGCCCGCTGT CGAATCTAGC GATGGCTGTT CTGTGGGGCG TGGTTTTGGT
 351  GCTGACTCCG TATGTCGGCG GGGCGTATCA GATGCCGTTG GCTCAAATGG
 401  CAAACTACGG TATTCTGATC AATGCGATTC TGTTCGCGCT CAACATCATC
 451  CCCATCCTGC CTTGGGACGG CGGCATTTTC ATCGACACCT TCCTGTCGGC
 501  GAAATATTCG CAAGCGTTCC GCAAAATCGA ACCTTATGGG ACGTGGATTA
 551  TCCTACTGCT GATGCTGACC GGGGTTTTGG GTGCGTTTAT TGCACCGATT
 601  GTGCGGCTGG TGATTGCGTT TGTGCAGATG TTCGTCTGA
```

This corresponds to the amino acid sequence <SEQ ID 48; ORF77-1>:

```
   1  MFQNFDLGVF LLAVLPVLLS ITVREVARGY TARYWGDNTA EQYGRLTLNP
  51  LPHIDLVGTI IVPLLTLMFT PFLFGWARPI PIDSRNFRNP RLAWRCVAAS
 101  GPLSNLAMAV LWGVVLVLTP YVGGAYQMPL AQMANYGILI NAILFALNII
 151  PILPWDGGIF IDTFLSAKYS QAFRKIEPYG TWIILLLMLT GVLGAFIAPI
 201  VRLVIAFVQM FV*
```

Computer analysis of this amino acid sequence reveals a putative leader sequence and several transmembrane domains.

A corresponding ORF from strain A of *N.meningitidis* was also identified:

Homology with a predicted ORF from *N.meningitidis* (strain A)

ORF77 shows 96.5% identity over a 173aa overlap with an ORF (ORF77a) from strain A of *N meningitidis:*

```
                10        20        30        40        50        60
orf77.pep   MFQNFDLGVFLLAVLPVLPSITVSHVARGYTARYWGDNTAEQYGRLTLNPLPHIDLVGTI
                                      ||||||||||||||||||||||||||||||||||
orf77a                                RGYTARYWGDNTAEQYGRLTLNPLPHIDLVGTI
                                             10        20        30

                70        80        90        100       110       120
orf77.pep   IVPLLTLMFTPFLFGWARPIPIDSRNFRNPRLAWRCVAASGPLSNLAMAVLWGVVLVLTP
            ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
orf77a      IVPLLTLMFTPFLFGWARPIPIDSRNFRNPRLAWRCVAASGPLSNLAMAVLWGVVLVLTP
                 40        50        60        70        80        90

               130       140       150       160       170       180
orf77.pep   YVGGAYQMPLAQMANYGILINAILFALNIIPILPWDGGIFIDTFLSAKYSQAFRKIEPYG
            |||||||||||||||| ||||||| ||||||||||||||||||||| ||||||||||
orf77a      YVGGAYQMPLAQMANYXILINAILXALNIIPILPWDGGIFIDTFLSAKXSQAFRKIEPYG
                100       110       120       130       140       150

                190       200       210       220
orf77.pep   TWIILLLMLTXVLGAFIAPIXRXRDCXCADVRLTGFQTAX
            |||| ||||| |||| ||||
orf77a      TWIIXLLMLTGVLGAXIAPIVQLVIAFVQMFVX
                160       170       180
```

ORF77-1 and ORF77a show 96.8% identity in 185 aa overlap:

```
                    10        20        30        40        50        60
orf77-1.pep  MFQNFDLGVFLLAVLPVLLSITVREVARGYTARYWGDNTAEQYGRLTLNPLPHIDLVGTI
                                          ||||||||||||||||||||||||||||||
orf77a                                    RGYTARYWGDNTAEQYGRLTLNPLPHIDLVGTI
                                                  10        20        30

                    70        80        90       100       110       120
orf77-1.pep  IVPLLTLMFTPFLFGWARPIPIDSRNFRNPRLAWRCVAASGPLSNLAMAVLWGVVLVLTP
             ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
orf77a       IVPLLTLMFTPFLFGWARPIPIDSRNFRNPRLAWRCVAASGPLSNLAMAVLWGVVLVLTP
                  40        50        60        70        80        90

                   130       140       150       160       170       180
orf77-1.pep  YVGGAYQMPLAQMANYGILINAILFALNIIPILPWDGGIFIDTFLSAKYSQAFRKIEPYG
             |||||||||||||||| ||||||| ||||||||||||||||||||| ||||||||||||
orf77a       YVGGAYQMPLAQMANYXILINAILXALNIIPILPWDGGIFIDTFLSAKXSQAFRKIEPYG
                 100       110       120       130       140       150

                   190       200       210
orf77-1.pep  TWIILLLMLTGVLGAFIAPIVRLVIAFVQMFVX
             |||| |||||||||| |||||:||||||||||
orf77a       TWIIXLLMLTGVLGAXIAPIVQLVIAFVQMFVX
                 160       170       180
```

A partial ORF77a nucleotide sequence <SEQ ID 49> was identified:

```
   1  ..CGCGGCTATA CAGCGCGCTA CTGGGGTGAC AACACTGCCG AACAATACGG
  51    CAGGCTGACA CTGAACCCCC TGCCCCATAT CGATTTGGTC GGCACAATCA
 101    TCGTACCGCT GCTTACTTTG ATGTTTACGC CCTTCCTGTT CGGCTGGGCG
 151    CGTCCGATTC CTATCGATTC GCGCAACTTC CGCAACCCGC GCCTTGCCTG
 201    GCGTTGCGTT GCCGCGTCCG GCCCGCTGTC GAATCTGGCG ATGGCTGTTC
 251    TGTGGGGCGT GGTTTTGGTG CTGACTCCGT ATGTCGGTGG GGCGTATCAG
 301    ATGCCGTTGG CNCAAATGGC AAACTACNNN ATTCTGATCA ATGCGATTCT
 351    GTNCGCGCTC AACATCATCC CCATCCTGCC TTGGGACGGC GGCATTTTCA
 401    TCGACACCTT CCTGTCGGCN AAATANTCGC AAGCGTTCCG CAAAATCGAA
 451    CCTTATGGGA CGTGGATTAT CCNGCTGCTT ATGCTGACCG GGGTTTTGGG
 501    TGCGTNTATT GCACCGATTG TGCAGCTGGT GATTGCGTTT GTGCAGATGT
 551    TCGTCTGA
```

This encodes a protein having amino acid sequence <SEQ ID 50>:

```
   1  ..RGYTARYWGD NTAEQYGRLT LNPLPHIDLV GTIIVPLLTL MFTPFLFGWA
  51    RPIPIDSRNF RNPRLAWRCV AASGPLSNLA MAVLWGVVLV LTPYVGGAYQ
 101    MPLAQMANYX ILINAILXAL NIIPILPWDG GIFIDTFLSA KXSQAFRKIE
 151    PYGTWIIXLL MLTGVLGAXI APIVQLVIAF VQMFV*
```

Based on this analysis, it is predicted that this protein from *N.meningitidis*, and its epitopes, could be useful antigens for vaccines or diagnostics.

Example 12

The following partial DNA sequence was identified in *N.meningitidis* <SEQ ID 51>

```
   1  ATGAACCTGA TTTCACGTTA CATCATCCGT CAAATGGCGG TTATGGCGGT
  51  TTACGCGCTC CTTGCCTTCC TCGCTTTGTA CAGCTTTTTT GAAATCCTGT
 101  ACGAAACCGG CAACCTCGGC AAAGGCAGTT ACGGCATATG GGAAATGCTG
 151  GGCTACACCG CCCTCAAAAT GCCCGCCCGC GCCTACGAAC TGATTCCCCT
 201  CGCCGTCCTT ATCGGCGGAC TGGTCTCCCT CAGCCAGCTT GCCGCCGGCA
 251  GCGAACTGAC CGTCATCAAA GCCAGCGGCA TGAGCACCAA AAAGCTGCTG
 301  TTGATTCTGT CGCAGTTCGG TTTTATTTTT GCTATTGCCA CCGTCGCGCT
 351  CGGCGAATGG GTTGCGCCCA CACTGAGCCA AAAAGCCGAA AACATCAAAG
```

```
401  CCGCCGCCAT CAACGGCAAA ATCAGCACCG GCAATACCGG CCTTTGGCTG
451  AAAGAAAAAA ACAGCGTGAT CAATGTGCGC GAAATGTTGC CCGACCAT..
```

This corresponds to the amino acid sequence <SEQ ID 52; ORF112>:

```
  1  MNLISRYIIR QMAVMAVYAL LAFLALYSFF EILYETGNLG KGSYGIWEML
 51  GYTALKMPAR AYELIPLAVL IGGLVSLSQL AAGSELTVIK ASGMSTKKLL
101  LILSQFGFIF AIATVALGEW VAPTLSQKAE NIKAAAINGK ISTGNTGLWL
151  KEKNSVINVR EMLPDH...
```

Further work revealed further partial nucleotide sequence <SEQ ID 53>:

```
  1  ATGAACCTGA TTTCACGTTA CATCATCCGT CAAATGGCGG TTATGGCGGT
 51  TTACGCGCTC CTTGCCTTCC TCGCTTTGTA CAGCTTTTTT GAAATCCTGT
101  ACGAAACCGG CAACCTCGGC AAAGGCAGTT ACGGCATATG GGAAATGCTG
151  gGCTACACCG CCCTCAAAAT GCCCGCCCGC GCCTACGAAC TGATTCCCCT
201  CGCCGTCCTT ATCGGCGGAC TGGTCTCCCT CAGCCAGCTT GCCGCCGGCA
251  GCGAACTGAC CGTCATCAAA GCCAGCGGCA TGAGCACCAA AAAGCTGCTG
301  TTGATTCTGT CGCAGTTCGG TTTTATTTTT GCTATTGCCA CCGTCGCGCT
351  CGGCGAATGG GTTGCGCCCA CACTGAGCCA AAAAGCCGAA AACATCAAAG
401  CCGCCGCCAT CAACGGCAAA ATCAGCACCG GCAATACCGG CCTTTGGCTG
451  AAAGAAAAAA ACAGCrTkAT CAATGTGCGC GAAATGTTGC CCGACCATAC
501  GCTTTTGGGC ATCAAAATTT GGGCGCGCAA CGATAAAAAC GAATTGGCAG
551  AGGCAGTGGA AGCCGATTCC GCCGTTTTGA ACAGCGACGG CAGTTGGCAG
601  TTGAAAAACA TCCGCCGCAG CACGCTTGGC GAAGACAAAG TCGAGGTCTC
651  TATTGCGGCT GAAGAAAACT GGCCGATTTC CGTCAAACGC AACCTGATGG
701  ACGTATTGCT CGTCAAACCC GACCAAATGT CCGTCGGCGA ACTGACCACC
751  TACATCCGCC ACCTCCAAAA CAACAGCCAA AACACCCGAA TCTACGCCAT
801  CGCATGGTGG CGCAAATTGG TTTACCCCGC CGCAGCCTGG GTGATGGCGC
851  TCGTCGCCTT TGCCTTTACC CCGCAAACCA CCCGCCACGG CAATATGGGC
901  TTAAAACTCT TCGGCGGCAT CTGTsTCGGA TTGCTGTTCC ACCTTGCCGG
951  ACGGCTCTTT GGGTTTACCA GCCAACTCGG...
```

This corresponds to the amino acid sequence <SEQ ID 54; ORF112-1>:

```
  1  MNLISRYIIR QMAVMAVYAL LAFLALYSFF EILYETGNLG KGSYGIWEML
 51  GYTALKMPAR AYELIPLAVL IGGLVSLSQL AAGSELTVIK ASGMSTKKLL
101  LILSQFGFIF AIATVALGEW VAPTLSQKAE NIKAAAINGK ISTGNTGLWL
151  KEKNSXINVR EMLPDHTLLG IKIWARNDKN ELAEAVEADS AVLNSDGSWQ
201  LKNIRRSTLG EDKVEVSIAA EENWPISVKR NLMDVLLVKP DQMSVGELTT
251  YIRHLQNNSQ NTRIYAIAWW RKLVYPAAAW VMALVAFAFT PQTTRHGNMG
301  LKLFGGICXG LLFHLAGRLF GFTSQL...
```

Computer analysis of this amino acid sequence predicts two transmembrane domains.

A corresponding ORF from strain A of *N.meningitidis* was also identified:

Homology with a predicted ORF from *N.meningitidis* (strain A)

ORF112 shows 96.4% identity over a 166aa overlap with an ORF (ORF112a) from strain A of *N. meningitidis*.

```
                10        20        30        40        50        60
orf112.pep      MNLISRYIIRQMAVMAVYALLAFLALYSFFEILYETGNLGKGSYGIWEMLGYTALKMPAR
                ||||||||||||||||||||||||||||||||||||||||||||||||| ||||||| ||
orf112a         MNLISRYIIRQMAVMAVYALLAFLALYSFFEILYETGNLGKGSYGIWEMXGYTALKMXAR
                10        20        30        40        50        60

                70        80        90        100       110       120
orf112.pep      AYELIPLAVLIGGLVSLSQLAAGSELTVIKASGMSTKKLLLLILSQFGFIFAIATVALGEW
                ||||:|||||||||||| |||||||||:|||||||||||||||||||||||||||||||||
orf112a         AYELMPLAVLIGGLVSXSQLAAGSELXVIKASGMSTKKLLLLILSQFGFIFAIATVALGEW
                70        80        90        100       110       120
```

```
                    130       140       150       160
orf112.pep  VAPTLSQKAENIKAAAINGKISTGNTGLWLKEKNSVINVREMLPDH
            ||||||||||||||||||||||||||||||||||||:||||||||||
orf112a     VAPTLSQKAENIKAAAINGKISTGNTGLWLKEKNSIINVREMLPDHTLLGIKIWARNDKN
                    130       140       150       160       170       180

orf112a     ELAEAVEADSAVLNSDGSWQLKNIRRSTLGEDKVEVSIAAEEXWPISVKRNLMDVLLVKP
                    190       200       210       220       230       240
```

A partial ORF112a nucleotide sequence <SEQ ID 55> was identified:

```
     1  ATGAACCTGA TTTCACGTTA CATCATCCGT CAAATGGCGG TTATGGCGGT
    51  TTACGCGCTC CTTGCCTTCC TCGCTTTGTA CAGCTTTTTT GAAATCCTGT
   101  ACGAAACCGG CAACCTCGGC AAAGGCAGTT ACGGCATATG GGAAATGNTG
   151  GGNTACACCG CCCTCAAAAT GNCCGCCCGC GCCTACGAAC TGATGCCCCT
   201  CGCCGTCCTT ATCGGCGGAC TGGTCTCTNT CAGCCAGCTT GCCGCCGGCA
   251  GCGAACTGAN CGTCATCAAA GCCAGCGGCA TGAGCACCAA AAAGCTGCTG
   301  TTGATTCTGT CGCAGTTCGG TTTTATTTTT GCTATTGCCA CCGTCGCGCT
   351  CGGCGAATGG GTTGCGCCCA CACTGAGCCA AAAAGCCGAA AACATCAAAG
   401  CCGCGGCCAT CAACGGCAAA ATCAGTACCG GCAATACCGG CCTTTGGCTG
   451  AAAGAAAAAA ACAGCATTAT CAATGTGCGC GAAATGTTGC CCGACCATAC
   501  CCTGCTGGGC ATTAAAATCT GGGCCCGCAA CGATAAAAAC GAACTGGCAG
   551  AGGCAGTGGA AGCCGATTCC GCCGTTTTGA ACAGCGACGG CAGTTGGCAG
   601  TTGAAAAACA TCCGCCGCAG CACGCTTGGC GAAGACAAAG TCGAGGTCTC
   651  TATTGCGGCT GAAGAAAANT GGCCGATTTC CGTCAAACGC AACCTGATGG
   701  ACGTATTGCT CGTCAAACCC GACCAAATGT CCGTCGGCGA ACTGACCACC
   751  TACATCCGCC ACCTCCAAAN NNACAGCCAA AACACCCGAA TCTACGCCAT
   801  CGCATGGTGG CGCAAATTGG TTTACCCCGC CGCAGCCTGG GTGATGGCGC
   851  TCGTCGCCTT TGCCTTTACC CCGCAAACCA CCCGCCACGG CAATATGGGC
   901  TTAAAANTCT TCGGCGGCAT CTGTCTCGGA TTGCTGTTCC ACCTTGCCGG
   951  NCGGCTCTTC NGGTTTACCA GCCAACTCTA CGGCATCCCG CCCTTCCTCG
  1001  NCGGCGCACT ACCTACCATA GCCTTCGCCT TGCTCGCCGT TTGGCTGATA
  1051  CGCAAACAGG AAAAACGCTA A
```

This encodes a protein having amino acid sequence <SEQ ID 56>:

```
     1  MNLISRYIIR QMAVMAVYAL LAFLALYSFF EILYETGNLG KGSYGIWEMX
    51  GYTALKMXAR AYELMPLAVL IGGLVSXSQL AAGSELXVIK ASGMSTKKLL
   101  LILSQFGFIF AIATVALGEW VAPTLSQKAE NIKAAAINGK ISTGNTGLWL
   151  KEKNSIINVR EMLPDHTLLG IKIWARNDKN ELAEAVEADS AVLNSDGSWQ
   201  LKNIRRSTLG EDKVEVSIAA EEXWPISVKR NLMDVLLVKP DQMSVGELTT
   251  YIRHLQXXSQ NTRIYAIAWW RKLVYPAAAW VMALVAFAFT PQTTRHGNMG
   301  LKXFGGICLG LLFHLAGRLF XFTSQLYGIP PFLXGALPTI AFALLAVWLI
   351  RKQEKR*
```

ORF112a and ORF112-1 show 96.3% identity in 326 aa overlap:

```
orf112a.pep   MNLISRYIIRQMAVMAVYALLAFLALYSFFEILYETGNLGKGSYGIWEMXGYTALKMXAR
              ||||||||||||||||||||||||||||||||||||||||||||||||||||||| ||||||| ||
orf112-1      MNLISRYIIRQMAVMAVYALLAFLALYSFFEILYETGNLGKGSYGIWEMLGYTALKMPAR

orf112a.pep   AYELMPLAVLIGGLVSXSQLAAGSELXVIKASGMSTKKLLLILSQFGFIFAIATVALGEW
              ||||:|||||||||||| |||||||||:|||||||||||||||||||||||||||||||
orf112-1      AYELIPLAVLIGGLVSLSQLAAGSELTVIKASGMSTKKLLLILSQFGFIFAIATVALGEW

orf112a.pep   VAPTLSQKAENIKAAAINGKISTGNTGLWLKEKNSIINVREMLPDHTLLGIKIWARNDKN
              |||||||||||||||||||||||||||||||| ||||||||||||||||||||||||||
orf112-1      VAPTLSQKAENIKAAAINGKISTGNTGLWLKEKNSXINVREMLPDHTLLGIKIWARNDKN

orf112a.pep   ELAEAVEADSAVLNSDGSWQLKNIRRSTLGEDKVEVSIAAEEXWPISVKRNLMDVLLVKP
              |||||||||||||||||||||||||||||||||||||||| ||||||||||||||||
orf112-1      ELAEAVEADSAVLNSDGSWQLKNIRRSTLGEDKVEVSIAAEENWPISVKRNLMDVLLVKP

orf112a.pep   DQMSVGELTTYIRHLQXXSQNTRIYAIAWWRKLVYPAAAWVMALVAFAFTPQTTRHGNMG
              ||||||||||||||| ||||||||||||||||||||||||||||||||||||||||||
orf112-1      DQMSVGELTTYIRHLQNNSQNTRIYAIAWWRKLVYPAAAWVMALVAFAFTPQTTRHGNMG

orf112a.pep   LKXFGGICLGLLFHLAGRLFXFTSQLYGIPPFLXGALPTIAFALLAVWLIRKQEKRX
              || ||||| |||||||||||| |||||
```

```
orf112-1        LKLFGGICXGLLFHLAGRLFGFTSQL
```

Based on this analysis, it is predicted that this protein from *N meningitidis*, and its epitopes, could be useful antigens for vaccines or diagnostics.

Example 13

The following partial DNA sequence was identified in *N.meningitidis* <SEQ ID 57>

```
   1  ..GCAGTAGCCG AAACTGCCAA CAGCCAGGGC AAAGGTAAAC AGGCAGGCAG
  51    TTCGGTTTCT GTTTCACTGA AAACTTCAGG CGACCTTTGC GGCAAACTCA
 101    AAACCACCCT TAAAACTTTG GTCTGCTCTT TGGTTTCCCT GAGTATGGTA
 151    TTGCCTGCCC ATGCCCAAAT TACCACCGAC AAATCAGCAC CTAAAAACCA
 201    GCAGGTCGTT ATCCTTAAAA CCAACACTGG TGCCCCCTTG GTGAATATCC
 251    AAACTCCGAA TGGACGCGGA TTGAGCCACA ACCGCTA.TA CGCATTTGAT
 301    GTTGACAACA AAGGGGCAGT GTTAAACAAC GACCGTAACA ATAATCCGTT
 351    TGTGGTCAAA GGCAGTGCGC AATTGATTTT GAACGAGGTA CGCGGTACGG
 401    CTAGCAAACT CAACGGCATC GTTACCGTAG GCGGTCAAAA GGCCGACGTG
 451    ATTATTGCCA ACCCCAACGG CATTACCGTT AATGGCGGCG GCTTTAAAAA
 501    TGTCGGTCGG GGCATCTTAA CTACCGGTGC GCCCCAAATC GGCAAAGACG
 551    GTGCACTGAC AGGATTTGAT GTGCGTCAAG GCACATTGgA CCGTAGrAGC
 601    AGCAGGTTGG AATGATAAAG GCGGAGCmrm yTACACCGGG GTACTTGCTC
 651    GTGCAGTTGC TTTGCAGGGG AAATTwmmGG GTAAA.AACT GGCGGTTTCT
 701    ACCGGTCCTC AGAAAGTAGA TTACGCCAGC GGCGAAATCA GTGCAGGTAC
 751    GGCAGCGGGT ACGAAACCGA CTATTGCCCT TGATACTGCC GCACTGGGCG
 801    GTATGTACGC CGACAGCATC ACACTGATTG CCAATGAAAA AGGCGTAGGC
 851    GTCTAA
```

This corresponds to the amino acid sequence <SEQ ID 58; ORF114>:

```
   1  ..AVAETANSQG KGKQAGSSVS VSLKTSGDLC GKLKTTLKTL VCSLVSLSMV
  51    LPAHAQITTD KSAPKNQQVV ILKTNTGAPL VNIQTPNGRG LSHNRXYAFD
 101    VDNKGAVLNN DRNNNPFVVK GSAQLILNEV RGTASKLNGI VTVGGQKADV
 151    IIANPNGITV NGGGFKNVGR GILTTGAPQI GKDGALTGFD VVKAHWTVXA
 201    AGWNDKGGAX YTGVLARAVA LQGKXXGKXL AVSTGPQKVD YASGEISAGT
 251    AAGTKPTIAL DTAALGGMYA DSITLIANEK GVGV*
```

Further work revealed the complete nucleotide sequence <SEQ ID 59>:

```
   1  ATGAATAAAG GTTTACATCG CATTATCTTT AGTAAAAAGC ACAGCACCAT
  51  GGTTGCAGTA GCCGAAACTG CCAACAGCCA GGGCAAAGGT AAACAGGCAG
 101  GCAGTTCGGT TTCTGTTTCA CTGAAAACTT CAGGCGACCT TTGCGGCAAA
 151  CTCAAAACCA CCCTTAAAAC TTTGGTCTGC TCTTTGGTTT CCCTGAGTAT
 201  GGTATTGCCT GCCCATGCCC AAATTACCAC CGACAAATCA GCACCTAAAA
 251  ACCAGCAGGT CGTTATCCTT AAAACCAACA CTGGTGCCCC CTTGGTGAAT
 301  ATCCAAACTC CGAATGGACG CGGATTGAGC CACAACCGCT ATACGCAGTT
 351  TGATGTTGAC AACAAAGGGG CAGTGTTAAA CAACGACCGT AACAATAATC
 401  CGTTTGTGGT CAAAGGCAGT GCGCAATTGA TTTTGAACGA GGTACGCGGT
 451  ACGGCTAGCA AACTCAACGG CATCGTTACC GTAGGCGGTC AAAAGGCCGA
 501  CGTGATTATT GCCAACCCCA ACGGCATTAC CGTTAATGGC GGCGGCTTTA
 551  AAAATGTCGG TCGGGGCATC TTAACTACCG GTGCGCCCCA AATCGGCAAA
 601  GACGGTGCAC TGACAGGATT TGATGTGCGT CAAGGCACAT TGACCGTAGG
 651  AGCAGCAGGT TGGAATGATA AAGGCGGAGC CGACTACACC GGGGTACTTG
 701  CTCGTGCAGT TGCTTTGCAG GGGAAATTAC AGGGTAAAAA CCTGGCGGTT
 751  TCTACCGGTC CTCAGAAAGT AGATTACGCC AGCGGCGAAA TCAGTGCAGG
 801  TACGGCAGCG GGTACGAAAC CGACTATTGC CCTTGATACT GCCGCACTGG
 851  GCGGTATGTA CGCCGACAGC ATCACACTGA TTGCCAATGA AAAAGGCGTA
 901  GGCGTCAAAA ATGCCGGCAC ACTCGAAGCG GCCAAGCAAT TGATTGTGAC
 951  TTCGTCAGGC CGCATTGAAA ACAGCGGCCG CATCGCCACC ACTGCCGACG
1001  GCACCGAAGC TTCACCGACT TATCTCTCCA TCGAAACCAC CGAAAAAGGA
1051  GCGGCAGGCA CATTTATCTC CAATGGTGGT CGGATCGAGA GCAAAGGCTT
1101  ATTGGTTATT GAGACGGGAG AAGATATCAG CTTGCGTAAC GGAGCCGTGG
1151  TGCAGAATAA CGGCAGTCGC CCAGCTACCA CGGTATTAAA TGCTGGTCAT
1201  AATTTGGTGA TTGAGAGCAA AACTAATGTG AACAATGCCA AAGGCCCGGC
```

```
1251  TACTCTGTCG GCCGACGGCC GTACCGTCAT CAAGGAGGCC AGTATTCAGA
1301  CTGGCACTAC CGTATACAGT TCCAGCAAAG GCAACGCCGA ATTAGGCAAT
1351  AACACACGCA TTACCGGGGC AGATGTTACC GTATTATCCA ACGGCACCAT
1401  CAGCAGTTCC GCCGTAATAG ATGCCAAAGA CACCGCACAC ATCGAAGCAG
1451  GCAAACCGCT TTCTTTGGAA GCTTCAACAG TTACCTCCGA TATCCGCTTA
1501  AACGGAGGCA GTATCAAGGG CGGCAAGCAG CTTGCTTTAC TGGCAGACGA
1551  TAACATTACT GCCAAAACTA CCAATCTGAA TACTCCCGGC AATCTGTATG
1601  TTCATACAGG TAAAGATCTG AATTTGAATG TTGATAAAGA TTTGTCTGCC
1651  GCCAGCATCC ATTTGAAATC GGATAACGCT GCCCATATTA CCGGCACCAG
1701  TAAAACCCTC ACTGCCTCAA AAGACATGGG TGTGGAGGCA GGCTCGCTGA
1751  ATGTTACCAA TACCAATCTG CGTACCAACT CGGGTAATCT GCACATTCAG
1801  GCAGCCAAAG GCAATATTCA GCTTCGCAAT ACCAAGCTGA ACGCAGCCAA
1851  GGCTCTCGAA ACCACCGCAT TGCAGGGCAA TATCGTTTCA GACGGCCTTC
1901  ATGCTGTTTC TGCAGACGGT CATGTATCCT TATTGGCCAA CGGTAATGCC
1951  GACTTTACCG GTCACAATAC CCTGACAGCC AAGGCCGATG TCAATGCAGG
2001  ATCGGTTGGT AAAGGCCGTC TGAAAGCAGA CAATACCAAT ATCACTTCAT
2051  CTTCAGGAGA TATTACGTTG GTTGCCGGCA ACGGTATTCA GCTTGGTGAC
2101  GGAAAACAAC GCAATTCAAT CAACGGAAAA CACATCAGCA TCAAAAACAA
2151  CGGTGGTAAT GCCGACTTAA AAAACCTTAA CGTCCATGCC AAAAGCGGGG
2201  CATTGAACAT TCATTCCGAC CGGGCATTGA GCATAGAAAA TACCAAGCTG
2251  GAGTCTACCC ATAATACGCA TCTTAATGCA CAACACGAGC GGGTAACGCT
2301  CAACCAAGTA GATGCCTACG CACACCGTCA TCTAAGCATT ACCGGCAGCC
2351  AGATTTGGCA AAACGACAAA CTGCCTTCTG CCAACAAGCT GGTGGCTAAC
2401  GGTGTATTGG CACTCAATGC GCGCTATTCC CAAATTGCCG ACAACACCAC
2451  GCTGAGAGCG GGTGCAATCA ACCTTACTGC CGGTACCGCC CTAGTCAAGC
2501  GCGGCAACAT CAATTGGAGT ACCGTTTCGA CCAAAACTTT GGAAGATAAT
2551  GCCGAATTAA AACCATTGGC CGGACGGCTG AATATTGAAG CAGGTAGCGG
2601  CACATTAACC ATCGAACCTG CCAACCGCAT CAGTGCGCAT ACCGACCTGA
2651  GCATCAAAAC AGGCGGAAAA TTGCTGTTGT CTGCAAAAGG AGGAAATGCA
2701  GGTGCGCCTA GTGCTCAAGT TTCCTCATTG GAAGCAAAAG GCAATATCCG
2751  TCTGGTTACA GGAGAAACAG ATTTAAGAGG TTCTAAAATT ACAGCCGGTA
2801  AAAACTTGGT TGTCGCCACC ACCAAAGGCA AGTTGAATAT CGAAGCCGTA
2851  AACAACTCAT TCAGCAATTA TTTTCCTACA CAAAAAGCGG CTGAACTCAA
2901  CCAAAAATCC AAAGAATTGG AACAGCAGAT TGCGCAGTTG AAAAAAAGCT
2951  CGCCTAAAAG CAAGCTGATT CCAACCCTGC AAGAAGAACG CGACCGTCTC
3001  GCTTTCTATA TTCAAGCCAT CAACAAGGAA GTTAAAGGTA AAAAACCCAA
3051  AGGCAAAGAA TACCTGCAAG CCAAGCTTTC TGCACAAAAT ATTGACTTGA
3101  TTTCCGCACA AGGCATCGAA ATCAGCGGTT CCGATATTAC CGCTTCCAAA
3151  AAACTGAACC TTCACGCCGC AGGCGTATTG CCAAAGGCAG CAGATTCAGA
3201  GGCGGCTGCT ATTCTGATTG ACGGCATAAC CGACCAATAT GAAATTGGCA
3251  AGCCCACCTA CAAGAGTCAC TACGACAAAG CTGCTCTGAA CAAGCCTTCA
3301  CGTTTGACCG GACGTACAGG GGTAAGTATT CATGCAGCTG CGGCACTCGA
3351  TGATGCACGT ATTATTATCG GTGCATCCGA AATCAAAGCT CCCTCAGGCA
3401  GCATAGACAT CAAAGCCCAT AGTGATATTG TACTGGAGGC TGGACAAAAC
3451  GATGCCTATA CCTTCTTAAA AACCAAAGGT AAAAGCGGCA AAATCATCAG
3501  AAAAACCAAG TTTACCAGCA CCCGCGACCA CCTGATTATG CCAGCCCCCG
3551  TCGAGCTGAC CGCCAACGGC ATAACGCTTC AGGCAGGCGG CAACATCGAA
3601  GCTAATACCA CCCGCTTCAA TGCCCCTGCA GGTAAAGTTA CCCTGGTTGC
3651  GGGTGAAGAG CTGCAACTGC TGGCAGAAGA AGGCATCCAC AAGCACGAGT
3701  TGGATGTCCA AAAAAGCCGC CGCTTTATCG GCATCAAGGT AGGCAAGAGC
3751  AATTACAGTA AAAACGAACT GAACGAAACC AAATTGCCTG TCCGCGTCGT
3801  CGCCCAAACT GCAGCCACCC GTTCAGGCTG GGATACCGTG CTCGAAGGTA
3851  CCGAATTCAA AACCACGCTG GCCGGTGCGG ACATTCAGGC AGGTGTAGGC
3901  GAAAAAGCCC GTGCCGATGC GAAAATTATC CTCAAAGGCA TTGTGAACCG
3951  TATCCAGTCG GAAGAAAAAT TAGAAACCAA CTCAACCGTA TGGCAGAAAC
4001  AGGCCGGACG CGGCAGCACT ATCGAAACGC TGAAACTGCC CAGCTTCGAA
4051  AGCCCTACTC CGCCCAAACT GACCGCCCCC GGTGGCTATA TCGTCGACAT
4101  TCCGAAAGGC AATTTGAAAA CCGAAATCGA AAAGCTGGCC AAACAGCCCG
4151  AGTATGCCTA TCTGAAACAG CTCCAAGTAG CGAAAAACGT CAACTGGAAC
4201  CAGGTGCAAC TGGCTTACGA TAAATGGGAC TATAAGCAGG AAGGCTTAAC
4251  CAGAGCCGGT GCAGCGATTG TTACCATAAT CGTAACCGCA CTGACTTATG
4301  GATACGGCGC AACCGCAGCG GGCGGTGTAG CCGCTTCAGG AAGTAGTACA
4351  GCCGCAGCTG CCGGAACAGC CGCCACAACG ACAGCAGCAG CTACTACCGT
4401  TTCTACAGCG ACTGCCATGC AAACCGCTGC TTTAGCCTCC TTGTATAGCC
4451  AAGCAGCTGT ATCCATCATC AATAATAAAG GTGATGTCGG CAAAGCGTTG
4501  AAAGATCTCG GCACCAGTGA TACGGTCAAG CAGATTGTCA CTTCTGCCCT
4551  GACGGCGGGT GCATTAAATC AGATGGGCGC AGATATTGCC CAATTGAACA
4601  GCAAGGTAAG AACCGAACTG TTCAGCAGTA CGGGCAATCA AACTATTGCC
4651  AACCTTGGAG GCAGACTGGC TACCAATCTC AGTAATGCAG GTATCTCAGC
4701  TGGTATCAAT ACCGCCGTCA ACGGCGGCAG CCTGAAAGAC AACTTAGGCA
4751  ATGCCGCATT AGGAGCATTG GTTAATAGCT TCCAAGGAGA AGCCGCCAGC
4801  AAAATCAAAA CAACCTTCAG CGACGATTAT GTTGCCAAAC AGTTCGCCCA
```

```
4851  CGCTTTGGCT GGGTGTGTTA GCGGATTGGT ACAAGGAAAA TGTAAAGACG
4901  GGGCAATTGG CGCAGCAGTT GGGGAAATCG TAGCCGACTC CATGCTTGGC
4951  GGCAGAAACC CTGCTACACT CAGCGATGCG GAAAAGCATA AGGTTATCAG
5001  TTACTCGAAG ATTATTGCCG GCAGCGTGGC GGCACTCAAC GGCGGCGATG
5051  TGAATACTGC GGCGAATGCG GCTGAGGTGG CGGTAGTGAA TAATGCTTTG
5101  AATTTTGACA GTACCCCTAC CAATGCGAAA AAGCATCAAC CGCAGAAGCC
5151  CGACAAAACC GCACTGGAAA AAATTATCCA AGGTATTATG CCTGCACATG
5201  CAGCAGGTGC GATGACTAAT CCGCAGGATA AGGATGCTGC CATTTGGATA
5251  AGCAATATCC GTAATGGCAT CACAGGCCCG ATTGTGATTA CCAGCTATGG
5301  GGTTTATGCT GCAGGTTGGA CAGCTCCGCT GATCGGTACA GCGGGTAAAT
5351  TAGCTATCAG CACCTGCATG GCTAATCCTT CTGGTTGTAC TGTCATGGTC
5401  ACTCAGGCTG CCGAAGCGGG CGCGGGAATC GCCACGGGTG CGGTAACGGT
5451  AGGCAACGCT TGGGAAGCGC CTGTGGGGGC GTTGTCGAAA GCGAAGGCGG
5501  CCAAGCAGGC TATACCAACC CAGACAGTTA AAGAACTTGA TGGCTTACTA
5551  CAAGAATCAA AAAATATAGG TGCTGTAAAT ACACGAATTA ATATAGCGAA
5601  TAGTACTACT CGATATACAC CAATGAGACA AACGGGACAA CCGGTATCTG
5651  CTGGCTTTGA GCATGTTCTT GAGGGGCACT TCCATAGGCC TATTGCGAAT
5701  AACCGTTCAG TTTTTACCAT CTCCCCAAAT GAATTGAAGG TTATACTTCA
5751  AAGTAATAAA GTAGTTTCTT CTCCCGTATC GATGACTCCT GATGGCCAAT
5801  ATATGCGGAC TGTCGATGTA GGAAAAGTTA TTGGTACTAC TTCTATTAAA
5851  GAAGGTGGAC AACCCACAAC TACAATTAAA GTATTTACAG ATAAGTCAGG
5901  AAATTTGATT ACTACATACC CAGTAAAAGG AAACTAA
```

This corresponds to the amino acid sequence <SEQ ID 60; ORF114-1>:

```
   1  MNKGLHRIIF SKKHSTMVAV AETANSQGKG KQAGSSVSVS LKTSGDLCGK
  51  LKTTLKTLVC SLVSLSMVLP AHAQITTDKS APKNQQVVIL KTNTGAPLVN
 101  IQTPNGRGLS HNRYTQFDVD NKGAVLNNDR NNNPFVVKGS AQLILNEVRG
 151  TASKLNGIVT VGGQKADVII ANPNGITVNG GGFKNVGRGI LTTGAPQIGK
 201  DGALTGFDVR QGTLTVGAAG WNDKGGADYT GVLARAVALQ GKLQGKNLAV
 251  STGPQKVDYA SGEISAGTAA GTKPTIALDT AALGGMYADS ITLIANEKGV
 301  GVKNAGTLEA AKQLIVTSSG RIENSGRIAT TADGTEASPT YLSIETTEKG
 351  AAGTFISNGG RIESKGLLVI ETGEDISLRN GAVVQNNGSR PATTVLNAGH
 401  NLVIESKTNV NNAKGPATLS ADGRTVIKEA SIQTGTTVYS SSKGNAELGN
 451  NTRITGADVT VLSNGTISSS AVIDAKDTAH IEAGKPLSLE ASTVTSDIRL
 501  NGGSIKGGKQ LALLADDNIT AKTTNLNTPG NLYVHTGKDL NLNVDKDLSA
 551  ASIHLKSDNA AHITGTSKTL TASKDMGVEA GSLNVTNTNL RTNSGNLHIQ
 601  AAKGNIQLRN TKLNAAKALE TTALQGNIVS DGLHAVSADG HVSLLANGNA
 651  DFTGHNTLTA KADVNAGSVG KGRLKADNTN ITSSSGDITL VAGNGIQLGD
 701  GKQRNSINGK HISIKNNGGN ADLKNLNVHA KSGALNIHSD RALSIENTKL
 751  ESTHNTHLNA QHERVTLNQV DAYAHRHLSI TGSQIWQNDK LPSANKLVAN
 801  GVLALNARYS QIADNTTLRA GAINLTAGTA LVKRGNINWS TVSTKTLEDN
 851  AELKPLAGRL NIEAGSGTLT IEPANRISAH TDLSIKTGGK LLLSAKGGNA
 901  GAPSAQVSSL EAKGNIRLVT GETDLRGSKI TAGKNLVVAT TKGKLNIEAV
 951  NNSFSNYFPT QKAAELNQKS KELEQQIAQL KKSSPKSKLI PTLQEERDRL
1001  AFYIQAINKE VKGKKPKGKE YLQAKLSAQN IDLISAQGIE ISGSDITASK
1051  KLNLHAAGVL PKAADSEAAA ILIDGITDQY EIGKPTYKSH YDKAALNKPS
1101  RLTGRTGVSI HAAAALDDAR IIIGASEIKA PSGSIDIKAH SDIVLEAGQN
1151  DAYTFLKTKG KSGKIIRKTK FTSTRDHLIM PAPVELTANG ITLQAGGNIE
1201  ANTTRFNAPA GKVTLVAGEE LQLLAEEGIH KHELDVQKSR RFIGIKVGKS
1251  NYSKNELNET KLPVRVVAQT AATRSGWDTV LEGTEFKTTL AGADIQAGVG
1301  EKARADAKII LKGIVNRIQS EEKLETNSTV WQKQAGRGST IETLKLPSFE
1351  SPTPPKLTAP GGYIVDIPKG NLKTEIEKLA KQPEYAYLKQ LQVAKNVNWN
1401  QVQLAYDKWD YKQEGLTRAG AAIVTIIVTA LTYGYGATAA GGVAASGSST
1451  AAAAGTAATT TAAATTVSTA TAMQTAALAS LYSQAAVSII NNKGDVGKAL
1501  KDLGTSDTVK QIVTSALTAG ALNQMGADIA QLNSKVRTEL FSSTGNQTIA
1551  NLGGRLATNL SNAGISAGIN TAVNGGSLKD NLGNAALGAL VNSFQGEAAS
1601  KIKTTFSDDY VAKQFAHALA GCVSGLVQGK CKDGAIGAAV GEIVADSMLG
1651  GRNPATLSDA EKHKVISYSK IIAGSVAALN GGDVNTAANA AEVAVVNNAL
1701  NFDSTPTNAK KHQPQKPDKT ALEKIIQGIM PAHAAGAMTN PQDKDAAIWI
1751  SNIRNGITGP IVITSYGVYA AGWTAPLIGT AGKLAISTCM ANPSGCTVMV
1801  TQAAEAGAGI ATGAVTVGNA WEAPVGALSK AKAAKQAIPT QTVKELDGLL
1851  QESKNIGAVN TRINIANSTT RYTPMRQTGQ PVSAGFEHVL EGHFHRPIAN
1901  NRSVFTISPN ELKVILQSNK VVSSPVSMTP DGQYMRTVDV GKVIGTTSIK
1951  EGGQPTTTIK VFTDKSGNLI TTYPVKGN*
```

Computer analysis of this amino acid sequence predicts a transmembrane region and also gives the following results:

Homology with a predicted ORF from *N.meningitidis* (strain A)

ORF114 shows 91.9% identity over a 284aa overlap with an ORF (ORF114a) from strain A of *N meningitidis:*

```
                                          10        20        30        40
orf114.pep                          AVAETANSQGKGKQAGSSVSVSLKTSGDLCGKLKTTLKTLVC
                                    ||||||||||||||||||||||||||||||||||||||||||
orf114a      MNKGLHRIIFSKKHSTMVAVAETANSQGKGKQAGSSVSVSLKTSGDLCGKLKTTLKTLVC
                    10        20        30        40        50        60

                     50        60        70        80        90       100
orf114.pep   SLVSLSMVLPAHAQITTDKSAPKNQQVVILKTNTGAPLVNIQTPNGRGLSHNRXYAFDVD
             |||||||     ||||||||||| |||||||||||||||||||||||||||    ||||
orf114a      SLVSLSMXXXXXXQITTDKSAPKNXQVVILKTNTGAPLVNIQTPNGRGLSHNRYTQFDVD
                    70        80        90       100       110       120

                   110       120       130       140       150       160
orf114.pep   NKGAVLNNDRNNNPFVVKGSAQLILNEVRGTASKLNGIVTVGGQKADVIIANPNGITVNG
             ||||||||||||||:|||||||||||||||||||||||||||||||||||||||||||||
orf114a      NKGAVLNNDRNNNPFLVKGSAQLILNEVRGTASKLNGIVTVGGQKADVIIANPNGITVNG
                    130       140       150       160       170       180

                   170       180       190       200       210       220
orf114.pep   GGFKNVGRGILTTGAPQIGKDGALTGFDVVKAHWTVXAAGWNDKGGAXYTGVLARAVALQ
             |||||||||||| |||||||||||||| ::  || |||||||||| ||||||||||||
orf114a      GGFKNVGRGILTIGAPQIGKDGALTGFDVRQGTLTVGAAGWNDKGGADYTGVLARAVALQ
                    190       200       210       220       230       240

                   230       240       250       260       270       280
orf114.pep   GKXXGKXLAVSTGPQKVDYASGEISAGTAAGTKPTIALDTAALGGMYADSITLIANEKGV
             ||  || ||||||||||||||||||||||||||||||||||||||||||||||||| ||||
orf114a      GKLQGKNLAVSTGPQKVDYASGEISAGTAAGTKPTIALDTAALGGMYADSITLIAXEKGV
                    250       260       270       280       290       300

orf114.pep   GVX
             ||
orf114a      GVKNAGTLEAAKQLIVTSSGRIENSGRIATTADGTEASPTYLXIETTEKGAXGTFISNGG
                    310       320       330       340       350       360
```

The complete length ORF114a nucleotide sequence <SEQ ID 61> is:

```
   1  ATGAATAAAG GTTTACATCG CATTATCTTT AGTAAAAAGC ACAGCACCAT
  51  GGTTGCAGTA GCCGAAACTG CCAACAGCCA GGGCAAAGGT AAACAGGCAG
 101  GCAGTTCGGT TTCTGTTTCA CTGAAAACTT CAGGCGACCT TTGCGGCAAA
 151  CTCAAAACCA CCCTTAAAAC CTTGGTCTGC TCTTTGGTTT CCCTGAGTAT
 201  GGNATTNCNN NNCNNTNCCC AAATTACCAC CGACAAATCA GCACCTAAAA
 251  ACCANCAGGT CGTTATCCTT AAAACCAACA CTGGTGCCCC CTTGGTGAAT
 301  ATCCAAACTC CGAATGGACG CGGATTGAGC CACAACCGCT ATACGCAGTT
 351  TGATGTTGAC AACAAAGGGG CAGTGTTAAA CAACGACCGT AACAATAATC
 401  CGTTTCTGGT CAAAGGCAGT GCGCAATTGA TTTTGAACGA GGTACGCGGT
 451  ACGGCTAGCA AACTCAACGG CATCGTTACC GTAGGCGGTC AAAAGGCCGA
 501  CGTGATTATT GCCAACCCCA ACGGCATTAC CGTTAATGGC GGCGGCTTTA
 551  AAAATGTCGG TCGGGGCATC TTAACTATCG GTGCGCCCCA AATCGGCAAA
 601  GACGGTGCAC TGACAGGATT TGATGTGCGT CAAGGCACAT TGACCGTAGG
 651  AGCAGCAGGT TGGAATGATA AAGGCGGAGC CGACTACACC GGGGTACTTG
 701  CTCGTGCAGT TGCTTTGCAG GGGAAATTAC AGGGTAAAAA CCTGGCGGTT
 751  TCTACCGGTC CTCAGAAAGT AGATTACGCC AGCGGCGAAA TCAGTGCAGG
 801  TACGGCAGCG GGTACGAAAC CGACTATTGC CCTTGATACT GCCGCACTGG
 851  GCGGTATGTA CGCCGACAGC ATCACACTGA TTGCCANTGA AAAAGGCGTA
 901  GGCGTCAAAA ATGCCGGCAC ACTCGAAGCG GCCAAGCAAT TGATTGTGAC
 951  TTCGTCAGGC CGCATTGAAA ACAGCGGCCG CATCGCCACC ACTGCCGACG
1001  GCACCGAAGC TTCACCGACT TATCTNNCNA TCGAAACCAC CGAAAAAGGA
1051  GCNNCAGGCA CATTTATCTC CAATGGTGGT CGGATCGAGA GCAAAGGCTT
1101  ATTGGTTATT GAGACGGGAG AAGATATCAN CTTGCGTAAC GGAGCCGTGG
1151  TGCAGAATAA CGGCAGTCGC CCAGCTACCA CGGTATTAAA TGCTGGTCAT
1201  AATTTGGTGA TTGAGAGTAA AACTAATGTG AACAATGCCA AAGGCTCGNC
```

```
1251  TAATCTGTCG GCCGGCGGTC GTACTACGAT CAATGATGCT ACTATTCAAG
1301  CGGGCAGTTC CGTGTACAGC TCCACCAAAG GCGATACTGA NTTGGGTGAA
1351  AATACCCGTA TTATTGCTGA AAACGTAACC GTATTATCTA ACGGTAGTAT
1401  TGGCAGTGCT GCTGTAATTG AGGCTAAAGA CACTGCACAC ATTGAATCGG
1451  GCAAACCGCT TTCTTTAGAA ACCTCGACCG TTGCCTCCAA CATCCGTTTG
1501  AACAACGGTA ACATTAAAGG CGGAAAGCAG CTTGCTTTAC TGGCAGACGA
1551  TAACATTACT GCCAAAACTA CCAATCTGAA TACTCCCGGC AATCTGTATG
1601  TTCATACAGG TAAAGATCTG AATTTGAATG TTGATAAAGA TTTGTCTGCC
1651  GCCAGCATCC ATTTGAAATC GGATAACGCT GCCCATATTA CCGGCACCAG
1701  TAAAACCCTC ACTGCCTCAA AAGACATGGG TGTGGAGGCA GGCTTGCTGA
1751  ATGTTACCAA TACCAATCTG CGTACCAACT CGGGTAATCT GCACATTCAG
1801  GCAGCCAAAG GCAATATTCA GCTTCGCAAT ACCAAGCTGA ACGCAGCCAA
1851  GGCTCTCGAA ACCACCGCAT TGCAGGGCAA TATCGTTTCA GACGGCCTTC
1901  ATGCTGTTTC TGCAGACGGT CATGTATCCT TATTGGCCAA CGGTAATGCC
1951  GACTTTACCG GTCACAATAC CCTGACAGCC AAGGCCGATG TCNATGCAGG
2001  ATCGGTTGGT AAAGGCCGTC TGAAAGCAGA CAATACCAAT ATCACTTCAT
2051  CTTCAGGAGA TATTACGTTG GTTGCCGNNN NCGGTATTCA GCTTGGTGAC
2101  GGAAAACAAC GCAATTCAAT CAACGGAAAA CACATCAGCA TCAAAAACAA
2151  CGGTGGTAAT GCCGACTTAA AAAACCTTAA CGTCCATGCC AAAAGCGGGG
2201  CATTGAACAT TCATTCCGAC CGGGCATTGA GCATAGAAAA TACNAAGCTG
2251  GAGTCTACCC ATAATACGCA TCTTAATGCA CAACACGAGC GGGTAACGCT
2301  CAACCAAGTA GATGCCTACG CACACCGTCA TCTAAGCATT ANCGGCAGCC
2351  AGATTTGGCA AAACGACAAA CTGCCTTCTG CCAACAAGCT GGTGGCTAAC
2401  GGTGTATTGG CANTCAATGC GCGCTATTCC CAAATTGCCG ACAACACCAC
2451  GCTGAGAGCG GGTGCAATCA ACCTTACTGC CGGTACCGCC CTAGTCAAGC
2501  GCGGCAACAT CAATTGGAGT ACCGTTTCGA CCAAGACTTT GGAAGATAAT
2551  GCCGAATTAA AACCATTGGC CGGACGGCTG AATATTGAAG CAGGTAGCGG
2601  CACATTAACC ATCGAACCTG CCAACCGCAT CAGTGCGCAT ACCGACCTGA
2651  GCATCAAAAC AGGCGGAAAA TTGCTGTTGT CTGCAAAAGG AGGAAATGCA
2701  GGTGCGCNTA GTGCTCAAGT TTCCTCATTG GAAGCAAAAG GCAATATCCG
2751  TCTGGTTACA GGAGNAACAG ATTTAAGAGG TTCTAAAATT ACAGCCGGTA
2801  AAAACTTGGT TGTCGCCACC ACCAAAGGCA AGTTGAATAT CGAAGCCGTA
2851  AACAACTCAT TCAGCAATTA TTTTCNTACA CAAAAAGNGN NNGNNCTCAA
2901  CCAAAAATCC AAAGAATTGG AACAGCAGAT TGCGCAGTTG AAAAAAAGCT
2951  CGCNTAAAAG CAAGCTGATT CCAACCCTGC AAGAAGAACG CGACCGTCTC
3001  GCTTTCTATA TTCAAGCCAT CAACAAGGAA GTTAAAGGTA AAAAACCCAA
3051  AGGCAAAGAA TACCTGCAAG CCAAGCTTTC TGCACAAAAT ATTGACTTGA
3101  TTTCCGCACA AGGCATCGAA ATCAGCGGTT CCGATATTAC CGCTTCCAAA
3151  AAACTGAACC TTCACGCCGC AGGCGTATTG CCAAAGGCAG CAGATTCAGA
3201  GGCGGCTGCT ATTCTGATTG ACGGCATAAC CGACCAATAT GAAATTGGCA
3251  AGCCCACCTA CAAGAGTCAC TACGACAAAG CTGCTCTGAA CAAGCCTTCA
3301  CGTTTGACCG GACGTACGGG GGTAAGTATT CATGCAGCTG CGGCACTCGA
3351  TGATGCACGT ATTATTATCG GTGCATCCGA AATCAAAGCT CCCTCAGGCA
3401  GCATAGACAT CAAAGCCCAT AGTGATATTG TACTGGAGGC TGGACAAAAC
3451  GATGCCTATA CCTTCTTANA AACCAAAGGT AAAAGCGGCA NAATNATCAG
3501  AAAAACNAAG TTTACCAGCA CCNGCGANCA CCTGATTATG CCAGCCCCNG
3551  TCGAGCTGAC CGCCAACGGT ATCACGCTTC AGGCAGGCGG CAACATCGAA
3601  GCTAATACCA CCCGCTTCAA TGCCCCTGCA GGTAAAGTTA CCCTGGTTGC
3651  GGGTGAANAG NTGCAACTGC TGGCAGAAGA AGGCATCCAC AAGCACGAGT
3701  TGGATGTCCA AAAAAGCCGC CGCTTTATCG GCATCAAGGT AGGTNAGAGC
3751  AATTACAGTA AAAACGAACT GAACGAAACC AAATTGCCTG TCCGCGTCGT
3801  CGCCCAAANT GCAGCCACCC GTTCAGGCTG GGATACCGTG CTCGAAGGTA
3851  CCGAATTCAA AACCACGCTG GCCGGTGCCG ACATTCAGGC AGGTGTANGC
3901  GAAAAAGCCC GTGTCGATGC GAAAATTATC CTCAAAGGCA TTGTGAACCG
3951  TATCCAGTCG GAAGAAAAAT TAGAAACCAA CTCAACCGTA TGGCAGAAAC
4001  AGGCCGGACG CGGCAGCACT ATCGAAACGC TAAAACTGCC CAGCTTCGAA
4051  AGCCCTACTC CGCCCAAATT GTCCGCACCC GGCGGNTATA TCGTCGACAT
4101  TCCGAAAGGC AATCTGAAAA CCGAAATCGA AAAGCTGTCC AAACAGCCCG
4151  AGTATGCCTA TCTGAAACAG CTCCAAGTAG CGAAAAACAT CAACTGGAAT
4201  CAGGTGCAGC TTGCTTACGA CAGATGGGAC TACAAACAGG AGGGCTTAAC
4251  CGAAGCAGGT GCGGCGATTA TCGCACTGGC CGTTACCGTG GTCACCTCAG
4301  GCGCAGGAAC CGGAGCCGTA TTGGGATTAA ACGGTGCGNC CGCCGCCGCA
4351  ACCGATGCAG CATTCGCCTC TTTGGCCAGC CAGGCTTCCG TATCGTTCAT
4401  CAACAACAAA GGCGATGTCG GCAAAACCCT GAAAGAGCTG GGCAGAAGCA
4451  GCACGGTGAA AAATCTGGTG GTTGCCGCCG CTACCGCAGG CGTAGCCGAC
4501  AAAATCGGCG CTTCGGCACT GANCAATGTC AGCGATAAGC AGTGGATCAA
4551  CAACCTGACC GTCAACCTAG CCAATGNCGG GCAGTGCCGC ACTGAttaa
```

This encodes a protein having amino acid sequence <SEQ ID 62>:

```
1  MNKGLHRIIF SKKHSTMVAV AETANSQGKG KQAGSSVSVS LKTSGDLCGK
```

```
  51  LKTTLKTLVC SLVSLSMXXX XXXQITTDKS APKNXQVVIL KTNTGAPLVN
 101  IQTPNGRGLS HNRYTQFDVD NKGAVLNNDR NNNPFLVKGS AQLILNEVRG
 151  TASKLNGIVT VGGQKADVII ANPNGITVNG GGFKNVGRGI LTIGAPQIGK
 201  DGALTGFDVR QGTLTVGAAG WNDKGGADYT GVLARAVALQ GKLQGKNLAV
 251  STGPQKVDYA SGEISAGTAA GTKPTIALDT AALGGMYADS ITLIAXEKGV
 301  GVKNAGTLEA AKQLIVTSSG RIENSGRIAT TADGTEASPT YLXIETTEKG
 351  AXGTFISNGG RIESKGLLVI ETGEDIXLRN GAVVQNNGSR PATTVLNAGH
 401  NLVIESKTNV NNAKGSXNLS AGGRTTINDA TIQAGSSVYS STKGDTXLGE
 451  NTRIIAENVT VLSNGSIGSA AVIEAKDTAH IESGKPLSLE TSTVASNIRL
 501  NNGNIKGGKQ LALLADDNIT AKTTNLNTPG NLYVHTGKDL NLNVDKDLSA
 551  ASIHLKSDNA AHITGTSKTL TASKDMGVEA GLLNVTNTNL RTNSGNLHIQ
 601  AAKGNIQLRN TKLNAAKALE TTALQGNIVS DGLHAVSADG HVSLLANGNA
 651  DFTGHNTLTA KADVXAGSVG KGRLKADNTN ITSSSGDITL VAXXGIQLGD
 701  GKQRNSINGK HISIKNNGGN ADLKNLNVHA KSGALNIHSD RALSIENTKL
 751  ESTHNTHLNA QHERVTLNQV DAYAHRHLSI XGSQIWQNDK LPSANKLVAN
 801  GVLAXNARYS QIADNTTLRA GAINLTAGTA LVKRGNINWS TVSTKTLEDN
 851  AELKPLAGRL NIEAGSGTLT IEPANRISAH TDLSIKTGGK LLLSAKGGNA
 901  GAXSAQVSSL EAKGNIRLVT GXTDLRGSKI TAGKNLVVAT TKGKLNIEAV
 951  NNSFSNYFXT QKXXXLNQKS KELEQQIAQL KKSSXKSKLI PTLQEERDRL
1001  AFYIQAINKE VKGKKPKGKE YLQAKLSAQN IDLISAQGIE ISGSDITASK
1051  KLNLHAAGVL PKAADSEAAA ILIDGITDQY EIGKPTYKSH YDKAALNKPS
1101  RLTGRTGVSI HAAAALDDAR IIIGASEIKA PSGSIDIKAH SDIVLEAGQN
1151  DAYTFLXTKG KSGXXIRKTK FTSTXXHLIM PAPVELTANG ITLQAGGNIE
1201  ANTTRFNAPA GKVTLVAGEX XQLLAEEGIH KHELDVQKSR RFIGIKVGXS
1251  NYSKNELNET KLPVRVVAQX AATRSGWDTV LEGTEFKTTL AGADIQAGVX
1301  EKARVDAKII LKGIVNRIQS EEKLETNSTV WQKQAGRGST IETLKLPSFE
1351  SPTPPKLSAP GGYIVDIPKG NLKTEIEKLS KQPEYAYLKQ LQVAKNINWN
1401  QVQLAYDRWD YKQEGLTEAG AAIIALAVTV VTSGAGTGAV LGLNGAXAAA
1451  TDAAFASLAS QASVSFINNK GDVGKTLKEL GRSSTVKNLV VAAATAGVAD
1501  KIGASALXNV SDKQWINNLT VNLANXGQCR TD*
```

ORF114-1 and ORF114a show 89.8% identity in 1564 aa overlap

```
orf114a.pep   MNKGLHRIIFSKKHSTMVAVAETANSQGKGKQAGSSVSVSLKTSGDLCGKLKTTLKTLVC
              ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
orf114-1      MNKGLHRIIFSKKHSTMVAVAETANSQGKGKQAGSSVSVSLKTSGDLCGKLKTTLKTLVC

orf114a.pep   SLVSLSMXXXXXXQITTDKSAPKNXQVVILKTNTGAPLVNIQTPNGRGLSHNRYTQFDVD
              |||||||      |||||||||||| ||||||||||||||||||||||||||||||||||
orf114-1      SLVSLSMVLPAHAQITTDKSAPKNQQVVILKTNTGAPLVNIQTPNGRGLSHNRYTQFDVD

orf114a.pep   NKGAVLNNDRNNNPFLVKGSAQLILNEVRGTASKLNGIVTVGGQKADVIIANPNGITVNG
              ||||||||||||||:|||||||||||||||||||||||||||||||||||||||||||||
orf114-1      NKGAVLNNDRNNNPFVVKGSAQLILNEVRGTASKLNGIVTVGGQKADVIIANPNGITVNG

orf114a.pep   GGFKNVGRGILTIGAPQIGKDGALTGFDVRQGTLTVGAAGWNDKGGADYTGVLARAVALQ
              |||||||||||| ||||||||||||||||||||||||||||||||||||||||||||||||
orf114-1      GGFKNVGRGILTTGAPQIGKDGALTGFDVRQGTLTVGAAGWNDKGGADYTGVLARAVALQ

orf114a.pep   GKLQGKNLAVSTGPQKVDYASGEISAGTAAGTKPTIALDTAALGGMYADSITLIAXEKGV
              ||||||||||||||||||||||||||||||||||||||||||||||||||||| ||||
orf114-1      GKLQGKNLAVSTGPQKVDYASGEISAGTAAGTKPTIALDTAALGGMYADSITLIANEKGV

orf114a.pep   GVKNAGTLEAAKQLIVTSSGRIENSGRIATTADGTEASPTYLXIETTEKGAXGTFISNGG
              |||||||||||||||||||||||||||||||||||||||||||| |||||||| ||||||||
orf114-1      GVKNAGTLEAAKQLIVTSSGRIENSGRIATTADGTEASPTYLSIETTEKGAAGTFISNGG

orf114a.pep   RIESKGLLVIETGEDIXLRNGAVVQNNGSRPATTVLNAGHNLVIESKTNVNNAKGSXNLS
              |||||||||||||||| |||||||||||||||||||||||||||||||||||  :||
orf114-1      RIESKGLLVIETGEDISLRNGAVVQNNGSRPATTVLNAGHNLVIESKTNVNNAKGPATLS

orf114a.pep   AGGRTTINDATIQAGSSVYSSTKGDTXLGENTRIIAENVTVLSNGSIGSAAVIEAKDTAH
              | |||:|::|:||:|::|||||:||:: ||:|||| : :|||||||:|:|:|||:|||||
orf114-1      ADGRTVIKEASIQTGTTVYSSSKGNAELGNNTRITGADVTVLSNGTISSSAVIDAKDTAH

orf114a.pep   IESGKPLSLETSTVASNIRLNNGNIKGGKQLALLADDNITAKTTNLNTPGNLYVHTGKDL
              ||:|||||||:|||:|:|||||:|:|||||||||||||||||||||||||||||||||||
orf114-1      IEAGKPLSLEASTVTSDIRLNGGSIKGGKQLALLADDNITAKTTNLNTPGNLYVHTGKDL

orf114a.pep   NLNVDKDLSAASIHLKSDNAAHITGTSKTLTASKDMGVEAGLLNVTNTNLRTNSGNLHIQ
              |||||||||||||||||||||||||||||||||||||||||| ||||||||||||||||||
```

```
orf114-1        NLNVDKDLSAASIHLKSDNAAHITGTSKTLTASKDMGVEAGSLNVTNTNLRTNSGNLHIQ

orf114a.pep     AAKGNIQLRNTKLNAAKALETTALQGNIVSDGLHAVSADGHVSLLANGNADFTGHNTLTA
                ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
orf114-1        AAKGNIQLRNTKLNAAKALETTALQGNIVSDGLHAVSADGHVSLLANGNADFTGHNTLTA

orf114a.pep     KADVXAGSVGKGRLKADNTNITSSSGDITLVAXXGIQLGDGKQRNSINGKHISIKNNGGN
                ||||  ||||||||||||||||||||||||||  ||||||||||||||||||||||||||
orf114-1        KADVNAGSVGKGRLKADNTNITSSSGDITLVAGNGIQLGDGKQRNSINGKHISIKNNGGN

orf114a.pep     ADLKNLNVHAKSGALNIHSDRALSIENTKLESTHNTHLNAQHERVTLNQVDAYAHRHLSI
                ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
orf114-1        ADLKNLNVHAKSGALNIHSDRALSIENTKLESTHNTHLNAQHERVTLNQVDAYAHRHLSI

orf114a.pep     XGSQIWQNDKLPSANKLVANGVLAXNARYSQIADNTTLRAGAINLTAGTALVKRGNINWS
                :|||||||||||||||||||||||  ||||||||||||||||||||||||||||||||||
orf114-1        TGSQIWQNDKLPSANKLVANGVLALNARYSQIADNTTLRAGAINLTAGTALVKRGNINWS

orf114a.pep     TVSTKTLEDNAELKPLAGRLNIEAGSGTLTIEPANRISAHTDLSIKTGGKLLLSAKGGNA
                ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
orf114-1        TVSTKTLEDNAELKPLAGRLNIEAGSGTLTIEPANRISAHTDLSIKTGGKLLLSAKGGNA

orf114a.pep     GAXSAQVSSLEAKGNIRLVTGXTDLRGSKITAGKNLVVATTKGKLNIEAVNNSFSNYFXT
                ||  |||||||||||||||||  ||||||||||||||||||||||||||||||||||  |
orf114-1        GAPSAQVSSLEAKGNIRLVTGETDLRGSKITAGKNLVVATTKGKLNIEAVNNSFSNYFPT

orf114a.pep     QKXXXLNQKSKELEQQIAQLKKSSXKSKLIPTLQEERDRLAFYIQAINKEVKGKKPKGKE
                ||   ||||||||||||||||  ||||||||||||||||||||||||||||||||||||
orf114-1        QKAAELNQKSKELEQQIAQLKKSSPKSKLIPTLQEERDRLAFYIQAINKEVKGKKPKGKE

orf114a.pep     YLQAKLSAQNIDLISAQGIEISGSDITASKKLNLHAAGVLPKAADSEAAAILIDGITDQY
                ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
orf114-1        YLQAKLSAQNIDLISAQGIEISGSDITASKKLNLHAAGVLPKAADSEAAAILIDGITDQY

orf114a.pep     EIGKPTYKSHYDKAALNKPSRLTGRTGVSIHAAAALDDARIIIGASEIKAPSGSIDIKAH
                ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
orf114-1        EIGKPTYKSHYDKAALNKPSRLTGRTGVSIHAAAALDDARIIIGASEIKAPSGSIDIKAH

orf114a.pep     SDIVLEAGQNDAYTFLXTKGKSGXXIRKTKFTSTXXHLIMPAPVELTANGITLQAGGNIE
                ||||||||||||||||  ||||||  |||||||||  ||||||||||||||||||||||||
orf114-1        SDIVLEAGQNDAYTFLKTKGKSGKIIRKTKFTSTRDHLIMPAPVELTANGITLQAGGNIE

orf114a.pep     ANTTRFNAPAGKVTLVAGEXXQLLAEEGIHKHELDVQKSRRFIGIKVGXSNYSKNELNET
                ||||||||||||||||||  ||||||||||||||||||||||||  ||||||||||||
orf114-1        ANTTRFNAPAGKVTLVAGEELQLLAEEGIHKHELDVQKSRRFIGIKVGKSNYSKNELNET

orf114a.pep     KLPVRVVAQXAATRSGWDTVLEGTEFKTTLAGADIQAGVXEKARVDAKIILKGIVNRIQS
                |||||||||:||||||||||||||||||||||||||||||  ||||:||||||||||||||
orf114-1        KLPVRVVAQTAATRSGWDTVLEGTEFKTTLAGADIQAGVGEKARADAKIILKGIVNRIQS

orf114a.pep     EEKLETNSTVWQKQAGRGSTIETLKLPSFESPTPPKLSAPGGYIVDIPKGNLKTEIEKLS
                |||||||||||||||||||||||||||||||||||||:|||||||||||||||||||||:
orf114-1        EEKLETNSTVWQKQAGRGSTIETLKLPSFESPTPPKLTAPGGYIVDIPKGNLKTEIEKLA

orf114a.pep     KQPEYAYLKQLQVAKNINWNQVQLAYDRWDYKQEGLTEAGAAIIALAVTVVTSGAGTGAV
                ||||||||||||||||:||||||||||:|||||||||:|||||::: ||::| | |: |:
orf114-1        KQPEYAYLKQLQVAKNVNWNQVQLAYDKWDYKQEGLTRAGAAIVTIIVTALTYGYGATAA

orf114a.pep     LGLNGA--------------XAAATD---------AAFASLASQASVSFINNKGDVGKTL 1477
                  |: ::               :||||          ||:||| |||:||:|||||||:|
orf114-1        GGVAASGSSTAAAAGTAATTTAAATTVSTATAMQTAALASLYSQAAVSIINNKGDVGKAL 1500

orf114a.pep     KELGRSSTVKNLVVAAATAGVADKIGA----------SALXNVSDKQWINNL----TVNL 1523
                |:|| |:|||::|::| |||: :::||          : | : : :| | ||    ::||
orf114-1        KDLGTSDTVKQIVTSALTAGALNQMGADIAQLNSKVRTELFSSTGNQTIANLGGRLATNL 1560

orf114a.pep     ANXGQCRTDX
                :| |
orf114-1        SNAGISAGINTAVN...
```

Homology with pspA putative secreted protein of *N.meningitidis* (accession number AF030941)

ORF114 and pspA protein show 36% aa identity in 302aa overlap:

```
Orf114: 1   AVAETANSQGKGKQAGSSVSVSL----KTSGDXXXXXXXXXXXXXXXXXXXXXPAHAQ 56
            AVAE  +  GK  Q   + SV +      S                       PA A
pspA:   19  AVAENVHRDGKSMQDSEAASVRVTGAASVSSARAAFGFRMAAFSVMLALGVAAFSPAPAS 78

Orf114: 57  -ITTDKSAPKNQQVVILKTNTGAPLVNIQTPNGRGLSHNRXYAFDVDNKGAVLNNDRNN- 114
             I  DKSAPKNQQ VIL+T  G P VNIQTP+ +G+S NR    FDVD KG +LNN R+N
pspA:   79  GIIADKSAPKNQQAVILQTANGLPQVNIQTPSSQGVSVNRFKQFDVDEKGVILNNSRSNT 138

Orf114: 115 ----------NPFVVKGSAQLILNEV-RGTASKLNGIVTVGGQKADVIIANPNGITVNGG 163
                     NP + +G A++I+N++      S LNG + VGG++A+V++ANP+GI VNGG
pspA:   139 QTQLGGWIQGNPHLARGEARVIVNQIDSSNPSLLNGYIEVGGKRAEVVVANPSGIRVNGG 198

Orf114: 164 GFKNVGRGILTTGAPQIGKDGALTGFDVVKAHWTVXAAGWNDKGGAXYTGVLARAVALQG 223
            G  N     LT+G P +  +G LTGFDV      +   G  D   A YT +L+RA  +
pspA:   199 GLINAASVTLTSGVPVL-NNGNLTGFDVSSGKVVIGGKGL-DTSDADYTRILSRAAEINA 256

Orf114: 224 KXXGKXLAVSTGPQKVDYASGEISAGTAAGTK----PTIALDTAALGGMYADSITLIANE 279
               GK + V +G  K+D+         +A  +     PT+A+DTA LGGMYAD ITLI+ +
pspA:   257 GVWGKDVKVVSGKNKLDFDGSLAKTASAPSSSDSVTPTVAIDTATLGGMYADKITLISTD 316

Orf114: 280 KG 281
             G
pspA:   317 NG 318
```

ORF114a is also homologous to pspA:

```
gi|2623258 (AF030941) putative secreted protein [Neisseria meningitidis] Length
= 2273
 Score =  261 bits (659), Expect = 3e-68
 Identities = 203/663 (30%), Positives = 314/663 (46%), Gaps = 76/663 (11%)

Query: 1   MNKGLHRIIFSKKHSTMVAVAETANSQGKGKQAGSSVSVSLK-----TSGDXXXXXXXXX 55
            MNK  +++IF+KK S M+AVAE  +  GK  Q   + SV +      +S
Sbjct: 1   MNKRCYKVIFNKKRSCMMAVAENVHRDGKSMQDSEAASVRVTGAASVSSARAAFGFRMAA 60

Query: 56  XXXXXXXXXXXXXXXXXQITTDKSAPKNXQVVILKTNTGAPLVNIQTPNGRGLSHNRYT 115
                          I  DKSAPKN Q VIL+T  G P VNIQTP+ +G+S NR+
Sbjct: 61  FSVMLALGVAAFSPAPASGIIADKSAPKNQQAVILQTANGLPQVNIQTPSSQGVSVNRFK 120

Query: 116 QFDVDNKGAVLNNDRNN-----------NPFLVKGSAQLILNEV-RGTASKLNGIVTVGG 163
            QFDVD KG +LNN R+N            NP L +G A++I+N++     S LNG + VGG
Sbjct: 121 QFDVDEKGVILNNSRSNTQTQLGGWIQGNPHLARGEARVIVNQIDSSNPSLLNGYIEVGG 180

Query: 164 QKADVIIANPNGITVNGGGFKNVGRGILTIGAPQIGKDGALTGFDVRQGTLTVGAAGWND 223
            ++A+V++ANP+GI VNGGG  N      LT G P +  +G LTGFDV  G + +G  G  D
Sbjct: 181 KRAEVVVANPSGIRVNGGGLINAASVTLTSGVPVL-NNGNLTGFDVSSGKVVIGGKGL-D 238

Query: 224 KGGADYTGVLARAVALQGKLQGKNLAVSTGPQKVDYASGEISAGTAAGTK----PTIALD 279
               ADYT +L+RA  +   + GK++ V +G  K+D+         +A  +     PT+A+D
Sbjct: 239 TSDADYTRILSRAAEINAGVWGKDVKVVSGKNKLDFDGSLAKTASAPSSSDSVTPTVAID 298

Query: 280 TAALGGMYADSITLIAXEKGVGVKNAGTLEAAK-QLIVTSSGRIENSGRIATTADGTEAS 338
            TA LGGMYAD ITLI+ + G  ++N G + AA   + +++ G++ NSG I        +A+
Sbjct: 299 TATLGGMYADKITLISTDNGAVIRNKGRIFAATGGVTLSADGKLSNSGSI-------DAA 351

Query: 339 PTYLXIETTEKGAXGTFISNGGRIESKGLLVIETGEDIXLRNGAVVQNNGSRPATTVLNA 398
              +  +T +         +  G I S     V++  + I  + G +     GS     + +
Sbjct: 352 EITISAQTVD--------NRQGFIRSGKGSVLKVSDGINNQAGLI----GSAGLLDIRDT 399

Query: 399 GHNLVIESKTNVNNAKGS----XNLSAGGRTTINDATIQAGSSVYSSTKGDTXLGENTRI 454
            G     +S  ++NN  G+      ++S   ++  ND  + A   V S +  D   G+
Sbjct: 400 G-----KSSLHINNTDGTIIAGKDVSLQAKSLDNDGILTAARDV-SVSLHDDFAGKRDIE 453

Query: 455 IAENVTVLSNGSIGSAAVIEAKDTAHIESGKPLSLETSTVASNIRLNNGNIKGGKQLALL 514
               +T  + G + +  +I+A DT  + + +  +  +  + S  R       G     L+
```

```
Sbjct: 454 AGRTLTFSTQGRLKNTRIIQAGDTVSLTAAQIDNTVSGKIQSGNRTGLNGKNGITNRGLI 513

Query: 515 ADDNIT-----AKTTNLNTPGNLYVHTGKDLNLNVDKDLSAASIHLKSDNAAHITGTSKT 569
             + IT     AK+ N  T G +Y   G  + +  D  L+          AA
Sbjct: 514 NSNGITLLQTEAKSDNAGT-GRIY---GSRVAVEADTLLNREETVNGETKAA-------V 562

Query: 570 LTASKDMGVEAGXXXXXXXXXXXSGNLHIQAA---KGNIQLRNTKL-NAAKALETTALQ 625
           + A + + + A              SG+LHI +A      +Q  NT L N + A+E++
Sbjct: 563 IAARERLDIGAREIENREAALLSSSGDLHIGSALNGSRQVQGANTSLHNRSAAIESS--- 619

Query: 626 GNI 628
           GNI
Sbjct: 620 GNI 622


 Score = 37.5 bits (85), Expect = 0.53
 Identities = 87/432 (20%), Positives = 159/432 (36%), Gaps = 62/432 (14%)

Query: 239  LQGKLQGKNLAVSTGPQKVDYASGEISAGTAAGTKPTIALDTAALGGMYADSITLIAXEK 298
             LQG LQGKN+  + G    +  +G I A  A   K        A   + + S T     +
Sbjct: 1023 LQGDLQGKNIFAAAGSDITN--TGSIGAENALLLK--------ASNNIESRSETRSNQNE 1072

Query: 299  GVGVKNAGTLEAAKQLIVTSSGRI--ENSGRIATTADGTEASPTYLXIETTEKGAXG-TF 355
               V+N G + A   L    +G +  +    I  TA            E T +   G T
Sbjct: 1073 QGSVRNIGRV-AGIYLTGRQNGSVLLDAGNNIVLTAS----------ELTNQSEDGQTV 1120

Query: 356  ISNGGRIESKGLLVIETGEDIXLRNGAVVQNNGSRPATTVLNAGHNLVIESK-------T 408
             ++ GG I S    +      I   +  V++   +   +T+   G NL + +K
Sbjct: 1121 LNAGGDIRSDTTGISRNQNTIFDSDNYVIRKEQNEVGSTIRTRG-NLSLNAKGDIRIRAA 1179

Query: 409  NVNNAKGSXNLSAGGRTTINDATIQAGSS--------VYSSTKGDTXLGENTRIIAENVT 460
              V + +G   L+AG       D  ++AG +          Y+   G     + TR +
Sbjct: 1180 EVGSEQGRLKLAAG-----RDIKVEAGKAHTETEDALKYTGRSGGGIKQKMTRHLKNQNG 1234

Query: 461  VLSNGSIGSAAVIEAKDTAHIESGKPLSLETSTVASNIRLNNGNIKGGKQLALLADDNIT 520
               +G++    +I         +G  +  +  T+ S    NN  +K  +  +  A+ N
Sbjct: 1235 QAVSGTLDGKEIILVSGRDITVTGSNIIADNHTILS--AKNNIVLKAAETRSRSAEMNKK 1292

Query: 521  AKTTNLNTPG-NLYVHTGKDLNLNVDKDLSAASIHLKSDN-------AAHITGTSKTLTA 572
              K+  + + G       + KD   N  + +S     + S N          H T T  T+++
Sbjct: 1293 EKSGLMGSGGIGFTAGSKKDTQTNRSETVSHTESVVGSLNGNTLISAGKHYTQTGSTISS 1352

Query: 573  SK-DMGVEAGXXXXXXXXXXXSGNLHIQAAKG-----NIQLRNTKLNAAKALETTALQG 626
              + D+G+ +G                +  +  KG      ++ + NT + A  A++     G
Sbjct: 1353 PQGDVGISSGKISIDAAQNRYSQESKQVYEQKGVTVAISVPVVNTVMGAVDAVKAVQTVG 1412

Query: 627  NIVSDGLHAVSA 638
                +  ++A++A
Sbjct: 1413 KSKNSRVNAMAA 1424
```

Amino acids 1-1423 of ORF114-1 were cloned in the pGex vector and expressed in *E.coli*, as described above. GST-fusion expression was visible using SDS-PAGE, and Figure 5 shows plots of hydrophilicity, antigenic index, and AMPHI regions for ORF114-1.

Based on these results, including the homology with the putative secreted protein of *N.meningitidis* and on the presence of a transmembrane domain, it is predicted that this protein from *N.meningitidis*, and its epitopes, could be useful antigens for vaccines or diagnostics.

Example 14

The following partial DNA sequence was identified in *N.meningitidis* <SEQ ID 63>

```
   1  ..CGCTTCATTC ATGATGAAGC AGTCGGCAGC AACATCGGCG GCGGCAAAAT
  51    GATTGTTGCA GCCGGGCAGG ATATCAATGT ACGCGGCAnA AGCCTTATTT
 101    CTGATAAGGG CATTGTTTTA AAAGCAGGAC ACGACATCGA TATTTCTACT
 151    GCCCATAATC GCTATACCGG CAATGAATAC CACGAGAGCA wAAAwTCAGG
 201    CGTCATGGGT ACTGGCGGAT TGGGCTTTAC TATCGGTAAC CGGAAAACTA
 251    CCGATGACAC TGATCGTACC AATATTGTsC ATACAGGCAG CATTATAGGC
 301    AGCCTGAaTG GAGACACCGT TACAGTTGCA GGAAACCGCT ACCGACAAAC
 351    CGGCAGTACC GTCTCCAGCC CCGAGGGGCG CAATACCGTC ACAGCCAAAw
 401    GCATAGATGT AGAGTTCGCA AACAACCGGT ATGCCACTGA CTACGcCCAT
 451    ACCCAgGGAA CAAAAAGGCC TTACCGTCGC CCTCAATGTC CCGGTTGTCC
 501    AAGCTGCACA AAACTTCATA CAAGCAGCCC AAAATGTGGG CAAAAGTAAA
 551    AATAAACGCG TTAATGCCAT GGCTGCAGCC AATGCTGCAT GGCAGAGTTA
 601    TCAAGCAACC CAACAAATGC AACAATTTGC TCCAAGCAGC AGTGCGGGAC
 651    AAGGTCAAAA CTACAATCAA AGCCCCAGTA TCAGTGTGTC CATTAC.TAC
 701    GGCGAACAGA AAAGTCGTAA CGAGCAAAAA AGACATTACA CCGAAgCGGC
 751    AgCAAGTCAA ATTATCGGCA AAGGGCAAAC CACACTTGCG GCAACAGGAA
 801    GTGGGGAGCA GTCCAATATC AATATTACAG GTTCCGATGT CATCGGCCAT
 851    GCAGGTACTC C.CTCATTGC CGACAACCAT ATCAGACTCC AATCTGCCAA
 901    ACAGGACGGC AGCGAGCAAA GCAAAAACAA AAGCAGTGGT TGGAATGCAG
 951    GCGTACGTnn CAAAATAGGC AACGGCATCA GGTTTGGAAT TACCGCCGGA
1001    GGAAATATCG GTAAAGGTAA AGAGCAAGGG GGAAGTACTA CCCACCGCCA
1051    CACCCATGTC GGCAGCACAA CCGGCAAAAC TACCATCCGA AGCGGCGGGG
1101    GATACCACCC TCAAAGGTGT GCAGCTCATC GGCAAAGGCA TACAGGCAGA
1151    TACGCGCAAC CTGCATATAG AAAGTGTTCA AGATACTGAA ACCTATCAGA
1201    GCAAACAGCA AAACGGCAAT GTCCAAGTTt ACTGTCGGTT ACGGATTCAG
1251    TGCAAGCGGC AGTTACCGCC AAAGCAAAGT CAAAGCAGAC CATGCCTCCG
1301    TAACCGGGCA AAgCGGTATT TATGCCGGAG AAGACGGCTA TCAAATyAAA
1351    GTyAGAGACA ACACAGACCT yAAGGGCGGT ATCATCACGT CTAGCCAAAG
1401    CGCAGAAGAT AAGGGCAAAA ACCTTTTTCA GACGGCCACC CTTACTGCCA
1451    GCGACATTCA AAACCACAGC CGCTACGAAG GCAGAAGCTT CGGCATAGGC
1501    GGCAGTTTCG ACCTGAACGG CGGCTGGGAC GGCACGGTTA CCGACAAACA
1551    AGGCAGGCCT ACCGACAGGA TAAGCCCGGC AGCCGGCTAC GGCAGCGACG
1601    GAGACAGCAA AAACAGCACC ACCCGCAGCG GCGTCAACAC CCACAACATA
1651    CACATCACCG ACGAAGCGGG ACAACTTGCC CGAACAGGCA GGACTGCAAA
1701    AGAAACCGAA GCGCGTATCT ACACCGGCAT CGACACCGAA ACTGCGGATC
1751    AACACTCAGG CCATCTGAAA AACAGCTTCG AC...
```

This corresponds to the amino acid sequence <SEQ ID 64; ORF116>:

```
   1  ..RFIHDEAVGS NIGGGKMIVA AGQDINVRGX SLISDKGIVL KAGHDIDIST
  51    AHNRYTGNEY HESXXSGVMG TGGLGFTIGN RKTTDDTDRT NIVHTGSIIG
 101    SLNGDTVTVA GNRYRQTGST VSSPEGRNTV TAKXIDVEFA NNRYATDYAH
 151    TQEQKGLTVA LNVPVVQAAQ NFIQAAQNVG KSKNKRVNAM AAANAAWQSY
 201    QATQQMQQFA PSSSAGQGQN YNQSPSISVS IXYGEQKSRN EQKRHYTEAA
 251    ASQIIGKGQT TLAATGSGEQ SNINITGSDV IGHAGTXLIA DNHIRLQSAK
 301    QDGSEQSKNK SSGWNAGVRX KIGNGIRFGI TAGGNIGKGK EQGGSTTHRH
 351    THVGSTTGKT TIRSGGDTTL KGVQLIGKGI QADTRNLHIE SVQDTETYQS
 401    KQQNGNVQVT VGYGFSASGS YRQSKVKADH ASVTGQSGIY AGEDGYQIKV
 451    RDNTDLKGGI ITSSQSAEDK GKNLFQTATL TASDIQNHSR YEGRSFGIGG
 501    SFDLNGGWDG TVTDKQGRPT DRISPAAGYG SDGDSKNSTT RSGVNTHNIH
 551    ITDEAGQLAR TGRTAKETEA RIYTGIDTET ADQHSGHLKN SFD...
```

Computer analysis of this amino acid sequence gave the following results:

Homology with pspA putative secreted protein of *N.meningitidis* (accession number AF030941)

ORF116 and pspA protein show 38% aa identity in 502aa overlap:

```
Orf116:  6    EAVGSNIGGGKMIVAAGQDINVRGXSLISDKGIVLKAGHDIDISTAHNRYTGNEYHESXX 65
              +AV   + G ++I+ +G+DI V G ++I+D   +L A ++I +  A  R    E ++
PspA:  1  235 QAVSGTLDGKEIILVSGRDITVTGSNIIADNHTILSAKNNIVLKAAETRSRSAEMNKKEK 1294

Orf116: 66    XXXXXXXXXXXXXXNRKXXXXXXRTNIVHTGSIIGSLNGDTVTVAGNRYRQTGSTVSSPE 125
                            ++K          + HT S++GSLNG+T+  AG  Y QTGST+SSP+
PspA:   1295 SGLMGSGGIGFTAGSKKDTQTNRSETVSHTESVVGSLNGNTLISAGKHYTQTGSTISSPQ 1354
```

```
Orf116: 126  GRNTVTAKXIDVEFANNRYATDYAHTQEQKGLTVALNVPXXXX---XXXXXXXXXXXGKS 182
             G   +++  I ++ A NRY+ +      EQKG+TVA++VP                GKS
PspA:   1355 GDVGISSGKISIDAAQNRYSQESKQVYEQKGVTVAISVPVVNTVMGAVDAVKAVQTVGKS 1414

Orf116. 183  KNKRVXXXXXXXXXWQSYQATQQMQQFA--PSSSAGQGQNYNQSPSISVSIXYGEQKSRN 240
             KN RV          +   +   +   A  P  +AGQG        ISVS+ YGEQK+ +
PspA:   1415 KNSRVNAMAAANALNKGVDSGVALYNAARNPKKAAGQG--------ISVSVTYGEQKNTS 1466

Orf116: 241  EQKRHYTEAAASQIIGKGQTTLAATGSGEQSNINITGSDVIGHAGTXLIADNHIRLQSAK 300
             E +   T+    +I G G+ +L A+G+G+ S I ITGSDV G  GT L A+N +++++A+
PspA:   1467 ESRIKGTQVQEGKITGGGKVSLTASGAGKDSRITITGSDVYGGKGTRLKAENAVQIEAAR 1526

Orf116: 301  QDGSEQSKNKSSGWNAGVRXKIGNGIRFGITAXXXXXXXXXXXXXSTTHRHTHVGSTTGKT 360
             Q   E+S+NKS+G+NAGV   I  GI FG TA             T +R++H+GS   +T
PspA:   1527 QTHQERSENKSAGFNAGVAIAINKGISFGFTAGANYGKGYGNGDETAYRNSHIGSKDSQT 1586

Orf116: 361  TIRSGGDTTLKGVQLIGKGIQADTRNLHIESVQDTETYQSKQQNGNVQVTVGYGFSASGS 420
              I SGGDT +KG QL GKG+      +LHIES+QDT  ++ KQ+N + QVTVGYGFS  GS
PspA:   1587 AIESGGDTVIKGGQLKGKGVGVTAESLHIESLQDTAVFKGKQENVSAQVTVGYGFSVGGS 1646

Orf116: 421  YRQSKVKADHASVTGQSGIYAGEDGYQIKVRDNTDLKGGIITSSQSAEDKGKNLFQTATL 480
             Y +SK  +D+ASV  QSGI+AG DGY+I+V   T L G  + S     DK KNL +T+ +
PspA:   1647 YNRSKSSSDYASVNEQSGIFAGGDGYRIRVNGKTGLVGAAVVSD---ADKSKNLLKTSEI 1703

Orf116: 481  TASDIQNHSRYEGRSFGIGGSF 502
                DIQNH+     + G+ G F
PspA:   1704 WHKDIQNHASAAASALGLSGGF 1725
```

Based on homology with pspA, it is predicted that this protein from *N.meningitidis*, and its epitopes, could be useful antigens for vaccines or diagnostics.

Example 15

The following partial DNA sequence was identified in *N.meningitidis* <SEQ ID 65>

```
   1  ..ACGACCGGCA GCCTCGGCGG CATACTGGCC GGCGGCGGCA CTTCCCTTGC
  51    CGCACCGTAT TTGGACAAAG CGGCGGAAAA CCTCGGTCCG GCGGGCAAAG
 101    CGGCGGTCAA CGCACTGGGC GGTGCGGCCA TCGGCTATGC AACTGGTGGT
 151    AGTGGTGGTG CTGTGGTGGG TGCGAATGTA GATTGGAACA ATAGGCAGCT
 201    GCATCCGAAA GAAATGGCGT TGGCCGACAA ATATGCCGAA GCCCTCAAGC
 251    GCGAAGTTGA AAAACGCGAA GGCAGAAAAA TCAGCAGCCA AGAAGCGGCA
 301    ATGAGAATCC GCAGGCAGAT ATGCGTTGGG TGGACAAAGG TTCCCAAGAC
 351    GGCTATACCG ACCAAAGCGT CATATCCCTT ATCGGAATGA
```

This corresponds to the amino acid sequence <SEQ ID 66; ORF118>:

```
   1  ..TTGSLGGILA GGGTSLAAPY LDKAAENLGP AGKAAVNALG GAAIGYATGG
  51    SGGAVVGANV DWNNRQLHPK EMALADKYAE ALKREVEKRE GRKISSQEAA
 101    MRIRRQICVG WTKVPKTAIP TKASYPLSE*
```

Computer analysis of this amino acid sequence reveals two putative transmembrane domains.

Based on this analysis, it is predicted that this protein from *N.meningitidis*, and its epitopes, could be useful antigens for vaccines or diagnostics.

Example 16

The following partial DNA sequence was identified in *N.meningitidis* <SEQ ID 67>

```
   1  ..CAATGCCGTC TGAAAAGCTC ACAATTTTAC AGACGGCATT TGTTATGCAA
```

```
  51   GTACATATAC AGATTCCCTA TATACTGCCC AGrkGCGTGC GTgGCTGAAG
 101   ACACCCCCTA CGCTTGCTAT TTGrAACAGC TCCAAGTCAC CAAAGACGTC
 151   AACTGGAACC AGGTACwACT GGCGTACGAC AAATGGGACT ATAAACAGGA
 201   AGGCTTAACC GGAGCCGGAG CAGCGATTAT TGCGCTGGCT GTTACCGTGG
 251   TTACTGCGGG CGCGGGAgCC GGAGCCGCAC TGGGcTTAAA CGGCGCGGCc
 301   GCAGCGGCAA CCGATGCCGC ATTCGCCTCG CTGGCCAGCC AGGcTTCCGT
 351   ATCGCTCATC AaCAACAAAG GCAATATCGG TAaCACCCTG AAAGAGCTGG
 401   GCAGAAGCAG CACGGTGAAA AATCTGATGG TTGCCGTCGc tACCGCAgGC
 451   GTagCcgaCA AAATCGGTGC TTCGGCACTG AACAATGTCA GCGATAAGCA
 501   GTGGATCAAC AACCTGACCG TCAACCTGGC CAATGCGGGC AGTGCCGCAC
 551   TGATTAATAC CGCTGTCAAC GGCGGCAGCc tgAAAGACAA TCTGGAAGCG
 601   AATATCCTTG CGGCTTTGGT GAATACTGCG CATGGAGAAG CAGCCAGTAA
 651   AATCAAACAG TTGGATCAGC ACTACATTAC CCACAAGATT GCCCaTGCCA
 701   TAGCGGGCTG TGCGGcTGCG GCGGCGAATA AGGGCAAGTG TCAGGATGGT
 751   GCGATAgGTG CGGCTGTGGG CGAGATAGTC GGGGAgGCTT TGACAAACGG
 801   CAAAAATCCT GACACTTTGA CAGCTAAAgA ACGCGaACAG ATTTTGGCAT
 851   ACAGCAAACT GGTTGCCGGT ACGGTAAGCG GTGTGGTCGG CGGCGATGTA
 901   AATGCGGCGG CGAATGCGGC TGAGGTAGCG GTGAAAAATA ATCAGCTTAG
 951   CGACAAAtGA
```

This corresponds to the amino acid sequence <SEQ ID 68; ORF41>:

```
  1   ..QCRLKSSQFY RRHLLCKYIY RFPIYCPXAC VAEDTPYACY LXQLQVTKDV
 51     NWNQVXLAYD KWDYKQEGLT GAGAAIIALA VTVVTAGAGA GAALGLNGAA
101     AAATDAAFAS LASQASVSLI NNKGNIGNTL KELGRSSTVK NLMVAVATAG
151     VADKIGASAL NNVSDKQWIN NLTVNLANAG SAALINTAVN GGSLKDNLEA
201     NILAALVNTA HGEAASKIKQ LDQHYITHKI AHAIAGCAAA AANKGKCQDG
251     AIGAAVGEIV GEALTNGKNP DTLTAKEREQ ILAYSKLVAG TVSGVVGGDV
301     NAAANAAEVA VKNNQLSDK*
```

Further work revealed the complete nucleotide sequence <SEQ ID 69>:

```
   1   ATGCAAGTAA ATATTCAGAT TCCCTATATA CTGCCCAGAT GCGTGCGTGC
  51   TGAAGACACC CCCTACGCTT GCTATTTGAA ACAGCTCCAA GTCACCAAAG
 101   ACGTCAACTG GAACCAGGTA CAACTGGCGT ACGACAAATG GGACTATAAA
 151   CAGGAAGGCT TAACCGGAGC CGGAGCAGCG ATTATTGCGC TGGCTGTTAC
 201   CGTGGTTACT GCGGGCGCGG GAGCCGGAGC CGCACTGGGC TTAAACGGCG
 251   CGGCCGCAGC GGCAACCGAT GCCGCATTCG CCTCGCTGGC CAGCCAGGCT
 301   TCCGTATCGC TCATCAACAA CAAAGGCAAT ATCGGTAACA CCCTGAAAGA
 351   GCTGGGCAGA AGCAGCACGG TGAAAAATCT GATGGTTGCC GTCGCTACCG
 401   CAGGCGTAGC CGACAAAATC GGTGCTTCGG CACTGAACAA TGTCAGCGAT
 451   AAGCAGTGGA TCAACAACCT GACCGTCAAC CTGGCCAATG CGGGCAGTGC
 501   CGCACTGATT AATACCGCTG TCAACGGCGG CAGCCTGAAA GACAATCTGG
 551   AAGCGAATAT CCTTGCGGCT TTGGTGAATA CTGCGCATGG AGAAGCAGCC
 601   AGTAAAATCA AACAGTTGGA TCAGCACTAC ATTACCCACA AGATTGCCCA
 651   TGCCATAGCG GGCTGTGCGG CTGCGGCGGC GAATAAGGGC AAGTGTCAGG
 701   ATGGTGCGAT AGGTGCGGCT GTGGGCGAGA TAGTCGGGGA GGCTTTGACA
 751   AACGGCAAAA ATCCTGACAC TTTGACAGCT AAAGAACGCG AACAGATTTT
 801   GGCATACAGC AAACTGGTTG CCGGTACGGT AAGCGGTGTG GTCGGCGGCG
 851   ATGTAAATGC GGCGGCGAAT GCGGCTGAGG TAGCGGTGAA AAATAATCAG
 901   CTTAGCGACA AAGAGGGTAG AGAATTTGAT AACGAAATGA CTGCATGCGC
 951   CAAACAGAAT AATCCTCAAC TGTGCAGAAA AAATACTGTA AAAAAGTATC
1001   AAAATGTTGC TGATAAAAGA CTTGCTGCTT CGATTGCAAT ATGTACGGAT
1051   ATATCCCGTA GTACTGAATG TAGAACAATC AGAAAACAAC ATTTGATCGA
1101   TAGTAGAAGC CTTCATTCAT CTTGGGAAGC AGGTCTAATT GGTAAAGATG
1151   ATGAATGGTA TAAATTATTC AGCAAATCTT ACACCCAAGC AGATTTGGCT
1201   TTACAGTCTT ATCATTTGAA TACTGCTGCT AAATCTTGGC TTCAATCGGG
1251   CAATACAAAG CCTTTATCCG AATGGATGTC CGACCAAGGT TATACACTTA
1301   TTTCAGGAGT TAATCCTAGA TTCATTCCAA TACCAAGAGG GTTTGTAAAA
1351   CAAAATACAC CTATTACTAA TGTCAAATAC CCGGAAGGCA TCAGTTTCGA
1401   TACAAACCTA AAAAGACATC TGGCAAATGC TGATGGTTTT AGTCAAAAAC
1451   AGGGCATTAA AGGAGCCCAT AACCGCACCA ATTTTATGGC AGAACTAAAT
1501   TCACGAGGAG GACGCGTAAA ATCTGAAACC CAAACTGATA TTGAAGGCAT
1551   TACCCGAATT AAATATGAGA TTCCTACACT AGACAGGACA GGTAAACCTG
1601   ATGGTGGATT TAAGGAAATT TCAAGTATAA AAACTGTTTA TAATCCTAAA
1651   AAATTTTCTG ATGATAAAAT ACTTCAAATG GCTCAAAATG CTGCTTCACA
1701   AGGATATTCA AAAGCCTCTA AAATTGCTCA AAATGAAAGA ACTAAATCAA
1751   TATCGGAAAG AAAAAATGTC ATTCAATTCT CAGAAACCTT TGACGGAATC
1801   AAATTTAGAT CATATTTTGA TGTAAATACA GGAAGAATTA CAAACATTCA
1851   CCCAGAATAA
```

This corresponds to the amino acid sequence <SEQ ID 70; ORF41-1>:

```
  1  MQVNIQIPYI LPRCVRAEDT PYACYLKQLQ VTKDVNWNQV QLAYDKWDYK
 51  QEGLTGAGAA IIALAVTVVT AGAGAGAALG LNGAAAAATD AAFASLASQA
101  SVSLINNKGN IGNTLKELGR SSTVKNLMVA VATAGVADKI GASALNNVSD
151  KQWINNLTVN LANAGSAALI NTAVNGGSLK DNLEANILAA LVNTAHGEAA
201  SKIKQLDQHY ITHKIAHAIA GCAAAAANKG KCQDGAIGAA VGEIVGEALT
251  NGKNPDTLTA KEREQILAYS KLVAGTVSGV VGGDVNAAAN AAEVAVKNNQ
301  LSDKEGREFD NEMTACAKQN NPQLCRKNTV KKYQNVADKR LAASIAICTD
351  ISRSTECRTI RKQHLIDSRS LHSSWEAGLI GKDDEWYKLF SKSYTQADLA
401  LQSYHLNTAA KSWLQSGNTK PLSEWMSDQG YTLISGVNPR FIPIPRGFVK
451  QNTPITNVKY PEGISFDTNL KRHLANADGF SQKQGIKGAH NRTNFMAELN
501  SRGGRVKSET QTDIEGITRI KYEIPTLDRT GKPDGGFKEI SSIKTVYNPK
551  KFSDDKILQM AQNAASQGYS KASKIAQNER TKSISERKNV IQFSETFDGI
601  KFRSYFDVNT GRITNIHPE*
```

Computer analysis of this amino acid sequence predicts a transmembrane domain, and homology with an ORF from *N.meningitidis* (strain A) was also found.

ORF41 shows 92.8% identity over a 279aa overlap with an ORF (ORF41a) from strain A of *N. meningitidis:*

```
                10        20        30        40        50        60        69
orf41.pep       YRRHLLCKYIYRFPIYCPXACVAEDTPYACYLXQLQVTKDVNWNQVXLAYDKWDYKQEGL
                                                  ||  ||||:|::||||| ||||:||||||||
orf41a                                            YLKQLQVAKNINWNQVQLAYDRWDYKQEGL
                                                          10        20        30

                70        80        90       100       110       120       129
orf41.pep       TGAGAAIIALAVTVVTAGAGAGAALGLNGAAAAATDAAFASLASQASVSLINNKGNIGNT
                | ||||||||||||||:|||:||:|||||| ||||||||||||||||||:||||||::|:|
orf41a          TEAGAAIIALAVTVVTSGAGTGAVLGLNGAXAAATDAAFASLASQASVSFINNKGDVGKT
                        40        50        60        70        80        90

               130       140       150       160       170       180       189
orf41.pep       LKELGRSSTVKNLMVAVATAGVADKIGASALNNVSDKQWINNLTVNLANAGSAALINTAV
                ||||||||||||||:||:|||||||||||||| ||||||||||||||||||||||||||||
orf41a          LKELGRSSTVKNLVVAAATAGVADKIGASALXNVSDKQWINNLTVNLANAGSAALINTAV
                       100       110       120       130       140       150

               190       200       210       220       230       240       249
orf41.pep       NGGSLKDNLEANILAALVNTAHGEAASKIKQLDQHYITHKIAHAIAGCAAAAANKGKCQD
                ||||||| |||||||||||||||||||||||||||||:||||||||||||||||||||||||
orf41a          NGGSLKDXLEANILAALVNTAHGEAASKIKQLDQHYIVHKIAHAIAGCAAAAANKGKCQD
                       160       170       180       190       200       210

               250       260       270       280       290       300       309
orf41.pep       GAIGAAVGEIVGEALTNGKNPDTLTAKEREQILAYSKLVAGTVSGVVGGDVNAAANAAEV
                ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
orf41a          GAIGAAVGEIVGEALTNGKNPDTLTAKEREQILAYSKLVAGTVSGVVGGDVNAAANAAEV
                       220       230       240       250       260       270

               310       320
orf41.pep       AVKNNQLSDKX
                |||||||||
orf41a          AVKNNQLSDXEGREFDNEMTACAKQNXPQLCRKNTVKKYQNVADKRLAASIAICTDISRS
                       280       290       300       310       320       330
```

A partial ORF41a nucleotide sequence <SEQ ID 71> is:

```
  1  ..TATCTGAAAC AGCTCCAAGT AGCGAAAAAC ATCAACTGGA ATCAGGTGCA
 51    GCTTGCTTAC GACAGATGGG ACTACAAACA GGAGGGCTTA ACCGAAGCAG
101    GTGCGGCGAT TATCGCACTG GCCGTTACCG TGGTCACCTC AGGCGCAGGA
151    ACCGGAGCCG TATTGGGATT AAACGGTGCG NCCGCCGCCG CAACCGATGC
```

```
 201  AGCATTCGCC TCTTTGGCCA GCCAGGCTTC CGTATCGTTC ATCAACAACA
 251  AAGGCGATGT CGGCAAAACC CTGAAAGAGC TGGGCAGAAG CAGCACGGTG
 301  AAAAATCTGG TGGTTGCCGC CGCTACCGCA GGCGTAGCCG ACAAAATCGG
 351  CGCTTCGGCA CTGANCAATG TCAGCGATAA GCAGTGGATC AACAACCTGA
 401  CCGTCAACCT AGCCAATGCG GGCAGTGCCG CACTGATTAA TACCGCTGTC
 451  AACGGCGGCA GCCTGAAAGA CANTCTGGAA GCGAATATCC TTGCGGCTTT
 501  GGTCAATACC GCGCATGGAG AAGCAGCCAG TAAAATCAAA CAGTTGGATC
 551  AGCACTACAT AGTCCACAAG ATTGCCCATG CCATAGCGGG CTGTGCGGCA
 601  GCGGCGGCGA ATAAGGGCAA GTGTCAGGAT GGTGCGATAG GTGCGGCTGT
 651  GGGCGAGATA GTCGGGGAGG CTTTGACAAA CGGCAAAAAT CCTGACACTT
 701  TGACAGCTAA AGAACGCGAA CAGATTTTGG CATACAGCAA ACTGGTTGCC
 751  GGTACGGTAA GCGGTGTGGT CGGCGGCGAT GTAAATGCGG CGGCGAATGC
 801  GGCTGAGGTA GCGGTGAAAA ATAATCAGCT TAGCGACNAA GAGGGTAGAG
 851  AATTTGATAA CGAAATGACT GCATGCGCCA AACAGAATAN TCCTCAACTG
 901  TGCAGAAAAA ATACTGTAAA AAAGTATCAA AATGTTGCTG ATAAAAGACT
 951  TGCTGCTTCG ATTGCAATAT GTACGGATAT ATCCCGTAGT ACTGAATGTA
1001  GAACAATCAG AAAACAACAT TTGATCGATA GTAGAAGCCT TCATTCATCT
1051  TGGGAAGCAG GTCTAATTGG TAAAGATGAT GAATGGTATA AATTATTCAG
1101  CAAATCTTAC ACCCAAGCAG ATTTGGCTTT ACAGTCTTAT CATTTGAATA
1151  CTGCTGCTAA ATCTTGGCTT CAATCGGGCA ATACAAAGCC TTTATCCGAA
1201  TGGATGTCCG ACCAAGGTTA TACACTTATT TCAGGAGTTA ATCCTAGATT
1251  CATTCCAATA CCAAGAGGGT TTGTAAAACA AAATACACCT ATTACTAATG
1301  TCAAATACCC GGAAGGCATC AGTTTCGATA CAAACCTANA AAGACATCTG
1351  GCAAATGCTG ATGGTTTTAG TCAAGAACAG GGCATTAAAG GAGCCCATAA
1401  CCGCACCAAT NTTATGGCAG AACTAAATTC ACGAGGAGGA NGNGTAAAAT
1451  CTGAAACCCA NACTGATATT GAAGGCATTA CCCGAATTAA ATATGAGATT
1501  CCTACACTAG ACAGGACAGG TAAACCTGAT GGTGGATTTA AGGAAATTTC
1551  AAGTATAAAA ACTGTTTATA ATCCTAAAAA NTTTTNNGAT GATAAAATAC
1601  TTCAAATGGC TCAANATGCT GNTTCACAAG GATATTCAAA AGCCTCTAAA
1651  ATTGCTCAAA ATGAAAGAAC TAAATCAATA TCGGAAAGAA AAAATGTCAT
1701  TCAATTCTCA GAAACCTTTG ACGGAATCAA ATTTAGANNN TATNTNGATG
1751  TAAATACAGG AAGAATTACA AACATTCACC CAGAATAA
```

This encodes a protein having the partial amino acid sequence <SEQ ID 72>:

```
   1  YLKQLQVAKN INWNQVQLAY DRWDYKQEGL TEAGAAIIAL AVTVVTSGAG
  51  TGAVLGLNGA XAAATDAAFA SLASQASVSF INNKGDVGKT LKELGRSSTV
 101  KNLVVAAATA GVADKIGASA LXNVSDKQWI NNLTVNLANA GSAALINTAV
 151  NGGSLKDXLE ANILAALVNT AHGEAASKIK QLDQHYIVHK IAHAIAGCAA
 201  AAANKGKCQD GAIGAAVGEI VGEALTNGKN PDTLTAKERE QILAYSKLVA
 251  GTVSGVVGGD VNAAANAAEV AVKNNQLSDX EGREFDNEMT ACAKQNXPQL
 301  CRKNTVKKYQ NVADKRLAAS IAICTDISRS TECRTIRKQH LIDSRSLHSS
 351  WEAGLIGKDD EWYKLFSKSY TQADLALQSY HLNTAAKSWL QSGNTKPLSE
 401  WMSDQGYTLI SGVNPRFIPI PRGFVKQNTP ITNVKYPEGI SFDTNLXRHL
 451  ANADGFSQEQ GIKGAHNRTN XMAELNSRGG XVKSETXTDI EGITRIKYEI
 501  PTLDRTGKPD GGFKEISSIK TVYNPKXFXD DKILQMAQXA XSQGYSKASK
 551  IAQNERTKSI SERKNVIQFS ETFDGIKFRX YXDVNTGRIT NIHPE*
```

ORF41a and ORF41-1 show 94.8% identity in 595 aa overlap:

```
                                          10        20        30
orf41a.pep                               YLKQLQVAKNINWNQVQLAYDRWDYKQEGLTEAGAA
                                         |||||||:|::||||||||||:||||||||| ||||
orf41-1      MQVNIQIPYILPRCVRAEDTPYACYLKQLQVTKDVNWNQVQLAYDKWDYKQEGLTGAGAA
                    10        20        30        40        50        60

                40        50        60        70        80        90
orf41a.pep   IIALAVTVVTSGAGTGAVLGLNGAXAAATDAAFASLASQASVSFINNKGDVGKTLKELGR
             ||||||||||:|||:||:|||||| ||||||||||||||||||:||||::|:||||||||
orf41-1      IIALAVTVVTAGAGAGAALGLNGAAAAATDAAFASLASQASVSLINNKGNIGNTLKELGR
                    70        80        90       100       110       120

               100       110       120       130       140       150
orf41a.pep   SSTVKNLVVAAATAGVADKIGASALXNVSDKQWINNLTVNLANAGSAALINTAVNGGSLK
             |||||||:||:|||||||||||||||| ||||||||||||||||||||||||||||||||
orf41-1      SSTVKNLMVAVATAGVADKIGASALNNVSDKQWINNLTVNLANAGSAALINTAVNGGSLK
                   130       140       150       160       170       180

               160       170       180       190       200       210
orf41a.pep   DXLEANILAALVNTAHGEAASKIKQLDQHYIVHKIAHAIAGCAAAAANKGKCQDGAIGAA
```

```
                | |||||||||||||||||||||||||||||||:|||||||||||||||||||||||||||
orf41-1         DNLEANILAALVNTAHGEAASKIKQLDQHYITHKIAHAIAGCAAAAAANKGKCQDGAIGAA
                       190       200       210       220       230       240

                  220       230       240       250       260       270
orf41a.pep      VGEIVGEALTNGKNPDTLTAKEREQILAYSKLVAGTVSGVVGGDVNAAANAAEVAVKNNQ
                ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
orf41-1         VGEIVGEALTNGKNPDTLTAKEREQILAYSKLVAGTVSGVVGGDVNAAANAAEVAVKNNQ
                       250       260       270       280       290       300

                  280       290       300       310       320       330
orf41a.pep      LSDXEGREFDNEMTACAKQNXPQLCRKNTVKKYQNVADKRLAASIAICTDISRSTECRTI
                ||| |||||||||||||||| |||||||||||||||||||||||||||||||||||||||
orf41-1         LSDKEGREFDNEMTACAKQNNPQLCRKNTVKKYQNVADKRLAASIAICTDISRSTECRTI
                       310       320       330       340       350       360

                  340       350       360       370       380       390
orf41a.pep      RKQHLIDSRSLHSSWEAGLIGKDDEWYKLFSKSYTQADLALQSYHLNTAAKSWLQSGNTK
                ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
orf41-1         RKQHLIDSRSLHSSWEAGLIGKDDEWYKLFSKSYTQADLALQSYHLNTAAKSWLQSGNTK
                       370       380       390       400       410       420

                  400       410       420       430       440       450
orf41a.pep      PLSEWMSDQGYTLISGVNPRFIPIPRGFVKQNTPITNVKYPEGISFDTNLXRHLANADGF
                |||||||||||||||||||||||||||||||||||||||||||||||||| |||||||||
orf41-1         PLSEWMSDQGYTLISGVNPRFIPIPRGFVKQNTPITNVKYPEGISFDTNLKRHLANADGF
                       430       440       450       460       470       480

                  460       470       480       490       500       510
orf41a.pep      SQEQGIKGAHNRTNXMAELNSRGGXVKSETXTDIEGITRIKYEIPTLDRTGKPDGGFKEI
                ||:||||||||||| ||||||||| ||||| |||||||||||||||||||||||||||||
orf41-1         SQKQGIKGAHNRTNFMAELNSRGGRVKSETQTDIEGITRIKYEIPTLDRTGKPDGGFKEI
                       490       500       510       520       530       540

                  520       530       540       550       560       570
orf41a.pep      SSIKTVYNPKXFXDDKILQMAQXAXSQGYSKASKIAQNERTKSISERKNVIQFSETFDGI
                |||||||||| | ||||||||| | |||||||||||||||||||||||||||||||||||
orf41-1         SSIKTVYNPKKFSDDKILQMAQNAASQGYSKASKIAQNERTKSISERKNVIQFSETFDGI
                       550       560       570       580       590       600

                  580       590
orf41a.pep      KFRXYXDVNTGRITNIHPEX
                ||| | ||||||||||||||
orf41-1         KFRSYFDVNTGRITNIHPEX
                       610       620
```

Amino acids 25-619 of ORF41-1 were amplified as described above. Figure 6 shows plots of hydrophilicity, antigenic index, and AMPHI regions for ORF41-1.

Based on this analysis, it is predicted that this protein from *N meningitidis*, and its epitopes, could be useful antigens for vaccines or diagnostics.

Example 17

The following DNA sequence was identified in *N.meningitidis* <SEQ ID 73>

```
  1  ATGGCAATCA TTACATTGTA TTATTCTGTC AATGGTATTT TAAATGTATG
 51  TGCAAAAGCA AAAAATATTC AAGTAGTTGC CAATAATAAG AATATGGTTC
101  TTTTTGGGTT TTTGGsmrGC ATCATCGGCG GTTCAACCAA TGCCATGTCT
151  CCCATATTGT TAATATTTTT GCTTAGCGAA ACAGAAAATA AAAATcgTAT
201  CGTAAAATCA AGCAATCTAT GCTATCTTTT GGCGAAAATT GTTCAAATAT
251  ATATGCTAAG AGACCAGTAT TGGTTATTAA ATAAGAGTGA ATACGdTTTA
301  ATATTTTTAC TGTCCGTATT GTCTGTTATT GGATTGTATG TTGGAATTCG
351  GTTAAGGACT AAGATTAGCC CAaATTTTTT TAAAATGTTA ATTTTTATTG
```

```
401  tTTTATTGGT ATTGGCtCTG AAAATCGGGC AttCGGGTTT AAtCAAACTT
451  TAA
```

This corresponds to the amino acid sequence <SEQ ID 74; ORF51>:

```
  1  MAIITLYYSV NGILNVCAKA KNIQVVANNK NMVLFGFLXX IIGGSTNAMS
 51  PILLIFLLSE TENKNRIVKS SNLCYLLAKI VQIYMLRDQY WLLNKSEYXL
101  IFLLSVLSVI GLYVGIRLRT KISPNFFKML IFIVLLVLAL KIGHSGLIKL
151  *
```

Further work revealed the complete nucleotide sequence <SEQ ID 75>:

```
  1  ATGCAAGAAA TAATGCAATC TATCGTTTTT GTTGCTGCCG CAATACTGCA
 51  CGGAATTACA GGCATGGGAT TTCCGATGCT CGGTACAACC GCATTGGCTT
101  TTATCATGCC ATTGTCTAAG GTTGTTGCCT TGGTGGCATT ACCAAGCCTG
151  TTAATGAGCT TGTTGGTTCT ATGCAGCAAT AACAAAAAGG GTTTTTGGCA
201  AGAGATTGTT TATTATTTAA AAACCTATAA ATTGCTTGCT ATCGGCAGCG
251  TCGTTGGCAG CATTTTGGGG GTGAAGTTGC TTTTGATACT TCCAGTGTCT
301  TGGCTGCTTT TACTGATGGC AATCATTACA TTGTATTATT CTGTCAATGG
351  TATTTTAAAT GTATGTGCAA AAGCAAAAAA TATTCAAGTA GTTGCCAATA
401  ATAAGAATAT GGTTCTTTTT GGGTTTTTGG CAGGCATCAT CGGCGGTTCA
451  ACCAATGCCA TGTCTCCCAT ATTGTTAATA TTTTTGCTTA GCGAAACAGA
501  AAATAAAAAT CGTATCGTAA AATCAAGCAA TCTATGCTAT CTTTTGGCGA
551  AAATTGTTCA AATATATATG CTAAGAGACC AGTATTGGTT ATTAAATAAG
601  AGTGAATACG GTTTAATATT TTTACTGTCC GTATTGTCTG TTATTGGATT
651  GTATGTTGGA ATTCGGTTAA GGACTAAGAT TAGCCCAAAT TTTTTTAAAA
701  TGTTAATTTT TATTGTTTTA TTGGTATTGG CTCTGAAAAT CGGGCATTCG
751  GGTTTAATCA AACTTTAA
```

This corresponds to the amino acid sequence <SEQ ID 76; ORF51-1>:

```
  1  MQEIMQSIVF VAAAILHGIT GMGFPMLGTT ALAFIMPLSK VVALVALPSL
 51  LMSLLVLCSN NKKGFWQEIV YYLKTYKLLA IGSVVGSILG VKLLLILPVS
101  WLLLLMAIIT LYYSVNGILN VCAKAKNIQV VANNKNMVLF GFLAGIIGGS
151  TNAMSPILLI FLLSETENKN RIVKSSNLCY LLAKIVQIYM LRDQYWLLNK
201  SEYGLIFLLS VLSVIGLYVG IRLRTKISPN FFKMLIFIVL LVLALKIGHS
251  GLIKL*
```

Computer analysis of this amino acid sequence reveals three putative transmembrane domains. A corresponding ORF from strain A of *N.meningitidis* was also identified:

Homology with a predicted ORF from *N.meningitidis* (strain A)

ORF51 shows 96.7% identity over a 150aa overlap with an ORF (ORF51a) from strain A of *N. meningitidis:*

```
                                                    10        20        30
orf51.pep                                           MAIITLYYSVNGILNVCAKAKNIQVVANNK
                                                    ||||||||||||||||||||||||||||||
orf51a      YKLLAIGSVVGSILGVKLLLILPVSWLLLLMAIITLYYSVNGILNVCAKAKNIQVVANNK
              80        90        100       110       120       130

                     40        50        60        70        80        90
orf51.pep   NMVLFGFLXXIIGGSTNAMSPILLIFLLSETENKNRIVKSSNLCYLLAKIVQIYMLRDQY
            ||||||||  ||||||||||||||||||||||||||||:||||||||||||||||||||
orf51a      NMVLFGFLAGIIGGSTNAMSPILLIFLLSETENKNRIAKSSNLCYLLAKIVQIYMLRDQY
              140       150       160       170       180       190

                   100       110       120       130       140       150
orf51.pep   WLLNKSEYXLIFLLSVLSVIGLYVGIRLRTKISPNFFKMLIFIVLLVLALKIGHSGLIKL
            ||||||||  ||||||||||||||||||||||||||||||||||||||||||||||||:|||||
orf51a      WLLNKSEYGLIFLLSVLSVIGLYVGIRLRTKISPNFFKMLIFIVLLVLALKIGYSGLIKL
              200       210       220       230       240       250
```

ORF51-1 and ORF51a show 99.2% identity in 255 aa overlap:

```
orf51a.pep   MQEIMQSIVFVAAAILHGITGMGFPMLGTTALAFIMPLSKVVALVALPSLLMSLLVLCSN
             ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
orf51-1      MQEIMQSIVFVAAAILHGITGMGFPMLGTTALAFIMPLSKVVALVALPSLLMSLLVLCSN

orf51a.pep   NKKGFWQEIVYYLKTYKLLAIGSVVGSILGVKLLLILPVSWLLLLMAIITLYYSVNGILN
             ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
orf51-1      NKKGFWQEIVYYLKTYKLLAIGSVVGSILGVKLLLILPVSWLLLLMAIITLYYSVNGILN

orf51a.pep   VCAKAKNIQVVANNKNMVLFGFLAGIIGGSTNAMSPILLIFLLSETENKNRIAKSSNLCY
             |||||||||||||||||||||||||||||||||||||||||||||||||||:|||||||
orf51-1      VCAKAKNIQVVANNKNMVLFGFLAGIIGGSTNAMSPILLIFLLSETENKNRIVKSSNLCY

orf51a.pep   LLAKIVQIYMLRDQYWLLNKSEYGLIFLLSVLSVIGLYVGIRLRTKISPNFFKMLIFIVL
             ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
orf51-1      LLAKIVQIYMLRDQYWLLNKSEYGLIFLLSVLSVIGLYVGIRLRTKISPNFFKMLIFIVL

orf51a.pep   LVLALKIGYSGLIKLX
             ||||||||:|||||||
orf51-1      LVLALKIGHSGLIKLX
```

The complete length ORF51a nucleotide sequence <SEQ ID 77> is:

```
   1  ATGCAAGAAA TAATGCAATC TATCGTTTTT GTTGCTGCCG CAATACTGCA
  51  CGGAATTACA GGCATGGGAT TTCCGATGCT CGGTACAACC GCATTGGCTT
 101  TTATCATGCC ATTGTCTAAG GTTGTTGCCT TGGTGGCATT ACCAAGCCTG
 151  TTAATGAGCT TGTTGGTTCT ATGCAGCAAT AACAAAAAGG GTTTTTGGCA
 201  AGAGATTGTT TATTATTTAA AAACCTATAA ATTGCTTGCT ATCGGCAGCG
 251  TCGTTGGCAG CATTTTGGGG GTGAAGTTGC TTTTGATACT TCCAGTGTCT
 301  TGGCTGCTTT TACTGATGGC AATCATTACA TTGTATTATT CTGTCAATGG
 351  TATTTTAAAT GTATGTGCAA AAGCAAAAAA TATTCAAGTA GTTGCCAATA
 401  ATAAGAATAT GGTTCTTTTT GGGTTTTTGG CAGGCATCAT CGGCGGTTCA
 451  ACCAATGCCA TGTCTCCCAT ATTGTTAATA TTTTTGCTTA GCGAAACAGA
 501  GAATAAAAAT CGTATCGCAA AATCAAGCAA TCTATGCTAT CTTTTGGCAA
 551  AAATTGTTCA AATATATATG CTAAGAGACC AGTATTGGTT ATTAAATAAG
 601  AGTGAATACG GTTTAATATT TTTACTGTCC GTATTGTCTG TTATTGGATT
 651  GTATGTTGGA ATTCGGTTAA GGACTAAGAT TAGCCCAAAT TTTTTTAAAA
 701  TGTTAATTTT TATTGTTTTA TTGGTATTGG CTCTGAAAAT CGGGTATTCA
 751  GGTTTAATCA AACTTTAA
```

This encodes a protein having amino acid sequence <SEQ ID 78>:

```
   1  MQEIMQSIVF VAAAILHGIT GMGFPMLGTT ALAFIMPLSK VVALVALPSL
  51  LMSLLVLCSN NKKGFWQEIV YYLKTYKLLA IGSVVGSILG VKLLLILPVS
 101  WLLLLMAIIT LYYSVNGILN VCAKAKNIQV VANNKNMVLF GFLAGIIGGS
 151  TNAMSPILLI FLLSETENKN RIAKSSNLCY LLAKIVQIYM LRDQYWLLNK
 201  SEYGLIFLLS VLSVIGLYVG IRLRTKISPN FFKMLIFIVL LVLALKIGYS
 251  GLIKL*
```

Based on this analysis, it is predicted that this protein from *N meningitidis*, and its epitopes, could be useful antigens for vaccines or diagnostics.

Example 18

The following partial DNA sequence was identified in *N.meningitidis* <SEQ ID 79>

```
   1  ATGAGACATA TGAAAATACA AAATTATTTA CTAGTATTTA TAGTTTTACA
  51  TATAGCCTTG ATAGTAATTA ATATAGTGTT TGGTTATTTT GTTTTTCTAT
 101  TTGATTTTTT TGCGTTTTTG TTTTTTGCAA ACGTCTTTCT TGCTGTAAAT
 151  TTATTATTTT TAGAAAAAAA CATAAAAAAC AAATTATTGT TTTTATTGCC
 201  GATTTCTATT ATTATATGGA TGGTAATTCA TATTAGTATG ATAAATATAA
 251  AATTTTATAA ATTTGAGCAT CAAATAAAGG AACAAAATAT ATCCTCGATT
 301  ACTGGGGTGA TAAAACCACA TGATAGTTAT AATTATGTTT ATGACTCAAA
```

```
351  TGGATATGCT AAATTAAAAG ATAATCATAG ATATGGTAGG GTAATTAGAG
401  AAACACCTTA TATTGATGTA GTTGCATCTG ATGTTAAAAA TAAATCCATA
451  AGATTAAGCT TGGTTTGTGG TATTCATTCA TATGCTCCAT GTGCCAATTT
501  TATAAAATTT GTCAGG..
```

This corresponds to the amino acid sequence <SEQ ID 80; ORF82>:

```
  1  MRHMKIQNYL LVFIVLHIAL IVINIVFGYF VFLFDFFAFL FFANVFLAVN
 51  LLFLEKNIKN KLLFLLPISI IIWMVIHISM INIKFYKFEH QIKEQNISSI
101  TGVIKPHDSY NYVYDSNGYA KLKDNHRYGR VIRETPYIDV VASDVKNKSI
151  RLSLVCGIHS YAPCANFIKF VR..
```

Further work revealed the complete nucleotide sequence <SEQ ID 81>:

```
  1  ATGAGACATA TGAAAAATAA AAATTATTTA CTAGTATTTA TAGTTTTACA
 51  TATAGCCTTG ATAGTAATTA ATATAGTGTT TGGTTATTTT GTTTTTCTAT
101  TTGATTTTTT TGCGTTTTTG TTTTTTGCAA ACGTCTTTCT TGCTGTAAAT
151  TTATTATTTT TAGAAAAAAA CATAAAAAAC AAATTATTGT TTTTATTGCC
201  GATTTCTATT ATTATATGGA TGGTAATTCA TATTAGTATG ATAAATATAA
251  AATTTTATAA ATTTGAGCAT CAAATAAAGG AACAAAATAT ATCCTCGATT
301  ACTGGGGTGA TAAAACCACA TGATAGTTAT AATTATGTTT ATGACTCAAA
351  TGGATATGCT AAATTAAAAG ATAATCATAG ATATGGTAGG GTAATTAGAG
401  AAACACCTTA TATTGATGTA GTTGCATCTG ATGTTAAAAA TAAATCCATA
451  AGATTAAGCT TGGTTTGTGG TATTCATTCA TATGCTCCAT GTGCCAATTT
501  TATAAAATTT GCAAAAAAAC CTGTTAAAAT TTATTTTTAT AATCAACCTC
551  AAGGAGATTT TATAGATAAT GTAATATTTG AAATTAATGA TGGAAACAAA
601  AGTTTGTACT TGTTAGATAA GTATAAAACA TTTTTTCTTA TTGAAAACAG
651  TGTTTGTATC GTATTAATTA TTTTATATTT AAAATTTAAT TTGCTTTTAT
701  ATAGGACTTA CTTCAATGAG TTGGAATAG
```

This corresponds to the amino acid sequence <SEQ ID 82; ORF82-1>:

```
  1  MRHMKNKNYL LVFIVLHIAL IVINIVFGYF VFLFDFFAFL FFANVFLAVN
 51  LLFLEKNIKN KLLFLLPISI IIWMVIHISM INIKFYKFEH QIKEQNISSI
101  TGVIKPHDSY NYVYDSNGYA KLKDNHRYGR VIRETPYIDV VASDVKNKSI
151  RLSLVCGIHS YAPCANFIKF AKKPVKIYFY NQPQGDFIDN VIFEINDGNK
201  SLYLLDKYKT FFLIENSVCI VLIILYLKFN LLLYRTYFNE LE*
```

Computer analysis of this amino acid sequence reveals a predicted leader peptide.

A corresponding ORF from strain A of *N.meningitidis* was also identified:

Homology with a predicted ORF from *N.meningitidis* (strain A)

ORF82 shows 97.1% identity over a 172aa overlap with an ORF (ORF82a) from strain A of *N. meningitidis:*

```
                10        20        30        40        50        60
orf82.pep       MRHMKIQNYLLVFIVLHIALIVINIVFGYFVFLFDFFAFLFFANVFLAVNLLFLEKNIKN
                ||||| :|||||||||||:|||||||||||||||||||||||||||||||||||||||||
orf82a          MRHMKNKNYLLVFIVLHITLIVINIVFGYFVFLFDFFAFLFFANVFLAVNLLFLEKNIKN
                10        20        30        40        50        60

                70        80        90        100       110       120
orf82.pep       KLLFLLPISIIIWMVIHISMINIKFYKFEHQIKEQNISSITGVIKPHDSYNYVYDSNGYA
                ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
orf82a          KLLFLLPISIIIWMVIHISMINIKFYKFEHQIKEQNISSITGVIKPHDSYNYVYDSNGYA
                70        80        90        100       110       120

               130       140       150       160       170
orf82.pep       KLKDNHRYGRVIRETPYIDVVASDVKNKSIRLSLVCGIHSYAPCANFIKFVR
                |||||||||||||||||||||||||||||||||||||||||||||||||||::
orf82a          KLKDNHRYGRVIRETPYIDVVASDVKNKSIRLSLVCGIHSYAPCANFIKFAKKPVKIYFY
                130       140       150       160       170       180
```

ORF82a and ORF82-1 show 99.2% identity in 242 aa overlap:

```
orf82a.pep   MRHMKNKNYLLVFIVLHITLIVINIVFGYFVFLFDFFAFLFFANVFLAVNLLFLEKNIKN
             ||||||||||||||||||:||||||||||||||||||||||||||||||||||||||||
orf82-1      MRHMKNKNYLLVFIVLHIALIVINIVFGYFVFLFDFFAFLFFANVFLAVNLLFLEKNIKN

orf82a.pep   KLLFLLPISIIIWMVIHISMINIKFYKFEHQIKEQNISSITGVIKPHDSYNYVYDSNGYA
             ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
orf82-1      KLLFLLPISIIIWMVIHISMINIKFYKFEHQIKEQNISSITGVIKPHDSYNYVYDSNGYA

orf82a.pep   KLKDNHRYGRVIRETPYIDVVASDVKNKSIRLSLVCGIHSYAPCANFIKFAKKPVKIYFY
             ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
orf82-1      KLKDNHRYGRVIRETPYIDVVASDVKNKSIRLSLVCGIHSYAPCANFIKFAKKPVKIYFY

orf82a.pep   NQPQGDFIDNVIFEINDGKKSLYLLDKYKTFFLIENSVCIVLIILYLKFNLLLYRTYFNE
             ||||||||||||||||||:||||||||||||||||||||||||||||||||||||||||
orf82-1      NQPQGDFIDNVIFEINDGNKSLYLLDKYKTFFLIENSVCIVLIILYLKFNLLLYRTYFNE

orf82a.pep   LEX
             |||
orf82-1      LEX
```

The complete length ORF82a nucleotide sequence <SEQ ID 83> is:

```
   1  ATGAGACATA TGAAAAATAA AAATTATTTA CTAGTATTTA TAGTTTTACA
  51  TATAACCTTG ATAGTAATTA ATATAGTGTT TGGTTATTTT GTTTTTCTAT
 101  TTGATTTTTT TGCGTTTTTG TTTTTTGCAA ACGTCTTTCT TGCTGTAAAT
 151  TTATTATTTT TAGAAAAAAA CATAAAAAAC AAATTATTGT TTTTATTGCC
 201  GATTTCTATT ATTATATGGA TGGTAATTCA TATTAGTATG ATAAATATAA
 251  AATTTTATAA ATTTGAGCAT CAAATAAAGG AACAAAATAT ATCCTCGATT
 301  ACTGGGGTGA TAAAACCACA TGATAGTTAT AATTATGTTT ATGACTCAAA
 351  TGGATATGCT AAATTAAAAG ATAATCATAG ATATGGTAGG GTAATTAGAG
 401  AAACACCTTA TATTGATGTA GTTGCATCTG ATGTTAAAAA TAAATCCATA
 451  AGATTAAGCT TGGTTTGTGG TATTCATTCA TATGCTCCAT GTGCCAATTT
 501  TATAAAATTT GCAAAAAAAC CTGTTAAAAT TTATTTTTAT AATCAACCTC
 551  AAGGAGATTT TATAGATAAT GTAATATTTG AAATTAATGA TGGAAAAAAA
 601  AGTTTGTACT TGTTAGATAA GTATAAAACA TTTTTTCTTA TTGAAAACAG
 651  TGTTTGTATC GTATTAATTA TTTTATATTT AAAATTTAAT TTGCTTTTAT
 701  ATAGGACTTA CTTCAATGAG TTGGAATAG
```

This encodes a protein having amino acid sequence <SEQ ID 84>:

```
   1  MRHMKNKNYL LVFIVLHITL IVINIVFGYF VFLFDFFAFL FFANVFLAVN
  51  LLFLEKNIKN KLLFLLPISI IIWMVIHISM INIKFYKFEH QIKEQNISSI
 101  TGVIKPHDSY NYVYDSNGYA KLKDNHRYGR VIRETPYIDV VASDVKNKSI
 151  RLSLVCGIHS YAPCANFIKF AKKPVKIYFY NQPQGDFIDN VIFEINDGKK
 201  SLYLLDKYKT FFLIENSVCI VLIILYLKFN LLLYRTYFNE LE*
```

Based on this analysis, it is predicted that this protein from *N.meningitidis*, and its epitopes, could be useful antigens for vaccines or diagnostics.

Example 19

The following partial DNA sequence was identified in *N.meningitidis* <SEQ ID 85>

```
   1  ..ACCCCCAACA GCGTGACCGT CTTGCCGTCT TTCGGCGGAT TCGGGCGTAC
  51    CGGCGCGACC ATCAATGCAG CAGGCGGGGT CGGCATGACT GCCTTTTCGA
 101    CAACCTTAAT TTCCGTAGCC GAGGGCGCGG TTGTAGAGCT GCAGGCCGTG
 151    AGAGCCAAAG CCGTCAATGC AACCGCCGCT TGCATTTTTA CGGTCTTGAG
 201    TAAGGACATT TTCGATTTCC TTTTTATTTT CCGTTTTCAG ACGGCTGACT
 251    TCCGCCTGTA TTTTCGCCAA AGCCATGCCG ACAGCGTGCG CCTTGACTTC
 301    ATATTTAAAA GCTTCCGCGC GTGCCAGTTC CAGTTCGCGC GCATAGTTTT
 351    GAGCCGACAA CAGCAGGGCT TGCGCCTTGT CGCGCTCCAT CTTGTCGATG
```

```
401    ACCGCCTGCA GCTTCGCAAA TGCCGACTTG TAGCCTTGAT GGTGCGACAC
451    AGCCAAGCCC GTGCCGACAA GCGCGATAAT GGCAATCGGT TGCCAGTAAT
501    TCGCCAGCAG TTTCACGAGA TTCATTCTCG ACCTCCTGAC GCTTCACGCT
551    GA
```

This corresponds to the amino acid sequence <SEQ ID 86; ORF124>.

```
  1  ..TPNSVTVLPS FGGFGRTGAT INAAGGVGMT AFSTTLISVA EGAVVELQAV
 51    RAKAVNATAA CIFTVLSKDI FDFLFIFRFQ TADFRLYFRQ SHADSVRLDF
101    IFKSFRACQF QFARIVLSRQ QQGLRLVALH LVDDRLQLRK CRLVALMVRH
151    SQARADKRDN GNRLPVIRQQ FHEIHSRPPD ASR*
```

Computer analysis of this amino acid sequence predicts a transmembrane domain.

Further work revealed the complete nucleotide sequence <SEQ ID 87>:

```
  1  ATGACTGCCT TTTCGACAAC CTTAATTTCC GTAGCCGAGG GCGCGGTTGT
 51  AGAGCTGCAG GCCGTGAGAG CCAAAGCCGT CAATGCAACC GCCGCTTGCA
101  TTTTTACGGT CTTGAGTAAG GACATTTTCG ATTTCCTTTT TATTTTCCGT
151  TTTCAGACGG CTGACTTCCG CCTGTTTTTT CGCCAAAGCC ATGCCGACAG
201  CGTGCGCCTT GACTTCATAT TTTTTAGCTT CCGCGCGTGC CAGTTCCAGT
251  TCGCGCGCAT AGTTTTGAGC CGACAACAGC AGGGCTTGCG CCTTGTCGCG
301  CTCCATCTTG TCGATGACCG CCTGCTGCTT CGCAAATGCC GACTTGTAGC
351  CTTGATGGTG CGACACAGCC AAGCCCGTGC CGACAAGCGC GATAATGGCA
401  ATCGGTTGCC AGTTATTCGC CAGCAGTTTC ACGAGATTCA TTCTCGACCT
451  CCTGACGCTT CACGCTGA
```

This corresponds to the amino acid sequence <SEQ ID 88; ORF124-1>:

```
  1  MTAFSTTLIS VAEGAVVELQ AVRAKAVNAT AACIFTVLSK DIFDFLFIFR
 51  FQTADFRLFF RQSHADSVRL DFIFFSFRAC QFQFARIVLS RQQQGLRLVA
101  LHLVDDRLLL RKCRLVALMV RHSQARADKR DNGNRLPVIR QQFHEIHSRP
151  PDASR*
```

A corresponding ORF from strain A of *N.meningitidis* was also identified:

Homology with a predicted ORF from *N.meningitidis* (strain A)

ORF124 shows 87.5% identity over a 152aa overlap with an ORF (ORF124a) from strain A of *N. meningitidis:*

```
                 10        20        30        40        50        60
orf124.pep   TPNSVTVLPSFGGFGRTGATINAAGGVGMTAFSTTLISVAEGAVVELQAVRAKAVNATAA
                                       ||||||||||||||||:|||||| |||||:|||
orf124a                                MTAFSTTLISVAEGALVELQAVMAKAVNTTAA
                                              10        20        30

                 70        80        90       100       110       120
orf124.pep   CIFTVLSKDIFDFLFIFRFQTADFRLYFRQSHADSVRLDFIFKSFRACQFQFARIVLSRQ
             ||||||||||||||||||||||||||:|||||||:|||||| |||:  |||| :||||||
orf124a      CIFTVLSKDIFDFLFIFRFQTADFRLFFRQSHADGVRLDFIFFSFRTRLFQFAGVVLSRQ
                   40        50        60        70        80        90

                130       140       150       160       170       180
orf124.pep   QQGLRLVALHLVDDRLQLRKCRLVALMVRHSQARADKRDNGNRLPVIRQQFHEIHSRPPD
             ||||||||||:::||| ||| |||||||| |:|||||:||||||||||||||||||||||
orf124a      QQGLRLVALHFLNDRLLLRKSRLVALMVRHRQTRADKRDDGNRLPVIRQQFHEIHSRPPD
                  100       110       120       130       140       150

orf124.pep   ASRX
             :
orf124a      VX
```

ORF124a and ORF124-1 show 89.5% identity in 152 aa overlap:

```
orf124-1.pep    MTAFSTTLISVAEGAVVELQAVRAKAVNATAACIFTVLSKDIFDFLFIFRFQTADFRLFF
                ||||||||||||||||:|||||| ||||||:||||||||||||||||||||||||||||||
orf124a         MTAFSTTLISVAEGALVELQAVMAKAVNTTAACIFTVLSKDIFDFLFIFRFQTADFRLFF

orf124-1.pep    RQSHADSVRLDFIFFSFRACQFQFARIVLSRQQQGLRLVALHLVDDRLLLRKCRLVALMV
                ||||||:|||||||||||:  |||| :|||||||||||||||:::|||||||  |||||||
orf124a         RQSHADGVRLDFIFFSFRTRLFQFAGVVLSRQQQGLRLVALHFLNDRLLLRKSRLVALMV

orf124-1.pep    RHSQARADKRDNGNRLPVIRQQFHEIHSRPPDASRX
                ||  |:||||||:||||||||||||||||||||:
orf124a         RHRQTRADKRDDGNRLPVIRQQFHEIHSRPPDVX
```

The complete length ORF124a nucleotide sequence <SEQ ID 89> is:

```
   1  ATGACCGCCT TTTCGACAAC CTTAATTTCC GTAGCCGAGG GCGCGCTTGT
  51  AGAGCTGCAA GCCGTGATGG CCAAAGCCGT CAATACAACC GCCGCCTGCA
 101  TTTTTACGGT CTTGAGTAAG GACATTTTCG ATTTCCTTTT TATTTTCCGT
 151  TTTCAGACGG CTGACTTCCG CCTGTTTTTT CGCCAAAGCC ATGCCGACGG
 201  CGTGCGCCTT GACTTCATAT TTTTTAGCTT CCGCACGCGC CTGTTCCAGT
 251  TCGCGGGCGT AGTTTTGAGC CGACAACAGC AGGGCTTGCG CCTTGTCGCG
 301  CTTCATTTTC TCAATGACCG CCTGCTGCTT CGCAAAAGCC GACTTGTAGC
 351  CTTGATGGTG CGACACCGCC AAACCCGTGC CGACAAGCGC GATGATGGCA
 401  ATCGGTTGCC AGTTATTCGC CAGCAGTTTC ACGAGATTCA TTCTCGACCT
 451  CCTGACGTTT GA
```

This encodes a protein having amino acid sequence <SEQ ID 90>:

```
   1  MTAFSTTLIS VAEGALVELQ AVMAKAVNTT AACIFTVLSK DIFDFLFIFR
  51  FQTADFRLFF RQSHADGVRL DFIFFSFRTR LFQFAGVVLS RQQQGLRLVA
 101  LHFLNDRLLL RKSRLVALMV RHRQTRADKR DDGNRLPVIR QQFHEIHSRP
 151  PDV*
```

ORF124-1 was amplified as described above. Figure 7 shows plots of hydrophilicity, antigenic index, and AMPHI regions for ORF124-1.

Based on this analysis, it is predicted that this protein from *N.meningitidis*, and its epitopes, could be useful antigens for vaccines or diagnostics.

It will be appreciated that the invention has been described by means of example only, and that modifications may be made whilst remaining within the spirit and scope of the invention.

TABLE I – PCR primers

ORF	Primer	Sequence	Restriction sites
ORF 38	Forward	CGCGGATCCCATATG-TCGCCGCAAAATTCCGA	BamHI-NdeI
	Reverse	CCCGCTCGAG-TTTTGCCGCGTTAAAAGC	XhoI
ORF 40	Forward	CGCGGATCCCATATG-ACCGTGAAGACCGCC	BamHI-NdeI
	Reverse	CCCGCTCGAG-CCACTGATAACCGACAGA	XhoI
ORF 41	Forward	CGCGGATCCCATATG-TATTTGAAACAGCTCCAAG	BamHI-NdeI
	Reverse	CCCGCTCGAG-TTCTGGGTGAATGTTA	XhoI
ORF 44	Forward	GCGGATCCCATATG-GGCACGGACAACCCC	BamHI-NdeI
	Reverse	CCCGCTCGAG-ACGTGGGGAACAGTCT	XhoI
ORF 51	Forward	GCGGATCCCATATG-AAAAATATTCAAGTAGTTGC	BamHI-NdeI
	Reverse	CCCGCTCGAG-AAGTTTGATTAAACCCG	XhoI
ORF 52	Forward	CGCGGATCCCATATG-TGCCAACCGCAATCCG	BamHI-NdeI
	Reverse	CCCGCTCGAG-TTTTTCCAGCTCCGGCA	XhoI
ORF 56	Forward	GCGGATCCCATATG-GTTATCGGAATATTACTCG	BamHI-NdeI
	Reverse	CCCGCTCGAG-GGCTGCAGAAGCTGG	XhoI
ORF 69	Forward	CGCGGATCCCATATG-CGGACGTGGTTGGTTTT	BamHI-NdeI
	Reverse	CCCGCTCGAG-ATATCTTCCGTTTTTTTCAC	XhoI
ORF 82	Forward	CGCGGATCCGCTAGC-GTAAATTTATTATTTTTAGAA	BamHI-NheI
	Reverse	CCCGCTCGAG-TTCCAACTCATTGAAGTA	XhoI
ORF 114	Forward	CGCGGATCCCATATG-AATAAAGGTTTACATCGCAT	BamHI-NheI
	Reverse	CCCGCTCGAG-AATCGCTGCACCGGCT	XhoI
ORF 124	Forward	CGCGGATCCCATATG-ACTGCCTTTTCGACA	BamHI-NheI
	Reverse	CCCGCTCGAG-GCGTGAAGCGTCAGGA	XhoI

TABLE II – Cloning, expression and purification

ORF	PCR/cloning	His-fusion expression	GST-fusion expression	Purification
orf 38	+	+	+	His-fusion
orf 40	+	+	+	His-fusion
orf 41	+	n.d.	n.d.	
orf 44	+	+	+	His-fusion
orf 51	+	n.d.	n.d.	
orf 52	+	n.d.	+	GST-fusion
orf 56	+	n.d.	n.d.	
orf 69	+	n.d.	n.d.	
orf 82	+	n.d.	n.d.	
orf 114	+	n.d.	+	GST-fusion
orf 124	+	n.d.	n.d.	

CLAIMS

1. A protein comprising an amino acid sequence selected from the group consisting of SEQ IDs 2, 4, and 6

2. A protein having 80% or greater sequence identity to a protein according to claim 1.

3. A nucleic acid molecule according to claim 1, comprising a nucleotide sequence selected from the group consisting of SEQ IDs 1, 3, and 5.

4. A nucleic acid molecule comprising a nucleotide sequences having 80% or greater sequence identity to a nucleic acid molecule according to any one of claims 1 to 3.