

IMIDAZOLE BASED LXR MODULATORS

CROSS-REFERENCE TO RELATED APPLICATIONS

This application claims priority to United States Provisional Application Number 60/694,372, filed June 27, 2005, and United States Provisional Application Number 60/736,120, filed November 10, 2005, both of which are hereby incorporated by reference in their entirety.

Field of the Invention

This invention relates to compounds that modulate the activity of liver X receptors (LXRs). The invention also provides pharmaceutical compositions comprising the compounds of the invention and methods of utilizing those compositions for modulating the activity of liver X receptor. In particular, imidazole isomers and derivatives are provided for modulating the activity of LXRs.

BACKGROUND OF THE INVENTION

Nuclear Receptors

Nuclear receptors are a superfamily of regulatory proteins that are structurally and functionally related and are receptors for, e.g., steroids, retinoids, vitamin D and thyroid hormones (see, e.g., Evans (1988) *Science* 240:889-895). These proteins bind to *cis*-acting elements in the promoters of their target genes and modulate gene expression in response to ligands for the receptors.

Nuclear receptors can be classified based on their DNA binding properties (see, e.g., Evans, *supra* and Glass (1994) *Endocr. Rev.* 15:391-407). For example, one class of nuclear receptors includes the glucocorticoid, estrogen, androgen, progestin and mineralocorticoid receptors which bind as homodimers to hormone response elements (HREs) organized as inverted repeats (see, e.g., Glass, *supra*). A second class of receptors, including those activated by retinoic acid, thyroid hormone, vitamin D₃, fatty acids/peroxisome proliferators (i.e., peroxisome proliferator activated receptors or PPARs) and ecdysone, bind to HREs as heterodimers with a common partner, the retinoid X receptors (i.e., RXRs, also known as the 9-*cis* retinoic acid receptors; see, e.g., Levin *et al.* (1992) *Nature* 355:359-361 and Heyman *et al.* (1992) *Cell* 68:397-406).

RXRs are unique among the nuclear receptors in that they bind DNA as a homodimer and are required as a heterodimeric partner for a number of additional nuclear receptors to bind DNA (see, e.g., Mangelsdorf *et al.* (1995) *Cell* 83:841-850). The latter receptors, termed the class II nuclear receptor subfamily, include many which are established or implicated as important regulators of gene expression.

There are three RXR genes (see, e.g., Mangelsdorf *et al.* (1992) *Genes Dev.* 6:329-344), coding for RXR α , β , and γ , all of which are able to heterodimerize with any of the class II receptors, although there appear to be preferences for distinct RXR subtypes by partner receptors *in vivo* (see, e.g., Chiba *et al.* (1997) *Mol. Cell. Biol.* 17:3013-3020). In the adult liver, RXR α is the most abundant of the three

RXRs (see, *e.g.*, Mangelsdorf *et al.* (1992) *Genes Dev.* 6:329-344), suggesting that it might have a prominent role in hepatic functions that involve regulation by class II nuclear receptors. See also, Wan *et al.* (2000) *Mol. Cell. Biol.* 20:4436-4444.

Orphan Nuclear Receptors

5 Included in the nuclear receptor superfamily of regulatory proteins are nuclear receptors for whom the ligand is known and those which lack known ligands. Nuclear receptors falling in the latter category are referred to as orphan nuclear receptors. The search for activators for orphan receptors has led to the discovery of previously unknown signaling pathways (see, *e.g.*, Levin *et al.*, (1992), *supra* and Heyman *et al.*, (1992), *supra*). For example, it has been reported that bile acids, which are involved in
10 physiological processes such as cholesterol catabolism, are ligands for the farnesoid X receptor (FXR).

Because it is known that products of intermediary metabolism act as transcriptional regulators in bacteria and yeast, such molecules may serve similar functions in higher organisms (see, *e.g.*, Tomkins (1975) *Science* 189:760-763 and O'Malley (1989) *Endocrinology* 125:1119-1120). For example, one biosynthetic pathway in higher eukaryotes is the mevalonate pathway, which leads to the synthesis of
15 cholesterol, bile acids, porphyrin, dolichol, ubiquinone, carotenoids, retinoids, vitamin D, steroid hormones and farnesylated proteins.

LXR α and LXR β

LXR α is found predominantly in the liver, with lower levels found in kidney, intestine, spleen and adrenal tissue (see, *e.g.*, Willy, *et al.* (1995) *Gene Dev.* 9(9):1033-1045). LXR β is ubiquitous in
20 mammals and was found in nearly all tissues examined. LXRs are activated by certain naturally occurring, oxidized derivatives of cholesterol (see, *e.g.*, Lehmann, *et al.* (1997) *J. Biol. Chem.* 272(6):3137-3140). LXR α is activated by oxysterol and promotes cholesterol metabolism (Peet *et al.* (1998) *Cell* 93:693-704). Thus, LXRs appear to play a role in, *e.g.*, cholesterol metabolism (see, *e.g.*, Janowski, *et al.* (1996) *Nature* 383:728-731).

Nuclear Receptors and Disease


25 Nuclear receptor activity has been implicated in a variety of diseases and disorders, including, but not limited to, hypercholesterolemia (see, *e.g.*, International Patent Application Publication No. WO 00/57915), osteoporosis and vitamin deficiency (see, *e.g.*, U.S. Patent No. 6,316,5103), hyperlipoproteinemia (see, *e.g.*, International Patent Application Publication No. WO 01/60818),
30 hypertriglyceridemia, lipodystrophy, hyperglycemia and diabetes mellitus (see, *e.g.*, International Patent Application Publication No. WO 01/82917), atherosclerosis and gallstones (see, *e.g.*, International Patent Application Publication No. WO 00/37077), disorders of the skin and mucous membranes (see, *e.g.*, U.S. Patent Nos. 6,184,215 and 6,187,814, and International Patent Application Publication No.

WO 98/32444), acne (see, *e.g.*, International Patent Application Publication No. WO 00/49992), and cancer, Parkinson's disease and Alzheimer's disease (see *e.g.*, International Patent Application Publication No. WO 00/17334). Activity of nuclear receptors, including LXRs, FXRs and PPARs, and orphan nuclear receptors, has been implicated in physiological processes including, but not limited to, bile acid biosynthesis, cholesterol metabolism or catabolism, and modulation of cholesterol 7 α -hydroxylase gene (CYP7A1) transcription (see, *e.g.*, Chiang et al. (2000) J. Biol. Chem. 275:10918-10924), HDL metabolism (see, *e.g.*, Urizar et al. (2000) J. Biol. Chem. 275:39313-39317 and International Patent Application Publication No. WO 01/03705), and increased cholesterol efflux and increased expression of ATP binding cassette transporter protein (ABC1) (see, *e.g.*, International Patent Application Publication No. WO 00/78972).


Thus, there is a need for compounds, compositions and methods of modulating the activity of nuclear receptors, including LXRs, FXRs, PPARs and orphan nuclear receptors. Such compounds are useful in the treatment, prevention, inhibition or amelioration of one or more symptoms of diseases or disorders in which nuclear receptor activity is implicated.

SUMMARY OF THE INVENTION


In one aspect, the present invention provides a compound according the following formulas IIa-d,




IIa



IIb



IIc



IIId

or a pharmaceutically acceptable salt, isomer, or prodrug thereof, which are useful as modulators of the activity of liver X receptors (LXR), where R¹, R², R²¹, R³, and G are defined herein.

Compounds for use in compositions and methods for modulating the activity of nuclear receptors are provided. In particular, compounds of the invention which are useful for modulating liver X receptors, LXR α and LXR β , FXR, PPAR and/or orphan nuclear receptors are provided.

In one embodiment, the compounds provided herein are agonists of LXR. In another embodiment, the compounds provided herein are antagonists of LXR. Agonists that exhibit low efficacy are, in certain embodiments, antagonists.

Another aspect of this invention is directed to methods of treating, preventing, inhibiting, or ameliorating the symptoms of a disease or disorder that is modulated or otherwise affected by nuclear receptor activity or in which nuclear receptor activity is implicated, comprising administering to a

subject in need thereof a therapeutically effective amount of a compound of formulae IIa, IIb, IIc, or IId, or a pharmaceutically acceptable derivative thereof.

Another aspect of this invention is directed to methods of reducing cholesterol levels in a subject in need thereof, comprising administering an effective cholesterol level-reducing amount of a compound of formulae IIa, IIb, IIc or IId, or a pharmaceutically acceptable derivative thereof.

Another aspect of this invention is directed to methods of treating, preventing, inhibiting, or ameliorating one or more symptoms of a disease or disorder which is affected by cholesterol, triglyceride, or bile acid levels, comprising administering to a subject in need thereof a therapeutically effective amount of a compound of formulae IIa, IIb, IIc or IId, or a pharmaceutically acceptable derivative thereof.

Another aspect of this invention is directed to methods of modulating nuclear receptor activity, comprising contacting the nuclear receptor with a compound of formulae IIa, IIb, IIc or IId, or a pharmaceutically acceptable derivative thereof.

Another aspect of this invention is directed to methods of modulating cholesterol metabolism, comprising administering an effective cholesterol metabolism-modulating amount of a compound of formulae IIa, IIb, IIc or IId, or a pharmaceutically acceptable derivative thereof.

Another aspect of this invention is directed to methods of treating, preventing, inhibiting or ameliorating one or more symptoms of hypocholesterolemia in a subject in need thereof, comprising administering a therapeutically effective amount of a compound of formulae IIa, IIb, IIc or IId, or a pharmaceutically acceptable derivative thereof.

Another aspect of this invention is directed to methods of increasing cholesterol efflux from cells of a subject, comprising administering an effective cholesterol efflux-increasing amount of a compound of formulae IIa, IIb, IIc or IId, or a pharmaceutically acceptable derivative thereof.

Another aspect of this invention is directed to methods of increasing the expression of ATP-Binding Cassette (ABC1) in the cells of a subject, comprising administering an effective ABC1 expression-increasing amount of a compound of formulae IIa, IIb, IIc or IId, or a pharmaceutically acceptable derivative thereof.

Another aspect of this invention is directed to in vitro methods for altering nuclear receptor activity, comprising contacting the nuclear receptor with a compound of formulae IIa, IIb, IIc or IId, or a pharmaceutically acceptable derivative thereof.

Another aspect of this invention is directed to methods of reducing cholesterol levels in a subject in need thereof, comprising administering an effective cholesterol level-reducing amount of a compound of formulae IIa, IIb, IIc or IId, or a pharmaceutically acceptable derivative thereof.


Another aspect of this invention is directed to pharmaceutical compositions comprising a pharmaceutically acceptable carrier, excipient and/or diluent and a compound of formulae IIa, IIb, IIc or IId.

Another aspect of this invention is directed to regulation of cholesterol transport and inflammatory signaling pathways that are implicated in human disease pathology including atherosclerosis and associated diseases such as myocardial infarction and ischemic stroke in a subject in need thereof, comprising administering an effective cholesterol transport and inflammatory signaling pathways regulating amount of a compound of formulae IIa, IIb, IIc or IId, or a pharmaceutically acceptable derivative thereof.


Another aspect of this invention is directed to treatment of the metabolic syndrome which comprises a constellation of disorders of the body's metabolism including obesity, hypertension and insulin resistance and diabetes including treatment of diseases resulting from compromised metabolism and immunity including atherosclerosis and diabetes as well as autoimmune disorders and diseases in a subject in need thereof, comprising administering a therapeutically effective amount of a compound of formulae IIa, IIb, IIc or IId, or a pharmaceutically acceptable derivative thereof.

DETAILED DESCRIPTION OF THE INVENTION


In a first embodiment, the present invention provides a compound according to one of the following formulas,




IIa



IIb



IIc



IId

or a pharmaceutically acceptable salt, isomer, or prodrug thereof, wherein,

(A) R¹ is -L¹-R⁵, wherein

L¹ is a bond, L⁵, L⁶, -L⁵-L⁶-L⁵-, or -L⁶-L⁵-L⁶-, wherein

each L⁵ is independently -[C(R¹⁵)₂]_m-, wherein

each R¹⁵ is independently hydrogen, halogen, (C₁-C₆)alkyl, (C₃-C₆)cycloalkyl, or (C₁-C₆)haloalkyl; and

each L⁶ is independently -CS-, -CO-, -SO₂-, -O-, -CON(R¹¹)-, -CONR¹¹N(R¹¹)-, -C(=NR¹¹)-, -C(=NOR¹¹)-, or -C(=NN(R¹¹)₂)-, -aryl-, -C₃-C₈cycloalkyl-, -heteroaryl-, or -heterocyclyl-,

wherein the aryl, cycloalkyl, heteroaryl, or heterocyclyl is optionally substituted with one or more R¹⁴;

or each L⁶ is independently C₂-C₆ aldiyl,

5 wherein the aldiyl chain is optionally interrupted by -C(R¹¹)₂-, -C(R¹¹)₂C(R¹¹)₂-, -C(R¹¹)=C(R¹¹)-, -C(R¹¹)₂O-, -C(R¹¹)₂NR¹¹-, -C≡C-, -O-, -S-, -N(R¹⁰)CO-, -N(R¹⁰)CO₂-, -CON(R¹⁰)-, -CO-, -CO₂-, -OC(=O)-, -OC(=O)N(R¹⁰)-, -SO₂-, -N(R¹⁰)SO₂-, or -SO₂N(R¹⁰); and

R⁵ is aryl, heterocyclyl, heteroaryl, -(C₃-C₆)cycloalkyl, -C, or -B-C, wherein

B is -[C(R¹⁵)₂]_m- or -C₃-C₈ cycloalkyl-; and

10 C is halogen, -C₁-C₆alkyl, -C₁-C₆haloalkyl, -SO₂R¹¹, -SR¹¹, -SO₂N(R¹¹)₂, -SO₂NR¹¹COR¹¹, -C≡N, -C(O)OR¹¹, -CON(R¹¹)₂, or -N(R¹¹)₂,

wherein R⁵ is optionally substituted with one or more R^{5a}, wherein

15 each R^{5a} is independently halogen, nitro, heteroaryl, heterocyclyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, (C₃-C₈ cycloalkyl)-C₁-C₆ alkyl-, (C₃-C₈ cycloalkenyl)-C₁-C₆ alkyl-, (C₃-C₈ cycloalkyl)-C₂-C₆ alkenyl-, aryl, arylalkyl, aryloxy, aryloxyaryl, arylC₁₋₆ alkoxy, C₁-C₆ alkyl, C₁-C₆ haloalkyl, SO₂R¹¹, OR¹¹, SR¹¹, N₃, SO₂R¹¹, COR¹¹, SO₂N(R¹¹)₂, SO₂NR¹¹COR¹¹, C≡N, C(O)OR¹¹, CON(R¹¹)₂, CON(R¹¹)OR¹¹, OCON(R¹¹)₂, NR¹¹COR¹¹, NR¹¹CON(R¹¹)₂, NR¹¹COOR¹¹, or N(R¹¹)₂, wherein

20 each R^{5a} is optionally substituted with one or more groups which independently are -halogen, -C₁-C₆ alkyl, aryloxy C₀₋₆ alkylSO₂R¹¹, C₀₋₆ alkylCOOR¹¹, C₀₋₆ alkoxyaryl, C₁-C₆ haloalkyl, -SO₂R¹¹, -OR¹¹, -SR¹¹, -N₃, -SO₂R¹¹, -COR¹¹, -SO₂N(R¹¹)₂, -SO₂NR¹¹COR¹¹, -C≡N, -C(O)OR¹¹, -CON(R¹¹)₂, -CON(R¹¹)OR¹¹, -OCON(R¹¹)₂, -NR¹¹COR¹¹, -NR¹¹CON(R¹¹)₂, -NR¹¹COOR¹¹, or -N(R¹¹)₂;

25 R² and R²¹ are -L³-R⁷, wherein

each L³ is independently a bond or -(CH₂)_m-V¹-(CH₂)_n-, wherein

n is 0-6; and

30 V¹ is -C(R¹¹)₂-, -C(R¹¹)₂C(R¹¹)₂-, -C(R¹¹)=C(R¹¹)-, -C(R¹¹)₂O-, -C(R¹¹)₂NR¹¹-, -C≡C-, -O-, -S-, -NR¹¹-, -N(R¹⁰)CO-, -N(R¹⁰)CO₂-, -OCO-, -CO-, -CS-, -CONR¹⁰-, -C(=N-R¹¹)-, -C(=N-OR¹¹)-, -C[=N-N(R¹¹)₂], -CO₂-, -OC(=O)-, -OC(=O)N(R¹⁰)-, -SO₂-, -N(R¹⁰)SO₂-, -SO₂N(R¹⁰)-, -NR¹⁰CONR¹⁰-, -NR¹⁰CSNR¹⁰-, C₃-C₆cyclo alkyl, or C₃-C₆ cyclohaloalkyl;

or each L³ is independently C₂-C₆ aldiyl,

wherein the alidiyl chain is optionally interrupted by $-C(R^{11})_2-$, $-C(R^{11})_2C(R^{11})_2-$, $-C(R^{11})=C(R^{11})-$, $-C(R^{11})_2O-$, $-C(R^{11})_2NR^{11}-$, $-C\equiv C-$, $-O-$, $-S-$, $-N(R^{10})CO-$, $-N(R^{10})CO_2-$, $-NR^{11}-$, $-CON(R^{10})-$, $-CO-$, $-CO_2-$, $-OC(=O)-$, $-OC(=O)N(R^{10})-$, $-SO_2-$, $-N(R^{10})SO_2-$, or $-SO_2N(R^{10})-$; and

5 each R^7 is independently hydrogen, halogen, nitro, aryl, heteroaryl, heterocyclyl, $-Z$, $-Y-Z$, or $-X-Y-Z$, wherein

X is $-O-$;

Y is $-[C(R^{15})_2]_m-$, C_2-C_6 alkenyl, or $-C_3-C_8$ cycloalkyl; and

10 Z is $-H$, halogen, $-OR^{11}$, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)N(R^{11})_2$, $-N(R^{11})_2$, $-C(=N-OH)R^{11}$, $-C(=S)N(R^{11})_2$, $-CN$, $-S(=O)_2N(R^{11})_2$, $-C(=O)N(R^{11})N(R^{11})_2$, $-C(=O)N(R^{11})(OR^{11})$, $-OC(=O)-R^{11}$, or $-OC(=O)-N(R^{11})_2$,

wherein R^7 is optionally substituted with one or more R^{7a} , wherein

R^{7a} is halogen, haloaryl, aryloxy, aralkyloxy, aryloxyalkyl, aryl C_0-C_6 alkylcarboxy, $C(R^{11})=C(R^{11})-COOH$, aryl, heteroaryl, heterocyclyl, heterocyclyloxy, heteroaryloxy, $-Z'$, $-Y'-Z'$, or $-X'-Y'-Z'$, wherein

15

X' is $-O-$;

Y' is $-[C(R^{15})_2]_m-$ or $-C_3-C_8$ cycloalkyl; and

20 Z' is $-H$, halogen, $-OR^{11}$, $-SR^{11}$, $-S(=O)_2R^{11}$, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)N(R^{11})_2$, $-N(R^{11})_2$, $-N(R^{11})C(=O)R^{11}$, $-S(=O)_2N(R^{11})C(=O)R^{11}$, $-CN$, $-S(=O)_2N(R^{11})_2$, $-C(=O)N(R^{11})N(R^{11})_2$, $-C(=O)N(R^{11})(OR^{11})$, $-OC(=O)-R^{11}$, $-OC(=O)-OR^{11}$, $-N(R^{11})C(=O)-R^{11}$, or $-N(R^{11})S(=O)_2R^{11}$,

20

wherein each R^{7a} is optionally substituted with one or more R^8 ,

wherein each R^8 is independently halogen, C_1-C_6 alkyl, C_1-C_6 alkoxy, C_1-C_6 haloalkyl, C_1-C_6 haloalkyl(OR^{11}), C_0-C_6 alkyl OR^{11} , C_0-C_6 alkyl $CON(R^{11})_2$, C_0-C_6 alkyl COR^{11} , C_0-C_6 alkyl $COOR^{11}$, or C_0-C_6 alkyl SO_2R^{11} ,

25

provided that R^2 and R^{21} are not simultaneously hydrogen,

R^3 is $-L-R^6$, wherein

L is a bond, $-X^3-(CH_2)_n-X^3-$, $-(CH_2)_m-X^3-(CH_2)_n-$ or $-(CH_2)_{1+w}-Y^3-(CH_2)_w-$, wherein

n is 0-6; each w is independently 0-5; and

30

each X^3 is independently a bond, $-C(R^{11})_2-$, $-C(R^{11})_2C(R^{11})_2-$, $-C(R^{11})=C(R^{11})-$, $-C\equiv C-$, $-CO-$, $-CS-$, $-CONR^{10}-$, $-C(=N)(R^{11})-$, $-C(=N-OR^{11})-$, $-C[=N-N(R^{11})_2]$, $-CO_2-$, $-SO_2-$, or $-SO_2N(R^{10})-$; and

Y^3 is $-O-$, $-S-$, $-NR^7-$, $-N(R^{10})CO-$, $-N(R^{10})CO_2-$, $-OCO-$, $-OC(=O)N(R^{10})-$,
 $-NR^{10}CONR^{10}-$, $-N(R^{10})SO_2-$, or $-NR^{10}CSNR^{10}-$;

or L is C_{2-6} aldiyl chain wherein the aldiyl chain is optionally interrupted by $-C(R^{11})_2-$,
 $-C(R^{11})_2C(R^{11})_2-$, $-C(R^{11})=C(R^{11})-$, $-C(R^{11})_2O-$, $-C(R^{11})_2NR^{11}-$, $-C\equiv C-$, $-O-$, $-S-$, $-N(R^{10})CO-$,
 $-N(R^{10})CO_2-$, $-CON(R^{10})-$, $-CO-$, $-CO_2-$, $-OC(=O)-$, $-OC(=O)N(R^{10})-$, $-SO_2-$, $-N(R^{10})SO_2-$, or
 $-SO_2N(R^{10})$; and

R^6 is C_1-C_6 alkyl, C_1-C_6 haloalkyl, aryl, C_3-C_8 cycloalkyl, heteroaryl, heterocyclyl, $-CN$, $-C(=O)R^{11}$,
 $-C(=O)OR^{11}$, $-C(=O)N(R^{11})_2$, $-N(R^{11})_2$, $-SO_2R^{11}$, $-S(=O)_2N(R^{11})_2$, $-C(=O)N(R^{11})N(R^{11})_2$,
 $-C(=O)N(R^{11})(OR^{11})$, wherein

the aryl, heteroaryl, cycloalkyl, or heterocyclyl is optionally substituted with with one or more
 R^{6a} , wherein

each R^{6a} is independently $-Z''$, $-Y''-Z''$, or $-X''-Y''-Z''$, wherein

X'' is $-O-$;


Y'' is $-[C(R^{15})_2]_m-$, $-C_2-C_6$ alkenyl, C_3-C_8 cycloalkyl, heterocyclyl, aryl, or heteroaryl,

wherein

the aryl, heteroaryl, cycloalkyl, or heterocyclyl is optionally substituted with at
least one group which is each independently Z'' ;

Z'' is $-H$, $-CN$, halogen, $-OR^{11}$, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)N(R^{11})_2$, $-N(R^{11})_2$, $-N_3$,
 $-SO_2R^{11}$, $-S(=O)_2N(R^{11})_2$, $-C(=O)N(R^{11})N(R^{11})_2$, $-N(R^{11})C(=O)N(R^{11})_2$,
 $-OC(=O)-OR^{11}$, $-C(=O)N(R^{11})(OR^{11})$, $-OC(=O)-R^{11}$, $-OC(=O)-N(R^{11})_2$, or
 $-N(R^{11})COOR^{11}$; and

G is a group of the formula,



J is aryl, heteroaryl, or absent;

K is aryl, heteroaryl, or absent;

provided that

(i) if K is absent, then q is 1 and R^4 is bonded directly to L^2 ; and

each R^4 and R^{41} is independently halogen, oxo, nitro, $CR^{11}=CR^{11}COOR^{11}$, aryloxy, aralkyloxy,
aryloxyalkyl, aryl C_0-C_6 alkylcarboxy, aryl, heteroaryl, heterocyclyl, heteroaryloxy,
heterocyclyloxy, $-G^1$, $-E-G^1$, or $-D-E-G^1$, wherein

D is $-O-$;

E is $-\text{[C(R}^{15})_2]_m-$ or $-\text{C}_3\text{-C}_8\text{cycloalkyl}$; and
 G^1 is $-\text{C}_1\text{-C}_6\text{alkyl}$, $-\text{C}_1\text{-C}_6\text{haloalkyl}$, $-\text{COR}^{11}$, $-\text{COOR}^{11}$, $-\text{CON(R}^{11})_2$, $-\text{C}\equiv\text{N}$, $-\text{OR}^{11}$,
 $-\text{OCON(R}^{11})_2$, $-\text{OCOOR}^{11}$, $-\text{N}_3$, $-\text{NR}^{11}\text{COR}^{11}$, $-\text{NR}^{11}\text{SO}_2\text{R}^{11}$, $-\text{N(R}^{11})_2$, $-\text{NR}^{11}\text{COOR}^{11}$,
 $-\text{SO}_2\text{R}^{11}$, $-\text{SO}_2\text{NR}^{11}\text{COR}^{11}$, $-\text{SO}_2\text{N(R}^{11})_2$, $-\text{SOR}^{11}$, or $-\text{SR}^{11}$,
 5 wherein each R^4 is optionally substituted with one or more R^{4a} ,
 wherein each R^{4a} is independently halogen, aryloxy, aralkyloxy, aryloxyalkyl,
 $\text{C}_1\text{-C}_6$ alkoxyaryl, aryl $\text{C}_0\text{-C}_6$ alkylcarboxy, $-\text{G}'$, $-\text{E}'\text{-G}'$, or $-\text{D}'\text{-E}'\text{-G}'$, wherein
 D' is $-\text{O}-$;
 E' is $-\text{[C(R}^{15})_2]_m-$ or $-\text{C}_3\text{-C}_8\text{cycloalkyl}$; and
 10 G' is $-\text{H}$, $-\text{halogen}$, $-\text{COR}^{11}$, $-\text{COOR}^{11}$, $-\text{C}\equiv\text{N}$, $-\text{OR}^{11}$, $-\text{NR}^{11}\text{SO}_2\text{R}^{11}$,
 $-\text{SO}_2\text{R}^{11}$, $-\text{SO}_2\text{N(R}^{11})_2$, or $-\text{SR}^{11}$,
 L^2 is a bond or $-\text{[C(R}^{15})_2]_m\text{-V}^2\text{-[C(R}^{15})_2]_n-$, wherein
 V^2 is independently $-\text{C(R}^{11})_2-$, $-\text{C(R}^{11})_2\text{C(R}^{11})_2-$, $-\text{C(R}^{11})=\text{C(R}^{11})-$, $-\text{C(R}^{11})_2\text{O}-$,
 $-\text{C(R}^{11})_2\text{NR}^{11}-$, $-\text{C}\equiv\text{C}-$, $-\text{O}-$, $-\text{S}-$, $-\text{SO}_2-$, $-\text{N(R}^{10})\text{CO}-$, $-\text{N(R}^{10})\text{CO}_2-$, $-\text{CON(R}^{10})-$,
 15 $-\text{CON(R}^{11})-$, $-\text{CON(R}^{11})\text{O}-$, $-\text{CO}-$, $-\text{CS}-$, $-\text{CO}_2-$, $-\text{OC(=O)-}$, $-\text{OC(=O)N(R}^{10})-$,
 $-\text{N(R}^{10})\text{SO}_2-$, $-\text{SO}_2\text{N(R}^{10})-$, $-\text{NR}^{10}\text{CONR}^{10}-$, $-\text{NR}^{10}\text{CSNR}^{10}-$, $\text{C}_3\text{-C}_6\text{cyclo alkyl-}$, or
 $\text{C}_3\text{-C}_6\text{cyclohaloalkyl}$,
 or V^2 is C_{2-6} alidiyl,
 wherein alidiyl chain is optionally interrupted by $-\text{C(R}^{11})_2-$, $-\text{C(R}^{11})_2\text{C(R}^{11})_2-$,
 20 $-\text{C(R}^{11})=\text{C(R}^{11})-$, $-\text{C(R}^{11})_2\text{O}-$, $-\text{C(R}^{11})_2\text{NR}^{11}-$, $-\text{C(R}^{11})_2\text{NR}^{11}-$, $-\text{C}\equiv\text{C}-$, $-\text{O}-$, $-\text{S}-$,
 $-\text{N(R}^{10})\text{CO}-$, $-\text{N(R}^{10})\text{CO}_2-$, $-\text{CON(R}^{10})-$, $-\text{CON(R}^{11})-$, $-\text{CON(R}^{11})\text{O}-$, $-\text{CO}-$, $-\text{CO}_2-$,
 $-\text{OC(=O)-}$, $-\text{OC(=O)N(R}^{10})-$, $-\text{SO}_2-$, $-\text{N(R}^{10})\text{SO}_2-$ or $-\text{SO}_2\text{N(R}^{10})-$;
 or V^2 is aryl, heteroaryl, or heterocyclyl, each of which is optionally substituted with
 one or more R^9 , wherein
 25 each R^9 is independently halogen, $\text{C}_1\text{-C}_6$ haloalkyl, $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_1\text{-C}_6$
 alkyloxy, $\text{C}_0\text{-C}_6$ alkyl or $\text{C}_1\text{-C}_6$ alkyl COOR^{11} ;
 each m is 0, 1, 2, 3, 4, 5, or 6;
 q is 0, 1, 2, 3, 4, or 5,
 provided that q is 0 if and only if K is not phenyl; and
 30 q' is 0, 1, 2, 3, or 4,
 each R^{10} is independently $-\text{R}^{11}$, $-\text{C(=O)R}^{11}$, $-\text{CO}_2\text{R}^{11}$, or $-\text{SO}_2\text{R}^{11}$;
 each R^{11} is independently $-\text{hydrogen}$, $-\text{C}_1\text{-C}_6$ alkyl, $\text{C}_2\text{-C}_6$ alkenyl, $\text{C}_2\text{-C}_6$ alkynyl, $\text{C}_3\text{-C}_8$ cycloalkyl, $(\text{C}_3\text{-C}_8$
 cycloalkyl)- $\text{C}_1\text{-C}_6$ alkyl-, $(\text{C}_3\text{-C}_8$ cycloalkenyl)- $\text{C}_1\text{-C}_6$ alkyl-, $(\text{C}_3\text{-C}_8$ cycloalkyl)- $\text{C}_2\text{-C}_6$ alkenyl-, $-\text{C}_1\text{-C}_6$ haloalkyl,

-N(R¹²)₂, aryl, -(C₁-C₆)alkyl-aryl, heteroaryl, -(C₁-C₆)alkyl-heteroaryl, heterocyclyl, or -(C₁-C₆)alkyl-heterocyclyl,

wherein any of R¹¹ is optionally substituted with one or more radicals of R¹²;

5 each R¹² is independently halogen, C₁-C₆haloalkyl, C₁-C₆ alkyl, C₁-C₆ alkoxy, (C₀-C₆ alkyl)C=O(OR¹³); C₀-C₆ alkylOR¹³, C₀-C₆ alkylCOR¹³, C₀-C₆ alkylSO₂R¹³, C₀-C₆ alkylCON(R¹³)₂, C₀-C₆ alkylCONR¹³OR¹³, C₀-C₆ alkylSO₂N(R¹³)₂, C₀-C₆ alkylSR¹³, C₀-C₆ haloalkylOR¹³, aryloxy, aralkyloxy, aryloxyalkyl, C₀-C₆alkoxyaryl, arylC₀-C₆ alkylcarboxy, -C₀-C₆ alkylN(R¹³)₂, -NR¹³SO₂R¹³, or -OC₀-C₆ alkylCOOR¹³;

10 each R¹³ is independently hydrogen, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, (C₃-C₈ cycloalkyl)-C₁-C₆ alkyl-, (C₃-C₈ cycloalkenyl)-C₁-C₆ alkyl-, or (C₃-C₈ cycloalkyl)-C₂-C₆ alkenyl-;

each R¹⁴ is independently C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen, C₁-C₆ haloalkyl, C₀-C₆ alkylCON(R¹¹)₂, C₀-C₆ alkylCONR¹¹OR¹¹, C₀-C₆ alkylOR¹¹, or C₀-C₆ alkylCOOR¹¹; and

(B) provided that

(i) when L² is a bond, both J and K are not absent;

15 (ii) if the compound is defined by formula IIa, then

a. if J is phenyl and K is thienyl, furyl, or thiazoyl and q is 0, then R¹ is not 4-(NH₂SO₂)phenyl, 4-(NH₂SO₂)-3-fluorophenyl, p-(CH₃SO₂)phenyl-, or 4p-(CH₃SO₂)-3-fluorophenyl-; and

20 b. if R⁵ is pyridyl or phenyl optionally substituted with one or more R^{5a} and L¹ is a bond, then G is not p-(NH₂SO₂)phenyl or p-(CH₃SO₂)phenyl-;

(iii) if the compound is defined by formula IIc or IId, then G is not p-(NH₂SO₂)phenyl or p-(CH₃SO₂)phenyl-;

(iv) the compound is not 1-(biphenyl-4-yl)-2,5-diphenyl-1H-imidazole.

In one embodiment, the invention provides the compound according to formula IIa, IIb, IIc, or

25 IId, wherein:

R¹ is -L¹-R⁵, wherein

L¹ is a bond, L⁵, L⁶, -L⁵-L⁶-L⁵-, or -L⁶-L⁵-L⁶-, wherein

each L⁵ is independently -[C(R¹⁵)₂]_m-, wherein

m is 0, 1, 2, 3, or 4; and

30 each R¹⁵ is independently hydrogen, halogen, (C₁-C₆)alkyl, or (C₁-C₆)haloalkyl;

and

L⁶ is -CO-, -SO₂-, -O-, -CON(R¹¹)-, -C₃-C₆cycloalkyl-, or -heterocyclyl-,

wherein the cycloalkyl, or heterocyclyl is optionally substituted with one or more R¹⁴; and

R⁵ is aryl, heterocyclyl, heteroaryl, -C, or -B-C, wherein

B is -[C(R¹⁵)₂]_m- or -C₃-C₆cycloalkyl-; and

C is halogen, -C₁-C₆alkyl, or -C₁-C₆haloalkyl;

wherein R⁵ is optionally substituted with one or more R^{5a}, wherein

5 each R^{5a} is independently halogen, nitro, heteroaryl, heterocyclyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, (C₃-C₈ cycloalkyl)-C₁-C₆ alkyl-, (C₃-C₈ cycloalkenyl)-C₁-C₆ alkyl-, (C₃-C₈ cycloalkyl)-C₂-C₆ alkenyl-; aryl, arylalkyl, aryloxy, aryloxyaryl, arylC₁₋₆ alkoxy, C₁-C₆ alkyl, C₁-C₆ haloalkyl, C₃-C₆cycloalkyl, SO₂R¹¹, OR¹¹, SR¹¹, N₃, SO₂R¹¹, COR¹¹, SO₂N(R¹¹)₂, SO₂NR¹¹COR¹¹, C≡N, C(O)OR¹¹, CON(R¹¹)₂, CON(R¹¹)OR¹¹, OCON(R¹¹)₂,
10 NR¹¹COR¹¹, NR¹¹CON(R¹¹)₂, NR¹¹COOR¹¹, or N(R¹¹)₂, wherein

each R^{5a} is optionally substituted with one or more groups which independently are -halogen, -C₁-C₆ alkyl, aryloxy C₀₋₆ alkylSO₂R¹¹, C₀₋₆ alkylCOOR¹¹, C₀₋₆ alkoxyaryl, -C₁-C₆ haloalkyl, -SO₂R¹¹, -OR¹¹, -SR¹¹, -N₃, -SO₂R¹¹, -COR¹¹, -SO₂N(R¹¹)₂, -SO₂NR¹¹COR¹¹, -C≡N, -C(O)OR¹¹,
15 -CON(R¹¹)₂, -CON(R¹¹)OR¹¹, -OCON(R¹¹)₂, -NR¹¹COR¹¹, -NR¹¹CON(R¹¹)₂, -NR¹¹COOR¹¹, or -N(R¹¹)₂;

R² is -L³-R⁷, wherein L³ is a bond; and

R⁷ is halogen, aryl, heteroaryl, heterocyclyl, -Z, or -Y-Z, wherein

Y is -[C(R¹⁵)₂]_m- or -C₃-C₆cycloalkyl; and

20 Z is -H, halogen, -OR¹¹, -C(=O)R¹¹, -C(=O)OR¹¹, -C(=O)N(R¹¹)₂, -N(R¹¹)₂, -C(=N-OH)R¹¹, -C(=S)N(R¹¹)₂, -CN, -S(=O)₂N(R¹¹)₂, -C(=O)N(R¹¹)N(R¹¹)₂, or -C(=O)N(R¹¹)(OR¹¹);

wherein R⁷ is optionally substituted with one or more R^{7a}, wherein

R^{7a} is halogen -Z', -Y'-Z', or -X'-Y'-Z', wherein


25 X' is -O-;

Y' is -[C(R¹⁵)₂]_m- or -C₃-C₆cycloalkyl; and

Z' is -H, halogen, -OR¹¹, -SR¹¹, -S(=O)₂R¹¹, -C(=O)R¹¹, -C(=O)OR¹¹, -C(=O)N(R¹¹)₂, -N(R¹¹)₂, -N(R¹¹)C(=O)R¹¹, -CN, -S(=O)₂N(R¹¹)₂, -C(=O)N(R¹¹)(OR¹¹), or -N(R¹¹)S(O)₂R¹¹;

30 R²¹ and R³ are each independently hydrogen, halogen, C₁-C₆alkyl, or C₁-C₆haloalkyl; and

G is a group of the formula,



J is aryl or heteroaryl;

K is aryl or heteroaryl;

each R^4 and R^{41} is independently halogen, aryloxy, aralkyloxy, aryloxyalkyl, aryl- $\text{C}_0\text{-C}_6$ alkylcarboxy, aryl, heteroaryl, heterocyclyl, heteroaryloxy, heterocyclyloxy, $-\text{G}^1$, $-\text{E}-\text{G}^1$, or $-\text{D}-\text{E}-\text{G}^1$, wherein

D is $-\text{O}-$;

E is $-\text{C}(\text{R}^{15})_2\text{---}$ or $-\text{C}_3\text{-C}_6\text{cycloalkyl}$; and

G^1 is $-\text{C}_1\text{-C}_6\text{alkyl}$, $-\text{C}_1\text{-C}_6\text{haloalkyl}$, $-\text{COR}^{11}$, $-\text{COOR}^{11}$, $-\text{CON}(\text{R}^{11})_2$, $-\text{C}\equiv\text{N}$, $-\text{OR}^{11}$, $-\text{OCON}(\text{R}^{11})_2$, $-\text{NR}^{11}\text{COR}^{11}$, $-\text{NR}^{11}\text{SO}_2\text{R}^{11}$, $-\text{N}(\text{R}^{11})_2$, $-\text{NR}^{11}\text{COOR}^{11}$, $-\text{SOR}^{11}$, $-\text{SO}_2\text{R}^{11}$, $-\text{SO}_2\text{NR}^{11}\text{COR}^{11}$, $-\text{SO}_2\text{N}(\text{R}^{11})_2$, or $-\text{SR}^{11}$,

L^2 is a bond;

q is 1, 2, or 3; and

q' is 0, 1, 2, or 3;

each R^{10} is independently $-\text{R}^{11}$, $-\text{C}(=\text{O})\text{R}^{11}$, $-\text{CO}_2\text{R}^{11}$, or $-\text{SO}_2\text{R}^{11}$;

each R^{11} is independently $-\text{hydrogen}$, $-\text{C}_1\text{-C}_6\text{ alkyl}$, $\text{C}_2\text{-C}_6\text{ alkenyl}$, $\text{C}_2\text{-C}_6\text{ alkynyl}$, $(\text{C}_3\text{-C}_8\text{ cycloalkyl})\text{-C}_1\text{-C}_6\text{ alkyl}$, $(\text{C}_3\text{-C}_8\text{ cycloalkenyl})\text{-C}_1\text{-C}_6\text{ alkyl}$, or $(\text{C}_3\text{-C}_8\text{ cycloalkyl})\text{-C}_2\text{-C}_6\text{ alkenyl}$; $\text{C}_1\text{-C}_6\text{ alkyl}$, $\text{C}_2\text{-C}_6\text{ alkenyl}$, $\text{C}_2\text{-C}_6\text{ alkynyl}$, $(\text{C}_3\text{-C}_8\text{ cycloalkyl})\text{-C}_1\text{-C}_6\text{ alkyl}$, $(\text{C}_3\text{-C}_8\text{ cycloalkenyl})\text{-C}_1\text{-C}_6\text{ alkyl}$, $(\text{C}_3\text{-C}_8\text{ cycloalkyl})\text{-C}_2\text{-C}_6\text{ alkenyl}$, $-\text{C}_3\text{-C}_8\text{ cycloalkyl}$, $-\text{C}_1\text{-C}_6\text{ haloalkyl}$, $-\text{N}(\text{R}^{12})_2$, aryl, $-(\text{C}_1\text{-C}_6)\text{alkyl-aryl}$, heteroaryl, $-(\text{C}_1\text{-C}_6)\text{alkyl-heteroaryl}$, heterocyclyl, or $-(\text{C}_1\text{-C}_6)\text{alkyl-heterocyclyl}$,

wherein any of R^{11} is optionally substituted with one or more radicals of R^{12} ;

each R^{12} is independently halogen, $\text{C}_0\text{-C}_6\text{alkylN}(\text{R}^{13})_2$, $\text{C}_1\text{-C}_6\text{haloalkyl}$, $\text{C}_1\text{-C}_6\text{ alkyl}$, $\text{C}_1\text{-C}_6\text{ alkoxy}$, $(\text{C}_0\text{-C}_6\text{ alkyl})\text{C}=\text{O}(\text{OR}^{13})$, $\text{C}_0\text{-C}_6\text{ alkylOR}^{13}$, $\text{C}_0\text{-C}_6\text{ alkylCOR}^{13}$, $\text{C}_0\text{-C}_6\text{ alkylSO}_2\text{R}^{13}$, $\text{C}_0\text{-C}_6\text{ alkylCON}(\text{R}^{13})_2$, $\text{C}_0\text{-C}_6\text{ alkylCONR}^{13}\text{OR}^{13}$, $\text{C}_0\text{-C}_6\text{ alkylSO}_2\text{N}(\text{R}^{13})_2$, $\text{C}_0\text{-C}_6\text{ alkylSR}^{13}$, $\text{C}_0\text{-C}_6\text{ haloalkylOR}^{13}$, aryloxy, aralkyloxy, aryloxyalkyl, $\text{C}_0\text{-C}_6\text{alkoxyaryl}$, aryl- $\text{C}_0\text{-C}_6\text{ alkylcarboxy}$, $\text{C}_0\text{-C}_6\text{ alkyl}$, $-\text{NR}^{13}\text{SO}_2\text{R}^{13}$, or $-\text{OC}_0\text{-C}_6\text{ alkylCOOR}^{13}$;

each R^{13} is independently hydrogen, $\text{C}_1\text{-C}_6\text{ alkyl}$, $\text{C}_2\text{-C}_6\text{ alkenyl}$, $\text{C}_2\text{-C}_6\text{ alkynyl}$, $(\text{C}_3\text{-C}_8\text{ cycloalkyl})\text{-C}_1\text{-C}_6\text{ alkyl}$, $(\text{C}_3\text{-C}_8\text{ cycloalkenyl})\text{-C}_1\text{-C}_6\text{ alkyl}$, or $(\text{C}_3\text{-C}_8\text{ cycloalkyl})\text{-C}_2\text{-C}_6\text{ alkenyl}$; and

each R^{14} is independently $\text{C}_1\text{-C}_6\text{ alkyl}$, $\text{C}_1\text{-C}_6\text{ alkoxy}$, halogen, $\text{C}_1\text{-C}_6\text{ haloalkyl}$, $\text{C}_0\text{-C}_6\text{ alkylCON}(\text{R}^{11})_2$, $\text{C}_0\text{-C}_6\text{ alkylCONR}^{11}\text{OR}^{11}$, $\text{C}_0\text{-C}_6\text{ alkylOR}^{11}$, or $\text{C}_0\text{-C}_6\text{ alkylCOOR}^{11}$.

In one embodiment, the invention provides the compound according to formula IIa, IIb, IIc, or IId, wherein:

R^1 is $-L^1-R^5$, wherein

L^1 is a bond, $-C_3-C_8$ cycloalkyl- or L^5 , wherein

5 each L^5 is independently $-[C(R^{15})_2]_m$, wherein

m is 0, 1, 2, or 3; and

each R^{15} is independently hydrogen, halogen, (C_1-C_6) alkyl, or (C_1-C_6) haloalkyl; and

R^5 is aryl, heterocyclyl, heteroaryl, $-C$, or $-B-C$, wherein

B is $-[C(R^{15})_2]_m$ or $-C_3-C_6$ cycloalkyl-; and

10 C is $-C_1-C_6$ alkyl or $-C_1-C_6$ haloalkyl;

wherein R^5 is optionally substituted with one or more R^{5a} , wherein

each R^{5a} is independently halogen, nitro, heteroaryl, heterocyclyl, C_2-C_6 alkenyl, C_2-C_6 alkynyl, $(C_3-C_8$ cycloalkyl)- C_1-C_6 alkyl-, $(C_3-C_8$ cycloalkenyl)- C_1-C_6 alkyl-, $(C_3-C_8$ cycloalkyl)- C_2-C_6 alkenyl-, aryl, arylalkyl, aryloxy, aryloxyaryl, aryl C_{1-6} alkoxy, C_1-C_6 alkyl, C_1-C_6 haloalkyl, C_3-C_6 cycloalkyl, SO_2R^{11} , OR^{11} , SR^{11} , N_3 , SO_2R^{11} , COR^{11} , $SO_2N(R^{11})_2$, $SO_2NR^{11}COR^{11}$, $C\equiv N$, $C(O)OR^{11}$, $CON(R^{11})_2$, $CON(R^{11})OR^{11}$, $OCON(R^{11})_2$, $NR^{11}COR^{11}$, $NR^{11}CON(R^{11})_2$, $NR^{11}COOR^{11}$, or $N(R^{11})_2$, wherein

each R^{5a} is optionally substituted with one or more groups which independently are -halogen, $-C_1-C_6$ alkyl, aryloxy, C_{0-6} alkyl SO_2R^{11} , C_{0-6} alkyl $COOR^{11}$, C_{0-6} alkoxyaryl, $-C_1-C_6$ haloalkyl, $-SO_2R^{11}$, $-OR^{11}$, $-SR^{11}$, $-N_3$, $-SO_2R^{11}$, $-COR^{11}$, $SO_2N(R^{11})_2$, $-SO_2NR^{11}COR^{11}$, $-C\equiv N$, $-C(O)OR^{11}$, $-CON(R^{11})_2$, $CON(R^{11})OR^{11}$, $-OCON(R^{11})_2$, $-NR^{11}COR^{11}$, $-NR^{11}CON(R^{11})_2$, $-NR^{11}COOR^{11}$, or $-N(R^{11})_2$;

R^2 is $-L^3-R^7$, wherein L^3 is a bond; and


25 R^7 is $-Z$ or $-Y-Z$, wherein

Y is $-[C(R^{15})_2]_m$ or $-C_3-C_6$ cycloalkyl; and

Z is $-H$, halogen, $-OR^{11}$, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)N(R^{11})_2$, $-N(R^{11})_2$, $-C(=N-OH)R^{11}$, $-C(=S)N(R^{11})_2$, $-CN$, $-S(=O)_2N(R^{11})_2$, $-OC(=O)-R^{11}$, or $-OC(=O)-N(R^{11})_2$;

30 R^{21} and R^3 are each independently hydrogen, halogen, C_1-C_6 alkyl, or C_1-C_6 haloalkyl; and

G is a group of the formula,



J is aryl or heteroaryl;

K is aryl or heteroaryl;

each R^4 and R^{41} is independently halogen, heteroaryl, heterocyclyl, $-G^1$, $-E-G^1$, or $-D-E-G^1$,
 5 wherein

D is $-O-$;

E is $-[C(R^{15})_2]_m-$ or $-C_3-C_6$ cycloalkyl; and

G^1 is $-C_1-C_6$ alkyl, $-C_1-C_6$ haloalkyl, $-COR^{11}$, $-COOR^{11}$, $-CON(R^{11})_2$, $-C\equiv N$, $-OR^{11}$, $-SOR^{11}$, $-SO_2R^{11}$, $-SO_2N(R^{11})_2$, or $-SR^{11}$;

10 L^2 is a bond;

q is 1, 2, or 3; and

q' is 0, 1, 2 or 3;

each R^{10} is independently $-R^{11}$, $-C(=O)R^{11}$, $-CO_2R^{11}$, or $-SO_2R^{11}$;

each R^{11} is independently $-hydrogen$, $-C_1-C_6$ alkyl, C_2-C_6 alkenyl, C_2-C_6 alkynyl, $(C_3-C_8$ cycloalkyl)- C_1-C_6
 15 alkyl-, $(C_3-C_8$ cycloalkenyl)- C_1-C_6 alkyl-, $(C_3-C_8$ cycloalkyl)- C_2-C_6 alkenyl-, $-C_3-C_8$ cycloalkyl,
 $-(C_1-C_6)alkyl-(C_3-C_8)cycloalkyl$, $-C_1-C_6$ haloalkyl, $-N(R^{12})_2$, aryl, $-(C_1-C_6)alkyl-aryl$, heteroaryl,
 $-(C_1-C_6)alkyl-heteroaryl$, heterocyclyl, or $-(C_1-C_6)alkyl-heterocyclyl$,

wherein any of R^{11} is optionally substituted with one or more radicals of R^{12} ;

each R^{12} is independently halogen, OR^{13} , $N(R^{13})_2$, C_1-C_6 haloalkyl, C_1-C_6 alkyl, C_1-C_6
 20 alkoxy, $(C_0-C_6$ alkyl) $C=O(OR^{13})$, C_0-C_6 alkyl OR^{13} , C_0-C_6 alkyl COR^{13} , C_0-C_6
 alkyl SO_2R^{13} , C_0-C_6 alkyl $CON(R^{13})_2$, C_0-C_6 alkyl $CONR^{13}OR^{13}$, C_0-C_6
 alkyl $SO_2N(R^{13})_2$, C_0-C_6 alkyl SR^{13} , C_0-C_6 haloalkyl OR^{13} , aryloxy, aralkyloxy,
 aryloxyalkyl, C_0-C_6 alkoxyaryl, aryl C_0-C_6 alkylcarboxy, C_0-C_6 alkyl, $-NR^{13}SO_2R^{13}$, or
 $-OC_0-C_6$ alkyl $COOR^{13}$;

each R^{13} is independently hydrogen, C_1-C_6 alkyl, C_2-C_6 alkenyl, C_2-C_6 alkynyl, $(C_3-C_8$ cycloalkyl)- C_1-C_6 alkyl-,
 25 $(C_3-C_8$ cycloalkenyl)- C_1-C_6 alkyl-, or $(C_3-C_8$ cycloalkyl)- C_2-C_6 alkenyl-;

each R^{14} is independently C_1-C_6 alkyl, C_1-C_6 alkoxy, halogen, C_1-C_6 haloalkyl, C_0-C_6 alkyl $CON(R^{11})_2$,
 C_0-C_6 alkyl $CONR^{11}OR^{11}$, C_0-C_6 alkyl OR^{11} , or C_0-C_6 alkyl $COOR^{11}$.

In one embodiment, the invention provides the compound according to formula IIa or IIb,

30 wherein:

R^1 is $-L^1-R^5$, wherein

L^1 is a bond, L^5 , L^6 , $-L^5-L^6-L^5$, or $-L^6-L^5-L^6$, wherein

each L^5 is independently $-[C(R^{15})_2]_m$, wherein

m is 0, 1, 2, 3, or 4; and

each R^{15} is independently hydrogen, halogen, (C_1-C_6) alkyl, or (C_1-C_6) haloalkyl; and

5 L^6 is $-CO-$, $-SO_2-$, $-O-$, $-CON(R^{11})-$, $-C_3-C_6$ cycloalkyl-, or -heterocyclyl-,

wherein the cycloalkyl, or heterocyclyl is optionally substituted with one or more R^{14} ; and

R^5 is aryl, heterocyclyl, heteroaryl, $-C$, or $-B-C$, wherein

B is $-[C(R^{15})_2]_m$ - or $-C_3-C_6$ cycloalkyl-; and

10 C is halogen, $-C_1-C_6$ alkyl, or $-C_1-C_6$ haloalkyl;

wherein R^5 is optionally substituted with one or more R^{5a} , wherein

each R^{5a} is independently halogen, nitro, heteroaryl, heterocyclyl, C_2-C_6 alkenyl, C_2-C_6 alkynyl, $(C_3-C_8$ cycloalkyl)- C_1-C_6 alkyl-, $(C_3-C_8$ cycloalkenyl)- C_1-C_6 alkyl-, $(C_3-C_8$ cycloalkyl)- C_2-C_6 alkenyl-, aryl, arylalkyl, aryloxy, aryloxyaryl, aryl C_{1-6} alkoxy, C_1-C_6 alkyl, C_1-C_6 haloalkyl, C_3-C_6 cycloalkyl, SO_2R^{11} , OR^{11} , SR^{11} , N_3 , SO_2R^{11} , COR^{11} , $SO_2N(R^{11})_2$, $SO_2NR^{11}COR^{11}$, $C\equiv N$, $C(O)OR^{11}$, $CON(R^{11})_2$, $CON(R^{11})OR^{11}$, $OCON(R^{11})_2$, $NR^{11}COR^{11}$, $NR^{11}CON(R^{11})_2$, $NR^{11}COOR^{11}$, or $N(R^{11})_2$, wherein

each R^{5a} is optionally substituted with one or more groups which independently are -halogen, $-C_1-C_6$ alkyl, aryloxy C_{0-6} alkyl SO_2R^{11} , C_{0-6} alkyl $COOR^{11}$, C_{0-6} alkoxyaryl, $-C_1-C_6$ haloalkyl, $-SO_2R^{11}$, $-OR^{11}$, $-SR^{11}$, $-N_3$, $-SO_2R^{11}$, $-COR^{11}$, $-SO_2N(R^{11})_2$, $-SO_2NR^{11}COR^{11}$, $-C\equiv N$, $-C(O)OR^{11}$, $-CON(R^{11})_2$, $CON(R^{11})OR^{11}$, $-OCON(R^{11})_2$, $-NR^{11}COR^{11}$, $-NR^{11}CON(R^{11})_2$, $-NR^{11}COOR^{11}$, or $-N(R^{11})_2$;

R^2 is $-L^3-R^7$, wherein

25 L^3 is a bond; and

R^7 is halogen, aryl, heteroaryl, heterocyclyl, $-Z$, or $-Y-Z$, wherein

Y is $-[C(R^{15})_2]_m$ - or $-C_3-C_6$ cycloalkyl-; and

Z is $-H$, halogen, $-OR^{11}$, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)N(R^{11})_2$, $-N(R^{11})_2$, $-C(=N-OH)R^{11}$, $-C(=S)N(R^{11})_2$, $-CN$, $-S(=O)_2N(R^{11})_2$, $-C(=O)N(R^{11})N(R^{11})_2$, or $-C(=O)N(R^{11})(OR^{11})$;

30 wherein R^7 is optionally substituted with one or more R^{7a} , wherein

R^{7a} is halogen, $-Z'$, $-Y'-Z'$, or $-X'-Y'-Z'$, wherein


X' is $-O-$;

Y' is $-\text{[C(R}^{15})_2]_m-$ or $-\text{C}_3\text{-C}_6\text{cycloalkyl}$; and

Z' is $-\text{H}$, halogen, $-\text{OR}^{11}$, $-\text{SR}^{11}$, $-\text{S(=O)}_2\text{R}^{11}$, $-\text{C(=O)R}^{11}$, $-\text{C(=O)OR}^{11}$,
 $-\text{C(=O)N(R}^{11})_2$, $-\text{N(R}^{11})_2$, $-\text{N(R}^{11})\text{C(=O)R}^{11}$, $-\text{CN}$, $-\text{S(=O)}_2\text{N(R}^{11})_2$,
 $-\text{C(=O)N(R}^{11})(\text{OR}^{11})$, or $-\text{N(R}^{11})\text{S(=O)}_2\text{R}^{11}$;

5 R^{21} is hydrogen, halogen, $\text{C}_1\text{-C}_6\text{alkyl}$, or $\text{C}_1\text{-C}_6\text{haloalkyl}$; and

G is a group of the formula,



J is aryl or heteroaryl;

K is aryl or heteroaryl;

10 each R^4 and R^{41} is independently halogen, aryloxy, aralkyloxy, aryloxyalkyl, aryl $\text{C}_0\text{-C}_6$
 alkylcarboxy, aryl, heteroaryl, heterocyclyl, heteroaryloxy, heterocyclyloxy, $-\text{G}^1$, $-\text{E-G}^1$, or $-\text{D-E-G}^1$, wherein

D is $-\text{O}-$;

E is $-\text{[C(R}^{15})_2]_m-$ or $-\text{C}_3\text{-C}_6\text{cycloalkyl}$; and

15 G^1 is $-\text{C}_1\text{-C}_6\text{alkyl}$, $-\text{C}_1\text{-C}_6\text{haloalkyl}$, $-\text{COR}^{11}$, $-\text{COOR}^{11}$, $-\text{CON(R}^{11})_2$, $-\text{C}\equiv\text{N}$, $-\text{OR}^{11}$,
 $-\text{OCON(R}^{11})_2$, $-\text{NR}^{11}\text{COR}^{11}$, $-\text{NR}^{11}\text{SO}_2\text{R}^{11}$, $-\text{N(R}^{11})_2$, $-\text{NR}^{11}\text{COOR}^{11}$, $-\text{SOR}^{11}$, $-\text{SO}_2\text{R}^{11}$,
 $-\text{SO}_2\text{NR}^{11}\text{COR}^{11}$, $-\text{SO}_2\text{N(R}^{11})_2$, or $-\text{SR}^{11}$,

L^2 is a bond;

q is 1, 2, or 3; and

20 q' is 0, 1, 2, or 3;

each R^{10} is independently $-\text{R}^{11}$, $-\text{C(=O)R}^{11}$, $-\text{CO}_2\text{R}^{11}$, or $-\text{SO}_2\text{R}^{11}$;

each R^{11} is independently $-\text{hydrogen}$, $-\text{C}_1\text{-C}_6\text{ alkyl}$, $\text{C}_2\text{-C}_6\text{ alkenyl}$, $\text{C}_2\text{-C}_6\text{ alkynyl}$, $(\text{C}_3\text{-C}_8\text{ cycloalkyl})\text{-C}_1\text{-C}_6$
 alkyl-, $(\text{C}_3\text{-C}_8\text{ cycloalkenyl})\text{-C}_1\text{-C}_6\text{ alkyl-}$, $(\text{C}_3\text{-C}_8\text{ cycloalkyl})\text{-C}_2\text{-C}_6\text{ alkenyl-}$, $-\text{C}_3\text{-C}_8\text{ cycloalkyl}$, $-\text{C}_1\text{-C}_6$
 haloalkyl, $-\text{N(R}^{12})_2$, aryl, $-(\text{C}_1\text{-C}_6)\text{alkyl-aryl}$, heteroaryl, $-(\text{C}_1\text{-C}_6)\text{alkyl-heteroaryl}$, heterocyclyl, or
 25 $-(\text{C}_1\text{-C}_6)\text{alkyl-heterocyclyl}$,

wherein any of R^{11} is optionally substituted with one or more radicals of R^{12} ;

each R^{12} is independently halogen, $\text{C}_0\text{-C}_6\text{alkylN(R}^{13})_2$, $\text{C}_1\text{-C}_6\text{haloalkyl}$, $\text{C}_1\text{-C}_6\text{ alkyl}$,
 $\text{C}_1\text{-C}_6\text{ alkoxy}$, $(\text{C}_0\text{-C}_6\text{ alkyl})\text{C=O(OR}^{13})$, $\text{C}_0\text{-C}_6\text{ alkylOR}^{13}$, $\text{C}_0\text{-C}_6\text{ alkylCOR}^{13}$, $\text{C}_0\text{-C}_6$
 alkyl SO_2R^{13} , $\text{C}_0\text{-C}_6\text{ alkylCON(R}^{13})_2$, $\text{C}_0\text{-C}_6\text{ alkylCONR}^{13}\text{OR}^{13}$, $\text{C}_0\text{-C}_6$
 30 alkyl $\text{SO}_2\text{N(R}^{13})_2$, $\text{C}_0\text{-C}_6\text{ alkylSR}^{13}$, $\text{C}_0\text{-C}_6\text{ haloalkylOR}^{13}$, aryloxy, aralkyloxy,

aryloxyalkyl, C₀₋₆alkoxyaryl, arylC₀₋₆ alkylcarboxy, C₀₋₆ alkyl, -NR¹³SO₂ R¹³, or -OC₀₋₆ alkylCOOR¹³;

each R¹³ is independently hydrogen C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, (C₃₋₈ cycloalkyl)-C₁₋₆ alkyl-, (C₃₋₈ cycloalkenyl)-C₁₋₆ alkyl-, or (C₃₋₈ cycloalkyl)-C₂₋₆ alkenyl-; and

5 each R¹⁴ is independently C₁₋₆ alkyl, C₁₋₆ alkoxy, halogen, C₁₋₆ haloalkyl, C₀₋₆ alkylCON(R¹¹)₂, C₀₋₆ alkylCONR¹¹OR¹¹, C₀₋₆ alkylOR¹¹, or C₀₋₆ alkylCOOR¹¹.

In one embodiment, the invention provides the compound according to formula IIa or IIb, wherein:

R¹ is -L¹-R⁵, wherein

10 L¹ is a bond, -C₃₋₈cycloalkyl-, or L⁵, wherein

each L⁵ is independently -[C(R¹⁵)₂]_m-, wherein

m is 0, 1, 2, or 3; and

each R¹⁵ is independently hydrogen, halogen, (C₁₋₆)alkyl, or (C₁₋₆)haloalkyl; and

R⁵ is aryl, heterocyclyl, heteroaryl, -C, or -B-C, wherein

15 B is -[C(R¹⁵)₂]_m- or -C₃₋₆cycloalkyl-; and

C is -C₁₋₆alkyl or -C₁₋₆haloalkyl;

wherein R⁵ is optionally substituted with one or more R^{5a}, wherein

20 each R^{5a} is independently halogen, nitro, heteroaryl, heterocyclyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, (C₃₋₈ cycloalkyl)-C₁₋₆ alkyl-, (C₃₋₈ cycloalkenyl)-C₁₋₆ alkyl-, (C₃₋₈ cycloalkyl)-C₂₋₆ alkenyl, aryl, arylalkyl, aryloxy, aryloxyaryl, arylC₁₋₆ alkoxy, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₃₋₆cycloalkyl, SO₂R¹¹, OR¹¹, SR¹¹, N₃, SO₂R¹¹, COR¹¹, SO₂N(R¹¹)₂, SO₂NR¹¹COR¹¹, C≡N, C(O)OR¹¹, CON(R¹¹)₂, CON(R¹¹)OR¹¹, OCON(R¹¹)₂, NR¹¹COR¹¹, NR¹¹CON(R¹¹)₂, NR¹¹COOR¹¹, or N(R¹¹)₂, wherein

25 each R^{5a} is optionally substituted with one or more groups which independently are -halogen, -C₁₋₆ alkyl, aryloxy, C₀₋₆ alkylSO₂R¹¹, C₀₋₆ alkylCOOR¹¹, C₀₋₆ alkoxyaryl, -C₁₋₆ haloalkyl, -SO₂R¹¹, -OR¹¹, -SR¹¹, -N₃, -SO₂R¹¹, -COR¹¹, -SO₂N(R¹¹)₂, -SO₂NR¹¹COR¹¹, -C≡N, -C(O)OR¹¹, -CON(R¹¹)₂, -CON(R¹¹)OR¹¹, -OCON(R¹¹)₂, -NR¹¹COR¹¹, -NR¹¹CON(R¹¹)₂, -NR¹¹COOR¹¹, or -N(R¹¹)₂;

30 R² is -L³-R⁷, wherein L³ is a bond; and


R⁷ is -Z or -Y-Z, wherein

Y is -[C(R¹⁵)₂]_m- or -C₃₋₆cycloalkyl; and

Z is -H, halogen, -OR¹¹, -C(=O)R¹¹, -C(=O)OR¹¹, -C(=O)N(R¹¹)₂, -N(R¹¹)₂,
-C(=N-OH)R¹¹, -C(=S)N(R¹¹)₂, -CN, -S(=O)₂N(R¹¹)₂, -OC(=O)-R¹¹, or
-OC(=O)-N(R¹¹)₂;

R²¹ is hydrogen, halogen, C₁-C₆ alkyl, or C₁-C₆ haloalkyl; and

5 G is a group of the formula,



J is aryl or heteroaryl;

K is aryl or heteroaryl;

each R⁴ and R⁴¹ is independently halogen, heteroaryl, heterocyclyl, -G¹, -E-G¹, or -D-E-G¹,

10 wherein

D is -O-;

E is -[C(R¹⁵)₂]_m- or -C₃-C₆cycloalkyl; and

G¹ is -C₁-C₆alkyl, -C₁-C₆haloalkyl, -COR¹¹, -COOR¹¹, -CON(R¹¹)₂, -C≡N, -OR¹¹, -
SOR¹¹, -SO₂R¹¹, -SO₂N(R¹¹)₂, or -SR¹¹;

15 L² is a bond;

q is 1, 2, or 3; and

q' is 0, 1, 2 or 3;

each R¹⁰ is independently -R¹¹, -C(=O)R¹¹, -CO₂R¹¹, or -SO₂R¹¹;

each R¹¹ is independently -hydrogen, -C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, (C₃-C₈ cycloalkyl)-C₁-C₆
20 alkyl-, (C₃-C₈ cycloalkenyl)-C₁-C₆ alkyl-, (C₃-C₈ cycloalkyl)-C₂-C₆ alkenyl-C₃-C₈ cycloalkyl, -C₁-C₆
haloalkyl, -N(R¹²)₂, aryl, -(C₁-C₆)alkyl-aryl, heteroaryl, -(C₁-C₆)alkyl-heteroaryl, heterocyclyl, or
-(C₁-C₆)alkyl-heterocyclyl,

wherein any of R¹¹ is optionally substituted with one or more radicals of R¹²;

each R¹² is independently halogen, OR¹³, N(R¹³)₂, C₁-C₆haloalkyl, C₁-C₆ alkyl, C₁-C₆
25 alkoxy, (C₀-C₆ alkyl)C=O(OR¹³); C₀-C₆ alkylOR¹³, C₀-C₆ alkylCOR¹³, C₀-C₆
alkylSO₂R¹³, C₀-C₆ alkylCON(R¹³)₂, C₀-C₆ alkylCONR¹³OR¹³, C₀-C₆
alkylSO₂N(R¹³)₂, C₀-C₆ alkylSR¹³, C₀-C₆ haloalkylOR¹³, aryloxy, aralkyloxy,
aryloxyalkyl, C₀-C₆alkoxyaryl, arylC₀-C₆ alkylcarboxy, C₀-C₆ alkyl, -NR¹³SO₂ R¹³, or
-OC₀-C₆ alkylCOOR¹³;

30 each R¹³ is independently hydrogen, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, (C₃-C₈ cycloalkyl)-C₁-C₆ alkyl-,
(C₃-C₈ cycloalkenyl)-C₁-C₆ alkyl-, or (C₃-C₈ cycloalkyl)-C₂-C₆ alkenyl-;


each R^{14} is independently C_1 - C_6 alkyl, C_1 - C_6 alkoxy, halogen, C_1 - C_6 haloalkyl, C_0 - C_6 alkylCON(R^{11})₂, C_0 - C_6 alkylCONR¹¹OR¹¹, C_0 - C_6 alkylOR¹¹, or C_0 - C_6 alkylCOOR¹¹.

In one embodiment, the invention provides the compound according to formula IIa, IIb, IIc, and II_d, wherein R^1 is L^1 - R^5 , wherein L^1 is a bond, and R^2 , R^{21} , R^3 , R^5 , and G are as defined for formulas IIa-d; such compounds are hereafter designated formulas IIIa-d.

In another embodiment, the invention provides the compound according to formulas IIa-d, wherein R^1 is L^1 - R^5 ,

wherein L^1 is a bond; and R^5 is aryl optionally substituted with one or more R^{5a} , and R^2 , R^{21} , R^3 , and G are as defined for formulas IIa-d; such compounds are hereafter designated formulas IVa-d.

In another embodiment, the invention provides the compound of formula IV,



or a pharmaceutically acceptable salt, isomer, or prodrug thereof, wherein R^1 is as defined for formula IVa-d, and R^2 , R^{21} , R^4 , R^{41} , J, K, L^2 , q, and q' are as defined for formula IIa-d.

In another embodiment, the invention provides the compound according to the formula IV, wherein L^2 is a bond.

In another embodiment, the invention provides the compound according to the formula IV, wherein K is phenyl.

In another embodiment, the invention provides the compound according to the formula IV, wherein L^2 is a bond and K is phenyl.


In another embodiment, the invention provides the compound according to the formula IV, wherein L^2 is a bond and J is phenyl.

In another embodiment, the invention provides the compound according to the formula IV, wherein J is phenyl.

In another embodiment, the invention provides the compound according to the formula IV, wherein J is phenyl, and K is phenyl.

In another embodiment, the invention provides the compound according to the formula IV, wherein J is phenyl, K is phenyl, and L^2 is a bond.


In another embodiment, the invention provides the compound of formula V,



(V),

or a pharmaceutically acceptable salt, isomer, or prodrug thereof, wherein R^1 is as defined for formula IVa-d, and R^2 , R^{21} , R^4 , R^{41} , q , and q' are as defined for formula IIa-d.

5 In another embodiment, the invention provides the compound of formula VII,



(VII),

or a pharmaceutically acceptable salt, isomer, or prodrug thereof, wherein R^{21} is $-H$, $-halogen$, $-C_1-C_6alkyl$, or $-C_1-C_6haloalkyl$; R^1 is as defined for formula IVa-d; and R^2 , R^4 , R^{41} , q , and q' are as defined for formula IIa-d.

10 In another embodiment, the invention provides the compound according of the formula VII, wherein each R^{41} is independently halogen, $-G^1-E-G^1$ or $-D-E-G^1$, wherein

D is $-O-$;

E is $-[C(R^{15})_2]_m-$;

15 G^1 is $-C_1-C_6alkyl$, $-C_1-C_6haloalkyl$, halogen, $-COR^{11}$, $-COOR^{11}$, $-CON(R^{11})_2$, $-C\equiv N$, $-OR^{11}$, $-NR^{11}COR^{11}$, $-NR^{11}SO_2R^{11}$, $-N(R^{11})_2$, $-SO_2R^{11}$, or $-SO_2N(R^{11})_2$.

In another embodiment, the invention provides the compound according of the formula VII, wherein each R^{41} is independently halogen, $-G^1$ or $-E-G^1$, wherein

E is $-[C(R^{15})_2]_m-$,

20 wherein each R^{15} is independently hydrogen or halogen; and

G^1 is $-C_1-C_6alkyl$, $-C_1-C_6haloalkyl$, or halogen,

such compounds are hereafter designated formula VIIa.

In another embodiment, the invention provides the compound according of the formula VII, wherein each R^4 is independently halogen, aryl, heteroaryl, heterocyclyl, $-G^1$, or $-E-G^1$, wherein

25 E is $-[C(R^{15})_2]_m-$; and

G^1 is $-C_1-C_6alkyl$, $-C_1-C_6haloalkyl$, halogen, $-COR^{11}$, $-COOR^{11}$, $-CON(R^{11})_2$, $-C\equiv N$, $-OR^{11}$, $-N(R^{11})_2$, $-SO_2R^{11}$, $-SO_2N(R^{11})_2$, or $-SR^{11}$.

In another embodiment, the invention provides the compound according of the formula VII, wherein each R^4 is independently halogen, aryl, heteroaryl, heterocyclyl, G^1 or $-E-G^1$, wherein

E is $-[C(R^{15})_2]_m-$, wherein

each R^{15} is independently hydrogen or halogen; and

5 G^1 is $-C_1-C_6$ alkyl, $-C_1-C_6$ haloalkyl, halogen, $-OR^{11}$, or $-SO_2R^{11}$,

such compounds are hereafter designated formula VIIb.

In another embodiment, the invention provides the compound according of the formula VII, wherein R^2 is $-L^3-R^7$, wherein L^3 is a bond; and

R^7 is hydrogen, halogen, nitro, aryl, heteroaryl, heterocyclyl, $-Z$, or $-Y-Z$, wherein

10 Y is $-[C(R^{15})_2]_m-$, $-(C_3-C_6)$ cycloalkyl-, or C_2-C_6 alkenyl;

Z is $-H$, halogen, $-OR^{11}$, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)N(R^{11})_2$, $-N(R^{11})_2$, $-C(=N-OH)R^{11}$, $-C(=S)N(R^{11})_2$, $-CN$, $-S(=O)_2N(R^{11})_2$, $-C(=O)N(R^{11})N(R^{11})_2$, $-C(=O)N(R^{11})(OR^{11})$, $-OC(=O)-R^{11}$, or $-OC(=O)-N(R^{11})_2$.

15 In another embodiment, the invention provides the compound according of the formula VII, wherein R^2 is $-L^3-R^7$, wherein L^3 is a bond; and

R^7 is hydrogen, halogen, $-Z$, or $-Y-Z$, wherein

Y is $-[C(R^{15})_2]_m-$, $-(C_3-C_6)$ cycloalkyl-, or C_2-C_6 alkenyl,

wherein each R^{15} is independently H, halogen, or (C_1-C_6) alkyl; and

20 Z is $-H$, halogen, $-OR^{11}$, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)N(R^{11})_2$, $-C(=N-OH)R^{11}$, or $-C(=S)N(R^{11})_2$,

such compounds are hereafter designated formula VIIc.

In another embodiment, the invention provides the compound according of the formula VIIa, wherein each R^4 is independently halogen, aryl, heteroaryl, heterocyclyl, $-G^1$ or $-E-G^1$, wherein E is $-[C(R^{15})_2]_m-$, wherein

25 each R^{15} is independently hydrogen or halogen; and

G^1 is $-C_1-C_6$ alkyl, $-C_1-C_6$ haloalkyl, halogen, $-OR^{11}$, or $-SO_2R^{11}$,

such compounds are hereafter designated formula VIId.

In another embodiment, the invention provides the compound according of the formula VIIb, wherein R^2 is $-L^3-R^7$, wherein L^3 is a bond; and

30 R^7 is hydrogen, halogen, $-Z$, or $-Y-Z$, wherein

Y is $-[C(R^{15})_2]_m-$, $-(C_3-C_6)$ cycloalkyl-, or C_2-C_6 alkenyl,

wherein each R^{15} is independently H, halogen, or (C_1-C_6) alkyl; and

Z is $-H$, halogen, $-OR^{11}$, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)N(R^{11})_2$, $-C(=N-OH)R^{11}$, or $-C(=S)N(R^{11})_2$,

such compounds are hereafter designated formula VIIe.

In another embodiment, the invention provides the compound according of the formula VIIc, wherein each R^{41} is independently halogen, $-G^1$ or $-E-G^1$, wherein

E is $-[C(R^{15})_2]_m$,

5 wherein each R^{15} is independently hydrogen or halogen; and

G^1 is $-C_1-C_6$ alkyl, $-C_1-C_6$ haloalkyl, or halogen,

such compounds are hereafter designated formula VIIf.

In another embodiment, the invention provides the compound according of the formula VIId, wherein R^2 is $-L^3-R^7$, wherein L^3 is a bond; and

10 R^7 is hydrogen, halogen, $-Z$, or $-Y-Z$, wherein

Y is $-[C(R^{15})_2]_m$, $-(C_3-C_6)$ cycloalkyl-, or C_2-C_6 alkenyl,

wherein each R^{15} is independently H, halogen, or (C_1-C_6) alkyl; and

Z is $-H$, halogen, $-OR^{11}$, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, -

$-C(=O)N(R^{11})_2$, $-C(=N-OH)R^{11}$, or $-C(=S)N(R^{11})_2$,

15 such compounds are hereafter designated formula VIIg.

In another embodiment, the invention provides the compound according of the formula VIIe, wherein each R^{5a} is independently halogen, C_1-C_6 alkyl, C_1-C_6 haloalkyl, C_3-C_8 cycloalkyl, $-OR^{11}$, $-SO_2R^{11}$, $-COR^{11}$, $-SO_2N(R^{11})_2$, $-C\equiv N$, $-C(O)OR^{11}$, $-CON(R^{11})_2$, $-NR^{11}COR^{11}$, or $-N(R^{11})_2$.

In another embodiment, the invention provides the compound according of the formula VIIe, 20 wherein each R^{5a} is independently halogen, C_1-C_6 alkyl, C_1-C_6 haloalkyl, C_3-C_8 cycloalkyl, $-OR^{11}$, $-SO_2R^{11}$, $-COR^{11}$, $-SO_2N(R^{11})_2$, $-C\equiv N$, $-C(O)OR^{11}$, $-CON(R^{11})_2$, $-NR^{11}COR^{11}$, or $-N(R^{11})_2$; and R^{21} is $-H$.

In another embodiment, the invention provides the compound according of the formula VIIf, wherein each R^{5a} is independently halogen, C_1-C_6 alkyl, or C_1-C_6 haloalkyl.

25 In another embodiment, the invention provides the compound according of the formula VIIf, wherein each R^{5a} is independently halogen, C_1-C_6 alkyl, or C_1-C_6 haloalkyl; and R^{21} is $-H$.

In another embodiment, the invention provides the compound according of the formula VII, wherein each R^{5a} is independently halogen, C_1-C_6 alkyl, C_1-C_6 haloalkyl, C_3-C_8 cycloalkyl, $-OR^{11}$, $-SO_2R^{11}$, $-COR^{11}$, $-SO_2N(R^{11})_2$, $-C\equiv N$, $-C(O)OR^{11}$, $-CON(R^{11})_2$, $-NR^{11}COR^{11}$, or $-N(R^{11})_2$.

30 In another embodiment, the invention provides the compound according of the formula VII, wherein each R^{5a} is independently halogen, C_1-C_6 alkyl, or C_1-C_6 haloalkyl.

In another embodiment, the invention provides the compound according of the formula VII, wherein each R^{5a} is independently halogen, C_1-C_6 alkyl, or C_1-C_6 haloalkyl; and R^{21} is $-H$.

In another embodiment, the invention provides the compound according of the formula VIIg, wherein each R^{5a} is independently halogen, C_1 - C_6 alkyl, or C_1 - C_6 haloalkyl.

In another embodiment, the invention provides the compound according of the formula VIIg, wherein each R^{5a} is independently halogen, C_1 - C_6 alkyl, or C_1 - C_6 haloalkyl; and R^{21} is -H.

5 In another embodiment, the invention provides the compound according to formula VII, wherein R^{21} is hydrogen, halogen, nitro, cyano, aryl, heteroaryl, heterocyclyl, $-C_1$ - C_6 alkyl-heterocyclyl, $-C_1$ - C_6 alkyl-heteroaryl, $-C_1$ - C_6 alkyl-aryl, -Z, -Y-Z, or -X-Y-Z, wherein

X is -O-;


Y is $-[C(R^{15})_2]_m$, $-C_2$ - C_6 alkenyl, or C_3 - C_8 cycloalkyl; and

10 Z is -H, -CN, halogen, $-OR^{11}$, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)N(R^{11})_2$, $-N(R^{11})_2$, -CN, $-N_3$, $-SO_2R^{11}$, $-S(=O)_2N(R^{11})_2$, $-C(=O)N(R^{11})N(R^{11})_2$, $-C(=O)N(R^{11})(OR^{11})$, $-OC(=O)-R^{11}$, $-OC(=O)-N(R^{11})_2$, or $-N(R^{11})COOR^{11}$.

In another embodiment, the invention provides the compound according to formula VII, wherein R^{21} is hydrogen, halogen, nitro, cyano, C_1 - C_6 alkyl, or C_1 - C_6 haloalkyl.

15 In another embodiment, the invention provides the compound according to any of formulas VIIa-VIIg, wherein R^{21} is hydrogen, halogen, nitro, cyano, C_1 - C_6 alkyl, or C_1 - C_6 haloalkyl.

In another embodiment, the invention provides the compound of formula VIII,



(VIII),

20 or a pharmaceutically acceptable salt, isomer, or prodrug thereof, wherein R^1 is as defined for formula IVa-d, and R^2 , R^{21} , R^4 , R^{41} , J, K, L^2 , q, and q' are as defined for formula IIa-d.

In another embodiment, the invention provides the compound according to formula VIII wherein K is phenyl or pyridyl.


25 In another embodiment, the invention provides the compound according to formula VIII wherein K is phenyl.

In another embodiment, the invention provides the compound according to formula VIII wherein K is pyridyl.

In another embodiment, the invention provides the compound according to formula VIII wherein J is phenyl.

In another embodiment, the invention provides the compound according to formula VIII wherein J is phenyl and K is phenyl.


In another embodiment, the invention provides the compound of formula IX,



(IX),

or a pharmaceutically acceptable salt, isomer, or prodrug thereof, wherein R^1 is as defined for formula IVa-d, and R^2 , R^{21} , R^4 , R^{41} , L^2 , q , and q' are as defined for formula IIa-d.

In another embodiment, the invention provides the compound of formula X,



(X),

or a pharmaceutically acceptable salt, isomer, or prodrug thereof, wherein R^1 is as defined for formula IVa-d, and R^2 , R^{21} , R^4 , R^{41} , L^2 , q , and q' are as defined for formula IIa-d.

In another embodiment, the invention provides the compound according to formula X, wherein L^2 is a bond; such compounds are designated hereafter as formula XI.

In another embodiment, the invention provides the compound according to formula XI, wherein each R^{5a} is independently halogen, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_3 - C_8 cycloalkyl, $-OR^{11}$, $-SO_2R^{11}$, $-COR^{11}$, $-SO_2N(R^{11})_2$, $-C\equiv N$, $-C(O)OR^{11}$, $-CON(R^{11})_2$, $-NR^{11}COR^{11}$, or $-N(R^{11})_2$.

In another embodiment, the invention provides the compound according to formula XI, wherein each R^{5a} is independently halogen, C_1 - C_6 alkyl, or C_1 - C_6 haloalkyl,

such compounds are designated hereafter as formula XIa.

In another embodiment, the invention provides the compound according to formula XI, wherein R^2 is $-L^3-R^7$, wherein L^3 is a bond; and

R^7 is hydrogen, halogen, nitro, aryl, heteroaryl, heterocyclyl, $-Z$, or $-Y-Z$, wherein

Y is $-[C(R^{15})_2]_m-$, $-(C_3-C_6)$ cycloalkyl-, or C_2-C_6 alkenyl;

Z is $-H$, halogen, $-OR^{11}$, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)N(R^{11})_2$, $-N(R^{11})_2$, $-C(=N-OH)R^{11}$, $-C(=S)N(R^{11})_2$, $-CN$, $-S(=O)_2N(R^{11})_2$, $-C(=O)N(R^{11})N(R^{11})_2$, $-C(=O)N(R^{11})(OR^{11})$, $-OC(=O)-R^{11}$, or $-OC(=O)-N(R^{11})_2$.

E is $-\text{[C(R}^{15})_2]_{m-}$, wherein

each R^{15} is independently hydrogen or halogen; and

G^1 is $-\text{C}_1\text{-C}_6\text{alkyl}$, $-\text{C}_1\text{-C}_6\text{haloalkyl}$, halogen, $-\text{OR}^{11}$, or $-\text{SO}_2\text{R}^{11}$.

In another embodiment, the invention provides the compound according to formula XI, wherein R^{21} is hydrogen, halogen, nitro, cyano, aryl, heteroaryl, heterocyclyl, $-\text{C}_1\text{-C}_6\text{ alkyl-heterocyclyl}$, $-\text{C}_1\text{-C}_6$ alkyl-heteroaryl, $-\text{C}_1\text{-C}_6\text{ alkyl-aryl}$, $-\text{Z}$, $-\text{Y-Z}$, or $-\text{X-Y-Z}$, wherein

X is $-\text{O}-$;

Y is $-\text{[C(R}^{15})_2]_{m-}$, $-(\text{C}_3\text{-C}_6)\text{cycloalkyl}$, $-\text{C}_2\text{-C}_6\text{ alkenyl}$, or $\text{C}_3\text{-C}_8\text{ cycloalkyl}$; and

Z is $-\text{H}$, $-\text{CN}$, halogen, $-\text{OR}^{11}$, $-\text{C(=O)R}^{11}$, $-\text{C(=O)OR}^{11}$, $-\text{C(=O)N(R}^{11})_2$, $-\text{N(R}^{11})_2$, $-\text{CN}$, $-\text{N}_3$, $-\text{SO}_2\text{R}^{11}$, $-\text{S(=O)}_2\text{N(R}^{11})_2$, $-\text{C(=O)N(R}^{11})\text{N(R}^{11})_2$, $-\text{C(=O)N(R}^{11})(\text{OR}^{11})$, $-\text{OC(=O)-R}^{11}$, $-\text{OC(=O)-N(R}^{11})_2$, or $-\text{N(R}^{11})\text{COOR}^{11}$.


In another embodiment, the invention provides the compound according to formula XI, wherein R^{21} is hydrogen, halogen, nitro, cyano, $\text{C}_1\text{-C}_6\text{alkyl}$, or $\text{C}_1\text{-C}_6\text{haloalkyl}$.

In another embodiment, the invention provides the compound according to any of formulas XIa-XId, wherein R^{21} is hydrogen, halogen, nitro, cyano, $\text{C}_1\text{-C}_6\text{alkyl}$, or $\text{C}_1\text{-C}_6\text{haloalkyl}$.

In another embodiment, the invention provides the compound according to formula VIII, wherein J is thienyl.

In another embodiment, the invention provides the compound according to formula VIII, wherein J is thienyl and K is phenyl.


In another embodiment, the invention provides the compound of formula XII,



(XII),

or a pharmaceutically acceptable salt, isomer, or prodrug thereof, where R^1 is as defined for formula IVa-d, and R^2 , R^{21} , R^4 , R^{41} , L^2 , q, and q' are as defined for formula IIa-d.


In another embodiment, the invention provides the compound of formula XIII,



(XIII)

or a pharmaceutically acceptable salt, isomer, or prodrug thereof, where R^1 is as defined for formula IVa-d, and $R^2, R^{21}, R^4, R^{41}, L^2, q$, and q' are as defined for formula IIa-d.

In another embodiment, the invention provides the compound of formula XIV,



(XIV)

or a pharmaceutically acceptable salt, isomer, or prodrug thereof, where R^1 is as defined for formula IVa-d, and $R^2, R^{21}, R^4, R^{41}, L^2, q$, and q' are as defined for formula IIa-d.

In another embodiment, the invention provides the compound according to formula XIV, wherein L^2 is a bond, such compounds are designated hereafter as formula XV.

In another embodiment, the invention provides the compound according to formula XV, wherein each R^{5a} is independently halogen, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_3 - C_8 cycloalkyl, $-OR^{11}$, $-SO_2R^{11}$, $-COR^{11}$, $-SO_2N(R^{11})_2$, $-C\equiv N$, $-C(O)OR^{11}$, $-CON(R^{11})_2$, $NR^{11}COR^{11}$, or $-N(R^{11})_2$.

In another embodiment, the invention provides the compound according to formula XV, wherein each R^{5a} is independently halogen, C_1 - C_6 alkyl, or C_1 - C_6 haloalkyl, such compounds are designated hereafter as formula XVa.

In another embodiment, the invention provides the compound according to formula XV, wherein R^2 is $-L^3-R^7$, wherein L^3 is a bond; and

R^7 is hydrogen, halogen, nitro, aryl, heteroaryl, heterocyclyl, $-Z$, or $-Y-Z$, wherein

Y is $-[C(R^{15})_2]_m$, $-(C_3-C_6)$ cycloalkyl-, or C_2 - C_6 alkenyl;

Z is $-H$, halogen, $-OR^{11}$, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)N(R^{11})_2$, $-N(R^{11})_2$, $-C(=N-OH)R^{11}$, $-C(=S)N(R^{11})_2$, $-CN$, $-S(=O)_2N(R^{11})_2$, $-C(=O)N(R^{11})N(R^{11})_2$, $-C(=O)N(R^{11})(OR^{11})$, $-OC(=O)-R^{11}$, or $-OC(=O)-N(R^{11})_2$.

In another embodiment, the invention provides the compound according to formula XV, wherein R^2 is $-L^3-R^7$, wherein L^3 is a bond; and

R^7 is hydrogen, halogen, $-Z$, or $-Y-Z$, wherein

Y is $-[C(R^{15})_2]_m$, $-(C_3-C_6)$ cycloalkyl-, or C_2 - C_6 alkenyl,

wherein each R^{15} is independently H , halogen, or (C_1-C_6) alkyl; and

Z is $-H$, halogen, $-OR^{11}$, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)N(R^{11})_2$, $-C(=N-OH)R^{11}$, or $-C(=S)N(R^{11})_2$.

such compounds are designated hereafter as formula XVb.

In another embodiment, the invention provides the compound according to formula XV, wherein each R^4 is independently halogen, aryl, heteroaryl, heterocyclyl, $-G^1$ or $-E-G^1$, wherein

E is $-[C(R^{15})_2]_m$; and

G^1 is $-C_1-C_6$ alkyl, $-C_1-C_6$ haloalkyl, halogen, $-COR^{11}$, $-COOR^{11}$, $-CON(R^{11})_2$, $-C\equiv N$, $-OR^{11}$, $-N(R^{11})_2$, $-SO_2R^{11}$, $-SO_2N(R^{11})_2$, or $-SR^{11}$.

In another embodiment, the invention provides the compound according to formula XV, wherein each R^4 is independently halogen, aryl, heteroaryl, heterocyclyl, $-G^1$ or $-E-G^1$, wherein

E is $-[C(R^{15})_2]_m$, wherein

each R^{15} is independently hydrogen or halogen; and

G^1 is $-C_1-C_6$ alkyl, $-C_1-C_6$ haloalkyl, halogen, $-OR^{11}$, or $-SO_2R^{11}$.

In another embodiment, the invention provides the compound according to formula XVa,

R^2 is $-L^3-R^7$, wherein L^3 is a bond; and

R^7 is hydrogen, halogen, $-Z$, or $-Y-Z$, wherein

Y is $-[C(R^{15})_2]_m$, $-(C_3-C_6)$ cycloalkyl-, or C_2-C_6 alkenyl,

wherein each R^{15} is independently H, halogen, or (C_1-C_6) alkyl; and

Z is $-H$, halogen, $-OR^{11}$, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)N(R^{11})_2$, $-C(=N-OH)R^{11}$, or $-C(=S)N(R^{11})_2$,

such compounds are designated hereafter as formula XVd.

In another embodiment, the invention provides the compound according to formula XVb, wherein each R^4 is independently halogen, aryl, heteroaryl, heterocyclyl, $-G^1$ or $-E-G^1$, wherein

E is $-[C(R^{15})_2]_m$, wherein

each R^{15} is independently hydrogen or halogen; and

G^1 is $-C_1-C_6$ alkyl, $-C_1-C_6$ haloalkyl, halogen, $-OR^{11}$, or $-SO_2R^{11}$.

In another embodiment, the invention provides the compound according to formula XVd, wherein each R^4 is independently halogen, aryl, heteroaryl, heterocyclyl, $-G^1$ or $-E-G^1$, wherein

E is $-[C(R^{15})_2]_m$, wherein

each R^{15} is independently hydrogen or halogen; and

G^1 is $-C_1-C_6$ alkyl, $-C_1-C_6$ haloalkyl, halogen, $-OR^{11}$, or $-SO_2R^{11}$.

In another embodiment, the invention provides the compound according to formula XV, wherein R^{21} is hydrogen, halogen, nitro, cyano, aryl, heteroaryl, heterocyclyl, $-C_1-C_6$ alkyl-heterocyclyl, $-C_1-C_6$ alkyl-heteroaryl, $-C_1-C_6$ alkyl-aryl, $-Z$, $-Y-Z$, or $-X-Y-Z$, wherein

X is $-O-$;

Y is $-[C(R^{15})_2]_m$, $-C_2-C_6$ alkenyl, or C_3-C_8 cycloalkyl; and


Z is -H, -CN, halogen, -OR¹¹, -C(=O)R¹¹, -C(=O)OR¹¹, -C(=O)N(R¹¹)₂, -N(R¹¹)₂, -CN, -N₃, -SO₂R¹¹, -S(=O)₂N(R¹¹)₂, -C(=O)N(R¹¹)N(R¹¹)₂, -C(=O)N(R¹¹)(OR¹¹), -OC(=O)-R¹¹, -OC(=O)-N(R¹¹)₂, or -N(R¹¹)COOR¹¹.

In another embodiment, the invention provides the compound according to formula XV,
 5 wherein R²¹ is hydrogen, halogen, nitro, cyano, C₁-C₆alkyl, or C₁-C₆haloalkyl.

In another embodiment, the invention provides the compound according to any of formulas XVa-XVd, wherein R²¹ is hydrogen, halogen, nitro, cyano, C₁-C₆alkyl, or C₁-C₆haloalkyl.

In another embodiment, the invention provides the compound according to formula VIII,
 wherein K is pyridinyl.


10 In another embodiment, the invention provides the compound according to formula XVI,



(XVI)

or a pharmaceutically acceptable salt, isomer, or prodrug thereof, where R¹ is as defined for formula IVa-d, and R², R²¹, R⁴, R⁴¹, L², q, and q' are as defined for formula IIa-d.


15 In another embodiment, the invention provides the compound according to formula XVII,



(XVII)

or a pharmaceutically acceptable salt, isomer, or prodrug thereof, where R¹ is as defined for formula IVa-d, and R², R²¹, R⁴, R⁴¹, L², q, and q' are as defined for formula IIa-d.

20 In another embodiment, the invention provides the compound according to formula XVIII,



(XVIII)

or a pharmaceutically acceptable salt, isomer, or prodrug thereof, where R¹ is as defined for formula IVa-d, and R², R²¹, R⁴, R⁴¹, L², q, and q' are as defined for formula IIa-d.

In another embodiment, the invention provides the compound according to formula XVIII, wherein L^2 is a bond, such compounds are designated hereafter as formula XIX.

In another embodiment, the invention provides the compound according to formula XIX, wherein each R^{5a} is independently halogen, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_3 - C_8 cycloalkyl, $-OR^{11}$,
5 $-SO_2R^{11}$, $-COR^{11}$, $-SO_2N(R^{11})_2$, $-C\equiv N$, $-C(O)OR^{11}$, $-CON(R^{11})_2$, $-NR^{11}COR^{11}$, or $-N(R^{11})_2$.

In another embodiment, the invention provides the compound according to formula XIX, wherein each R^{5a} is independently halogen, C_1 - C_6 alkyl, or C_1 - C_6 haloalkyl, such compounds are designated hereafter as formula XIXa.

In another embodiment, the invention provides the compound according to formula XIX,
10 wherein R^2 is $-L^3-R^7$, wherein L^3 is a bond; and
 R^7 is hydrogen, halogen, nitro, aryl, heteroaryl, heterocyclyl, $-Z$, or $-Y-Z$, wherein

Y is $-[C(R^{15})_2]_m-$, $-(C_3-C_6)$ cycloalkyl-, or C_2 - C_6 alkenyl; and
 Z is $-H$, halogen, $-OR^{11}$, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)N(R^{11})_2$, $-N(R^{11})_2$,
 $-C(=N-OH)R^{11}$, $-C(=S)N(R^{11})_2$, $-CN$, $-S(=O)_2N(R^{11})_2$, $-C(=O)N(R^{11})N(R^{11})_2$,
15 $-C(=O)N(R^{11})(OR^{11})$, $-OC(=O)-R^{11}$, or $-OC(=O)-N(R^{11})_2$.

In another embodiment, the invention provides the compound according to formula XIX, wherein R^2 is $-L^3-R^7$, wherein L^3 is a bond; and

R^7 is hydrogen, halogen, $-Z$, or $-Y-Z$, wherein
 Y is $-[C(R^{15})_2]_m-$, $-(C_3-C_6)$ cycloalkyl-, or C_2 - C_6 alkenyl; and
20 wherein each R^{15} is independently H , halogen, or (C_1-C_6) alkyl; and
 Z is $-H$, halogen, $-OR^{11}$, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)N(R^{11})_2$, $-C(=N-OH)R^{11}$, or
 $-C(=S)N(R^{11})_2$,

such compounds are designated hereafter as formula XIXb.

In another embodiment, the invention provides the compound according to formula XIX,
25 wherein each R^4 is independently halogen, aryl, heteroaryl, heterocyclyl, $-G^1$ or $-E-G^1$, wherein

E is $-[C(R^{15})_2]_m-$; and
 G^1 is $-C_1-C_6$ alkyl, $-C_1-C_6$ haloalkyl, halogen, $-COR^{11}$, $-COOR^{11}$, $-CON(R^{11})_2$,
 $-C\equiv N$, $-OR^{11}$, $-N(R^{11})_2$, $-SO_2R^{11}$, $-SO_2N(R^{11})_2$, or $-SR^{11}$.

In another embodiment, the invention provides the compound according to formula XIX,
30 wherein each R^4 is independently halogen, aryl, heteroaryl, heterocyclyl, $-G^1$ or $-E-G^1$, wherein

E is $-[C(R^{15})_2]_m-$, wherein
each R^{15} is independently hydrogen or halogen; and
 G^1 is $-C_1-C_6$ alkyl, $-C_1-C_6$ haloalkyl, halogen, $-OR^{11}$, or $-SO_2R^{11}$.

In another embodiment, the invention provides the compound according to formula XIXa, wherein R^2 is $-L^3-R^7$, wherein L^3 is a bond; and

R^7 is hydrogen, halogen, $-Z$, or $-Y-Z$, wherein

Y is $-[C(R^{15})_2]_m-$, $-(C_3-C_6)\text{cycloalkyl-}$, or $C_2-C_6\text{alkenyl}$,

5 wherein each R^{15} is independently H, halogen, or $(C_1-C_6)\text{alkyl}$; and

Z is $-H$, halogen, $-OR^{11}$, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)N(R^{11})_2$, $-C(=N-OH)R^{11}$, or $-C(=S)N(R^{11})_2$

such compounds are designated hereafter as formula XIXd.

In another embodiment, the invention provides the compound according to formula XIXb, wherein each R^4 is independently halogen, aryl, heteroaryl, heterocyclyl, $-G^1$ or $-E-G^1$, wherein E is $-[C(R^{15})_2]_m-$, wherein

each R^{15} is independently hydrogen or halogen; and

G^1 is $-C_1-C_6\text{alkyl}$, $-C_1-C_6\text{haloalkyl}$, halogen, $-OR^{11}$, or $-SO_2R^{11}$.

In another embodiment, the invention provides the compound according to formula XIXd, wherein each R^4 is independently halogen, aryl, heteroaryl, heterocyclyl, $-G^1$ or $-E-G^1$, wherein E is $-[C(R^{15})_2]_m-$, wherein

each R^{15} is independently hydrogen or halogen; and

G^1 is $-C_1-C_6\text{alkyl}$, $-C_1-C_6\text{haloalkyl}$, halogen, $-OR^{11}$, or $-SO_2R^{11}$.

In another embodiment, the invention provides the compound according to formula XIX, wherein R^{21} is hydrogen, halogen, nitro, cyano, aryl, heteroaryl, heterocyclyl, $-C_1-C_6\text{alkyl-heterocyclyl}$, $-C_1-C_6\text{alkyl-heteroaryl}$, $-C_1-C_6\text{alkyl-aryl}$, $-Z$, $-Y-Z$, or $-X-Y-Z$, wherein

X is $-O-$;

Y is $-[C(R^{15})_2]_m-$, $-C_2-C_6\text{alkenyl}$, or $C_3-C_8\text{cycloalkyl}$; and

25 Z is $-H$, $-CN$, halogen, $-OR^{11}$, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)N(R^{11})_2$, $-N(R^{11})_2$, $-CN$, $-N_3$, $-SO_2R^{11}$, $-S(=O)_2N(R^{11})_2$, $-C(=O)N(R^{11})N(R^{11})_2$, $-C(=O)N(R^{11})(OR^{11})$, $-OC(=O)-R^{11}$, $-OC(=O)-N(R^{11})_2$, or $-N(R^{11})COOR^{11}$.

In another embodiment, the invention provides the compound according to formula XIX, wherein R^{21} is hydrogen, halogen, nitro, cyano, $C_1-C_6\text{alkyl}$, or $C_1-C_6\text{haloalkyl}$.

In another embodiment, the invention provides the compound according to any of formulas XIXa-XIXd, wherein R^{21} is hydrogen, halogen, nitro, cyano, $C_1-C_6\text{alkyl}$, or $C_1-C_6\text{haloalkyl}$.

In another embodiment, the invention provides the compound according to formula IIIa-d, wherein R^1 is heteroaryl optionally substituted with one or more R^{5a} .

In another embodiment, the invention provides the compound according to formulas IIIa-d, wherein R¹ is thienyl, furyl, pyrrolyl, pyridinyl, pyrimidinyl, pyrazidinyl, pyrazolyl, quinolinyl, or isoquinolinyl, all of which are optionally substituted with one or more R^{5a}.

5 In another embodiment, the invention provides the compound according to formulas IIIa-d, wherein R₁ is thienyl, furyl, pyrrolyl, pyridinyl, pyrimidinyl, pyrazidinyl, pyrazolyl, quinolinyl, or isoquinolinyl, all of which are optionally substituted with one or more R^{5a}; and J is phenyl.

In another embodiment, the invention provides the compound according to formula IIIa-d, wherein R¹ is thienyl, furyl, pyrrolyl, pyridinyl, pyrimidinyl, pyrazidinyl, pyrazolyl, quinolinyl, or isoquinolinyl, all of which are optionally substituted with one or more R^{5a}; and K is phenyl.

10 In another embodiment, the invention provides the compound according to formula IIIa-d, wherein R¹ is thienyl, furyl, pyrrolyl, pyridinyl, pyrimidinyl, pyrazidinyl, pyrazolyl, quinolinyl, or isoquinolinyl, all of which are optionally substituted with one or more R^{5a}; J is phenyl; and K is phenyl.

In another embodiment, the invention provides the compound according to formula IIIa-d, wherein R¹ is thienyl, furyl, pyrrolyl, pyridinyl, pyrimidinyl, pyrazidinyl, pyrazolyl, quinolinyl, or isoquinolinyl, all of which are optionally substituted with one or more R^{5a}; J is phenyl; K is phenyl; and L² is a bond.

In another embodiment, the invention provides the compound according to formula IIIa-d, wherein R¹ is thienyl, furyl, pyrrolyl, pyridinyl, pyrimidinyl, pyrazidinyl, pyrazolyl, quinolinyl, or isoquinolinyl, all of which are optionally substituted with at least one or more R^{5a};

20 J is phenyl; K is phenyl; L² is a bond;
and R² is -L³-R⁷, wherein L³ is a bond; and

R⁷ is hydrogen, halogen, -Z, or -Y-Z, wherein

Y is -[C(R¹⁵)₂]_m-, -(C₃-C₆)cycloalkyl-, or C₂-C₆alkenyl,

wherein each R¹⁵ is independently H, halogen, or (C₁-C₆)alkyl; and

25 Z is -H, halogen, -OR¹¹, -C(=O)R¹¹, -C(=O)OR¹¹, -C(=O)N(R¹¹)₂, -C(=N-OH)R¹¹, or -C(=S)N(R¹¹)₂.

In another embodiment, the invention provides the compound according to formula IIIa-d, wherein R¹ is thienyl, furyl, pyrrolyl, pyridinyl, pyrimidinyl, pyrazidinyl, pyrazolyl, quinolinyl, or isoquinolinyl, all of which are optionally substituted with one or more R^{5a};

30 J is phenyl; K is phenyl; L² is a bond;
and each R⁴ is independently halogen, aryl, heteroaryl, heterocyclyl, -G¹ or -E-G¹, wherein E is -[C(R¹⁵)₂]_m-, wherein

each R¹⁵ is independently hydrogen or halogen; and

G¹ is -C₁-C₆alkyl, -C₁-C₆haloalkyl, halogen, -OR¹¹, or -SO₂R¹¹.

In another embodiment, the invention provides the compound according to formula IIIa-d, wherein R^1 is thienyl, furyl, pyrrolyl, pyridinyl, pyrimidinyl, pyrazidinyl, pyrazolyl, quinolinyl, or isoquinolinyl, all of which are optionally substituted with one or more R^{5a} ;

J is phenyl; K is phenyl; L^2 is a bond;

5 and each R^{41} is independently halogen, $-G^1$ or $-E-G^1$, wherein

E is $-[C(R^{15})_2]_{m'}$,

wherein each R^{15} is independently hydrogen or halogen; and

G^1 is $-C_1-C_6$ alkyl, $-C_1-C_6$ haloalkyl, or halogen.

10 In another embodiment, the invention provides the compound according to formula IIIa-d, wherein R^1 is thienyl, furyl, pyrrolyl, pyridinyl, pyrimidinyl, pyrazidinyl, pyrazolyl, quinolinyl, or isoquinolinyl, all of which are optionally substituted with one or more R^{5a} ;

J is phenyl; K is phenyl; L^2 is a bond;

and R^{21} is hydrogen, halogen, nitro, cyano, C_1-C_6 alkyl, or C_1-C_6 haloalkyl.


15 In another embodiment, the invention provides the compound according to formula IIa-d, wherein L^1 is $-[C(R^{15})_2]_{m'}$ or $-C_3-C_8$ cycloalkyl, wherein

m' is any of 1 to 4; and

each R^{15} is independently hydrogen, halogen, (C_1-C_6) alkyl, or (C_1-C_6) haloalkyl,

and R^2 , R^{21} , R^3 , R^5 , and G are as defined for formulas IIa-d.

In another embodiment, the invention provides the compound of formula XX,



or a pharmaceutically acceptable salt, isomer, or prodrug thereof, wherein

L^1 is $-[C(R^{15})_2]_{m'}$ or $-C_3-C_8$ cycloalkyl, wherein

m' is any of 1 to 3; and

25 each R^{15} is independently hydrogen, halogen, (C_1-C_6) alkyl, or (C_1-C_6) haloalkyl,

and R^2 , R^{21} , R^4 , R^{41} , R^5 , L^2 , J, K, q, and q' are as defined for formula IIa-d.


In another embodiment, the invention provides the compound according to formula XX, wherein J is phenyl.

In another embodiment, the invention provides the compound according to formula XX, wherein K is phenyl.

In another embodiment, the invention provides the compound according to formula XX, wherein J is phenyl and K is phenyl.

5 In another embodiment, the invention provides the compound according to formula XX, wherein J is phenyl; K is phenyl; and L^2 is a bond.


In another embodiment, the invention provides the compound of formula XXI,



(XXI)

10 or a pharmaceutically acceptable salt, isomer, or prodrug thereof, where L^1 is as defined in formula XX, and R^2 , R^{21} , R^4 , R^{41} , R^5 , q , and q' are as defined for formula IIa-d.

In another embodiment, the invention provides the compound according to formula XXII,



(XXII)

15 or a pharmaceutically acceptable salt, isomer, or prodrug thereof, where L^1 is as defined in formula XX, and R^2 , R^{21} , R^4 , R^{41} , R^5 , q , and q' are as defined for formula IIa-d.

In another embodiment, the invention provides the compound according to formula XXII, wherein each R^{15} is independently $-H$ or $-(C_1-C_2)alkyl$; m' is 1 or 2; and R^5 is phenyl optionally substituted with one or more R^{5a} ,

20 such compounds are designated hereafter as formula XXIII.

In another embodiment, the invention provides the compound according to formula XXIII, wherein R^2 is $-L^3-R^7$, wherein L^3 is a bond; and

R^7 is hydrogen, halogen, $-Z$, or $-Y-Z$, wherein

Y is $-[C(R^{15})_2]_{m'}$, $-(C_3-C_6)cycloalkyl$ -, or $C_2-C_6alkenyl$,

25 wherein each R^{15} is independently H , halogen, $-(C_3-C_6)cycloalkyl$ -, or $(C_1-C_6)alkyl$; and

Z is -H, halogen, -OR¹¹, -C(=O)R¹¹, -C(=O)OR¹¹, -C(=O)N(R¹¹)₂, -C(=N-OH)R¹¹, or -C(=S)N(R¹¹)₂.

In another embodiment, the invention provides the compound according to formula XXIII, wherein each R⁴ is independently halogen, aryl, heteroaryl, heterocyclyl, -G¹ or -E-G¹, wherein E is
5 -[C(R¹⁵)₂]_m, wherein

each R¹⁵ is independently hydrogen or halogen; and

G¹ is -C₁-C₆alkyl, -C₁-C₆haloalkyl, halogen, -OR¹¹, or -SO₂R¹¹.

In another embodiment, the invention provides the compound according to formula XXIII, wherein each R⁴¹ is independently halogen, -G¹ or -E-G¹, wherein

10 E is -[C(R¹⁵)₂]_m,

wherein each R¹⁵ is independently hydrogen or halogen; and

G¹ is -C₁-C₆alkyl, -C₁-C₆haloalkyl, or halogen.

In another embodiment, the invention provides the compound according to formula XXIII, wherein R²¹ is hydrogen, halogen, nitro, cyano, C₁-C₆alkyl, or C₁-C₆haloalkyl.

15 In another embodiment, the invention provides the compound according to formula XXII, wherein R¹⁵ is -H; m is 1, 2, or 3; and R⁵ is heterocyclyl optionally substituted with one or more R^{5a}, such compounds are designated hereafter as formula XXIV.

In another embodiment, the invention provides the compound according to formula XXIV, wherein R² is -L³-R⁷, wherein L³ is a bond; and

20 R⁷ is hydrogen, halogen, -Z, or -Y-Z, wherein

Y is -[C(R¹⁵)₂]_m, -(C₃-C₆)cycloalkyl-, or C₂-C₆alkenyl,

wherein each R¹⁵ is independently H, halogen, -(C₃-C₆)cycloalkyl-, or (C₁-C₆)alkyl; and

25 Z is -H, halogen, -OR¹¹, -C(=O)R¹¹, -C(=O)OR¹¹, -C(=O)N(R¹¹)₂, -C(=N-OH)R¹¹, or -C(=S)N(R¹¹)₂.

In another embodiment, the invention provides the compound according to formula XXIV, wherein each R⁴ is independently halogen, aryl, heteroaryl, heterocyclyl, -G¹ or -E-G¹, wherein E is -[C(R¹⁵)₂]_m, wherein

each R¹⁵ is independently hydrogen or halogen; and

30 G¹ is -C₁-C₆alkyl, -C₁-C₆haloalkyl, halogen, -OR¹¹, or -SO₂R¹¹.

In another embodiment, the invention provides the compound according to formula XXIV, wherein each R⁴¹ is independently halogen, -G¹ or -E-G¹, wherein

E is -[C(R¹⁵)₂]_m,

wherein each R¹⁵ is independently hydrogen or halogen; and

G^1 is $-C_1-C_6$ alkyl, $-C_1-C_6$ haloalkyl, or halogen.

In another embodiment, the invention provides the compound according to formula XXIV, wherein R^{21} is hydrogen, halogen, nitro, cyano, C_1-C_6 alkyl, or C_1-C_6 haloalkyl.

In another embodiment, the invention provides the compound according to formula XXII, wherein each R^{15} is independently $-H$ or $-(C_1-C_2)$ alkyl; m is 1, 2, 3, or 4; and R^5 is heterocyclyl optionally substituted with one or more R^{5a} , such compounds are designated hereafter as formula XXIVa.

In another embodiment, the invention provides the compound according to formula XXIVa, wherein R^2 is $-L^3-R^7$, wherein L^3 is a bond; and

R^7 is hydrogen, halogen, $-Z$, or $-Y-Z$, wherein

Y is $-[C(R^{15})_2]_m$, $-(C_3-C_6)$ cycloalkyl-, or C_2-C_6 alkenyl,

wherein each R^{15} is independently H , halogen, or (C_1-C_6) alkyl; and

Z is $-H$, halogen, $-OR^{11}$, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)N(R^{11})_2$, $-C(=N-OH)R^{11}$, or $-C(=S)N(R^{11})_2$.

In another embodiment, the invention provides the compound according to formula XXIVa, wherein each R^4 is independently halogen, aryl, heteroaryl, heterocyclyl, $-G^1$ or $-E-G^1$, wherein E is $-[C(R^{15})_2]_m$, wherein

each R^{15} is independently hydrogen or halogen; and

G^1 is $-C_1-C_6$ alkyl, $-C_1-C_6$ haloalkyl, halogen, $-OR^{11}$, or $-SO_2R^{11}$.

In another embodiment, the invention provides the compound according to formula XXIVa, wherein each R^{41} is independently halogen, $-G^1$ or $-E-G^1$, wherein


E is $-[C(R^{15})_2]_m$,

wherein each R^{15} is independently hydrogen or halogen; and

G^1 is $-C_1-C_6$ alkyl, $-C_1-C_6$ haloalkyl or halogen.

In another embodiment, the invention provides the compound according to formula XXIVa, wherein R^{21} is hydrogen, halogen, nitro, cyano, C_1-C_6 alkyl, or C_1-C_6 haloalkyl.

In another embodiment, the invention provides the compound of formula XXVI,



or a pharmaceutically acceptable salt, isomer, or prodrug thereof, wherein

L^1 is $-[C(R^{15})_2]_{m'}$ or $-C_3-C_8$ cycloalkyl, wherein

m' is any of 1 to 3; and

each R^{15} is independently hydrogen, halogen, (C_1-C_6) alkyl, (C_3-C_6) cycloalkyl, or (C_1-C_6) haloalkyl,

5 and $R^2, R^{21}, R^4, R^{41}, R^5, L^2, J, K, q$, and q' are as defined for formula IIa-d.

In another embodiment, the invention provides the compound according to formula XXVI, wherein J is aryl.

In another embodiment, the invention provides the compound according to formula XXVI, wherein J is phenyl.

10 In another embodiment, the invention provides the compound according to formula XXVI, wherein J is heteroaryl, such compounds are referred to hereafter as compounds of formula XXVIa.

In another embodiment, the invention provides the compound according to formula XXVIa wherein K is aryl.

15 In another embodiment, the invention provides the compound according to formula XXVIa wherein K is phenyl.

In another embodiment, the invention provides the compound according to formula XXVI, wherein K is aryl.

In another embodiment, the invention provides the compound according to formula XXVI, wherein K is heteroaryl, such compounds are referred to hereafter as compounds of formula XXVIb.

20 In another embodiment, the invention provides the compound according to formula XXVIb wherein J is aryl.


In another embodiment, the invention provides the compound according to formula XXVIb wherein J is phenyl.

25 In another embodiment, the invention provides the compound according to formula XXVI, wherein K is phenyl.

In another embodiment, the invention provides the compound according to formula XXVI, wherein J is phenyl and K is phenyl.

In another embodiment, the invention provides the compound according to formula XXVI, wherein J is phenyl; K is phenyl; and L^2 is a bond.


30 In another embodiment, the invention provides the compound of formula XXVII,



(XXVII)

or a pharmaceutically acceptable salt, isomer, or prodrug thereof, where L^1 is as defined in formula XXVI, and $R^2, R^{21}, R^4, R^{41}, R^5, q$, and q' are as defined for formula IIa-d.

5 In another embodiment, the invention provides the compound according to formula XXVIII,



(XXVIII)

or a pharmaceutically acceptable salt, isomer, or prodrug thereof, where L^1 is as defined in formula XXVI, and $R^2, R^{21}, R^4, R^{41}, R^5, q$, and q' are as defined for formula IIa-d.

10 In another embodiment, the invention provides the compound according to formula XXVIII, wherein R^2 is $-L^3-R^7$, wherein L^3 is a bond; and

R^7 is hydrogen, halogen, $-Z$, or $-Y-Z$, wherein

Y is $-[C(R^{15})_2]_m$, $-(C_3-C_6)\text{cycloalkyl}$ -, or $C_2-C_6\text{alkenyl}$,

wherein each R^{15} is independently H, halogen, $-(C_3-C_6)\text{cycloalkyl}$ -, or

15 $(C_1-C_6)\text{alkyl}$; and

Z is $-H$, halogen, $-OR^{11}$, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)N(R^{11})_2$, $-C(=N-OH)R^{11}$, or $-C(=S)N(R^{11})_2$.

In another embodiment, the invention provides the compound according to formula XXVIII, wherein each R^4 is independently halogen, aryl, heteroaryl, heterocyclyl, $-G^1$ or $-E-G^1$, wherein E is

20 $-[C(R^{15})_2]_m$, wherein

each R^{15} is independently hydrogen or halogen; and

G^1 is $-C_1-C_6\text{alkyl}$, $-C_1-C_6\text{haloalkyl}$, halogen, $-OR^{11}$, or $-SO_2R^{11}$.

In another embodiment, the invention provides the compound according to formula XXVIII, wherein each R^{41} is independently halogen, $-G^1$ or $-E-G^1$, wherein

25 E is $-[C(R^{15})_2]_m$,

wherein each R^{15} is independently hydrogen or halogen; and

G^1 is $-C_1-C_6\text{alkyl}$, $-C_1-C_6\text{haloalkyl}$, or halogen.

In another embodiment, the invention provides the compound according to formula XXVII, wherein each R^{15} is independently $-H$, $-(C_1-C_6)alkyl$, or $-(C_3-C_6)cycloalkyl$; m' is 1 or 2; and R^5 is phenyl optionally substituted with one or more R^{5a} , such compounds are designated hereafter as formula XXIX.

5 In another embodiment, the invention provides the compound according to formula XXIX, wherein R^2 is $-L^3-R^7$, wherein L^3 is a bond; and

R^7 is hydrogen, halogen, $-Z$, or $-Y-Z$, wherein

Y is $-[C(R^{15})_2]_{m'}$, $-(C_3-C_6)cycloalkyl$ -, or $C_2-C_6alkenyl$,

10 wherein each R^{15} is independently H , halogen, $-(C_3-C_6)cycloalkyl$ -, or $(C_1-C_6)alkyl$; and

Z is $-H$, halogen, $-OR^{11}$, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)N(R^{11})_2$, $-C(=N-OH)R^{11}$, or $-C(=S)N(R^{11})_2$.

In another embodiment, the invention provides the compound according to formula XXIX, wherein each R^4 is independently halogen, aryl, heteroaryl, heterocyclyl, $-G^1$ or $-E-G^1$, wherein E is
15 $-[C(R^{15})_2]_{m'}$, wherein

each R^{15} is independently hydrogen or halogen; and

G^1 is $-C_1-C_6alkyl$, $-C_1-C_6haloalkyl$, halogen, $-OR^{11}$, or $-SO_2R^{11}$.

In another embodiment, the invention provides the compound according to formula XXIX, wherein each R^{41} is independently halogen, $-G^1$ or $-E-G^1$, wherein

20 E is $-[C(R^{15})_2]_{m'}$,

wherein each R^{15} is independently hydrogen or halogen; and

G^1 is $-C_1-C_6alkyl$, $-C_1-C_6haloalkyl$, or halogen.

In another embodiment, the invention provides the compound according to formula XXIX, wherein R^{21} is hydrogen, halogen, nitro, cyano, C_1-C_6alkyl , or $C_1-C_6haloalkyl$.

25 In another embodiment, the invention provides the compound according to formula XXVIII, wherein R^{15} is $-H$; m is 1, 2, or 3; and R^5 is heterocyclyl optionally substituted with one or more R^{5a} , such compounds are designated hereafter as formula XXX.

In another embodiment, the invention provides the compound according to formula XXX, wherein R^2 is $-L^3-R^7$, wherein L^3 is a bond; and

30 R^7 is hydrogen, halogen, $-Z$, or $-Y-Z$, wherein

Y is $-[C(R^{15})_2]_{m'}$, $-(C_3-C_6)cycloalkyl$ -, or $C_2-C_6alkenyl$,

wherein each R^{15} is independently H , halogen, $-(C_3-C_6)cycloalkyl$ -, or $(C_1-C_6)alkyl$; and

Z is -H, halogen, -OR¹¹, -C(=O)R¹¹, -C(=O)OR¹¹, -C(=O)N(R¹¹)₂, -C(=N-OH)R¹¹, or -C(=S)N(R¹¹)₂.

In another embodiment, the invention provides the compound according to formula XXX, wherein each R⁴ is independently halogen, aryl, heteroaryl, heterocyclyl, -G¹ or -E-G¹, wherein E is
5 -[C(R¹⁵)₂]_m, wherein

each R¹⁵ is independently hydrogen or halogen; and

G¹ is -C₁-C₆alkyl, -C₁-C₆haloalkyl, halogen, -OR¹¹, or -SO₂R¹¹.

In another embodiment, the invention provides the compound according to formula XXX, wherein each R⁴¹ is independently halogen, -G¹ or -E-G¹, wherein

10 E is -[C(R¹⁵)₂]_m,

wherein each R¹⁵ is independently hydrogen or halogen; and

G¹ is -C₁-C₆alkyl, -C₁-C₆haloalkyl, or halogen.

In another embodiment, the invention provides the compound according to formula XXX, wherein R²¹ is hydrogen, halogen, nitro, cyano, C₁-C₆alkyl, or C₁-C₆haloalkyl.

15 In another embodiment, the invention provides the compound according to formula XXVIII, wherein each R¹⁵ is independently -H or -(C₁-C₂)alkyl; m is 1, 2, 3, or 4; and R⁵ is heterocyclyl optionally substituted with one or more R^{5a}, such compounds are designated hereafter as formula XXXI.

In another embodiment, the invention provides the compound according to formula XXXI, wherein R² is -L³-R⁷, wherein L³ is a bond; and

20 R⁷ is hydrogen, halogen, -Z, or -Y-Z, wherein

Y is -[C(R¹⁵)₂]_m, -(C₃-C₆)cycloalkyl-, or C₂-C₆alkenyl,

wherein each R¹⁵ is independently H, halogen, -(C₃-C₆)cycloalkyl-, or (C₁-C₆)alkyl; and

Z is -H, halogen, -OR¹¹, -C(=O)R¹¹, -C(=O)OR¹¹, -C(=O)N(R¹¹)₂, -C(=N-OH)R¹¹, or
25 -C(=S)N(R¹¹)₂.

In another embodiment, the invention provides the compound according to formula XXXI, wherein each R⁴ is independently halogen, aryl, heteroaryl, heterocyclyl, -G¹ or -E-G¹, wherein E is
-[C(R¹⁵)₂]_m, wherein

each R¹⁵ is independently hydrogen or halogen; and

30 G¹ is -C₁-C₆alkyl, -C₁-C₆haloalkyl, halogen, -OR¹¹, or -SO₂R¹¹.

In another embodiment, the invention provides the compound according to formula XXXI, wherein each R⁴¹ is independently halogen, -G¹ or -E-G¹, wherein

E is -[C(R¹⁵)₂]_m,

wherein each R¹⁵ is independently hydrogen or halogen; and

G^1 is $-C_1-C_6$ alkyl, $-C_1-C_6$ haloalkyl or halogen.

In another embodiment, the invention provides the compound according to formula XXXI, wherein R^{21} is hydrogen, halogen, nitro, cyano, C_1-C_6 alkyl, or C_1-C_6 haloalkyl.

In another embodiment, the invention provides the compound according to formula XXII, wherein each R^{15} is independently $-H$ or $-(C_1-C_2)$ alkyl; m' is 1 or 2; and R^5 is heteroaryl optionally substituted with one or more R^{5a} .

In another embodiment, the invention provides the compound according to formula XXII, wherein each R^{15} is independently $-H$ or $-(C_1-C_2)$ alkyl; m' is 1 or 2; R^5 is heteroaryl optionally substituted with one or more R^{5a} ; and R^2 is $-L^3-R^7$, wherein

L^3 is a bond; and

R^7 is hydrogen, halogen, $-Z$, or $-Y-Z$, wherein

Y is $-[C(R^{15})_2]_{m'}$ or C_2-C_6 alkenyl,

wherein each R^{15} is independently H , halogen, or (C_1-C_6) alkyl; and

Z is $-H$, halogen, $-OR^{11}$, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)N(R^{11})_2$, $-C(=N-OH)R^{11}$, or $-C(=S)N(R^{11})_2$.

In another embodiment, the invention provides the compound according to formula XXII, wherein each R^{15} is independently $-H$ or $-(C_1-C_2)$ alkyl; m' is 1 or 2;

R^5 is heteroaryl optionally substituted with one or more R^{5a} ; and

each R^4 is independently halogen, aryl, heteroaryl, heterocyclyl, $-G^1$, or $-E-G^1$, wherein

E is $-[C(R^{15})_2]_{m'}$, wherein

each R^{15} is independently hydrogen or halogen; and

G^1 is $-C_1-C_6$ alkyl, $-C_1-C_6$ haloalkyl, halogen, $-OR^{11}$, or $-SO_2R^{11}$.

In another embodiment, the invention provides the compound according to formula XXII, wherein each R^{15} is independently $-H$ or $-(C_1-C_2)$ alkyl; m' is 1 or 2; R^5 is heteroaryl optionally

substituted with one or more R^{5a} ; and each R^{41} is independently halogen, $-G^1$, or $-E-G^1$, wherein

E is $-[C(R^{15})_2]_{m'}$,

wherein each R^{15} is independently hydrogen or halogen; and

G^1 is $-C_1-C_6$ alkyl, $-C_1-C_6$ haloalkyl, or halogen.

In another embodiment, the invention provides the compound according to formula XXVII, wherein each R^{15} is independently $-H$ or $-(C_1-C_2)$ alkyl; m' is 1 or 2; and R^5 is heteroaryl optionally

substituted with one or more R^{5a} .

In another embodiment, the invention provides the compound according to formula XXVII, wherein each R^{15} is independently $-H$ or $-(C_1-C_2)$ alkyl; m' is 1 or 2; R^5 is heteroaryl optionally substituted with one or more R^{5a} ; and

R^2 is $-L^3-R^7$, wherein

L^3 is a bond; and

R^7 is hydrogen, halogen, $-Z$, or $-Y-Z$, wherein

Y is $-[C(R^{15})_2]_{m'}$ or C_2-C_6 alkenyl,

5 wherein each R^{15} is independently H, halogen, or (C_1-C_6) alkyl; and

Z is $-H$, halogen, $-OR^{11}$, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)N(R^{11})_2$, $-C(=N-OH)R^{11}$, or $-C(=S)N(R^{11})_2$.

In another embodiment, the invention provides the compound according to formula XXVII, wherein each R^{15} is independently $-H$ or $-(C_1-C_2)$ alkyl; m' is 1 or 2; R^5 is heteroaryl optionally substituted with one or more R^{5a} ; and each R^4 is independently halogen, aryl, heteroaryl, heterocyclyl, $-G^1$, or $-E-G^1$, wherein

10 E is $-[C(R^{15})_2]_{m'}$, wherein

each R^{15} is independently hydrogen or halogen; and

G^1 is $-C_1-C_6$ alkyl, $-C_1-C_6$ haloalkyl, halogen, $-OR^{11}$, or $-SO_2R^{11}$.

15 In another embodiment, the invention provides the compound according to formula XXVII, wherein each R^{15} is independently $-H$ or $-(C_1-C_2)$ alkyl; m' is 1 or 2; R^5 is heteroaryl optionally substituted with one or more R^{5a} ; and


each R^{41} is independently halogen, $-G^1$, or $-E-G^1$, wherein

E is $-[C(R^{15})_2]_{m'}$,

20 wherein each R^{15} is independently hydrogen or halogen; and

G^1 is $-C_1-C_6$ alkyl, $-C_1-C_6$ haloalkyl, or halogen.

In another embodiment, the invention provides the compound according to formulas XXXII and XXXIII,



25

(XXXII)

(XXXIII)

or a pharmaceutically acceptable salt, isomer, or prodrug thereof, wherein R^3 is hydrogen, C_1-C_6 alkyl, or C_1-C_6 haloalkyl, and R^1 , R^2 , R^4 , R^{41} , L^2 , q , and q' are as defined for formulas IIa-d.

In another embodiment, the invention provides the compound according to formulas XXXII and XXXIII, wherein J is aryl; K is aryl; and L^2 is a bond.

In another embodiment, the invention provides the compound according to formulas XXXII and XXXIII, wherein J is phenyl; K is phenyl; and L^2 is a bond.

In another embodiment, the invention provides the compound according to formulas XXXII and XXXIII, wherein J is phenyl; K is phenyl; L^2 is a bond; and R^1 is phenyl optionally substituted with one or more R^{5a} .

In another embodiment, the invention provides the compound according to formulas XXXII and XXXIII, wherein J is phenyl; K is phenyl; L^2 is a bond; R^1 is phenyl optionally substituted with one or more R^{5a} ; and each R^4 is independently halogen, aryl, heteroaryl, heterocyclyl, $-G^1$ or $-E-G^1$, wherein

E is $-[C(R^{15})_2]_m-$; and

G^1 is $-C_1-C_6$ alkyl, $-C_1-C_6$ haloalkyl, halogen, $-COR^{11}$, $-COOR^{11}$, $-CON(R^{11})_2$, $-C\equiv N$, $-OR^{11}$, $-N(R^{11})_2$, $-SO_2R^{11}$, $-SO_2N(R^{11})_2$, or $-SR^{11}$.

In another embodiment, the invention provides the compound according to formulas XXXII and XXXIII, wherein J is phenyl; K is phenyl; L^2 is a bond; R^1 is phenyl optionally substituted with one or more R^{5a} ; and each R^4 is independently halogen, $-G^1$ or $-E-G^1$, wherein

E is $-[C(R^{15})_2]_m-$,

wherein each R^{15} is independently hydrogen or halogen; and

G^1 is $-C_1-C_6$ alkyl, $-C_1-C_6$ haloalkyl, or halogen.

In another embodiment, the invention provides the compound according to formulas XXXII and XXXIII, wherein J is phenyl; K is phenyl; L^2 is a bond; R^1 is phenyl optionally substituted with one or more R^{5a} ; and R^2 is $-L^3-R^7$, wherein

L^3 is a bond; and

R^7 is hydrogen, halogen, $-Z$, or $-Y-Z$, wherein

Y is $-[C(R^{15})_2]_m-$, $-(C_3-C_6)$ cycloalkyl-, or C_2-C_6 alkenyl-,

wherein each R^{15} is independently H, halogen, or (C_1-C_6) alkyl; and

Z is $-H$, halogen, $-OR^{11}$, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)N(R^{11})_2$, $-C(=N-OH)R^{11}$, or $-C(=S)N(R^{11})_2$.

In another embodiment, the invention provides the compound according to formulas XXXII and XXXIII, wherein J is phenyl; K is phenyl; L^2 is a bond; R^1 is phenyl optionally substituted with one or more R^{5a} ; and each R^{5a} is independently halogen, C_1-C_6 alkyl, C_1-C_6 alkoxy, or C_1-C_6 haloalkyl.

In another embodiment, the invention provides the compound according to formulas XXXII and XXXIII, wherein J is heteroaryl; K is aryl; and L^2 is a bond.

In another embodiment, the invention provides the compound according to formulas XXXII and XXXIII, wherein J is thienyl, pyrrolyl, furanyl, pyridyl, pyrimidyl, or pyrazinyl; K is phenyl; and L² is a bond.

5 In another embodiment, the invention provides the compound according to formulas XXXII and XXXIII, wherein J is thienyl, pyrrolyl, furanyl, pyridyl, pyrimidyl, or pyrazinyl; K is phenyl; L² is a bond; and R¹ is phenyl optionally substituted with one or more R^{5a}.

In another embodiment, the invention provides the compound according to formulas XXXII and XXXIII, wherein J is thienyl, pyrrolyl, furanyl, pyridyl, pyrimidyl, or pyrazinyl; K is phenyl; L² is a bond; R¹ is phenyl optionally substituted with one or more R^{5a}; and each R⁴ is independently halogen, aryl, heteroaryl, heterocyclyl, -G¹ or -E-G¹, wherein

E is -[C(R¹⁵)₂]_m-; and

G¹ is -C₁-C₆alkyl, -C₁-C₆haloalkyl, halogen, -COR¹¹, -COOR¹¹, -CON(R¹¹)₂, -C≡N, -OR¹¹, -N(R¹¹)₂, -SO₂R¹¹, -SO₂N(R¹¹)₂, or -SR¹¹.

15 In another embodiment, the invention provides the compound according to formulas XXXII and XXXIII, wherein J is thienyl, pyrrolyl, furanyl, pyridyl, pyrimidyl, or pyrazinyl; K is phenyl; L² is a bond; R¹ is phenyl optionally substituted with one or more R^{5a}; and each R⁴¹ is independently halogen, -G¹ or -E-G¹, wherein

E is -[C(R¹⁵)₂]_m-,

wherein each R¹⁵ is independently hydrogen or halogen; and

20 G¹ is -C₁-C₆alkyl, -C₁-C₆haloalkyl, or halogen.

In another embodiment, the invention provides the compound according to formulas XXXII and XXXIII, wherein J is thienyl, pyrrolyl, furanyl, pyridyl, pyrimidyl, or pyrazinyl; K is phenyl; L² is a bond; R¹ is phenyl optionally substituted with one or more R^{5a}; and R² is -L³-R⁷, wherein

L³ is a bond; and

25 R⁷ is hydrogen, halogen, -Z, or -Y-Z, wherein

Y is -[C(R¹⁵)₂]_m-, -(C₃-C₆)cycloalkyl-, or C₂-C₆alkenyl, ,

wherein each R¹⁵ is independently H, halogen, or (C₁-C₆)alkyl; and

Z is -H, halogen, -OR¹¹, -C(=O)R¹¹, -C(=O)OR¹¹, -C(=O)N(R¹¹)₂, -C(=N-OH)R¹¹, or -C(=S)N(R¹¹)₂.

30 In another embodiment, the invention provides the compound according to formulas XXXII and XXXIII, wherein J is thienyl, pyrrolyl, furanyl, pyridyl, pyrimidyl, or pyrazinyl; K is phenyl; L² is a bond; R¹ is phenyl optionally substituted with one or more R^{5a}; and each R^{5a} is independently halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, or C₁-C₆ haloalkyl.

In another embodiment, the invention provides the compound according to formulas XXXII and XXXIII, wherein J is heteroaryl; K is heteroaryl; and L^2 is a bond.

In another embodiment, the invention provides the compound according to formulas XXXII and XXXIII, wherein J is thienyl, pyrrolyl, furanyl, pyridyl, pyrimidyl, or pyrazinyl; K is pyridyl, pyrimidyl, or pyrazinyl; and L^2 is a bond.

In another embodiment, the invention provides the compound according to formulas XXXII and XXXIII, wherein J is thienyl, pyrrolyl, furanyl, pyridyl, pyrimidyl, or pyrazinyl; K is pyridyl, pyrimidyl, or pyrazinyl; L^2 is a bond; and R^1 is phenyl optionally substituted with one or more R^{5a} .

In another embodiment, the invention provides the compound according to formulas XXXII and XXXIII, wherein J is thienyl, pyrrolyl, furanyl, pyridyl, pyrimidyl, or pyrazinyl; K is pyridyl, pyrimidyl, or pyrazinyl; L^2 is a bond; R^1 is phenyl optionally substituted with one or more R^{5a} ; and each R^4 is independently halogen, aryl, heteroaryl, heterocyclyl, $-G^1$ or $-E-G^1$, wherein

E is $-[C(R^{15})_2]_m$; and

G^1 is $-C_1-C_6$ alkyl, $-C_1-C_6$ haloalkyl, halogen, $-COR^{11}$, $-COOR^{11}$, $-CON(R^{11})_2$, $-C\equiv N$, $-OR^{11}$, $-N(R^{11})_2$, $-SO_2R^{11}$, $-SO_2N(R^{11})_2$, or $-SR^{11}$.

In another embodiment, the invention provides the compound according to formulas XXXII and XXXIII, wherein J is thienyl, pyrrolyl, furanyl, pyridyl, pyrimidyl, or pyrazinyl; K is pyridyl, pyrimidyl, or pyrazinyl; L^2 is a bond; R^1 is phenyl optionally substituted with one or more R^{5a} ; and each R^{41} is independently halogen, $-G^1$ or $-E-G^1$, wherein

E is $-[C(R^{15})_2]_m$,

wherein each R^{15} is independently hydrogen or halogen; and

G^1 is $-C_1-C_6$ alkyl, $-C_1-C_6$ haloalkyl, or halogen.

In another embodiment, the invention provides the compound according to formulas XXXII and XXXIII, wherein J is thienyl, pyrrolyl, furanyl, pyridyl, pyrimidyl, or pyrazinyl; K is pyridyl, pyrimidyl, or pyrazinyl; L^2 is a bond; R^1 is phenyl optionally substituted with one or more R^{5a} ; and R^2 is $-L^3-R^7$, wherein

L^3 is a bond; and

R^7 is hydrogen, halogen, $-Z$, or $-Y-Z$, wherein



Y is $-[C(R^{15})_2]_m$, $-(C_3-C_6)$ cycloalkyl-, or C_2-C_6 alkenyl, ,

wherein each R^{15} is independently H, halogen, or (C_1-C_6) alkyl; and

Z is $-H$, halogen, $-OR^{11}$, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)N(R^{11})_2$, $-C(=N-OH)R^{11}$, or $-C(=S)N(R^{11})_2$.

In another embodiment, the invention provides the compound according to formulas XXXII and XXXIII, wherein J is thienyl, pyrrolyl, furanyl, pyridyl, pyrimidyl, or pyrazinyl; K is pyridyl, pyrimidyl, or pyrazinyl; L² is a bond; R¹ is phenyl optionally substituted with one or more R^{5a}; and each R^{5a} is independently halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, or C₁-C₆ haloalkyl.

5 In another embodiment, the invention provides the compound according to formulas XXXIV and XXXV,



10 or a pharmaceutically acceptable salt, isomer, or prodrug thereof, wherein R³ is hydrogen, C₁-C₆alkyl, or C₁-C₆haloalkyl, and R¹, R², R⁴, R⁴¹, L², q, and q' are as defined for formulas IIa-d.

In another embodiment, the invention provides the compound according to formulas XXXIV and XXXV, wherein J is aryl; K is aryl; and L² is a bond.

In another embodiment, the invention provides the compound according to formulas XXXIV and XXXV, wherein J is phenyl; K is phenyl; and L² is a bond.

15 In another embodiment, the invention provides the compound according to formulas XXXIV and XXXV, wherein J is phenyl; K is phenyl; and L² is a bond; each R¹⁵ is independently -H or - (C₁-C₂)alkyl; m' is 1 or 2; and R⁵ is heteroaryl optionally substituted with one or more R^{5a}.

20 In another embodiment, the invention provides the compound according to formulas XXXIV and XXXV, wherein J is phenyl; K is phenyl; and L² is a bond; each R¹⁵ is independently -H or - (C₁-C₂)alkyl; m' is 1 or 2; R⁵ is heteroaryl optionally substituted with one or more R^{5a}; and R² is -L³-R⁷, wherein

L³ is a bond; and

R⁷ is hydrogen, halogen, -Z, or -Y-Z, wherein

Y is -[C(R¹⁵)₂]_m- or C₂-C₆alkenyl,

25 wherein each R¹⁵ is independently H, halogen, or (C₁-C₆)alkyl; and

Z is -H, halogen, -OR¹¹, -C(=O)R¹¹, -C(=O)OR¹¹, -C(=O)N(R¹¹)₂, -C(=N-OH)R¹¹, or -C(=S)N(R¹¹)₂.

In another embodiment, the invention provides the compound according to formulas XXXIV and XXXV, wherein J is phenyl; K is phenyl; and L² is a bond; each R¹⁵ is independently -H or -

(C₁-C₂)alkyl; m' is 1 or 2; R⁵ is heteroaryl optionally substituted with one or more R^{5a}; and each R⁴ is independently halogen, aryl, heteroaryl, heterocyclyl, -G¹, or -E-G¹, wherein

E is -[C(R¹⁵)₂]_m, wherein

each R¹⁵ is independently hydrogen or halogen; and

5 G¹ is -C₁-C₆alkyl, -C₁-C₆haloalkyl, halogen, -OR¹¹, or -SO₂R¹¹.

In another embodiment, the invention provides the compound according to formulas XXXIV and XXXV, wherein J is phenyl; K is phenyl; and L² is a bond; each R¹⁵ is independently -H or - (C₁-C₂)alkyl; m' is 1 or 2; R⁵ is heteroaryl optionally substituted with one or more R^{5a}; and each R⁴¹ is independently halogen, -G¹, or -E-G¹, wherein

10 E is -[C(R¹⁵)₂]_m,

wherein each R¹⁵ is independently hydrogen or halogen; and

G¹ is -C₁-C₆alkyl, -C₁-C₆haloalkyl, or halogen.

In another embodiment, the invention provides the compound according to formulas XXXIV and XXXV, wherein J is phenyl; K is phenyl; L² is a bond; and each R¹⁵ is independently -H or - (C₁-C₂)alkyl; m' is 1 or 2; and R⁵ is phenyl optionally substituted with one or more R^{5a}.

In another embodiment, the invention provides the compound according to formulas XXXIV and XXXV, wherein J is phenyl; K is phenyl; L² is a bond; each R¹⁵ is independently -H or - (C₁-C₂)alkyl; m' is 1 or 2; and R⁵ is phenyl optionally substituted with one or more R^{5a}; and each R⁴ is independently halogen, aryl, heteroaryl, heterocyclyl, -G¹ or -E-G¹, wherein

20 E is -[C(R¹⁵)₂]_m; and

G¹ is -C₁-C₆alkyl, -C₁-C₆haloalkyl, halogen, -COR¹¹, -COOR¹¹, -CON(R¹¹)₂, -C≡N, -OR¹¹, -N(R¹¹)₂, -SO₂R¹¹, -SO₂N(R¹¹)₂, or -SR¹¹.

In another embodiment, the invention provides the compound according to formulas XXXIV and XXXV, wherein J is phenyl; K is phenyl; L² is a bond; each R¹⁵ is independently -H or - (C₁-C₂)alkyl; m' is 1 or 2; and R⁵ is phenyl optionally substituted with one or more R^{5a}; and each R⁴¹ is independently halogen, -G¹ or -E-G¹, wherein

E is -[C(R¹⁵)₂]_m,

wherein each R¹⁵ is independently hydrogen or halogen; and

G¹ is -C₁-C₆alkyl, -C₁-C₆haloalkyl, or halogen.

30 In another embodiment, the invention provides the compound according to formulas XXXIV and XXXV, wherein J is phenyl; K is phenyl; L² is a bond; each R¹⁵ is independently -H or - (C₁-C₂)alkyl; m' is 1 or 2; and R⁵ is phenyl optionally substituted with one or more R^{5a}; and R² is -L³-R⁷, wherein L³ is a bond; and

R⁷ is hydrogen, halogen, -Z, or -Y-Z, wherein

Y is -[C(R¹⁵)₂]_m-, -(C₃-C₆)cycloalkyl-, or C₂-C₆alkenyl,

wherein each R¹⁵ is independently H, halogen, or (C₁-C₆)alkyl; and

Z is -H, halogen, -OR¹¹, -C(=O)R¹¹, -C(=O)OR¹¹, -C(=O)N(R¹¹)₂, -C(=N-OH)R¹¹, or

-C(=S)N(R¹¹)₂.

In another embodiment, the invention provides the compound according to formulas XXXIV and XXXV, wherein J is phenyl; K is phenyl; L² is a bond; each R¹⁵ is independently -H or - (C₁-C₂)alkyl; m' is 1 or 2; and R⁵ is phenyl optionally substituted with one or more R^{5a}, and each R^{5a} is independently halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, or C₁-C₆ haloalkyl.

In another embodiment, the invention provides the compound according to formulas XXXIV and XXXV, wherein J is phenyl; K is phenyl; L² is a bond; R¹⁵ is -H; m is 1, 2, or 3; and R⁵ is heterocyclyl optionally substituted with one or more R^{5a}.

In another embodiment, the invention provides the compound according to formulas XXXIV and XXXV, wherein J is phenyl; K is phenyl; L² is a bond; R¹⁵ is -H; m is 1, 2, or 3;

R⁵ is heterocyclyl optionally substituted with one or more R^{5a}, and each R⁴ is independently halogen, aryl, heteroaryl, heterocyclyl, -G¹ or -E-G¹, wherein

E is -[C(R¹⁵)₂]_m-, and

G¹ is -C₁-C₆alkyl, -C₁-C₆haloalkyl, halogen, -COR¹¹, -COOR¹¹, -CON(R¹¹)₂, -C≡N, -OR¹¹, -N(R¹¹)₂, -SO₂R¹¹, -SO₂N(R¹¹)₂, or -SR¹¹.

In another embodiment, the invention provides the compound according to formulas XXXIV and XXXV, wherein J is phenyl; K is phenyl; L² is a bond; R¹⁵ is -H; m is 1, 2, or 3; R⁵ is heterocyclyl optionally substituted with one or more R^{5a}; and each R⁴¹ is independently halogen, -G¹ or -E-G¹, wherein

E is -[C(R¹⁵)₂]_m-,

wherein each R¹⁵ is independently hydrogen or halogen; and

G¹ is -C₁-C₆alkyl, -C₁-C₆haloalkyl, or halogen.

In another embodiment, the invention provides the compound according to formulas XXXIV and XXXV, wherein J is phenyl; K is phenyl; L² is a bond; R¹⁵ is -H; m is 1, 2, or 3;

R⁵ is heterocyclyl optionally substituted with one or more R^{5a}; and R² is -L³-R⁷, wherein

L³ is a bond; and

R⁷ is hydrogen, halogen, -Z, or -Y-Z, wherein

Y is -[C(R¹⁵)₂]_m-, -(C₃-C₆)cycloalkyl-, or C₂-C₆alkenyl, ,

wherein each R¹⁵ is independently H, halogen, or (C₁-C₆)alkyl; and

Z is -H, halogen, -OR¹¹, -C(=O)R¹¹, -C(=O)OR¹¹, -C(=O)N(R¹¹)₂, -C(=N-OH)R¹¹, or -C(=S)N(R¹¹)₂.

In another embodiment, the invention provides the compound according to formulas XXXIV and XXXV, wherein J is phenyl; K is phenyl; L² is a bond; R¹⁵ is -H; m is 1, 2, or 3; R⁵ is heterocyclyl
5 optionally substituted with one or more R^{5a}; and each R^{5a} is independently halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, or C₁-C₆ haloalkyl.

In another embodiment, the invention provides the compound according to formula IIa-d, wherein J is aryl or heteroaryl.

In another embodiment, the invention provides the compound according to formula IIa-d,
10 wherein J is phenyl, pyridyl, thienyl, pyrrolyl, furanyl, pyrimidinyl, pyrazinyl, imidazolyl, pyrazolyl, oxazolyl, thiazolyl, isoxazolyl, isothiazolyl, triazolyl, triazinyl, tetrazolyl, or tetrazinyl.

In another embodiment, the invention provides the compound according to formula IIa-d, wherein J is phenyl, pyridyl, thienyl, pyrrolyl, or furanyl.

In another embodiment, the invention provides the compound according to formula IIa-d,
15 wherein J is phenyl, pyridyl, or thienyl.

In another embodiment, the invention provides the compound according to formula IIa-d, wherein J is phenyl.

In another embodiment, the invention provides the compound according to formula IIa-d, wherein J is pyridyl.

20 In another embodiment, the invention provides the compound according to formula IIa-d, wherein J is thienyl.

In another embodiment, the invention provides the compound according to formula IIa-d, wherein K is aryl or heteroaryl.

In another embodiment, the invention provides the compound according to formula IIa-d,
25 wherein K is phenyl, pyridyl, thienyl, pyrrolyl, furanyl, pyrimidinyl, pyrazinyl, imidazolyl, pyrazolyl, oxazolyl, thiazolyl, isoxazolyl, isothiazolyl, triazolyl, triazinyl, tetrazolyl, or tetrazinyl.

In another embodiment, the invention provides the compound according to formula IIa-d, wherein K is phenyl, pyridyl, thienyl, pyrrolyl, or furanyl.

In another embodiment, the invention provides the compound according to formula IIa-d,
30 wherein K is phenyl or pyridyl.

In another embodiment, the invention provides the compound according to formula IIa-d, wherein K is pyridyl.

In another embodiment, the invention provides the compound according to formula IIa-d, wherein K is phenyl.

In another embodiment, the invention provides the compound according to formula IIa-d, wherein L^2 is a bond.

In another embodiment, the invention provides the compound according to formula IIa-d, wherein J is aryl or heteroaryl; and K is aryl or heteroaryl.

5 In another embodiment, the invention provides the compound according to formula IIa-d, wherein J is aryl or heteroaryl; K is aryl or heteroaryl; and L^2 is a bond.

In another embodiment, the invention provides the compound according to formula IIa-d, wherein

10 J is phenyl, pyridyl, thienyl, pyrrolyl, furanyl, pyrimidinyl, pyrazinyl, imidazolyl, pyrazolyl, oxazolyl, thiazolyl, isoxazolyl, isothiazolyl, triazolyl, triazinyl, tetrazolyl, or tetrazinyl;

K is phenyl, pyridyl, thienyl, pyrrolyl, furanyl, pyrimidinyl, pyrazinyl, imidazolyl, pyrazolyl, oxazolyl, thiazolyl, isoxazolyl, isothiazolyl, triazolyl, triazinyl, tetrazolyl, or tetrazinyl; and

15 L^2 is a bond.

In another embodiment, the invention provides the compound according to formula IIa-d, wherein J is phenyl, pyridyl, thienyl, pyrrolyl, or furanyl; K is phenyl, pyridyl, thienyl, pyrrolyl, or furanyl; and L^2 is a bond.

20 In another embodiment, the invention provides the compound according to formula IIa-d, wherein J is phenyl, pyridyl, or thienyl; K is phenyl or pyridyl; and L^2 is a bond.

In another embodiment, the invention provides the compound according to formula IIa-d, wherein J is phenyl; K is phenyl; and L^2 is a bond.

In another embodiment, the invention provides the compound according to formula IIa-d, wherein J is pyridyl; K is phenyl; and L^2 is a bond.

25 In another embodiment, the invention provides the compound according to formula IIa-d, wherein J is thienyl; K is phenyl; and L^2 is a bond.

In another embodiment, the invention provides the compound according to formula IIa-d, wherein J is phenyl; K is pyridyl; and L^2 is a bond.

30 In another embodiment, the invention provides the compound according to formula IIa-d, wherein J is pyridyl; K is pyridyl; and L^2 is a bond.

In another embodiment, the invention provides the compound according to formula IIa-d, wherein J is thienyl; K is pyridyl; and L^2 is a bond.

In another embodiment, the invention provides the compound according to formula IIa-d, wherein R^5 is aryl, heterocyclyl, or heteroaryl, wherein R^5 is optionally substituted with one or more R^{5a} .

In another embodiment, the invention provides the compound according to formula IIa-d, wherein R^5 is aryl or heteroaryl, wherein R^5 is optionally substituted with one or more R^{5a} .

In another embodiment, the invention provides the compound according to formula IIa-d, wherein R^5 is phenyl, pyridyl, thienyl, pyrrolyl, furanyl, pyrimidinyl, pyrazinyl, imidazolyl, pyrazoyl, oxazolyl, thiazoyl, isoxazolyl, isothiazoyl, triazolyl, triazinyl, tetrazoyl, or tetrazinyl wherein R^5 is optionally substituted with one or more R^{5a} .

In another embodiment, the invention provides the compound according to formula IIa-d, wherein L^1 is a bond; and R^5 is phenyl, pyridyl, thienyl, pyrrolyl, furanyl, pyrimidinyl, pyrazinyl, imidazolyl, pyrazoyl, oxazolyl, thiazoyl, isoxazolyl, isothiazoyl, triazolyl, triazinyl, tetrazoyl, or tetrazinyl, wherein R^5 is optionally substituted with one or more R^{5a} .

In another embodiment, the invention provides the compound according to formula IIa-d, wherein L^1 is a bond; and R^5 is phenyl, pyridyl, thienyl, pyrrolyl, or furanyl, wherein R^5 is optionally substituted with one or more R^{5a} .

In another embodiment, the invention provides the compound according to formula IIa-d, wherein L^1 is a bond; and R^5 is phenyl optionally substituted with one or more R^{5a} .

In another embodiment, the invention provides the compound according to formula IIa-d, wherein L^1 is a bond; and R^5 is pyridyl optionally substituted with one or more R^{5a} .

In another embodiment, the invention provides the compound according to formula IIa-d, wherein L^1 is a bond; and R^5 is thienyl optionally substituted with one or more R^{5a} .

In another embodiment, the invention provides the compound according to formula IIa-d wherein R^{21} is hydrogen, halogen, nitro, cyano, aryl, heteroaryl, heterocyclyl, $-C_1-C_6$ alkyl-heterocyclyl, $-C_1-C_6$ alkyl-heteroaryl, $-C_1-C_6$ alkyl-aryl, $-Z$, $-Y-Z$, or $-X-Y-Z$, wherein

X is $-O-$;

Y is $-[C(R^{15})_2]_m-$, $-C_2-C_6$ alkenyl, or C_3-C_8 cycloalkyl; and

Z is $-H$, $-CN$, halogen, $-OR^{11}$, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)N(R^{11})_2$, $-N(R^{11})_2$, $-CN$, $-N_3$, $-SO_2R^{11}$, $-S(=O)_2N(R^{11})_2$, $-C(=O)N(R^{11})N(R^{11})_2$, $-C(=O)N(R^{11})(OR^{11})$, $-OC(=O)-R^{11}$, $-OC(=O)-N(R^{11})_2$, or $-N(R^{11})COOR^{11}$.

In another embodiment, the invention provides the compound according to formula IIa-d wherein R^{21} is hydrogen, halogen, nitro, cyano, C_1-C_6 alkyl, or C_1-C_6 haloalkyl.

In another embodiment, the invention provides the compound according to formula IIa-d wherein R^3 is hydrogen, aryl, heteroaryl, heterocyclyl, $-C_1-C_6$ alkyl-heterocyclyl, $-C_1-C_6$ alkyl-heteroaryl, $-C_1-C_6$ alkyl-aryl, $-Z$, or $-Y-Z$ wherein

Y is $-[C(R^{15})_2]_m-$, $-C_2-C_6$ alkenyl, or C_3-C_8 cycloalkyl; and

Z is -H, -CN, halogen, -OR¹¹, -C(=O)R¹¹, -C(=O)OR¹¹, -C(=O)N(R¹¹)₂, -N(R¹¹)₂, -CN, -N₃, -SO₂R¹¹, -S(=O)₂N(R¹¹)₂, -C(=O)N(R¹¹)N(R¹¹)₂, -C(=O)N(R¹¹)(OR¹¹), -OC(=O)-R¹¹, -OC(=O)-N(R¹¹)₂, or -N(R¹¹)COOR¹¹.

In another embodiment, the invention provides the compound according to formula IIa-d
5 wherein R³ is hydrogen, C₁-C₆alkyl, or C₁-C₆haloalkyl.

In another embodiment, the invention provides the compound according to any of the formulas IIa-d, IIIa-d, IVa-d, IV-XXIV, and XXVI-XXXV wherein R² is -L³-R⁷, wherein

L³ is a bond; and

R⁷ is hydrogen, halogen, -Z, or -Y-Z, wherein

10 Y is -[C(R¹⁵)₂]_m, -(C₃-C₆)cycloalkyl-, or C₂-C₆alkenyl,
wherein each R¹⁵ is independently H, halogen, or (C₁-C₆)alkyl; and
Z is -H, halogen, -OR¹¹, -C(=O)R¹¹, -C(=O)OR¹¹, -C(=O)N(R¹¹)₂,
-C(=N-OH)R¹¹, or -C(=S)N(R¹¹)₂.

In another embodiment, the invention provides the compound according to any of the formulas
15 IIa-d, IIIa-d, IVa-d, IV-XXIV, and XXVI-XXXV, wherein R² is -L³-R⁷, wherein

L³ is a bond; and

R⁷ is hydrogen, halogen, or -[C(R¹⁵)₂]-Z, wherein

each R¹⁵ is independently H, halogen, or (C₁-C₂)alkyl; and

20 Z is -H, halogen, -OR^{11''}, -C(=O)R^{11''}, -C(=O)OR^{11''}, -C(=O)N(R^{11''})₂,
-C(=N-OH)R^{11''}, or -C(=S)N(R^{11''})₂,
wherein R^{11''} is -H or -(C₁-C₆ alkyl).

In another embodiment, the invention provides the compound according to any of the formulas IIa-d, IIIa-d, IVa-d, IV-XXIV, and XXVI-XXXV, wherein R² is -halogen, -CF₃, -CH₂OH, -CH₂SO₂Me, -C(CH₃)₂OH, or -C(CH₃)₂SO₂Me.

25 In another embodiment, the invention provides the compound according to any of the formulas IIa-d, IIIa-d, IVa-d, IV-XXIV, and XXVI-XXXV, wherein R² is -halogen, -CF₃, -CH₂OH, or -C(CH₃)₂OH.

In another embodiment, the invention provides the compound according to any of the formulas IIa-d, IIIa-d, IVa-d, IV-XXIV, and XXVI-XXXV, wherein R² is -CF₃ or -C(CH₃)₂OH.

30 In another embodiment, the invention provides the compound according to any of the formulas IIa-d, IIIa-d, IVa-d, IV-XXIV, and XXVI-XXXV, wherein

each R⁴ is independently halogen, aryl, heteroaryl, heterocyclyl, -G¹ or -E-G¹, wherein

E is -[C(R¹⁵)₂]_m, wherein

each R¹⁵ is independently hydrogen or halogen; and

G^1 is $-C_1-C_6$ alkyl, $-C_1-C_6$ haloalkyl, halogen, $-OR^{11}$, or $-SO_2R^{11}$.

In another embodiment, the invention provides the compound according to any of the formulas IIa-d, IIIa-d, IVa-d, IV-XXIV, and XXVI-XXXV, wherein

each R^4 is independently halogen, $-CH_2-G^1$, $-C(H)(F)-G^1$, $-CF_2-G^1$, wherein

5 G^1 is $-C_1-C_6$ alkyl, $-C_1-C_6$ haloalkyl, $-F$, $-OR^{11'}$, or $-SO_2R^{11'}$

wherein $R^{11'}$ is $-H$ or $-C_1-C_6$ alkyl.

In another embodiment, the invention provides the compound according to any of the formulas IIa-d, IIIa-d, IVa-d, IV-XXIV, and XXVI-XXXV, wherein each R^4 is independently $-CH_3$, $-CF_3$, $-CF_2H$, $-CH_2F$, $-OH$, $-OMe$, $-CH_2OH$, or $-SO_2(C_1-C_3$ alkyl).

10 In one embodiment, the invention provides the compound according to any of the formulas IIa-d, IIIa-d, IVa-d, IV-XXIV, and XXVI-XXXV, wherein

each R^{41} is independently halogen, $-G^1$ or $-E-G^1$, wherein

E is $-[C(R^{15})_2]_m$,

wherein each R^{15} is independently hydrogen or halogen; and

15 G^1 is $-C_1-C_6$ alkyl, $-C_1-C_6$ haloalkyl, or halogen.

In another embodiment, the invention provides the compound according to formulas IIa-d, IIIa-d, IVa-d, IV-XXIV, and XXVI-XXXV, wherein wherein each R^{41} is independently halogen, methyl or trifluoromethyl.

In another embodiment, the invention provides the compound according to formulas to formulas IIa-d, IIIa-d, IVa-d, IV-XXIV, and XXVI-XXXV, wherein R^{21} is hydrogen, halogen, nitro, cyano, aryl, heteroaryl, heterocyclyl, $-C_1-C_6$ alkyl-heterocyclyl, $-C_1-C_6$ alkyl-heteroaryl, $-C_1-C_6$ alkyl-aryl, $-Z$, $-Y-Z$, or $-X-Y-Z$, wherein

X is $-O-$;

Y is $-[C(R^{15})_2]_m$, $-C_2-C_6$ alkenyl, or C_3-C_8 cycloalkyl;

25 Z is $-H$, $-CN$, halogen, $-OR^{11}$, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)N(R^{11})_2$, $-N(R^{11})_2$, $-CN$, $-N_3$, $-SO_2R^{11}$, $-S(=O)_2N(R^{11})_2$, $-C(=O)N(R^{11})N(R^{11})_2$, $-C(=O)N(R^{11})(OR^{11})$, $-OC(=O)-R^{11}$, $-OC(=O)-N(R^{11})_2$, or $-N(R^{11})COOR^{11}$;

In another embodiment, the invention provides the compound according to formulas IIa-d, IIIa-d, IVa-d, IV-XXIV, and XXVI-XXXV, wherein R^{21} is hydrogen, halogen, nitro, cyano, C_1-C_6 alkyl, or C_1-C_6 haloalkyl.

30 In another embodiment, the invention provides the compound according to formula IIa-d, wherein R^1 is $-L^1-R^5$, wherein L^1 is a bond, $-[C(R^{15})_2]_m$, or $-C_3-C_8$ cycloalkyl-; and

R^5 is phenyl or pyridyl, each optionally substituted with one or two R^{5a} , wherein

each R^{5a} is independently -halogen, -C₁-C₆ alkoxy, -C₁-C₆ alkyl, or -C₁-C₆ haloalkyl;

R^2 , R^{21} , and R^3 are each independently -H, $-[C(R^{15})_2]_m-OH$, -C₁-C₆ alkyl, -C₁-C₆ haloalkyl, halogen, $-C(O)N(R^{11})_2$, or $-COOR^{11}$;

5 L^2 is a bond;

J is phenyl, pyridyl, or thienyl;

K is phenyl, or pyridyl;

each R^{41} is -halogen, -C₁-C₆ alkyl, or C₁-C₆ haloalkyl; and

10 each R^4 is -halogen, $-[C(R^{15})_2]_m-OH$, $-SO_2R^{11}$, $-SO_2N(R^{11})_2$, $-C(O)N(R^{11})_2$, $-COOR^{11}$, -C₁-C₆ alkyl, or -C₁-C₆ haloalkyl.

In another embodiment, the invention provides the compound according to formula IVa-d, wherein

R^1 is $-L^1-R^5$, wherein

L^1 is a bond, $-[C(R^{15})_2]_m$, or -C₃-C₈ cycloalkyl; and

15 R^5 is phenyl or pyridyl, each optionally substituted with one or two R^{5a} , wherein

each R^{5a} is independently -halogen, -C₁-C₆ alkoxy, -C₁-C₆ alkyl, or -C₁-C₆ haloalkyl;

R^2 , R^{21} , and R^3 are each independently -H, $-[C(R^{15})_2]_m-OH$, -C₁-C₆ alkyl, -C₁-C₆ haloalkyl, halogen, $-C(O)N(R^{11})_2$, or $-COOR^{11}$;

20 L^2 is a bond;





J is phenyl, pyridyl, or thienyl;

K is phenyl, or pyridyl;

each R^{41} is -halogen, -C₁-C₆ alkyl, or C₁-C₆ haloalkyl; and

25 each R^4 is -halogen, $-[C(R^{15})_2]_m-OH$, $-SO_2R^{11}$, $-SO_2N(R^{11})_2$, $-C(O)N(R^{11})_2$, $-COOR^{11}$, -C₁-C₆ alkyl, or -C₁-C₆ haloalkyl.

In a second aspect, the invention provides intermediate compounds according to one of the formulas XXVa-d,



or a pharmaceutically acceptable salt, isomer, or prodrug thereof, wherein,

R¹ is -L¹-R⁵, wherein

L¹ is a bond, L⁵, L⁶, -L⁵-L⁶-L⁵-, or -L⁶-L⁵-L⁶-, wherein

each L⁵ is independently -[C(R¹⁵)₂]_m-, wherein

each R¹⁵ is independently hydrogen, halogen, (C₁-C₆)alkyl, or (C₁-C₆)haloalkyl; and

5 each L⁶ is independently -CS-, -CO-, -SO₂-, -O-, -CON(R¹¹)-, -CONR¹¹N(R¹¹)-,
-C(=NR¹¹)-, -C(=NOR¹¹)-, or -C(=NN(R¹¹)₂)-, aryl-, -C₃-C₈cycloalkyl-, heteroaryl-, or
-heterocyclyl-;

wherein the aryl, cycloalkyl, heteroaryl, or heterocyclyl is optionally substituted
with one or more R¹⁴;

10 or each L⁶ is independently C₂-C₆ alidiyl,

wherein the alidiyl chain is optionally interrupted by -C(R¹¹)₂-,
-C(R¹¹)₂C(R¹¹)₂-, -C(R¹¹)=C(R¹¹)-, -C(R¹¹)₂O-, -C(R¹¹)₂NR¹¹-, -C≡C-, -O-,
-S-, -N(R¹⁰)CO-, -N(R¹⁰)CO₂-, -CON(R¹⁰)-, -CO-, -CO₂-, -OC(=O)-,
-OC(=O)N(R¹⁰)-, -SO₂-, -N(R¹⁰)SO₂-, or -SO₂N(R¹⁰); and

15 R⁵ is aryl, heterocyclyl, heteroaryl, -C, or -B-C,

wherein

B is -[C(R¹⁵)₂]_m- or -C₃-C₈ cycloalkyl-; and

C is halogen, -C₁-C₆alkyl, -C₁-C₆haloalkyl, -SO₂R¹¹, -SR¹¹, -SO₂N(R¹¹)₂, -SO₂NR¹¹COR¹¹,
-C≡N, -C(O)OR¹¹, -CON(R¹¹)₂, or -N(R¹¹)₂,

20 wherein R⁵ is optionally substituted with one or more R^{5a}, wherein

each R^{5a} is independently halogen, nitro, heteroaryl, heterocyclyl, C₂-C₆ alkenyl, C₂-C₆
alkynyl, C₃-C₈ cycloalkyl, (C₃-C₈ cycloalkyl)-C₁-C₆ alkyl-, (C₃-C₈ cycloalkenyl)-C₁-C₆ alkyl-,
(C₃-C₈ cycloalkyl)-C₂-C₆ alkenyl-, aryl, arylalkyl, aryloxy, aryloxyaryl, arylC₁₋₆ alkoxy,
C₁-C₆ alkyl, C₁-C₆ haloalkyl, SO₂R¹¹, OR¹¹, SR¹¹, N₃, SO₂R¹¹, COR¹¹, SO₂N(R¹¹)₂,
25 SO₂NR¹¹COR¹¹, C≡N, C(O)OR¹¹, CON(R¹¹)₂, CON(R¹¹)OR¹¹, OCON(R¹¹)₂,
NR¹¹COR¹¹, NR¹¹CON(R¹¹)₂, NR¹¹COOR¹¹, or N(R¹¹)₂, wherein

each R^{5a} is optionally substituted with one or more groups which
independently are -halogen, -C₁-C₆ alkyl, aryloxy C₀₋₆ alkylSO₂R¹¹, C₀₋₆
alkylCOOR¹¹, C₀₋₆ alkoxyaryl, C₁-C₆ haloalkyl, -SO₂R¹¹, -OR¹¹, -SR¹¹, -N₃,
30 -SO₂R¹¹, -COR¹¹, -SO₂N(R¹¹)₂, -SO₂NR¹¹COR¹¹, -C≡N, -C(O)OR¹¹,
-CON(R¹¹)₂, -CON(R¹¹)OR¹¹, -OCON(R¹¹)₂, -NR¹¹COR¹¹, -NR¹¹CON(R¹¹)₂,
-NR¹¹COOR¹¹, or -N(R¹¹)₂;

R² and R²¹ are -L³-R⁷, wherein

each L^3 is independently a bond or $-(CH_2)_m-V^1-(CH_2)_n-$ wherein

n is 0-6; and

V^1 is $-C(R^{11})_2-$, $-C(R^{11})_2C(R^{11})_2-$, $-C(R^{11})=C(R^{11})-$, $-C(R^{11})_2O-$, $-C(R^{11})_2NR^{11}-$, $-C\equiv C-$,
-O-, -S-, $-NR^{11}-$, $-N(R^{10})CO-$, $-N(R^{10})CO_2-$, $-OCO-$, $-CO-$, $-CS-$, $-CONR^{10}-$,
-C(=N- R^{11})-, $-C(=N-OR^{11})-$, $-C[=N-N(R^{11})_2]$, $-CO_2-$, $-OC(=O)-$, $-OC(=O)N(R^{10})-$,
-SO₂-, $-N(R^{10})SO_2-$, $-SO_2N(R^{10})-$, $-NR^{10}CONR^{10}-$, $-NR^{10}CSNR^{10}-$, C₃-C₆cyclo alkyl,
or C₃-C₆ cyclohaloalkyl;

or each L^3 is independently C₂-C₆ alidiyl,

wherein the alidiyl chain is optionally interrupted by $-C(R^{11})_2-$, $-C(R^{11})_2C(R^{11})_2-$,
-C(=N- R^{11})-, $-C(R^{11})_2O-$, $-C(R^{11})_2NR^{11}-$, $-C\equiv C-$, -O-, -S-, $-N(R^{10})CO-$,
-N(R¹⁰)CO₂-, $-NR^{11}-$, $-CON(R^{10})-$, $-CO-$, $-CO_2-$, $-OC(=O)-$, $-OC(=O)N(R^{10})-$, $-SO_2-$,
-N(R¹⁰)SO₂-, or $-SO_2N(R^{10})-$; and

each R^7 is independently hydrogen, halogen, nitro, aryl, heteroaryl, heterocyclyl, -Z-, -Y-Z, or -
X-Y-Z, wherein

X is -O-;

Y is $-[C(R^{15})_2]_m-$ or C₂-C₆alkenyl, C₃-C₈cycloalkyl; and

Z is -H, halogen, $-OR^{11}$, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)N(R^{11})_2$, $-N(R^{11})_2$,
-C(=N-OH) R^{11} , $-C(=S)N(R^{11})_2$, -CN, $-S(=O)_2N(R^{11})_2$, $-C(=O)N(R^{11})N(R^{11})_2$,
 $-C(=O)N(R^{11})(OR^{11})$, $-OC(=O)-R^{11}$, or $-OC(=O)-N(R^{11})_2$,

wherein R^7 is optionally substituted with one or more R^{7a} , wherein

R^{7a} is halogen, haloaryl, aryloxy, aralkyloxy, aryloxyalkyl, arylC₀-C₆ alkylcarboxy,
 $C(R^{11})=C(R^{11})-COOH$, aryl, heteroaryl, heterocyclyl, heterocycliloxy, heteroaryloxy,
-Z', -Y'-Z', or -X'-Y'-Z', wherein

X' is -O-;

Y' is $-[C(R^{15})_2]_m-$ or C₃-C₈cycloalkyl; and

Z' is -H, halogen, $-OR^{11}$, $-SR^{11}$, $-S(=O)_2R^{11}$, $-C(=O)R^{11}$, $-C(=O)OR^{11}$,
 $-C(=O)N(R^{11})_2$, $-N(R^{11})_2$, $-N(R^{11})C(=O)R^{11}$, $-S(=O)_2N(R^{11})C(=O)R^{11}$, -CN,
 $-S(=O)_2N(R^{11})_2$, $-C(=O)N(R^{11})N(R^{11})_2$, $-C(=O)N(R^{11})(OR^{11})$, $-OC(=O)-R^{11}$,
 $-OC(=O)-OR^{11}$, $-N(R^{11})C(=O)-R^{11}$, or $-N(R^{11})S(=O)_2R^{11}$,

wherein each R^{7a} is optionally substituted with one or more R^8 ,

wherein each R^8 is independently halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkyl,
C₁-C₆ haloalkyl(OR^{11}), C₀-C₆ alkyl OR^{11} , C₀-C₆ alkyl $CON(R^{11})_2$, C₀-C₆ alkyl COR^{11} ,
C₀-C₆ alkyl $COOR^{11}$, or C₀-C₆ alkyl SO_2R^{11} ,

R³ is -L-R⁶, wherein

n is 0-6; each w is independently 0–5; and

Y³ is -O-, -S-, -NR⁷-, -N(R¹⁰)CO-, -N(R¹⁰)CO₂-, -OCO-, -OC(=O)N(R¹⁰)-, -NR¹⁰CONR¹⁰-, -N(R¹⁰)SO₂-, or -NR¹⁰CSNR¹⁰-;

R^6 is C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, aryl, C_3 - C_8 cycloalkyl, heteroaryl, heterocyclyl, $-CN$, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)N(R^{11})_2$, $-N(R^{11})_2$, $-SO_2R^{11}$, $-S(=O)_2N(R^{11})_2$, $-C(=O)N(R^{11})N(R^{11})_2$, or $-C(=O)N(R^{11})(OR^{11})$, wherein

each R^{6a} is independently -Z'', -Y''-Z'', or -X''-Y''-Z'', wherein

Y" is -[C(R¹⁵)₂]_m-, -C₂-C₆ alkenyl, C₃-C₈ cycloalkyl, heterocyclyl, aryl, or heteroaryl, wherein

25 Z'' is -H, -CN, halogen, $-OR^{11}$, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)N(R^{11})_2$, $-N(R^{11})_2$, $-N_3$, $-SO_2R^{11}$, $-S(=O)_2N(R^{11})_2$, $-C(=O)N(R^{11})N(R^{11})_2$, $-N(R^{11})C(=O)N(R^{11})_2$, $-OC(=O)-OR^{11}$, $-C(=O)N(R^{11})(OR^{11})$, $-OC(=O)-R^{11}$, $-OC(=O)-N(R^{11})_2$, or $-N(R^{11})COOR^{11}$; and

Hal is halogen:

J is aryl or heteroaryl;

each R^{41} is independently -halogen, nitro, $CR^{11}=CR^{11}COOR^{11}$, aryloxy, aralkyloxy, aryloxyalkyl, aryl- C_0 - C_6 alkylcarboxy, aryl, heteroaryl, heterocyclyl, heteroaryloxy, heterocyclyloxy, $-G^1-E-G^1$ or $-D-E-G^1$, wherein

5 D is -O-;

E is $-[C(R^{15})_2]_m-$ or $-C_3$ - C_8 cycloalkyl; and

G^1 is $-C_1$ - C_6 alkyl, $-C_1$ - C_6 haloalkyl, $-COR^{11}$, $-COOR^{11}$, $-CON(R^{11})_2$, $-C\equiv N$, $-OR^{11}$, $-OCON(R^{11})_2$, $-OCOOR^{11}$, $-N_3$, $-NR^{11}COR^{11}$, $-NR^{11}SO_2R^{11}$, $-N(R^{11})_2$, $-NR^{11}COOR^{11}$, $-SOR^{11}$, $-SO_2R^{11}$, $-SO_2NR^{11}COR^{11}$, $-SO_2N(R^{11})_2$, or $-SR^{11}$;

10 wherein each R^4 is optionally substituted with one or more R^{4a} ,

wherein each R^{4a} is independently halogen, aryloxy, aralkyloxy, aryloxyalkyl, C_1 - C_6 alkoxyaryl, aryl- C_0 - C_6 alkylcarboxy, $-G'$, $-E'-G'$, or $-D'-E'-G'$, wherein

D' is -O-;

E' is $-[C(R^{15})_2]_m-$ or $-C_3$ - C_8 cycloalkyl; and

15 G' is $-H$, -halogen, $-COR^{11}$, $-COOR^{11}$, $-C\equiv N$, $-OR^{11}$, $-NR^{11}SO_2R^{11}$, $-SO_2R^{11}$, $-SO_2N(R^{11})_2$, or $-SR^{11}$;

each m is 0, 1, 2, 3, 4, 5, or 6; and

q' is 0, 1, 2, 3, or 4,

each R^{10} is independently $-R^{11}$, $-C(=O)R^{11}$, $-CO_2R^{11}$, or $-SO_2R^{11}$;

20 each R^{11} is independently -hydrogen, $-C_1$ - C_6 alkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, C_3 - C_8 cycloalkyl, $(C_3$ - C_8 cycloalkyl)- C_1 - C_6 alkyl-, $(C_3$ - C_8 cycloalkenyl)- C_1 - C_6 alkyl-, $(C_3$ - C_8 cycloalkyl)- C_2 - C_6 alkenyl-, $-C_1$ - C_6 haloalkyl, $-N(R^{12})_2$, aryl, $-(C_1$ - C_6)alkyl-aryl, heteroaryl, $-(C_1$ - C_6)alkyl-heteroaryl, heterocyclyl, or $-(C_1$ - C_6)alkyl-heterocyclyl,

wherein any of R^{11} is optionally substituted with one or more radicals of R^{12} ;

25 each R^{12} is independently halogen, C_1 - C_6 haloalkyl, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, $(C_0$ - C_6 alkyl) $C=O(OR^{13})$, C_0 - C_6 alkyl OR^{13} , C_0 - C_6 alkyl COR^{13} , C_0 - C_6 alkyl SO_2R^{13} , C_0 - C_6 alkyl $CON(R^{13})_2$, C_0 - C_6 alkyl $CONR^{13}OR^{13}$, C_0 - C_6 alkyl $SO_2N(R^{13})_2$, C_0 - C_6 alkyl SR^{13} , C_0 - C_6 haloalkyl OR^{13} , aryloxy, aralkyloxy, aryloxyalkyl, C_0 - C_6 alkoxyaryl, aryl- C_0 - C_6 alkylcarboxy, $-C_0$ - C_6 alkyl $IN(R^{13})_2$, $-NR^{13}SO_2R^{13}$, or $-OC_0$ - C_6 alkyl $COOR^{13}$;

30 each R^{13} is independently hydrogen, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, C_3 - C_8 cycloalkyl, $(C_3$ - C_8 cycloalkyl)- C_1 - C_6 alkyl-, $(C_3$ - C_8 cycloalkenyl)- C_1 - C_6 alkyl-, or $(C_3$ - C_8 cycloalkyl)- C_2 - C_6 alkenyl-;

each R^{14} is independently C_1 - C_6 alkyl, C_1 - C_6 alkoxy, halogen, C_1 - C_6 haloalkyl, C_0 - C_6 alkyl $CON(R^{11})_2$, C_0 - C_6 alkyl $CONR^{11}OR^{11}$, C_0 - C_6 alkyl OR^{11} , or C_0 - C_6 alkyl $COOR^{11}$.

In another embodiment, the invention provides the compound according to formula XXVa-d, wherein Hal is -Cl, -Br, or -I.

In another embodiment, the invention provides the compound according to formula XXVa-d, wherein Hal is -Cl.

5 In another embodiment, the invention provides the compound according to formula XXVa-d, wherein Hal is -Br.

In another embodiment, the invention provides the compound according to formula XXVa-d, wherein Hal is -I.

10 In another embodiment, the invention provides the compound according to formula XXVa-d, wherein J is aryl or heteroaryl.

In another embodiment, the invention provides the compound according to formula XXVa-d, wherein J is phenyl, pyridyl, thienyl, pyrrolyl, furanyl, pyrimidinyl, pyrazinyl, imidazolyl, pyrazolyl, oxazolyl, thiazolyl, isoxazolyl, isothiazolyl, triazolyl, triazinyl, tetrazolyl, or tetrazinyl.

15 In another embodiment, the invention provides the compound according to formula XXVa-d, wherein J is phenyl, pyridyl, thienyl, pyrrolyl, or furanyl.

In another embodiment, the invention provides the compound according to formula XXVa-d, wherein J is phenyl, pyridyl, or thienyl.

In another embodiment, the invention provides the compound according to formula XXVa-d, wherein J is phenyl.

20 In another embodiment, the invention provides the compound according to formula XXVa-d, wherein J is pyridyl.

In another embodiment, the invention provides the compound according to formula XXVa-d, wherein J is thienyl.

25 In another embodiment, the invention provides the compound according to formula XXVa-d, wherein R⁵ is aryl, heterocyclyl, or heteroaryl, wherein R⁵ is optionally substituted with one or more R^{5a}.

In another embodiment, the invention provides the compound according to formula XXVa-d, wherein R⁵ is aryl or heteroaryl, wherein R⁵ is optionally substituted with one or more R^{5a}.

30 In another embodiment, the invention provides the compound according to formula XXVa-d, wherein R⁵ is phenyl, pyridyl, thienyl, pyrrolyl, furanyl, pyrimidinyl, pyrazinyl, imidazolyl, pyrazolyl, oxazolyl, thiazolyl, isoxazolyl, isothiazolyl, triazolyl, triazinyl, tetrazolyl, or tetrazinyl wherein R⁵ is optionally substituted with one or more R^{5a}.

In another embodiment, the invention provides the compound according to formula XXVa-d, wherein L¹ is a bond.

In another embodiment, the invention provides the compound according to formula XXVa-d, wherein L^1 is a bond; and R^5 is phenyl, pyridyl, thienyl, pyrrolyl, furanyl, pyrimidinyl, pyrazinyl, imidazolyl, pyrazoyl, oxazolyl, thiazoyl, isoxazolyl, isothiazoyl, triazolyl, triazinyl, tetrazoyl, or tetrazinyl, wherein R^5 is optionally substituted with one or more R^{5a} .

5 In another embodiment, the invention provides the compound according to formula XXVa-d, wherein L^1 is a bond; and R^5 is phenyl, pyridyl, thienyl, pyrrolyl, or furanyl, wherein R^5 is optionally substituted with one or more R^{5a} .

In another embodiment, the invention provides the compound according to formula XXVa-d, wherein L^1 is a bond; and R^5 is phenyl optionally substituted with one or more R^{5a} .

10 In another embodiment, the invention provides the compound according to formula XXVa-d, wherein L^1 is a bond; and R^5 is pyridyl optionally substituted with one or more R^{5a} .

In another embodiment, the invention provides the compound according to formula XXVa-d, wherein L^1 is a bond; and R^5 is thienyl optionally substituted with one or more R^{5a} .

In another embodiment, the invention provides the compound according to formulas XXVa-d, 15 wherein R^2 is $-L^3-R^7$, wherein L^3 is a bond; and

R^7 is hydrogen, halogen, $-Z$, or $-Y-Z$, wherein

Y is $-[C(R^{15})_2]_m-$, $-(C_3-C_6)\text{cycloalkyl}-$, or $C_2-C_6\text{alkenyl}$,

wherein each R^{15} is independently H, halogen, or $(C_1-C_6)\text{alkyl}$; and

Z is $-H$, halogen, $-OR^{11}$, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)N(R^{11})_2$, $-C(=N-OH)R^{11}$, or 20 $-C(=S)N(R^{11})_2$.

In another embodiment, the invention provides the compound according to formulas XXVa-d, wherein R^2 is $-L^3-R^7$, wherein L^3 is a bond; and

R^7 is hydrogen, halogen, or $-[C(R^{15})_2]-Z$, wherein

each R^{15} is independently H, halogen, or $(C_1-C_2)\text{alkyl}$; and

25 Z is $-H$, halogen, $-OR^{11''}$, $-C(=O)R^{11''}$, $-C(=O)OR^{11''}$, $-C(=O)N(R^{11''})_2$, $-C(=N-OH)R^{11''}$, or $-C(=S)N(R^{11''})_2$,

wherein $R^{11''}$ is $-H$ or $-(C_1-C_6)\text{alkyl}$.

In another embodiment, the invention provides the compound according to formulas XXVa-d, wherein R^2 is $-\text{halogen}$, $-\text{CF}_3$, $-\text{CH}_2\text{OH}$, $-\text{CH}_2\text{SO}_2\text{Me}$, $-\text{C}(\text{CH}_3)_2\text{OH}$, or $-\text{C}(\text{CH}_3)_2\text{SO}_2\text{Me}$.

30 In another embodiment, the invention provides the compound according to formulas XXVa-d, wherein R^2 is $-\text{halogen}$, $-\text{CF}_3$, $-\text{CH}_2\text{OH}$, or $-\text{C}(\text{CH}_3)_2\text{OH}$.

In another embodiment, the invention provides the compound according to formulas XXVa-d, wherein R^2 is $-\text{CF}_3$ or $-\text{C}(\text{CH}_3)_2\text{OH}$.

In one embodiment, the invention provides the compound according to formulas XXVa-d, wherein each R^{41} is independently halogen, $-G^1$ or $-E-G^1$, wherein

E is $-[C(R^{15})_2]_m-$,

wherein each R^{15} is independently hydrogen or halogen; and

5 G^1 is $-C_1-C_6$ alkyl, $-C_1-C_6$ haloalkyl, or halogen.

In another embodiment, the invention provides the compound according to formula XXVa-d, wherein R^1 is L^1-R^5 , wherein L^1 is a bond; and

R^5 is phenyl or pyridyl, each optionally substituted with one or two R^{5a} , wherein

each R^{5a} is independently halogen, $-CH_3$, or $-CF_3$;

10 R^2 is $-H$, $-C(R^{20})_2OH$, $-CH_3$, $-CF_3$, or halogen, wherein

each R^{20} is independently $-H$, $-F$, $-CH_3$, or $-CF_3$;

J is phenyl, pyridyl, or thienyl; and

each R^{41} is halogen, $-CH_3$, $-CH_2CH_3$, $-CF_3$, $-CF_2CF_3$, or $-CH_2CF_3$.

In another embodiment, the invention provides the compound according to formula XXVa-d, 15 wherein q' is 0 or 1; R^1 is L^1-R^5 , wherein L^1 is a bond; and

R^5 is phenyl optionally substituted with one or two R^{5a} , wherein

each R^{5a} is independently halogen, $-CH_3$, or $-CF_3$;

each R^2 is $-H$, $-C(R^{20})_2OH$, $-CH_3$, $-CF_3$, or halogen, wherein

each R^{20} is independently $-H$, $-F$, $-CH_3$, or $-CF_3$; and

20 R^{41} is halogen, $-CH_3$, $-CH_2CH_3$, $-CF_3$, $-CF_2CF_3$, or $-CH_2CF_3$.

In another embodiment, the invention provides the compound according to formulas XXVa-d, wherein wherein each R^{41} is independently halogen, methyl or trifluoromethyl. In another embodiment, the invention provides the compound according any of the previous embodiments wherein R^{21} is hydrogen.

25 In a third aspect, the invention provides a pharmaceutical composition comprising a compound of any of formulas IIa-d, IIIa-d, IVa-d, XXVa-d, or IV-XXXV, or a pharmaceutically acceptable derivative thereof, in a pharmaceutically acceptable carrier.

In another embodiment, the invention provides a pharmaceutical composition comprising a compound of formula IV, or a pharmaceutically acceptable derivative thereof, in a pharmaceutically 30 acceptable carrier.

In another embodiment, the invention provides a pharmaceutical composition comprising a compound of formula V, or a pharmaceutically acceptable derivative thereof, in a pharmaceutically acceptable carrier.

In another embodiment, the invention provides a pharmaceutical composition comprising a compound of formula VI, or a pharmaceutically acceptable derivative thereof, in a pharmaceutically acceptable carrier

5 In another embodiment, the invention provides a pharmaceutical composition comprising a compound of formula VII, or a pharmaceutically acceptable derivative thereof, in a pharmaceutically acceptable carrier.

In another embodiment, the invention provides a pharmaceutical composition comprising a compound of formula IIa-d, or a pharmaceutically acceptable derivative thereof, in a pharmaceutically acceptable carrier.

10 In a fourth aspect, the invention provides a kit, comprising a packaging material and a compound of any of formula IIa-d, IIIa-d, IVa-d, XXVa-d, or IV-XXXV, or a pharmaceutically acceptable derivative thereof, which is effective for modulating the activity of a nuclear receptor or for treatment, prevention, inhibition, or amelioration of one or more symptoms of nuclear receptor mediated diseases or disorders.

15 In another embodiment, the invention provides a kit, comprising a packaging material, and a compound of formula IIa-d, or a pharmaceutically acceptable derivative thereof, which is effective for modulating the activity of a nuclear receptor or for treatment, prevention, inhibition, or amelioration of one or more symptoms of nuclear receptor mediated diseases or disorders.

20 In another embodiment, the invention provides a kit, comprising a packaging material, a compound of formula IIa-d, or a pharmaceutically acceptable derivative thereof, which is effective for modulating the activity of a nuclear receptor or for treatment, prevention, inhibition, or amelioration of one or more symptoms of nuclear receptor mediated diseases or disorders, further comprising a label that indicates that the compound of formula IIa-d, or pharmaceutically acceptable derivative thereof, is used for modulating the activity of a nuclear receptor or for treatment, prevention or amelioration of one
25 or more symptoms of nuclear receptor mediated diseases or disorders, or diseases or disorders in which nuclear receptor activity is implicated.

In a sixth aspect, the invention provides a method of treating, preventing, inhibiting, or ameliorating the symptoms of a disease or disorder that is modulated or otherwise affected by nuclear receptor activity or in which nuclear receptor activity is implicated, comprising administering to a
30 subject in need thereof a therapeutically effective amount of a compound of any of formula IIa-d, IIIa-d, IVa-d, XXVa-d, or IV-XXXV.

In a preferred embodiment of the sixth aspect, the invention provides a method of treating, preventing, inhibiting, or ameliorating the symptoms of a disease or disorder that is modulated or otherwise affected by nuclear receptor activity or in which nuclear receptor activity is implicated,

comprising administering to a subject in need thereof a therapeutically effective amount of a compound according to part (A) of formulas IIa-d.

When part (A) of formulas IIa-d is referenced herein with respect to methods of using compounds of the invention, such as for treatment, prevention, inhibition, or amelioration of disease, or
5 for use in preparation of a medicament for the treatment, prevention, or amelioration of disease, it is meant that all compounds defined by part (A) are included and the provisos of part (B) of the same formulas are not to be considered when determining the scope of the compounds defined for the uses therein.

In a preferred embodiment of the sixth aspect, the invention provides the method wherein the
10 disease or disorder is hypercholesterolemia, hyperlipoproteinemia, hypertriglyceridemia, lipodystrophy, hyperglycemia, diabetes mellitus, dyslipidemia, atherosclerosis, gallstone disease, acne vulgaris, acneiform skin conditions, diabetes, Parkinson's disease, cancer, Alzheimer's disease, inflammation, immunological disorders, lipid disorders, obesity, conditions characterized by a perturbed epidermal barrier function, conditions of disturbed differentiation or excess proliferation of the epidermis or
15 mucous membrane, or cardiovascular disorders.

In a seventh aspect, the invention provides a method of reducing cholesterol levels in a subject in need thereof, comprising administering an effective cholesterol level-reducing amount of a compound of any of formula IIa-d, IIIa-d, IVa-d, XXVa-d, or IV-XXXV.

In a preferred embodiment of the seventh aspect, the invention provides a method of reducing
20 cholesterol levels in a subject in need thereof, comprising administering an effective cholesterol level-reducing amount of a compound according to part (A) of formulas IIa-d.

In an eighth aspect, the invention provides a method of treating, preventing, or ameliorating one or more symptoms of a disease or disorder which is affected by cholesterol, triglyceride, or bile acid levels, comprising administering to a subject in need thereof a therapeutically effective amount of a
25 compound of any of formula IIa-d, IIIa-d, IVa-d, XXVa-d, or IV-XXXV.

In a preferred embodiment of the eighth aspect, the invention provides a method of treating, preventing, or ameliorating one or more symptoms of a disease or disorder which is affected by cholesterol, triglyceride, or bile acid levels, comprising administering to a subject in need thereof a therapeutically effective amount of a compound according to part (A) of formulas IIa-d.

30 In a ninth aspect, the invention provides a method of modulating nuclear receptor activity, comprising contacting the nuclear receptor with a compound of any of formula IIa-d, IIIa-d, IVa-d, XXVa-d, or IV-XXXV.

In a preferred embodiment of the ninth aspect, the invention provides a method of modulating nuclear receptor activity, comprising contacting the nuclear receptor with a compound according to part (A) of formulas IIa-d.

5 In an embodiment of the ninth aspect, the invention provides the method wherein the nuclear receptor is an orphan nuclear receptor.

In an embodiment of the ninth aspect, the invention provides the method wherein the nuclear receptor is a liver X receptor.

In a preferred embodiment of the ninth aspect, the invention provides the method wherein the nuclear receptor is a liver X receptor, wherein the liver X receptor is LXR α or LXR β .

10 In an eleventh aspect, the invention provides a method of modulating cholesterol metabolism, comprising administering an effective cholesterol metabolism-modulating amount of a compound of any of formula IIa-d, IIIa-d, IVa-d, XXVa-d, or IV-XXXV.

In a preferred embodiment of the eleventh aspect, the invention provides a method of modulating cholesterol metabolism, comprising administering an effective cholesterol metabolism-modulating amount of a compound according to part (A) of formulas IIa-d.

15 In a twelfth aspect, the invention provides a method of treating, preventing or ameliorating one or more symptoms of hypocholesterolemia in a subject in need thereof, comprising administering a therapeutically effective amount of a compound of any of formula IIa-d, IIIa-d, IVa-d, XXVa-d, or IV-XXXV.

20 In a preferred embodiment of the twelfth aspect, the invention provides a method of treating, preventing or ameliorating one or more symptoms of hypocholesterolemia in a subject in need thereof, comprising administering a therapeutically effective amount of a compound according to part (A) of formulas IIa-d.

25 In a thirteenth aspect, the invention provides a method of increasing cholesterol efflux from cells of a subject, comprising administering an effective cholesterol efflux-increasing amount of a compound of any of formula IIa-d, IIIa-d, IVa-d, XXVa-d, or IV-XXXV.


In a preferred embodiment of the thirteenth aspect, the invention provides a method of increasing cholesterol efflux from cells of a subject, comprising administering an effective cholesterol efflux-increasing amount of a compound according to part (A) of formulas IIa-d.

30 In a fourteenth aspect, the invention provides a method of increasing the expression of ATP-Binding Cassette (ABC1) in the cells of a subject, comprising administering an effective ABC1 expression-increasing amount of a compound of any of formula IIa-d, IIIa-d, IVa-d, XXVa-d, or IV-XXXV.

In the following embodiments of the first aspect, it is understood that the following provisos apply:

- (i) when L^2 is a bond, both J and K are not absent;
- (ii) if the compound is defined by formula IIaa, then
 - a. if J is phenyl and K is thienyl, furyl, or thiazoyl and q is 0, then R^1 is not 4-(NH_2SO_2)phenyl, 4-(NH_2SO_2)-3-fluorophenyl, p-(CH_3SO_2)phenyl-, or 4p-(CH_3SO_2)-3-fluorophenyl-; and
 - b. if R^5 is pyridyl or phenyl optionally substituted with one or more R^{5a} and L^1 is a bond, then G is not p-(NH_2SO_2)phenyl or p-(CH_3SO_2)phenyl-;
- (iii) if the compound is defined by formula IIcc or IIdd, then G is not p-(NH_2SO_2)phenyl or p-(CH_3SO_2)phenyl-
- (iv) the compound is not 1-(biphenyl-4-yl)-2,5-diphenyl-1H-imidazole.

One embodiment of the invention relates to compounds represented by formulae IIaa, IIbb, IIcc or IIdd:



- as an isomer, a mixture of stereoisomers, a racemic mixture thereof of stereoisomers, or as a tautomer; or as a pharmaceutically acceptable salt, prodrug, solvate or polymorph thereof.

Each R^1 substituent is independently selected from the group consisting of R^5 and $-L_1-R^5$.

- Another embodiment is that R^1 substituent is R^5 ; Preferred R^5 is selected from the group consisting of 5-12 membered aromatic or non-aromatic ring, 5-12 membered heterocyclyl or heteroaryl having one or more heteroatoms N, O or S; R^5 is optionally substituted at a substitutable position with one or more radicals of R^{5a} . R^5 is preferably thienyl, furyl, piperidinyl, pyrrolidinyl, piperazinyl, morpholinyl, thiazolyl, indolyl, oxazolyl, isoxazolyl, pyridinyl, pyrimidinyl, imidazolyl and phenyl.

- Examples of R^{5a} groups include halogen, C_{1-6} haloalkyl, hydroxy, nitro, C_{1-6} aliphatic group, C_{1-6} alkoxy, C_{0-6} alkylOR¹¹, OCOR¹¹, OCON(R¹¹)₂, NR¹¹COR¹¹, NR¹¹CON(R¹¹)₂, C_{0-6} alkylSO₂R¹¹, C_{0-6} alkylSR¹¹, C_{0-6} alkylSO₂N(R¹¹)₂; arylalkyl, aryloxyaryl, aryl C_{1-6} alkoxy, OC₁₋₆ alkylCOR¹¹, OC₁₋₆ alkylIN(R¹¹)₂, C_{0-6} alkylIN(R¹¹)₂; C_{0-6} alkylOR¹¹, C_{0-6} alkylIN(R¹¹)₂, C_{0-6} alkylCOOR¹¹, C_{0-6} alkylCOON(R¹¹)₂, C_{0-6} alkylCON(R¹¹)₂; C_{0-6} alkylC≡N, OC₀₋₆ alkylCOOR¹¹, C_{0-6} alkylOCON(R¹¹)₂, or C_{1-6} alkylOC₁₋₆ alkyl. R^{5a} is optionally substituted at substitutable position with SO₂R¹¹, C_{0-6} alkoxyaryl, 5-12 membered aromatic or non-aromatic ring, or 5-12 membered heterocyclyl or heteroaryl having one or more heteroatoms N, O or S. Preferably, R^{5a} is Cl, Br, F, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{1-6} alkoxy, OC₀₋₆

alkylCOOR¹¹, NR¹¹COR¹¹, C₀₋₆ alkylCON(R¹¹)₂, NO₂, and OC₁₋₆ alkylCON(R¹¹)₂. Examples of R^{5a} include OCH₂C(CH₃)₃, Cl, F, Br, OCH₂CH(CH₃)₂, OCH₂CH₃, CF₃, COOH, OCH₃, OH, NO₂, OCOCH(CH₃)₂, OCOC(CH₃)₃, NHCOCH₃, OCON(CH₃)₂, OCONHCH₃, OCON(CH₂)₂CH₃, OCONHCH(CH₃)₂, O(CH₂)₂, CONH₂, O(CH)(CH₃)₂, C₁₋₆ alkyl, OCH₂COOH, OCH₂COOC(CH₃)₃,
 5 O(CH₂)₂N(CH₂CH₃)₂, OC(CH₃)₂COOC(CH₃)₃, or OCH₂CH₂OH. More preferably, R^{5a} is Cl, F, or CF₃,

Another embodiment is that R¹ substituent is -L₁-R⁵. Preferred R⁵ is selected from the group consisting of 5-12 membered aromatic or non-aromatic ring, 5-12 membered heterocyclyl or heteroaryl having one or more heteroatoms N, O or S; R⁵ is optionally substituted at a substitutable position with one or more radicals of R^{5a}. Examples of preferred R⁵ include phenyl, pyridinyl, oxazolyl, thienyl,
 10 thiazolyl, morpholinyl, furanyl, imidazolyl, piperazinyl, pyrimidinyl, isoxazolyl or piperidinyl, more preferably, oxazolyl, pyridinyl, phenyl, furanyl, thienyl or thiazolyl.

Embodiments for L₁ include a direct bond, C₁₋₆alkoxy, carbonyl, SO₂, CS, CON(R¹¹)₂, CONR¹¹OR¹¹, CONR¹¹N(R¹¹)₂, -C(=NR¹¹)-, -C(=NOR¹¹)-, -C(=NN(R¹¹)₂)-, 5-12 membered aromatic or non-aromatic ring, 5-12 membered heteroaryl or heterocyclyl having one or more heteroatoms N, O, or S which is optionally substituted at a substitutable position with one or more radicals of R¹⁴. Another
 15 embodiment for L₁ is -(CH₂)_m-V-(CH₂)_n-; or V-(CH₂)_n-V; m is 0-6; n is 0-6; V is independently -C(R¹¹)₂-, -C(R¹¹)₂C(R¹¹)₂-, -C(R¹¹)=C(R¹¹)-, -C(R¹¹)₂O-, -C(R¹¹)₂NR¹¹-, -C≡C-, -O-, -S-, -N(R¹⁰)CO-, -N(R¹⁰)CO₂-, -CON(R¹⁰)-, -CO-, -CO₂-, -OC(=O)-, -OC(=O)N(R¹⁰)-, -CONR¹¹OR¹¹-, -CONR¹¹NR¹¹-, -CONR¹¹-, -SO₂-, -N(R¹⁰)SO₂-, -SO₂N(R¹⁰)-, cycloC₃₋₆ alkyl, -NR¹⁰CSNR¹⁰-, cycloC₃₋₆haloalkyl or
 20 -NR¹⁰CONR¹⁰-. A preferred L₁ is selected from the group consisting of CONH, C₁₋₆ aldiyl, CO, SO₂, CH₂, CH₂O, CH₂CH₂, C=O, CONH, CONHC(CH₃)₂, CONH(CH₂)₃OCH₂, OCH₂CH₂, OCH₂CO, OCH₂CH₂N(CH₃)₂, and CONHCH₂CH₂N(CH₃)₂. More preferred L₁ is selected from the group consisting of CH₂, CH₂O, CH₂CH₂, C=O, SO₂, CONH, CONHC(CH₃)₂, CONH(CH₂)₃OCH₂, CONHCH₂CH₂N(CH₃)₂, OCH₂ and OCH₂CH₂. Preferred R⁵ is selected from the group consisting of
 25 phenyl, pyridinyl, oxazolyl, thienyl, thiazolyl, furanyl, morpholinyl, imidazolyl, piperazinyl, pyrimidinyl, isoxazolyl and piperidinyl. Other preferred R^{5a} includes halogen, haloalkyl, OCH₂CON(CH₃)₂, OCH₂COOC(CH₃)₃, OCH₂CH₂N(CH₂CH₃)₂, OCH₂COOH, OC(CH₃)₂COOC(CH₃)₃, OCON(CH₃)₂, OCONHCH₃, OCH₂CH₂OH, OCONHCH₂CHCH₃, or NHCOCH₃.

Another embodiment is that R² is independently selected from the group consisting of R⁷ and L₃-R⁷; Each R⁷ is independently selected from hydrogen, C₁₋₆ alkyl, halogen, C₁₋₆ haloalkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkylOR¹¹, C₀₋₆ alkylCOOR¹¹, C₀₋₆ alkylCON(R¹¹)₂, C₀₋₆ alkylN(R¹¹)₂, C₀₋₆ alkylOR¹¹, C₀₋₆ alkylC≡N, cycloC₃₋₆ alkylC≡N, C₀₋₆ alkyl CONR¹¹N(R¹¹)₂, C₀₋₆ alkylCONR¹¹OR¹¹, C₀₋₆ alkylOCOR¹¹,
 30

C_{0-6} alkyl $SO_2N(R^{11})_2$, cyclo C_{3-6} alkyl, cyclo C_{3-6} alkyl OR^{11} , C_{0-6} alkyl COR^{11} ; 5-12 membered aromatic or non-aromatic ring; or 5-12 membered heteroaryl and heterocyclyl having one or more heteroatoms N, O or S; R^7 is optionally substituted at a substitutable position with one or more radicals of R^{7a} ;

Another embodiment is that R^2 is R^7 , selected from the group consisting of 5-12 membered aromatic or non-aromatic ring; 5-12 membered heteroaryl and heterocyclyl having one or more heteroatoms N, O or S. R^7 is optionally substituted at a substitutable position with one or more radicals of R^{7a} ;

Preferred R^7 is phenyl, pyridinyl, thienyl, furanyl, piperidinyl, pyrrolidinyl, piperazinyl, morpholinyl, thiazolyl, indolyl, oxazolyl, pyridinyl, isoxazolyl, pyrimidinyl, naphthyl, hydrogen, CF_3 , C_{1-6} alkyl $C\equiv N$, CH_2OH , $COOCH_3$, $COON(R^{11})_2$ or $COOR^{11}$. Other examples of R^7 include trifluoromethyl, $CH_2C\equiv N$, $C(CH_3)_2C\equiv N$, $COOCH_3$, CH_2OH , $CONHCH_2CH_3$, $CONHOCH_2CH(OH)CH_2OH$, $CONHCH_2CH_2N(CH_3)_2$, $CONHCH_2CH_2OCH_3$, $CONHCH_2CH_2OCH_3$, CH_2COOCH_3 , $CON(CH_3)_2$, $COOCH(CH_3)_2$, $CONHCH_2CH_2CH_2OCH_3$, $OCOCH(CH_3)_2$, $OCH_2CON(CH_3)_2$, $CH_2CONHCH_2(CH_3)$, $C(CH_3)_2OH$, $COOH$, nitro, cyclo C_{3-6} alkyl, cyclo C_{3-6} alkyl OR^{11} , cyclo C_{3-6} alkyl $C\equiv N$, or $COOCH(CH_3)_2$. More preferably, R^7 is CF_3 , $COOCH_3$, $COOH$, or $CONHCH_2CH_3$. When R^7 is phenyl or pyridinyl, preferred R^{7a} is selected from the group consisting of halogen, C_{1-6} alkyl, C_{1-6} alkoxy, and C_{1-6} haloalkyl. Examples of R^{7a} include halogen, trifluoromethyl, C_{1-6} alkyl, C_{1-6} alkoxy, $CH=CHCOOH$, CH_2COOH , OCH_2COOH , $OCONHCH(CH_3)_2$, $NHCOCH_3$, OH , OCH_3 , $COOH$, $COOCH_3$, $OCH_2C(CH_3)_3$, $OCH_2CH(CH_3)_2$, $OCH(CH_3)_2OCOCH(CH_3)_2$, $OCONHCH_3$, OCH_2CH_3 , and $OCH(CH_3)_2$.

Another embodiment is that R^2 is L_3-R^7 . A preferred L_3 is independently selected from a direct bond, $-CS-$, $-CO-$, $-CONH-$, $-CONR^{11}-$, $-C(=NR^{11})-$, $-C(=NOR^{11})-$, $-C(=NN(R^{11})_2)-$, $(CH_2)_m-V_1-(CH_2)_n-$ or $-V_1-(CH_2)_n-V_1-$; m is 0-6; n is 0-6; V_1 is independently $-C(R^{11})_2-$, $-C(R^{11})_2C(R^{11})_2-$, $-C(R^{11})=C(R^{11})-$, $-C(R^{11})_2O-$, $-C(R^{11})_2NR^{11}-$, $-C\equiv C-$, $-O-$, $-S-$, $-NR^{11}-$, $-CR^{11}NR^{11}-$, $-N(R^{10})CO-$, $-N(R^{10})CO_2-$, $-CON(R^{10})-$, $-OCO-$, $-CO-$, $-CO_2-$, $-OC(=O)-$, $-OC(=O)N(R^{10})-$, $-SO_2-$, $-N(R^{10})SO_2-$, $-NR^{10}CONR^{10}-$, cyclo C_{3-6} alkyl, $-NR^{10}CSNR^{10}-$, cyclo C_{3-6} haloalkyl or $-SO_2N(R^{10})$. More preferably, L_3 is CH_2 , CO , OCH_2 , CH_2OCH_2 , $CONH$, CH_2OCOCH_2 , CH_2NHCH_2 , $CH_2NC(CH_3)_2$, $CH_2N(CH_3)CH_2$, CH_2COCH_3 , $CH_2N(CH_3)_2CH_2$, cyclohexamine or cyclopropanamine.

Each R^{7a} is independently a halogen, C_{1-6} alkyl, $CR^{11}=CR^{11}COOR^{11}$, C_{1-6} alkoxy, C_{0-6} alkyl OR^{11} , C_{0-6} alkyl $OCOOR^{11}$, C_{0-6} alkyl $NR^{11}COR^{11}$, C_{0-6} alkyl SO_2NCOR^{11} , C_{0-6} alkyl $SO_2N(R^{11})_2$; C_{0-6} alkyl SR^{11} , $(C_{0-6}$ alkyl) $C=O(OR^{11})$, $OVOR^{11}$, C_{1-6} haloalkyl, OC_{1-6} haloalkyl, haloaryl, aryloxy, aralkyloxy, aryloxyalkyl, C_{1-6} alkoxyaryl, aryl C_{0-6} alkylcarboxy, $NR^{11}SO_2R^{11}$, OC_{1-6} alkyl, OC_{0-6} alkyl $COOR^{11}$, C_{0-6} alkoxyheteroaryl, C_{0-6} alkoxyheterocyclyl, cyclo C_{3-6} alkyl $COOR^{11}$,

cycloC₃₋₆alkylamine; 5-12 membered aromatic or non-aromatic ring, or 5-12 membered heteroaryl or heterocyclyl having one or more heteroatoms N, O or S; Examples of R^{7a} is selected from the group consisting of halogen, trifluoromethyl, C₁₋₆alkyl, C₁₋₆alkoxy, CH=CHCOOH, CH₂COOH, OCH₂COOH, OCONHCH(CH₃)₂, NHCOCH₃, OH, OCH₃, COOH, COOCH₃, OCH₂C(CH₃)₃,
 5 OCH₂CH(CH₃)₂, OCH(CH₃)₂OCOCH(CH₃)₂, OCONHCH₃, OCH₂CH₃, and OCH(CH₃)₂.

Each R^{7a} may be substituted at a substitutable position with one or more radicals of R⁸; Each R⁸ is independently C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkyl, C₁₋₆ haloalkylOR¹¹, C₀₋₆ alkylOR¹¹, C₀₋₆ alkylCON(R¹¹)₂, C₀₋₆ alkylCOR¹¹, C₀₋₆ alkylCOOR¹¹, NR¹¹COOR¹¹, or C₀₋₆ alkylSO₂R¹¹.

Examples of R^{7a} are selected from the group consisting of halogen, trifluoromethyl, C₁₋₆alkyl,
 10 C₁₋₆alkoxy, CH=CHCOOH, CH₂COOH, OCH₂COOH, OCONHCH(CH₃)₂, NHCOCH₃, OH, OCH₃, COOH, COOCH₃, OCH₂C(CH₃)₃, OCH₂CH(CH₃)₂, OCH(CH₃)₂OCOCH(CH₃)₂, OCONHCH₃, OCH₂CH₃, and OCH(CH₃)₂.

Each R³ is independently selected from the group consisting of R⁶ and -L-R⁶. One embodiment is that R³ is R⁶ and is independently hydrogen, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkyl, C₀₋₆ haloalkylOR¹¹,
 15 C₀₋₆ alkylOR¹¹, C₀₋₆ alkylCON(R¹¹)₂, OCON(R¹¹)₂, C₀₋₆ alkylCOR¹¹, CONR¹¹OR¹¹, C₀₋₆ alkylCOOR¹¹; 5-12 membered aromatic or non-aromatic ring; 5-12 membered heteroaryl or heterocyclyl having one or more heteroatoms N, O or S; Each R⁶ may be substituted at a substitutable position with one or more radicals of R^{6a}; Preferred R⁶ is hydrogen or optionally substituted phenyl.

Each R^{6a} is independently halogen, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkyl, C₀₋₆ haloalkylOR¹¹,
 20 CON(R¹¹)₂, CONR¹¹OR¹¹, C₀₋₆ alkylCOOR¹¹, CR¹¹=CR¹¹COOR¹¹, C₀₋₆ alkylOR¹¹, C₀₋₆ alkylCOR¹¹, C₀₋₆ alkylSO₂R¹¹, C₀₋₆ alkylOCOOR¹¹, C₀₋₆ alkylNR¹¹OR¹¹, C₀₋₆ alkyl SO₂NR¹¹COR¹¹, C₀₋₆ alkyl SO₂N(R¹¹)₂; C₀₋₆ alkylSR¹¹, (C₀₋₆ alkyl)C=O(OR¹¹), OVOR¹¹, OC₁₋₆ haloalkyl, aryloxy, aralkyloxy, aryloxyalkyl, C₁₋₆ alkoxyaryl, arylC₀₋₆ alkylcarboxy, NR¹¹SO₂R¹¹, OC₁₋₆ alkyl, OC₀₋₆ alkylCOOR¹¹, C₀₋₆ alkoxyheteroaryl, C₀₋₆alkoxyheterocyclyl, cycloalkylCOOR¹¹.

Another embodiment is that R³ is L-R⁶, L is independently selected from direct bond, -CO-,
 25 -CONH-, -CONR¹¹-, -C(=NR¹¹)-, -C(=NOR¹¹)-, -CS-, -C(=NN(R¹¹)₂)-, C₂₋₆ alidiyl chain wherein the alidiyl chain is optionally interrupted by -C(R¹¹)₂-, -C(R¹¹)₂C(R¹¹)₂-, -C(R¹¹)=C(R¹¹)-, -C≡C-, -O-, -S-, -N(R¹⁰)CO-, -N(R¹⁰)CO₂-, -NR¹¹-, -OR¹¹-, -CON(R¹⁰)-, -CO-, -CO₂-, -OC(=O)-, -OC(=O)N(R¹⁰)-, -SO₂-, -N(R¹⁰)SO₂-, or -SO₂N(R¹⁰)-; -(CH₂)_m-V₀-(CH₂)_n- or -V₀-(CH₂)_n-V₀; m is 0-6; n is 0-6; V₀ is
 30 independently -C(R¹¹)₂-, -C(R¹¹)₂C(R¹¹)₂-, -C(R¹¹)=C(R¹¹)-, -C(R¹¹)₂O-, -C(R¹¹)₂NR¹¹-, -C≡C-, -O-, -S-, -NR¹¹-, -CR¹¹NR¹¹-, -N(R¹⁰)CO-, -N(R¹⁰)CO₂-, -CON(R¹⁰)-, -OCO-, -COR¹¹-, -COOR¹¹-, -CO-, -CO₂-, -OC(=O), -OC(=O)N(R¹⁰)-, -SO₂-, -N(R¹⁰)SO₂-, -NR¹⁰COR¹⁰-, -NR¹⁰CSNR¹⁰-,

cycloC₃₋₆haloalkyl or -SO₂N(R¹⁰)-. Examples of L include O, CH₂, CH₂O, CH₂CH₂, C=O, SO₂, CONH, CONHC(CH₃)₂, CONH(CH₂)₃OCH₂, CONHCH₂CH₂N(CH₃)₂ or OCH₂CH₂.

Each R⁴ is independently selected from C₁₋₆ alkyl, CR¹¹=CR¹¹COOH, C₁₋₆ alkoxy, C₀₋₆ alkylOR¹¹, C₀₋₆ alkylCOR¹¹, C₀₋₆ alkylSO₂R¹¹, C₀₋₆ alkylOCOOR¹¹, C₀₋₆ alkylNR¹¹COR¹¹, C₀₋₆ alkylSO₂NR¹¹COR¹¹, C₀₋₆ alkylSO₂N(R¹¹)₂, C₀₋₆ alkylSR¹¹, (C₀₋₆ alkyl)C=O(OR¹¹), OVOR¹¹, halogen, C₁₋₆haloalkyl, C₀₋₆ alkylC≡N, OC₁₋₆ haloalkyl, aryloxy, aralkyloxy, aryloxyalkyl, C₁₋₆ alkoxyaryl, arylC₀₋₆ alkylCOOR¹¹, NR¹¹SO₂R¹¹, OC₁₋₆ alkyl, OC₀₋₆ alkylCOOR¹¹, C₀₋₆ alkoxyheteroaryl, C₀₋₆ alkoxyheterocyclyl, cycloalkylCOOR¹¹, a 5-12 membered aromatic ring or non-aromatic ring, or 5-12 membered heteroaryl or heterocyclyl having one or more heteroatoms N, O or S. Preferred R⁴ is selected from the group consisting of OH, CN, C(CH₃)₂OH, SO₂CH₃, SO₂C(CH₃)₃, SO₂CH₂CH₃, SCH₂CH₃, SCH₃, OCH₃, C₁₋₆ alkyl, CH₂COOH, C(CH₃)₂COOH, NHSO₂CH₃, F, Cl, Br, cyclobutane-COOH, OC(CH₃)₂COOH, CF₃, C(CH₃)₂COOH, CH₂COOCH₃, CH₂CH₂COOH, OCH₂COOCH₃, and COCH₃. More preferably, R⁴ is SO₂CH₃, SO₂CH₂CH₃, SCH₂CH₃, or SCH₃.

Each R⁴ is optionally substituted at a substitutable position with one or more radicals of R^{4a}. Each R^{4a} is independently selected from C₁₋₆ alkyl, (C₁₋₆ alkyl)C=O(OR¹¹); C₁₋₆ alkoxy, C₀₋₆ alkylOR¹¹, C₀₋₆ alkylCOR¹¹, C₀₋₆ alkylSO₂R¹¹, C₀₋₆ alkyl SO₂N(R¹¹)₂; C₀₋₆ alkylSR¹¹, (C₀₋₆ alkyl)C=O(OR¹¹), halogen, C₁₋₆ haloalkyl, aryloxy, aralkyloxy, aryloxyalkyl, C₁₋₆ alkoxyaryl, arylC₀₋₆ alkylCOOR¹¹, C₀₋₆ alkylC≡N, NR¹¹SO₂R¹¹, OC₁₋₆ alkyl, or OC₀₋₆ alkylCOOR¹¹.

Each R¹⁰ is independently selected from R¹¹, C(=O)R¹¹, CO₂R¹¹, SO₂R¹¹.

Each R¹¹ is independently selected from hydrogen or substituted or unsubstituted C₁₋₈ aliphatic group; C₁₋₆haloalkyl; N(R¹²)₂; 5-12 membered aromatic or non-aromatic ring, or 5-12 membered heteroaryl or heterocyclyl having one or more heteroatoms, N, S or O; which is optionally substituted at a substitutable position with one or more radicals of R¹².

Each R¹² is independently halogen, C₁₋₆haloalkyl, C₁₋₆ alkyl, C₁₋₆ alkoxy, (C₁₋₆ alkyl)C=O(OR¹³); C₀₋₆ alkylCOR¹³, C₀₋₆ alkylSO₂R¹³, C₀₋₆ alkylCON(R¹³)₂, C₀₋₆ alkylCONR¹³OR¹³, C₀₋₆ alkylOR¹³, C₀₋₆ alkyl SO₂N(R¹³)₂, C₀₋₆ alkylSR¹³, (C₀₋₆ alkyl)C=O(OR¹³), C₀₋₆ haloalkylOR¹³, aryloxy, aralkyloxy, aryloxyalkyl, C₀₋₆ alkoxyaryl, arylC₀₋₆ alkylcarboxy, C₀₋₆ alkylNR¹³SO₂ R¹³, OC₁₋₆ alkyl, or OC₁₋₆ alkylCOOH.


Each R¹³ is independently hydrogen or substituted or unsubstituted C₁₋₈ aliphatic group.

Each R¹⁴ is independently C₁₋₆ alkyl, C₁₋₆ alkoxy, halogen, C₁₋₆ haloalkyl, C₀₋₆ alkylCON(R¹¹)₂, C₀₋₆ alkyl CONR¹¹OR¹¹, C₀₋₆ alkylOR¹¹, or C₀₋₆ alkylCOOR¹¹.


Another embodiment of the invention is that G is independently G1, G2 or G3;



G1



G2



G3

Each Ring J or Ring K may be independently absent, same or different and is independently selected from a 5-12 membered aromatic or non-aromatic ring, or 5-12 membered heterocyclyl or heteroaryl having one or more hetero atoms, N, S or O.

Each Ring J or Ring K independently is optionally substituted at a substitutable position with one or more radicals of R^4 . Ring J is preferably a phenyl ring or a 5-6 membered heteroaryl ring. Examples of Ring J include phenyl, pyridinyl, thienyl, furanyl, morpholinyl, thiazolyl, indolyl, oxazolyl, biphenyl, naphthyl, piperidinyl, piperazinyl, or imidazolyl. A preferred Ring J is thienyl or phenyl. Ring J is optionally substituted at a substitutable position with one or more radicals of R^4 . Suitable Ring J substituents designated as R^4 include methylsulfonyl, or C_{1-6} aliphatic group or substituents selected from the group consisting of $CR^{11}=CR^{11}COOR^{11}$, C_{1-6} alkyl, C_{1-6} alkoxy, $C_{0-6}alkylOR^{11}$, $C_{1-6}alkylCOR^{11}$, $C_{0-6}alkylSO_2R^{11}$, $C_{0-6}alkylOCOOR^{11}$, $C_{0-6}alkylCON(R^{11})_2$, $OC_{1-6}alkylCON(R^{11})_2$, $C_{0-6}alkylC\equiv N$, $C_{0-6}alkylNR^{11}COR^{11}$, $C_{0-6}alkylSO_2NR^{11}COR^{11}$, $C_{0-6}alkylSO_2N(R^{11})_2$, $C_{0-6}alkylSR^{11}$, $(C_{0-6}alkyl)C=O(OR^{11})$, $OVOR^{11}$, halogen, $C_{1-6}haloalkyl$, $OC_{1-6}haloalkyl$, aryloxy, aralkyloxy, aryloxyalkyl, $C_{1-6}alkoxyaryl$, $arylC_{0-6}alkylCOOR^{11}$, $NR^{11}SO_2R^{11}$, $OC_{1-6}alkyl$, $OC_{0-6}alkylCOOR^{11}$, $C_{0-6}alkoxyheteroaryl$, $C_{0-6}alkoxyheterocyclyl$, $cycloalkylCOOR^{11}$, a 5-12 membered aromatic ring or non-aromatic ring, and 5-12 membered heteroaryl or heterocyclyl having one or more heteroatoms N, O or S. Examples of preferred R^4 include OH, CN, $C(CH_3)_2OH$, SO_2CH_3 , SO_2NH_2 , $SO_2CH_2CH_3$, $SO_2C(CH_3)_3$, SCH_2CH_3 , SCH_3 , OCH_3 , $C_{1-6}alkyl$, CH_2COOH , $C(CH_3)_2COOH$, $NHSO_2CH_3$, F, Cl, Br, $C(CH_2CH_3)_2COOH$, CH_2COOCH_3 , $C(CH_3)_2COOCH_3$, CH_2CH_2COOH , $CH=CHCOOH$, OCH_2COOCH_3 , $COCH_3$, OCH_3 , $COOC(CH_3)_3$, cyclobutane-COOH, $OC(CH_3)_2COOH$, CH_2CH_3 , CH_3 , $CH(CH_3)_2$, CH_2COOCH_3 , $OCON(CH_2CH_3)_2$, $NHCOCH_3$, or CF_3 .

Examples of Ring K include phenyl, pyridinyl, thienyl, furanyl, morpholinyl, thiazolyl, indolyl, oxazolyl, biphenyl, naphthyl, piperidinyl, piperazinyl, isoxazolyl, pyrimidinyl, or imidazolyl. Ring K is optionally substituted at a substitutable position with one or more radicals of R^4 . Suitable Ring K substituents designated as R^4 include methylsulfonyl, or C_{1-6} aliphatic group or substituents selected from the group consisting of $CR^{11}=CR^{11}COOR^{11}$, $C_{1-6}alkyl$, $C_{1-6}alkoxy$, $C_{0-6}alkylOR^{11}$, $C_{1-6}alkylCOR^{11}$, $C_{0-6}alkylSO_2R^{11}$, $C_{0-6}alkylOCOOR^{11}$, $C_{0-6}alkylNR^{11}COR^{11}$, $C_{0-6}alkylSO_2NR^{11}COR^{11}$, $C_{0-6}alkylSO_2N(R^{11})_2$, $C_{0-6}alkylSR^{11}$, $(C_{0-6}alkyl)C=O(OR^{11})$, $OVOR^{11}$, halogen, $C_{1-6}haloalkyl$, $OC_{1-6}haloalkyl$, $C_{0-6}alkylC\equiv N$, aryloxy, aralkyloxy, aryloxyalkyl, $C_{1-6}alkoxyaryl$, $arylC_{0-6}alkylcarboxy$, $NR^{11}SO_2R^{11}$, $OC_{1-6}alkyl$, $OC_{0-6}alkylCOOR^{11}$, $C_{0-6}alkoxyheteroaryl$, $C_{0-6}alkoxyheterocyclyl$, $cycloalkylCOOR^{11}$, a 5-12 membered aromatic ring or non-aromatic ring, and 5-12 membered heteroaryl or

heterocyclyl having one or more heteroatoms N, O or S. Preferably, Ring K is phenyl, pyridinyl, thienyl, furanyl, morpholinyl, thiazolyl, indolyl, oxazolyl, biphenyl, naphthyl, piperidinyl, piperazinyl, isoxazolyl, pyrimidinyl, or imidazolyl. When Ring K is a phenyl or pyridinyl, it is preferably substituted by methylsulfonyl.

- 5 Examples of preferred R^4 groups include include OH, CN, $C(CH_3)_2OH$, SO_2CH_3 , SO_2NH_2 , $SO_2CH_2CH_3$, $SO_2C(CH_3)_3$, SCH_2CH_3 , SCH_3 , OCH_3 , C_{1-6} alkyl, CH_2COOH , $C(CH_3)_2COOH$, $NHSO_2CH_3$, F, Cl, Br, $C(CH_2CH_3)_2COOH$, CH_2COOCH_3 , $C(CH_3)_2COOCH_3$, CH_2CH_2COOH , $CH=CHCOOH$, OCH_2COOCH_3 , $COCH_3$, OCH_3 , $COOC(CH_3)_3$, cyclobutane- $COOH$, $OC(CH_3)_2COOH$, CH_2CH_3 , CH_3 , $CH(CH_3)_2$, CH_2COOCH_3 , $OCON(CH_2CH_3)_2$, $NHCOCH_3$, or CF_3 .
- 10 $-L_2$ is $-(CH_2)_m-V_2-(CH_2)_n-$ or $-V_2-(CH_2)_m-V_2-$; m is 0-6; n is 0-6; V_2 is independently $-C(R^{11})_2-$, $-C(R^{11})_2C(R^{11})_2-$, $-C(R^{11})=C(R^{11})-$, $-C(R^{11})_2O-$, $-C(R^{11})_2NR^{11}-$, $-C\equiv C-$, $-O-$, $-S-$, $-N(R^{10})CO-$, $-N(R^{10})CO_2-$, $-CON(R^{10})-$, $-CON(R^{11})-$, $-CON(R^{11})O-$, $-CO-$, $-CO_2-$, $-OR^{11}N-$, $-OR^{11}COO-$, $-OC(=O)-$, $-OC(=O)N(R^{10})-$, $-SO_2-$, $-N(R^{10})SO_2-$, $-NR^{10}CONR^{10}-$, cyclo C_{3-6} alkyl, $-NR^{10}CSNR^{10}-$, cyclo C_{3-6} haloalkyl or $-SO_2N(R^{10})-$; C_{0-6} alidiyl chain wherein alidiyl chain is optionally interrupted by $-$
- 15 $C(R^{11})_2-$, $-C(R^{11})_2C(R^{11})_2-$, $-C(R^{11})=C(R^{11})-$, $-C(R^{11})_2O-$, $-C\equiv C-$, $-O-$, $-S-$, $-N(R^{10})CO-$, $-N(R^{10})CO_2-$, $-CON(R^{10})-$, $-CON(R^{11})-$, $-CON(R^{11})O-$, $-CO-$, $-CO_2-$, $-OC(=O)-$, $-OC(=O)N(R^{10})-$, $-SO_2-$, $-N(R^{10})SO_2-$, cyclo C_{3-6} alkyl or $-SO_2N(R^{10})-$; 5-12 membered aromatic or non-aromatic ring, 5-12 membered heteroaryl or heterocyclyl having one or more heteroatoms, N, S or O which is optionally substituted at a substitutable position with one or more radicals of R^9 . Alternatively, L_2 is a direct bond,
- 20 $-CS-$, $-C_{1-6}$ alkyl-, $-C_{1-6}$ alkoxy-, $-C_{0-6}$ alkyl $COO-$, $-CH=CHCOO-$, $-C_{0-6}$ alkyl $CONR^{11}-$, $-OC_{0-6}$ alkyl $COO-$, $-C_{0-6}$ alkyl SO_2- , $-C_{0-6}$ alkyl $N(R^{11})-$, $-C_{0-6}$ alkyl $N(R^{11})-$, $-C_{0-6}$ alkyl $CO-$, $-cycloalkylamine-$, $-C(=NR^{11})-$, $-C(=NOR^{11})-$, $-C(=NN(R^{11})_2)-$; 5-12 membered aromatic or non-aromatic ring, or 5-12 membered heteroaryl or heterocyclyl having one or more heteroatoms, N, S or O which is optionally substituted at a substitutable position with one or more radicals of R^9 . A
- 25 preferred L_2 is selected from the group consisting of $-CONH-$, $-CONHCH_2-$, $-CH_2O-$, $-OCH_2COOCH_2-$, $-CONHCH_2-$, and $-C\equiv C-$.

Another embodiment is that G is G1, R^1 is R^5 and R^2 is R^7 . When G of formulae IIaa, IIbb, IIcc, or IIdd is G1, a more preferred embodiment of this invention relates to a compound having one or more features selected from the group consisting of:

- 30 a) R^1 is phenyl, pyridinyl, thienyl, furanyl, morpholinyl, thiazolyl, indolyl, oxazolyl, isoxazolyl, pyrimidinyl, or imidazolyl; R^5 is optionally substituted at a substitutable position with one or more radicals of R^{5a} ;

- b) R^{5a} is halogen, trifluoromethyl, $OCONHCH(CH_3)_2$, $NHCOCH_3$, OH , OCH_3 , $COOH$, $COOCH_3$, $OCH_2C(CH_3)_3$, $OCH_2CH(CH_3)_2$, $OCH_2CH_2N(CH_3)_2$, $OCH(CH_3)_2OCOCH(CH_3)_2$, $OCONHCH_3$, OCH_2CH_3 , or $OCH(CH_3)_2$;
- c) R^2 is trifluoromethyl, $COOCH_3$, CH_2OH , $CONHCH_2CH_3$, $CONHOCH_2CH(OH)CH_2OH$, $CONHCH_2CH_2N(CH_3)_2$, $CONHCH_2CH_2OCH_3$, $CONHCH_2CH_2OCH_3$, CH_2COOCH_3 , $CON(CH_3)_2$, $COOCH(CH_3)_2$, $CONHCH_2CH_2CH_2OCH_3$, $OCOCH(CH_3)_2$, $OCH_2CON(CH_3)_2$, $CH_2CONHCH_2(CH_3)$, $C(CH_3)_2OH$, $COOH$, nitro or $COOCH(CH_3)_2$;
- d) R^3 is hydrogen or optionally substituted phenyl;
- e) Ring J is thienyl, thiazolyl, furanyl, pyridinyl or phenyl;
- a) Ring K is optionally substituted phenyl or pyridinyl; and
- b) R^4 is OH , CN , $C(CH_3)_2OH$, SO_2CH_3 , $SO_2C(CH_3)_3$, CH_3 , SO_2NH_2 , $SO_2CH_2CH_3$, SCH_2CH_3 , SCH_3 , OCH_3 , CF_3 , OCF_3 , CH_2CF_3 , C_{1-6} alkyl, halogen or CH_2COOH .

Another embodiment is that G is $G1$, R^1 is R^5 and R^2 is R^7 . When G of formulae IIaa, IIbb, IIcc, or IIdd is $G1$, a more preferred embodiment of this invention relates to a compound having one or more features selected from the group consisting of:

- a) R^1 is thienyl, furanyl, morpholinyl, thiazolyl, indolyl, oxazolyl, pyridinyl, imidazolyl, isoxazolyl, pyrimidinyl or phenyl; R^5 is optionally substituted at a substitutable position with one or more radicals of R^{5a} ;
- b) R^{5a} is halogen, trifluoromethyl, $OCONH(CH_2)_2CH_3$, $OCONH(CH_2CH_3)_2$, $NHCOCH_3$, OH , OCH_3 , $COOH$, $COOCH_3$, $OCH_2C(CH_3)_3$, $OCH_2CH(CH_3)_2$, $OCH(CH_3)_2OCOCH(CH_3)_2$, $OCONHCH_3$, OCH_2CH_3 , or $OCH(CH_3)_2$;
- c) R^2 is R^7 independently selected from $CH_2C\equiv N$, $C(CH_3)_2C\equiv N$, cyclo C_{3-6} alkyl $C\equiv N$, thienyl, furanyl, morpholinyl, thiazolyl, indolyl, oxazolyl, pyridinyl, imidazolyl, isoxazolyl, pyrimidinyl or phenyl; R^7 is optionally substituted at a substitutable position with one or more radicals of R^{7a} ;
- d) R^{7a} is selected from the group consisting of halogen, trifluoromethyl, C_{1-6} alkyl, C_{1-6} alkoxy, $CH=CHCOOH$, CH_2COOH , OCH_2COOH , $OCONHCH(CH_3)_2$, $NHCOCH_3$, OH , OCH_3 , $COOH$, $COOCH_3$, $OCH_2C(CH_3)_3$, $OCH_2CH(CH_3)_2$, $OCH(CH_3)_2OCOCH(CH_3)_2$, $OCONHCH_3$, OCH_2CH_3 , and $OCH(CH_3)_2$;
- e) R^3 is hydrogen or optionally substituted phenyl;
- f) Ring J is thienyl, thiazolyl, furanyl, pyridinyl, or phenyl;
- g) Ring K is optionally substituted phenyl or pyridinyl; and

- h) R^4 is OH, CN, $C(CH_3)_2OH$, $CH=CHCOOH$, SO_2CH_3 , SO_2NH_2 , $SO_2CH_2CH_3$, SCH_2CH_3 , $SO_2C(CH_3)_3$, SCH_3 , OCH_3 , C_{1-6} alkyl, CF_3 , F, Cl, or Br.

Another embodiment is that G is G1, R^1 is L_1-R^5 and R^2 is R^7 . When G of formulae IIaa, IIbb, IIcc, or IIdd is G1, a more preferred embodiment of this invention relates to a compound having one or

5 more features selected from the group consisting of:

- a) R^1 is L_1-R^5 ; R^5 is phenyl, pyridinyl, morpholinyl, oxazolyl, furanyl, thiazolyl or thienyl; R^5 is optionally substituted with R^{5a} ;
- b) R^{5a} is halogen or trifluoromethyl;
- c) L_1 is CS, CH_2 , CH_2O , CH_2CH_2 , $C=O$, SO_2 , CONH, $CONHC(CH_3)_2$, $CONH(CH_2)_3OCH_2$, OCH_2 , OCH_2CO , or OCH_2CH_2 ;
- d) R^2 is trifluoromethyl, $CONHCH_2CH_2N(CH_3)_2$, $CONHCH_2CH_2CH_2N(CH_3)_2$, or $CONHCH_2CH_2CH_2OCH_3$;
- e) R^3 is hydrogen or phenyl optionally substituted with R^{6a} ;
- f) Ring J is thienyl, pyridinyl, furanyl, thiazolyl or phenyl; Ring K is substituted phenyl or pyridinyl; and
- g) R^4 is SO_2CH_3 , SO_2NH_2 , $SO_2CH_2CH_3$, SCH_2CH_3 , SCH_3 , OCH_3 , C_{1-6} alkyl, halogen or CH_2COOH .

Another embodiment is that G is G1, R^1 is R^5 and R^2 is L_3R^7 . When G of formulae IIaa, IIbb, IIcc, or IIdd is G1, a more preferred embodiment of this invention relates to a compound having one or

20 more features selected from the group consisting of:

- a) R^1 is R^5 selected from the group consisting of thienyl, furanyl, morpholinyl, thiazolyl, indolyl, oxazolyl, pyridinyl, imidazolyl, isoxazolyl, pyrimidinyl and phenyl; R^5 is optionally substituted at a substitutable position with one or more radicals of R^{5a} ;
- b) R^{5a} is $OCH_2C(CH_3)_3$, Cl, F, Br, $OCH_2CH(CH_3)_2$, OCH_2CH_3 , CF_3 , $COOH$, OCH_3 , OH, NO_2 , $OCOCH(CH_3)_2$, $NHCOCH_3$, $CONHCH(CH_3)_2$, $O(CH_2)_2$, $CONH_2$, $O(CH)(CH_3)_2$, C_{1-6} alkyl, OCH_2COOH , $OCH_2COOC(CH_3)_3$, $O(CH_2)_2N(CH_2CH_3)_2$, $OCOC(CH_3)_3$, $OC(CH_2)_2COOH$, $OCONH(CH_3)_2$, $OCONCH_3$, $OCONHCH_2CH_2CH_3$, $OC(CH_3)_2COOC(CH_3)_3$, and $O(CH_2)_2OH$;
- c) R^2 is L_3-R^7 ; R^7 is phenyl, pyridinyl, thienyl, furanyl, morpholinyl, thiazolyl, oxazolyl, piperidinyl, imidazolyl, piperazinyl, or pyridinyl;
- d) L_3 is $-CO-$, $-C_{1-6}$ alidyl-, $-CONH-$, $-CONR^{11}-$, $-CONR^{11}NR^{11}-$, $-CH_2OCH_2-$, $-CH_2OCH_2CH_2-$, $-OCH_2-$, $-CH_2N(CH_3)_2-$, $-CH_2NHCH_2-$, $-CONR^{11}O-$, $-CH_2OCOCH_2-$, $-CH_3N(CH_3)(CH_2)-$, $-CH_2N(cyclopropane)CH_2-$, $-CH_2NC(CH_3)_2CH_2-$, $-CH_2N(cyclohexane)CH_2-$, $-CH_2NCH(CH_3)_2CH_2-$, $-CH_2N(CF_3)(CH_2)_2-$,

-CH₂N(CH₃)(CH₂)CH₂OCOCH₂CH₂-, -CONHCH₂CH₂N(CH₃)₂-, or
 -CH₂N(CH₂C≡N)CH₂-, -CS-;

e) R^{7a} is selected from the group consisting of halogen, C₁₋₆alkyl, C₁₋₆ alkoxy, CF₃, OCH₂CH₂COOH, CH₂COOH, COOCH₃, CH₂OH and OCH₃.

5 f) R³ is hydrogen or phenyl optionally substituted with R^{6a};

g) Ring J is thienyl, pyridinyl, thiazolyl or phenyl;

h) Ring K is substituted phenyl or pyridinyl; and

i) R⁴ is SO₂CH₃, SO₂CH₂CH₃, SCH₂CH₃, SCH₃, SO₂NH₂, OCH₃, C₁₋₆ alkyl, CH₂COOH, C(CH₃)₂COOH, NHSO₂CH₃, F, Cl, Br, CF₃ or COCH₃;

10 Another embodiment is that G is G1, R¹ is L₁-R⁵ and R² is L₃-R⁷. When G of formulae IIaa, IIbb, IIcc, or IIdd is G1, a more preferred embodiment of this invention relates to a compound having one or more features selected from the group consisting of:

a) R⁵ is L₁-R⁵; R⁵ is selected from the group consisting of thienyl, furanyl, morpholinyl, thiazolyl, indolyl, imidazolyl, piperazinyl, piperidinyl, oxazolyl, pyridinyl, isoxazolyl, pyrimidinyl, imidazolyl and phenyl; R⁵ is optionally substituted at a substitutable position with one or more radicals of R^{5a};

b) R^{5a} is C₁₋₆ alkyl, C₁₋₆ alkoxy, COOH, halogen or trifluoromethyl;

c) L₁ is CS, CH₂, CH₂O, CH₂CH₂, OCH₂CH₂, OCH₂CO, C=O, SO₂, CONH, CONHC(CH₃)₂, CONH(CH₂)₃OCH₂, or CONHCH₂CHN(CH₃)₂;

20 d) R² is L₃-R⁷; R⁷ is selected from the group consisting of phenyl, pyridinyl, thienyl, furanyl, morpholinyl, thiazolyl, oxazolyl, pyridinyl, isoxazolyl, pyrimidinyl, imidazolyl, CF₃, cycloC₃₋₆ alkylC≡N, C₀₋₆ alkylC≡N and COOCH₃; R⁷ is optionally substituted at a substitutable position with one or more radicals of R^{7a};

25 e) L₃ is CH₂, CS, CH₂OCH₂, NC(CH₃)₂, CH₂NH(CH₂)₂, CONH, CO, CONR¹¹, OCH₂, CH₂N(CH₃)₂CH₂, CH₂OCOCH₂, CH₂CONHCH₂, CH₂CONHCH₂CH₂, cycloalkylamine, CH₂N(CH₃)CH₂, or CH₂NCH(CH₃)₂CH₂;

30 f) R^{7a} is selected from the group consisting of halogen, trifluoromethyl, C₁₋₆alkyl, C₁₋₆alkoxy, CH=CHCOOH, CH₂COOH, OCH₂COOH, OCONHCH(CH₃)₂, NHCOCH₃, OH, OCH₃, COOH, COOCH₃, OCH₂C(CH₃)₃, OCH₂CH(CH₃)₂, OCH(CH₃)₂OCOCH(CH₃)₂, OCONHCH₃, OCH₂CH₃, CH₂N(CH₂)CH₂CF₃, and OCH(CH₃)₂;

g) R³ is hydrogen or phenyl optionally substituted with R^{6a};

h) Ring J is thienyl, thiazolyl, pyridinyl, furanyl or phenyl; Ring K is optionally substituted phenyl or pyridinyl; and

- i) R^4 is SO_2CH_3 , $SO_2CH_2CH_3$, SCH_2CH_3 , SCH_3 , OCH_3 , C_{1-6} alkyl, CH_2COOH , $C(CH_3)_2COOH$, $NHSO_2CH_3$, F , Cl , or Br .

Another embodiment is that G is $G2$ and R^1 is R^5 and R^2 is R^7 . When G of formulae IIaa, IIbb, IIcc, or IIdd is $G2$, a more preferred embodiment of this invention relates to a compound having one or more features selected from the group consisting of:

- a) R^1 is R^5 selected from the group consisting of thienyl, furanyl, morpholinyl, thiazolyl, indolyl, oxazolyl, pyridinyl, isoxazolyl, pyrimidinyl, imidazolyl and phenyl; R^5 is optionally substituted at a substitutable position with one or more radicals of R^{5a} ;
- b) R^2 is R^7 selected from the group consisting of phenyl, pyridinyl, thienyl, furanyl, morpholinyl, thiazolyl, oxazolyl, pyridinyl, isoxazolyl, pyrimidinyl, imidazolyl, CF_3 , cyclo C_{3-6} alkyl $C\equiv N$, C_{0-6} alkyl $C\equiv N$, and $COOCH_3$; R^7 is optionally substituted at a substitutable position with one or more radicals of R^{7a} ;
- c) R^3 is hydrogen or optionally substituted phenyl;
- d) L_2 is selected from the group consisting of $-CONH-$, $-CONHCH_2-$, $-CH_2O-$, $-OCH_2COOCH_2-$, $-O-$, $C\equiv C-$, $-OCH_2CH_2-$, and $-CONHOCH_2CH(OH)CH_2O-$, $-CS-$;
- e) Ring J or K is substituted phenyl, biphenyl, pyridinyl, piperidinyl, piperazinyl, morpholinyl, furanyl, thienyl, or naphthyl; and
- f) R^4 is selected from the group consisting of OH , CN , $C(CH_3)_2OH$, SO_2CH_3 , $SO_2CH_2CH_3$, $SO_2CH_2CH_2CH_3$, SCH_2CH_3 , SCH_3 , OCH_3 , C_{1-6} alkyl, CH_2COOH , $C(CH_3)_2COOH$, $NHSO_2CH_3$, F , Cl , Br , $C(CH_3)_2COOH$, CH_2COOCH_3 , $C(CH_3)_2COOCH_3$, CH_2CH_2COOH , $OCH_2CON(R^{11})_2$, $OCH_2CH_2N(CH_3)_2$, OCH_2COOH , OCH_2COOCH_3 , CH_2OH , $COCH_3$, $COOC(CH_3)_3$, cyclobutane- $COOH$, $OC(CH_3)_2COOH$ and CF_3 .

Another embodiment is that G is $G2$, R^1 is L_1-R^5 and R^2 is R^7 . When G of formulae IIaa, IIbb, IIcc, or IIdd is $G2$, a more preferred embodiment of this invention relates to a compound having one or more features selected from the group consisting of:

- a) R^1 is L_1-R^5 ; R^5 is substituted phenyl or pyridinyl;
- b) R^{5a} is halogen, trifluoromethyl, C_{1-6} alkyl, C_{1-6} haloalkyl, nitro, C_{1-6} alkoxy, or $OCON(C_{1-6}alkyl)_2$;
- c) L_1 is CH_2 , CH_2O , CH_2CH_2 , $C=O$, SO_2 , $CONH$, $CONHC(CH_3)_2$, $CONH(CH_2)_3OCH_2$, $CONHCH_2CH_2N(CH_3)_2$ or OCH_2CH_2 , $-CS-$;
- d) R^2 is R^7 is selected from the group consisting of phenyl, pyridinyl, thienyl, furanyl, morpholinyl, thiazolyl, oxazolyl, pyridinyl, CF_3 , cyclo C_{3-6} alkyl $C\equiv N$, C_{0-6} alkyl $C\equiv N$ or $COOCH_3$;

- e) R^3 is hydrogen or phenyl optionally substituted with R^{6a} ;
- f) Ring J or K is substituted phenyl, thienyl, furanyl, piperazinyl, piperidinyl or pyridinyl;
- g) L_2 is CONH, CONHCH₂, CH₂O, OCH₂COOCH₂, O, C≡C, -CS-, OCH₂CH₂ or CONHOCH₂CH(OH)CH₂O; and
- 5 h) R^4 is selected from the group consisting of halogen, C₁₋₆ haloalkyl, C₁₋₆ alkylCOOR¹¹, and methyl sulfonyl.

Another embodiment is that G is G2, R^1 is R^5 and R^2 is L_3R^7 . When G of formulae IIaa, IIbb, IIcc, or IIdd is G2, a more preferred embodiment of this invention relates to a compound having one or more features selected from the group consisting of:

- 10 a) R^1 is R^5 selected from the group consisting of thienyl, furanyl, morpholinyl, thiazolyl, indolyl, oxazolyl, pyridinyl, imidazolyl, isoxazolyl, pyrimidinyl and phenyl; R^5 is optionally substituted at a substitutable position with one or more radicals of R^{5a} ;
- b) R^{5a} is halogen or trifluoromethyl;
- c) R^2 is L_3R^7 ; R^7 is selected from the group consisting of phenyl, pyridinyl, thienyl, furanyl, morpholinyl, thiazolyl, oxazolyl, pyridinyl, phenyl, imidazolyl, isoxazole, pyrimidinyl, CF₃, cycloC₃₋₆ alkylC≡N, C₀₋₆ alkylC≡N and COOCH₃; R^7 is optionally substituted at a substitutable position with one or more radicals of R^{7a} ;
- 15 d) L_3 is -CS-, CH₂, CH₂OCH₂, NCH₂ (CH₂)₂, CH₂N(CH₂)₂, CH₂CN, CONH, CO, or CONHCH₂;
- e) R^3 is hydrogen or optionally substituted phenyl;
- f) Ring J or K is substituted phenyl, furanyl, thienyl, pyridinyl, biphenyl or naphthyl;
- g) L_2 is CONH, CS, CONHCH₂, CH₂O, OCH₂COOCH₂, OCH₂CH₂, or or OCH₂; and
- h) R^4 is SO₂CH₃, SO₂CH₂CH₃, SCH₂CH₃, CH₂COOH, C(CH₃)₂COOH, NHSO₂CH₃, F, Cl, Br, SCH₃, OCH₃, C₁₋₆ alkyl, COOCH₂CO, OCH₃, CH₂COOH, CH₂COOCH₃, CH(CH₃)₂COOH, OC(CH₃)₂COOH, COOC(CH₃)₃, cyclobutane-COOH, C(CH₃)₂COOH, OCH₂COOCH₃, and CF₃.
- 25

Another embodiment is that G is G3, R^1 is R^5 and R^2 is R^7 . When G of formulae IIaa, IIbb, IIcc, or IIdd is G3, a more preferred embodiment of this invention relates to a compound having one or more features selected from the group consisting of:

- 30 a) R^1 is R^5 selected from the group consisting of thienyl, furanyl, morpholinyl, thiazolyl, indolyl, oxazolyl, pyridinyl, imidazolyl, isoxazole, pyrimidinyl and phenyl; R^5 is optionally substituted at a substitutable position with one or more radicals of R^{5a} ;


- b) R^2 is R^7 selected from the group consisting of phenyl, pyridinyl, thienyl, furanyl, morpholinyl, thiazolyl, oxazolyl, pyridinyl, imidazolyl, isoxazole, pyrimidinyl, CF_3 , halogen, cyclo C_{3-6} alkyl $C\equiv N$, C_{0-6} alkyl $C\equiv N$ and $COOCH_3$; R^7 is optionally substituted at a substitutable position with one or more radicals of R^{7a} ;
- 5 c) R^3 is hydrogen or optionally substituted phenyl;
- d) L_2 is selected from the group consisting of $-CONH-$, $-CONHCH_2-$, $-CS-$, $-CH_2O-$, $-OCH_2COOCH_2-$, $-COOCH_2-$, $-CO-$, $-OCH_2-$, $-OCO-$, $-NHCON-$, $-O-$, $-OCH_2CH_2-$, $-OCON-$, and $-SO_2-$;
- 10 e) Ring J or K is substituted phenyl, biphenyl, pyridinyl, piperidinyl, piperazinyl, morpholinyl, thienyl, pyrimidinyl or naphthyl;
- f) R^4 is methylsulfonyl, halogen, haloalkyl, CH_2COOH , OCH_2 -phenyl, CH_2COO -phenyl, OCH_2COOH , or $OCH_2CHN(CH_3)_2$; and
- 15 g) R^{5a} is $OCH_2C(CH_3)_3$, Cl, F, Br, $OCH_2CH(CH_3)_2$, OCH_2CH_3 , CF_3 , $COOH$, OCH_3 , OH, NO_2 , $OCOCH(CH_3)_2$, $NHCOCH_3$, $OCONHCH(CH_3)_2$, $O(CH_2)_2$, $CONH_2$, $O(CH)(CH_3)_2$, C_{1-6} alkyl, OCH_2COOH , $OCH_2COOC(CH_3)_3$, $O(CH_2)_2N(CH_2CH_3)_2$, $OCOC(CH_3)_3$, $OC(CH_2)_2COOH$, $OCONH(CH_3)_2$, $OCONCH_3$, $OCONHCH_2CH_2CH_3$, $OC(CH_3)_2COOC(CH_3)_3$, and $O(CH_2)_2OH$.
- Another embodiment is that G is G3, R^1 is L_1-R^5 and R^2 is R^7 . When G of formulae IIaa, IIbb, IIcc, or IIdd is G3, a more preferred embodiment of this invention relates to a compound
- 20 having one or more features selected from the group consisting of:
- a) R^1 is L_1-R^5 ; R^5 is substituted phenyl or pyridinyl;
- b) R^{5a} is halogen or trifluoromethyl;
- c) L_1 is CH_2 , CH_2O , CH_2CH_2 , $C=O$, SO_2 , $CONH$, $CONHC(CH_3)_2$, CS , $CONH(CH_2)_3OCH_2$, $CONHCH_2CH_2N(CH_3)_2$ or OCH_2CH_2 ;
- 25 d) R^2 is halogen, C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} carboxyalkyl, or CF_3 ;
- e) R^3 is hydrogen or phenyl optionally substituted with R^{6a} ;
- f) Ring J or K is phenyl, pyridinyl, thienyl, furanyl, piperidinyl, pyrrolidinyl, piperazinyl, morpholinyl, thiazolyl, indolyl, oxazolyl, isoxazolyl, pyrimidinyl, imidazolyl, or biphenyl; and
- 30 g) L_2 is CS , $CONH$, $CONHCH_2$, CH_2O , OCH_2COOCH_2 , OCH_2 or OCH_2CH_2 ;
- h) R^4 is selected from the group consisting of halogen, C_{1-6} haloalkyl, C_{1-6} alkyl $COOR^{11}$, and methyl sulfonyl.

Another embodiment is that G is G3, R¹ is R⁵ and R² is L₃R⁷. When G of formulae IIaa, IIbb, IIcc, or IIdd is G3, a more preferred embodiment of this invention relates to a compound having one or more features selected from the group consisting of:


- a) R¹ is selected from the group consisting of thienyl, furanyl, morpholinyl, thiazolyl, indolyl, oxazolyl, pyridinyl, isoxazolyl, imidazolyl, pyrimidinyl and phenyl; R⁵ is optionally substituted at a substitutable position with one or more radicals of R^{5a};
- b) R² is L₃-R⁷; R⁷ is phenyl, pyridinyl, thienyl, furanyl, morpholinyl, thiazolyl, oxazolyl, piperidinyl, imidazolyl, piperazinyl, pyridinyl, isoxazolyl, imidazolyl, pyrimidinyl, CF₃, cycloC₃₋₆ alkylC≡N, C₀₋₆ alkylC≡N and COOCH₃; R⁷ is optionally substituted at a substitutable position with one or more radicals of R^{7a};
- c) L₃ is -CO-, -CS-, -C₁₋₆ alidyl-, -CONH-, -CONR¹¹-, -CONR¹¹NR¹¹-, -CH₂OCH₂-, -CH₂OCH₂CH₂-, -OCH₂-, -CH₂N(CH₃)₂-, -CH₂NHCH₂-, -CONR¹¹O-, -CH₂OCOCH₂-, -CH₃N(CH₃)(CH₂)-, -CH₂N(cyclopropane)CH₂-, -CH₂NC(CH₃)₂CH₂-, -CH₂N(cyclohexane)CH₂-, -CH₂NCH(CH₃)₂CH₂-, -CH₂N(CF₃)(CH₂)₂-, -CH₂N(CH₃)(CH₂)CH₂OCOCH₂CH₂-, -CONHCH₂CH₂N(CH₃)₂-, or -CH₂N(CH₂C≡N)CH₂-;
- d) R³ is hydrogen or optionally substituted phenyl;
- e) Ring J or K is substituted phenyl, furanyl, thienyl, pyridinyl, biphenyl or naphthyl; and
- f) L₂ is CONH, CONHCH₂, CH₂O, OCH₂COOCH₂, CS or CONHCH₂;
- g) R⁴ is SO₂CH₃, SO₂CH₂CH₃, SCH₂CH₃, SCH₃, SO₂NH₂, OCH₃, C₁₋₆ alkyl, CH₂COOH, C(CH₃)₂COOH, NHSO₂CH₃, F, Cl, Br, CF₃ or COCH₃.

Another embodiment of this invention relates to a composition comprising a compound of the present invention or a tautomer or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier. It will be appreciated that the compounds of this invention may be derivatized at functional groups to provide prodrug derivatives which are capable of conversion back to the parent compounds *in vivo*. Examples of such prodrugs include the physiologically acceptable and metabolically labile ester derivatives, such as methoxymethyl esters, methylthiomethyl esters, or pivaloyloxymethyl esters derived from a hydroxyl group of the compound or a carbamoyl moiety derived from an amino group of the compound. Additionally, any physiologically acceptable equivalents of the compounds of the present invention, similar to metabolically labile esters or carbamates, which are capable of producing the parent compounds of the present invention *in vivo*, are within the scope of this invention.


Another embodiment of this invention relates to compounds represented by formulae IIaa-1, IIaa-2, IIaa-3 or IIaa-4 (Embodiment IIaa):




Another embodiment of this invention relates to compounds represented by formulae IIbb-1, IIbb-2, IIbb-3, or IIbb-4 (Embodiment IIb):



Another embodiment of this invention relates to compounds represented by formulae IIcc-1, IIcc-2, IIcc-3, or IIcc-4 (Embodiment IIcc):



Another embodiment of this invention relates to compounds represented by formulae IIdd-1, IIdd-2, IIdd-3, or IIdd-4 (Embodiment IIdd):



Of the above embodiments IIaa-IIdd, R^1 is R^5 selected from the group consisting of thienyl, furanyl, morpholinyl, thiazolyl, indolyl, oxazolyl, pyridinyl, isoxazolyl, pyrimidinyl, imidazolyl, and phenyl; R^5 is optionally substituted at a substitutable position with one or more radicals of R^{5a} . Preferred R^5 is phenyl or pyridinyl optionally substituted with R^{5a} .

R^2 is R^7 selected from the group consisting of trifluoromethyl, COOCH_3 , CH_2OH , $\text{CONHCH}_2\text{CH}_3$, $\text{CONHOCH}_2\text{CH}(\text{OH})\text{CH}_2\text{OH}$, $\text{CONHCH}_2\text{CH}_2\text{N}(\text{CH}_3)_2$, $\text{CONHCH}_2\text{CH}_2\text{OCH}_3$, $\text{CONHCH}_2\text{CH}_2\text{OCH}_3$, $\text{CH}_2\text{COOCH}_3$, $\text{CON}(\text{CH}_3)_2$, $\text{COOCH}(\text{CH}_3)_2$, $\text{CONHCH}_2\text{CH}_2\text{CH}_2\text{OCH}_3$, $\text{OCOCH}(\text{CH}_3)_2$, $\text{OCH}_2\text{CON}(\text{CH}_3)_2$, $\text{CH}_2\text{CONHCH}_2(\text{CH}_3)$, $\text{C}(\text{CH}_3)_2\text{OH}$, COOH , nitro or $\text{COOCH}(\text{CH}_3)_2$, $\text{CH}_2\text{C}\equiv\text{N}$, $\text{C}(\text{CH}_3)_2\text{C}\equiv\text{N}$, cyclo C_{3-6} alkyl $\text{C}\equiv\text{N}$, thienyl, furanyl, morpholinyl, thiazolyl,

indolyl, oxazolyl, pyridinyl, imidazolyl, isoxazolyl, pyrimidinyl and phenyl; R⁷ is optionally substituted at a substitutable position with one or more radicals of R^{7a}.

L₁ is independently selected from direct bond, -CO-, -CONH-, -CONR¹¹-, -C(=NR¹¹)-, -C(=NOR¹¹)-, -C(=NN(R¹¹)₂)-, C₂₋₆ alidiyl chain wherein the alidiyl chain is optionally interrupted by
 5 -C(R¹¹)₂-, -C(R¹¹)₂C(R¹¹)₂-, -C(R¹¹)=C(R¹¹)-, -C≡C-, -O-, -S-, -N(R¹⁰)CO-, -N(R¹⁰)CO₂-, -NR¹¹-, -OR¹¹-,
 -CON(R¹⁰)-, -CO-, -CO₂-, -OC(=O)-, -OC(=O)N(R¹⁰)-, -SO₂-, -N(R¹⁰)SO₂-, or -SO₂N(R¹⁰)-;
 -(CH₂)_m-V₀-(CH₂)_n- or -V₀-(CH₂)_n-V₀-; m is 0-6; n is 0-6; V₀ is independently -C(R¹¹)₂-,
 -C(R¹¹)₂C(R¹¹)₂-, -C(R¹¹)=C(R¹¹)-, -C(R¹¹)₂O-, -C(R¹¹)₂NR¹¹-, -C≡C-, -O-, -S-, -NR¹¹-, -CR¹¹NR¹¹-,
 -N(R¹⁰)CO-, -N(R¹⁰)CO₂-, -CON(R¹⁰)-, -OCO-, -COR¹¹-, -COOR¹¹-, -CO-, -CO₂-, -OC(=O),
 10 -OC(=O)N(R¹⁰)-, -SO₂-, -N(R¹⁰)SO₂-, -NR¹⁰COR¹⁰-, -NR¹⁰CSNR¹⁰-, cycloC₃₋₆haloalkyl or
 -SO₂N(R¹⁰)-. More specifically, L₁ is selected from the group consisting of -CONH-, -C₁₋₆ alkyl-, -C₁₋₆
 alkoxy-, -CO-, -SO₂-, -CH₂-, -CH₂O-, -CH₂CH₂-, -C=O-, -CONH-, -CONHC(CH₃)₂-,
 -CONH(CH₂)₃OCH₂-, -OCH₂CH₂-, -OCH₂CH₂N(CH₃)₂-, and -CONHCH₂CH₂N(CH₃)₂-.

L₃ is independently selected from direct bond, -CO-, -CONH-, -CONR¹¹-, -C(=NR¹¹)-,
 15 -C(=NOR¹¹)-, -C(=NN(R¹¹)₂)-, C₂₋₆ alidiyl chain wherein the alidiyl chain is optionally interrupted by
 -C(R¹¹)₂-, -C(R¹¹)₂C(R¹¹)₂-, -C(R¹¹)=C(R¹¹)-, -C≡C-, -O-, -S-, -N(R¹⁰)CO-, -N(R¹⁰)CO₂-, -NR¹¹-, -OR¹¹-,
 -CON(R¹⁰)-, -CO-, -CO₂-, -OC(=O)-, -OC(=O)N(R¹⁰)-, -SO₂-, -N(R¹⁰)SO₂-, or -SO₂N(R¹⁰)-;
 -(CH₂)_m-V₀-(CH₂)_n- or -V₀-(CH₂)_n-V₀-; m is 0-6; n is 0-6; V₀ is independently -C(R¹¹)₂-,
 -C(R¹¹)₂C(R¹¹)₂-, -C(R¹¹)=C(R¹¹)-, -C(R¹¹)₂O-, -C(R¹¹)₂NR¹¹-, -C≡C-, -O-, -S-, -NR¹¹-, -CR¹¹NR¹¹-,
 20 -N(R¹⁰)CO-, -N(R¹⁰)CO₂-, -CON(R¹⁰)-, -OCO-, -COR¹¹-, -COOR¹¹-, -CO-, -CO₂-, -OC(=O),
 -OC(=O)N(R¹⁰)-, -SO₂-, -N(R¹⁰)SO₂-, -NR¹⁰COR¹⁰-, -NR¹⁰CSNR¹⁰-, cycloC₃₋₆haloalkyl or
 -SO₂N(R¹⁰)-. More specifically, L₃ is -CO-, -C₁₋₆ alidiyl-, -CONH-, -CONR¹¹-, -CONR¹¹NR¹¹-,
 -CH₂OCH₂-, -CH₂OCH₂CH₂-, -OCH₂-, -CH₂N(CH₃)₂-, -CH₂NHCH₂-, -CONR¹¹O-, -CH₂OCOCH₂-,
 -CH₃N(CH₃)(CH₂)-, -CH₂N(cyclopropane)CH₂-, -CH₂NC(CH₃)₂CH₂-, -CH₂N(cyclohexane)CH₂-,
 25 -CH₂NCH(CH₃)₂CH₂-, -CH₂N(CF₃)(CH₂)₂-, -CH₂N(CH₃)(CH₂)CH₂OCOCH₂CH₂-,
 -CONHCH₂CH₂N(CH₃)₂-, or -CH₂N(CH₂C≡N)CH₂-.


R^{7a} is selected from the group consisting of halogen, trifluoromethyl, C₁₋₆alkyl, C₁₋₆alkoxy,
 CH=CHCOOH, CH₂COOH, OCH₂COOH, OCONHCH(CH₃)₂, NHCOCH₃, OH, OCH₃, COOH,
 COOCH₃, OCH₂C(CH₃)₃, OCH₂CH(CH₃)₂, OCH(CH₃)₂OCOCH(CH₃)₂, OCONHCH₃, OCH₂CH₃, or
 30 OCH(CH₃)₂.

L₂ is independently selected from direct bond, -CO-, -CONH-, -CONR¹¹-, -C(=NR¹¹)-,
 -C(=NOR¹¹)-, -C(=NN(R¹¹)₂)-, C₂₋₆ alidiyl chain wherein the alidiyl chain is optionally interrupted by
 -C(R¹¹)₂-, -C(R¹¹)₂C(R¹¹)₂-, -C(R¹¹)=C(R¹¹)-, -C≡C-, -O-, -S-, -N(R¹⁰)CO-, -N(R¹⁰)CO₂-, -NR¹¹-, -OR¹¹-,

-CON(R¹⁰)-, -CO-, -CO₂-, -OC(=O)-, -OC(=O)N(R¹⁰)-, -SO₂-, -N(R¹⁰)SO₂-, or -SO₂N(R¹⁰)-;
 -(CH₂)_m-V₀-(CH₂)_n- or -V₀-(CH₂)_n-V₀-; m is 0-6; n is 0-6; V₀ is independently -C(R¹¹)₂-,
 -C(R¹¹)₂C(R¹¹)₂-, -C(R¹¹)=C(R¹¹)-, -C(R¹¹)₂O-, -C(R¹¹)₂NR¹¹-, -C≡C-, -O-, -S-, -NR¹¹-, -CR¹¹NR¹¹-,
 -N(R¹⁰)CO-, -N(R¹⁰)CO₂-, -CON(R¹⁰)-, -OCO-, -COR¹¹-, -COOR¹¹-, -CO-, -CO₂-, -OC(=O),
 5 -OC(=O)N(R¹⁰)-, -SO₂-, -N(R¹⁰)SO₂-, -NR¹⁰COR¹⁰-, -NR¹⁰CSNR¹⁰-, cycloC₃₋₆haloalkyl or
 -SO₂N(R¹⁰)-. More specifically, L₂ is selected from the group consisting of -CONH-, -CONHCH₂-,
 -CH₂O-, -OCH₂COOCH₂-, -O-, C≡C-, -OCH₂CH₂-, and -CONHOCH₂CH(OH)CH₂O-.

R^{5a} is independently selected from the group consisting of OCH₂C(CH₃)₃, Cl, F, Br,
 OCH₂CH(CH₃)₂, OCH₂CH₃, CF₃, COOH, OCH₃, OH, NO₂, OCOCH(CH₃)₂, OCOC(CH₃)₃,
 10 NHCOCH₃, OCON(CH₃)₂, OCONHCH₃, OCON(CH₂)₂CH₃, OCONHCH(CH₃)₂, O(CH₂)₂, CONH₂,
 O(CH)(CH₃)₂, C₁₋₆ alkyl, OCH₂COOH, OCH₂COOC(CH₃)₃, O(CH₂)₂N(CH₂CH₃)₂,
 OC(CH₃)₂COOC(CH₃)₃, and OCH₂CH₂OH. R⁴ is selected from the group consisting of SO₂CH₃,
 SO₂C(CH₃)₃, SO₂NH₂, SO₂CH₂CH₃, SCH₂CH₃, SCH₃, OCH₃, C₁₋₆ alkyl, CH₂COOH, C(CH₃)₂COOH,
 NHSO₂CH₃, F, Cl, Br, C(CH₃)₂COOH, CH₂COOCH₃, C(CH₃)₂COOCH₃, CH₂CH₂COOH,
 15 OCH₂COOCH₃, COCH₃, COOC(CH₃)₃, cyclobutane-COOH, OC(CH₃)₂COOH, COOCH₂CH₃,
 OCF₃, and CF₃.

Another embodiment of this invention relates to compounds as described above wherein G is
 selected from the group consisting of:



20 Of the above compounds, R is selected from the group consisting of C₀₋₆ alidiyl chain wherein
 the alidiyl chain is optionally interrupted by -C(R¹¹)₂-, -C(R¹¹)₂C(R¹¹)₂-, -C(R¹¹)=C(R¹¹)-, -C(R¹¹)₂O-,
 -C(R¹¹)₂NR¹¹-, -C≡C-, -O-, -S-, -N(R¹⁰)CO-, -N(R¹⁰)CO₂-, -CON(R¹⁰)-, -CO-, -CO₂-, -OC(=O)-,
 -OC(=O)N(R¹⁰)-, -SO₂-, -N(R¹⁰)SO₂-, or -SO₂N(R¹⁰)-.

Each R⁴ is independently selected from, C₁₋₆ alkyl, CR¹¹=CR¹¹COOR¹¹, C₁₋₆ alkoxy, C₀₋₆
 25 alkylOR¹¹, C₀₋₆ alkylCOR¹¹, C₀₋₆ alkylSO₂R¹¹, C₀₋₆ alkylOCOOR¹¹, C₀₋₆ alkylNR¹¹COR¹¹, C₀₋₆
 alkylSO₂NR¹¹COR¹¹, C₀₋₆ alkyl SO₂N(R¹¹)₂, C₀₋₆ alkylSR¹¹, (C₀₋₆ alkyl)C=O(OR¹¹), OVOR¹¹, halogen,
 C₁₋₆haloalkyl, C₁₋₆haloalkylOR¹¹, OC₁₋₆haloalkyl, aryloxy, aralkyloxy, aryloxyalkyl, C₁₋₆ alkoxyaryl,

arylC₀₋₆ alkylcarboxy, NR¹¹SO₂R¹¹, OC₁₋₆ alkyl, OC₀₋₆ alkylCOOR¹¹, C₀₋₆ alkylC≡N, C₀₋₆ alkoxyheteroaryl, C₀₋₆ alkoxyheterocyclyl, cycloalkylCOOR¹¹, a 5-12 membered aromatic ring or non-aromatic ring, or 5-12 membered heteroaryl or heterocyclyl having one or more heteroatoms N, O or S. Preferred R⁴ is selected from the group consisting of SO₂CH₃, SO₂C(CH₃)₃, SO₂CH₂CH₃, SCH₂CH₃, SCH₃, OCH₃, C₁₋₆ alkyl, CH₂COOH, C(CH₃)₂COOH, NHSO₂CH₃, F, Cl, Br, cyclobutane-COOH, OC(CH₃)₂COOH, CF₃, C(CH₃)₂COOH, CH₂COOCH₃, CH₂CH₂COOH, OCH₂COOCH₃, and COCH₃. More preferably, R⁴ is SO₂CH₃, SO₂CH₂CH₃, SCH₂CH₃, or SCH₃.

X is selected from the group consisting of S, NR¹¹ and O.

Each R⁴ is optionally substituted at a substitutable position with one or more radicals of R^{4a}. Each R^{4a} is independently selected from, C₁₋₆ alkyl, (C₁₋₆ alkyl)C=O(OR¹¹); C₁₋₆ alkoxy, C₀₋₆ alkylOR¹¹, C₀₋₆ alkylCOR¹¹, C₀₋₆ alkylSO₂R¹¹, C₀₋₆ alkylSO₂N(R¹¹)₂; C₀₋₆ alkylSR¹¹, (C₀₋₆ alkyl)OC=O(OR¹¹), halogen, C₁₋₆ haloalkyl, aryloxy, aralkyloxy, aryloxyalkyl, C₁₋₆ alkoxyaryl, arylC₀₋₆ alkylcarboxy, NR¹¹SO₂R¹¹, OC₁₋₆ alkyl, C₀₋₆ alkylC≡N, or OC₀₋₆ alkylCOOR¹¹.

Definitions

The following definitions apply to the terms used herein, unless expressly stated to the contrary. So, for example, "alkyl" is defined hereinbelow as containing from 1 to 12 carbon atoms, but a substituent defined as C₁₋₆alkyl is limited to an alkyl moiety of from 1 to 6 carbons. All selections of any variables in connection with any of the general structures or formulas herein are understood to be proper only when said selection yields a stable chemical structure as recognized by one skilled in the art.

When particular embodiments are referred to by structure only, all otherwise unnamed chemical groups making up that structure are as defined in each individual embodiment of that structure. So, for example, when it is stated, "In another embodiment, the invention provides the compound according to any one of formulas Ia-d, wherein K is phenyl or pyridyl," it is meant that another embodiment of the invention comprises each embodiment of any one of formulas Ia-d described in the specification in which K is phenyl or pyridyl and all other moieties are as defined in the particular embodiments.

For simplicity, chemical moieties are defined and referred to throughout primarily as univalent chemical moieties (e.g., alkyl, aryl, etc.). Nevertheless, such terms are also used to convey corresponding multivalent moieties under the appropriate structural circumstances clear to those skilled in the art. For example, while an "alkyl" moiety generally refers to a monovalent radical (e.g. CH₃-CH₂-), in certain circumstances a bivalent linking moiety can be "alkyl," in which case those skilled in the art will understand the alkyl to be a divalent radical (e.g., -CH₂-CH₂-), which is equivalent to the term "alkylene." (Similarly, in circumstances in which a divalent moiety is required and is stated as being "aryl," those skilled in the art will understand that the term "aryl" refers to the corresponding divalent

moiety, arylene.) All atoms are understood to have their normal number of valences for bond formation (i.e., 4 for carbon, 3 for N, 2 for O, and 2, 4, or 6 for S, depending on the oxidation state of the S). On occasion a moiety may be defined, for example, as (A)_a-B-, wherein a is 0 or 1. In such instances, when a is 0 the moiety is B- and when a is 1 the moiety is A-B-. Similarly, C₀₋₆ alkylOR¹¹ includes both -OR¹¹ and C₁-C₆-OR¹¹, and -[C(R¹⁵)₂]_m- is a bond when m is 0. In the instances when a moiety is a divalent radical, there is no implied limitation on the location of the two bonds connecting the linking radical to its two supporting chemical units. For example, for a divalent cyclohexyl radical, the cyclohexyl can be connected either through two separate chemical bonds to two distinct carbons atoms within the ring; or the two bonds can be connected to the same carbon atom in the ring. In an illustrative example, if a divalent cyclopropyl radical connects two phenyl rings together, this definition encompasses both 1,2-diphenylcyclopropyl and 1,1-diphenylcyclopropyl units.

As used herein the singular forms "a", "an", and "the" include plural referents unless the context clearly dictates otherwise. For example, "a compound" refers to one or more of such compounds, while "the enzyme" includes a particular enzyme as well as other family members and equivalents thereof as known to those skilled in the art. As used in the specification and appended claims, unless specified to the contrary, the following terms have the meaning indicated.

The term "absent" as used herein means the group is replaced by a single bond. If replacing the group with a bond results in two connected moieties both defined as bonds, then -bond-bond- groups are understood to reduce to a single bond.

The term "interrupted by" as used herein means the group specified is inserted at any point within the specified chain, but not at the termini. For example, if a C₃-alkyl chain, as defined herein, is interrupted by -O-, then the following groups would be encompassed: -CH₂-O-CH₂CH₂-, -CH₂CH₂-O-CH₂-, -CH(CH₃)-O-CH₂-, and -CH₂-O-CH(CH₃)-.

The terms "aliphatic" and "aliphatic group" as used herein means straight-chain, branched or cyclic C₁-C₁₂ (unless stated otherwise) hydrocarbon radicals which are completely saturated or which contain one or more units of unsaturation but which are not aromatic. For example, suitable aliphatic groups include substituted or unsubstituted linear, branched or cyclic alkyl, alkenyl, alkynyl groups and hybrids thereof such as (cycloalkyl)alkyl, (cycloalkenyl)alkyl or (cycloalkyl)alkenyl.

The terms "alkyl", "alkoxy", "hydroxyalkyl", "alkoxyalkyl", and "alkoxycarbonyl", used alone or as part of a larger moiety include both straight and branched chains containing one to twelve carbon atoms.

The terms "alkenyl" and "alkynyl" used alone or as part of a larger moiety include both straight and branched chains containing two to twelve carbon atoms.

The term "alkoxy" refers to an -O-alkyl radical, where alkyl is defined herein.

"Alkyl" refers to a straight or branched hydrocarbon chain radical consisting solely of carbon and hydrogen atoms, containing no unsaturation, having from one to twelve carbon atoms, preferably one to eight, and which is attached to the rest of the molecule by a single bond, *e.g.*, methyl, ethyl, *n*-propyl, 1-methylethyl (iso-propyl), *n*-butyl, *n*-pentyl, 1,1-dimethylethyl (*t*-butyl), and the like. Unless
5 stated otherwise specifically in the specification, the alkyl radical is optionally substituted by one or more substituents selected from the group consisting of halo, cyano, nitro, -OR¹¹, -N(R¹¹)₂, -COR¹¹, -COOR¹¹, -CON(R¹¹)₂, -N(R¹¹)COOR¹⁰, -N(R¹¹)COR¹¹, -NSO₂ R¹¹, -N(R¹¹)SO₂ R¹¹, -SO₂OR¹¹, -SO₂R¹¹, and -SO₂N(R¹¹)₂ where each R¹⁰ and R¹¹ are as defined above in the First aspect of the invention. Unless stated otherwise specifically in the specification, it is understood that for radicals, as
10 defined below, that contain a substituted alkyl group that the substitution can occur on any carbon of the alkyl group.

"Alkenyl" refers to a straight or branched hydrocarbon chain radical consisting solely of carbon and hydrogen atoms, containing at least one double bond, having from two to eight carbon atoms, and which is attached to the rest of the molecule by a single bond or a double bond, *e.g.*, ethenyl, prop-1-
15 enyl, but-1-enyl, pent-1-enyl, penta-1,4-dienyl, and the like. Unless stated otherwise specifically in the specification, the alkenyl radical is optionally substituted by one or more substituents selected from the group consisting of halo, cyano, nitro, -OR¹¹, -N(R¹¹)₂, -COR¹¹, -COOR¹¹, -CON(R¹¹)₂, -N(R¹¹)COOR¹⁰, -N(R¹¹)COR¹¹, -NSO₂R¹¹, -N(R¹¹)SO₂R¹¹, -SO₂OR¹¹, -SO₂R¹¹, and -SO₂N(R¹¹)₂ where each R¹⁰ and R¹¹ are as defined above in the First aspect of the invention. Unless stated otherwise
20 specifically in the specification, it is understood that for radicals, as defined below, that contain a substituted alkenyl group that the substitution can occur on any carbon of the alkenyl group.

"Aryl" refers to aromatic monocyclic or multicyclic ring system containing from 6 to 19 carbon atoms, where the ring system is optionally partially or fully saturated. Aryl groups include, but are not limited to groups such as fluorenyl, phenyl and naphthyl. Unless stated otherwise specifically in the
25 specification, the term "aryl" is meant to include aryl radicals optionally substituted by one or more substituents selected from the group consisting of alkyl, alkenyl, halogen, haloalkyl, haloalkenyl, cyano, nitro, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heteroaryl, heteroarylalkyl, heterocyclyl, heterocyclylalkyl, R⁰-OR¹¹, R⁰-N(R¹¹)₂, R⁰-COR¹¹, R⁰-COOR¹¹, R⁰-CON(R¹¹)₂, R⁰-N(R¹¹)COOR¹⁰, R⁰-N(R¹¹)COR¹¹, R⁰-NSO₂R¹¹, R⁰-N(R¹¹)SO₂R¹¹, R⁰-SO₂OR¹¹, R⁰-SO₂R¹¹, and R⁰-SO₂N(R¹¹)₂ where
30 each R⁰ is independently selected from a substituted or an unsubstituted aliphatic group, an unsubstituted heteroaryl or heterocyclic ring, phenyl (Ph), substituted Ph, -OPh, substituted -OPh, or substituted -CH₂Ph. Examples of substituents on the aliphatic group or phenyl ring of R⁰ include amino, alkylamino, dialkylamino, aminocarbonyl, halogen, alkyl, alkylaminocarbonyl, dialkylaminocarbonyl,

alkylaminocarbonyloxy, dialkylaminocarbonyloxy, alkoxy, nitro, cyano, carboxy, alkoxy carbonyl, alkyl carbonyl, hydroxy, haloalkoxy or haloalkyl.

An aliphatic group or non-aromatic heterocyclic ring may contain one or more substituents. Examples of suitable substituents on the saturated carbon of an aliphatic group or of non-aromatic heterocyclic ring include those listed above for unsaturated carbon of an aryl or heteroaryl group and including the following: $=O$, $=S$, $=NNHR^0$, $=NN(R^0)_2$, $=N-$, $=NNHC(O)R^0$, $=NNHCO_2$ (alkyl), $=NNHSO_2$ (alkyl), or $=NR^0$, where R^0 is independently selected from hydrogen, unsubstituted or substituted aliphatic group, an unsubstituted heteroaryl or heterocyclic ring, phenyl (Ph), substituted Ph, -OPh, substituted -OPh, $-CH_2Ph$ or substituted $-CH_2Ph$. Examples of substituents on the aliphatic group include amino, alkylamino, dialkylamino, aminocarbonyl, halogen, alkyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylaminocarbonyloxy, dialkylaminocarbonyloxy, alkoxy, nitro, cyano, carboxy, alkoxy carbonyl, alkyl carbonyl, hydroxy, haloalkoxy or haloalkyl.

Suitable substituents on the nitrogen of a non-aromatic heterocyclic ring include $-R^0$, $-N(R^0)_2$, $-C(O)R^0$, CO_2R^0 , $-C(O)C(O)R^0$, $-SO_2R$, $-SO_2N(R^0)_2$, $-C(=S)N(R^0)_2$, $-C(=NH)-N(R^0)_2$, and $NR^0RSO_2R^0$ wherein each R^0 is independently selected from hydrogen, unsubstituted or substituted aliphatic group, an unsubstituted heteroaryl or heterocyclic ring, phenyl (Ph), substituted Ph, -OPh, substituted -OPh, or substituted $-CH_2Ph$. Examples of substituents on the aliphatic group or the phenyl ring include amino, alkylamino, dialkylamino, aminocarbonyl, halogen, alkyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylaminocarbonyloxy, dialkylaminocarbonyloxy, alkoxy, nitro, cyano, carboxy, alkoxy carbonyl, alkyl carbonyl, hydroxy, haloalkoxy or haloalkyl.

The term "alkoxyaryl" as used herein means an aryl group, as defined herein, substituted with one or more alkoxy groups, as defined herein. Examples of alkoxyaryl groups include, but are not limited to, methoxyphenyl, butyloxyphenyl, and dimethoxynaphthyl.

"Aralkyl" or "arylalkyl" refers to a radical of the formula $-RaRb$ where Ra is an alkyl radical as defined above and Rb is one or more aryl radicals as defined above, *e.g.*, benzyl, diphenylmethyl and the like. The aryl radical(s) and the alkyl radical is optionally substituted as described above.

The term "aralkyloxy" or "arylalkoxy" as used herein, means an aralkyl group, as defined herein, appended to the parent molecule through a oxygen atom. Examples of aralkyloxy include, but are not limited to, benzyloxy, 2-phenylethoxy, 4-phenylbutoxy, 9-fluorenylmethoxy, and the like.

The term "arylalkylcarboxy" as used herein, means an arylalkyl group, as defined herein, appended to the parent molecule through a carboxy group, as defined herein. The carboxy group can be bonded in either sense; either with the carbonyl carbon bonded to the arylalkyl group and the oxygen bonded to the parent molecule; or the carbonyl bonded to the parent molecule and the oxygen bonded to

the arylalkyl group. Examples of arylalkylcarboxy groups include, but are not limited to, benzylacetoxo, (benzyloxy)carbonyl, (2-phenylethoxy)carbonyl, phenyl-acetyloxy, and 1-oxo-5-phenyl-pentyloxy.

5 The term "aryloxy" as used herein, means an aryl group, as defined herein, appended to a parent molecule through an oxygen atom. Examples of "aryloxy" groups include, but are not limited to phenoxy, 1-naphthyloxy, and 2-naphthyloxy.

"Alkylene" and "alkylene chain" refer to a straight or branched divalent hydrocarbon chain, linking the rest of the molecule to a radical group, consisting solely of carbon and hydrogen, containing no unsaturation and having from one to twelve carbon atoms, preferably having from one to eight carbons, *e.g.*, methylene, ethylene, propylene, *n*-butylene, and the like. The alkylene chain may be
10 attached to the rest of the molecule and to the radical group through one carbon within the chain or through any two carbons within the chain. The alkylene chain is optionally substituted by one or more substituents selected from the group consisting of halo, cyano, nitro, $-OR^{11}$, $-N(R^{11})_2$, $-COR^{11}$, $-COOR^{11}$, $-CON(R^{11})_2$, $-N(R^{11})COOR^{10}$, $-N(R^{11})COR^{11}$, $-NSO_2R^{11}$, $-N(R^{11})SO_2R^{11}$, $-SO_2OR^{11}$, $-SO_2R^{11}$, and $-SO_2N(R^{11})_2$ where each R^{10} and R^{11} are as defined above in the First aspect of the invention. The
15 alkylene chain may be attached to the rest of the molecule through any two carbons within the chain.

"Alkenylene" and "alkenylene chain" refer to a straight or branched divalent hydrocarbon chain linking the rest of the molecule to a radical group, consisting solely of carbon and hydrogen, containing at least one double bond and having from two to twelve carbon atoms, *e.g.*, ethenylene, propenylene, *n*-butenylene, and the like. The alkenylene chain is attached to the rest of the molecule through a single
20 bond and to the radical group through a double bond or a single bond. The points of attachment of the alkenylene chain to the rest of the molecule and to the radical group can be through one carbon or any two carbons within the chain. The alkenylene chain is optionally substituted by one or more substituents selected from the group consisting of halo, cyano, nitro, $-OR^{11}$, $-N(R^{11})_2$, $-COR^{11}$, $-COOR^{11}$, $-CON(R^{11})_2$, $-N(R^{11})COOR^{10}$, $-N(R^{11})COR^{11}$, $-NSO_2R^{11}$, $-N(R^{11})SO_2R^{11}$, $-SO_2OR^{11}$, $-SO_2R^{11}$, and $-SO_2N(R^{11})_2$ where each R^{10} and R^{11} are as defined above in the First aspect of the invention.
25

The term "aryloxyalkyl" as used herein, means an alkyl group appended to the parent molecule, wherein the alkyl group is substituted with one aryloxy group, as defined herein. Examples of aryloxyalkyl groups include, but are not limited to phenoxymethyl, naphthyloxybutyl, and phenoxyhexyl.

30 The term "aryloxyaryl" as used herein, means an aryl group appended to the parent molecule, wherein the aryl group is substituted with one aryloxy group, as defined herein. Examples of aryloxyaryl groups include, but are not limited to phenoxyphenyl, naphthyloxyphenyl, and phenoxy-naphthyl.

The term "carbonyl" as used herein, means a $-C(=O)-$ group.

The term "carboxy" as used herein, means a $-C(=O)O-$ group.

"Cycloalkyl" refers to a stable monovalent monocyclic or bicyclic hydrocarbon radical consisting solely of carbon and hydrogen atoms, having from three to ten carbon atoms (unless stated otherwise), and which is saturated or includes one more unsaturated units (but is not aromatic) and is attached to the rest of the molecule by a single bond, e.g., cyclopropyl, cyclobutyl, cyclopentyl, cyclopent-1-enyl, cyclohexyl, cyclohex-2,4-dienyl, decalyl and the like. Unless otherwise stated specifically in the specification, the term "cycloalkyl" is meant to include cycloalkyl radicals which are optionally substituted by one or more substituents independently selected from the group consisting of alkyl, alkenyl, halo, haloalkyl, haloalkenyl, cyano, nitro, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-OR^{11}$, $-N(R^{11})_2$, $-COR^{11}$, $-COOR^{11}$, $-CON(R^{11})_2$, $-N(R^{11})COOR^{10}$, $-N(R^{11})COR^{11}$, $-NSO_2R^{11}$, $-N(R^{11})SO_2R^{11}$, $-SO_2OR^{11}$, $-SO_2R^{11}$, and $-SO_2N(R^{11})_2$ where each R^{10} and R^{11} are as defined above in the first aspect of the invention.

"Cycloalkylalkyl" refers to a radical of the formula $-RaRd$ where Ra is an alkyl radical as defined above and Rd is a cycloalkyl radical as defined above. The alkyl radical and the cycloalkyl radical may be optionally substituted as defined above.

The term "cyclohaloalkyl" as used herein means a cycloalkyl group, as defined herein which is substituted by one or more halo groups, as defined herein. Examples of "cyclohaloalkyl" groups include, but are not limited to, bromocyclohexyl, trifluorocyclopentyl, dichlorocyclohexyl and the like.

"Halo" or "Halogen" refers to bromo, chloro, fluoro or iodo.

"Haloalkyl" refers to an alkyl radical, as defined above, that is substituted by one or more halo radicals, as defined above, e.g., trifluoromethyl, difluoromethyl, trichloromethyl, 2,2,2-trifluoroethyl, 1-fluoromethyl-2-fluoroethyl, 3-bromo-2-fluoropropyl, 1-bromomethyl-2-bromoethyl, and the like.

"Haloalkenyl" refers to an alkenyl radical, as defined above, that is substituted by one or more halo radicals, as defined above, e.g., 2-bromoethenyl, 3-bromoprop-1-enyl, and the like.

The term "haloaryl" as used herein, means an aryl group, as defined herein, substituted with one or more halo groups. Examples of haloaryl groups include, but are not limited to, bromophenyl, fluorophenyl, pentafluorophenyl, chloronaphthyl, chloro-iodophenyl, and the like.

"Heterocyclyl" refers to a stable 3- to 18-membered non-aromatic ring radical which consists of carbon atoms and from one to five heteroatoms selected from the group consisting of nitrogen, oxygen and sulfur. For purposes of this invention, the heterocyclyl radical may be a monocyclic, bicyclic, tricyclic or tetracyclic ring system, which may include fused or bridged ring systems; and the nitrogen, carbon or sulfur atoms in the heterocyclyl radical is optionally oxidized; the nitrogen atom is optionally quaternized; and the heterocyclyl radical may be partially or fully saturated. Examples of such heterocyclyl radicals include, but are not limited to, dioxolanyl, decahydroisoquinolyl, imidazolyl,

imidazolidinyl, isothiazolidinyl, isoxazolidinyl, morpholinyl, octahydroindolyl, octahydroisoindolyl, 2-oxopiperazinyl, 2-oxopiperidinyl, 2-oxopyrrolidinyl, oxazolidinyl, piperidinyl, piperazinyl, 4-piperidonyl, pyrrolidinyl, pyrazolidinyl, thiazolidinyl, tetrahydrofuranyl, trithianyl, tetrahydropyranyl, thiomorpholinyl, thiamorpholinyl, 1-oxo-thiomorpholinyl, and 1,1-dioxo-thiomorpholinyl. Unless
5 stated otherwise specifically in the specification, the term "heterocyclyl" is meant to include heterocyclyl radicals as defined above which are optionally substituted by one or more substituents selected from the group consisting of alkyl, alkenyl, halo, haloalkyl, haloalkenyl, nitro, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-OR^{11}$, $-N(R^{11})_2$, $-COR^{11}$, $-COOR^{11}$, $-CON(R^{11})_2$, $-N(R^{11})COOR^{10}$, $-N(R^{11})COR^{11}$, $-NSO_2R^{11}$, $-N(R^{11})SO_2R^{11}$, $-SO_2OR^{11}$, $-SO_2R^{11}$, and $-SO_2N(R^{11})_2$ where each R^{10} and R^{11} are as defined above in the First aspect of the
10 invention.

"Heterocyclylalkyl" refers to a radical of the formula $-RaRe$ where Ra is an alkyl radical as defined above and Re is a heterocyclyl radical as defined above, and if the heterocyclyl is a nitrogen-containing heterocyclyl, the heterocyclyl may be attached to the alkyl radical at the nitrogen atom. The
15 heterocyclyl radical and the alkyl radical are optionally substituted as defined above.

The term "heterocyclyloxy" as used herein, means a heterocyclyl group, as defined herein, appended to a parent molecule through an oxygen atom. Examples of "heterocyclyloxy" groups include, but are not limited to piperidinylloxy, tetrahydrofuranylloxy, tetrahydrotheinyloxy tetrahydropyranylloxy, dihydropyranylloxy, pyrrolidinylloxy, oxetanylloxy, and oxiranylloxy.

"Heteroaryl" refers to a 3- to 18-membered aromatic ring radical which consists of carbon atoms and from one to five heteroatoms selected from the group consisting of nitrogen, oxygen and sulfur. For purposes of this invention, the heteroaryl radical may be a monocyclic, bicyclic, tricyclic or tetracyclic ring system, which may include fused or bridged ring systems; and the nitrogen, carbon or sulfur atoms in the heteroaryl radical is optionally oxidized; the nitrogen atom is optionally quaternized.
20 Examples include, but are not limited to, azepinyl, acridinyl, benzimidazolyl, benzthiazolyl, benzindolyl, benzothiadiazolyl, benzonaphthofuranyl, benzoxazolyl, benzodioxolyl, benzodioxinyl, benzopyranyl, benzopyranonyl, benzofuranyl, benzofuranonyl, benzothieryl (benzothiophenyl), benzotriazolyl, benzo[4,6]imidazo[1,2-a]pyridinyl, carbazolyl, cinnoliny, dibenzofuranyl, furanyl, furanonyl, isothiazolyl, imidazolyl, indolyl, indazolyl, isoindolyl, indoliny, isoindoliny, indoliziny, isoxazolyl, naphthyridinyl, oxadiazolyl, 2-oxoazepinyl, oxazolyl, oxiranyl, phenazinyl, phenothiazinyl,
25 phenoxazinyl, phthalazinyl, phthalimidyl pteridinyl, purinyl, pyrrolyl, pyrazolyl, pyridinyl, pyrazinyl, pyrimidinyl, pyridazinyl, quinazoliny, quinoxaliny, quinoliny, quinuclidiny, isoquinoliny, thiazolyl, thiadiazolyl, triazolyl, tetrazolyl, triazinyl, and thiophenyl. Unless stated otherwise specifically in the specification, the term "heteroaryl" is meant to include heteroaryl radicals as defined above which are
30

optionally substituted by one or more substituents selected from the group consisting of alkyl, alkenyl, halo, haloalkyl, haloalkenyl, nitro, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, -OR¹¹, -N(R¹¹)₂, -COR¹¹, -COOR¹¹, -CON(R¹¹)₂, -N(R¹¹)COOR¹⁰, -N(R¹¹)COR¹¹, -NSO₂R¹¹, -N(R¹¹)SO₂R¹¹, -SO₂OR¹¹, -SO₂R¹¹, and -SO₂N(R¹¹)₂ where each R¹⁰ and R¹¹ are as defined above in the First aspect of the invention. For purposes of this invention, the term "N-heteroaryl" refers to heteroaryl radicals as defined above containing at least one nitrogen atom in ring.

The term "heteroaryloxy" as used herein, means a heteroaryl group, as defined herein, appended to a parent molecule through an oxygen atom. Examples of "heteroaryloxy" groups include, but are not limited to pyridyloxy, indolyloxy, and quinolyloxy.

"Heteroarylalkyl" refers to a radical of the formula -R_aR_f where R_a is an alkyl radical as defined above and R_f is a heteroaryl radical as defined above, and if the heteroaryl is a nitrogen-containing heteroaryl, the heteroaryl may be attached to the alkyl radical at the nitrogen atom. The heteroaryl radical and the alkyl radical are optionally substituted as defined above.

The term "linker group" or "linker" means an organic moiety that connects two parts of a compound. Linkers are typically comprised of an atom such as oxygen or sulfur, a unit such as -NH-, -CH₂-, -CO-, -CONH-, or a chain of atoms, such as an alidyl chain. The molecular mass of a linker is typically in the range of about 14 to 200, preferably in the range of 14 to 96 with a length of up to about six atoms. Examples of linkers include a saturated or unsaturated C₁₋₆ alidyl chain which is optionally substituted, and wherein one or two saturated carbons of the chain are optionally replaced by -CO-, -COCO-, -CONH-, -CONHNH-, -CO₂-, -NHCO₂-, -O-, -NHCONH-, -OCONH-, -NHNH-, -NHCO-, -S-, -SO-, -SO₂-, -NH-, -SO₂NH-, or -NHSO₂-.

The term "alidyl" or "alidyl chain" refers to an optionally substituted, straight or branched carbon chain that may be fully saturated or have one or more units of unsaturation. The optional substituents are as described above for an aliphatic group. Alidyl chain used herein may include alidyl chains containing 0-4 fluorine substituents. An "agonist for a nuclear receptor" is an agent that, when bound to the nuclear receptor, activates nuclear receptor activity to activate or repress gene function. In some cases, nuclear receptors can act through second messenger signaling pathways, and the invention would apply to these actions as well. The activation can be similar in degree to that provided by a natural hormone for the receptor, or can be stronger (optionally referred to as a "strong agonist"), or can be weaker (optionally referred to as a "weak agonist" or "partial agonist"). An example of a hormone for a nuclear receptor is thyroid hormone, which is a natural hormone for the thyroid receptor. A "putative agonist" is an agent to be tested for agonist activity.

Partial agonists or partial antagonists bind to receptors and yield a response less than that of a full agonist at saturating ligand concentrations. A partial agonist will block binding of a full agonist and suppress receptor activity to the level induced by the partial agonist alone. For example, partial agonists bind to receptors and induce only part of the changes in the receptors that are induced by agonists. The differences can be qualitative or quantitative. Thus, a partial agonist can induce some of the conformation changes induced by agonists, but not others, or it may only induce certain changes to a limited extent. Some of these compounds are naturally produced. For example, many plant estrogens (phytoestrogens), such as genistein, can behave as partial estrogen receptor agonists.

An "antagonist for a nuclear receptor" is an agent that reduces or blocks activity mediated by the receptor in response to an agonist of the receptor. The activity of the antagonist can be mediated, *e.g.*, by blocking binding of the agonist to the receptor, or by altering receptor configuration and/or activity of the receptor. A "putative antagonist" is an agent to be tested for antagonist activity.

A "nuclear receptor" is a receptor that activates or represses transcription of one or more genes in the nucleus (but can also have second messenger signaling actions), typically in conjunction with other transcription factors. The nuclear receptor is activated by the natural cognate ligand for the receptor. Nuclear receptors are ordinarily found in the cytoplasm or nucleus, rather than being membrane-bound. Nuclear receptor is a member of a superfamily of regulatory proteins that are receptors for, *e.g.*, steroids, retinoids, vitamin D and thyroid hormones. These proteins bind to cis-acting elements in the promoters of their target genes and modulate gene expression in response to a ligand therefor. Nuclear receptors may be classified based on their DNA binding properties. For example, the glucocorticoid, estrogen, androgen, progestin and mineralocorticoid receptors bind as homodimers to hormone response elements (HREs) organized as inverted repeats. Another example are receptors, including those activated by retinoic acid, thyroid hormone, vitamin D₃, fatty acids/peroxisome proliferators and ecdysone, that bind to HREs as heterodimers with a common partner, the retinoid X receptor (RXR). Among the latter receptors is LXR.

As used herein, an orphan nuclear receptor is a nuclear receptor for which the natural ligand is unknown.

As used herein, liver X receptor or LXR refers to a nuclear receptor implicated in cholesterol biosynthesis. As used herein, the term LXR refers to both LXR α and LXR β , two forms of the protein found in mammals. Liver X receptor- α , or LXR α refers to the receptor described in U.S. Pat. Nos. 5,571,696, 5,696,233 and 5,710,004, and Willy et al. (1995) *Gene Dev.* 9(9):1033-1045. Liver X receptor- β or LXR β refers to the receptor described in Peet et al. (1998) *Curr. Opin. Genet. Dev.*

8(5):571-575; Song et al. (1995) Ann. N.Y. Acad. Sci. 761:38-49; Alberti et al. (2000) Gene 243(1-2):93-103; and references cited therein; and in U.S. Pat. Nos. 5,571,696, 5,696,233 and 5,710,004.

As used herein, compounds which are "commercially available" may be obtained from standard commercial sources including Acros Organics (Pittsburgh PA), Aldrich Chemical (Milwaukee WI, including Sigma Chemical and Fluka), Apin Chemicals Ltd. (Milton Park UK), Avocado Research (Lancashire U.K.), BDH Inc. (Toronto, Canada), Bionet (Cornwall, U.K.), Chemservice Inc. (West Chester PA), Crescent Chemical Co. (Hauppauge NY), Eastman Organic Chemicals, Eastman Kodak Company (Rochester NY), Fisher Scientific Co. (Pittsburgh PA), Fisons Chemicals (Leicestershire UK), Frontier Scientific (Logan UT), ICN Biomedicals, Inc. (Costa Mesa CA), Key Organics (Cornwall U.K.), Lancaster Synthesis (Windham NH), Maybridge Chemical Co. Ltd. (Cornwall U.K.), Parish Chemical Co. (Orem UT), Pfaltz & Bauer, Inc. (Waterbury CN), Polyorganix (Houston TX), Pierce Chemical Co. (Rockford IL), Riedel de Haen AG (Hannover, Germany), Spectrum Quality Product, Inc. (New Brunswick, NJ), TCI America (Portland OR), Trans World Chemicals, Inc. (Rockville MD), and Wako Chemicals USA, Inc. (Richmond VA).

As used herein, "suitable conditions" for carrying out a synthetic step are explicitly provided herein or may be discerned by reference to publications directed to methods used in synthetic organic chemistry. The reference books and treatise set forth above that detail the synthesis of reactants useful in the preparation of compounds of the present invention, will also provide suitable conditions for carrying out a synthetic step according to the present invention.

As used herein, "methods known to one of ordinary skill in the art" may be identified through various reference books and databases. Suitable reference books and treatise that detail the synthesis of reactants useful in the preparation of compounds of the present invention, or provide references to articles that describe the preparation, include for example, "Synthetic Organic Chemistry", John Wiley & Sons, Inc., New York; S. R. Sandler *et al.*, "Organic Functional Group Preparations," 2nd Ed., Academic Press, New York, 1983; H. O. House, "Modern Synthetic Reactions", 2nd Ed., W. A. Benjamin, Inc. Menlo Park, Calif. 1972; T. L. Gilchrist, "Heterocyclic Chemistry", 2nd Ed., John Wiley & Sons, New York, 1992; J. March, "Advanced Organic Chemistry: Reactions, Mechanisms and Structure", 4th Ed., Wiley-Interscience, New York, 1992. Specific and analogous reactants may also be identified through the indices of known chemicals prepared by the Chemical Abstract Service of the American Chemical Society, which are available in most public and university libraries, as well as through on-line databases (the American Chemical Society, Washington, D.C. may be contacted for more details). Chemicals that are known but not commercially available in catalogs may be prepared by custom chemical synthesis houses, where many of the standard chemical supply houses (e.g., those listed above) provide custom synthesis services.

"Prodrugs" is meant to indicate a compound that may be converted under physiological conditions or by solvolysis to a biologically active compound of the invention. Thus, the term "prodrug" refers to a metabolic precursor of a compound of the invention that is pharmaceutically acceptable. A prodrug may be inactive when administered to a subject in need thereof, but is converted *in vivo* to an active compound of the invention. Prodrugs are typically rapidly transformed *in vivo* to yield the parent compound of the invention, for example, by hydrolysis in blood. The prodrug compound often offers advantages of solubility, tissue compatibility or delayed release in a mammalian organism (see, Bundgard, H., *Design of Prodrugs* (1985), pp. 7-9, 21-24 (Elsevier, Amsterdam). A discussion of prodrugs is provided in Higuchi, T., *et al.*, "Pro-drugs as Novel Delivery Systems," A.C.S. Symposium Series, Vol. 14, and in *Bioreversible Carriers in Drug Design*, ed. Edward B. Roche, American Pharmaceutical Association and Pergamon Press, 1987, both of which are incorporated in full by reference herein. The term "prodrug" is also meant to include any covalently bonded carriers which release the active compound of the invention *in vivo* when such prodrug is administered to a mammalian subject. Prodrugs of a compound of the invention may be prepared by modifying functional groups present in the compound of the invention in such a way that the modifications are cleaved, either in routine manipulation or *in vivo*, to the parent compound of the invention. By virtue of knowledge of pharmacodynamic processes and drug metabolism *in vivo*, those of skill in this art, once a pharmaceutically active compound is known, can design prodrugs of the compound (see, e.g., Nogrady (1985) *Medicinal Chemistry A Biochemical Approach*, Oxford University Press, New York, pages 388-392). Prodrugs include compounds of the invention wherein a hydroxy, amino or mercapto group is bonded to any group that, when the prodrug of the compound of the invention is administered to a mammalian subject, cleaves to form a free hydroxy, free amino or free mercapto group, respectively. Examples of prodrugs include, but are not limited to, acetate, formate and benzoate derivatives of alcohol and amine functional groups in the compounds of the invention and the like.

"Polymorph" refers to the different crystal forms of a compound, resulting from the possibility of at least two different arrangements of the molecules of the compound in the solid state. Polymorphs of a given compound will be different in crystal structure but identical in liquid or vapor states. Different polymorphic forms of a given substance may differ from each other with respect to one or more physical properties, such as solubility and dissociation, true density, crystal shape, compaction behavior, flow properties, and/or solid state stability.

"Stable compound" and "stable structure" are meant to indicate a compound that is sufficiently robust to survive isolation to a useful degree of purity from a reaction mixture, and formulation into an efficacious therapeutic agent.

"Mammal" includes humans and domestic animals, such as cats, dogs, swine, cattle, sheep, goats, horses, rabbits, and the like.

"Optional" or "optionally" means that the subsequently described event of circumstances may or may not occur, and that the description includes instances where said event or circumstance occurs and instances in which it does not. For example, "optionally substituted aryl" means that the aryl radical may or may not be substituted and that the description includes both substituted aryl radicals as defined herein and aryl radicals having no substitution.

"Pharmaceutically acceptable carrier, diluent or excipient" includes without limitation any adjuvant, carrier, excipient, glidant, sweetening agent, diluent, preservative, dye/colorant, flavor enhancer, surfactant, wetting agent, dispersing agent, suspending agent, stabilizer, isotonic agent, solvent, or emulsifier which has been approved by the United States Food and Drug Administration as being acceptable for use in humans or domestic animals.

"Pharmaceutically acceptable salt" includes both acid and base addition salts.

"Pharmaceutically acceptable acid addition salt" refers to those salts which retain the biological effectiveness and properties of the free bases, which are not biologically or otherwise undesirable, and which are formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid and the like, and organic acids such as acetic acid, trifluoroacetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, maleic acid, malonic acid, succinic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, salicylic acid, and the like.

"Pharmaceutically acceptable base addition salt" refers to those salts which retain the biological effectiveness and properties of the free acids, which are not biologically or otherwise undesirable. These salts are prepared from addition of an inorganic base or an organic base to the free acid. Salts derived from inorganic bases include, but are not limited to, the sodium, potassium, lithium, ammonium, calcium, magnesium, iron, zinc, copper, manganese, aluminum salts and the like. Preferred inorganic salts are the ammonium, sodium, potassium, calcium, and magnesium salts. Salts derived from organic bases include, but are not limited to, salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins, such as isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine, ethanolamine, 2-dimethylaminoethanol, 2-diethylaminoethanol, dicyclohexylamine, lysine, arginine, histidine, caffeine, procaine, hydrabamine, choline, betaine, ethylenediamine, glucosamine, methylglucamine, theobromine, purines, piperazine, piperidine, N-ethylpiperidine, polyamine resins and the like. Particularly preferred organic bases are isopropylamine, diethylamine, ethanolamine, trimethylamine, dicyclohexylamine, choline and caffeine.

"Pharmaceutically acceptable derivative" refers to pharmaceutically acceptable salts as defined herein and also includes esters, prodrugs, solvates and polymorphs of the compounds of the invention.

"Therapeutically effective amount" refers to that amount of a compound of the invention which, when administered to a mammal, preferably a human, is sufficient to effect treatment, as defined below, for a disease-state associated with nuclear receptor activity. The amount of a compound of the invention which constitutes a "therapeutically effective amount" will vary depending on the compound, the condition and its severity, and the age of the mammal to be treated, but can be determined routinely by one of ordinary skill in the art having regard to his own knowledge and to this disclosure.

"Modulating" or "modulate" refers to the treating, prevention, suppression, enhancement or induction of a function or condition. For example, the compounds of the present invention can modulate hyperlipidemia by lowering cholesterol in a human, thereby suppressing hyperlipidemia.

"Treating" or "treatment" as used herein covers the treatment of a disease or condition associated with the nuclear receptor activity as disclosed herein, in a mammal, preferably a human, and includes:

- i. Preventing a disease or condition associated with the nuclear receptor activity from occurring in a mammal, in particular, when such mammal is predisposed to the disease or condition but has not yet been diagnosed as having it;
- ii. inhibiting a disease or condition associated with the nuclear receptor activity, *i.e.*, arresting its development; or
- iii. relieving a disease or condition associated with the nuclear receptor activity, *i.e.*, causing regression of the condition.

The compounds of formulae IIaa, IIbb, IIcc, or IIdd or their pharmaceutically acceptable salts may contain one or more asymmetric centers and may thus give rise to enantiomers, diastereomers, and other stereoisomeric forms that may be defined, in terms of absolute stereochemistry, as (*R*)- or (*S*)- or, as (*D*)- or (*L*)- for amino acids. The present invention is meant to include all such possible isomers, as well as, their racemic and optically pure forms. Optically active (+) and (-), (*R*)- and (*S*)-, or (*D*)- and (*L*)- isomers may be prepared using chiral synthons or chiral reagents, or resolved using conventional techniques, such as reverse phase HPLC. When the compounds described herein contain olefinic double bonds or other centers of geometric asymmetry, and unless specified otherwise, it is intended that the compounds include both *E* and *Z* geometric isomers. It will be apparent to one skilled in the art that certain compounds of this invention may exist in tautomeric forms, all such tautomeric forms of the compounds being within the scope of the invention. Unless otherwise stated, structures depicted herein are also meant to include all stereochemical forms of the structure; *i.e.*, the *R* and *S* configurations for each asymmetric center. Therefore, single stereochemical isomers as well as enantiomeric and

diastereomeric mixtures of the present compounds are within the scope of the invention. Unless otherwise stated, structures depicted herein are also meant to include compounds which differ only in the presence of one or more isotopically enriched atoms. For example, compounds having the present structure except for the replacement of a hydrogen atom by a deuterium or tritium, or the replacement of
5 a carbon atom by a ^{13}C - or ^{14}C -enriched carbon are within the scope of this invention.

The chemical naming protocol and structure diagrams used herein employ and rely on the chemical naming features as utilized by the ChemDraw program (available from CambridgeSoft Corp., Cambridge, MA). In particular, the compound names were derived from the structures using the Autonom program as utilized by Chemdraw Ultra or ISIS base (MDL Corp.).

10 The term "atherosclerosis" refers to process whereby atherosclerotic plaques form within the inner lining of the artery wall leading to atherosclerotic cardiovascular diseases. Atherosclerotic cardiovascular diseases can be recognized and understood by physicians practicing in the relevant fields of medicine, and include without limitation, restenosis, coronary heart disease (also known as coronary artery heart disease or ischemic heart disease), cerebrovascular disease including ischemic stroke, multi-
15 infarct dementia, and peripheral vessel disease, including intermittent claudication, and erectile dysfunction.

The term "dyslipidemia" refers to abnormal levels of lipoproteins in blood plasma including both depressed and/or elevated levels of lipoproteins (*e.g.*, elevated levels of Low Density Lipoprotein, (LDL), Very Low Density Lipoprotein (VLDL) and depressed levels of High Density Lipoprotein
20 (HDL) (less than 40 mg/dL)).

As used herein, "EC₅₀" refers to a dosage, concentration or amount of a particular test compound that elicits a dose-dependent response at 50% of maximal expression of a particular response that is induced, provoked or potentiated by the particular test compound.

25 The term "cholesterol" refers to a steroid alcohol that is an essential component of cell membranes and myelin sheaths and, as used herein, incorporates its common usage. Cholesterol also serves as a precursor for steroid hormones and bile acids.

The term "triglyceride(s)" ("TGs"), as used herein, incorporates its common usage. TGs consist of three fatty acid molecules esterified to a glycerol molecule and serve to store fatty acids which are used by muscle cells for energy production or are taken up and stored in adipose tissue.

30 The term "hyperlipidemia" refers to the presence of an abnormally elevated level of lipids in the blood. Hyperlipidemia can appear in at least three forms (ES): (1) hypercholesterolemia, *i.e.*, an elevated LDL cholesterol level (120 mg/dL and above); (2) hypertriglyceridemia, *i.e.*, an elevated triglyceride level; (150 mg/dL and above) and (3) combined hyperlipidemia, *i.e.*, a combination of hypercholesterolemia and hypertriglyceridemia.

Exemplary Primary Hyperlipidemia include, but are not limited to, the following:

(1) Familial Hyperchylomicronemia, a rare genetic disorder which causes a deficiency in an enzyme, LP lipase, that breaks down fat molecules. The LP lipase deficiency can cause the accumulation of large quantities of fat or lipoproteins in the blood;

(2) Familial Hypercholesterolemia, a relatively common genetic disorder caused where the underlying defect is a series of mutations in the LDL receptor gene that result in malfunctioning LDL receptors and/or absence of the LDL receptors. This brings about ineffective clearance of LDL by the LDL receptors resulting in elevated LDL and total cholesterol levels in the plasma;

(3) Familial Combined Hyperlipidemia, also known as multiple lipoprotein-type hyperlipidemia; an inherited disorder where patients and their affected first-degree relatives can at various times manifest high cholesterol and high triglycerides. Levels of HDL cholesterol are often moderately decreased;

(4) Familial Defective Apolipoprotein B-100 is a relatively common autosomal dominant genetic abnormality. The defect is caused by a single nucleotide mutation that produces a substitution of glutamine for arginine which can cause reduced affinity of LDL particles for the LDL receptor. Consequently, this can cause high plasma LDL and total cholesterol levels;

Familial Dysbetalipoproteinemia, also referred to as Type III Hyperlipoproteinemia, is an uncommon inherited disorder resulting in moderate to severe elevations of serum triglyceride (TG) and cholesterol levels with abnormal apolipoprotein E function. HDL levels are usually normal; and

Familial Hypertriglyceridemia, is a common inherited disorder in which the concentration of plasma VLDL is elevated. This can cause mild to moderately elevated triglyceride levels (and usually not cholesterol levels) and can often be associated with low plasma HDL levels.

Risk factors in exemplary Secondary Hyperlipidemia include, but are not limited to, the following: (1) disease risk factors, such as a history of type 1 diabetes, type 2 diabetes, Cushing's syndrome, hypothyroidism and certain types of renal failure; (2) drug risk factors, which include, birth control pills; hormones, such as estrogen, and corticosteroids; certain diuretics; and various beta-blockers; (3) dietary risk factors include dietary fat intake per total calories greater than 40%; saturated fat intake per total calories greater than 10%; cholesterol intake greater than 300 mg per day; habitual and excessive alcohol use; and obesity; and (4) non-genetic dyslipidemias.

The methods of the present invention can be used effectively in combination with one or more additional active diabetes agents depending on the desired target therapy (see, *e.g.*, Turner, N. et al. *Prog. Drug Res.* (1998) 51:33-94; Haffner, S. *Diabetes Care* (1998) 21: 160178; and DeFronzo, R. et al. (eds.), *Diabetes Reviews* (1997) Vol. 5 No. 4). A number of studies have investigated the benefits of

combination therapies with oral agents (see, *e.g.*, Mahler, R., J. Clin. Endocrinol. Metab. (1999)84:1165-71; United Kingdom Prospective Diabetes Study Group: UKPDS 28, Diabetes Care (1998)21:87-92; Bardin, C.W.(ed.), CURRENT THERAPY IN ENDOCRINOLOGY AND METABOLISM, 6th Edition (Mosby--Year Book, Inc., St. Louis, Mo. 1997); Chiasson, J. et al., Ann. Intern. Med. (1994) 121: 928-935; Coniff, R. et al., Clin.Ther. (1997) 19: 16-26; Coniff, R. et al., Am. J. Med. (1995) 98: 443-451; and Iwamoto, Y. et al, Diabet. Med. (1996)13: 365-370; Kwitterovich, P. Am. J. Cardiol (1998) 82(12A):3U-17U). These studies indicate that diabetes and hyperlipidemia modulation can be further improved by the addition of a second agent to the therapeutic regimen. As used herein, "IC₅₀" refers to an amount, concentration or dosage of a particular test compound that achieves a 50% inhibition of a maximal response, such as modulation of nuclear receptor, including the LXR α or LXR β activity, in an assay that measures such response.

As used herein, "LXR α " (LXR alpha) refers to all mammalian forms of such receptor including, for example, alternative splice isoforms and naturally occurring isoforms. Representative LXR α species include, without limitation the rat (Genbank Accession NM_031627), mouse (Genbank Accession BC012646), and human (GenBank Accession No. U22662) forms of the receptor.

As used herein, "LXR β " (LXR beta) refers to all mammalian forms of such receptor including, for example, alternative splice isoforms and naturally occurring isoforms. Representative LXR β species include, without limitation the rat (GenBank Accession NM_031626), mouse (Genbank Accession NM_009473), and human (GenBank Accession No. U07132) forms of the receptor.

As used herein "LXR" or "LXRs" refers to both LXR α and LXR β .

The terms "obese" and "obesity" refers to a Body Mass Index (BMI) greater than 27.8 kg/m² for men and 27.3 kg/m² for women (BMI equals weight (kg)/(height)²(m²).

Use of the Compounds of the Invention

The compounds of the invention exhibit valuable pharmacological properties in mammals, and are particularly useful as selective LXR agonists, antagonists, inverse agonists, partial agonists and antagonists, for the treatment, or prevention of diseases associated with, or symptoms arising from the complications of, altered cholesterol transport, cholesterol reverse transport, fatty acid metabolism, cholesterol absorption, cholesterol re-absorption, cholesterol secretion, cholesterol excretion, or cholesterol metabolism.

These diseases include, for example, hyperlipidemia, dyslipidemia, hypercholesterolemia, atherosclerosis, atherosclerotic cardiovascular diseases, hyperlipoproteinemia, (see, *e.g.*, Patent Application Publication Nos. WO 00/57915 and WO 00/37077), hyperglycemia, insulin resistance,

diabetes, lipodystrophy, obesity, syndrome X (US Patent Application No. 20030073614, International Patent Application Publication No. WO 01/82917), excess lipid deposition in peripheral tissues such as skin (xanthomas) (see, e.g., U.S. Patent Nos. 6,184,215 and 6,187,814), stroke, peripheral occlusive disease, memory loss (*Brain Research* (1997), Vol. 752, pp. 189-196), optic nerve and retinal pathologies (i.e., macular degeneration, retinitis pigmentosa), repair of traumatic damage to the central or peripheral nervous system (*Trends in Neurosciences* (1994), Vol. 17, pp. 525-530), prevention of the degenerative process due to aging (*American Journal of Pathology* (1997), Vol. 151, pp. 1371-1377), Parkinson's disease or Alzheimer's disease (see, e.g., International Patent Application Publication No. WO 00/17334; *Trends in Neurosciences* (1994), Vol. 17, pp. 525-530), prevention of degenerative neuropathies occurring in diseases such as diabetic neuropathies (see, e.g., International Patent Application Publication No. WO 01/82917), multiple sclerosis (*Annals of Clinical Biochem.* (1996), Vol. 33, No. 2, pp. 148-150), and autoimmune diseases (*J. Lipid Res.* (1998), Vol. 39, pp. 1740-1743).

Also provided, are methods of increasing the expression of ATP-Binding Cassette (ABCA1), (see, e.g., International Patent Application Publication No. WO 00/78972) thereby increasing reverse cholesterol transport in mammalian cells using the claimed compounds and compositions.

Accordingly in another aspect, the invention also includes methods to remove cholesterol from tissue deposits such as atherosclerotic plaques or xanthomas in a patient with atherosclerosis or atherosclerotic cardiovascular disease manifest by clinical signs of such disease, wherein the methods comprise administering to the patient a therapeutically effective amount of a compound or composition of the present invention. Additionally, the instant invention also provides a method for preventing or reducing the risk of a first or subsequent occurrence of an atherosclerotic cardiovascular disease event including ischemic heart disease, ischemic stroke, multi-infarct dementia, and intermittent claudication comprising the administration of a prophylactically effective amount of a compound or composition of the present invention to a patient at risk for such an event. The patient may already have atherosclerotic cardiovascular disease at the time of administration, or may be at risk for developing it. Risk factors for developing atherosclerotic cardiovascular disease events include increasing age (65 and over), male gender, a family history of atherosclerotic cardiovascular disease events, high blood cholesterol (especially LDL or "bad" cholesterol over 100 mg/dL), cigarette smoking and exposure to tobacco smoke, high blood pressure, diabetes, obesity and physical inactivity.

Also contemplated herein is the use of a compound of the invention, or a pharmaceutically acceptable derivative thereof, in combination with one or more of the following therapeutic agents in treating atherosclerosis: antihyperlipidemic agents, plasma HDL-raising agents, antihypercholesterolemic agents, cholesterol biosynthesis inhibitors (such as HMG CoA reductase inhibitors, such as lovastatin, simvastatin, pravastatin, fluvastatin, atorvastatin and rivastatin), acyl-

coenzyme A:cholesterol acyltransferase (ACAT) inhibitors, probucol, raloxifene, nicotinic acid, niacinamide, cholesterol absorption inhibitors, bile acid sequestrants (such as anion exchange resins, or quaternary amines (*e.g.*, cholestyramine or colestipol)), low density lipoprotein receptor inducers, clofibrate, fenofibrate, benzofibrate, ciprofibrate, gemfibrozil, vitamin B₆, vitamin B₁₂, anti-oxidant
5 vitamins, β -blockers, anti-diabetes agents, angiotensin II antagonists, angiotensin converting enzyme inhibitors, platelet aggregation inhibitors, fibrinogen receptor antagonists, aspirin or fibric acid derivatives.

In one embodiment compounds of the invention are used in combination with a cholesterol biosynthesis inhibitor, particularly an HMG-CoA reductase inhibitor. The term HMG-CoA reductase
10 inhibitor is intended to include all pharmaceutically acceptable salt, ester, free acid and lactone forms of compounds which have HMG-CoA reductase inhibitory activity and, therefore, the use of such salts, esters, free acids and lactone forms is included within the scope of this invention. Compounds which have inhibitory activity for HMG-CoA reductase can be readily identified using assays well-known in the art. For instance, suitable assays are described or disclosed in U.S. Patent No. 4,231,938 and WO
15 84/02131. Examples of suitable HMG-CoA reductase inhibitors include, but are not limited to, lovastatin (MEVACOR®; *see*, U.S. Patent No. 4,231,938); simvastatin (ZOCOR®; *see*, U.S. Patent No. 4,444,784); pravastatin sodium (PRAVACHOL®; *see*, U.S. Patent No. 4,346,227); fluvastatin sodium (LESCOL®; *see*, U.S. Patent No. 5,354,772); atorvastatin calcium (LIPITOR®; *see*, U.S. Patent No. 5,273,995) and rivastatin (also known as cerivastatin; *see*, U.S. Patent No. 5,177,080). The
20 structural formulas of these and additional HMG-CoA reductase inhibitors that can be used in combination with the compounds of the invention are described at page 87 of M. Yalpani, "Cholesterol Lowering Drugs," *Chemistry & Industry*, pp. 85-89 (5 February 1996). In presently preferred embodiments, the HMG-CoA reductase inhibitor is selected from lovastatin and simvastatin.

The compounds of the present invention can also be used in methods for decreasing
25 hyperglycemia and insulin resistance, *i.e.*, in methods for treating diabetes (International Patent Application Publication No. WO 01/82917), and in methods of treatment, prevention, or amelioration of disorders related to, or arising as complications of diabetes, hyperglycemia or insulin resistance including the cluster of disease states, conditions or disorders that make up "Syndrome X" (See US Patent Application 20030073614) comprising the administration of a therapeutically effective amount of
30 a compound or composition of the present invention to a patient in need of such treatment. Additionally, the instant invention also provides a method for preventing or reducing the risk of developing hyperglycemia, insulin resistance, diabetes or syndrome X in a patient, comprising the administration of a prophylactically effective amount of a compound or composition of the present invention to a patient at risk for such an event.

Diabetes mellitus, commonly called diabetes, refers to a disease process derived from multiple causative factors and characterized by elevated levels of plasma glucose, referred to as hyperglycemia. See, *e.g.*, LeRoith, D. et al., (eds.), DIABETES MELLITUS (Lippincott-Raven Publishers, Philadelphia, Pa. U.S.A. 1996). According to the American Diabetes Association, diabetes mellitus is estimated to affect approximately 6% of the world population. Uncontrolled hyperglycemia is associated with increased and premature mortality due to an increased risk for macrovascular and macrovascular diseases, including nephropathy, neuropathy, retinopathy, hypertension, cerebrovascular disease and coronary heart disease. Therefore, control of glucose homeostasis is a critically important approach for the treatment of diabetes.

There are two major forms of diabetes: type 1 diabetes (formerly referred to as insulin-dependent diabetes or IDDM); and type 2 diabetes (formerly referred to as noninsulin dependent diabetes or NIDDM).

Type 2 diabetes is a disease characterized by insulin resistance accompanied by relative, rather than absolute, insulin deficiency. Type 2 diabetes can range from predominant insulin resistance with relative insulin deficiency to predominant insulin deficiency with some insulin resistance. Insulin resistance is the diminished ability of insulin to exert its biological action across a broad range of concentrations. In insulin resistant individuals, the body secretes abnormally high amounts of insulin to compensate for this defect. When inadequate amounts of insulin are present to compensate for insulin resistance and adequate control of glucose, a state of impaired glucose tolerance develops. In a significant number of individuals, insulin secretion declines further and the plasma glucose level rises, resulting in the clinical state of diabetes. Type 2 diabetes can be due to a profound resistance to insulin stimulating regulatory effects on glucose and lipid metabolism in the main insulin-sensitive tissues: muscle, liver and adipose tissue. This resistance to insulin responsiveness results in insufficient insulin activation of glucose uptake, oxidation and storage in muscle and inadequate insulin repression of lipolysis in adipose tissue and of glucose production and secretion in liver. In Type 2 diabetes, free fatty acid levels are often elevated in obese and some non-obese patients and lipid oxidation is increased.

Premature development of atherosclerosis and increased rate of cardiovascular and peripheral vascular diseases are characteristic features of patients with diabetes. Hyperlipidemia is an important precipitating factor for these diseases. Hyperlipidemia is a condition generally characterized by an abnormal increase in serum lipids, *e.g.*, cholesterol and triglyceride, in the bloodstream and is an important risk factor in developing atherosclerosis and heart disease. For a review of disorders of lipid metabolism, see, *e.g.*, Wilson, J. et al., (ed.), Disorders of Lipid Metabolism, Chapter 23, Textbook of Endocrinology, 9th Edition, (W. B. Sanders Company, Philadelphia, Pa. U.S.A. 1998). Hyperlipidemia is usually classified as primary or secondary hyperlipidemia. Primary hyperlipidemia is generally

caused by genetic defects, while secondary hyperlipidemia is generally caused by other factors, such as various disease states, drugs, and dietary factors. Alternatively, hyperlipidemia can result from both a combination of primary and secondary causes of hyperlipidemia. Elevated cholesterol levels are associated with a number of disease states, including coronary artery disease, angina pectoris, carotid artery disease, strokes, cerebral arteriosclerosis, and xanthoma.

Dyslipidemia, or abnormal levels of lipoproteins in blood plasma, is a frequent occurrence among diabetics, and has been shown to be one of the main contributors to the increased incidence of coronary events and deaths among diabetic subjects (see, *e.g.*, Joslin, E. Ann. Chim. Med. (1927), Vol. 5, pp. 1061-1079). Epidemiological studies since then have confirmed the association and have shown a several-fold increase in coronary deaths among diabetic subjects when compared with non-diabetic subjects (see, *e.g.*, Garcia, M. J. et al., Diabetes (1974), Vol. 23, pp. 105-11 (1974); and Laakso, M. and Lehto, S., Diabetes Reviews (1997), Vol. 5, No. 4, pp. 294-315). Several lipoprotein abnormalities have been described among diabetic subjects (Howard B., et al., Arteriosclerosis (1978), Vol. 30, pp. 153-162).

The compounds of the invention can also be used effectively in combination with one or more additional active diabetes agents depending on the desired target therapy (see, *e.g.*, Turner, N. et al., Prog. Drug Res. (1998), Vol. 51, pp.33-94; Haffner, S., Diabetes Care (1998), Vol. 21, pp. 160-178; and DeFronzo, R. et al. (eds.), Diabetes Reviews (1997), Vol. 5, No. 4). A number of studies have investigated the benefits of combination therapies with oral agents (see, *e.g.*, Mahler, R., J. Clin. Endocrinol. Metab. (1999), Vol. 84, pp. 1165-71; United Kingdom Prospective Diabetes Study Group: UKPDS 28, Diabetes Care (1998), Vol. 21, pp. 87-92; Bardin, C. W.(ed.), CURRENT THERAPY IN ENDOCRINOLOGY AND METABOLISM, 6th Edition (Mosby--Year Book, Inc., St. Louis, Mo. 1997); Chiasson, J. et al., Ann. Intern. Med. (1994), Vol. 121, pp. 928-935; Coniff, R. et al., Clin. Ther. (1997), Vol.19, pp. 16-26; Coniff, R. et al., Am. J. Med. (1995), Vol. 98, pp. 443-451; Iwamoto, Y. et al., Diabet. Med. (1996), Vol. 13, pp. 365-370; Kwiterovich, P., Am. J. Cardiol (1998), Vol. 82 (12A), pp. 3U-17U). These studies indicate that diabetes and hyperlipidemia modulation can be further improved by the addition of a second agent to the therapeutic regimen.

Accordingly, the compounds of the invention may be used in combination with one or more of the following therapeutic agents in treating diabetes: sulfonylureas (such as chlorpropamide, tolbutamide, acetohexamide, tolazamide, glyburide, glipizide, glynnase, glimepiride, and glipizide), biguanides (such as metformin), thiazolidinediones (such as ciglitazone, pioglitazone, troglitazone, and rosiglitazone), and related insulin sensitizers, such as selective and non-selective activators of PPAR α , PPAR β and PPAR γ ; dehydroepiandrosterone (also referred to as DHEA or its conjugated sulphate ester,

DHEA-SO₄); antilglucocorticoids; TNF α inhibitors; α -glucosidase inhibitors (such as acarbose, miglitol, and voglibose), pramlintide (a synthetic analog of the human hormone amylin), other insulin secretagogues (such as repaglinide, gliquidone, and nateglinide), insulin, as well as the therapeutic agents discussed above for treating atherosclerosis.

5 Further provided by this invention are methods of using the compounds of the invention to treat obesity, as well as the complications of obesity. Obesity is linked to a variety of medical conditions including diabetes and hyperlipidemia. Obesity is also a known risk factor for the development of type 2 diabetes (See, *e.g.*, Barrett-Conner, E., *Epidemol. Rev.* (1989), Vol. 11, pp. 172-181; and Knowler, et al., *Am. J Clin. Nutr.* (1991), Vol. 53, pp. 1543-1551).

10 In addition, the compounds of the invention can be used in combination with agents used in treated obesity or obesity-related disorders. Such agents, include, but are not limited to, phenylpropanolamine, phentermine, diethylpropion, mazindol, fenfluramine, dexfenfluramine, phentramine, β_3 adrenoceptor agonist agents; sibutramine, gastrointestinal lipase inhibitors (such as orlistat), and leptins. Other agents used in treating obesity or obesity-related disorders include
15 neuropeptide Y, enterostatin, cholecystokinin, bombesin, amylin, histamine H₃ receptors, dopamine D₂ receptor modulators, melanocyte stimulating hormone, corticotrophin releasing factor, galanin and gamma amino butyric acid (GABA).

Evaluation of the Use of the Compounds of the Invention

Standard physiological, pharmacological and biochemical procedures are available for testing
20 the compounds to identify those that possess biological activities that modulate the activity of nuclear receptors, including the LXRs (LXR α and LXR β). Such assays include, for example, biochemical assays such as binding assays, fluorescence polarization assays, FRET based coactivator recruitment assays (see, generally, Glickman et al., *J. Biomolecular Screening* (2002), Vol. 7, No. 1, pp. 3-10, as well as cell based assays including the co-transfection assay, the use of LBD-Gal 4 chimeras and
25 protein-protein interaction assays, (see, Lehmann. et al., *J. Biol Chem.* (1997), Vol. 272, No. 6, pp. 3137-3140).

High throughput screening systems are commercially available (see, *e.g.*, Zymark Corp., Hopkinton, MA; Air Technical Industries, Mentor, OH; Beckman Instruments Inc., Fullerton, CA; Precision Systems, Inc., Natick, MA) that enable these assays to be run in a high throughput mode.
30 These systems typically automate entire procedures, including all sample and reagent pipetting, liquid dispensing timed incubations, and final readings of the microplate in detector(s) appropriate for the assay. These configurable systems provide high throughput and rapid start up as well as a high degree of flexibility and customization. The manufacturers of such systems provide detailed protocols for

various high throughput systems. Thus, for example, Zymark Corp. provides technical bulletins describing screening systems for detecting the modulation of gene transcription, ligand binding, and the like.

Assays that do not require washing or liquid separation steps are preferred for such high throughput screening systems and include biochemical assays such as fluorescence polarization assays (see, for example, Owicki, J., *Biomol. Screen* (2000 October), Vol. 5, No. 5, pp. 297), scintillation proximity assays (SPA) (see, for example, Carpenter et al., *Methods Mol. Biol.* (2002), Vol 190, pp. 31-49) and fluorescence resonance energy transfer (FRET) or time resolved FRET based coactivator recruitment assays (Mukherjee et al., *J. Steroid Biochem. Mol. Biol.* (2002 July); Vol. 81, No. 3, pp. 217-25; (Zhou et al., *Mol. Endocrinol.* (1998 October), Vol. 12, No. 10, pp. 1594-604). Generally such assays can be preformed using either the full length receptor, or isolated ligand binding domain (LBD). In the case of LXR α , the LBD comprises amino acids 188-447, for LXR β the LBD comprises amino acids 198-461, and for FXR, the LBD comprises amino acids 244 to 472 of the full length sequence.

If a fluorescently labeled ligand is available, fluorescence polarization assays provide a way of detecting binding of compounds to the nuclear receptor of interest by measuring changes in fluorescence polarization that occur as a result of the displacement of a trace amount of the label ligand by the compound. Additionally this approach can also be used to monitor the ligand dependent association of a fluorescently labeled coactivator peptide to the nuclear receptor of interest to detect ligand binding to the nuclear receptor of interest.

The ability of a compound to bind to a receptor, or heterodimer complex with RXR, can also be measured in a homogeneous assay format by assessing the degree to which the compound can compete off a radiolabelled ligand with known affinity for the receptor using a scintillation proximity assay (SPA). In this approach, the radioactivity emitted by a radiolabelled compound (for example, [^3H] 24,25 Epoxycholesterol) generates an optical signal when it is brought into close proximity to a scintillant such as a YSI-copper containing bead, to which the nuclear receptor is bound. If the radiolabelled compound is displaced from the nuclear receptor the amount of light emitted from the nuclear receptor bound scintillant decreases, and this can be readily detected using standard microplate liquid scintillation plate readers such as, for example, a Wallac MicroBeta reader.

The heterodimerization of LXR with RXR α can also be measured by fluorescence resonance energy transfer (FRET), or time resolved FRET, to monitor the ability of the compounds provided herein to bind to LXR or other nuclear receptors. Both approaches rely upon the fact that energy transfer from a donor molecule to an acceptor molecule only occurs when donor and acceptor are in

close proximity. Typically the purified LBD of the nuclear receptor of interest is labeled with biotin then mixed with stoichiometric amounts of europium labeled streptavidin (Wallac Inc.), and the purified LBD of RXR α is labeled with a suitable fluorophore such as CY5TM. Equimolar amounts of each modified LBD are mixed together and allowed to equilibrate for at least 1 hour prior to addition to either variable or constant concentrations of the sample for which the affinity is to be determined. After equilibration, the time-resolved fluorescent signal is quantitated using a fluorescent plate reader. The affinity of the compound can then be estimated from a plot of fluorescence versus concentration of compound added.

This approach can also be exploited to measure the ligand dependent interaction of a co-activator peptide with a nuclear receptor in order to characterize the agonist or antagonist activity of the compounds disclosed herein. Typically the assay in this case involves the use a recombinant Glutathione-S-transferase (GST)-nuclear receptor ligand binding domain (LBD) fusion protein and a synthetic biotinylated peptide sequenced derived from the receptor interacting domain of a co-activator peptide such as the steroid receptor coactivator 1 (SRC-1). Typically GST-LBD is labeled with a europium chelate (donor) via a europium-tagged anti-GST antibody, and the coactivator peptide is labeled with allophycocyanin via a streptavidin-biotin linkage.

In the presence of an agonist for the nuclear receptor, the peptide is recruited to the GST-LBD bringing europium and allophycocyanin into close proximity to enable energy transfer from the europium chelate to the allophycocyanin. Upon excitation of the complex with light at 340 nm excitation energy absorbed by the europium chelate is transmitted to the allophycocyanin moiety resulting in emission at 665 nm. If the europium chelate is not brought in to close proximity to the allophycocyanin moiety there is little or no energy transfer and excitation of the europium chelate results in emission at 615 nm. Thus the intensity of light emitted at 665 nm gives an indication of the strength of the protein-protein interaction. The activity of a nuclear receptor antagonist can be measured by determining the ability of a compound to competitively inhibit (*i.e.*, IC₅₀) the activity of an agonist for the nuclear receptor

In addition, a variety of cell based assay methodologies may be successfully used in screening assays to identify and profile the specificity of compounds of the present invention. These approaches include the co-transfection assay, translocation assays, complementation assays and the use of gene activation technologies to over express endogenous nuclear receptors.

Three basic variants of the co-transfection assay strategy exist, co-transfection assays using full-length nuclear receptor, co transfection assays using chimeric nuclear receptors comprising the ligand

binding domain of the nuclear receptor of interest fused to a heterologous DNA binding domain, and assays based around the use of the mammalian two hybrid assay system.

The basic co-transfection assay is based on the co-transfection into the cell of an expression plasmid to express the nuclear receptor of interest in the cell with a reporter plasmid comprising a reporter gene whose expression is under the control of DNA sequence that is capable of interacting with that nuclear receptor (see, for example, US Patents Nos. 5,071,773; 5,298,429 and 6,416,957). Treatment of the transfected cells with an agonist for the nuclear receptor increases the transcriptional activity of that receptor which is reflected by an increase in expression of the reporter gene which may be measured by a variety of standard procedures.

For those receptors that function as heterodimers with RXR, such as the LXRs, the co-transfection assay typically includes the use of expression plasmids for both the nuclear receptor of interest and RXR. Typical co-transfection assays require access to the full length nuclear receptor and suitable response elements that provide sufficient screening sensitivity and specificity to the nuclear receptor of interest.

Typically, the expression plasmid comprises: (1) a promoter, such as an SV40 early region promoter, HSV tk promoter or phosphoglycerate kinase (pgk) promoter, CMV promoter, Sro₃ promoter or other suitable control elements known in the art, (2) a cloned polynucleotide sequence, such as a cDNA encoding a receptor, co-factor, or fragment thereof, ligated to the promoter in sense orientation so that transcription from the promoter will produce a RNA that encodes a functional protein, and (3) a polyadenylation sequence. For example and not limitation, an expression cassette of the invention may comprise the cDNA expression cloning vectors, or other preferred expression vectors known and commercially available from vendors such as Invitrogen, (CA), Stratagene, (CA) or Clontech, (CA). Alternatively expression vectors developed by academic groups such as the pCMX vectors originally developed in the Evans lab (Willey et al. *Genes & Development* 9 1033-1045 (1995)) may also be used.

The transcriptional regulatory sequences in an expression cassette are selected by the practitioner based on the intended application; depending upon the specific use, transcription regulation can employ inducible, repressible, constitutive, cell-type specific, developmental stage-specific, sex-specific, or other desired type of promoter or control sequence.

Alternatively, the expression plasmid may comprise an activation sequence to activate or increase the expression of an endogenous chromosomal sequence. Such activation sequences include for example, a synthetic zinc finger motif (for example, see US Patents 6,534,261 and 6,503,717) or a strong promoter or enhancer sequence together with a targeting sequence to enable homologous or non-homologous recombination of the activating sequence upstream of the gene of interest.

Genes encoding the following full-length previously described proteins, which are suitable for use in the co-transfection studies and profiling the compounds described herein, include human LXR α (accession U22662), human LXR β (accession U07132), rat FXR (accession U18374), human FXR (accession NM_005123), human RXR α (accession NM_002957), human RXR β (accession XM_042579), human RXR γ (accession XM_053680), human PPAR α (accession X57638) and human PPAR δ (accession U10375). All accession numbers in this application refer to GenBank accession numbers.

Reporter plasmids may be constructed using standard molecular biological techniques by placing cDNA encoding for the reporter gene downstream from a suitable minimal promoter. For example luciferase reporter plasmids may be constructed by placing cDNA encoding firefly luciferase (typically with SV40 small t intron and poly-A tail, (de Wet et al., (1987) Mol. Cell. Biol. 7 725-735) down stream from the herpes virus thymidine kinase promoter (located at nucleotides residues-105 to +51 of the thymidine kinase nucleotide sequence, obtained for example, from the plasmid pBLCAT2 (Luckow & Schutz (1987) Nucl. Acid. Res.15 5490-5494)) which is linked in turn to the appropriate response element (RE).

The choice of hormone response element is dependent upon the type of assay to be used. In the case of the use of the full-length LXR α or LXR β a reporter plasmid comprising a known LXR RE would typically be used, such as for example in a reporter plasmid such as LXREx1-tk-luciferase, (see U.S. patent No. 5,747,661, which is hereby incorporated by reference). In the case of a LXR α or LXR β -LBD-Gal4 fusion, GAL4 Upstream Activating Sequences (UAS) would be used. Typically the GAL4 UAS would comprise the sequence 5'CGGRNNRCYNYNCNCCG-3', where Y = C or T, R = A or G, and N = A, C, T or G, and would be present as a tandem repeat of 4 copies.

Numerous methods of co-transfecting the expression and reporter plasmids are known to those of skill in the art and may be used for the co-transfection assay to introduce the plasmids into a suitable cell type. Typically such a cell will not endogenously express nuclear receptors that interact with the response elements used in the reporter plasmid.

Numerous reporter gene systems are known in the art and include, for example, alkaline phosphatase (see, Berger, J., et al., Gene (1988), Vol. 66, pp. 1-10; and Kain, S.R., Methods. Mol. Biol. (1997), Vol. 63, pp. 49-60), β -galactosidase (See, U.S. Patent No. 5,070,012, issued Dec, 3, 1991 to Nolan et al., and Bronstein, I., et al., J. Chemilum. Biolum. (1989), Vol. 4, pp. 99-111), chloramphenicol acetyltransferase (See, Gorman et al., Mol. Cell Biol. (1982), Vol. 2, pp. 1044-51), β -glucuronidase, peroxidase, β -lactamase (U.S. Patent Nos. 5,741,657 and 5,955,604), catalytic antibodies, luciferases

(U.S. Patents 5,221,623; 5,683,888; 5,674,713; 5,650,289; and 5,843,746) and naturally fluorescent proteins (Tsien, R.Y., *Annu. Rev. Biochem.* (1998), Vol. 67, pp. 509-44).

The use of chimeras comprising the ligand binding domain (LBD) of the nuclear receptor of interest to a heterologous DNA binding domain (DBD) expands the versatility of cell based assays by directing activation of the nuclear receptor in question to defined DNA binding elements recognized by defined DNA binding domain (see WO95/18380). This assay expands the utility of cell based co-transfection assays in cases where the biological response or screening window using the native DNA binding domain is not satisfactory.

In general the methodology is similar to that used with the basic co-transfection assay, except that a chimeric construct is used in place of the full length nuclear receptor. As with the full length nuclear receptor, treatment of the transfected cells with an agonist for the nuclear receptor LBD increases the transcriptional activity of the heterologous DNA binding domain which is reflected by an increase in expression of the reporter gene as described above. Typically for such chimeric constructs, the DNA binding domains from defined nuclear receptors, or from yeast or bacterially derived transcriptional regulators such as members of the GAL 4 and Lex A / UmuD super families are used.

A third cell based assay of utility for screening compounds of the present invention is a mammalian two-hybrid assay that measures the ability of the nuclear hormone receptor to interact with a cofactor in the presence of a ligand (see, for example, US Patent Nos. US 5,667,973, 5,283,173 and 5,468,614). The basic approach is to create three plasmid constructs that enable the interaction of the nuclear receptor with the interacting protein to be coupled to a transcriptional readout within a living cell. The first construct is an expression plasmid for expressing a fusion protein comprising the interacting protein, or a portion of that protein containing the interacting domain, fused to a GAL4 DNA binding domain. The second expression plasmid comprises DNA encoding the nuclear receptor of interest fused to a strong transcription activation domain such as VP16, and the third construct comprises the reporter plasmid comprising a reporter gene with a minimal promoter and GAL4 upstream activating sequences.

Once all three plasmids are introduced into a cell, the GAL4 DNA binding domain encoded in the first construct allows for specific binding of the fusion protein to GAL4 sites upstream of a minimal promoter. However because the GAL4 DNA binding domain typically has no strong transcriptional activation properties in isolation, expression of the reporter gene occurs only at a low level. In the presence of a ligand, the nuclear receptor-VP16 fusion protein can bind to the GAL4-interacting protein fusion protein bringing the strong transcriptional activator VP16 in close proximity to the GAL4 binding sites and minimal promoter region of the reporter gene. This interaction significantly enhances the transcription of the reporter gene which can be measured for various reporter genes as described above.

Transcription of the reporter gene is thus driven by the interaction of the interacting protein and nuclear receptor of interest in a ligand dependent fashion.

Any compound which is a candidate for activation of LXR α or LXR β may be tested by these methods. Generally, compounds are tested at several different concentrations to optimize the chances
5 that activation of the receptor will be detected and recognized if present. Typically assays are performed in triplicate and vary within experimental error by less than 15%. Each experiment is typically repeated three or more times with similar results.

Activity of the reporter gene can be conveniently normalized to the internal control and the data plotted as fold activation relative to untreated cells. A positive control compound (agonist) may be
10 included along with DMSO as high and low controls for normalization of the assay data. Similarly, antagonist activity can be measured by determining the ability of a compound to competitively inhibit the activity of an agonist.

Additionally the compounds and compositions can be evaluated for their ability to increase or decrease the expression of genes known to be modulated by LXR α or LXR β and other nuclear
15 receptors *in vivo*, using Northern-blot, RT PCR or oligonucleotide microarray analysis to analyze RNA levels. Western-blot analysis can be used to measure expression of proteins encoded by LXR target genes. Genes that are known to be regulated by the LXRs include the ATP binding cassette transporters ABCA1, ABCG1, ABCG5, ABCG8, the sterol response element binding protein 1c (SREBP1c) gene, stearyl CoA desaturase 1 (SCD-1) and the apolipoprotein apoE gene (ApoE).

Established animal models exist for a number of diseases of direct relevance to the claimed
20 compounds and these can be used to further profile and characterize the claimed compounds. These model systems include diabetic dislipidemia using Zucker (fa/fa) rats or (db/db) mice, spontaneous hyperlipidemia using apolipoprotein E deficient mice (ApoE^{-/-}), diet-induced hyperlipidemia, using low density lipoprotein receptor deficient mice (LDLR^{-/-}) and atherosclerosis using both the Apo E^{-/-} and
25 LDLR^{-/-} mice fed a western diet. (21% fat, 0.05% cholesterol). Additionally LXR or FXR animal models (*e.g.*, knockout mice) can be used to further evaluate the present compounds and compositions *in vivo* (see, for example, Peet, et al., Cell (1998), Vol. 93, pp. 693-704, and Sinal, et al., Cell (2000), Vol. 102, pp. 731-744).

Administration of the Compounds of the Invention

Administration of the compounds of the invention, or their pharmaceutically acceptable salts, in
30 pure form or in an appropriate pharmaceutical composition, can be carried out via any of the accepted modes of administration of agents for serving similar utilities. The pharmaceutical compositions of the invention can be prepared by combining a compound of the invention with an appropriate

pharmaceutically acceptable carrier, diluent or excipient, and may be formulated into preparations in solid, semi-solid, liquid or gaseous forms, such as tablets, capsules, powders, granules, ointments, solutions, suppositories, injections, inhalants, gels, microspheres, and aerosols. Typical routes of administering such pharmaceutical compositions include, without limitation, oral, topical, transdermal, inhalation, parenteral, sublingual, rectal, vaginal, and intranasal. The term parenteral as used herein includes subcutaneous injections, intravenous, intramuscular, intrasternal injection or infusion techniques. Pharmaceutical compositions of the invention are formulated so as to allow the active ingredients contained therein to be bioavailable upon administration of the composition to a patient. Compositions that will be administered to a subject or patient take the form of one or more dosage units, where for example, a tablet may be a single dosage unit, and a container of a compound of the invention in aerosol form may hold a plurality of dosage units. Actual methods of preparing such dosage forms are known, or will be apparent, to those skilled in this art; for example, see Remington's Pharmaceutical Sciences, 18th Ed., (Mack Publishing Company, Easton, Pennsylvania, 1990). The composition to be administered will, in any event, contain a therapeutically effective amount of a compound of the invention, or a pharmaceutically acceptable salt thereof, for treatment of a disease-state associated with the activity of a nuclear receptor in accordance with the teachings of this invention.

A pharmaceutical composition of the invention may be in the form of a solid or liquid. In one aspect, the carrier(s) are particulate, so that the compositions are, for example, in tablet or powder form. The carrier(s) may be liquid, with the compositions being, for example, an oral syrup, injectable liquid or an aerosol, which is useful in, *e.g.*, inhalatory administration.

When intended for oral administration, the pharmaceutical composition is preferably in either solid or liquid form, where semi-solid, semi-liquid, suspension and gel forms are included within the forms considered herein as either solid or liquid.

As a solid composition for oral administration, the pharmaceutical composition may be formulated into a powder, granule, compressed tablet, pill, capsule, chewing gum, wafer or the like form. Such a solid composition will typically contain one or more inert diluents or edible carriers. In addition, one or more of the following may be present: binders such as carboxymethylcellulose, ethyl cellulose, microcrystalline cellulose, gum tragacanth or gelatin; excipients such as starch, lactose or dextrins, disintegrating agents such as alginic acid, sodium alginate, Primogel, corn starch and the like; lubricants such as magnesium stearate or Sterotex; glidants such as colloidal silicon dioxide; sweetening agents such as sucrose or saccharin; a flavoring agent such as peppermint, methyl salicylate or orange flavoring; and a coloring agent.

When the pharmaceutical composition is in the form of a capsule, *e.g.*, a gelatin capsule, it may contain, in addition to materials of the above type, a liquid carrier such as polyethylene glycol or oil.

The pharmaceutical composition may be in the form of a liquid, *e.g.*, an elixir, syrup, solution, emulsion or suspension. The liquid may be for oral administration or for delivery by injection, as two examples. When intended for oral administration, preferred composition contain, in addition to the present compounds, one or more of a sweetening agent, preservatives, dye/colorant and flavor enhancer.

5 In a composition intended to be administered by injection, one or more of a surfactant, preservative, wetting agent, dispersing agent, suspending agent, buffer, stabilizer and isotonic agent may be included.

The liquid pharmaceutical compositions of the invention, whether they be solutions, suspensions or other like form, may include one or more of the following adjuvants: sterile diluents such as water for injection, saline solution, preferably physiological saline, Ringer's solution, isotonic sodium
10 chloride, fixed oils such as synthetic mono or diglycerides which may serve as the solvent or suspending medium, polyethylene glycols, glycerin, propylene glycol or other solvents; antibacterial agents such as benzyl alcohol or methyl paraben; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. The parenteral preparation
15 can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic. Physiological saline is a preferred adjuvant. An injectable pharmaceutical composition is preferably sterile.

A liquid pharmaceutical composition of the invention intended for either parenteral or oral administration should contain an amount of a compound of the invention such that a suitable dosage will
20 be obtained. Typically, this amount is at least 0.01% of a compound of the invention in the composition. When intended for oral administration, this amount may be varied to be between 0.1 and about 70% of the weight of the composition. Preferred oral pharmaceutical compositions contain between about 4% and about 50% of the compound of the invention. Preferred pharmaceutical compositions and preparations according to the present invention are prepared so that a parenteral dosage unit contains
25 between 0.01 to 1% by weight of the compound of the invention.

The pharmaceutical composition of the invention may be intended for topical administration, in which case the carrier may suitably comprise a solution, emulsion, ointment or gel base. The base, for example, may comprise one or more of the following: petrolatum, lanolin, polyethylene glycols, bee wax, mineral oil, diluents such as water and alcohol, and emulsifiers and stabilizers. Thickening agents
30 may be present in a pharmaceutical composition for topical administration. If intended for transdermal administration, the composition may include a transdermal patch or iontophoresis device. Topical formulations may contain a concentration of the compound of the invention from about 0.1 to about 10% w/v (weight per unit volume).

The pharmaceutical composition of the invention may be intended for rectal administration, in the form, *e.g.*, of a suppository, which will melt in the rectum and release the drug. The composition for rectal administration may contain an oleaginous base as a suitable nonirritating excipient. Such bases include, without limitation, lanolin, cocoa butter and polyethylene glycol.

5 The pharmaceutical composition of the invention may include various materials, which modify the physical form of a solid or liquid dosage unit. For example, the composition may include materials that form a coating shell around the active ingredients. The materials that form the coating shell are typically inert, and may be selected from, for example, sugar, shellac, and other enteric coating agents. Alternatively, the active ingredients may be encased in a gelatin capsule.

10 The pharmaceutical composition of the invention in solid or liquid form may include an agent that binds to the compound of the invention and thereby assists in the delivery of the compound. Suitable agents that may act in this capacity include a monoclonal or polyclonal antibody, a protein or a liposome.

15 The pharmaceutical composition of the invention may consist of dosage units that can be administered as an aerosol. The term aerosol is used to denote a variety of systems ranging from those of colloidal nature to systems consisting of pressurized packages. Delivery may be by a liquefied or compressed gas or by a suitable pump system that dispenses the active ingredients. Aerosols of compounds of the invention may be delivered in single phase, bi-phasic, or tri-phasic systems in order to deliver the active ingredient(s). Delivery of the aerosol includes the necessary container, activators,
20 valves, subcontainers, and the like, which together may form a kit. One skilled in the art, without undue experimentation may determine preferred aerosols.

25 The pharmaceutical compositions of the invention may be prepared by methodology well known in the pharmaceutical art. For example, a pharmaceutical composition intended to be administered by injection can be prepared by combining a compound of the invention with sterile, distilled water so as to form a solution. A surfactant may be added to facilitate the formation of a homogeneous solution or suspension. Surfactants are compounds that non-covalently interact with the compound of the invention so as to facilitate dissolution or homogeneous suspension of the compound in the aqueous delivery system.

30 The compounds of the invention, or their pharmaceutically acceptable salts, are administered in a therapeutically effective amount, which will vary depending upon a variety of factors including the activity of the specific compound employed; the metabolic stability and length of action of the compound; the age, body weight, general health, sex, and diet of the patient; the mode and time of administration; the rate of excretion; the drug combination; the severity of the particular disorder or condition; and the subject undergoing therapy. Generally, a therapeutically effective daily dose is from

about 0.1 mg to about 20 mg/kg of body weight per day of a compound of the invention, or a pharmaceutically acceptable salt thereof; preferably, from about 0.1 mg to about 10 mg/kg of body weight per day; and most preferably, from about 0.1 mg to about 7.5 mg/kg of body weight per day.

Compounds of the invention, or pharmaceutically acceptable derivatives thereof, may also be administered simultaneously with, prior to, or after administration of one or more of the therapeutic agents described above in the Utility of the Compounds of the Invention. Such combination therapy includes administration of a single pharmaceutical dosage formulation which contains a compound of the invention and one or more additional active agents, as well as administration of the compound of the invention and each active agent in its own separate pharmaceutical dosage formulation. For example, a compound of the invention and an HMG-CoA reductase inhibitor can be administered to the patient together in a single oral dosage composition such as a tablet or capsule, or each agent administered in separate oral dosage formulations. Where separate dosage formulations are used, the compounds of the invention and one or more additional active agents can be administered at essentially the same time, *i.e.*, concurrently, or at separately staggered times, *i.e.*, sequentially; combination therapy is understood to include all these regimens.

Dosage information for HMG-CoA reductase inhibitors is well known in the art, since several HMG-CoA reductase inhibitors are marketed in the U.S. In particular, the daily dosage amounts of the HMG-CoA reductase inhibitor may be the same or similar to those amounts which are employed for anti-hypercholesterolemic treatment and which are described in the Physicians' Desk Reference (PDR). For example, see the 50th Ed. of the PDR, 1996 (Medical Economics Co); in particular, see at page 216 the heading "Hypolipidemics," sub-heading "HMG-CoA Reductase Inhibitors," and the reference pages cited therein. Preferably, the oral dosage amount of HMG-CoA reductase inhibitor is from about 1 to 200 mg/day and, more preferably, from about 5 to 160 mg/day. However, dosage amounts will vary depending on the potency of the specific HMG-CoA reductase inhibitor used as well as other factors as noted above. An HMG-CoA reductase inhibitor which has sufficiently greater potency may be given in sub-milligram daily dosages.

As examples, the daily dosage amount for simvastatin may be selected from 5 mg, 10 mg, 20 mg, 40 mg, 80 mg and 160 mg for lovastatin, 10 mg, 20 mg, 40 mg and 80 mg; for fluvastatin sodium, 20 mg, 40 mg and 80 mg; and for pravastatin sodium, 10 mg, 20 mg, and 40 mg. The daily dosage amount for atorvastatin calcium may be in the range of from 1 mg to 160 mg and, more particularly, from 5 mg to 80 mg. Oral administration may be in a single or divided doses of two, three, or four times daily, although a single daily dose of the HMG-CoA reductase inhibitor is preferred.

Preparation of the Compounds of the Invention

It is understood that in the following description, combinations of substituents and/or variables of the depicted formulae are permissible only if such contributions result in stable compounds.

It will also be appreciated by those skilled in the art that in the processes described below the functional groups of intermediate compounds may need to be protected by suitable protecting groups.

5 Such functional groups include hydroxy, amino, mercapto and carboxylic acid. Suitable protecting groups for hydroxy include trialkylsilyl or diarylalkylsilyl (*e.g.*, *t*-butyldimethylsilyl, *t*-butyldiphenylsilyl or trimethylsilyl), tetrahydropyranyl, benzyl, and the like. Suitable protecting groups for 1,2-dihydroxys include ketal- and acetal-forming groups. Suitable protecting groups for amino, amidino and guanidino include *t*-butoxycarbonyl, benzyloxycarbonyl, and the like. Suitable protecting groups for mercapto
10 include -C(O)-R (where R is alkyl, aryl or aralkyl), *p*-methoxybenzyl, trityl and the like. Suitable protecting groups for carboxylic acid include alkyl, aryl or aralkyl esters.

Protecting groups may be added or removed in accordance with standard techniques, which are well-known to those skilled in the art and as described herein. The use of protecting groups is described in detail in Green, T.W. and P.G.M. Wutz, *Protective Groups in Organic Synthesis* (1991), 2nd Ed.,
15 Wiley-Interscience. The protecting group may also be a polymer resin such as a Wang resin or a 2-chlorotrityl chloride resin.

It will also be appreciated by those skilled in the art, although such protected derivatives of compounds of the invention, as described above in the First aspect of the invention, may not possess pharmacological activity as such, they may be administered to a mammal having a disease associated
20 with defects in cholesterol transport, glucose metabolism, fatty acid metabolism and cholesterol metabolism, and thereafter metabolized in the body to form compounds of the invention which are pharmacologically active. Such derivatives may therefore be described as "prodrugs". All prodrugs of compounds of the invention are included within the scope of the invention.

It is understood that one of ordinary skill in the art would be able to make the compounds of the
25 invention not specifically prepared herein in light of the following disclosure, including the Preparations and Examples, and information known to those of ordinary skill in the chemical synthesis field.

Starting materials in the synthesis examples provided herein are either available from commercial sources or via literature procedures or by methods disclosed herein. All commercially available compounds were used without further purification unless otherwise indicated. Deuterated
30 solvents such as DMSO or CDCl₃ (99.8% D, Cambridge Isotope Laboratories) were used in all experiments as indicated. ¹H NMR spectra were recorded on a Bruker Avance 400 MHz NMR spectrometer. Significant peaks are tabulated and typically include: number of protons, multiplicity (s, singlet; d, double; t, triplet; q, quartet; m, multiplet; br s, broad singlet) and coupling constant(s) in Hertz. Chemical shifts are reported as parts per million (δ) relative to tetramethylsilane. Mass spectra were

recorded on a Perkin-Elmer SCIEX HPLC/MS instrument using reverse-phase conditions (acetonitrile/water, 0.05% trifluoroacetic acid) and electrospray (ES) ionization. Abbreviations used in the examples below have their accepted meanings in the chemical literature. For example, CH₂Cl₂ (dichloromethane), C₆H₆ (benzene), TFA (trifluoroacetic acid), EtOAc (Ethyl Acetate), Et₂O (diethyl ether), DMAP (4-dimethylaminopyridine), DMF (N,N-dimethylformamide) and THF (tetrahydrofuran). Flash chromatography was performed using Merck Silica Gel 60 (230-400 mesh).


For purposes of illustration only, most of the formulae in the following Reaction Schemes are directed to specific embodiments of the compounds of invention. However, one of ordinary skill in the art, in view of the teachings of this specification would reasonably be expected to be able to prepare all the compounds of the invention in the First aspect of the invention utilizing the appropriately-substituted starting materials and methods known to one skilled in the art.

In the general descriptions immediately following each Reaction Scheme, the phrase "standard isolation procedures" is meant to include one or more of the following techniques familiar to one schooled in the art of organic chemistry: organic extraction, washing of organic solutions with dilute aqueous acid or base, use of drying agents, filtration, concentration *in vacuo*, followed by purification using distillation, crystallization, or solid-liquid phase chromatography. The phrase "elevated temperature" refers to a temperature above ambient temperature and the phrase "reduced temperature" refers to a temperature below ambient temperature.

The following specific Preparations (for intermediates) and Examples (for compounds, pharmaceutical compositions and methods of use of the invention) are provided as a guide to assist in the practice of the invention, and are not intended as a limitation on the scope of the invention. Without further description, it is believed that one of ordinary skill in the art can, using the preceding description and the following illustrative examples, make and utilize the compounds of the present invention and practice the claimed methods. It should be understood that the foregoing discussion and examples merely present a detailed description of certain preferred embodiments. It will be apparent to one of ordinary skill in the art that various modifications and equivalents can be made without departing from the spirit and scope of the invention.

Unless otherwise indicated, all compounds associated with NMR and/or mass spectra data were prepared and the NMR and mass spectra measured.

Scheme 1



Reaction and conditions: (a) NaHMDS, THF; 0°C- rt; (b) 3-bromo-1,1,1-trifluoroacetone, NaHCO₃, 2-
 5 PrOH, 80°C; (c) TSA, toluene, reflux; (d) ArB(OH)₂, K₂CO₃, PdCl₂(dppf), DME/H₂O, 80°C.


In general, compounds of formula (0016a) are prepared by first reacting an aniline of formula (0012a) with aryl nitriles (0011a), such as 5-bromo-thiophene-2-carbonitrile, in the presence of base to give compounds of formula (0013a) after standard isolation procedures. In a subsequent step, exposure of amidine (0013a) to haloketone, such as 1-bromo-3,3,3-trifluoroacetone, under basic condition at
 10 elevated temperature provides 1H-imidazol-4-ol of formula (0014a) after standard isolation procedures. Conversion of compounds of formula (0014a) to compounds of formula (0015a) is then accomplished by treating the compounds of formula (0014a) with a catalyst acid such as *p*-toluenesulfonic acid at reflux. In a palladium-mediated coupling reaction, for example, a Suzuki reaction, compounds of formula (0015a) are then reacted with a boronate or boronic acid reagent to give compounds of formula
 15 (0016a) after standard isolation procedures.

Example 1

1-(2,5-Dichloro-phenyl)-2-[5-(3-methanesulfonyl-phenyl)-thiophen-2-yl]-4-trifluoromethyl-1H-imidazole

Example 1a


Preparation of 5-Bromo-N-(2,5-dichloro-phenyl)-thiophene-2-carboxamidine



Under N₂ atmosphere, to a solution of 2,5-dichloroaniline (341 mg, 2.1 mmol) in 2 mL dry THF was added sodium bis(trimethylsilyl)amide (2.1 mL, 1M solution in THF, 2.1 mmol). After the mixture was stirred at ambient temperature for 40 min, a solution of 5-bromo-thiophene-2-carbonitrile (376 mg, 2 mmol) in 2 mL dry THF was added dropwise. The reaction mixture was stirred overnight, and then poured into 100 mL ice water. The orange precipitate was collected by filtration, washed with a solution of ether and hexane (3/7, v/v), and air dried to give a light orange solid (653 mg, 94% yield). ¹H-NMR (400MHz, MeOH-*d*₄): δ 6.94 (d, 1H), 6.99 (dd, 1H), 7.06 (d, 1H), 7.32 (d, 1H), 7.37 (d, 1H).

Example 1b


Preparation of 2-(5-Bromo-thiophen-2-yl)-1-(2,5-dichloro-phenyl)-4-trifluoromethyl-4,5-dihydro-1H-imidazol-4-ol



To a suspension of 5-bromo-N-(2,5-dichloro-phenyl)-thiophene-2-carboxamidine (634 mg, 1.8 mmol) and sodium bicarbonate (227 mg, 2.7 mmol) in 8 mL 2-propanol was added 3-bromo-1,1,1-trifluoroacetone (520 mg, 2.7 mmol). The reaction mixture was heated at 50°C for 1.5 hrs, then at 80°C for 4 hrs. After cooling to room temperature, the residue was filtered out and washed with dichloromethane. The filtrate and washings were combined together and concentrated under vacuum to give a light pink solid that was used directly for the next step (820 mg, 99% yield). ¹H-NMR (400MHz, DMSO-*d*₆): δ 3.09 (d, 1H), 3.95 (dd, 1H), 6.31 (d, 1H), 6.91 (d, 1H), 7.15 (m, 1H), 7.33 (dd, 1H), 7.44 (m, 2H).

Example 1c

Preparation of 2-(5-Bromo-thiophen-2-yl)-1-(2,5-dichloro-phenyl)-4-trifluoromethyl-1H-imidazole




The mixture of 2-(5-bromo-thiophen-2-yl)-1-(2,5-dichloro-phenyl)-4-trifluoromethyl-4,5-dihydro-1H-imidazol-4-ol (822 mg, 1.8 mmol) and *p*-toluenesulfonic acid monohydrate (172 mg, 0.9 mmol) in 10 mL toluene was heated at 120°C for 6 hrs. The solvent was removed, and the residue was redissolved into dichloromethane, then washed with saturated NaHCO₃ and brine, dried over Na₂SO₄, and concentrated *in vacuo*. The crude product was purified by chromatography on silica gel

(hexane/EtOAc 85/15) to give an off-white solid (512 mg, 64% yield). $^1\text{H-NMR}$ (400MHz, CDCl_3): δ 6.68 (d, 1H), 6.89 (d, 1H), 7.31 (dd, 1H), 7.47 (dd, 1H), 7.55 (d, 1H), 7.56 (s, 1H).

Example 1d

Preparation of 1-(2,5-Dichloro-phenyl)-2-[5-(3-methanesulfonyl-phenyl)-thiophen-2-yl]-4-trifluoromethyl-1H-imidazole




2-(5-Bromo-thiophen-2-yl)-1-(2,5-dichloro-phenyl)-4-trifluoromethyl-1H-imidazole (90 mg, 0.2 mmol), (3-methylsulfonyl)phenylboronic acid (60 mg, 0.3 mmol), potassium carbonate (110 mg, 0.8 mmol), and $\text{PdCl}_2(\text{dppf}) \cdot \text{CH}_2\text{Cl}_2$ (16 mg, 0.02 mmol) were mixed with 2 mL 9:1 DME/ H_2O (v/v), then heated at 80°C overnight. All solvent was removed *in vacuo*. The crude product was purified by chromatography on silica gel (Hexane/EtOAc, 6/4) to give a white solid (76 mg, 74% yield). $^1\text{H-NMR}$ (400MHz, CDCl_3): δ 3.08 (s, 3H), 6.79 (d, 1H), 7.22 (d, 1H), 7.36 (d, 1H), 7.52 (d, 1H), 7.58 (m, 3H), 7.78 (m, 1H), 7.86 (m, 1H), 8.10 (t, 1H).

All the following compounds were prepared in similar manner as described in Scheme 1 and mass and NMR spectra measured. The boronic acid or boronate reagents for Suzuki coupling, if not commercially available, were made using standard techniques that are readily apparent to one skilled in the art.

- 1-(2,5-dichlorophenyl)-2-{5-[3-(methylsulfonyl)phenyl]-2-thienyl}-4-(trifluoromethyl)-1H-imidazole; MS (ES): 517.3 $[\text{M}+\text{H}]^+$;
- 5-{5-[1-(2,5-dichlorophenyl)-4-(trifluoromethyl)-1H-imidazol-2-yl]-2-thienyl}-3-methyl-2-(methylthio)pyridine; MS (ES): 500.4 $[\text{M}+\text{H}]^+$;
- 5-{5-[1-(2,5-dichlorophenyl)-4-(trifluoromethyl)-1H-imidazol-2-yl]-2-thienyl}-2-(ethylthio)-3-methylpyridine; MS (ES): 514.2 $[\text{M}+\text{H}]^+$;
- 4-(5-{5-[1-(2,5-dichlorophenyl)-4-(trifluoromethyl)-1H-imidazol-2-yl]-2-thienyl}pyridin-2-yl)morpholine; MS (ES): 525.4, 527.3 $[\text{M}+\text{H}]^+$;
- 1,1-dimethylethyl 4-(5-{5-[1-(2,5-dichlorophenyl)-4-(trifluoromethyl)-1H-imidazol-2-yl]-2-thienyl}pyridin-2-yl)piperazine-1-carboxylate; MS (ES): 624.5, 626.3 $[\text{M}+\text{H}]^+$;
- 1-(2-chlorophenyl)-2-{5-[3-(methylsulfonyl)phenyl]-2-thienyl}-4-(trifluoromethyl)-1H-imidazole; MS (ES): 483.2 $[\text{M}+\text{H}]^+$;

- 5-{5-[1-(2-chlorophenyl)-4-(trifluoromethyl)-1H-imidazol-2-yl]-2-thienyl}-3-methyl-2-(methylthio)pyridine; MS (ES): 466.2 [M+H]⁺;
- 5-{5-[1-(2-chlorophenyl)-4-(trifluoromethyl)-1H-imidazol-2-yl]-2-thienyl}-2-(ethylthio)-3-methylpyridine; MS (ES): 480.2 [M+H]⁺;
- 5 • methyl (4-{5-[1-(2-chlorophenyl)-4-(trifluoromethyl)-1H-imidazol-2-yl]-2-thienyl}-3-methylphenyl)acetate; MS (ES): 491.2 [M+H]⁺;
- 3-{5-[1-(2-chlorophenyl)-4-(trifluoromethyl)-1H-imidazol-2-yl]-2-thienyl}benzenesulfonamide; MS (ES): 484.0 [M+H]⁺;
- 10 • 4-{5-[1-(2-chlorophenyl)-4-(trifluoromethyl)-1H-imidazol-2-yl]-2-thienyl}benzenesulfonamide; MS (ES): 484.1 [M+H]⁺;


Scheme 2



The starting materials were prepared in similar manner as Scheme 1, followed by further transformations to make the final products as described in Scheme 2.

Example 2

Preparation of 5-{5-[1-(2,5-dichlorophenyl)-4-(trifluoromethyl)-1H-imidazol-2-yl]-2-thienyl}-2-(methylsulfonyl)-3-methylpyridine




5-{5-[1-(2,5-Dichloro-phenyl)-4-trifluoromethyl-1H-imidazol-2-yl]-thiophen-2-yl}-3-methyl-2-methylsulfonyl-pyridine (80 mg, 0.16 mmol) was dissolved in 8 mL mixture of dichloromethane and methanol (5:1, V/V). MMPP (magnesium monoperoxyphthalate hexahydrate, 200 mg, 0.35 mmol, 80% tech.) was added then. The mixture was stirred at room temperature for 2 hrs, then diluted with dichloromethane, and filtered. The filtrate was washed with saturated NaHCO₃, and brine, dried over Na₂SO₄, and concentrated *in vacuo*. The crude product was purified by column chromatography on silica gel (15 → 60% EtOAc/Hexane) to give a white solid (52 mg, 61% yield). ¹H-NMR (400MHz,

DMSO-*d*₆): δ 2.61 (s, 3H), 3.37 (s, 3H), 6.58 (d, *J* = 4.0, 1H), 7.63 (d, *J* = 4.0, 1H), 7.86 – 7.79 (m, 2H), 8.20 – 8.17 (m, 2H), 8.31 – 8.30 (m, 1H), 8.80 (m, 1H). MS (ES): 532.2, 536.2 [M+H]⁺

The following compounds were made in similar manner by oxidation of appropriate sulfides.

- 5 • 5-{5-[1-(2,5-dichlorophenyl)-4-(trifluoromethyl)-1H-imidazol-2-yl]-2-thienyl}-2-(ethylsulfonyl)-3-methylpyridine; MS (ES): 546.2 [M+H]⁺
- 5-{5-[1-(2,5-dichlorophenyl)-4-(trifluoromethyl)-1H-imidazol-2-yl]-2-thienyl}-3-methyl-2-(methylsulfonyl)pyridine; MS (ES): 532.2 [M+H]⁺
- 5-{5-[1-(2-chlorophenyl)-4-(trifluoromethyl)-1H-imidazol-2-yl]-2-thienyl}-3-methyl-2-(methylsulfonyl)pyridine; MS (ES): 498.4 [M+H]⁺
- 10 • 5-{5-[1-(2-chlorophenyl)-4-(trifluoromethyl)-1H-imidazol-2-yl]-2-thienyl}-2-(ethylsulfonyl)-3-methylpyridine; MS (ES): 512.2 [M+H]⁺


Scheme 3



15 The starting materials were prepared in similar manner as Scheme 1, followed by further transformations to make the final products as described in Scheme 3.

Example 3


1-(5-{5-[1-(2,5-Dichloro-phenyl)-4-trifluoromethyl-1H-imidazol-2-yl]-thiophen-2-yl}-pyridin-2-yl)-piperazine



20 4-(5-{5-[1-(2,5-Dichloro-phenyl)-4-trifluoromethyl-1H-imidazol-2-yl]-thiophen-2-yl}-pyridin-2-yl)-piperazine-1-carboxylic acid *tert*-butyl ester (104 mg, 0.17 mmol) was mixed with 4 mL 50% trifluoromethylacetic acid in dichloromethane, and stirred at room temperature for 2 hrs. All solvent was removed; the residue was redissolved in dichloromethane and neutralized to pH 7 by saturated NaHCO₃. The organic layer was washed with brine, dried over Na₂SO₄, and concentrated *in vacuo*.
 25 The residue was precipitated from dichloromethane / Hexane mixture, then filtered and washed several

times with dichloromethane to give yellow solid (50 mg, 57% yield). ¹H-NMR (400MHz, CDCl₃): δ 3.30 – 2.97 (m, 4H), 3.57 – 3.54 (m, 4H), 6.63 (d, *J* = 8.7, 1H), 6.78 (d, *J* = 4.0, 1H), 6.96 (d, *J* = 4.0, 1H), 7.31(s, 1H), 7.60 – 7.50 (m, 4H), 8.37 (m, 1H). MS (ES): 524.3, 526.5, [M+H]⁺


Scheme 4




The starting materials were prepared in similar manner as Scheme 1, followed by further transformations to make the final products as described in Scheme 4.

Example 4

(4-{5-[1-(2-Chloro-phenyl)-4-trifluoromethyl-1H-imidazol-2-yl]-thiophen-2-yl}-3-methyl-phenyl)-acetic acid



(4-{5-[1-(2-Chloro-phenyl)-4-trifluoromethyl-1H-imidazol-2-yl]-thiophen-2-yl}-3-methyl-phenyl)-acetic acid methyl ester (123 mg, 0.25 mmol) was dissolved in 6 mL mixture of THF and water (3:1, V/V). Lithium hydroxide monohydrate (2.3 mg, 0.55 mmol) was added then. The mixture was stirred at room temperature for 3 hrs. The mixture was neutralized to pH 7 by 1N HCl, and then extracted with ethyl acetate. The combined organic layers were washed with brine, dried over Na₂SO₄, and concentrated *in vacuo*. The crude product was purified by column chromatography on silica gel (0 → 5% MeOH/DCM) to give a white solid (47 mg, 39% yield). ¹H-NMR (400MHz, CDCl₃): δ 2.35 (s, 3H), 3.64 (s, 2H), 6.83 (s, 2H), 7.13 – 7.10 (m, 1H), 7.16 (m, 1H), 7.29 – 7.27 (m, 1H), 7.35 (m, 1H), 7.47 - 7.46 (m, 2H), 7.57 - 7.52 (m, 1H), 7.64 – 7.62 (m, 1H). MS (ES): 477.1, [M+H]⁺



(a) Me_3Al , PhMe , 90°C ; (b) ethyl bromopyruvate, NaHCO_3 , 2-PrOH , 95°C ; (c) ArB(OH)_2 , $\text{Cl}_2\text{Pd(dppf)}$, K_2CO_3 , $\text{H}_2\text{O/DME}$, 80°C ; (d) R^4MgBr , THF , 0°C-rt .


In general, compounds of formula (0056) can be prepared as depicted in Scheme 5. An aryl nitrile (0051) and an arylamine (0052) can react in the presence of a Lewis acid, such as trimethylaluminum, to give the corresponding amidine (0053). Heteroaryl nitriles 0051 and/or heteroaryl amines 0052 also can be utilized for the syntheses of their respective amidines 0053. Intermediate 0053 then can react with ethyl bromopyruvate in the presence of a weak base, such as sodium bicarbonate, followed by dehydration at elevated temperature to yield the corresponding imidazole (0054). This intermediate 0054 can undergo cross-coupling reactions, such as with an arylboronic acid under typical Suzuki conditions, to afford the corresponding 2-biaryl-imidazole (0055). 0055 can react with a Grignard reagent, such as an alkylmagnesium bromide, to afford the desired product 0056.

Example 5

2-[1-(2,6-dichlorophenyl)-2-[5-(3-methanesulfonyl-phenyl)-thiophen-2-yl]-1H-imidazol-4-yl]-propan-2-ol

Example 5a


Preparation of 5-bromo-N-(2,6-dichlorophenyl)-thiophene-2-carboxamidine



To a stirred solution of 2,6-dichloroaniline (1.75 g, 10.8 mmol) in toluene (25 mL, anhyd) was added dropwise a 2.0M solution of trimethylaluminum in toluene (8.0 mL, 16 mmol). After 90 min the reaction mixture was charged with a solution of 5-bromothiophene-2-carbonitrile (3.00 g, 16 mmol) in toluene (20 mL, anhyd) and then heated at 90°C. After 5 h the reaction mixture was allowed to cool to ambient temperature and quenched by addition to a stirred slurry of silica (25 g) in 3:1 CHCl₃/MeOH (100 mL). After 30 min the resulting mixture was filtered and the solids rinsed with MeOH and DCM. The combined filtrates were concentrated to afford the title compound (3.9 g, quant) as a pale yellow solid, which was used in the next step without purification. GC-MS(EI): 348, 350, 352.

Example 5b


10 Preparation of 2-(5-bromothiophen-2-yl)-1-(2,6-dichlorophenyl)-1H-imidazole-4-carboxylic acid ethyl ester



A stirred mixture of 5-bromo-N-(2,6-dichlorophenyl)-thiophene-2-carboxamidine (3.9 g, 11 mmol) and sodium bicarbonate (1.85 g, 22 mmol) in isopropanol (55 mL) was charged with ethyl bromopyruvate (2.5 mL, 20 mmol) and then heated at 95°C. After 17 h the reaction mixture was allowed to cool to ambient temperature, filtered and the solids rinsed with EtOAc. The combined filtrates were concentrated under reduced pressure and purified by chromatography (silica, EtOAc/Hex, 15:85 to 65:35) to give the title compound (3.3 g, 67%) as a tacky yellow solid, which was used in the next step without further purification. ¹H-NMR (DCM-*d*₂): δ 7.60 (s, 1H), 7.49-7.59 (m, 3H), 6.89 (d, 1H), 6.56 (d, 1H), 4.37 (q, 2H), 1.38 (t, 3H).

Example 5c

Preparation of 1-(2,6-dichlorophenyl)-2-[5-(3-methanesulfonyl-phenyl)-thiophen-2-yl]-1H-imidazole-4-carboxylic acid ethyl ester




A mixture of 2-(5-bromothiophen-2-yl)-1-(2,6-dichlorophenyl)-1H-imidazole-4-carboxylic acid ethyl ester (335 mg, 0.75 mmol), 3-methanesulfonyl-phenylboronic acid (180 mg, 0.90 mmol), K₂CO₃ (0.32 g, 2.3 mmol), Cl₂Pd(dppf)·DCM (30 mg, 5 mol%) and H₂O (0.4 mL) in DME (4 mL) was sparged with Argon for 5 min and then heated at 80°C as a sealed flask. After 2 h the reaction mixture

was allowed to cool to ambient temperature, filtered (Celite™) and the filter agent rinsed with EtOAc. The combined filtrates were concentrated under reduced pressure and purified by chromatography (silica, EtOAc/Hex, 30:70 to 80:20) to give the title compound (313 mg, 80%) as a pale yellow solid, which was used in the next step without further purification. ¹H-NMR (DCM-*d*₂): δ 8.08 (m, 1H), 7.82 (m, 2H), 7.64 (s, 1H), 7.51-7.62 (m, 4H), 7.24 (d, 1H), 6.72 (d, 1H), 4.38 (q, 2H), 3.05 (s, 3H), 1.40 (t, 3H).

Example 5d

Preparation of 2-{1-(2,6-dichlorophenyl)-2-[5-(3-methanesulfonyl-phenyl)-thiophen-2-yl]-1H-imidazol-4-yl}-propan-2-ol



To a stirred solution of 1-(2,6-dichlorophenyl)-2-[5-(3-methanesulfonyl-phenyl)-thiophen-2-yl]-1H-imidazole-4-carboxylic acid ethyl ester (300 mg, 0.58 mmol) in THF (3 mL) at 0°C was added a 3.0M solution of methylmagnesium bromide in Et₂O (0.80 mL, 2.4 mmol). After addition was completed the flask was removed from the ice-water bath and allowed to warm to ambient temperature. At 40 min the reaction mixture was quenched by addition of satd NH₄Cl, extracted with EtOAc, dried (Na₂SO₄), concentrated and purified by chromatography (silica, EtOAc/Hex, 40:60 to 90:10) to give the title compound (0.17 g, 59%) as a white solid. ¹H-NMR (DCM-*d*₂): δ 8.07 (m, 1H), 7.81 (m, 2H), 7.46-7.60 (m, 4H), 7.21 (d, 1H), 6.85 (s, 1H), 6.62 (d, 1H), 3.05 (s, 3H), 2.68 (br s, 1H), 1.61 (s, 6H); MS(ES): 507, 509 [M+H]⁺.

In a similar manner, the following imidazoles were prepared from appropriate reagents. Optionally the order of Suzuki cross-coupling and addition of Grignard reagent was switched.

- 2-(1-(2,6-dichlorophenyl)-2-(5-(3-(ethylsulfonyl)phenyl)thiophen-2-yl)-1H-imidazol-4-yl)propan-2-ol; MS(ES): 521, 523 [M+H]⁺
- 2-(1-(2,6-dichlorophenyl)-2-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; MS(ES): 515, 517 [M+H]⁺
- 2-(1-(2,6-dichlorophenyl)-2-(3'-(ethylsulfonyl)-3-methylbiphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; MS(ES): 529, 531 [M+H]⁺
- 2-(1-(2-isopropyl-6-methylphenyl)-2-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; MS(ES): 503 [M+H]⁺
- 2-(2-(3'-(ethylsulfonyl)-3-methylbiphenyl-4-yl)-1-(2-isopropyl-6-methylphenyl)-1H-imidazol-4-yl)propan-2-ol; MS(ES): 517 [M+H]⁺


- 2-(1-(2-isopropylphenyl)-2-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; MS(ES): 489 [M+H]⁺
- 2-(2-(3'-(ethylsulfonyl)-3-methylbiphenyl-4-yl)-1-(2-isopropylphenyl)-1H-imidazol-4-yl)propan-2-ol; MS(ES): 503 [M+H]⁺
- 5 • 2-(1-(2,6-dichlorophenyl)-2-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; MS(ES): 501, 503 [M+H]⁺
- 2-(1-(2,6-dichlorophenyl)-2-(3'-(ethylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; MS(ES): 515, 517 [M+H]⁺
- 2-(2-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-1-(2,6-dichlorophenyl)-1H-imidazol-4-yl)propan-2-ol; MS(ES): 535, 537 [M+H]⁺
- 10 • 2-(2-(3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl)-1-(2,6-dichlorophenyl)-1H-imidazol-4-yl)propan-2-ol; MS(ES): 549, 551 [M+H]⁺

Example 6

15 2-[5-chloro-2-(3-chloro-3'-methanesulfonyl-biphenyl-4-yl)-1-(2,6-dichlorophenyl)-1H-imidazol-4-yl]-propan-2-ol

Example 6a


Preparation of 2-[2-(4-bromo-2-chlorophenyl)-1-(2,6-dichlorophenyl)-1H-imidazol-4-yl]-propan-2-ol



20 To a 3.0 M solution of MeMgBr in Et₂O (7.0 mL, 21.0 mmol) at 0°C added slowly a solution of 2-(4-bromo-2-chlorophenyl)-1-(2,6-dichlorophenyl)-1H-imidazole-4-carboxylic acid ethyl ester (2.80 g, 5.90 mmol) in THF (30 mL, anhyd). After addition was completed the flask was removed from the ice-water bath and allowed to warm to ambient temperature. At 50 min the reaction mixture was quenched by addition of satd NH₄Cl, extracted with EtOAc, dried (Na₂SO₄), concentrated and purified by chromatography (silica, EtOAc/Hex, 15:85 to 55:45) to give the title compound (2.1 g, 77%) as a white solid. ¹H-NMR (DCM-*d*₂): δ 7.58 (d, 1H), 7.37-7.41 (m, 2H), 7.26-7.32 (m, 2H), 7.11 (d, 1H), 25 6.93 (s, 1H), 2.73 (br s, 1H), 1.61 (s, 6H).

Example 6b


Preparation of 2-[2-(4-bromo-2-chlorophenyl)-5-chloro-1-(2,6-dichlorophenyl)-1H-imidazol-4-yl]-propan-2-ol



A suspension of 2-[2-(4-bromo-2-chlorophenyl)-1-(2,6-dichlorophenyl)-1H-imidazol-4-yl]-propan-2-ol (230 mg, 0.50 mmol) in MeCN (4 mL, anhyd) was charged with N-chlorosuccinimide (70 mg, 0.52 mg) and then heated at 70°C. After 22 h the reaction mixture was charged with additional N-chlorosuccinimide (67 mg, 0.50 mmol) and heated at 85°C. After 40 h (total reaction time) the reaction mixture was allowed to cool to ambient temperature, concentrated and purified by chromatography (silica, EtOAc/Hex, 0:100 to 30:70) to give the title compound (0.19 g, 77%) as a white solid. ¹H-NMR (DCM-*d*₂): δ 7.61 (d, 1H), 7.42-7.45 (m, 2H), 7.36 (m, 1H), 7.27 (dd, 1H), 7.10 (d, 1H), 3.33 (s, 1H), 1.65 (s, 6H).


Example 6c

Preparation of 2-[5-chloro-2-(3-chloro-3'-methanesulfonyl-biphenyl-4-yl)-1-(2,6-dichlorophenyl)-1H-imidazol-4-yl]-propan-2-ol



A mixture of 2-[2-(4-bromo-2-chlorophenyl)-5-chloro-1-(2,6-dichlorophenyl)-1H-imidazol-4-yl]-propan-2-ol (184 mg, 0.37 mmol), 3-methanesulfonyl-phenylboronic acid (90 mg, 0.45 mmol), K₂CO₃ (0.15 g, 1.1 mmol), Cl₂Pd(dppf)·DCM (15 mg, 5 mol%) and H₂O (0.2 mL) in DME (2 mL) was sparged with Argon for 5 min and then heated at 60°C as a sealed flask. After 70 min the reaction mixture was allowed to cool to ambient temperature, filtered (Celite™) and the filter agent rinsed with EtOAc. The combined filtrates were concentrated under reduced pressure and purified by chromatography (silica, EtOAc/Hex, 20:80 to 60:40) to give the title compound (165 mg, 78%) as a white solid. ¹H-NMR (DCM-*d*₂): δ 8.08 (m, 1H), 7.93 (m, 1H), 7.84 (m, 1H), 7.73 (d, 1H), 7.67 (m, 1H), 7.33-7.47 (m, 5H), 3.42 (br s, 1H), 3.06 (s, 3H), 1.68 (s, 6H); MS(ES): 591, 593, 595 [M+Na]⁺.

Scheme 6



(a) Et_3N , THF; (b) SOCl_2 , 90°C ; concentrate and add to NH_3/MeOH ; (c) ethyl bromopyruvate, NaHCO_3 , EtOH, μW 175°C ; (d) R^4MgBr , THF, 0°C -rt; (e) $\text{ArB}(\text{OH})_2$, $\text{Cl}_2\text{Pd}(\text{dppf})$, K_2CO_3 , $\text{H}_2\text{O}/\text{DME}$, μW 120°C .


In general, compounds of formula **00616** can be prepared as shown in Scheme 6. First, an amide can be generated from reacting a benzoyl chloride (**00611**) and a benzylamine (**00612**) under basic conditions. The resulting amide can be treated with thionyl chloride at elevated temperature to give an imidoyl chloride, which can be converted directly to its corresponding amidine (**00613**) upon addition to a solution of ammonia. Amidine **00613** and ethyl bromopyruvate can react under basic conditions and at elevated temperature to yield the corresponding imidazole (**00614**). This intermediate ester **00614** can react with a Grignard reagent, such as an alkylmagnesium bromide, to afford the respective carbinol (**00615**). Last intermediate **00615** can undergo cross-couplings reactions, such as a Suzuki-Miyaura reaction with boronic acids (esters), to yield the target imidazole **00616**.

Example 7

2-[1-(2-chlorobenzyl)-2-(3'-methanesulfonyl-biphenyl-4-yl)-1H-imidazol-4-yl]-propan-2-ol

Example 7a


Preparation of 4-bromo-N-(2-chlorobenzyl)-benzamide



To a stirred solution of 2-chlorobenzylamine (2.42 mL, 20 mmol) in THF (80 mL, anhyd) at 0°C added a solution of 4-bromobenzoyl chloride (4.39 g, 20 mmol) in THF (20 mL, anhyd) and then triethylamine (2.93 mL, 21 mmol). After 12 h the resulting mixture was filtered and the solids rinsed with EtOAc. The combined filtrates were diluted with EtOAc (100 mL), washed with 1N HCl, satd NaHCO₃ and brine, then dried (anhyd Na₂SO₄) and concentrated to afford the title compound (6.36 g, 98%) as a white solid, which was used in the next step without purification. GC-MS(EI): 322, 324.

Example 7b


Preparation of 4-bromo-N-(2-chlorobenzyl)-benzamide



A mixture of 4-bromo-N-(2-chlorobenzyl)-benzamide (1.62 g, 5.0 mmol) and thionyl chloride (0.73 mL, 10 mmol) was heated at 90°C. After 150 min the reaction mixture was cooled and concentrated under reduced pressure. The resulting residue was diluted with toluene (10 mL, anhyd) and concentrated to give the corresponding imidoyl chloride [conversion confirmed by NMR] as a pale yellow solid. To a stirred 2M solution of NH₃ in MeOH (8 mL, 16 mmol) was added this intermediate in small portions. After 12 h the reaction mixture was added to satd NaHCO₃ and extracted with EtOAc (2 x 50 mL). The combined extracts were washed with brine, dried (Na₂SO₄) and concentrated under reduced pressure to afford the title compound (1.5 g, 93%), which was used in the next step without purification. GC-MS(EI): 321, 323.

Example 7c


Preparation of 2-(4-bromophenyl)-1-(2-chlorobenzyl)-1H-imidazole-4-carboxylic acid ethyl ester



A mixture of 4-bromo-N-(2-chlorobenzyl)-benzimidine (1.4 g, 4.3 mmol), sodium bicarbonate (0.72 g, 8.6 mmol) and ethyl bromopyruvate (0.98 mL, 7.8 mmol) in EtOH (14 mL) was heated in a microwave unit (Biotage Initiator™) at 175°C for 12 min. The resulting mixture was decanted and the solids rinsed with EtOAc. The combined filtrates were concentrated and purified by chromatography (silica, EtOAc/Hex, 0:100 to 40:60) to give the title compound (0.24 g, 13%) as a pale brown residue. ¹H-NMR (DCM-*d*₂): δ 7.58-7.62 (m, 3H), 7.43-7.49 (m, 3H), 7.34 (m, 1H), 7.28 (m, 1H), 6.96 (d, 1H), 5.30 (s, 2H), 4.32 (q, 2H), 1.35 (t, 3H); MS(ES): 419, 421 [M+H]⁺.

Example 7d


Preparation of 2-[2-(4-bromophenyl)-1-(2-chlorobenzyl)-1H-imidazol-4-yl]-propan-2-ol



To a 3.0 M solution of MeMgBr in Et₂O (1.0 mL, 3.0 mmol) at 0°C added slowly a solution of
5 2-(4-bromophenyl)-1-(2-chlorobenzyl)-1H-imidazole-4-carboxylic acid ethyl ester (235 mg, 0.56 mmol) in THF (3 mL, anhyd). After addition was completed the flask was removed from the ice-water bath and allowed to warm to ambient temperature. At 40 min the reaction mixture was quenched by addition of satd NH₄Cl, extracted with EtOAc, dried (MgSO₄), concentrated and purified by chromatography (silica, EtOAc/Hex, 20:80 to 60:40) to give the title compound (135 mg, 59%) as a
10 pale amber solid. ¹H-NMR (DCM-*d*₂): δ 7.54 (d, 2H), 7.38-7.45 (m, 3H), 7.24-7.33 (m, 2H), 6.89 (d, 1H), 6.82 (s, 1H), 5.25 (s, 2H), 2.86 (br s, 1H), 1.54 (s, 6H); MS(ES): 405, 407 [M+H]⁺.

Example 7e


Preparation of 2-[1-(2-chlorobenzyl)-2-(3'-methanesulfonyl-biphenyl-4-yl)-1H-imidazol-4-yl]-propan-2-ol



A mixture of 2-[2-(4-bromophenyl)-1-(2-chlorobenzyl)-1H-imidazol-4-yl]-propan-2-ol (130 mg, 0.32 mmol), 3-methanesulfonyl-phenylboronic acid (80 mg, 0.40 mmol), Cl₂Pd(dppf)·DCM (13 mg, 5 mol%) and a 2.8M aqueous solution of K₂CO₃ (0.35 mL, 0.98 mmol) in DME (2 mL) was heated in a microwave unit (Biotage Initiator™) at 120°C for 4 min. The reaction mixture was concentrated and purified by chromatography (silica, EtOAc/Hex, 45:55 to 90:10) to give the title compound (118 mg, 77%) as a white solid. ¹H-NMR (DCM-*d*₂): δ 8.08 (m, 1H), 7.84 (dd, 2H), 7.55-7.64 (m, 5H), 7.36 (m, 1H), 7.17-7.26 (m, 2H), 6.85 (d, 1H), 6.77 (s, 1H), 5.24 (s, 2H), 3.00 (s, 3H), 2.86 (br s, 1H), 1.49 (s, 6H); MS(ES): 481 [M+H]⁺.

In a similar manner 2-[1-(2,3-dichlorobenzyl)-2-(3'-methanesulfonyl-biphenyl-4-yl)-1H-imidazol-4-yl]-propan-2-ol was prepared by replacing 2-chlorobenzylamine with 2,3-dichlorobenzylamine. MS(ES): 515, 517 [M+H]⁺.

Scheme 7



Reaction and conditions: (a) NaOAc, NH₄OH, MeOH; (b) ArB(OH)₂, [Cu(OH).TMEDA]₂Cl₂, DCM; (c) Ar'B(OH)₂, K₂CO₃, PdCl₂(dppf), DME/H₂O, 80°C.


In general, compounds of formula (0075b) are prepared by first reacting 3,3-dihaloketone such as 3,3-dibromo-1,1,1-trifluoroacetone with aqueous sodium acetate to generate a glyoxal which react *in situ* with aldehyde of formula (0071b) and ammonia to yield an imidazole of formula (0073b). Conversion of compounds of formula (0073b) to compounds of formula (0074b) is then accomplished by means of N-arylation such as a Buchwald reaction or Cu(II) catalyzed coupling with boronic acid regents. In a palladium mediated coupling reaction, for example a Suzuki reaction, compounds of formula (0074b) are then reacted with a boronate or boronic acid to give compounds of formula (0075b) after standard isolation procedures.

Example 8

2-(2-Chloro-phenyl)-1-(3'-methanesulfonyl-biphenyl-4-yl)-4-trifluoromethyl-1H-imidazole

Example 8a

Preparation of 2-(2-Chloro-phenyl)-4-trifluoromethyl-1H-imidazole




3,3-Dibromo-1,1,1-trifluoroacetone (6.75 g, 25 mmol) was added to a solution of NaOAc (4.11 g, 50 mmol) in 15 mL water, and the mixture was heated at 100°C for 60 min. After cooling to room temperature, 2-chlorobenzaldehyde (2.8 g, 20 mmol) dissolved in MeOH (45 mL) was added, followed by conc. NH₄OH (10 mL). The mixture was stirred at room temperature overnight. After methanol was removed by evaporation, water was added. The precipitate was collected by filtration and washed with

water, and dried under vacuum to give a light yellow solid (4.7 g, 76%). $^1\text{H-NMR}$ (400MHz, CDCl_3): δ 7.37 (dd, 1H), 7.40 (dd, 1H), 7.46 (dd, 1H), 7.50 (m, 1H), 8.33 (dd, 1H).

Example 8b


Preparation of 1-(4-Bromo-phenyl)-2-(2-chloro-phenyl)-4-trifluoromethyl-1H-imidazole



2-(2-Chloro-phenyl)-4-trifluoromethyl-1H-imidazole (370 mg, 1.5 mmol), 4-bromo-phenylboronic acid (600 mg, 3 mmol), and $[\text{Cu}(\text{OH})\cdot\text{TMEDA}]_2\text{Cl}_2$ (140 mg, 0.3 mmol) were mixed with 6 mL dry dichloromethane, and stirred at room temperature under atmosphere of dioxygen for 2 days. The reaction mixture was diluted with dichloromethane and filtered through celite. The filtrate was concentrated under vacuum, and the residue was purified by chromatography on silica gel (Hexane/EtOAc 8/2) to give a solid (151 mg, 25% yield). $^1\text{H-NMR}$ (400MHz, CDCl_3): δ 7.02 (d, 2H), 7.34 (m, 3H), 7.47 (d, 2H), 7.53 (dd, 1H), 7.56 (dd, 1H).

Example 8c

Preparation of 2-(2-Chloro-phenyl)-1-(3'-methanesulfonyl-biphenyl-4-yl)-4-trifluoromethyl-1H-imidazole



1-(4-Bromo-phenyl)-2-(2-chloro-phenyl)-4-trifluoromethyl-1H-imidazole (100 mg, 0.25 mmol), (3-methylsulfonyl)phenylboronic acid (100 mg, 0.5 mmol), potassium carbonate (155 mg, 1.13 mmol), and $\text{PdCl}_2(\text{dppf})\cdot\text{CH}_2\text{Cl}_2$ (20 mg, 0.025 mmol) were mixed with 3 mL 9:1 DME/ H_2O (v/v), then heated at 80°C overnight. All solvent was removed *in vacuo*. The residue was purified by chromatography on silica gel (Hexane/EtOAc, 6/4), then by reverse phase HPLC (30% acetonitrile in water) to give a white solid (55 mg, 46% yield). $^1\text{H-NMR}$ (400MHz, CDCl_3): δ 3.10 (s, 3H), 7.26 (m, 2H), 7.37 (m, 3H), 7.62 (m, 5H), 7.84 (m, 1H), 7.94 (m, 1H), 8.12 (t, 1H).

In similar manner, all the following compounds were prepared using appropriate aldehydes and boronic acid reagents.


- 2-(2-chlorophenyl)-1-[3'-(methylsulfonyl)biphenyl-4-yl]-4-(trifluoromethyl)-1H-imidazole;
MS(ES): 477.0 $[\text{M}+\text{H}]^+$;

- 4'-[2-(2-chlorophenyl)-4-(trifluoromethyl)-1H-imidazol-1-yl]biphenyl-3-sulfonamide; MS(ES): 478.0 [M+H]⁺;
- 1-(4-bromophenyl)-4-(trifluoromethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazole; MS(ES): 435.0 [M+H]⁺;
- 5 • 1-[3'-(methylsulfonyl)biphenyl-4-yl]-4-(trifluoromethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazole; MS(ES): 511.3 [M+H]⁺;
- 4'-{4-(trifluoromethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-3-sulfonamide; MS(ES): 512.3 [M+H]⁺;

10 The following compounds were prepared in similar manner as described in Scheme 7, except Cu (II) catalyzed coupling reaction to make compounds of formula (0074b) was replaced by base, like sodium hydride, catalyzed replacement reaction with aryl bromids bearing appropriate leaving groups, such as 2,5-dibromopyridine.

- 5-[3-(methylsulfonyl)phenyl]-2-{4-(trifluoromethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}pyridine; MS(ES): 512.3 [M+H]⁺;
- 15 • 2-[2-(2-chlorophenyl)-4-(trifluoromethyl)-1H-imidazol-1-yl]-5-[3-(methylsulfonyl)phenyl]pyridine; MS(ES): 478.0 [M+H]⁺;

Scheme 8



20 (a) Method A: Me_3Al , Toluene, 0°C- 80°C; Method B: $NaHMDS$, THF; 0°C- rt; (b) Ethyl bromopyruvate, $NaHCO_3$, 2-PrOH, 80°C; (c) $NaHCO_3$ / 2-PrOH or $HOAc$ / 2-PrOH, reflux; (d) R_2MgBr , THF, 0°C- rt; (e) $ArB(OH)_2$, K_2CO_3 , $PdCl_2(dppf)$, DME/ H_2O , 80°C.

In general, compounds of formula (0087) are prepared by first reacting aniline of formula (0082) with aryl nitriles (0081), in the presence of either Lewis Acid (Method A) or base (Method B), to

give compounds of formula (0083) after standard isolation procedures. In a subsequent step, exposure of amidine (0083) to haloester, such as Ethyl α -bromopyruvate, under basic condition at elevated temperature provides intermediate 1*H*-imidazol-4-ol of formula (0084), which can be converted to compounds of formula (0085) *in situ*, or by means of acid catalyzed dehydration. Compounds of formula (0085) are then subjected to functionality transformation, such as from ester to carbinol. In a palladium mediated coupling reaction, for example Suzuki reactions, compounds of formula (0086) are then reacted with a diboronate or boronic acid reagent to give compounds of formula (0087) after standard isolation procedures.


Example 9

2-[2-(2,6-Difluoro-phenyl)-1-(3'-methanesulfonyl-biphenyl-4-yl)-1*H*-imidazol-4-yl]-propan-2-ol

Example 9a

(Method A)

Preparation of *N*-(4-bromophenyl)-2,6-difluorobenzimidamide




N-(4-Bromo-phenyl)-2,6-difluoro-benzimidine: 4-bromoaniline (1.72 g, 10 mmol) was placed into a three-neck reactor, and the reactor was flushed with argon for 20min. 50 mL anhydrous toluene was added. After cooling to 0°C, trimethylaluminum (7.5 mL, 2.0M solution in toluene, 15 mmol) was added slowly to keep the inner temperature below 18 °C. After addition, the reaction mixture was warmed to room temperature and then stirred for additional 2 hrs. A solution of 2, 6-difluorobenzonitrile (2.78 g, 20 mmol) in 50 mL anhydrous toluene was added, and the reaction mixture was heated to 80 °C. After 18 hrs, the reaction mixture was cooled to room temperature and poured over a slurry of silica gel in CHCl₃/MeOH (4/1, V/V). After being stirring for 20 min, the mixture was filtered and the residue was washed with a mixture of CHCl₃/MeOH (4/1, V/V). The combined filtrates were concentrated *in vacuo*, and the resulting solid was stirred with a solution of Hexane/Ether (3/1, V/V). The intermediate was filtered and washed with additional Hexane/Ether (3/1, V/V) to give a white solid (2.4 g, 79% yield), which was used directly for the next step without further purification.

Example 9aa

(Method B)


Preparation of *N*-(4-bromophenyl)-2-chloro-6-methylbenzimidamide



N-(4-Bromo-phenyl)-2-chloro-6-methyl-benzamidine: Under N₂ atmosphere at room temperature to a solution of lithium bis(trimethylsilyl)amide (9.7 mL, 1.0 M solution in THF, 9.7 mmol) was added a solution of 4-bromoaniline (1.67 g, 9.7 mmol) in 10 mL anhydrous THF. After the mixture was stirred at ambient temperature for 40 min, a solution of 2-chloro-6-methylbenzonitrile (1.55 mg, 10.2 mmol) in 10 mL anhydrous THF was added. The reaction mixture was stirred at room temperature overnight, and then poured into 300 mL ice water. The orange precipitate was collected by filtration, washed with hexane and air dried to give a light orange solid (2.19 g, 66% yield), which was used directly for the next step without further purification.

Example 9b


Preparation of ethyl 1-(4-bromophenyl)-2-(2,6-difluorophenyl)-1H-imidazole-4-carboxylate



To a mixture of N-(4-Bromo-phenyl)-2,6-difluorobenzamidine (2.4 g, 7.7 mmol) and sodium bicarbonate (1.3 g, 15.4 mmol) in 40 mL 2-propanol was added ethyl α -bromopyruvate (3.34 g, 15.4 mmol, 90% tech). The mixture was heated at 80 – 85 °C overnight. After cooling to room temperature, the solvent was removed and the residue was redissolved into dichloromethane. The precipitate was filtered out, and the filtrate was concentrated down. The crude product was purified by column chromatography on silica gel (30 → 50% EtOAc/Hexane) to give a yellow solid (1.82 g, 58% yield).

Example 9c


Preparation of 2-(1-(4-bromophenyl)-2-(2,6-difluorophenyl)-1H-imidazol-4-yl)propan-2-ol



Methylmagnesium bromide (11.2 mL, 15.7 mmol, 1.4M in toluene/THF 75:25) was placed into a three-neck flask. Under nitrogen at 0 °C to it was added a solution of 1-(4-Bromo-phenyl)-2-(2,6-difluoro-phenyl)-1H-imidazole-4-carboxylic acid ethyl ester in 15 mL THF. After the addition, the ice bath was removed and the reaction mixture was stirred at room temperature for 2.5 hrs. The reaction was quenched with sat. Ammonium chloride. Two layers were separated, and the aqueous layer was extracted three times with ethyl acetate. The combined organic layers were washed with brine, dried over Na₂SO₄, and concentrated *in vacuo*. The crude product was purified by column chromatography on silica gel (30 → 50% EtOAc/Hexane) to give an off-white solid (0.9 g, 51% yield).

Example 9d

Preparation of 2-(2-(2,6-difluorophenyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol



- 5 2-[1-(4-Bromo-phenyl)-2-(2,6-difluoro-phenyl)-1H-imidazol-4-yl]-propan-2-ol (250 mg, 0.64 mmol), (3-methylsulfonylphenyl)boronic acid (254 mg, 1.27 mmol), and potassium carbonate (398 mg, 2.88 mmol) were placed into a flask. 6 mL of mixture of DME (1,2-dimethoxyethane) and water (9:1, V/V) was added. The flask was flushed with argon for 15 min, and then Pd catalyst (Dichloro[1,1'-bis (diphenylphosphino)ferrocene]palladium (II) dichloromethane adduct, 52 mg, 0.064 mmol) was added.
- 10 The mixture was heated to 80 °C overnight. After cooling to room temperature, all solvent was removed. The crude product was purified by column chromatography on silica gel (45 → 99% EtOAc/Hexane) to give an off-white solid (230 mg, 77% yield). ¹H-NMR (400MHz, CDCl₃): δ 1.70 (s, 6H), 2.71 (s, 1H), 3.09 (s, 3H), 6.92 – 6.88 (m, 2H), 7.21 (s, 1H), 7.39 – 7.28 (m, 3H), 7.59 – 7.57 (m, 2H), 7.66 (t, 1H, *J*=7.8), 7.86 – 7.83 (M, 1H), 7.95 - 7.92 (m, 1H), 8.12 (t, 1H, *J*=1.7). MS (ES): 491.4
- 15 [M+Na]⁺

In similar manner, all the following compounds were made using either Method A (Example 9a) or Method B (Example 9aa) to prepare amidines. The order of last two steps of Grignard addition and Suzuki coupling were switched in some cases. The boronic acid or boronate reagents for Suzuki coupling, if not commercially available, were made using standard techniques that are readily apparent

20 to one skilled in the art.

- ethyl 1-[3'-(methylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazole-4-carboxylate; MS(ES): 515.3 [M+H]⁺;
- 2-{1-[3'-(methylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS(ES): 501.4 [M+H]⁺, 523.3 [M+Na]⁺;

25 • 5-{3-[(1-methylethyl)sulfonyl]phenyl}-2-{4-(trifluoromethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}pyridine; MS(ES): 540.3 [M+H]⁺;

- 1-[3'-[(1-methylethyl)sulfonyl]biphenyl-4-yl]-4-(trifluoromethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazole; MS(ES): 539.3 [M+H]⁺;
- N-(4'-{4-(trifluoromethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-3-

30 yl)methanesulfonamide; MS(ES): 526.5 [M+H]⁺;

- ethyl 1-[3'-(methylsulfonyl)biphenyl-4-yl]-2-phenyl-1H-imidazole-4-carboxylate; MS(ES): 447.3 [M+H]⁺;
- 2-{1-[3'-(methylsulfonyl)biphenyl-4-yl]-2-phenyl-1H-imidazol-4-yl}propan-2-ol;
- 2-{2-(2-chlorophenyl)-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol;
- 5 MS(ES): 467.4 [M+H]⁺;
- 2-{2-(2-fluorophenyl)-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol;
- MS(ES): 451.1 [M+H]⁺;
- 1-{2-(2-fluorophenyl)-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}ethanone; MS(ES): 457.1 [M+Na]⁺;
- 10 • 2-{2-(4-fluorophenyl)-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol;
- MS(ES): 451.3 [M+H]⁺;
- 2-{2-(2-chloro-6-methylphenyl)-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS(ES): 481.1 [M+H]⁺;
- 2-{1-[4'-(methylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS(ES): 501.5 [M+H]⁺;
- 15 • 2-{1-[3'-(ethylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS(ES): 515.5 [M+H]⁺;
- 2-{1-[3'-(methylsulfonyl)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS(ES): 501.4 [M+H]⁺;
- 20 • 2-{1-[4'-(methylsulfonyl)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS(ES): 501.4 [M+H]⁺;
- 2-{2-(2,6-difluorophenyl)-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS(ES): 491.4 [M+Na]⁺;
- 2-(1-{3'-[(1-methylethyl)sulfonyl]biphenyl-4-yl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl)propan-2-ol; MS(ES): 529.3 [M+H]⁺;
- 25 • 2-(1-{3'-[(1-methylethyl)sulfonyl]biphenyl-3-yl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl)propan-2-ol; MS(ES): 529.3 [M+H]⁺;
- 2-{2-(2-chlorophenyl)-1-[3'-(ethylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS(ES): 481.3 [M+H]⁺;
- 30 • 2-{2-(2-chloro-6-fluorophenyl)-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS(ES): 485.3 [M+H]⁺;
- 2-{2-(2,6-difluorophenyl)-1-[3'-(ethylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS(ES): 483.4 [M+H]⁺;

- 2-(1-{5-[3-(methylsulfonyl)phenyl]pyridin-2-yl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl)propan-2-ol; MS(ES): 502.5 [M+H]⁺;
- 2-(1-{6-[3-(methylsulfonyl)phenyl]pyridin-3-yl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl)propan-2-ol; MS(ES): 502.3 [M+H]⁺;
- 5 • 2-{2-(2,6-dichlorophenyl)-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS(ES): 523.3 [M+Na]⁺;
- 2-{2-[2-fluoro-6-(trifluoromethyl)phenyl]-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS(ES): 541.3 [M+Na]⁺;
- 2-{2-(2-methylphenyl)-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS(ES): 447.0 [M+H]⁺;
- 10 • 2-(1-[3'-(methylsulfonyl)biphenyl-4-yl]-2-{2-[(trifluoromethyl)oxy]phenyl}-1H-imidazol-4-yl)propan-2-ol; MS(ES): 517.3 [M+H]⁺, 539.3 [M+Na]⁺;
- 2-{2-[2-(dimethylamino)-6-fluorophenyl]-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS(ES): 516.3 [M+Na]⁺;
- 15 • 2-{2-(2,6-dichlorophenyl)-1-[3'-(ethylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS(ES): 515.2 [M+H]⁺;
- 2-{2-(2-chloro-6-fluorophenyl)-1-[3'-(ethylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS(ES): 499.5 [M+H]⁺, 521.3 [M+Na]⁺;
- 2-{2-(2,6-dichlorophenyl)-1-[4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS(ES): 531.1, 533.3 [M+H]⁺;
- 20 • 2-{2-(2-chloro-6-fluorophenyl)-1-[4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS(ES): 515.7, 517.3 [M+H]⁺;
- 2-(2-(3-chloro-3'-(methylsulfonyl)biphenyl-2-yl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; MS (ES): 621.5 [M+H]⁺
- 25 • 2-{2-(2,3-dichlorophenyl)-1-[3'-(hydroxymethyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 453 [M+H]⁺.
- 2-{2-(2,3-dichlorophenyl)-1-[4'-(hydroxymethyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 453 [M+H]⁺.
- 2-{1-[3-methyl-3'-(methylsulfonyl)biphenyl-4-yl]-2-(3-thienyl)-1H-imidazol-4-yl}propan-2-ol; MS (ES): 453 [M+H]⁺.
- 30 • 2-{1-[3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl]-2-(2-chloro-6-fluorophenyl)-1H-imidazol-4-yl}propan-2-ol; MS (ES): 535 [M+H]⁺.

- 2-{1-[3-chloro-3'-(methylsulfonyl)biphenyl-4-yl]-2-(2,6-dichlorophenyl)-1H-imidazol-4-yl}propan-2-ol; MS (ES): 537 [M+H]⁺.
- 2-{1-[3'-(ethylsulfonyl)-3-methylbiphenyl-4-yl]-2-isoquinolin-5-yl-1H-imidazol-4-yl}propan-2-ol; MS (ES): 512 [M+H]⁺.
- 5 • 2-{2-isoquinolin-5-yl-1-[3-methyl-3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 498 [M+H]⁺.
- 2-{2-(1,5-dimethyl-1H-pyrrol-2-yl)-1-[3-methyl-3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 464 [M+H]⁺.
- 2-{1-[3-methyl-3'-(methylsulfonyl)biphenyl-4-yl]-2-(2-thienyl)-1H-imidazol-4-yl}propan-2-ol; MS
10 (ES): 453 [M+H]⁺.
- 2-{1-[3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl]-2-isoquinolin-1-yl-1H-imidazol-4-yl}propan-2-ol; MS (ES): 529 [M+H]⁺.
- 2-{1-[3-chloro-3'-(methylsulfonyl)biphenyl-4-yl]-2-isoquinolin-1-yl-1H-imidazol-4-yl}propan-2-ol; MS (ES): 518 [M+H]⁺.
- 15 • 2-{2-(3-chloro-2-methylphenyl)-1-[3'-(ethylsulfonyl)-3-fluorobiphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 513 [M+H]⁺.
- 2-{2-(3-chloro-2-methylphenyl)-1-[3-fluoro-3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 499 [M+H]⁺.
- 2-{2-(3-chloro-2-methylphenyl)-1-[3'-(ethylsulfonyl)-3-methylbiphenyl-4-yl]-1H-imidazol-4-
20 yl}propan-2-ol; MS (ES): 509 [M+H]⁺.
- 2-{2-(3-chloro-2-methylphenyl)-1-[3-methyl-3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 495 [M+H]⁺.
- 2-{1-[3'-(methylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 501 [M+H]⁺.
- 25 • 2-{1-[3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl]-2-naphthalen-1-yl-1H-imidazol-4-yl}propan-2-ol; MS (ES): 531 [M+H]⁺.
- 2-{1-[3'-(ethylsulfonyl)-3-methylbiphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 529 [M+H]⁺.
- 2-{1-[3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl]-2-(3-chloro-2-methylphenyl)-1H-imidazol-4-
30 yl}propan-2-ol; MS (ES): 529 [M+H]⁺.
- 2-{2-(3-chloro-2-methylphenyl)-1-[3-chloro-3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 515 [M+H]⁺.

- 2-{1-[3-methyl-3'-(methylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 515 [M+H]⁺.
- 2-{1-[3-chloro-3'-(methylsulfonyl)biphenyl-4-yl]-2-naphthalen-1-yl-1H-imidazol-4-yl}propan-2-ol; MS (ES): 517 [M+H]⁺.
- 5 • 2-{1-[3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl]-2-(2-chlorophenyl)-1H-imidazol-4-yl}propan-2-ol; MS (ES): 515 [M+H]⁺.
- 2-{1-[3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 549 [M+H]⁺.
- 2-{1-[3-chloro-3'-(methylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 535 [M+H]⁺.
- 10 • 2-{1-[3'-(ethylsulfonyl)-2-fluorobiphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 533 [M+H]⁺.
- 2-{1-[2-fluoro-3'-(methylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 519 [M+H]⁺.
- 15 • 2-{1-[3'-(ethylsulfonyl)-3-fluorobiphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 533 [M+H]⁺.
- 2-{1-[3-fluoro-3'-(methylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 519 [M+H]⁺.
- 2-(2-(2-chloro-6-methylphenyl)-1-(3-fluoro-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; MS (ES): 499.3 [M+H]⁺.
- 20 • 2-(2-(2-chloro-6-methylphenyl)-1-(3'-(ethylsulfonyl)-3-fluorobiphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; MS (ES): 513.3 [M+H]⁺.
- 2-(2-(2,6-dichlorophenyl)-1-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; MS (ES): 515.3, 517.3 [M+H]⁺.
- 25 • 2-(2-(2,6-dichlorophenyl)-1-(3'-(ethylsulfonyl)-3-methylbiphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; MS (ES): 529.3, 531.3 [M+H]⁺.
- 2-(1-(3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-1H-imidazol-4-yl)propan-2-ol; MS (ES): 549.3, 551.3, 553.3 [M+H]⁺.
- 2-(1-(3-chloro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-1H-imidazol-4-yl)propan-2-ol; MS (ES): 565.0, 567.0, 569.2 [M+H]⁺.
- 30 • 2-(1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2-chloro-6-methylphenyl)-1H-imidazol-4-yl)propan-2-ol; MS (ES): 515.2, 517.3 [M+H]⁺.

- 2-(1-(3-chloro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2-chloro-6-methylphenyl)-1H-imidazol-4-yl)propan-2-ol; MS (ES): 545.3, 547.3 $[M+H]^+$
- 2-(1-(3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl)-2-(2-chloro-6-methylphenyl)-1H-imidazol-4-yl)propan-2-ol; MS (ES): 529.2, 531.2 $[M+H]^+$
- 5 • 2-(2-(2,6-dichlorophenyl)-1-(3-fluoro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; MS (ES): 549.3, 551.3 $[M+H]^+$, 571.2, 573.2 $[M+Na]^+$
- 2-(2-(2-chloro-6-methylphenyl)-1-(3'-(ethylsulfonyl)-3-methylbiphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; MS (ES): 509.3, 511.3 $[M+H]^+$
- 2-(2-(2-chloro-6-methylphenyl)-1-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; MS (ES): 495.4, 497.4 $[M+H]^+$
- 10 • 2-(2-(2-chloro-6-methylphenyl)-1-(3'-(ethylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; MS (ES): 495.4 $[M+H]^+$
- 2-(2-(2-chloro-6-methylphenyl)-1-(4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; MS (ES): 511.3 $[M+H]^+$
- 15 • 2-(2-(2,6-dichlorophenyl)-1-(3-fluoro-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; MS (ES): 519.3, 521.3 $[M+H]^+$
- 2-(2-(2,6-dichlorophenyl)-1-(3'-(ethylsulfonyl)-3-methylbiphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; MS (ES): 529.3, 531.3 $[M+H]^+$, 551.3, 553.3 $[M+Na]^+$
- 2-(1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-1H-imidazol-4-yl)propan-2-ol; MS (ES): 535.3, 537.3, 539.3 $[M+H]^+$
- 20 • 2-(2-(2,6-dichlorophenyl)-1-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; MS (ES): 537.0 $[M+Na]^+$
- 2-(2-(2-chloro-6-methylphenyl)-1-(4'-(hydroxymethyl)-3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; MS (ES): 525.5 $[M+H]^+$, 547.3 $[M+Na]^+$
- 25 • 2-(2-(2-chloro-6-methylphenyl)-1-(3-fluoro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; MS (ES): 529.3, 531.3 $[M+H]^+$
- ethyl 1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-1H-imidazole-4-carboxylate; MS (ES): 549.0, 551.0, 553.0 $[M+H]^+$
- 2-(1-[2'-(trifluoromethyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl)propan-2-ol; MS (ES): 491 $[M+H]^+$
- 30 • 2-(1-(2',3'-difluorobiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl)propan-2-ol; MS (ES): 459 $[M+H]^+$

- 2-(1-{4'-[(1-methylethyl)oxy]biphenyl-4-yl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl)propan-2-ol; MS (ES): 481 [M+H]⁺;
- 2-{1-(4'-fluoro-3'-methylbiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 455 [M+H]⁺;
- 5 • 2-{1-(3'-fluorobiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 441 [M+H]⁺;
- 2-{1-(2',4',5'-trifluorobiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 477 [M+H]⁺;
- 2-{1-[5'-fluoro-2'-(methyloxy)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 471 [M+H]⁺;
- 10 • 2-{1-(4'-chlorobiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 457 [M+H]⁺;
- 2-{1-(4'-pyrimidin-5-ylphenyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 425 [M+H]⁺;
- 15 • 2-{1-[4'-(hydroxymethyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 453 [M+H]⁺;
- 2-{1-[4'-(dimethylamino)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 466 [M+H]⁺;
- 2-{1-[4'-(trifluoromethyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 491 [M+H]⁺;
- 20 • 2-{1-[4'-(1-methylethyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 465 [M+H]⁺;
- 2-{1-[4'-chloro-2'-(trifluoromethyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 525 [M+H]⁺;
- 25 • 2-{1-(2',3',4'-trifluorobiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 477 [M+H]⁺;
- 2-{1-(3',4'-difluorobiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 459 [M+H]⁺;
- 2-{1-(2'-chloro-6'-fluoro-3'-methylbiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 489 [M+H]⁺;
- 30 • 2-{1-[5'-chloro-2'-(methyloxy)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 487 [M+H]⁺;

- 2-{1-[2'-fluoro-5'-(trifluoromethyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 509 [M+H]⁺;
- 2-{1-[2'-(methylthio)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 469 [M+H]⁺;
- 5 • 4'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-4-carboxylic acid; MS (ES): 467 [M+H]⁺;
- 2-{1-[4-(1,3-benzodioxol-5-yl)phenyl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 467 [M+H]⁺;
- 2-(1-{4-[6-(methyloxy)pyridin-3-yl]phenyl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl)propan-2-ol; MS (ES): 454 [M+H]⁺;
- 10 • 2-(1-{4-[(1E)-3,3-dimethylbut-1-en-1-yl]phenyl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl)propan-2-ol; MS (ES): 429 [M+H]⁺;
- 2-{1-(3'-chlorobiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 457 [M+H]⁺;
- 15 • 2-{1-[2'-fluoro-5'-(methyloxy)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 471 [M+H]⁺;
- ethyl 4'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-4-carboxylate; MS (ES): 495 [M+H]⁺;
- 2-{1-(4-{2-[(1-methylethyl)oxy]pyridin-3-yl}phenyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 482 [M+H]⁺;
- 20 • 2-{1-[3'-chloro-4'-(methyloxy)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 487 [M+H]⁺;
- 2-{1-[2'-fluoro-3'-(methyloxy)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 471 [M+H]⁺;
- 25 • 4'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-3-carboxamide; MS (ES): 466 [M+H]⁺;
- 4'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-4-carboxamide; MS (ES): 466 [M+H]⁺;
- 2-(1-{4'-[(trifluoromethyl)oxy]biphenyl-4-yl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl)propan-2-ol; MS (ES): 507 [M+H]⁺;
- 30 • 2-{1-[4'-fluoro-3'-(trifluoromethyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 509 [M+H]⁺;

- 2-{1-(4'-propylbiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 465 [M+H]⁺;
- 2-{1-[4'-(ethyloxy)-3'-(trifluoromethyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 535 [M+H]⁺;
- 5 • 2-{1-[2'-[(1-methylethyl)oxy]biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 481 [M+H]⁺;
- 2-{1-[3'-chloro-4'-[(1-methylethyl)oxy]biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 515 [M+H]⁺;
- 2-{1-[4-(1H-indol-4-yl)phenyl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 462 [M+H]⁺;
- 10 • 2-{1-[4'-(methylthio)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 469 [M+H]⁺;
- 2-{1-[3'-(hydroxymethyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 453 [M+H]⁺;
- 15 • 2-{1-[3'-(ethyloxy)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 467 [M+H]⁺;
- 2-{1-[4'-ethylbiphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 451 [M+H]⁺;
- 2-{1-[2',4'-difluorobiphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 459 [M+H]⁺;
- 20 • 2-{1-[3',4'-dichlorobiphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 491 [M+H]⁺;
- 2-{1-[2'-chloro-4'-(trifluoromethyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 525 [M+H]⁺;
- 25 • 2-{1-[4-naphthalen-2-yl]phenyl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 473 [M+H]⁺;
- 2-{1-[3'-(methyloxy)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 453 [M+H]⁺;
- 4'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-3-ol; MS (ES): 439 [M+H]⁺;
- 30 • 2-{1-[3',4',5'-trifluorobiphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 477 [M+H]⁺;

- 1-[5-(4-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}phenyl)-2-thienyl]ethanone; MS (ES): 471 [M+H]⁺;
- 2-{1-(3',5'-difluorobiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 459 [M+H]⁺;
- 5 • 2-{1-(3'-chloro-4'-fluorobiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 475 [M+H]⁺;
- 2-{1-[5'-methyl-2'-(methyloxy)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 467 [M+H]⁺;
- 2-{1-(2',5'-difluorobiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 459 [M+H]⁺;
- 10 • 2-{1-[3'-(butoxy)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 495 [M+H]⁺;
- 2-{1-[5'-chloro-2'-(ethyloxy)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 501 [M+H]⁺;
- 15 • 2-(1-{3'-[(trifluoromethyl)oxy]biphenyl-4-yl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl)propan-2-ol; MS (ES): 507 [M+H]⁺;
- 2-{1-(2',3',5'-trifluorobiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 477 [M+H]⁺;
- 2-{1-[3'-(ethylthio)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 483 [M+H]⁺;
- 20 • 2-(1-{3'-[(1-methylethyl)oxy]biphenyl-4-yl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl)propan-2-ol; MS (ES): 481 [M+H]⁺;
- 2-{1-[4-(1-benzothien-3-yl)phenyl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 479 [M+H]⁺;
- 25 • 2-{1-[4-(4-methylnaphthalen-1-yl)phenyl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 487 [M+H]⁺;
- 2-{1-(2',4'-dichlorobiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 491 [M+H]⁺;
- 2-{1-[3',4'-bis(methyloxy)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 483 [M+H]⁺;
- 30 • 2-{2-[2-(trifluoromethyl)phenyl]-1-(2',4',5'-trimethylbiphenyl-4-yl)-1H-imidazol-4-yl}propan-2-ol; MS (ES): 465 [M+H]⁺;

- 4-fluoro-4'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-2-ol; MS (ES): 457 [M+H]⁺;
- 2-{1-[2'-(hydroxymethyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 453 [M+H]⁺;
- 5 • 2-{1-[3'-(trifluoromethyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 491 [M+H]⁺;
- 2-{1-(2'-chloro-6'-fluorobiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 475 [M+H]⁺;
- 2-{1-[3',5'-difluoro-2'-(methyloxy)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 489 [M+H]⁺;
- 10 • 2-(1-{4-[2-(methyloxy)pyridin-3-yl]phenyl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl)propan-2-ol; MS (ES): 454 [M+H]⁺;
- 2-{1-[2'-methyl-5'-(methyloxy)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 467 [M+H]⁺;
- 15 • 2-{1-(2'-ethylbiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 451 [M+H]⁺;
- 2-(1-{2'-methyl-4'-[(1-methylethyl)oxy]biphenyl-4-yl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl)propan-2-ol; MS (ES): 495 [M+H]⁺;
- 2-{1-[4'-(ethylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 515 [M+H]⁺;
- 20 • 2-{1-(5'-fluoro-2'-methylbiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 455 [M+H]⁺;
- 2-{1-[3'-chloro-4'-(trifluoromethyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 525 [M+H]⁺;
- 25 • 2-{1-(2',5'-dimethylbiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 451 [M+H]⁺;
- 2-{1-[2'-(ethyloxy)-5'-(trifluoromethyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 535 [M+H]⁺;
- 2-{1-(3'-fluoro-4'-methylbiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 455 [M+H]⁺;
- 30 • methyl (2E)-3-(4'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-4-yl)prop-2-enoate; MS (ES): 507 [M+H]⁺;

- N-ethyl-4'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-3-carboxamide; MS (ES): 494 [M+H]⁺;
- 2-{1-[4-(2,3-dihydro-1-benzofuran-5-yl)phenyl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 465 [M+H]⁺;
- 5 • N-butyl-4'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-3-sulfonamide; MS (ES): 558 [M+H]⁺;
- N-(1,1-dimethylethyl)-4'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}-6-methylbiphenyl-3-sulfonamide; MS (ES): 572 [M+H]⁺;
- 4'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}-N-methylbiphenyl-3-sulfonamide; MS (ES): 516 [M+H]⁺;
- 10 • N-ethyl-4'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-3-sulfonamide; MS (ES): 530 [M+H]⁺;
- N-(1,1-dimethylethyl)-4'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-3-sulfonamide; MS (ES): 558 [M+H]⁺;
- 15 • 2-{1-[2'-amino-5'-(trifluoromethyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 506 [M+H]⁺;
- 2-{1-[3'-fluoro-5'-(trifluoromethyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 509 [M+H]⁺;
- 2-{1-[4'-chloro-3'-(trifluoromethyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 525 [M+H]⁺;
- 20 • 3-chloro-4'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-4-carboxylic acid; MS (ES): 501 [M+H]⁺;
- 2-(2-[2-(trifluoromethyl)phenyl]-1-{3'-[(trifluoromethyl)thio]biphenyl-4-yl}-1H-imidazol-4-yl)propan-2-ol; MS (ES): 523 [M+H]⁺;
- 25 • 2-{1-[3'-(morpholin-4-ylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 572 [M+H]⁺;
- 2-(1-{4-[5-(hydroxymethyl)-1,3-thiazol-2-yl]phenyl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl)propan-2-ol; MS (ES): 460 [M+H]⁺;
- 2-{1-[2'-methyl-5'-(piperidin-1-ylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 584 [M+H]⁺;
- 30 • 1-[4-(4'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}phenyl)-2-thienyl]ethanone; MS (ES): 471 [M+H]⁺;

- 2-(1-{4-[5-(hydroxymethyl)-3-thienyl]phenyl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl)propan-2-ol; MS (ES): 459 [M+H]⁺;
- 2-chloro-4'-[2-(2,6-dichlorophenyl)-4-(1-hydroxy-1-methylethyl)-1H-imidazol-1-yl]-3'-methylbiphenyl-4-carboxylic acid; MS (ES): 515 [M+H]⁺;
- 5 • 2-(1-{3-[6-(methyloxy)pyridin-3-yl]phenyl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl)propan-2-ol; MS (ES): 454 [M+H]⁺;
- 3'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}-N-(1-methylethyl)biphenyl-3-sulfonamide; MS (ES): 544 [M+H]⁺;
- 3'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-4-
10 carboxamide; MS (ES): 466 [M+H]⁺;
- 2-(1-{3-[2-(cyclopentyloxy)pyridin-3-yl]phenyl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl)propan-2-ol; MS (ES): 508 [M+H]⁺;
- 2-{1-(3'-chlorobiphenyl-3-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS
(ES): 457 [M+H]⁺;
- 15 • 2-{1-[2'-fluoro-5'-(methyloxy)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 471 [M+H]⁺;
- 3'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-3-carboxamide; MS (ES): 466 [M+H]⁺;
- 2-{1-[3',4'-bis(methyloxy)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-
20 2-ol; MS (ES): 483 [M+H]⁺;
- N-(3'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-2-yl)methanesulfonamide; MS (ES): 516 [M+H]⁺;
- 2-{1-[3'-(trifluoromethyl)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 491 [M+H]⁺;
- 25 • 3-fluoro-3'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-4-carboxylic acid; MS (ES): 485 [M+H]⁺;
- 4-chloro-3'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-3-carboxamide; MS (ES): 500 [M+H]⁺;
- 2-{1-(3-{2-[(1-methylethyl)oxy]pyridin-3-yl}phenyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-
30 4-yl}propan-2-ol; MS (ES): 482 [M+H]⁺;
- N-(3'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-3-yl)acetamide; MS (ES): 480 [M+H]⁺;

- 2-{1-[2'-methyl-5'-(methyloxy)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 467 [M+H]⁺;
- N-(3'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-4-yl)acetamide; MS (ES): 480 [M+H]⁺;
- 5 • 2-{1-(4'-chlorobiphenyl-3-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 457 [M+H]⁺;
- 2-{1-[4'-(phenyloxy)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 515 [M+H]⁺;
- 2-{1-[5'-fluoro-2'-(methyloxy)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 471 [M+H]⁺;
- 10 • 2-{1-(2',3'-dichlorobiphenyl-3-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 491 [M+H]⁺;
- 2-(1-{3'-[(trifluoromethyl)oxy]biphenyl-3-yl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl)propan-2-ol; MS (ES): 507 [M+H]⁺;
- 15 • 3'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-3-sulfonamide; MS (ES): 502 [M+H]⁺;
- 2-{1-[3',5'-bis(trifluoromethyl)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 559 [M+H]⁺;
- 2-{1-(3',5'-dichlorobiphenyl-3-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 491 [M+H]⁺;
- 20 • 3-chloro-3'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}-N-(1-methylethyl)biphenyl-4-carboxamide; MS (ES): 542 [M+H]⁺;
- N,N-diethyl-3'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-3-carboxamide; MS (ES): 522 [M+H]⁺;
- 25 • 4-chloro-3'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}-N-(1-methylethyl)biphenyl-3-carboxamide; MS (ES): 542 [M+H]⁺;
- 2-{1-[3'-(methyloxy)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 453 [M+H]⁺;
- 3'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-3-carboxylic acid; MS (ES): 467 [M+H]⁺;
- 30 • N-ethyl-3'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-3-carboxamide; MS (ES): 494 [M+H]⁺;

- 4-chloro-N-ethyl-3'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-3-carboxamide; MS (ES): 528 [M+H]⁺;
- 2-{1-(2',5'-difluorobiphenyl-3-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 459 [M+H]⁺;
- 5 • 2-(1-{3'-[(1-methylethyl)oxy]biphenyl-3-yl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl)propan-2-ol; MS (ES): 481 [M+H]⁺;
- 2-{1-[2'-fluoro-5'-(trifluoromethyl)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 509 [M+H]⁺;
- 2-{1-(3',4'-dichlorobiphenyl-3-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
- 10 MS (ES): 491 [M+H]⁺;
- 2-{1-[3'-(ethyloxy)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 467 [M+H]⁺;
- 2-(1-{2'-methyl-4'-[(1-methylethyl)oxy]biphenyl-3-yl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl)propan-2-ol; MS (ES): 495 [M+H]⁺;
- 15 • 2-{1-[3-(1-methyl-1H-indol-5-yl)phenyl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 476 [M+H]⁺;
- 2-{1-[4'-(ethyloxy)-3'-(trifluoromethyl)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 535 [M+H]⁺;
- 2-{1-[3'-(ethylsulfonyl)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
- 20 MS (ES): 515 [M+H]⁺;
- 2-{1-[2'-(trifluoromethyl)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 491 [M+H]⁺;
- 2-{1-[3'-(hydroxymethyl)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 453 [M+H]⁺;
- 25 • 2-{1-[3-(1H-indol-4-yl)phenyl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 462 [M+H]⁺;
- 3'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-4-carboxylic acid; MS (ES): 467 [M+H]⁺;
- 1-[5-(3'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}phenyl)-2-thienyl]ethanone; MS (ES): 471 [M+H]⁺;
- 30 • 2-{1-(5'-chloro-2'-methylbiphenyl-3-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 471 [M+H]⁺;

- 2-(1-{4'-[(trifluoromethyl)oxy]biphenyl-3-yl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl)propan-2-ol; MS (ES): 507 [M+H]⁺;
- 2-{1-[2'-chloro-4'-(trifluoromethyl)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 525 [M+H]⁺;
- 5 • 2-{1-(2',5'-dichlorobiphenyl-3-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 491 [M+H]⁺;
- 2-{1-[2'-(ethyloxy)-5'-(trifluoromethyl)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 535 [M+H]⁺;
- 3'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}-N-(1-methylethyl)biphenyl-4-carboxamide; MS (ES): 508 [M+H]⁺;
- 10 • 2-{1-(2',4'-dichlorobiphenyl-3-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 491 [M+H]⁺;
- 2-{1-[4'-(ethylsulfonyl)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 515 [M+H]⁺;
- 15 • 2-{1-[4'-fluoro-3'-(trifluoromethyl)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 509 [M+H]⁺;
- 2-{1-(3'-fluoro-4'-methylbiphenyl-3-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 455 [M+H]⁺;
- 2-{2-(2-chloro-6-fluorophenyl)-1-[3'-(ethylsulfonyl)-3-fluorobiphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 539(M+Na);
- 20 • 2-{2-(2-chloro-6-fluorophenyl)-1-[3-fluoro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol MS (ES): 533[M+H]⁺
- 2-{2-(2,6-difluorophenyl)-1-[3-fluoro-3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 487 [M+H]⁺
- 25 • 2-{2-(2,6-difluorophenyl)-1-[3'-(ethylsulfonyl)-3-fluorobiphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 501 [M+H]⁺
- 2-{2-(2,6-difluorophenyl)-1-[3-fluoro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol MS (ES): 517[M+H]⁺
- 2-{2-(2-chloro-6-fluorophenyl)-1-[3-fluoro-3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol MS (ES): 503 [M+H]⁺
- 30 • 2-{2-(2,6-difluorophenyl)-1-[4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol MS (ES): 499[M+H]⁺

- 2-{2-(2-chloro-6-fluorophenyl)-1-[3-chloro-3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol MS (ES): 519[M+H]⁺
- 2-{1-[3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl]-2-(2-chloro-6-fluorophenyl)-1H-imidazol-4-yl}propan-2-ol MS (ES): 533[M+H]⁺
- 5 • 2-{1-[3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl]-2-(2-chloro-6-fluorophenyl)-1H-imidazol-4-yl}propan-2-ol MS (ES): 533[M+H]⁺
- 2-{2-(2-chloro-6-fluorophenyl)-1-[3-chloro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol MS (ES): 549[M+H]⁺
- 2-{2-(2-chloro-6-fluorophenyl)-1-[3-methyl-3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol MS (ES): 499[M+H]⁺
- 10 • 2-{2-(2-chloro-6-fluorophenyl)-1-[3'-(ethylsulfonyl)-3-methylbiphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol MS (ES): 513[M+H]⁺
- 2-{2-(2-chloro-6-fluorophenyl)-1-[4'-(hydroxymethyl)-3-methyl-3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol MS (ES): 529[M+H]⁺
- 15 • 2-{1-[3'-(methylsulfonyl)biphenyl-4-yl]-2-[4-(trifluoromethyl)pyridin-3-yl]-1H-imidazol-4-yl}propan-2-ol MS (ES): 502[M+H]⁺
- 2-{1-[3'-(ethylsulfonyl)biphenyl-4-yl]-2-[4-(trifluoromethyl)pyridin-3-yl]-1H-imidazol-4-yl}propan-2-ol MS (ES): 516[M+H]⁺
- 2-{1-[4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl]-2-[4-(trifluoromethyl)pyridin-3-yl]-1H-imidazol-4-yl}propan-2-ol MS (ES): 532[M+H]⁺
- 20 • 2-{1-[3'-(methylsulfonyl)biphenyl-4-yl]-2-(2-morpholin-4-ylethyl)-1H-imidazol-4-yl}propan-2-ol MS (ES): 470[M+H]⁺
- 2-{1-[3-chloro-3'-(methylsulfonyl)biphenyl-4-yl]-2-(2,6-difluorophenyl)-1H-imidazol-4-yl}propan-2-ol MS (ES): 503[M+H]⁺
- 25 • 2-{1-[3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl]-2-(2,6-difluorophenyl)-1H-imidazol-4-yl}propan-2-ol MS (ES): 517[M+H]⁺
- 2-{1-[3-chloro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl]-2-(2,6-difluorophenyl)-1H-imidazol-4-yl}propan-2-ol MS (ES): 533[M+H]⁺
- 2-{2-(2,6-difluorophenyl)-1-[3-methyl-3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol MS (ES): 483[M+H]⁺
- 30 • 2-{2-(2,6-difluorophenyl)-1-[3'-(ethylsulfonyl)-3-methylbiphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol MS (ES): 497[M+H]⁺

- 2-{2-(2,6-difluorophenyl)-1-[4'-(hydroxymethyl)-3-methyl-3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol MS (ES): 513[M+H]⁺
- 2-{2-(2,3-dichlorophenyl)-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS(ES): 501, 503 [M+H]⁺
- 5 • 2-{2-(2-chloro-3-fluorophenyl)-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS(ES): 485 [M+H]⁺
- 2-{2-(2-chlorophenyl)-1-[3-fluoro-3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS(ES): 485 [M+H]⁺
- 2-{2-(2-chlorophenyl)-1-[2-fluoro-3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS(ES): 485 [M+H]⁺
- 10 • 2-{1-[3'-(methylsulfonyl)biphenyl-4-yl]-2-[3-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS(ES): 501 [M+H]⁺
- 2-{2-(3-chlorophenyl)-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS(ES): 467 [M+H]⁺
- 15 • 2-{1-[3-chloro-3'-(methylsulfonyl)biphenyl-4-yl]-2-(2-chlorophenyl)-1H-imidazol-4-yl}propan-2-ol; MS(ES): 501, 503 [M+H]⁺
- 2-{2-(2-chlorophenyl)-1-[3-methyl-3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS(ES): 481 [M+H]⁺
- 2-{2-(2,3-dichlorophenyl)-1-[3'-(ethylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS(ES): 515, 517 [M+H]⁺
- 20 • 2-{2-(2-chlorophenyl)-1-[3'-(ethylsulfonyl)-3-fluorobiphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS(ES): 499 [M+H]⁺
- 2-{2-(2-chlorophenyl)-1-[3'-(ethylsulfonyl)-2-fluorobiphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS(ES): 499 [M+H]⁺
- 25 • 2-{2-(3-chloro-2-methylphenyl)-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS(ES): 481 [M+H]⁺
- 2-{2-(2-chlorophenyl)-1-[2-methyl-3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS(ES): 481 [M+H]⁺
- 2-{2-(3-chloro-2-methylphenyl)-1-[3'-(ethylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS(ES): 495 [M+H]⁺
- 30 • 2-(2-(2-chlorophenyl)-1-(3-ethyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; MS(ES): 495 [M+H]⁺

- 2-(1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2,3-dichlorophenyl)-1H-imidazol-4-yl)propan-2-ol; MS(ES): 535, 537 [M+H]⁺
- 2-(1-(3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl)-2-(2,3-dichlorophenyl)-1H-imidazol-4-yl)propan-2-ol; MS(ES): 549, 551 [M+H]⁺
- 5 • 2-(2-(2,3-dichlorophenyl)-1-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; MS(ES): 515, 517 [M+H]⁺
- 2-(2-(2,3-dichlorophenyl)-1-(3'-(ethylsulfonyl)-3-methylbiphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; MS(ES): 529, 531 [M+H]⁺
- 2-(2-(2,3-dichlorophenyl)-1-(3-fluoro-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-10 2-ol; MS(ES): 519, 521 [M+H]⁺
- 2-(2-(2,3-dichlorophenyl)-1-(3'-(ethylsulfonyl)-3-fluorobiphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; MS(ES): 533, 535 [M+H]⁺
- 2-(2-(2-chloro-3-fluorophenyl)-1-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; MS(ES): 499 [M+H]⁺
- 15 • 2-(2-(2-chloro-3-fluorophenyl)-1-(3'-(ethylsulfonyl)-3-methylbiphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; MS(ES): 513 [M+H]⁺
- 2-(2-(2,3-dichlorophenyl)-1-(4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; MS(ES): 531, 533 [M+H]⁺
- 2-(2-(2,3-dichlorophenyl)-1-(3-fluoro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; MS(ES): 549, 551 [M+H]⁺
- 20 • 2-(2-(2-chloro-3-fluorophenyl)-1-(4'-(hydroxymethyl)-3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; MS(ES): 529 [M+H]⁺
- 2-(1-(3-chloro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2,3-dichlorophenyl)-1H-imidazol-4-yl)propan-2-ol; MS(ES): 565, 567 [M+H]⁺
- 25 • 2-(2-(2,3-dichlorophenyl)-1-(4'-(hydroxymethyl)-3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; MS(ES): 545, 547 [M+H]⁺
- 2-(1-(3-chloro-5-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2-chlorophenyl)-1H-imidazol-4-yl)propan-2-ol; MS(ES): 537, 539 [M+Na]⁺
- 2-(1-(3-chloro-3'-(ethylsulfonyl)-5-methylbiphenyl-4-yl)-2-(2-chlorophenyl)-1H-imidazol-4-yl)propan-2-ol; MS(ES): 551, 553 [M+Na]⁺
- 30 • 2-(1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2-chloro-3-fluorophenyl)-1H-imidazol-4-yl)propan-2-ol; MS(ES): 541, 543 [M+Na]⁺


- 2-(1-(3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl)-2-(2-chloro-3-fluorophenyl)-1H-imidazol-4-yl)propan-2-ol; MS(ES): 555, 557 [M+Na]⁺
- 2-(2-(2-chloro-3-fluorophenyl)-1-(3-fluoro-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; MS(ES): 525 [M+Na]⁺
- 5 • 2-(2-(2-chloro-3-fluorophenyl)-1-(3'-(ethylsulfonyl)-3-fluorobiphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; MS(ES): 517 [M+H]⁺
- 2-(2-(2-chloro-3-fluorophenyl)-1-(3-fluoro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; S(ES): 555 [M+Na]⁺
- 2-(2-(2-chloro-3-fluorophenyl)-1-(3-chloro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; MS(ES): 571, 573 [M+Na]⁺
- 10 • 2-(2-(2-chloro-3-fluorophenyl)-1-(3'-(ethylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; MS(ES): 499 [M+H]⁺

In a similar manner, the following compounds were prepared by replacing methylmagnesium bromide with ethylmagnesium bromide:

- 15 • 3-(2-(2,3-dichlorophenyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)pentan-3-ol; MS(ES): 529, 531 [M+H]⁺;
- 3-(2-(2,3-dichlorophenyl)-1-(3'-(ethylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)pentan-3-ol; MS(ES): 543, 545 [M+H]⁺;
- 3-{2-(2,6-dichlorophenyl)-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}pentan-3-ol; 529.3, [M+H]⁺.
- 20

In a similar manner, the following compound was prepared by replacing ethyl bromopyruvate with ethyl 3-bromo-2-ketobutyrate: 2-{5-methyl-1-[3'-(methylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS(ES): 515 [M+H]⁺

Example 10



Preparation of 2-[5-bromo-1-(3'-methanesulfonyl-biphenyl-4-yl)-2-(2-trifluoromethyl-phenyl)-1H-imidazol-4-yl]-propan-2-ol

To a solution of 2-[1-(3'-methanesulfonyl-biphenyl-4-yl)-2-(2-trifluoromethyl-phenyl)-1H-imidazol-4-yl]-propan-2-ol (100 mg, 0.20 mmol) in MeCN (2 mL, anhyd) was added N-bromosuccinimide (45 mg, 0.25 mmol). After 2h, the reaction mixture was diluted with EtOAc,

washed with saturated NaHCO₃, water (3x 10 mL) and brine, then dried (Na₂SO₄) and concentrated. The residue was purified by chromatography (silica), eluting with EtOAc/hexanes (65:40 to 95:5) to yield the title compound (100 g, 86%) as a white solid. ¹H-NMR (CD₂Cl₂): δ 8.10 (1H, m), 7.92 (1H, m), 7.86 (1H, m), 7.60-7.74 (4H, m), 7.47 (2H, m), 7.25-7.31 (3H, m), 3.56 (1H, s), 3.06 (3H, s), 1.66 (6H, s); MS (ES): 579, 581 [M+H]⁺.

In a similar manner, the following compound was prepared by replacing N-bromosuccinimide with N-chlorosuccinimide: 2-{5-chloro-1-[3'-(methylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS(ES): 535 [M+H]⁺


In a similar manner, the following compound was prepared by treating the appropriate imidazole-4-carboxylic acid ethyl ester intermediate with 1-chloromethyl-4-fluoro-1,4-diazoniabicyclo[2.2.2]octane bis(tetrafluoroborate) (SelectfluorTM) followed by the previously described sequence of Suzuki cross-coupling and reaction with Grignard reagent: 2-{5-fluoro-1-[3'-(methylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS(ES): 519 [M+H]⁺

The following compounds were made in a similar manner, except boronic acid or boronate reagents for Suzuki coupling were prepared as described in Examples 23-29.

- 6-{3-Chloro-4-[2-(2-chloro-6-fluorophenyl)-4-(1-hydroxy-1-methylethyl)-imidazol-1-yl]-phenyl}-1,1-dioxo-2,3-dihydro-1H-1λ⁶-benzo[b]thiophen-3-ol; MS (ES): 547 [M+H]⁺.
- 2-{2-(2,3-dichlorophenyl)-1-[4'-(methoxy)-3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 531, 533 each [M+H]⁺.
- 2-{2-(2-chloro-6-fluorophenyl)-1-[3-chloro-4'-(methoxy)-3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 549, 551 each [M+H]⁺.
- ethyl 3'-chloro-4'-[2-(2-chloro-6-fluorophenyl)-4-(1-hydroxy-1-methylethyl)-1H-imidazol-1-yl]-3-(methylsulfonyl)biphenyl-4-carboxylate; MS (ES): 591, 593 each [M+H]⁺.
- 2-{2-(2,3-dichlorophenyl)-1-[4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 531, 533 each [M+H]⁺.
- 2-{4'-[2-(2,3-dichlorophenyl)-4-(1-hydroxy-1-methylethyl)-1H-imidazol-1-yl]-3-(methylsulfonyl)biphenyl-4-yl}propan-2-ol; MS (ES): 559 [M+H]⁺.
- 2-{3'-chloro-4'-[2-(2-chloro-6-fluorophenyl)-4-(1-hydroxy-1-methylethyl)-1H-imidazol-1-yl]-3-(methylsulfonyl)biphenyl-4-yl}propan-2-ol; MS (ES): 577 [M+H]⁺.
- 2-{1-[3-chloro-4'-(1-hydroxy-1-methylethyl)-3'-(methylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 593 [M+H]⁺.

- 2-{3'-chloro-4'-[2-(2-chlorophenyl)-4-(1-hydroxy-1-methylethyl)-1H-imidazol-1-yl]-3-(methylsulfonyl)biphenyl-4-yl}propan-2-ol; MS (ES): 559 [M+H]⁺.
- 2-{2-(3-chloro-2-methylphenyl)-1-[4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 511 [M+H]⁺.
- 5 • 2-{1-[3-chloro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl]-2-(3-chloro-2-methylphenyl)-1H-imidazol-4-yl}propan-2-ol; MS (ES): 545 [M+H]⁺.
- 2-{1-[3-fluoro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 549 [M+H]⁺.
- 2-{2-(3-chloro-2-methylphenyl)-1-[3-fluoro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 529 [M+H]⁺.
- 10 • 2-{2-(3-chloro-2-methylphenyl)-1-[4'-(hydroxymethyl)-3-methyl-3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 525 [M+H]⁺.
- 2-{1-[3-chloro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl]-2-naphthalen-1-yl-1H-imidazol-4-yl}propan-2-ol; MS (ES): 547 [M+H]⁺.
- 15 • 2-{1-[4'-(hydroxymethyl)-3-methyl-3'-(methylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 545 [M+H]⁺.
- 2-{1-[3-chloro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 565 [M+H]⁺.
- 2-{1-[3-chloro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl]-2-(2-chlorophenyl)-1H-imidazol-4-yl}propan-2-ol; MS (ES): 531 [M+H]⁺.
- 20 • ethyl 4'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}-5-(methylsulfonyl)biphenyl-3-carboxylate; MS (ES): 573 [M+H]⁺.
- 2-{1-[4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 531 [M+H]⁺.
- 25 • 2-{1-[4'-(methyloxy)-3'-(methylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 531 [M+H]⁺.
- 2-(2-(2,6-dichlorophenyl)-1-(3'-methyl-5'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; MS(ES): 515, 517 [M+H]⁺.
- 2-(1-(2'-chloro-5'-(methylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-1H-imidazol-4-yl)propan-2-ol; MS(ES): 535, 537 [M+H]⁺.
- 30 • 2-(1-(2',3'-dichloro-5'-(methylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-1H-imidazol-4-yl)propan-2-ol; MS(ES): 591, 593, 595 [M+Na]⁺.

Scheme 9



(a) Me_3Al , Toluene, 0°C - 80°C ; (b) Ethyl bromopyruvate, NaHCO_3 , EtOH , 80°C ; (c) R_3MgBr , THF , 0°C -rt; (d) $\text{ArB}(\text{OH})_2$, K_2CO_3 , $\text{PdCl}_2(\text{dppf})$, $\text{DME}/\text{H}_2\text{O}$, 80°C .


In general, compounds of formula (0096) are prepared by first reacting aniline of formula (0092) with benzyl cyanide (0091), such as 2, 3-dichlorobenzyl cyanide, in the presence of base to give compounds of formula (0093) after standard isolation procedures. In a subsequent step, exposure of amidine (0093) to haloester, such as ethyl α -bromopyruvate, under basic condition at elevated temperature provides 1H-imidazole of formula (0094) after standard isolation procedures. Compounds of formula (0094) are then subjected to functionality transformation, such as from ester to carbinol. In a palladium mediated coupling reaction, for example Suzuki reactions, compounds of formula (0095) are then reacted with a diboronate or boronic acid reagent to give compounds of formula (0096) after standard isolation procedures.

Example 11

2-{2-[(2-fluorophenyl)methyl]-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol

Example 11a

Preparation of N-(4-bromophenyl)-2-(2-chlorophenyl)acetimidamide




4-Bromoaniline (6.4 g, 37 mmol) was placed into a three-neck reactor, and the reactor was flushed with argon for 20min. 60 mL anhydrous toluene was added. After cooling to 0°C , trimethylaluminum (28 mL, 2.0M solution in toluene, 56 mmol) was added slowly to keep the inner temperature below 20°C . Over 0.5 h, the addition was completed. Ice bath was removed, and the reaction mixture was warmed to room temperature and then stirred for additional 2.5 hrs. A solution of

2-chlorobenzyl cyanide (8.4 g, 55 mmol) in 35 mL anhydrous toluene was added, and the reaction mixture was heated to 80 °C. After 18 hrs, the reaction mixture was cooled to room temperature and poured over a slurry of silica gel in CHCl₃/MeOH (4/1, 400 mL). After stirring for 30 min, the mixture was filtered and the residue was washed with a mixture of CHCl₃/MeOH (4/1). The combined filtrates
5 were concentrated *in vacuo*, and the resulting solid was washed with hexane to give a light pink solid (5.78 g, 48% yield).

Example 11b


Preparation of ethyl 1-(4-bromophenyl)-2-(2-chlorobenzyl)-1H-imidazole-4-carboxylate



10 To a mixture of N-(4-Bromo-phenyl)-2-(2-chloro-phenyl)-acetamidine (5.78 g, 17.9 mmol, from above reaction) and sodium bicarbonate (3.0 g, 36 mmol) in 50 mL 2-propanol was added ethyl α-bromopyruvate (7.8 g, 36 mmol, 90% tech.). The mixture was heated to 80 °C overnight, and then cooled to room temperature. The supernatant was decanted and concentrated *in vacuo*. The residue was redissolved into dichloromethane, and then filtered. The precipitate was washed several times with
15 dichloromethane. The combined filtrates were concentrated down. The crude product was purified by column chromatography on silica gel (30 → 50% EtOAc/Hexane) to give a brown solid (1.75 g, 23% yield).

Example 11c

Preparation of 2-(1-(4-bromophenyl)-2-(2-chlorobenzyl)-1H-imidazol-4-yl)propan-2-ol




20 Methylmagnesium bromide (13 mL, 18.6 mmol, 1.4M in toluene/THF 75:25) was placed into a three-neck flask. Under nitrogen at 0 °C to it was added a solution of 1-(4-Bromo-phenyl)-2-(2-chloro-benzyl)-1H-imidazole-4-carboxylic acid ethyl ester (1.74 g, 4.1 mmol) in 15 mL THF. After the addition, the ice bath was removed and the reaction mixture was stirred at room temperature for 3 hrs.
25 The reaction was quenched with sat. Ammonium chloride. Two layers were separated, and the aqueous layer was extracted three times with ethyl acetate. The combined organic layers were washed with

brine, dried over Na₂SO₄, and concentrated *in vacuo*. The crude product was purified by column chromatography on silica gel (30 → 50% EtOAc/Hexane) to give a yellow solid (600 mg, 36% yield).

Example 11d

Preparation of 2-(2-(2-chlorobenzyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol

5



2-[1-(4-Bromo-phenyl)-2-(2-chloro-benzyl)-1H-imidazol-4-yl]-propan-2-ol (202 mg, 0.5 mmol), (3-methylsulfonyl)phenylboronic acid (200 mg, 1.0 mmol), potassium carbonate (311 mg, 2.3 mmol), and PdCl₂(dppf).CH₂Cl₂ (42 mg, 0.05 mmol) were mixed with 5 mL 9:1 DME/H₂O (v/v), then heated at 80°C overnight. All solvent was removed *in vacuo*. The crude product was purified by chromatography on silica gel (50 → 95 % EtOAc/Hexane) to give a white solid (158 mg, 66% yield). ¹H-NMR (400MHz, *d*-DMSO): δ 1.43 (s, 6H), 3.30 (s, 3H), 4.13 (s, 2H), 4.73 (s, 1H), 7.11 – 7.09 (m, 1H), 7.13 (m, 1H), 7.24 – 7.20 (m, 2H), 7.37 – 7.34 (m, 1H), 7.47 (m, 2H), 7.75 (t, 1H, *J*=7.8), 7.88 – 7.86 (m, 2H), 7.94 – 7.92 (m, 1H), 8.08 – 8.06 (m, 1H), 8.18 (t, 1H, *J*=1.9). MS (ES): 481.0 [M+H]⁺.

15

All the following compounds were made in similar manner using appropriate nitriles and anilines. The boronic acid or boronate reagents for Suzuki coupling, if not commercially available, were made using standard techniques that are readily apparent to one skilled in the art.

- 2-[2-[(2-fluorophenyl)methyl]-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl]propan-2-ol; MS (ES): 465.0 [M+H]⁺;

20

- 2-[1-[3'-(methylsulfonyl)biphenyl-4-yl]-2-{[2-(trifluoromethyl)phenyl]methyl}-1H-imidazol-4-yl]propan-2-ol; MS (ES): 515.3 [M+H]⁺;

- 2-[2-[(2-chlorophenyl)methyl]-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl]propan-2-ol; MS (ES): 481.0 [M+H]⁺;

- 2-[2-[(2-chlorophenyl)methyl]-1-[3'-(ethylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl]propan-2-ol; MS (ES): 495.4 [M+H]⁺;

25

- 2-[2-[(2-chlorophenyl)methyl]-1-[4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl]propan-2-ol; MS (ES): 511.4 [M+H]⁺;

- 2-[2-[(2,6-dichlorophenyl)methyl]-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl]propan-2-ol; MS (ES): 515.3 [M+H]⁺;


- 2-{2-[(2,6-dichlorophenyl)methyl]-1-[3'-(ethylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 529.3 [M+H]⁺;
- 2-{2-[(2,6-dichlorophenyl)methyl]-1-[3'-(ethylsulfonyl)-4'-(hydroxymethyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 545.3 [M+H]⁺;
- 5 • 2-{2-[(2-chloro-6-fluorophenyl)methyl]-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 499.5 [M+H]⁺;
- 2-{2-[(2-chloro-6-fluorophenyl)methyl]-1-[3'-(ethylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 513.5 [M+H]⁺;
- 2-{2-[(2,3-dichlorophenyl)methyl]-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 529.3 [M+H]⁺;
- 10 • 2-{2-[(2,3-dichlorophenyl)methyl]-1-[3'-(ethylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 529.3 [M+H]⁺;
- 3-{2-[(2-chlorophenyl)methyl]-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}pentan-3-ol; MS (ES): 509.3 [M+H]⁺;
- 15 • 3-{2-[(2-chlorophenyl)methyl]-1-[3'-(ethylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}pentan-3-ol; MS (ES): 523.3 [M+H]⁺;
- 3-{2-[(2-chlorophenyl)methyl]-1-[4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}pentan-3-ol; MS (ES): 539.2 [M+H]⁺;
- 2-(2-(5-chloro-2-(trifluoromethyl)benzyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; MS (ES): 549.3 [M+H]⁺;
- 20 • 2-(2-(5-chloro-2-(trifluoromethyl)benzyl)-1-(3'-(ethylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; MS (ES): 563.3 [M+H]⁺;
- 2-(2-(5-chloro-2-(trifluoromethyl)benzyl)-1-(4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; MS (ES): 579.2 [M+H]⁺;
- 25 • 2-(2-(2-(2-chlorophenyl)propan-2-yl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; MS (ES): 509.0 [M+H]⁺;
- 2-(2-(2-(2-chlorophenyl)propan-2-yl)-1-(3'-(ethylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; MS (ES): 523.3 [M+H]⁺;
- 2-(2-(2-(2,3-dichlorophenyl)propan-2-yl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; MS (ES): 543.3 [M+H]⁺;
- 30 • 2-(2-(2-(2,3-dichlorophenyl)propan-2-yl)-1-(3'-(ethylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; MS (ES): 557.3 [M+H]⁺;

- 2-(2-(2,3-dichlorobenzyl)-1-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; MS (ES): 529.3 [M+H]⁺;
- 2-(2-(2,3-dichlorobenzyl)-1-(3'-(ethylsulfonyl)-3-methylbiphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; MS (ES): 543.3 [M+H]⁺;
- 5 • 2-(1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2,3-dichlorobenzyl)-1H-imidazol-4-yl)propan-2-ol; MS (ES): 549.3 [M+H]⁺;
- 2-(1-(3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl)-2-(2,3-dichlorobenzyl)-1H-imidazol-4-yl)propan-2-ol; MS (ES): 565.2 [M+H]⁺;
- 2-(2-(2,3-dichlorobenzyl)-1-(3-fluoro-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; MS (ES): 533.0 [M+H]⁺;
- 10 • 2-(2-(2,3-dichlorobenzyl)-1-(3'-(ethylsulfonyl)-3-fluorobiphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; MS (ES): 547.3 [M+H]⁺;
- 2-(1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2-(2,3-dichlorophenyl)propan-2-yl)-1H-imidazol-4-yl)propan-2-ol; MS (ES): 579.3 [M+H]⁺;
- 15 • 2-(1-(3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl)-2-(2-(2,3-dichlorophenyl)propan-2-yl)-1H-imidazol-4-yl)propan-2-ol; MS (ES): 593.3, 595.3 [M+H]⁺;
- 2-(2-(5-chloro-2-fluorobenzyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; MS (ES): 499.3, [M+H]⁺; 521.3 [M+Na]⁺;
- 2-(2-(5-chloro-2-fluorobenzyl)-1-(3'-(ethylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; MS (ES): 535.0 [M+Na]⁺;
- 20 • 2-(1-(2'-chloro-3-methyl-5'-(methylsulfonyl)biphenyl-4-yl)-2-(2,3-dichlorobenzyl)-1H-imidazol-4-yl)propan-2-ol; MS(ES): 563, 565 [M+H]⁺.

The following compounds were made in a similar manner, except the boronic acid or boronate reagents for the Suzuki coupling were made as described in Examples 23-29.

- 25 • 2-{1-[3-chloro-4'-(methyloxy)-3'-(methylsulfonyl)biphenyl-4-yl]-2-[(2,3-dichlorophenyl)methyl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 579, 581, 583 each [M+H]⁺.
- 2-{2-[(2,3-dichlorophenyl)methyl]-1-[3-methyl-4'-(methyloxy)-3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 559, 561 [each [M+H]⁺].
- 2-(1-(2'-chloro-3-methyl-5'-(methylsulfonyl)biphenyl-4-yl)-2-(2,3-dichlorobenzyl)-1H-imidazol-4-yl)propan-2-ol; MS(ES): 563, 565 [M+H]⁺
- 30

The following compounds were made in similar manner, except appropriate propionitrile was used as starting material, instead of benzyl cyanide.




- 2-{2-[2-(2-chlorophenyl)ethyl]-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 495.4 [M+H]⁺;
- 2-{2-[2-(2-chlorophenyl)ethyl]-1-[3'-(ethylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 509.3 [M+H]⁺;
- 2-{2-[2-(2-chlorophenyl)ethyl]-1-[4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 525.3 [M+H]⁺.

Example 12

1,1,1-trifluoro-2-(1-(3'-(methylsulfonyl)biphenyl-4-yl)-2-(2-(trifluoromethyl)phenyl)-1H-imidazol-4-yl)propan-2-ol

Example 12a


Preparation of 2-[1-(4-bromophenyl)-2-(2-trifluoromethyl-phenyl)-1H-imidazol-4-yl]-1,1,1-trifluoro-propan-2-ol



To a solution of 1-[1-(4-bromophenyl)-2-(2-trifluoromethyl-phenyl)-1H-imidazol-4-yl]-ethanone (0.24 g, 0.59 mmol) and trifluoromethyl-trimethylsilane (0.10 mL, 0.68 mmol) in toluene (3 mL, anhyd) at 0°C was added a 1.0M solution of tetrabutylammonium fluoride (TBAF) in THF (0.12 mL, 20 mol%, dried over 4Å molecular sieves). After 24 h the reaction mixture was charged with additional trifluoromethyl-trimethylsilane (0.10 mL) and 1.0M TBAF (0.12 mL) and then heated at 50°C. After 48 h (total) the reaction mixture was allowed to cool to ambient temperature, quenched with H₂O and extracted with DCM (2 x 30 mL). The combined extracts were dried (Na₂SO₄), concentrated and purified by chromatography (silica, EtOAc/Hex, 20:80 to 60:40) to yield the title compound (0.11 g, 39%) as a white solid. ¹H-NMR (DCM-d₂): δ 7.76 (d, 1H), 7.50-7.60 (m, 2H), 7.46 (d, 2H), 7.27 (m, 2H), 7.02 (d, 2H), 4.36 (s, 1H), 1.72 (s, 3H).


Example 12b

Preparation of 1,1,1-trifluoro-2-(1-(3'-(methanesulfonyl)biphenyl-4-yl)-2-(2-(trifluoromethyl)phenyl)-1H-imidazol-4-yl)propan-2-ol



A mixture of 2-[1-(4-bromophenyl)-2-(2-(trifluoromethyl)phenyl)-1H-imidazol-4-yl]-1,1,1-trifluoro-2-propanol (0.11 g, 0.23 mmol), 3-methanesulfonylphenylboronic acid (54 mg, 0.27 mmol), K₂CO₃ (95 mg, 0.69 mmol), Cl₂Pd(dppf)-DCM (15 mg, 8 mol%) and H₂O (0.25 mL) in DME (2.5 mL) was sparged with Argon for 5 min and then heated at 80°C as a sealed flask. After 90 min the reaction mixture was allowed to cool to ambient temperature, filtered (Celite™) and the filter agent rinsed with EtOAc. The combined filtrates were concentrated under reduced pressure and purified by chromatography (silica, EtOAc/Hex, 35:65 to 80:20) to give the title compound (116 mg, 89%) as a white solid. ¹H-NMR (DCM-*d*₂): δ 8.09 (m, 1H), 7.91 (m, 1H), 7.84 (m, 1H), 7.78 (d, 1H), 7.66 (m, 1H), 7.59 (d, 2H), 7.49-7.57 (m, 2H), 7.33 (m, 2H), 7.25 (d, 2H), 4.40 (s, 1H), 3.06 (s, 3H), 1.75 (s, 3H); MS(ES): 555 [M+H]⁺.

Scheme 10



(a) NaHCO₃, 2-PrOH, μ W 180°C; (b) LiBH₄, 10-20 mol% B(OMe)₃, THF; (c) ArB(OH)₂, Cl₂Pd(dppf), K₂CO₃, H₂O/DME, μ W 120°C.

In general, compounds of formula **01017** can be prepared as depicted in Scheme 10. First an amidine (**01013**) and a bromoketone (**01014**) can react under basic conditions and at elevated temperature to give the corresponding imidazole (**01015**). This intermediate ester **01015** can be reduced under standard conditions, such as with lithium borohydride and, optionally, catalytic trimethyl borate,


to afford the corresponding carbinol (**01016**). Last intermediate **01016** can undergo cross-couplings reactions, such as a Suzuki-Miyaura reaction with boronic acids (esters), to yield the target imidazole **01017**.

Example 13

5 2-[2-(2,6-dichlorophenyl)-1-(3'-methanesulfonyl-biphenyl-4-yl)-1H-imidazol-4-yl]-2-methyl-propan-1-ol

Example 13a

Preparation of 2-[1-(4-bromophenyl)-2-(2,6-dichlorophenyl)-1H-imidazol-4-yl]-2-methyl-propionic acid methyl ester




A mixture of N-(4-bromophenyl)-2,6-dichlorobenzamidine (1.03 g, 3.0 mmol), methyl 4-bromo-2,2-dimethylacetoacetate (1.78 g, 8.0 mmol) and NaHCO₃ (0.76 g, 9.0 mmol) in isopropanol (12 mL) was heated in a microwave unit (Biotage Initiator™) at 170°C for 25 min. The resulting mixture was decanted and the solids rinsed with EtOAc. The combined filtrates were concentrated and purified by chromatography (silica, EtOAc/Hex, 15:85 to 50:50) to give the title compound (0.80 g, 57%) as a pale yellow solid. ¹H-NMR (DCM-*d*₂): δ 7.44 (d, 2H), 7.27-7.34 (m, 3H), 7.09 (m, 2H), 3.67 (s, 3H), 1.61 (s, 6H); MS(ES): 467, 469, 471 [M+H]⁺.

15

Example 13b

Preparation of 2-[1-(4-bromophenyl)-2-(2,6-dichlorophenyl)-1H-imidazol-4-yl]-2-methyl-propan-1-ol




To a stirred suspension of 2-[1-(4-bromophenyl)-2-(2,6-dichlorophenyl)-1H-imidazol-4-yl]-2-methyl-propionic acid methyl ester (0.65 g, 1.4 mmol) in Et₂O (3.5 mL, anhyd) added dropwise 2.0M solution of lithium borohydride in THF (1.4 mL, 2.8 mmol) and then added trimethyl borate (22 μL, 15 mol%). After 3 h the reaction mixture was quenched with 1N NaOH (4 mL) and extracted with EtOAc (2 x 20 mL). The combined extracts were washed with satd NH₄Cl and brine, dried (Na₂SO₄), concentrated and purified by chromatography (silica, EtOAc/Hex, 15:85 to 50:50) to yield a white solid (0.21 g) consisting of a 2:1 mixture of the title compound, MS(ES): 439, 441, 443 [M+H]⁺, and 2-[2-(2,6-dichlorophenyl)-1-phenyl-1H-imidazol-4-yl]-2-methyl-propan-1-ol, MS(ES): 361, 363 [M+H]⁺.

25

The latter resulted from over-reduction of the title compound under the given conditions. The isolated product was used in the next step without further purification.


Example 13c

Preparation of 2-[2-(2,6-dichlorophenyl)-1-(3'-methanesulfonyl-biphenyl-4-yl)-1H-imidazol-4-yl]-2-methyl-propan-1-ol



A stirred mixture of 2-[1-(4-bromophenyl)-2-(2,6-dichlorophenyl)-1H-imidazol-4-yl]-2-methyl-propan-1-ol (202 mg, 0.46 mmol), 3-methanesulfonyl-phenylboronic acid (92 mg, 0.46 mmol), potassium carbonate (0.19 g, 1.4 mmol), $\text{Cl}_2\text{Pd}(\text{dppf})\cdot\text{DCM}$ (19 mg, 5 mol%) and H_2O (0.25 mL) in DME (2.5 mL) was sparged with Argon for 5 min and then heated at 60°C as a sealed flask. After 30 min the reaction mixture was concentrated under reduced pressure and purified by chromatography (silica, EtOAc/Hex, 40:60 to 75:25) to give the title compound (35 mg) as a white solid. $^1\text{H-NMR}$ (DCM-d_2): δ 8.10 (m, 1H), 7.91 (d, 1H), 7.85 (d, 1H), 7.66 (m, 1H), 7.60 (d, 2H), 7.29-7.37 (m, 5H), 7.10 (s, 1H), 3.79 (br s, 1H), 3.62 (s, 2H), 3.06 (s, 3H), 1.36 (s, 6H); MS(ES): 515, 517 $[\text{M}+\text{H}]^+$.

Scheme 11.



(a) aq NaOH, MeOH; (b) $(\text{COCl})_2$, DCM, DMF; (c) $\text{R}_3\text{R}_4\text{NH}$, CHCl_3 , DIEA; (d) PdCl_2dppf , K_2CO_3 , DME, H_2O

In general, amide containing compounds of formula (0115) can be synthesized following the methodology shown in Scheme 11. The ester group in compound (0111) can be hydrolyzed under aqueous basic conditions to give the carboxylic acid (0112). Acid (0112) can be converted to an acid chloride (0113) using known procedures followed by reaction with a primary or secondary amine to


provide amide intermediate (0114). The amide intermediate (0114) can undergo cross coupling transformations to afford the corresponding 2-biaryl-imidazole (0115).

Example 14

*2-(2-Chloro-phenyl)-1-(3'-methanesulfonyl-biphenyl-4-yl)-1H-imidazole-4-carboxylic acid
(2,2,2-trifluoro-ethyl)-amide*

Example 14a


Preparation of 1-(4-Bromo-phenyl)-2-(2-chloro-phenyl)-1H-imidazole-4-carboxylic acid



To a 100 mL round bottom flask was added 1-(4-Bromo-phenyl)-2-(2-chloro-phenyl)-1H-imidazole-4-carboxylic acid ethyl ester (1.03 g, 2.54 mmol), MeOH (20 mL), and 1N aq NaOH (10 mL). The reaction solution was allowed to stir at 45 °C for 1 hr prior to analysis by LCMS. The reaction solution was diluted with EtOAc (150 mL), poured into a separatory funnel and the organic phase was partitioned. The aqueous phase was neutralized by the addition of aq 1 N HCl and extracted with EtOAc (70 mL x 2). The combined organic phase was dried over Na₂SO₄, filtered into a round bottom flask and concentrated on the Rotavapor. The crude residue was chromatographed thru a 25 g SiO₂ column using a mobile phase gradient of 100 % Hx to 85 % EtOAc to afford 845 mg (88 % yield) carboxylic acid intermediate. MS (ES): 378 [M+H]⁺.

Example 14b

Preparation of 1-(4-Bromo-phenyl)-2-(2-chloro-phenyl)-1H-imidazole-4-carboxylic acid (2,2,2-trifluoro-ethyl)-amide




To a dry, N₂ purged 100 mL round bottom flask was added 1-(4-Bromo-phenyl)-2-(2-chloro-phenyl)-1H-imidazole-4-carboxylic acid (845 mg, 2.24 mmol), and (10 mL) anhydrous DCM. The solution was cooled to 0 °C prior to addition of oxalyl chloride (590 μL, 6.7 mmol) and several drops anhydrous DMF. The reaction solution was allowed to stir warming to rt over 2 hrs. The solvent and

excess reagent was removed *in vacuo*. To the crude acid chloride residue was added anhydrous CHCl_3 (15 mL), 2,2,2-trifluoroethylamine (700 mg, 6.70 mmol) and DIEA (1.3 mL, 7.46 mmol). The reaction solution was allowed to stir at 40 °C for approx 1 hr. The reaction solution was diluted with DCM (70 mL) and transferred to a separatory funnel. The solution was washed with aq NH_4Cl (50 mL x 2) and with aq NaCl (50 mL). The organic phase was dried over Na_2SO_4 , filtered, concentrated on the Rotavapor and chromatographed through a 25 g SiO_2 column using a mobile phase gradient of 100 % Hx to 70 % EtOAc to afford 840 mg (82 % yield) of amide product. MS (ES): 461 $[\text{M}+\text{H}]^+$.

Example 14c

Preparation of 2-(2-Chloro-phenyl)-1-(3'-methanesulfonyl-biphenyl-4-yl)-1H-imidazole-4-carboxylic acid (2,2,2-trifluoro-ethyl)-amide



To a 50 mL round bottom flask attached with Vigreux column and magnetic stir bar was added 1-(4-Bromo-phenyl)-2-(2-chloro-phenyl)-1H-imidazole-4-carboxylic acid (2,2,2-trifluoro-ethyl)-amide (249 mg, 540 μmol), 3-methylsulfonylphenyl boronic acid (130 mg, 650 μmol), PdCl_2dppf (40 mg, 10 mol %), K_2CO_3 (260 mg, 1.90 mmol), 1,2-dimethoxyethane (14 mL) and H_2O (2 mL). The reaction solution was allowed to stir at 75 °C for 2 hrs. The reaction solution was diluted with EtOAc (150 mL) and filtered through a Celite padded Buchner funnel to remove spent Pd. The filtrate was transferred to a separatory funnel and washed with aq NH_4Cl (100 mL) and aq NaCl (100 mL). The organic phase was dried over Na_2SO_4 , filtered, concentrated on the Rotavapor and chromatographed through a 25 g SiO_2 column using a mobile phase gradient of 2% EtOAc to 90 % EtOAc to afford 212 mg (74 % yield) of the title compound. MS (ES): 534 $[\text{M}+\text{H}]^+$, 557 $[\text{M}+\text{Na}]^+$. ^1H NMR (400 MHz, CDCl_3): δ 8.13 (s, 1H), 7.99 (s, 1H), 7.92 (d, 1H), 7.65 (t, 1H), 7.57 (d, 2H), 7.52 (d, 2H), 7.41-7.34 (m, 3H), 7.24 (d, 2H), 4.11 (q, 2H), 3.08 (s, 3H); ^{19}F NMR (400 MHz, CDCl_3) δ -72.4 ppm.


The following compounds were prepared in a manner similar to that described in experimental procedure Example 14:

- 2-(2-isopropylphenyl)-1-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-N-(2,2,2-trifluoroethyl)-1H-imidazole-4-carboxamide; MS (ES): 556.3 $[\text{M}+\text{H}]^+$.
- 2-(2,6-dichlorophenyl)-1-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-N-(2,2,2-trifluoroethyl)-1H-imidazole-4-carboxamide; MS (ES): 582.2, 584.2 $[\text{M}+\text{H}]^+$.

- 2-(2,6-dichlorophenyl)-1-(3'-(ethylsulfonyl)-3-methylbiphenyl-4-yl)-N-(2,2,2-trifluoroethyl)-1H-imidazole-4-carboxamide; MS (ES): 596.3, 598.3 $[M+H]^+$.
- 2-(2,6-dichlorophenyl)-N,N-diethyl-1-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazole-4-carboxamide; MS (ES): 556.3, 558.3 $[M+H]^+$.
- 5 • 2-(2,6-dichlorophenyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-N-(2,2,2-trifluoroethyl)-1H-imidazole-4-carboxamide; MS (ES): 568.2, 570.2 $[M+H]^+$.
- 2-(2,6-dichlorophenyl)-N-(2-fluoroethyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazole-4-carboxamide; MS (ES): 532, 534 $[M+H]^+$.
- 2-(2,6-dichlorophenyl)-N-ethyl-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazole-4-carboxamide; MS (ES): 514.3, 516.3 $[M+H]^+$.
- 10 • (2-(2,6-dichlorophenyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)(4-methylpiperazin-1-yl)methanone; MS (ES): 569.2, 571.3 $[M+H]^+$.
- 2-(2,6-dichlorophenyl)-N-(2-hydroxyethyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazole-4-carboxamide; MS (ES): 530.3, 530.3 $[M+H]^+$.
- 15 • 1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-N-(2,2,2-trifluoroethyl)-1H-imidazole-4-carboxamide; MS (ES): 602.0, 604.0, 606.3 $[M+H]^+$.
- 1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-N-(2-fluoroethyl)-1H-imidazole-4-carboxamide; MS (ES): 566.3, 568.3, 570.2 $[M+H]^+$.
- 2-(1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-1H-imidazole-4-carboxamido)acetic acid; MS (ES): 578.3, 580.3, 582.3 $[M+H]^+$.
- 20 • 1-(3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-N-(2,2,2-trifluoroethyl)-1H-imidazole-4-carboxamide; MS (ES): 616.0, 618.0, 620.0 $[M+H]^+$.
- 1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-N-(1-hydroxy-2-methylpropan-2-yl)-1H-imidazole-4-carboxamide; MS (ES): 592.0, 594.0, 596.2 $[M+H]^+$.
- 25 • 1-(3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-N-(1-hydroxy-2-methylpropan-2-yl)-1H-imidazole-4-carboxamide; MS (ES): 606.3, 608.3, 610.0 $[M+H]^+$.
- N-tert-butyl-1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-1H-imidazole-4-carboxamide; MS (ES): 576.3, 578.3, 580.3 $[M+H]^+$.
- 1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-N-(2-hydroxyethyl)-1H-imidazole-4-carboxamide; MS (ES): 564.2, 566.2, 568.3 $[M+H]^+$.
- 30 • 1-(3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-N-(2-hydroxyethyl)-1H-imidazole-4-carboxamide; MS (ES): 578.3, 580.3, 582.2 $[M+H]^+$.


- 1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-N-(2-hydroxy-2-methylpropyl)-1H-imidazole-4-carboxamide; MS (ES): 592.3, 594.3, 596.0 [M+H]⁺
- 1-(3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-N-(2-hydroxy-2-methylpropyl)-1H-imidazole-4-carboxamide; MS (ES): 606.2, 608.2, 610.2 [M+H]⁺
- 5 • 2-(2,6-dichlorophenyl)-N-(1-hydroxy-2-methylpropan-2-yl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazole-4-carboxamide; MS (ES): 558.3, 560.3 [M+H]⁺
- 2-(2,6-dichlorophenyl)-1-(3'-(ethylsulfonyl)biphenyl-4-yl)-N-(1-hydroxy-2-methylpropan-2-yl)-1H-imidazole-4-carboxamide; MS (ES): 572.3, 574.3 [M+H]⁺
- N-tert-butyl-2-(2,6-dichlorophenyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazole-4-
- 10 carboxamide; MS (ES): 542.3, 544.3 [M+H]⁺

Scheme 12



In general, carboxylic acid amides of formula (0125) can be converted to the thioamide derivative (0126) as shown in Scheme 12. The functional transformation can be carried out using a suitable thiation reagent, such as Lawesson's reagents or related compound.

Example 15



Preparation of 2-(2-Chloro-phenyl)-1-(3'-methanesulfonyl-biphenyl-4-yl)-1H-imidazole-4-carbothioic acid (2,2,2-trifluoro-ethyl)-amide


20 To a 25 mL round bottom flask attached with Vigreux column and magnetic stir bar was added 2-(2-Chloro-phenyl)-1-(3'-methanesulfonyl-biphenyl-4-yl)-1H-imidazole-4-carboxylic acid (2,2,2-trifluoro-ethyl)-amide (65 mg, 127 μ mol), Lawesson's Reagent (230 mg, 570 μ mol), and anhydrous toluene (6 mL). The reaction solution was allowed to stir at reflux 5 hrs prior to TLC analysis showing

full transformation of starting material. The reaction solution was allowed to cool to room temperature prior to addition of a 1:1 mixture of benzene and Et₂O. The resulting precipitate was removed by vacuum filtration through a Buchner funnel. The filtrate was concentrated on the Rotavapor and the crude residue was chromatographed through a 12 g SiO₂ column using a mobile phase gradient of 100 % Hx to 50 % EtOAc to afford 55 mg (79 % yield) of the title compound. ¹H NMR (400 MHz, CDCl₃) δ 9.21 (t, 1H), 8.11 (s, 1H), 8.04 (t, 1H), 7.85 (d, 1H), 7.75 (d, 1H), 7.59 (t, 1H), 7.53-7.46 (m, 4H), 7.34-7.311 (m, 3H), 7.18 (d, 2H), 4.56 (m, 2H), 3.02 (s, 3H); ¹⁹F NMR (400 MHz, CDCl₃) δ -70.8 ppm; MS (ES): 550.3, 552.3 [M+H]⁺.

The following compounds were prepared in a manner similar to that described in experimental procedure Example 15:

- 2-(2-isopropylphenyl)-1-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-N-(2,2,2-trifluoroethyl)-1H-imidazole-4-carbothioamide; MS (ES): 572.2 [M+H]⁺
- 1-(4-bromophenyl)-2-(2-chlorophenyl)-N-(2,2,2-trifluoroethyl)-1H-imidazole-4-carbothioamide; MS (ES): 474.0, 476.0 [M+H]⁺

Scheme 13



(a) (i) Me₃Al, toluene, MeNHCH₂CH₂NHMe, reflux, (ii) aq HCl; (b) PdCl₂dppf, K₂CO₃, DME / H₂O; (c) H₂NOH-HCl, 2:1 MeOH/H₂O, NaOAc, reflux


In general, oxime containing compounds of formula (**0139**) can be prepared as shown in Scheme 13. The imidazole-ester intermediate (**0131**) can be converted to the methylketone substituted compound (**0137**) using trimethylaluminium and a diamine chelant, such as N,N-dimethylethylenediamine. Compound (**0137**) can undergo cross-coupling to install the D-ring to yield final compound (**0138**). The oxime substituted product (**0139**) can be prepared by reaction with hydroxylamine under known conditions.

Example 16

1-[2-(2-Chloro-phenyl)-1-(3'-methanesulfonyl-biphenyl-4-yl)-1H-imidazol-4-yl]-ethanone oxime

Example 16a


Preparation of 1-[1-(4-Bromo-phenyl)-2-(2-chloro-phenyl)-1H-imidazol-4-yl]-ethanone



5 To a dried, N₂ purged 100 mL RB flask, attached with Vigreux column was added anh. toluene (20 mL) and N,N-dimethylethylenediamine (2.2 mL, 1.1 equiv). The solution was cooled to 0 °C prior to dropwise addition of a 2.0 M solution of Me₃Al in hexanes (28 mL, 3.1 equiv). The reaction solution was stirred at room temperature for 40 min prior to the addition of 1-(4-Bromo-phenyl)-2-(2-chloro-phenyl)-1H-imidazole-4-carboxylic acid ethyl ester (7.41 g, 18.3 mmol) in a solution of toluene (10 mL). The reaction solution was allowed to stir at reflux for 2 hrs. The reaction solution was cooled to room temperature and quenched by the addition of 1N aq HCl (10 mL). The reaction solution was dilute with EtOAc, partitioned, washed with aq NaCl, dried over Na₂SO₄, filtered, concentrated on the Rotavapor and chromatographed through an 80 g SiO₂ column using a mobile phase gradient of 100 % Hx to 75 % EtOAc to afford 3.10 g (45 % yield) of the methyl ketone intermediate. MS (ES): 376
15 [M+H]⁺.

Example 16b

Preparation of 1-[2-(2-Chloro-phenyl)-1-(3'-methanesulfonyl-biphenyl-4-yl)-1H-imidazol-4-yl]-ethanone



20 To a 50 mL round bottom flask attached with Vigreux column and magnetic stir bar was added 1-[1-(4-Bromo-phenyl)-2-(2-chloro-phenyl)-1H-imidazol-4-yl]-ethanone (270 mg, 719 μmol), 3-methylsulfonylphenyl boronic acid (220 mg, 1.08 mmol), PdCl₂dppf (55 mg, 10 mol %), K₂CO₃ (300 mg, 2.16 mmol), 1,2-dimethoxyethane (11 mL) and H₂O (1 mL). The reaction solution was allowed to stir at 70 °C for 14 hrs. The reaction solution was diluted with EtOAc (150 mL) and filtered through a
25 Celite padded Buchner funnel to remove spent Pd. The filtrate was transferred to a separatory funnel and washed with aq NH₄Cl (100 mL) and aq NaCl (100 mL). The organic phase was dried over Na₂SO₄, filtered, concentrated on the Rotavapor and chromatographed through a 25 g SiO₂ column


using a mobile phase gradient of 100% Hx to 75 % EtOAc to afford 170 mg (52 % yield) of product. MS (ES): 451 [M+H]⁺, 473 [M+Na]⁺.

The following compounds were synthesized in a manner similar to the above procedure:

- 1-{1-[3'-(methylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}ethanone; MS (ES): 485.4 [M+H]⁺
- 1-(2-(2,6-dichlorophenyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)ethanone; MS (ES): 485.3, 487.3 [M+H]⁺

Example 16c

Preparation of 1-[2-(2-Chloro-phenyl)-1-(3'-methanesulfonyl-biphenyl-4-yl)-1H-imidazol-4-yl]-ethanone oxime




To a 50 mL RB flask attached with Vigreux column was added compound 1-[2-(2-Chloro-phenyl)-1-(3'-methanesulfonyl-biphenyl-4-yl)-1H-imidazol-4-yl]-ethanone (162 mg, 359 μ mol), hydroxylamine-HCl (200 mg, 2.87 mmol), sodium acetate (238 mg, 2.90 mmol), MeOH (12 mL) and H₂O (6 mL). The reaction solution was allowed to stir at reflux for 2 hrs prior to TLC analysis. The reaction solution was diluted with EtOAc (150 mL), washed with aq. NaCl (100 mL), partitioned, dried over Na₂SO₄, filtered, and concentrated on the Rotavapor. The crude product was chromatographed through a 25 g SiO₂ column using a mobile phase gradient of 100 % Hx to 80 % EtOAc to afford 85 mg (51 % yield) of the title compound as a mixture of *E* and *Z* isomers. The isomers were separated by chromatography with a Reverse Phase HPLC using a C₁₈ Preparative column. Characterization for one single oxime isomer: ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.86 (s, 1H), 8.15 (t, 1H), 8.02 (d, 1H), 7.90 (d, 1H), 7.85 (s, 1H), 7.80 (d, 2H), 7.72 (t, 1H), 7.64 (d, 1H), 7.47-7.44 (m, 3H), 7.35 (d, 2H); MS (ES): 466.1 [M+H]⁺.

The following compounds were prepared in a manner similar to that described in experimental procedure Example 16:

- 1-{1-[3'-(methylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}ethanone oxime; MS (ES): 500.3 [M+H]⁺
- 1-{2-(2-chlorophenyl)-1-[3'-(ethylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}ethanone oxime; MS (ES): 480.0, 482.0 [M+H]⁺

- 1-{2-(2-chlorophenyl)-1-[3'-(ethylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}ethanone oxime; MS (ES): 480.0, 482.0 $[M+H]^+$
- 1-{2-(2-chlorophenyl)-1-[3'-(1-methylethyl)biphenyl-4-yl]-1H-imidazol-4-yl}ethanone oxime; MS (ES): 430.0, 432.0 $[M+H]^+$
- 5 • 1-{2-(2-chlorophenyl)-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}ethanone oxime; MS (ES): 466.0, 468.0 $[M+H]^+$
- 1-{2-(2-chlorophenyl)-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}ethanone oxime; MS (ES): 466.0, 468.0 $[M+H]^+$
- 1-{2-(2-chlorophenyl)-1-[4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}ethanone oxime; MS (ES): 495.3, 497.3 $[M+H]^+$
- 10 • (E)-1-(2-(2,6-dichlorophenyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)ethanone oxime; MS (ES): 500.4, 502.3 $[M+H]^+$
- (E)-1-(2-(2-ethylphenyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)ethanone oxime; MS (ES): 460.3 $[M+H]^+$
- 15 • (E)-1-(2-(2-isopropylphenyl)-1-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)ethanone oxime; MS (ES): 488.3 $[M+H]^+$


Scheme 14



(a) HCl, dioxane, Δ or HOAc, toluene, Δ

In general, carbinol substituted imidazole compounds of formula (01419) can be dehydrated using known methods to provide the corresponding alkene (01420) (Scheme 14). The dehydration reaction can be carried out using several different acid catalysts, such as HCl or acetic acid, in a heated and anhydrous solvent system.

Example 17




Preparation of 2-(2,6-Dichloro-phenyl)-4-isopropenyl-1-(3'-methanesulfonyl-biphenyl-4-yl)-1H-imidazole

To a 25 mL round bottom flask attached with Vigreux column was added 2-[2-(2,6-Dichloro-phenyl)-1-(3'-methanesulfonyl-biphenyl-4-yl)-1H-imidazol-4-yl]-propan-2-ol (383 mg, 764 μ mol), anhydrous toluene (8 mL) and acetic acid (3.5 mL). The reaction solution was allowed to stir at reflux for 1.5 hr. The cooled reaction solution was concentrated *in vacuo*, and the residue was taken into EtOAc (200 mL) and washed with aq. NaHCO₃ (100 mL x 2) and aq. NaCl (100 mL). The organic phase was partitioned, dried over Na₂SO₄, filtered and concentrated *in vacuo*. The crude product was chromatographed through a 25 g SiO₂ column using a mobile gradient of 100 % Hx to 80 % EtOAc to provide 288 mg (78 % yield) of title compound. ¹H NMR (400 MHz, DMSO-*d*₆): δ 8.15 (br s, 1H), 8.08-7.97 (m, 2H), 7.92 (d, 1H), 7.89 (d, 2H), 7.75 (t, 1H), 7.64-7.57 (m, 3H), 7.43 (d, 2H), 5.77 (br s, 1H), 5.10 (br s, 1H), 3.29 (s, 3H), 2.11 (s, 3H); MS (ES): 483.1, 485.1 [M+H]⁺.

The following compound was made according to this scheme:

- 2-(2,3-dichlorophenyl)-1-(3'-(ethylsulfonyl)biphenyl-4-yl)-4-(prop-1-en-2-yl)-1H-imidazole;

Scheme 15



(a) CuCN, DMF, reflux; (b) (i) Me₃Al, toluene, (ii) reflux 1 d; (c) (i) EtOH, NaHCO₃, (ii) HOAc, reflux; (d) 4 equiv MeMgBr; (e) PdCl₂dppf, K₂CO₃, DME / H₂O

In general, compounds of formula (01515), with alkyl A-ring substituents larger than Me or Et, can be synthesized following the sequence shown in Scheme 15. By example, the methodology in Scheme 15 can afford imidazole compounds with 2-iso-propyl groups on the phenyl A-ring. In general, the methodology can also afford compounds with either mono- or disubstituted phenyl A-rings. An arylbromide (01510) can be converted to an aryl nitrile (01511) by reaction with copper(I) cyanide. The aryl nitrile can be reacted with aniline in the presence of a Lewis acid, such as trimethylaluminum, to

give the corresponding amidine (**01512**). Amidine (**01512**) can be reacted with ethyl bromopyruvate in the presence of sodium bicarbonate, followed by dehydration to afford the cyclized imidazole intermediate (**01513**). Carbinol formation to provide (**01514**) and Suzuki cross coupling can be used to afford compound (**01515**) in a manner similar to that described in Scheme 8.

- 5 The following compound was made according to this scheme: 2-(2,3-dichlorophenyl)-1-(3'-(ethylsulfonyl)biphenyl-4-yl)-4-(prop-1-en-2-yl)-1H-imidazole; MS(ES): 497, 499 [M+H]⁺


Example 18

2-[2-(2-Isopropyl-phenyl)-1-(3'-methanesulfonyl-biphenyl-4-yl)-1H-imidazol-4-yl]-propan-2-ol

10

Example 18a


Preparation of 2-isopropylbenzonitrile



- To a dry, N₂ purged 100 mL round bottom flask attached with condenser was added 1-bromo-2-isopropylbenzene (10.0 g, 50.2 mmol), anhydrous DMF (26 mL), and CuCN (5.85 g, 65.3 mmol).
 15 The reaction slurry was stirred at reflux under N₂ for 4 hrs prior to analysis by GCMS. The reaction mixture was cooled to room temperature and poured into an ice / aq NH₄Cl solution. The resulting precipitates were removed by vacuum filtration through a Buchner funnel. The aqueous filtrate was extracted with EtOAc (150 mL x 3). The combined organic phase was dried over Na₂SO₄, filtered and concentrated on the Rotavapor to afford 5.97 g (82 % yield) of 2-isopropylbenzonitrile. The product
 20 was used in the next step without further purification. GCMS m/z = 145 [M⁺].

Example 18b

Preparation of N-(4-Bromo-phenyl)-2-isopropyl-benzamidine




- To a dry, N₂ purged 100 mL round bottom flask was added 4-bromo-aniline (2.73g, 15.9
 25 mmol) and anhydrous toluene (40 mL). To the solution at 0 °C was added, dropwise, a 2.0 M solution of Me₃Al in hexanes (9.4 mL). The solution was allowed to stir, warming to room temperature for approximately 1 hr. To the reaction solution was added 2-isopropylbenzonitrile (3.00 g, 20.7 mmol) in a toluene solution (15 mL). The reaction solution was allowed to stir at 80 °C for approximately 24 hrs.

The reaction solution was allowed to cool to room temperature prior to quenching by pouring the reaction solution into an Erlenmeyer containing a 2:1 CHCl_3 / MeOH solution and 100 g of silica. The slurry was allowed to stir 30 min prior to filtration into a Buchner funnel under vacuum. The filtrate was concentrated on the Rotavapor and the resulting residue was reprecipitated using a 10:1 Hx / Et_2O mixture. The resulting white precipitates were isolated by vacuum filtration to afford 1.98 g (40 % yield) of amidine product. GCMS $m/z = 317$ [M^+].

Example 18c


Preparation of 1-(4-Bromo-phenyl)-2-(2-isopropyl-phenyl)-1H-imidazole-4-carboxylic acid ethyl ester



To a 250 mL round bottom flask attached with condenser was added N-(4-Bromo-phenyl)-2-isopropyl-benzimidine (1.94 g, 6.11 mmol), ethyl bromopyruvate (1.54 mL, 12.2 mmol), sodium bicarbonate (1.02 g, 12.2 mmol), and iso-propanol (45 mL). The reaction slurry was allowed to stir at 80 °C for approximately 2 hrs. The reaction solution was decanted into a clean round bottom flask and concentrated *in vacuo*. The resulting residue was dissolved in acetic acid (25 mL), and the solution was allowed to stir at reflux for approximately 2 hrs. The solution was concentrated *in vacuo*, and the product residue was taken into EtOAc (200 mL) and washed with aq NaCl (200 mL x 2) and aq NaHCO_3 (100 mL). The organic phase was partitioned, dried over Na_2SO_4 , filtered, concentrated, and chromatographed through a 40 g SiO_2 column using a mobile gradient of 100 % Hx to 70 % EtOAc to afford 1.90 g (75 % yield) of title compound. MS (ES): 414.0 [$\text{M}+\text{H}$] $^+$.

Example 18d

Preparation of 2-[1-(4-Bromo-phenyl)-2-(2-isopropyl-phenyl)-1H-imidazol-4-yl]-propan-2-ol




To a dry, N_2 purged round bottom flask containing 1-(4-Bromo-phenyl)-2-(2-isopropyl-phenyl)-1H-imidazole-4-carboxylic acid ethyl ester (1.87 g, 4.52 mmol) was added anhydrous THF (55 mL). The solution was cooled to 0 °C prior to the addition of a 3.0 M solution of methylmagnesium

bromide (5.3 mL). The reaction was stirred 1 hr at room temperature prior to quenching with aq NH_4Cl (30 mL). The organic phase was partitioned, dried over Na_2SO_4 , filtered, and concentrated *in vacuo*. The crude product was chromatographed through a SiO_2 column using a mobile gradient to 100 % Hx to 80 % EtOAc to afford 1.19 g (66 % yield) of 369-45.

5

Example 18e

Preparation of 2-[2-(2-Isopropyl-phenyl)-1-(3'-methanesulfonyl-biphenyl-4-yl)-1H-imidazol-4-yl]-propan-2-ol




To a 50 mL round bottom flask attached with Vigreux column and magnetic stir bar was added
 10 2-[1-(4-Bromo-phenyl)-2-(2-isopropyl-phenyl)-1H-imidazol-4-yl]-propan-2-ol (297 mg, 743 μmol), 3-methylsulfonylphenyl boronic acid (193 mg, 968 μmol), PdCl_2dppf (60 mg, 10 mol %), K_2CO_3 (310 mg, 2.23 mmol), 1,2-dimethoxyethane (14 mL) and H_2O (4 mL). The reaction solution was allowed to stir at 80 $^\circ\text{C}$ for 2 hrs. The reaction solution was diluted with EtOAc (150 mL) and filtered through a Celite padded Buchner funnel to remove spent Pd. The filtrate was transferred to a separatory funnel
 15 and washed with aq NH_4Cl (100 mL) and aq NaCl (100 mL). The organic phase was dried over Na_2SO_4 , filtered, concentrated on the Rotavapor and chromatographed through a 25 g SiO_2 column using a mobile phase gradient of 100 % Hx to 90 % EtOAc to afford 303 mg (86 % yield) of the title compound. ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 8.14 (s, 1H), 8.02 (d, 1 H), 7.89 (d, 1H), 7.89 (d, 1H), 7.78 (d, 2H), 7.72 (t, 1H), 7.37-7.34 (m, 3H), 7.26 (d, 2H), 7.18-7.16 (m, 2H), 4.84 (s, 1H), 3.26 (s, 3H),
 20 2.85 (sept, 1H), 1.50 (6H), 0.96 (d, 6H); MS (ES): 475.4 $[\text{M}+\text{H}]^+$, 497.5 $[\text{M}+\text{Na}]^+$.

The following compounds were prepared in a manner similar to that described in experimental procedure Example 18:

- 2-(1-(3'-(ethylsulfonyl)biphenyl-4-yl)-2-(2-isopropylphenyl)-1H-imidazol-4-yl)propan-2-ol; MS (ES): 489.4 $[\text{M}+\text{H}]^+$, 511.4 $[\text{M}+\text{Na}]^+$
- 25 • 2-(1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2-isopropylphenyl)-1H-imidazol-4-yl)propan-2-ol; MS (ES): 509.3, 511.3 $[\text{M}+\text{H}]^+$
- 2-(2-(2-isopropylphenyl)-1-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol; MS (ES): 489.3, 511.4 $[\text{M}+\text{Na}]^+$

- 2-(1-(3'-(ethylsulfonyl)-3-methylbiphenyl-4-yl)-2-(2-isopropylphenyl)-1H-imidazol-4-yl)propan-2-ol; MS (ES): 503.4 [M+H]⁺

Scheme 16



(a) Suzuki Coupling; (b) 4 equiv MeMgBr, toluene

- 5 In general, compounds of formula (01618) can be synthesized as shown in Scheme 16. The method involves use of a methylester substituted D-ring compound, such as depicted with compound (01617). The D-ring ester (01617) can be converted to a dimethylcarbinol compound (01618) by reaction with a molar excess of methylmagnesium bromide. The transformation can be carried out when R₂ is either an ester group or a carbinol group. When R₂ is an ester group, four equivalents of
- 10 methylmagnesium bromide can be used to prepare the bis-carbinol version of compound (01618).


Example 19

2-{2-(2-Chloro-phenyl)-1-[3'-(1-hydroxy-1-methyl-ethyl)-biphenyl-4-yl]-1H-imidazol-4-yl}-propan-2-ol

Example 19a

Preparation of 4'-[2-(2-Chloro-phenyl)-4-(1-hydroxy-1-methyl-ethyl)-imidazol-1-yl]-biphenyl-3-carboxylic acid methyl ester

15




- To a 50 mL round bottom flask attached with Vigreux column and magnetic stir bar was added 2-[1-(4-Bromo-phenyl)-2-(2-chloro-phenyl)-1H-imidazol-4-yl]-propan-2-ol (275 mg, 702 μmol), 3-methoxycarbonylphenyl boronic acid (164 mg, 913 μmol), PdCl₂dppf (51 mg, 10 mol %), K₂CO₃ (290 mg, 2.11 mmol), 1,2-dimethoxyethane (13 mL) and H₂O (1.5 mL). The reaction solution was allowed to stir at 70 °C for approximately 2 hrs. The reaction solution was diluted with EtOAc (150 mL) and filtered through a Celite padded Buchner funnel to remove spent Pd. The filtrate was transferred to a separatory funnel and washed with aq NH₄Cl (100 mL) and aq NaCl (100 mL). The organic phase was dried over Na₂SO₄, filtered, concentrated on the Rotavapor and chromatographed through a 25 g SiO₂
- 20

column using a mobile phase gradient of 100 % Hx to 90 % EtOAc to afford 122 mg (39 % yield) of product. MS (ES): 446.3, 448.3 [M+H]⁺.

Example 19b

Preparation of Preparation of 2-{2-(2-Chloro-phenyl)-1-[3'-(1-hydroxy-1-methyl-ethyl)-biphenyl-4-yl]-1H-imidazol-4-yl}-propan-2-ol




To a dry, N₂ purged 50 mL round bottom flask was added 4'-[2-(2-Chloro-phenyl)-4-(1-hydroxy-1-methyl-ethyl)-imidazol-1-yl]-biphenyl-3-carboxylic acid methyl ester (100 mg, 225 μmol), anhydrous and toluene (7 mL). To the solution at 0 °C was added a 1.4 M solution of methylmagnesium bromide (640 μL). The reaction solution was allowed to stir at room temperature for approximately 1hr. The reaction was quenched by the addition of aq NH₄Cl (20 mL). The organic phase was partitioned, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. The crude residue was chromatographed through a 12 g SiO₂ column using a gradient of 100 % Hx to 100 % EtOAc to afford 45 mg (45 % yield) of title compound. ¹H NMR (400 MHz, CDCl₃): δ 7.70 (s, 1H), 7.55 (d, 2H), 7.48-7.38 (m, 5H), 7.35 (s, 1H), 7.33 (m, 1H), 7.18 (d, 2H), 7.16 (s, 1H), 1.73 (s, 6H), 1.62 (s, 6H); MS (ES): 447.4 [M+H]⁺, 469.3 [M+Na]⁺

The following compounds were prepared in a manner similar to that described in experimental procedure Example 19:

- 1-(3-chloro-3'-(2-hydroxypropan-2-yl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-N-(1-hydroxy-2-methylpropan-2-yl)-1H-imidazole-4-carboxamide; MS (ES): 572.3, 574.3, 576.3 [M+H]⁺
- 2-{2-(2,3-dichlorophenyl)-1-[3'-(1-hydroxy-1-methylethyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 481 [M+H]⁺, 503 [M+Na]⁺

Example 20

Preparation of 3'-chloro-4'-[2-(2-chloro-6-fluorophenyl)-4-(1-hydroxy-1-methylethyl)-1H-imidazol-1-yl]-N-methyl-N-(methoxy)-3-(methylsulfonyl)biphenyl-4-carboxamide.



Ethyl 3'-chloro-4'-[2-(2-chloro-6-fluorophenyl)-4-(1-hydroxy-1-methylethyl)-1H-imidazol-1-yl]-3-(methylsulfonyl)biphenyl-4-carboxylate was prepared as described in Example 9. Into a 500 mL flask was weighed 11.19 g (18.92 mmol) of ester, 100 mL of THF, 100 mL of methanol, and 19.0 mL of 3.0 M LiOH-H₂O. The resulting suspension was stirred at room temperature for 21 h then was concentrated *in vacuo*. The residue was treated with ethyl acetate and 1.0 M HCl and the resulting suspension was filtered of solids. The filtrate was placed into a separatory funnel and the ethyl acetate was separated, washed with brine, was dried (Na₂SO₄), and concentrated *in vacuo*. The solids collected by filtration were added and the combined solids were dried under high vacuum to afford 3'-chloro-4'-[2-(2-chloro-6-fluorophenyl)-4-(1-hydroxy-1-methylethyl)-1H-imidazol-1-yl]-3-(methylsulfonyl)biphenyl-4-carboxylic acid as a colorless solid, yield: 7.36 g (69%); MS (ES): 563 [M+H]⁺.

Also recovered as a minor byproduct was 3'-chloro-4'-[2-(2-chloro-6-fluorophenyl)-4-(1-methylethenyl)-1H-imidazol-1-yl]-3-(methylsulfonyl)biphenyl-4-carboxylic acid; MS (ES): 545 and 547 each [M+H]⁺.

Into an 8 mL vial was weighed 68 mg (697 μmol) of *N,O*-Dimethyl-hydroxylamine hydrochloride, and 157 mg (279 μmol) of acid. The solids were then treated with 710 μL of a 0.5 M *O*-(Benzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate solution in DMF and 200 μL of diisopropylethylamine was added. The reaction was stirred at room temperature for 20 h then was washed into a separatory funnel with ethyl acetate and 1 M HCl. The ethyl acetate was separated, washed with brine, was dried (Na₂SO₄), and was concentrated *in vacuo*. The product was purified by silica gel flash chromatography (Biotage, 12.5 x 150 mm SiO₂, gradient elution from 100% hexanes to 100% ethyl acetate over 0.5 h). Appropriate fractions were combined and concentrate *in vacuo* to afford the product as a colorless solid, yield: 85 mg (47%); ¹H NMR (400MHz, DMSO-*d*₆): δ 8.26(s, 1H), 8.19(d, *J* = 8 Hz, 1H), 8.15(s, 1H), 7.87(d, *J* = 6 Hz, 1H), 7.72(d, *J* = 8 Hz, 1H), 7.53(q, *J* = 6 Hz, 1H),


7.4-7.5(m, 3H), 7.31(t, $J = 8$ Hz, 1H), 3.39(s, 3H), 3.32 and 3.36(each s, 3H), 2.75(s, 3H), 1.58(s, 6H); MS (ES): 606 $[M+H]^+$.

In a manner similar to that described, the following examples of the invention were prepared by substituting an appropriate reagent.

- 5 • 3'-Chloro-4'-[2-(2-chloro-6-fluorophenyl)-4-(1-hydroxy-1-methylethyl)-1H-imidazol-1-yl]-3-(methylsulfonyl)biphenyl-4-carboxamide; MS (ES): 562 $[M+H]^+$.
- 3'-Chloro-4'-[2-(2-chloro-6-fluorophenyl)-4-(1-hydroxy-1-methylethyl)-1H-imidazol-1-yl]-*N,N*-dimethyl-3-(methylsulfonyl)biphenyl-4-carboxamide; MS (ES): 590 $[M+H]^+$.
- 3'-Chloro-4'-[2-(2-chloro-6-fluorophenyl)-4-(1-hydroxy-1-methylethyl)-1H-imidazol-1-yl]-*N*-ethyl-3-(methylsulfonyl)biphenyl-4-carboxamide; MS (ES): 590 $[M+H]^+$.
- 10 • 2-{2-(2-Chloro-6-fluorophenyl)-1-[3-chloro-3'-(methylsulfonyl)-4'-(morpholin-4-ylcarbonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol; MS (ES): 632 $[M+H]^+$.
- 3'-Chloro-4'-[2-(2-chloro-6-fluorophenyl)-4-(1-hydroxy-1-methylethyl)-1H-imidazol-1-yl]-3-(methylsulfonyl)-*N*-(phenylmethyl)biphenyl-4-carboxamide; MS (ES): 652 $[M+H]^+$.

Scheme 17

2,4,5-Trisubstituted imidazole (Isomer IIc and tautomer II d)



- Reactions and conditions: (a) TMSCN, ZnI_2 rt; (b) 5-bromothiophene-2-carboxaldehyde, LiHMDS, THF, $-78^\circ C$; (c) Bi_2O_3 , HOAc, $90^\circ C$; (d) NH_4OAc , $CF_3CH(OEt)(OH)$, AcOH, reflux; (e) $ArB(OH)_2$, K_2CO_3 , $PdCl_2(dppf)$, DME/ H_2O , $80^\circ C$.

- The subject imidazole compounds of formula (0176c) are synthesized according to the sequence outlined in Scheme 3. Aldehyde (0171c) such as 2-chlorobenzaldehyde is converted to the protected cyanohydrin of formula (0172c) by reaction with trialkylsilyl cyanide, such as trimethylsilyl cyanide (TMSCN), in the presence of a catalyst such as zinc iodide. Reaction of cyanohydrin (0172c) with aldehyde such as 5-bromothiophene-2-carboxaldehyde, under strong base produces bezoin of


formula (0173c). Oxidation of bezoin with a suitable oxidant such as bismuth oxide provides the diketone of formula (0174c), which subsequently reacts with ammonium acetate and appropriate aldehyde or its equivalent such as trifluoroacetaldehyde ethyl hemiacetal to give 1*H*-imidazole of formula (0175c). In a palladium mediated coupling reaction, for example a Suzuki reaction, compounds of formula (0175c) are then reacted with a boronate or boronic acid reagent to give compounds of formula (0176c) after standard isolation procedures.

Example 21

4-(2-chlorophenyl)-5-(5-(3-(methylsulfonyl)phenyl)thiophen-2-yl)-2-(trifluoromethyl)-1*H*-imidazole

Example 21a


Preparation of (2-Chloro-phenyl)-trimethylsilanyloxy-acetonitrile



Under an atmosphere of nitrogen, anhydrous zinc iodide (30 mg, 0.1 mmol) was placed into a dry flask. At 0°C trimethylsilylcyanide (5.2 g, 52.5 mmol) was added, followed by 2-chlorobenzaldehyde (7.0 g, 50 mmol). The mixture was stirred at 0°C for 3 hrs, then at room temperature overnight. Ethyl ether was added, and the mixture was washed with water and brine, dried over Na₂SO₄, and concentrated *in vacuo* to give a brown oil (10.95 g, 91% yield), which was used directly for the next step without further purification. ¹H-NMR (400MHz, CDCl₃): δ 5.80 (s, 1H), 7.38 (m, 3H), 7.73 (dd, 1H).

Example 21b

Preparation of 2-(5-Bromo-thiophen-2-yl)-1-(2-chloro-phenyl)-2-hydroxy-ethanone




Under an atmosphere of nitrogen, to a cold (-78°C) solution of lithium bis(trimethylsilyl)amide (8.05 mL, 1.0M solution in THF, 8.05 mmol) was added a solution of (2-chloro-phenyl)-trimethylsilanyloxy-acetonitrile (1.69 g, 7 mmol) in tetrahydrofuran (10 mL) dropwise over 10 min. After stirring for another 30min, a solution of 5-bromothiophen-2-carboxaldehyde in tetrahydrofuran (10 mL) was added dropwise. The reaction was stirred overnight from -78°C to reach room temperature, then was quenched by adding 3N HCl (20 mL) and warming at 40°C for 5 hrs. After cooling, the mixture was partitioned between ethyl ether and water. The organic layer was stirred with 1M NaOH (30 mL) for 2 hrs, and the layers were separated. The organic layer was washed with brine, dried over Na₂SO₄, and concentrated *in vacuo*. The crude product was purified by chromatography on silica gel

(Hexane/EtOAc, 8/2) to give a brown oil (320 mg, 14% yield). ¹H-NMR (400MHz, CDCl₃): δ 4.35 (d, 1H), 6.12 (d, 1H), 7.02 (d, 1H), 7.26 (m, 2H), 7.41 (d, 1H), 7.44 (dd, 1H).

Example 21c


Preparation of 1-(5-Bromo-thiophen-2-yl)-2-(2-chloro-phenyl)-ethane-1,2-dione



A mixture of benzoin prepared from above (316 mg, 0.95 mmol), and bismuth oxide (560 mg, 1.2 mmol) in acetic acid (5 mL) was heated at 90°C for 5 hrs. The hot reaction mixture was filtered through celite and the filtrate was concentrated *in vacuo*. The brown residue was redissolved into hot methanol, and then filtered. The filtrate was evaporated to give a brown oil (311 mg, 99% yield), which is used directly for the next step without further purification. ¹H-NMR (400MHz, CDCl₃): δ 7.21 (d, 1H), 7.43 (m, 1H), 7.45 (m, 1H), 7.54 (m, 1H), 7.70 (d, 1H), 7.77 (dd, 1H).

Example 21d


Preparation of 5-(5-Bromo-thiophen-2-yl)-4-(2-chloro-phenyl)-2-trifluoromethyl-1H-imidazole



A mixture of diketone (311 mg, 0.95 mmol), ammonium acetate (366 mg, 4.75 mmol), and trifluoroacetaldehyde ethyl hemiacetal (684 mg, 4.75 mmol) in 5 mL acetic acid was heated at 110°C for 10 hrs under nitrogen atmosphere. All solvent was removed under *vacuo*, and the residue was redissolved into dichloromethane and filtered. The filtrate was concentrated, and then purified by chromatography on silica gel (Hexane/EtOAc, 8/2) to give a light pink solid (141 mg, 36% yield). ¹H-NMR (400MHz, CDCl₃): δ 6.81 (d, 1H), 6.87 (d, 1H), 7.38 (t, 1H), 7.49 (m, 2H), 7.56 (d, 1H), 9.55 (s, 1H).

Example 21e

Preparation of 4-(2-Chloro-phenyl)-5-[5-(3-methanesulfonyl-phenyl)-thiophen-2-yl]-2-trifluoromethyl-1H-imidazole




5-(5-Bromo-thiophen-2-yl)-4-(2-chloro-phenyl)-2-trifluoromethyl-1*H*-imidazole (70 mg, 0.17 mmol), (3-methylsulfonyl)phenylboronic acid (69 mg, 0.34 mmol), potassium carbonate (106 mg, 0.77 mmol), and PdCl₂(dppf).CH₂Cl₂ (14 mg, 0.017 mmol) were mixed with 2 mL 9:1 (v/v) DME/H₂O, then heated at 80 °C overnight. All solvent was removed *in vacuo*. The residue was purified by chromatography on silica gel (Hexane/EtOAc, 6/4) to give a white solid (25 mg, 30% yield). ¹H-NMR (400MHz, CDCl₃): δ 3.07 (s, 3H), 7.00 (d, 1H), 7.12 (m, 1H), 7.51 (m, 5H), 7.81 (m, 2H), 8.10 (t, 1H), 9.65 (s, 1H).

All the following compounds were prepared in similar manner using appropriate aldehydes as starting materials

- 10 • 4-(2-chlorophenyl)-5-[3'-(methylsulfonyl)biphenyl-4-yl]-2-(trifluoromethyl)-1*H*-imidazole; 477.1, 479.3 [M+H]⁺;
- 4'-[4-(2-chlorophenyl)-2-(trifluoromethyl)-1*H*-imidazol-5-yl]biphenyl-3-sulfonamide; 478.0 [M+H]⁺;
- 15 • 3-{5-[4-(2-chlorophenyl)-2-(trifluoromethyl)-1*H*-imidazol-5-yl]-2-thienyl}benzenesulfonamide; 484.0 [M+H]⁺.

Example 22

Preparation of 2-(4-methoxy-3-(methylsulfonyl)phenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane




a) NaHCO₃, Na₂SO₃, H₂O, 85 °C; b) Bis(pinacolato)diboron, Pd(dppf), KOAc, DMSO, 100 °C.

20 Into a 1L flask was weighed 41.4 g of sodium sulfite, 29 g of sodium bicarbonate, and 175 mL of water. The suspension was stirred at 80-85 °C and sulfonyl chloride (50 g) was added portionwise over 3 h. Heating was continued for 3 h then the reaction was allowed to stand at room temperature for 3 days. The intermediate sulfinate was collected by filtration with added water then was dried under high vacuum. The dry solids (45 g) were returned to a 1 L flask along with 28.0 g of sodium bicarbonate, 25 mL of dimethylsulfate, and 63.75 mL of water. The resulting suspension was heated at 25 120-125 °C, where it became a solution, for 20 h then was cooled and washed into a separatory funnel with ethyl acetate and water. The ethyl acetate was separated, washed with brine, was dried (Na₂SO₄), and concentrated *in vacuo*. The product was precipitated from dichloromethane with hexanes and was dried under high vacuum to afford the intermediate 4-Bromo-2-methanesulfonyl-1-methoxy-benzene as 30 a colorless powder, yield: 31.1 g (67%). ¹H NMR (400MHz, CDCl₃): δ 8.08(2, 1H), 7.69(d, *J*= 8 Hz, 1H), 6.96(d, *J*= 8 Hz, 1H), 4.00(s, 3H), 3.21(s, 3H).

Into a 500 mL flask was weighed 15.48 g (58.4 mmol) of bromide, 23 g of boronate, 21 g of potassium acetate, 5 g of dichloro[1,1'-bis(diphenylphosphino)ferrocene]palladium (II) dichloromethane adduct, and 150 mL of DMSO. The resulting suspension was heated at ~100 °C for 20 h then was cooled and diluted with 200 mL of ethyl acetate and 200 mL of water. The suspension was filtered through celite to remove solids and the filtrate was transferred to a separatory funnel. The aqueous phase was separated and washed with ethyl acetate. The ethyl acetate washings were combined, washed with brine, were dried (Na₂SO₄) and concentrated *in vacuo*. The residue was purified by silica gel flash chromatography (Biotage, 65 x 200 mm SiO₂, gradient elution from 100% hexanes to 100% ethyl acetate over 1 h). Appropriate fractions were combined and concentrated *in vacuo*. The partially purified product was dissolved in ethyl acetate and was precipitated with hexanes. The 2-(3-Methanesulfonyl-4-methoxy-phenyl)-4,4,5,5-tetramethyl-[1,3,2]dioxaborolane was recovered as a faintly yellow powder, yield: 12.56 g (77%). ¹H NMR (400MHz, CDCl₃): δ 8.43(s, 1H), 8.01(d, *J*=8 Hz, 1H), 7.03(d, *J*=8 Hz, 1H), 4.02(s, 3H), 3.20(s, 3H), 1.33(s, 12H).

Example 23

Preparation of ethyl 2-(methylsulfonyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate



a) EDCI, DMAP, EtOH, CH₂Cl₂, 45 °C; b) NaSMe, THF, 80 °C; c) MCPBA, CH₂Cl₂, 25 °C; d) Bis(pinacolato)diboron, Pd(dppf), KOAc, DMSO, 85 °C.

Into a 1 L flask was weighed 24.66 g (113 mmol) of acid, 26.5 g (138 mmol) of EDCI, 1.7 g of DMAP, 425 mL of dichloromethane, and 25 mL of ethanol. The resulting solution was heated at 40-45 °C for 24 h then was concentrated *in vacuo* to remove dichloromethane. The residue was washed into a separatory funnel with ethyl acetate and 1 M HCl. The ethyl acetate was separated, washed with brine, was dried (Na₂SO₄) and was concentrated *in vacuo*. The intermediate 4-Bromo-2-fluoro-benzoic acid ethyl ester was recovered as a colorless oil, yield: 24.99 g (89.8%).

The ester was treated with 12.2 g of sodium thiomethoxide and 200 mL of THF and the resulting suspension was heated at 80-85 °C for 5 h. The reaction was then concentrated to remove THF and was washed into a separatory funnel with ethyl acetate and 1 M HCl. The ethyl acetate was separated, washed with brine, was dried (Na₂SO₄), and concentrated *in vacuo* affording the intermediate 4-Bromo-2-methylsulfonyl-benzoic acid ethyl ester as a light gray solid, yield: 27.5 g (99%). ¹H NMR


(400MHz, CDCl₃): δ 7.86(d, J = 8 Hz, 1H), 7.36(s, 1H), 7.28(d, J = 8 Hz, 1H), 4.38(q, J = 7 Hz, 2H), 2.45(s, 3H), 1.39(t, J = 7 Hz, 3H).

Into a 1 L flask was weighed 15.0 g of 4-Bromo-2-methylsulfonyl-benzoic acid ethyl ester (54.5 mmol), 200 mL of dichloromethane, and 28.0 g of MCPBA (77% max., Aldrich) was added portionwise at room temperature. The resulting suspension was stirred at room temperature for three days then was concentrated *in vacuo* to remove dichloromethane. The residue was washed into a separatory funnel with ethyl acetate and 1.0 M NaOH. The ethyl acetate was separated, washed with brine, was dried (Na₂SO₄), and concentrated *in vacuo*. The intermediate 4-Bromo-2-methanesulfonyl-benzoic acid ethyl ester was recovered as a colorless oil which crystallized on standing, yield: 16.3 g (97%). ¹H NMR (400MHz, CDCl₃): δ 8.27(s, 1H), 7.82(d, J = 8 Hz, 1H), 7.60(d, J = 8 Hz, 1H), 4.44(q, J = 7 Hz, 2H), 3.38(s, 3H), 1.41(t, J = 7 Hz, 3H).

The 4-Bromo-2-methanesulfonyl-benzoic acid ethyl ester (16.3 g, 53 mmol) was weighed into a flask with 21 g of bis(pinacolato)diboron, 19 g of potassium acetate, 5 g of dichloro[1,1'-bis(diphenylphosphino)ferrocene]palladium (II) dichloromethane adduct, and 150 mL of DMSO. The resulting suspension was heated at 80-85 °C for 20 h then was diluted with 200 mL of water, 200 mL of ethyl acetate, and the reaction mixture was filtered through celite to remove solids. The filtrate was transferred to a separatory funnel and the aqueous phase was separated and washed with ethyl acetate. The ethyl acetate washings were combined, washed with brine, were dried (Na₂SO₄) and concentrated *in vacuo*. The residue was purified by silica gel flash chromatography (Biotage, 65 x 200 mm SiO₂, gradient elution from 100% hexanes to 40% ethyl acetate over 1 h). Appropriate fractions were combined and concentrated *in vacuo* affording the product as a colorless solid, yield: 12.65 g (67%). ¹H-NMR (400MHz, CDCl₃): δ 8.52(s, 1H), 8.08(d, J = 8 Hz, 1H), 7.65(d, J = 8 Hz, 1H), 4.45(q, J = 7 Hz, 2H), 3.33(s, 3H), 1.42(t, J = 7 Hz, 3H), 1.35(s, 12H).

Example 24

Preparation of (2-(methylsulfonyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)methanol



a) LiBH₄, THF, 85 °C; b) MCPBA, CH₂Cl₂, 25 °C; c) Bis(pinacolato)diboron, Pd(dppe), KOAc, DMSO, 100 °C.

The 4-Bromo-2-methanesulfonyl-benzoic acid ethyl ester was prepared as described in Example 23. Into a 1 L flask was weighed 27.5 g of ester (99.9 mmol) and 150 mL of THF. A solution of 2.0 M LiBH₄ in THF (50 mL, 100 mmol) was then added and the reaction was heated to 80-85 °C where it remained for 23h. The reaction was then removed from heat and was cooled in an ice bath as it

was quenched by addition of acetone. The reaction was then concentrated *in vacuo* and was washed into a separatory funnel with ethyl acetate and 1 M HCl. The ethyl acetate was separated, washed with brine, was dried (Na₂SO₄), and concentrated *in vacuo*. The intermediate (4-Bromo-2-methylsulfonyl-phenyl)-methanol was recovered as a colorless oil that solidified on standing, yield: 25.5 g (100+%).

5 ¹H NMR (400MHz, CDCl₃): δ 7.24-7.34(m, 3H), 4.69(s, 2H), 2.50(s, 3H).


The alcohol was then dissolved in 250 mL of dichloromethane, was cooled to 0-3 °C in an ice bath, and 44 g of 3-chloroperbenzoic acid (77% max., Aldrich) was added portionwise. The reaction was then allowed to warm to room temperature where it remained for 22 h. The reaction was then concentrated *in vacuo* to remove dichloromethane and the residue was washed into a separatory funnel
10 with ethyl acetate and 1 M NaOH. The ethyl acetate was separated, washed with 1 M NaOH, was dried (Na₂SO₄), and concentrated *in vacuo*. The residue was purified by silica gel flash chromatography (Biotage, 65 x 200 mm SiO₂, gradient elution from 100% hexanes to 100% ethyl acetate over 1 h). Appropriate fractions were combined and concentrated *in vacuo* to afford the intermediate (4-Bromo-2-methanesulfonyl-phenyl)-methanol as a colorless, semi-crystalline solid, yield: 17.13 g (65%). ¹H
15 NMR (400MHz, CDCl₃): δ 8.18(s, 1H), 7.77(d, *J* = 8 Hz, 1H), 7.46(d, *J* = 8 Hz, 1H), 4.92(s, 2H), 3.19(s, 3H), 2.94(br s, 1H).

Into a 1 L flask was weighed 17.13 g of bromide, 25 g of bis(pinacolato)diboron, 5.0 g of dichloro[1,1'-bis(diphenylphosphino)ferrocene]palladium (II) dichloromethane adduct, 23 g of potassium acetate, and 175 mL of DMSO. The resulting suspension was heated at 98-102 °C for 18 h then was
20 diluted with 200 mL of ethyl acetate and 200 mL of water. The resulting suspension was filtered through celite to remove solids and the filtrate was transferred to a separatory funnel. The aqueous phase was separated and washed with ethyl acetate. The ethyl acetate washings were combined, washed with brine, were dried (Na₂SO₄) and concentrated *in vacuo*. The residue was purified by silica gel flash chromatography (Biotage, 65 x 200 mm SiO₂, gradient elution from 100% hexanes to 40%
25 ethyl acetate over 1 h). Appropriate fractions were combined and concentrated *in vacuo*. The partially purified product was dissolved in dichloromethane and was precipitated with hexanes. The [2-Methanesulfonyl-4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-methanol was recovered as an off-white powder, yield: 8.78 g (43%). ¹H NMR (400MHz, CDCl₃): δ 8.45(s, 1H), 8.04(d, *J* = 8
Hz, 1H), 7.57(d, *J* = 8 Hz, 1H), 4.96(s, 1H), 3.17(s, 3H), 1.35(s, 6H), 1.24(s, 6H).

30

Example 25

Preparation of 6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-2,3-dihydro-1-benzothiophene-3-ol 1,1-dioxide



a) DMSO, oxalyl chloride, CH_2Cl_2 , $-78\text{ }^\circ\text{C}$; b) MCPBA, CH_2Cl_2 , $25\text{ }^\circ\text{C}$; c) Bis(pinacolato)diboron, $\text{Pd}(\text{dppf})$, KOAc, DMSO, $100\text{ }^\circ\text{C}$.

The (4-Bromo-2-methylsulfanyl-phenyl)-methanol was prepared as described in Example 24.

5 Into a 500 mL flask was placed 220 mL of dichloromethane and 5.66 mL of oxalyl chloride under nitrogen. The solution was cooled to -70 to $-78\text{ }^\circ\text{C}$ and 9.80 mL of DMSO was added dropwise. The resulting suspension was stirred at $-78\text{ }^\circ\text{C}$ for 15 minutes. A solution of alcohol (10.0 g, 42.9 mmol) in 30 mL of dichloromethane was prepared and was added to the suspension via cannula. The reaction was stirred at $-78\text{ }^\circ\text{C}$ for 0.5 h then 30 mL of triethylamine was added via syringe. The reaction was stirred at $-78\text{ }^\circ\text{C}$ for 30 minutes then was quenched by addition of saturated ammonium chloride. The mixture was washed into a separatory funnel with added water and dichloromethane. The dichloromethane was separated, dried (Na_2SO_4), and concentrated *in vacuo*. The residue was purified by silica gel flash chromatography (Biotage, 65 x 200 mm SiO_2 , gradient elution from 100% hexanes to 20% ethyl acetate over 1 h). Appropriate fractions were combined and concentrated *in vacuo* to afford the product as a lemon yellow solid, yield: 6.78 g (88% based on recovered starting material). ^1H NMR (400MHz, CDCl_3): δ 10.19(s, 1H), 7.65(d, $J=8\text{ Hz}$, 1H), 7.40-7.45(m, 2H), 2.50(s, 3H).


15 Into a 250 mL flask was weighed 6.78 g (25.8 mmol) of aldehyde and 60 mL of dichloromethane. The reaction was cooled to 0 - $3\text{ }^\circ\text{C}$ in an ice bath and 10 g of MCPBA (77% max., Aldrich) was added portionwise. The reaction was allowed to warm to room temperature where it remained for 21 h. The reaction was then concentrated *in vacuo* to remove dichloromethane and was washed into a separatory funnel with ethyl acetate and 1 M NaOH. The ethyl acetate was separated, washed with 1.0 M NaOH, brine, was dried (Na_2SO_4), and was concentrated *in vacuo*. The residue was purified by silica gel flash chromatography (Biotage, 45 x 150 mm SiO_2 , gradient elution from 100% hexanes to 20% ethyl acetate over 1 h). Appropriate fractions were combined and concentrated *in vacuo* to afford the product as a colorless solid, yield: 5.32 g (89%). ^1H NMR (400MHz, CDCl_3): δ 10.71(s, 1H), 8.30(s, 1H), 7.95(s, 2H), 3.29(s, 3H).

25 Into a 250 mL flask was weighed 5.32 g (20.2 mmol) of aldehyde, 7.13 g of potassium acetate, 7.85 g of bis(pinacolato)diboron, 1.88 g of dichloro[1,1'-bis(diphenylphosphino)ferrocene]palladium (II) dichloromethane adduct, and 55 mL of DMSO. The resulting suspension was heated at 98 - $102\text{ }^\circ\text{C}$ for 20 h then was diluted with 100 mL of ethyl acetate and 100 mL of water. The resulting suspension was filtered through celite to remove solids and the filtrate was transferred to a separatory funnel. The

aqueous phase was separated and washed with ethyl acetate. The ethyl acetate washings were combined, washed with brine, were dried (Na_2SO_4) and concentrated *in vacuo*. The residue was purified by silica gel flash chromatography (Biotage, 65 x 200 mm SiO_2 , gradient elution from 100% hexanes to 40% ethyl acetate over 0.5 h). Appropriate fractions were combined and concentrated *in vacuo*. The partially purified product was dissolved in dichloromethane and was precipitated with hexanes. The 1,1-Dioxo-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-2,3-dihydro-1H-1 λ *6*-benzo[b] thiophen-3-ol was recovered as a tan powder, yield: 1.88 g (30%). ^1H NMR (400MHz, CDCl_3): δ 8.21(s, 1H), 8.08(d, J = 8 Hz, 1H), 7.64(d, J = 8 Hz, 1H), 5.49(m, 1H), 3.80(dd, J = 7, 13 Hz, 1H), 3.45(dd, J = 5, 13 Hz, 1H), 2.79(d, J = 8 Hz, 1H), 1.34(s, 12H).

Example 26

Preparation of 2-(2-(methylsulfonyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)propan-2-ol



a) MeMgBr , THF, $-78\text{ }^\circ\text{C}$; b) MCPBA, CH_2Cl_2 , $50\text{ }^\circ\text{C}$; c) Bis(pinacolato)diboron, $\text{Pd}(\text{dppf})$, KOAc, DMSO, $100\text{ }^\circ\text{C}$.

The 4-Bromo-2-methanesulfonyl-benzoic acid ethyl ester was prepared as described in Example 25. Into a 500 mL flask was weighed 10.2 g (37.1 mmol) of ester and 100 mL of anhydrous THF. The resulting solution was cooled to $-78\text{ }^\circ\text{C}$ and 80 mL of 1.4 M MeMgBr in THF (Aldrich) was added portionwise. The reaction was allowed to warm to room temperature where it remained for 3 h. The reaction was then quenched by addition of saturated ammonium chloride and the mixture was washed into a separatory funnel with ethyl acetate and saturated ammonium chloride. The ethyl acetate was separated, washed with brine, was dried (Na_2SO_4), and concentrated *in vacuo*. The intermediate 2-(4-Bromo-2-methylsulfonyl-phenyl)-propan-2-ol was carried into the subsequent step without additional purification.


The crude alcohol was dissolved in 100 mL of dichloromethane, was cooled to $0-3\text{ }^\circ\text{C}$ in an ice bath, and 20 g of MCPBA (77% max., Aldrich) was added portionwise. The reaction was then removed to an oil bath where it was heated at $\sim 50\text{ }^\circ\text{C}$ for 18 h. The reaction was then cooled and concentrated *in vacuo* to remove dichloromethane. The residue was washed into a separatory funnel with ethyl acetate and 1.0 M NaOH. The ethyl acetate was separated, washed with 1.0 M NaOH, brine,

was dried (Na₂SO₄), and concentrated *in vacuo*. The residue was purified by silica gel flash chromatography (Biotage 45 x 150 mm SiO₂, gradient elution from 100% hexanes to 40% ethyl acetate over 1 h). Appropriate fractions were combined and concentrated *in vacuo* affording the intermediate 2-(4-Bromo-2-methanesulfonyl-phenyl)-propan-2-ol as a colorless solid, yield: 10.02 g (92% for both
 5 steps). ¹H NMR (400MHz, CDCl₃): δ 8.33(s, 1H), 7.68(d, *J* = 8 Hz, 1H), 7.33(d, *J* = 8 Hz, 1H), 3.41(s, 3H), 1.76(s, 1H), 1.69(s, 6H).

Into a 500 mL flask was weighed 9.88 g (33.7 mmol) of bromide, 12 g of potassium acetate, 13.1 g of bis(pinacolato)diboron, 3.0 g of dichloro[1,1'-bis(diphenylphosphino)ferrocene]palladium (II) dicloromethane adduct, and 90 mL of DMSO. The resulting suspension was heated at 98-102 °C for 20
 10 h then was diluted with 200 mL of ethyl acetate and 200 mL of water. The resulting suspension was filtered through celite to remove solids and the filtrate was transferred to a separatory funnel. The aqueous phase was separated and washed with ethyl acetate. The ethyl acetate washings were combined, washed with brine, were dried (Na₂SO₄) and concentrated *in vacuo*. The residue was purified by silica gel flash chromatography (Biotage, 65 x 200 mm SiO₂, gradient elution from 100%
 15 hexanes to 100% ethyl acetate over 1 h). Appropriate fractions were combined and concentrated *in vacuo* affording the 2-[2-Methanesulfonyl-4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-propan-2-ol as a gray solid, yield: 10.0 g (87%). ¹H NMR (400MHz, CDCl₃): δ 8.61(s, 1H), 7.96(d, *J* = 8 Hz, 1H), 7.45(d, *J* = 8 Hz, 1H), 3.39(s, 3H), 1.70(s, 6H), 1.34(s, 12H).

Several different boronates can be synthesized by the general methods outlined in Scheme 18
 20 and exemplified in Example 27. A suitable aromatic substrate, such as a 2-halopyridine, can react to produce an intermediate with an 'directing' group attached. Such a system can then be halogenated to afford a *para*-disposed halogen. Such a system can then be converted to a boronate using methods such as those described below. Alternatively, several commercially available intermediates could be used such as 2,4-Dibromo-3-methylpyridine, 4-Bromo-2-chlorpyrimidine, and several substituted benzene
 25 rings.


Scheme 18



Q = Used to identify any aromatic ring system that can be suitably functionalized as would be readily apparent to one skilled in the art.
 X = halogen or preexisting 'directing' group
 R' = 'directing group'
 R = any suitable functionality that will not interfere with the directing effect of the R' group

Example 27

Preparation of 2-(methylthio)-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridine



Reaction condition a) NaSMe, DMF, 160 °C; b) Bis(pinacolato)diboron, Pd(dppf), KOAc,

5 DMSO, 80 °C.

2,5-Dibromo-pyridine (3.0g, 12.7mmol) and sodium thiomethoxide (0.84g, 12mmol) were dissolved in 18ml anhydrous *N, N*-dimethylformamide. The mixture was heated at 160°C under nitrogen for 6 hrs. After cooling to room temperature, water and ethyl acetate were added to the reaction mixture. The aqueous layer was extracted with ethyl acetate several times. The combined extract was washed with brine, dried over sodium sulfate, and concentrated *in vacuo*. The crude product was purified by column chromatography on silica gel (0→6% EtOAc/Hexane) to give 5-Bromo-2-methylsulfanyl-pyridine as a white solid (2.18g, 84% yield). ¹H-NMR (400MHz, CDCl₃): δ 2.55 (s, 3H), 7.09 (dd, 1H, *J*=0.7, *J*=8.6), 7.59 (dd, 1H, *J*=2.4, *J*=8.6), 8.50 (m, 1H).


Palladium catalyst ([1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium (II) complex with dichloromethane (1:1), 167mg, 0.21mmol), potassium acetate (1.81g, 18.5mmol, Aldrich), and bis(pinacolato)diboron (1.56g, 6.1mmol) were placed into a vial and degassed with stream of nitrogen for 20min. In a separate vial, 5-bromo-2-methylsulfanylpyridine (no. (2)) (836mg, 4.1mmol) was dissolved in 8ml anhydrous DMSO and degassed with stream of nitrogen for 20min. This DMSO solution was added to the "catalyst" vial, and then heated at 80°C overnight. After cooling to room temperature, water and ethyl acetate were added to the reaction mixture. The aqueous layer was extracted with ethyl acetate. The combined organic layers were washed with brine, dried over sodium sulfate, and concentrated *in vacuo*. The crude product was purified by column chromatography on silica gel (10→30% EtOAc/Hexane, 025% Et₃N in hexane) to give the 2-Methylsulfanyl-5-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)pyridine as a colorless oil (1.00g, 97% yield). ¹H NMR (400MHz, CDCl₃): δ 1.35 (s, 12H), 2.58 (s, 3H), 7.16 (dd, 1H, *J*=1.0, *J*=8.0), 7.83 (dd, 1H, *J*=1.8, *J*=8.0), 8.50 (dd, 1H, *J*=1.7, *J*=1.0)

In a manner similar to that stated the following examples of the invention were prepared by substituting an appropriate reagent.

- 2-Methylsulfanyl-3-methyl-5-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)pyridine. ¹H-NMR (400MHz, CDCl₃): δ 1.35 (s, 12H), 2.24 (s, 3H), 2.59 (s, 3H), 7.64 (d, *J*=0.8, 1H), 8.63 (d, *J*=1.0, 1H).

- 2-Ethylsulfonyl-3-methyl-5-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)pyridine. ¹H-NMR (400MHz, CDCl₃): δ 1.35 (s, 12H), 1.38 (t, *J*=7.4, 3H), 2.22 (s, 3H), 3.23 (q, *J*=7.4, 2H), 7.64 (d, *J*=0.8, 1H), 8.61 (d, *J*=1.0, 1H). MS (ES): 280.2, 282.1 [M+H]⁺.

Scheme 19




a) NaO₂SR², 10 mol% CuI, 20 mol% proline, 20 mol% NaOH, DMSO, μW 210°C; b) bis(pinacolato)-diboron, KOAc, Cl₂Pd(dppf), DMSO, μW 130°C.

In general, compounds of formula 1,3- (or 3,5-) disubstituted aryl systems can be prepared as depicted in Scheme 19. A dibromoarene can react with a sulfinic acid sodium salt (NaO₂SR²) in the presence of catalytic copper iodide and the sodium salt of proline at elevated temperature to give the corresponding sulfone. This intermediate then can be converted into the corresponding arylboronate or boronic acid under standard conditions apparent to one skilled in the art.

Example 28


Preparation of 1-bromo-3-methanesulfonyl-5-methyl-benzene



A mixture of 3,5-dibromotoluene (2.0 g, 8.0 mmol), sodium methanesulfinate (0.92 g, 9.0 mmol), copper(I) iodide (152 mg, 0.8 mmol), proline (0.18 g, 1.6 mmol) and sodium hydroxide (64 mg, 1.6 mmol) in DMSO (15 mL, anhyd) was heated in a microwave unit (Biotage Initiator™) at 210°C for 20 min. The reaction was repeated two more times and the reaction mixtures were combined, diluted with H₂O and extracted with EtOAc. The combined extracts were washed with H₂O and brine, then dried (Na₂SO₄), concentrated and purified by chromatography (silica, EtOAc/Hex, 0:100 to 40:60) to give the title compound (1.2 g, 20% combined yield) as a white solid. GC-MS(EI): 248, 250.

2-(3-Methanesulfonyl-5-methyl-phenyl)-4,4,5,5-tetramethyl-[1,3,2]dioxaborolane was prepared from 1-bromo-3-methanesulfonyl-5-methyl-benzene in a similar manner as other boronate syntheses previously described.

Scheme 20.




a) $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$, conc HCl, DME/EtOH; b) NaNO_2 , 48% HBr, H_2O ; then CuBr; c) bis(pinacolato)-diboron, KOAc, $\text{Cl}_2\text{Pd}(\text{dppf})$, DMSO, μW 130°C .

5 In general, several different boronates can be prepared as depicted in Scheme 20. A nitroarene can be reduced under standard conditions, such as with tin chloride, to yield the corresponding arylamine, which can be converted to the corresponding bromoarene via the Sandmeyer reaction. Aryl bromides can easily be converted to boronates using conditions apparent to one skilled in the art.

Example 29

Preparation of 2-bromo-1-chloro-4-methanesulfonyl-benzene

10



15 To a stirred mixture of 1-chloro-4-methanesulfonyl-2-nitrobenzene (5.2 g, 22 mmol) and 3:4 DME/EtOH (77 mL) was added dropwise a solution of tin(II) chloride dihydrate (15.3 g, 68 mmol) in 9N HCl (36 mL). After 3 h the reaction mixture was poured onto ice and treated with 10% NaOH until pH 11 had been achieved. The resulting mixture was extracted with EtOAc (4 x 75 mL) and the combined extracts were washed with brine, dried (Na_2SO_4) and concentrated to afford the title compound (4.5 g, quant) as a white solid, which was used without purification in the next step. GC-MS(EI): 205.

20 To a slurry of 2-chloro-5-methanesulfonyl-aniline (4.50 g, 21.9 mmol) in 48% HBr (12 mL) at 0°C was added dropwise a solution of sodium nitrite (2.28 g, 33 mmol) in H_2O (8 mL) at such a rate that the temperature never exceeds 5°C . The resulting yellow mixture was allowed to stir 20 min at 0°C and then charged with copper(I) bromide (3.3 g, 23 mmol). After 30 min the resulting mixture was extracted with EtOAc (2 x 100 mL) and the combined extracts were washed successively with satd NH_4Cl , 3N

HCl and brine, then dried (Na₂SO₄) and concentrated under reduced pressure. The resulting residue was purified by chromatography (silica, EtOAc/Hex, 0:100 to 50:50) to give the title compound (4.92 g, 83%) as a white solid. GC-MS(EI): 268, 270.

2-(2-Chloro-5-methanesulfonyl-phenyl)-4,4,5,5-tetramethyl-[1,3,2]dioxaborolane was prepared from 2-bromo-1-chloro-4-methanesulfonyl-benzene in a similar manner as other boronate syntheses previously described.

Example 30

The following compounds of the invention, in Tables 1 and 2, were prepared according to one of the previous Example 1-29.

10

Table 1

Cpd #	Structure	Name	Cpd #	Structure	Name
1		2-(2-(2-fluorobenzyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;	7		2-(2-(2,6-dichlorobenzyl)-1-(3'-(ethylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
2		2-(1-(3'-(methylsulfonyl)biphenyl-4-yl)-2-(2-(trifluoromethyl)benzyl)-1H-imidazol-4-yl)propan-2-ol;	8		2-(2-(2,6-dichlorobenzyl)-1-(3'-(ethylsulfonyl)-4'-(hydroxymethyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
3		2-(2-(2-chlorobenzyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;	9		2-(2-(2-chloro-6-fluorobenzyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
4		2-(2-(2-chlorobenzyl)-1-(3'-(ethylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;	10		2-(2-(2-chloro-6-fluorobenzyl)-1-(3'-(ethylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
5		2-(2-(2-chlorobenzyl)-1-(4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;	11		2-(2-(2,3-dichlorobenzyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
6		2-(2-(2,6-dichlorobenzyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;	12		2-(2-(2,3-dichlorobenzyl)-1-(3'-(ethylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;

Cpd #	Structure	Name
13		3-(2-(2-chlorobenzyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)pentan-3-ol;
14		3-(2-(2-chlorobenzyl)-1-(3'-(ethylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)pentan-3-ol;
15		3-(2-(2-chlorobenzyl)-1-(4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)pentan-3-ol;
16		2-(2-(5-chloro-2-(trifluoromethyl)benzyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
17		2-(2-(5-chloro-2-(trifluoromethyl)benzyl)-1-(3'-(ethylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
18		2-(2-(5-chloro-2-(trifluoromethyl)benzyl)-1-(4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
19		2-(2-(2-chlorophenyl)propan-2-yl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
20		2-(2-(2-chlorophenyl)propan-2-yl)-1-(3'-(ethylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;

Cpd #	Structure	Name
21		2-(2-(2-(2,3-dichlorophenyl)propan-2-yl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
22		2-(2-(2-(2,3-dichlorophenyl)propan-2-yl)-1-(3'-(ethylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
23		2-(2-(2,3-dichlorobenzyl)-1-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
24		2-(2-(2,3-dichlorobenzyl)-1-(3'-(ethylsulfonyl)-3-methylbiphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
25		2-(1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2,3-dichlorobenzyl)-1H-imidazol-4-yl)propan-2-ol;
26		2-(1-(3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl)-2-(2,3-dichlorobenzyl)-1H-imidazol-4-yl)propan-2-ol;
27		2-(2-(2-chlorophenethyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
28		2-(2-(2-chlorophenethyl)-1-(3'-(ethylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;

Cpd #	Structure	Name
29		2-(2-(2-chlorophenethyl)-1-(4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
30		2-(1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(naphthalen-1-yl)-1H-imidazol-4-yl)propan-2-ol;
31		2-(1-(3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl)-2-(naphthalen-1-yl)-1H-imidazol-4-yl)propan-2-ol;
32		2-(1-(3-chloro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-2-(naphthalen-1-yl)-1H-imidazol-4-yl)propan-2-ol;
33		2-(1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(isoquinolin-1-yl)-1H-imidazol-4-yl)propan-2-ol;
34		2-(1-(3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl)-2-(isoquinolin-1-yl)-1H-imidazol-4-yl)propan-2-ol;
35		2-(2-(isoquinolin-5-yl)-1-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
36		2-(1-(3'-(ethylsulfonyl)-3-methylbiphenyl-4-yl)-2-(isoquinolin-5-yl)-1H-imidazol-4-yl)propan-2-ol;

Cpd #	Structure	Name
37		2-(1-(3'-(methylsulfonyl)biphenyl-4-yl)-2-(4-(trifluoromethyl)pyridin-3-yl)-1H-imidazol-4-yl)propan-2-ol;
38		2-(1-(3'-(ethylsulfonyl)biphenyl-4-yl)-2-(4-(trifluoromethyl)pyridin-3-yl)-1H-imidazol-4-yl)propan-2-ol;
39		2-(1-(4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-2-(4-(trifluoromethyl)pyridin-3-yl)-1H-imidazol-4-yl)propan-2-ol;
40		2-(1-(3'-(methylsulfonyl)biphenyl-4-yl)-2-(2-morpholinoethyl)-1H-imidazol-4-yl)propan-2-ol;
41		2-(1-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-2-(thiophen-2-yl)-1H-imidazol-4-yl)propan-2-ol;
42		2-(2-(1,5-dimethyl-1H-pyrrol-2-yl)-1-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
43		2-(2-(1,5-dimethyl-1H-pyrrol-2-yl)-1-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
44		2-(1-(2,6-dichlorophenyl)-2-(5-(3-(methylsulfonyl)phenyl)thiophen-2-yl)-1H-imidazol-4-yl)propan-2-ol;

Cpd #	Structure	Name
45		2-(1-(2,6-dichlorophenyl)-2-(5-(3-(ethylsulfonyl)phenyl)thiophen-2-yl)-1H-imidazol-4-yl)propan-2-ol;
46		1-(2,5-dichlorophenyl)-2-{5-[3-(methylsulfonyl)phenyl]-2-thienyl}-4-(trifluoromethyl)-1H-imidazole;
47		1-(2-chlorophenyl)-2-{5-[3-(methylsulfonyl)phenyl]-2-thienyl}-4-(trifluoromethyl)-1H-imidazole;
48		methyl 4-{5-[1-(2-chlorophenyl)-4-(trifluoromethyl)-1H-imidazol-2-yl]-2-thienyl}-3-methylphenylacetate;
49		5-{5-[1-(2,5-dichlorophenyl)-4-(trifluoromethyl)-1H-imidazol-2-yl]-2-thienyl}-2-(ethylthio)-3-methylpyridine;
50		5-{5-[1-(2,5-dichlorophenyl)-4-(trifluoromethyl)-1H-imidazol-2-yl]-2-thienyl}-2-(ethylsulfonyl)-3-methylpyridine;
51		5-{5-[1-(2,5-dichlorophenyl)-4-(trifluoromethyl)-1H-imidazol-2-yl]-2-thienyl}-3-methyl-2-(methylsulfonyl)pyridine;
52		4-{5-[5-[1-(2,5-dichlorophenyl)-4-(trifluoromethyl)-1H-imidazol-2-yl]-2-thienyl]pyridin-2-yl)morpholine;

Cpd #	Structure	Name
53		1,1-dimethylethyl 4-{5-[5-[1-(2,5-dichlorophenyl)-4-(trifluoromethyl)-1H-imidazol-2-yl]-2-thienyl]pyridin-2-yl}piperazine-1-carboxylate;
54		1-{5-[5-[1-(2,5-dichlorophenyl)-4-(trifluoromethyl)-1H-imidazol-2-yl]-2-thienyl]pyridin-2-yl}piperazine;
55		5-{5-[1-(2-chlorophenyl)-4-(trifluoromethyl)-1H-imidazol-2-yl]-2-thienyl}-2-(ethylthio)-3-methylpyridine;
56		5-{5-[1-(2-chlorophenyl)-4-(trifluoromethyl)-1H-imidazol-2-yl]-2-thienyl}-3-methyl-2-(methylsulfonyl)pyridine;
57		5-{5-[1-(2-chlorophenyl)-4-(trifluoromethyl)-1H-imidazol-2-yl]-2-thienyl}-2-(ethylsulfonyl)-3-methylpyridine;
58		2-(1-(2,6-dichlorophenyl)-2-(3-(ethylsulfonyl)-3-methylbiphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
59		2-(1-(2-isopropyl-6-methylphenyl)-2-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
60		2-(2-(3-(ethylsulfonyl)-3-methylbiphenyl-4-yl)-1-(2-isopropyl-6-methylphenyl)-1H-imidazol-4-yl)propan-2-ol;

Cpd #	Structure	Name
61		2-(1-(2-isopropylphenyl)-2-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
62		2-(2-(3'-(ethylsulfonyl)-3-methylbiphenyl-4-yl)-1-(2-isopropylphenyl)-1H-imidazol-4-yl)propan-2-ol;
63		2-(1-(2,6-dichlorophenyl)-2-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
64		2-(1-(2,6-dichlorophenyl)-2-(3'-(ethylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
65		2-(2-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-1-(2,6-dichlorophenyl)-1H-imidazol-4-yl)propan-2-ol;
66		2-(2-(3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl)-1-(2,6-dichlorophenyl)-1H-imidazol-4-yl)propan-2-ol;
67		1,1,1-trifluoro-2-(1-(3'-(methylsulfonyl)biphenyl-4-yl)-2-(2-(trifluoromethyl)phenyl)-1H-imidazol-4-yl)propan-2-ol;
68		ethyl 1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-1H-imidazole-4-carboxylate;

Cpd #	Structure	Name
69		1-(2-(2,6-dichlorophenyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)ethanone;
70		(E)-1-(2-(2,6-dichlorophenyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)ethanone oxime;
71		(E)-1-(2-(2-ethylphenyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)ethanone oxime;
72		3-(2-(2,3-dichlorophenyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)pentan-3-ol;
73		3-(2-(2,3-dichlorophenyl)-1-(3'-(ethylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)pentan-3-ol;
74		3-(2-(2,6-dichlorophenyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)pentan-3-ol;
75		2-(2-isopropylphenyl)-1-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-N-(2,2,2-trifluoroethyl)-1H-imidazole-4-carbothioamide;
76		(E)-1-(2-(2-isopropylphenyl)-1-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)ethanone oxime;

Cpd #	Structure	Name
77		2-(2,3-dichlorophenyl)-1-(3'-(ethylsulfonyl)biphenyl-4-yl)-4-(prop-1-en-2-yl)-1H-imidazole;
78		2-(2,6-dichlorophenyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-4-(prop-1-en-2-yl)-1H-imidazole;
79		3'-chloro-4'-(2-(2-chloro-6-fluorophenyl)-4-(prop-1-en-2-yl)-1H-imidazol-1-yl)-3-(methylsulfonyl)biphenyl-4-carboxylic acid;
80		N-benzyl-3'-chloro-4'-(2-(2-chloro-6-fluorophenyl)-4-(2-hydroxypropan-2-yl)-1H-imidazol-1-yl)-3-(methylsulfonyl)biphenyl-4-carboxamide;
81		3'-chloro-4'-(2-(2-chloro-6-fluorophenyl)-4-(2-hydroxypropan-2-yl)-1H-imidazol-1-yl)-N-ethyl-3-(methylsulfonyl)biphenyl-4-carboxamide;
82		2-(2,6-dichlorophenyl)-N-ethyl-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazole-4-carboxamide;
83		2-(2,6-dichlorophenyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazole-4-yl(4-methylpiperazin-1-yl)methanone;

Cpd #	Structure	Name
84		2-(2,6-dichlorophenyl)-N-(2-hydroxyethyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazole-4-carboxamide;
85		2-(2,6-dichlorophenyl)-1-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-N-(2,2,2-trifluoroethyl)-1H-imidazole-4-carboxamide;
86		2-(2,6-dichlorophenyl)-N,N-diethyl-1-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazole-4-carboxamide;
87		2-(2,6-dichlorophenyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-N-(2,2,2-trifluoroethyl)-1H-imidazole-4-carboxamide;
88		2-(2,6-dichlorophenyl)-N-(2-fluoroethyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazole-4-carboxamide;
89		2-(2-isopropylphenyl)-1-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-N-(2,2,2-trifluoroethyl)-1H-imidazole-4-carboxamide;
90		2-(2-(dimethylamino)-6-fluorophenyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazole-4-ylpropan-2-ol;

Cpd #	Structure	Name
91		2-(2-(2,6-dichlorophenyl)-1-(3'-(ethylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
92		2-(2-(2-chloro-6-fluorophenyl)-1-(3'-(ethylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
93		2-(2-(2-ethylphenyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
94		2-(2-(2-isopropylphenyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
95		2-(2-(2-chloro-3-fluorophenyl)-1-(3'-(ethylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
96		2-(1-(3'-(ethylsulfonyl)biphenyl-4-yl)-2-(2-isopropylphenyl)-1H-imidazol-4-yl)propan-2-ol;
97		2-(2-(2-chloro-6-methylphenyl)-1-(3'-(ethylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
98		2-(2-(2,3-dichlorophenyl)-1-(4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;

Cpd #	Structure	Name
99		2-(2-(2,6-dichlorophenyl)-1-(4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
100		2-(2-(2-chloro-6-fluorophenyl)-1-(4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
101		2-(2-(3-chloro-2-methylphenyl)-1-(4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
102		2-(2-(2-chloro-6-methylphenyl)-1-(4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
103		2-(2-(2,6-difluorophenyl)-1-(4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
104		2-(2,6-dichlorophenyl)-1-(3'-(ethylsulfonyl)-3-methylbiphenyl-4-yl)-N-(2,2,2-trifluoroethyl)-1H-imidazole-4-carboxamide;
105		2-(2-(2,3-dichlorophenyl)-1-(4'-(2-hydroxypropan-2-yl)-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;

Cpd #	Structure	Name
106		2-(2-(2,6-dichlorophenyl)-1-(3'-methyl-5'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
107		2-(1-(2'-chloro-5'-(methylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-1H-imidazol-4-yl)propan-2-ol;
108		4-(2-(2,6-dichlorophenyl)-4-(2-hydroxypropan-2-yl)-1H-imidazol-1-yl)-N-methylbiphenyl-3-sulfonamide;
109		2-(1-(3-fluoro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2-(trifluoromethyl)phenyl)-1H-imidazol-4-yl)propan-2-ol;
110		2-(1-(3'-(ethylsulfonyl)-3-fluorobiphenyl-4-yl)-2-(2-(trifluoromethyl)phenyl)-1H-imidazol-4-yl)propan-2-ol;
111		2-(1-(2-fluoro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2-(trifluoromethyl)phenyl)-1H-imidazol-4-yl)propan-2-ol;
112		2-(1-(3'-(ethylsulfonyl)-2-fluorobiphenyl-4-yl)-2-(2-(trifluoromethyl)phenyl)-1H-imidazol-4-yl)propan-2-ol;
113		2-(2-(2,3-dichlorophenyl)-1-(3-fluoro-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;

Cpd #	Structure	Name
114		2-(2-(2,3-dichlorophenyl)-1-(3'-(ethylsulfonyl)-3-fluorobiphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
115		2-(2-(2,6-dichlorophenyl)-1-(3-fluoro-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
116		2-(2-(2,6-dichlorophenyl)-1-(3'-(ethylsulfonyl)-3-fluorobiphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
117		2-(2-(2,3-dichlorophenyl)-1-(3-fluoro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
118		2-(2-(3-chloro-2-methylphenyl)-1-(3-fluoro-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
119		2-(2-(3-chloro-2-methylphenyl)-1-(3'-(ethylsulfonyl)-3-fluorobiphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
120		2-(2-(3-chloro-2-methylphenyl)-1-(3-fluoro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
121		2-(1-(3-fluoro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2-(trifluoromethyl)phenyl)-1H-imidazol-4-yl)propan-2-ol;

Cpd #	Structure	Name
122		2-(2-(2,6-dichlorophenyl)-1-(3-fluoro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
123		2-(2-(2-chloro-3-fluorophenyl)-1-(3-fluoro-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
124		2-(2-(2-chloro-3-fluorophenyl)-1-(3'-(ethylsulfonyl)-3-fluorobiphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
125		2-(2-(2-chloro-3-fluorophenyl)-1-(3-fluoro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
126		2-(2-(2-chloro-6-fluorophenyl)-1-(3'-(ethylsulfonyl)-3-fluorobiphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
127		2-(2-(2-chloro-6-fluorophenyl)-1-(3-fluoro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
128		2-(2-(2-chloro-6-methylphenyl)-1-(3-fluoro-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;

Cpd #	Structure	Name
129		2-(2-(2-chloro-6-methylphenyl)-1-(3'-(ethylsulfonyl)-3-fluorobiphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
130		2-(2-(2,6-difluorophenyl)-1-(3-fluoro-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
131		2-(2-(2,6-difluorophenyl)-1-(3'-(ethylsulfonyl)-3-fluorobiphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
132		2-(2-(2,6-difluorophenyl)-1-(3'-(ethylsulfonyl)-3-fluorobiphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
133		2-(2-(2-chloro-6-methylphenyl)-1-(3-fluoro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
134		2-(2-(2-chloro-6-fluorophenyl)-1-(3-fluoro-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
135		2-(1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2-(trifluoromethyl)phenyl)-1H-imidazol-4-yl)propan-2-ol;
136		2-(1-(3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl)-2-(2-(trifluoromethyl)phenyl)-1H-imidazol-4-yl)propan-2-ol;

Cpd #	Structure	Name
137		2-(1-(3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl)-2-(2-chlorophenyl)-1H-imidazol-4-yl)propan-2-ol;
138		2-(1-(3-chloro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2-chlorophenyl)-1H-imidazol-4-yl)propan-2-ol;
139		2-(1-(3-chloro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2-(trifluoromethyl)phenyl)-1H-imidazol-4-yl)propan-2-ol;
140		2-(1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2,3-dichlorophenyl)-1H-imidazol-4-yl)propan-2-ol;
141		2-(1-(3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl)-2-(2,3-dichlorophenyl)-1H-imidazol-4-yl)propan-2-ol;
142		2-(1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-1H-imidazol-4-yl)propan-2-ol;
143		2-(2-(3-chloro-2-methylphenyl)-1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
144		2-(2-(3-chloro-2-methylphenyl)-1-(3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;

Cpd #	Structure	Name
145		2-(1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2-chloro-6-fluorophenyl)-1H-imidazol-4-yl)propan-2-ol;
146		2-(1-(3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl)-2-(2-chloro-6-fluorophenyl)-1H-imidazol-4-yl)propan-2-ol;
147		2-(1-(3-chloro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2-chloro-6-fluorophenyl)-1H-imidazol-4-yl)propan-2-ol;
148		2-(1-(3-chloro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2,3-dichlorophenyl)-1H-imidazol-4-yl)propan-2-ol;
149		2-(1-(3-chloro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-1H-imidazol-4-yl)propan-2-ol;
150		2-(1-(3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-1H-imidazol-4-yl)propan-2-ol;
151		2-(1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2-chloro-3-fluorophenyl)-1H-imidazol-4-yl)propan-2-ol;
152		2-(1-(3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl)-2-(2-chloro-3-fluorophenyl)-1H-imidazol-4-yl)propan-2-ol;

Cpd #	Structure	Name
153		2-(2-(2-chloro-3-fluorophenyl)-1-(3-chloro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
154		2-(1-(3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl)-2-(2-chloro-6-methylphenyl)-1H-imidazol-4-yl)propan-2-ol;
155		2-(1-(3-chloro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2-chloro-6-methylphenyl)-1H-imidazol-4-yl)propan-2-ol;
156		2-(1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2-chloro-6-methylphenyl)-1H-imidazol-4-yl)propan-2-ol;
157		2-(1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2,6-difluorophenyl)-1H-imidazol-4-yl)propan-2-ol;
158		2-(1-(3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl)-2-(2,6-difluorophenyl)-1H-imidazol-4-yl)propan-2-ol;
159		2-(1-(3-chloro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2,6-difluorophenyl)-1H-imidazol-4-yl)propan-2-ol;

Cpd #	Structure	Name
160		2-(2-(3-chloro-2-methylphenyl)-1-(3-chloro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
161		2-(1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2-isopropylphenyl)-1H-imidazol-4-yl)propan-2-ol;
162		2-(3'-chloro-4'-(2-(2-chlorophenyl)-4-(2-hydroxypropan-2-yl)-1H-imidazol-1-yl)-3-(methylsulfonyl)biphenyl-4-yl)propan-2-ol;
163		2-(1-(3-chloro-4'-(2-hydroxypropan-2-yl)-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2-(trifluoromethyl)phenyl)-1H-imidazol-4-yl)propan-2-ol;
164		2-(3'-chloro-4'-(2-(2-chloro-6-fluorophenyl)-4-(2-hydroxypropan-2-yl)-1H-imidazol-1-yl)-3-(methylsulfonyl)biphenyl-4-yl)propan-2-ol;
165		ethyl 3'-chloro-4'-(2-(2-chloro-6-fluorophenyl)-4-(2-hydroxypropan-2-yl)-1H-imidazol-1-yl)-3-(methylsulfonyl)biphenyl-4-carboxylate;
166		(3'-chloro-4'-(2-(2-chloro-6-fluorophenyl)-4-(2-hydroxypropan-2-yl)-1H-imidazol-1-yl)-3-(methylsulfonyl)biphenyl-4-yl)morpholine methanone;

Cpd #	Structure	Name
167		3'-chloro-4'-(2-(2-chloro-6-fluorophenyl)-4-(2-hydroxypropan-2-yl)-1H-imidazol-1-yl)-N,N-dimethyl-3-(methylsulfonyl)biphenyl-4-carboxamide;
168		2-(1-(3-chloro-5-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2-chlorophenyl)-1H-imidazol-4-yl)propan-2-ol;
169		2-(1-(3-chloro-3'-(ethylsulfonyl)-5-methylbiphenyl-4-yl)-2-(2-chlorophenyl)-1H-imidazol-4-yl)propan-2-ol;
170		2-(2-(2-chlorophenyl)-1-(3-ethyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
171		2-(1-(4'-(hydroxymethyl)-3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2-(trifluoromethyl)phenyl)-1H-imidazol-4-yl)propan-2-ol;
172		2-(2-(2-chloro-6-fluorophenyl)-1-(4'-(hydroxymethyl)-3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
173		2-(2-(2-chloro-3-fluorophenyl)-1-(4'-(hydroxymethyl)-3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;

Cpd #	Structure	Name
174		2-(2-(2,6-dichlorophenyl)-1-(4'-(hydroxymethyl)-3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
175		2-(2-(3-chloro-2-methylphenyl)-1-(4'-(hydroxymethyl)-3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
176		2-(2-(2,3-dichlorophenyl)-1-(4'-(hydroxymethyl)-3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
177		2-(2-(2-chloro-6-methylphenyl)-1-(4'-(hydroxymethyl)-3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
178		2-(2-(2,6-difluorophenyl)-1-(4'-(hydroxymethyl)-3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
179		2-(2-(2,3-dichlorophenyl)-1-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
180		2-(2-(2,3-dichlorophenyl)-1-(3'-(ethylsulfonyl)-3-methylbiphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;

Cpd #	Structure	Name
181		2-(2-(2,6-dichlorophenyl)-1-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
182		2-(2-(2,6-dichlorophenyl)-1-(3'-(ethylsulfonyl)-3-methylbiphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
183		2-(1-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2-(trifluoromethyl)phenyl)-1H-imidazol-4-yl)propan-2-ol;
184		2-(1-(3'-(ethylsulfonyl)-3-methylbiphenyl-4-yl)-2-(2-(trifluoromethyl)phenyl)-1H-imidazol-4-yl)propan-2-ol;
185		2-(2-(2-chloro-3-fluorophenyl)-1-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
186		2-(2-(2-chloro-3-fluorophenyl)-1-(3'-(ethylsulfonyl)-3-methylbiphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
187		2-(2-(2-chloro-6-fluorophenyl)-1-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
188		2-(2-(2-chloro-6-fluorophenyl)-1-(3'-(ethylsulfonyl)-3-methylbiphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;

Cpd #	Structure	Name
189		2-(2-(3-chloro-2-methylphenyl)-1-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
190		2-(2-(3-chloro-2-methylphenyl)-1-(3'-(ethylsulfonyl)-3-methylbiphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
191		2-(2-(2-chloro-6-methylphenyl)-1-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
192		2-(2-(2-chloro-6-methylphenyl)-1-(3'-(ethylsulfonyl)-3-methylbiphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
193		2-(2-(2,6-difluorophenyl)-1-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
194		2-(2-(2,6-difluorophenyl)-1-(3'-(ethylsulfonyl)-3-methylbiphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
195		2-(2-(2-isopropylphenyl)-1-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
196		2-(1-(3'-(ethylsulfonyl)-3-methylbiphenyl-4-yl)-2-(2-isopropylphenyl)-1H-imidazol-4-yl)propan-2-ol;

Cpd #	Structure	Name
197		2-(1-(2,6-dichlorophenyl)-2-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
198		2-{1-[3'-(methylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
199		2-{1-[4'-(methylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
200		2-{1-[3'-(ethylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
201		2-{1-[3'-(methylsulfonyl)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
202		2-{1-[2'-(trifluoromethyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
203		2-{1-[2'-(difluorobiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
204		2-(1-{4'-[(1-methylethyl)oxy]biphenyl-4-yl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl)propan-2-ol;
205		2-{1-[4'-fluoro-3'-methylbiphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;

Cpd #	Structure	Name
206		2-{1-(3'-fluorobiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
207		2-{1-(2',4',5'-trifluorobiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
208		2-{1-[5'-fluoro-2'-(methoxy)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
209		2-{1-[4'-chlorobiphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
210		2-{1-[4-pyrimidin-5-ylphenyl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
211		2-{1-[4'-(hydroxymethyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
212		2-{1-[4'-(dimethylamino)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
213		2-{1-[4'-(trifluoromethyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
214		2-{1-[4'-(1-methylethyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;

Cpd #	Structure	Name
215		2-{1-[4'-chloro-2'-(trifluoromethyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
216		2-{1-(2',3',4'-trifluorobiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
217		2-{1-(3',4'-difluorobiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
218		2-{1-[2'-chloro-6'-fluoro-3'-methylbiphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
219		2-{1-[5'-chloro-2'-(methoxy)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
220		2-{1-[2'-fluoro-5'-(trifluoromethyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
221		2-{1-[2'-(methylthio)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
222		4'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-4-carboxylic acid;
223		2-{1-[4-(1,3-benzodioxol-5-yl)phenyl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;

Cpd #	Structure	Name
224		2-(1-{4-[6-(methoxy)pyridin-3-yl]phenyl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
225		2-(1-{4-[(1E)-3,3-dimethylbut-1-en-1-yl]phenyl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
226		2-{1-(3'-chlorobiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
227		2-{1-[2'-fluoro-5'-(methoxy)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
228		ethyl 4'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-4-carboxylate;
229		2-{1-(4-[2-(1-methylethyl)oxy]pyridin-3-yl]phenyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
230		2-{1-[3'-chloro-4'-(methoxy)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
232		2-{1-[2'-fluoro-3'-(methoxy)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
233		4'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-3-carboxamide;

Cpd #	Structure	Name
234		4'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}biphenyl-4-carboxamide;
235		2-(1-{4'-[(trifluoromethyl)oxy]biphenyl-4-yl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
236		2-{1-[4'-fluoro-3'-(trifluoromethyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
237		2-{1-(4'-propylbiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
238		2-{1-[4'-(ethyloxy)-3'-(trifluoromethyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
239		2-(1-{2'-[(1-methylethyl)oxy]biphenyl-4-yl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
240		2-(1-{3'-chloro-4'-[(1-methylethyl)oxy]biphenyl-4-yl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
241		2-{1-[4-(1H-indol-4-yl)phenyl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
242		2-{1-[4'-(methylthio)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;

Cpd #	Structure	Name
243		2-{1-[3'-(hydroxymethyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
244		2-{1-[3'-(ethyloxy)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
245		2-{1-(4'-ethylbiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
246		2-{1-(2',4'-difluorobiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
247		2-{1-(3',4'-dichlorobiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
248		2-{1-[2'-chloro-4'-(trifluoromethyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
249		2-{1-(4-naphthalen-2-ylphenyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
250		2-{1-[3'-(methyloxy)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
251		4'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-3-ol;
252		2-{1-(3',4',5'-trifluorobiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;

Cpd #	Structure	Name
253		1-[5-(4-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}phenyl)-2-thienyl]ethanone;
254		2-{1-(3',5'-difluorobiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
255		2-{1-(3'-chloro-4'-fluorobiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
256		2-{1-[5'-methyl-2'-(methyloxy)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
257		2-{1-(2',5'-difluorobiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
258		2-{1-[3'-(butyloxy)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
259		2-{1-[5'-chloro-2'-(ethyloxy)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
260		2-{1-[3'-[(trifluoromethyl)oxy]biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
261		2-{1-(2',3',5'-trifluorobiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;

Cpd #	Structure	Name
262		2-{1-[3'-(ethylthio)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
263		2-{1-[3'-(1-methylethoxy)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
264		2-{1-[4-(1-benzothien-3-yl)phenyl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
265		2-{1-[4-(4-methylnaphthalen-1-yl)phenyl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
266		2-{1-(2',4'-dichlorobiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
267		2-{1-[3',4'-bis(methyloxy)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
268		2-{2-[2-(trifluoromethyl)phenyl]-1-(2',4',5'-trimethylbiphenyl-4-yl)-1H-imidazol-4-yl}propan-2-ol;
269		4-fluoro-4'-[4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl]biphenyl-2-ol;
270		2-{1-[2'-(hydroxymethyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;

Cpd #	Structure	Name
271		2-[1-[3'-(trifluoromethyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl]propan-2-ol;
272		2-[1-(2'-chloro-6'-fluorobiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl]propan-2-ol;
273		2-[1-[3',5'-difluoro-2-(methoxy)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl]propan-2-ol;
274		2-[1-[4-[2-(methoxy)pyridin-3-yl]phenyl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl]propan-2-ol;
275		2-[1-[2'-methyl-5'-(methoxy)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl]propan-2-ol;
276		2-[1-(2'-ethylbiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl]propan-2-ol;
277		2-[1-(2'-methyl-4'-[(1-methylethyl)oxy]biphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl]propan-2-ol;
278		2-[1-[4'-(ethylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl]propan-2-ol;
279		2-[1-(5'-fluoro-2'-methylbiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl]propan-2-ol;

Cpd #	Structure	Name
280		2-[1-[3'-chloro-4'-(trifluoromethyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl]propan-2-ol;
281		2-[1-(5'-chloro-2'-methylbiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl]propan-2-ol;
282		2-[1-[2'-(ethyloxy)-5'-(trifluoromethyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl]propan-2-ol;
283		2-[1-(3'-fluoro-4'-methylbiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl]propan-2-ol;
284		methyl (2E)-3-(4'-(4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl)biphenyl-4-yl)prop-2-enoate;
285		N-ethyl-4'-(4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl)biphenyl-3-carboxamide;
286		2-[1-[4-(2,3-dihydro-1-benzofuran-5-yl)phenyl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl]propan-2-ol;
287		2-[1-[3-[6-(methoxy)pyridin-3-yl]phenyl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl]propan-2-ol;
288		3'-(4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl)-N-(1-methylethyl)biphenyl-3-sulfonamide;

Cpd #	Structure	Name
289		3'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-4-carboxamide;
290		2-(1-{3-[2-(cyclopentyloxy)pyridin-3-yl]phenyl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
291		2-{1-(3'-chlorobiphenyl-3-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
292		2-{1-[2'-fluoro-5-(methyloxy)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
293		3'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-3-carboxamide;
294		2-{1-[3',4'-bis(methyloxy)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
295		N-(3'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-2-yl)methanesulfonamide;
296		2-{1-[3'-(trifluoromethyl)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
297		3-fluoro-3'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-4-carboxylic acid;

Cpd #	Structure	Name
298		4-chloro-3'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-3-carboxamide;
299		2-{1-(3-[2-((1-methylethyl)oxy)pyridin-3-yl]phenyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
300		N-(3'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-3-yl)acetamide;
301		2-{1-[2'-methyl-5-(methyloxy)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
302		N-(3'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-4-yl)acetamide;
303		2-{1-(4'-chlorobiphenyl-3-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
304		2-{1-[4'-(phenyloxy)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
305		2-{1-[5'-fluoro-2-(methyloxy)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
306		2-{1-(2',3'-dichlorobiphenyl-3-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;

Cpd #	Structure	Name
307		2-(1-{3'-[(trifluoromethyl)oxy]biphenyl-3-yl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
308		3'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-3-sulfonamide;
309		2-{1-[3',5'-bis(trifluoromethyl)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
310		2-{1-(3',5'-dichlorobiphenyl-3-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
311		3-chloro-3'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}-N-(1-methylethyl)biphenyl-4-carboxamide;
312		N,N-diethyl-3'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-3-carboxamide;
313		4-chloro-3'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}-N-(1-methylethyl)biphenyl-3-carboxamide;
314		2-{1-[3'-(methyloxy)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;

Cpd #	Structure	Name
315		3'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-3-carboxylic acid;
316		N-ethyl-3'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-3-carboxamide;
317		4-chloro-N-ethyl-3'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-3-carboxamide;
318		2-{1-(2',5'-difluorobiphenyl-3-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
319		2-(1-{3'-[(1-methylethyl)oxy]biphenyl-3-yl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
320		2-{1-[2'-fluoro-5'-(trifluoromethyl)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
321		2-{1-[3',4'-dichlorobiphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
322		2-{1-[3'-(ethyloxy)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
323		2-(1-{2'-methyl-4'-[(1-methylethyl)oxy]biphenyl-3-yl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;

Cpd #	Structure	Name
324		2-{1-[3-(1-methyl-1H-indol-5-yl)phenyl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
325		2-{1-[4'-(ethoxy)-3'-(trifluoromethyl)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
326		2-{1-[3'-(ethylsulfonyl)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
327		2-{1-[2'-(trifluoromethyl)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
328		2-{1-[3'-(hydroxymethyl)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
329		2-{1-[3-(1H-indol-4-yl)phenyl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
330		3'-(4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl)biphenyl-4-carboxylic acid;
331		1-[5-(3-(4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl)phenyl)-2-thienyl]ethanone;
332		2-{1-[5'-(chloro)-2'-methylbiphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;

Cpd #	Structure	Name
333		2-(1-{4'-[(trifluoromethyl)oxy]biphenyl-3-yl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
334		2-{1-[2'-chloro-4'-(trifluoromethyl)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
335		2-{1-[2,5'-dichlorobiphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
336		2-{1-[2'-(ethoxy)-5'-(trifluoromethyl)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
337		3'-(4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl)-N-(1-methylethyl)biphenyl-4-carboxamide;
338		2-{1-[2,4'-dichlorobiphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
339		2-{1-[4'-(ethylsulfonyl)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
340		2-{1-[4'-fluoro-3'-(trifluoromethyl)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
341		2-{1-[3'-fluoro-4'-methylbiphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;

Cpd #	Structure	Name
342		N-butyl-4'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-3-sulfonamide;
343		N-(1,1-dimethylethyl)-4'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}-6-methylbiphenyl-3-sulfonamide;
344		4'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}-N-methylbiphenyl-3-sulfonamide;
345		N-ethyl-4'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-3-sulfonamide;
346		N-(1,1-dimethylethyl)-4'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-3-sulfonamide;
347		2-{1-[2-amino-5'-(trifluoromethyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
348		2-{1-[3'-fluoro-5'-(trifluoromethyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
349		2-{1-[4'-chloro-3'-(trifluoromethyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;

Cpd #	Structure	Name
350		3-chloro-4'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-4-carboxylic acid;
351		2-(2-[2-(trifluoromethyl)phenyl]-1-{3'[(trifluoromethyl)thio]biphenyl-4-yl}-1H-imidazol-4-yl)propan-2-ol;
352		2-{1-[3'-(morpholin-4-yl)sulfonyl]biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
353		2-(1-{4-[5-(hydroxymethyl)-1,3-thiazol-2-yl]phenyl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl)propan-2-ol;
354		2-{1-[2'-methyl-5'-(piperidin-1-yl)sulfonyl]biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
355		1-[4-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}phenyl]-2-thienyl]ethanone;
356		2-(1-{4-[5-(hydroxymethyl)-3-thienyl]phenyl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl)propan-2-ol;
357		4-(2-chlorophenyl)-5-{5-[3-(methylsulfonyl)phenyl]-2-thienyl}-2-(trifluoromethyl)-1H-imidazole;

Cpd #	Structure	Name
358		2-(2-chlorophenyl)-1-[3'-(methylsulfonyl)biphenyl-4-yl]-4-(trifluoromethyl)-1H-imidazole;
359		4-(2-chlorophenyl)-5-[3'-(methylsulfonyl)biphenyl-4-yl]-2-(trifluoromethyl)-1H-imidazole;
360		4'-[2-(2-chlorophenyl)-4-(trifluoromethyl)-1H-imidazol-1-yl]biphenyl-3-sulfonamide;
361		4'-[4-(2-chlorophenyl)-2-(trifluoromethyl)-1H-imidazol-5-yl]biphenyl-3-sulfonamide;
362		3-[5-[4-(2-chlorophenyl)-2-(trifluoromethyl)-1H-imidazol-5-yl]-2-thienyl]benzenesulfonamide;
364		1-[3'-(methylsulfonyl)biphenyl-4-yl]-4-(trifluoromethyl)-2-(trifluoromethyl)phenyl-1H-imidazole;
365		4'-[4-(trifluoromethyl)-2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl]biphenyl-3-sulfonamide;
366		ethyl 1-(4-bromophenyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazole-4-carboxylate;
367		ethyl 1-[3'-(methylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazole-4-carboxylate;

Cpd #	Structure	Name
369		5-[3-(methylsulfonyl)phenyl]-2-[4-(trifluoromethyl)-2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl]pyridine;
370		2-[2-(2-chlorophenyl)-4-(trifluoromethyl)-1H-imidazol-1-yl]-5-[3-(methylsulfonyl)phenyl]pyridine;
371		2-[1-[3'-(methylsulfonyl)biphenyl-4-yl]-2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl]propan-2-ol;
372		5-[3-[(1-methylethyl)sulfonyl]phenyl]-2-[4-(trifluoromethyl)-2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl]pyridine;
373		1-[3'-(1-methylethyl)sulfonyl]biphenyl-4-yl]-4-(trifluoromethyl)-2-(trifluoromethyl)phenyl-1H-imidazole;
375		N-(4'-[4-(trifluoromethyl)-2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl]biphenyl-3-methanesulfonamide;
376		ethyl 1-[3'-(methylsulfonyl)biphenyl-4-yl]-2-phenyl-1H-imidazole-4-carboxylate;
378		2-[1-[3'-(methylsulfonyl)biphenyl-4-yl]-2-phenyl-1H-imidazol-4-yl]propan-2-ol;

Cpd #	Structure	Name
379		2-{2-(2-chlorophenyl)-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol;
380		2-{2-(2-fluorophenyl)-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol;
381		1-{2-(2-fluorophenyl)-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}ethanone;
382		2-{2-(2,3-dichlorophenyl)-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol;
383		2-{2-(2-chloro-3-fluorophenyl)-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol;
384		4'-[2-(2-chlorophenyl)-4-(1-hydroxy-1-methylethyl)-1H-imidazol-1-yl]biphenyl-3-sulfonamide;
385		2-{2-(4-fluorophenyl)-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol;
386		2-{2-(2-chlorophenyl)-1-[3'-fluoro-3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol;
387		2-{2-(2-chlorophenyl)-1-[2-fluoro-3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol;

Cpd #	Structure	Name
388		2-{2-(2-chlorophenyl)-1-[3'-(1-methylethyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol;
389		2-{2-(2-chlorophenyl)-1-[4'-(1,1-dimethylethyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol;
390		methyl 4'-[2-(2-chlorophenyl)-4-(1-hydroxy-1-methylethyl)-1H-imidazol-1-yl]biphenyl-3-carboxylate;
391		N-[4'-[2-(2-chlorophenyl)-4-(1-hydroxy-1-methylethyl)-1H-imidazol-1-yl]biphenyl-3-yl]-1,3-benzodioxole-5-carboxamide;
392		2-{1-[3'-(methylsulfonyl)biphenyl-4-yl]-2-[3-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
393		2-{2-(3-chlorophenyl)-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol;
394		2-{2-(2-chloro-6-methylphenyl)-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol;
395		4'-[2-(2-chlorophenyl)-4-(1-hydroxy-1-methylethyl)-1H-imidazol-1-yl]biphenyl-3-carboxylic acid;

Cpd #	Structure	Name
397		2-(2-chlorophenyl)-1-[3'-(methylsulfonyl)biphenyl-4-yl]-N-(2,2,2-trifluoroethyl)-1H-imidazole-4-carboxamide;
398		2-(2-chlorophenyl)-1-[3'-(methylsulfonyl)biphenyl-4-yl]-N-(2,2,2-trifluoroethyl)-1H-imidazole-4-carbothioamide;
399		2-{1-[4'-(methoxy)-3'-(methylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
400		2-{1-[4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
401		ethyl 4'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}-5-(methylsulfonyl)biphenyl-3-carboxylate;
402		2-{5-bromo-1-[3'-(methylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
403		2-{5-chloro-1-[3'-(methylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
404		2-{1-[4'-(1-hydroxy-1-methylethyl)sulfonyl]biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;








Cpd #	Structure	Name
405		2-[1-(3'-aminobiphenyl-4-yl)-2-(2-chlorophenyl)-1H-imidazol-4-yl]propan-2-ol;
406		2-{5-fluoro-1-[3'-(methylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
407		2-{1-[4'-(1-hydroxy-1-methylethyl)sulfonyl]biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
408		1-{2-(2-chlorophenyl)-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}ethanone oxime;
409		2-{2-(2-chlorophenyl)-1-[3'-(1-hydroxy-1-methylethyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol;
410		2-{2-(2,6-difluorophenyl)-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol;
411		1-{1-[3'-(methylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}ethanone;
412		1-{1-[3'-(methylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}ethanone oxime;
413		2-{1-[3'-(1-methylethyl)sulfonyl]biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;









Cpd #	Structure	Name
414		2-(1-{3'-[(1-methylethyl)sulfonyl]biphenyl-3-yl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl)propan-2-ol;
415		2-(2-(2-chlorophenyl)-1-[3'-(ethylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl)propan-2-ol;
416		2-(2-(2-chloro-6-fluorophenyl)-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl)propan-2-ol;
417		2-(1-[3-chloro-3'-(methylsulfonyl)biphenyl-4-yl]-2-(2-chlorophenyl)-1H-imidazol-4-yl)propan-2-ol;
418		2-(2-(2-chlorophenyl)-1-[3'-methyl-3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl)propan-2-ol;
419		2-(2-(2,6-difluorophenyl)-1-[3'-(ethylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl)propan-2-ol;
420		2-(1-{5-[3-(methylsulfonyl)phenyl]pyridin-2-yl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl)propan-2-ol;
421		2-(1-{6-[3-(methylsulfonyl)phenyl]pyridin-3-yl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl)propan-2-ol;
422		2-(2-(2,6-dichlorophenyl)-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl)propan-2-ol;

Cpd #	Structure	Name
424		1-{2-(2-chlorophenyl)-1-[3'-(ethylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}ethanone oxime;
425		1-{2-(2-chlorophenyl)-1-[3'-(ethylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}ethanone oxime;
426		1-{2-(2-chlorophenyl)-1-[3'-(1-methylethyl)biphenyl-4-yl]-1H-imidazol-4-yl}ethanone oxime;
427		2-(2-(2,3-dichlorophenyl)-1-[3'-(ethylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl)propan-2-ol;
428		2-(5-methyl-1-[3'-(methylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl)propan-2-ol;
429		2-(2-(2-chlorophenyl)-1-[3'-(ethylsulfonyl)-3-fluorobiphenyl-4-yl]-1H-imidazol-4-yl)propan-2-ol;
430		2-(2-[2-fluoro-6-(trifluoromethyl)phenyl]-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl)propan-2-ol;
431		2-(2-(2-methylphenyl)-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl)propan-2-ol;
432		2-(1-[3'-(methylsulfonyl)biphenyl-4-yl]-2-[2-[(trifluoromethyl)oxy]phenyl]-1H-imidazol-4-yl)propan-2-ol;

Cpd #	Structure	Name
433		1-{2-(2-chlorophenyl)-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}ethanone oxime;
434		1-{2-(2-chlorophenyl)-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}ethanone oxime;
435		1-{2-(2-chlorophenyl)-1-[4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}ethanone oxime;
436		2-(1-(2',5'-dimethylbiphenyl-4-yl)-2-(2-trifluoromethyl)phenyl)-1H-imidazol-4-yl}propan-2-ol;
437		2-{2-(2-chlorophenyl)-1-[3'-(ethylsulfonyl)-2-fluorobiphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol;
438		2-{2-(3-chloro-2-methylphenyl)-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol;
439		2-{2-(2-chlorophenyl)-1-[2-methyl-3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol;
440		2-{2-(3-chloro-2-methylphenyl)-1-[3'-(ethylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol;
443		2-(1-(2-chlorobenzyl)-2-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol

Cpd #	Structure	Name
444		2-(1-(2,3-dichlorobenzyl)-2-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol
445		2-(5-chloro-2-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-1-(2,6-dichlorophenyl)-1H-imidazol-4-yl)propan-2-ol
446		2-(5-chloro-1-(2,6-dichlorophenyl)-2-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol
447		3'-chloro-4'-(2-(2-chloro-6-fluorophenyl)-4-(2-hydroxypropan-2-yl)-1H-imidazol-1-yl)-3-(methylsulfonyl)biphenyl-4-carboxylic acid
448		3'-chloro-4'-(2-(2-chloro-6-fluorophenyl)-4-(2-hydroxypropan-2-yl)-1H-imidazol-1-yl)-3-methoxy-N-methyl-3-(methylsulfonyl)biphenyl-4-carboxamide
449		2-(2-(2,3-dichlorobenzyl)-1-(3-fluoro-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol
450		2-(2-(2,3-dichlorobenzyl)-1-(3'-(ethylsulfonyl)-3-fluorobiphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol

Cpd #	Structure	Name
451		2-(1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2-(2,3-dichlorophenyl)propan-2-yl)-1H-imidazol-4-yl)propan-2-ol
452		2-(1-(3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl)-2-(2-(2,3-dichlorophenyl)propan-2-yl)-1H-imidazol-4-yl)propan-2-ol
453		3'-chloro-4'-(2-(2-chloro-6-fluorophenyl)-4-(2-hydroxypropan-2-yl)-1H-imidazol-1-yl)-3-(methylsulfonyl)biphenyl-4-carboxamide
454		2-(1-(3-chloro-4'-methoxy-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2-chloro-6-fluorophenyl)-1H-imidazol-4-yl)propan-2-ol
455		2-(2-(2,3-dichlorophenyl)-1-(4'-methoxy-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol
456		2-(2-(2,3-dichlorobenzyl)-1-(4'-methoxy-3'-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol
457		2-(1-(3-chloro-4'-methoxy-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2,3-dichlorobenzyl)-1H-imidazol-4-yl)propan-2-ol

Cpd #	Structure	Name
458		2-(2-(3-chloro-3'-(methylsulfonyl)biphenyl-2-yl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol
459		1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-N-(2,2,2-trifluoroethyl)-1H-imidazole-4-carboxamide
460		1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-N-(2-fluoroethyl)-1H-imidazole-4-carboxamide
461		2-(1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-1H-imidazole-4-carboxamido)acetic acid
462		1-(3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-N-(2,2,2-trifluoroethyl)-1H-imidazole-4-carboxamide
463		1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-N-(1-hydroxy-2-methylpropan-2-yl)-1H-imidazole-4-carboxamide
464		1-(3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-N-(1-hydroxy-2-methylpropan-2-yl)-1H-imidazole-4-carboxamide
465		N-tert-butyl-1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-1H-imidazole-4-carboxamide

Cpd #	Structure	Name
466		2-chloro-4'-(2-(2,6-dichlorophenyl)-4-(2-hydroxypropan-2-yl)-1H-imidazol-1-yl)-3'-methylbiphenyl-4-carboxylic acid
467		2-(1-(2',3-dichloro-5'-(methylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-1H-imidazol-4-yl)propan-2-ol
468		2-(1-(2'-chloro-3-methyl-5'-(methylsulfonyl)biphenyl-4-yl)-2-(2,3-dichlorobenzyl)-1H-imidazol-4-yl)propan-2-ol
469		6-(3-chloro-4-[2-(2-chloro-6-fluorophenyl)-4-(1-hydroxy-1-methylethyl)-1H-imidazol-1-yl]phenyl)-2,3-dihydro-1-benzothiophene-3-ol 1,1-dioxide
470		1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-N-(2-hydroxyethyl)-1H-imidazole-4-carboxamide
471		1-(3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-N-(2-hydroxyethyl)-1H-imidazole-4-carboxamide
472		1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-N-(2-hydroxy-2-methylpropyl)-1H-imidazole-4-carboxamide

Cpd #	Structure	Name
473		1-(3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-N-(2-hydroxy-2-methylpropyl)-1H-imidazole-4-carboxamide
474		2-(2-(5-chloro-2-fluorobenzyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol
475		2-(2-(2,3-dichlorophenyl)-1-(4'-(hydroxymethyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol
476		2-(2-(2,3-dichlorophenyl)-1-(3'-(hydroxymethyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol
477		2-(2-(2,3-dichlorophenyl)-1-(4'-(2-hydroxypropan-2-yl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol
478		2-(2-(2,3-dichlorophenyl)-1-(3'-(2-hydroxypropan-2-yl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol
479		2-(2-(2,6-dichlorophenyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)-2-methylpropan-1-ol

Cpd #	Structure	Name
480		1-(3-chloro-3'-(2-hydroxypropan-2-yl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-N-(1-hydroxy-2-methylpropan-2-yl)-1H-imidazole-4-carboxamide
481		2-(2,6-dichlorophenyl)-N-(1-hydroxy-2-methylpropan-2-yl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazole-4-carboxamide
482		2-(2-(5-chloro-2-fluorobenzyl)-1-(3'-(ethylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol
483		2-(2-(3-chloro-2-fluorobenzyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol

Cpd #	Structure	Name
484		2-(1-(3'-(methylsulfonyl)biphenyl-4-yl)-2-(3-(trifluoromethyl)benzyl)-1H-imidazol-4-yl)propan-2-ol
485		2-(1-(3'-(ethylsulfonyl)biphenyl-4-yl)-2-(3-(trifluoromethyl)benzyl)-1H-imidazol-4-yl)propan-2-ol
486		2-(2,6-dichlorophenyl)-1-(3'-(ethylsulfonyl)biphenyl-4-yl)-N-(1-hydroxy-2-methylpropan-2-yl)-1H-imidazole-4-carboxamide
487		N-tert-butyl-2-(2,6-dichlorophenyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazole-4-carboxamide

Table 2

363		1-(4-bromophenyl)-4-(trifluoromethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazole;
368		5-bromo-2-{4-(trifluoromethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}pyridine;
374		2-{1-(4-bromophenyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
377		ethyl 1-(4-bromophenyl)-2-(2-chlorophenyl)-1H-imidazole-4-carboxylate;
396		1-(4-bromophenyl)-2-(2-chlorophenyl)-N-(2,2,2-trifluoroethyl)-1H-imidazole-4-carbothioamide;

423	 The structure shows an imidazole ring with a 4-bromophenyl group at position 1 and a 2-chlorophenyl group at position 2. At position 4, there is an ethanone oxime group (-CH2-C(=N-OH)CH3).	1-[1-(4-bromophenyl)-2-(2-chlorophenyl)-1H-imidazol-4-yl]ethanone oxime;
441	 The structure shows an imidazole ring with a 5-bromo-2-thienyl group at position 1, a 2,5-dichlorophenyl group at position 2, and a trifluoromethyl group (-CF3) at position 4.	2-(5-bromo-2-thienyl)-1-(2,5-dichlorophenyl)-4-(trifluoromethyl)-1H-imidazole;
442	 The structure shows an imidazole ring with a 5-bromo-2-thienyl group at position 1, a 2-chlorophenyl group at position 2, and a trifluoromethyl group (-CF3) at position 4.	2-(5-bromo-2-thienyl)-1-(2-chlorophenyl)-4-(trifluoromethyl)-1H-imidazole;

Example 31

FRET Coactivator assay

The FRET coactivator assay measures the ability of LXR ligands to promote protein-protein interactions between the ligand binding domain (LBD) of LXR and transcriptional coactivator proteins. The assay involves the use a recombinant Glutathione-S-transferase (GST)-nuclear receptor ligand binding domain (LBD) fusion protein and a synthetic biotinylated peptide sequence derived from the receptor interacting domain of a co-activator peptide such as the steroid receptor coactivator 1 (SRC-1). Typically GST-LBD is labeled with a europium chelate (donor) via a europium-tagged anti-GST antibody, and the coactivator peptide is labeled with allophycocyanin via a streptavidin-biotin linkage.

In the presence of an agonist for the nuclear receptor, the peptide is recruited to the GST-LBD bringing europium and allophycocyanin into close proximity to enable energy transfer from the europium chelate to the allophycocyanin. Upon excitation of the complex with light at 340 nm excitation energy absorbed by the europium chelate is transmitted to the allophycocyanin moiety resulting in emission at 665 nm. If the europium chelate is not brought into close proximity to the allophycocyanin moiety there is little or no energy transfer and excitation of the europium chelate results in emission at 615 nm. Thus the intensity of light emitted at 665 nm gives an indication of the strength of the protein-protein interaction.

Required Materials:

Partially purified recombinant protein comprising glutathione-S-transferase fused in frame to the LXR-ligand binding domain (comprising amino acids 188-447 of human LXR α , or amino acids 198-461 of human LXR β). Biotinylated peptide containing a SRC-1 LXXLL receptor interaction motif (B-SRC-1). Anti-GST antibody conjugated to a Europium chelate (α GST-K) (From Wallace/PE Life Sciences Cat# AD0064). Streptavidin linked allophycocyanin (SA-APC) (From Wallace/PE Life Sciences Cat# AD0059A). 1x FRET Buffer: (20 mM KH₂PO₄/K₂HPO₄ pH 7.3, 150 mM NaCl, 2.5

mM CHAPS, 2 mM EDTA, 1 mM DTT (add fresh)). 96 well or 384 well black multiwell plates (from LJI)

Stock Solutions:

0.5 M $\text{KH}_2\text{PO}_4/\text{K}_2\text{HPO}_4$: pH 7.3; 5 M NaCl; 80 mM (5%) CHAPS; 0.5 M EDTA pH 8.0; 1 M DTT (keep at -20°C)

Preparation of Screening Reagents:

Prepare reaction mixture for the appropriate number of wells by combining the following reagents 5 nM/well GST-hLXR α LBD, 5 nM/well GST-hLXR β LBD, 5 nM/well Anti-GST antibody (Eu), 12 nM/well biotin-SRC-1 peptide, 12 nM/well APC-SA adjust the volume to 10 μL /well with 1x-FRET buffer.

Procedure:

Add 0.5 μL of a 1 mM stock compound (for approx. 10 μM final concentration) or solvent to each well in a 96 well or 384 well black plate (LJI). Add 10 μL reaction mixture (prepared above) to each well of the multiwell plate. Incubate covered or in the dark (the APC is light sensitive) at ambient temperature for 1-4 hours. After this time if reactions are not read they can be stored at 4°C for several more hours without too much loss of signal.

Read the plate using an LJI Analyst, or similar instrument, using the following conditions: Channel 1: Excitation is 330 nm and emission is 615. This is for Eu chelate; Channel 2: Excitation is 330 nm and emission is 665. This is for APC; For channel 1: Flashes per well = 100; Integration time = 1000 μs ; interval between flashes = 1x10 ms; Delay after flash = 200 μs ; For channel 2: Flashes per well = 100; Integration time = 100 μs ; interval between flashes = 1x10 ms; Delay after flashes = 65 μs .

Example 32

Scintillation proximity assay (SPA)

The SPA assay measures the radioactive signal generated by the binding of ^3H -24,25-epoxycholesterol to LXR α or LXR β . The basis of the assay is the use of SPA beads containing a scintillant, such that when binding to the receptor brings the labeled ligand into proximity with the bead, the energy from the label stimulates the scintillant to emit light. The light is measured using a standard microplate scintillation reader. The ability of a ligand to bind to a receptor can be measured by assessing the degree to which the compound can compete off a radiolabelled ligand with known affinity for the receptor.

Required Materials:

Label: ^3H -24,25-epoxy-cholesterol (Amersham)

LXR α lysate: Baculovirus expressed LXR α /RXR heterodimer with RXR having a 6-HIS tag produced as a crude lysate

LXR β lysate: Baculovirus expressed LXR β /RXR heterodimer with RXR having a 6-HIS tag produced as a crude lysate

SPA beads: Ysi copper His-tag SPA beads (Amersham)

Plates: Non-binding surface 96-well plate (Corning)

Protein lysate dilution buffer: (20 mM Tris-HCl pH 7.9, 500 mM NaCl, 5 mM Imidazole). 2x SPA

Buffer: (40 mM K₂HPO₄/KH₂PO₄ pH7.3, 100 mM NaCl, 0.05% Tween 20, 20% Glycerol, 4 mM EDTA) 2x SPA Buffer w/o EDTA: (40 mM K₂HPO₄/KH₂PO₄ pH7.3, 100mM NaCl, 0.05% Tween 20, 20% Glycerol)

Stock Solutions

0.5 M K₂HPO₄/KH₂PO₄ pH 7.3; 0.5 M EDTA pH 8.0; 5 M NaCl; 10% Tween-20; Glycerol

Preparation of protein lysates

Baculovirus expression plasmids for human RXR α (accession No NM_002957) LXR α (accession No U22662), LXR β (accession No U07132) were made by cloning the appropriate full-length cDNAs into the pBacPakhis1 vector (Clontech, CA) following standard procedures. Insertion of the cDNAs into the pBacPakhis1 vector polylinker created an in frame fusion to the cDNA to an N-terminal poly-His tag present in pBacPakhis1. Correct cloning was confirmed by restriction mapping, and /or sequencing.

Cell lysates were prepared by infecting healthy, Sf9 insect cells at a density of approximately 1.25×10^6 /ml at 27°C, in a total volume of 500 mL per 1L sized spinner flasks, cultured under standard conditions. To prepare LXR α lysate, insect cells were co-transfected with the LXR α expression cassette at an M.O.I of 0.5 to 0.8 and with the RXR expression cassette at a M.O.I. of approximately 1.6. To prepare LXR β lysate, insect cells were co-transfected with the LXR β expression cassette at an M.O.I of approximately 1.6 and with the RXR expression cassette at a M.O.I. of approximately 1.6. In both cases cells were incubated for 48 hours at 27°C with constant shaking prior to harvesting.

After incubation, cells were harvested by centrifugation and pelleted. Cell pellets were resuspended in two volumes of ice-cold freshly prepared extraction buffer (20mM Tris pH 8.0, 10mM Imidazole, 400mM NaCl, containing one EDTA free protease inhibitor tablet (Roche Catalog No: 1836170) per 10 ml of extraction buffer). Cells were homogenized slowly on ice using a Douncer to achieve 80-90% cell lysis. The homogenate was centrifuged in a pre-chilled rotor (Ti50 or Ti70, or equivalent) at 45,000 rpm for 30 minutes at 4°C. Aliquots of the supernatant were frozen on dry ice and stored frozen at -80°C until quantification and quality control. Aliquots of the lysates were tested in the

SPA assay to ensure lot to lot consistency, and via SDS-PAGE analysis after purification using Ni-NTA Resin (Qiagen) and adjusted for protein concentration and expression level prior to use in screening assays.

Preparation of Screening Reagents

[³H] 24,25 Epoxycholesterol (EC) solution: For a single 384-well plate (or 400 wells), 21 µL of [³H] EC (specific activity 76.5 Ci/mmol, concentration 3.2 mCi/mL) was added to 4.4 mL of 2x SPA buffer to provide for a final concentration of 200 nM. For each additional 384-well plate, an additional 19.1 µL of [³H] EC was added to 4.0 mL of additional 2x SPA buffer. The final concentration of [³H] EC in the well was 50 nM. LXRα lysate (prepared as above) was diluted with protein lysate dilution buffer. 1400 µL of diluted LXRα lysate was prepared per 384-well plate, (or 200 wells) and 1120 µL of diluted LXRα lysate was prepared for each additional 384-well plate. LXRβ lysate (prepared as above) was diluted with protein lysate dilution buffer. 1400 µL of diluted LXRβ lysate was prepared per 384-well plate, (or 200 wells) and 1120 µL of diluted LXRβ lysate was prepared for each additional 384-well plate. SPA bead solution: For a 384-well plate (or 400 wells), 3.75 mL of 2x SPA buffer w/o EDTA, 2.25 mL of H₂O, and 1.5 mL of Ysi His-tag SPA beads (vortex well before taking) were mixed together. For each additional 384-well plate, an additional 3.5 mL of 2x SPA buffer w/o EDTA, 2.1 mL of H₂O, and 1.4 mL of Ysi His-tag SPA beads were mixed together.

Procedure:

Appropriate dilutions of each compound were prepared and pipetted into the appropriate wells of a multiwell plate. 9.1 µL of [³H] EC was added to each well of column 2-23 of the multiwell plate. 5 µL of diluted LXRα lysate was added to each well of column 2-23 on odd rows of the multiwell plate. 5 µL of diluted LXRβ lysate was added to each well of column 2-23 on even rows of the multiwell plate. 17.5 µL of SPA bead solution was added to each well of column 2-23 of the multiwell plate.

The plates were covered with clear sealer and placed in an incubator at ambient temperature for 1 hour. After incubation plates were analyzed using a luminescent plate reader (MicroBeta, Wallac) using the program n ABASE 3H_384DPM. The setting for n ABASE 3H_384DPM was: Counting Mode: DPM; Sample Type: SPA; ParaLux Mode: low background; Count time: 30 sec.

Assays for LXRα and LXRβ were performed in the identical manner. The determined Ki represents the average of at least two independent dose response experiments. The binding affinity for each compound may be determined by non-linear regression analysis using the one site competition formula to determine the IC₅₀ where:

$$Y = \text{Bottom} + \frac{(\text{Top} - \text{Bottom})}{(1 + 10^{X - \log \text{IC}_{50}})}$$

The K_i is then calculated using the Cheng and Prusoff equation where:

$$K_i = IC_{50} / (1 + [\text{Concentration of Ligand}] / K_d \text{ of Ligand})$$

For this assay, typically the Concentration of Ligand = 50 nM and the K_d of EC for the receptor is 200 nM as determined by saturation binding.

The compounds of the invention demonstrated the ability to bind to LXR α and/or LXR β when tested in this assay.

Example 33

Co-Transfection Assay

To measure the ability of compounds to activate or inhibit the transcriptional activity of LXR in a cell based assay, the co-transfection assay was used. It has been shown that LXR functions as a heterodimer with RXR. For the co-transfection assay, expression plasmids for LXR and RXR are introduced via transient transfection into mammalian cells along with a luciferase reporter plasmid that contains one copy of a DNA sequence that is bound by LXR-RXR heterodimers (*LXRE*; Willy, P. *et.al.* 1995). Treatment of transfected cells with an LXR agonist increases the transcriptional activity of LXR, which is measured by an increase in luciferase activity. Similarly, LXR antagonist activity can be measured by determining the ability of a compound to competitively inhibit the activity of a LXR agonist.

Required Materials

CV-1 African Green Monkey Kidney Cells Co-transfection expression plasmids, comprising full-length LXR α (pCMX-h LXR α), LXR β (pCMX-hLXR β), or RXR α (pCMX-RXR), reporter plasmid (LXREx1-Tk-Luciferase), and control (pCMX-Galactosidase expression vector) (Willey et al. *Genes & Development* 9 1033-1045 (1995)). Transfection reagent such as FuGENE6 (Roche). 1x Cell lysis buffer (1 % Triton X 100 (JT Baker X200-07), 10% Glycerol (JT Baker M778-07), 5 mM Dithiothreitol (Quantum Bioprobe DTT03; add fresh before lysing), 1 mM EGTA (Ethylene Glycol-bis (B-Amino ethyl ether)-N,N,N',N'-Tetracetic Acid) (Sigma E-4378), 25 mM Tricine (ICN 807420) pH 7.8) 1x Luciferase assay buffer (pH at 7.8) (0.73 mM ATP, 22.3 mM Tricine, 0.11 mM EDTA, 33.3 mM DTT) 1x Luciferin/CoA (11 mM Luciferin, 3.05 mM Coenzyme A, 10 mM HEPES)

Preparation of Screening Reagents

CV-1 cells were prepared 24 hours prior to the experiment by plating them into T-175 flasks or 500 cm² dishes in order to achieve 70-80% confluency on the day of the transfection. The number of cells to be transfected was determined by the number of plates to be screened. Each 384 well plate requires 1.92x10⁶ cells or 5000 cells per well. DNA Transfection Reagent was prepared by mixing the

required plasmid DNAs with a cationic lipid transfection reagent FuGENE6 (Roche) by following the instructions provided with the reagents. Optimal DNA amounts were determined empirically per cell line and size of vessel to be transfected. 10-12 mL of media was added to the DNA Transfection Reagent and this mixture was added to the cells after aspirating media from the T175 cm² flask. Cells were then incubated at least 5 hours at 37°C to prepare screening cells.

Luciferase assay reagent was prepared by combining before use (per 10 mL): 10 mL 1x Luciferase assay buffer; 0.54 mL of 1x Luciferrin/CoA; 0.54 mL of 0.2 M Magnesium sulfate

Procedure

Assay plates were prepared by dispensing 5 µL of compound per well of a 384 well plate to achieve final compound concentration of 10 µM and no more than 1% DMSO. Media was removed from the screening cells, the cells trypsinized, harvested cells by centrifugation, counted, and plated at a density of approximately 5000 cells per well in the 384 well assay plate prepared above in a volume of about 45 µL. Assay plates containing both compounds and screening cells (50 µL in total volume) were incubated for 20 hours at 37°C.

After incubation with compounds, media was removed from the cells and lysis buffer (30 µL/well) added. After 30 minutes at ambient temperature, luciferase assay buffer (30 µL/well) was added and the assay plates read on a luminometer (PE Biosystems Northstar reader with on-board injectors, or equivalent). Plates were read immediately after addition of luciferase assay buffer.

The LXR/LXRE co-transfection assay can be used to establish the EC₅₀/IC₅₀ values for potency and percent activity or inhibition for efficacy. Efficacy defines the activity of a compound relative to a high control ((N-(3-((4-fluorophenyl)-(naphthalene-2-sulfonyl)amino)propyl)-2,2-dimethylpropionamide)) or a low control (DMSO/vehicle). The dose response curves are generated from an 8 point curve with concentrations differing by ½ LOG units. Each point represents the average of 4 wells of data from a 384 well plate.

The data from this assay is fitted to the following equation, from the EC₅₀ value may be solved:

$$Y = \text{Bottom} + (\text{Top} - \text{Bottom}) / (1 + 10^{((\log \text{EC}_{50} - X) * \text{HillSlope})})$$

The EC₅₀/IC₅₀ is therefore defined as the concentration at which an agonist or antagonist elicits a response that is half way between the Top (maximum) and Bottom (baseline) values. The EC₅₀/IC₅₀ values represented are the averages of at least 3 independent experiments. The determination of the relative efficacy or % control for an agonist is by comparison to the maximum response achieved by ((N-(3-((4-fluorophenyl)-(naphthalene-2-sulfonyl)-amino)propyl)-2,2-dimethylpropionamide) that is measured individually in each dose response experiment.

For the antagonist assay, a LXR agonist can be added to each well of a 384 well plate to elicit a response. The % inhibition for each antagonist is therefore a measurement of the inhibition of the

activity of the agonist. In this example, 100% inhibition would indicate that the activity of a specific concentration of LXR agonist has been reduced to baseline levels, defined as the activity of the assay in the presence of DMSO only.

Compounds of the invention, when tested in this assay, demonstrated the ability to modulate the activity of LXR α and/or LXR β . Preferably, the active compounds modulate the activity of LXR with a EC₅₀ or IC₅₀ of about 10 μ M or less. More preferably, the EC₅₀ or IC₅₀ of the preferred active compounds is about 1 μ M or less.

Example 34

In vivo Studies

In order to evaluate direct regulation of key target genes by the compounds of the invention, animals are administered a single oral dose of the test compound and tissues collected at various time points. Male C57BL/6 mice (n=8) are dosed by oral gavage with vehicle or compound. At various time points after the dose, animals are bled via the retro orbital sinus for plasma collection. Animals are then euthanized and tissues, such as liver and intestinal mucosa are collected and snap frozen for further analysis. Plasma is analyzed for a lipid parameters, such as total cholesterol, HDL cholesterol and triglyceride levels. RNA is extracted for frozen tissues and can be analyzed by quantitative real time PCR for regulation of key target genes. To identify specificity of target gene regulation by LXR subtypes, LXR deficient mice (LXR α -/- or LXR β -/-) and C57BL/6 wild-type controls are used in this same protocol.

Plasma Lipid Evaluation:

To compare the effects of compounds on plasma cholesterol and triglycerides, animals are dosed with compound for one week and plasma lipid levels are monitored throughout the study. Male C57BL/6 mice (n=8) are dosed daily by oral gavage with vehicle or compound. Plasma samples are taken on day -1 (in order to group animals), day 1, 3, and 7. Samples are collected three hours after the daily dose. On day 7 of the study, following plasma collection, animals are euthanized and tissues, such as liver and intestinal mucosa are collected and snap frozen for further analysis. Plasma is analyzed for lipid parameters, such as total cholesterol, HDL cholesterol and triglyceride levels. RNA is extracted for frozen tissues and can be analyzed by quantitative real time PCR for regulation of key target genes. To identify specificity of target gene regulation by LXR subtypes, LXR deficient mice (LXR α -/- or LXR β -/-) and C57BL/6 wild-type controls are used in this same protocol.

Example 35

Measured EC₅₀ or IC₅₀ for LXR for compounds of the invention

Compounds of the invention, when tested as described in Example 33, demonstrated the ability to modulate the activity of LXR $_{\alpha}$ and/or LXR $_{\beta}$. LXR activities for various compounds of the invention are presented in the following table; those compounds with EC₅₀ or IC₅₀ values < 10 μ M for at least one of LXR $_{\alpha}$ and LXR $_{\beta}$ are considered to be active. In the following Table, IC₅₀ or EC₅₀ data is represented as follows: A = < 1 μ M, B = 1 – 10 μ M, and C = > 10 μ M

Cpd #	activity	44	B	100	B	142	A	186	A	379	A	443	A
		45	B	101	A	143	A	187	A	380	B	444	A
1	A	58	A	102	A	144	A	188	A	381	B	445	A
2	A	59	A	103	B	145	A	189	A	382	A	448	C
3	A	60	A	104	A	146	A	190	A	383	A	449	A
4	A	61	A	105	A	147	A	191	A	384	B	450	A
5	A	62	A	106	B	148	A	192	A	385	B	451	A
6	A	63	A	107	A	149	A	193	A	386	A	452	A
7	A	64	A	108	A	150	A	194	A	387	B	453	B
8	A	65	A	109	A	151	A	195	A	388	B	454	B
9	A	66	A	110	A	152	A	196	A	389	B	455	A
10	A	67	A	111	A	153	A	197	A	390	B	456	A
11	A	68	A	112	A	154	A	198	A	391	B	457	A
12	A	69	A	113	A	155	A	199	B	392	A	458	B
13	A	70	A	114	A	156	A	200	A	393	A	459	A
14	A	71	A	115	A	157	A	201	A	394	A	460	A
15	A	72	A	116	A	158	A	220	A	395	B	461	C
16	A	73	A	117	A	159	A	274	B	396	B	462	A
17	A	74	A	118	A	160	A	281	A	397	A	463	A
18	A	75	A	119	A	161	A	344	A	398	A	464	A
19	A	76	A	120	A	162	A	357	A	399	B	465	A
20	A	77	A	121	A	163	A	358	A	400	A	466	C
21	A	78	A	122	A	164	B	359	A	401	B	467	A
22	A	80	B	123	A	165	B	360	A	402	A	468	A
23	A	82	A	124	A	168	B	361	B	403	A	469	B
24	A	83	B	125	A	169	A	362	A	404	B	470	A
25	A	84	A	126	A	170	A	363	B	405	B	471	A
26	A	85	A	127	B	171	A	364	A	406	A	472	A
27	A	86	A	128	A	172	A	365	A	407	A	473	A
28	A	87	A	129	A	173	A	366	B	408	A	474	A
29	A	88	A	130	B	174	A	367	A	409	B	475	B
30	A	89	A	131	B	175	A	368	B	410	B	476	B
31	A	90	A	132	B	176	A	369	A	411	A	477	B
32	A	91	B	133	A	177	A	370	A	412	A	479	A
33	A	92	B	134	A	178	A	371	A	413	A	480	A
34	A	93	A	135	A	179	A	372	A	415	B	481	A
35	B	94	A	136	A	180	A	373	A	416	B	482	A
36	B	95	A	137	A	181	A	374	B	417	A		
37	C	96	A	138	A	182	A	375	A	418	A		
38	C	97	B	139	A	183	A	376	A	419	B		
41	B	98	A	140	A	184	A	377	B	420	B		
42	A	99	A	141	A	185	A	378	B	421	B		

It is understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be suggested to persons skilled in the art and are to be incorporated within the spirit and purview of this application and scope of


the appended claims. All publications, patents, and patent applications cited herein are hereby incorporated herein by reference for all purposes.

All of the U.S. patents, U.S. patent application publications, U.S. patent applications, foreign patents, foreign patent applications and non-patent publications referred to in this specification and/or listed in the Application Data Sheet, are incorporated herein by reference, in their entirety.

From the foregoing it will be appreciated that, although specific embodiments of the invention have been described herein for purposes of illustration, various modifications may be made without deviating from the spirit and scope of the invention. Accordingly, the invention is not limited except as by the appended claims. This invention also encompasses all combinations of alternative aspects of the invention noted herein. It is understood that any and all embodiments of the present invention may be taken in conjunction with any other embodiment to describe additional embodiments of the present invention. Furthermore, any elements of an embodiment may be combined with any and all other elements from any of the embodiments to describe additional embodiments.

Claims

1. A compound according to the formula,



or a pharmaceutically acceptable salt thereof, wherein

R^1 is $-L^1-R^5$, wherein

L^1 is a bond; and

R^5 is aryl optionally substituted with one or more R^{5a} ; wherein

each R^{5a} is independently halogen, nitro, heteroaryl, heterocyclyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, C_3 - C_8 cycloalkyl, $(C_3$ - C_8 cycloalkyl)- C_1 - C_6 alkyl-, $(C_3$ - C_8 cycloalkenyl)- C_1 - C_6 alkyl-, $(C_3$ - C_8 cycloalkyl)- C_2 - C_6 alkenyl-, aryl, arylalkyl, aryloxy, aryloxyaryl, aryl C_{1-6} alkoxy, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, SO_2R^{11} , OR^{11} , SR^{11} , N_3 , SO_2R^{11} , COR^{11} , $SO_2N(R^{11})_2$, $SO_2NR^{11}COR^{11}$, $C\equiv N$, $C(O)OR^{11}$, $CON(R^{11})_2$, $CON(R^{11})OR^{11}$, $OCON(R^{11})_2$, $NR^{11}COR^{11}$, $R^{11}CON(R^{11})_2$, $NR^{11}COOR^{11}$, or $N(R^{11})_2$, wherein

each R^{5a} is optionally substituted with one or more groups which independently are -halogen, $-C_1$ - C_6 alkyl, aryloxy C_{0-6} alkyl SO_2R^{11} , C_{0-6} alkyl $COOR^{11}$, C_{0-6} alkoxyaryl, C_1 - C_6 haloalkyl, $-SO_2R^{11}$, $-OR^{11}$, $-SR^{11}$, $-N_3$, $-SO_2R^{11}$, $-COR^{11}$, $-SO_2N(R^{11})_2$, $-SO_2NR^{11}COR^{11}$, $-C\equiv N$, $-C(O)OR^{11}$, $-CON(R^{11})_2$, $-CON(R^{11})OR^{11}$, $-OCON(R^{11})_2$, $-NR^{11}COR^{11}$, $-NR^{11}CON(R^{11})_2$, $-NR^{11}COOR^{11}$, or $-N(R^{11})_2$;

R^{21} is hydrogen, halogen, nitro, cyano, C_1 - C_6 alkyl, or C_1 - C_6 haloalkyl;

R^2 is $-H$, $-[C(R^{15})_2]_m-OH$, $-C_1$ - C_6 alkyl, $-C_1$ - C_6 haloalkyl, halogen, $-C(O)N(R^{11})_2$, or $-COOR^{11}$;

provided that R^2 and R^{21} are not simultaneously hydrogen;

J is aryl or heteroaryl;

K is aryl or heteroaryl;

each R^4 and R^{41} is independently halogen, oxo, nitro, $CR^{11}=CR^{11}COOR^{11}$, aryloxy, aralkyloxy, aryloxyalkyl, aryl C_{0-6} alkylcarboxy, aryl, heteroaryl, heterocyclyl, heteroaryloxy, heterocyclioxy, $-G^1$, $-E-G^1$ or $-D-E-G^1$, wherein

D is $-O-$;

E is $-\text{[C(R}^{15})_2\text{]}_m-$ or $-\text{C}_3\text{-C}_8\text{cycloalkyl}$; and

G^1 is $-\text{C}_1\text{-C}_6\text{alkyl}$, $-\text{C}_1\text{-C}_6\text{haloalkyl}$, $-\text{COR}^{11}$, $-\text{COOR}^{11}$, $-\text{CON(R}^{11})_2$, $-\text{C}\equiv\text{N}$, $-\text{OR}^{11}$, $-\text{OCON(R}^{11})_2$, $-\text{OCOOR}^{11}$, $-\text{N}_3$, $-\text{NR}^{11}\text{COR}^{11}$, $-\text{NR}^{11}\text{SO}_2\text{R}^{11}$, $-\text{N(R}^{11})_2$, $-\text{NR}^{11}\text{COOR}^{11}$, $-\text{SO}_2\text{R}^{11}$, $-\text{SO}_2\text{NR}^{11}\text{COR}^{11}$, $-\text{SO}_2\text{N(R}^{11})_2$, $-\text{SOR}^{11}$, or $-\text{SR}^{11}$;

wherein each R^4 is optionally substituted with one or more R^{4a} ,

wherein each R^{4a} is independently halogen, aryloxy, aralkyloxy, aryloxyalkyl, $\text{C}_1\text{-C}_6$ alkoxyaryl, aryl $\text{C}_0\text{-C}_6$ alkylcarboxy, $-\text{G}'$, $-\text{E}'\text{-G}'$, or $-\text{D}'\text{-E}'\text{-G}'$, wherein

D' is $-\text{O}-$;

E' is $-\text{[C(R}^{15})_2\text{]}_m-$ or $-\text{C}_3\text{-C}_8\text{cycloalkyl}$; and

G' is $-\text{H}$, $-\text{halogen}$, $-\text{COR}^{11}$, $-\text{COOR}^{11}$, $-\text{C}\equiv\text{N}$, $-\text{OR}^{11}$, $-\text{NR}^{11}\text{SO}_2\text{R}^{11}$, $-\text{SO}_2\text{R}^{11}$, $-\text{SO}_2\text{N(R}^{11})_2$, or $-\text{SR}^{11}$;

L^2 is a bond;

each m is 0, 1, 2, 3, 4, 5, or 6;

q is 0, 1, 2, 3, 4, or 5, provided that q is 0 if and only if K is not phenyl; and

q' is 0, 1, 2, 3, or 4,

each R^{10} is independently $-\text{R}^{11}$, $-\text{C(=O)R}^{11}$, $-\text{CO}_2\text{R}^{11}$, or $-\text{SO}_2\text{R}^{11}$;

each R^{11} is independently $-\text{hydrogen}$, $-\text{C}_1\text{-C}_6$ alkyl, $\text{C}_2\text{-C}_6$ alkenyl, $\text{C}_2\text{-C}_6$ alkynyl, $\text{C}_3\text{-C}_8$ cycloalkyl, $(\text{C}_3\text{-C}_8$ cycloalkyl)- $\text{C}_1\text{-C}_6$ alkyl-, $(\text{C}_3\text{-C}_8$ cycloalkenyl)- $\text{C}_1\text{-C}_6$ alkyl-, $(\text{C}_3\text{-C}_8$ cycloalkyl)- $\text{C}_2\text{-C}_6$ alkenyl-, $-\text{C}_1\text{-C}_6$ haloalkyl, $-\text{N(R}^{12})_2$, aryl, $-(\text{C}_1\text{-C}_6)\text{alkyl-aryl}$, heteroaryl, $-(\text{C}_1\text{-C}_6)\text{alkyl-heteroaryl}$, heterocyclyl, or $-(\text{C}_1\text{-C}_6)\text{alkyl-heterocyclyl}$,


wherein any of R^{11} is optionally substituted with one or more radicals of R^{12} ;

each R^{12} is independently halogen, $\text{C}_1\text{-C}_6\text{haloalkyl}$, $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_1\text{-C}_6$ alkoxy, $(\text{C}_0\text{-C}_6$ alkyl) $\text{C=O(OR}^{13})$, $\text{C}_0\text{-C}_6$ alkyl OR^{13} , $\text{C}_0\text{-C}_6$ alkyl COR^{13} , $\text{C}_0\text{-C}_6$ alkyl SO_2R^{13} , $\text{C}_0\text{-C}_6$ alkyl $\text{CON(R}^{13})_2$, $\text{C}_0\text{-C}_6$ alkyl $\text{CONR}^{13}\text{OR}^{13}$, $\text{C}_0\text{-C}_6$ alkyl $\text{SO}_2\text{N(R}^{13})_2$, $\text{C}_0\text{-C}_6$ alkyl SR^{13} , $\text{C}_0\text{-C}_6$ haloalkyl OR^{13} , aryloxy, aralkyloxy, aryloxyalkyl, $\text{C}_0\text{-C}_6\text{alkoxyaryl}$, aryl $\text{C}_0\text{-C}_6$ alkylcarboxy, $-\text{C}_0\text{-C}_6$ alkyl $\text{N(R}^{13})_2$, $-\text{NR}^{13}\text{SO}_2\text{R}^{13}$, or $-\text{OC}_0\text{-C}_6$ alkyl COOR^{13} ;

each R^{13} is independently hydrogen, $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_2\text{-C}_6$ alkenyl, $\text{C}_2\text{-C}_6$ alkynyl, $\text{C}_3\text{-C}_8$ cycloalkyl, $(\text{C}_3\text{-C}_8$ cycloalkyl)- $\text{C}_1\text{-C}_6$ alkyl-, $(\text{C}_3\text{-C}_8$ cycloalkenyl)- $\text{C}_1\text{-C}_6$ alkyl-, or $(\text{C}_3\text{-C}_8$ cycloalkyl)- $\text{C}_2\text{-C}_6$ alkenyl-;


and each R^{14} is independently hydrogen, halogen, $(\text{C}_1\text{-C}_6)\text{alkyl}$, $(\text{C}_3\text{-C}_6)\text{cycloalkyl}$, or $(\text{C}_1\text{-C}_6)\text{haloalkyl}$.

2. The compound according to claim 1, wherein at least one of J and K is phenyl.
3. The compound according to claim 2, of the formula,



or a pharmaceutically acceptable salt thereof.


4. The compound according to claim 3, of the formula,



or a pharmaceutically acceptable salt thereof, wherein R^{21} is $-H$, $-halogen$, $-C_1-C_6alkyl$, or $-C_1-C_6haloalkyl$.

5. The compound according to claim 3, wherein each R^4 is independently halogen, aryl, heteroaryl, heterocyclyl, $-G^1$, or $-E-G^1$, wherein E is $-[C(R^{15})_2]_m-$, wherein each R^{15} is independently hydrogen or halogen; and G^1 is $-C_1-C_6alkyl$, $-C_1-C_6haloalkyl$, $-OR^{11}$, or $-SO_2R^{11}$.
6. The compound according to claim 3, wherein each R^{41} is independently halogen, $-G^2$, or $-E-G^2$, wherein E is $-[C(R^{15})_2]_m-$, wherein each R^{15} is independently hydrogen or halogen; and G^2 is $-C_1-C_6alkyl$, $-C_1-C_6haloalkyl$, or halogen; and each R^4 is independently halogen, aryl, heteroaryl, heterocyclyl, $-G^1$, or $-E-G^1$, wherein E is $-[C(R^{15})_2]_m-$, wherein each R^{15} is independently hydrogen or halogen; and G^1 is $-C_1-C_6alkyl$, $-C_1-C_6haloalkyl$, $-OR^{11}$, or $-SO_2R^{11}$.
7. The compound according to claim 4, wherein each R^{5a} is independently halogen, C_1-C_6alkyl , $C_1-C_6haloalkyl$, $C_3-C_8cycloalkyl$, OR^{11} , SO_2R^{11} , COR^{11} , $SO_2N(R^{11})_2$, $C\equiv N$, $C(O)OR^{11}$, $CON(R^{11})_2$, $NR^{11}COR^{11}$, or $N(R^{11})_2$.

8. A compound according to the formula,



or a pharmaceutically acceptable salt thereof, wherein

L¹ is -[C(R¹⁵)₂]_m-, or -C₃-C₈ cycloalkyl; wherein m' is any of 1 to 3; and

each R¹⁵ is independently hydrogen, halogen, (C₁-C₆)alkyl, or (C₁-C₆)haloalkyl;

R⁵ is aryl, heterocyclyl, heteroaryl, -(C₃-C₆)cycloalkyl, -C, or -B-C, wherein

B is -[C(R¹⁵)₂]_m-, -C₃-C₈cycloalkyl; and

C is -C₁-C₆alkyl or -C₁-C₆haloalkyl;

C is halogen, -C₁-C₆alkyl, -C₁-C₆haloalkyl, -SO₂R¹¹, -SR¹¹, -SO₂N(R¹¹)₂, -SO₂NR¹¹COR¹¹, -C≡N, -C(O)OR¹¹, -CON(R¹¹)₂, or -N(R¹¹)₂,

wherein R⁵ is optionally substituted with one or more R^{5a}, wherein

each R^{5a} is independently halogen, nitro, heteroaryl, heterocyclyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, (C₃-C₈ cycloalkyl)-C₁-C₆ alkyl-, (C₃-C₈ cycloalkenyl)-C₁-C₆ alkyl-, (C₃-C₈ cycloalkyl)-C₂-C₆ alkenyl-, aryl, arylalkyl, aryloxy, aryloxyaryl, arylC₁₋₆ alkoxy, C₁-C₆ alkyl, C₁-C₆ haloalkyl, SO₂R¹¹, OR¹¹, SR¹¹, N₃, SO₂R¹¹, COR¹¹, SO₂N(R¹¹)₂, SO₂NR¹¹COR¹¹, C≡N, C(O)OR¹¹, CON(R¹¹)₂, CON(R¹¹)OR¹¹, OCON(R¹¹)₂, NR¹¹COR¹¹, NR¹¹CON(R¹¹)₂, NR¹¹COOR¹¹, or N(R¹¹)₂, wherein

each R^{5a} is optionally substituted with one or more groups which independently are -halogen, -C₁-C₆ alkyl, aryloxy C₀₋₆ alkylSO₂R¹¹, C₀₋₆ alkylCOOR¹¹, C₀₋₆ alkoxyaryl, C₁-C₆ haloalkyl, -SO₂R¹¹, -OR¹¹, -SR¹¹, -N₃, -SO₂R¹¹, -COR¹¹, -SO₂N(R¹¹)₂, -SO₂NR¹¹COR¹¹, -C≡N, -C(O)OR¹¹, -CON(R¹¹)₂, -CON(R¹¹)OR¹¹, -OCON(R¹¹)₂, -NR¹¹COR¹¹, -NR¹¹CON(R¹¹)₂, -NR¹¹COOR¹¹, or -N(R¹¹)₂;

R²¹ is hydrogen, halogen, nitro, cyano, C₁-C₆alkyl, or C₁-C₆haloalkyl;

R² is -H, -[C(R¹⁵)₂]_m-OH, -C₁-C₆alkyl, -C₁-C₆haloalkyl, halogen, -C(O)N(R¹¹)₂, or -COOR¹¹;

provided that R² and R²¹ are not simultaneously hydrogen;

J and K are both phenyl;

each R^4 and R^{41} is independently halogen, oxo, nitro, $CR^{11}=CR^{11}COOR^{11}$, aryloxy, aralkyloxy, aryloxyalkyl, arylC₀-C₆ alkylcarboxy, aryl, heteroaryl, heterocyclyl, heteroaryloxy, heterocyclyloxy, -G¹, -E-G¹ or -D-E-G¹, wherein

D is -O-;

E is $-[C(R^{15})_2]_m-$ or -C₃-C₈cycloalkyl; and

G¹ is -C₁-C₆alkyl, -C₁-C₆haloalkyl, -COR¹¹, -COOR¹¹, -CON(R¹¹)₂, -C≡N, -OR¹¹, -OCON(R¹¹)₂, -OCOOR¹¹, -N₃, -NR¹¹COR¹¹, -NR¹¹SO₂R¹¹, -N(R¹¹)₂, -NR¹¹COOR¹¹, -SO₂R¹¹, -SO₂NR¹¹COR¹¹, -SO₂N(R¹¹)₂, -SOR¹¹, or -SR¹¹;

wherein each R^4 is optionally substituted with one or more R^{4a} ,

wherein each R^{4a} is independently halogen, aryloxy, aralkyloxy, aryloxyalkyl,

C₁-C₆ alkoxyaryl, arylC₀-C₆ alkylcarboxy, -G', -E'-G', or -D'-E'-G', wherein

D' is -O-;

E' is $-[C(R^{15})_2]_m-$ or -C₃-C₈cycloalkyl; and

G' is -H, -halogen, -COR¹¹, -COOR¹¹, -C≡N, -OR¹¹, -NR¹¹SO₂R¹¹, -SO₂R¹¹, -SO₂N(R¹¹)₂, or -SR¹¹;

L² is a bond or $-[C(R^{15})_2]_m-V^2-[C(R^{15})_2]_n-$, wherein

V² is independently -C(R¹¹)₂-, -C(R¹¹)₂C(R¹¹)₂-, -C(R¹¹)=C(R¹¹)-, -C(R¹¹)₂O-, -C(R¹¹)₂NR¹¹-, -C≡C-, -O-, -S-, -SO₂-, -N(R¹⁰)CO-, -N(R¹⁰)CO₂-, -CON(R¹⁰)-, -CON(R¹¹)-, -CON(R¹¹)O-, -CO-, -CS-, -CO₂-, -OC(=O)-, -OC(=O)N(R¹⁰)-, -N(R¹⁰)SO₂-, -SO₂N(R¹⁰)-, -NR¹⁰CONR¹⁰-, -NR¹⁰CSNR¹⁰-, C₃-C₆cycloalkyl-, or C₃-C₆cyclohaloalkyl,

or V² is C₂₋₆ alidiyl,

wherein alidiyl chain is optionally interrupted by -C(R¹¹)₂-, -C(R¹¹)₂C(R¹¹)₂-, -C(R¹¹)=C(R¹¹)-, -C(R¹¹)₂O-, -C(R¹¹)₂NR¹¹-, -C(R¹¹)₂NR¹¹-, -C≡C-, -O-, -S-, -N(R¹⁰)CO-, -N(R¹⁰)CO₂-, -CON(R¹⁰)-, -CON(R¹¹)-, -CON(R¹¹)O-, -CO-, -CO₂-, -OC(=O)-, -OC(=O)N(R¹⁰)-, -SO₂-, -N(R¹⁰)SO₂- or -SO₂N(R¹⁰)-;

or V² is aryl, heteroaryl, or heterocyclyl, each of which is optionally substituted with one or more R^9 , wherein

each R^9 is independently halogen, C₁-C₆ haloalkyl, C₁-C₆ alkyl, C₁-C₆ alkyloxy, C₀-C₆ alkyl or C₁-C₆ alkylCOOR¹¹;

each m is 0, 1, 2, 3, 4, 5, or 6;

q is 1, 2, 3, 4, or 5,

q' is 0, 1, 2, 3, or 4,

each R¹⁰ is independently -R¹¹, -C(=O)R¹¹, -CO₂R¹¹, or -SO₂R¹¹;

each R¹¹ is independently -hydrogen, -C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, (C₃-C₈ cycloalkyl)-C₁-C₆ alkyl-, (C₃-C₈ cycloalkenyl)-C₁-C₆ alkyl-, (C₃-C₈ cycloalkyl)-C₂-C₆ alkenyl-, -C₁-C₆ haloalkyl, -N(R¹²)₂, aryl, -(C₁-C₆)alkyl-aryl, heteroaryl, -(C₁-C₆)alkyl-heteroaryl, heterocyclyl, or -(C₁-C₆)alkyl-heterocyclyl,


wherein any of R¹¹ is optionally substituted with one or more radicals of R¹²;

each R¹² is independently halogen, C₁-C₆haloalkyl, C₁-C₆ alkyl, C₁-C₆ alkoxy, (C₀-C₆ alkyl)C=O(OR¹³); C₀-C₆ alkylOR¹³, C₀-C₆ alkylCOR¹³, C₀-C₆ alkylSO₂R¹³, C₀-C₆ alkylCON(R¹³)₂, C₀-C₆ alkylCONR¹³OR¹³, C₀-C₆ alkylSO₂N(R¹³)₂, C₀-C₆ alkylSR¹³, C₀-C₆ haloalkylOR¹³, aryloxy, aralkyloxy, aryloxyalkyl, C₀₋₆alkoxyaryl, arylC₀₋₆ alkylcarboxy, -C₀-C₆ alkylN(R¹³)₂, -NR¹³SO₂ R¹³, or -OC₀₋₆ alkylCOOR¹³;
and

each R¹³ is independently hydrogen, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, (C₃-C₈ cycloalkyl)-C₁-C₆ alkyl-, (C₃-C₈ cycloalkenyl)-C₁-C₆ alkyl-, or (C₃-C₈ cycloalkyl)-C₂-C₆ alkenyl-.

9. The compound according to claim 8, wherein L² is a bond.

10. The compound according to claim 9, of the formula,



or a pharmaceutically acceptable salt thereof.

11. The compound according to claim 10, wherein

each R¹⁵ is independently -H or -(C₁-C₂)alkyl;

m' is 1 or 2; and

R⁵ is phenyl optionally substituted with one or more R^{5a}.


12. The compound according to claim 11, wherein

each R⁴ is independently halogen, aryl, heteroaryl, heterocyclyl, -G¹, or -E-G¹, wherein


E is -[C(R¹⁵)₂]_m-, wherein each R¹⁵ is independently hydrogen or halogen; and

G^1 is $-C_1-C_6$ alkyl, $-C_1-C_6$ haloalkyl, halogen, $-OR^{11}$, or $-SO_2R^{11}$.


13. The compound according to claim 11, wherein each R^{41} is independently halogen, $-G^1$, or $-E-G^1$, wherein E is $-[C(R^{15})_2]_m-$, wherein each R^{15} is independently hydrogen or halogen; and G^1 is $-C_1-C_6$ alkyl or $-C_1-C_6$ haloalkyl.
14. The compound according to claim 10, wherein R^{15} is $-H$; m is 1, 2, or 3; and R^5 is heterocyclyl optionally substituted with one or more R^{5a} .
15. The compound according to claim 14, wherein each R^4 is independently halogen, aryl, heteroaryl, heterocyclyl, $-G^1$, or $-E-G^1$, wherein E is $-[C(R^{15})_2]_m-$, wherein each R^{15} is independently hydrogen or halogen; and G^1 is $-C_1-C_6$ alkyl, $-C_1-C_6$ haloalkyl, $-OR^{11}$, or $-SO_2R^{11}$.
16. The compound according to claim 14, wherein each R^{41} is independently halogen, $-G^1$, or $-E-G^1$, wherein E is $-[C(R^{15})_2]_m-$, wherein each R^{15} is independently hydrogen or halogen; and G^1 is $-C_1-C_6$ alkyl or $-C_1-C_6$ haloalkyl.
17. A compound according to one of the following formulas,




IIa



IIb



IIc



IIId


or a pharmaceutically acceptable salt thereof, wherein,

R^1 is $-L^1-R^5$, wherein

L^1 is a bond, $-[C(R^{15})_2]_m-$, or $-C_3-C_8$ cycloalkyl-; and

R^5 is phenyl or pyridyl, each optionally substituted with one or two R^{5a} , wherein

each R^{5a} is independently -halogen, -C₁-C₆ alkoxy, -C₁-C₆ alkyl, or -C₁-C₆ haloalkyl;
 R^2 , R^{21} , and R^3 are each independently -H, $-[C(R^{15})_2]_m-OH$, -C₁-C₆ alkyl, -C₁-C₆ haloalkyl, halogen, $-C(O)N(R^{11})_2$, or $-COOR^{11}$;
 provided that R^2 and R^{21} are not simultaneously hydrogen;
 G is a group of the formula,



L^2 is a bond;
 J is phenyl, pyridyl, or thienyl;
 K is phenyl or pyridyl;
 q is 0, 1, 2, 3, 4 or 5, provided that q is 0 if and only if K is not phenyl;
 q' is 0, 1, 2, 3, or 4,
 each R^{41} is -halogen, -C₁-C₆ alkyl, or C₁-C₆ -haloalkyl; and
 each R^4 is -halogen, $-[C(R^{15})_2]_m-OH$, $-SO_2R^{11}$, $-SO_2N(R^{11})_2$, $-C(O)N(R^{11})_2$, $-COOR^{11}$, -C₁-C₆ alkyl, or -C₁-C₆ haloalkyl;
 each m is 0, 1, 2, 3, 4, 5, or 6;
 each R^{15} is independently hydrogen, halogen, (C₁-C₆)alkyl, (C₃-C₆)cycloalkyl, or (C₁-C₆)haloalkyl;
 and
 each R^{11} is independently -hydrogen, -C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, (C₃-C₈ cycloalkyl)-C₁-C₆ alkyl-, (C₃-C₈ cycloalkenyl)-C₁-C₆ alkyl-, (C₃-C₈ cycloalkyl)-C₂-C₆ alkenyl-, -C₁-C₆ haloalkyl, $-N(R^{12})_2$, heterocyclyl, or -(C₁-C₆)alkyl-heterocyclyl, wherein any of R^{11} is optionally substituted with one or more radicals of R^{12} ;
 each R^{12} is independently halogen, C₁-C₆haloalkyl, C₁-C₆ alkyl, C₁-C₆ alkoxy, (C₀-C₆ alkyl)C=O(OR¹³); C₀-C₆ alkylOR¹³, C₀-C₆ alkylCOR¹³, C₀-C₆ alkylSO₂R¹³, C₀-C₆ alkylCON(R¹³)₂, C₀-C₆ alkylCONR¹³OR¹³, C₀-C₆ alkylSO₂N(R¹³)₂, C₀-C₆ alkylSR¹³, C₀-C₆ haloalkylOR¹³, aryloxy, aralkyloxy, aryloxyalkyl, C₀₋₆alkoxyaryl, arylC₀₋₆ alkylcarboxy, $-C_0-C_6$ alkylN(R¹³)₂, $-NR^{13}SO_2R^{13}$, or $-OC_{0-6}$ alkylCOOR¹³; and

each R^{13} is independently hydrogen, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, C_3 - C_8 cycloalkyl, $(C_3$ - C_8 cycloalkyl)- C_1 - C_6 alkyl-, $(C_3$ - C_8 cycloalkenyl)- C_1 - C_6 alkyl-, or $(C_3$ - C_8 cycloalkyl)- C_2 - C_6 alkenyl-.

18. The compound according to claim 10, wherein

each R^{15} is independently $-H$ or $-(C_1-C_2)$ alkyl;

m' is 1 or 2; and

R^5 is heteroaryl optionally substituted with one or more R^{5a} .

19. The compound according to claim 18, wherein

each R^4 is independently halogen, aryl, heteroaryl, heterocyclyl, $-G^1$, or $-E-G^1$,
wherein

E is $-[C(R^{15})_2]_{m'}$, wherein

each R^{15} is independently hydrogen or halogen; and

G^1 is $-C_1$ - C_6 alkyl, $-C_1$ - C_6 haloalkyl, $-OR^{11}$, or $-SO_2R^{11}$.

20. The compound according to claim 18, wherein


each R^{41} is independently halogen, $-G^1$, or $-E-G^1$, wherein

E is $-[C(R^{15})_2]_{m'}$,

wherein each R^{15} is independently hydrogen or halogen; and

G^1 is $-C_1$ - C_6 alkyl or $-C_1$ - C_6 haloalkyl.

21. A compound according to the formula,



or a pharmaceutically acceptable salt thereof, wherein

L_1 is $-[C(R^{15})_2]_{m'}$ or $-C_3$ - C_8 cycloalkyl; wherein m' is 1 or 2; and

R^5 is heteroaryl optionally substituted with one or more R^{5a} , wherein

each R^{5a} is independently halogen, nitro, heteroaryl, heterocyclyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, C_3 - C_8 cycloalkyl, $(C_3$ - C_8 cycloalkyl)- C_1 - C_6 alkyl-, $(C_3$ - C_8 cycloalkenyl)- C_1 - C_6 alkyl-, $(C_3$ - C_8 cycloalkyl)- C_2 - C_6 alkenyl-, aryl, arylalkyl, aryloxy, aryloxyaryl, aryl C_{1-6} alkoxy, C_1 - C_6

alkyl, C₁-C₆ haloalkyl, SO₂R¹¹, OR¹¹, SR¹¹, N₃, SO₂R¹¹, COR¹¹, SO₂N(R¹¹)₂, SO₂NR¹¹COR¹¹, C≡N, C(O)OR¹¹, CON(R¹¹)₂, CON(R¹¹)OR¹¹, OCON(R¹¹)₂, NR¹¹COR¹¹, R¹¹CON(R¹¹)₂, NR¹¹COOR¹¹, or N(R¹¹)₂, wherein

each R^{5a} is optionally substituted with one or more groups which independently are -halogen, -C₁-C₆ alkyl, aryloxy C₀₋₆ alkylSO₂R¹¹, C₀₋₆ alkylCOOR¹¹, C₀₋₆ alkoxyaryl, C₁-C₆ haloalkyl, -SO₂R¹¹, -OR¹¹, -SR¹¹, -N₃, -SO₂R¹¹, -COR¹¹, -SO₂N(R¹¹)₂, -SO₂NR¹¹COR¹¹, -C≡N, -C(O)OR¹¹, -CON(R¹¹)₂, -CON(R¹¹)OR¹¹, -OCON(R¹¹)₂, -NR¹¹COR¹¹, -NR¹¹CON(R¹¹)₂, -NR¹¹COOR¹¹, or -N(R¹¹)₂;

R²¹ is hydrogen, halogen, nitro, cyano, C₁-C₆alkyl, or C₁-C₆haloalkyl;

R² is -H, -[C(R¹⁵)₂]_m-OH, -C₁-C₆alkyl, -C₁-C₆ haloalkyl, halogen, -C(O)N(R¹¹)₂, or -COOR¹¹;

provided that R² and R²¹ are not simultaneously hydrogen;

each m is 0, 1, 2, 3, 4, 5, or 6;

q is 1, 2, 3, 4, or 5,

q' is 0, 1, 2, 3, or 4, and

each R⁴ and R⁴¹ is independently halogen, oxo, nitro, CR¹¹=CR¹¹COOR¹¹, aryloxy, aralkyloxy, aryloxyalkyl, arylC₀₋₆ alkylcarboxy, aryl, heteroaryl, heterocyclyl, heteroaryloxy, heterocycloxy, -G¹, -E-G¹, or -D-E-G¹, wherein

D is -O-;

E is -[C(R¹⁵)₂]_m- or -C₃-C₈cycloalkyl; and

G¹ is -C₁-C₆alkyl, -C₁-C₆haloalkyl, -COR¹¹, -COOR¹¹, -CON(R¹¹)₂, -C≡N, -OR¹¹, -OCON(R¹¹)₂, -OCOOR¹¹, -N₃, -NR¹¹COR¹¹, -NR¹¹SO₂R¹¹, -N(R¹¹)₂, -NR¹¹COOR¹¹, -SO₂R¹¹, -SO₂NR¹¹COR¹¹, -SO₂N(R¹¹)₂, -SOR¹¹, or -SR¹¹;

wherein each R⁴ is optionally substituted with one or more R^{4a},

wherein each R^{4a} is independently halogen, aryloxy, aralkyloxy, aryloxyalkyl, C₁-C₆ alkoxyaryl, arylC₀₋₆ alkylcarboxy, -G', -E'-G', or -D'-E'-G', wherein

D' is -O-;

E' is -[C(R¹⁵)₂]_m- or -C₃-C₈cycloalkyl; and

G' is -H, -halogen, -COR¹¹, -COOR¹¹, -C≡N, -OR¹¹, -NR¹¹SO₂R¹¹, -SO₂R¹¹, -SO₂N(R¹¹)₂, or -SR¹¹;

each R¹¹ is independently -hydrogen, -C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, (C₃-C₈ cycloalkyl)-C₁-C₆ alkyl-, (C₃-C₈ cycloalkenyl)-C₁-C₆ alkyl-, (C₃-C₈ cycloalkyl)-C₂-C₆ alkenyl-,

-C₁-C₆ haloalkyl, -N(R¹²)₂, aryl, -(C₁-C₆)alkyl-aryl, heteroaryl, -(C₁-C₆)alkyl-heteroaryl, heterocyclyl, or -(C₁-C₆)alkyl-heterocyclyl,

wherein any of R¹¹ is optionally substituted with one or more radicals of R¹²;

each R¹² is independently halogen, C₁-C₆haloalkyl, C₁-C₆ alkyl, C₁-C₆ alkoxy, (C₀-C₆ alkyl)C=O(OR¹³); C₀-C₆ alkylOR¹³, C₀-C₆ alkylCOR¹³, C₀-C₆ alkylSO₂R¹³, C₀-C₆ alkylCON(R¹³)₂, C₀-C₆ alkylCONR¹³OR¹³, C₀-C₆ alkylSO₂N(R¹³)₂, C₀-C₆ alkylSR¹³, C₀-C₆ haloalkylOR¹³, aryloxy, aralkyloxy, aryloxyalkyl, C₀₋₆alkoxyaryl, arylC₀₋₆ alkylcarboxy, -C₀-C₆ alkylN(R¹³)₂, -NR¹³SO₂ R¹³, or -OC₀₋₆ alkylCOOR¹³;

each R¹³ is independently hydrogen, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, (C₃-C₈ cycloalkyl)-C₁-C₆ alkyl-, (C₃-C₈ cycloalkenyl)-C₁-C₆ alkyl-, or (C₃-C₈ cycloalkyl)-C₂-C₆ alkenyl-;

and each R¹⁵ is independently H, halogen or (C₁-C₆)alkyl.


22. The compound according to claim 21, wherein

each R⁴ is independently halogen, aryl, heteroaryl, heterocyclyl, -G¹, or -E-G¹, wherein

E is -[C(R¹⁵)₂]_m-, wherein each R¹⁵ is independently hydrogen or halogen; and

G¹ is -C₁-C₆alkyl, -C₁-C₆haloalkyl, -OR¹¹, or -SO₂R¹¹.

23. A compound according to one of the following formulas,



or a pharmaceutically acceptable salt thereof, wherein,

R¹ is -L¹-R⁵, wherein

L¹ is a bond, -C₃-C₈ cycloalkyl- or L⁵, wherein

each L⁵ is independently -[C(R¹⁵)₂]_m-, wherein

m is 0, 1, 2, or 3; and

each R¹⁵ is independently hydrogen, halogen, (C₁-C₆)alkyl, or (C₁-C₆)haloalkyl;

and

R⁵ is aryl, heterocyclyl, heteroaryl, -C, or -B-C, wherein

B is $-\text{C}(\text{R}^{15})_2\text{m}-$, $-\text{C}_3\text{-C}_6\text{cycloalkyl}-$; and

C is $-\text{C}_1\text{-C}_6\text{alkyl}$ or $-\text{C}_1\text{-C}_6\text{haloalkyl}$;

wherein R^5 is optionally substituted with one or more R^{5a} , wherein

each R^{5a} is independently halogen, nitro, heteroaryl, heterocyclyl, $\text{C}_2\text{-C}_6$ alkenyl, $\text{C}_2\text{-C}_6$ alkynyl, $(\text{C}_3\text{-C}_8 \text{ cycloalkyl})\text{-C}_1\text{-C}_6 \text{ alkyl}-$, $(\text{C}_3\text{-C}_8 \text{ cycloalkenyl})\text{-C}_1\text{-C}_6 \text{ alkyl}-$, $(\text{C}_3\text{-C}_8 \text{ cycloalkyl})\text{-C}_2\text{-C}_6 \text{ alkenyl}-$, aryl, arylalkyl, aryloxy, aryloxyaryl, aryl C_{1-6} alkoxy, $\text{C}_1\text{-C}_6 \text{ alkyl}$, $\text{C}_1\text{-C}_6 \text{ haloalkyl}$, $\text{C}_3\text{-C}_6\text{cycloalkyl}$, SO_2R^{11} , OR^{11} , SR^{11} , N_3 , SO_2R^{11} , COR^{11} , $\text{SO}_2\text{N}(\text{R}^{11})_2$, $\text{SO}_2\text{NR}^{11}\text{COR}^{11}$, $\text{C}\equiv\text{N}$, $\text{C}(\text{O})\text{OR}^{11}$, $\text{CON}(\text{R}^{11})_2$, $\text{CON}(\text{R}^{11})\text{OR}^{11}$, $\text{OCON}(\text{R}^{11})_2$, $\text{NR}^{11}\text{COR}^{11}$, $\text{NR}^{11}\text{CON}(\text{R}^{11})_2$, $\text{NR}^{11}\text{COOR}^{11}$, or $\text{N}(\text{R}^{11})_2$, wherein


each R^{5a} is optionally substituted with one or more groups which independently are -halogen, $-\text{C}_1\text{-C}_6 \text{ alkyl}$, aryloxy, $\text{C}_{0-6} \text{ alkylSO}_2\text{R}^{11}$, $\text{C}_{0-6} \text{ alkylCOOR}^{11}$, $\text{C}_{0-6} \text{ alkoxyaryl}$, $-\text{C}_1\text{-C}_6 \text{ haloalkyl}$, $-\text{SO}_2\text{R}^{11}$, $-\text{OR}^{11}$, $-\text{SR}^{11}$, $-\text{N}_3$, $-\text{SO}_2\text{R}^{11}$, $-\text{COR}^{11}$, $\text{SO}_2\text{N}(\text{R}^{11})_2$, $-\text{SO}_2\text{NR}^{11}\text{COR}^{11}$, $-\text{C}\equiv\text{N}$, $-\text{C}(\text{O})\text{OR}^{11}$, $-\text{CON}(\text{R}^{11})_2$, $-\text{CON}(\text{R}^{11})\text{OR}^{11}$, $-\text{OCON}(\text{R}^{11})_2$, $-\text{NR}^{11}\text{COR}^{11}$, $-\text{NR}^{11}\text{CON}(\text{R}^{11})_2$, $-\text{NR}^{11}\text{COOR}^{11}$, or $-\text{N}(\text{R}^{11})_2$;

R^2 is $-\text{H}$, $-\text{C}(\text{R}^{15})_2\text{m}-\text{OH}$, $-\text{C}_1\text{-C}_6 \text{ alkyl}$, $-\text{C}_1\text{-C}_6 \text{ haloalkyl}$, halogen, $-\text{C}(\text{O})\text{N}(\text{R}^{11})_2$, or $-\text{COOR}^{11}$;

R^{21} is hydrogen, halogen, $\text{C}_1\text{-C}_6 \text{ alkyl}$, or $\text{C}_1\text{-C}_6 \text{ haloalkyl}$; and

provided that R^2 and R^{21} are not simultaneously hydrogen;

G is a group of the formula,



J is aryl or heteroaryl;

K is aryl or heteroaryl;

each R^4 and R^{41} is independently halogen, heteroaryl, heterocyclyl, $-\text{G}^1$, $-\text{E}-\text{G}^1$, or $-\text{D}-\text{E}-\text{G}^1$,

wherein

D is $-\text{O}-$;

E is $-\text{C}(\text{R}^{15})_2\text{m}-$ or $-\text{C}_3\text{-C}_6\text{cycloalkyl}$; and

G^1 is $-\text{C}_1\text{-C}_6\text{alkyl}$, $-\text{C}_1\text{-C}_6\text{haloalkyl}$, $-\text{COR}^{11}$, $-\text{COOR}^{11}$, $-\text{CON}(\text{R}^{11})_2$, $-\text{C}\equiv\text{N}$, $-\text{OR}^{11}$, $-\text{SOR}^{11}$, $-\text{SO}_2\text{R}^{11}$, $-\text{SO}_2\text{N}(\text{R}^{11})_2$, or $-\text{SR}^{11}$,

L^2 is a bond;

q is 1, 2, or 3, and

q' is 0, 1, 2 or 3,

each R^{11} is independently -hydrogen, -C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, (C₃-C₈ cycloalkyl)-C₁-C₆ alkyl-, (C₃-C₈ cycloalkenyl)-C₁-C₆ alkyl-, (C₃-C₈ cycloalkyl)-C₂-C₆ alkenyl-, -C₃-C₈ cycloalkyl, -(C₁-C₆)alkyl-(C₃-C₈)cycloalkyl, -C₁-C₆ haloalkyl, -N(R¹²)₂, aryl, -(C₁-C₆)alkyl-aryl, heteroaryl, -(C₁-C₆)alkyl-heteroaryl, heterocyclyl, or -(C₁-C₆)alkyl-heterocyclyl,

wherein any of R^{11} is optionally substituted with one or more radicals of R^{12} ;

each R^{12} is independently halogen, OR¹³, N(R¹³)₂, C₁-C₆haloalkyl, C₁-C₆ alkyl, C₁-C₆ alkoxy, C₀-C₆ alkylOR¹³, C₀-C₆ alkylCOR¹³, C₀-C₆ alkylSO₂R¹³, C₀-C₆ alkylCON(R¹³)₂, C₀-C₆ alkylCONR¹³OR¹³, C₀-C₆ alkylSO₂N(R¹³)₂, C₀-C₆ alkylSR¹³, C₀-C₆ haloalkylOR¹³, aryloxy, aralkyloxy, aryloxyalkyl, C₀₋₆alkoxyaryl, arylC₀₋₆ alkylcarboxy, C₀-C₆ alkyl, -NR¹³SO₂ R¹³, or -OC₀₋₆ alkylCOOR¹³;

each R^{13} is independently hydrogen C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, (C₃-C₈ cycloalkyl)-C₁-C₆ alkyl-, (C₃-C₈ cycloalkenyl)-C₁-C₆ alkyl-, or (C₃-C₈ cycloalkyl)-C₂-C₆ alkenyl-.

24. A compound which is one of the species selected from the group consisting of:

Cpd #	Structure	Name
1		2-(2-(2-fluorobenzyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
2		2-(1-(3'-(methylsulfonyl)biphenyl-4-yl)-2-(2-(trifluoromethyl)benzyl)-1H-imidazol-4-yl)propan-2-ol;

Cpd #	Structure	Name
3		2-(2-(2-chlorobenzyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
4		2-(2-(2-chlorobenzyl)-1-(3'-(ethylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
5		2-(2-(2-chlorobenzyl)-1-(4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
6		2-(2-(2,6-dichlorobenzyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
7		2-(2-(2,6-dichlorobenzyl)-1-(3'-(ethylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
8		2-(2-(2,6-dichlorobenzyl)-1-(3'-(ethylsulfonyl)-4'-(hydroxymethyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
9		2-(2-(2-chloro-6-fluorobenzyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
10		2-(2-(2-chloro-6-fluorobenzyl)-1-(3'-(ethylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;

Cpd #	Structure	Name
11		2-(2-(2,3-dichlorobenzyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
12		2-(2-(2,3-dichlorobenzyl)-1-(3'-(ethylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
13		3-(2-(2-chlorobenzyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)pentan-3-ol;
14		3-(2-(2-chlorobenzyl)-1-(3'-(ethylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)pentan-3-ol;
15		3-(2-(2-chlorobenzyl)-1-(4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)pentan-3-ol;
16		2-(2-(5-chloro-2-(trifluoromethyl)benzyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
17		2-(2-(5-chloro-2-(trifluoromethyl)benzyl)-1-(3'-(ethylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;

Cpd #	Structure	Name
18		2-(2-(5-chloro-2-(trifluoromethyl)benzyl)-1-(4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
19		2-(2-(2-(2-chlorophenyl)propan-2-yl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
20		2-(2-(2-(2-chlorophenyl)propan-2-yl)-1-(3'-(ethylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
21		2-(2-(2-(2,3-dichlorophenyl)propan-2-yl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
22		2-(2-(2-(2,3-dichlorophenyl)propan-2-yl)-1-(3'-(ethylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
23		2-(2-(2,3-dichlorobenzyl)-1-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
24		2-(2-(2,3-dichlorobenzyl)-1-(3'-(ethylsulfonyl)-3-methylbiphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;

Cpd #	Structure	Name
25		2-(1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2,3-dichlorobenzyl)-1H-imidazol-4-yl)propan-2-ol;
26		2-(1-(3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl)-2-(2,3-dichlorobenzyl)-1H-imidazol-4-yl)propan-2-ol;
27		2-(2-(2-chlorophenethyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
28		2-(2-(2-chlorophenethyl)-1-(3'-(ethylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
29		2-(2-(2-chlorophenethyl)-1-(4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
30		2-(1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(naphthalen-1-yl)-1H-imidazol-4-yl)propan-2-ol;
31		2-(1-(3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl)-2-(naphthalen-1-yl)-1H-imidazol-4-yl)propan-2-ol;

Cpd #	Structure	Name
32		2-(1-(3-chloro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-2-(naphthalen-1-yl)-1H-imidazol-4-yl)propan-2-ol;
33		2-(1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(isoquinolin-1-yl)-1H-imidazol-4-yl)propan-2-ol;
34		2-(1-(3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl)-2-(isoquinolin-1-yl)-1H-imidazol-4-yl)propan-2-ol;
35		2-(2-(isoquinolin-5-yl)-1-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
36		2-(1-(3'-(ethylsulfonyl)-3-methylbiphenyl-4-yl)-2-(isoquinolin-5-yl)-1H-imidazol-4-yl)propan-2-ol;
37		2-(1-(3'-(methylsulfonyl)biphenyl-4-yl)-2-(4-(trifluoromethyl)pyridin-3-yl)-1H-imidazol-4-yl)propan-2-ol;
38		2-(1-(3'-(ethylsulfonyl)biphenyl-4-yl)-2-(4-(trifluoromethyl)pyridin-3-yl)-1H-imidazol-4-yl)propan-2-ol;
39		2-(1-(4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-2-(4-(trifluoromethyl)pyridin-3-yl)-1H-imidazol-4-yl)propan-2-ol;

Cpd #	Structure	Name
40		2-(1-(3'-(methylsulfonyl)biphenyl-4-yl)-2-(2-morpholinoethyl)-1H-imidazol-4-yl)propan-2-ol;
41		2-(1-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-2-(thiophen-2-yl)-1H-imidazol-4-yl)propan-2-ol;
42		2-(2-(1,5-dimethyl-1H-pyrrol-2-yl)-1-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
43		2-(2-(1,5-dimethyl-1H-pyrrol-2-yl)-1-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
44		2-(1-(2,6-dichlorophenyl)-2-(5-(3-(methylsulfonyl)phenyl)thiophen-2-yl)-1H-imidazol-4-yl)propan-2-ol;
45		2-(1-(2,6-dichlorophenyl)-2-(5-(3-(ethylsulfonyl)phenyl)thiophen-2-yl)-1H-imidazol-4-yl)propan-2-ol;
46		1-(2,5-dichlorophenyl)-2-{5-[3-(methylsulfonyl)phenyl]-2-thienyl}-4-(trifluoromethyl)-1H-imidazole;

Cpd #	Structure	Name
47		1-(2-chlorophenyl)-2-{5-[3-(methylsulfonyl)phenyl]-2-thienyl}-4-(trifluoromethyl)-1H-imidazole;
48		methyl 4-{5-[1-(2-chlorophenyl)-4-(trifluoromethyl)-1H-imidazol-2-yl]-2-thienyl}-3-methylphenylacetate;
49		5-{5-[1-(2,5-dichlorophenyl)-4-(trifluoromethyl)-1H-imidazol-2-yl]-2-thienyl}-2-(ethylthio)-3-methylpyridine;
50		5-{5-[1-(2,5-dichlorophenyl)-4-(trifluoromethyl)-1H-imidazol-2-yl]-2-thienyl}-2-(ethylsulfonyl)-3-methylpyridine;
51		5-{5-[1-(2,5-dichlorophenyl)-4-(trifluoromethyl)-1H-imidazol-2-yl]-2-thienyl}-3-methyl-2-(methylsulfonyl)pyridine;
52		4-(5-{5-[1-(2,5-dichlorophenyl)-4-(trifluoromethyl)-1H-imidazol-2-yl]-2-thienyl}pyridin-2-yl)morpholine;
53		1,1-dimethylethyl 4-(5-{5-[1-(2,5-dichlorophenyl)-4-(trifluoromethyl)-1H-imidazol-2-yl]-2-thienyl}pyridin-2-yl)piperazine-1-carboxylate;

Cpd #	Structure	Name
54		1-(5-{5-[1-(2,5-dichlorophenyl)-4-(trifluoromethyl)-1H-imidazol-2-yl]-2-thienyl}pyridin-2-yl)piperazine;
55		5-{5-[1-(2-chlorophenyl)-4-(trifluoromethyl)-1H-imidazol-2-yl]-2-thienyl}-2-(ethylthio)-3-methylpyridine;
56		5-{5-[1-(2-chlorophenyl)-4-(trifluoromethyl)-1H-imidazol-2-yl]-2-thienyl}-3-methyl-2-(methylsulfonyl)pyridine;
57		5-{5-[1-(2-chlorophenyl)-4-(trifluoromethyl)-1H-imidazol-2-yl]-2-thienyl}-2-(ethylsulfonyl)-3-methylpyridine;
58		2-(1-(2,6-dichlorophenyl)-2-(3'-(ethylsulfonyl)-3-methylbiphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
59		2-(1-(2-isopropyl-6-methylphenyl)-2-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
60		2-(2-(3'-(ethylsulfonyl)-3-methylbiphenyl-4-yl)-1-(2-isopropyl-6-methylphenyl)-1H-imidazol-4-yl)propan-2-ol;
61		2-(1-(2-isopropylphenyl)-2-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;

Cpd #	Structure	Name
62		2-(2-(3'-(ethylsulfonyl)-3-methylbiphenyl-4-yl)-1-(2-isopropylphenyl)-1H-imidazol-4-yl)propan-2-ol;
63		2-(1-(2,6-dichlorophenyl)-2-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
64		2-(1-(2,6-dichlorophenyl)-2-(3'-(ethylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
65		2-(2-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-1-(2,6-dichlorophenyl)-1H-imidazol-4-yl)propan-2-ol;
66		2-(2-(3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl)-1-(2,6-dichlorophenyl)-1H-imidazol-4-yl)propan-2-ol;
67		1,1,1-trifluoro-2-(1-(3'-(methylsulfonyl)biphenyl-4-yl)-2-(2-(trifluoromethyl)phenyl)-1H-imidazol-4-yl)propan-2-ol;
68		ethyl 1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-1H-imidazole-4-carboxylate;
72		3-(2-(2,3-dichlorophenyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)pentan-3-ol;

Cpd #	Structure	Name
73		3-(2-(2,3-dichlorophenyl)-1-(3'-(ethylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)pentan-3-ol;
74		3-(2-(2,6-dichlorophenyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)pentan-3-ol;
80		N-benzyl-3'-(2-chloro-6-fluorophenyl)-4-(2-hydroxypropan-2-yl)-1H-imidazol-1-yl)-3-(methylsulfonyl)biphenyl-4-carboxamide;
81		3'-chloro-4'-(2-(2-chloro-6-fluorophenyl)-4-(2-hydroxypropan-2-yl)-1H-imidazol-1-yl)-N-ethyl-3-(methylsulfonyl)biphenyl-4-carboxamide;
82		2-(2,6-dichlorophenyl)-N-ethyl-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazole-4-carboxamide;
83		(2-(2,6-dichlorophenyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)(4-methylpiperazin-1-yl)methanone;
84		2-(2,6-dichlorophenyl)-N-(2-hydroxyethyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazole-4-carboxamide;

Cpd #	Structure	Name
85		2-(2,6-dichlorophenyl)-1-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-N-(2,2,2-trifluoroethyl)-1H-imidazole-4-carboxamide;
86		2-(2,6-dichlorophenyl)-N,N-diethyl-1-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazole-4-carboxamide
87		2-(2,6-dichlorophenyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-N-(2,2,2-trifluoroethyl)-1H-imidazole-4-carboxamide;
88		2-(2,6-dichlorophenyl)-N-(2-fluoroethyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazole-4-carboxamide;
89		2-(2-isopropylphenyl)-1-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-N-(2,2,2-trifluoroethyl)-1H-imidazole-4-carboxamide;
90		2-(2-(2-(dimethylamino)-6-fluorophenyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
91		2-(2-(2,6-dichlorophenyl)-1-(3'-(ethylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;

Cpd #	Structure	Name
92		2-(2-(2-chloro-6-fluorophenyl)-1-(3'-(ethylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
93		2-(2-(2-ethylphenyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
94		2-(2-(2-isopropylphenyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
95		2-(2-(2-chloro-3-fluorophenyl)-1-(3'-(ethylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
96		2-(1-(3'-(ethylsulfonyl)biphenyl-4-yl)-2-(2-isopropylphenyl)-1H-imidazol-4-yl)propan-2-ol;
97		2-(2-(2-chloro-6-methylphenyl)-1-(3'-(ethylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
98		2-(2-(2,3-dichlorophenyl)-1-(4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
99		2-(2-(2,6-dichlorophenyl)-1-(4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;

Cpd #	Structure	Name
100		2-(2-(2-chloro-6-fluorophenyl)-1-(4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
101		2-(2-(3-chloro-2-methylphenyl)-1-(4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
102		2-(2-(2-chloro-6-methylphenyl)-1-(4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
103		2-(2-(2,6-difluorophenyl)-1-(4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
104		2-(2-(2,6-dichlorophenyl)-1-(3'-(ethylsulfonyl)-3-methylbiphenyl-4-yl)-N-(2,2,2-trifluoroethyl)-1H-imidazole-4-carboxamide;
105		2-(2-(2,3-dichlorophenyl)-1-(4'-(2-hydroxypropan-2-yl)-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
106		2-(2-(2,6-dichlorophenyl)-1-(3'-methyl-5'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
107		2-(1-(2'-chloro-5'-(methylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-1H-imidazol-4-yl)propan-2-ol;

Cpd #	Structure	Name
108		4'-(2-(2,6-dichlorophenyl)-4-(2-hydroxypropan-2-yl)-1H-imidazol-1-yl)-N-methylbiphenyl-3-sulfonamide;
109		2-(1-(3-fluoro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2-(trifluoromethyl)phenyl)-1H-imidazol-4-yl)propan-2-ol;
110		2-(1-(3'-(ethylsulfonyl)-3-fluorobiphenyl-4-yl)-2-(2-(trifluoromethyl)phenyl)-1H-imidazol-4-yl)propan-2-ol;
111		2-(1-(2-fluoro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2-(trifluoromethyl)phenyl)-1H-imidazol-4-yl)propan-2-ol;
112		2-(1-(3'-(ethylsulfonyl)-2-fluorobiphenyl-4-yl)-2-(2-(trifluoromethyl)phenyl)-1H-imidazol-4-yl)propan-2-ol;
113		2-(2-(2,3-dichlorophenyl)-1-(3-fluoro-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
114		2-(2-(2,3-dichlorophenyl)-1-(3'-(ethylsulfonyl)-3-fluorobiphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;

Cpd #	Structure	Name
115		2-(2-(2,6-dichlorophenyl)-1-(3-fluoro-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
116		2-(2-(2,6-dichlorophenyl)-1-(3'-(ethylsulfonyl)-3-fluorobiphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
117		2-(2-(2,3-dichlorophenyl)-1-(3-fluoro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
118		2-(2-(3-chloro-2-methylphenyl)-1-(3-fluoro-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
119		2-(2-(3-chloro-2-methylphenyl)-1-(3'-(ethylsulfonyl)-3-fluorobiphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
120		2-(2-(2,3-dichlorophenyl)-1-(3-fluoro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
121		2-(1-(3-fluoro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2-(trifluoromethyl)phenyl)-1H-imidazol-4-yl)propan-2-ol;

Cpd #	Structure	Name
122		2-(2-(2,6-dichlorophenyl)-1-(3-fluoro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
123		2-(2-(2-chloro-3-fluorophenyl)-1-(3-fluoro-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
124		2-(2-(2-chloro-3-fluorophenyl)-1-(3'-(ethylsulfonyl)-3-fluorobiphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
125		2-(2-(2-chloro-3-fluorophenyl)-1-(3-fluoro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
126		2-(2-(2-chloro-6-fluorophenyl)-1-(3'-(ethylsulfonyl)-3-fluorobiphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
127		2-(2-(2-chloro-6-fluorophenyl)-1-(3-fluoro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
128		2-(2-(2-chloro-6-methylphenyl)-1-(3-fluoro-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;

Cpd #	Structure	Name
129		2-(2-(2-chloro-6-methylphenyl)-1-(3'-(ethylsulfonyl)-3-fluorobiphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
130		2-(2-(2,6-difluorophenyl)-1-(3-fluoro-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
131		2-(2-(2,6-difluorophenyl)-1-(3'-(ethylsulfonyl)-3-fluorobiphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
132		2-(2-(2,6-difluorophenyl)-1-(3'-(ethylsulfonyl)-3-fluorobiphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
133		2-(2-(2-chloro-6-methylphenyl)-1-(3-fluoro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
134		2-(2-(2-chloro-6-fluorophenyl)-1-(3-fluoro-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
135		2-(1-(3-chloro-3'-(trifluoromethyl)phenyl)-2-(2-(methylsulfonyl)biphenyl-4-yl)-2-(2-(trifluoromethyl)phenyl)-1H-imidazol-4-yl)propan-2-ol;

Cpd #	Structure	Name
136		2-(1-(3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl)-2-(2-(trifluoromethyl)phenyl)-1H-imidazol-4-yl)propan-2-ol;
137		2-(1-(3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl)-2-(2-chlorophenyl)-1H-imidazol-4-yl)propan-2-ol;
138		2-(1-(3-chloro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2-chlorophenyl)-1H-imidazol-4-yl)propan-2-ol;
139		2-(1-(3-chloro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2-(trifluoromethyl)phenyl)-1H-imidazol-4-yl)propan-2-ol;
140		2-(1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2,3-dichlorophenyl)-1H-imidazol-4-yl)propan-2-ol;
141		2-(1-(3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl)-2-(2,3-dichlorophenyl)-1H-imidazol-4-yl)propan-2-ol;
142		2-(1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-1H-imidazol-4-yl)propan-2-ol;

Cpd #	Structure	Name
143		2-(2-(3-chloro-2-methylphenyl)-1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
144		2-(2-(3-chloro-2-methylphenyl)-1-(3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
145		2-(1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2-chloro-6-fluorophenyl)-1H-imidazol-4-yl)propan-2-ol;
146		2-(1-(3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl)-2-(2-chloro-6-fluorophenyl)-1H-imidazol-4-yl)propan-2-ol;
147		2-(1-(3-chloro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2-chloro-6-fluorophenyl)-1H-imidazol-4-yl)propan-2-ol;
148		2-(1-(3-chloro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2,3-dichlorophenyl)-1H-imidazol-4-yl)propan-2-ol;
149		2-(1-(3-chloro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-1H-imidazol-4-yl)propan-2-ol;

Cpd #	Structure	Name
150		2-(1-(3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-1H-imidazol-4-yl)propan-2-ol;
151		2-(1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2-chloro-3-fluorophenyl)-1H-imidazol-4-yl)propan-2-ol;
152		2-(1-(3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl)-2-(2-chloro-3-fluorophenyl)-1H-imidazol-4-yl)propan-2-ol;
153		2-(2-(2-chloro-3-fluorophenyl)-1-(3-chloro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
154		2-(1-(3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl)-2-(2-chloro-6-methylphenyl)-1H-imidazol-4-yl)propan-2-ol;
155		2-(1-(3-chloro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2-chloro-6-methylphenyl)-1H-imidazol-4-yl)propan-2-ol;
156		2-(1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2-chloro-6-methylphenyl)-1H-imidazol-4-yl)propan-2-ol;

Cpd #	Structure	Name
157		2-(1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2,6-difluorophenyl)-1H-imidazol-4-yl)propan-2-ol;
158		2-(1-(3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl)-2-(2,6-difluorophenyl)-1H-imidazol-4-yl)propan-2-ol;
159		2-(1-(3-chloro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2,6-difluorophenyl)-1H-imidazol-4-yl)propan-2-ol;
160		2-(2-(3-chloro-2-methylphenyl)-1-(3-chloro-4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
161		2-(1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2-isopropylphenyl)-1H-imidazol-4-yl)propan-2-ol;
162		2-(3'-chloro-4'-(2-(2-chlorophenyl)-4-(2-hydroxypropan-2-yl)-1H-imidazol-1-yl)-3-(methylsulfonyl)biphenyl-4-yl)propan-2-ol;
163		2-(1-(3-chloro-4'-(2-hydroxypropan-2-yl)-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2-(trifluoromethyl)phenyl)-1H-imidazol-4-yl)propan-2-ol;

Cpd #	Structure	Name
164		2-(3'-chloro-4'-(2-(2-chloro-6-fluorophenyl)-4-(2-hydroxypropan-2-yl)-1H-imidazol-1-yl)-3-(methylsulfonyl)biphenyl-4-yl)propan-2-ol;
165		ethyl 3'-chloro-4'-(2-(2-chloro-6-fluorophenyl)-4-(2-hydroxypropan-2-yl)-1H-imidazol-1-yl)-3-(methylsulfonyl)biphenyl-4-carboxylate;
166		(3'-chloro-4'-(2-(2-chloro-6-fluorophenyl)-4-(2-hydroxypropan-2-yl)-1H-imidazol-1-yl)-3-(methylsulfonyl)biphenyl-4-yl)(morpholino)methanone;
167		3'-chloro-4'-(2-(2-chloro-6-fluorophenyl)-4-(2-hydroxypropan-2-yl)-1H-imidazol-1-yl)-N,N-dimethyl-3-(methylsulfonyl)biphenyl-4-carboxamide;
168		2-(1-(3-chloro-5-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2-chlorophenyl)-1H-imidazol-4-yl)propan-2-ol;
169		2-(1-(3-chloro-3'-(ethylsulfonyl)-5-methylbiphenyl-4-yl)-2-(2-chlorophenyl)-1H-imidazol-4-yl)propan-2-ol;
170		2-(2-(2-chlorophenyl)-1-(3-ethyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;

Cpd #	Structure	Name
171		2-(1-(4'-(hydroxymethyl)-3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2-(trifluoromethyl)phenyl)-1H-imidazol-4-yl)propan-2-ol;
172		2-(2-(2-chloro-6-fluorophenyl)-1-(4'-(hydroxymethyl)-3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
173		2-(2-(2-chloro-3-fluorophenyl)-1-(4'-(hydroxymethyl)-3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
174		2-(2-(2,6-dichlorophenyl)-1-(4'-(hydroxymethyl)-3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
175		2-(2-(3-chloro-2-methylphenyl)-1-(4'-(hydroxymethyl)-3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
176		2-(2-(2,3-dichlorophenyl)-1-(4'-(hydroxymethyl)-3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
177		2-(2-(2-chloro-6-methylphenyl)-1-(4'-(hydroxymethyl)-3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;

Cpd #	Structure	Name
178		2-(2-(2,6-difluorophenyl)-1-(4'-(hydroxymethyl)-3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
179		2-(2-(2,3-dichlorophenyl)-1-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
180		2-(2-(2,3-dichlorophenyl)-1-(3'-(ethylsulfonyl)-3-methylbiphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
181		2-(2-(2,6-dichlorophenyl)-1-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
182		2-(2-(2,6-dichlorophenyl)-1-(3'-(ethylsulfonyl)-3-methylbiphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
183		2-(1-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2-(trifluoromethyl)phenyl)-1H-imidazol-4-yl)propan-2-ol;
184		2-(1-(3'-(ethylsulfonyl)-3-methylbiphenyl-4-yl)-2-(2-(trifluoromethyl)phenyl)-1H-imidazol-4-yl)propan-2-ol;
185		2-(2-(2-chloro-3-fluorophenyl)-1-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;

Cpd #	Structure	Name
194		2-(2-(2,6-difluorophenyl)-1-(3'-(ethylsulfonyl)-3-methylbiphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
195		2-(2-(2-isopropylphenyl)-1-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
196		2-(1-(3'-(ethylsulfonyl)-3-methylbiphenyl-4-yl)-2-(2-isopropylphenyl)-1H-imidazol-4-yl)propan-2-ol;
197		2-(1-(2,6-dichlorophenyl)-2-(3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol;
198		2-{1-[3'-(methylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
199		2-{1-[4'-(methylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
200		2-{1-[3'-(ethylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
201		2-{1-[3'-(methylsulfonyl)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;

Cpd #	Structure	Name
202		2-{1-[2'-(trifluoromethyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
203		2-{1-(2',3'-difluorobiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
204		2-(1-{4'-[(1-methylethyl)oxy]biphenyl-4-yl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl)propan-2-ol;
205		2-{1-(4'-fluoro-3'-methylbiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
206		2-{1-(3'-fluorobiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
207		2-{1-(2',4',5'-trifluorobiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
208		2-{1-[5'-fluoro-2'-(methoxy)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;

Cpd #	Structure	Name
209		2-{1-(4'-chlorobiphenyl-4-yl)-2-[(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
210		2-{1-(4-pyrimidin-5-ylphenyl)-2-[(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
211		2-{1-[4'-(hydroxymethyl)biphenyl-4-yl]-2-[(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
212		2-{1-[4'-(dimethylamino)biphenyl-4-yl]-2-[(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
213		2-{1-[4'-(trifluoromethyl)biphenyl-4-yl]-2-[(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
214		2-{1-[4'-(1-methylethyl)biphenyl-4-yl]-2-[(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
215		2-{1-[4'-chloro-2'-(trifluoromethyl)biphenyl-4-yl]-2-[(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
216		2-{1-(2',3',4'-trifluorobiphenyl-4-yl)-2-[(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;

Cpd #	Structure	Name
217		2-{1-(3',4'-difluorobiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
218		2-{1-(2'-chloro-6'-fluoro-3'-methylbiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
219		2-{1-[5'-chloro-2'-(methoxy)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
220		2-{1-[2'-fluoro-5'-(trifluoromethyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
221		2-{1-[2'-(methylthio)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
222		4'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-4-carboxylic acid;
223		2-{1-[4-(1,3-benzodioxol-5-yl)phenyl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
224		2-(1-{4-[6-(methoxy)pyridin-3-yl]phenyl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl)propan-2-ol;

Cpd #	Structure	Name
226		2-{1-(3'-chlorobiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
227		2-{1-[2'-fluoro-5'-(methoxy)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
228		ethyl 4'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-4-carboxylate;
229		2-{1-(4-{2-[(1-methylethyl)oxy]pyridin-3-yl}phenyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
230		2-{1-[3'-chloro-4'-(methoxy)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
232		2-{1-[2'-fluoro-3'-(methoxy)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
233		4'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-3-carboxamide;
234		4'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-4-carboxamide;

Cpd #	Structure	Name
235		2-(1-{4'-[(trifluoromethyl)oxy]biphenyl-4-yl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl)propan-2-ol;
236		2-{1-[4'-fluoro-3'-(trifluoromethyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
237		2-{1-(4'-propylbiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
238		2-{1-[4'-(ethoxy)-3'-(trifluoromethyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
239		2-(1-{2'-(1-methylethyl)oxy}biphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl)propan-2-ol;
240		2-(1-{3'-chloro-4'-[(1-methylethyl)oxy]biphenyl-4-yl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl)propan-2-ol;
241		2-{1-[4-(1H-indol-4-yl)phenyl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
242		2-{1-[4-(methylthio)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;

Cpd #	Structure	Name
243		2-{1-[3'-(hydroxymethyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
244		2-{1-[3'-(ethoxy)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
245		2-{1-(4'-ethylbiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
246		2-{1-(2',4'-difluorobiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
247		2-{1-(3',4'-dichlorobiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
248		2-{1-[2'-chloro-4'-(trifluoromethyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
249		2-{1-(4-naphthalen-2-ylphenyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
250		2-{1-[3'-(methoxy)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol

Cpd #	Structure	Name
251		4'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl} biphenyl-3-ol;
252		2-{1-(3',4',5'-trifluorobiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
253		1-[5-(4-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}phenyl)-2-thienyl]ethanone;
254		2-{1-(3',5'-difluorobiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
255		2-{1-(3'-chloro-4'-fluorobiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
256		2-{1-[5'-methyl-2'-(methyloxy)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
257		2-{1-(2',5'-difluorobiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
258		2-{1-[3'-(butyloxy)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;

Cpd #	Structure	Name
259		2-{1-[5'-chloro-2'-(ethyloxy)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
260		2-(1-{3'-[(trifluoromethyl)oxy]biphenyl-4-yl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl)propan-2-ol;
261		2-{1-(2',3',5'-trifluorobiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
262		2-{1-[3'-(ethylthio)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
263		2-(1-{3'-[(1-methylethyl)oxy]biphenyl-4-yl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl)propan-2-ol;
264		2-{1-[4-(1-benzothien-3-yl)phenyl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
265		2-{1-[4-(4-methylnaphthalen-1-yl)phenyl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
266		2-{1-(2',4'-dichlorobiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;

Cpd #	Structure	Name
267		2-{1-[3',4'-bis(methoxy)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
268		2-{2-[2-(trifluoromethyl)phenyl]-1-(2',4',5'-trimethylbiphenyl-4-yl)-1H-imidazol-4-yl}propan-2-ol;
269		4-fluoro-4'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-2-ol;
270		2-{1-[2'-(hydroxymethyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
271		2-{1-[3'-(trifluoromethyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
272		2-{1-(2'-chloro-6'-fluorobiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
273		2-{1-[3',5'-difluoro-2'-(methoxy)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
274		2-(1-{4-[2-(methoxy)pyridin-3-yl]phenyl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl)propan-2-ol;

Cpd #	Structure	Name
275		2-{1-[2'-methyl-5'-(methoxy)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
276		2-{1-(2'-ethylbiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
277		2-(1-{2'-methyl-4'-[(1-methylethyl)oxy]biphenyl-4-yl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl)propan-2-ol;
278		2-{1-[4'-(ethylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
279		2-{1-(5'-fluoro-2'-methylbiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
280		2-{1-[3'-chloro-4'-(trifluoromethyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
281		2-{1-(5'-chloro-2'-methylbiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
282		2-{1-[2'-(ethoxy)-5'-(trifluoromethyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;

Cpd #	Structure	Name
283		2-{1-(3'-fluoro-4'-methylbiphenyl-4-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
284		methyl (2E)-3-(4'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-4-yl)prop-2-enoate;
285		N-ethyl-4'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-3-carboxamide;
286		2-{1-[4-(2,3-dihydro-1-benzofuran-5-yl)phenyl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
287		2-(1-{3-[6-(methoxy)pyridin-3-yl]phenyl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl)propan-2-ol;
288		3'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}-N-(1-methylethyl)biphenyl-3-sulfonamide;
289		3'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-4-carboxamide;

Cpd #	Structure	Name
290		2-(1-{3-[2-(cyclopentyloxy)pyridin-3-yl]phenyl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
291		2-{1-(3'-chlorobiphenyl-3-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
292		2-{1-[2'-fluoro-5'-(methyloxy)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
293		3'-(4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-3-carboxamide;
294		2-{1-[3',4'-bis(methyloxy)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
295		N-(3'-(4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-2-yl)methanesulfonamide ;
296		2-{1-[3'-(trifluoromethyl)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
297		3-fluoro-3'-(4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-4-carboxylic acid;

Cpd #	Structure	Name
298		4-chloro-3'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-3-carboxamide;
299		2-{1-(3-{2-[(1-methylethyl)oxy]pyridin-3-yl}phenyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
300		N-(3'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-3-yl)acetamide;
301		2-{1-[2'-methyl-5'-(methyloxy)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
302		N-(3'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-4-yl)acetamide;
303		2-{1-(4'-chlorobiphenyl-3-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
304		2-{1-[4'-(phenyloxy)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
305		2-{1-[5'-fluoro-2'-(methyloxy)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;

Cpd #	Structure	Name
306		2-{1-(2',3'-dichlorobiphenyl-3-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
307		2-(1-{3'-[(trifluoromethyl)oxy]biphenyl-3-yl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
308		3'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-3-sulfonamide;
309		2-{1-[3',5'-bis(trifluoromethyl)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
310		2-{1-(3',5'-dichlorobiphenyl-3-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
311		3-chloro-3'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}-N-(1-methylethyl)biphenyl-4-carboxamide;
312		N,N-diethyl-3'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-3-carboxamide;

Cpd #	Structure	Name
313		4-chloro-3'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}-N-(1-methylethyl)biphenyl-3-carboxamide;
314		2-{1-[3'-(methyloxy)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
315		3'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-3-carboxylic acid;
316		N-ethyl-3'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-3-carboxamide;
317		4-chloro-N-ethyl-3'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-3-carboxamide;
318		2-{1-(2',5'-difluorobiphenyl-3-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
319		2-(1-{3'-[(1-methylethyl)oxy]biphenyl-3-yl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl)propan-2-ol;

Cpd #	Structure	Name
320		2-{1-[2'-fluoro-5'-(trifluoromethyl)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
321		2-{1-(3',4'-dichlorobiphenyl-3-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
322		2-{1-[3'-(ethyloxy)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
323		2-(1-{2'-methyl-4'-[(1-methylethyl)oxy]biphenyl-3-yl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl)propan-2-ol;
324		2-{1-[3-(1-methyl-1H-indol-5-yl)phenyl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
325		2-{1-[4'-(ethyloxy)-3'-(trifluoromethyl)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
326		2-{1-[3'-(ethylsulfonyl)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
327		2-{1-[2'-(trifluoromethyl)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;

Cpd #	Structure	Name
328		2-{1-[3'-(hydroxymethyl)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
329		2-{1-[3-(1H-indol-4-yl)phenyl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
330		3'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-4-carboxylic acid;
331		1-[5-(3-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}phenyl)-2-thienyl]ethanone;
332		2-{1-(5'-chloro-2'-methylbiphenyl-3-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
333		2-(1-{4'-[(trifluoromethyl)oxy]biphenyl-3-yl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl)propan-2-ol;
334		2-{1-[2'-chloro-4'-(trifluoromethyl)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
335		2-{1-(2',5'-dichlorobiphenyl-3-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;

Cpd #	Structure	Name
336		2-{1-[2'-(ethyloxy)-5'-(trifluoromethyl)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
337		3'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}-N-(1-methylethyl)biphenyl-4-carboxamide;
338		2-{1-(2',4'-dichlorobiphenyl-3-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
339		2-{1-[4'-(ethylsulfonyl)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
340		2-{1-[4'-fluoro-3'-(trifluoromethyl)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
341		2-{1-(3'-fluoro-4'-methylbiphenyl-3-yl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
342		N-butyl-4'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-3-sulfonamide;

Cpd #	Structure	Name
343		N-(1,1-dimethylethyl)-4'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}-6-methylbiphenyl-3-sulfonamide;
344		4'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}-N-methylbiphenyl-3-sulfonamide;
345		N-ethyl-4'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-3-sulfonamide;
346		N-(1,1-dimethylethyl)-4'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-3-sulfonamide;
347		2-{1-[2'-amino-5'-(trifluoromethyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
348		2-{1-[3'-fluoro-5'-(trifluoromethyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
349		2-{1-[4'-chloro-3'-(trifluoromethyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;

Cpd #	Structure	Name
350		3-chloro-4'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-4-carboxylic acid;
351		2-(2-[2-(trifluoromethyl)phenyl]-1-{3'-[(trifluoromethyl)thio]biphenyl-4-yl}-1H-imidazol-4-yl)propan-2-ol;
352		2-{1-[3'-(morpholin-4-ylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
353		2-(1-{4-[5-(hydroxymethyl)-1,3-thiazol-2-yl]phenyl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl)propan-2-ol;
354		2-{1-[2'-methyl-5'-(piperidin-1-ylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
355		1-[4-(4-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}phenyl)-2-thienyl]ethanone;
356		2-(1-{4-[5-(hydroxymethyl)-3-thienyl]phenyl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl)propan-2-ol;

Cpd #	Structure	Name
358		2-(2-chlorophenyl)-1-[3'-(methylsulfonyl)biphenyl-4-yl]-4-(trifluoromethyl)-1H-imidazole;
360		4'-[2-(2-chlorophenyl)-4-(trifluoromethyl)-1H-imidazol-1-yl]biphenyl-3-sulfonamide;
364		1-[3'-(methylsulfonyl)biphenyl-4-yl]-4-(trifluoromethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazole;
365		4'-{4-(trifluoromethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-3-sulfonamide;

Cpd #	Structure	Name
367		ethyl 1-[3'-(methylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazole-4-carboxylate;
369		5-[3-(methylsulfonyl)phenyl]-2-{4-(trifluoromethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}pyridine;
370		2-[2-(2-chlorophenyl)-4-(trifluoromethyl)-1H-imidazol-1-yl]-5-[3-(methylsulfonyl)phenyl]pyridine;
371		2-{1-[3'-(methylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
372		5-{3-[(1-methylethyl)sulfonyl]phenyl}-2-{4-(trifluoromethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}pyridine;
373		1-{3'[(1-methylethyl)sulfonyl]biphenyl-4-yl}-4-(trifluoromethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazole;
375		N-(4'-{4-(trifluoromethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}biphenyl-3-yl)methanesulfonamide ;
376		ethyl 1-[3'-(methylsulfonyl)biphenyl-4-yl]-2-phenyl-1H-imidazole-4-carboxylate;

Cpd #	Structure	Name
378		2-{1-[3'-(methylsulfonyl)biphenyl-4-yl]-2-phenyl-1H-imidazol-4-yl}propan-2-ol;
379		2-{2-(2-chlorophenyl)-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol;
380		2-{2-(2-fluorophenyl)-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol;
382		2-{2-(2,3-dichlorophenyl)-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol;
383		2-{2-(2-chloro-3-fluorophenyl)-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol;
384		4'-[2-(2-chlorophenyl)-4-(1-hydroxy-1-methylethyl)-1H-imidazol-1-yl]biphenyl-3-sulfonamide;
385		2-{2-(4-fluorophenyl)-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol;
386		2-{2-(2-chlorophenyl)-1-[3-fluoro-3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol;
387		2-{2-(2-chlorophenyl)-1-[2-fluoro-3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol;

Cpd #	Structure	Name
388		2-{2-(2-chlorophenyl)-1-[3'-(1-methylethyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol;
389		2-{2-(2-chlorophenyl)-1-[4'-(1,1-dimethylethyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol;
390		methyl 4'-[2-(2-chlorophenyl)-4-(1-hydroxy-1-methylethyl)-1H-imidazol-1-yl]biphenyl-3-carboxylate;
391		N-{4'-[2-(2-chlorophenyl)-4-(1-hydroxy-1-methylethyl)-1H-imidazol-1-yl]biphenyl-3-yl}-1,3-benzodioxole-5-carboxamide;
392		2-{1-[3'-(methylsulfonyl)biphenyl-4-yl]-2-[3-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
393		2-{2-(3-chlorophenyl)-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol;
394		2-{2-(2-chloro-6-methylphenyl)-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol;
395		4'-[2-(2-chlorophenyl)-4-(1-hydroxy-1-methylethyl)-1H-imidazol-1-yl]biphenyl-3-carboxylic acid;

Cpd #	Structure	Name
397		2-(2-chlorophenyl)-1-[3'-(methylsulfonyl)biphenyl-4-yl]-N-(2,2,2-trifluoroethyl)-1H-imidazole-4-carboxamide;
399		2-{1-[4'-(methoxy)-3'-(methylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
400		2-{1-[4'-(hydroxymethyl)-3'-(methylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
401		ethyl 4'-{4-(1-hydroxy-1-methylethyl)-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-1-yl}-5-(methylsulfonyl)biphenyl-3-carboxylate;
402		2-{5-bromo-1-[3'-(methylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
403		2-{5-chloro-1-[3'-(methylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
404		2-{1-[4'-(methylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;

Cpd #	Structure	Name
405		2-[1-(3'-aminobiphenyl-4-yl)-2-(2-chlorophenyl)-1H-imidazol-4-yl]propan-2-ol;
406		2-{5-fluoro-1-[3'-(methylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
407		2-{1-[4'-(methylsulfonyl)biphenyl-3-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
409		2-{2-(2-chlorophenyl)-1-[3'-(1-hydroxy-1-methylethyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol;
410		2-{2-(2,6-difluorophenyl)-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol;
413		2-(1-{3'-[(1-methylethyl)sulfonyl]biphenyl-4-yl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl)propan-2-ol;
414		2-(1-{3'-[(1-methylethyl)sulfonyl]biphenyl-3-yl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl)propan-2-ol;
415		2-{2-(2-chlorophenyl)-1-[3'-(ethylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol;
416		2-{2-(2-chloro-6-fluorophenyl)-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol;

Cpd #	Structure	Name
417		2-{1-[3-chloro-3'-(methylsulfonyl)biphenyl-4-yl]-2-(2-chlorophenyl)-1H-imidazol-4-yl}propan-2-ol;
418		2-{2-(2-chlorophenyl)-1-[3-methyl-3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol;
419		2-{2-(2,6-difluorophenyl)-1-[3'-(ethylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol;
420		2-(1-{5-[3-(methylsulfonyl)phenyl]pyridin-2-yl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl)propan-2-ol;
421		2-(1-{6-[3-(methylsulfonyl)phenyl]pyridin-3-yl}-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl)propan-2-ol;
422		2-{2-(2,6-dichlorophenyl)-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol;
427		2-{2-(2,3-dichlorophenyl)-1-[3'-(ethylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol;
428		2-{5-methyl-1-[3'-(methylsulfonyl)biphenyl-4-yl]-2-[2-(trifluoromethyl)phenyl]-1H-imidazol-4-yl}propan-2-ol;
429		2-{2-(2-chlorophenyl)-1-[3'-(ethylsulfonyl)-3-fluorobiphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol;

Cpd #	Structure	Name
430		2-{2-[2-fluoro-6-(trifluoromethyl)phenyl]-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol;
431		2-{2-(2-methylphenyl)-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol;
432		2-(1-[3'-(methylsulfonyl)biphenyl-4-yl]-2-{2-[(trifluoromethyl)oxy]phenyl}-1H-imidazol-4-yl)propan-2-ol;
436		2-(1-(2',5'-dimethylbiphenyl-4-yl)-2-(2-trifluoromethyl)phenyl)-1H-imidazol-4-yl)propan-2-ol;
437		2-{2-(2-chlorophenyl)-1-[3'-(ethylsulfonyl)-2-fluorobiphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol;
438		2-{2-(3-chloro-2-methylphenyl)-1-[3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol;
439		2-{2-(2-chlorophenyl)-1-[2-methyl-3'-(methylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol;
440		2-{2-(3-chloro-2-methylphenyl)-1-[3'-(ethylsulfonyl)biphenyl-4-yl]-1H-imidazol-4-yl}propan-2-ol;

Cpd #	Structure	Name
443		2-(1-(2-chlorobenzyl)-2-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol
444		2-(1-(2,3-dichlorobenzyl)-2-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol
445		2-(5-chloro-2-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-1-(2,6-dichlorophenyl)-1H-imidazol-4-yl)propan-2-ol
446		2-(5-chloro-1-(2,6-dichlorophenyl)-2-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol
447		3'-(chloro-4'-(2-(2-chloro-6-fluorophenyl)-4-(2-hydroxypropan-2-yl)-1H-imidazol-1-yl)-3-(methylsulfonyl)biphenyl-4-carboxylic acid
448		3'-(chloro-4'-(2-(2-chloro-6-fluorophenyl)-4-(2-hydroxypropan-2-yl)-1H-imidazol-1-yl)-3-(methylsulfonyl)biphenyl-4-carboxamide
449		2-(2-(2,3-dichlorobenzyl)-1-(3-fluoro-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol

Cpd #	Structure	Name
450		2-(2-(2,3-dichlorobenzyl)-1-(3'-(ethylsulfonyl)-3-fluorobiphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol
451		2-(1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2-(2,3-dichlorophenyl)propan-2-yl)-1H-imidazol-4-yl)propan-2-ol
452		2-(1-(3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl)-2-(2-(2,3-dichlorophenyl)propan-2-yl)-1H-imidazol-4-yl)propan-2-ol
453		3'-(3-chloro-4'-(2-(2-chloro-6-fluorophenyl)-4-(2-hydroxypropan-2-yl)-1H-imidazol-1-yl)-3-(methylsulfonyl)biphenyl-4-carboxamide
454		2-(1-(3-chloro-4'-(methoxy-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2-chloro-6-fluorophenyl)-1H-imidazol-4-yl)propan-2-ol
455		2-(2-(2,3-dichlorophenyl)-1-(4'-(methoxy-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol
456		2-(2-(2,3-dichlorobenzyl)-1-(4'-(methoxy-3-methyl-3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol

Cpd #	Structure	Name
457		2-(1-(3-chloro-4'-(methoxymethyl)-3'-methoxy-3'-methylsulfonyl)biphenyl-4-yl)-2-(2,3-dichlorobenzyl)-1H-imidazol-4-ylpropan-2-ol
458		2-(2-(3-chloro-3'-(methoxymethyl)-2'-methylsulfonyl)biphenyl-2-yl)-1-(3'-(methoxymethyl)-3'-methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-ylpropan-2-ol
459		1-(3-chloro-3'-(methoxymethyl)-3'-methylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-N-(2,2,2-trifluoroethyl)-1H-imidazole-4-carboxamide
460		1-(3-chloro-3'-(methoxymethyl)-3'-methylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-N-(2-fluoroethyl)-1H-imidazole-4-carboxamide
461		2-(1-(3-chloro-3'-(methoxymethyl)-3'-methylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-1H-imidazole-4-carboxamido)acetic acid
462		1-(3-chloro-3'-(ethoxymethyl)-3'-methylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-N-(2,2,2-trifluoroethyl)-1H-imidazole-4-carboxamide
463		1-(3-chloro-3'-(methoxymethyl)-3'-methylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-N-(1-hydroxy-2-methylpropan-2-yl)-1H-imidazole-4-carboxamide

Cpd #	Structure	Name
464		1-(3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-N-(1-hydroxy-2-methylpropan-2-yl)-1H-imidazole-4-carboxamide
465		N-tert-butyl-1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-1H-imidazole-4-carboxamide
466		2-chloro-4'-(2-(2,6-dichlorophenyl)-4-(2-hydroxypropan-2-yl)-1H-imidazol-1-yl)-3'-methylbiphenyl-4-carboxylic acid
467		2-(1-(2',3-dichloro-5'-(methylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-1H-imidazol-4-yl)propan-2-ol
468		2-(1-(2'-chloro-3-methyl-5'-(methylsulfonyl)biphenyl-4-yl)-2-(2,3-dichlorobenzyl)-1H-imidazol-4-yl)propan-2-ol
469		6-{3-chloro-4-[2-(2-chloro-6-fluorophenyl)-4-(1-hydroxy-1-methylethyl)-1H-imidazol-1-yl]phenyl}-2,3-dihydro-1-benzothiophene-3-ol 1,1-dioxide
470		1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-N-(2-hydroxyethyl)-1H-imidazole-4-carboxamide


Cpd #	Structure	Name
471		1-(3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-N-(2-hydroxyethyl)-1H-imidazole-4-carboxamide
472		1-(3-chloro-3'-(methylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-N-(2-hydroxy-2-methylpropyl)-1H-imidazole-4-carboxamide
473		1-(3-chloro-3'-(ethylsulfonyl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-N-(2-hydroxy-2-methylpropyl)-1H-imidazole-4-carboxamide
474		2-(2-(5-chloro-2-fluorobenzyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol
475		2-(2-(2,3-dichlorophenyl)-1-(4'-(hydroxymethyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol
476		2-(2-(2,3-dichlorophenyl)-1-(3'-(hydroxymethyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol
477		2-(2-(2,3-dichlorophenyl)-1-(4'-(2-hydroxypropan-2-yl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol

Cpd #	Structure	Name
478		2-(2-(2,3-dichlorophenyl)-1-(3'-(2-hydroxypropan-2-yl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol
479		2-(2-(2,6-dichlorophenyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)-2-methylpropan-1-ol
480		1-(3-chloro-3'-(2-hydroxypropan-2-yl)biphenyl-4-yl)-2-(2,6-dichlorophenyl)-N-(1-hydroxy-2-methylpropan-2-yl)-1H-imidazole-4-carboxamide
481		2-(2-(2,6-dichlorophenyl)-N-(1-hydroxy-2-methylpropan-2-yl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazole-4-carboxamide
482		2-(2-(5-chloro-2-fluorobenzyl)-1-(3'-(ethylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol
483		2-(2-(3-chloro-2-fluorobenzyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazol-4-yl)propan-2-ol
484		2-(1-(3'-(methylsulfonyl)biphenyl-4-yl)-2-(3-(trifluoromethyl)benzyl)-1H-imidazol-4-yl)propan-2-ol

Cpd #	Structure	Name
485		2-(1-(3'-(ethylsulfonyl)biphenyl-4-yl)-2-(3-(trifluoromethyl)benzyl)-1H-imidazol-4-yl)propan-2-ol
486		2-(2,6-dichlorophenyl)-1-(3'-(ethylsulfonyl)biphenyl-4-yl)-N-(1-hydroxy-2-methylpropan-2-yl)-1H-imidazole-4-carboxamide, and
487		N-tert-butyl-2-(2,6-dichlorophenyl)-1-(3'-(methylsulfonyl)biphenyl-4-yl)-1H-imidazole-4-carboxamide.

25. A pharmaceutical composition comprising a compound of claim 8, or a pharmaceutically acceptable salt thereof in a pharmaceutically acceptable carrier.


26. The compound according to claim 17, of the formula,



or a pharmaceutically acceptable salt thereof.

27. The compound according to claim 4, wherein R^2 is -halogen, -CF₃, -CH₂OH, or -C(CH₃)₂OH.

28. A compound which is



or a pharmaceutically acceptable salt thereof.

29. The compound of claim 23, wherein the compound is according to formula IIa.

30. The compound of claim 23, wherein the compound is according to formula IIb.

31. A pharmaceutical composition comprising a compound of claim 28, or a pharmaceutically acceptable salt thereof in a pharmaceutically acceptable carrier.