28. ~~A method of treating cancer in a subject, the method comprising administering to a subject~~ Use of a composition of any one of claims 1-27 in the manufacture of a medicament for treating cancer in a subject.

29. The composition of any one of claims 11-16, wherein the peptide is an antibody or antibody fragment comprising:

a V_H chain with identity to SEQ ID NO:222, wherein regions corresponding to CDR1, CDR2, and CDR3 comprise CDR1, CDR2, and CDR3 of the V_H of antibody ID-10 shown in table 1 having 5 or fewer conservative amino acid substitutions within the CDR1, CDR2, and CDR3 regions, and regions within SEQ ID NO:74 corresponding to FR1, FR2, FR3, FR4, comprise amino acid sequences with at least 80%, 85%, 90%, 95%, 96%, 97%, 98, 99%, or 100% identity to FR1, FR2, FR3, FR4 of the V_H of antibody ID-10 shown in table 1; and

a V_L chain with identity to SEQ ID NO:224, wherein regions corresponding to CDR1, CDR2, and CDR3 comprise CDR1, CDR2, and CDR3 of the V_L of antibody ID-10 shown in table 1 having 5 or fewer conservative amino acid substitutions within the CDR1, CDR2, and CDR3 regions, and regions within SEQ ID NO:224 corresponding to FR1, FR2, FR3, FR4, comprise amino acid sequences with at least 80%, 85%, 90%, 95%, 96%, 97%, 98, 99%, or 100% identity to FR1, FR2, FR3, FR4 of the V_L of antibody ID-10 shown in table 1.

30. The composition of claim 23, wherein the peptide is an antibody or antibody fragment comprising a V_H chain comprising SEQ ID NO:222 and a V_L chain comprising SEQ ID NO:224.

31. The composition of any one of claims 1-8 and 11-16 further comprising an histone deacetylase inhibitor (HDAC) selected from the group consisting of hydroxamic acid, vorinostat, suberoylanilide hydroxamic acid (SAHA, trichostatin A (TSA), LAQ824, panobinostat (LBH589), belinostat (PXD101), ITF2357 italfarmaco SpA, cyclic tetrapeptide, depsipeptide (romidepsin, FK228), benzamide; entinostat (SNDX-