

## Anti-IL-11R $\alpha$ Antibodies

### Cross-Reference to Related Applications

This application claims priority to U.S. Provisional Application No. 63/238,443, filed August 30, 2021; and U.S. Provisional Application No. 63/334,923, filed April 26, 2022, each of which is incorporated by reference in its entirety.

### Statement Regarding Sequence Listing

The Sequence Listing XML associated with this application is provided in XML file format and is hereby incorporated by reference into the specification. The name of the XML file containing the Sequence Listing XML is LASS\_005\_02WO\_ST26.xml. The XML file is about 328,887 bytes, was created on August 15, 2022, and is being submitted electronically via USPTO Patent Center.

### Background

#### *Technical Field*

The present disclosure relates to antibodies and antigen binding fragments thereof that bind to human interleukin-11 receptor subunit  $\alpha$  (IL-11R $\alpha$ ) and related compositions, which may be used in any of a variety of therapeutic or diagnostic methods, including the treatment or diagnosis of cancers, inflammatory diseases, autoimmune diseases, and others.

#### *Description of the Related Art*

Interleukin-11 (IL-11) is a member of the IL-6 family, and plays a prominent role in chronic inflammation, autoimmunity, cancer, and other diseases. It also plays a critical role in the initiation and maintenance of chronic fibrotic responses. Thus, IL-11 signaling inhibitors represent a promising therapeutic approach for treating a variety of diseases, including inflammatory diseases, autoimmune diseases, chronic fibrotic diseases, and cancers.

Exemplary IL-11 signaling inhibitors under development include antibodies that bind to the interleukin-11 receptor subunit  $\alpha$  (IL-11R $\alpha$ ) (see, for example, U.S. Patent Nos. 9,796,782; and 9,340,618). However, there is a need in the art for anti-IL-11R $\alpha$  antibodies with increased potency and optimal developability characteristics.

### Brief Summary

Embodiments of the present disclosure include an isolated antibody, or an antigen binding fragment thereof, which binds to interleukin-11 receptor subunit  $\alpha$  (IL-11R $\alpha$ ), wherein the at least one antibody, or antigen binding fragment thereof, comprises:

a heavy chain variable region ( $V_H$ ) that comprises complementary determining region  $V_HCDR1$ ,  $V_HCDR2$ , and  $V_HCDR3$  sequences selected from **Table A1** and variants thereof which specifically bind to IL-11R $\alpha$ ; and

a light chain variable region ( $V_L$ ) that comprises complementary determining region  $V_LCDR1$ ,  $V_LCDR2$ , and  $V_LCDR3$  sequences selected from **Table A1** and variants thereof which specifically bind to IL-11R $\alpha$ .

In some embodiments:

the  $V_HCDR1$ ,  $V_HCDR2$ , and  $V_HCDR3$  sequences comprise SEQ ID NOs: 1-3, respectively, and the  $V_LCDR1$ ,  $V_LCDR2$ , and  $V_LCDR3$  sequences comprise SEQ ID NOs: 4-6, respectively;

the  $V_HCDR1$ ,  $V_HCDR2$ , and  $V_HCDR3$  sequences comprise SEQ ID NOs: 7-9, respectively, and the  $V_LCDR1$ ,  $V_LCDR2$ , and  $V_LCDR3$  sequences comprise SEQ ID NOs: 10-12, respectively;

the  $V_HCDR1$ ,  $V_HCDR2$ , and  $V_HCDR3$  sequences comprise SEQ ID NOs: 13-15, respectively, and the  $V_LCDR1$ ,  $V_LCDR2$ , and  $V_LCDR3$  sequences comprise SEQ ID NOs: 16-18, respectively;

the  $V_HCDR1$ ,  $V_HCDR2$ , and  $V_HCDR3$  sequences comprise SEQ ID NOs: 19-21, respectively, and the  $V_LCDR1$ ,  $V_LCDR2$ , and  $V_LCDR3$  sequences comprise SEQ ID NOs: 22-24, respectively;

the  $V_HCDR1$ ,  $V_HCDR2$ , and  $V_HCDR3$  sequences comprise SEQ ID NOs: 25-27, respectively, and the  $V_LCDR1$ ,  $V_LCDR2$ , and  $V_LCDR3$  sequences comprise SEQ ID NOs: 28-30, respectively;

the  $V_HCDR1$ ,  $V_HCDR2$ , and  $V_HCDR3$  sequences comprise SEQ ID NOs: 31-33, respectively, and the  $V_LCDR1$ ,  $V_LCDR2$ , and  $V_LCDR3$  sequences comprise SEQ ID NOs: 34-36, respectively;

the  $V_HCDR1$ ,  $V_HCDR2$ , and  $V_HCDR3$  sequences comprise SEQ ID NOs: 34-39, respectively, and the  $V_LCDR1$ ,  $V_LCDR2$ , and  $V_LCDR3$  sequences comprise SEQ ID NOs: 40-42, respectively;

the  $V_HCDR1$ ,  $V_HCDR2$ , and  $V_HCDR3$  sequences comprise SEQ ID NOs: 43-45, respectively, and the  $V_LCDR1$ ,  $V_LCDR2$ , and  $V_LCDR3$  sequences comprise SEQ ID NOs: 46-48, respectively;

the  $V_HCDR1$ ,  $V_HCDR2$ , and  $V_HCDR3$  sequences comprise SEQ ID NOs: 49-51, respectively, and the  $V_LCDR1$ ,  $V_LCDR2$ , and  $V_LCDR3$  sequences comprise SEQ ID NOs: 52-54, respectively;

the  $V_HCDR1$ ,  $V_HCDR2$ , and  $V_HCDR3$  sequences comprise SEQ ID NOs: 55-57, respectively, and the  $V_LCDR1$ ,  $V_LCDR2$ , and  $V_LCDR3$  sequences comprise SEQ ID NOs: 58-60, respectively;

the  $V_HCDR1$ ,  $V_HCDR2$ , and  $V_HCDR3$  sequences comprise SEQ ID NOs: 61-63, respectively, and the  $V_LCDR1$ ,  $V_LCDR2$ , and  $V_LCDR3$  sequences comprise SEQ ID NOs: 64-66, respectively;

the  $V_HCDR1$ ,  $V_HCDR2$ , and  $V_HCDR3$  sequences comprise SEQ ID NOs: 67-69, respectively, and the  $V_LCDR1$ ,  $V_LCDR2$ , and  $V_LCDR3$  sequences comprise SEQ ID NOs: 70-72, respectively;

the  $V_HCDR1$ ,  $V_HCDR2$ , and  $V_HCDR3$  sequences comprise SEQ ID NOs: 73-75, respectively, and the  $V_LCDR1$ ,  $V_LCDR2$ , and  $V_LCDR3$  sequences comprise SEQ ID NOs: 76-78, respectively;

the  $V_HCDR1$ ,  $V_HCDR2$ , and  $V_HCDR3$  sequences comprise SEQ ID NOs: 79-81, respectively, and the  $V_LCDR1$ ,  $V_LCDR2$ , and  $V_LCDR3$  sequences comprise SEQ ID NOs: 82-84, respectively;

the  $V_HCDR1$ ,  $V_HCDR2$ , and  $V_HCDR3$  sequences comprise SEQ ID NOs: 85-87, respectively, and the  $V_LCDR1$ ,  $V_LCDR2$ , and  $V_LCDR3$  sequences comprise SEQ ID NOs: 88-90, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 91-93, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 94-96, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 97-99, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 100-102, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 103-105, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 106-108, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 109-111, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 112-114, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 115-117, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 118-120, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 121-123, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 124-126, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 127-129, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 130-132, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 133-135, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 136-138, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 139-141, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 142-144, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 145-147, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 148-150, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 151-153, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 154-156, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 157-159, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 160-162, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 163-165, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 166-168, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 169-171, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 172-174, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 175-177, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 178-180, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 181-183, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 184-186, respectively; or

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 187-189, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 190-192, respectively.

In some embodiments, the V<sub>H</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to a sequence selected from **Table A2**, optionally wherein the V<sub>H</sub> has 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 alterations in the framework regions. In some embodiments, the V<sub>L</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to a sequence selected from **Table A2**, optionally wherein the V<sub>L</sub> has 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 alterations in the framework regions.

In certain embodiments:

the V<sub>H</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 193, and the V<sub>L</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 194;

the V<sub>H</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 195, and the V<sub>L</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 196;

the V<sub>H</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 197, and the V<sub>L</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 198;

the V<sub>H</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 199, and the V<sub>L</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 200;

the V<sub>H</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 201, and the V<sub>L</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 202;

the V<sub>H</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 203, and the V<sub>L</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 204;





the V<sub>H</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 253, and the V<sub>L</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 254; or

the V<sub>H</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 255, and the V<sub>L</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 256.

In some embodiments, the isolated antibody, or antigen binding fragment thereof, binds to human IL-11R $\alpha$  (see **Table B1**). In specific embodiments, the isolated antibody, or antigen binding fragment thereof, binds to a fibronectin domain III of human IL-11R $\alpha$ , or approximately residues 112-219 of SEQ ID NO: 257. In some embodiments, the isolated antibody, or antigen binding fragment thereof, has one or more of the following characteristics:

a binding affinity for human IL-11R $\alpha$  of less than about 500, 400, 300, 200, 100, 90, 80, 70, 60, 50, 40, or 30 pM, and optionally has increased binding affinity for human IL-11R $\alpha$  relative to that of the TS7 and 8E2 antibodies; antagonizes the binding and/or signaling activity between IL-11R $\alpha$  and IL-11, and optionally has increased potency as an IL-11 signaling antagonist relative to that of the TS7 and 8E2 antibodies; reduces IL-11R $\alpha$ /gp130 dimerization or complex formation, optionally in a cell-based assay; and/or reduced N-linked glycosylation in V<sub>L</sub>CDR3, optionally relative to the TS7 and 8E2 antibodies.

In some embodiments, an isolated antibody, or antigen binding fragment thereof, comprises an IgA (including subclasses IgA1 and IgA2), IgD, IgE, IgG (including subclasses IgG1, IgG2, IgG3, and IgG4), or IgM Fc domain, optionally a human Fc domain, or a hybrid and/or variant thereof. In some embodiments, an isolated antibody, or antigen binding fragment thereof, comprises an IgG Fc domain with high effector function in humans, optionally an IgG1 or IgG3 Fc domain. In some embodiments, an isolated antibody, or antigen binding fragment thereof, comprises an IgG Fc domain with low effector function in humans, optionally an IgG2 or IgG4 Fc domain. In specific embodiments, an isolated antibody, or antigen binding fragment thereof, comprises a human IgG1 or IgG4 Fc domain, optionally selected from **Table F1**.

In some embodiments, an isolated antibody, or antigen binding fragment thereof, is a monoclonal antibody. In some embodiments, an isolated antibody, or antigen binding fragment thereof, comprises a humanized antibody, including wherein the antibody, or antigen binding fragment thereof, is a humanized monoclonal antibody that comprises a human IgG4 Fc domain with an S228P mutation (EU numbering). In some embodiments, an isolated antibody, or antigen binding fragment thereof, is selected from an Fv fragment, a single chain Fv (scFv) polypeptide, an adnectin, an anticalin, an aptamer, an avimer, a camelid antibody, a designed ankyrin repeat protein (DARPin), a minibody, a nanobody, and a unibody.

Also included are isolated polynucleotide(s) encoding an isolated antibody, or antigen binding fragment thereof, as described herein, expression vector(s) comprising the isolated polynucleotide(s), and an isolated host cell(s) comprising the vector(S).

Certain embodiments relate to pharmaceutical compositions, comprising an isolated antibody, or antigen binding fragment thereof, described herein, and a pharmaceutically-acceptable carrier. In some embodiments, the composition has a purity of at least about 80%, 85%, 90%, 95%, 98%, or 99% on a protein basis with respect to the at least one antibody or antigen binding fragment, and is substantially aggregate- and endotoxin-free. In some embodiments, the composition has reduced or undetectable heterogeneity of N-linked glycosylation (optionally relative to the TS7 and 8E2 antibodies), optionally in the V<sub>L</sub>CDR3 sequence. In some embodiments, the composition is a sterile, injectable solution, optionally suitable for intravenous, intramuscular, subcutaneous, or intraperitoneal administration.

Also included are methods of treating a disease or condition in a subject in need thereof, comprising administering to the subject a pharmaceutical composition described herein. In some embodiments, the disease or condition is an IL-11-associated or IL-11-mediated disease or condition. In some embodiments, the disease or condition is a cancer, an inflammatory disease, an autoimmune disease, a wasting disease, a bone disease, or fibrosis.

In some embodiments, the disease is a cancer, optionally a cancer that expresses or overexpresses IL-11R $\alpha$  and/or IL-11, optionally wherein the cancer displays IL-11R $\alpha$ /IL-11-dependent growth, adhesion, migration, invasion, and/or chemoresistance. In some embodiments, the cancer is selected from one or more of bone cancer, prostate cancer, melanoma (e.g., metastatic melanoma), pancreatic cancer, small cell lung cancer, non-small cell lung cancer (NSCLC), mesothelioma, leukemia (e.g., lymphocytic leukemia, chronic myelogenous leukemia, acute myeloid leukemia, relapsed acute myeloid leukemia, hairy cell leukemias, acute lymphoblastic leukemias), lymphoma (e.g., non-Hodgkin's lymphomas, Hodgkin's lymphoma), hepatoma (hepatocellular carcinoma), sarcoma, B-cell malignancy, breast cancer, ovarian cancer, colorectal cancer, glioma, glioblastoma multiforme, meningioma, pituitary adenoma, vestibular schwannoma, primary CNS lymphoma, primitive neuroectodermal tumor (medulloblastoma), kidney cancer (e.g., renal cell carcinoma), bladder cancer, uterine cancer, esophageal cancer, brain cancer, head and neck cancers, cervical cancer, testicular cancer, thyroid cancer, and stomach cancer. In some embodiments, the cancer is a metastatic cancer, optionally a metastatic cancer which has metastasized to the bone.

In particular embodiments, the inflammatory disease is selected from one or more of airway or lung inflammation (e.g., inflammatory lung disease), asthma, rhinitis, chronic obstructive pulmonary disorder (COPD), dermatitis, psoriasis, hepatitis, gastric inflammation, irritable bowel syndrome (IBS), ulcerative colitis, Crohn's disease, colitis, diverticulitis, lupus erythematosus, nephritis, Parkinson's disease, multiple sclerosis (MS), Alzheimer's disease, arthritis, rheumatoid

arthritis, sepsis, infection-induced inflammation, cardiovascular diseases such as atherosclerosis and vasculitis, diabetes, and gout.

In certain embodiments, the autoimmune disease is selected from one or more of arthritis (including rheumatoid arthritis, reactive arthritis), systemic lupus erythematosus (SLE), psoriasis, inflammatory bowel disease (IBD), ulcerative colitis, Crohn's disease, encephalomyelitis, uveitis, myasthenia gravis, multiple sclerosis, insulin dependent diabetes, Addison's disease, celiac disease, chronic fatigue syndrome, autoimmune hepatitis, autoimmune alopecia, ankylosing spondylitis, fibromyalgia, pemphigus vulgaris, Sjogren's syndrome, Kawasaki's Disease, hyperthyroidism/Graves' disease, hypothyroidism/Hashimoto's disease, endometriosis, scleroderma, pernicious anemia, Goodpasture syndrome, Guillain-Barré syndrome, Wegener's disease, glomerulonephritis, aplastic anemia (including multiply transfused aplastic anemia patients), paroxysmal nocturnal hemoglobinuria, myelodysplastic syndrome, idiopathic thrombocytopenic purpura, autoimmune hemolytic anemia, Evan's syndrome, Factor VIII inhibitor syndrome, systemic vasculitis, dermatomyositis, polymyositis, rheumatic fever, autoimmune lymphoproliferative syndrome (ALPS), autoimmune bullous pemphigoid, Parkinson's disease, sarcoidosis, vitiligo, primary biliary cirrhosis, and autoimmune myocarditis.

In some embodiments, the wasting disease is selected from one or more of cachexia, optionally cachexia associated with cancer or renal failure, and sarcopenia.

In some embodiments, the bone disease is selected from osteoporosis (including post-menopausal osteoporosis), bone fracture, Paget's disease of bone, and bone resorption/damage associated with cancer or cancer therapy, including chemotherapy, hormone ablation, and hormone inhibition.

In some embodiments, the fibrosis is selected from fibrosis of the lungs, cardiovascular system, liver, brain, joints (optionally knee, hip, ankle, foot joints, shoulder, elbow, wrist, hand joints, or spinal vertebrae), intestine, skin, kidney, liver, thyroid, bone marrow, retroperitoneum, and eye. In certain embodiments, the fibrosis of the lungs is selected from fibrothorax, pulmonary fibrosis (optionally cystic fibrosis or interstitial lung disease (ILD)), autosomal recessive genetic disease optionally Hermansky-Pudlak syndrome, and radiation-induced lung injury. In specific embodiments, the ILD is idiopathic ILD, optionally selected from idiopathic pulmonary fibrosis (IPF), desquamative interstitial pneumonia (DIP), acute interstitial pneumonia (AIP) or Hamman-Rich syndrome, nonspecific interstitial pneumonia (NSIP), respiratory bronchiolitis-associated interstitial lung disease (RB-ILD), cryptogenic organizing pneumonia (COP), and lymphoid interstitial pneumonia (LIP). In certain embodiments, the ILD is secondary ILD, optionally selected from ILD related to connective tissue and autoimmune diseases (optionally sarcoidosis, rheumatoid arthritis, systemic lupus erythematosus, systemic sclerosis, polymyositis, dermatomyositis, or antisynthetase syndrome), inhaled substances (optionally silicosis, asbestosis, berylliosis, industrial printing chemicals, or chronic hypersensitivity pneumonitis), drugs (optionally antibiotics, chemotherapeutics, or anti-

arrhythmic agents), infections (optionally SARS CoV-2, atypical pneumonia, pneumocystis pneumonia, tuberculosis, Chlamydia trachomatis, or respiratory syncytial virus), malignancies (optionally lymphangitic carcinomatosis), and pediatric ILDs (optionally developmental disorders, growth abnormalities deficient alveolarization, infant conditions of undefined cause, and ILD related to alveolar surfactant region).

In some embodiments, the fibrosis of the cardiovascular system is myocardial fibrosis (for example, interstitial fibrosis or replacement fibrosis).

### **Brief Description of the Figures**

**Figures 1A-1B** shows the gross appearance of lung tissue from Donor 1 (**1A**) and Donor 2 (**1B**).

**Figures 2A-2B** show the hematoxylin and eosin staining of the fibrotic lung tissue from donor 1 (**2A**, top subpleural; bottom central) and Donor 2 (**2B**, top subpleural; bottom central). Staining was done on tissue adjacent to the PCLS tissue cores. Representative overview pictures are shown.

**Figures 3A-3B** show the tissue viability of untreated (medium) and LPS treated Precision-Cut Lung Slices (PCLS) from Donor 1 (**3A**) and Donor 2 (**3B**). LDH release is given as % of Triton-lysed control (set to 100%). Bars show mean  $\pm$  SD, with dots representing technical replicates (duplicate wells with 2 PCLS each). For donor 2 (**3B**), supernatants of duplicate wells were pooled before measurement. Region 1 is subpleural, region 2 is central.

**Figures 4A-4B** show IL-1 $\beta$  upregulation in LPS treated PCLS from Donor 1 (**4A**) and Donor 2 (**4B**). IL-1 $\beta$  was determined in PCLS lysates by ELISA and normalized to total protein concentration of the tissue. Data are shown as mean  $\pm$  SD. For each condition, lysates from duplicate wells (with 2 PCLS each) were pooled before measurement. Region 1 is subpleural, region 2 is central.

**Figure 5** shows type VI collagen formation evaluated by PRO-C6 across all donors and regions. Region 1 is subpleural, region 2 is central. Data are presented as percentage of the no treatment (media) control and shown as individual data points for three replicates from each region from each donor (total of 12) and with indication of the median and interquartile range.

### **Detailed Description**

The present disclosure relates to antibodies, and antigen binding fragments thereof, which specifically bind to interleukin-11 receptor subunit  $\alpha$  (IL-11R $\alpha$ ), in particular antibodies having epitopic specificity and improved characteristics. Examples of such improved characteristics include increased binding affinity for IL-11R $\alpha$ , increased potency as interleukin-11 (IL-11) signaling inhibitors/antagonists, and improved developability, such as increased manufacturing homogeneity, for example, due to decreased heterogeneity at potential N-linked glycosylation sites. Some

embodiments thus include specific humanized antibodies and fragments thereof capable of binding to IL-11R $\alpha$ , blocking IL-11R $\alpha$  binding with its ligand IL-11, and inhibiting downstream cell signaling and biological effects. In certain embodiments, an anti-IL-11R $\alpha$  antibody, or antigen binding fragment thereof, is a IL-11R $\alpha$  antagonist or inhibitor.

IL-11R $\alpha$  antagonist antibodies described herein are useful in the treatment and prevention of various diseases and conditions, such as cancers, autoimmune diseases, and inflammatory diseases, including diseases and conditions associated with or mediated by IL-11 signaling. Some embodiments thus relate to the use of anti- IL-11R $\alpha$  antibodies, or antigen binding fragments thereof, for the diagnosis, assessment, and treatment of diseases and conditions, including those associated with IL-11 activity or aberrant expression thereof.

The practice of the present disclosure will employ, unless indicated specifically to the contrary, conventional methods of virology, immunology, microbiology, molecular biology and recombinant DNA techniques within the skill of the art, many of which are described below for the purpose of illustration. Such techniques are explained fully in the literature. See, *e.g.*, *Current Protocols in Molecular Biology* or *Current Protocols in Immunology*, John Wiley & Sons, New York, N.Y.(2009); Ausubel *et al.*, *Short Protocols in Molecular Biology*, 3<sup>rd</sup> ed., Wiley & Sons, 1995; Sambrook and Russell, *Molecular Cloning: A Laboratory Manual* (3rd Edition, 2001); Maniatis *et al.* *Molecular Cloning: A Laboratory Manual* (1982); *DNA Cloning: A Practical Approach*, vol. I & II (D. Glover, ed.); *Oligonucleotide Synthesis* (N. Gait, ed., 1984); *Nucleic Acid Hybridization* (B. Hames & S. Higgins, eds., 1985); *Transcription and Translation* (B. Hames & S. Higgins, eds., 1984); *Animal Cell Culture* (R. Freshney, ed., 1986); Perbal, *A Practical Guide to Molecular Cloning* (1984) and other like references.

As used in this specification and the appended claims, the singular forms “a,” “an” and “the” include plural references unless the content clearly dictates otherwise.

By “about” is meant a quantity, level, value, number, frequency, percentage, dimension, size, amount, weight or length that varies by as much as 20, 15, 10, 9, 8, 7, 6, 5, 4, 3, 2 or 1% to a reference quantity, level, value, number, frequency, percentage, dimension, size, amount, weight or length.

The term “antigen” refers to a molecule or a portion of a molecule capable of being bound by a selective binding agent, such as an antibody, and additionally capable of being used in an animal to produce antibodies capable of binding to an epitope of that antigen. An antigen may have one or more epitopes. As used herein, the term “antigen” includes substances that are capable, under appropriate conditions, of inducing an immune response to the substance and of reacting with the products of the immune response. For example, an antigen can be recognized by antibodies (humoral immune response) or sensitized T-lymphocytes (T helper or cell-mediated immune response), or both. Antigens can be soluble substances, such as toxins and foreign proteins, or particulates, such as bacteria and tissue cells; however, only the portion of the protein or polysaccharide molecule known as the antigenic determinant (epitopes) combines with the antibody or a specific receptor on a

lymphocyte. More broadly, the term “antigen” includes any substance to which an antibody binds, or for which antibodies are desired, regardless of whether the substance is immunogenic. For such antigens, antibodies can be identified by recombinant methods, independently of any immune response.

An “antagonist” refers to an agent (e.g., antibody) that interferes with or otherwise reduces the physiological action of another agent or molecule. In some instances, the antagonist specifically binds to the other agent or molecule. Included are full and partial antagonists.

An “agonist” refers to an agent (e.g., antibody) that increases or enhances the physiological action of another agent or molecule. In some instances, the agonist specifically binds to the other agent or molecule. Included are full and partial agonists.

As used herein, the term “amino acid” is intended to mean both naturally occurring and non-naturally occurring amino acids as well as amino acid analogs and mimetics. Naturally-occurring amino acids include the 20 (L)-amino acids utilized during protein biosynthesis as well as others such as 4-hydroxyproline, hydroxylysine, desmosine, isodesmosine, homocysteine, citrulline and ornithine, for example. Non-naturally occurring amino acids include, for example, (D)-amino acids, norleucine, norvaline, p-fluorophenylalanine, ethionine and the like, which are known to a person skilled in the art. Amino acid analogs include modified forms of naturally and non-naturally occurring amino acids. Such modifications can include, for example, substitution or replacement of chemical groups and moieties on the amino acid or by derivatization of the amino acid. Amino acid mimetics include, for example, organic structures which exhibit functionally similar properties such as charge and charge spacing characteristic of the reference amino acid. For example, an organic structure which mimics arginine (Arg or R) would have a positive charge moiety located in similar molecular space and having the same degree of mobility as the  $\epsilon$ -amino group of the side chain of the naturally occurring Arg amino acid. Mimetics also include constrained structures so as to maintain optimal spacing and charge interactions of the amino acid or of the amino acid functional groups. Those skilled in the art know or can determine what structures constitute functionally equivalent amino acid analogs and amino acid mimetics.

As used herein, the term “antibody” encompasses not only intact polyclonal or monoclonal antibodies, but also fragments thereof (such as dAb, Fab, Fab', F(ab')<sub>2</sub>, Fv), single chain (scFv), synthetic variants thereof, naturally occurring variants, fusion proteins comprising an antibody portion with an antigen binding fragment of the required specificity, humanized antibodies, chimeric antibodies, and any other modified configuration of the immunoglobulin molecule that comprises an antigen binding site or fragment (epitope recognition site) of the required specificity. Certain features and characteristics of antibodies (and antigen binding fragments thereof) are described in greater detail herein.

An antibody or antigen binding fragment can be of essentially any type. As is well known in the art, an antibody is an immunoglobulin molecule capable of specific binding to a target, such as an

immune checkpoint molecule, through at least one epitope recognition site, located in the variable region of the immunoglobulin molecule.

The term “antigen binding fragment” as used herein refers to a polypeptide fragment that contains at least one CDR of an immunoglobulin heavy and/or light chain that binds to the antigen of interest. In this regard, an antigen binding fragment of the herein described antibodies may comprise 1, 2, 3, 4, 5, or all 6 CDRs of a  $V_H$  and  $V_L$  sequence from antibodies that bind to a target molecule.

The binding properties of antibodies and antigen binding fragments thereof can be quantified using methods well known in the art (see Davies et al., Annual Rev. Biochem. 59:439-473, 1990). In some embodiments, an antibody or antigen binding fragment thereof specifically binds to a target molecule, for example, an IL-11R $\alpha$  polypeptide or an epitope or complex thereof, with an equilibrium dissociation constant that is about or ranges from about  $\leq 10^{-7}$  M to about  $10^{-8}$  M. In some embodiments, the equilibrium dissociation constant is about or ranges from about  $\leq 10^{-9}$  M to about  $\leq 10^{-10}$  M. In certain illustrative embodiments, an antibody or antigen binding fragment thereof has an affinity ( $K_D$  or  $EC_{50}$ ) for a target molecule (to which it specifically binds) of about, at least about, or less than about, 0.01, 0.05, 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 40, or 50 nM.

A molecule such as a polypeptide or antibody is said to exhibit “specific binding” or “preferential binding” if it reacts or associates more frequently, more rapidly, with greater duration and/or with greater affinity with a particular cell, substance, or particular epitope than it does with alternative cells or substances, or epitopes. An antibody “specifically binds” or “preferentially binds” to a target molecule or epitope if it binds with greater affinity, avidity, more readily, and/or with greater duration than it binds to other substances or epitopes, for example, by a statistically significant amount. Typically one member of the pair of molecules that exhibit specific binding has an area on its surface, or a cavity, which specifically binds to and is therefore complementary to a particular spatial and/or polar organization of the other member of the pair of molecules. Thus, the members of the pair have the property of binding specifically to each other. For instance, an antibody that specifically or preferentially binds to a specific epitope is an antibody that binds that specific epitope with greater affinity, avidity, more readily, and/or with greater duration than it binds to other epitopes. It is also understood by reading this definition that, for example, an antibody (or moiety or epitope) that specifically or preferentially binds to a first target may or may not specifically or preferentially bind to a second target. The term is also applicable where, for example, an antibody is specific for a particular epitope which is carried by a number of antigens, in which case the specific binding member carrying the antigen binding fragment or domain will be able to bind to the various antigens carrying the epitope; for example, it may be cross reactive to a number of different forms of a target antigen from multiple species that share a common epitope.

Immunological binding generally refers to the non-covalent interactions of the type which occur between an immunoglobulin molecule and an antigen for which the immunoglobulin is specific,

for example by way of illustration and not limitation, as a result of electrostatic, ionic, hydrophilic and/or hydrophobic attractions or repulsion, steric forces, hydrogen bonding, van der Waals forces, and other interactions. The strength, or affinity of immunological binding interactions can be expressed in terms of the dissociation constant ( $K_D$ ) of the interaction, wherein a smaller  $K_D$  represents a greater affinity. Immunological binding properties of selected polypeptides can be quantified using methods well known in the art. One such method entails measuring the rates of antigen binding site/antigen complex formation and dissociation, wherein those rates depend on the concentrations of the complex partners, the affinity of the interaction, and on geometric parameters that equally influence the rate in both directions. Thus, both the “on rate constant” ( $K_{on}$ ) and the “off rate constant” ( $K_{off}$ ) can be determined by calculation of the concentrations and the actual rates of association and dissociation. The ratio of  $K_{off}/K_{on}$  enables cancellation of all parameters not related to affinity, and is thus equal to the dissociation constant  $K_D$ . As used herein, the term “affinity” includes the equilibrium constant for the reversible binding of two agents and is expressed as  $K_D$  or  $EC_{50}$ . Affinity of a binding protein to a ligand such as affinity of an antibody for an epitope can be, for example, from about 100 nanomolar (nM) to about 0.1 nM, from about 100 nM to about 1 picomolar (pM), or from about 100 nM to about 1 femtomolar (fM). As used herein, the term “avidity” refers to the resistance of a complex of two or more agents to dissociation after dilution. In some embodiments, affinity is expressed in the terms of the half maximal effective concentration ( $EC_{50}$ ), which refers to the concentration of an agent, such as an antibody, or an anti-IL-11R $\alpha$  antibody, as disclosed herein, which induces a response halfway between the baseline and maximum after a specified exposure time. The  $EC_{50}$  is commonly used as a measure of an antibody’s potency.

Antibodies can be prepared by any of a variety of techniques known to those of ordinary skill in the art. See, e.g., Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988. Monoclonal antibodies specific for a polypeptide of interest can be prepared, for example, using the technique of Kohler and Milstein, *Eur. J. Immunol.* 6:511-519, 1976, and improvements thereto. Also included are methods that utilize transgenic animals such as mice to express human antibodies. See, e.g., Neuberger et al., *Nature Biotechnology* 14:826, 1996; Lonberg et al., *Handbook of Experimental Pharmacology* 113:49-101, 1994; and Lonberg et al., *Internal Review of Immunology* 13:65-93, 1995. Particular examples include the VELOCIMMUNE® platform by REGENEREX® (see, e.g., U.S. Patent No. 6,596,541).

Antibodies can also be generated or identified by the use of phage display or yeast display libraries (see, e.g., U.S. Patent No. 7,244,592; Chao et al., *Nature Protocols*. 1:755-768, 2006). Non-limiting examples of available libraries include cloned or synthetic libraries, such as the Human Combinatorial Antibody Library (HuCAL), in which the structural diversity of the human antibody repertoire is represented by seven heavy chain and seven light chain variable region genes. The combination of these genes gives rise to 49 frameworks in the master library. By superimposing highly variable genetic cassettes (CDRs = complementarity determining regions) on these

frameworks, the vast human antibody repertoire can be reproduced. Also included are human libraries designed with human-donor-sourced fragments encoding a light-chain variable region, a heavy-chain CDR-3, synthetic DNA encoding diversity in heavy-chain CDR-1, and synthetic DNA encoding diversity in heavy-chain CDR-2. Other libraries suitable for use will be apparent to persons skilled in the art.

In certain embodiments, antibodies and antigen binding fragments thereof as described herein include a heavy chain and a light chain CDR set, respectively interposed between a heavy chain and a light chain framework region (FR) set which provide support to the CDRs and define the spatial relationship of the CDRs relative to each other. As used herein, the term “CDR set” refers to the three hypervariable regions of a heavy or light chain V region. Proceeding from the N-terminus of a heavy or light chain, these regions are denoted as “CDR1,” “CDR2,” and “CDR3” respectively. An antigen binding site, therefore, includes six CDRs, comprising the CDR set from each of a heavy and a light chain V region. A polypeptide comprising a single CDR, (e.g., a CDR1, CDR2 or CDR3) is referred to herein as a “molecular recognition unit.” Crystallographic analysis of a number of antigen-antibody complexes has demonstrated that the amino acid residues of CDRs form extensive contact with bound antigen, wherein the most extensive antigen contact is with the heavy chain CDR3. Thus, the molecular recognition units are primarily responsible for the specificity of an antigen binding site.

As used herein, the term “FR set” refers to the four flanking amino acid sequences which frame the CDRs of a CDR set of a heavy or light chain V region. Some FR residues may contact bound antigen; however, FRs are primarily responsible for folding the V region into the antigen binding site, particularly the FR residues directly adjacent to the CDRs. Within FRs, certain amino residues and certain structural features are very highly conserved. In this regard, most V region sequences contain an internal disulfide loop of around 90 amino acid residues. When the V regions fold into a binding-site, the CDRs are displayed as projecting loop motifs which form an antigen binding surface. It is generally recognized that there are conserved structural regions of FRs which influence the folded shape of the CDR loops into certain “canonical” structures—regardless of the precise CDR amino acid sequence. Further, certain FR residues are known to participate in non-covalent interdomain contacts which stabilize the interaction of the antibody heavy and light chains.

The structures and locations of immunoglobulin variable domains may be determined by reference to Kabat, E. A. et al., Sequences of Proteins of Immunological Interest. 4th Edition. US Department of Health and Human Services. 1987, and updates thereof.

Also include are “monoclonal” antibodies, which refer to a homogeneous antibody population wherein the monoclonal antibody is comprised of amino acids (naturally occurring and non-naturally occurring) that are involved in the selective binding of an epitope. Monoclonal antibodies are highly specific, being directed against a single epitope. The term “monoclonal antibody” encompasses not only intact monoclonal antibodies and full-length monoclonal antibodies, but also fragments thereof (such as Fab, Fab', F(ab')<sub>2</sub>, Fv), single chain (scFv), variants thereof, fusion proteins comprising an

antigen binding portion, humanized monoclonal antibodies, chimeric monoclonal antibodies, and any other modified configuration of the immunoglobulin molecule that comprises an antigen binding fragment (epitope recognition site) of the required specificity and the ability to bind to an epitope. It is not intended to be limited as regards the source of the antibody or the manner in which it is made (e.g., by hybridoma, phage selection, recombinant expression, transgenic animals). The term includes whole immunoglobulins as well as the fragments etc. described above under the definition of “antibody.”

The proteolytic enzyme papain preferentially cleaves IgG molecules to yield several fragments, two of which (the F(ab) fragments) each comprise a covalent heterodimer that includes an intact antigen binding site. The enzyme pepsin is able to cleave IgG molecules to provide several fragments, including the F(ab')<sub>2</sub> fragment which comprises both antigen binding sites. An Fv fragment for use according to certain embodiments can be produced by preferential proteolytic cleavage of an IgM, and on rare occasions of an IgG or IgA immunoglobulin molecule. Fv fragments are, however, more commonly derived using recombinant techniques known in the art. The Fv fragment includes a non-covalent VH::VL heterodimer including an antigen binding site which retains much of the antigen recognition and binding capabilities of the native antibody molecule (Inbar et al., PNAS USA. 69:2659-2662, 1972; Hochman et al., Biochem. 15:2706-2710, 1976; and Ehrlich et al., Biochem. 19:4091-4096, 1980). In some embodiments, Fvs are stabilized by other means, for example, incorporation of at least one disulfide bond (Wörn & Pluckthun, J. Mol. Biol. 305, 989-1010, 2001)

In certain embodiments, single chain Fv (scFv) antibodies are contemplated. For example, Kappa bodies (Ill et al., Prot. Eng. 10:949-57, 1997); minibodies (Martin et al., EMBO J 13:5305-9, 1994); diabodies (Holliger et al., PNAS 90: 6444-8, 1993); or Janusins (Traunecker et al., EMBO J 10: 3655-59, 1991; and Traunecker et al., Int. J. Cancer Suppl. 7:51-52, 1992), may be prepared using standard molecular biology techniques following the teachings of the present application with regard to selecting antibodies having the desired specificity.

A single chain Fv (scFv) polypeptide is a covalently linked VH::VL heterodimer which is expressed from a gene fusion including VH- and VL-encoding genes linked by a peptide-encoding linker. Huston et al. (PNAS USA. 85(16):5879-5883, 1988). A number of methods have been described to discern chemical structures for converting the naturally aggregated—but chemically separated—light and heavy polypeptide chains from an antibody V region into an scFv molecule which will fold into a three dimensional structure substantially similar to the structure of an antigen binding site. See, e.g., U.S. Pat. Nos. 5,091,513 and 5,132,405, to Huston et al.; and U.S. Pat. No. 4,946,778, to Ladner et al.

In certain embodiments, the antibodies or antigen binding fragments described herein are in the form of a “diabody.” Diabodies are multimers of polypeptides, each polypeptide comprising a first domain comprising a binding region of an immunoglobulin light chain and a second domain

comprising a binding region of an immunoglobulin heavy chain, the two domains being linked (e.g., by a peptide linker) but unable to associate with each other to form an antigen binding site: antigen binding sites are formed by the association of the first domain of one polypeptide within the multimer with the second domain of another polypeptide within the multimer (WO94/13804). A dAb fragment of an antibody consists of a VH domain (Ward et al., *Nature* 341:544-546, 1989). Diabodies and other multivalent or multispecific fragments can be constructed, for example, by gene fusion (see WO94/13804; and Holliger et al., *PNAS USA* 90:6444-6448, 1993)).

Minibodies comprising a scFv joined to a CH3 domain are also included (see Hu et al., *Cancer Res.* 56:3055-3061, 1996). See also Ward et al., *Nature* 341:544-546, 1989; Bird et al., *Science* 242:423-426, 1988; Huston et al., *PNAS USA* 85:5879-5883, 1988); PCT/US92/09965; WO94/13804; and Reiter et al., *Nature Biotech.* 14:1239-1245, 1996.

Where bispecific antibodies are to be used, these may be conventional bispecific antibodies, which can be manufactured in a variety of ways (Holliger and Winter, *Current Opinion Biotechnol.* 4:446-449, 1993), e.g., prepared chemically or from hybrid hybridomas, or may be any of the bispecific antibody fragments mentioned above.

Bispecific diabodies, as opposed to bispecific whole antibodies, may also be particularly useful because they can be readily constructed and expressed in *E. coli*. Diabodies (and many other polypeptides such as antibody fragments) of appropriate binding specificities can be readily selected using phage display (WO94/13804) from libraries. If one arm of the diabody is to be kept constant, for instance, with a specificity directed against antigen X, then a library can be made where the other arm is varied and an antibody of appropriate specificity selected. Bispecific whole antibodies may be made by a number of methods (Brinkman & Kontermann, *mAbs* 9:182-212, 2017) including knobs-into-holes engineering (Ridgeway et al., *Protein Eng.* 9:616-621, 1996).

In certain embodiments, the antibodies or antigen binding fragments described herein are in the form of a UniBody®. A UniBody® is an IgG4 antibody with the hinge region removed (see GenMab Utrecht, The Netherlands; see also, e.g., US20090226421). This antibody technology creates a stable, smaller antibody format with an anticipated longer therapeutic window than current small antibody formats. IgG4 antibodies are considered inert and thus do not interact with the immune system. Fully human IgG4 antibodies may be modified by eliminating the hinge region of the antibody to obtain half-molecule fragments having distinct stability properties relative to the corresponding intact IgG4 (GenMab, Utrecht). Halving the IgG4 molecule leaves only one area on the UniBody® that can bind to cognate antigens (e.g., disease targets) and the UniBody® therefore binds univalently to only one site on target cells. For certain cancer cell surface antigens, this univalent binding may not stimulate the cancer cells to grow as may be seen using bivalent antibodies having the same antigen specificity, and hence UniBody® technology may afford treatment options for some types of cancer that may be refractory to treatment with conventional antibodies. The small size of the

UniBody® can be a great benefit when treating some forms of cancer, allowing for better distribution of the molecule over larger solid tumors and potentially increasing efficacy.

In certain embodiments, the antibodies and antigen binding fragments described herein are in the form of a nanobody. Nanobodies are encoded by single genes and are efficiently produced in almost all prokaryotic and eukaryotic hosts, for example, *E. coli* (see U.S. Pat. No. 6,765,087), molds (for example *Aspergillus* or *Trichoderma*) and yeast (for example *Saccharomyces*, *Kluyveromyces*, *Hansenula* or *Pichia* (see U.S. Pat. No. 6,838,254). The production process is scalable and multi-kilogram quantities of nanobodies have been produced. Nanobodies may be formulated as a ready-to-use solution having a long shelf life. The Nanoclone method (see WO 06/079372) is a proprietary method for generating Nanobodies against a desired target, based on automated high-throughput selection of B-cells.

In some embodiments, the antibodies or antigen binding fragments described herein are in the form of an aptamer (see, e.g., Ellington et al., *Nature*. 346, 818-22, 1990; and Tuerk et al., *Science*. 249, 505-10, 1990, incorporated by reference). Examples of aptamers included nucleic acid aptamers (e.g., DNA aptamers, RNA aptamers) and peptide aptamers. Nucleic acid aptamers refer generally to nucleic acid species that have been engineered through repeated rounds of *in vitro* selection or equivalent method, such as SELEX (systematic evolution of ligands by exponential enrichment), to bind to various molecular targets such as small molecules, proteins, nucleic acids, and even cells, tissues and organisms. See, e.g., U.S. Patent Nos. 6,376,190; and 6,387,620, incorporated by reference.

Peptide aptamers typically include a variable peptide loop attached at both ends to a protein scaffold, a double structural constraint that typically increases the binding affinity of the peptide aptamer to levels comparable to that of an antibody's (e.g., in the nanomolar range). In certain embodiments, the variable loop length may be composed of about 10-20 amino acids (including all integers in between), and the scaffold may include any protein that has good solubility and compacity properties. Certain exemplary embodiments utilize the bacterial protein Thioredoxin-A as a scaffold protein, the variable loop being inserted within the reducing active site (-Cys-Gly-Pro-Cys- loop in the wild protein), with the two cysteines lateral chains being able to form a disulfide bridge. Methods for identifying peptide aptamers are described, for example, in U.S. Application No. 2003/0108532, incorporated by reference. Peptide aptamer selection can be performed using different systems known in the art, including the yeast two-hybrid system.

In some embodiments, the antibodies or antigen binding fragments described herein are in the form of an avimer. Avimers refer to multimeric binding proteins or peptides engineered using *in vitro* exon shuffling and phage display. Multiple binding domains are linked, resulting in greater affinity and specificity compared to single epitope immunoglobulin domains. See, e.g., Silverman et al., *Nature Biotechnology*. 23:1556-1561, 2005; U.S. Patent No. 7,166,697; and U.S. Application Nos.

2004/0175756, 2005/0048512, 2005/0053973, 2005/0089932 and 2005/0221384, incorporated by reference.

In some embodiments, the antibodies or antigen binding fragments described herein are in the form of an adnectin. Adnectins refer to a class of targeted biologics derived from human fibronectin, an abundant extracellular protein that naturally binds to other proteins. *See, e.g.*, U.S. Application Nos. 2007/0082365; 2008/0139791; and 2008/0220049, incorporated by reference. Adnectins typically consists of a natural fibronectin backbone, as well as the multiple targeting domains of a specific portion of human fibronectin. The targeting domains can be engineered to enable an adnectin to specifically recognize an IL-11R $\alpha$  polypeptide or an epitope thereof.

In some embodiments, the antibodies or antigen binding fragments described herein are in the form of an anticalin. Anticalins refer to a class of antibody mimetics that are typically synthesized from human lipocalins, a family of binding proteins with a hypervariable loop region supported by a structurally rigid framework. *See, e.g.*, U.S. Application No. 2006/0058510. Anticalins typically have a size of about 20 kDa. Anticalins can be characterized by a barrel structure formed by eight antiparallel  $\beta$ -strands (a stable  $\beta$ -barrel scaffold) that are pairwise connected by four peptide loops and an attached  $\alpha$ -helix. In certain aspects, conformational deviations to achieve specific binding are made in the hypervariable loop region(s). *See, e.g.*, Skerra, FEBS J. 275:2677-83, 2008, incorporated by reference.

In some embodiments, the antibodies or antigen binding fragments described herein are in the form of a designed ankyrin repeat protein (DARPin). DARPins include a class of non-immunoglobulin proteins that can offer advantages over antibodies for target binding in drug discovery and drug development. Among other uses, DARPins are ideally suited for *in vivo* imaging or delivery of toxins or other therapeutic payloads because of their favorable molecular properties, including small size and high stability. The low-cost production in bacteria and the rapid generation of many target-specific DARPins make the DARPin approach useful for drug discovery. Additionally, DARPins can be easily generated in multispecific formats, offering the potential to target an effector DARPin to a specific organ or to target multiple receptors with one molecule composed of several DARPins. *See, e.g.*, Stumpp et al., Curr Opin Drug Discov Devel. 10:153-159, 2007; U.S. Application No. 2009/0082274; and PCT/EP2001/10454, incorporated by reference.

Also included are heavy chain dimers, such as antibodies from camelids and sharks. Camelid and shark antibodies comprise a homodimeric pair of two chains of V-like and C-like domains (neither has a light chain). Since the V<sub>H</sub> of a heavy chain dimer IgG in a camelid does not have to make hydrophobic interactions with a light chain, the region in the heavy chain that normally contacts a light chain is changed to hydrophilic amino acid residues in a camelid. V<sub>H</sub> domains of heavy-chain dimer IgGs are called VHH domains. Shark Ig-NARs comprise a homodimer of one variable domain (termed a V-NAR domain) and five C-like constant domains (C-NAR domains).

In camelids, the diversity of antibody repertoire is determined by the complementary determining regions (CDR) 1, 2, and 3 in the VH or VHH regions. The CDR3 in the camelid VHH region is characterized by its relatively long length averaging 16 amino acids (Muyldermans et al., 1994, *Protein Engineering* 7(9): 1129). This is in contrast to CDR3 regions of antibodies of many other species. For example, the CDR3 of mouse VH has an average of 9 amino acids. Libraries of camelid-derived antibody variable regions, which maintain the in vivo diversity of the variable regions of a camelid, can be made by, for example, the methods disclosed in U.S. Patent Application Ser. No. 20050037421, published Feb. 17, 2005

In certain embodiments, the antibodies or antigen binding fragments thereof are humanized. These embodiments refer to a chimeric molecule, generally prepared using recombinant techniques, having an antigen binding site derived from an immunoglobulin from a non-human species and the remaining immunoglobulin structure of the molecule based upon the structure and/or sequence of a human immunoglobulin. The antigen binding site may comprise either complete variable domains fused onto constant domains or only the CDRs (entire or in part) grafted onto appropriate framework regions in the variable domains. Epitope binding sites may be wild type or modified by one or more amino acid substitutions. This eliminates the constant region as an immunogen in human individuals, but the possibility of an immune response to the foreign variable region remains (LoBuglio et al., *PNAS USA* 86:4220-4224, 1989; Queen et al., *PNAS USA* 86:10029-10033, 1988; Riechmann et al., *Nature* 332:323-327, 1988). Illustrative methods for humanization of antibodies include the methods described in U.S. Patent No. 7,462,697.

Another approach focuses not only on providing human-derived constant regions, but modifying the variable regions as well so as to reshape them as closely as possible to human form. It is known that the variable regions of both heavy and light chains contain three complementarity-determining regions (CDRs) which vary in response to the epitopes in question and determine binding capability, flanked by four framework regions (FRs) which are relatively conserved in a given species and which putatively provide a scaffolding for the CDRs. When nonhuman antibodies are prepared with respect to a particular epitope, the variable regions can be “reshaped” or “humanized” by grafting CDRs derived from nonhuman antibody on the FRs present in the human antibody to be modified. Application of this approach to various antibodies has been reported by Sato et al., *Cancer Res.* 53:851-856, 1993; Riechmann et al., *Nature* 332:323-327, 1988; Verhoeyen et al., *Science* 239:1534-1536, 1988; Kettleborough et al., *Protein Engineering* 4:773-778, 1991; Maeda et al., *Human Antibodies Hybridoma* 2:124-134, 1991; Gorman et al., *PNAS USA* 88:4181-4185, 1991; Tempest et al., *Bio/Technology* 9:266-271, 1991; Co et al., *PNAS USA* 88:2869-2873, 1991; Carter et al., *PNAS USA* 89:4285-4289, 1992; and Co et al., *J Immunol.* 148:1149-1154, 1992. In some embodiments, humanized antibodies preserve all CDR sequences (for example, a humanized mouse antibody which contains all six CDRs from the mouse antibodies). In some embodiments, only some of the CDR sequences are grafted from the nonhuman antibody (Bowers et al., *J. Biol. Chem.*

288:7688-7696, 2013). In certain embodiments, humanized antibodies have one or more CDRs (one, two, three, four, five, six) which are altered with respect to the original antibody, which are also termed one or more CDRs “derived from” one or more CDRs from the original antibody.

In certain embodiments, the antibodies are “chimeric” antibodies. In this regard, a chimeric antibody is comprised of an antigen binding fragment of an antibody operably linked or otherwise fused to a heterologous Fc portion of a different antibody. In certain embodiments, the Fc domain or heterologous Fc domain is of human origin. In certain embodiments, the Fc domain or heterologous Fc domain is of mouse origin. In other embodiments, the heterologous Fc domain may be from a different Ig class from the parent antibody, including IgA (including subclasses IgA1 and IgA2), IgD, IgE, IgG (including subclasses IgG1, IgG2, IgG3, and IgG4), and IgM. In further embodiments, the heterologous Fc domain may be comprised of CH2 and CH3 domains from one or more of the different Ig classes. As noted above with regard to humanized antibodies, the antigen binding fragment of a chimeric antibody may comprise only one or more of the CDRs of the antibodies described herein (e.g., 1, 2, 3, 4, 5, or 6 CDRs of the antibodies described herein), or may comprise an entire variable domain (VL, VH or both).

The term “binding” refers to a direct association between two molecules, due to, for example, covalent, electrostatic, hydrophobic, and ionic and/or hydrogen-bond interactions, including interactions such as salt bridges and water bridges.

By “coding sequence” is meant any nucleic acid sequence that contributes to the code for the polypeptide product of a gene. By contrast, the term “non-coding sequence” refers to any nucleic acid sequence that does not directly contribute to the code for the polypeptide product of a gene.

Throughout this specification, unless the context requires otherwise, the word “comprise”, or variations such as “comprises” or “comprising”, will be understood to imply the inclusion of a stated element or integer or group of elements or integers but not the exclusion of any other element or integer or group of elements or integers.

By “consisting of” is meant including, and limited to, whatever follows the phrase “consisting of.” Thus, the phrase “consisting of” indicates that the listed elements are required or mandatory, and that no other elements may be present. By “consisting essentially of” is meant including any elements listed after the phrase, and limited to other elements that do not interfere with or contribute to the activity or action specified in the disclosure for the listed elements. Thus, the phrase “consisting essentially of” indicates that the listed elements are required or mandatory, but that other elements are optional and may or may not be present depending upon whether or not they materially affect the activity or action of the listed elements.

The term “effector function”, or “ADCC effector function” in the context of antibodies refers to the ability of that antibody to engage with other arms of the immune system, including for example, the activation of the classical complement pathway, or through engagement of Fc receptors. Complement dependent pathways are primarily driven by the interaction of C1q with the C1 complex

with clustered antibody Fc domains. Antibody dependent cellular cytotoxicity (ADCC), is primarily driven by the interaction of Fc receptors (FcRs) on the surface of effector cells (natural killer cells, macrophages, monocytes and eosinophils) which bind to the Fc region of an IgG which itself is bound to a target cell. Fc receptors (FcRs) are key immune regulatory receptors connecting the antibody mediated (humoral) immune response to cellular effector functions. Receptors for all classes of immunoglobulins have been identified, including Fc $\gamma$ R (IgG), Fc $\epsilon$ RI (IgE), Fc $\alpha$ RI (IgA), Fc $\mu$ R (IgM) and Fc $\delta$ R (IgD). There are at least three classes of receptors for human IgG found on leukocytes: CD64 (Fc $\gamma$ RI), CD32 (Fc $\gamma$ RIIa, Fc $\gamma$ RIIb and Fc $\gamma$ RIIc) and CD16 (Fc $\gamma$ RIIIa and Fc $\gamma$ RIIIb). Fc $\gamma$ RI is classed as a high affinity receptor (nanomolar range  $K_D$ ) while Fc $\gamma$ RII and Fc $\gamma$ RIII are low to intermediate affinity (micromolar range  $K_D$ ). Upon Fc binding a signaling pathway is triggered which results in the secretion of various substances, such as lytic enzymes, perforin, granzymes and tumor necrosis factor, which mediate in the destruction of the target cell. The level of ADCC effector function varies for human IgG subtypes. Although this is dependent on the allotype and specific Fc $\gamma$ R, in simple terms ADCC effector function is “high” for human IgG1 and IgG3, and “low” for IgG2 and IgG4.

The term “endotoxin free” or “substantially endotoxin free” relates generally to compositions, solvents, and/or vessels that contain at most trace amounts (*e.g.*, amounts having no clinically adverse physiological effects to a subject) of endotoxin, and preferably undetectable amounts of endotoxin. Endotoxins are toxins associated with certain micro-organisms, such as bacteria, typically gram-negative bacteria, although endotoxins may be found in gram-positive bacteria, such as *Listeria monocytogenes*. The most prevalent endotoxins are lipopolysaccharides (LPS) or lipo-oligosaccharides (LOS) found in the outer membrane of various Gram-negative bacteria, and which represent a central pathogenic feature in the ability of these bacteria to cause disease. Small amounts of endotoxin in humans may produce fever, a lowering of the blood pressure, and activation of inflammation and coagulation, among other adverse physiological effects.

Therefore, in pharmaceutical production, it is often desirable to remove most or all traces of endotoxin from drug products and/or drug containers, because even small amounts may cause adverse effects in humans. A depyrogenation oven may be used for this purpose, as temperatures in excess of 300°C are typically required to break down most endotoxins. For instance, based on primary packaging material such as syringes or vials, the combination of a glass temperature of 250°C and a holding time of 30 minutes is often sufficient to achieve a 3 log reduction in endotoxin levels. Other methods of removing endotoxins are contemplated, including, for example, chromatography and filtration methods, as described herein and known in the art.

Endotoxins can be detected using routine techniques known in the art. For example, the Limulus Amoebocyte Lysate assay, which utilizes blood from the horseshoe crab, is a very sensitive assay for detecting presence of endotoxin. In this test, very low levels of LPS can cause detectable coagulation of the limulus lysate due a powerful enzymatic cascade that amplifies this reaction.

Endotoxins can also be quantitated by enzyme-linked immunosorbent assay (ELISA). To be substantially endotoxin free, endotoxin levels may be less than about 0.001, 0.005, 0.01, 0.02, 0.03, 0.04, 0.05, 0.06, 0.08, 0.09, 0.1, 0.5, 1.0, 1.5, 2, 2.5, 3, 4, 5, 6, 7, 8, 9, or 10 EU/mg of active compound. Typically, 1 ng lipopolysaccharide (LPS) corresponds to about 1-10 EU.

The term “epitope” includes any determinant, preferably a polypeptide determinant, capable of specific binding to an immunoglobulin or T-cell receptor. An epitope includes a region of an antigen that is bound by an antibody. In certain embodiments, epitope determinants include chemically active surface groupings of molecules such as amino acids, sugar side chains, phosphoryl or sulfonyl, and may in certain embodiments have specific three-dimensional structural characteristics, and/or specific charge characteristics. Epitopes can be contiguous or non-contiguous in relation to the primary structure of the antigen, for example, an IL-11R $\alpha$  polypeptide. In particular embodiments, an epitope comprises, consists, or consists essentially of about, at least about, or no more than about 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 contiguous amino acids (i.e., a linear epitope) or non-contiguous amino acids (i.e., conformational epitope) of a reference sequence (see, e.g., **Table B1**) or target molecule described herein.

An “epitope” includes that portion of an antigen or other macromolecule capable of forming a binding interaction that interacts with the variable region binding pocket of a binding protein. Such binding interaction can be manifested as an intermolecular contact with one or more amino acid residues of a CDR. Antigen binding can involve a CDR3 or a CDR3 pair. An epitope can be a linear peptide sequence (i.e., “continuous”) or can be composed of noncontiguous amino acid sequences (i.e., “conformational” or “discontinuous”). A binding protein can recognize one or more amino acid sequences; therefore an epitope can define more than one distinct amino acid sequence. Epitopes recognized by binding protein can be determined by peptide mapping and sequence analysis techniques well known to one of skill in the art. A “cryptic epitope” or a “cryptic binding site” is an epitope or binding site of a protein sequence that is not exposed or substantially protected from recognition within an unmodified polypeptide, but is capable of being recognized by a binding protein of a denatured or proteolyzed polypeptide. Amino acid sequences that are not exposed, or are only partially exposed, in the unmodified polypeptide structure are potential cryptic epitopes. If an epitope is not exposed, or only partially exposed, then it is likely that it is buried within the interior of the polypeptide. Candidate cryptic epitopes can be identified, for example, by examining the three-dimensional structure of an unmodified polypeptide.

The term “half maximal effective concentration” or “EC<sub>50</sub>” refers to the concentration of an agent (e.g., antibody) as described herein at which it induces a response halfway between the baseline and maximum after some specified exposure time; the EC<sub>50</sub> of a graded dose response curve therefore represents the concentration of a compound at which 50% of its maximal effect is observed. EC<sub>50</sub> also represents the plasma concentration required for obtaining 50% of a maximum effect in vivo. Similarly, the “EC<sub>90</sub>” refers to the concentration of an agent or composition at which 90% of its

maximal effect is observed. The “EC<sub>90</sub>” can be calculated from the “EC<sub>50</sub>” and the Hill slope, or it can be determined from the data directly, using routine knowledge in the art. In some embodiments, the EC<sub>50</sub> of an agent (e.g., antibody) is less than about 0.01, 0.05, 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 40, 50, 60, 70, 80, 90, 100, 200 or 500 nM. In some embodiments, an agent will have an EC<sub>50</sub> value of about 1nM or less.

“Immune response” means any immunological response originating from immune system, including responses from the cellular and humeral, innate and adaptive immune systems. Exemplary cellular immune cells include for example, lymphocytes, macrophages, T cells, B cells, NK cells, neutrophils, eosinophils, dendritic cells, mast cells, monocytes, and all subsets thereof. Cellular responses include for example, effector function, cytokine release, phagocytosis, efferocytosis, translocation, trafficking, proliferation, differentiation, activation, repression, cell-cell interactions, apoptosis, etc. Humeral responses include for example IgG, IgM, IgA, IgE, responses and their corresponding effector functions.

The “half-life” of an agent such as an antibody can refer to the time it takes for the agent to lose half of its pharmacologic, physiologic, or other activity, relative to such activity at the time of administration into the serum or tissue of an organism, or relative to any other defined time-point. “Half-life” can also refer to the time it takes for the amount or concentration of an agent to be reduced by half of a starting amount administered into the serum or tissue of an organism, relative to such amount or concentration at the time of administration into the serum or tissue of an organism, or relative to any other defined time-point. The half-life can be measured in serum and/or any one or more selected tissues.

The terms “modulating” and “altering” include “increasing,” “enhancing” or “stimulating,” as well as “decreasing,” “reducing,” or “inhibiting,” typically in a statistically significant or a physiologically significant amount or degree relative to a control. An “increased,” “stimulated” or “enhanced” amount is typically a “statistically significant” amount, and may include an increase that is 1.1, 1.2, 1.5, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 30, 40, 50, 60, 70, 80, 90, 100 or more times (e.g., 500, 1000 times) (including all integers and ranges in between e.g., 1.5, 1.6, 1.7, 1.8, etc.) the amount produced by no composition (e.g., the absence of agent) or a control composition. A “decreased” or “reduced” or “inhibited” amount is typically a “statistically significant” amount, and may include a 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 100% decrease (including all integers and ranges in between) in the amount produced by no composition (e.g., the absence of an agent) or a control composition. Examples of comparisons and “statistically significant” amounts are described herein.

The terms “polypeptide,” “protein” and “peptide” are used interchangeably and mean a polymer of amino acids not limited to any particular length. The term “enzyme” includes polypeptide or protein catalysts. The terms include modifications such as myristoylation, sulfation, glycosylation,

phosphorylation and addition or deletion of signal sequences. The terms “polypeptide” or “protein” means one or more chains of amino acids, wherein each chain comprises amino acids covalently linked by peptide bonds, and wherein said polypeptide or protein can comprise a plurality of chains non-covalently and/or covalently linked together by peptide bonds, having the sequence of native proteins, that is, proteins produced by naturally-occurring and specifically non-recombinant cells, or genetically-engineered or recombinant cells, and comprise molecules having the amino acid sequence of the native protein, or molecules having deletions from, additions to, and/or substitutions of one or more amino acids of the native sequence. In certain embodiments, the polypeptide is a “recombinant” polypeptide, produced by recombinant cell that comprises one or more recombinant DNA molecules, which are typically made of heterologous polynucleotide sequences or combinations of polynucleotide sequences that would not otherwise be found in the cell.

The term “polynucleotide” and “nucleic acid” includes mRNA, RNA, cRNA, cDNA, and DNA. The term typically refers to polymeric form of nucleotides of at least 10 bases in length, either ribonucleotides or deoxynucleotides or a modified form of either type of nucleotide. The term includes single and double stranded forms of DNA. The terms “isolated DNA” and “isolated polynucleotide” and “isolated nucleic acid” refer to a molecule that has been isolated free of total genomic DNA of a particular species. Therefore, an isolated DNA segment encoding a polypeptide refers to a DNA segment that contains one or more coding sequences yet is substantially isolated away from, or purified free from, total genomic DNA of the species from which the DNA segment is obtained. Also included are non-coding polynucleotides (*e.g.*, primers, probes, oligonucleotides), which do not encode a polypeptide. Also included are recombinant vectors, including, for example, expression vectors, viral vectors, plasmids, cosmids, phagemids, phage, viruses, and the like.

Additional coding or non-coding sequences may, but need not, be present within a polynucleotide described herein, and a polynucleotide may, but need not, be linked to other molecules and/or support materials. Hence, a polynucleotide or expressible polynucleotides, regardless of the length of the coding sequence itself, may be combined with other sequences, for example, expression control sequences.

“Expression control sequences” include regulatory sequences of nucleic acids, or the corresponding amino acids, such as promoters, leaders, enhancers, introns, recognition motifs for RNA, or DNA binding proteins, polyadenylation signals, terminators, internal ribosome entry sites (IRES), secretion signals, subcellular localization signals, and the like, which have the ability to affect the transcription or translation, or subcellular, or cellular location of a coding sequence in a host cell. Exemplary expression control sequences are described in Goeddel; *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, Calif. (1990).

A “promoter” is a DNA regulatory region capable of binding RNA polymerase in a cell and initiating transcription of a downstream (3' direction) coding sequence. As used herein, the promoter sequence is bounded at its 3' terminus by the transcription initiation site and extends upstream (5'

direction) to include the minimum number of bases or elements necessary to initiate transcription at levels detectable above background. A transcription initiation site (conveniently defined by mapping with nuclease S1) can be found within a promoter sequence, as well as protein binding domains (consensus sequences) responsible for the binding of RNA polymerase. Eukaryotic promoters can often, but not always, contain “TATA” boxes and “CAT” boxes. Prokaryotic promoters contain Shine- Dalgarno sequences in addition to the -10 and -35 consensus sequences.

A large number of promoters, including constitutive, inducible and repressible promoters, from a variety of different sources are well known in the art. Representative sources include for example, viral, mammalian, insect, plant, yeast, and bacterial cell types), and suitable promoters from these sources are readily available, or can be made synthetically, based on sequences publicly available on line or, for example, from depositories such as the ATCC as well as other commercial or individual sources. Promoters can be unidirectional (i.e., initiate transcription in one direction) or bi-directional (i.e., initiate transcription in either a 3' or 5' direction). Non-limiting examples of promoters include, for example, the T7 bacterial expression system, pBAD (araA) bacterial expression system, the cytomegalovirus (CMV) promoter, the SV40 promoter, the RSV promoter. Inducible promoters include the Tet system, (US Patents 5,464,758 and 5,814,618), the Ecdysone inducible system (No et al., Proc. Natl. Acad. Sci. (1996) 93 (8): 3346-3351; the T-REx<sup>TM</sup> system (Invitrogen Carlsbad, CA), LacSwitch<sup>®</sup> (Stratagene, (San Diego, CA) and the Cre-ERT tamoxifen inducible recombinase system (Indra et al. Nuc. Acid. Res. (1999) 27 (22): 4324-4327; Nuc. Acid. Res. (2000) 28 (23): e99; US Patent No. 7,112,715; and Kramer & Fussenegger, Methods Mol. Biol. (2005) 308: 123-144) or any promoter known in the art suitable for expression in the desired cells.

An “expressible polynucleotide” includes a cDNA, RNA, mRNA or other polynucleotide that comprises at least one coding sequence and optionally at least one expression control sequence, for example, a transcriptional and/or translational regulatory element, and which can express an encoded polypeptide upon introduction into a cell, for example, a cell in a subject.

Various viral vectors that can be utilized to deliver an expressible polynucleotide include adenoviral vectors, herpes virus vectors, vaccinia virus vectors, adeno-associated virus (AAV) vectors, and retroviral vectors. In some instances, the retroviral vector is a derivative of a murine or avian retrovirus, or is a lentiviral vector. Examples of retroviral vectors in which a single foreign gene can be inserted include, but are not limited to: Moloney murine leukemia virus (MoMuLV), Harvey murine sarcoma virus (HaMuSV), murine mammary tumor virus (MuMTV), SIV, BIV, HIV and Rous Sarcoma Virus (RSV). A number of additional retroviral vectors can incorporate multiple genes. All of these vectors can transfer or incorporate a gene for a selectable marker so that transduced cells can be identified and generated. By inserting a polypeptide sequence of interest into the viral vector, along with another gene that encodes the ligand for a receptor on a specific target cell, for example, the vector may be made target specific. Retroviral vectors can be made target specific by inserting, for example, a polynucleotide encoding a protein. Illustrative targeting may be accomplished by using an

antibody to target the retroviral vector. Those of skill in the art will know of, or can readily ascertain without undue experimentation, specific polynucleotide sequences which can be inserted into the retroviral genome to allow target specific delivery of the retroviral vector.

In particular embodiments, the expressible polynucleotide is a modified RNA or modified mRNA polynucleotide, for example, a non-naturally occurring RNA analog. In certain embodiments, the modified RNA or mRNA polypeptide comprises one or more modified or non-natural bases, for example, a nucleotide base other than adenine (A), guanine (G), cytosine (C), thymine (T), and/or uracil (U). In some embodiments, the modified mRNA comprises one or more modified or non-natural internucleotide linkages. Expressible RNA polynucleotides for delivering an encoded therapeutic polypeptide are described, for example, in Kormann et al., *Nat Biotechnol.* 29:154-7, 2011; and U.S. Application Nos. 2015/0111248; 2014/0243399; 2014/0147454; and 2013/0245104, which are incorporated by reference in their entireties.

The term “isolated” polypeptide or protein referred to herein means that a subject protein (1) is free of at least some other proteins with which it would typically be found in nature, (2) is essentially free of other proteins from the same source, *e.g.*, from the same species, (3) is expressed by a cell from a different species, (4) has been separated from at least about 50 percent of polynucleotides, lipids, carbohydrates, or other materials with which it is associated in nature, (5) is not associated (by covalent or non-covalent interaction) with portions of a protein with which the “isolated protein” is associated in nature, (6) is operably associated (by covalent or non-covalent interaction) with a polypeptide with which it is not associated in nature, or (7) does not occur in nature. Such an isolated protein can be encoded by genomic DNA, cDNA, mRNA or other RNA, of may be of synthetic origin, or any combination thereof. In certain embodiments, the isolated protein is substantially free from proteins or polypeptides or other contaminants that are found in its natural environment that would interfere with its use (therapeutic, diagnostic, prophylactic, research or otherwise).

In certain embodiments, the “purity” of any given agent (*e.g.*, antibody) in a composition may be defined. For instance, certain compositions may comprise an agent that is at least 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% pure on a protein basis or a weight-weight basis, including all decimals and ranges in between, as measured, for example and by no means limiting, by high performance liquid chromatography (HPLC), a well-known form of column chromatography used frequently in biochemistry and analytical chemistry to separate, identify, and quantify compounds.

The term “reference sequence” refers generally to a nucleic acid coding sequence, or amino acid sequence, to which another sequence is being compared. All polypeptide and polynucleotide sequences described herein are included as reference sequences, including those described by name and those described in the Tables and the Sequence Listing.

Certain embodiments include biologically active “variants” and “fragments” of the polypeptides (e.g., antibodies) described herein, and the polynucleotides that encode the same. “Variants” contain one or more substitutions, additions, deletions, and/or insertions relative to a reference polypeptide or polynucleotide (see, e.g., the Tables and the Sequence Listing). A variant polypeptide or polynucleotide comprises an amino acid or nucleotide sequence with at least about 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more sequence identity or similarity or homology to a reference sequence, as described herein, and substantially retains the activity of that reference sequence. Also included are sequences that consist of or differ from a reference sequences by the addition, deletion, insertion, or substitution of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150 or more amino acids or nucleotides and which substantially retain the activity of that reference sequence. In certain embodiments, the additions or deletions include C-terminal and/or N-terminal additions and/or deletions.

The terms “sequence identity” or, for example, comprising a “sequence at least 50% identical to,” as used herein, refer to the extent that sequences are identical on a nucleotide-by-nucleotide basis or an amino acid-by-amino acid basis over a window of comparison. Thus, a “percentage of sequence identity” may be calculated by comparing two optimally aligned sequences over the window of comparison, determining the number of positions at which the identical nucleic acid base (e.g., A, T, C, G, I) or the identical amino acid residue (e.g., Ala, Pro, Ser, Thr, Gly, Val, Leu, Ile, Phe, Tyr, Trp, Lys, Arg, His, Asp, Glu, Asn, Gln, Cys and Met) occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison (i.e., the window size), and multiplying the result by 100 to yield the percentage of sequence identity. Optimal alignment of sequences for aligning a comparison window may be conducted by computerized implementations of algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package Release 7.0, Genetics Computer Group, 575 Science Drive Madison, Wis., USA) or by inspection and the best alignment (i.e., resulting in the highest percentage homology over the comparison window) generated by any of the various methods selected. Reference also may be made to the BLAST family of programs as for example disclosed by Altschul et al., Nucl. Acids Res. 25:3389, 1997.

The term “solubility” refers to the property of an agent (e.g., antibody) provided herein to dissolve in a liquid solvent and form a homogeneous solution. Solubility is typically expressed as a concentration, either by mass of solute per unit volume of solvent (g of solute per kg of solvent, g per dL (100 mL), mg/ml, etc.), molarity, molality, mole fraction or other similar descriptions of concentration. The maximum equilibrium amount of solute that can dissolve per amount of solvent is the solubility of that solute in that solvent under the specified conditions, including temperature, pressure, pH, and the nature of the solvent. In certain embodiments, solubility is measured at physiological pH, or other pH, for example, at pH 5.0, pH 6.0, pH 7.0, pH 7.4, pH 7.6, pH 7.8, or pH

8.0 (e.g., about pH 5-8). In certain embodiments, solubility is measured in water or a physiological buffer such as PBS or NaCl (with or without NaPO<sub>4</sub>). In specific embodiments, solubility is measured at relatively lower pH (e.g., pH 6.0) and relatively higher salt (e.g., 500mM NaCl and 10mM NaPO<sub>4</sub>). In certain embodiments, solubility is measured in a biological fluid (solvent) such as blood or serum. In certain embodiments, the temperature can be about room temperature (e.g., about 20, 21, 22, 23, 24, 25°C) or about body temperature (37°C). In certain embodiments, an agent has a solubility of at least about 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 40, 50, 60, 70, 80, 90 or 100 mg/ml at room temperature or at 37°C.

A “subject” or a “subject in need thereof” or a “patient” or a “patient in need thereof” includes a mammalian subject such as a human subject.

“Substantially” or “essentially” means nearly totally or completely, for instance, 95%, 96%, 97%, 98%, 99% or greater of some given quantity.

By “statistically significant,” it is meant that the result was unlikely to have occurred by chance. Statistical significance can be determined by any method known in the art. Commonly used measures of significance include the p-value, which is the frequency or probability with which the observed event would occur, if the null hypothesis were true. If the obtained p-value is smaller than the significance level, then the null hypothesis is rejected. In simple cases, the significance level is defined at a p-value of 0.05 or less.

“Therapeutic response” refers to improvement of symptoms (whether or not sustained) based on administration of one or more therapeutic agents.

As used herein, the terms “therapeutically effective amount”, “therapeutic dose,” “prophylactically effective amount,” or “diagnostically effective amount” is the amount of an agent (e.g., anti-IL-11R $\alpha$  antibody, immunotherapy agent) needed to elicit the desired biological response following administration.

As used herein, “treatment” of a subject (e.g., a mammal, such as a human) or a cell is any type of intervention used in an attempt to alter the natural course of the individual or cell. Treatment includes, but is not limited to, administration of a pharmaceutical composition, and may be performed either prophylactically or subsequent to the initiation of a pathologic event or contact with an etiologic agent. Also included are “prophylactic” treatments, which can be directed to reducing the rate of progression of the disease or condition being treated, delaying the onset of that disease or condition, or reducing the severity of its onset. “Treatment” or “prophylaxis” does not necessarily indicate complete eradication, cure, or prevention of the disease or condition, or associated symptoms thereof.

The term “wild-type” refers to a gene or gene product (e.g., a polypeptide) that is most frequently observed in a population and is thus arbitrarily designed the “normal” or “wild-type” form of the gene.

Each embodiment in this specification is to be applied *mutatis mutandis* to every other embodiment unless expressly stated otherwise.

### Anti-IL-11R $\alpha$ Antibodies

Certain embodiments include antibodies, and antigen binding fragments thereof, which bind to interleukin-11 receptor subunit  $\alpha$  (IL-11R $\alpha$ ). In some embodiments, an antibody or antigen binding fragment thereof modulates (e.g., interferes with, antagonizes, inhibits) binding of IL-11R $\alpha$  to its ligand, interleukin 11 (IL-11). In certain embodiments, an antibody or antigen binding fragment thereof is characterized by or comprises a heavy chain variable region (V<sub>H</sub>) that comprises complementary determining region V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences, and a light chain variable region (V<sub>L</sub>) that comprises complementary determining region V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences. Exemplary V<sub>H</sub>, V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, V<sub>H</sub>CDR3, V<sub>L</sub>, V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences are provided in **Table A1** and **Table A2** below.

| <b>Table A1: Exemplary CDR Sequences</b> |                   |                   |
|------------------------------------------|-------------------|-------------------|
| <b>Description</b>                       | <b>Sequence</b>   | <b>SEQ ID NO:</b> |
| <b>mAb1</b>                              |                   |                   |
| V <sub>H</sub> CDR1                      | WYSMT             | 1                 |
| V <sub>H</sub> CDR2                      | SIVPSGGHTQYADSVKG | 2                 |
| V <sub>H</sub> CDR3                      | GPDWGSFDL         | 3                 |
| V <sub>L</sub> CDR1                      | QASQDINNLYN       | 4                 |
| V <sub>L</sub> CDR2                      | DASNLQT           | 5                 |
| V <sub>L</sub> CDR3                      | QQHESQSPT         | 6                 |
| <b>mAb2</b>                              |                   |                   |
| V <sub>H</sub> CDR1                      | WYSMT             | 7                 |
| V <sub>H</sub> CDR2                      | SIVPSGGHTQYADSVKG | 8                 |
| V <sub>H</sub> CDR3                      | GPDWGSFDL         | 9                 |
| V <sub>L</sub> CDR1                      | QASQDINNLYN       | 10                |
| V <sub>L</sub> CDR2                      | DASNLQT           | 11                |
| V <sub>L</sub> CDR3                      | QQHEFQSPT         | 12                |
| <b>mAb3</b>                              |                   |                   |
| V <sub>H</sub> CDR1                      | WYSMT             | 13                |
| V <sub>H</sub> CDR2                      | SIVPYGDLTQYADSVKG | 14                |
| V <sub>H</sub> CDR3                      | GPDWGSFDL         | 15                |
| V <sub>L</sub> CDR1                      | QASQDINNLYN       | 16                |
| V <sub>L</sub> CDR2                      | DASNLQT           | 17                |
| V <sub>L</sub> CDR3                      | QQHESQSPT         | 18                |
| <b>mAb4</b>                              |                   |                   |
| V <sub>H</sub> CDR1                      | WYSMT             | 19                |
| V <sub>H</sub> CDR2                      | SIVPYGDLTQYADSVKG | 20                |
| V <sub>H</sub> CDR3                      | GPDWGSFDL         | 21                |
| V <sub>L</sub> CDR1                      | QASQDINNLYN       | 22                |
| V <sub>L</sub> CDR2                      | DASNLQT           | 23                |
| V <sub>L</sub> CDR3                      | QQHEFQSPT         | 24                |
| <b>mAb5</b>                              |                   |                   |
| V <sub>H</sub> CDR1                      | WYSMT             | 25                |
| V <sub>H</sub> CDR2                      | SIVPYGDLTQYADSVKG | 26                |
| V <sub>H</sub> CDR3                      | GPGWGSFDL         | 27                |
| V <sub>L</sub> CDR1                      | QASQDINNLYN       | 28                |
| V <sub>L</sub> CDR2                      | DASNLQT           | 29                |
| V <sub>L</sub> CDR3                      | QQHESQSPT         | 30                |
| <b>mAb6</b>                              |                   |                   |
| V <sub>H</sub> CDR1                      | WYSMT             | 31                |
| V <sub>H</sub> CDR2                      | SIVPYGDLTQYADSVKG | 32                |

|                     |                   |    |
|---------------------|-------------------|----|
| V <sub>H</sub> CDR3 | GPGWGSFDL         | 33 |
| V <sub>L</sub> CDR1 | QASQDINNLYN       | 34 |
| V <sub>L</sub> CDR2 | DASNLQT           | 35 |
| V <sub>L</sub> CDR3 | QQHEFQSPT         | 36 |
| <b>mAb7</b>         |                   |    |
| V <sub>H</sub> CDR1 | NYAMS             | 37 |
| V <sub>H</sub> CDR2 | SIVPYGDLTQYADSVKG | 38 |
| V <sub>H</sub> CDR3 | GPGWGSFDL         | 39 |
| V <sub>L</sub> CDR1 | QASQDINNLYN       | 40 |
| V <sub>L</sub> CDR2 | DASNLQT           | 41 |
| V <sub>L</sub> CDR3 | QQHESQSPT         | 42 |
| <b>mAb8</b>         |                   |    |
| V <sub>H</sub> CDR1 | SYAMS             | 43 |
| V <sub>H</sub> CDR2 | SIVPYGDLTQYADSVKG | 44 |
| V <sub>H</sub> CDR3 | GPGWGSFDL         | 45 |
| V <sub>L</sub> CDR1 | QASQDINNLYN       | 46 |
| V <sub>L</sub> CDR2 | DASNLQT           | 47 |
| V <sub>L</sub> CDR3 | QQHESQSPT         | 48 |
| <b>mAb9</b>         |                   |    |
| V <sub>H</sub> CDR1 | WYSMT             | 49 |
| V <sub>H</sub> CDR2 | GIVPYGDLTQYADSVKG | 50 |
| V <sub>H</sub> CDR3 | GPGWGSFDL         | 51 |
| V <sub>L</sub> CDR1 | QASQDINNLYN       | 52 |
| V <sub>L</sub> CDR2 | DASNLQT           | 53 |
| V <sub>L</sub> CDR3 | QQHESQSPT         | 54 |
| <b>mAb10</b>        |                   |    |
| V <sub>H</sub> CDR1 | WYSMT             | 55 |
| V <sub>H</sub> CDR2 | SIVAYGDLTQYADSVKG | 56 |
| V <sub>H</sub> CDR3 | GPGWGSFDL         | 57 |
| V <sub>L</sub> CDR1 | QASQDINNLYN       | 58 |
| V <sub>L</sub> CDR2 | DASNLQT           | 59 |
| V <sub>L</sub> CDR3 | QQHESQSPT         | 60 |
| <b>mAb11</b>        |                   |    |
| V <sub>H</sub> CDR1 | WYSMT             | 61 |
| V <sub>H</sub> CDR2 | SIVDYGDLTQYADSVKG | 62 |
| V <sub>H</sub> CDR3 | GPGWGSFDL         | 63 |
| V <sub>L</sub> CDR1 | QASQDINNLYN       | 64 |
| V <sub>L</sub> CDR2 | DASNLQT           | 65 |
| V <sub>L</sub> CDR3 | QQHESQSPT         | 66 |
| <b>mAb12</b>        |                   |    |
| V <sub>H</sub> CDR1 | WYSMT             | 67 |
| V <sub>H</sub> CDR2 | SIVPYGDLTQYADSVKG | 68 |
| V <sub>H</sub> CDR3 | GPGWYSFDL         | 69 |
| V <sub>L</sub> CDR1 | QASQDINNLYN       | 70 |
| V <sub>L</sub> CDR2 | DASNLQT           | 71 |
| V <sub>L</sub> CDR3 | QQHESQSPT         | 72 |
| <b>mAb13</b>        |                   |    |
| V <sub>H</sub> CDR1 | WYSMT             | 73 |
| V <sub>H</sub> CDR2 | SIVPYGDLTQYADSVKG | 74 |
| V <sub>H</sub> CDR3 | PEDWGRFDL         | 75 |
| V <sub>L</sub> CDR1 | QASQDINNLYN       | 76 |
| V <sub>L</sub> CDR2 | DASNLQT           | 77 |
| V <sub>L</sub> CDR3 | QQHESQSPT         | 78 |
| <b>mAb14</b>        |                   |    |
| V <sub>H</sub> CDR1 | WYSMT             | 79 |
| V <sub>H</sub> CDR2 | SIVPYGDLTQYAESVKG | 80 |
| V <sub>H</sub> CDR3 | GPGWGSFDL         | 81 |
| V <sub>L</sub> CDR1 | QASQDINNLYN       | 82 |

|                     |                   |     |
|---------------------|-------------------|-----|
| V <sub>L</sub> CDR2 | DASNLQT           | 83  |
| V <sub>L</sub> CDR3 | QQHESQSPT         | 84  |
| <b>mAb15</b>        |                   |     |
| V <sub>H</sub> CDR1 | WYSMT             | 85  |
| V <sub>H</sub> CDR2 | SIVPYGDLTQYADSVKG | 86  |
| V <sub>H</sub> CDR3 | GPGWGSFDL         | 87  |
| V <sub>L</sub> CDR1 | QASQDINNLYN       | 88  |
| V <sub>L</sub> CDR2 | DASNLQT           | 89  |
| V <sub>L</sub> CDR3 | QQHETQTPT         | 90  |
| <b>mAb16</b>        |                   |     |
| V <sub>H</sub> CDR1 | AYSMT             | 91  |
| V <sub>H</sub> CDR2 | SIVPYGDLTQYADSVKG | 92  |
| V <sub>H</sub> CDR3 | GPGWGSFDL         | 93  |
| V <sub>L</sub> CDR1 | QASQDINNLYN       | 94  |
| V <sub>L</sub> CDR2 | DASNLQT           | 95  |
| V <sub>L</sub> CDR3 | QQHESQSPT         | 96  |
| <b>mAb17</b>        |                   |     |
| V <sub>H</sub> CDR1 | DYSMT             | 97  |
| V <sub>H</sub> CDR2 | SIVPYGDLTQYADSVKG | 98  |
| V <sub>H</sub> CDR3 | GPGWGSFDL         | 99  |
| V <sub>L</sub> CDR1 | QASQDINNLYN       | 100 |
| V <sub>L</sub> CDR2 | DASNLQT           | 101 |
| V <sub>L</sub> CDR3 | QQHESQSPT         | 102 |
| <b>mAb18</b>        |                   |     |
| V <sub>H</sub> CDR1 | EYSMT             | 103 |
| V <sub>H</sub> CDR2 | SIVPYGDLTQYADSVKG | 104 |
| V <sub>H</sub> CDR3 | GPGWGSFDL         | 105 |
| V <sub>L</sub> CDR1 | QASQDINNLYN       | 106 |
| V <sub>L</sub> CDR2 | DASNLQT           | 107 |
| V <sub>L</sub> CDR3 | QQHESQSPT         | 108 |
| <b>mAb19</b>        |                   |     |
| V <sub>H</sub> CDR1 | FYSMT             | 109 |
| V <sub>H</sub> CDR2 | SIVPYGDLTQYADSVKG | 110 |
| V <sub>H</sub> CDR3 | GPGWGSFDL         | 111 |
| V <sub>L</sub> CDR1 | QASQDINNLYN       | 112 |
| V <sub>L</sub> CDR2 | DASNLQT           | 113 |
| V <sub>L</sub> CDR3 | QQHESQSPT         | 114 |
| <b>mAb20</b>        |                   |     |
| V <sub>H</sub> CDR1 | GYSMT             | 115 |
| V <sub>H</sub> CDR2 | SIVPYGDLTQYADSVKG | 116 |
| V <sub>H</sub> CDR3 | GPGWGSFDL         | 117 |
| V <sub>L</sub> CDR1 | QASQDINNLYN       | 118 |
| V <sub>L</sub> CDR2 | DASNLQT           | 119 |
| V <sub>L</sub> CDR3 | QQHESQSPT         | 120 |
| <b>mAb21</b>        |                   |     |
| V <sub>H</sub> CDR1 | HYSMT             | 121 |
| V <sub>H</sub> CDR2 | SIVPYGDLTQYADSVKG | 122 |
| V <sub>H</sub> CDR3 | GPGWGSFDL         | 123 |
| V <sub>L</sub> CDR1 | QASQDINNLYN       | 124 |
| V <sub>L</sub> CDR2 | DASNLQT           | 125 |
| V <sub>L</sub> CDR3 | QQHESQSPT         | 126 |
| <b>mAb22</b>        |                   |     |
| V <sub>H</sub> CDR1 | IYSMT             | 127 |
| V <sub>H</sub> CDR2 | SIVPYGDLTQYADSVKG | 128 |
| V <sub>H</sub> CDR3 | GPGWGSFDL         | 129 |
| V <sub>L</sub> CDR1 | QASQDINNLYN       | 130 |
| V <sub>L</sub> CDR2 | DASNLQT           | 131 |
| V <sub>L</sub> CDR3 | QQHESQSPT         | 132 |

| mAb23               |                   |     |
|---------------------|-------------------|-----|
| V <sub>H</sub> CDR1 | KYSMT             | 133 |
| V <sub>H</sub> CDR2 | SIVPYGDLTQYADSVKG | 134 |
| V <sub>H</sub> CDR3 | GPGWGSFDL         | 135 |
| V <sub>L</sub> CDR1 | QASQDINNLYN       | 136 |
| V <sub>L</sub> CDR2 | DASNLQT           | 137 |
| V <sub>L</sub> CDR3 | QQHESQSPT         | 138 |
| mAb24               |                   |     |
| V <sub>H</sub> CDR1 | LYSMT             | 139 |
| V <sub>H</sub> CDR2 | SIVPYGDLTQYADSVKG | 140 |
| V <sub>H</sub> CDR3 | GPGWGSFDL         | 141 |
| V <sub>L</sub> CDR1 | QASQDINNLYN       | 142 |
| V <sub>L</sub> CDR2 | DASNLQT           | 143 |
| V <sub>L</sub> CDR3 | QQHESQSPT         | 144 |
| mAb25               |                   |     |
| V <sub>H</sub> CDR1 | MYSMT             | 145 |
| V <sub>H</sub> CDR2 | SIVPYGDLTQYADSVKG | 146 |
| V <sub>H</sub> CDR3 | GPGWGSFDL         | 147 |
| V <sub>L</sub> CDR1 | QASQDINNLYN       | 148 |
| V <sub>L</sub> CDR2 | DASNLQT           | 149 |
| V <sub>L</sub> CDR3 | QQHESQSPT         | 150 |
| mAb26               |                   |     |
| V <sub>H</sub> CDR1 | NYSMT             | 151 |
| V <sub>H</sub> CDR2 | SIVPYGDLTQYADSVKG | 152 |
| V <sub>H</sub> CDR3 | GPGWGSFDL         | 153 |
| V <sub>L</sub> CDR1 | QASQDINNLYN       | 154 |
| V <sub>L</sub> CDR2 | DASNLQT           | 155 |
| V <sub>L</sub> CDR3 | QQHESQSPT         | 156 |
| mAb27               |                   |     |
| V <sub>H</sub> CDR1 | PYSMT             | 157 |
| V <sub>H</sub> CDR2 | SIVPYGDLTQYADSVKG | 158 |
| V <sub>H</sub> CDR3 | GPGWGSFDL         | 159 |
| V <sub>L</sub> CDR1 | QASQDINNLYN       | 160 |
| V <sub>L</sub> CDR2 | DASNLQT           | 161 |
| V <sub>L</sub> CDR3 | QQHESQSPT         | 162 |
| mAb28               |                   |     |
| V <sub>H</sub> CDR1 | RYSMT             | 163 |
| V <sub>H</sub> CDR2 | SIVPYGDLTQYADSVKG | 164 |
| V <sub>H</sub> CDR3 | GPGWGSFDL         | 165 |
| V <sub>L</sub> CDR1 | QASQDINNLYN       | 166 |
| V <sub>L</sub> CDR2 | DASNLQT           | 167 |
| V <sub>L</sub> CDR3 | QQHESQSPT         | 168 |
| mAb29               |                   |     |
| V <sub>H</sub> CDR1 | SYSMT             | 169 |
| V <sub>H</sub> CDR2 | SIVPYGDLTQYADSVKG | 170 |
| V <sub>H</sub> CDR3 | GPGWGSFDL         | 171 |
| V <sub>L</sub> CDR1 | QASQDINNLYN       | 172 |
| V <sub>L</sub> CDR2 | DASNLQT           | 173 |
| V <sub>L</sub> CDR3 | QQHESQSPT         | 174 |
| mAb30               |                   |     |
| V <sub>H</sub> CDR1 | TYSMT             | 175 |
| V <sub>H</sub> CDR2 | SIVPYGDLTQYADSVKG | 176 |
| V <sub>H</sub> CDR3 | GPGWGSFDL         | 177 |
| V <sub>L</sub> CDR1 | QASQDINNLYN       | 178 |
| V <sub>L</sub> CDR2 | DASNLQT           | 179 |
| V <sub>L</sub> CDR3 | QQHESQSPT         | 180 |
| mAb31               |                   |     |
| V <sub>H</sub> CDR1 | VYSMT             | 181 |

|                     |                   |     |
|---------------------|-------------------|-----|
| V <sub>H</sub> CDR2 | SIVPYGDLTQYADSVKG | 182 |
| V <sub>H</sub> CDR3 | GPGWGSFDL         | 183 |
| V <sub>L</sub> CDR1 | QASQDINNYLN       | 184 |
| V <sub>L</sub> CDR2 | DASNLQT           | 185 |
| V <sub>L</sub> CDR3 | QQHESQSPT         | 186 |
| <b>mAb32</b>        |                   |     |
| V <sub>H</sub> CDR1 | YYSMT             | 187 |
| V <sub>H</sub> CDR2 | SIVPYGDLTQYADSVKG | 188 |
| V <sub>H</sub> CDR3 | GPGWGSFDL         | 189 |
| V <sub>L</sub> CDR1 | QASQDINNYLN       | 190 |
| V <sub>L</sub> CDR2 | DASNLQT           | 191 |
| V <sub>L</sub> CDR3 | QQHESQSPT         | 192 |

Thus, in certain embodiments, an antibody or antigen binding fragment thereof comprises a V<sub>H</sub> sequence that comprises complementary determining region V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences selected from **Table A1** and variants thereof which bind to IL-11R $\alpha$ ; and a V<sub>L</sub> sequence that comprises complementary determining region V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences selected from **Table A1** and variants thereof which bind to IL-11R $\alpha$ .

In certain embodiments, the CDR sequences are as follows:

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 1-3, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 4-6, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 7-9, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 10-12, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 13-15, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 16-18, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 19-21, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 22-24, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 25-27, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 28-30, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 31-33, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 34-36, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 37-39, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 40-42, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 43-45, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 46-48, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 49-51, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 52-54, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 55-57, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 58-60, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 61-63, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 64-66, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 67-69, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 70-72, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 73-75, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 76-78, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 79-81, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 82-84, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 85-87, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 88-90, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 91-93, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 94-96, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 97-99, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 100-102, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 103-105, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 106-108, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 109-111, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 112-114, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 115-117, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 118-120, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 121-123, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 124-126, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 127-129, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 130-132, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 133-135, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 136-138, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 139-141, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 142-144, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 145-147, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 148-150, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 151-153, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 154-156, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 157-159, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 160-162, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 163-165, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 166-168, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 169-171, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 172-174, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 175-177, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 178-180, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 181-183, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 184-186, respectively; or

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 187-189, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 190-192, respectively.

Also included are minor variants the foregoing CDRs. Exemplary variants bind to IL-11R $\alpha$  and have 1, 2, or 3 total alterations in any one or more of the individual CDRs, for example, any one or more the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, V<sub>H</sub>CDR3, V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and/or V<sub>L</sub>CDR3 sequences described herein. Exemplary “alterations” include amino acid substitutions, additions, and deletions.

Exemplary V<sub>H</sub> and V<sub>L</sub> sequences are provided in **Table A2** below.

| <b>Table A2: Exemplary Polypeptide Sequences</b> |                                                                                                                                |                   |
|--------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------|-------------------|
| <b>Description</b>                               | <b>Sequence</b>                                                                                                                | <b>SEQ ID NO:</b> |
| <b>mAb1</b>                                      |                                                                                                                                |                   |
| Heavy chain variable region (V <sub>H</sub> )    | EVQLLESGGGLVQPGGSLRLSCAASGFTFSWYSMTWVRQAPGKGLEWVS<br>SIVPSGGHTQYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAK<br>GPDWGSFDLWGRGTLTVSS | 193               |
| Light chain variable region (V <sub>L</sub> )    | DIQMTQSPSSLSASVGDRVTITCQASQDINNYLNWYQQKPGKAPKLLIY<br>DASNLQGTGVPSRFSGSGSGTDFFTISSLPEDATYYCQQHESQSPTF<br>GPGTKVDIK              | 194               |
| <b>mAb2</b>                                      |                                                                                                                                |                   |
| Heavy chain variable region (V <sub>H</sub> )    | EVQLLESGGGLVQPGGSLRLSCAASGFTFSWYSMTWVRQAPGKGLEWVS<br>SIVPSGGHTQYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAK<br>GPDWGSFDLWGRGTLTVSS | 195               |

|                                               |                                                                                                                                |     |
|-----------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------|-----|
| Light chain variable region (V <sub>L</sub> ) | DIQMTQSPSSLSASVGDRVTITCQASQDINNYLNWYQQKPGKAPKLLIY<br>DASNLQGTGVPSRFSGSGSGTDFTFTISSLQPEDATYYCQQHEFQSPTF<br>GPGTKVDIK            | 196 |
| <b>mAb3</b>                                   |                                                                                                                                |     |
| Heavy chain variable region (V <sub>H</sub> ) | EVQLLESGGGLVQPGGSLRLSCAASGFTFSWYSMTWVRQAPGKGLEWVS<br>SIVPYGDLTQYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAK<br>GPDWGSFDLWGRGTLTVSS | 197 |
| Light chain variable region (V <sub>L</sub> ) | DIQMTQSPSSLSASVGDRVTITCQASQDINNYLNWYQQKPGKAPKLLIY<br>DASNLQGTGVPSRFSGSGSGTDFTFTISSLQPEDATYYCQQHESQSPTF<br>GPGTKVDIK            | 198 |
| <b>mAb4</b>                                   |                                                                                                                                |     |
| Heavy chain variable region (V <sub>H</sub> ) | EVQLLESGGGLVQPGGSLRLSCAASGFTFSWYSMTWVRQAPGKGLEWVS<br>SIVPYGDLTQYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAK<br>GPDWGSFDLWGRGTLTVSS | 199 |
| Light chain variable region (V <sub>L</sub> ) | DIQMTQSPSSLSASVGDRVTITCQASQDINNYLNWYQQKPGKAPKLLIY<br>DASNLQGTGVPSRFSGSGSGTDFTFTISSLQPEDATYYCQQHEFQSPTF<br>GPGTKVDIK            | 200 |
| <b>mAb5</b>                                   |                                                                                                                                |     |
| Heavy chain variable region (V <sub>H</sub> ) | EVQLLESGGGLVQPGGSLRLSCAASGFTFSWYSMTWVRQAPGKGLEWVS<br>SIVPYGDLTQYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAK<br>GPGWGSFDLWGRGTLTVSS | 201 |
| Light chain variable region (V <sub>L</sub> ) | DIQMTQSPSSLSASVGDRVTITCQASQDINNYLNWYQQKPGKAPKLLIY<br>DASNLQGTGVPSRFSGSGSGTDFTFTISSLQPEDATYYCQQHESQSPTF<br>GPGTKVDIK            | 202 |
| <b>mAb6</b>                                   |                                                                                                                                |     |
| Heavy chain variable region (V <sub>H</sub> ) | EVQLLESGGGLVQPGGSLRLSCAASGFTFSWYSMTWVRQAPGKGLEWVS<br>SIVPYGDLTQYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAK<br>GPGWGSFDLWGRGTLTVSS | 203 |
| Light chain variable region (V <sub>L</sub> ) | DIQMTQSPSSLSASVGDRVTITCQASQDINNYLNWYQQKPGKAPKLLIY<br>DASNLQGTGVPSRFSGSGSGTDFTFTISSLQPEDATYYCQQHEFQSPTF<br>GPGTKVDIK            | 204 |
| <b>mAb7</b>                                   |                                                                                                                                |     |
| Heavy chain variable region (V <sub>H</sub> ) | EVQLLESGGGLVQPGGSLRLSCAASGFTFSNYAMSWVRQAPGKGLEWVS<br>SIVPYGDLTQYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAK<br>GPGWGSFDLWGRGTLTVSS | 205 |
| Light chain variable region (V <sub>L</sub> ) | DIQMTQSPSSLSASVGDRVTITCQASQDINNYLNWYQQKPGKAPKLLIY<br>DASNLQGTGVPSRFSGSGSGTDFTFTISSLQPEDATYYCQQHESQSPTF<br>GPGTKVDIK            | 206 |
| <b>mAb8</b>                                   |                                                                                                                                |     |
| Heavy chain variable region (V <sub>H</sub> ) | EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVS<br>SIVPYGDLTQYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAK<br>GPGWGSFDLWGRGTLTVSS | 207 |
| Light chain variable region (V <sub>L</sub> ) | DIQMTQSPSSLSASVGDRVTITCQASQDINNYLNWYQQKPGKAPKLLIY<br>DASNLQGTGVPSRFSGSGSGTDFTFTISSLQPEDATYYCQQHESQSPTF<br>GPGTKVDIK            | 208 |
| <b>mAb9</b>                                   |                                                                                                                                |     |

|                                               |                                                                                                                                |     |
|-----------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------|-----|
| Heavy chain variable region (V <sub>H</sub> ) | EVQLLESGGGLVQPGGSLRLSCAASGFTFSWYSMTWVRQAPGKGLEWVS<br>GIVPYGDLTQYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAK<br>GPGWGSFDLWGRGTLTVSS | 209 |
| Light chain variable region (V <sub>L</sub> ) | DIQMTQSPSSLSASVGDRVTITCQASQDINNYLNWYQQKPGKAPKLLIY<br>DASNLQGTGVPSRFSGSGSGTDFTFTISSLPEDIATYYCQQHESQSPTF<br>GPGTKVDIK            | 210 |
| <b>mAb10</b>                                  |                                                                                                                                |     |
| Heavy chain variable region (V <sub>H</sub> ) | EVQLLESGGGLVQPGGSLRLSCAASGFTFSWYSMTWVRQAPGKGLEWVS<br>SIVAYGDLTQYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAK<br>GPGWGSFDLWGRGTLTVSS | 211 |
| Light chain variable region (V <sub>L</sub> ) | DIQMTQSPSSLSASVGDRVTITCQASQDINNYLNWYQQKPGKAPKLLIY<br>DASNLQGTGVPSRFSGSGSGTDFTFTISSLPEDIATYYCQQHESQSPTF<br>GPGTKVDIK            | 212 |
| <b>mAb11</b>                                  |                                                                                                                                |     |
| Heavy chain variable region (V <sub>H</sub> ) | EVQLLESGGGLVQPGGSLRLSCAASGFTFSWYSMTWVRQAPGKGLEWVS<br>SIVDYGDLTQYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAK<br>GPGWGSFDLWGRGTLTVSS | 213 |
| Light chain variable region (V <sub>L</sub> ) | DIQMTQSPSSLSASVGDRVTITCQASQDINNYLNWYQQKPGKAPKLLIY<br>DASNLQGTGVPSRFSGSGSGTDFTFTISSLPEDIATYYCQQHESQSPTF<br>GPGTKVDIK            | 214 |
| <b>mAb12</b>                                  |                                                                                                                                |     |
| Heavy chain variable region (V <sub>H</sub> ) | EVQLLESGGGLVQPGGSLRLSCAASGFTFSWYSMTWVRQAPGKGLEWVS<br>SIVPYGDLTQYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAK<br>GPGWYSFDLWGRGTLTVSS | 215 |
| Light chain variable region (V <sub>L</sub> ) | DIQMTQSPSSLSASVGDRVTITCQASQDINNYLNWYQQKPGKAPKLLIY<br>DASNLQGTGVPSRFSGSGSGTDFTFTISSLPEDIATYYCQQHESQSPTF<br>GPGTKVDIK            | 216 |
| <b>mAb13</b>                                  |                                                                                                                                |     |
| Heavy chain variable region (V <sub>H</sub> ) | EVQLLESGGGLVQPGGSLRLSCAASGFTFSWYSMTWVRQAPGKGLEWVS<br>SIVPYGDLTQYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAK<br>PEDWGRFDLWGRGTLTVSS | 217 |
| Light chain variable region (V <sub>L</sub> ) | DIQMTQSPSSLSASVGDRVTITCQASQDINNYLNWYQQKPGKAPKLLIY<br>DASNLQGTGVPSRFSGSGSGTDFTFTISSLPEDIATYYCQQHESQSPTF<br>GPGTKVDIK            | 218 |
| <b>mAb14</b>                                  |                                                                                                                                |     |
| Heavy chain variable region (V <sub>H</sub> ) | EVQLLESGGGLVQPGGSLRLSCAASGFTFSWYSMTWVRQAPGKGLEWVS<br>SIVPYGDLTQYAESVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAK<br>GPGWGSFDLWGRGTLTVSS | 219 |
| Light chain variable region (V <sub>L</sub> ) | DIQMTQSPSSLSASVGDRVTITCQASQDINNYLNWYQQKPGKAPKLLIY<br>DASNLQGTGVPSRFSGSGSGTDFTFTISSLPEDIATYYCQQHESQSPTF<br>GPGTKVDIK            | 220 |
| <b>mAb15</b>                                  |                                                                                                                                |     |
| Heavy chain variable region (V <sub>H</sub> ) | EVQLLESGGGLVQPGGSLRLSCAASGFTFSWYSMTWVRQAPGKGLEWVS<br>SIVPYGDLTQYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAK<br>GPGWGSFDLWGRGTLTVSS | 221 |

|                                               |                                                                                                                                 |     |
|-----------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------|-----|
| Light chain variable region (V <sub>L</sub> ) | DIQMTQSPSSLSASVGDRVTITCQASQDINNYLNWYQQKPGKAPKLLIY<br>DASNLQGTGVPSRFSGSGSGTDFTFTISSLPEDIATYYCQQHETQTPTF<br>GPGTKVDIK             | 222 |
| <b>mAb16</b>                                  |                                                                                                                                 |     |
| Heavy chain variable region (V <sub>H</sub> ) | EVQLLESGGGLVQPGGSLRLSCAASGFTFSAYSMTWVRQAPGKGLEWVS<br>SIVPYGDLTQYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAK<br>GPGWGSFDLWGRGTLTVSS  | 223 |
| Light chain variable region (V <sub>L</sub> ) | DIQMTQSPSSLSASVGDRVTITCQASQDINNYLNWYQQKPGKAPKLLIY<br>DASNLQGTGVPSRFSGSGSGTDFTFTISSLPEDIATYYCQQHESQSPTF<br>GPGTKVDIK             | 224 |
| <b>mAb17</b>                                  |                                                                                                                                 |     |
| Heavy chain variable region (V <sub>H</sub> ) | EVQLLESGGGLVQPGGSLRLSCAASGFTFSAYSMTWVRQAPGKGLEWVS<br>SIVPYGDLTQYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAK<br>GPGWGSFDLWGRGTLTVSS  | 225 |
| Light chain variable region (V <sub>L</sub> ) | DIQMTQSPSSLSASVGDRVTITCQASQDINNYLNWYQQKPGKAPKLLIY<br>DASNLQGTGVPSRFSGSGSGTDFTFTISSLPEDIATYYCQQHESQSPTF<br>GPGTKVDIK             | 226 |
| <b>mAb18</b>                                  |                                                                                                                                 |     |
| Heavy chain variable region (V <sub>H</sub> ) | EVQLLESGGGLVQPGGSLRLSCAASGFTFSEYSMTWVRQAPGKGLEWVS<br>SIVPYGDLTQYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAK<br>GPGWGSFDLWGRGTLTVSS  | 227 |
| Light chain variable region (V <sub>L</sub> ) | DIQMTQSPSSLSASVGDRVTITCQASQDINNYLNWYQQKPGKAPKLLIY<br>DASNLQGTGVPSRFSGSGSGTDFTFTISSLPEDIATYYCQQHESQSPTF<br>GPGTKVDIK             | 228 |
| <b>mAb19</b>                                  |                                                                                                                                 |     |
| Heavy chain variable region (V <sub>H</sub> ) | EVQLLESGGGLVQPGGSLRLSCAASGFTFSFYSM TWVRQAPGKGLEWVS<br>SIVPYGDLTQYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAK<br>GPGWGSFDLWGRGTLTVSS | 229 |
| Light chain variable region (V <sub>L</sub> ) | DIQMTQSPSSLSASVGDRVTITCQASQDINNYLNWYQQKPGKAPKLLIY<br>DASNLQGTGVPSRFSGSGSGTDFTFTISSLPEDIATYYCQQHESQSPTF<br>GPGTKVDIK             | 230 |
| <b>mAb20</b>                                  |                                                                                                                                 |     |
| Heavy chain variable region (V <sub>H</sub> ) | EVQLLESGGGLVQPGGSLRLSCAASGFTFSGYSM TWVRQAPGKGLEWVS<br>SIVPYGDLTQYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAK<br>GPGWGSFDLWGRGTLTVSS | 231 |
| Light chain variable region (V <sub>L</sub> ) | DIQMTQSPSSLSASVGDRVTITCQASQDINNYLNWYQQKPGKAPKLLIY<br>DASNLQGTGVPSRFSGSGSGTDFTFTISSLPEDIATYYCQQHESQSPTF<br>GPGTKVDIK             | 232 |
| <b>mAb21</b>                                  |                                                                                                                                 |     |
| Heavy chain variable region (V <sub>H</sub> ) | EVQLLESGGGLVQPGGSLRLSCAASGFTFSHYSM TWVRQAPGKGLEWVS<br>SIVPYGDLTQYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAK<br>GPGWGSFDLWGRGTLTVSS | 233 |
| Light chain variable region (V <sub>L</sub> ) | DIQMTQSPSSLSASVGDRVTITCQASQDINNYLNWYQQKPGKAPKLLIY<br>DASNLQGTGVPSRFSGSGSGTDFTFTISSLPEDIATYYCQQHESQSPTF<br>GPGTKVDIK             | 234 |
| <b>mAb22</b>                                  |                                                                                                                                 |     |

|                                               |                                                                                                                                 |     |
|-----------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------|-----|
| Heavy chain variable region (V <sub>H</sub> ) | EVQLLESGGGLVQPGGSLRLSCAASGFTFSIYSMTWVRQAPGKGLEWVS<br>SIVPYGDLTQYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAK<br>GPGWGSFDLWGRGTLTVSS  | 235 |
| Light chain variable region (V <sub>L</sub> ) | DIQMTQSPSSLSASVGDRVTITCQASQDINNYLNWYQQKPGKAPKLLIY<br>DASNLQGTGVPSRFSGSGSGTDFTFTISSLPEDIATYYCQQHESQSPTF<br>GPGTKVDIK             | 236 |
| <b>mAb23</b>                                  |                                                                                                                                 |     |
| Heavy chain variable region (V <sub>H</sub> ) | EVQLLESGGGLVQPGGSLRLSCAASGFTFSKY SMTWVRQAPGKGLEWVS<br>SIVPYGDLTQYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAK<br>GPGWGSFDLWGRGTLTVSS | 237 |
| Light chain variable region (V <sub>L</sub> ) | DIQMTQSPSSLSASVGDRVTITCQASQDINNYLNWYQQKPGKAPKLLIY<br>DASNLQGTGVPSRFSGSGSGTDFTFTISSLPEDIATYYCQQHESQSPTF<br>GPGTKVDIK             | 238 |
| <b>mAb24</b>                                  |                                                                                                                                 |     |
| Heavy chain variable region (V <sub>H</sub> ) | EVQLLESGGGLVQPGGSLRLSCAASGFTFSLYSMTWVRQAPGKGLEWVS<br>SIVPYGDLTQYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAK<br>GPGWGSFDLWGRGTLTVSS  | 239 |
| Light chain variable region (V <sub>L</sub> ) | DIQMTQSPSSLSASVGDRVTITCQASQDINNYLNWYQQKPGKAPKLLIY<br>DASNLQGTGVPSRFSGSGSGTDFTFTISSLPEDIATYYCQQHESQSPTF<br>GPGTKVDIK             | 240 |
| <b>mAb25</b>                                  |                                                                                                                                 |     |
| Heavy chain variable region (V <sub>H</sub> ) | EVQLLESGGGLVQPGGSLRLSCAASGFTFSMYSMTWVRQAPGKGLEWVS<br>SIVPYGDLTQYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAK<br>GPGWGSFDLWGRGTLTVSS  | 241 |
| Light chain variable region (V <sub>L</sub> ) | DIQMTQSPSSLSASVGDRVTITCQASQDINNYLNWYQQKPGKAPKLLIY<br>DASNLQGTGVPSRFSGSGSGTDFTFTISSLPEDIATYYCQQHESQSPTF<br>GPGTKVDIK             | 242 |
| <b>mAb26</b>                                  |                                                                                                                                 |     |
| Heavy chain variable region (V <sub>H</sub> ) | EVQLLESGGGLVQPGGSLRLSCAASGFTFSNYSMTWVRQAPGKGLEWVS<br>SIVPYGDLTQYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAK<br>GPGWGSFDLWGRGTLTVSS  | 243 |
| Light chain variable region (V <sub>L</sub> ) | DIQMTQSPSSLSASVGDRVTITCQASQDINNYLNWYQQKPGKAPKLLIY<br>DASNLQGTGVPSRFSGSGSGTDFTFTISSLPEDIATYYCQQHESQSPTF<br>GPGTKVDIK             | 244 |
| <b>mAb27</b>                                  |                                                                                                                                 |     |
| Heavy chain variable region (V <sub>H</sub> ) | EVQLLESGGGLVQPGGSLRLSCAASGFTFSPYSMTWVRQAPGKGLEWVS<br>SIVPYGDLTQYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAK<br>GPGWGSFDLWGRGTLTVSS  | 245 |
| Light chain variable region (V <sub>L</sub> ) | DIQMTQSPSSLSASVGDRVTITCQASQDINNYLNWYQQKPGKAPKLLIY<br>DASNLQGTGVPSRFSGSGSGTDFTFTISSLPEDIATYYCQQHESQSPTF<br>GPGTKVDIK             | 246 |
| <b>mAb28</b>                                  |                                                                                                                                 |     |
| Heavy chain variable region (V <sub>H</sub> ) | EVQLLESGGGLVQPGGSLRLSCAASGFTFSRYSMTWVRQAPGKGLEWVS<br>SIVPYGDLTQYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAK<br>GPGWGSFDLWGRGTLTVSS  | 247 |

|                                               |                                                                                                                               |     |
|-----------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------|-----|
| Light chain variable region (V <sub>L</sub> ) | DIQMTQSPSSLSASVGDRVTITCQASQDINNYLNWYQQKPGKAPKLLIY<br>DASNLQGTGVPSRFSGSGSGTDFTFTISSLQPEDATYYCQQHESQSPTF<br>GPGTKVDIK           | 248 |
| <b>mAb29</b>                                  |                                                                                                                               |     |
| Heavy chain variable region (V <sub>H</sub> ) | EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYSMTWVRQAPGKLEWVS<br>SIVPYGDLTQYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAK<br>GPGWGSFDLWGRGTLTVSS | 249 |
| Light chain variable region (V <sub>L</sub> ) | DIQMTQSPSSLSASVGDRVTITCQASQDINNYLNWYQQKPGKAPKLLIY<br>DASNLQGTGVPSRFSGSGSGTDFTFTISSLQPEDATYYCQQHESQSPTF<br>GPGTKVDIK           | 250 |
| <b>mAb30</b>                                  |                                                                                                                               |     |
| Heavy chain variable region (V <sub>H</sub> ) | EVQLLESGGGLVQPGGSLRLSCAASGFTFSTYSMTWVRQAPGKLEWVS<br>SIVPYGDLTQYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAK<br>GPGWGSFDLWGRGTLTVSS | 251 |
| Light chain variable region (V <sub>L</sub> ) | DIQMTQSPSSLSASVGDRVTITCQASQDINNYLNWYQQKPGKAPKLLIY<br>DASNLQGTGVPSRFSGSGSGTDFTFTISSLQPEDATYYCQQHESQSPTF<br>GPGTKVDIK           | 252 |
| <b>mAb31</b>                                  |                                                                                                                               |     |
| Heavy chain variable region (V <sub>H</sub> ) | EVQLLESGGGLVQPGGSLRLSCAASGFTFSVYSMTWVRQAPGKLEWVS<br>SIVPYGDLTQYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAK<br>GPGWGSFDLWGRGTLTVSS | 253 |
| Light chain variable region (V <sub>L</sub> ) | DIQMTQSPSSLSASVGDRVTITCQASQDINNYLNWYQQKPGKAPKLLIY<br>DASNLQGTGVPSRFSGSGSGTDFTFTISSLQPEDATYYCQQHESQSPTF<br>GPGTKVDIK           | 254 |
| <b>mAb32</b>                                  |                                                                                                                               |     |
| Heavy chain variable region (V <sub>H</sub> ) | EVQLLESGGGLVQPGGSLRLSCAASGFTFSYYSMTWVRQAPGKLEWVS<br>SIVPYGDLTQYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAK<br>GPGWGSFDLWGRGTLTVSS | 255 |
| Light chain variable region (V <sub>L</sub> ) | DIQMTQSPSSLSASVGDRVTITCQASQDINNYLNWYQQKPGKAPKLLIY<br>DASNLQGTGVPSRFSGSGSGTDFTFTISSLQPEDATYYCQQHESQSPTF<br>GPGTKVDIK           | 256 |

Thus, in certain embodiments, an antibody, or antigen binding fragment thereof, binds to IL-11R $\alpha$  and comprises a V<sub>H</sub> sequence and a corresponding V<sub>L</sub> sequence selected from **Table A2**. In certain embodiments, the V<sub>H</sub> comprises a sequence least 80, 85, 90, 95, 97, 98, 99, or 100% identical to a sequence selected from **Table A2**, including, for example, wherein the V<sub>H</sub> has 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 alterations in one or more framework regions. In some embodiments, the V<sub>L</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to a sequence selected from **Table A2**, including, for example, wherein the V<sub>L</sub> has 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 alterations in one or more framework regions.

In some embodiments, the V<sub>H</sub> and V<sub>L</sub> of an antibody or antigen binding fragment are as follows:

the V<sub>H</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 193, and the V<sub>L</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 194;

the V<sub>H</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 195, and the V<sub>L</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 196;

the V<sub>H</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 197, and the V<sub>L</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 198;

the V<sub>H</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 199, and the V<sub>L</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 200;

the V<sub>H</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 201, and the V<sub>L</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 202;

the V<sub>H</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 203, and the V<sub>L</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 204;

the V<sub>H</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 205, and the V<sub>L</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 206;

the V<sub>H</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 207, and the V<sub>L</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 208;

the V<sub>H</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 209, and the V<sub>L</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 210;

the V<sub>H</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 211, and the V<sub>L</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 212;

the V<sub>H</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 213, and the V<sub>L</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 214;

the V<sub>H</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 215, and the V<sub>L</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 216;



the V<sub>H</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 241, and the V<sub>L</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 242;

the V<sub>H</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 243, and the V<sub>L</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 244;

the V<sub>H</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 245, and the V<sub>L</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 246;

the V<sub>H</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 247, and the V<sub>L</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 248;

the V<sub>H</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 249, and the V<sub>L</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 250;

the V<sub>H</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 251, and the V<sub>L</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 252;

the V<sub>H</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 253, and the V<sub>L</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 254; or

the V<sub>H</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 255, and the V<sub>L</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 256.

Also included are variants thereof that bind to IL-11R $\alpha$ , for example, variants having 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 alterations in one or more framework regions of any one or more of the foregoing V<sub>H</sub> and/or V<sub>L</sub> sequences. Exemplary “alterations” include amino acid substitutions, additions, and deletions.

As noted above, an antibody or antigen binding fragment thereof, described herein, binds to IL-11R $\alpha$ , for example, membrane-bound IL-11R $\alpha$ . In certain embodiments, an antibody or an antigen binding fragment thereof binds to human IL-11R $\alpha$ , or a region or fragment thereof. The amino acid sequence of human IL-11R $\alpha$  is provided in **Table B1** below.

| <b>Table B1. Exemplary IL-11R<math>\alpha</math> Sequences</b> |                                                                                                                                                                                                                      |            |
|----------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------|
| Name                                                           | Sequence                                                                                                                                                                                                             | SEQ ID NO: |
| Human IL-11R $\alpha$ membrane precursor                       | MSSSCSGLSRVLVAVATALVSASSPCPQAWGPPGVQYGGQPGRSVKLCCPG<br>VTAGDPVSWFRDGEKLLQGPDGSLGHELVLAQADSTDEGTYYICQTLDGA<br>LGGTVTLQLGYPPARPVVSCQAADYENFSCWSPSQISGLPTRYLTSYRK<br>KTVLGADSQRRSPSTGWPWPCPDPLGAARCVVHGAEFWSQYRINVTEVNP | 257        |

|  |                                                                                                                                                                                                                                                 |  |
|--|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--|
|  | LGASTRLLDVSLQSIILRPDPPQGLRVESVPGYPRRLRASWTYPASWPCQP<br>HFLKFRLOQYRPAQHFAWSTVEPAGLEEVITDAVAGLPHAVRVSARDFLD<br>AGTWSTWSPEAWGTPSTGTIPKEIPAWGQLHTQPEVEPQVDSAPPRLSL<br>QPHPRLLDHRDSVEQVAVLASLGLLSFLGLVAGALALGLWLRRLRRGGKDG<br>SPKPGFLASVIPVDRRPGAPNL |  |
|--|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--|

Thus, in certain embodiments, an antibody or antigen binding fragment thereof binds to an IL-11R $\alpha$  sequence in **Table B1**, for example, a human IL-11R $\alpha$  sequence. In particular embodiments, an antibody or antigen binding fragment thereof binds to a fibronectin type-III domain in human IL-11R $\alpha$ , composed, for example, of approximately residues 112-219 of SEQ ID NO: 257.

In some embodiments, an antibody or antigen binding fragment thereof binds to human IL-11R $\alpha$  with a binding affinity of about 1 pM to about 10 pM to about 500 pM, or about, at least about, or less than about 1, 5, 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 300, 400, or 500 pM, or optionally with an affinity that ranges from about 1 pM to about 500 pM, about 1 pM to about 400 pM, about 1 pM to about 300 pM, about 1 pM to about 200 pM, about 1 pM to about 100 pM, about 1 pM to about 50 pM, about 1 pM to about 40 pM, about 1 pM to about 30 pM, about 1 pM to about 20 pM, about 1 pM to about 10 pM, about 1 pM to about 5 pM, about 5 pM to about 500 pM, about 5 pM to about 400 pM, about 5 pM to about 300 pM, about 5 pM to about 200 pM, about 5 pM to about 100 pM, about 5 pM to about 50 pM, about 5 pM to about 40 pM, about 5 pM to about 30 pM, about 5 pM to about 20 pM, about 5 pM to about 10 pM, about 10 pM to about 500 pM, about 10 pM to about 400 pM, about 10 pM to about 300 pM, about 10 pM to about 200 pM, about 10 pM to about 100 pM, about 10 pM to about 50 pM, about 10 pM to about 40 pM, about 10 pM to about 30 pM, about 10 pM to about 20 pM, or about 20 pM to about 500 pM, about 20 pM to about 400 pM, about 20 pM to about 300 pM, about 20 pM to about 200 pM, about 20 pM to about 100 pM, about 20 pM to about 50 pM, about 20 pM to about 40 pM, about 20 pM to about 30 pM, or about 30 pM to about 500 pM, about 30 pM to about 400 pM, about 30 pM to about 300 pM, about 30 pM to about 200 pM, about 30 pM to about 100 pM, about 30 pM to about 50 pM, or about 30 pM to about 40 pM. In certain embodiments, an antibody or antigen binding fragment thereof, has increased binding affinity for human IL-11R $\alpha$  relative to that of the TS7 and 8E2 antibodies (see, for example, U.S. Patent Nos. 9,796,782; 9,340,618).

In some embodiments, an antibody, or antigen binding fragment thereof, is an IL-11R $\alpha$  antagonist. In some instances, an antibody, or antigen binding fragment thereof, antagonizes the binding and/or signaling activity between IL-11R $\alpha$  and its ligand, IL-11. In some embodiments, an antibody, or antigen binding fragment thereof, antagonizes or reduces the binding and/or signaling activity between IL-11R $\alpha$  and IL-11 by about or at least about 10-1000% (e.g., about 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 90, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000% or more), for example, in a cell-based assay. In some embodiments, an anti-IL-11R $\alpha$  antibody or antigen binding fragment thereof reduces IL-11-mediated STAT3 phosphorylation. In an exemplary assay, cells expressing IL-11R $\alpha$  and gp130 are cultured in the presence of IL-11 in the presence or absence

of the IL-11R $\alpha$ -binding protein. The level of STAT3 phosphorylation is then assessed by Western blotting or FACS using an antibody specific for phosphorylated STAT3. An exemplary assay making use of FACS is described in Dams-Kozłowska et al., *BMC Biotechnol*, 12: 8, 2012. In certain embodiments, an antibody or antigen binding fragment thereof, has increased potency as an IL-11 signaling antagonist relative to that of the TS7 and 8E2 antibodies (see, for example, U.S. Patent Nos. 9,796,782; 9,340,618). In certain embodiments, an antibody or antigen binding fragment thereof has no detectable agonist activity with respect to IL-11 signaling.

In some embodiments, an antibody, or antigen binding fragment thereof, inhibits or otherwise reduces IL-11R $\alpha$  dimerization or complex formation, for example, with gp130. In certain embodiments, an antibody, or antigen binding fragment thereof, inhibits or otherwise reduces IL-11R $\alpha$  dimerization or complex formation by about or at least about 10-1000% (e.g., about 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 90, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000% or more), for example, in a cell-based assay.

In some embodiments, an anti-IL-11R $\alpha$  antibody or antigen binding fragment thereof reduces proliferation of cells (e.g., BaF3 cells, B9 cells, T10 cells) expressing IL-11R $\alpha$  and gp130 (e.g., cells naturally-expressing or modified to express both proteins) which are cultured in the presence of IL-11. Methods for assessing cell proliferation are known in the art and include, for example, MTT reduction and/or thymidine incorporation. Assays with B9 cells or T10 cells are described (see Dams-Kozłowska et al., *BMC Biotechnol*, 12: 8, 2012; and Yokote et al., *J AOAC*, 83: 1053-1057, 2000). For T10 cells, proliferation can be measured by colorimetrically detecting reduction of the tetrazolium compound, 4-[3-(4-iodophenyl)-2-(4-nitrophenyl)-2H-5-tetrazolio]-1,3-benzene disulfonate (WST-1). An IL-11R $\alpha$ -binding protein that reduces the level of proliferation compared to the level observed in the absence of the IL-11R $\alpha$ -binding protein is considered to reduce or otherwise reduce IL-11R signaling.

In some embodiments, an anti-IL-11R $\alpha$  antibody or antigen binding fragment thereof reduces IL-11-mediated proliferation of cancer cells (for example, gastric cancer cells, acute myelogenous leukemia (AML) cells). In an exemplary assay, cancer cells (e.g., AGN, MKN45 gastric cancer cells) are cultured in the presence of IL-11 in the presence or absence of the IL-11R $\alpha$ -binding protein. For AML cells, the cells may also be cultured in the presence of G-CSF. Proliferation of the cells is then measured using standard techniques, e.g., as discussed herein and/or by assessing formation of L-CFU in the case of AML cells. Exemplary assays adaptable to the present disclosure are included in Zhang et al., *Int J Biol Sci*, 8: 383-393, 2012 and Kimura et al., *Leukemia*, 13: 1018-1027, 1999.

Merely for illustrative purposes, the binding interactions between IL-11R $\alpha$  an antibody, or antigen binding fragment thereof, described herein, or the binding/signaling between IL-11R $\alpha$  and IL-11, can be detected and quantified using a variety of routine methods, including Biacore® assays (for example, with appropriately tagged soluble reagents, bound to a sensor chip), FACS analyses with cells expressing IL-11R $\alpha$  on the cell surface (either native, or recombinant), immunoassays,

fluorescence staining assays, ELISA assays, and microcalorimetry approaches such as ITC (Isothermal Titration Calorimetry). Similarly, the functional properties of anti-IL-11R $\alpha$  antibodies may be assessed using a variety of methods known to the skilled person affinity/binding assays (for example, surface plasmon resonance, competitive inhibition assays); cytotoxicity assays, cell viability assays, cell proliferation or differentiation assays, cancer cell and/or tumor growth inhibition using *in vitro* or *in vivo* models. Other assays may test the ability of antibodies described herein to block normal IL-11R $\alpha$ -mediated responses. The antibodies described herein may also be tested for *in vitro* and *in vivo* efficacy. Such assays may be performed using well-established protocols known to the skilled person (see *e.g.*, Current Protocols in Molecular Biology (Greene Publ. Assoc. Inc. & John Wiley & Sons, Inc., NY, NY); Current Protocols in Immunology (Edited by: John E. Coligan, Ada M. Kruisbeek, David H. Margulies, Ethan M. Shevach, Warren Strober 2001 John Wiley & Sons, NY, NY); or commercially available kits.

In particular embodiments, the Fc region of an antibody, or antigen binding fragment thereof, comprises, consists, or consists essentially an IgA (including subclasses IgA1 and IgA2), IgD, IgE, IgG (including subclasses IgG1, IgG2, IgG3, and IgG4), or IgM Fc domain, optionally a human Fc domain, or a hybrid and/or variant thereof. In particular embodiments, the Fc region comprises, consists, or consists essentially of the Fc from human IgG1 or IgG4 (see, *e.g.*, Allberse and Schuurman, Immunology. 105:9-19, 2002), or a fragment or variant thereof. **Table F1** below provides exemplary sequences (CH1, hinge (underlined), CH2, and CH3 regions) from human IgG4. Examples of variant IgG4 sequences that can be employed include the S228P/S241P variant.

| <b>Table F1. Exemplary IgG4 Fc Sequences</b> |                                                                                                                                                                                                                                                                                                                                                                     |                   |
|----------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------|
| <b>Name</b>                                  | <b>Sequence</b>                                                                                                                                                                                                                                                                                                                                                     | <b>SEQ ID NO:</b> |
| Wild-type IgG4                               | ASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALT<br>SGVHTFPAVLQSSGLYSLSSVVTVPSSSLGKTYTCNVDPKPSNTKV<br>DKRVESKYGPPCPSCPAPEFLGGPSVFLFPPKPKDTLMISRTPEVTC<br>VVVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVVSVLTV<br>LHQDWLNGKEYKCKVSNKGLPSSIEKTIKAKGQPREPQVYTLPPSQ<br>EEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDG<br>SFFLYSRLTVDKSRWQEGNVFSCVMHEALHNHYTQKSLSLGLK          | 258               |
| S228P (EU)<br>S241P (Kabat)                  | ASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALT<br>SGVHTFPAVLQSSGLYSLSSVVTVPSSSLGKTYTCNVDPKPSNTKV<br>DKRVESKYGPPCP <b>PC</b> PAPEFLGGPSVFLFPPKPKDTLMISRTPEVTC<br>VVVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVVSVLTV<br>LHQDWLNGKEYKCKVSNKGLPSSIEKTIKAKGQPREPQVYTLPPSQ<br>EEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDG<br>SFFLYSRLTVDKSRWQEGNVFSCVMHEALHNHYTQKSLSLGLK | 259               |

In certain embodiments, an antibody or antigen binding fragment thereof comprises variant or otherwise modified Fc region(s), including those having altered properties or biological activities relative to wild-type Fc region(s). Examples of modified Fc regions include those having mutated sequences, for instance, by substitution, insertion, deletion, or truncation of one or more amino acids relative to a wild-type sequence, hybrid Fc polypeptides composed of domains from different

immunoglobulin classes/subclasses, Fc polypeptides having altered glycosylation/sialylation patterns, and Fc polypeptides that are modified or derivatized, for example, by biotinylation (*see, e.g.*, US Application No. 2010/0209424), phosphorylation, sulfation, etc., or any combination of the foregoing. Such modifications can be employed to alter (*e.g.*, increase, decrease) the binding properties of the Fc region to one or more particular FcRs (*e.g.*, FcγRI, FcγRIIa, FcγRIIb, FcγRIIc, FcγRIIIa, FcγRIIIb, FcRn), its pharmacokinetic properties (*e.g.*, stability or half-life, bioavailability, tissue distribution, volume of distribution, concentration, elimination rate constant, elimination rate, area under the curve (AUC), clearance,  $C_{max}$ ,  $t_{max}$ ,  $C_{min}$ , fluctuation), its immunogenicity, its complement fixation or activation, and/or the CDC/ADCC/ADCP-related activities of the Fc region, among other properties described herein, relative to a corresponding wild-type Fc sequence of an antibody or antigen binding fragment thereof. Included are modified Fc regions of human and/or mouse origin.

In certain embodiments, an antibody or antigen binding fragment thereof comprises a hybrid Fc region, for example, an Fc region that comprises a combination of Fc domains (*e.g.*, hinge, CH<sub>2</sub>, CH<sub>3</sub>, CH<sub>4</sub>) from immunoglobulins of different species (*e.g.*, human, mouse), different Ig classes, and/or different Ig subclasses. Also included are antibodies or antigen binding fragments thereof that comprise derivatized or otherwise modified Fc regions. In certain aspects, the Fc region is modified by phosphorylation, sulfation, acrylation, glycosylation, methylation, farnesylation, acetylation, amidation, and the like, for instance, relative to a wild-type or naturally-occurring Fc region. In certain embodiments, the Fc region comprises wild-type or native glycosylation patterns, or alternatively, it comprises increased glycosylation relative to a native form, decreased glycosylation relative to a native form, or it is entirely deglycosylated. As one example of a modified Fc glycoform, decreased glycosylation of an Fc region reduces binding to the C1q region of the first complement component C1, a decrease in ADCC-related activity, and/or a decrease in CDC-related activity. Certain embodiments thus employ a deglycosylated or aglycosylated Fc region. *See, e.g.*, WO 2005/047337 for the production of exemplary aglycosylated Fc regions. Another example of an Fc region glycoform is generated by substituting the Q295 position with a cysteine residue (*see, e.g.*, U.S. Application No. 2010/0080794), according to the Kabat *et al.* numbering system. Certain embodiments include Fc regions where about 80-100% of the glycoprotein in Fc region comprises a mature core carbohydrate structure that lacks fucose (*see, e.g.*, U.S. Application No. 2010/0255013). Some embodiments include Fc regions that are optimized by substitution or deletion to reduce the level of fucosylation, for instance, to increase affinity for FcγRI, FcγRIa, or FcγRIIIa, and/or to improve phagocytosis by FcγRIIIa-expressing cells (*see* U.S. Application Nos. 2010/0249382 and 2007/0148170).

As another example of a modified Fc glycoform, an Fc region of an antibody or antigen binding fragment thereof may comprise oligomannose-type N-glycans, and optionally have one or more of the following: increased ADCC effector activity, increased binding affinity for FcγRIIIA (and

certain other FcRs), similar or increased binding specificity for the target of the IL-11R $\alpha$  polypeptide, similar or higher binding affinity for the target of the IL-11R $\alpha$  polypeptide, and/or similar or lower binding affinity for mannose receptor, relative to a corresponding Fc region that contains complex-type N-glycans (see, e.g., U.S. Application No. 2007/0092521 and U.S. Patent No. 7,700,321). As another example, enhanced affinity of Fc regions for Fc $\gamma$ Rs has been achieved using engineered glycoforms generated by expression of antibodies in engineered or variant cell lines (see, e.g., Umana et al., Nat Biotechnol. 17:176-180, 1999; Davies et al., Biotechnol Bioeng. 74:288-294, 2001; Shields et al., J Biol Chem. 277:26733-26740, 2002; Shinkawa et al., 2003, J Biol Chem. 278:3466-3473, 2003; and U.S. Application No. 2007/0111281). Certain Fc region glycoforms comprise an increased proportion of N-glycoside bond type complex sugar chains, which do not have the 1-position of fucose bound to the 6-position of N-acetylglucosamine at the reducing end of the sugar chain (see, e.g., U.S. Application No. 2010/0092997). Particular embodiments may include IgG Fc region that is glycosylated with at least one galactose moiety connected to a respective terminal sialic acid moiety by an  $\alpha$ -2,6 linkage, optionally where the Fc region has a higher anti-inflammatory activity relative to a corresponding, wild-type Fc region (see U.S. Application No. 2008/0206246). Certain of these and related altered glycosylation approaches have generated substantial enhancements of the capacity of Fc regions to selectively bind FcRs such as Fc $\gamma$ RIII, to mediate ADCC, and to alter other properties of Fc regions, as described herein.

Certain variant, fragment, hybrid, or otherwise modified Fc regions of an antibody or antigen binding fragment thereof may have altered binding to one or more FcRs, and/or corresponding changes to effector function, relative to a corresponding, wild-type Fc sequence (e.g., same species, same Ig class, same Ig subclass). For instance, such Fc regions may have increased binding to one or more of Fc $\gamma$  receptors, Fc $\alpha$  receptors, Fc $\epsilon$  receptors, and/or the neonatal Fc receptor, relative to a corresponding, wild-type Fc sequence. In other embodiments, variant, fragment, hybrid, or modified Fc regions may have decreased binding to one or more of Fc $\gamma$  receptors, Fc $\alpha$  receptors, Fc $\epsilon$  receptors, and/or the neonatal Fc receptor, relative to a corresponding, wild-type Fc sequence. Specific FcRs are described elsewhere herein.

In some embodiments, an antibody comprises an Fc domain, comprising one or more mutations to increase binding to one or more of Fc $\gamma$  receptors, Fc $\alpha$  receptors, Fc $\epsilon$  receptors, and/or the neonatal Fc receptor, relative to a corresponding, wild-type Fc sequence. In some embodiments, an antibody comprises an IgG1 or IgG3 Fc domain, comprising one or more mutations to increase binding to one or more of Fc $\gamma$  receptors, Fc $\alpha$  receptors, Fc $\epsilon$  receptors, and/or the neonatal Fc receptor, relative to a corresponding, wild-type Fc sequence. In some embodiments, an antibody comprises an Fc domain, comprising one or more mutations to increase effector function. In some embodiments the at least one antibody comprises an Fc domain selected from a human IgG1 and IgG3, comprising one or more mutations to increase effector function.

In some embodiments, an antibody is a blocking antibody that comprises an Fc domain with high effector activity. In some embodiments, the blocking antibody comprises an Fc domain selected from a human IgG1 and IgG3, comprising one or more mutations to increase effector function. In some embodiments, an antibody is a partial-blocking antibody that comprises an Fc domain with high effector activity. In some embodiments, the partial-blocking antibody comprises an Fc domain selected from a human IgG1 and IgG3, comprising one or more mutations to increase effector function. In some embodiments, an antibody is a non-blocking antibody that comprises an Fc domain with high effector activity. In some embodiments, the non-blocking antibody comprises an Fc domain selected from a human IgG1 or IgG3, comprising one or more mutations to increase effector function.

In some embodiments, an antibody comprises an Fc domain, comprising one or more mutations to decrease binding to one or more of Fc $\gamma$  receptors, Fc $\alpha$  receptors, Fc $\epsilon$  receptors, and/or the neonatal Fc receptor, relative to a corresponding, wild-type Fc sequence. In some embodiments, an antibody comprises an IgG1 or IgG3 Fc domain, comprising one or more mutations to decrease binding to one or more of Fc $\gamma$  receptors, Fc $\alpha$  receptors, Fc $\epsilon$  receptors, and/or the neonatal Fc receptor, relative to a corresponding, wild-type Fc sequence. In some embodiments, an antibody comprises an Fc domain, comprising one or more mutations to decrease effector function. In some embodiments, an antibody comprises an Fc domain selected from a human IgG2 and IgG4, comprising one or more mutations to decrease effector function.

In some embodiments, an antibody is a blocking antibody comprising an Fc domain with low effector activity. In some embodiments, the blocking antibody comprises an Fc domain selected from a human IgG2 and IgG4, comprising one or more mutations to decrease effector function. In some embodiments, an antibody is a partial-blocking antibody comprising an Fc domain with low effector activity. In some embodiments, the partial-blocking antibody comprises an Fc domain selected from a human IgG2 and IgG4, comprising one or more mutations to decrease effector function. In some embodiments, an antibody is a non-blocking antibody comprising an Fc domain with low effector activity. In some embodiments, the non-blocking antibody comprises an Fc domain selected from a human IgG2 and IgG4, comprising one or more mutations to decrease effector function.

Specific examples of Fc variants having altered (e.g., increased, decreased) effector function/FcR binding can be found, for example, in U.S. Pat. Nos. 5,624,821 and 7,425,619; U.S. Application Nos. 2009/0017023, 2009/0010921, and 2010/0203046; and WO 2000/42072 and WO 2004/016750. Certain examples include human Fc regions having a one or more substitutions at position 298, 333, and/or 334, for example, S298A, E333A, and/or K334A (based on the numbering of the EU index of Kabat et al.), which have been shown to increase binding to the activating receptor Fc $\gamma$ RIIIa and reduce binding to the inhibitory receptor Fc $\gamma$ RIIb. These mutations can be combined to obtain double and triple mutation variants that have further improvements in binding to FcRs. Certain embodiments include a S298A/E333A/K334A triple mutant, which has increased binding to Fc $\gamma$ RIIIa, decreased binding to Fc $\gamma$ RIIb, and increased ADCC (see, e.g., Shields et al., J Biol Chem. 276:6591-

6604, 2001; and Presta et al., Biochem Soc Trans. 30:487-490, 2002). See also engineered Fc glycoforms that have increased binding to FcRs, as disclosed in Umana et al., supra; and U.S. Patent No. 7,662,925. Some embodiments include Fc regions that comprise one or more substitutions selected from 434S, 252Y/428L, 252Y/434S, and 428L/434S (see U.S. Application Nos. 2009/0163699 and 20060173170), based on the EU index of Kabat et al.

Certain variant, fragment, hybrid, or modified Fc regions may have altered effector functions, relative to a corresponding, wild-type Fc sequence. For example, such Fc regions may have increased complement fixation or activation, increased C1q binding affinity, increased CDC-related activity, increased ADCC-related activity, and/or increased ADCP-related activity, relative to a corresponding, wild-type Fc sequence. In other embodiments, such Fc regions may have decreased complement fixation or activation, decreased C1q binding affinity, decreased CDC-related activity, decreased ADCC-related activity, and/or decreased ADCP-related activity, relative to a corresponding, wild-type Fc sequence. As merely one illustrative example, an Fc region may comprise a deletion or substitution in a complement-binding site, such as a C1q-binding site, and/or a deletion or substitution in an ADCC site. Examples of such deletions/substitutions are described, for example, in U.S. Patent No. 7,030,226. Many Fc effector functions, such as ADCC, can be assayed according to routine techniques in the art. (see, e.g., Zuckerman et al., CRC Crit Rev Microbiol. 7:1-26, 1978). Useful effector cells for such assays includes, but are not limited to, natural killer (NK) cells, macrophages, and other peripheral blood mononuclear cells (PBMC). Alternatively, or additionally, certain Fc effector functions may be assessed *in vivo*, for example, by employing an animal model described in Clynes et al. PNAS. 95:652-656, 1998.

Certain variant hybrid, or modified Fc regions may have altered stability or half-life relative to a corresponding, wild-type Fc sequence. In certain embodiments, such Fc regions may have increased half-life relative to a corresponding, wild-type Fc sequence. In other embodiments, variant hybrid, or modified Fc regions may have decreased half-life relative to a corresponding, wild-type Fc sequence. Half-life can be measured *in vitro* (e.g., under physiological conditions) or *in vivo*, according to routine techniques in the art, such as radiolabeling, ELISA, or other methods. *In vivo* measurements of stability or half-life can be measured in one or more bodily fluids, including blood, serum, plasma, urine, or cerebrospinal fluid, or a given tissue, such as the liver, kidneys, muscle, central nervous system tissues, bone, etc. As one example, modifications to an Fc region that alter its ability to bind the FcRn can alter its half-life *in vivo*. Assays for measuring the *in vivo* pharmacokinetic properties (e.g., *in vivo* mean elimination half-life) and non-limiting examples of Fc modifications that alter its binding to the FcRn are described, for example, in U.S. Pat. Nos. 7,217,797 and 7,732,570; and U.S. Application Nos. US 2010/0143254 and 2010/0143254.

Additional non-limiting examples of modifications to alter stability or half-life include substitutions/deletions at one or more of amino acid residues selected from 251-256, 285-290, and 308-314 in the CH<sub>2</sub> domain, and 385-389 and 428-436 in the CH<sub>3</sub> domain, according to the

numbering system of Kabat *et al.* See U.S. Application No. 2003/0190311. Specific examples include substitution with leucine at position 251, substitution with tyrosine, tryptophan or phenylalanine at position 252, substitution with threonine or serine at position 254, substitution with arginine at position 255, substitution with glutamine, arginine, serine, threonine, or glutamate at position 256, substitution with threonine at position 308, substitution with proline at position 309, substitution with serine at position 311, substitution with aspartate at position 312, substitution with leucine at position 314, substitution with arginine, aspartate or serine at position 385, substitution with threonine or proline at position 386, substitution with arginine or proline at position 387, substitution with proline, asparagine or serine at position 389, substitution with methionine or threonine at position 428, substitution with tyrosine or phenylalanine at position 434, substitution with histidine, arginine, lysine or serine at position 433, and/or substitution with histidine, tyrosine, arginine or threonine at position 436, including any combination thereof. Such modifications optionally increase affinity of the Fc region for the FcRn and thereby increase half-life, relative to a corresponding, wild-type Fc region.

Certain variant hybrid, or modified Fc regions may have altered solubility relative to a corresponding, wild-type Fc sequence. In certain embodiments, such Fc regions may have increased solubility relative to a corresponding, wild-type Fc sequence. In other embodiments, variant hybrid, or modified Fc regions may have decreased solubility relative to a corresponding, wild-type Fc sequence. Solubility can be measured, for example, *in vitro* (e.g., under physiological conditions) according to routine techniques in the art. Exemplary solubility measurements are described elsewhere herein.

Variant Fc regions can also have one or more mutated hinge regions, as described, for example, in U.S. Application No. 2003/0118592. For instance, one or more cysteines in a hinge region can be deleted or substituted with a different amino acid. The mutated hinge region can comprise no cysteine residues, or it can comprise 1, 2, or 3 fewer cysteine residues than a corresponding, wild-type hinge region. In some embodiments, an Fc region having a mutated hinge region of this type exhibits a reduced ability to dimerize, relative to a wild-type Ig hinge region.

In particular embodiments, an antibody or antigen binding fragment thereof has a biological half life at about pH 7.4, at about a physiological pH, at about 25°C or room temperature, and/or at about 37°C or human body temperature (e.g., *in vivo*, in serum, in a given tissue, in a given species such as rat, mouse, monkey, or human), of about or at least about 30 minutes, about 1 hour, about 2 hours, about 3 hours, about 4 hours, about 5 hours, about 6 hours, about 12 hours, about 18 hours, about 20 hours, about 24 hours, about 30 hours, about 36 hours, about 40 hours, about 48 hours, about 50 hours, about 60 hours, about 70 hours, about 72 hours, about 80 hours, about 84 hours, about 90 hours, about 96 hours, about 120 hours, or about 144 hours or more, or about 1 week, or about 2 weeks, or about 3 weeks, or about 4 weeks, or about 5 weeks, or about 6 weeks or more, or any intervening half-life, including all ranges in between.

In some embodiments, an antibody or antigen binding fragment thereof has a  $T_m$  of about or at least about 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, or 75°C. In some embodiments, an antibody or antigen binding fragment thereof has a  $T_m$  of about 65°C or greater, for example, in PBS (phosphate buffered saline).

In some embodiments, an antibody or antigen binding fragment thereof conjugated to one or more cytotoxic or chemotherapeutic agents. General examples of cytotoxic or chemotherapeutic agents include, without limitation, alkylating agents, anti-metabolites, anthracyclines, anti-tumor antibiotics, platinum, type I topoisomerase inhibitors, type II topoisomerase inhibitors, vinca alkaloids, and taxanes. Specific examples of cytotoxic or chemotherapeutic agents include, without limitation, cyclophosphamide, cilengitide, lomustine (CCNU), melphalan, procarbazine, carmustine (BCNU), enzastaurin, busulfan, daunorubicin, doxorubicin, gefitinib, erlotinib, idarubicin, temozolomide, epirubicin, mitoxantrone, bleomycin, cisplatin, carboplatin, oxaliplatin, camptothecins, irinotecan, topotecan, amsacrine, etoposide, etoposide phosphate, teniposide, temsirolimus, everolimus, vincristine, vinblastine, vinorelbine, vindesine, CT52923, paclitaxel, imatinib, dasatinib, sorafenib, pazopanib, sunitinib, vatalanib, gefitinib, erlotinib, AEE-788, dichloroacetate, tamoxifen, fasudil, SB-681323, semaxanib, donepezil, galantamine, memantine, rivastigmine, tacrine, rasagiline, naltrexone, lubiprostone, safinamide, istradefylline, pimavanserin, pitolisant, isradipine, pridopidine (ACR16), tetrabenazine, bexarotene, glatirimer acetate, fingolimod, and mitoxantrone, including pharmaceutically acceptable salts and acids thereof. Further examples of cytotoxic or chemotherapeutic agents include alkylating agents such as thiotepa, cyclophosphamide (CYTOXAN<sup>TM</sup>); alkyl sulfonates such as busulfan, improsulfan and piposulfan; aziridines such as benzodopa, carboquone, meturedopa, and uredopa; ethylenimines and methylamelamines including altretamine, triethylenemelamine, triethylenephosphoramide, triethylenethiophosphoramide and trimethylolomelamine; nitrogen mustards such as chlorambucil, chlornaphazine, cholophosphamide, estramustine, ifosfamide, mechlorethamine, mechlorethamine oxide hydrochloride, melphalan, novembichin, phenesterine, prednimustine, trofosfamide, uracil mustard; nitrosureas such as carmustine, chlorozotocin, fotemustine, lomustine, nimustine, ranimustine; antibiotics such as aclacinomysins, actinomycin, anthramycin, azaserine, bleomycins, cactinomycin, calicheamicin, carabycin, carminomycin, carzinophilin, chromomycins, dactinomycin, daunorubicin, detorubicin, 6-diazo-5-oxo-L-norleucine, doxorubicin, epirubicin, esorubicin, idarubicin, marcellomycin, mitomycins, mycophenolic acid, nogalamycin, olivomycins, peplomycin, potfiromycin, puromycin, quelamycin, rodorubicin, streptonigrin, streptozocin, tubercidin, ubenimex, zinostatin, zorubicin; anti-metabolites such as methotrexate and 5-fluorouracil (5-FU); folic acid analogues such as denopterin, methotrexate, pteropterin, trimetrexate; purine analogs such as fludarabine, 6-mercaptopurine, thiamiprine, thioguanine; pyrimidine analogs such as ancitabine, azacitidine, 6-azauridine, carmofur, cytarabine, dideoxyuridine, doxifluridine, enocitabine, floxuridine, 5-FU; androgens such as calusterone, dromostanolone propionate, epitostanol, mepitostane, testolactone; anti-adrenals such as

aminoglutethimide, mitotane, trilostane; folic acid replenisher such as frolinic acid; aceglatone; aldophosphamide glycoside; aminolevulinic acid; amsacrine; bestrabucil; bisantrene; edatraxate; defofamine; demecolcine; diaziquone; elformithine; elliptinium acetate; etoglucid; gallium nitrate; hydroxyurea; lentinan; lonidamine; mitoguazone; mitoxantrone; mopidamol; nitracrine; pentostatin; phenamet; pirarubicin; podophyllinic acid; 2-ethylhydrazide; procarbazine; PSK; razoxane; sizofiran; spirogermanium; tenuazonic acid; triaziquone; 2,2',2''-trichlorotriethylamine; urethan; vindesine; dacarbazine; mannomustine; mitobronitol; mitolactol; pipobroman; gacytosine; arabinoside ("Ara-C"); cyclophosphamide; thiotepa; taxoids, e.g., paclitaxel (TAXOL®; Bristol-Myers Squibb Oncology, Princeton, N.J.) and doxetaxel (TAXOTERE®, Rhne-Poulenc Rorer, Antony, France); chlorambucil; gemcitabine; 6-thioguanine; mercaptopurine; methotrexate; platinum analogs such as cisplatin and carboplatin; vinblastine; platinum; etoposide (VP-16); ifosfamide; mitomycin C; mitoxantrone; vincristine; vinorelbine; navelbine; novantrone; teniposide; daunomycin; aminopterin; xeloda; ibandronate; CPT-11; topoisomerase inhibitor RFS 2000; difluoromethylomithine (DMFO); retinoic acid derivatives such as Targretin™ (bexarotene), Panretin™ (alitretinoin); ONTAK™ (denileukin diftitox); esperamicins; capecitabine; and pharmaceutically acceptable salts, acids or derivatives of any of the above.

In some embodiments, a herein-disclosed antibody is conjugated or operably linked to a radioisotope to form a radioconjugate and/or macrocyclic chelators useful for conjugating radiometal ions. A variety of radioactive isotopes are available for the production of radioconjugate antibodies. Examples include, but are not limited to <sup>90</sup>Y, <sup>123</sup>I, <sup>125</sup>I, <sup>131</sup>I, <sup>186</sup>Re, <sup>188</sup>Re, <sup>211</sup>At, and <sup>212</sup>Bi. In certain embodiments, the macrocyclic chelator is 1,4,7,10-tetraazacyclododecane-N,N',N'',N'''-tetraacetic acid (DOTA) which can be attached to the antibody via a linker molecule. Such linker molecules are commonly known in the art and described in Denardo *et al.*, 1998, Clin Cancer Res. 4:2483-90; Peterson *et al.*, 1999, Bioconjug. Chem. 10:553; and Zimmerman *et al.*, 1999, Nucl. Med. Biol. 26:943-50.

Other modifications of the antibodies (and polypeptides) of the disclosure are also contemplated herein. For example, in some embodiments the antibody is linked to one of a variety of nonproteinaceous polymers, e.g., polyethylene glycol, polypropylene glycol, polyoxyalkylenes, or copolymers of polyethylene glycol and polypropylene glycol. In some embodiments, the antibody is entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization (for example, hydroxymethylcellulose or gelatin-microcapsules and poly-(methylmethacrylate)microcapsules, respectively), in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles and nanocapsules), or in macroemulsions. Such techniques are disclosed in Remington's Pharmaceutical Sciences, 16th edition, Oslo, A., Ed., (1980).

The antibodies or antigen binding fragments thereof can be used in any of the compositions, methods, and/or kits described herein, and combined with one or more of the additional agents described herein.

### **Methods of Use and Pharmaceutical Compositions**

Certain embodiments relate to methods of treating, ameliorating the symptoms of, and/or reducing the progression of, a disease or condition in a subject in need thereof, comprising administering to the subject an antibody or antigen binding fragment thereof that binds to interleukin-11 receptor subunit  $\alpha$  (IL-11R $\alpha$ ), as described herein, or a pharmaceutical composition comprising the same. In some instances, the antibody or antigen binding fragment thereof antagonizes the binding/signaling activity between the IL-11R $\alpha$  and its ligand, IL-11. In some embodiments, the disease or condition is an IL-11-associated or IL-11-mediated disease or condition. In some embodiments, the disease or condition is a cancer, an inflammatory disease, an autoimmune disease, a wasting disease, a bone disease, or a fibrotic disease.

In some embodiments, as noted above, the disease or condition is a cancer or tumor. For example, in some instances, the cancer expresses or overexpresses IL-11R $\alpha$  and/or IL-11, and in some instances, the cancer displays IL-11R $\alpha$ /IL-11-dependent growth, adhesion, migration, invasion, and/or chemoresistance. In some instances, the cancer is a primary cancer. In some instances, the cancer is a metastatic cancer.

Exemplary cancers include, without limitation, bone cancer, prostate cancer, melanoma (e.g., metastatic melanoma), pancreatic cancer, small cell lung cancer, non-small cell lung cancer (NSCLC), mesothelioma, leukemia (e.g., lymphocytic leukemia, chronic myelogenous leukemia, acute myeloid leukemia, relapsed acute myeloid leukemia, hairy cell leukemias, acute lymphoblastic leukemias), lymphoma (e.g., non-Hodgkin's lymphomas, Hodgkin's lymphoma), hepatoma (hepatocellular carcinoma), sarcoma, B-cell malignancy, breast cancer, ovarian cancer, colorectal cancer, glioma, glioblastoma multiforme, meningioma, pituitary adenoma, vestibular schwannoma, primary CNS lymphoma, primitive neuroectodermal tumor (medulloblastoma), kidney cancer (e.g., renal cell carcinoma), bladder cancer, uterine cancer, esophageal cancer, brain cancer, head and neck cancers, cervical cancer, testicular cancer, thyroid cancer, and stomach cancer. In specific embodiments, the cancer is a metastatic cancer, for example, which has metastasized to the bone.

In some embodiments, the methods and compositions described herein increase median survival time of a subject by 4 weeks, 5 weeks, 6 weeks, 7 weeks, 8 weeks, 9 weeks, 10 weeks, 15 weeks, 20 weeks, 25 weeks, 30 weeks, 40 weeks, or longer. In certain embodiments, the methods and compositions described herein increase median survival time of a subject by 1 year, 2 years, 3 years, or longer. In some embodiments, the methods and compositions described herein increase progression-free survival by 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 7 weeks, 8 weeks, 9

weeks, 10 weeks or longer. In certain embodiments, the methods and compositions described herein increase progression-free survival by 1 year, 2 years, 3 years, or longer.

In certain embodiments, the methods and compositions described herein are sufficient to result in tumor regression, as indicated by a statistically significant decrease in the amount of viable tumor, for example, at least a 10%, 20%, 30%, 40%, 50% or greater decrease in tumor mass, or by altered (e.g., decreased with statistical significance) scan dimensions. In some embodiments, the methods and compositions described herein reduce the growth rate (e.g., *in vivo* or *in vitro*, including cancer cells isolated from a biopsy or other sample and grown *in vitro*) of the cancer by about or at least about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 2000% or more relative to an untreated control. In some instances, the methods and compositions described herein reduce cancer cell initiation, migration, adhesion, invasiveness, and/or metastasis by about or at least about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 2000% or more relative to an untreated control. In some instances, the methods and compositions described herein reduce angiogenesis in the tumor environment by about or at least about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 2000% or more relative to an untreated control.

In certain embodiments, the disease or condition is an inflammatory disease. Non-limiting examples of inflammatory diseases and conditions include airway or lung inflammation (e.g., inflammatory lung disease), asthma, rhinitis, chronic obstructive pulmonary disorder (COPD), dermatitis, psoriasis, hepatitis, gastric inflammation, irritable bowel syndrome (IBS), ulcerative colitis, Crohn's disease, colitis, diverticulitis, lupus erythematosus, nephritis, Parkinson's disease, multiple sclerosis (MS), Alzheimer's disease, arthritis, rheumatoid arthritis, sepsis, infection-induced inflammation, cardiovascular diseases such as atherosclerosis and vasculitis, diabetes, and gout.

In certain embodiments, the disease or condition is an autoimmune disease. Non-limiting examples of autoimmune diseases and conditions include arthritis (including rheumatoid arthritis, reactive arthritis), systemic lupus erythematosus (SLE), psoriasis, inflammatory bowel disease (IBD), ulcerative colitis, Crohn's disease, encephalomyelitis, uveitis, myasthenia gravis, multiple sclerosis, insulin dependent diabetes, Addison's disease, celiac disease, chronic fatigue syndrome, autoimmune hepatitis, autoimmune alopecia, ankylosing spondylitis, fibromyalgia, pemphigus vulgaris, Sjogren's syndrome, Kawasaki's Disease, hyperthyroidism/Graves' disease, hypothyroidism/Hashimoto's disease, endometriosis, scleroderma, pernicious anemia, Goodpasture syndrome, Guillain-Barré syndrome, Wegener's disease, glomerulonephritis, aplastic anemia (including multiply transfused aplastic anemia patients), paroxysmal nocturnal hemoglobinuria, myelodysplastic syndrome, idiopathic thrombocytopenic purpura, autoimmune hemolytic anemia, Evan's syndrome, Factor VIII inhibitor syndrome, systemic vasculitis, dermatomyositis, polymyositis, rheumatic fever, autoimmune lymphoproliferative syndrome (ALPS), autoimmune bullous pemphigoid, Parkinson's disease, sarcoidosis, vitiligo, primary biliary cirrhosis, and autoimmune myocarditis.

In certain embodiments, the disease or condition is a wasting disease. Non-limiting examples of wasting diseases and conditions include cachexia, including cachexia associated with cancer or renal failure, and sarcopenia. In certain embodiments, the disease or condition is a bone disease. Non-limiting examples of bone diseases and conditions include osteoporosis (including post-menopausal osteoporosis), bone fracture, Paget's disease of bone, and bone resorption/damage associated with cancer or cancer therapy, including chemotherapy, hormone ablation, and hormone inhibition.

In some embodiments, the disease or condition is fibrosis, or a fibrotic disease. Examples include fibrosis of the lungs, cardiovascular system, liver, brain, joints (e.g., knee, hip, ankle, foot joints, shoulder, elbow, wrist, hand joints, spinal vertebrae), intestine, skin, kidney, liver, thyroid, bone marrow, retroperitoneum, eye (see, for example, Schafer et al., *Nature*. 552: 110-115, 2017; and Ng et al., *Sci Transl Med*. 2019 Sep 25;11(511)).

In some embodiments, the fibrosis of the lungs is selected from fibrothorax, pulmonary fibrosis (for example, cystic fibrosis, interstitial lung disease (ILD), autosomal recessive genetic disease such as Hermansky-Pudlak syndrome type 1, 2, 3, 4, 5, 6, 7, or 8), and radiation-induced lung injury. In some embodiments, the pulmonary fibrosis is related to ILD, for example, idiopathic or secondary ILD. Examples of idiopathic ILD include idiopathic pulmonary fibrosis (IPF), desquamative interstitial pneumonia (DIP), acute interstitial pneumonia (AIP) also known as Hamman-Rich syndrome, nonspecific interstitial pneumonia (NSIP), respiratory bronchiolitis-associated interstitial lung disease (RB-ILD), cryptogenic organizing pneumonia (COP), and lymphoid interstitial pneumonia (LIP). General examples of secondary ILD include ILD related to connective tissue and autoimmune diseases (for example, sarcoidosis, rheumatoid arthritis, systemic lupus erythematosus, systemic sclerosis, polymyositis, dermatomyositis, antisynthetase syndrome), inhaled substances (for example, silicosis, asbestosis, berylliosis, industrial printing chemicals, chronic hypersensitivity pneumonitis), drugs (drug-induced ILD, for example, antibiotics, chemotherapeutics, anti-arrhythmic agents), infections (for example, SARS CoV-2, atypical pneumonia, pneumocystis pneumonia, tuberculosis, Chlamydia trachomatis, respiratory syncytial virus), malignancies (lymphangitic carcinomatosis), and pediatric ILDs such as diffuse developmental disorders, growth abnormalities deficient alveolarization, infant conditions of undefined cause, and ILD related to alveolar surfactant region.

In particular embodiments, the fibrosis of the cardiovascular system is myocardial fibrosis (for example, interstitial fibrosis, replacement fibrosis).

In some embodiments, the antibodies described herein are selective for both mouse and human IL-11R $\alpha$ , block IL-11 signaling through both the STAT3 and ERK pathways, and abrogate both cis- and trans- IL-11 signaling. Following TGF $\beta$  stimulation, certain anti-huIL-11R $\alpha$  antibodies described herein have been shown to reduce collagen expression in primary IPF patient fibroblasts and decrease expression of procollagen I and TIMP by PCLS from healthy donors. Similar reductions in procollagen release have been observed in response to anti-huIL-11R $\alpha$  in preliminary PCLS studies

from IPF patients. In mouse bleomycin lung fibrosis studies, blockade of IL-11R $\alpha$  signaling reduces lung fibrosis and decreases BAL inflammatory cells when administered prophylactically or therapeutically after the onset of lung fibrosis.

In certain embodiments, the methods and compositions described herein are sufficient to result in stable disease. In certain embodiments, the methods and compositions described herein are sufficient to result in clinically relevant reduction in symptoms of a particular disease indication known to the skilled clinician.

For in vivo use, certain embodiments include pharmaceutical compositions, comprising an antibody or antigen binding fragment thereof, as described herein, and a pharmaceutically-acceptable carrier. To prepare a therapeutic or pharmaceutical composition, an effective or desired amount of one or more agents is mixed with any pharmaceutical carrier(s) or excipient known to those skilled in the art to be suitable for the particular agent and/or mode of administration. A pharmaceutical carrier may be liquid, semi-liquid or solid. Solutions or suspensions used for parenteral, intradermal, intraocular, subcutaneous, direct instillation into the bladder, or topical application may include, for example, a sterile diluent (such as water), saline solution (*e.g.*, phosphate buffered saline; PBS), fixed oil, polyethylene glycol, glycerin, propylene glycol or other synthetic solvent; antimicrobial agents (such as benzyl alcohol and methyl parabens); antioxidants (such as ascorbic acid and sodium bisulfite) and chelating agents (such as ethylenediaminetetraacetic acid (EDTA)); buffers (such as acetates, citrates and phosphates). If administered intravenously (*e.g.*, by IV infusion), suitable carriers include physiological saline or phosphate buffered saline (PBS), and solutions containing thickening and solubilizing agents, such as glucose, polyethylene glycol, polypropylene glycol and mixtures thereof.

Administration of agents described herein, in pure form or in an appropriate therapeutic or pharmaceutical composition, can be carried out via any of the accepted modes of administration of agents for serving similar utilities. The therapeutic or pharmaceutical compositions can be prepared by combining an agent-containing composition with an appropriate physiologically acceptable carrier, diluent or excipient, and may be formulated into preparations in solid, semi-solid, liquid or gaseous forms, such as tablets, capsules, powders, granules, ointments, solutions, suppositories, injections, inhalants, gels, microspheres, and aerosols. In addition, other pharmaceutically active ingredients (including other small molecules as described elsewhere herein) and/or suitable excipients such as salts, buffers and stabilizers may, but need not, be present within the composition.

Administration may be achieved by a variety of different routes, including oral, parenteral, nasal, intravenous, intraocular, intradermal, intramuscular, subcutaneous, installation into the bladder, or topical. Preferred modes of administration depend upon the nature of the condition to be treated or prevented. Particular embodiments include administration by IV infusion.

Carriers can include, for example, pharmaceutically- or physiologically-acceptable carriers, excipients, or stabilizers that are non-toxic to the cell or mammal being exposed thereto at the dosages and concentrations employed. Often the physiologically-acceptable carrier is an aqueous pH buffered

solution. Examples of physiologically acceptable carriers include buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid; low molecular weight (less than about 10 residues) polypeptide; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, arginine, histidine, and/or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; salt-forming counterions such as sodium; and/or nonionic surfactants such as polysorbate 20 (TWEEN™) polyethylene glycol (PEG), and poloxamers (PLURONICS™), and the like.

In some embodiments, one or more agents can be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization (for example, hydroxymethylcellulose or gelatin-microcapsules and poly-(methylmethacrylate)microcapsules, respectively), in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles and nanocapsules), or in macroemulsions. Such techniques are disclosed in Remington's Pharmaceutical Sciences, 16th edition, Oslo, A., Ed., (1980). The particle(s) or liposomes may further comprise other therapeutic or diagnostic agents.

The precise dosage and duration of treatment is a function of the disease being treated and may be determined empirically using known testing protocols or by testing the compositions in model systems known in the art and extrapolating therefrom. Controlled clinical trials may also be performed. Dosages may also vary with the severity of the condition to be alleviated. A pharmaceutical composition is generally formulated and administered to exert a therapeutically useful effect while minimizing undesirable side effects. The composition may be administered one time, or may be divided into a number of smaller doses to be administered at intervals of time. For any particular subject, specific dosage regimens may be adjusted over time according to the individual need.

Typical routes of administering these and related therapeutic or pharmaceutical compositions thus include, without limitation, oral, topical, transdermal, inhalation, parenteral, sublingual, buccal, ocular, rectal, vaginal, and intranasal. The term parenteral as used herein includes subcutaneous injections, intravenous, instillation into the bladder, intramuscular, intrasternal injection or infusion techniques. Therapeutic or pharmaceutical compositions according to certain embodiments of the present disclosure are formulated so as to allow the active ingredients contained therein to be bioavailable upon administration of the composition to a subject or patient. Compositions that will be administered to a subject or patient may take the form of one or more dosage units, where for example, a tablet may be a single dosage unit, and a container of a herein described agent in aerosol form may hold a plurality of dosage units. Actual methods of preparing such dosage forms are known, or will be apparent, to those skilled in this art; for example, see *Remington: The Science and Practice of Pharmacy*, 20th Edition (Philadelphia College of Pharmacy and Science, 2000). The composition

to be administered will typically contain a therapeutically effective amount of an agent described herein, for treatment of a disease or condition of interest.

A therapeutic or pharmaceutical composition can be in the form of a solid or liquid. In some embodiments, the carrier(s) are particulate, so that the compositions are, for example, in tablet or powder form. The carrier(s) can be liquid, with the compositions being, for example, an oral oil, injectable liquid or an aerosol, which is useful in, for example, inhalatory administration. When intended for oral administration, the pharmaceutical composition is preferably in either solid or liquid form, where semi-solid, semi-liquid, suspension and gel forms are included within the forms considered herein as either solid or liquid. Certain embodiments include sterile, injectable solutions.

As a solid composition for oral administration, the pharmaceutical composition may be formulated into a powder, granule, gel, compressed tablet, pill, capsule, chewing gum, wafer or the like. Such a solid composition will typically contain one or more inert diluents or edible carriers. In addition, one or more of the following may be present: binders such as carboxymethylcellulose, ethyl cellulose, microcrystalline cellulose, gum tragacanth or gelatin; excipients such as starch, lactose or dextrins, disintegrating agents such as alginic acid, sodium alginate, Primogel, corn starch and the like; lubricants such as magnesium stearate or Sterotex; glidants such as colloidal silicon dioxide; sweetening agents such as sucrose or saccharin; a flavoring agent such as peppermint, methyl salicylate or orange flavoring; and a coloring agent. When the pharmaceutical composition is in the form of a capsule, for example, a gelatin capsule, it may contain, in addition to materials of the above type, a liquid carrier such as polyethylene glycol or oil.

The therapeutic or pharmaceutical composition may be in the form of a liquid, for example, an elixir, syrup, solution, gel, emulsion or suspension. The liquid may be for oral administration or for delivery by injection, as two examples. When intended for oral administration, preferred composition contain, in addition to the present compounds, one or more of a sweetening agent, preservatives, dye/colorant and flavor enhancer. In a composition intended to be administered by injection, one or more of a surfactant, preservative, wetting agent, dispersing agent, suspending agent, buffer, stabilizer and isotonic agent may be included.

The liquid therapeutic or pharmaceutical compositions, whether they be solutions, suspensions or other like form, may include one or more of the following adjuvants: sterile diluents such as water for injection, saline solution, preferably physiological saline, Ringer's solution, isotonic sodium chloride, fixed oils such as synthetic mono or diglycerides which may serve as the solvent or suspending medium, polyethylene glycols, glycerin, propylene glycol or other solvents; antibacterial agents such as benzyl alcohol or methyl paraben; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made

of glass or plastic. Physiological saline is a preferred adjuvant. An injectable pharmaceutical composition is preferably sterile.

A liquid therapeutic or pharmaceutical composition intended for either parenteral, intraocular, or oral administration should contain an amount of an agent such that a suitable dosage will be obtained. Typically, this amount is at least 0.01% of the agent of interest in the composition. When intended for oral administration, this amount may be varied to be between 0.1 and about 70% of the weight of the composition. Certain oral therapeutic or pharmaceutical compositions contain between about 4% and about 75% of the agent of interest. In certain embodiments, therapeutic or pharmaceutical compositions and preparations are prepared so that a parenteral dosage unit contains between 0.01 to 10% by weight of the agent of interest prior to dilution.

The therapeutic or pharmaceutical compositions may be intended for topical administration, in which case the carrier may suitably comprise a solution, emulsion, ointment or gel base. The base, for example, may comprise one or more of the following: petrolatum, lanolin, polyethylene glycols, bee wax, mineral oil, diluents such as water and alcohol, and emulsifiers and stabilizers. Thickening agents may be present in a therapeutic or pharmaceutical composition for topical administration. If intended for transdermal administration, the composition may include a transdermal patch or iontophoresis device.

The therapeutic or pharmaceutical compositions may be intended for rectal administration, in the form, for example, of a suppository, which will melt in the rectum and release the drug. The composition for rectal administration may contain an oleaginous base as a suitable nonirritating excipient. Such bases include, without limitation, lanolin, cocoa butter, and polyethylene glycol.

The therapeutic or pharmaceutical composition may include various materials, which modify the physical form of a solid or liquid dosage unit. For example, the composition may include materials that form a coating shell around the active ingredients. The materials that form the coating shell are typically inert, and may be selected from, for example, sugar, shellac, and other enteric coating agents. Alternatively, the active ingredients may be encased in a gelatin capsule. The therapeutic or pharmaceutical compositions in solid or liquid form may include a component that binds to agent and thereby assists in the delivery of the compound. Suitable components that may act in this capacity include monoclonal or polyclonal antibodies, one or more proteins or a liposome.

The therapeutic or pharmaceutical composition may consist essentially of dosage units that can be administered as an aerosol. The term aerosol is used to denote a variety of systems ranging from those of colloidal nature to systems consisting of pressurized packages. Delivery may be by a liquefied or compressed gas or by a suitable pump system that dispenses the active ingredients. Aerosols may be delivered in single phase, bi-phasic, or tri-phasic systems in order to deliver the active ingredient(s). Delivery of the aerosol includes the necessary container, activators, valves, subcontainers, and the like, which together may form a kit. One of ordinary skill in the art, without undue experimentation may determine preferred aerosols.

The compositions described herein may be prepared with carriers that protect the agents against rapid elimination from the body, such as time release formulations or coatings. Such carriers include controlled release formulations, such as, but not limited to, implants and microencapsulated delivery systems, and biodegradable, biocompatible polymers, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, polyorthoesters, polylactic acid and others known to those of ordinary skill in the art.

The pharmaceutical compositions may be prepared by methodology well known in the pharmaceutical art. For example, a therapeutic or pharmaceutical composition intended to be administered by injection may comprise one or more of salts, buffers and/or stabilizers, with sterile, distilled water so as to form a solution. A surfactant may be added to facilitate the formation of a homogeneous solution or suspension. Surfactants are compounds that non-covalently interact with the agent so as to facilitate dissolution or homogeneous suspension of the agent in the aqueous delivery system.

The therapeutic or pharmaceutical compositions may be administered in a therapeutically effective amount, which will vary depending upon a variety of factors including the activity of the specific compound employed; the metabolic stability and length of action of the compound; the age, body weight, general health, sex, and diet of the subject; the mode and time of administration; the rate of excretion; the drug combination; the severity of the particular disorder or condition; and the subject undergoing therapy. In some instances, a therapeutically effective daily dose is (for a 70 kg mammal) from about 0.001 mg/kg (i.e., ~ 0.07 mg) to about 100 mg/kg (i.e., ~ 7.0 g); preferably a therapeutically effective dose is (for a 70 kg mammal) from about 0.01 mg/kg (i.e., ~ 0.7 mg) to about 50 mg/kg (i.e., ~ 3.5 g); more preferably a therapeutically effective dose is (for a 70 kg mammal) from about 1 mg/kg (i.e., ~ 70 mg) to about 25 mg/kg (i.e., ~ 1.75 g). In some embodiments, the therapeutically effective dose is administered on a weekly, bi-weekly, or monthly basis. In specific embodiments, the therapeutically effective dose is administered on a weekly, bi-weekly, or monthly basis, for example, at a dose of about 1-10 or 1-5 mg/kg, or about 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 mg/kg.

Also included are patient care kits, comprising (a) an antibody or antigen binding fragment thereof that binds to IL-11R $\alpha$ , as described herein; and optionally (b) at least one additional therapeutic agent. In certain kits, (a) and (b) are in separate therapeutic compositions. In some kits, (a) and (b) are in the same therapeutic composition.

The kits herein may also include a one or more additional therapeutic agents or other components suitable or desired for the indication being treated, or for the desired diagnostic application. The kits herein can also include one or more syringes or other components necessary or desired to facilitate an intended mode of delivery (e.g., stents, implantable depots, etc.).

In some embodiments, a patient care kit contains separate containers, dividers, or compartments for the composition(s) and informational material(s). For example, the composition(s) can be contained in a bottle, vial, or syringe, and the informational material(s) can be contained in

association with the container. In some embodiments, the separate elements of the kit are contained within a single, undivided container. For example, the composition is contained in a bottle, vial or syringe that has attached thereto the informational material in the form of a label. In some embodiments, the kit includes a plurality (e.g., a pack) of individual containers, each containing one or more unit dosage forms (e.g., a dosage form described herein) of an antibody and optionally at least one additional therapeutic agent. For example, the kit includes a plurality of syringes, ampules, foil packets, or blister packs, each containing a single unit dose of an antibody and optionally at least one additional therapeutic agent. The containers of the kits can be air tight, waterproof (e.g., impermeable to changes in moisture or evaporation), and/or light-tight.

The patient care kit optionally includes a device suitable for administration of the composition, e.g., a syringe, inhalant, dropper (e.g., eye dropper), swab (e.g., a cotton swab or wooden swab), or any such delivery device. In some embodiments, the device is an implantable device that dispenses metered doses of the agent(s). Also included are methods of providing a kit, e.g., by combining the components described herein.

### Expression and Purification Systems

Certain embodiments include methods and related compositions for expressing and purifying an anti-IL-11R $\alpha$  antibody or antigen binding fragment thereof described herein. Such recombinant anti-IL-11R $\alpha$  antibodies can be conveniently prepared using standard protocols as described for example in Sambrook, *et al.*, (1989, *supra*), in particular Sections 16 and 17; Ausubel *et al.*, (1994, *supra*), in particular Chapters 10 and 16; and Coligan *et al.*, Current Protocols in Protein Science (John Wiley & Sons, Inc. 1995-1997), in particular Chapters 1, 5 and 6. As one general example, anti-IL-11R $\alpha$  antibodies may be prepared by a procedure including one or more of the steps of: (a) preparing a construct that comprises a polynucleotide sequence which encodes an anti-IL-11R $\alpha$  antibody heavy chain and/or light chains, and which is operably linked to a regulatory element; (b) introducing the construct into a host cell; (c) culturing the host cell to express the anti-IL-11R $\alpha$  antibody; and (d) isolating the anti-IL-11R $\alpha$  from the host cell.

Certain embodiments thus include polynucleotides that encode an anti-IL-11R $\alpha$  antibody or antigen binding fragment thereof described herein, including vectors comprising said polynucleotides, and host cells comprising the polynucleotides and/or vectors. In order to express a desired polypeptide, a nucleotide sequence encoding an anti-IL-11R $\alpha$ , or a functional equivalent, may be inserted into appropriate expression vector, *i.e.*, a vector which contains the necessary elements for the transcription and translation of the inserted coding sequence. Methods which are well known to those skilled in the art may be used to construct expression vectors containing sequences encoding a polypeptide of interest and appropriate transcriptional and translational control elements. These methods include *in vitro* recombinant DNA techniques, synthetic techniques, and *in vivo* genetic

recombination. Such techniques are described in Sambrook *et al.*, Molecular Cloning, A Laboratory Manual (1989), and Ausubel *et al.*, Current Protocols in Molecular Biology (1989).

A variety of expression vector/host systems are known and may be utilized to contain and express polynucleotide sequences. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with virus expression vectors (*e.g.*, baculovirus); plant cell systems transformed with virus expression vectors (*e.g.*, cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or with bacterial expression vectors (*e.g.*, Ti or pBR322 plasmids); or animal cell systems, including mammalian cell and more specifically human cell systems.

The “control elements” or “regulatory sequences” present in an expression vector are those non-translated regions of the vector--enhancers, promoters, 5' and 3' untranslated regions--which interact with host cellular proteins to carry out transcription and translation. Such elements may vary in their strength and specificity. Depending on the vector system and host utilized, any number of suitable transcription and translation elements, including constitutive and inducible promoters, may be used. For example, when cloning in bacterial systems, inducible promoters such as the hybrid lacZ promoter of the PBLUESCRIPT phagemid (Stratagene, La Jolla, Calif.) or PSORT1 plasmid (Gibco BRL, Gaithersburg, Md.) and the like may be used. In mammalian cell systems, promoters from mammalian genes or from mammalian viruses are generally preferred. If it is necessary to generate a cell line that contains multiple copies of the sequence encoding a polypeptide, vectors based on SV40 or EBV may be advantageously used with an appropriate selectable marker.

In bacterial systems, a number of expression vectors may be selected depending upon the use intended for the expressed polypeptide. For example, when large quantities are needed, vectors which direct high level expression of fusion proteins that are readily purified may be used. Such vectors include, but are not limited to, the multifunctional *E. coli* cloning and expression vectors such as BLUESCRIPT (Stratagene), in which the sequence encoding the polypeptide of interest may be ligated into the vector in frame with sequences for the amino-terminal Met and the subsequent 7 residues of  $\beta$ -galactosidase so that a hybrid protein is produced; pIN vectors (Van Heeke & Schuster, *J. Biol. Chem.* 264:5503-5509 (1989)); and the like. pGEX Vectors (Promega, Madison, Wis.) may also be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione-agarose beads followed by elution in the presence of free glutathione. Proteins made in such systems may be designed to include heparin, thrombin, or factor XA protease cleavage sites so that the cloned polypeptide of interest can be released from the GST moiety at will.

Certain embodiments may employ *E. coli*-based expression systems (*see, e.g.*, Structural Genomics Consortium *et al.*, *Nature Methods*. 5:135-146, 2008). These and related embodiments may rely partially or totally on ligation-independent cloning (LIC) to produce a suitable expression vector.

In specific embodiments, protein expression may be controlled by a T7 RNA polymerase (*e.g.*, pET vector series). These and related embodiments may utilize the expression host strain BL21(DE3), a  $\lambda$ DE3 lysogen of BL21 that supports T7-mediated expression and is deficient in lon and ompT proteases for improved target protein stability. Also included are expression host strains carrying plasmids encoding tRNAs rarely used in *E. coli*, such as ROSETTA™ (DE3) and Rosetta 2 (DE3) strains. Cell lysis and sample handling may also be improved using reagents sold under the trademarks BENZONASE® nuclease and BUGBUSTER® Protein Extraction Reagent. For cell culture, auto-inducing media can improve the efficiency of many expression systems, including high-throughput expression systems. Media of this type (*e.g.*, OVERNIGHT EXPRESS™ Autoinduction System) gradually elicit protein expression through metabolic shift without the addition of artificial inducing agents such as IPTG. Particular embodiments employ hexahistidine tags (such as those sold under the trademark HIS•TAG® fusions), followed by immobilized metal affinity chromatography (IMAC) purification, or related techniques. In certain aspects, however, clinical grade proteins can be isolated from *E. coli* inclusion bodies, without or without the use of affinity tags (*see, e.g.*, Shimp *et al.*, *Protein Expr Purif.* 50:58-67, 2006). As a further example, certain embodiments may employ a cold-shock induced *E. coli* high-yield production system, because over-expression of proteins in *Escherichia coli* at low temperature improves their solubility and stability (*see, e.g.*, Qing *et al.*, *Nature Biotechnology.* 22:877-882, 2004).

Also included are high-density bacterial fermentation systems. For example, high cell density cultivation of *Ralstonia eutropha* allows protein production at cell densities of over 150 g/L, and the expression of recombinant proteins at titers exceeding 10 g/L.

In the yeast *Saccharomyces cerevisiae*, a number of vectors containing constitutive or inducible promoters such as alpha factor, alcohol oxidase, and PGH may be used. For reviews, see Ausubel *et al.* (*supra*) and Grant *et al.*, *Methods Enzymol.* 153:516-544 (1987). Also included are *Pichia pandoris* expression systems (*see, e.g.*, Li *et al.*, *Nature Biotechnology.* 24, 210 – 215, 2006; and Hamilton *et al.*, *Science*, 301:1244, 2003). Certain embodiments include yeast systems that are engineered to selectively glycosylate proteins, including yeast that have humanized N-glycosylation pathways, among others (*see, e.g.*, Hamilton *et al.*, *Science.* 313:1441-1443, 2006; Wildt *et al.*, *Nature Reviews Microbiol.* 3:119-28, 2005; and Gerngross *et al.*, *Nature-Biotechnology.* 22:1409 -1414, 2004; U.S. Patent Nos. 7,629,163; 7,326,681; and 7,029,872). Merely by way of example, recombinant yeast cultures can be grown in Fernbach Flasks or 15L, 50L, 100L, and 200L fermentors, among others.

In cases where plant expression vectors are used, the expression of sequences encoding polypeptides may be driven by any of a number of promoters. For example, viral promoters such as the 35S and 19S promoters of CaMV may be used alone or in combination with the omega leader sequence from TMV (Takamatsu, *EMBO J.* 6:307-311 (1987)). Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters may be used (Coruzzi *et al.*, *EMBO J.*

3:1671-1680 (1984); Broglie et al., *Science* 224:838-843 (1984); and Winter et al., *Results Probl. Cell Differ.* 17:85-105 (1991)). These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. Such techniques are described in a number of generally available reviews (see, e.g., Hobbs in McGraw Hill, *Yearbook of Science and Technology*, pp. 191-196 (1992)).

An insect system may also be used to express a polypeptide of interest. For example, in one such system, *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes in *Spodoptera frugiperda* cells or in *Trichoplusia* cells. The sequences encoding the polypeptide may be cloned into a non-essential region of the virus, such as the polyhedrin gene, and placed under control of the polyhedrin promoter. Successful insertion of the polypeptide-encoding sequence will render the polyhedrin gene inactive and produce recombinant virus lacking coat protein. The recombinant viruses may then be used to infect, for example, *S. frugiperda* cells or *Trichoplusia* cells in which the polypeptide of interest may be expressed (Engelhard et al., *Proc. Natl. Acad. Sci. U.S.A.* 91:3224-3227 (1994)). Also included are baculovirus expression systems, including those that utilize SF9, SF21, and T. ni cells (see, e.g., Murphy and Piwnica-Worms, *Curr Protoc Protein Sci.* Chapter 5:Unit5.4, 2001). Insect systems can provide post-translation modifications that are similar to mammalian systems.

In mammalian host cells, a number of viral-based expression systems are generally available. For example, in cases where an adenovirus is used as an expression vector, sequences encoding a polypeptide of interest may be ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain a viable virus which is capable of expressing the polypeptide in infected host cells (Logan & Shenk, *Proc. Natl. Acad. Sci. U.S.A.* 81:3655-3659 (1984)). In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells.

Examples of useful mammalian host cell lines include monkey kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651); human embryonic kidney line (293 or 293 cells sub-cloned for growth in suspension culture, Graham et al., *J. Gen Virol.* 36:59 (1977)); baby hamster kidney cells (BHK, ATCC CCL 10); mouse sertoli cells (TM4, Mather, *Biol. Reprod.* 23:243-251 (1980)); monkey kidney cells (CV1 ATCC CCL 70); African green monkey kidney cells (VERO-76, ATCC CRL-1587); human cervical carcinoma cells (HELA, ATCC CCL 2); canine kidney cells (MDCK, ATCC CCL 34); buffalo rat liver cells (BRL 3A, ATCC CRL 1442); human lung cells (W138, ATCC CCL 75); human liver cells (Hep G2, HB 8065); mouse mammary tumor (MMT 060562, ATCC CCL51); TR1 cells (Mather et al., *Annals N.Y. Acad. Sci.* 383:44-68 (1982)); MRC 5 cells; FS4 cells; and a human hepatoma line (Hep G2). Other useful mammalian host cell lines include Chinese hamster ovary (CHO) cells, including DHFR-CHO cells (Urlaub et al., *PNAS USA* 77:4216 (1980)); and myeloma cell lines such as NSO and Sp2/0. For a review of certain mammalian host cell lines

suitable for antibody production, see, e.g., Yazaki and Wu, *Methods in Molecular Biology*, Vol. 248 (B. K.C Lo, ed., Humana Press, Totowa, N.J., 2003), pp. 255-268. Certain preferred mammalian cell expression systems include CHO and HEK293-cell based expression systems. Mammalian expression systems can utilize attached cell lines, for example, in T-flasks, roller bottles, or cell factories, or suspension cultures, for example, in 1L and 5L spinners, 5L, 14L, 40L, 100L and 200L stir tank bioreactors, or 20/50L and 100/200L WAVE bioreactors, among others known in the art.

Also included is the cell-free expression of proteins. These and related embodiments typically utilize purified RNA polymerase, ribosomes, tRNA and ribonucleotides; these reagents may be produced by extraction from cells or from a cell-based expression system.

Specific initiation signals may also be used to achieve more efficient translation of sequences encoding a polypeptide of interest. Such signals include the ATG initiation codon and adjacent sequences. In cases where sequences encoding the polypeptide, its initiation codon, and upstream sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a portion thereof, is inserted, exogenous translational control signals including the ATG initiation codon should be provided. Furthermore, the initiation codon should be in the correct reading frame to ensure translation of the entire insert. Exogenous translational elements and initiation codons may be of various origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers which are appropriate for the particular cell system which is used, such as those described in the literature (Scharf. et al., *Results Probl. Cell Differ.* 20:125-162 (1994)).

In addition, a host cell strain may be chosen for its ability to modulate the expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, post-translational modifications such as acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" form of the protein may also be used to facilitate correct insertion, folding and/or function. Different host cells such as yeast, CHO, HeLa, MDCK, HEK293, and W138, in addition to bacterial cells, which have or even lack specific cellular machinery and characteristic mechanisms for such post-translational activities, may be chosen to ensure the correct modification and processing of the foreign protein.

For long-term, high-yield production of recombinant proteins, stable expression is generally preferred. For example, cell lines which stably express a polynucleotide of interest may be transformed using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells may be allowed to grow for about 1-2 days in an enriched media before they are switched to selective media. The purpose of the selectable marker is to confer resistance to selection, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resistant clones of stably transformed cells may be proliferated

using tissue culture techniques appropriate to the cell type. Transient production, such as by transient transfection or infection, can also be employed. Exemplary mammalian expression systems that are suitable for transient production include HEK293 and CHO-based systems.

Any number of selection systems may be used to recover transformed or transduced cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase (Wigler et al., *Cell* 11:223-232 (1977)) and adenine phosphoribosyltransferase (Lowy et al., *Cell* 22:817-823 (1990)) genes which can be employed in tk- or aprt- cells, respectively. Also, antimetabolite, antibiotic or herbicide resistance can be used as the basis for selection; for example, dhfr which confers resistance to methotrexate (Wigler et al., *PNAS USA* 77:3567-70 (1980)); npt, which confers resistance to the aminoglycosides, neomycin and G-418 (Colbere-Garapin et al., *J. Mol. Biol.* 150:1-14 (1981)); and als or pat, which confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively (Murry, *supra*). Additional selectable genes have been described, for example, trpB, which allows cells to utilize indole in place of tryptophan, or hisD, which allows cells to utilize histinol in place of histidine (Hartman & Mulligan, *Proc. Natl. Acad. Sci. U.S.A.* 85:8047-51 (1988)). The use of visible markers has gained popularity with such markers as green fluorescent protein (GFP) and other fluorescent proteins (e.g., RFP, YFP), anthocyanins,  $\beta$ -glucuronidase and its substrate GUS, and luciferase and its substrate luciferin, being widely used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system (see, e.g., Rhodes et al., *Methods Mol. Biol.* 55:121-131 (1995)).

Also included are high-throughput protein production systems, or micro-production systems. Certain aspects may utilize, for example, hexa-histidine fusion tags for protein expression and purification on metal chelate-modified slide surfaces or MagneHis Ni-Particles (see, e.g., Kwon et al., *BMC Biotechnol.* 9:72, 2009; and Lin et al., *Methods Mol Biol.* 498:129-41, 2009)). Also included are high-throughput cell-free protein expression systems (see, e.g., Sitaraman et al., *Methods Mol Biol.* 498:229-44, 2009). These and related embodiments can be used, for example, to generate microarrays of antibodies which can then be used for screening libraries to identify antibodies and antigen binding domains that interact with the IL-11R $\alpha$  polypeptide(s) of interest.

A variety of protocols for detecting and measuring the expression of polynucleotide-encoded products, using binding agents or antibodies such as polyclonal or monoclonal antibodies specific for the product, are known in the art. Examples include enzyme-linked immunosorbent assay (ELISA), western immunoblots, radioimmunoassays (RIA), and fluorescence activated cell sorting (FACS). These and other assays are described, among other places, in Hampton et al., *Serological Methods, a Laboratory Manual* (1990) and Maddox et al., *J. Exp. Med.* 158:1211-1216 (1983).

A wide variety of labels and conjugation techniques are known by those skilled in the art and may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides include oligolabeling, nick translation, end-labeling or PCR amplification using a labeled nucleotide. Alternatively, the

sequences, or any portions thereof may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by addition of an appropriate RNA polymerase such as T7, T3, or SP6 and labeled nucleotides. These procedures may be conducted using a variety of commercially available kits. Suitable reporter molecules or labels, which may be used include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Host cells transformed with a polynucleotide sequence of interest may be cultured under conditions suitable for the expression and recovery of the protein from cell culture. Certain specific embodiments utilize serum free cell expression systems. Examples include HEK293 cells and CHO cells that can grow on serum free medium (see, e.g., Rosser et al., *Protein Expr. Purif.* 40:237–43, 2005; and U.S. Patent number 6,210,922).

An antibody, or antigen binding fragment thereof, produced by a recombinant cell may be secreted or contained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides may be designed to contain signal sequences which direct secretion of the encoded polypeptide through a prokaryotic or eukaryotic cell membrane. Other recombinant constructions may be used to join sequences encoding a polypeptide of interest to nucleotide sequence encoding a polypeptide domain which will facilitate purification and/or detection of soluble proteins. Examples of such domains include cleavable and non-cleavable affinity purification and epitope tags such as avidin, FLAG tags, poly-histidine tags (e.g., 6xHis), cMyc tags, V5-tags, glutathione S-transferase (GST) tags, and others.

The protein produced by a recombinant cell can be purified and characterized according to a variety of techniques known in the art. Exemplary systems for performing protein purification and analyzing protein purity include fast protein liquid chromatography (FPLC) (e.g., AKTA and Bio-Rad FPLC systems), high-pressure liquid chromatography (HPLC) (e.g., Beckman and Waters HPLC). Exemplary chemistries for purification include ion exchange chromatography (e.g., Q, S), size exclusion chromatography, salt gradients, affinity purification (e.g., Ni, Co, FLAG, maltose, glutathione, protein A/G), gel filtration, reverse-phase, ceramic HYPERD® ion exchange chromatography, and hydrophobic interaction columns (HIC), among others known in the art. Also included are analytical methods such as SDS-PAGE (e.g., Coomassie, silver stain), immunoblot, Bradford, and ELISA, which may be utilized during any step of the production or purification process, typically to measure the purity of the protein composition.

Also included are methods of concentrating anti-IL-11R $\alpha$  antibodies and antigen binding fragments thereof, and composition comprising concentrated soluble proteins. In certain aspects, concentrated solutions of anti-IL-11R $\alpha$  antibodies comprise proteins at a concentration of about 5 mg/mL; or about 8 mg/mL; or about 10 mg/mL; about 15 mg/mL; or about 20 mg/mL or more.

In some aspects, the compositions are substantially monodisperse, for example, where the anti-IL-11R $\alpha$  antibody exists primarily (i.e., at least about 90%, or greater) in one apparent molecular weight form when assessed, for example, by size exclusion chromatography, dynamic light scattering, and/or analytical ultracentrifugation.

In some aspects, the compositions have a purity (on a protein basis) of at least about 90%, or in some aspects at least about 95% purity, or in some embodiments, at least about 98% purity. Purity can be determined via any routine analytical method as known in the art.

In some aspects, the compositions have a high molecular weight aggregate content of less than about 10%, less than about 5%, less than about 3%, less than about 1%. High molecular weight aggregate content can be determined by a variety of analytical techniques including for example, by size exclusion chromatography, dynamic light scattering, and/or analytical ultracentrifugation.

Examples of concentration approaches contemplated herein include lyophilization, which is typically employed when the solution contains few soluble components other than the protein of interest. Lyophilization is often performed after HPLC run, and can remove most or all volatile components from the mixture. Also included are ultrafiltration techniques, which typically employ one or more selective permeable membranes to concentrate a protein solution. The membrane allows water and small molecules to pass through and retains the protein; the solution can be forced against the membrane by mechanical pump, gas pressure, or centrifugation, among other techniques.

In certain embodiments, the anti-IL-11R $\alpha$  antibodies, reagents, or related agents have a purity of at least about 90%, as measured according to routine techniques in the art. In certain embodiments, an anti-IL-11R $\alpha$  composition has a purity of at least about 95%. In specific embodiments, such as therapeutic or pharmaceutical compositions, an anti-IL-11R $\alpha$  antibody composition has a purity of at least about 97% or 98% or 99%. In some embodiments, such as when being used as reference or research reagents, anti-IL-11R $\alpha$  antibodies can be of lesser purity, and may have a purity of at least about 50%, 60%, 70%, or 80%. Purity can be measured overall or in relation to selected components, such as other proteins, *e.g.*, purity on a protein basis.

Purified antibodies can also be characterized according to their biological characteristics. Binding affinity and binding kinetics can be measured according to a variety of techniques known in the art, such as Biacore® and related technologies that utilize surface plasmon resonance (SPR), an optical phenomenon that enables detection of unlabeled interactants in real time. SPR-based biosensors can be used in determination of active concentration, screening and characterization in terms of both affinity and kinetics. The presence or levels of one or more canonical or non-canonical biological activities can be measured according to cell-based assays, including those that utilize a cellular binding partner of a selected anti-IL-11R $\alpha$  antibody, which is functionally coupled to a readout or indicator, such as a fluorescent or luminescent indicator of biological activity, as described herein.

In certain embodiments, as noted above, a composition is substantially endotoxin free, including, for example, about or at least about 95% endotoxin free, about or at least about 99% endotoxin free, or about or at least about 99.99% endotoxin free. The presence of endotoxins can be detected according to routine techniques in the art, as described herein. In specific embodiments, a composition is made from a eukaryotic cell such as a mammalian or human cell in substantially serum free media. In certain embodiments, as noted herein, a composition has an endotoxin content of less than about 10 EU/mg of antibody, or less than about 5 EU/mg of antibody, less than about 3 EU/mg of antibody, or less than about 1 EU/mg of antibody.

In certain embodiments, a composition comprises less than about 10% wt/wt high molecular weight aggregates, or less than about 5% wt/wt high molecular weight aggregates, or less than about 2% wt/wt high molecular weight aggregates, or less than about or less than about 1% wt/wt high molecular weight aggregates.

Also included are protein-based analytical assays and methods, which can be used to assess, for example, protein purity, size, solubility, and degree of aggregation, among other characteristics. Protein purity can be assessed a number of ways. For instance, purity can be assessed based on primary structure, higher order structure, size, charge, hydrophobicity, and glycosylation. Examples of methods for assessing primary structure include N- and C-terminal sequencing and peptide-mapping (see, e.g., Allen et al., *Biologicals*. 24:255-275, 1996)). Examples of methods for assessing higher order structure include circular dichroism (see, e.g., Kelly et al., *Biochim Biophys Acta*. 1751:119-139, 2005), fluorescent spectroscopy (see, e.g., Meagher et al., *J. Biol. Chem.* 273:23283-89, 1998), FT-IR, amide hydrogen-deuterium exchange kinetics, differential scanning calorimetry, NMR spectroscopy, immunoreactivity with conformationally sensitive antibodies. Higher order structure can also be assessed as a function of a variety of parameters such as pH, temperature, or added salts. Examples of methods for assessing protein characteristics such as size include analytical ultracentrifugation and size exclusion HPLC (SEC-HPLC), and exemplary methods for measuring charge include ion-exchange chromatography and isoelectric focusing. Hydrophobicity can be assessed, for example, by reverse-phase HPLC and hydrophobic interaction chromatography HPLC. Glycosylation can affect pharmacokinetics (e.g., clearance), conformation or stability, receptor binding, and protein function, and can be assessed, for example, by mass spectrometry and nuclear magnetic resonance (NMR) spectroscopy.

As noted above, certain embodiments include the use of SEC-HPLC to assess protein characteristics such as purity, size (e.g., size homogeneity) or degree of aggregation, and/or to purify proteins, among other uses. SEC, also including gel-filtration chromatography (GFC) and gel-permeation chromatography (GPC), refers to a chromatographic method in which molecules in solution are separated in a porous material based on their size, or more specifically their hydrodynamic volume, diffusion coefficient, and/or surface properties. The process is generally used to separate biological molecules, and to determine molecular weights and molecular weight

distributions of polymers. Typically, a biological or protein sample (such as a protein extract produced according to the protein expression methods provided herein and known in the art) is loaded into a selected size-exclusion column with a defined stationary phase (the porous material), preferably a phase that does not interact with the proteins in the sample. In certain aspects, the stationary phase is composed of inert particles packed into a dense three-dimensional matrix within a glass or steel column. The mobile phase can be pure water, an aqueous buffer, an organic solvent, or a mixture thereof. The stationary-phase particles typically have small pores and/or channels which only allow molecules below a certain size to enter. Large particles are therefore excluded from these pores and channels, and their limited interaction with the stationary phase leads them to elute as a “totally-excluded” peak at the beginning of the experiment. Smaller molecules, which can fit into the pores, are removed from the flowing mobile phase, and the time they spend immobilized in the stationary-phase pores depends, in part, on how far into the pores they penetrate. Their removal from the mobile phase flow causes them to take longer to elute from the column and results in a separation between the particles based on differences in their size. A given size exclusion column has a range of molecular weights that can be separated. Overall, molecules larger than the upper limit will not be trapped by the stationary phase, molecules smaller than the lower limit will completely enter the solid phase and elute as a single band, and molecules within the range will elute at different rates, defined by their properties such as hydrodynamic volume. For examples of these methods in practice with pharmaceutical proteins, see Bruner et al., *Journal of Pharmaceutical and Biomedical Analysis*. 15: 1929-1935, 1997.

Protein purity for clinical applications is also discussed, for example, by Anicetti et al. (*Trends in Biotechnology*. 7:342-349, 1989). More recent techniques for analyzing protein purity include, without limitation, the LabChip GXII, an automated platform for rapid analysis of proteins and nucleic acids, which provides high throughput analysis of titer, sizing, and purity analysis of proteins. In certain non-limiting embodiments, clinical grade proteins such as protein fragments and antibodies can be obtained by utilizing a combination of chromatographic materials in at least two orthogonal steps, among other methods (see, e.g., *Therapeutic Proteins: Methods and Protocols*. Vol. 308, Eds., Smales and James, Humana Press Inc., 2005). Typically, protein agents (e.g., antibodies and antigen binding fragments) are substantially endotoxin-free, as measured according to techniques known in the art and described herein.

Protein solubility assays are also included. Such assays can be utilized, for example, to determine optimal growth and purification conditions for recombinant production, to optimize the choice of buffer(s), and to optimize the choice of antibodies or antigen binding fragments thereof. Solubility or aggregation can be evaluated according to a variety of parameters, including temperature, pH, salts, and the presence or absence of other additives. Examples of solubility screening assays include, without limitation, microplate-based methods of measuring protein solubility using turbidity or other measure as an end point, high-throughput assays for analysis of the

solubility of purified recombinant proteins (see, e.g., Stenvall et al., *Biochim Biophys Acta*. 1752:6-10, 2005), assays that use structural complementation of a genetic marker protein to monitor and measure protein folding and solubility in vivo (see, e.g., Wigley et al., *Nature Biotechnology*. 19:131-136, 2001), and electrochemical screening of recombinant protein solubility in *Escherichia coli* using scanning electrochemical microscopy (SECM) (see, e.g., Nagamine et al., *Biotechnology and Bioengineering*. 96:1008-1013, 2006), among others. Antibodies with increased solubility (or reduced aggregation) can be identified or selected for according to routine techniques in the art, including simple in vivo assays for protein solubility (see, e.g., Maxwell et al., *Protein Sci.* 8:1908-11, 1999).

Protein solubility and aggregation can also be measured by dynamic light scattering techniques. Aggregation is a general term that encompasses several types of interactions or characteristics, including soluble/insoluble, covalent/noncovalent, reversible/irreversible, and native/denatured interactions and characteristics. For protein therapeutics, the presence of aggregates is typically considered undesirable because of the concern that aggregates may cause an immunogenic reaction (e.g., small aggregates), or may cause adverse events on administration (e.g., particulates). Dynamic light scattering refers to a technique that can be used to determine the size distribution profile of small particles in suspension or polymers such as proteins in solution. This technique, also referred to as photon correlation spectroscopy (PCS) or quasi-elastic light scattering (QELS), uses scattered light to measure the rate of diffusion of the protein particles. Fluctuations of the scattering intensity can be observed due to the Brownian motion of the molecules and particles in solution. This motion data can be conventionally processed to derive a size distribution for the sample, wherein the size is given by the Stokes radius or hydrodynamic radius of the protein particle. The hydrodynamic size depends on both mass and shape (conformation). Dynamic scattering can detect the presence of very small amounts of aggregated protein (<0.01% by weight), even in samples that contain a large range of masses. It can also be used to compare the stability of different formulations, including, for example, applications that rely on real-time monitoring of changes at elevated temperatures. Accordingly, certain embodiments include the use of dynamic light scattering to analyze the solubility and/or presence of aggregates in a sample that contains an antibody of the present disclosure.

Although the foregoing embodiments have been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be readily apparent to one of ordinary skill in the art in light of the teachings of this disclosure that certain changes and modifications may be made thereto without departing from the spirit or scope of the appended claims. The following examples are provided by way of illustration only and not by way of limitation. Those of skill in the art will readily recognize a variety of noncritical parameters that could be changed or modified to yield essentially similar results.

## Examples

### Example 1

#### Anti-IL-11R $\alpha$ Antibodies

Studies were performed to generate improved anti-IL-11R $\alpha$  antibodies, relative, for example, to the antibodies described in U.S. Patent Nos. 9,796,782; and 9,340,618 – including the antibodies designated as 8E2 and TS7. For example, a potential N-linked glycosylation in V<sub>L</sub>CDR3 of the TS7 antibody was eliminated to reduce heterogeneity and improve developability, and other sequence changes were made to improve binding affinity and potency. The CDR sequences of the antibodies are provided in **Table A1** (supra). The antibodies were expressed in HEK293 cells and purified using protein A based chromatography using standard methodology. All antibodies were expressed in human IgG4 (S228P by EU numbering; S241P by Kabat numbering) format with kappa light chains.

Relative binding activity of the antibodies was tested by competitive ELISA using biotinylated mAbs 8E2 or TS7. Antibodies 8E2 and TS7 were biotinylated using sulfo-NHS Biotin (Thermo Fisher) per the manufacturer's instructions. IL-11R was coated onto ELISA plates (Nunc, MaxiSorp) at 2  $\mu$ g/mL in PBS, and plates were blocked with 2% BSA in PBS. After washing the plate with water, antibodies to be tested were titrated across the plate in a final volume of 90  $\mu$ L followed by addition of 10  $\mu$ L of biotinylated 8E2 or biotinylated TS7, as indicated in **Table E1** (0.1  $\mu$ g/mL final concentration) to each well. Plates were incubated for 2 hours on a plate-shaker, followed by washing and addition of 100  $\mu$ L/well of streptavidin-HRP (Jackson ImmunoResearch, 1/1000 dilution). After a further 1 hour incubation, plates were washed, developed with Ultra-TMB HRP substrate (Thermo Fisher), and stopped by the addition of 50  $\mu$ L/well of 2 M sulfuric acid. Plates were read at 450 nm on an iD5 plate reader (Molecular Devices).

As shown in **Table E1** below, many of the tested antibodies were shown to have increased binding to human IL-11R $\alpha$  relative to that of the 8E2 antibody and the TS7 antibody.

| <b>Table E1. Competitive ELISA for binding strength relative to 8E2 and TS7</b> |                                |                                |
|---------------------------------------------------------------------------------|--------------------------------|--------------------------------|
| Competitor                                                                      | 8E2-bio                        | TS7-bio                        |
|                                                                                 | IC <sub>50</sub> ( $\mu$ g/mL) | IC <sub>50</sub> ( $\mu$ g/mL) |
| 8E2                                                                             | 22                             |                                |
| TS7                                                                             | 3.1                            | 6.0                            |
| mAb1                                                                            | >10                            |                                |
| mAb2                                                                            | >30                            |                                |
| mAb3                                                                            | 2.8                            |                                |
| mAb4                                                                            | 5.5                            |                                |
| mAb5                                                                            | 1.4                            | 2.1                            |
| mAb6                                                                            | 1.5                            |                                |
| mAb7                                                                            |                                | 6.9                            |
| mAb8                                                                            |                                | 6.2                            |
| mAb9                                                                            |                                | 2.3                            |
| mAb10                                                                           |                                | >15                            |
| mAb11                                                                           |                                | >15                            |

|       |  |     |
|-------|--|-----|
| mAb12 |  | 3.6 |
| mAb13 |  | 2.3 |
| mAb14 |  | 2.0 |
| mAb15 |  | 1.7 |

Kinetic binding analysis of mAbs was tested to determine antigen binding affinity using biolayer interferometry on an Octet RED96e instrument. mAbs were loaded onto protein G biosensors (FortéBio) in 10x kinetics buffer consisting of PBS, 0.1% BSA, 0.02% Tween 20 for 120 seconds to achieve a spectral shift value of 0.8 to 1.2 nm. Association was carried out in the presence of a 2-fold dilution series of hIL-11R $\alpha$  and was allowed to proceed for 120 seconds; dissociation was measured for 300 to 1200 seconds. Dilution series started at 100 nM for weaker variants or 10 nM for the most potent mAbs.

As shown in **Table E2** below, mAb5, mAb6, mAb7, mAb8, mAb9, mAb12, mAb13, mAb14, and mAb15 had very high binding affinity, beyond the ability of the instrument to determine an accurate off-rate (kd).

| <b>Table E2. Kinetic binding analysis of mAbs to IL-11R using biolayer interferometry</b> |                  |                 |                          |
|-------------------------------------------------------------------------------------------|------------------|-----------------|--------------------------|
| Antibody                                                                                  | <b>ka (1/Ms)</b> | <b>kd (1/s)</b> | <b>K<sub>D</sub> (M)</b> |
| mAb5                                                                                      | 6.37E+04         | <1.0E-07        | <1.0E-12                 |
| mAb6                                                                                      | 8.00E+05         | <1.0E-07        | <1.0E-12                 |
| mab7                                                                                      | 6.35E+04         | <1.0E-07        | <1.0E-12                 |
| mab8                                                                                      | 6.51E+04         | <1.0E-07        | <1.0E-12                 |
| mab9                                                                                      | 6.22E+04         | <1.0E-07        | <1.0E-12                 |
| mab10                                                                                     | 8.40E+05         | 4.16E-04        | 4.96E-10                 |
| mab11                                                                                     | 3.29E+04         | 1.81E-02        | 5.50E-07                 |
| mab12                                                                                     | 5.81E+04         | <1.0E-07        | <1.0E-12                 |
| mab13                                                                                     | 6.67E+04         | <1.0E-07        | <1.0E-12                 |
| mab14                                                                                     | 3.37E+04         | <1.0E-07        | <1.0E-12                 |
| mab15                                                                                     | 8.40E+04         | <1.0E-07        | <1.0E-12                 |

Further testing of the lead mAbs was carried out using a cell-based reporter assay. IL-11 is reported to signal through the STAT3 and ERK pathways. Therefore, a STAT3 reporter cell line in which firefly luciferase gene expression is driven by STAT3 response elements located upstream of a minimal TATA promoter was selected to analyze mAb functional activity. In these cells, IL-11 can activate the endogenous STAT3 allowing it to bind to the STAT3 response elements, inducing transcription of the luciferase reporter gene and resulting in readily detectable luciferase expression. Antibodies to IL-11R which block IL-11 signaling should inhibit expression of IL-11 driven luciferase in these cells.

The STAT3 Reporter (Luciferase) - HEK293 cell line (BPS Bioscience, catalog #79800-P) was grown, passaged, and assayed as per manufacturer's protocols. Cells were added at 25,000 cells

per well in a 96-well microtiter dish and incubated at 37 °C for 30-45 min. in 5% CO<sub>2</sub> humidified air. Anti-IL-11R mAbs were added as a 3-fold dilution series in duplicate columns or rows, and plates were returned to the incubator for an additional 1 h at 37 °C after which 40 ng/mL IL-11 was added to each well to initiate the signal transduction cascade and luciferase production. Plates were incubated for 18-24 hours at 37°C and luciferase was detected using a One-Step Luciferase Assay System (BPS Bioscience, cat. no. 60690-1) as per manufacturer's instructions.

As shown in **Table E3**, TS7 is significantly more potent than 8E2, and mAb5, mAb9, and mAb13 are more potent at inhibiting IL-11 signaling than either of TS7 or 8E2. In agreement with the ELISA data, mAb7, mAb8, mAb10, and mAb11 were less potent than TS7, and mAb12, mAb14, and mAb15 were comparable in potency to TS7. The results for mAb6 were variable in this assay, but the ELISA and Octet data above suggest mAb6 is also a potent mAb.

| <b>Table E3. Cell-based reporter assay measuring relative potency for inhibiting IL-11</b> |                |                |             |
|--------------------------------------------------------------------------------------------|----------------|----------------|-------------|
| <b>IC<sub>50</sub> (μg/mL)</b>                                                             |                |                |             |
| <b>Antibody</b>                                                                            | <b>Assay 1</b> | <b>Assay 2</b> | <b>Mean</b> |
| 8E2                                                                                        | >50            | 22.8           | >36.4       |
| TS7                                                                                        | 0.15           | 0.42           | 0.28        |
| mAb5                                                                                       | 0.16           | 0.18           | 0.17        |
| mAb6                                                                                       | 11.0           | 0.39           | 5.70        |
| mab7                                                                                       | 1.00           |                | 1.00        |
| mab8                                                                                       | 0.67           | 1.25           | 0.96        |
| mab9                                                                                       | 0.11           | 0.10           | 0.10        |
| mab10                                                                                      | 2.00           |                | 2.00        |
| mab11                                                                                      | >100           |                | >100        |
| mab12                                                                                      | 0.42           | 0.39           | 0.41        |
| mab13                                                                                      | 0.05           | 0.19           | 0.12        |
| mab14                                                                                      | 0.39           | 0.28           | 0.34        |
| mab15                                                                                      | 0.38           | 0.33           | 0.36        |

A further series of antibodies were designed in which the residue at Kabat position 31 of the heavy chain was varied (mAbs 16-32). In this case an alternate biolayer interferometry assay was used to initially rank the antibodies compared to mAb5, the closest relative in sequence to mAbs 16-32. MAbs were loaded onto anti-human constant domain (AHC) biosensors (FortéBio) in 10x kinetics buffer consisting of PBS, 0.1% BSA, 0.02% Tween 20 for 120 s to achieve a spectral shift value of 0.8 to 1.2 nm. Association was carried out in the presence of 2-fold dilution series of hIL-11R and was allowed to proceed for 120 s. Dilution series started 10 nM. The dissociation in 10x kinetics buffer for 1200 s. Binding kinetic measurements were taken on an Octet RED96e instrument.

As shown in **Table E4**, initial qualitative assessment identified additional antibodies with high potency either similar to or better than mAb5. More detailed kinetic measurements were taken for mAb21, mAb23, mAb29, and mAb31 compared to mAb5, which confirmed further improved

potency for these mAbs. Further testing of these mAbs by ELISA confirmed the improved potency of the selected mAbs.

| <b>Table E4</b> |                               |                           |                             |                            |
|-----------------|-------------------------------|---------------------------|-----------------------------|----------------------------|
| <b>mAb</b>      | <b>qualitative assessment</b> | <b>K<sub>D</sub> (pM)</b> | <b>k<sub>a</sub> (1/Ms)</b> | <b>k<sub>d</sub> (1/s)</b> |
| 5               | comparator                    | 50 pM                     | 4.90E+05                    | 2.50E-05                   |
| 16              | similar                       | ND                        | ND                          | ND                         |
| 17              | decreased                     | ND                        | ND                          | ND                         |
| 18              | decreased                     | ND                        | ND                          | ND                         |
| 19              | similar                       | ND                        | ND                          | ND                         |
| 20              | similar                       | ND                        | ND                          | ND                         |
| 21              | improved/<br>increased        | <30 pM                    | 7.50E+05                    | off-scale                  |
| 22              | similar                       | ND                        | ND                          | ND                         |
| 23              | improved/<br>increased        | <30 pM                    | 7.20E+05                    | off-scale                  |
| 24              | similar                       | ND                        | ND                          | ND                         |
| 25              | similar                       | ND                        | ND                          | ND                         |
| 26              | decreased                     | ND                        | ND                          | ND                         |
| 27              | decreased                     | ND                        | ND                          | ND                         |
| 28              | similar                       | ND                        | ND                          | ND                         |
| 29              | improved/<br>increased        | 30 pM                     | 6.10E+05                    | 1.80E-05                   |
| 30              | decreased                     | ND                        | ND                          | ND                         |
| 31              | improved/<br>increased        | <30 pM                    | 6.90E+05                    | off-scale                  |
| 32              | similar                       | ND                        | ND                          | ND                         |

All publications, patent applications, and issued patents cited in this specification are herein incorporated by reference as if each individual publication, patent application, or issued patent were specifically and individually indicated to be incorporated by reference.

## Example 2

### Activity of Anti-IL-11R $\alpha$ Antibodies in a Fibrotic Human Precision-Cut Lung Slice Model

Studies were performed to evaluate the effect of the mAb5 antibody on biomarkers reflecting ECM remodeling in a fibrotic human precision-cut lung slice (PCLS) model. The anti-fibrotic effect was investigated at three different concentrations in human ex vivo lung tissue slices prepared from pulmonary fibrosis patients.

#### Materials and Methods

Tissue slices were prepared from the subpleural and central region of the fibrotic lung from two human end-stage donors with pulmonary fibrosis (IPF and secondary fibrosis) (see, for example, Hess et al., Toxicol In Vitro. 32:347-61, 2016). For both donors, lung tissue samples from two

different regional areas (subpleural and central) were used to generate tissue cores used for PCLS. Adjacent cores were used to achieve serial PCLS for the analysis of one treatment as compared to medium treated PCLS from the same cores. This enables the comparison of densely packed fibrotic areas with fibrotic foci with less affected areas. To ensure the fibrotic status, clinical/demographic information for each patient is provided.

Experimental treatments included are listed in **Table E5**. Each treatment was tested with three replicates for each lung region and for each donor.

| <b>Table E5</b>                       |           |                        |
|---------------------------------------|-----------|------------------------|
| Treatment                             | Timepoint | Samples                |
| mAb5 IL11RA Ab 0.3 µg/mL              | 48 hours  | Supernatant and Tissue |
| mAb5 IL11RA Ab 3.0 µg/mL              | 48 hours  | Supernatant and Tissue |
| mAb5 IL11RA Ab 30 µg/mL               | 48 hours  | Supernatant and Tissue |
| Isotype control 30 µg/mL              | 48 hours  | Supernatant and Tissue |
| Nintedanib 1 µM                       | 48 hours  | Supernatant and Tissue |
| Medium control (PCLS w/o treatment)   | 48 hours  | Supernatant and Tissue |
| DMEM/F12 (Medium background w/o PCLS) | 48 hours  | Supernatant and Tissue |

**Preparation and incubation of precision-cut lung slices.** Lung lobes were canuled and filled with 37 °C warm 2% low-melting agarose/medium solution. Filled lobes were cooled on ice to polymerise agarose. Tissue cores with a diameter of 8 mm were prepared and cut into approximately 300 µm thick slices using a microtome (Krumdieck tissue slicer, Alabama Research and Development, Muniford, AL, USA or Vibratome OTS-5000, Science Services GmbH, Munich) in Earle's Balanced Salts Solution (EBSS). Tissue slices were incubated in Dulbecco's modified Eagle's medium/nutrient mixture F-12 Ham (DMEM) with L-glutamine and 15 mM HEPES without phenol red. Tissue slices were transferred into petri dishes and washed with DMEM for 2 hours. Medium was changed four times every 30 minutes in order to remove cell debris.

Two PCLS per well were left over night in 500 µL DMEM in 24-well plates for a final washing step. Next day, PCLS were treated with experimental treatments or controls (see **Table E5**) in 250µl DMEM/F-12 culture medium for 48 hours under normal cell culture conditions (37°C, 5% CO<sub>2</sub>). Medium contained 100 units/mL penicillin and 100 µg/mL streptomycin and was not supplemented with fetal calf serum.

After incubation for 48 hours, supernatants were collected, proteinase inhibitor cocktail (0.2%) was added and samples were frozen to -80°C. Additionally, the two PCLS from each well were transferred to a 2 mL safe-lock tube, snap-frozen in liquid nitrogen immediately, and stored at -80°C. Samples from the three replicates for each condition were not pooled.

The viability and responsiveness of the tissue to immune stimulators was confirmed in a quality control sample where PCLS was incubated with the mitogen 100 ng/mL lipopolysaccharide (LPS). Following 48 hours incubation, assessment of viability by lactate dehydrogenase (LDH) activity and pro-inflammatory response by IL-1β was performed.

***Tissue viability.*** The viability of the lung tissue was assessed to ensure sufficient quality of the PCLS. For quality control samples, LDH activity was determined with a commercial enzymatic assay in supernatants after incubation. An increase of the amount of dead or plasma membrane-damaged cells in PCLS will result in an increase of LDH enzyme activity in the culture supernatant. Triton X-100 (1 % in PBS) treated PCLS were investigated as reference control. 50  $\mu$ L samples were incubated with 50  $\mu$ L LDH reagent for 20 minutes at room temperature. Absorbance was measured at 490 nm and 630 nm as reference wavelength using a microplate reader. All tests were performed as single measurements per well. Due to the dense tissue material of fibrotic PCLS, absorbance values measured for the Triton X-100 control were at maximum. Therefore, 1:10 dilutions of these controls were tested in parallel to achieve appropriate values within the range of the detection spectra for calculation of the relative LDH release of the samples.

***Tissue responsiveness.*** The general responsiveness of the lung tissue to treatment was assessed to ensure optimal condition of the PCLS. For quality control samples, PCLS were lysed with Triton X-100 (1% in PBS) for one hour, whereafter intrinsic IL-1 $\beta$  levels were assessed using ELISA from R&D according to the manufacturer's instructions. The photometric measurement was performed at 450 nm and a reference wavelength of 570 nm using a microplate reader. The IL-1 $\beta$  concentration was normalized to total protein concentration of PCLS determined by BCA assay.

***Tissue appearance.*** Macroscopic appearance of the tissue as well as location of cores used to generate PCLS were recorded by photography. Tissue cores adjacent to the ones used for PCLS preparation were fixed in 10% formalin and subsequently embedded in paraffin. Paraffin-embedded tissue sections were further processed into slices for staining with hematoxylin and eosin (H&E) to assess fibrotic remodeling.

***Extracellular matrix remodeling evaluation.*** Biomarkers that reflect ECM remodeling were assessed in supernatant by Nordic Bioscience using the following competitive ELISA: PRO-C6 (C-terminal of type VI collagen C5 domain released as type VI collagen is incorporated into the matrix; measured on mesoscale platform [hsPRO-C6]): Reflects type VI collagen formation. Type VI collagen is a beaded filament collagen mainly found in the interface between the interstitial matrix and the basement membrane.

***Statistics.*** Biomarker data are presented as scatter plots with indication of median and interquartile range (IQR). Data are presented as raw biomarker values for each region and donor. Pooled data are presented as percentage of the “no treatment” control, calculated for individual regions and donors. Statistical significant differences were evaluated on pooled data by Kruskal-Wallis with Dunn’s multiple comparisons test, comparing to the “no treatment” control. Adjusted p-values are shown as: \* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$ , \*\*\*\* $p < 0.0001$ .

## Results

**Fibrotic Status.** Human PCLS were generated from lung tissue from two donors with pulmonary fibrosis. The pathological report provided after the completion of the study stated that the fibrotic tissue represents a clear UIP pattern and would be compatible with IPF (see **Table E6**).

| <b>Table E6</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <p><b>Donor 1</b><br/>62 years old male<br/>Bilateral lung explant with:</p> <ul style="list-style-type: none"> <li>- advanced interstitial fibrosis with architectural disturbance, bronchiolisation, myogenic metaplasia and multifocal fibroblastic foci with emphasis on the pulmonary sublobes</li> <li>- mild chronic and focal (lower lobe left) catarrhal bronchitis and bronchiolitis</li> <li>- still low-grade pulmonary artery sclerosis (see commentary).</li> </ul> <p>Comment:<br/>The present findings fulfil the morphological criteria of a UIP pattern (definite UIP according to ATS / ERS) and are - especially with clear accentuation of the pulmonary basement - suitable for an idiopathic pulmonary fibrosis (IPF). No evidence of malignancy.</p> | <p><b>Donor 2</b><br/>56 years old male<br/>Bilateral lung explant with:</p> <ul style="list-style-type: none"> <li>- advanced interstitial fibrosis with numerous fibroblastic foci, architectural disorder of the lung parenchyma, bronchial metaplasia and myogenic metaplasia corresponding to a UIP pattern,</li> <li>- chronic and florid catarrhal bronchitis/olitis with signs of secretion retention and multifocal transition to florid bronchopneumonia,</li> <li>- mild pulmonary artery sclerosis,</li> <li>- dust granulomas in the hilar lymph nodes.</li> </ul> <p>Comment:<br/>The findings are well compatible with idiopathic pulmonary fibrosis (IPF) in a far advanced stage. No evidence of malignancy.</p> |

**Tissue Appearance.** The PCLS were prepared from different regions of the fibrotic lung. **Figures 1A-1B** shows images of the lungs of each donor and the regions used, where drawn circles marks the areas of tissue that were used to generate PCLS. Adjacent tissue cores from the same lung region as that used for PCLS were subsequently prepared for H&E staining to assess fibrotic remodeling of the tissue (see **Figures 2A-2B**). Donor 2 appears to have more extensive tissue changes than donor 1 which is in accordance with the statement in the pathological report that donor 2 tissue is compatible with IPF in a far advanced stage.

**Tissue Viability.** Tissue viability of quality control samples (LPS) was measured by LDH release after 48 hours incubation. LDH release <20% of the triton X-lysed control was observed for all samples, indicating that the tissue was viable over the experimental period of 48 hours (see **Figures 3A-3B**).

**Tissue Responsiveness.** Tissue responsiveness of quality control sample (LPS) was analysed by IL-1 $\beta$  expression. Analysis showed that human fibrotic lung tissue from both donors responded to the pro-inflammatory stimuli LPS, identified by the upregulated expression of IL-1 $\beta$  after 48 hours (see **Figures 4A-4B**).

**Extracellular Matrix Remodeling.** Type VI collagen formation was evaluated by PRO-C6. Medium background levels of PRO-C6 were low and at the level of lower limit of detection for the

assay. Donor 1 showed lower levels of PRO-C6 as compared to donor 2, with levels quite close to the medium background.

As shown in **Figure 5**, the pooled PRO-C6 data for all donors and regions showed that median levels of PRO-C6 for nintedanib and all concentrations of mAb5 (anti-IL11R) were decreased as compared to the no treatment (media) control. mAb5 reduced PRO-C6 in a dose-dependent manner. The isotype control did not show a change in median PRO-C6 levels as compared to the no treatment (media) control, and even tended to increase slightly.

Based on the effects of nintedanib, the lung slices were responsive to anti-fibrotic treatment as ECM remodeling could be reduced. Overall, tendencies for effects on ECM remodeling were observed with the mAb5 anti-IL-11R $\alpha$  antibody. Especially interesting were the dose-related reductions of type VI collagen formation (PRO-C6) by the mAb5 antibody. In several cases, the mAb5 antibody showed better effects on ECM remodeling relative to the approved therapy nintedanib, strengthening the conclusion that this antibody has anti-fibrotic effects.

## Claims

1. An isolated antibody, or an antigen binding fragment thereof, which binds to interleukin-11 receptor subunit  $\alpha$  (IL-11R $\alpha$ ), wherein the at least one antibody, or antigen binding fragment thereof, comprises:

a heavy chain variable region (V<sub>H</sub>) that comprises complementary determining region V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences selected from **Table A1** and variants thereof which specifically bind to IL-11R $\alpha$ ; and

a light chain variable region (V<sub>L</sub>) that comprises complementary determining region V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences selected from **Table A1** and variants thereof which specifically bind to IL-11R $\alpha$ .

2. The isolated antibody, or antigen binding fragment thereof, of claim 1, wherein:

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 1-3, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 4-6, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 7-9, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 10-12, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 13-15, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 16-18, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 19-21, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 22-24, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 25-27, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 28-30, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 31-33, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 34-36, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 34-39, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 40-42, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 43-45, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 46-48, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 49-51, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 52-54, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 55-57, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 58-60, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 61-63, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 64-66, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 67-69, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 70-72, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 73-75, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 76-78, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 79-81, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 82-84, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 85-87, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 88-90, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 91-93, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 94-96, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 97-99, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 100-102, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 103-105, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 106-108, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 109-111, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 112-114, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 115-117, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 118-120, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 121-123, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 124-126, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 127-129, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 130-132, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 133-135, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 136-138, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 139-141, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 142-144, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 145-147, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 148-150, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 151-153, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 154-156, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 157-159, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 160-162, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 163-165, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 166-168, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 169-171, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 172-174, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 175-177, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 178-180, respectively;

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 181-183, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 184-186, respectively; or

the V<sub>H</sub>CDR1, V<sub>H</sub>CDR2, and V<sub>H</sub>CDR3 sequences comprise SEQ ID NOs: 187-189, respectively, and the V<sub>L</sub>CDR1, V<sub>L</sub>CDR2, and V<sub>L</sub>CDR3 sequences comprise SEQ ID NOs: 190-192, respectively.

3. The isolated antibody, or antigen binding fragment thereof, of claim 1 or 2, wherein the V<sub>H</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to a sequence selected from **Table A2**, optionally wherein the V<sub>H</sub> has 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 alterations in the framework regions.

4. The isolated antibody, or antigen binding fragment thereof, of any one of claims 1-3, wherein the V<sub>L</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to a sequence selected from **Table A2**, optionally wherein the V<sub>L</sub> has 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 alterations in the framework regions.

5. The isolated antibody, or antigen binding fragment thereof, of any one of claims 1-4, wherein:

the V<sub>H</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 193, and the V<sub>L</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 194;

the V<sub>H</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 195, and the V<sub>L</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 196;





the V<sub>H</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 245, and the V<sub>L</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 246;

the V<sub>H</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 247, and the V<sub>L</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 248;

the V<sub>H</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 249, and the V<sub>L</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 250;

the V<sub>H</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 251, and the V<sub>L</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 252;

the V<sub>H</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 253, and the V<sub>L</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 254; or

the V<sub>H</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 255, and the V<sub>L</sub> comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 256.

6. The isolated antibody, or antigen binding fragment thereof, of any one of claims 1-5, which binds to human IL-11R $\alpha$  (see **Table B1**).

7. The isolated antibody, or antigen binding fragment thereof, of claim 6, which binds to a fibronectin domain III of human IL-11R $\alpha$ , or approximately residues 112-219 of SEQ ID NO: 257.

8. The isolated antibody, or antigen binding fragment thereof, of any one of claims 1-7, which has one or more of the following characteristics:

has a binding affinity for human IL-11R $\alpha$  of less than about 500, 400, 300, 200, 100, 90, 80, 70, 60, 50, 40, or 30 pM, and optionally has increased binding affinity for human IL-11R $\alpha$  relative to that of the TS7 and 8E2 antibodies;

antagonizes the binding and/or signaling activity between IL-11R $\alpha$  and IL-11, and optionally has increased potency as an IL-11 signaling antagonist relative to that of the TS7 and 8E2 antibodies;

reduces IL-11R $\alpha$ /gp130 dimerization or complex formation, optionally in a cell-based assay; and/or

has reduced N-linked glycosylation in V<sub>L</sub>CDR3, optionally relative to the TS7 and 8E2 antibodies.

9. The isolated antibody, or antigen binding fragment thereof, of any one of claims 1-8, comprising an IgA (including subclasses IgA1 and IgA2), IgD, IgE, IgG (including subclasses IgG1, IgG2, IgG3, and IgG4), or IgM Fc domain, optionally a human Fc domain, or a hybrid and/or variant thereof.
10. The isolated antibody, or antigen binding fragment thereof, of any one of claims 1-9, comprising an IgG Fc domain with high effector function in humans, optionally an IgG1 or IgG3 Fc domain.
11. The isolated antibody, or antigen binding fragment thereof, of any one of claims 1-9, comprising an IgG Fc domain with low effector function in humans, optionally an IgG2 or IgG4 Fc domain.
12. The isolated antibody, or antigen binding fragment thereof, of any one of claims 1-9, comprising a human IgG1 or IgG4 Fc domain, optionally selected from **Table F1**.
13. The isolated antibody, or antigen binding fragment thereof, of any one of claims 1-12, which is a monoclonal antibody.
14. The isolated antibody, or antigen binding fragment thereof, of any one of claims 1-13, which is a humanized antibody,  
optionally wherein the antibody, or antigen binding fragment thereof, is a humanized monoclonal antibody that comprises a human IgG4 Fc domain with an S228P mutation (EU numbering).
15. The isolated antibody, or antigen binding fragment thereof, of any one of claims 1-14, which is selected from an Fv fragment, a single chain Fv (scFv) polypeptide, an adnectin, an anticalin, an aptamer, an avimer, a camelid antibody, a designed ankyrin repeat protein (DARPin), a minibody, a nanobody, and a unibody.
16. An isolated polynucleotide encoding the isolated antibody, or antigen binding fragment thereof, according to any one of claims 1-15, an expression vector comprising the isolated polynucleotide, or an isolated host cell comprising the vector.
17. A pharmaceutical composition, comprising the isolated antibody, or antigen binding fragment thereof, of any one of claims 1-15, and a pharmaceutically-acceptable carrier.

18. The pharmaceutical composition of claim 17, wherein the composition has a purity of at least about 80%, 85%, 90%, 95%, 98%, or 99% on a protein basis with respect to the at least one antibody or antigen binding fragment, and is substantially aggregate- and endotoxin-free.

19. The pharmaceutical composition of claim 17 or 18, wherein the composition has reduced or undetectable heterogeneity of N-linked glycosylation (optionally relative to the TS7 and 8E2 antibodies), optionally in the V<sub>L</sub>CDR3 sequence.

20. The pharmaceutical composition of any one of claims 17-19, wherein the composition is a sterile, injectable solution, optionally suitable for intravenous, intramuscular, subcutaneous, or intraperitoneal administration.

21. A method of treating a disease or condition in a subject in need thereof, comprising administering to the subject a pharmaceutical composition of any one of claims 17-20.

22. The method of claim 21, wherein the disease or condition is an IL-11-associated or IL-11-mediated disease or condition.

23. The method of claim 21 or 22, wherein the disease or condition is a cancer, an inflammatory disease, an autoimmune disease, a wasting disease, a bone disease, or fibrosis.

24. The method of claim 23, wherein the disease is a cancer, optionally a cancer that expresses or overexpresses IL-11R $\alpha$  and/or IL-11, optionally wherein the cancer displays IL-11R $\alpha$ /IL-11-dependent growth, adhesion, migration, invasion, and/or chemoresistance.

25. The method of claim 24, wherein the cancer is selected from one or more of bone cancer, prostate cancer, melanoma (e.g., metastatic melanoma), pancreatic cancer, small cell lung cancer, non-small cell lung cancer (NSCLC), mesothelioma, leukemia (e.g., lymphocytic leukemia, chronic myelogenous leukemia, acute myeloid leukemia, relapsed acute myeloid leukemia, hairy cell leukemias, acute lymphoblastic leukemias), lymphoma (e.g., non-Hodgkin's lymphomas, Hodgkin's lymphoma), hepatoma (hepatocellular carcinoma), sarcoma, B-cell malignancy, breast cancer, ovarian cancer, colorectal cancer, glioma, glioblastoma multiforme, meningioma, pituitary adenoma, vestibular schwannoma, primary CNS lymphoma, primitive neuroectodermal tumor (medulloblastoma), kidney cancer (e.g., renal cell carcinoma), bladder cancer, uterine cancer, esophageal cancer, brain cancer, head and neck cancers, cervical cancer, testicular cancer, thyroid cancer, and stomach cancer.

26. The method of claim 24 or 25, wherein the cancer is a metastatic cancer, optionally a metastatic cancer which has metastasized to the bone.

27. The method of claim 23, wherein the inflammatory disease is selected from one or more of airway or lung inflammation (e.g., inflammatory lung disease), asthma, rhinitis, chronic obstructive pulmonary disorder (COPD), dermatitis, psoriasis, hepatitis, gastric inflammation, irritable bowel syndrome (IBS), ulcerative colitis, Crohn's disease, colitis, diverticulitis, lupus erythematosus, nephritis, Parkinson's disease, multiple sclerosis (MS), Alzheimer's disease, arthritis, rheumatoid arthritis, sepsis, infection-induced inflammation, cardiovascular diseases such as atherosclerosis and vasculitis, diabetes, and gout.

28. The method of claim 23, wherein the autoimmune disease is selected from one or more of arthritis (including rheumatoid arthritis, reactive arthritis), systemic lupus erythematosus (SLE), psoriasis, inflammatory bowel disease (IBD), ulcerative colitis, Crohn's disease, encephalomyelitis, uveitis, myasthenia gravis, multiple sclerosis, insulin dependent diabetes, Addison's disease, celiac disease, chronic fatigue syndrome, autoimmune hepatitis, autoimmune alopecia, ankylosing spondylitis, fibromyalgia, pemphigus vulgaris, Sjogren's syndrome, Kawasaki's Disease, hyperthyroidism/Graves' disease, hypothyroidism/Hashimoto's disease, endometriosis, scleroderma, pernicious anemia, Goodpasture syndrome, Guillain-Barré syndrome, Wegener's disease, glomerulonephritis, aplastic anemia (including multiply transfused aplastic anemia patients), paroxysmal nocturnal hemoglobinuria, myelodysplastic syndrome, idiopathic thrombocytopenic purpura, autoimmune hemolytic anemia, Evan's syndrome, Factor VIII inhibitor syndrome, systemic vasculitis, dermatomyositis, polymyositis, rheumatic fever, autoimmune lymphoproliferative syndrome (ALPS), autoimmune bullous pemphigoid, Parkinson's disease, sarcoidosis, vitiligo, primary biliary cirrhosis, and autoimmune myocarditis.

29. The method of claim 23, wherein the wasting disease is selected from one or more of cachexia, optionally cachexia associated with cancer or renal failure, and sarcopenia.

30. The method of claim 23, wherein the bone disease is selected from osteoporosis (including post-menopausal osteoporosis), bone fracture, Paget's disease of bone, and bone resorption/damage associated with cancer or cancer therapy, including chemotherapy, hormone ablation, and hormone inhibition.

31. The method of claim 23, wherein the fibrosis is selected from fibrosis of the lungs, cardiovascular system, liver, brain, joints (optionally knee, hip, ankle, foot joints, shoulder, elbow,

wrist, hand joints, or spinal vertebrae), intestine, skin, kidney, liver, thyroid, bone marrow, retroperitoneum, and eye.

32. The method of claim 31, wherein the fibrosis of the lungs is selected from fibrothorax, pulmonary fibrosis (optionally cystic fibrosis or interstitial lung disease (ILD)), autosomal recessive genetic disease optionally Hermansky-Pudlak syndrome, and radiation-induced lung injury.

33. The method of claim 32, wherein the ILD is idiopathic ILD, optionally selected from idiopathic pulmonary fibrosis (IPF), desquamative interstitial pneumonia (DIP), acute interstitial pneumonia (AIP) or Hamman-Rich syndrome, nonspecific interstitial pneumonia (NSIP), respiratory bronchiolitis-associated interstitial lung disease (RB-ILD), cryptogenic organizing pneumonia (COP), and lymphoid interstitial pneumonia (LIP).

34. The method of claim 32, wherein the ILD is secondary ILD, optionally selected from ILD related to connective tissue and autoimmune diseases (optionally sarcoidosis, rheumatoid arthritis, systemic lupus erythematosus, systemic sclerosis, polymyositis, dermatomyositis, or antisynthetase syndrome), inhaled substances (optionally silicosis, asbestosis, berylliosis, industrial printing chemicals, or chronic hypersensitivity pneumonitis), drugs (optionally antibiotics, chemotherapeutics, or anti-arrhythmic agents), infections (optionally SARS CoV-2, atypical pneumonia, pneumocystis pneumonia, tuberculosis, Chlamydia trachomatis, or respiratory syncytial virus), malignancies (optionally lymphangitic carcinomatosis), and pediatric ILDs (optionally developmental disorders, growth abnormalities deficient alveolarization, infant conditions of undefined cause, and ILD related to alveolar surfactant region).

35. The method of claim 31, wherein the fibrosis of the cardiovascular system is myocardial fibrosis (optionally interstitial fibrosis or replacement fibrosis).