

SUGAR CHAIN-RELATED GENE AND USE THEREOF

ABSTRACT

5 As a result of dedicated studies, the present inventors succeeded in discovering, for the
first time, that fibrogenesis could be suppressed at the physiological tissue level by inhibiting
sulfation at position 4 or 6 of GalNAc, which is a sugar that constitutes sugar chains.
Furthermore, the present inventors conducted studies using various disease model animals, and
as a result, successfully demonstrated that inhibitors of sulfation at position 4 or 6 of GalNAc
10 had therapeutic effects on diseases caused by tissue fibrogenesis (tissue fibrogenic disorders).

(No Suitable Figures)

SUGAR CHAIN-RELATED GENE AND USE THEREOF

ABSTRACT

5 As a result of dedicated studies, the present inventors succeeded in
discovering, for the first time, that fibrogenesis could be suppressed at the
physiological tissue level by inhibiting sulfation at position 4 or 6 of GalNAc, which
is a sugar that constitutes sugar chains. Furthermore, the present inventors conducted
studies using various disease model animals, and as a result, successfully
10 demonstrated that inhibitors of sulfation at position 4 or 6 of GalNAc had therapeutic
effects on diseases caused by tissue fibrogenesis (tissue fibrogenic disorders).

(No Suitable Figures)