a Raman spectrum with peaks at 1623, 1503, 1436, 1337, 901, 853, 779, 744, 708, and 634 ± 2 cm^{-1}.

5. The crystalline N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide hydrate of claim 4, wherein the degree of hydration ranges from about 0.1 molar equivalent of water to about 1 molar equivalent of water relative to N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide.

6. A pharmaceutical composition comprising a therapeutically effective amount of the crystalline N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide hydrate of claim 1 and a pharmaceutically acceptable excipient.

7. Use of the crystalline N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide hydrate of claim 1 in the manufacture of a medicament for treating cancer in a subject.

8. The use of claim 7 wherein the subject is a human.

9. The use of claim 7, wherein the cancer being treated is selected from the group consisting of renal cancer, gastric cancer, head and neck cancer, lung cancer, breast cancer, prostate cancer, colorectal cancer, squamous cell myeloid leukemia, hemangiomas, melanomas, squamous cell carcinoma, hepatocellular carcinoma and brain cancer.

10. The use of claim 9, wherein the cancer being treated is selected from the group consisting of papillary renal cell carcinoma, squamous cell carcinoma and metastatic gastric carcinoma.

11. The use of claim 9, wherein the cancer is hepatocellular carcinoma.

12. Use of the crystalline N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide hydrate of claim 3 in the manufacture of a medicament for treating cancer in a subject, wherein the cancer being treated is selected from the group consisting of renal cancer, gastric cancer, head and neck cancer, lung cancer, breast cancer, prostate cancer, colorectal cancer, squamous cell myeloid leukemia, hemangiomas, melanomas, squamous cell carcinoma, hepatocellular carcinoma and brain cancer.

13. The use of claim 12 wherein the subject is human.

14. The use of claim 12, wherein the cancer being treated is selected from the group consisting of cell carcinoma, squamous cell carcinoma and metastatic gastric carcinoma.

15. The use of claim 12, wherein the cancer is hepatocellular carcinoma.

16. A method of preparing the crystalline N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide hydrate of claim 1 comprising the steps of:

dissolving N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide in an aqueous solvent, and

crystallizing the crystalline N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide hydrate from the aqueous solution.

17. A method of preparing the crystalline N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide hydrate of claim 1 comprising the steps of:

dissolving N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide in an aqueous solvent,

crystallizing the crystalline N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide hydrate from the aqueous solution, and

placing athe crystalline N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide hydrate in a humidity chamber under conditions and for a time sufficient to increase or decrease the degree of hydration of crystalline N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide hydrate.