

PRODRUGS OF FUMARATES AND THEIR USE IN TREATING VARIOUS DISEASES

RELATED APPLICATIONS

This application claims priority to U.S. Provisional Application No. 61/782,445, filed
5 March 14, 2013, and U.S. Provisional Application No. 61/934,365, filed January 31, 2014,
the contents of which are incorporated herein by reference in their entireties.

FIELD OF THE INVENTION

The present invention relates to various prodrugs of monomethyl fumarate. In
10 particular, the present invention relates to derivatives of monomethyl fumarate which offer
improved properties relative to dimethyl fumarate. The invention also relates to methods of
treating various diseases.

BACKGROUND OF THE INVENTION

15 Fumaric acid esters (FAEs) are approved in Germany for the treatment of psoriasis,
are being evaluated in the United States for the treatment of psoriasis and multiple sclerosis,
and have been proposed for use in treating a wide range of immunological, autoimmune, and
inflammatory diseases and conditions.

FAEs and other fumaric acid derivatives have been proposed for use in treating a
20 wide-variety of diseases and conditions involving immunological, autoimmune, and/or
inflammatory processes including psoriasis (Joshi and Strebel, WO 1999/49858; U.S. Pat.
No. 6,277,882; Mrowietz and Asadullah, *Trends Mol Med* 2005, 111(1), 43-48; and Yazdi
and Mrowietz, *Clinics Dermatology* 2008, 26, 522-526); asthma and chronic obstructive
pulmonary diseases (Joshi et al., WO 2005/023241 and US 2007/0027076); cardiac
25 insufficiency including left ventricular insufficiency, myocardial infarction and angina
pectoris (Joshi et al., WO 2005/023241; Joshi et al., US 2007/0027076); mitochondrial and
neurodegenerative diseases such as Parkinson's disease, Alzheimer's disease, Huntington's
disease, retinopathia pigmentosa and mitochondrial encephalomyopathy (Joshi and Strebel,
WO 2002/055063, US 2006/0205659, U.S. Pat. No. 6,509,376, U.S. Pat. No. 6,858,750, and
30 U.S. Pat. No. 7,157,423); transplantation (Joshi and Strebel, WO 2002/055063, US
2006/0205659, U.S. Pat. No. 6,359,003, U.S. Pat. No. 6,509,376, and U.S. Pat. No.
7,157,423; and Lehmann et al., *Arch Dermatol Res* 2002, 294, 399-404); autoimmune
diseases (Joshi and Strebel, WO 2002/055063, U.S. Pat. No. 6,509,376, U.S. Pat. No.
7,157,423, and US 2006/0205659) including multiple sclerosis (MS) (Joshi and Strebel, WO

1998/52549 and U.S. Pat. No. 6,436,992; Went and Lieberburg, US 2008/0089896; Schimrigk et al., *Eur J Neurology* 2006, 13, 604-610; and Schilling et al., *Clin Experimental Immunology* 2006, 145, 101-107); ischemia and reperfusion injury (Joshi et al., US 2007/0027076); AGE-induced genome damage (Heidland, WO 2005/027899); inflammatory
5 bowel diseases such as Crohn's disease and ulcerative colitis; arthritis; and others (Nilsson et al., WO 2006/037342 and Nilsson and Muller, WO 2007/042034).

Fumaderm®, an enteric coated tablet containing a salt mixture of monoethyl fumarate and dimethyl fumarate (DMF) which is rapidly hydrolyzed to monomethyl fumarate, regarded as the main bioactive metabolite, was approved in Germany in 1994 for the
10 treatment of psoriasis. Fumaderm® is dosed TID with 1-2 grams/day administered for the treatment of psoriasis. Fumaderm® exhibits a high degree of interpatient variability with respect to drug absorption and food strongly reduces bioavailability. Absorption is thought to occur in the small intestine with peak levels achieved 5-6 hours after oral administration. Significant side effects occur in 70-90% of patients (Brewer and Rogers, *Clin Expt'l*
15 *Dermatology* 2007, 32, 246-49; and Hoefnagel et al., *Br J Dermatology* 2003, 149, 363-369). Side effects of current FAE therapy include gastrointestinal upset including nausea, vomiting, diarrhea and/or transient flushing of the skin.

Multiple sclerosis (MS) is an autoimmune disease with the autoimmune activity directed against central nervous system (CNS) antigens. The disease is characterized by
20 inflammation in parts of the CNS, leading to the loss of the myelin sheathing around neuronal axons (gradual demyelination), axonal loss, and the eventual death of neurons, oligodendrocytes and glial cells.

Dimethyl fumarate (DMF) is the active component of the experimental therapeutic, BG-12, studied for the treatment of relapsing-remitting MS (RRMS). In a Phase IIb RRMS
25 study, BG-12 significantly reduced gadolinium-enhancing brain lesions. In preclinical studies, DMF administration has been shown to inhibit CNS inflammation in murine and rat EAE. It has also been found that DMF can inhibit astrogliosis and microglial activations associated with EAE. *See, e.g.*, US Published Application No. 2012/0165404.

There are four major clinical types of MS: 1) relapsing-remitting MS (RRMS),
30 characterized by clearly defined relapses with full recovery or with sequelae and residual deficit upon recovery; periods between disease relapses characterized by a lack of disease progression; 2) secondary progressive MS (SPMS), characterized by initial relapsing remitting course followed by progression with or without occasional relapses, minor remissions, and plateaus; 3) primary progressive MS (PPMS), characterized by disease

progression from onset with occasional plateaus and temporary minor improvements allowed; and 4) progressive relapsing MS (PRMS), characterized by progressive disease onset, with clear acute relapses, with or without full recovery; periods between relapses characterized by continuing progression.

5 Clinically, the illness most often presents as a relapsing-remitting disease and, to a lesser extent, as steady progression of neurological disability. Relapsing-remitting MS (RRMS) presents in the form of recurrent attacks of focal or multifocal neurologic dysfunction. Attacks may occur, remit, and recur, seemingly randomly over many years. Remission is often incomplete and as one attack follows another, a stepwise downward
10 progression ensues with increasing permanent neurological deficit. The usual course of RRMS is characterized by repeated relapses associated, for the majority of patients, with the eventual onset of disease progression. The subsequent course of the disease is unpredictable, although most patients with a relapsing-remitting disease will eventually develop secondary progressive disease. In the relapsing-remitting phase, relapses alternate with periods of
15 clinical inactivity and may or may not be marked by sequelae depending on the presence of neurological deficits between episodes. Periods between relapses during the relapsing-remitting phase are clinically stable. On the other hand, patients with progressive MS exhibit a steady increase in deficits, as defined above and either from onset or after a period of episodes, but this designation does not preclude the further occurrence of new relapses.

20 Notwithstanding the above, dimethyl fumarate is also associated with significant drawbacks.

For example, dimethyl fumarate is known to cause side effects upon oral administration, such as flushing and gastrointestinal events including, nausea, diarrhea, and/or upper abdominal pain in subjects. *See, e.g., Gold et al., N. Eng. J. Med.*, 2012,
25 367(12), 1098-1107. Dimethyl fumarate is dosed BID or TID with a total daily dose of about 480 mg to about 1 gram or more.

Further, in the use of a drug for long-term therapy it is desirable that the drug be formulated so that it is suitable for once- or twice-daily administration to aid patient compliance. A dosing frequency of once-daily or less is even more desirable.

30 Another problem with long-term therapy is the requirement of determining an optimum dose which can be tolerated by the patient. If such a dose is not determined this can lead to a diminution in the effectiveness of the drug being administered.

Accordingly, it is an object of the present invention to provide compounds and/or compositions which are suitable for long-term administration.

It is a further object of the present invention to provide the use of a pharmaceutical active agent in a manner which enables one to achieve a tolerable steady state level for the drug in a subject being treated therewith.

Because of the disadvantages of dimethyl fumarate described above, there continues to be a need to decrease the dosing frequency, reduce side-effects and/or improve the physicochemical properties associated with DMF. There remains, therefore, a real need in the treatment of neurological diseases, such as MS, for a product which retains the pharmacological advantages of DMF but overcomes its flaws in formulation and/or adverse effects upon administration. The present invention addresses these needs.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 depicts the hydrolysis of Compound 16 at pH 7.9, 25°C, showing vinylic region, as observed by NMR over 90 minutes.

Figure 2 depicts the hydrolysis of Compound 16 at pH 7.9, 25°C, showing vinylic region, as observed by NMR over 19 hours.

Figure 3 depicts the hydrolysis of Compound 16 at pH 7.9, 25°C, showing aliphatic region, as observed by NMR over 19 hours.

Figure 4 depicts the hydrolysis of Reference Compound A at pH 7.9, 37°C, showing vinylic region, as observed by NMR over 15 hours.

Figure 5 depicts the hydrolysis of Reference Compound A at pH 7.9, 37°C, showing aliphatic region, as observed by NMR over 15 hours.

Figure 6 depicts a plot of weight loss vs time for Compound 14 and DMF.


Figure 7 depicts the unit cell for crystalline Compound 14.

SUMMARY OF THE INVENTION

This invention is directed to the surprising and unexpected discovery of novel prodrugs and related methods useful in the treatment of neurological diseases. The methods and compositions described herein comprise one or more prodrugs (*e.g.*, aminoalkyl prodrugs) of monomethyl fumarate (MMF). The methods and compositions provide for a therapeutically effective amount of an active moiety in a subject for a time period of at least about 8 hours to at least about 24 hours.

More specifically, the compounds of the invention can be converted *in vivo*, upon oral administration, to monomethyl fumarate. Upon conversion, the active moiety (*i.e.*, monomethyl fumarate) is effective in treating subjects suffering from a neurological disease.

The present invention provides, in part, a compound of Formula (I), or a pharmaceutically acceptable salt, polymorph, hydrate, solvate or co-crystal thereof:



wherein:

5 R₁ is unsubstituted C₁-C₆ alkyl;

 L_a is substituted or unsubstituted C₁-C₆ alkyl linker, substituted or unsubstituted C₃-C₁₀ carbocycle, substituted or unsubstituted C₆-C₁₀ aryl, substituted or unsubstituted heterocycle comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S, or substituted or unsubstituted heteroaryl comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S; and

 R₂ and R₃ are each, independently, H, substituted or unsubstituted C₁-C₆ alkyl, substituted or unsubstituted C₂-C₆ alkenyl, substituted or unsubstituted C₂-C₆ alkynyl, substituted or unsubstituted C₆-C₁₀ aryl, substituted or unsubstituted C₃-C₁₀ carbocycle, substituted or unsubstituted heterocycle comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S, or substituted or unsubstituted heteroaryl comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S;

 or alternatively, R₂ and R₃, together with the nitrogen atom to which they are attached, form a substituted or unsubstituted heteroaryl comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S or a substituted or unsubstituted heterocycle comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S. The present invention also provides pharmaceutical compositions comprising one or more compounds of any of the formulae described herein and one or more pharmaceutically acceptable carriers.

 The present invention also provides methods of treating a neurological disease by administering to a subject in need thereof, a therapeutically effective amount of a compound of any of the formulae described herein, or a pharmaceutically acceptable salt, polymorph, hydrate, solvate or co-crystal thereof, such that the disease is treated.

 The present invention also provides methods of treating multiple sclerosis by administering to a subject in need thereof, a therapeutically effective amount of a compound

of any of the formulae described herein, or a pharmaceutically acceptable salt, polymorph, hydrate, solvate or co-crystal thereof, such that the multiple sclerosis is treated.

The present invention also provides methods of treating relapsing-remitting multiple sclerosis (RRMS) by administering to a subject in need thereof, a therapeutically effective
5 amount of a compound of any of the formulae described herein, or a pharmaceutically acceptable salt, polymorph, hydrate, solvate or co-crystal thereof, such that the multiple sclerosis is treated.

The present invention also provides methods of treating secondary progressive multiple sclerosis (SPMS) by administering to a subject in need thereof, a therapeutically effective
10 amount of a compound of any of the formulae described herein, or a pharmaceutically acceptable salt, polymorph, hydrate, solvate or co-crystal thereof, such that the multiple sclerosis is treated.

The present invention also provides methods of treating primary progressive multiple sclerosis (PPMS) by administering to a subject in need thereof, a therapeutically effective
15 amount of a compound of any of the formulae described herein, or a pharmaceutically acceptable salt, polymorph, hydrate, solvate or co-crystal thereof, such that the multiple sclerosis is treated.

The present invention also provides methods of treating progressive relapsing multiple sclerosis (PRMS) by administering to a subject in need thereof, a therapeutically effective
20 amount of a compound of any of the formulae described herein, or a pharmaceutically acceptable salt, polymorph, hydrate, solvate or co-crystal thereof, such that the multiple sclerosis is treated.

The present invention also provides methods of treating Alzheimer's disease by administering to a subject in need thereof, a therapeutically effective amount of a compound
25 of any of the formulae described herein, or a pharmaceutically acceptable salt, polymorph, hydrate, solvate or co-crystal thereof, such that the Alzheimer's disease is treated.

The present invention also provides methods of treating cerebral palsy by administering to a subject in need thereof, a therapeutically effective amount of a compound of any of the formulae described herein, or a pharmaceutically acceptable salt, polymorph,
30 hydrate, solvate or co-crystal thereof, such that the cerebral palsy is treated.

The present invention also provides compounds and compositions that enable improved oral, controlled- or sustained-release formulations. Specifically, dimethyl fumarate is administered twice or three times daily for the treatment of relapsing-remitting multiple sclerosis. In contrast, the compounds and compositions of the present invention may enable

formulations with a modified duration of therapeutic efficacy for reducing relapse rates in subjects with multiple sclerosis. For example, the present compounds and compositions provide therapeutically effective amounts of monomethyl fumarate in subjects for at least about 8 hours, at least about 12 hours, at least about 16 hours, at least about 20 hours or at least about 24 hours.

The present invention also provides compounds, compositions and methods which may result in decreased side effects upon administration to a subject relative to dimethyl fumarate. For example, gastric irritation and flushing are known side effects of oral administration of dimethyl fumarate in some subjects. The compounds, compositions and methods of the present invention can be utilized in subjects that have experienced or are at risk of developing such side effects.

The present invention also provides for compounds and compositions which exhibit improved physical stability relative to dimethyl fumarate. Specifically, dimethyl fumarate is known in the art to undergo sublimation at ambient and elevated temperature conditions. The compounds of the invention possess greater physical stability than dimethyl fumarate under controlled conditions of temperature and relative humidity. Specifically, in one embodiment, the compounds of the formulae described herein exhibit decreased sublimation relative to dimethyl fumarate.

Further, dimethyl fumarate is also known to be a contact irritant. *See e.g.*, Material Safety Data Sheet for DMF. In one embodiment, the compounds of the present invention exhibit reduced contact irritation relative to dimethyl fumarate. For example, the compounds of the formulae described herein exhibit reduced contact irritation relative to dimethyl fumarate.

The present invention also provides for compounds and compositions which exhibit decreased food effect relative to dimethyl fumarate. The bioavailability of dimethyl fumarate is known in the art to be reduced when administered with food. Specifically, in one embodiment, the compounds of the formulae described herein exhibit decreased food effect relative to dimethyl fumarate.

Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. In the specification, the singular forms also include the plural unless the context clearly dictates otherwise. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, suitable methods and materials are described below. All publications, patent applications, patents and

other references mentioned herein are incorporated by reference. The references cited herein are not admitted to be prior art to the claimed invention. In the case of conflict, the present specification, including definitions, will control. In addition, the materials, methods and examples are illustrative only and are not intended to be limiting.

- 5 Other features and advantages of the invention will be apparent from the following detailed description and claims.

DETAILED DESCRIPTION OF THE INVENTION

10 The present invention provides novel compounds and methods of treating a neurological disease by administering a compound of Formula (I), (Ia), (Ib), (II), (III), or (IV), synthetic methods for making a compound of Formula (I), (Ia), (Ib), (II), (III), or (IV), and pharmaceutical compositions containing a compound of Formula (I), (Ia), (Ib), (II), (III), or (IV).

15 The present invention also provides compounds and methods for the treatment of psoriasis by administering to a subject in need thereof, a therapeutically effective amount of a compound of Formula (I), (Ia), (Ib), (II), (III), or (IV), or a pharmaceutically acceptable salt, polymorph, hydrate, solvate or co-crystal thereof.


20 The present invention provides, in part, methods for the treatment of a neurological disease by administering to a subject in need thereof, a therapeutically effective amount of a compound of Formula (I), (Ia), (Ib), (II), (III), or (IV), or a pharmaceutically acceptable salt, polymorph, hydrate, solvate or co-crystal thereof. The neurological disease can be multiple sclerosis. The present invention further provides the use of a compound of Formula (I), (Ia), (Ib), (II), (III), or (IV), or a pharmaceutically acceptable salt, polymorph, hydrate, solvate or co-crystal thereof, for the preparation of a medicament useful for the treatment of a
25 neurological disease.

 According to the present invention, a neurological disease is a disorder of the brain, spinal cord or nerves in a subject. In one embodiment, the neurological disease is characterized by demyelination, or degeneration of the myelin sheath, of the central nervous system. The myelin sheath facilitates the transmission of nerve impulses through a nerve
30 fiber or axon. In another embodiment, the neurological disease is selected from the group consisting of multiple sclerosis, Alzheimer's disease, cerebral palsy, spinal cord injury, Amyotrophic lateral sclerosis (ALS), stroke, Huntington's disease, Parkinson's disease, optic neuritis, Devic disease, transverse myelitis, acute disseminated encephalomyelitis, adrenoleukodystrophy and adrenomyeloneuropathy, acute inflammatory demyelinating

polyneuropathy (AIDP), chronic inflammatory demyelinating polyneuropathy (CIDP), acute transverse myelitis, progressive multifocal leucoencephalopathy (PML), acute disseminated encephalomyelitis (ADEM), and other hereditary disorders, such as leukodystrophies, Leber's optic atrophy, and Charcot-Marie-Tooth disease. In some embodiments, the neurological disorder is an auto-immune disease. In one embodiment, the neurological disease is multiple sclerosis. In another embodiment, the neurological disease is stroke. In another embodiment, the neurological disease is Alzheimer's disease. In another embodiment, the neurological disease is cerebral palsy. In another embodiment, the neurological disease is spinal cord injury. In another embodiment, the neurological disease is ALS. In another embodiment, the neurological disease is Huntington's disease. *See, e.g.*, US Patent No. 8,007,826, WO2005/099701 and WO2004/082684, which are incorporated by reference in their entireties.

In a further embodiment, the present invention provides methods for the treatment of a disease or a symptom of a disease described herein by administering to a subject in need thereof, a therapeutically effective amount of a compound of Formula (I), (Ia), (Ib), (II), (III), or (IV), or a pharmaceutically acceptable salt, polymorph, hydrate, solvate or co-crystal thereof. The present invention further provides the use of a compound of Formula (I), (Ia), (Ib), (II), (III), or (IV), or a pharmaceutically acceptable salt, polymorph, hydrate, solvate or co-crystal thereof, for the preparation of a medicament useful for the treatment of a disease or a symptom of a disease described herein.

In another embodiment, the present invention provides a compound of Formula (I), or a pharmaceutically acceptable salt, polymorph, hydrate, solvate or co-crystal thereof, or a method for the treatment of a neurological disease by administering to a subject in need thereof, a therapeutically effective amount of a compound of Formula (I), or a pharmaceutically acceptable salt, polymorph, hydrate, solvate or co-crystal thereof:



wherein:

R₁ is unsubstituted C₁-C₆ alkyl;

L_a is substituted or unsubstituted C_1 - C_6 alkyl linker, substituted or unsubstituted C_3 - C_{10} carbocycle, substituted or unsubstituted C_6 - C_{10} aryl, substituted or unsubstituted heterocycle comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S, or substituted or unsubstituted heteroaryl comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S; and

R_2 and R_3 are each, independently, H, substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_2 - C_6 alkenyl, substituted or unsubstituted C_2 - C_6 alkynyl, substituted or unsubstituted C_6 - C_{10} aryl, substituted or unsubstituted C_3 - C_{10} carbocycle, substituted or unsubstituted heterocycle comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S, or substituted or unsubstituted heteroaryl comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S;

or alternatively, R_2 and R_3 , together with the nitrogen atom to which they are attached, form a substituted or unsubstituted heteroaryl comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S or a substituted or unsubstituted heterocycle comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S.

In one aspect of the compound of Formula (I), or a pharmaceutically acceptable salt, polymorph, hydrate, solvate or co-crystal thereof:

R_1 is unsubstituted C_1 - C_6 alkyl;

L_a is unsubstituted C_1 - C_6 alkyl linker, unsubstituted C_3 - C_{10} carbocycle, unsubstituted C_6 - C_{10} aryl, unsubstituted heterocycle comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S, or unsubstituted heteroaryl comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S; and

R_2 and R_3 are each, independently, H, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_6 - C_{10} aryl, C_3 - C_{10} carbocycle, heterocycle comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S, or heteroaryl comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S, wherein the alkyl, alkenyl, aryl, carbocycle, heterocycle, or heteroaryl groups may be optionally independently substituted one or more times with C_1 - C_3 -alkyl, OH, $O(C_1$ - C_4 alkyl), carbonyl, halo, NH_2 , $N(H)(C_1$ - C_6 alkyl), $N(C_1$ - C_6 alkyl) $_2$, SO_2H , $SO_2(C_1$ - C_6 alkyl), CHO, CO_2H , $CO_2(C_1$ - C_6 alkyl), or CN;

or alternatively, R_2 and R_3 , together with the nitrogen atom to which they are attached, form a heteroaryl comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S; or a heterocycle comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S, wherein the heteroaryl or heterocycle may be

optionally substituted one or more times with C₁-C₆ alkyl, CN, OH, halo, O(C₁-C₆ alkyl), CHO, carbonyl, thione, NO, or NH₂.

In one embodiment of this aspect, at least one of R₁ and R² is H.

In another embodiment of this aspect, L_a is (CH₂)₂.

5 In another embodiment of Formula (I), R₂ and R₃ together with the nitrogen to which they are attached form a heteroaryl, wherein the heteroaryl ring is a pyrrole ring, a pyrazole ring, an imidazole ring, a benzimidazole ring, a thiazole ring, a 1*H*-1,2,4-triazole ring, a 1*H*-1,2,3-triazole ring, a 1*H*-tetrazole ring, a pyrimidinone ring, an indole ring, or a benzoisothiazole ring, wherein all of the rings may be optionally substituted one or more
10 times with C₁-C₆ alkyl, CN, OH, O(C₁-C₆ alkyl), CHO, NO₂, or NH₂.

In still another embodiment of Formula (I), R₂ and R₃ together with the nitrogen to which they are attached form a heterocycle, wherein the heterocycle is a morpholine ring, a thiomorpholine ring, a pyrrolidine ring, a 2,5-dihydropyrrole ring, a 1,2-dihydropyridine ring, a piperazine ring, a succinimide ring, an isoindoline ring, a 2,5-dihydro-1*H*-tetrazole ring, an
15 azetidine ring, a piperidine ring, a hexahydropyrimidine ring, a 2,3,3a,4,7,7a-hexahydro-1*H*-4,7-epoxyisoindole ring, a 3,4-dihydroquinazoline ring, a 1,2,3,4-tetrahydroquinazoline ring, an oxazolidine ring, an oxazolidinone ring, an imidazolidinone ring, a 1,3-dihydro-2*H*-imidazol-2-one ring, an imidizolidine thione ring, or an isothiazolidine ring, wherein all of the rings may be optionally substituted one or more times with C₁-C₆ alkyl, CO₂(C₁-C₆ alkyl),
20 OH, (CH₂)₁₋₄OH, O(C₁-C₆ alkyl), halo, NH₂, (CH₂)₁₋₄NH₂, (CH₂)₁₋₄NH(C₁-C₄ alkyl), (CH₂)₁₋₄N(C₁-C₄ alkyl)₂, carbonyl, or thione.

In one embodiment of Formula (I):

R₁ is unsubstituted C₁-C₃ alkyl;

L_a is (CH₂)₁₋₆; and

25 R₂ and R₃ are each, independently: H, methyl, ethyl, isopropyl, butyl, *tert*-butyl, cyclohexyl, cyclohexenyl, phenyl, benzyl, benzodioxole, pyridinyl, (CH₂)₂N(CH₃)₂, (CH₂)₃SO₂H, (CH₂)₂SO₂Me, CH₂CO₂H, or (CH₂)₂CN,

or alternatively, R₂ and R₃, together with the nitrogen atom to which they are attached, form a morpholine ring optionally substituted one or more times with C₁-C₄ alkyl,
30 carbonyl, or (CH₂)₁₋₃N(C₁-C₄ alkyl)₂; an 8-oxa-3-azabicyclo[3.2.1]octane ring; a 1,4-dioxo-8-azaspiro[4.5]decane ring; a thiomorpholine ring substituted one or more times with carbonyl or thione; a piperazine ring optionally substituted with C₁-C₄ alkyl, halo, (CH₂)₂OH, C₁-C₄ alkyl ester; a pyrrolidine ring optionally substituted one or more times with C₁-C₄ alkyl or carbonyl; a 2,5-dihydropyrrole ring optionally substituted one or more times with C₁-C₄ alkyl

or carbonyl; a succinimide ring optionally substituted one or more times with C₁-C₄ alkyl; a 3-azabicyclo[3.1.0]hexane-2,4-dione ring; a hexahydropyrimidine ring optionally substituted one or more times with C₁-C₄ alkyl or carbonyl; a pyrimidinone ring optionally substituted one or more times with C₁-C₄ alkyl; a pyrrole ring optionally substituted one or more times with C₁-C₄ alkyl, halo, C(O)NH₂, or NO₂; a pyrazole ring optionally substituted one or more times with C₁-C₄ alkyl, C(O)NH₂, or NO₂; an imidazole ring optionally substituted one or more times with C₁-C₄ alkyl or NO₂; a 1,3-dihydro-2H-imidazol-2-one ring; a benzimidazole ring; a thiazole ring; an isoindoline ring substituted with carbonyl; a 1H-tetrazole ring; a 1H-2,5-dihydro-1H-tetrazole ring substituted with thione; a 1H-1,2,4-triazole ring; a 1H-1,2,3-triazole ring; an azetidine ring substituted with carbonyl; an piperidine ring optionally substituted one or more times with C₁-C₄ alkyl, carbonyl, halo, OH, or (CH₂)₁₋₄OH; a pyridinone ring optionally substituted one or more times with C₁-C₄ alkyl, OH, or CN; a 1,2-dihydropyridine ring substituted with carbonyl; a pyrimidinone ring optionally substituted one or more times with C₁-C₄ alkyl; an oxazolidine ring optionally substituted one or more times with C₁-C₄ alkyl; an oxazolidinone ring; an imidazolidinone ring optionally substituted one or more times with C₁-C₄ alkyl or carbonyl; an imidizolidine thione ring; an isothiazolidine ring optionally substituted one or more times with carbonyl; an indole ring; a 2,3,3a,4,7,7a-hexahydro-1H-4,7-epoxyisoindole ring optionally substituted one or more times with carbonyl; a 3,4-dihydroquinazoline ring substituted with carbonyl; 1,2,3,4-tetrahydroquinazoline ring substituted one or more times with carbonyl; or a benzoisothiazole ring optionally substituted one or more times with carbonyl.

In another embodiment of Formula (I):

R₁ is unsubstituted C₁-C₃ alkyl;

L_a is (CH₂)₁₋₆; and

R₂ and R₃ are each, independently: H, methyl, ethyl, isopropyl, butyl, *tert*-butyl, cyclohexyl, phenyl, benzyl, benzodioxole, pyridinyl, (CH₂)₂N(CH₃)₂, (CH₂)₃SO₂H, (CH₂)₂SO₂Me, CH₂CO₂H, or (CH₂)₂CN;

or alternatively, R₂ and R₃, together with the nitrogen atom to which they are attached, form a morpholine ring optionally substituted one or more times with C₁-C₄ alkyl, carbonyl, or (CH₂)₁₋₃N(C₁-C₄ alkyl)₂; an 8-oxa-3-azabicyclo[3.2.1]octane ring; a thiomorpholine ring substituted one or more times with carbonyl or thione; a piperazine ring substituted with C₁-C₄ alkyl ester; a pyrrolidine ring optionally substituted one or more times with C₁-C₄ alkyl or carbonyl; a 2,5-dihydropyrrole ring optionally substituted one or more times with C₁-C₄ alkyl or carbonyl; a succinimide ring optionally substituted one or more

times with C₁-C₄ alkyl; a 3-azabicyclo[3.1.0]hexane-2,4-dione ring; a hexahydropyrimidine ring optionally substituted one or more times with C₁-C₄ alkyl or carbonyl; a pyrimidinone ring optionally substituted one or more times with C₁-C₄ alkyl; an imidazole ring substituted with NO₂; an isoindoline ring substituted with carbonyl; an azetidine ring substituted with carbonyl; an piperidine ring optionally substituted one or more times with C₁-C₄ alkyl, carbonyl, halo, OH, or (CH₂)₁₋₄OH; a pyridinone ring optionally substituted one or more times with C₁-C₄ alkyl, OH, or CN; a pyrimidinone ring optionally substituted one or more times with C₁-C₄ alkyl; an oxazolidine ring optionally substituted one or more times with C₁-C₄ alkyl; an oxazolidinone ring; an imidazolidinone ring optionally substituted one or more times with C₁-C₄ alkyl or carbonyl; an imidizolidine thione ring; an isothiazolidine ring optionally substituted one or more times with carbonyl; or a benzoisothiazole ring optionally substituted one or more times with carbonyl.

In one aspect of the compound of Formula (I), or a pharmaceutically acceptable salt, polymorph, hydrate, solvate or co-crystal thereof:

R₁ is unsubstituted C₁-C₆ alkyl;

L_a is unsubstituted C₁-C₆ alkyl linker, unsubstituted C₃-C₁₀ carbocycle, unsubstituted C₆-C₁₀ aryl, unsubstituted heterocycle comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S, or unsubstituted heteroaryl comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S; and

or R₂ and R₃, together with the nitrogen atom to which they are attached, form a heteroaryl comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S; or a heterocycle comprising a 5-member ring and 1-3 heteroatoms selected from N, O and S, wherein the heteroaryl or heterocycle may be optionally substituted one or more times with C₁-C₆ alkyl, CN, OH, halo, O(C₁-C₆ alkyl), CHO, carbonyl, thione, NO, or NH₂.

In one embodiment of this aspect, at least one of R₂ and R₃ is H.

In another embodiment of this aspect, L_a is (CH₂)₂.

In still another embodiment of Formula (I), R₂ and R₃ together with the nitrogen to which they are attached form a heterocycle, wherein the heterocycle is, a thiomorpholine ring, a pyrrolidine ring, a 2,5-dihydropyrrole ring, a 1,2-dihydropyridine ring, a piperazine ring, a succinimide ring, an isoindoline ring, a 2,5-dihydro-1*H*-tetrazole ring, an azetidine ring, a piperidine ring, a hexahydropyrimidine ring, a 2,3,3a,4,7,7a-hexahydro-1*H*-4,7-epoxyisoindole ring, a 3,4-dihydroquinazoline ring, a 1,2,3,4-tetrahydroquinazoline ring, an oxazolidine ring, an oxazolidinone ring, an imidazolidinone ring, a 1,3-dihydro-2*H*-imidazol-2-one ring, an imidizolidine thione ring, or an isothiazolidine ring, wherein all of the rings

may be optionally substituted one or more times with C₁-C₆ alkyl, CO₂(C₁-C₆ alkyl), OH, (CH₂)₁₋₄OH, O(C₁-C₆ alkyl), halo, NH₂, (CH₂)₁₋₄NH₂, (CH₂)₁₋₄NH(C₁-C₄ alkyl), (CH₂)₁₋₄N(C₁-C₄ alkyl)₂, carbonyl, or thione.

In one embodiment of Formula (I):

- 5 R₁ is unsubstituted C₁-C₃ alkyl;
 L_a is (CH₂)₁₋₆; and
 R₂ and R₃, together with the nitrogen atom to which they are attached, form a morpholine ring substituted one or more times with C₁-C₄ alkyl, carbonyl, or (CH₂)₁₋₃N(C₁-C₄ alkyl)₂; an 8-oxa-3-azabicyclo[3.2.1]octane ring; a 1,4-dioxo-8-azaspiro[4.5]decane ring; a
 10 thiomorpholine ring substituted one or more times with carbonyl or thione; a piperazine ring optionally substituted with C₁-C₄ alkyl, halo, (CH₂)₂OH, C₁-C₄ alkyl ester; a pyrrolidine ring optionally substituted one or more times with C₁-C₄ alkyl or carbonyl; a 2,5-dihydropyrrole ring optionally substituted one or more times with C₁-C₄ alkyl or carbonyl; a succinimide ring optionally substituted one or more times with C₁-C₄ alkyl; a 3-azabicyclo[3.1.0]hexane-2,4-
 15 dione ring; a hexahydropyrimidine ring optionally substituted one or more times with C₁-C₄ alkyl or carbonyl; a pyrimidinone ring optionally substituted one or more times with C₁-C₄ alkyl; a pyrrole ring optionally substituted one or more times with C₁-C₄ alkyl, halo, C(O)NH₂, or NO₂; a pyrazole ring optionally substituted one or more times with C₁-C₄ alkyl, C(O)NH₂, or NO₂; an imidazole ring optionally substituted one or more times with C₁-C₄
 20 alkyl or NO₂; a 1,3-dihydro-2H-imidazol-2-one ring; a benzimidazole ring; a thiazole ring; an isoindoline ring substituted with carbonyl; a 1H-tetrazole ring; a 1H-2,5-dihydro-1H-tetrazole ring substituted with thione; a 1H-1,2,4-triazole ring; a 1H-1,2,3-triazole ring; an azetidine ring substituted with carbonyl; an piperidine ring optionally substituted one or more times with C₁-C₄ alkyl, carbonyl, halo, OH, or (CH₂)₁₋₄OH; a pyridinone ring optionally substituted
 25 one or more times with C₁-C₄ alkyl, OH, or CN; a 1,2-dihydropyridine ring substituted with carbonyl; a pyrimidinone ring optionally substituted one or more times with C₁-C₄ alkyl; an oxazolidine ring optionally substituted one or more times with C₁-C₄ alkyl; an oxazolidinone ring; an imidazolidinone ring optionally substituted one or more times with C₁-C₄ alkyl or carbonyl; an imidizolidine thione ring; an isothiazolidine ring optionally substituted one or
 30 more times with carbonyl; an indole ring; a 2,3,3a,4,7,7a-hexahydro-1H-4,7-epoxyisoindole ring optionally substituted one or more times with carbonyl; a 3,4-dihydroquinazoline ring substituted with carbonyl; 1,2,3,4-tetrahydroquinazoline ring substituted one or more times with carbonyl; or a benzoisothiazole ring optionally substituted one or more times with carbonyl.

In some embodiments of Formula (I), at least one of R^1 and R^2 is H.

In other embodiments of Formula (I), L_a is $(CH_2)_2$.

In a particular embodiment of Formula (I):

R_1 is methyl;

5 L_a is $(CH_2)_2$; and

R_2 and R_3 , together with the nitrogen atom to which they are attached, form a succinimide ring.

In another embodiment of Formula (I):

R_1 is methyl;

10 L_a is $(CH_2)_3$; and

R_2 and R_3 , together with the nitrogen atom to which they are attached, form a succinimide ring.

In still another embodiment of Formula (I):

R_1 is methyl;

15 L_a is $(CH_2)_4$; and

R_2 and R_3 , together with the nitrogen atom to which they are attached, form a succinimide ring.

For example, the neurological disease is multiple sclerosis.

For example, the neurological disease is relapsing-remitting multiple sclerosis
20 (RRMS).

For example, the compound of Formula (I) is a compound listed in Table 1 herein.

For example, in the compound of Formula (I), R_1 is methyl.

For example, in the compound of Formula (I), R_1 is ethyl.

For example, in the compound of Formula (I), L_a is substituted or unsubstituted C_1 - C_6
25 alkyl linker.

For example, in the compound of Formula (I), L_a is substituted or unsubstituted C_1 - C_3 alkyl linker.

For example, in the compound of Formula (I), L_a is substituted or unsubstituted C_2 alkyl linker.

30 For example, in the compound of Formula (I), L_a is methyl substituted or unsubstituted C_2 alkyl linker.

For example, in the compound of Formula (I), L_a is di-methyl substituted or unsubstituted C_2 alkyl linker.

For example, in the compound of Formula (I), L_a is methyl or di-methyl substituted C_2 alkyl linker.

For example, in the compound of Formula (I), L_a is unsubstituted C_2 alkyl linker.

For example, in the compound of Formula (I), R_2 is substituted or unsubstituted C_1 - C_6 alkyl.

For example, in the compound of Formula (I), R_2 is unsubstituted C_1 - C_6 alkyl.

For example, in the compound of Formula (I), R_2 is unsubstituted C_1 - C_3 alkyl.

For example, in the compound of Formula (I), R_2 is unsubstituted C_1 - C_2 alkyl.

For example, in the compound of Formula (I), R_2 is $C(O)OR_a$ -substituted C_1 - C_6 alkyl, wherein R_a is H or unsubstituted C_1 - C_6 alkyl.

For example, in the compound of Formula (I), R_2 is $S(O)(O)R_b$ -substituted C_1 - C_6 alkyl, wherein R_b is unsubstituted C_1 - C_6 alkyl.

For example, in the compound of Formula (I), R_3 is H.

For example, in the compound of Formula (I), R_3 is substituted or unsubstituted C_1 - C_6 alkyl.

For example, in the compound of Formula (I), R_3 is unsubstituted C_1 - C_6 alkyl.

For example, in the compound of Formula (I), R_2 and R_3 , together with the nitrogen atom to which they are attached, form a substituted or unsubstituted heteroaryl comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S, or a substituted or unsubstituted heterocycle comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S.

For example, in the compound of Formula (I), R_2 and R_3 , together with the nitrogen atom to which they are attached, form a substituted or unsubstituted heterocycle comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S.

For example, in the compound of Formula (I), R_2 and R_3 , together with the nitrogen atom to which they are attached, form a substituted or unsubstituted pyrrolidinyl, imidazolidinyl, pyrazolidinyl, oxazolidinyl, isoxazolidinyl, triazolidinyl, tetrahydrofuranyl, piperidinyl, piperazinyl, or morpholinyl ring.

For example, in the compound of Formula (I), R_2 and R_3 , together with the nitrogen atom to which they are attached, form a substituted or unsubstituted piperidinyl ring.

For example, in the compound of Formula (I), R_2 and R_3 , together with the nitrogen atom to which they are attached, form an unsubstituted piperidinyl ring.

For example, in the compound of Formula (I), R_2 and R_3 , together with the nitrogen atom to which they are attached, form a halogen substituted piperidinyl ring.

For example, in the compound of Formula (I), R₂ and R₃, together with the nitrogen atom to which they are attached, form a 4-halogen substituted piperidinyl ring.

For example, in the compound of Formula (I), R₂ and R₃, together with the nitrogen atom to which they are attached, form an unsubstituted morpholinyl ring.

5 For example, in the compound of Formula (I), R₂ and R₃, together with the nitrogen atom to which they are attached, form a morpholino N-oxide ring.

For example, in the compound of Formula (I), R₂ and R₃, together with the nitrogen atom to which they are attached, form an unsubstituted pyrrolidinyl ring.

10 For example, in the compound of Formula (I), R₂ and R₃, together with the nitrogen atom to which they are attached, form a substituted or unsubstituted heteroaryl comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S.

For example, in the compound of Formula (I), R₂ is substituted or unsubstituted C₆-C₁₀ aryl.


For example, in the compound of Formula (I), R₂ is unsubstituted C₆-C₁₀ aryl.

15 For example, in the compound of Formula (I), R₂ is unsubstituted phenyl.

For example, in the compound of Formula (I), R₂ is unsubstituted benzyl.

In another embodiment, the present invention provides a compound of Formula (Ia), or a pharmaceutically acceptable salt, polymorph, hydrate, solvate or co-crystal thereof, or a method for the treatment of a neurological disease by administering to a subject in need thereof a therapeutically effective amount of a compound of Formula (Ia), or a pharmaceutically acceptable salt, polymorph, hydrate, solvate or co-crystal thereof:

20



wherein:

25 R₁ is unsubstituted C₁-C₆ alkyl;

L_a is substituted or unsubstituted C₁-C₆ alkyl linker, substituted or unsubstituted C₃-C₁₀ carbocycle, substituted or unsubstituted C₆-C₁₀ aryl, substituted or unsubstituted heterocycle comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from

N, O and S, or substituted or unsubstituted heteroaryl comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S; and

R₂ is H, substituted or unsubstituted C₁-C₆ alkyl, substituted or unsubstituted C₂-C₆ alkenyl, substituted or unsubstituted C₂-C₆ alkynyl, substituted or unsubstituted C₆-C₁₀ aryl, substituted or unsubstituted C₃-C₁₀ carbocycle, substituted or unsubstituted heterocycle comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S, or substituted or unsubstituted heteroaryl comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S.

For example, the neurological disease is multiple sclerosis.

For example, the neurological disease is relapsing-remitting multiple sclerosis (RRMS).

For example, in the compound of Formula (Ia), R₁ is methyl.

For example, in the compound of Formula (Ia), R₁ is ethyl.

For example, in the compound of Formula (Ia), L_a is substituted or unsubstituted C₁-C₆ alkyl linker.

For example, in the compound of Formula (Ia), L_a is substituted or unsubstituted C₁-C₃ alkyl linker.

For example, in the compound of Formula (Ia), L_a is substituted or unsubstituted C₂ alkyl linker.

For example, in the compound of Formula (Ia), L_a is methyl substituted or unsubstituted C₂ alkyl linker.

For example, in the compound of Formula (Ia), L_a is di-methyl substituted or unsubstituted C₂ alkyl linker.

For example, in the compound of Formula (Ia), L_a is methyl or di-methyl substituted C₂ alkyl linker.

For example, in the compound of Formula (Ia), L_a is unsubstituted C₂ alkyl linker.

For example, in the compound of Formula (Ia), R₂ is substituted or unsubstituted C₁-C₆ alkyl.

For example, in the compound of Formula (Ia), R₂ is unsubstituted C₁-C₆ alkyl.

For example, in the compound of Formula (Ia), R₂ is methyl.


For example, in the compound of Formula (Ia), R₂ is unsubstituted C₁-C₃ alkyl.

For example, in the compound of Formula (Ia), R₂ is unsubstituted C₁-C₂ alkyl.

For example, in the compound of Formula (Ia), R₂ is C(O)OR_a-substituted C₁-C₆ alkyl, wherein R_a is H or unsubstituted C₁-C₆ alkyl.

For example, in the compound of Formula (Ia), R_2 is $S(O)(O)R_b$ -substituted C_1 - C_6 alkyl, wherein R_b is unsubstituted C_1 - C_6 alkyl.

In another embodiment, the present invention provides a compound of Formula (Ib), or a pharmaceutically acceptable polymorph, hydrate, solvate or co-crystal thereof, or a method for the treatment of a neurological disease by administering to a subject in need thereof a therapeutically effective amount of a compound of Formula (Ib), or a pharmaceutically acceptable polymorph, hydrate, solvate or co-crystal thereof:



A^- is a pharmaceutically acceptable anion;

R_1 is unsubstituted C_1 - C_6 alkyl;

L_a is substituted or unsubstituted C_1 - C_6 alkyl linker, substituted or unsubstituted C_3 - C_{10} carbocycle, substituted or unsubstituted C_6 - C_{10} aryl, substituted or unsubstituted heterocycle comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S, or substituted or unsubstituted heteroaryl comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S;

R_3' is substituted or unsubstituted C_1 - C_6 alkyl; and

R_2 and R_3 are each, independently, H, substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_2 - C_6 alkenyl, substituted or unsubstituted C_2 - C_6 alkynyl, substituted or unsubstituted C_6 - C_{10} aryl, substituted or unsubstituted C_3 - C_{10} carbocycle, substituted or unsubstituted heterocycle comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S, or substituted or unsubstituted heteroaryl comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S;

or alternatively, R_2 and R_3 , together with the nitrogen atom to which they are attached, form a substituted or unsubstituted heteroaryl comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S, or a substituted or unsubstituted heterocycle comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S.

For example, the neurological disease is multiple sclerosis.

For example, the neurological disease is relapsing-remitting multiple sclerosis (RRMS).

For example, in the compound of Formula (Ib), R₁ is methyl.

For example, in the compound of Formula (Ib), R₁ is ethyl.

5 For example, in the compound of Formula (Ib), L_a is substituted or unsubstituted C₁-C₆ alkyl linker.

For example, in the compound of Formula (Ib), L_a is substituted or unsubstituted C₁-C₃ alkyl linker.

10 For example, in the compound of Formula (Ib), L_a is substituted or unsubstituted C₂ alkyl linker.

For example, in the compound of Formula (Ib), L_a is methyl substituted or unsubstituted C₂ alkyl linker.

For example, in the compound of Formula (Ib), L_a is di-methyl substituted or unsubstituted C₂ alkyl linker.

15 For example, in the compound of Formula (Ib), L_a is methyl or di-methyl substituted C₂ alkyl linker.

For example, in the compound of Formula (Ib), L_a is unsubstituted C₂ alkyl linker.

For example, in the compound of Formula (Ib), R₂ is substituted or unsubstituted C₁-C₆ alkyl.

20 For example, in the compound of Formula (Ib), R₂ is unsubstituted C₁-C₆ alkyl.

For example, in the compound of Formula (Ib), R₂ is unsubstituted C₁-C₃ alkyl.

For example, in the compound of Formula (Ib), R₂ is unsubstituted C₁-C₂ alkyl.

For example, in the compound of Formula (Ib), R₂ is C(O)OR_a-substituted C₁-C₆ alkyl, wherein R_a is H or unsubstituted C₁-C₆ alkyl.

25 For example, in the compound of Formula (Ib), R₂ is S(O)(O)R_b-substituted C₁-C₆ alkyl, wherein R_b is unsubstituted C₁-C₆ alkyl.

For example, in the compound of Formula (Ib), R₃ is H.

For example, in the compound of Formula (Ib), R₃ is substituted or unsubstituted C₁-C₆ alkyl.

30 For example, in the compound of Formula (Ib), R₃ is unsubstituted C₁-C₆ alkyl.

For example, in the compound of Formula (Ib), R₂ and R₃, together with the nitrogen atom to which they are attached, form a substituted or unsubstituted heteroaryl comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S, or a

substituted or unsubstituted heterocycle comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S.

For example, in the compound of Formula (Ib), R₂ and R₃, together with the nitrogen atom to which they are attached, form a substituted or unsubstituted heterocycle comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S.

For example, in the compound of Formula (Ib), R₂ and R₃, together with the nitrogen atom to which they are attached, form a substituted or unsubstituted pyrrolidinyl, imidazolidinyl, pyrazolidinyl, oxazolidinyl, isoxazolidinyl, triazolidinyl, tetrahydrofuranyl, piperidinyl, piperazinyl, or morpholinyl ring.

For example, in the compound of Formula (Ib), R₂ and R₃, together with the nitrogen atom to which they are attached, form a substituted or unsubstituted piperidinyl ring.

For example, in the compound of Formula (Ib), R₂ and R₃, together with the nitrogen atom to which they are attached, form an unsubstituted piperidinyl ring.

For example, in the compound of Formula (Ib), R₂ and R₃, together with the nitrogen atom to which they are attached, form a halogen substituted piperidinyl ring.

For example, in the compound of Formula (Ib), R₂ and R₃, together with the nitrogen atom to which they are attached, form a 4-halogen substituted piperidinyl ring.

For example, in the compound of Formula (Ib), R₂ and R₃, together with the nitrogen atom to which they are attached, form an unsubstituted morpholinyl ring.

For example, in the compound of Formula (Ib), R₂ and R₃, together with the nitrogen atom to which they are attached, form an unsubstituted pyrrolidinyl ring.

For example, in the compound of Formula (Ib), R₂ and R₃, together with the nitrogen atom to which they are attached, form a substituted or unsubstituted heteroaryl comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S.

For example, in the compound of Formula (Ib), R₂ is substituted or unsubstituted C₆-C₁₀ aryl.

For example, in the compound of Formula (Ib), R₂ is unsubstituted C₆-C₁₀ aryl.

For example, in the compound of Formula (Ib), R₂ is unsubstituted phenyl.

For example, in the compound of Formula (Ib), R₂ is unsubstituted benzyl.


For example, in the compound of Formula (Ib), R₃' is unsubstituted C₁-C₆ alkyl.

For example, in the compound of Formula (Ib), R₃' is unsubstituted C₁-C₃ alkyl.

For example, in the compound of Formula (Ib), R₃' is methyl.

In one embodiment, the present invention provides a compound of Formula (II), or a pharmaceutically acceptable salt, polymorph, hydrate, solvate or co-crystal thereof, or a

method for the treatment of a neurological disease by administering to a subject in need thereof a therapeutically effective amount of a compound of Formula (II), or a pharmaceutically acceptable salt, polymorph, hydrate, solvate or co-crystal thereof:



5 wherein:

R₁ is unsubstituted C₁-C₆ alkyl;

R₄ and R₅ are each, independently, H, substituted or unsubstituted C₁-C₆ alkyl, substituted or unsubstituted C₂-C₆ alkenyl, substituted or unsubstituted C₂-C₆ alkynyl, substituted or unsubstituted C₆-C₁₀ aryl, substituted or unsubstituted C₃-C₁₀ carbocycle, substituted or unsubstituted heterocycle comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S, or substituted or unsubstituted heteroaryl comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S;

R₆, R₇, R₈ and R₉ are each, independently, H, substituted or unsubstituted C₁-C₆ alkyl, substituted or unsubstituted C₂-C₆ alkenyl, substituted or unsubstituted C₂-C₆ alkynyl or C(O)OR_a; and

R_a is H or substituted or unsubstituted C₁-C₆ alkyl.

In one embodiment of Formula (II),

R₁ is methyl;

R₄ and R₅ are each methyl; and

20 R₆, R₇, R₈ and R₉ are each, independently, H or methyl.

For example, the neurological disease is multiple sclerosis.

For example, the neurological disease is relapsing-remitting multiple sclerosis (RRMS).

For example, in the compound of Formula (II), R₁ is methyl.

25 For example, in the compound of Formula (II), R₁ is ethyl.

For example, in the compound of Formula (II), R₄ is substituted or unsubstituted C₁-C₆ alkyl.

For example, in the compound of Formula (II), R₄ is unsubstituted C₁-C₆ alkyl.

For example, in the compound of Formula (II), R₄ is unsubstituted C₁-C₃ alkyl.

For example, in the compound of Formula (II), R₄ is unsubstituted C₁-C₂ alkyl.

For example, in the compound of Formula (II), R₄ is C(O)OR_a-substituted C₁-C₆ alkyl, wherein R_a is H or unsubstituted C₁-C₆ alkyl.

For example, in the compound of Formula (II), R₄ is S(O)(O)R_b-substituted C₁-C₆ alkyl, wherein R_b is unsubstituted C₁-C₆ alkyl.

For example, in the compound of Formula (II), R₅ is H.

For example, in the compound of Formula (II), R₅ is substituted or unsubstituted C₁-C₆ alkyl.

For example, in the compound of Formula (II), R₅ is unsubstituted C₁-C₆ alkyl.

For example, in the compound of Formula (II), R₄ is substituted or unsubstituted C₆-C₁₀ aryl.

For example, in the compound of Formula (II), R₄ is unsubstituted C₆-C₁₀ aryl.

For example, in the compound of Formula (II), R₄ is unsubstituted phenyl.

For example, in the compound of Formula (II), R₄ is unsubstituted benzyl.

For example, in the compound of Formula (II), R₆, R₇, R₈ and R₉ are each H.

For example, in the compound of Formula (II), R₆ is substituted or unsubstituted C₁-C₆ alkyl and R₇, R₈ and R₉ are each H.

For example, in the compound of Formula (II), R₆ is unsubstituted C₁-C₆ alkyl and R₇, R₈ and R₉ are each H.

For example, in the compound of Formula (II), R₈ is substituted or unsubstituted C₁-C₆ alkyl and R₆, R₇ and R₉ are each H.

For example, in the compound of Formula (II), R₈ is unsubstituted C₁-C₆ alkyl and R₆, R₇ and R₉ are each H.

For example, in the compound of Formula (II), R₆ and R₈ are each, independently, substituted or unsubstituted C₁-C₆ alkyl and R₇ and R₉ are each H.

For example, in the compound of Formula (II), R₆ and R₈ are each, independently, unsubstituted C₁-C₆ alkyl and R₇ and R₉ are each H.


For example, in the compound of Formula (II), R₆ and R₇ are each, independently, substituted or unsubstituted C₁-C₆ alkyl and R₈ and R₉ are each H.

For example, in the compound of Formula (II), R₆ and R₇ are each, independently, unsubstituted C₁-C₆ alkyl and R₈ and R₉ are each H.

For example, in the compound of Formula (II), R₈ and R₉ are each, independently, substituted or unsubstituted C₁-C₆ alkyl and R₆ and R₇ are each H.


For example, in the compound of Formula (II), R_8 and R_9 are each, independently, unsubstituted C_1 - C_6 alkyl and R_6 and R_7 are each H.

In one embodiment, the present invention provides a compound of Formula (III), or a pharmaceutically acceptable salt, polymorph, hydrate, solvate or co-crystal thereof, or a method for the treatment of a neurological disease by administering to a subject in need thereof a therapeutically effective amount of a compound of Formula (III), or a pharmaceutically acceptable salt, polymorph, hydrate, solvate or co-crystal thereof:




wherein:

R_1 is unsubstituted C_1 - C_6 alkyl;



is selected from the group consisting of:



X is N, O, S, or SO_2 ;

Z is C or N;

m is 0, 1, 2, or 3;

n is 1 or 2;

w is 0, 1, 2 or 3;

t is 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10;

R_6 , R_7 , R_8 and R_9 are each, independently, H, substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_2 - C_6 alkenyl, substituted or unsubstituted C_2 - C_6 alkynyl or $C(O)OR_a$; and

R_a is H or substituted or unsubstituted C_1 - C_6 alkyl; and

5 each R_{10} is, independently, H, halogen, substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_2 - C_6 alkenyl, substituted or unsubstituted C_2 - C_6 alkynyl, substituted or unsubstituted C_3 - C_{10} carbocycle, substituted or unsubstituted heterocycle comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S, or substituted or unsubstituted heteroaryl comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S;

or, alternatively, two R_{10} 's attached to the same carbon atom, together with the carbon atom to which they are attached, form a carbonyl, substituted or unsubstituted C_3 - C_{10} carbocycle, substituted or unsubstituted heterocycle comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S, or substituted or unsubstituted heteroaryl comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S;


or, alternatively, two R_{10} 's attached to different atoms, together with the atoms to which they are attached, form a substituted or unsubstituted C_3 - C_{10} carbocycle, substituted or unsubstituted heterocycle comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S, or substituted or unsubstituted heteroaryl comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S.




For example, the neurological disease is multiple sclerosis.

For example, the neurological disease is relapsing-remitting multiple sclerosis (RRMS).

25 For example, in the compound of Formula (III), R_1 is methyl.

For example, in the compound of Formula (III), R_1 is ethyl.





For example, in the compound of Formula (III), R_6 is substituted or unsubstituted C_1 - C_6 alkyl and R_7 , R_8 and R_9 are each H.

For example, in the compound of Formula (III), R_6 is unsubstituted C_1 - C_6 alkyl and
10 R_7 , R_8 and R_9 are each H.

For example, in the compound of Formula (III), R_8 is substituted or unsubstituted C_1 - C_6 alkyl and R_6 , R_7 and R_9 are each H.

For example, in the compound of Formula (III), R_8 is unsubstituted C_1 - C_6 alkyl and
15 R_6 , R_7 and R_9 are each H.

For example, in the compound of Formula (III), R_6 and R_8 are each, independently,
substituted or unsubstituted C_1 - C_6 alkyl and R_7 and R_9 are each H.

For example, in the compound of Formula (III), R_6 and R_8 are each, independently, unsubstituted C_1 - C_6 alkyl and R_7 and R_9 are each H.

For example, in the compound of Formula (III), R_6 and R_7 are each, independently, substituted or unsubstituted C_1 - C_6 alkyl and R_8 and R_9 are each H.


5 For example, in the compound of Formula (III), R_6 and R_7 are each, independently, unsubstituted C_1 - C_6 alkyl and R_8 and R_9 are each H.

For example, in the compound of Formula (III), R_8 and R_9 are each, independently, substituted or unsubstituted C_1 - C_6 alkyl, and R_6 and R_7 are each H.

10 For example, in the compound of Formula (III), R_8 and R_9 are each, independently, unsubstituted C_1 - C_6 alkyl, and R_6 and R_7 are each H.

In one embodiment of Formula (III):

R_1 is unsubstituted C_1 - C_6 alkyl;



15 m is 0, 1, 2, or 3;


t is 2, 4, or 6;

R_6 , R_7 , R_8 and R_9 are each, independently, H, unsubstituted C_1 - C_6 alkyl, or $C(O)OR_a$, wherein R_a is H or unsubstituted C_1 - C_6 alkyl; and

20 two R_{10} 's attached to the same carbon atom, together with the carbon atom to which they are attached, form a carbonyl.

In another embodiment, the present invention provides a compound of Formula (IV), or a pharmaceutically acceptable salt, polymorph, hydrate, solvate or co-crystal thereof, or a method for the treatment of a neurological disease by administering to a subject in need thereof, a therapeutically effective amount of a compound of Formula (IV), or a

25 pharmaceutically acceptable salt, polymorph, hydrate, solvate or co-crystal thereof:



wherein:

R₁ is unsubstituted C₁-C₆ alkyl;

L_a is substituted or unsubstituted C₁-C₆ alkyl linker;

- 5 R₂ and R₃ are each, independently, H, substituted or unsubstituted acyl, NR₁₄R₁₅, C(S)R₁₁, C(S)SR₁₁, C(S)NR₁₁R₁₂, C(S)NR₁₁NR₁₃R₁₄, C(NR₁₃)NR₁₁R₁₂, substituted or unsubstituted C₁-C₆ alkyl, substituted or unsubstituted C₂-C₆ alkenyl, substituted or unsubstituted C₂-C₆ alkynyl, substituted or unsubstituted C₆-C₁₀ aryl, substituted or unsubstituted C₃-C₁₀ carbocycle, substituted or unsubstituted heterocycle comprising one or
10 two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S, or substituted or unsubstituted heteroaryl comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S;

- R₁₁ and R₁₂ are each, independently, H, substituted or unsubstituted C₁-C₆ alkyl, substituted or unsubstituted C₂-C₆ alkenyl, substituted or unsubstituted C₂-C₆ alkynyl,
15 substituted or unsubstituted C₆-C₁₀ aryl, substituted or unsubstituted C₃-C₁₀ carbocycle, substituted or unsubstituted heterocycle comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S, or substituted or unsubstituted heteroaryl comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S;

 R₁₃ is H or substituted or unsubstituted C₁-C₆ alkyl; and

- 20 R₁₄ and R₁₅ are each, independently, H, substituted or unsubstituted acyl, substituted or unsubstituted C₁-C₆ alkyl, substituted or unsubstituted C₂-C₆ alkenyl, substituted or unsubstituted C₂-C₆ alkynyl, substituted or unsubstituted C₆-C₁₀ aryl, substituted or unsubstituted C₃-C₁₀ carbocycle, substituted or unsubstituted heterocycle comprising one or
25 two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S, or substituted or unsubstituted heteroaryl comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S;

 wherein at least one of R₂ and R₃ is substituted or unsubstituted acyl, NR₁₄R₁₅, C(S)R₁₁, C(S)SR₁₁, C(S)NR₁₁R₁₂, C(S)NR₁₁NR₁₃R₁₄, or C(NR₁₃)NR₁₁R₁₂.

 In one embodiment of Formula (IV),

R_1 is C_1 - C_6 alkyl;

L_a is substituted or unsubstituted C_1 - C_4 alkyl linker; and

one of R_2 and R_3 is $CO_2(C_1$ - C_6 alkyl), CO_2CH_2Ph , CO_2Ph , CO_2Py , pyridinyl-*N*-oxide ester, $C(O)CH_2$ (imidazole), $C(S)NHPh$, or $C(NH)NH_2$, wherein Ph or imidazole groups are

5 optionally substituted with NO_2 .

In another embodiment of Formula (IV),

R_1 is C_1 - C_4 alkyl;

L_a is substituted or unsubstituted C_1 - C_4 alkyl linker; and

10 R_2 and R_3 are each, independently, H, methyl, ethyl, isopropyl, butyl, *tert*-butyl, cyclohexyl, cyclohexenyl, phenyl, benzyl, benzodioxole, pyridinyl, $(CH_2)_2N(CH_3)_2$, $(CH_2)_3SO_2H$, $(CH_2)_2SO_2Me$, CHO, CH_2CO_2H , $C(O)(CH_2)_2CO_2H$, NO, $C(O)NH_2$, $(CH_2)_2CN$, *tert*-butyl ester, benzyl ester, pyridinyl ester, pyridinyl-*N*-oxide ester, $C(O)CH_2$ (2-nitro-1*H*-imidazol-1-yl), $C(S)NHPh$, $C(NH)NH_2$, ethyl substituted with carbonyl, propyl substituted with carbonyl, or phenyl ester substituted with NO_2 , wherein the phenyl and benzyl groups
15 can be optionally substituted one or more times with methyl, NH_2 , NO_2 , OH, or CHO;

wherein at least one of R_2 and R_3 is substituted or unsubstituted acyl, $NR_{14}R_{15}$, $C(S)R_{11}$, $C(S)SR_{11}$, $C(S)NR_{11}R_{12}$, $C(S)NR_{11}NR_{13}R_{14}$, or $C(NR_{13})NR_{11}R_{12}$.

In one embodiment of Formula (IV),

R_1 is C_1 - C_6 alkyl;

20 L_a is $(CH_2)_{1-4}$;

R_2 is H or $C(O)C_1$ - C_6 alkyl; and



R_3 is H or $C(O)C_1$ - C_6 alkyl












wherein at least one of R_2 and R_3 is $C(O)C_{1-6}$ alkyl.












For example, the compound is a compound listed in Table 1 herein.











25 Representative compounds of the present invention include compounds listed in Table 1 and in Table 2.











Table 1.












1	
2	











3	
4	 A ⁻
5	
6	
7	
8	
9	
10	
11	
12	
13	












14	
15	
16	
17	
18	
19	
20	
21	
22	
23	
24	












25	
26	
27	
28	
29	
30	
31	
32	
33	
34	











35	
36	
37	
38	
39	
40	
41	
42	
43	
44	













45	
46	
47	
48	
49	
50	
51	
52	
53	
54	
55	












56	
57	
58	
59	
60	
61	
62	
63	
64	
65	










66	
67	
68	
69	
70	
71	
72	
73	
74	
75	
76	

77	
78	
79	
80	
81	
82	
83	
84	
85	
86	
87	

88	
89	
90	
91	
92	
93	
94	
95	
96	
97	


98	
99	
100	
101	
102	
103	
104	
105	
106	
107	
108	
109	




110	
111	
112	
113	
114	
115	
116	
117	
118	
119	
120	

121	
122	
123	
124	
125	
126	
127	
128	
129	

A⁻ is a pharmaceutically acceptable anion.

Table 2.

130	
-----	---

131	
132	
133	

The present invention also provides pharmaceutical compositions comprising one or more compounds of Formula (I), (Ia), (Ib), (II), (III), or (IV) and one or more pharmaceutically acceptable carriers.

- 5 In one embodiment, the pharmaceutical composition is a controlled release composition comprising a compound of Formula (I), (Ia), (Ib), (II), (III), or (IV) and one or more pharmaceutically acceptable carriers, wherein the controlled release composition provides a therapeutically effective amount of monomethyl fumarate to a subject. In another embodiment, the pharmaceutical composition is a controlled release composition comprising
- 10 a compound of Formula (I), (Ia), (Ib), (II), (III), or (IV) and one or more pharmaceutically acceptable carriers, wherein the controlled release composition provides a therapeutically effective amount of monomethyl fumarate to a subject for at least about 8 hours to at least about 24 hours. In another embodiment, the pharmaceutical composition is a controlled release composition comprising a compound of Formula (I), (Ia), (Ib), (II), (III), or (IV) and
- 15 one or more pharmaceutically acceptable carriers, wherein the controlled release composition provides a therapeutically effective amount of monomethyl fumarate to a subject for at least about 8 hours, at least about 10 hours, at least about 12 hours, at least about 13 hours, at least about 14 hours, at least about 15 hours, at least about 16 hours, at least about 17 hours, at least about 18 hours, at least about 19 hours, at least about 20 hours, at least about 21 hours,
- 20 at least about 22 hours, at least about 23 hours or at least about 24 hours or longer. For example, at least about 18 hours. For example, at least about 12 hours. For example, greater than 12 hours. For example, at least about 16 hours. For example, at least about 20 hours. For example, at least about 24 hours.

In another embodiment, a compound of Formula (I), (Ia), (Ib), (II), (III), or (IV) is efficiently converted to the active species, *i.e.*, monomethyl fumarate, upon oral administration. For example, about 50 mole percent, about 55 mole percent, about 60 mole percent, about 65 mole percent, about 70 mole percent, about 75 mole percent, about 80 mole percent, about 85 mole percent, about 90 mole percent, or greater than 90 mole percent of the total dose of a compound of Formula (I), (Ia), (Ib), (II), (III), or (IV) administered is converted to monomethyl fumarate upon oral administration. In another embodiment, a compound of Formula (I), (Ia), (Ib), (II), (III), or (IV) is converted to the active species, *i.e.*, monomethyl fumarate, upon oral administration more efficiently than dimethyl fumarate. In another embodiment, a compound of Formula (I), (Ia), (Ib), (II), (III), or (IV) is converted to the active species, *i.e.*, monomethyl fumarate, upon oral administration more efficiently than one or more of the compounds described in US 8,148,414. For example, a compound of Formula (I), (Ia), (Ib), (II), (III), or (IV) is essentially completely converted to the active species, *i.e.*, monomethyl fumarate, upon oral administration. US 8,148,414 is expressly incorporated by reference herein.

In another embodiment, any one of Compounds 1 – 133 is efficiently converted to the active species, *i.e.*, monomethyl fumarate, upon oral administration. For example, about 50 percent, about 55 percent, about 60 percent, about 65 percent, about 70 percent, about 75 percent, about 80 percent, about 85 percent, about 90 percent, or greater than 90 percent of the total dose of any one of Compounds 1 – 133 administered is converted to monomethyl fumarate upon oral administration. In another embodiment, any one of Compounds 1 – 133 is converted to the active species, *i.e.*, monomethyl fumarate, upon oral administration more efficiently than dimethyl fumarate. In another embodiment, any one of Compounds 1 – 133 is converted to the active species, *i.e.*, monomethyl fumarate, upon oral administration more efficiently than one or more of the compounds described in US 8,148,414. For example, any one of Compounds 1 – 133 is completely converted to the active species, *i.e.*, monomethyl fumarate, upon oral administration.

For a drug to achieve its therapeutic effect, it is necessary to maintain the required level of blood or plasma concentration. Many drugs, including dimethyl fumarate, must be administered multiple times a day to maintain the required concentration. Furthermore, even with multiple administrations of such a drug per day, the blood or plasma concentrations of the active ingredient may still vary with time, *i.e.*, at certain time points between administrations there are higher concentrations of the active ingredient than at other times. Thus, at certain time points of a 24-hour period, a patient may receive therapeutically

effective amounts of the active ingredient, while at other time points the concentration of the active ingredient in the blood may fall below therapeutic levels. Additional problems with such drugs include that multiple dosing a day often adversely affects patient compliance with the treatment. Therefore, it is desirable to have a drug dosage form wherein the active ingredient is delivered in such a controlled manner that a constant or substantially constant level of blood or plasma concentration of the active ingredient can be achieved by one or at most two dosing per day. Accordingly, the present invention provides controlled-release formulations as described below. In general, such formulations are known to those skilled in the art or are available using conventional methods.

As used herein, "controlled-release" means a dosage form in which the release of the active agent is controlled or modified over a period of time. Controlled can mean, for example, sustained, delayed or pulsed-release at a particular time. For example, controlled-release can mean that the release of the active ingredient is extended for longer than it would be in an immediate-release dosage form, i.e., at least over several hours.

As used herein, "immediate-release" means a dosage form in which greater than or equal to about 75% of the active ingredient is released within two hours, or, more specifically, within one hour, of administration. Immediate-release or controlled-release may also be characterized by their dissolution profiles.

Formulations may also be characterized by their pharmacokinetic parameters. As used herein, "pharmacokinetic parameters" describe the in vivo characteristics of the active ingredient over time, including for example plasma concentration of the active ingredient. As used herein, " $C_{\max}$ " means the measured concentration of the active ingredient in the plasma at the point of maximum concentration. " $T_{\max}$ " refers to the time at which the concentration of the active ingredient in the plasma is the highest. "AUC" is the area under the curve of a graph of the concentration of the active ingredient (typically plasma concentration) vs. time, measured from one time to another.

The controlled-release formulations provided herein provide desirable properties and advantages. For example, the formulations can be administered once daily, which is particularly desirable for the subjects described herein. The formulation can provide many therapeutic benefits that are not achieved with corresponding shorter acting, or immediate-release preparations. For example, the formulation can maintain lower, more steady plasma peak values, for example, $C_{\max}$, so as to reduce the incidence and severity of possible side effects.

Sustained-release dosage forms release their active ingredient into the gastro-intestinal tract of a patient over a sustained period of time following administration of the dosage form to the patient. Particular dosage forms include: (a) those in which the active ingredient is embedded in a matrix from which it is released by diffusion or erosion; (b) those
5 in which the active ingredient is present in a core which is coated with a release rate-controlling membrane; (c) those in which the active ingredient is present in a core provided with an outer coating impermeable to the active ingredient, the outer coating having an aperture (which may be drilled) for release of the active ingredient; (d) those in which the active ingredient is released through a semi-permeable membrane, allowing the drug to
10 diffuse across the membrane or through liquid filled pores within the membrane; and (e) those in which the active ingredient is present as an ion exchange complex.

It will be apparent to those skilled in the art that some of the above means of achieving sustained-release may be combined, for example a matrix containing the active compound may be formed into a multiparticulate and/or coated with an impermeable coating
15 provided with an aperture.

Pulsed-release formulations release the active compound after a sustained period of time following administration of the dosage form to the patient. The release may then be in the form of immediate- or sustained-release. This delay may be achieved by releasing the drug at particular points in the gastro-intestinal tract or by releasing drug after a pre-
20 determined time. Pulsed-release formulations may be in the form of tablets or multiparticulates or a combination of both. Particular dosage forms include: (a) osmotic potential triggered release (see U.S. Pat. No. 3,952,741); (b) compression coated two layer tablets (see U.S. Pat. No. 5,464,633); (c) capsules containing an erodible plug (see U.S. Pat. No. 5,474,784); sigmoidal releasing pellets (referred to in U.S. Pat. No. 5,112,621); and (d)
25 formulations coated with or containing pH-dependent polymers including shellac, phthalate derivatives, polyacrylic acid derivatives and crotonic acid copolymers.

Dual release formulations can combine the active ingredient in immediate release form with additional active ingredient in controlled-release form. For example, a bilayer tablet can be formed with one layer containing immediate release active ingredient and the
30 other layer containing the active ingredient embedded in a matrix from which it is released by diffusion or erosion. Alternatively, one or more immediate release beads can be combined with one or more beads which are coated with a release rate-controlling membrane in a capsule to give a dual release formulation. Sustained release formulations in which the active ingredient is present in a core provided with an outer coating impermeable to the active

ingredient, the outer coating having an aperture (which may be drilled) for release of the active ingredient, can be coated with drug in immediate release form to give a dual release formulation. Dual release formulations can also combine drug in immediate release form with additional drug in pulsed release form. For example, a capsule containing an erodible
5 plug could liberate drug initially and, after a predetermined period of time, release additional drug in immediate- or sustained-release form.

In some embodiments, the dosage forms to be used can be provided as controlled-release with respect to one or more active ingredients therein using, for example, hydroxypropylmethyl cellulose, other polymer matrices, gels, permeable membranes, osmotic
10 systems, multilayer coatings, microparticles, liposomes, or microspheres or a combination thereof to provide the desired release profile in varying proportions. Suitable controlled-release formulations known to those of ordinary skill in the art, including those described herein, can be readily selected for use with the pharmaceutical compositions of the invention. Thus, single unit dosage forms suitable for oral administration, such as tablets, capsules,
15 gelcaps, and caplets that are adapted for controlled-release are encompassed by the present invention.

Most controlled-release formulations are designed to initially release an amount of drug that promptly produces the desired therapeutic effect, and gradually and continually release of additional amounts of drug to maintain this level of therapeutic effect over an
20 extended period of time. In order to maintain this constant level of drug in the body, the drug must be released from the dosage form at a rate that will replace the amount of drug being metabolized and excreted from the body.

Controlled-release of an active ingredient can be stimulated by various inducers, for example pH, temperature, enzymes, concentration, or other physiological conditions or
25 compounds.

Powdered and granular formulations of a pharmaceutical preparation of the invention may be prepared using known methods. Such formulations may be administered directly to a subject, used, for example, to form tablets, to fill capsules, or to prepare an aqueous or oily suspension or solution by addition of an aqueous or oily vehicle thereto. Each of these
30 formulations may further comprise one or more of a dispersing agent, wetting agent, suspending agent, and a preservative. Additional excipients, such as fillers, sweeteners, flavoring, or coloring agents, may also be included in these formulations.

A formulation of a pharmaceutical composition of the invention suitable for oral administration may be prepared or packaged in the form of a discrete solid dose unit

including, but not limited to, a tablet, a hard or soft capsule, a cachet, a troche, or a lozenge, each containing a predetermined amount of the active ingredient. In one embodiment, a formulation of a pharmaceutical composition of the invention suitable for oral administration is coated with an enteric coat.

5 A tablet comprising the active ingredient may, for example, be made by compressing or molding the active ingredient, optionally with one or more additional ingredients. Compressed tablets may be prepared by compressing, in a suitable device, the active ingredient in a free flowing form such as a powder or granular preparation, optionally mixed with one or more of a binder, a lubricant, an excipient, a surface-active agent, and a
10 dispersing agent. Molded tablets may be made by molding, in a suitable device, a mixture of the active ingredient, a pharmaceutically acceptable carrier, and at least sufficient liquid to moisten the mixture. Pharmaceutically acceptable excipients used in the manufacture of tablets include, but are not limited to, inert diluents, granulating and disintegrating agents, binding agents, and lubricating agents. Known dispersing agents include, but are not limited
15 to, potato starch and sodium starch glycolate. Known surface-active agents include, but are not limited to, sodium lauryl sulphate and poloxamers. Known diluents include, but are not limited to, calcium carbonate, sodium carbonate, lactose, microcrystalline cellulose, calcium phosphate, calcium hydrogen phosphate, and sodium phosphate. Known granulating and disintegrating agents include, but are not limited to, corn starch and alginic acid. Known
20 binding agents include, but are not limited to, gelatin, acacia, pre-gelatinized maize starch, polyvinylpyrrolidone, and hydroxypropyl methylcellulose. Known lubricating agents include, but are not limited to, magnesium stearate, stearic acid, silica, and talc.

Tablets may be non-coated or they may be coated using known methods to achieve delayed disintegration in the gastrointestinal tract of a subject, thereby providing sustained
25 release and absorption of the active ingredient. By way of example, a material such as glyceryl monostearate or glyceryl distearate may be used to coat tablets. Further by way of example, tablets may be coated using methods described in U.S. Pat. Nos. 4,256,108; 4,160,452; and 4,265,874 to form osmotically-controlled release tablets, optionally, with laser drilling. Tablets may further comprise a sweetener, a flavoring agent, a coloring agent, a
30 preservative, or some combination of these in order to provide for pharmaceutically elegant and palatable formulations.

Hard capsules comprising the active ingredient may be made using a physiologically degradable composition, such as gelatin or HPMC. Such hard capsules comprise the active

ingredient, and may further comprise additional ingredients including, for example, an inert solid diluent such as calcium carbonate, calcium phosphate, or kaolin.

Soft gelatin capsules comprising the active ingredient may be made using a physiologically degradable composition, such as gelatin. Such soft capsules comprise the active ingredient, which may be mixed with water or an oil medium such as peanut oil, liquid paraffin, or olive oil.

As used herein, “alkyl”, “C₁, C₂, C₃, C₄, C₅ or C₆ alkyl” or “C₁-C₆ alkyl” is intended to include C₁, C₂, C₃, C₄, C₅ or C₆ straight chain (linear) saturated aliphatic hydrocarbon groups and C₃, C₄, C₅ or C₆ branched saturated aliphatic hydrocarbon groups. For example, C₁-C₆ alkyl is intended to include C₁, C₂, C₃, C₄, C₅ and C₆ alkyl groups. Examples of alkyl include, moieties having from one to six carbon atoms, such as, but not limited to, methyl, ethyl, n-propyl, i-propyl, n-butyl, s-butyl, t-butyl, n-pentyl, s-pentyl, or n-hexyl.

In certain embodiments, a straight chain or branched alkyl has six or fewer carbon atoms (*e.g.*, C₁-C₆ for straight chain, C₃-C₆ for branched chain), and in another embodiment, a straight chain or branched alkyl has four or fewer carbon atoms.

As used herein, “alkyl linker” is intended to include C₁, C₂, C₃, C₄, C₅, or C₆ straight chain (linear) saturated aliphatic hydrocarbon groups and C₃, C₄, C₅, or C₆ branched saturated aliphatic hydrocarbon groups. For example, C₁-C₆ alkyl linker is intended to include C₁, C₂, C₃, C₄, C₅, and C₆ alkyl linker groups. Examples of alkyl linker include, moieties having from one to six carbon atoms, such as, but not limited to, methyl (-CH₂-), ethyl (-CH₂CH₂-), n-propyl (-CH₂CH₂CH₂-), i-propyl (-CHCH₃CH₂-), n-butyl (-CH₂CH₂CH₂CH₂-), s-butyl (-CHCH₃CH₂CH₂-), i-butyl (-C(CH₃)₂CH₂-), n-pentyl (-CH₂CH₂CH₂CH₂CH₂-), s-pentyl (-CHCH₃CH₂CH₂CH₂-) or n-hexyl (-CH₂CH₂CH₂CH₂CH₂CH₂-). The term “substituted alkyl linker” refers to alkyl linkers having substituents replacing one or more hydrogen atoms on one or more carbons of the hydrocarbon backbone. Such substituents do not alter the sp³-hybridization of the carbon atom to which they are attached and include those listed below for “substituted alkyl.”

“Heteroalkyl” groups are alkyl groups, as defined above, that have an oxygen, nitrogen, sulfur or phosphorous atom replacing one or more hydrocarbon backbone carbon atoms.

As used herein, the term “cycloalkyl”, “C₃, C₄, C₅, C₆, C₇ or C₈ cycloalkyl” or “C₃-C₈ cycloalkyl” is intended to include hydrocarbon rings having from three to eight carbon atoms in their ring structure. In one embodiment, a cycloalkyl group has five or six carbons in the ring structure.

The term “substituted alkyl” refers to alkyl moieties having substituents replacing one or more hydrogen atoms on one or more carbons of the hydrocarbon backbone. Such substituents can include, for example, alkyl, alkenyl, alkynyl, halogen, hydroxyl, alkylcarbonyloxy, arylcarbonyloxy, alkoxycarbonyloxy, aryloxy carbonyloxy, carboxylate, 5 alkylcarbonyl, arylcarbonyl, alkoxycarbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylthiocarbonyl, alkoxyl, phosphate, phosphonato, phosphinato, amino (including alkylamino, dialkylamino, arylamino, diarylamino, and alkylaryl amino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl, and ureido), amidino, imino, sulfhydryl, alkylthio, arylthio, thiocarboxylate, sulfates, alkylsulfinyl, 10 sulfonato, sulfamoyl, sulfonamido, nitro, trifluoromethyl, cyano, azido, heterocyclyl, alkylaryl, or an aromatic or heteroaromatic moiety. Cycloalkyls can be further substituted, *e.g.*, with the substituents described above. An “alkylaryl” or an “aralkyl” moiety is an alkyl substituted with an aryl (*e.g.*, phenylmethyl(benzyl)).

Unless the number of carbons is otherwise specified, “lower alkyl” includes an alkyl 15 group, as defined above, having from one to six, or in another embodiment from one to four, carbon atoms in its backbone structure. “Lower alkenyl” and “lower alkynyl” have chain lengths of, for example, two to six or of two to four carbon atoms.

“Aryl” includes groups with aromaticity, including “conjugated”, or multicyclic, systems with at least one aromatic ring. Examples include phenyl, benzyl, naphthyl, etc. 20 “Heteroaryl” groups are aryl groups, as defined above, having from one to four heteroatoms in the ring structure, and may also be referred to as “aryl heterocycles” or “heteroaromatics”. As used herein, the term “heteroaryl” is intended to include a stable 5-, 6-, or 7-membered monocyclic or 7-, 8-, 9-, 10-, 11- or 12-membered bicyclic aromatic heterocyclic ring which consists of carbon atoms and one or more heteroatoms, *e.g.*, 1 or 1-2 or 1-3 or 1-4 or 1-5 or 1- 25 6 heteroatoms, independently selected from the group consisting of nitrogen, oxygen and sulfur. The nitrogen atom may be substituted or unsubstituted (*i.e.*, N or NR wherein R is H or other substituents, as defined). The nitrogen and sulfur heteroatoms may optionally be oxidized (*i.e.*, N→O and S(O)_p, where p=1 or 2). It is to be noted that total number of S and O atoms in the heteroaryl is not more than 1.

30 Examples of heteroaryl groups include pyrrole, furan, thiophene, thiazole, isothiazole, imidazole, triazole, tetrazole, pyrazole, oxazole, isoxazole, pyridine, pyrazine, pyridazine, pyrimidine, and the like.

As used herein, “Ph” refers to phenyl, and “Py” refers to pyridinyl.

Furthermore, the terms “aryl” and “heteroaryl” include multicyclic aryl and heteroaryl groups, *e.g.*, tricyclic, bicyclic, *e.g.*, naphthalene, benzoxazole, benzodioxazole, benzothiazole, benzoimidazole, benzothiophene, methylenedioxyphenyl, quinoline, isoquinoline, naphthrydine, indole, benzofuran, purine, benzofuran, deazapurine, or
5 indolizine.

In the case of multicyclic aromatic rings, only one of the rings needs to be aromatic (*e.g.*, 2,3-dihydroindole), although all of the rings may be aromatic (*e.g.*, quinoline). The second ring can also be fused or bridged.

The aryl or heteroaryl aromatic ring can be substituted at one or more ring positions
10 with such substituents as described above, for example, alkyl, alkenyl, alkynyl, halogen, hydroxyl, alkoxy, alkylcarbonyloxy, arylcarbonyloxy, alkoxy carbonyloxy, aryloxy carbonyloxy, carboxylate, alkylcarbonyl, alkylaminocarbonyl, aralkylaminocarbonyl, alkenylaminocarbonyl, alkylcarbonyl, arylcarbonyl, aralkylcarbonyl, alkenylcarbonyl, alkoxy carbonyl, aminocarbonyl, alkylthiocarbonyl, phosphate, phosphonate, phosphinato,
15 amino (including alkylamino, dialkylamino, arylamino, diarylamino, and alkylaryl amino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl, and ureido), amidino, imino, sulfhydryl, alkylthio, arylthio, thiocarboxylate, sulfates, alkylsulfinyl, sulfonate, sulfamoyl, sulfonamido, nitro, trifluoromethyl, cyano, azido, heterocyclyl, alkylaryl, or an aromatic or heteroaromatic moiety. Aryl groups can also be fused or bridged
20 with alicyclic or heterocyclic rings, which are not aromatic so as to form a multicyclic system (*e.g.*, tetralin, methylenedioxyphenyl).

As used herein, “carbocycle” or “carbocyclic ring” is intended to include any stable monocyclic, bicyclic or tricyclic ring having the specified number of carbons, any of which may be saturated, unsaturated, or aromatic. For example, a C₃-C₁₄ carbocycle is intended to
25 include a monocyclic, bicyclic or tricyclic ring having 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13 or 14 carbon atoms. Examples of carbocycles include, but are not limited to, cyclopropyl, cyclobutyl, cyclobutenyl, cyclopentyl, cyclopentenyl, cyclohexyl, cycloheptenyl, cycloheptyl, cycloheptenyl, adamantyl, cyclooctyl, cyclooctenyl, cyclooctadienyl, fluorenyl, phenyl, naphthyl, indanyl, adamantyl, and tetrahydronaphthyl. Bridged rings are also included in the
30 definition of carbocycle, including, for example, [3.3.0]bicyclooctane, [4.3.0]bicyclononane, [4.4.0]bicyclodecane and [2.2.2]bicyclooctane. A bridged ring occurs when one or more carbon atoms link two non-adjacent carbon atoms. In one embodiment, bridge rings are one or two carbon atoms. It is noted that a bridge always converts a monocyclic ring into a tricyclic ring. When a ring is bridged, the substituents recited for the ring may also be

present on the bridge. Fused (e.g., naphthyl, tetrahydronaphthyl) and spiro rings are also included.

As used herein, “heterocycle” includes any ring structure (saturated or partially unsaturated) which contains at least one ring heteroatom (e.g., N, O or S). Examples of heterocycles include, but are not limited to, morpholine, pyrrolidine, tetrahydrothiophene, piperidine, piperazine, and tetrahydrofuran.

Examples of heterocyclic groups include, but are not limited to, acridinyl, azocinyl, benzimidazolyl, benzofuranyl, benzothiofuranyl, benzothiophenyl, benzoxazolyl, benzoxazoliny, benzthiazolyl, benztriazolyl, benztetrazolyl, benzisoxazolyl, benzisothiazolyl, benzimidazoliny, carbazolyl, 4aH-carbazolyl, carboliny, chromanyl, chromenyl, cinnoliny, decahydroquinoliny, 2H,6H-1,5,2-dithiazinyl, dihydrofuro[2,3-b]tetrahydrofuran, furanyl, furazanyl, imidazolidinyl, imidazoliny, imidazolyl, 1H-indazolyl, indolenyl, indoliny, indoliziny, indolyl, 3H-indolyl, isatinoyl, isobenzofuranyl, isochromanyl, isoindazolyl, isoindoliny, isoindolyl, isoquinoliny, isothiazolyl, isoxazolyl, methylenedioxyphenyl, morpholiny, naphthyridiny, octahydroisoquinoliny, oxadiazolyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, 1,2,4-oxadiazol5(4H)-one, oxazolidinyl, oxazolyl, oxindolyl, pyrimidinyl, phenanthridiny, phenanthroliny, phenaziny, phenothiaziny, phenoxathiny, phenoxaziny, phthalaziny, piperaziny, piperidinyl, piperidonyl, 4-piperidonyl, piperonyl, pteridinyl, puriny, pyranly, pyraziny, pyrazolidiny, pyrazoliny, pyrazolyl, pyridaziny, pyridooxazole, pyridoimidazole, pyridothiazole, pyridiny, pyridyl, pyrimidinyl, pyrrolidinyl, pyrroliny, 2H-pyrrolyl, pyrrolyl, quinazoliny, quinoliny, 4H-quinoliziny, quinoxaliny, quinuclidiny, tetrahydrofuranyl, tetrahydroisoquinoliny, tetrahydroquinoliny, tetrazolyl, 6H-1,2,5-thiadiaziny, 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,2,5-thiadiazolyl, 1,3,4-thiadiazolyl, thianthrenyl, thiazolyl, thienyl, thienothiazolyl, thienooxazolyl, thienoimidazolyl, thiophenyl, triazinyl, 1,2,3-triazolyl, 1,2,4-triazolyl, 1,2,5-triazolyl, 1,3,4-triazolyl, and xanthenyl.

The term “substituted”, as used herein, means that any one or more hydrogen atoms on the designated atom is replaced with a selection from the indicated groups, provided that the designated atom's normal valency is not exceeded, and that the substitution results in a stable compound. When a substituent is keto (*i.e.*, =O), then 2 hydrogen atoms on the atom are replaced. Keto substituents are not present on aromatic moieties. Ring double bonds, as used herein, are double bonds that are formed between two adjacent ring atoms (*e.g.*, C=C, C=N or N=N). “Stable compound” and “stable structure” are meant to indicate a compound

that is sufficiently robust to survive isolation to a useful degree of purity from a reaction mixture, and formulation into an efficacious therapeutic agent.

The term “acyl”, as used herein, includes moieties that contain the acyl radical (–C(O)–) or a carbonyl group. “Substituted acyl” includes acyl groups where one or more of the hydrogen atoms are replaced by, for example, alkyl groups, alkynyl groups, halogen, hydroxyl, alkylcarbonyloxy, arylcarbonyloxy, alkoxy carbonyloxy, aryloxy carbonyloxy, carboxylate, alkylcarbonyl, arylcarbonyl, alkoxy carbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylthiocarbonyl, alkoxy, phosphate, phosphonato, phosphinato, amino (including alkylamino, dialkylamino, arylamino, diarylamino and alkylaryl amino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido), amidino, imino, sulfhydryl, alkylthio, arylthio, thiocarboxylate, sulfates, alkylsulfinyl, sulfonato, sulfamoyl, sulfonamido, nitro, trifluoromethyl, cyano, azido, heterocyclyl, alkylaryl, or an aromatic or heteroaromatic moiety.

The description of the disclosure herein should be construed in congruity with the laws and principals of chemical bonding. For example, it may be necessary to remove a hydrogen atom in order accommodate a substituent at any given location. Furthermore, it is to be understood that definitions of the variables (*i.e.*, “R groups”), as well as the bond locations of the generic formulae of the invention (*e.g.*, Formulas I, Ia, Ib, II III, and IV), will be consistent with the laws of chemical bonding known in the art. It is also to be understood that all of the compounds of the invention described above will further include bonds between adjacent atoms and/or hydrogens as required to satisfy the valence of each atom. That is, bonds and/or hydrogen atoms are added to provide the following number of total bonds to each of the following types of atoms: carbon: four bonds; nitrogen: three bonds; oxygen: two bonds; and sulfur: two-six bonds.

As used herein, a “subject in need thereof” is a subject having a neurological disease. In one embodiment, a subject in need thereof has multiple sclerosis. A “subject” includes a mammal. The mammal can be *e.g.*, any mammal, *e.g.*, a human, primate, bird, mouse, rat, fowl, dog, cat, cow, horse, goat, camel, sheep or a pig. In one embodiment, the mammal is a human.

The present invention provides methods for the synthesis of the compounds of each of the formulae described herein. The present invention also provides detailed methods for the synthesis of various disclosed compounds of the present invention according to the following schemes and as shown in the Examples.

Throughout the description, where compositions are described as having, including, or comprising specific components, it is contemplated that compositions also consist essentially of, or consist of, the recited components. Similarly, where methods or processes are described as having, including, or comprising specific process steps, the processes also
5 consist essentially of, or consist of, the recited processing steps. Further, it should be understood that the order of steps or order for performing certain actions is immaterial so long as the invention remains operable. Moreover, two or more steps or actions can be conducted simultaneously.

The synthetic processes of the invention can tolerate a wide variety of functional
10 groups; therefore various substituted starting materials can be used. The processes generally provide the desired final compound at or near the end of the overall process, although it may be desirable in certain instances to further convert the compound to a pharmaceutically acceptable salt, polymorph, hydrate, solvate or co-crystal thereof.

Compounds of the present invention can be prepared in a variety of ways using
15 commercially available starting materials, compounds known in the literature, or from readily prepared intermediates, by employing standard synthetic methods and procedures either known to those skilled in the art, or which will be apparent to the skilled artisan in light of the teachings herein. Standard synthetic methods and procedures for the preparation of organic molecules and functional group transformations and manipulations can be obtained from the
20 relevant scientific literature or from standard textbooks in the field. Although not limited to any one or several sources, classic texts such as Smith, M. B., March, J., March's Advanced Organic Chemistry Reactions, Mechanisms, and Structure, 5th edition, John Wiley & Sons: New York, 2001; and Greene, T. W., Wuts, P. G. M., Protective Groups in Organic
Synthesis, 3rd edition, John Wiley & Sons: New York, 1999, incorporated by reference
25 herein, are useful and recognized reference textbooks of organic synthesis known to those in the art. The following descriptions of synthetic methods are designed to illustrate, but not to limit, general procedures for the preparation of compounds of the present invention.

Compounds of the present invention can be conveniently prepared by a variety of methods familiar to those skilled in the art. The compounds of this invention with each of the
30 formulae described herein may be prepared according to the following procedures from commercially available starting materials or starting materials which can be prepared using literature procedures. These procedures show the preparation of representative compounds of this invention.

EXPERIMENTAL

General Procedure 1

To a mixture of monomethyl fumarate (MMF) (1.0 equivalent) and HBTU (1.5 equivalents) in DMF (25 ml per g of MMF) was added Hünigs base (2.0 equivalents). The dark brown solution was stirred for 10 minutes, where turned into a brown suspension, before addition of the alcohol (1.0 – 1.5 equivalents). The reaction was stirred for 18 hours at room temperature. Water was added and the product extracted into ethyl acetate three times. The combined organic layers were washed with water three times, dried with magnesium sulphate, filtered and concentrated in vacuo at 45 °C to give the crude product. The crude product was purified by silica chromatography and in some cases further purified by trituration with diethyl ether to give the clean desired ester product. All alcohols were either commercially available or made following known literature procedures.

As an alternative to HBTU (*N,N,N',N'*-Tetramethyl-*O*-(1*H*-benzotriazol-1-yl)uronium hexafluorophosphate), any one of the following coupling reagents can be used: EDCI/HOBt (*N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride/hydroxybenzotriazole hydrate); COMU ((1-cyano-2-ethoxy-2-oxoethylidenaminoxy)dimethylamino-morpholinocarbenium hexafluorophosphate); TBTU (*O*-(benzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium tetrafluoroborate); TATU (*O*-(7-azabenzotriazole-1-yl)-1,1,3,3-tetramethyluronium tetrafluoroborate); Oxyma (ethyl (hydroxyimino)cyanoacetate); PyBOP ((benzotriazol-1-yloxy)tripyrrolidinophosphonium hexafluorophosphate); HOTT (*S*-(1-oxido-2-pyridyl)-*N,N,N',N'*-tetramethylthiuronium hexafluorophosphate); FDPP (pentafluorophenyl diphenylphosphinate); T3P (propylphosphonic anhydride); DMTMM (4-(4,6-dimethoxy-1,3,5-triazin-2-yl)-4-methylmorpholinium tetrafluoroborate); PyOxim ([ethyl cyano(hydroxyimino)acetato-*O*²]tri-1-pyrrolidinylphosphonium hexafluorophosphate); TSTU (*N,N,N',N'*-tetramethyl-*O*-(*N*-succinimidyl)uronium tetrafluoroborate); TDBTU (*O*-(3,4-dihydro-4-oxo-1,2,3-benzotriazin-3-yl)-*N,N,N',N'*-tetramethyluronium tetrafluoroborate); TPTU (*O*-(2-oxo-1(2*H*)pyridyl)-*N,N,N',N'*-tetramethyluronium tetrafluoroborate); TOTU (*O*-[(ethoxycarbonyl)cyanomethylenamino]-*N,N,N',N'*-tetramethyluronium tetrafluoroborate); IIDQ (isobutyl 1,2-dihydro-2-isobutoxy-1-quinolinecarboxylate); or PyCIU (chlorodipyrrolidinocarbenium hexafluorophosphate),

As an alternative to Hünig's base (diisopropylethylamine), any one of the following amine bases can be used: triethylamine; tributylamine; triphenylamine; pyridine; lutidine (2,6-dimethylpyridine); collidine (2,4,6-trimethylpyridine); imidazole; DMAP (4-(dimethylamino)pyridine); DABCO (1,4-diazabicyclo[2.2.2]octane); DBU (1,8-

diazabicyclo[5.4.0]undec-7-ene); DBN (1,5-diazabicyclo[4.3.0]non-5-ene); or proton sponge® (*N,N,N',N'*-tetramethyl-1,8-naphthalenediamine).

General Procedure 2- Conversion of the Ester Product into the Hydrochloride Salt

To a mixture of the ester product in diethyl ether (25 ml per g) was added 2M HCl in diethyl ether (1.5 equivalents). The mixture was stirred at room temperature for two hours. The solvent was decanted, more diethyl ether added and the solvent decanted again. The remaining mixture was then concentrated in vacuo at 45 °C and further dried in a vacuum oven at 55 °C for 18 hours to give the solid HCl salt.

General Procedure 3

To a 100 mL, one-necked, round-bottomed flask, fitted with a magnetic stirrer and nitrogen inlet/outlet, were added 11 mL of an MTBE solution containing freshly prepared mono-methyl fumaryl chloride (4.9 g, 33 mmol) and 50 mL of additional MTBE at 20 °C. The resulting yellow solution was cooled to <20 °C with an ice water bath. Then, the alcohol, (33 mmol, 1 eq) was added dropwise, via syringe, over approximately 10 minutes. The reaction mixture was allowed to stir at <20 °C for 10 minutes after which time the cooling bath was removed and the reaction was allowed to warm to 20 °C and stir at 20 °C temperature for 16 hours. The reaction was deemed complete by TLC after 16 hours at RT. The reaction mixture was filtered through a medium glass fritted funnel to collect the off-white solids. The solids were dried in a vacuum oven at 25 °C overnight to afford the final product as an HCl salt. All alcohols were either commercially available or made following known literature procedures.

General Procedure 4- Alkylation with an Appropriate Alkyl Mesylate

A mixture of monomethyl fumarate (MMF) (1.3 equivalent), the alkyl mesylate (1 equivalent), and potassium carbonate (1.5 equivalent) in acetonitrile (50 ml per g of MMF) was heated at reflux overnight. The mixture was partitioned between ethyl acetate and saturated aqueous sodium hydrogen carbonate, and the organic phase dried (MgSO₄).

Filtration and removal of the solvent under reduced pressure gave the crude product which was purified in each case by silica chromatography.

General Procedure 5- Alkylation with an Appropriate Alkyl Chloride


A mixture of monomethyl fumarate (MMF) (1.3 equivalent), the alkyl chloride (1 equivalent), and potassium carbonate (1.5 equivalent) in acetonitrile or dimethylformamide (50 ml per g of MMF) was heated at 20 to 65 °C overnight. The mixture was partitioned between ethyl acetate and saturated aqueous sodium hydrogen carbonate, and the organic phase dried (MgSO₄). Filtration and removal of the solvent under reduced pressure gave the crude product which was further purified by silica chromatography.

Chemical Analysis/Procedures

The NMR spectra described herein were obtained with a Varian 400 MHz NMR spectrometer using standard techniques known in the art.


ExamplesExample 1

(E)-2,2'-((2-((4-methoxy-4-oxobut-2-enyl)oxy)ethyl)azanediyl)diacetic acid hydrochloride (1)



To a solution of 2-(bis(2-(tert-butoxy)-2-oxoethyl)amino)ethyl methyl fumarate (2.52 g, 6.2 mmol) in dioxane (25 ml) was added 2M HCl in dioxane (30 ml) and the mixture stirred for 90 hours. The precipitate was filtered, washed with diethyl ether and dried in a vacuum oven at 55 °C for 18 hours to give (E)-2,2'-((2-((4-methoxy-4-oxobut-2-enyl)oxy)ethyl)azanediyl)diacetic acid hydrochloride, a white solid (1.31 g, 65 %).


¹H NMR (300 MHz, MeOD): δ 6.87 (2H, dd, J = 16.1 Hz); 4.46-4.53 (2H, m); 4.09 (4H, s); 3.79 (3H, s); 3.57-3.63 (2H, m). [M+H]⁺ = 290.12.

Methyl (2-(methyl(2-(methylsulfonyl)ethyl)amino)ethyl) fumarate hydrochloride (2)

Methyl (2-(N-methylmethylsulfonamido)ethyl) fumarate **2** was synthesized following general procedure 1 and was converted to the HCl salt methyl (2-(methyl(2-(methylsulfonyl)ethyl)amino)ethyl) fumarate hydrochloride (procedure 2) (1.39 g, 95 %).

^1H NMR (400 MHz, DMSO): δ 11.51 (1H, m); 6.83 (2H, dd, J = 15.8 Hz); 4.48 (1H, bs); 3.24-3.90 (7H, m); 3.07 (3H, s); 2.78 (2H, bs). $[\text{M}+\text{H}]^+ = 294.09$.


2-(dimethylamino)propyl methyl fumarate hydrochloride (3)



2-(dimethylamino)propyl methyl fumarate **3** was synthesized following general procedure 1 and was converted to the HCl salt: 2-(dimethylamino)propyl methyl fumarate hydrochloride (procedure 2) (329 mg, 92 %).

^1H NMR (300 MHz, DMSO): δ 10.40 (1H, bs); 6.86 (2H, dd, J = 15.8 Hz); 4.25-4.46 (2H, m); 3.71 (3H, s); 3.34 (1H, s); 2.69 (6H, s); 1.24 (3H, s). $[\text{M}+\text{H}]^+ = 216.14$.


(E)-2-((4-methoxy-4-oxobut-2-enyl)oxy)-N,N,N-trimethylethanaminium iodide (4)



To a solution of 2-(dimethylamino)ethyl methyl fumarate **19** (760 mg, 3.7 mmol) in diethyl ether (20 ml) was added methyl iodide (246 μl , 3.9 mmol). The mixture was stirred at room temperature for 18 hours where a precipitate slowly formed. The mixture was filtered, washed with diethyl ether and dried in a vacuum oven at 55 $^{\circ}\text{C}$ for 18 hours to give (E)-2-((4-methoxy-4-oxobut-2-enyl)oxy)-N,N,N-trimethylethanaminium iodide, a white solid (1.15 g, 90 %).


^1H NMR (300 MHz, DMSO): δ 6.80 (2H, dd, J = 16.1 Hz); 4.56 (2H, bs); 3.66-3.75 (5H, m); 3.11 (9H, s). $[\text{M}+\text{H}]^+ = 216.14$.

2-(4,4-difluoropiperidin-1-yl)ethyl methyl fumarate hydrochloride (5)




2-(4,4-difluoropiperidin-1-yl)ethyl methyl fumarate **5** was synthesized following general procedure 1 and was converted to the HCl salt: 2-(4,4-difluoropiperidin-1-yl)ethyl methyl fumarate hydrochloride (procedure 2) (780 mg, 87 %).

^1H NMR (300 MHz, DMSO): δ 11.25 (1H, bs); 6.84 (2H, dd, J = 16.1 Hz); 4.50 (2H, bs); 3.35-4.00 (8H, m); 3.05-3.30 (2H, m); 2.20-2.45 (3H, s). $[\text{M}+\text{H}]^+ = 278.16$.

1-(dimethylamino)propan-2-yl methyl fumarate hydrochloride (6)


5 1-(dimethylamino)propan-2-yl methyl fumarate **6** was synthesized following general procedure 1 and was converted to the HCl salt 1-(dimethylamino)propan-2-yl methyl fumarate hydrochloride (procedure 2) (690 mg, 72 %).

¹H NMR (300 MHz, DMSO): δ 10.41 (1H, bs); 6.80 (2H, dd, J = 15.8 Hz); 5.18-5.33 (1H, m); 3.20-3.55 (2H, m); 3.72 (3H, s); 2.60-2.80 (7H, m); 1.18-1.28 (3H, m). [M+H]⁺ =
10 216.14.

Methyl (2-thiomorpholinoethyl) fumarate hydrochloride (7)


15 Methyl (2-thiomorpholinoethyl) fumarate **7** was synthesized following general procedure 1 and was converted to the HCl salt, methyl (2-thiomorpholinoethyl) fumarate hydrochloride (procedure 2) (623 mg, 93 %).

¹H NMR (300 MHz, DMSO): δ 11.03 (1H, bs); 6.83 (2H, dd, J = 15.6 Hz); 4.50 (2H, s); 3.00-3.80 (11H, m); 2.70-2.80 (2H, m). [M+H]⁺ = 216.14. [M+H]⁺ = 260.11.

Methyl (2-(phenylamino)ethyl) fumarate hydrochloride (8)


25 Methyl (2-(phenylamino)ethyl) fumarate **8** was synthesized following general procedure 1 and was converted to the HCl salt methyl (2-(phenylamino)ethyl) fumarate hydrochloride (procedure 2) (1.80 g, quantitative).

¹H NMR (300 MHz, DMSO): δ 6.50-6.80 (9H, m); 4.29 (2H, t, 4.4 Hz); 3.72 (3H, s); 3.45 (2H, t, J = 4.5 Hz). [M+H]⁺ = 250.13.

2-(dimethylamino)-2-methylpropyl methyl fumarate hydrochloride (9)


2-(dimethylamino)-2-methylpropyl methyl fumarate **9** was synthesized following
 5 general procedure 1 and was converted to the HCl salt, 2-(dimethylamino)-2-methylpropyl
 methyl fumarate hydrochloride (procedure 2) (883 mg, 76 %).

^1H NMR (300 MHz, DMSO): δ 10.20 (1H, bs); 6.91 (2H, dd, J = 15.6 Hz); 4.29 (2H, s); 3.73 (3H, s); 2.57-2.80 (6H, m); 1.32 (6H, s). $[\text{M}+\text{H}]^+ = 230.16$.

Methyl (2-(methylsulfonyl)ethyl) fumarate (10)


Methyl (2-(methylsulfonyl)ethyl) fumarate **10** was synthesized following general
 procedure 1 and (1.01 g, 37 %).

15 ^1H NMR (400 MHz, CDCl_3): δ 6.88 (2H, dd, J = 16.0 Hz); 4.66 (2H, t, J = 5.8 Hz);
 3.82 (3H, s); 3.38 (2H, t, J = 6.0 Hz); 2.99 (3H, s). $[\text{M}+\text{H}]^+ = 236.97$.

2-(1,1-dioxidothiomorpholino)ethyl methyl fumarate hydrochloride (11)

20 2-(1,1-dioxidothiomorpholino)ethyl methyl fumarate **11** was synthesized following
 general procedure 1 and was converted to the HCl salt 2-(1,1-dioxidothiomorpholino)ethyl
 methyl fumarate hydrochloride (procedure 2) (1.33 g, 87 %).


25 ^1H NMR (400 MHz, DMSO): δ 6.79 (2H, dd, J = 15.8 Hz); 4.34 (2H, bs); 3.72 (4H, s);
 2.90-3.70 (11H, m). $[\text{M}+\text{H}]^+ = 292.11$.

Methyl (2-(methyl(phenyl)amino)ethyl) fumarate hydrochloride (12)

Methyl (2-(methyl(phenyl)amino)ethyl) fumarate **12** was synthesized following general procedure 1 and was converted to the HCl salt methyl (2-(methyl(phenyl)amino)ethyl) fumarate hydrochloride (procedure 2) (1.76 g, 97 %).

5 ^1H NMR (400 MHz, DMSO): δ 6.72-7.40 (5H, m); 6.64 (2H, dd, J = 16.0 Hz); 4.27 (2H, s); 3.70 (5H, s); 2.97 (3H, s). $[\text{M}+\text{H}]^+ = 264.14$.

2-(benzyl(methyl)amino)ethyl methyl fumarate hydrochloride (13)




10

2-(benzyl(methyl)amino)ethyl methyl fumarate **13** was synthesized following general procedure 1 and was converted to the HCl salt 2-(benzyl(methyl)amino)ethyl methyl fumarate hydrochloride (procedure 2) (2.70 g, 96 %).

15 ^1H NMR (400 MHz, DMSO): δ 10.65 (1H, bs); 7.39-7.60 (5H, m); 6.82 (2H, dd, J = 15.8 Hz); 4.20-4.60 (4H, m); 3.73 (3H, s); 3.27-3.50 (2H, m); 2.69 (3H, s). $[\text{M}+\text{H}]^+ = 278.16$.


2-(2,5-dioxopyrrolidin-1-yl)ethyl methyl fumarate (14)



20 2-(2,5-dioxopyrrolidin-1-yl)ethyl methyl fumarate **14** was synthesized following general procedure 1 (1.03 g, 35 %).

^1H NMR (400 MHz, DMSO): δ 6.81 (2H, dd, J = 15.8 Hz); 4.36 (2H, t, J = 5.3 Hz); 3.84 (2H, t, J = 5.1 Hz); 3.80 (3H, s); 2.73 (4H, s). $[\text{M}+\text{H}]^+ = 256.07$.


25 Methyl (2-(piperidin-1-yl)ethyl) fumarate hydrochloride (15)



Methyl (2-(piperidin-1-yl)ethyl) fumarate hydrochloride **15** was synthesized following general procedure 3.

^1H NMR (400 MHz, DMSO- d_6) δ 10.76 (s, 1H), 6.94 – 6.77 (m, 2H), 4.58 – 4.51 (m, 2H), 3.76 (s, 3H), 3.48 – 3.36 (m, 4H), 2.94 (dddd, J = 15.9, 12.1, 9.2, 4.4 Hz, 2H), 1.91 – 1.64 (m, 5H), 1.37 (dt, J = 16.4, 11.3, 4.9 Hz, 1H). $[\text{M}+\text{H}]^+ = 241.93$.


5 Methyl (2-morpholinoethyl) fumarate hydrochloride (16)



Methyl (2-morpholinoethyl) fumarate hydrochloride **16** was synthesized following general procedure 3.

10 ^1H NMR (400 MHz, DMSO- d_6) δ 11.36 (s, 1H), 6.92 (d, J = 15.9 Hz, 1H), 6.82 (d, J = 15.9 Hz, 1H), 4.60 – 4.52 (m, 2H), 4.00 – 3.77 (m, 6H), 3.76 (s, 3H), 3.22 – 3.04 (m, 4H). $[\text{M}+\text{H}]^+ = 244.00$.


15 2-(1,4-dioxo-8-azaspiro[4.5]decan-8-yl)ethyl methyl fumarate hydrochloride (17)



20 2-(1,4-dioxo-8-azaspiro[4.5]decan-8-yl)ethyl methyl fumarate hydrochloride **17** was synthesized following general procedure 3.

^1H NMR (400 MHz, DMSO- d_6) δ 11.26 (s, 1H), 6.91 (d, J = 15.9 Hz, 1H), 6.82 (d, J = 15.9 Hz, 1H), 4.58 – 4.51 (m, 2H), 3.93 (s, 4H), 3.76 (s, 3H), 3.57 – 3.43 (m, 4H), 3.22 – 3.03 (m, 2H), 2.20 – 2.02 (m, 2H), 1.89 – 1.79 (m, 2H). $[\text{M}+\text{H}]^+ = 300.00$.


25 Methyl (2-(pyrrolidin-1-yl)ethyl) fumarate hydrochloride (18)



Methyl (2-(pyrrolidin-1-yl)ethyl) fumarate hydrochloride **18** was synthesized following general procedure 3.

^1H NMR (400 MHz, DMSO- d_6) δ 11.12 (s, 1H), 6.94 (d, J = 15.8 Hz, 1H), 6.82 (d, J = 15.8 Hz, 1H), 4.53 – 4.46 (m, 2H), 3.76 (s, 3H), 3.61 – 3.45 (m, 4H), 3.11 – 2.94 (m, 2H), 2.06 – 1.79 (m, 4H). $[\text{M}+\text{H}]^+ = 228.46$.


5 2-(dimethylamino)ethyl methyl fumarate hydrochloride (19)



2-(dimethylamino)ethyl methyl fumarate hydrochloride **19** was synthesized following general procedure 3.

10 ^1H NMR (500 MHz, DMSO- d_6) δ 10.87 (s, 1H), 6.93 (d, J = 15.9 Hz, 1H), 6.80 (d, J = 15.9 Hz, 1H), 4.53 – 4.45 (m, 2H), 3.75 (s, 3H), 3.44 – 3.38 (m, 2H), 2.77 (s, 5H). $[\text{M}+\text{H}]^+ = 201.84$.

2-(diethylamino)ethyl methyl fumarate hydrochloride (20)




15

2-(diethylamino)ethyl methyl fumarate hydrochloride **20** was synthesized following general procedure 3.

20 ^1H NMR (400 MHz, DMSO- d_6) δ 10.85 (s, 1H), 6.90 (d, J = 15.8 Hz, 1H), 6.81 (d, J = 15.9 Hz, 1H), 4.56 – 4.48 (m, 2H), 3.76 (s, 3H), 3.48 – 3.38 (m, 2H), 3.15 (qq, J = 9.7, 5.5, 4.9 Hz, 4H), 1.24 (t, J = 7.3 Hz, 6H). $[\text{M}+\text{H}]^+ = 230.59$.

2-(3,3-difluoropyrrolidin-1-yl)ethyl methyl fumarate hydrochloride (21)




25

2-(3,3-Difluoropyrrolidin-1-yl)ethyl methyl fumarate **21** was synthesised from 2-(3,3-difluoropyrrolidin-1-yl)ethanol following general procedure 1.

2-(3,3-difluoropyrrolidin-1-yl)ethyl methyl fumarate was converted to 2-(3,3-difluoropyrrolidin-1-yl)ethyl methyl fumarate hydrochloride following general procedure 2 (0.55 g, 69 %).

^1H NMR (300 MHz, DMSO); δ 6.79 (2H, d); 4.20-4.39 (2H, m), 3.81 (2H, t), 3.66 (3H, s), 3.53-3.65 (4H, m), 2.54 (2H, sep). m/z $[\text{M}+\text{H}]^+ = 264.14$.

2-(bis(2-methoxyethyl)amino)ethyl methyl fumarate hydrochloride (24)



5


2-(Bis(2-methoxyethyl)amino)ethyl methyl fumarate **24** was synthesised from 2-(bis(2-methoxyethyl)amino)ethanol following general procedure 1.

2-(Bis(2-methoxyethyl)amino)ethyl methyl fumarate was converted to 2-(bis(2-methoxyethyl)amino)ethyl methyl fumarate hydrochloride following general procedure 2

10 (1.00 g, 27 %).


^1H NMR (300 MHz, DMSO); δ 12.84 (1H, br s), 6.90 (2H, d), 4.73 (2H, t), 3.92 (4H, t), 3.81 (3H, s), 3.62 (2H, br s), 3.51-3.36 (4H, m), 3.34 (6H, s). m/z $[\text{M}+\text{H}]^+ = 290.12$.

2-(2,4-Dioxo-3-azabicyclo[3.1.0]hexan-3-yl)ethyl methyl fumarate (22)




15

3-oxabicyclo[3.1.0]hexane-2,4-dione (1.0 g, 8.9 mmol) and ethanolamine (545 mg, 8.9 mmol) were heated neat at 200 °C for 2 hours. The crude reaction mixture was purified by silica chromatography (EtOAc) giving 3-(2-Hydroxyethyl)-3-azabicyclo[3.1.0]hexane-2,4-dione (1.06 g, 77%).


20 ^1H NMR (300 MHz, CDCl_3): δ 3.71 (2H, t), 3.56 (2H, t), 2.51 (2H, dd), 1.95 (1H, br s), 1.59-1.43 (2H, m).25 2-(2,4-dioxo-3-azabicyclo[3.1.0]hexan-3-yl)ethyl methyl fumarate **22** was synthesised from 3-(2-Hydroxyethyl)-3-azabicyclo[3.1.0]hexane-2,4-dione following general procedure 1 (452 mg, 53 %).

^1H NMR (300 MHz, CDCl_3): δ 6.81 (2H, d), 4.28 (2H, t), 3.80 (3H, s), 3.69 (2H, t), 2.48 (2H, dd), 1.59-1.49 (1H, m), 1.44-1.38 (1H, m). m/z $[\text{M}+\text{H}]^+ = 268.11$.

2-(2,2-Dimethyl-5-oxopyrrolidin-1-yl)ethyl methyl fumarate (24)


Tert-butyl acrylate (19.7 mL, 134.8 mmol) was added dropwise over 10 minutes to a
 5 refluxing solution of 2-nitropropane and Triton B (40% in methanol) (440 μ L) in ethanol (50
 mL). The reaction was heated at reflux overnight. The reaction solvent was removed under
 reduced pressure giving a crude residue that was dissolved in ethanol (200 mL) and
 hydrogenated overnight (300 psi) using Raney nickel (approximately 15 g). The reaction was
 filtered through celite. The solvent was removed under reduced pressure giving tert-butyl 4-
 10 amino-4-methylpentanoate (15.82 g, 63% yield).

^1H NMR (300 MHz, CDCl_3): δ 2.26 (2H, t), 1.65 (2H, t), 1.43 (9H, s), 1.68 (6H, s).



To a solution of tert-butyl 4-amino-4-methylpentanoate (3.0 g, 16.04 mmol) in
 methanol (100 mL) was added chloroacetaldehyde (45% in H_2O) (6.7 mL, 38.4 mmol)
 15 followed by acetic acid (2 mL, 35.0 mmol). After 1.5 hours sodium cyanoborohydride (1.51
 g, 24.0 mmol) was added and the mixture stirred at room temperature for 3 hours. The
 reaction was partitioned between saturated aqueous sodium hydrogen carbonate (100 mL)
 and dichloromethane (300 mL). The organic phase was dried (MgSO_4). Filtration and
 removal of the solvent under reduced pressure gave tert-butyl 4-((2-chloroethyl)amino)-4-
 20 methylpentanoate (3.90 g, 98% yield).


^1H NMR (300 MHz, CDCl_3): δ 3.63 (2H, t), 2.85 (2H, t), 2.24 (2H, t), 1.67 (2H, t),
 1.44 (9H, s), 1.07 (6H, s).



A mixture of tert-butyl 4-((2-chloroethyl)amino)-4-methylpentanoate (3.9 g, 15.7
 25 mmol) and trifluoroacetic acid (27 mL) in dichloromethane (80 mL) were stirred at room
 temperature overnight. The reaction mixture was concentrated under reduced pressure. The
 residue was dissolved in further dichloromethane and concentrated again. This was repeated a
 further 3 times until the majority of the excess trifluoroacetic acid had been removed. The
 residue was dissolved in dichloromethane (500 mL) and N-(3-Dimethylaminopropyl)-N'-

ethylcarbodiimide hydrochloride (4.61 g, 24.1 mmol), hydroxybenzotriazole hydrate (3.25 g, 24.1 mmol) and diisopropylethylamine (21 mL, 120 mmol) added. The mixture was stirred at room temperature overnight. The reaction was washed with water (300 mL) and dried (MgSO₄). Filtration and removal of the solvent under reduced pressure gave a crude residue that was purified by silica chromatography (heptane to ethyl acetate) giving 1-(2-chloroethyl)-5,5-dimethylpyrrolidin-2-one (1.24 g, 44% yield).


¹H NMR (300 MHz, CDCl₃): δ 3.61 (2H, t), 3.41 (2H, t), 2.38 (2H, t), 1.88 (2H, t), 1.24 (6H, s).



2-(2,2-Dimethyl-5-oxopyrrolidin-1-yl)ethyl methyl fumarate **24** was synthesised from 1-(2-chloroethyl)-5,5-dimethylpyrrolidin-2-one following general procedure 5 (1.02 g, 41 %).

¹H NMR (300 MHz, CDCl₃); 6.85 (2H, d), 4.33 (2H, t), 3.80 (3H, s), 3.41 (2H, t), 2.39 (2H, t), 1.88 (2H, t), 1.23 (6H, s). *m/z* [M+H]⁺ = 270.17.

(E)-4-(2-((4-methoxy-4-oxobut-2-enyl)oxy)ethyl)morpholine 4-oxide (26)




To a solution of methyl (2-morpholinoethyl) fumarate (1.1 g, 4.5 mmol) [synthesised from 4-(2-chloroethyl)morpholine following general procedure 5] in dichloromethane was added *m*-chloroperbenzoic acid (1.87 g, 5.4 mmol) and the reaction mixture stirred for 1 h.

The reaction mixture was diluted with water (25 mL) and washed with dichloromethane (3 x 50 mL). The aqueous phase was lyophilized giving (E)-4-(2-((4-methoxy-4-oxobut-2-enyl)oxy)ethyl)morpholine 4-oxide **26** (0.19 g, 16 %).

¹H NMR (300 MHz, CDCl₃); 6.87 (1H, d), 6.81 (1H, d), 4.92-4.88 (2H, M), 4.44 (2H, t), 3.78-3.73 (2H, m), 3.54-3.48 (2H, m), 3.34 (2H, t), 3.15 (2H, d). *m/z* [M+H]⁺ = 260.2


2-(3,5-dioxomorpholino)ethyl methyl fumarate (27)



To a solution of diglycolic anhydride (2.0 g, 17 mmol) in pyridine (10 mL) was added ethanolamine (2.1 g, 34 mmol) and heated at reflux for 2 h. The volatiles were removed *in*

vacuo and the residue heated at 180 °C for 2 h and then 220 °C for 90 min. The reaction mixture was cooled and the residue purified on silica eluting with dichloromethane/ethyl acetate (4:1) giving 4-(2-hydroxyethyl)morpholine-3,5-dione (1.05 g, 38%).

¹H NMR (300 MHz, CDCl₃); 4.39 (4H, s), 4.02 (2H, t), 3.80 (2H, t).




2-(3,5-dioxomorpholino)ethyl methyl fumarate **27** was synthesised from 4-(2-hydroxyethyl)morpholine-3,5-dione following general procedure 1 (0.82 g, 96%).

¹H NMR (300 MHz, CDCl₃); 6.83 (1H, d), 6.75 (1H, d), 4.39-4.43 (6H, m), 4.12 (2H, t), 3.79 (3H, s).

10


2-(2,2-dimethylmorpholino)ethyl methyl fumarate hydrochloride (28)



To a solution of 2,2-dimethylmorpholine (1.0 g, 8.7 mmol) in dichloromethane (35 mL) was added chloroacetaldehyde (50% in water, 1.65 mL, 13.0 mmol), followed by sodium triacetoxyborohydride (2.8 g, 13.0 mmol). The reaction mixture was stirred for 90 min, diluted with 1 M aqueous sodium hydroxide (40 mL) and the organic phase separated. The aqueous phase was extracted with dichloromethane (2 x 30 mL) and the organic phases combined. After being dried over MgSO₄ the volatiles were removed *in vacuo* giving 4-(2-chloroethyl)-2,2-dimethylmorpholine (1.45 g, 94%).

¹H NMR (300 MHz, CDCl₃); 3.73 (2H, dd), 3.55 (2H, t), 2.64 (2H, t), 2.43 (2H, dd), 2.25 (2H, s), 1.24 (6H, s).

20




2-(2,2-Dimethylmorpholino)ethyl methyl fumarate **28** was synthesised from 4-(2-chloroethyl)-2,2-dimethylmorpholine following general procedure 5 (0.71 g, 93%).

4-(2-chloroethyl)-2,2-dimethylmorpholine was converted to 4-(2-chloroethyl)-2,2-dimethylmorpholine hydrochloride following general procedure 2 (0.69 g, 87 %).

25


¹H NMR (300 MHz, CDCl₃); 6.85 (1H, d), 6.77 (1H, d), 4.52-4.47 (2H, m), 3.93-3.85 (2H, m), 3.70 (3H, s), 3.48-3.43 (2H, m), 3.32-3.00 (4H, m), 1.24 (6H, s). *m/z* [M+H]⁺ = 272.2

2-(2,6-dimethylmorpholino)ethyl methyl fumarate hydrochloride (29)



To a solution of 2,6-dimethylmorpholine (1.0 g, 9.0 mmol) in dichloromethane (40 mL) was added chloroacetaldehyde (50% in water, 1.02 mL, 13.5 mmol) and acetic acid (0.75 mL, 13.5 mmol) followed by sodium triacetoxyborohydride (2.8 g, 13.5 mmol). The reaction mixture was stirred for 4 h, diluted with dichloromethane (20 mL) and washed with saturated aqueous sodium hydrogen carbonate (30 mL). The organic phase separated, dried over MgSO₄ the volatiles were removed *in vacuo*. The residue was further purified by silica chromatography eluting with heptanes/ethyl acetate (1:1) giving 4-(2-chloroethyl)-2,6-dimethylmorpholine (0.44 g, 30%).

¹H NMR (300 MHz, CDCl₃); 3.75-3.62 (2H, m), 3.58 (2H, t), 2.65-2.79 (4H, m), 1.83 (2H, t), 1.15 (6H, d).




2-(2,6-dimethylmorpholino)ethyl methyl fumarate **29** was synthesised from 4-(2-chloroethyl)-2,6-dimethylmorpholine following general procedure 5 (0.54 g, 71%).

2-(2,6-dimethylmorpholino)ethyl methyl fumarate was converted to 2-(2,6-dimethylmorpholino)ethyl methyl fumarate hydrochloride following general procedure 2 (0.19 g, 64 %).

¹H NMR (300 MHz, CDCl₃); 6.83 (1H, d), 6.75 (1H, d), 4.47-4.43 (2H, m), 3.93-3.82 (2H, m), 3.67 (3H, s), 3.46-3.40 (2H, m), 2.72 (2H, t), 1.10 (6H, d). *m/z* [M+H]⁺ = 272.2


Methyl (2-(3-oxomorpholino)ethyl) fumarate (30)



A mixture of potassium *tert*-butoxide (5.9 g, 52.3 mmol) and toluene (50 mL) was heated at 75 °C for 30 min and then diethanolamine (5.0 g, 47.6 mmol) added. The reaction mixture was heated a further 30 min and then methyl chloroacetate (4.4 mL, 50.0 mmol)

added. After a further 2 h heating the reaction was diluted with methanol (21 mL) and cooled to room temperature. The reaction mixture was filtered, washed with toluene and the mother liquor evaporated. The residue was further purified by silica flash column chromatography giving 4-(2-hydroxyethyl)morpholin-3-one (0.65 g, 9%).


5 ^1H NMR (300 MHz, CDCl_3); 4.19 (2H, s), 3.89 (2H, t), 3.81 (2H, t), 3.57 (2H, t), 3.48 (2H, t), 2.89 (1H, s).



10 Methyl (2-(3-oxomorpholino)ethyl) fumarate **30** was synthesised from 4-(2-hydroxyethyl)morpholin-3-one following general procedure 1 (0.71 g, 62%).

^1H NMR (300 MHz, DMSO); 6.72 (2H, s), 4.28 (2H, t), 3.98 (2H, s), 3.77 (2H, t), 3.71 (3H, t), 3.59 (2H, t), 3.38 (2H, t). m/z $[\text{M}+\text{H}]^+ = 258.1$

Methyl (2-(2-oxomorpholino)ethyl) fumarate hydrochloride (31)




15 Methyl (2-(2-oxomorpholino)ethyl) fumarate **31** was synthesised from 4-(2-hydroxyethyl)morpholin-2-one following general procedure 1 (0.53 g, 34%).

Methyl (2-(2-oxomorpholino)ethyl) fumarate was converted to methyl (2-(2-oxomorpholino)ethyl) fumarate hydrochloride following general procedure 2 (0.20 g, 34%).

20 ^1H NMR (300 MHz, DMSO); 3.75 (2H, s), 4.29-4.23 (4H, m), 3.71 (3H, s), 3.34 (2H, s), 2.73 (2H, t), 2.68 (2H, t). m/z $[\text{M}+\text{H}]^+ = 258.15$

2-(8-Oxa-3-azabicyclo[3.2.1]octan-3-yl)ethyl methyl fumarate hydrochloride (32)




25 2-(8-Oxa-3-azabicyclo[3.2.1]octan-3-yl)ethyl methyl fumarate **32** was synthesised from 3-(2-chloroethyl)-8-oxa-3-azabicyclo[3.2.1]octane following general procedure 5 (0.25 g, 50%).

2-(8-Oxa-3-azabicyclo[3.2.1]octan-3-yl)ethyl methyl fumarate was converted to 2-(8-oxa-3-azabicyclo[3.2.1]octan-3-yl)ethyl methyl fumarate hydrochloride following general procedure 2 (0.20 g, 73%).

¹H NMR (300 MHz, D₂O); 6.82 (1H, d), 6.75 (1H, d), 4.52-4.42 (4H, m), 3.69 (3H, s), 3.45-3.37 (4H, m), 3.26-3.19 (2H, m), 2.10-1.85 (4H, m). *m/z* [M+H]⁺ = 270.0

2-(2-((Dimethylamino)methyl)morpholino)ethyl methyl fumarate hydrochloride (33)




2-(2-((Dimethylamino)methyl)morpholino)ethyl methyl fumarate **33** was synthesised from 1-(4-(2-chloroethyl)morpholin-2-yl)-N,N-dimethylmethanamine following general procedure 5 (0.17 g, 16%).

2-(2-((Dimethylamino)methyl)morpholino)ethyl methyl fumarate was converted to 2-(2-((Dimethylamino)methyl)morpholino)ethyl methyl fumarate hydrochloride following general procedure 2 (0.17 g, 95%).

¹H NMR (300 MHz, D₂O); 6.84 (1H, d), 6.77 (1H, d), 4.50-4.45 (2H, m), 4.21-4.06 (2H, m), 3.87-3.77 (1H, m), 3.68 (3H, s), 3.56-3.47 (2H, m), 3.25-3.09 (3H, m), 2.94 (1H, dd), 2.81 (6H, bs). *m/z* [M+H]⁺ = 301.2


2-((3S,5S)-3,5-Dimethylmorpholino)ethyl methyl fumarate hydrochloride (34)




2-((3S,5S)-3,5-Dimethylmorpholino)ethyl methyl fumarate **34** was synthesised from (3S,5S)-4-(2-chloroethyl)-3,5-dimethylmorpholine following general procedure 5 (0.11 g, 25%).

2-((3S,5S)-3,5-Dimethylmorpholino)ethyl methyl fumarate was converted to 2-((3S,5S)-3,5-dimethylmorpholino)ethyl methyl fumarate hydrochloride following general procedure 2 (0.08 g, 68%).

¹H NMR (300 MHz, D₂O); 7.15-7.00 (2H, m), 4.77-4.70 (2H, m), 4.20-4.08 (2H, m), 4.01-3.85 (8H, m), 3.68-3.58 (1H, m). *m/z* [M+H]⁺ = 272.3


2-(2,5-Dioxomorpholino)ethyl methyl fumarate (35)

2-(2,5-Dioxomorpholino)ethyl methyl fumarate **35** was synthesised from 4-(2-hydroxyethyl)morpholine-2,5-dione following general procedure 1 (0.27 g, 65%). ¹H NMR (300 MHz, DMSO); 6.75 (1H, d), 6.71 (1H, d), 4.72 (2H, s), 4.30 (2H, s), 4.26 (2H, t), 3.72 (3H, s), 3.60 (2H, t). *m/z* [M+H]⁺ = 272.2

(E)-Methyl 3-(4-methyl-2,5,7-trioxabicyclo[2.2.2]octan-1-yl)acrylate (130)


Methyl ((3-methyloxetan-3-yl)methyl) fumarate was synthesised from 3-methyl-3-oxetane methanol following general procedure 1 (0.86 g, 89%).

¹H NMR (300 MHz, CDCl₃); 6.88 (2H, s), 4.52 (2H, d), 4.40 (2H, d), 4.30 (2H, s), 3.82 (3H, s), 1.35 (3H, s).




To a solution of methyl ((3-methyloxetan-3-yl)methyl) fumarate **130** (0.20 g, 0.93 mmol) in dichloromethane (5 mL) at 5 °C was added borontrifluoride diethyletherate (0.058 mL, 0.47 mmol). After 1 h a further portion of borontrifluoride diethyletherate (0.058 mL, 0.47 mmol) was added and the reaction mixture warmed to 20 °C over 1h. To the reaction mixture was added triethylamine (0.13 mL, 0.93 mmol) and then this was loaded directly onto a silica column. The desired product was eluted with heptane/ethyl acetate (6:4) containing triethylamine (2.5% v/v) giving (E)-methyl 3-(4-methyl-2,5,7-trioxabicyclo[2.2.2]octan-1-yl)acrylate (0.12 g, 60%).

¹H NMR (300 MHz, CDCl₃); 6.66 (1H, d), 6.25 (1H, d), 3.97 (6H, s), 3.73 (3H, s), 0.84 (3H, s). *m/z* [M+H]⁺ = 215.2

Methyl prop-2-yn-1-yl fumarate (131)

Methyl prop-2-yn-1-yl fumarate **131** was synthesized from propargyl alcohol following general procedure 1 (0.51 g, 68%).


- 5 ^1H NMR (300 MHz, DMSO); 6.85-6.70 (2H, m), 4.81 (2H, d), 3.72 (3H, s), 3.60 (1H, t).

2-(1,3-Dioxoisindolin-2-yl)ethyl methyl fumarate (36)

- 10 2-(1,3-Dioxoisindolin-2-yl)ethyl methyl fumarate **36** was synthesised from 2-(2-hydroxyethyl)isindoline-1,3-dione following general procedure 1 (0.63 g, 79%).


^1H NMR (300 MHz, MeOD); 7.87-7.77 (4H, m), 6.74-6.73 (2H, m), 4.45-4.40 (2H, m), 4.01-3.96 (2H, m), 3.76 (3H, s). m/z $[\text{M}+\text{H}]^+ = 304.1$

- 15 4-(2,5-Dioxopyrrolidin-1-yl)butyl methyl fumarate (132)



4-(2,5-Dioxopyrrolidin-1-yl)butyl methyl fumarate **132** was synthesised from 1-(4-hydroxybutyl)pyrrolidine-2,5-dione following general procedure 1 (0.77 g, 79%).


- 20 ^1H NMR (300 MHz, MeOD); 6.81-6.79 (2H, m), 4.20 (2H, t), 3.78 (3H, s), 3.50 (2H, t), 2.67 (4H, s), 1.71-1.62 (4H, m). m/z $[\text{M}+\text{H}]^+ = 284.2$

2-(3,3-Dimethyl-2,5-dioxopyrrolidin-1-yl)ethyl methyl fumarate (36)

2-(3,3-Dimethyl-2,5-dioxopyrrolidin-1-yl)ethyl methyl fumarate **36** was synthesised from 1-(2-hydroxyethyl)-3,3-dimethylpyrrolidine-2,5-dione following general procedure 1 (0.72 g, 74%).

¹H NMR (300 MHz, CDCl₃); 6.83 (1H, d), 6.77 (1H, d), 4.38 (2H, t), 3.82 (1H, t),
 5 3.80 (3H, s), 2.55 (2H, s), 1.31 (6H, s). *m/z* [M+H]⁺ = 284.1


3-(2,5-Dioxopyrrolidin-1-yl)propyl methyl fumarate (133)



3-(2,5-Dioxopyrrolidin-1-yl)propyl methyl fumarate **133** was synthesised from 1-(3-hydroxypropyl)pyrrolidine-2,5-dione following general procedure 1 (0.64 g, 69%).

¹H NMR (300 MHz, MeOD); 6.82 (2H, s), 4.17 (2H, t), 3.79 (3H, s), 3.59 (2H, t),
 2.67 (4H, s), 1.95 (2H, dt). *m/z* [M+H]⁺ = 270.2

Methyl (2-(2-oxopyrrolidin-1-yl)ethyl) fumarate (38)




15

Methyl (2-(2-oxopyrrolidin-1-yl)ethyl) fumarate **38** was synthesised from 1-(2-hydroxyethyl)pyrrolidin-2-one following general procedure 1 (0.68 g, 73%).

¹H NMR (300 MHz, MeOD); 6.85 (2H, s), 4.33 (2H, t), 3.80 (3H, s), 3.59 (2H, t),
 3.46 (2H, t), 2.37 (2H, t), 2.03 (2H, dt). [M+H]⁺ = 242.1


20

Methyl (2-(2-oxooxazolidin-3-yl)ethyl) fumarate (39)




Methyl (2-(2-oxooxazolidin-3-yl)ethyl) fumarate **39** was synthesised from 3-(2-hydroxyethyl)oxazolidin-2-one following general procedure 1 (0.77 g, 92%).

¹H NMR (300 MHz, MeOD); 6.82 (2H, s), 4.39-4.30 (4H, m), 3.78 (3H, s), 3.72-3.67
 25 (2H, m), 3.58-3.54 (2H, m). *m/z* [M+H]⁺ = 244.2

2-(4,4-Dimethyl-2,5-dioxoimidazolidin-1-yl)ethyl methyl fumarate (42)


2-(4,4-Dimethyl-2,5-dioxoimidazolidin-1-yl)ethyl methyl fumarate **42** was synthesised from 3-(2-hydroxyethyl)-5,5-dimethylimidazolidine-2,4-dione following general procedure 1 (0.33 g, 33%).

^1H NMR (300 MHz, CDCl_3); 6.82 (2H, s), 5.50 (NH), 4.40 (2H, t), 3.86-3.76 (5H, m), 1.43 (6H, s). m/z $[\text{M}+\text{H}]^+ = 285.2$

Methyl (2-(N-propionylpropionamido)ethyl) fumarate (42)


Methyl (2-propionamidoethyl) fumarate **41** was synthesised from N-(2-hydroxyethyl)propionamide following general procedure 1 (1.7 g, 96%).

^1H NMR (300 MHz, CDCl_3); 6.87 (2H, s), 4.29 (2H, t), 3.81 (3H, s), 3.58 (2H, q), 2.21 (2H, q), 1.15 (3H, t).




A mixture of methyl (2-propionamidoethyl) fumarate (1.7 g, 7.4 mmol), propionic anhydride (36 mL) and sodium propionate (1.0 g, 10.4 mmol) was heated at 150 °C for 16 h. The reaction was cooled, concentrated to 1/3rd volume and then loaded onto a silica column and eluted with 0-20% ethyl acetate/dichloromethane. The product containing fractions were combined, evaporated and re-purified by silica flash chromatography eluting with 10-50% ethyl acetate/heptanes giving methyl (2-(N-propionylpropionamido)ethyl) fumarate **42** (0.18 g, 21%).

^1H NMR (300 MHz, CDCl_3); 6.83-6.82 (2H, m), 4.34 (2H, t), 4.01 (2H, t), 3.81 (3H, s), 2.75 (4H, q), 1.16 (6H, t).

2-((3R,4S)-3,4-Dimethyl-2,5-dioxopyrrolidin-1-yl)ethyl methyl fumarate (23)


Racemic 2-((3R,4S)-3,4-dimethyl-2,5-dioxopyrrolidin-1-yl)ethyl methyl fumarate **23** was synthesised from racemic (3R,4S)-1-(2-hydroxyethyl)-3,4-dimethylpyrrolidine-2,5-dione following general procedure 1 (0.54 g, 44%).

^1H NMR (300 MHz, CDCl_3); 6.81-6.80 (2H, m), 4.37 (2H, t), 3.82 (2H, t), 3.80 (3H, s), 3.00-2.88 (2H, m), 1.25-1.18 (6H, m). m/z $[\text{M}+\text{H}]^+ = 284.2$

2-Acetamidoethyl methyl fumarate (43)


2-Acetamidoethyl methyl fumarate was synthesised from N-(2-hydroxyethyl)acetamide **43** following general procedure 1 (0.23 g, 70%).

^1H NMR (300 MHz, CDCl_3); 6.87 (2H, s), 5.80 (NH), 4.29 (2H, t), 3.81 (3H, s), 3.57 (2H, q), 2.00 (3H, s). m/z $[\text{M}+\text{H}]^+ = 216.14$

2-(N-Acetylacetamido)ethyl methyl fumarate (44)

A mixture of 2-acetamidoethyl methyl fumarate (0.62 g, 2.9 mmol), acetic anhydride (15 mL) and sodium acetate (0.33 g, 4.0 mmol) was heated at reflux for 20 h. The reaction mixture was evaporated and the residue suspended in dichloromethane. The supernatant was loaded onto a silica column and eluted with 0-20% ethyl acetate/dichloromethane giving 2-(N-Acetylacetamido)ethyl methyl fumarate **44** (0.36 g, 48%).


^1H NMR (300 MHz, CDCl_3); 6.87 (1H, d), 6.82 (1H, d), 4.36 (2H, d), 4.00 (2H, d), 3.81 (3H, s), 2.44 (3H, s).

2-((*tert*-butoxycarbonyl)amino)ethyl methyl fumarate (48)

To a suspension of monomethyl fumarate (MMF) (1.0 equivalent) in dichloromethane (11 mL per g of MMF) was added diisopropylethylamine (3 equivalents), 2-((*tert*-
 5 butoxycarbonyl)amino)ethanol (1.02 equivalents) and *N,N,N',N'*-tetramethyl-*O*-(1*H*-benzotriazol-1-yl)uronium tetrafluoroborate (1.5 equivalents). The reaction was stirred for 1-18 hours at <10 °C. The reaction was quenched with 1M hydrochloric acid (0.6 mL/mL of DCM). The organic layer was washed with 10% (w/w) aqueous sodium bicarbonate solution (0.6 mL/mL of DCM) followed by 37% (w/w) sodium chloride solution (0.6 mL/mL of
 10 DCM). The organic layer was dried over sodium sulfate, filtered to remove the drying agent, and the solution added to a silica plug (~6 g of silica gel/g of MMF) and the plug flushed with DCM until no more product eluted. ~80% of the DCM was removed under reduced pressure at 30 °C after which time 25 mL of MTBE/g of MMF were added and the solution further concentrated at 30 °C until ~10 mL/g of MMF remained. The resulting suspension
 15 was cooled to 5 °C for at least 1 hour and then the resulting solids were collected by filtration to give the desired MMF ester prodrug. (3.8 g, 91 %).

¹H NMR (400 MHz, DMSO-*d*₆) δ 7.08 (t, *J* = 5.4 Hz, 1H), 6.89 (d, *J* = 15.8 Hz, 1H), 6.79 (d, *J* = 15.8 Hz, 1H), 4.18 (t, *J* = 5.3 Hz, 2H), 3.81 (s, 3H), 3.28 (q, *J* = 5.4 Hz, 2H), 1.43 (s, 9H). *m/z* [M+H]⁺ = 274.3.

20

2-(2,5-dioxo-2,5-dihydro-1*H*-pyrrol-1-yl)ethyl methyl fumarate (55)


To a suspension of monomethyl fumarate (MMF) (1.0 equivalent) in dichloromethane (11 ml per g of MMF) was added diisopropylethylamine (3 equivalents), the desired alcohol
 25 (1.02 equivalents) and *N,N,N',N'*-tetramethyl-*O*-(1*H*-benzotriazol-1-yl)uronium tetrafluoroborate (1.5 equivalents). The reaction was stirred for 1-18 hours at <10 °C. The reaction was quenched with 1M hydrochloric acid (0.6 mL/mL of DCM). The organic layer was washed with 10% (w/w) aqueous sodium bicarbonate solution (0.6 mL/mL of DCM) followed by 37% (w/w) sodium chloride solution (0.6 mL/mL of DCM). The organic layer
 30 was dried over sodium sulfate, filtered to remove the drying agent, and the solution added to a silica plug (~6 g of silica gel/g of MMF) and the plug flushed with DCM until no more

product eluted. ~80% of the DCM was removed under reduced pressure at 30 °C after which time 25 mL of MTBE/g of MMF were added and the solution further concentrated at 30 °C until ~10 mL/g of MMF remained. The resulting suspension was cooled to 5 °C for at least 1 hour and then the resulting solids were collected by filtration to give the desired MMF ester produg. (2.4 g, 67 %).

¹H NMR (400 MHz, Chloroform-*d*) δ 6.82 (d, J = 2.8 Hz, 2H), 6.74 (s, 2H), 4.36 (t, J = 5.3 Hz, 2H), 3.86 (t, J = 5.3 Hz, 2H), 3.81 (s, 3H). *m/z* [M+H]⁺ = 254.2.

Reference Compound A

2-(diethylamino)-2-oxoethyl methyl fumarate



2-(diethylamino)-2-oxoethyl methyl fumarate was synthesized following general procedure 3 and conformed to reported data in US Patent No. 8,148,414.

Example 2- Aqueous Chemical Stability of Several Compounds

Stock solutions of the compounds in acetonitrile or acetonitrile/methanol were prepared at 20 mg/mL and 20 μL, spiked into 3mL of buffer phosphate (100mM) and incubated at 37 °C. Aliquots (50 μL) were sampled at different time points and diluted 20 fold with ammonium formate (pH 3.5)/acetonitrile. The diluted samples were analyzed by HPLC. The peak areas corresponding to the compounds were plotted against time and the data were fitted to a first-order mono-exponential decay where the rate constant and the half-life were determined (Table 3). In some cases, in which the half life is too long (>360min), an estimated value of the half life is reported using the initial slope at low conversion (<10%).

Table 3.

Compound	pH 8 (t _{1/2} , min)
1	15
4	45
5	24
6	2.0
7	26.0
8	36.0
9	7.0

Compound	pH 8 (t $\frac{1}{2}$, min)
10	67.0
11	>240
12	396
14	144
15	3.0
16	20.0
17	11.0
18	5.0
19	6.0
20	5.0
Reference Compound A	120

Stock solutions of the compounds in acetonitrile or acetonitrile/MeOH were prepared at 0.05M. A 0.010 mL aliquot of the stock was spiked into 1 mL of 50 mM buffer phosphate pH 8 and incubated at 37 °C. Typically, aliquots (0.010 mL) were sampled at different time points and immediately injected in the HPLC with UV detection (211nm). The peak areas corresponding to the compounds were plotted against time and the data were fitted to a first-order mono-exponential decay where the rate constant and the half-life were determined from the slope (Table 4).

10

Table 4.

Compound	pH 8 (t $\frac{1}{2}$, min)
1	15
4	30
5	24
6	2
19	117
22	144
23	186
26	129
27	37
28	<10
29	<10
30	229
31	26
32	13
33	115
35	
37	182
38	201

Compound	pH 8 (t ^{1/2} , min)
39	183
40	203
42	158
43	177.5
44	145
48	220
130	1010
131	96
133	246

Example 3- Evaluation of Aqueous Chemical Stability with NMR

The chemical hydrolysis was followed by dissolving the ester in phosphate buffered D₂O (pH 7.9) in an NMR tube, heating the NMR tube to 37° C and periodically recording the spectra. These various species produced by hydrolysis of the diesters were followed over time. See Figures 1-5.

Example 4- Delivery of MMF in Rats Upon Oral Administration of Prodrugs

Rats were obtained commercially and were pre-cannulated in the jugular vein. Animals were conscious at the time of the experiment. All animals were fasted overnight and until 4 hours post-dosing of a prodrug in the disclosure.

Blood samples (0.25 mL/sample) were collected from all animals at different time-points up to 24 hours post-dose into tubes containing sodium fluoride/sodium EDTA. Samples were centrifuged to obtain plasma. Plasma samples were transferred to plain tubes and stored at or below -70° C prior to analysis.

To prepare analysis standards, 20 uL of rat plasma standard was quenched with 60 uL of internal standard. The sample tubes were vortexed for at least 1 min and then centrifuged at 3000 rpm for 10 min. 50 uL of supernatant was then transferred to 96-well plates containing 100 uL water for analysis by LC-MS-MS.

LC-MS/MS analysis was performed using an API 4000 equipped with HPLC and autosampler. The following HPLC column conditions were used: HPLC column: Waters Atlantis T3; flow rate 0.5 mL/min; run time 5 min; mobile phase A: 0.1% formic acid in water; mobile phase B: 0.1% formic acid in acetonitrile (ACN); gradient: 98% A/2% B at 0.0 min; 98% A/2% B at 1 min; 5% A/95% B at 3 min; 5% A/95% B at 3.75 min; 97% A/3% B at 4 min; and 98% A/2% B at 5.0 min. MMF was monitored in positive ion mode.

MMF, DMF or MMF prodrug was administered by oral gavage to groups of two to six adult male Sprague-Dawley rats (about 250 g). Animals were conscious at the time of the experiment. MMF, DMF or MMF prodrug was orally administered in an aqueous solution of 0.5% hydroxypropyl methyl cellulose (HPMC), 0.02% polysorbate 80, and 20 mM citrate buffer (pH 5), at a dose of 10 mg-equivalents MMF per kg body weight.

The percent absolute bioavailability (F%) of MMF was determined by comparing the area under the MMF concentration vs time curve (AUC) following oral administration of MMF, DMF or MMF prodrug with the AUC of the MMF concentration vs time curve following intravenous administration of MMF on a dose normalized basis.

The MMF prodrugs, when administered orally to rats at a dose of 10 mg/kg MMF-equivalents in the aqueous vehicle, exhibited an absolute oral bioavailability (relative to IV) ranging from about 3% to about 96% (See Tables 5 and 6). Tables 5 and 6 show data from two independent studies.

Table 5.

Compound No.	Percent Absolute Bioavailability (F%)
MMF	43%
DMF	53%
16	60-82%
4	3%
14	96%
10	73%

Table 6.

Compound No.	Percent Absolute Bioavailability (F%)
MMF	69.6
DMF	69.6
132	60.3
40	70.4
39	91.4
5	81.1
11	71.4

Example 5- Delivery of MMF in Dogs Upon Oral Administration of Prodrugs

Male Beagle dogs were obtained from the test facility's colony of non-native animals. All animals were fasted overnight prior to dose administration.

Oral doses were administered via oral gavage. The gavage tube was flushed with 10 mL of water prior to removal.

All animals were observed at dosing and at each scheduled collection. All abnormalities were recorded.

Blood samples were collected in Sodium Fluoride/ Na_2EDTA tubes and stored on wet ice until processed to plasma by centrifugation (300 rpm at 5°C) within 30 minutes of collection. All plasma samples were transferred into separate 96-well plates (matrix tubes) and stored at -80 °C until concentration analysis was performed via LC/MS/MS using an RGA 3 assay.

Extraction Procedure:

Note: Thawed test samples at 4°C. (Kept in ice while on bench).

1. Aliquoted 20uL of study sample, standard, and QC samples into labeled 96-well plate.
2. Added 120uL of appropriate internal standard solution (125ng/mL mouse embryo fibroblasts (MEF)) to each tube, except for the double blank to which 120uL of appropriate acetonitrile:FA (100:1) was added.
3. Sealed and vortexed for one minute.
4. Centrifuged at 3000 rpm for 10 minutes.
5. Transferred 100uL of supernatant to a clean 96-well plate containing 100uL water.
6. Sealed and vortexed gently for 2 minutes.

The percent absolute bioavailability (F%) of MMF was determined by comparing the area under the MMF concentration vs time curve (AUC) following oral administration of MMF prodrug with the AUC of the MMF concentration vs time curve following intravenous administration of MMF on a dose normalized basis.

The MMF prodrugs, when administered orally to dogs at a dose of 10 mg/kg MMF-equivalents in the aqueous vehicle, exhibited an absolute oral bioavailability (relative to IV) ranging from about 31% to about 78% (See Table 7).

Table 7

Compound No.	Percent Absolute Bioavailability (F%)
16	54%
16 (capsule)	54%
14	78%
10	31%

Example 6- Physical Stability of the Instant Prodrugs and DMF in Crystalline Form

The physical stability of compounds of the present invention and DMF were measured via thermogravimetric analysis (TGA). Figure 6 shows a plot of weight loss at 60 °C vs time for Compound 14 (12.15 mg), no change, and DMF (18.40 mg), ~100 % weight loss in less than 4 hours. These data indicate that DMF undergoes sublimation while Compound 14 is physically stable under similar conditions.

Example 7- Single Crystal X-ray Data for Compound 14

Compound 14 produced by the method described in Example 1 was analyzed. Figure 7 depicts the unit cell. The single crystal x-ray data are included below:

Single crystal data:

Empirical formula: C₁₁ H₁₃ N O₆

Formula weight: 255.22

Temperature: 173(2) K

Wavelength: 1.54178 Å

Space group: P-1

Unit cell dimensions: a = 6.07750(10) Å α = 84.9390(10)°.

b = 7.96290(10) Å β = 80.0440(10)°.

c = 12.7850(2) Å γ = 71.9690(10)°.

Volume: 579.080(15) Å³

Z: 2

Density (calculated): 1.464 Mg/m³

Absorption coefficient: 1.034 mm⁻¹

F(000): 268

Crystal size: 0.37 x 0.15 x 0.15 mm³

Reflections collected: 8446

Independent reflections: 2229 [$R(\text{int}) = 0.0249$]

Refinement method: Full-matrix least-squares on F^2

Goodness-of-fit on F^2 : 1.049


Final R indices [$I > 2\sigma(I)$] $R1 = 0.0317$, $wR2 = 0.0850$

5 R indices (all data): $R1 = 0.0334$, $wR2 = 0.0864$

CLAIMS

What is claimed is:

1. A compound of Formula (I), or a pharmaceutically acceptable salt thereof:



wherein:

R₁ is unsubstituted C₁-C₆ alkyl;

- L_a is unsubstituted C₁-C₆ alkyl linker, unsubstituted C₃-C₁₀ carbocycle, unsubstituted C₆-C₁₀ aryl, unsubstituted heterocycle comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S, or unsubstituted heteroaryl comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S; and

- R₂ and R₃ are each, independently, H, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₆-C₁₀ aryl, C₃-C₁₀ carbocycle, heterocycle comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S, or heteroaryl comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S, wherein the alkyl, alkenyl, aryl, carbocycle, heterocycle, or heteroaryl groups may be optionally independently substituted one or more times with C₁-C₃-alkyl, OH, O(C₁-C₄ alkyl), carbonyl, halo, NH₂, N(H)(C₁-C₆ alkyl), N(C₁-C₆ alkyl)₂, SO₂H, SO₂(C₁-C₆ alkyl), CHO, CO₂H, CO₂(C₁-C₆ alkyl), or CN;

- or alternatively, R₂ and R₃, together with the nitrogen atom to which they are attached, form a heteroaryl comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S; or a heterocycle comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S, wherein the heteroaryl or heterocycle may be optionally substituted one or more times with C₁-C₆ alkyl, CN, OH, halo, O(C₁-C₆ alkyl), CHO, carbonyl, thione, NO, or NH₂.


2. The compound of claim 1, wherein R₁ is methyl.

3. The compound according to any of the preceding claims, wherein L_a is an unsubstituted C_1 - C_6 alkyl linker.

4. The compound according to any of the preceding claims, wherein R_2 and R_3 together with the nitrogen to which they are attached form a heterocycle, wherein the heterocycle is a morpholine ring, a thiomorpholine ring, a pyrrolidine ring, a 2,5-dihydropyrrole ring, a 1,2-dihydropyridine ring, a piperazine ring, a succinimide ring, an isoindoline ring, a 2,5-dihydro-1*H*-tetrazole ring, an azetidine ring, a piperidine ring, a hexahydropyrimidine ring, a 2,3,3a,4,7,7a-hexahydro-1*H*-4,7-epoxyisoindole ring, a 3,4-dihydroquinazoline ring, a 1,2,3,4-tetrahydroquinazoline ring, an oxazolidine ring, an oxazolidinone ring, an imidazolidinone ring, a 1,3-dihydro-2*H*-imidazol-2-one ring, an imidizolidine thione ring, or an isothiazolidine ring, wherein all of the rings may be optionally substituted one or more times with C_1 - C_6 alkyl, $CO_2(C_1$ - C_6 alkyl), OH, $(CH_2)_{1-4}OH$, $O(C_1$ - C_6 alkyl), halo, NH_2 , $(CH_2)_{1-4}NH_2$, $(CH_2)_{1-4}NH(C_1$ - C_4 alkyl), $(CH_2)_{1-4}N(C_1$ - C_4 alkyl) $_2$, carbonyl, or thione.

5. The compound according to any of the preceding claims, wherein R_2 and R_3 , together with the nitrogen atom to which they are attached, form a substituted or unsubstituted heteroaryl comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S or a substituted or unsubstituted heterocycle comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S.

6. A compound of Formula (Ia), or a pharmaceutically acceptable salt thereof:



wherein:


R_1 is unsubstituted C_1 - C_6 alkyl;

L_a is substituted or unsubstituted C_1 - C_6 alkyl linker, substituted or unsubstituted C_3 - C_{10} carbocycle, substituted or unsubstituted C_6 - C_{10} aryl, substituted or unsubstituted heterocycle comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from

N, O and S, or substituted or unsubstituted heteroaryl comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S; and

R_2 is H, substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_2 - C_6 alkenyl, substituted or unsubstituted C_2 - C_6 alkynyl, substituted or unsubstituted C_6 - C_{10} aryl, substituted or unsubstituted C_3 - C_{10} carbocycle, substituted or unsubstituted heterocycle comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S, or substituted or unsubstituted heteroaryl comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S.

7. A compound of Formula (Ib):



wherein:

A^- is a pharmaceutically acceptable anion;

R_1 is unsubstituted C_1 - C_6 alkyl;

L_a is substituted or unsubstituted C_1 - C_6 alkyl linker, substituted or unsubstituted C_3 - C_{10} carbocycle, substituted or unsubstituted C_6 - C_{10} aryl, substituted or unsubstituted heterocycle comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S, or substituted or unsubstituted heteroaryl comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S;


R_3' is substituted or unsubstituted C_1 - C_6 alkyl; and

R_2 and R_3 are each, independently, H, substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_2 - C_6 alkenyl, substituted or unsubstituted C_2 - C_6 alkynyl, substituted or unsubstituted C_6 - C_{10} aryl, substituted or unsubstituted C_3 - C_{10} carbocycle, substituted or unsubstituted heterocycle comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S, or substituted or unsubstituted heteroaryl comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S;

or alternatively, R_2 and R_3 , together with the nitrogen atom to which they are attached, form a substituted or unsubstituted heteroaryl comprising one or two 5- or 6-

member rings and 1-4 heteroatoms selected from N, O and S, or a substituted or unsubstituted heterocycle comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S.

- 5 8. A compound of Formula (II), or a pharmaceutically acceptable salt thereof:




wherein:

R₁ is unsubstituted C₁-C₆ alkyl;

- R₄ and R₅ are each, independently, H, substituted or unsubstituted C₁-C₆ alkyl,
 10 substituted or unsubstituted C₂-C₆ alkenyl, substituted or unsubstituted C₂-C₆ alkynyl,
 substituted or unsubstituted C₆-C₁₀ aryl, substituted or unsubstituted C₃-C₁₀ carbocycle,
 substituted or unsubstituted heterocycle comprising one or two 5- or 6-member rings and 1-4
 heteroatoms selected from N, O and S, or substituted or unsubstituted heteroaryl comprising
 one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S;
 15 R₆, R₇, R₈ and R₉ are each, independently, H, substituted or unsubstituted C₁-C₆ alkyl,
 substituted or unsubstituted C₂-C₆ alkenyl, substituted or unsubstituted C₂-C₆ alkynyl or
 C(O)OR_a; and


R_a is H or substituted or unsubstituted C₁-C₆ alkyl.

- 20 9. A compound of Formula (III), or a pharmaceutically acceptable salt thereof:



wherein:

R₁ is unsubstituted C₁-C₆ alkyl;



X is N, O, S or SO₂;

5 Z is C or N;

m is 0, 1, 2, or 3;

n is 1 or 2;

w is 0, 1, 2 or 3;

t is 0, 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10;

10 R₆, R₇, R₈ and R₉ are each, independently, H, substituted or unsubstituted C₁-C₆ alkyl, substituted or unsubstituted C₂-C₆ alkenyl, substituted or unsubstituted C₂-C₆ alkynyl or C(O)OR_a; and

R_a is H or substituted or unsubstituted C₁-C₆ alkyl; and

each R₁₀ is, independently, H, halogen, substituted or unsubstituted C₁-C₆ alkyl,


15 substituted or unsubstituted C₂-C₆ alkenyl, substituted or unsubstituted C₂-C₆ alkynyl, substituted or unsubstituted C₃-C₁₀ carbocycle, substituted or unsubstituted heterocycle comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S, or substituted or unsubstituted heteroaryl comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S;

20 or, alternatively, two R₁₀'s attached to the same carbon atom, together with the carbon atom to which they are attached, form a carbonyl, substituted or unsubstituted C₃-C₁₀ carbocycle, substituted or unsubstituted heterocycle comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S, or substituted or unsubstituted

heteroaryl comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S;

or, alternatively, two R₁₀'s attached to different atoms, together with the atoms to which they are attached, form a substituted or unsubstituted C₃-C₁₀ carbocycle, substituted or unsubstituted heterocycle comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S, or substituted or unsubstituted heteroaryl comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S.

10. A method of treating a neurological disease by administering to a subject in need thereof a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof:



wherein:

R₁ is unsubstituted C₁-C₆ alkyl;

15 L_a is unsubstituted C₁-C₆ alkyl linker, unsubstituted C₃-C₁₀ carbocycle, unsubstituted C₆-C₁₀ aryl, unsubstituted heterocycle comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S, or unsubstituted heteroaryl comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S; and

20 R₂ and R₃ are each, independently, H, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₆-C₁₀ aryl, C₃-C₁₀ carbocycle, heterocycle comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S, or heteroaryl comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S, wherein the alkyl, alkenyl, aryl, carbocycle, heterocycle, or heteroaryl groups may be optionally independently substituted one or more times with C₁-C₃-alkyl, OH, O(C₁-C₄ alkyl), carbonyl, halo, NH₂, N(H)(C₁-C₆ alkyl), N(C₁-C₆ alkyl)₂, SO₂H, SO₂(C₁-C₆ alkyl), CHO, CO₂H, CO₂(C₁-C₆ alkyl), or CN;

or alternatively, R₂ and R₃, together with the nitrogen atom to which they are attached, form a heteroaryl comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S; or a heterocycle comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S, wherein the heteroaryl or heterocycle may be

optionally substituted one or more times with C₁-C₆ alkyl, CN, OH, halo, O(C₁-C₆ alkyl), CHO, carbonyl, thione, NO, or NH₂.


11. The method of claim 10, wherein the neurological disease is multiple sclerosis.

12. The method of claim 10 or 11, wherein the neurological disease is relapsing-remitting multiple sclerosis.

13. The method of any one of claims 10-12, wherein the compound of formula (I) is a pharmaceutically acceptable salt.

14. A pharmaceutical composition comprising:

(i) a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof:



wherein:

R₁ is unsubstituted C₁-C₆ alkyl;

L_a is unsubstituted C₁-C₆ alkyl linker, unsubstituted C₃-C₁₀ carbocycle, unsubstituted C₆-C₁₀ aryl, unsubstituted heterocycle comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S, or unsubstituted heteroaryl comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S; and

R₂ and R₃ are each, independently, H, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₆-C₁₀ aryl, C₃-C₁₀ carbocycle, heterocycle comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S, or heteroaryl comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S, wherein the alkyl, alkenyl, aryl, carbocycle, heterocycle, or heteroaryl groups may be optionally independently substituted one or more times with C₁-C₃-alkyl, OH, O(C₁-C₄ alkyl), carbonyl, halo, NH₂, N(H)(C₁-C₆ alkyl), N(C₁-C₆ alkyl)₂, SO₂H, SO₂(C₁-C₆ alkyl), CHO, CO₂H, CO₂(C₁-C₆ alkyl), or CN;

or alternatively, R₂ and R₃, together with the nitrogen atom to which they are attached, form a heteroaryl comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S; or a heterocycle comprising one or two 5- or 6-member rings and 1-4 heteroatoms selected from N, O and S, wherein the heteroaryl or heterocycle may be
5 optionally substituted one or more times with C₁-C₆ alkyl, CN, OH, halo, O(C₁-C₆ alkyl), CHO, carbonyl, thione, NO, or NH₂; and

(ii) a pharmaceutically acceptable carrier or excipient.

15. The composition of claim 14, wherein the therapeutically effective amount is
10 sufficient for the treatment of multiple sclerosis.

15