

1455 116

PATENT SPECIFICATION

(11) 1 455 116

- (21) Application No. 57970/72 (22) Filed 15 Dec. 1972
 (21) Application No. 43478/73 (22) Filed 17 Sept. 1973
 (23) Complete Specification filed 30 Nov. 1973
 (44) Complete Specification published 10 Nov. 1976
 (51) INT CL^a C07C 103/44; A61K 31/045, 31/085, 31/135, 31/16,
 31/17; C07C 103/82, 127/15, 127/17, 127/19 // 87/14
 (52) Index at acceptance

C2C 1173 1175 200 20Y 220 221 222 225 226 227 22Y
 280 282 29X 29Y 302 30Y 311 313 314 31Y 321
 322 323 326 32Y 332 338 339 340 341 342 34Y
 360 361 362 364 365 366 368 36Y 385 450 456
 45Y 502 50Y 510 512 51X 530 533 534 53X 581
 583 591 593 620 623 624 626 627 62X 630 633
 634 643 644 650 652 660 662 682 695 699 70X
 70Y 71X 790 79Y KB KD KH KJ LF SG SJ
 A5B 22Y 29X 29Y 303 30Y 334 33Y 382 384 386 387 38Y
 392 39X 420 422 426 42Y 430 431 433 43Y 480
 482 483 484 48Y 491 492 493 49Y 586 58Y 595
 596 597 59Y 605 606 60Y 640 642 646 64Y 77Y


(72) Inventor LESLIE HAROLD SMITH

(54) ARYLOXYALKANOLAMINE DERIVATIVES, THEIR
 PREPARATION AND USE IN PHARMACEUTICAL
 COMPOSITIONS


(71) We, IMPERIAL CHEMICAL INDUSTRIES LIMITED, Imperial Chemical House, Millbank, London, SW1P 3JF, a British Company, do hereby declare the invention, for which we pray that a patent may be granted to us, and the method by which it is to be performed, to be particularly described in and by the following statement:—

This invention relates to new alkanolamine derivatives which possess β -adrenergic blocking activity.

According to the invention there is provided a new alkanolamine derivative of the formula:—



wherein A stands for an alkylene radical of from 2 to 12 carbon atoms; wherein R¹ stands for the hydrogen atom or for an alkyl, halogenoalkyl, alkenyl or cycloalkyl radical each of up to 10 carbon atoms, or for an aryl radical of the formula:—



wherein R², R³, R¹² and R¹³, which may be the same or different, each stands for a hydrogen or halogen atom, a hydroxy, amino, nitro or cyano radical, an alkyl, cycloalkyl, alkenyl, alkynyl, alkoxy, alkylthio, cycloalkoxy, alkynyloxy, alkynylthio or alkanoyl radical each of up to 6 carbon atoms, or an aryl, aryloxy or aralkoxy radical each of up to 12 carbon atoms; or wherein R² and R³ together, and/or R¹² and R¹³ together, form the trimethylene, tetramethylene, 1-oxotetramethylene, propenylene, but-2-enylene or buta-1,3-dienylene radical such that together with the adjacent



benzene ring they form respectively the indanyl, 5,6,7,8-tetrahydronaphthyl, 5-oxo-5,6,7,8-tetrahydronaphthyl, indenyl, 5,8-dihydronaphthyl or naphthyl radical; wherein R^4 stands for the hydrogen atom or for the hydroxy or hydroxymethyl radical or for an aralkoxy radical of up to 12 carbon atoms; wherein R^{14} stands for the hydrogen atom or for the amino radical or for a dialkylamino radical of up to 12 carbon atoms; wherein X stands for the carbonyl ($-\text{CO}-$) or sulphonyl ($-\text{SO}_2-$) radical and wherein Y stands for a direct link, or for an alkylene, oxyalkylene or alkyleneoxy radical each of up to 6 carbon atoms, or for the imino ($-\text{NH}-$) radical, or for an alkylimino, iminoalkylene, iminoalkyleneoxy or iminoalkylenecarbonyloxy radical each of up to 6 carbon atoms, or (except where R^1 stands for the hydrogen atom) for the oxygen atom; or an acid-addition salt thereof.

It will be observed that the alkanolamine derivative of the invention possesses an asymmetric carbon atom, namely the carbon atom of the $-\text{CHOH}-$ group in the alkanolamine sidechain, and it can therefore exist in racemic and optically-active forms. It is to be understood that this invention encompasses the racemic form of the alkanolamine derivative and any optically-active form which possesses β -adrenergic blocking activity, it being a matter of common general knowledge how a racemic compound may be resolved into optically-active forms, and how the β -adrenergic blocking activity of these forms may be determined. It is further to be understood that β -adrenergic blocking activity usually predominates in that optically-active form which has the "S" absolute configuration of the said $-\text{CHOH}-$ group.

A suitable value for the alkylene radical A is, for example, an alkylene radical of from 2 to 10 carbon atoms, particularly of from 2 to 6 carbon atoms. A may be, for example, the ethylene, trimethylene, tetramethylene, hexamethylene, dodecamethylene, 1-methylethylene, 2-methylethylene or 1,1-dimethylethylene radical. A is preferably the ethylene, 1-methylethylene or 1,1-dimethylethylene radical.

A suitable value for R^1 when it stands for an alkyl, halogenoalkyl, alkenyl or cycloalkyl radical is, for example, the methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, t-butyl, n-pentyl, n-octyl, trifluoromethyl, allyl, cyclopropyl, cyclopentyl or cyclohexyl radical.

A suitable value for R^2 , R^3 , R^{12} or R^{13} when it stands for a halogen atom is the fluorine, chlorine, bromine or iodine atom.

A suitable value for R^2 , R^3 , R^{12} or R^{13} when it stands for an alkyl, cycloalkyl, alkenyl, alkynyl, alkoxy, alkylthio, cycloalkoxy, alkenyloxy, alkynyloxy or alkanoyl radical is, for example, the methyl, ethyl, n-propyl, cyclopropyl, cyclopentyl, allyl, ethynyl, methoxy, isopropoxy, methylthio, cyclopentyloxy allyloxy propargyloxy, formyl or acetyl radical.

A suitable value for R^2 , R^3 , R^{12} or R^{13} when it stands for an aryl or aryloxy radical is, for example, the phenyl or phenoxy radical.

A suitable value for R^2 , R^3 , R^4 , R^{12} or R^{13} when it stands for an aralkoxy radical is, for example the benzyloxy radical.

A suitable value for R^{14} when it stands for a dialkylamino radical is, for example, the dimethylamino radical.

A suitable value for Y when it stands for an alkylene, oxyalkylene or alkyleneoxy radical is, for example, the methylene, ethylene, oxymethylene, methyleneoxy, ethyleneoxy, trimethyleneoxy, 1-methylethylideneoxy or 1-methylpropylideneoxy radical.

A suitable value for Y when it stands for an alkylimino, iminoalkylene, iminoalkyleneoxy or iminoalkylenecarbonyloxy radical is, for example, the methylimino, iminomethylene, iminomethyleneoxy or iminomethylenecarbonyloxy radical.

A suitable acid-addition salt of an alkanolamine derivative of the invention is, for example, a salt derived from an inorganic acid, for example a hydrochloride, hydrobromide, phosphate or sulphate, or a salt derived from an organic acid, for example an oxalate, lactate, tartrate, acetate, salicylate, citrate, benzoate, β -naphthoate, adipate or 1,1-methylene-bis-(2-hydroxy-3-naphthoate), or a salt derived from an acidic synthetic resin, for example a sulphonated polystyrene resin.

A preferred alkanolamine derivative of the invention is a compound of the formula given above wherein A stands for the ethylene, 1-methylethylene or 1,1-dimethylethylene radical and wherein either

(a) R^1 stands for the hydrogen atom or for an alkyl, alkenyl or cycloalkyl radical each of up to 6 carbon atoms, or for the trifluoromethyl radical, or for a monohydroxyphenyl or dihydroxyphenyl radical, Y stands for a direct link or for an imino radical, X stands for the carbonyl radical, R^2 stands for the hydrogen atom or for a chloro, cyano, nitro, methyl, allyl, methoxy or allyloxy substituent which is in the *ortho*-

position of the benzene ring and R³ and R⁴ both stand for hydrogen atoms; or

(b) R¹ stands for a phenyl radical wherein R¹² stands for the hydrogen atom or for a chloro, nitro, methyl or methoxy substituent and R¹³ and R¹⁴ both stand for hydrogen atoms, Y stands for the methylene, ethylene, methyleneoxy or imino radical, X stands for the carbonyl radical, R² stands for the hydrogen atom or for a chloro, cyano, nitro, methyl, allyl, methoxy or allyloxy substituent which is in the *ortho*- or *meta*-position of the benzene ring and R³ and R⁴ both stand for hydrogen atoms; or

(c) R¹ stands for the hydrogen atom or for an alkyl radical of up to 6 carbon atoms, Y stands for the methyleneoxy radical, X stands for the carbonyl radical and R², R³ and R⁴ have the meanings stated in paragraph (a) above; or

(d) R¹ stands for an alkyl radical of up to 6 carbon atoms or for a phenyl radical wherein R¹² stands for the hydrogen atom or for a chloro, amino, nitro or methyl substituent and R¹³ and R¹⁴ both stand for hydrogen atoms, Y stands for a direct link, X stands for the sulphonyl radical and R², R³ and R⁴ have the meanings stated in paragraph (a) above; or

(e) R¹ stands for an alkyl or cycloalkyl radical each of up to 6 carbon atoms, or for a phenyl radical wherein R¹² stands for the hydrogen atom or for a methyl, methoxy or phenyl substituent and R¹³ and R¹⁴ both stand for hydrogen atoms, Y stands for a direct link or for the methylene or imino radical, X stands for the carbonyl or sulphonyl radical, R² stands for the hydrogen atom or for the hydroxy radical, R³ stands for the hydrogen atom and R⁴ stands for the hydroxy radical; or

(f) R¹ stands for an alkyl radical of up to 6 carbon atoms, Y stands for the oxygen atom or for the methyleneoxy, methylimino, iminomethyleneoxy or iminomethylene-carbonyloxy radical, X stands for the carbonyl or sulphonyl radical and R², R³ and R⁴ have the meanings stated in paragraph (e) above;
or an acid-addition salt thereof.

Specific alkanolamine derivatives of the invention are those hereinafter described in the Examples. Of these, preferred compounds by virtue of their high cardio-selective β -adrenergic blocking activity (as hereinafter defined) are:

- 1-phenoxy-3- β -benzamidoethylamino-2-propanol;
 - 1-phenoxy-3- β -benzenesulphonamidoethylamino-2-propanol;
 - 1-phenoxy-3- β -phenylacetamidoethylamino-2-propanol;
 - 1-phenoxy-3- β -phenoxyacetamidoethylamino-2-propanol;
 - 1-phenoxy-3- β -(3-phenylureido)ethylamino-2-propanol;
 - 1-phenoxy-3- β -(α -methyl- β -phenylacetamidoethyl)amino-2-propanol;
 - 1-phenoxy-3- β -pivalamidoethylamino-2-propanol;
 - 1-phenoxy-3- β -(β -isobutyramido- α -methylethyl)amino-2-propanol;
 - 1-(2-nitrophenoxy)-3- β -isobutyramidoethylamino-2-propanol;
 - 1-phenoxy-3- β -propionamidoethylamino-2-propanol;
 - 1-phenoxy-3- β -hydroxyacetamidoethylamino-2-propanol;
 - 1-phenoxy-3- β -(2-nitrobenzenesulphonamidoethyl)amino-2-propanol;
 - 1-(2-cyanophenoxy)-3- β -propionamidoethylamino-2-propanol;
 - 1-phenoxy-3- β -isopropylsulphonamidoethylamino-2-propanol;
 - 1-phenoxy-3- β -(2-chlorophenylacetamidoethyl)amino-2-propanol;
 - 1-(2-cyanophenoxy)-3- β -(3-phenylureidoethyl)amino-2-propanol;
 - 1-phenoxy-3- β -butyramidoethylamino-2-propanol;
 - 1-phenoxy-3- β -(3-n-butylureidoethyl)amino-2-propanol;
 - 1-phenoxy-3- β -isobutyramidoethylamino-2-propanol;
 - 1-phenoxy-3- β -(4-methoxyphenylacetamidoethyl)amino-2-propanol;
 - 1-phenoxy-3- β -acetamidoethylamino-2-propanol
 - 1-phenoxy-3- β -(n-propylsulphonamidoethyl)amino-2-propanol;
 - 1-(2-methoxyphenoxy)-3- β -isobutyramidoethylamino-2-propanol;
 - 1-phenoxy-3- β -(α , α -dimethyl- β -isobutyramidoethyl)amino-2-propanol; and
 - 1-phenoxy-3- β -(α , α -dimethyl- β -phenylacetamidoethyl)amino-2-propanol;
- and the acid-addition salts thereof.


Preferred compounds by virtue of their high level of cardiac stimulant activity (as hereinafter defined) are:—

- 1-(4-hydroxyphenoxy)-3-β-(3-phenylureido)ethylamino-2-propanol;
 1-(3,4-dihydroxyphenoxy)-3-β-isobutyramidoethylamino-2-propanol;
 1-(3,4-dihydroxyphenoxy)-3-β-(phenylureido)ethylamino-2-propanol;
 1-(3-hydroxyphenoxy)-3-β-(3-phenylureido)ethylamino-2-propanol;
 1-(3-hydroxyphenoxy)-3-β-benzenesulphonamidoethylamino-2-propanol;
 1-(4-hydroxyphenoxy)-3-β-(3,3-dimethylsulphamido)ethylamino-2-propanol;
 1-(4-hydroxyphenoxy)-3-β-(3-methoxymethylureido)ethylamino-2-propanol; and
 1-(4-hydroxyphenoxy)-3-β-(3-n-butyloxycarbonylmethylureido)ethylamino-2-propanol;
 and the acid-addition salts thereof.

The alkanolamine derivative of the invention may be manufactured by any chemical process publically known to be useful for the manufacture of chemically-analogous compounds.


According to a further feature of the invention there is provided a process for the manufacture of the alkanolamine derivative of the invention which comprises assembling in sequence, by chemical methods known to be suitable for this purpose, the five radicals:

- (i) an aryloxy radical of the formula:—



wherein R², R³ and R⁴ have the meanings stated above;

- (ii) an oxygenated three carbon radical of the formula:—



- wherein R⁵ stands for hydrogen or for a protecting group;

- (iii) an imino radical of the formula —NR⁶—, wherein R⁶ stands for hydrogen or for a protecting group;

- (iv) a radical of the formula:—



- wherein A has the meaning stated above and wherein R⁷ stands for hydrogen or for a protecting group; and


- (v) a radical of the formula:—




- wherein R¹, X and Y have the meanings stated above; whereafter if one or more of R², R³ and R⁴ stands for a protecting group, the one or more protecting groups are removed.

The various stages of the assembly may be carried out in any possible order. Thus, for example:


(a) a phenol of the formula:—



wherein R^2 , R^3 and R^4 have the meanings stated above, may first be reacted with an oxygenated three-carbon derivative, for example a compound of the formula:—



wherein R^5 has the meaning stated above, wherein Z^1 stands for a displaceable radical and wherein Z^2 stands for the hydroxy radical or for a displaceable radical. If Z^2 stands for the hydroxy radical, the intermediate compound obtained is further reacted with a reagent which will replace the primary hydroxy radical Z^2 with a displaceable radical Z^1 . The resulting product, which is a compound of the formula:—



wherein R^2 , R^3 and R^4 have the meanings stated above and wherein Z^3 stands for the group




wherein R^5 and Z^1 have the meanings stated above, or which may be, when R^5 stands for hydrogen, a mixture of such compounds wherein Z^3 has both meanings stated above, is then reacted with an amine of the formula:—



wherein A, R^1 , R^4 , R^7 , X and Y have the meanings stated above, or with a precursor of such an amine.

(b) An oxygenated three-carbon derivative, for example a compound of the formula:—




wherein R^5 , Z^1 and Z^2 have the meanings stated above, is reacted with an amine of the formula:—



wherein A, R¹, R², R', X and Y have the meanings stated above, or with a precursor of such an amine. If Z² stands for the hydroxy radical the intermediate compound obtained is further reacted with a reagent which will replace the primary hydroxy radical Z² with a displaceable radical Z'. The resulting product, which is a compound of the formula:—




wherein A, R¹, R², R', X, Y and Z² have the meanings stated above, or which may be, when R² stands for hydrogen, a mixture of such compounds wherein Z² has both meanings stated above, is then reacted with a phenol of the formula:—




wherein R², R³ and R⁴ have the meanings stated above. Alternatively, the compound of the formula:—



may be converted, by heating, into the azetidinol derivative of the formula:—



When R² stands for hydrogen, the azetidinol salt is converted into its free base form and then reacted with a phenol of the formula stated above. When R² stands for a protecting group, the azetidinium salt is reacted directly with the said phenol. The azetidinol derivative may alternatively be obtained by the reaction of a compound of the formula:—



wherein R² and Z¹ have the meanings stated above, with an amine of the formula:—



wherein A, R¹, R², R', X and Y have the meanings stated above.

A suitable value for Z¹, or for Z² when it stands for a displaceable radical, is, for example, a halogen atom, for example the chlorine or bromine atom, or a sulphonyloxy radical, for example an alkanesulphonyloxy radical of up to 6 carbon atoms or an arenesulphonyloxy radical of up to 10 carbon atoms, for example the methanesulphonyloxy, benzenesulphonyloxy or toluene-*p*-sulphonyloxy radical.

A suitable reagent which will replace the primary hydroxy radical Z² with a displaceable radical Z¹ is, for example, a halogenating agent, for example a thionyl halide, for example thionyl chloride or thionyl bromide, or a sulphonylating agent, for example an alkanesulphonyl halide or an arenesulphonyl halide, for example methanesulphonyl chloride, benzenesulphonyl chloride or toluene-*p*-sulphonyl chloride.

The reaction involving a phenol reactant may be carried out in the presence of an acid-binding agent, for example an alkali metal hydroxide, for example sodium hydroxide, or an organic base, for example piperidine. Alternatively, an alkali metal derivative of the phenol reactant, for example the sodium or potassium derivative,

may be used as starting material. The reaction may be carried out in a diluent or solvent, for example methanol or ethanol, and it may be accelerated or completed by the application of heat, for example by heating to the boiling point of the diluent or solvent.

The reaction involving an amine of the formula:—



may be carried out at ambient temperature or it may be accelerated or completed by the application of heat, for example by heating to a temperature of 90—110°C.; it may be carried out at atmospheric or at an elevated pressure, for example by heating in a sealed vessel; and it may be carried out in an inert diluent or solvent, for example methanol, ethanol or n-propanol, or an excess of the amine may be used as diluent or solvent.


(c) The series of reactions described under (a) or (b) above may be carried out except that an amine of the formula R^aNH_2 is used in place of an amine of the formula:—



it being understood that when R^a stands for hydrogen the amine is ammonia. The radical



may then be inserted as a separate step, for example either by the reaction of the final product from the series of reactions described under (a) or (b) above when an amine of the formula R^aNH_2 is used with a compound of the formula:—



wherein A, R^1 , R' , X, Y and Z^1 have the meanings stated above, or, when R^a stands for hydrogen, by the reaction under reducing conditions of the final product from the series of reactions described under (a) or (b) above when an amine of the formula R^aNH_2 is used with a carbonyl compound of the formula:—




wherein R^1 , R' , X and Y have the meanings stated above and wherein A^1 stands for hydrogen or for an alkyl radical and A^2 stands for an alkylene radical such that the radical



has the same meaning as is stated above for A.

The reaction involving a compound of the formula:—



may conveniently be carried out in the presence of a base, for example sodium or potassium carbonate, in a diluent or solvent, for example ethanol or isopropanol, at an elevated temperature, for example at the boiling point of the diluent or solvent.

Suitable reducing conditions for the reaction involving the carbonyl compound are those provided by the presence of hydrogen and a hydrogenation catalyst, for example palladium or platinum, in an inert diluent or solvent, for example in one or more solvents selected from water, ethanol and an excess of the carbonyl compound used as starting material; or by the presence of an alkali metal borohydride, for example sodium borohydride or lithium cyanoborohydride, in an inert diluent or solvent, for example in one or more solvents selected from water, ethanol, methanol and an excess of the carbonyl compound used as starting material. It is to be understood that when in the starting material R^1 stands for an alkenyl radical, or one or more of R^2 , R^3 , R^4 , R^{12} and R^{13} stands for a halogen atom or for a nitro, cyano, alkenyl, alkynyl, alkoxy, alkenyloxy, alkynyloxy or α -aralkoxy radical, hydrogen and a hydrogenation catalyst are preferably not used to provide the reducing conditions,

in order to prevent the radical R^1 , R^2 , R^3 , R^4 , R^{1a} or R^{1b} from being affected by catalytic hydrogenation.


(d) The series of reactions described under (a) or (b) above may be carried out except that an amine of the formula:—



wherein R^6 , R^7 and A have the meanings stated above, is used in place of an amine of the formula:—



or the reaction described under (c) above may be carried out except that the radical $-A-NHR^7$ is inserted in place of the radical $-A-NR^7-X-Y-R^1$. The amidic linkage $-NR^7-X-$ may then be formed as a separate step by reaction of the resulting product, which is a compound of the formula:—



wherein R^2 , R^3 , R^4 , R^5 , R^6 , R^7 and A have the meanings stated above, with a compound of the formula:—



wherein R^1 , X, Y and Z^1 have the meanings stated above, or, when X stands for the carbonyl radical and Y stands for the imino radical, with an isocyanate of the formula:—



wherein R^1 has the meaning stated above.

(e) A compound wherein one or more of R^5 , R^6 and R^7 stands for a protecting group may be prepared by the series of reactions described under (a) or (b) or (c) or (d) above. Alternatively, a suitable protecting group may be introduced by conventional means into an intermediate compound at any stage preceding the final stage.

A suitable value for R^5 when it stands for a protecting group is, for example, a hydrogenolysable radical, for example an α -arylalkyl, α -aralkoxy-carbonyl or α -arylalkoxymethyl radical, for example the benzyl benzyloxycarbonyl or benzyloxymethyl radical, or an acyl radical, for example an alkanoyl radical of up to 20 carbon atoms, for example the acetyl, t-butoxycarbonyl or 2,2,2-trichloroethoxycarbonyl radical, or an aroyl radical of up to 10 carbon atoms, for example the benzoyl radical, or an α -alkoxyalkyl radical (that is, a radical which forms with the oxygenated three-carbon radical an acetal radical), for example the tetrahydropyranyl radical, or a tertiary alkyl radical, for example the t-butyl radical.

A suitable value for R^6 when it stands for a protecting group is, for example, a hydrogenolysable or tertiary alkyl radical as defined for R^5 , or a relatively easily hydrolysable acyl radical, for example the 2,2,2-trichloroethoxycarbonyl or t-butoxycarbonyl radical. It is to be understood that when R^6 stands for an acyl radical, this radical must be removable under conditions which will not destroy the amidic linkage $-NR^7-X$.

Alternatively, R^5 and R^6 may be joined together so that one protecting group serves to protect both the oxygen and nitrogen atoms. Such a protecting group may be, for example, a radical of the formula $-CHR^8-$, wherein R^8 stands for hydrogen, or for an alkyl radical of up to 4 carbon atoms or an aryl radical of up to 10 carbon atoms, such that it forms, together with the adjacent oxygen and nitrogen atoms and two carbon atoms of the three-carbon radical, an oxazolidine nucleus.

A suitable value for R^7 when it stands for a protecting group is, for example, a hydrogenolysable or tertiary alkyl group as defined for R^5 or R^6 .

The hydrogenolysable protecting group R^5 , R^6 or R^7 may be removed, for example, by catalytic hydrogenolysis, for example by hydrogenation in the presence of a palladium-on-charcoal catalyst, in an inert diluent or solvent, for example ethanol or aqueous ethanol. The process may be accelerated or completed by the presence of an acidic catalyst, for example hydrochloric or oxalic acid.


The acyl protecting group R^5 or R^6 may be removed by hydrolysis in the presence of a base, for example an alkali metal hydroxide, in a diluent or solvent, for example water, methanol, ethanol or a mixture thereof. It is to be understood that the hydrolytic conditions used must be sufficiently mild to avoid hydrolysis of the amidic linkage $-NR^7-X$.

The α -alkoxyalkyl protecting group R^5 or the protecting group $-R^6CH-$ formed by R^5 and R^6 taken together may be removed by hydrolysis in the presence of an acid, for example a mineral acid, for example aqueous hydrochloric acid, and the hydrolysis may be carried out at a temperature of up to 100°C .

The tertiary alkyl protecting group R^5 , R^6 or R^7 , or the acyl protecting group R^5 or R^6 when it stands for a tertiary alkoxycarbonyl radical, for example the *t*-butoxycarbonyl radical, may be removed by treatment with an acid, for example hydrogen chloride, under anhydrous conditions, for example in an ethereal solution.

A compound wherein one or more of R^2 , R^3 , R^4 , R^{12} and R^{13} stands for an α -arylalkoxy radical, for example the benzyloxy radical, may be converted into the corresponding compound wherein one or more of R^2 , R^3 , R^4 , R^{12} and R^{13} stands for the hydroxy radical by hydrogenolysis.

A preferred process for the manufacture of the alkanolamine derivative of the invention comprises the reaction of a compound of the formula:—



wherein R^2 , R^3 and R^4 have the meanings stated above (both of which compounds may be obtained by the reaction of the corresponding phenol with epichlorohydrin), with an amine of the formula:—



wherein A, R^1 , X and Y have the meanings stated above and wherein R^6 stands for hydrogen or for the benzyl radical, whereafter if R^6 stands for the benzyl radical this radical is removed by hydrogenolysis.

Optically-active enantiomorphs of the alkanolamine derivative of the invention may be obtained by the resolution by conventional means of the corresponding racemic alkanolamine derivative of the invention.

The said resolution may be carried out by reacting the racemic alkanolamine derivative with an optically-active acid, followed by fractional crystallisation of the diastereoisomeric mixture of salts thus obtained from a diluent or solvent, for example ethanol, whereafter the optically-active alkanolamine derivative is liberated from the salt by treatment with a base. A suitable optically-active acid is, for example, (+)- or (-)-O,O-di-*p*-toluoyltartaric acid or (-)-2,3,4,5-di-*O*-isopropylidene-2-keto-L-gulonic acid.

The resolution process may be facilitated by treating the partially resolved alkanolamine derivative in free base form obtained after a single fractional crystallisation of the diastereoisomeric mixture of salts with a solubilising agent, for example a primary amine, for example allylamine, in a relatively non-polar diluent or solvent, for example petroleum ether.

The alkanolamine derivative of the invention in free base form may be converted into an acid-addition salt thereof by reaction with an acid by conventional means.

As stated above, the alkanolamine derivative of the invention or an acid-addition salt thereof possesses β -adrenergic blocking activity, and furthermore this activity is cardioselective. This activity may be determined by the reversal of isoprenaline-induced tachycardia in rats or cats, a standard test for the determination of β -adrenergic blocking activity, and by relative freedom from antagonism of isoprenaline-induced vasodilation in cats or of the relief produced by isoprenaline of histamine-induced bronchospasm in guinea-pigs. Compounds exhibiting this cardioselective action show a greater degree of specificity in blocking the cardiac β -receptors than the β -receptors in peripheral blood vessels and bronchial muscle. Thus, a dose may be selected for such a compound at which the compound blocks the cardiac inotropic and chronotropic actions of a catecholamine such as isoprenaline but does not block the relaxation of tracheal smooth muscle produced by isoprenaline or the peripheral vasodilator action of isoprenaline. Because of this selective action, one of those compounds may advantageously be used together with a sympathomimetic bronchodilator, for example isoprenaline, orciprenaline, adrenaline or ephedrine, in the treatment of asthma and other obstructive airways diseases, inasmuch as the cardioselective compound will substantially inhibit the unwanted stimulatory effects of the bronchodilator on the heart but will not hinder the desirable therapeutic effect of the bronchodilator. A preferred alkanolamine derivative of the invention is three to ten times more active as a cardioselective β -adrenergic blocking agent than practolol. At doses of an alkanolamine derivative of the invention which produce effective β -adrenergic blockade in rats or cats, no symptoms of toxicity are apparent.

Some of the alkanolamine derivatives of the invention wherein one or more of the substituents R^2 , R^3 and R^4 stands for the hydroxy radical, and in particular those wherein R^4 stands for a hydroxy radical in the 3- or 4-position of the benzene nucleus, R^2 stands for the hydrogen atom or for a hydroxy radical in the 3-position when R^4 is in the 4-position of the benzene nucleus and R^3 stands for the hydrogen atom possess, in addition to β -adrenergic blocking activity, substantial cardiac stimulant activity. This may be demonstrated in either conscious or pentobarbitone-anaesthetised dogs, where the alkanolamine derivative or salt thereof produces an increase in heart rate, and/or an increase in force of contraction of the heart and an increase in the speed of conduction of electrical activity through the tissues of the heart. Unlike isoprenaline, a known cardiac stimulating agent, a preferred stimulant alkanolamine derivative of the invention or a salt thereof is well absorbed when administered orally and has a substantial duration of action. At doses of an alkanolamine derivative of the invention which produce effective cardiac stimulation in dogs, no symptoms of toxicity are apparent.

The alkanolamine derivative of the invention may be administered to warm-blooded animals, including man, in the form of a pharmaceutical composition comprising as active ingredient at least one alkanolamine derivative of the invention, or an acid-addition salt thereof, in association with a pharmaceutically-acceptable diluent or carrier therefor.

A suitable composition is, for example, a tablet, capsule, aqueous or oily solution or suspension, emulsion, injectable aqueous or oily solution or suspension, dispersible powder, spray or aerosol formulation.

The pharmaceutical composition may contain, in addition to the alkanolamine derivative of the invention, one or more drugs selected from sedatives, for example phenobarbitone, meprobamate, chlorpromazine and the benzodiazepine sedative drugs, for example chlordiazepoxide and diazepam; vasodilators, for example glycerol trinitrate, pentaerythritol tetranitrate and isosorbide dinitrate, diuretics, for example chlorothiazide; hypotensive agents, for example reserpine, bethanidine and guanethidine; cardiac membrane stabilising agents, for example quinidine; agents used in the treatment of Parkinson's disease and other tremors, for example benzhexol; cardiotonic agents, for example digitalis preparations; α -adrenergic blocking agents, for example phentolamine and sympathomimetic bronchodilators, for example isoprenaline, orciprenaline, adrenaline and ephedrine.

When used for the treatment of heart diseases, for example angina pectoris and cardiac arrhythmias, or for the treatment of hypertension or anxiety states in man, it is expected that the alkanolamine derivative would be given to man at a total oral dose of between 20 mg. and 600 mg. daily, at doses spaced at 6—8 hourly intervals, or at an intravenous dose of between 1 mg. and 20 mg.

When used for the treatment of acute or chronic heart failure in man, it is expected that a cardiac stimulant alkanolamine derivative would be given to man at

a total oral dose of between 10 mg. and 200 mg. daily, at doses spaced at 6—8 hourly intervals or at an intravenous dose of between 1 mg. and 100 mg.

Preferred oral dosage forms are tablets or capsules containing between 10 and 100 mg., and preferably 10 mg. or 50 mg. of active ingredient. Preferred intravenous dosage forms are sterile aqueous solutions of the alkanolamine derivative or of a non-toxic acid-addition salt thereof, containing between 0.05% and 1% w/v of active ingredient, and more particularly containing 0.1% w/v of active ingredient.

The invention is illustrated but not limited by the following Examples:—

Example 1.

A mixture of 3.0 g. of 2,3-epoxy-1-phenoxypropane, 50 ml. of n-propanol, 4.0 g. of β -benzamidoethylamine hydrochloride and a solution of 0.8 g. of sodium hydroxide in 5 ml. of water is heated under reflux for 18 hours. The mixture is evaporated to dryness under reduced pressure and the residue is stirred with 40 ml. of aqueous 2N-hydrochloric acid. The mixture is filtered and the solid residue is crystallised from ethanol. There is thus obtained 3 - β - benzamidoethylamino - 1 - phenoxy - 2 - propanol hydrochloride, m.p. 198—199°C.

Example 2.

A mixture of 0.75 g. of 2,3-epoxy-1-phenoxypropane, 25 ml. of n-propanol, 1.18 g. of β -benzenesulphonamidoethylamine hydrochloride and a solution of 0.42 g. of sodium bicarbonate in 5 ml. of water is heated under reflux for 18 hours. The mixture is evaporated to dryness under reduced pressure and the residue is stirred with 25 ml. of aqueous N-sodium hydroxide solution. The mixture is washed three times with 25 ml. of ether each time, and the aqueous alkaline phase is neutralised with glacial acetic acid. The mixture is extracted three times with 25 ml. of ethyl acetate each time, and the combined extracts are dried over anhydrous magnesium sulphate and filtered. The filtrate is added to a solution of 0.6 g. of oxalic acid in 30 ml. of ethyl acetate. The mixture is filtered and the solid residue is washed with ethyl acetate and crystallised from acetonitrile. There is thus obtained 1 - phenoxy - 3 - β - benzene-sulphonamidoethylamino - 2 - propanol oxalate, m.p. 135—137°C.

Example 3.

A mixture of 1.5 g. of 2,3-epoxy-1-phenoxypropane, 50 ml. of n-propanol, 2.31 g. of β -phenoxyacetamidoethylamine hydrochloride and a solution of 0.4 g. of sodium hydroxide in 5 ml. of water is heated under reflux for 18 hours. The mixture is evaporated to dryness under reduced pressure and the residue is stirred with 20 ml. of aqueous N-sodium hydroxide solution. The mixture is extracted three times with 20 ml. of ethyl acetate each time, and the combined extracts are dried over anhydrous magnesium sulphate and filtered. The filtrate is added to a solution of 2.5 g. of oxalic acid in 100 ml. of ether. The mixture is filtered and the solid residue is crystallised from acetonitrile. There is thus obtained 1 - phenoxy - 3 - β - phenoxyacetamidoethylamino - 2 - propanol oxalate, m.p. 131—133°C.

The β -phenoxyacetamidoethylamine hydrochloride used as starting material may be obtained as follows:—

A mixture of 8.3 g. of methyl phenoxyacetate and 10.8 ml. of ethylenediamine is heated at 90°C. for 18 hours and then cooled and stirred with 70 ml. water. The mixture is filtered and the filtrate is evaporated to dryness under reduced pressure. The residue is dissolved in ethyl acetate, dried over anhydrous magnesium sulphate and filtered, and the filtrate is acidified with ethereal hydrochloric acid. The mixture is filtered and the solid residue is crystallised from ethanol. There is thus obtained β -phenoxyacetamidoethylamine hydrochloride, m.p. 159—160°C.

Example 4.


A mixture of 1.5 g. of 2,3-epoxy-1-phenoxypropane, 50 ml. of n-propanol, 2.15 g. of β -phenylacetamidoethylamine hydrochloride and a solution of 0.4 g. of sodium hydroxide in 5 ml. of water is heated under reflux for 18 hours. The mixture is evaporated to dryness under reduced pressure and the residue is stirred with 20 ml. of aqueous N-sodium hydroxide solution. The mixture is extracted three times with 20 ml. of ethyl acetate each time and the combined extracts are dried over anhydrous magnesium sulphate and filtered. The filtrate is evaporated to dryness under reduced pressure and the residue is crystallised from ethyl acetate. There is thus obtained 1 - phenoxy - 3 - β - phenylacetamidoethylamino - 2 - propanol, m.p. 124—125°C.

Example 5.

A mixture of 1.5 g. of 2,3-epoxy-1-phenoxypropane, 50 ml. of n-propanol and 1.79 g. of β -(3-phenylureido)ethylamine is heated under reflux for 18 hours. The mixture is evaporated to dryness under reduced pressure and the residue is crystallised from ethyl acetate. There is thus obtained 1 - phenoxy - 3 - β - (3 - phenylureido)-ethylamino - 2 - propanol, m.p. 144—145°C.

Example 6.

The process described in Example 1 is repeated except that the appropriate 2,3-epoxy-1-phenoxypropane (or the corresponding 3-chloro-1-phenoxy-2-propanol) and the appropriate β -carboxamidoethylamine are used as starting materials. There are thus obtained the compounds described in the following table:—



R^1	R^2	R^3	m.p. (°C.)	Crystallisation solvent
methyl	H	H	103–105	ethyl acetate
ethyl	H	H	126–127	acetonitrile
n-propyl	H	H	86–87	ethyl acetate
isopropyl	H	H	125–126	ethyl acetate
n-octyl	H	H	95–96	ethyl acetate
cyclopropyl	H	H	125–126	ethyl acetate
cyclopentyl	H	H	133–135	ethyl acetate
cyclohexyl	H	H	138–139	acetonitrile
2-chlorophenyl	H	H	hydrochloride 182–183	acetonitrile
4-tolyl	H	H	hydrochloride 211–212	acetonitrile
2-nitrophenyl	H	H	131–132	ethyl acetate/ petroleum ether
2-methoxy- phenyl	H	H	hydrogen oxalate 164–165	ethanol
2-dimethyl- aminophenyl	H	H	82–84	ethyl acetate
isopropyl	2-cyano	H	109–110	ethyl acetate
isopropyl	2-nitro	H	108–110	ethyl acetate
isobutyl	H	H	114–115	ethyl acetate
t-butyl	H	H	hydrogen oxalate 167–168	ethanol
n-pentyl	H	H	101–102	ethyl acetate
ethyl	2-cyano	H	96–97	ethyl acetate/ petroleum ether
isopropyl	2-chloro	H	129–130	ethyl acetate
isopropyl	2-methyl	H	120–122	ethyl acetate
isopropyl	2-methoxy	H	142–143	acetonitrile
isopropyl	2-chloro	4-methyl	119–120	ethyl acetate
isopropyl	2-chloro	5-chloro	152–153	acetonitrile
isopropyl	2-nitro	4-methoxy	122–124	ethyl acetate

R ¹	R ²	R ³	m.p. (°C.)	Crystallisation solvent
methyl	3-cyano	H	121-122	ethyl acetate
4-benzyloxy-phenyl	H	H	124-126	acetonitrile
isopropyl	2-benzyl-oxy	H	137-138	acetonitrile
3,4-dibenzyl-oxyphenyl	H	H	hydrochloride 223-224	ethanol

Many of the β -carboxamidoethylamines used as starting materials are novel compounds, and may be prepared by a process exemplified by the following preparation of β -cyclopentanecarboxamidoethylamine:—

A mixture of 12.8 g. of methane cyclopentanecarboxylate and 26.6 ml. of ethylenediamine is heated at 90°C. for 18 hours and then cooled and stirred with 100 ml. of water. The mixture is filtered and the filtrate is evaporated to dryness under reduced pressure. The residue is extracted with 50 ml. of toluene and the extract is evaporated to dryness under reduced pressure. The residue is then dissolved in 200 ml. of ethyl acetate and the solution is added to a solution of 2.4 g. of oxalic acid and in a mixture of 250 ml. of acetone and 250 ml. of ethyl acetate. The mixture is filtered and the solid residue is crystallised from ethanol. There is thus obtained β -cyclopentanecarboxamidoethylamine hydrogen oxalate, m.p. 164-165°C.

The following amines may similarly be prepared from the appropriate methyl or ethyl ester and ethylenediamine:—



R ¹	Salt	m.p. (°C.)	Crystallisation solvent
n-octyl	hydrochloride	217-221	ethanol
cyclopropyl	hydrogen oxalate	147-149	ethanol
cyclohexyl	hydrogen oxalate	154-155	ethanol/ethyl acetate
2-chlorophenyl	hydrochloride	164-166	ethanol/ethyl acetate
2-methoxy-phenyl	hydrogen oxalate	163-164	ethanol
2-dimethyl-aminophenyl	dihydrochloride	220-221	ethanol
ethyl	hydrogen oxalate	133-134	ethanol

R ¹	Salt	m.p. (°C.)	Crystallisation solvent
n-propyl	hydrogen oxalate	120-121	ethanol/ethyl acetate
isopropyl	hydrogen oxalate	130-131	ethanol
isobutyl	hydrogen oxalate	100-102	ethanol/ethyl acetate
t-butyl	hydrogen oxalate	188-189 (with decomposition)	ethanol
n-pentyl	(free base)	b.p. 142°C./0.2 mm.Hg	—
4-benzyloxy-phenyl	(free base)	118-120	toluene
3,4-dibenzyl-oxyphenyl	(free base)	128-130	ethanol

Example 7.

The process described in Example 2 is repeated except that the appropriate 2,3-epoxy-1-phenoxypropane and the appropriate β -sulphonamidoethylamine are used as starting materials. There are thus obtained the compounds described in the following table:—




R ¹	R ²	m.p. (°C.)	Crystallisation solvent
methyl	H	155–156	methanol
4-tolyl	H	145–148	ethanol
4-chlorophenyl	H	166–169	ethanol
2-nitrophenyl	H	186–188	acetonitrile
3-nitrophenyl	H	166–167	methanol
isopropyl	H	oxalate 181–183	ethanol
n-propyl	H	87–88	ethyl acetate/petroleum ether
phenyl	2-cyano	hydrogen oxalate monohydrate 136–138 (with decomposition)	ethanol
4-aminophenyl	H	oxalate 158–160 (with decomposition)	ethanol
3-amino-4-methyl-5-nitrophenyl	H	hemioxalate 223–225 (with decomposition)	ethyl acetate

The β -methanesulphonamidoethylamine used as starting material may be obtained as follows:—

Methanesulphonyl chloride (5.7 g.) is added dropwise during 10 minutes to a cooled solution of 30 g. of ethylenediamine in 100 ml. of water and the mixture is stirred for a further 30 minutes and then neutralised with 4.2 g. of sodium bicarbonate. The mixture is evaporated to dryness under reduced pressure and the residue is extracted three times with 200 ml. of boiling acetonitrile each time. The combined extracts are dried over anhydrous magnesium sulphate and evaporated to dryness under reduced pressure and the residue is dissolved in a mixture of 90 ml. of ethyl acetate and 10 ml. of methanol. The solution is added to a solution of 6 g. of oxalic acid in 100 ml. of ethyl acetate, the mixture is filtered and the solid residue is washed with ethyl acetate. There is thus obtained 2-methanesulphonamidoethylamine hydrogen oxalate, m.p. 170–174°C. (with decomposition).


Other novel β -sulphonamidoethylamines which may similarly be prepared and which have been characterised are described in the following table:—



R ¹	Salt	m.p. (°C.)	Crystallisation solvent
isopropyl	hydrogen oxalate	166-167	ethanol
n-propyl	hydrogen oxalate hemihydrate	145-147	ethanol
3-amino-4-methyl-5-nitrophenyl	hydrochloride	222-225	ethanol

Example 8.

The process described in Example 4 is repeated except that the appropriate 2,3-epoxy-1-phenoxypropane and the appropriate β -carboxamidoethylamine are used as starting materials. There are thus obtained the compounds described in the following table:—



R ¹	Y	R ²	m.p. (°C.)	Crystallisation solvent
4-chlorophenyl	-CH ₂ -	H	119-120	acetonitrile
2-chlorophenyl	-CH ₂ -	H	137-138	ethyl acetate
phenyl	-CH ₂ CH ₂ -	H	112-113	ethyl acetate/petroleum ether
4-chlorophenyl	-CH ₂ O-	H	163-164	methanol
phenyl	-CH ₂ CH ₂ CH ₂ O-	H	89-91	ethyl acetate/petroleum ether
ethyl	-O-	H	87-88	ethyl acetate/petroleum ether
H	-CH ₂ O-	H	hydrogen oxalate 128-129	ethanol
methyl	-CH ₂ O-	H	hydrogen oxalate 135-137 (with decomposition)	ethanol
2-nitrophenyl	-CH ₂ -	H	130-131	ethyl acetate
4-nitrophenyl	-CH ₂ -	H	135-136	ethyl acetate
4-methoxyphenyl	-CH ₂ -	H	122-123	ethyl acetate
2-chlorophenyl	-CH ₂ -	2-cyano	120-122	acetonitrile
2-chlorophenyl	-CH ₂ -	2-chloro	144-146	ethyl acetate
4-chlorophenyl	-C(CH ₃) ₂ O-	H	hydrogen oxalate 172-174	ethanol
2-methoxyphenyl	-CH ₂ O-	H	hydrogen oxalate 156-158	acetonitrile

The β-carboxamidoethylamines used as starting material may be prepared by a similar process to that described in the second part of Example 6. Those amines which have been characterised are described in the following table:—



R^1-Y-	Salt	m.p. (°C.)	Crystallisation solvent
4-chlorophenyl-CH ₂ -	hydrochloride	194-196	ethanol/ethyl acetate
2-chlorophenyl-CH ₂ -	hydrochloride	162-163	ethanol/ethyl acetate
phenyl-CH ₂ CH ₂ -	hydrochloride	134-137	ethanol/ethyl acetate
4-chlorophenyl-OCH ₂ -	<i>p</i> -chlorophenoxy-acetate	147-148	acetonitrile
phenyl-OCH ₂ CH ₂ CH ₂ -	hydrochloride	137-140	acetonitrile
methyl-OCH ₂ -	hydrogen oxalate	137-138	ethanol
4-methoxyphenyl-CH ₂ -	(free base)	130-140	toluene
4-chlorophenyl-OC(CH ₃) ₂ -	hydrochloride	98-100	acetonitrile

The β -hydroxyacetamidoethylamine used as starting material may be obtained by heating 2-hydroxymethylimidazoline (m.p. 87-88°C., obtained from ethyl glycolate and ethylenediamine), under reflux with water for 15 minutes.

Example 9.

The process described in Example 5 is repeated except that the appropriate 2,3-epoxy-1-phenoxypropane and the appropriate β -ureidoethylamine are used as starting materials. There are thus obtained the compounds described in the following table:—




R ¹	R ²	m.p. (°C.)	Crystallisation solvent
ethyl	H	118–120	ethyl acetate
isopropyl	H	145–147	acetonitrile
n-butyl	H	116–117	ethyl acetate
allyl	H	131–132	ethyl acetate
4-tolyl	H	147–149	ethyl acetate
4-methoxyphenyl	H	126–128	ethyl acetate
phenyl	2-chloro	157–158	acetonitrile
phenyl	2-nitro	159–161	acetonitrile
phenyl	2-methyl	160–161	acetonitrile
phenyl	2-cyano	155–156	acetonitrile
n-butyl	2-chloro	149–150	acetonitrile
n-butyl	2-methyl	136–137	acetonitrile
n-butyl	2-cyano	145–147	acetonitrile

The β -(3-n-butylureido)ethylamine used as starting material may be obtained as follows:—

A solution of 22.6 ml. of n-butyl isocyanate in 50 ml. of chloroform is added dropwise during 45 minutes to a solution of 26.6 ml. of ethylenediamine in 50 ml. of chloroform which is cooled to between -10°C . and 0°C . The mixture is then stirred at laboratory temperature for 2 hours and filtered, and the filtrate is evaporated to dryness under reduced pressure. The residue is dissolved in 150 ml. of toluene and the mixture is evaporated to dryness. The residue is dissolved in 50 ml. of ethyl acetate and the solution is added to a solution of 12.4 g. of oxalic acid in 100 ml. of ethyl acetate. The ethyl acetate is removed by decantation and the residue is triturated with 50 ml. of acetonitrile. The mixture is filtered and the solid residue is crystallised from ethanol. There is thus obtained β -(3-n-butylureido)ethylamine hydrogen oxalate, m.p. $138-139^\circ\text{C}$.


Other novel β -ureidoethylamines which may similarly be prepared and which have been characterised are described in the following table:—



R ¹	Salt	m.p. (°C.)	Crystallisation solvent
ethyl	hydrochloride	122-124	isopropanol ethyl acetate
isopropyl	hydrogen oxalate	146-148	ethanol acetonitrile
allyl	hydrochloride	128-130	isopropanol
4-tolyl	hydrochloride	234-236	ethyl acetate
4-methoxyphenyl	(free base)	104-106	ethyl acetate

Example 10.

The process described in Example 5 is repeated except that the appropriate 2,3-epoxy-1-phenoxypropane and the appropriate β -carboxamidoalkylamine are used as starting materials. There are thus obtained the compounds described in the following table:—



R ¹	Y	A	R ²	m.p. (°C.)	Crystallisation solvent
isopropyl	direct	-(CH ₂) ₄ -	H	hydrogen oxalate 128-130	acetonitrile
methyl	direct	-(CH ₂) ₆ -	H	hydrogen oxalate 111-113	acetonitrile
phenyl	-CH ₂ -	-CHCH ₂ - CH ₃	H	124-126	ethyl acetate
phenyl	-CH ₂ -	-CHCH ₂ - CH ₃	2-cyano	124-128	ethyl acetate
isopropyl	direct	-CHCH ₂ - CH ₃	H	124-126	ethyl acetate
n-pentyl	direct	-CHCH ₂ - CH ₃	H	102-103	ethyl acetate
isopropyl	direct	-(CH ₂) ₃ -	H	94-95	ethyl acetate
n-pentyl	direct	-(CH ₂) ₆ -	H	85-86	ethyl acetate
phenyl	-CH ₂ -	-CHCH ₂ - CH ₃	2-nitro	115-117	ethyl acetate
phenyl	-CH ₂ -	-CHCH ₂ - CH ₃	2-allyloxy	102-105	ethyl acetate

The 1-methyl-2-(phenylacetamido)ethylamine used as starting material may be obtained as follows:—

A mixture of 32.8 g. of ethyl phenylacetate and 44.4 g. of 1,2-diaminopropane is heated at 90°C. for 18 hours and then dissolved in 100 ml. of toluene. The solution is evaporated to dryness under reduced pressure, the residue is triturated with 100 ml. of acetonitrile and 200 ml. of ether and the mixture is filtered. The solid residue is crystallised from acetonitrile and then stirred with a mixture of 10 ml. of aqueous 2N-sodium hydroxide solution and 20 ml. of saturated aqueous sodium chloride solution. The mixture is extracted four times with 50 ml. of chloroform each time and the combined extracts are dried over anhydrous magnesium sulphate and evaporated to dryness under reduced pressure. The residue is triturated with ether and petroleum ether (b.p. 60–80°C.) and the mixture is filtered. The solid residue is washed with petroleum ether (b.p. 60–80°C.) and there is thus obtained 1-methyl-2-(phenylacetamido)ethylamine, m.p. 46–48°C.

There may similarly be obtained:—

from ethyl isobutyrate and 1,2-diaminopropane, 1-methyl-2-isobutyramidoethylamine, b.p. 106°C./0.3 mm. Hg (hydrogen oxalate m.p. 141–143°C. after crystallisation from a mixture of ethanol and ethyl acetate);

from ethyl hexanoate and 1,2-diaminopropane, 1-methyl-3-hexanamidoethylamine, b.p. 136°C./0.15 mm. Hg;

from methyl isobutyrate and 1,3-diaminopropane, γ -isobutyramidopropylamine, b.p. 120–122°C./0.2 mm. Hg;

from ethyl isobutyrate and 1,4-diaminobutane, 4-isobutyramidobutylamine (hydrochloride m.p. 153–154°C. after crystallisation from a mixture of ethanol and ethyl acetate);

from ethyl acetate and 1,6-diaminohexane, 6-acetamidohexylamine, b.p. 123–130°C./0.13 mm. Hg.

Example 11.

A mixture of 0.64 g. of 3-benzylamino-1-phenoxy-2-propanol, 1.44 g. of β -(2-methyl-2-*p*-chlorophenoxybutyramido)ethyl chloride and 0.42 g. of sodium bicarbonate is heated at 120°C. for 18 hours and then cooled and stirred with 25 ml. of ethyl acetate and 25 ml. of water. The ethyl acetate phase is separated, dried over anhydrous magnesium sulphate and evaporated to dryness under reduced pressure. There is thus obtained as an oil, which is used without further purification, 1-phenoxy-3-[N-benzyl-N- β -(2-methyl-2-*p*-chlorophenoxybutyramido)ethyl]amino-2-propanol.

A mixture of 1.9 g. of the above compound, 40 ml. of ethanol, 1 ml. of 11 N-aqueous hydrochloric acid and 0.2 g. of a 30% w/w palladium-on-charcoal catalyst is shaken with hydrogen at laboratory temperature and atmospheric pressure until the uptake of hydrogen ceases. The mixture is filtered and the filtrate is evaporated to dryness under reduced pressure. The residue is dissolved in 50 ml. of ether and the solution is added to a solution of 0.6 g. of oxalic acid in 25 ml. of ethyl acetate. The mixture is filtered and the solid residue is crystallised from acetonitrile. There is thus obtained 1-phenoxy-3- β -(2-methyl-2-phenoxybutyramido)ethylamino-2-propanol oxalate, m.p. 112–113°C. (the chlorine atom being removed during the hydrogenolysis reaction).

Example 12.


A solution of 2.5 g. of 1-phenoxy-3- β -(*p*-benzyloxybenzamido)ethylamino-2-propanol (Example 6) in 50 ml. of a 1:1 v/v mixture of ethanol and acetic acid is shaken for 1 hour with hydrogen at laboratory temperature and atmospheric pressure in the presence of 400 mg. of a 5% w/w palladium-on-charcoal, when 180 ml. of hydrogen have been absorbed. The mixture is filtered, the filtrate is evaporated to dryness under reduced pressure and the residue is dissolved in water. The solution is extracted with ethyl acetate and the extract is dried and evaporated to dryness. The residue is dissolved in ethyl acetate and a solution of oxalic acid in acetone is added. The mixture is filtered and the solid residue is crystallised from ethanol. There is thus obtained 1-phenoxy-3- β -(*p*-hydroxybenzamido)ethylamino-2-propanol hydrogen oxalate, m.p. 152–154°C. (with decomposition).

Example 13.

A mixture of 2.5 g. of 1-*p*-benzyloxyphenoxy-2,3-epoxypropane, 1.79 g. of β -(3-phenylureido)ethylamine and 20 ml. of isopropanol is heated under reflux for

3 hours and then cooled and evaporated to dryness under reduced pressure. The residue is suspended in acetic acid and shaken for 30 minutes in the presence of a 30% w/w palladium-on-charcoal catalyst and in an atmosphere of hydrogen, at laboratory temperature and atmospheric pressure, when 255 ml. of hydrogen have been absorbed. The mixture is filtered, the filtrate is evaporated to dryness under reduced pressure and the residue is dissolved in methanol. A solution of oxalic acid in ether is added and the mixture is filtered. The solid residue is crystallised from a mixture of methanol and ether and there is thus obtained 1 - *p* - hydroxyphenoxy - 3 - β - (3-phenylureido)ethylamino - 2 - propanol hemioxalate, m.p. 164—165°C.

The process described above is repeated except that the appropriate 1-benzyloxyphenoxy-2,3-epoxypropane (prepared by conventional means from the corresponding benzyloxyphenol and epichlorohydrin) and the appropriate β -substituted-ethylamine derivative (prepared as hereinbefore described) are used as starting materials, and there are thus obtained the compounds described in the following tables:—



position of HO-substituent	R ¹ -Y-	base or salt	m.p. (°C.)	Crystallisation solvent
2-	phenyl-NH-	base	154	ethanol
4-	ethyl-	hemi-oxalate	149	ethanol/ether
4-	methoxy-CH ₂ -	oxalate	199 (with decomposition)	ethanol
4-	phenyl-	acetate	145-146	ethanol
4-	phenyl-CH ₂ -	hemi-oxalate hydrate	110 (with decomposition)	ethanol
3-	isopropyl	base	(oil)	—
2-	isopropyl	hydrogen oxalate	168-170	ethanol/acetonitrile



position of HO-substituent	R ¹	base or salt	m.p. (°C.)	Crystallisation solvent
4-	phenyl	hemi-oxalate	158-159	methanol
3-	phenyl	oxalate hydrate	(glass)	—
4-	methyl	oxalate	119.5-120 (with decomposition)	ethanol

1 - (*p* - Hydroxyphenoxy) - 3 - γ - isobutyramidopropylamino - 2 - propanol hemioxalate hemihydrate has m.p. 179—180°C. (crystallised from a mixture of methanol and ether).

Example 14.

p-Tolyl isocyanate (2.9 g.) is added slowly to a solution of 8.12 g. of 1 - *p*-benzyloxyphenoxy - 3 - (*N* - benzyl - *N* - β - aminoethyl)amino - 2 - propanol in 50 ml. of toluene which is maintained at laboratory temperature. The mixture is diluted with petroleum ether and filtered and the solid residue is crystallised from a mixture of ethyl acetate and petroleum ether. There is thus obtained 1 - *p* - benzyloxyphenoxy - 3 - [*N* - benzyl - *N* - β - (3 - *p* - tolylureido)ethyl]amino - 2 - propanol.

A solution of 6.55 g. of the above product in acetic acid is shaken with hydrogen in the presence of a 30% w/w palladium-on-charcoal catalyst at laboratory temperature and atmospheric pressure until 500 ml. of hydrogen have been absorbed and the uptake of hydrogen ceases. The mixture is filtered, the filtrate is evaporated to dryness and the residue is converted into an oxalate salt which is crystallised from water. There is thus obtained 1 - *p* - hydroxyphenoxy - 3 - β - (3 - *p* - tolylureido)ethylamino - 2 - propanol hemioxalate, m.p. 173—174°C. (which contains one quarter of a mole of water of crystallisation).


The 1 - *p* - benzyloxyphenoxy - 3 - (*N* - benzyl - *N* - β - aminoethyl)amino - 2 - propanol used as starting material may be obtained as follows:—

A mixture of 51.2 g. of 1 - *p* - benzyloxyphenoxy - 2,3 - epoxypropane, 51.3 g. of *N* - benzyl - *N* - β - isobutyramidoethylamino hydrochloride, 200 ml. of aqueous *N* - sodium hydroxide solution and 600 ml. of isopropanol is heated under reflux for 17 hours and then evaporated to dryness under reduced pressure. The residue is shaken with a mixture of chloroform and water and the chloroform layer is dried and evaporated to dryness.

The residue is added to a solution of 200 g. of potassium hydroxide in 400 ml. of ethanol and the mixture is heated under reflux for 96 hours and then diluted with water and extracted with ether. The ethereal extract is washed with water, dried and evaporated to dryness. There is thus obtained as oily residue 1 - *p* - benzyloxyphenoxy - 3 - (*N* - benzyl - *N* - β - aminoethyl)amino - 2 - propanol which is used without further purification.

Example 15.

The process described in Example 14 is repeated except that the appropriate isocyanate is used in place of *p*-tolyl isocyanate. There are thus obtained the compounds described in the following table:—



R ¹	base or salt	m.p. (°C.)	Crystallisation solvent
isopropyl	hemioxalate hemihydrate	192—193 (with decomposition)	methanol/water/ether
n-butyl	hydrochloride dihydrate	151—152	ethanol/ether
n-octyl	hydrochloride hemihydrate	117	ethanol/ether
cyclohexyl	hydrochloride tetrahydrate	145—150	ethanol/ether
<i>p</i> -methoxyphenyl	hemioxalate hemihydrate	180—181	water

Example 16.

A mixture of 5.12 g. of 1 - *m* - benzyloxyphenoxy - 2,3 - epoxypropane, 5.38 g. of *N* - benzyl - *N* - β - (3 - phenylureido)ethylamine and 50 ml. of isopropanol is heated under reflux for 17 hours, cooled and evaporated to dryness under reduced pressure. The residue is dissolved in acetic acid and shaken for 19 hours in the presence of a 30% w/w palladium-on-charcoal catalyst and in an atmosphere of hydrogen at laboratory temperature and atmospheric pressure, when 830 ml. of hydrogen have been absorbed. The mixture is filtered, the filtrate is evaporated to dryness and the residue is dissolved in ethanol. A solution of oxalic acid in ether is added, the mixture is filtered and the solid residue is crystallised from water. There is thus obtained 1 - *m* - hydroxyphenoxy - 3 - β - (3 - phenylureido)ethylamino - 2 - propanol hemioxalate hemihydrate, m.p. 145—146°C.


The process described above is repeated except that 1 - (3,4 - dibenzyloxyphenoxy) - 2,3 - epoxypropane is used in place of 1 - *m* - benzyloxyphenoxy - 2,3 - epoxypropane. There is thus obtained 1 - (3,4 - dihydroxyphenoxy) - 3 - β - (3 - phenylureido)ethylamino - 2 - propanol hemioxalate, m.p. 122°C. with decomposition (crystallised from isopropanol).

The *N* - benzyl - *N* - β - (3 - phenylureido)ethylamine used as starting material may be obtained as follows:—

Benzaldehyde (13 g.) is added dropwise to a stirred solution of *N* - β - (3 - phenylureido)ethylamine (20 g.) in ethanol (100 ml.) which is maintained at 5°C. The mixture is then stirred for 1 hour at laboratory temperature, ethanol (200 ml.) is added and the mixture is again stirred and cooled to 5°C. Sodium borohydride (5 g.) is added portionwise at this temperature, and the mixture is then stirred for 2 hours at laboratory temperature. Aqueous acetic acid is then added to destroy the excess of sodium borohydride, the mixture is evaporated to dryness under reduced pressure and the residue is suspended in aqueous sodium hydroxide solution. The suspension is extracted with ethyl acetate, and the extract is then shaken with aqueous 2*N*-hydrochloric acid. The mixture is filtered and the solid residue is washed with water and with ethyl acetate and dried. There is thus obtained *N* - benzyl - *N* - β - (3-phenylureido)ethylamine hydrochloride, m.p. 165—166°C.

Example 17.


The process described in Example 1 is repeated except that the appropriate 2,3-epoxy-1-phenoxypropane and the appropriate β -carboxamidoethylamine are used as starting materials. There are thus obtained the compounds described in the following table:—



R¹	R²	m.p. (°C.)	Crystallisation solvent
t-butyl	2-cyano	152—154	acetonitrile
t-butyl	2-nitro	165—166	acetonitrile
isopropyl	2-allyl	112—113	ethyl acetate/cyclohexane
isopropyl	2-allyloxy	119—120	ethyl acetate

Example 18.


The process described in Example 2 is repeated except that the appropriate 2,3-epoxy-1-phenoxypropane and the appropriate β -sulphonamidoethylamine are used as starting materials. There are thus obtained the compounds described in the following table:—



R ¹	m.p. (°C.)	Crystallisation solvent
1-naphthyl	oxalate 199–201	water
2-tolyl	153–155	ethanol

Example 19.


5 The process described in Example 5 is repeated except that the appropriate 2,3-epoxy-1-phenoxypropane and the appropriate β -ureidoethylamine are used as starting materials. There are thus obtained the compounds described in the following table:—




R ¹	m.p. (°C.)	Crystallisation solvent
H	108–110	acetonitrile
cyclohexyl	156–158	acetonitrile

Example 20.

10 The process described in Example 5 is repeated except that the appropriate 2,3-epoxy-1-phenoxypropane and the appropriate β -amidoalkylamine are used as starting materials. There are thus obtained the compounds described in the following tables:—



R ¹	-X-Y-	R ²	m.p. (°C.)	Crystallisation solvent
phenyl	-COCH ₂	3-methyl	hydrogen fumarate 87–90	acetonitrile
phenyl	-COCH ₂	3-bromo	118–119	ethyl acetate
2-chlorophenyl	-COCH ₂ -	2-nitro	98–101	toluene
phenyl	-COCH ₂	3-methoxy	hydrogen fumarate 85–90	ethyl acetate
2-nitrophenyl	-SO ₂ -	H	(oil)	—
n-butyl	-CONH-	H	(oil)	—



R ¹	-X-Y-	A	m.p. (°C.)	Crystallisation solvent
isopropyl	-CO-	-C(CH ₃) ₂ CH ₂ -	(oil)	-
phenyl	-COCH ₂ -	-C(CH ₃) ₂ CH ₂ -	(oil)	-
phenyl	-SO ₂ -	-(CH ₂) ₁₂ -	86-87	ethyl acetate

The compounds isolated only as oils are purified by thick-layer chromatography, and their structures are confirmed by proton magnetic resonance spectroscopy.

The various β -amidoalkylamine derivatives used as starting materials may be obtained by a similar process to that described in the second part of Example 10. Those that are characterised may be obtained as follows:—

from 2-nitrobenzenesulphonyl chloride and 1,2-diaminopropane, 1-methyl-2-(2-nitrobenzenesulphonamido)ethylamine, m.p. 178—179°C. after crystallisation from ethanol; from ethylisobutyrate and 1,2-diamino-2-methylpropane, 1,1-dimethyl-2-isobutyramidoethylamine (hydrochloride m.p. 269—270°C. after crystallisation from ethanol); from ethyl phenylacetate and 1,2-diamino-2-methylpropane, 1,1-dimethyl-2-phenylacetamidoethylamine (hydrochloride m.p. 268—270°C. after crystallisation from ethanol);


from benzenesulphonyl chloride and 1,12-diaminododecane, 12-benzenesulphonamidododecylamine, m.p. 77—80°C. after crystallisation from a mixture of ethyl acetate and petroleum ether.

• Example 21.

The process described in Example 12 is repeated except that 1 - phenoxy - 3 - β - (2,5 - dibenzyloxybenzamido)ethylamino - 2 - propanol (oxalate m.p. 176—178°C., prepared as described in Example 1 from β - (2,5 - dibenzyloxybenzamido)ethylamine, m.p. 96—98°C.) or 1 - phenoxy - 3 - β - (3,4 - dibenzyloxybenzamido)ethylamino - 2 - propanol (Example 6) is used as starting material. There are thus obtained respectively 1 - phenoxy - 3 - β - (2,5 - dihydroxybenzamido)ethylamino - 2 - propanol (hemioxalate m.p. 195—197°C. after crystallisation from a mixture of ethanol and water) and 1 - phenoxy - 3 - β - (3,4 - dihydroxybenzamido)ethylamino - 2 - propanol, m.p. 178—180°C. after crystallisation from ethanol.

Example 22.


The process described in Example 13 is repeated except that the appropriate β -substituted ethylamine is used in place of β -(3-phenylureido)ethylamine. There are thus obtained the compounds described in the following table:—



R'	-X-Y-	A	m.p. (°C.)	Crystallisation solvent
isopropyl	-CO-	-CH ₂ CH ₂ -	hemioxalate quarter-hydrate 213	ethanol
isopropyl	-CO-	$\begin{array}{c} \text{CH}_3 \\ \\ -\text{CH}-\text{CH}_2- \end{array}$	hemioxalate quarter-hydrate 220 (with decomposition)	water
ethyl	-CO-O-	-CH ₂ CH ₂ -	hemioxalate 189	aqueous ethanol

Example 23.


The process described in Example 14 is repeated except that the appropriate isocyanate is used in place of *p*-tolyl isocyanate. There are thus obtained the compounds described in the following table:—



R	Base or salt	m.p. (°C.)	Crystallisation solvent
<i>t</i> -butyl	oxalate hemihydrate	133–135 (with decomposition)	ethanol/ether
<i>o</i> -tolyl	acetate	109–111	petroleum ether (b.p. 60–80°C.)
<i>o</i> -methoxyphenyl	oxalate sesquihydrate	68–71 (with decomposition)	ethanol
2,4-dimethylphenyl	acetate quarter hydrate	133–135	isopropanol
1-naphthyl	hemioxalate hydrate	159–162	ethanol
4-biphenyl	hemioxalate hemihydrate	211 (with decomposition)	methanol
methoxymethyl	oxalate	(hygroscopic)	ethanol/ether
<i>n</i> -butyloxycarbonyl- methyl	oxalate	(hygroscopic)	—

Example 24.

The process described in Example 16 is repeated except that the appropriate 1 - benzyloxyphenoxy - 2,3 - epoxyp propane and the appropriate *N* - benzyl - *N* - β - amidoethylamine are used as starting materials. There are thus obtained the compounds described in the following table:—



R ¹	-X-Y-	R ²	R ⁴	m.p. (°C.)	Crystallisation solvent
phenyl	-SO ₂ -	3-methoxy	4-hydroxy	oxalate hemihydrate 127-130	ethanol
phenyl	-CONH-	3-methoxy	4-hydroxy	hemioxalate quarter hydrate 167-168 (with decomposition)	water
isopropyl	-CO-	3-hydroxy	4-hydroxy	hemioxalate 168 (with decomposition)	isopropanol/methanol
isopropyl	-CO-	3-hydroxy	5-hydroxy	hydrochloride (oil)	-
isopropyl	-CO-	4-hydroxy	3-hydroxy-methyl	hemioxalate hemihydrate 148-149	ethanol
phenyl	-SO ₂ -	4-hydroxy	3-hydroxy-methyl	oxalate 117-119	ethanol
phenyl	-SO ₂ -	3-hydroxy	4-hydroxy	oxalate hemihydrate 134 (with decomposition)	ethanol/isopropanol
phenyl	-CONH-	4-hydroxy	3-hydroxy-methyl	hemioxalate hemihydrate 135-137	methanol/ether

The *N* - benzyl - *N* - β - amidoethylamine derivatives used as starting material may be obtained by a similar process to that described in the last part of Example 16, from benzaldehyde and the appropriate β -amidoethylamine. *N* - benzyl - *N* - β - isobutyramidoethylamine hydrochloride has m.p. 197-199°C., and *N* - benzyl - *N* - β - benzenesulphonamidoethylamine hydrochloride has m.p. 173-175°C.

Example 25.

The process described in Example 16 is repeated except that 1 - *p* - benzyloxyphenoxy - 2,3 - epoxyp propane and *N* - benzyl - *N* - [1 - methyl - 2 - (3 - phenylureido)]ethylamine are used as starting materials. There is thus obtained 1 - *p* -

hydroxyphenoxy - 3 - [1 - methyl - 2 - (3 - phenylureido)]ethylamino - 2 - propanol, characterised as the oxalate hydrate m.p. 81°C. (with decomposition) after crystallisation from ethanol.

The *N* - benzyl - *N* - [1 - methyl - 2 - (3 - phenylureido)]ethylamine used as starting material may be obtained as follows:—

A mixture of ethyl phenylcarbamate (50 g.) and 1,2-diaminopropane (105 g.) is heated at 125°C. for 17 hours and cooled, and the excess of amine is removed by evaporation under reduced pressure. The residue is dissolved in ethyl acetate and the solution is washed with saturated brine, dried and evaporated to dryness. The residue is triturated with toluene and there is thus obtained as solid residue 1-methyl-2-(3-phenylureido)ethylamine, which is used without further purification.

Benzaldehyde (6.05 g.) and the above amine (10 g.) are condensed and the product reduced under similar conditions to those described in the last part of Example 16. There is thus obtained *N* - benzyl - *N* - [1 - methyl - 2 - (3 - phenylureido)]-ethylamine, m.p. 118—120°C. after crystallisation from isopropanol.

Example 26.

A mixture of dimethylcarbamoyl chloride (2.15 g.), 1 - *p* - benzyloxyphenoxy-3 - (*N* - benzyl - *N* - β - aminoethyl)amino - 2 - propanol (8.12 g.), anhydrous potassium carbonate (2.76 g.) and toluene (60 ml.) is stirred at laboratory temperature for 2 hours and then diluted with ethyl acetate. The solution is washed with water, dried and evaporated to dryness and the residue is dissolved in acetic acid. The solution is shaken with hydrogen by a similar process to that described in the second part of Example 14, and there is thus obtained 1 - *p* - hydroxyphenoxy - 3 - β - (3,3-dimethylureido)ethylamino - 2 - propanol oxalate hemihydrate, m.p. 181°C. after crystallisation from methanol.

The process described above is repeated except that dimethylsulphamoyl chloride is used in place of dimethylcarbamoyl chloride, and that the dibenzyl intermediate is purified by chromatography, on silica gel using a 1:1 v/v mixture of ethyl acetate and chloroform as eluant, before removal of the benzyl groups by hydrogenolysis. There is thus obtained 1 - *p* - hydroxyphenoxy - 3 - β - (3,3 - dimethylsulphamido)ethylamino - 2 - propanol hemioxalate hemihydrate, m.p. 171—172°C. (with decomposition) after crystallisation from ethanol.

Example 27.

A solution of 2.1 g. of trifluoroacetic anhydride in 5 ml. of toluene is added during 15 minutes to an ice-cooled solution of 3.0 g. of 1 - phenoxy - 3 - (*N* - benzyl - *N* - β - aminoethyl)amino - 2-propanol and 1.01 g. of triethylamine in 30 ml. of toluene and the mixture is stirred for a further 30 minutes and then washed four times with 20 ml. of water each time. The toluene phase is separated, dried over anhydrous magnesium sulphate and evaporated to dryness under reduced pressure. The residue is dissolved in 50 ml. of ethanol and the solution is shaken with hydrogen in the presence of 0.5 g. of a 30% w/w palladium-on-charcoal catalyst at laboratory temperature and atmospheric pressure until 230 ml. of hydrogen have been absorbed. The mixture is filtered, the filtrate is evaporated to dryness and the residue is crystallised from a mixture of ethyl acetate and petroleum ether (b.p. 60—80°C.). There is thus obtained 1-phenoxy-3- β -trifluoroacetamidoethylamino-2-propanol, m.p. 106—108°C.

The 1 - phenoxy - 3 - (*N* - benzyl - *N* - β - aminoethyl)amino - 2 - propanol used as starting material may be obtained in a similar manner to that described in the third part of Example 14, except that 1 - phenoxy - 2,3 - epoxypropane is used in place of the 1 - *p* - benzyloxyphenoxy - 2,3 - epoxypropane. There is thus obtained 1 - phenoxy - 3 - (*N* - benzyl - *N* - β - aminoethyl)amino - 2 - propanol, m.p. 209—210°C. after crystallisation from ethanol.

Example 28.

A solution of 3.74 of 1 - phenoxy - 3 - (*N* - benzyl - *N* - β - aminoethyl)amino-2-propanol, 0.6 g. of methyl formate and 1.68 g. of sodium bicarbonate in 40 ml. of *n*-propanol is heated under reflux for 18 hours and then evaporated to dryness under reduced pressure. The residue is shaken with hydrogen by a similar process to that described in Example 27, and there is thus obtained 1 - phenoxy - 3 - β - formamidoethylamino - 2 - propanol, m.p. 107—109°C. after crystallisation from ethyl acetate.

Example 29.

A mixture of 1,2-diaminopropane (44.4 g.) and ethyl acetate (17.6 g.) is heated under reflux for 48 hours and then distilled, the fraction having a boiling point of 112—113°C./0.35 mm. Hg being collected. This material is dissolved in acetonitrile (200 ml.) and ethereal hydrogen chloride solution is added. The mixture is filtered and there is thus obtained as solid residue a mixture of the hydrochlorides of *N*-(2-amino-1-methylethyl)acetamide and *N*-(2-aminopropyl)acetamide.

Benzenesulphonyl chloride (8.8 g.) is added dropwise during 1 hour to a stirred mixture of the above mixture of hydrochlorides (7.5 g.), water (50 ml.) and sodium bicarbonate (8.4 g.) and the mixture is stirred at laboratory temperature for a further 18 hours and then extracted twice with chloroform (50 ml. each time). The combined chloroform extracts are dried and evaporated to dryness under reduced pressure, and the residue is heated under reflux with aqueous 5*N*-hydrochloric acid (50 ml.) for 18 hours. The mixture is cooled, washed with chloroform (50 ml.) and then evaporated to dryness under reduced pressure. There is thus obtained as solid residue a mixture of the hydrochlorides of *N*-(2-aminopropyl)benzenesulphonamide and *N*-(2-amino-1-methylethyl)benzenesulphonamide.

A mixture of the above mixture of hydrochlorides (2.5 g.), *n*-propanol (40 ml.), sodium bicarbonate (0.84 g.) and 1-phenoxy-2,3-epoxypropane (1.5 g.) is heated under reflux for 18 hours, cooled and filtered and the filtrate is evaporated to dryness under reduced pressure. The residue is extracted three times with chloroform (30 ml. each time) and the combined extracts are dried and evaporated to dryness. The residue is chromatographed on silica gel plates (20 cm. × 20 cm. × 0.5 mm., Merck Kieselgel GF 60F254) using a 4:1 v/v mixture of chloroform and methanol as developing solvent, and the band having an *R_F* value of 0.5 is removed from each plate and extracted with methanol. The methanol extract is evaporated to dryness and the residue is reappplied to similar plates. The plates are developed with a 9:1 v/v mixture of chloroform and methanol, and are then re-developed 5 times with this solvent. Two distinct bands are obtained, having *R_F* values of 0.33 and 0.37, and these bands are separately removed from the plates and extracted with methanol. The methanol extract from the band having *R_F* 0.33 is evaporated to dryness and the residue is dissolved in ethyl acetate (10 ml.) and added to a solution of oxalic acid (0.4 g.) in ethyl acetate (10 ml.). The mixture is filtered and the residue is crystallised from acetonitrile. There is thus obtained 1-phenoxy-3-(2-methyl-2-benzenesulphonamidoethyl)-amino-2-propanol oxalate, m.p. 122—123°C.

The methanol extract from the band having *R_F* 0.37 is evaporated to dryness and there is thus obtained as oily residue 1-phenoxy-3-(1-methyl-2-benzenesulphonamidoethyl)amino-2-propanol, from which no crystalline salt has been obtained.

Example 30.

The process described in Example 4 is repeated except that 2,3-epoxy-1-(naphth-1-yloxy)propane and β-2-chlorophenylacetamidoethylamine hydrochloride are used as starting materials. There is thus obtained 1-(naphth-1-yloxy)-3-β-(2-chlorophenyl)acetamidoethylamino-2-propanol, characterised as the hydrochloride salt, m.p. 176—177°C.

Example 31.

A solution of 2.8 g. of racemic 1-phenoxy-3-β-isobutyramidoethylamino-2-propanol in 25 ml. of ethanol is added to a solution of 2.9 g. of (–)-2,3,4,6-di-*O*-isopropylidene-2-keto-L-gulonic acid hydrate in 25 ml. of ethanol and the mixture is kept at 4°C. for 18 hours and then filtered. The solid residue is crystallised four times from 10 ml. of ethanol each time, and there is thus obtained (–)-1-phenoxy-3-β-isobutyramidoethylamino-2-propanol (–)-2,3,4,6-di-*O*-isopropylidene-2-keto-L-gulonate, m.p. 170—172°C.

The above salt is stirred with aqueous 2*N*-sodium hydroxide solution (3 ml.) and the mixture is extracted twice with ethyl acetate (5 ml. each time). The combined extracts are dried through a molecular sieve and are added to a solution of 0.6 g. of oxalic acid in 10 ml. of ethyl acetate. The mixture is filtered and the solid residue is washed with ether. There is thus obtained (–)-1-phenoxy-3-β-isobutyramidoethylamino-2-propanol oxalate, m.p. 148—150°C. (with decomposition), α_D^{25} –12.0° (c, 7.6% in water).

Example 32.

A mixture of 1.86 g. of 1 - chloro - 3 - phenoxy - 2 - propanol, 1.3 g. of β -isobutyramidoethylamine, 40 ml. of n-propanol and 0.84 g. of sodium bicarbonate is heated under reflux for 18 hours, cooled and filtered and the filtrate is evaporated to dryness under reduced pressure. The residue is stirred with 20 ml. of water and the mixture is extracted twice with 20 ml. of chloroform each time. The combined chloroform extracts are dried over anhydrous magnesium sulphate and evaporated to dryness. The residue is crystallised from ethyl acetate and there is thus obtained 1-phenoxy - 3 - β - isobutyramidoethylamino - 2 - propanol, m.p. 125—126°C.

Example 33.


A mixture of 3.12 g. of 1 - chloro - 3 - β - isobutyramidoethylamino - 2 - propanol oxalate, 0.94 g. of phenol, 1.6 g. of sodium hydroxide, 5 ml. of water and 30 ml. of isopropanol is heated under reflux for 18 hours, cooled and filtered and the filtrate is evaporated to dryness under reduced pressure. The residue is stirred with 20 ml. of water and the mixture is extracted twice with 20 ml. of chloroform each time. The combined chloroform extracts are dried over anhydrous magnesium sulphate and evaporated to dryness. The residue is crystallised from ethyl acetate and there is thus obtained 1 - phenoxy - 3 - β - isobutyramidoethylamino - 2 - propanol, m.p. 125—126°C.

The 1 - chloro - 3 - β - isobutyramidoethylamino - 2 - propanol used as starting material may be obtained as follows:—


A mixture of 6.5 g. of β -isobutyramidoethylamine, 3.9 ml. of epichlorohydrin and 60 ml. of isopropanol is stirred at laboratory temperature for 18 hours and is then added to a solution of 6.3 g. of oxalic acid in 100 ml. of ethyl acetate. The mixture is stirred for 30 minutes and filtered and the solid residue is crystallised from 100 ml. of ethanol. There is thus obtained 1 - chloro - 3 - β - isobutyramidoethylamino - 2 - propanol oxalate, m.p. 129—130°C.

WHAT WE CLAIM IS:—

1. An alkanolamine derivative of the formula:—



wherein A stands for an alkylene radical of from 2 to 12 carbon atoms; wherein R¹ stands for the hydrogen atom or for an alkyl, halogenoalkyl, alkenyl or cycloalkyl radical each of up to 10 carbon atoms, or for an aryl radical of the formula:—



wherein R², R³, R¹² and R¹³, which may be the same or different, each stands for a hydrogen or halogen atom, a hydroxy, amino, nitro or cyano radical, an alkyl, cycloalkyl, alkenyl, alkynyl, alkoxy, alkylthio, cycloalkoxy, alkenyloxy, alkynyloxy or alkanoyl radical each of up to 6 carbon atoms, or an aryl, aryloxy or aralkoxy radical each of up to 12 carbon atoms; or wherein R² and R³ together, and/or R¹² and R¹³ together, form the trimethylene, tetramethylene, 1-oxotetramethylene, propenylene, but-2-enylene or buta-1,3-dienylene radical such that together with the adjacent benzene ring they form respectively the indanyl, 5,6,7,8-tetrahydronaphthyl, 5-oxo-5,6,7,8-tetrahydronaphthyl, indenyl, 5,8-dihydronaphthyl or naphthyl radical; wherein R⁴ stands for the hydrogen atom or for the hydroxy or hydroxymethyl radical or for an aralkoxy radical of up to 12 carbon atoms; wherein R¹⁴ stands for the hydrogen atom or for the amino radical or for a dialkylamino radical of up to 12 carbon atoms; wherein X stands for the carbonyl (—CO—) or sulphonyl (—SO₂—) radical and wherein Y stands for a direct link, or for an alkylene, oxyalkylene or alkyleneoxy radical each of up to 6 carbon atoms, or for the imino (—NH—) radical, or for an



Example 32.

A mixture of 1.86 g. of 1 - chloro - 3 - phenoxy - 2 - propanol, 1.3 g. of β -isobutyramidoethylamine, 40 ml. of n-propanol and 0.84 g. of sodium bicarbonate is heated under reflux for 18 hours, cooled and filtered and the filtrate is evaporated to dryness under reduced pressure. The residue is stirred with 20 ml. of water and the mixture is extracted twice with 20 ml. of chloroform each time. The combined chloroform extracts are dried over anhydrous magnesium sulphate and evaporated to dryness. The residue is crystallised from ethyl acetate and there is thus obtained 1-phenoxy - 3 - β - isobutyramidoethylamino - 2 - propanol, m.p. 125—126°C.

Example 33.


A mixture of 3.12 g. of 1 - chloro - 3 - β - isobutyramidoethylamino - 2 - propanol oxalate, 0.94 g. of phenol, 1.6 g. of sodium hydroxide, 5 ml. of water and 30 ml. of isopropanol is heated under reflux for 18 hours, cooled and filtered and the filtrate is evaporated to dryness under reduced pressure. The residue is stirred with 20 ml. of water and the mixture is extracted twice with 20 ml. of chloroform each time. The combined chloroform extracts are dried over anhydrous magnesium sulphate and evaporated to dryness. The residue is crystallised from ethyl acetate and there is thus obtained 1 - phenoxy - 3 - β - isobutyramidoethylamino - 2 - propanol, m.p. 125—126°C.

The 1 - chloro - 3 - β - isobutyramidoethylamino - 2 - propanol used as starting material may be obtained as follows:—


A mixture of 6.5 g. of β -isobutyramidoethylamine, 3.9 ml. of epichlorohydrin and 60 ml. of isopropanol is stirred at laboratory temperature for 18 hours and is then added to a solution of 6.3 g. of oxalic acid in 100 ml. of ethyl acetate. The mixture is stirred for 30 minutes and filtered and the solid residue is crystallised from 100 ml. of ethanol. There is thus obtained 1 - chloro - 3 - β - isobutyramidoethylamino - 2 - propanol oxalate, m.p. 129—130°C.

WHAT WE CLAIM IS:—

1. An alkanolamine derivative of the formula:—



wherein A stands for an alkylene radical of from 2 to 12 carbon atoms; wherein R¹ stands for the hydrogen atom or for an alkyl, halogenoalkyl, alkenyl or cycloalkyl radical each of up to 10 carbon atoms, or for an aryl radical of the formula:—



wherein R², R³, R¹² and R¹³, which may be the same or different, each stands for a hydrogen or halogen atom, a hydroxy, amino, nitro or cyano radical, an alkyl, cycloalkyl, alkenyl, alkynyl, alkoxy, alkylthio, cycloalkoxy, alkenyloxy, alkynyloxy or alkanoyl radical each of up to 6 carbon atoms, or an aryl, aryloxy or aralkoxy radical each of up to 12 carbon atoms; or wherein R² and R³ together, and/or R¹² and R¹³ together, form the trimethylene, tetramethylene, 1-oxotetramethylene, propenylene, but-2-enylene or buta-1,3-dienylene radical such that together with the adjacent benzene ring they form respectively the indanyl, 5,6,7,8-tetrahydronaphthyl, 5-oxo-5,6,7,8-tetrahydronaphthyl, indenyl, 5,8-dihydronaphthyl or naphthyl radical; wherein R⁴ stands for the hydrogen atom or for the hydroxy or hydroxymethyl radical or for an aralkoxy radical of up to 12 carbon atoms; wherein R¹⁴ stands for the hydrogen atom or for the amino radical or for a dialkylamino radical of up to 12 carbon atoms; wherein X stands for the carbonyl (—CO—) or sulphonyl (—SO₂—) radical and wherein Y stands for a direct link, or for an alkylene, oxyalkylene or alkyleneoxy radical each of up to 6 carbon atoms, or for the imino (—NH—) radical, or for an




1,455,116

45
33


alkylimino, iminoalkylene, iminoalkyleneoxy or iminoalkylenecarbonyloxy radical each of up to 6 carbon atoms, or (except where R¹ stands for the hydrogen atom) for the oxygen atom; or an acid-addition salt thereof.

2. An alkanolamine derivative as claimed in claim 1 wherein A stands for the ethylene, trimethylene, tetramethylene, hexamethylene, dodecamethylene, 1-methylethylene, 2-methylethylene or 1,1-dimethylethylene radical, wherein R¹ stands for the hydrogen atom, or for the methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, t-butyl, n-pentyl, n-octyl, trifluoromethyl, allyl, cyclopropyl, cyclopentyl or cyclohexyl radical, or for a radical of the formula:—




wherein R¹, R², R¹² and R¹³, which may be the same or different, each stands for a hydrogen, fluorine, chlorine, bromine or iodine atom or for the hydroxy, amino, nitro, cyano, methyl, ethyl, n-propyl, cyclopropyl, cyclopentyl, allyl, ethynyl, methoxy, isopropoxy, methylthio, cyclopentyl, allyloxy propargyloxy, formyl, acetyl, phenyl, phenoxy or benzyloxy radical, or wherein R² and R³ together and/or R¹² and R¹³ together form the trimethylene, tetramethylene, 1-oxotetramethylene, propenylene, but-2-enylene or buta-1,3-dienylene radical; wherein R⁴ stands for the hydrogen atom or for the hydroxy, hydroxymethyl or benzyloxy radical; wherein R¹⁴ stands for the hydrogen atom or for the amino or dimethylamino radical; wherein X stands for the carbonyl or sulphonyl radical and wherein Y stands for a direct link, or for the methylene, ethylene, oxymethylene, methyleneoxy, ethyleneoxy, trimethyleneoxy, 1-methylethylideneoxy, 1-methylpropylideneoxy, imino, methylimino, iminomethylene, iminomethyleneoxy or iminomethylenecarbonyloxy radical, or (except when R¹ stands for the hydrogen atom) for the oxygen atom; or an acid-addition salt thereof.

3. An alkanolamine derivative of the formula:—



wherein A stands for an alkylene radical of from 2 to 10 carbon atoms, wherein R¹ stands for the hydrogen atom or for an alkyl, alkenyl or cycloalkyl radical each of up to 10 carbon atoms, or for an aryl radical of the formula:—



wherein R², R³, R¹² and R¹³, which may be the same or different, each stands for a hydrogen or halogen atom, a hydroxy, amino, nitro or cyano radical, an alkyl, alkenyl, alkoxy or alkenyloxy radical each of up to 6 carbon atoms, or an aralkoxy radical of up to 12 carbon atoms; wherein R⁴ stands for the hydrogen atom or for the hydroxy radical or for an aralkoxy radical of up to 12 carbon atoms; wherein R¹⁴ stands for the hydrogen atom or for the amino radical or for a dialkylamino radical of up to 12 carbon atoms; wherein X stands for the carbonyl (—CO—) or sulphonyl (—SO₂—) radical and wherein Y stands for a direct link, or for an alkylene or alkyleneoxy radical each of up to 6 carbon atoms, or for the imino (—NH—) radical or (except when R¹ stands for hydrogen) for the oxygen atom, or an acid-addition salt thereof.


4. An alkanolamine derivative of the formula:—




34

34

1,455,116



wherein A stands for an alkylene radical of from 2 to 6 carbon atoms, wherein R¹ stands for an alkyl, alkenyl or cycloalkyl radical each of up to 6 carbon atoms, or for an aryl radical of the formula:—



wherein R², R³, R¹² and R¹³, which may be the same or different, each stands for a hydrogen or halogen atom, a hydroxy, nitro or cyano radical, an alkyl, alkenyl, alkoxy or alkenyloxy radical each of up to 6 carbon atoms, or an aralkoxy radical of up to 12 carbon atoms; wherein R⁴ stands for the hydrogen atom or for the hydroxy radical or for an aralkoxy radical of up to 12 carbon atoms; wherein X stands for the carbonyl (—CO—) or sulphonyl (—SO₂—) radical and wherein Y stands for a direct link, or for an alkylene or alkyleneoxy radical each of up to 6 carbon atoms, or for the imino (—NH—) radical; or an acid-addition salt thereof.

5. An alkanolamine derivative as claimed in claim 1 wherein A stands for the ethylene, 1-methylethylene or 1,1-dimethylethylene radical, R² stands for the hydrogen atom or for an alkyl, alkenyl or cycloalkyl radical each of up to 6 carbon atoms, or for the trifluoromethyl radical or for a monohydroxyphenyl or dihydroxyphenyl radical, Y stands for a direct link or for an imino radical, X stands for the carbonyl radical, R³ stands for the hydrogen atom or for a chloro, cyano, nitro, methyl, allyl, methoxy or allyloxy substituent which is in the *ortho* position of the benzene ring and R³ and R⁴ both stand for hydrogen atoms; or an acid-addition salt thereof.

6. An alkanolamine derivative as claimed in claim 1 wherein A stands for the ethylene, 1-methylethylene or 1,1-dimethylethylene radical, R¹ stands for a phenyl radical, wherein R¹² stands for the hydrogen atom or for a chloro, nitro, methyl or methoxy substituent and R¹³ and R¹⁴ both stand for hydrogen atoms, Y stands for the methylene, ethylene, methyleneoxy or imino radical, X stands for the carbonyl radical, R² stands for the hydrogen atom or for a chloro, cyano, nitro, methyl, allyl, methoxy or allyloxy substituent which is in the *ortho* or *meta*- position of the benzene ring and R³ and R⁴ both stand for hydrogen atoms; or an acid-addition salt thereof.

7. An alkanolamine derivative as claimed in claim 1 wherein A stands for the ethylene, 1-methylethylene or 1,1-dimethylethylene radical, R¹ stands for the hydrogen atom or for an alkyl radical of up to 6 carbon atoms, Y stands for the methyleneoxy radical, X stands for the carbonyl radical and R², R³ and R⁴ have the meanings stated in claim 5; or an acid-addition salt thereof.

8. An alkanolamine derivative as claimed in claim 1 wherein A stands for the ethylene, 1-methylethylene or 1,1-dimethylethylene radical, R¹ stands for an alkyl radical of up to 6 carbon atoms or for a phenyl radical wherein R¹² stands for the hydrogen atom or for a chloro, amino, nitro or methyl substituent and R¹³ and R¹⁴ both stand for hydrogen atoms, Y stands for a direct link, X stands for the sulphonyl radical and R², R³ and R⁴ have the meanings stated in claim 5; or an acid-addition salt thereof.

9. An alkanolamine derivative as claimed in claim 1 wherein A stands for the ethylene, 1-methylethylene or 1,1-dimethylethylene radical, R¹ stands for an alkyl or cycloalkyl radical each of up to 6 carbon atoms, or for a phenyl radical wherein R¹² stands for the hydrogen atom or for a methyl, methoxy or phenyl substituent and R¹³ and R¹⁴ both stand for hydrogen atoms, Y stands for a direct link or for the methylene or imino radical, X stands for the carbonyl or sulphonyl radical, R² stands for the hydrogen atom or for the hydroxy radical, R³ stands for the hydrogen atom and R⁴ stands for the hydroxy radical, or an acid-addition salt thereof.



* C L M S *

1,455,116

35

10. An alkanolamine derivative as claimed in claim 1 wherein A stands for the ethylene, 1-methylethylene or 1,1-dimethylethylene radical, R¹ stands for an alkyl radical of up to 6 carbon atoms, Y stands for the oxygen atom or for the methyleneoxy, methylimino, iminomethyleneoxy or iminomethylenecarbonyloxy radical, X stands for the carbonyl or sulphonyl radical and R², R³ and R⁴ have the meanings stated in claim 9; or an acid-addition salt thereof.

11. An alkanolamine derivative as claimed in claim 3 wherein A stands for the ethylene or 1-methylethylene radical, R¹ stands for an alkyl, alkenyl or cycloalkyl radical each of up to 6 carbon atoms, Y stands for a direct link or for an imino radical, X stands for the carbonyl radical, R² stands for hydrogen or for a chloro, cyano, nitro, methyl, methoxy or allyloxy substituent which is in the *ortho* position of the benzene ring and R³ and R⁴ both stand for hydrogen; or an acid-addition salt thereof.

12. An alkanolamine derivative as claimed in claim 3 wherein A stands for the ethylene or 1-methylethylene radical, R¹ stands for a phenyl radical wherein R¹² stands for the hydrogen atom or for a chloro, nitro or methoxy substituent and R¹³ and R¹⁴ both stand for hydrogen atoms, Y stands for the methylene, ethylene, methyleneoxy or imino radical, X stands for the carbonyl radical and R², R³ and R⁴ have the meanings stated in claim 11; or an acid-addition salt thereof.

13. An alkanolamine derivative as claimed in claim 3 wherein A stands for the ethylene or 1-methylethylene radical, R¹ stands for the hydrogen atom or for an alkyl radical of up to 6 carbon atoms, Y stands for the methyleneoxy radical, X stands for the carbonyl radical and R², R³ and R⁴ have the meanings stated in claim 11; or an acid-addition salt thereof.

14. An alkanolamine derivative as claimed in claim 3 wherein A stands for the ethylene or 1-methylethylene radical, R¹ stands for an alkyl radical of up to 6 carbon atoms or for a phenyl radical wherein R¹² stands for the hydrogen atom or for a chloro, nitro, or methyl substituent and R¹³ and R¹⁴ both stand for hydrogen atoms, Y stands for a direct link, X stands for the sulphonyl radical and R², R³ and R⁴ all stand for hydrogen; or an acid-addition salt thereof.

15. An alkanolamine derivative as claimed in claim 3 wherein A stands for the ethylene or 1-methylethylene radical, R¹ stands for an alkyl radical of up to 6 carbon atoms, or for a phenyl radical wherein R¹² stands for the hydrogen atom or for a methyl or methoxy substituent and R¹³ and R¹⁴ both stand for hydrogen atoms, either Y stands for a direct link and X stands for the carbonyl or sulphonyl radical or Y stands for the imino radical and X stands for the carbonyl radical, R² stands for the hydroxy radical and R³ and R⁴ both stand for hydrogen atoms; or an acid-addition salt thereof.

16. An alkanolamine derivative as claimed in claim 4 wherein A stands for the ethylene radical, R¹ stands for the phenyl radical, R², R³, R⁴, R¹² and R¹³ all stand for hydrogen atoms, and X and Y have the meanings stated in claim 4.

17. An alkanolamine derivative as claimed in claim 4 wherein A stands for the ethylene radical, R¹ stands for an alkyl, alkenyl or cycloalkyl radical each of up to 6 carbon atoms, R², R³ and R⁴ all stand for hydrogen atoms, X stands for the carbonyl or sulphonyl radical and Y stands for a direct link or for the imino radical; or an acid-addition salt thereof.

18. An alkanolamine derivative as claimed in claim 17 wherein X stands for the carbonyl radical; or an acid-addition salt thereof.

19. The compound:—

1-phenoxy-3- β -benzamidoethylamino-2-propanol;
1-phenoxy-3- β -benzenesulphonamidoethylamino-2-propanol;
1-phenoxy-3- β -phenylacetamidoethylamino-2-propanol;
1-phenoxy-3- β -phenoxyacetamidoethylamino-2-propanol; or
1-phenoxy-3- β -(3-phenylureido)ethylamino-2-propanol;
or an acid-addition salt thereof.

20. The compound:—

1-phenoxy-3-(α -methyl- β -phenylacetamidoethyl)amino-2-propanol;
1-phenoxy-3- β -pivalamidoethylamino-2-propanol;
1-phenoxy-3-(β -isobutyramido- α -methylethyl)amino-2-propanol;
1-(2-nitrophenoxy)-3- β -isobutyramidoethylamino-2-propanol;
1-phenoxy-3- β -propionamidoethylamino-2-propanol;
1-phenoxy-3- β -hydroxyacetamidoethylamino-2-propanol;
1-phenoxy-3- β -(2-nitrobenzenesulphonamidoethyl)amino-2-propanol;
1-(2-cyanophenoxy)-3- β -propionamidoethylamino-2-propanol;



- 1-phenoxy-3- β -isopropylsulphonamidoethylamino-2-propanol;
 1-phenoxy-3- β -(2-chlorophenylacetamidoethyl)amino-2-propanol;
 1-(2-cyanophenoxy)-3- β -(3-phenylureidoethyl)amino-2-propanol;
 1-phenoxy-3- β -butyramidoethylamino-2-propanol;
 1-phenoxy-3- β -(3-n-butylureidoethyl)amino-2-propanol;
 1-phenoxy-3- β -isobutyramidoethylamino-2-propanol;
 1-phenoxy-3- β -(4-methoxyphenylacetamidoethyl)amino-2-propanol;
 1-phenoxy-3- β -acetamidoethylamino-2-propanol or
 1-phenoxy-3- β -(n-propylsulphonamidoethyl)amino-2-propanol;
 or an acid-addition salt thereof.

21. The compound:—

- 1-(2-methoxyphenoxy)-3- β -isobutyramidoethylamino-2-propanol;
 1-phenoxy-3-(α,α -dimethyl- β -isobutyramidoethyl)amino-2-propanol; or
 1-phenoxy-3-(α,α -dimethyl- β -phenylacetamidoethyl)amino-2-propanol;
 or an acid-addition salt thereof.


22. The compound:—

- 1-(4-hydroxyphenoxy)-3- β -(3-phenylureido)ethylamino-2-propanol;
 1-(3,4-dihydroxyphenoxy)-3- β -isobutyramidoethylamino-2-propanol;
 1-(3,4-dihydroxyphenoxy)-3- β -(3-phenylureido)ethylamino-2-propanol;
 1-(3-hydroxyphenoxy)-3- β -(3-phenylureido)ethylamino-2-propanol;
 1-(3-hydroxyphenoxy)-3- β -benzenesulphonamidoethylamino-2-propanol;
 1-(4-hydroxyphenoxy)-3- β -(3,3-dimethylsulphamido)ethylamino-2-propanol;
 1-(4-hydroxyphenoxy)-3- β -(3-methoxymethylureido)ethylamino-2-propanol; or
 1-(4-hydroxyphenoxy)-3- β -(3-n-butyloxycarbonylmethylureido)ethylamino-
 2-propanol;
 or an acid-addition salt thereof.

23. An acid-addition salt as claimed in any of claims 1 to 22 which is a hydrochloride, hydrobromine, phosphate, sulphate, oxalate, lactate, tartrate, acetate, salicylate, citrate, benzoate, β -naphthoate, adipate or 1,1-methylene-bis-(2-hydroxy-3-naphthoate), or a salt derived from a sulphonated polystyrene resin.


24. A process for the manufacture of an alkanolamine derivative or an acid-addition salt thereof, claimed in any of claims 1 to 23, which comprises assembling in sequence, by chemical methods known to be suitable for this purpose, the five radicals:

- (i) an aryloxy radical of the formula:—



wherein R^2 , R^3 and R^4 have the meanings stated in any of claims 1 to 18;

- (ii) an oxygenated three carbon radical of the formula:—



wherein R^5 stands for hydrogen or for a protecting group;

- (iii) an imino radical of the formula $-\text{NR}^6-$, wherein R^6 stands for hydrogen or for a protecting group;

- (iv) a radical of the formula:—





1,455,116

37

wherein A has the meaning stated in any of claims 1 to 18 and wherein R' stands for hydrogen or for a protecting group; and

(v) a radical of the formula:—




5 wherein R¹, X and Y have the meanings stated in any of claims 1 to 18; whereafter if one or more of R², R³ and R' stands for a protecting group, the one or more protecting groups are removed; and whereafter if desired:—

10 (a) an alkanolamine derivative wherein one or more of R², R³, R⁴, R¹² and R¹³ stands for an α-arylalkoxy radical may be converted into the corresponding compound wherein one or more of R², R³, R⁴, R¹² and R¹³ stands for the hydroxy radical by hydrogenolysis;

(b) a racemic alkanolamine derivative may be resolved into its optically-active enantiomorphs; or

15 (c) an alkanolamine derivative in free base form may be converted into an acid-addition salt thereof by reaction with an acid.

25. A process as claimed in claim 24 which comprises the reaction of a compound of the formula:—




wherein R², R³ and R⁴ have the meanings stated in any of claims 1 to 18 with an amine of the formula:—



25 wherein A, R¹, X and Y have the meanings stated in any of claims 1 to 18 and wherein R⁴ stands for hydrogen or for the benzyl radical, whereafter if R⁴ stands for the benzyl radical this radical is removed by hydrogenolysis, and whereafter any of the processes described in paragraphs (a), (b) and (c) of claim 24 may be carried out if desired.

26. A process as claimed in claim 24 which comprises the reaction of a compound of the formula:—



wherein R², R³, R⁴ and A have the meanings stated in any of claims 1 to 18 and R⁵, R⁶ and R⁷ have the meanings stated in claim 24, with a compound of the formula:—





1,455,116

38

wherein R¹, X and Y have the meanings stated in any of claims 1 to 18 and wherein Z¹ stands for a displaceable radical; or, when X stands for the carbonyl radical and Y stands for the imino radical, with an isocyanate of the formula:—



- 5 wherein R¹ has the meaning stated in any of claims 1 to 18; whereafter if one or more
of R¹, R² and R³ stands for a protecting group the one or more protecting groups are
removed, and whereafter any of the processes described in paragraphs (a), (b) and (c)
of claim 24 may be carried out if desired.
- 10 27. A pharmaceutical composition comprising as active ingredient at least one
alkanolamine derivative or an acid-addition salt thereof, claimed in any of claims 1 to
23, in association with a pharmaceutically-acceptable diluent or carrier therefor.
- 15 28. A composition as claimed in claim 27 which is in the form of a tablet, capsule,
aqueous or oily solution or suspension, emulsion, injectable aqueous or oily solution
or suspension, dispersible powder, spray or aerosol formulation.
- 20 29. A composition as claimed in claim 27 or 28 which contains, in addition to
the alkanolamine derivative, one or more drugs selected from sedatives, vasodilators,
diuretics, hypotensive agents, cardiac membrane stabilising agents, agents used in the
treatment of Parkinson's disease and other tremors, cardiotonic agents, α -adrenergic
blocking agents and sympathomimetic bronchodilators.
- 25 30. An alkanolamine derivative as claimed in any of claims 4 to 16 and 19 as
hereinbefore particularly described in any of Examples 1 to 5.
31. An alkanolamine derivative as claimed in any of claims 3, 4 and 11 to 20
as hereinbefore particularly described in any of Examples 6 to 16.
32. An alkanolamine derivative as claimed in any of claims 1 to 23 as hereinbefore
particularly described in any of Examples 17 to 33.
33. A process for the manufacture of alkanolamine derivatives, claimed in any of
claims 24 to 26, as hereinbefore particularly described in any of Examples 1 to 33.

R. P. SLATCHER,
Agent for the Applicants.

Printed for Her Majesty's Stationery Office by the Courier Press, Leamington Spa, 1976.
Published by the Patent Office, 25 Southampton Buildings, London, WC2A 1AY, from
which copies may be obtained.