



2. The process according to claim 1, for the preparation of a compound of formula (Ia)



5

3. The process according to claim 1 or 2, wherein the compound of the formula (I) or (Ia) is 1-((R)2,2-dimethyl-[1,3]dioxolan-4-ylmethyl)-1H-pyrazol-3-ylamine.




4. The process according to claim 1, wherein step a) is performed with a
10 carboxylating agent at a reaction temperature between 20 and 100°C.


5. The process according to claim 4, wherein ~~the reaction~~step a) is performed in a solvent selected from tetrahydrofuran, acetic acid, water, iso-propyl acetate or ethyl acetate.

15

6. The process according to claim 4, wherein the carboxylating agent is acetic anhydride, acetyl chloride, benzoic anhydride, benzoyl chloride or pivaloyl chloride.




2. The process according to claim 1, for the preparation of a compound of formula (Ia)



5

3. The process according to claim 1 or 2, wherein the compound of the formula (I) or (Ia) is 1-((R)2,2-dimethyl-[1,3]dioxolan-4-ylmethyl)-1H-pyrazol-3-ylamine.



4. The process according to claim 1, wherein step a) is performed with a
10 carboxylating agent at a reaction temperature between 20 and 100°C.

5. The process according to claim 4, wherein step a) is performed in a solvent selected from tetrahydrofuran, acetic acid, water, iso-propyl acetate or ethyl acetate.

15 6. The process according to claim 4, wherein the carboxylating agent is acetic anhydride, acetyl chloride, benzoic anhydride, benzoyl chloride or pivaloyl chloride.