

ULTRA MILD SKIN CLEANSING COMPOSITION

Leonard E. Small

Paul H. Garrison, Jr.

William M. Winkler

Sharon A. Seaman

Alyce J. Papa

5

10 This invention relates to mild skin cleansers. More particularly, this invention relates to skin cleansers comprising synthetic surfactants, moisturizers, polymeric skin feel and mildness aids and soap.

15 The cleansing of skin with surface-active cleansing preparations has become a focus of great interest. Many people wash and scrub their skin with various surface-active preparations several times a day. Ideal skin cleansers should cleanse the skin gently, causing little or no irritation, without defatting and over-drying the skin or leaving it taut after frequent routine use.

20 Most lathering soaps, liquids and bars included, fail in this respect.

 Synthetic detergent bars, frequently referred to as "syndet bars," are well known and are becoming increasingly popular. However, widespread replacement of soap bars by syndet bars

25 has not so far been possible for a variety of reasons, primarily the poor physical characteristics of syndet bars as compared to soap bars, e.g., smear or bar messiness and lather quality.

 Certain synthetic surfactants are particularly mild. However, a major drawback of most mild synthetic surfactant systems

30 when formulated for skin cleansing is poor lather performance, when compared to the highest bar soap standards (bars which are rich in coconut soap and superfatted). On the other side, the use of known high sudsing anionic surfactants with lather boosters can yield acceptable lather volume. Unfortunately, however,

35 the highest sudsing anionic surfactants are, in fact, poor in clinical skin mildness. Surfactants that are among the mildest, such as sodium lauryl glyceryl ether sulfonate, (AGS), are

marginal in lather. It will be appreciated that these two factors make the surfactant selection, the lather and the skin feel benefit formulation process, a delicate balancing act.

5 Similarly, optimization of bar firmness also requires a delicate balancing act with respect to maintaining acceptable lather and mildness, although as a single variable, bar firmness is accomplished through a fairly straightforward process with adjustments to levels of fillers, binders, etc., such as free fatty acids.

10 It is known that moisturizers provide skin conditioning benefits. For example, it is known that glycerin and/or free fatty acids are added to bars or liquid cleansing products for skin benefits.

15 Likewise, polymeric skin feel aids are known to those knowledgeable in the art for providing unique tactile characteristics to both the lather and the skin during rinsing.

For background, examples of liquid cleansing compositions are disclosed in the following references. U.S. Pat. No. 4,338,211, Stiros, issued July 6, 1982, discloses liquid skin cleanser with 2.3% to 3% AGS, the polymer JR-400 and small amounts of free fatty acid plus a fatty acid alkylolamide as lather boosting agents. Compositions containing the surfactants AGS and sarcosinate are not disclosed. Also, U.S. Pat. No. 4,491,539, James J. Hoskins and Adriaan Kessler, issued Jan. 1, 1985, discloses liquid cleansing products comprising about 5% to 25 30% of surfactant, about 0.1% to about 1.0% of guar material, about 0.15% to about 1.0% of nonionic carboxyvinyl polymer, and water. Exemplary compositions containing mild surfactants in general and, specifically, the surfactants AGS and sarcosinate are not disclosed. Another background reference is British Pat. No. 30 2,103,236A, Colgate, Feb. 16, 1984, which discloses a liquid detergent containing guar gum, a ternary surfactant mixture including betaine. AGS is not used.

35 U.S. Pat. No. 2,894,912, Geitz, issued July 14, 1959, for "Isethionate Detergent Bar," discloses a detergent bar consisting essentially of from 30-70% of water-soluble alkali metal detergent

salts of esters of isethionic acid with mixed aliphatic fatty acids having from 6 to 18 carbon atoms and an iodine value of less than 20, of which mixed acids at least 75% have from 12 to 18 carbon atoms and up to 25% have from 6 to 10 carbon atoms, from 2-10%
5 of at least one water-soluble suds-boosting detergent salt selected from the group consisting of alkali metal and organic amine higher aliphatic fatty alcohol sulfates, alkyl aryl sulfonates, and higher aliphatic fatty acid taurides, from about 1% to about 9% water, from about 2.5% to about 25% of water-soluble higher fatty acid
10 soap, and from 10-40% of at least one higher fatty acid having from about 12 to about 25 carbon atoms as a binder and plasticizer, said bar having a pH within the range from 6 to 8, measured as a 10% aqueous solution of the bar composition at 35°C. This patent does not teach the use of polymeric skin feel aids nor
15 the use of other mild synthetics.

U.S. Pat. No. 4,234,464, Morshauser, issued Nov. 18, 1980, for "Detergent Bar Composition and Binder Therefor," discloses a detergent bar in Example 6 which comprises: 45% sodium cocoyl isethionate, 5% alkyl amide, 37.5% stearic acid, 5.0% hydrogenated
20 tallow glycerides, and 1% Polymer JR. Morshauser teaches that his detergent bars can contain up to 5% soap "without substantial detriment." This patent also teaches the use of up to 1.5% cationic polymer.

U.S. Pat. No. 4,012,341, Orshitzer et al., issued Mar. 15,
25 1977, for a "Unique All Synthetic Detergent Shampoo Bar," discloses a bar comprising a mixture of anionic and nonionic detergents. Examples 2 and 4 use 1% JR-400 in bars primarily based on sodium lauryl sulfate, which is an unacceptable surfactant for the present invention.

30 U.S. Pat. No. 3,761,418, Parran, Jr., issued Sept. 25, 1973, for "Detergent Compositions Containing Particle Deposition Enhancing Agents," discloses detergent compositions including a bar which contains 1% JR-400. However, the main surfactant is alkyl sulfate, which is unacceptable for the mild skin cleanser of
35 the present invention.

Thus, in view of the above, it will be appreciated that rather stringent requirements for skin cleansers limit the choice

of surface-active agents and final formulations represent some degree of compromise. Mildness is often obtained at the expense of effective cleansing and lathering. Conversely, mildness may be sacrificed for either preferred lathering characteristics, bar firmness, product stability, or all of these.

This invention represents a skin cleansing composition which provides ultra skin mildness, excellent lather and bar firmness. Therefore, one object of this invention is the development of skin cleansing compositions which exhibit superior skin mildness and moisturization while maintaining acceptable lather and bar properties.

Another object of the present invention is the development of skin cleansers having skin feel and mildness benefits.

Other objects will become apparent from the detailed description below.

Fig. 1 shows a static diffusion cell diagram.

Figs. 2-5 show comparative clinical mildness results of various cleansing compositions. Fig. 6 shows lather volume as a function of selected soap levels.

The invention relates to an ultra mild skin cleansing composition comprising

A. 20-70% by weight of mild synthetic surfactant as defined herein below;

B. 10-40% by weight moisturizer;

C. 0.1-5% by weight of polymeric skin feel and mildness aid as defined hereinbelow; and

D. 5.5-25% by weight soap;

wherein said synthetic surfactant to said soap have a weight ratio of from 2:1 to 12:1; and wherein said moisturizer to soap have a weight ratio of from 1:1 to 7:1; and

wherein said soap to polymer have a stoichiometric ratio (i.e. ratio of g.ion equivalents) of from 1:1 to 14:1.

The invention relates to an ultra mild skin cleansing composition with superior moisturization, excellent skin feel benefits

and lather performance. This mild cleansing composition is believed to provide superior moisturization, less skin irritation and facial tautness than commercially available toilet soap bars, synthetic toilet bars or other known surfactant based specialty skin cleansing products such as liquids and creams.

The composition of this invention leaves the skin feeling soft and smooth after washing and is also clinically milder than washing with water alone. The percentages herein are on a total composition weight basis unless otherwise specified.

A preferred composition contains: 10-40% moisturizer, e.g., free fatty acids; 40-70% of mild surfactants, e.g., alkyl glyceryl ether sulfonate (AGS) plus a co-surfactant selected from anionic alkyloyl(acyl) sarcosinates; 0.1-4% polymeric skin feel and skin mildness aids selected from cationic guar gums, cationic cellululosic resins, polysaccharides; homopolymers and copolymers of dimethyldiallylammonium chloride and nonionic guar gums, and soap at a level of 5.5-20%, preferably 6-15%.

The Surfactant

A mild surfactant as defined herein includes those which have a Relative Skin Barrier Penetration Value as defined hereinbelow of as close to zero as possible, preferably from 7 to 75 and more preferably less than 50. The mild synthetic surfactant is present in the composition at a level of 20-70%, preferably 40-70% and more preferably 50-65%. Also the synthetic surfactant and soap have ratios of 2:1 to 12:1, preferably 4:1 to 10:1 and more preferably 6:1 to 9:1.

The mild surfactants used in this invention are selected from alkyl glyceryl ether sulfonates(AGS), anionic acyl sarcosinates, methyl acyl taurates, N-acyl glutamates, alkyl glucosides, acyl isethionates, alkyl sulfosuccinates, alkyl phosphate esters, alkylether sulfates, ethoxylated alkyl phosphate esters, methyl glucose esters, protein condensates, mixtures of ethoxylated alkyl sulfates and alkyl amine oxides, betaines, sultaines and mixtures thereof.

Included in the surfactants are the alkyl ether sulfates with 1 to 12 ethoxy groups, especially ammonium and sodium lauryl ether sulfates. Alkyl chains for these surfactants are C_8 - C_{22} , preferably C_{10} - C_{18} .

A preferred primary mild surfactant is sodium coco glyceryl ether sulfonate which is mild and relatively nonirritating to the skin. This has been demonstrated in in vitro nonclinical mildness testing. While desirable to incorporate into a skin cleanser for its mildness properties, this coco AGS alone does not provide optimum lather creaminess. A sodium 90/10 coconut/tallow alkyl AGS distribution is preferred for creaminess. Salts other than the sodium salt such as TEA-, ammonium- and K-AGS and chain length distributions other than 90/10 coconut/tallow are usable at moderate levels. Also, some soap is added to improve lather volume and speed of lathering. Certain secondary co-surfactants used in combination with AGS can also provide a creamier and more stable lather. These secondary surfactants must also be intrinsically mild. One secondary surfactant that has been found to be especially desirable is sodium lauroyl sarcosinate (trade name Hamposyl L, made by Hampshire Chemical).

A preferred mild surfactant herein is an AGS/sarcosinate mix having a weight ratio of 1:1 to 5:1, preferably 2:1 to 4:1.

The amphoteric betaines and sultaines can be used as the sole surfactant, but are more preferred as a co-surfactant. Nonionics cannot be used as the sole surfactant in this product because of their low foaming ability; however, they can be incorporated as a co-surfactant. The mild surfactant and co-surfactant preferably have a ratio of 1:1 to 5:1.

In Vitro Skin Barrier Penetration Test

Introduction

The skin barrier penetration test (see test apparatus in Fig. 1) is used to screen mild surfactants from nonmild surfactants. In this test the milder the surfactant, the lesser the skin barrier 4 is destroyed. Skin barrier destruction is measured by the relative amount of radio-labeled water ($^3\text{H}-\text{H}_2\text{O}$) which passes from the test solution 3 through the skin epidermis 5 into the distilled water contained in the diffusate chamber 6. (This test is also described by T. J. Franz in the J. Invest. Derm., 64, pp. 190-195, 1975.)

Test Apparatus

The skin barrier penetration test utilizes a static diffusion cell 1 diagrammed in Fig. 1. When studying surfactants which are not soluble at room temperature, water-jacketed cell tops (not

shown) are used to hold the test solution 3 reservoir 2 at a different temperature than the diffusate chamber 6.

Human skin epidermis 5 is placed on the flat area of the diffusate chamber 6 and the diffusate chamber 6 is filled with distilled water 11 for collection of the transported radio-labeled water. The diffusate chamber 6 is surrounded by a water jacket 12. Temperature control is provided by circulating water bath (outflow 8 and inflow 9) pumped from a water bath at a predetermined temperature. When applicable, a jacketed top is connected to a second circulator/temperature bath. The diffusate chamber 6 is typically maintained at 25°C. An O-ring joint clamp secures the test solution reservoir 2, sandwiching the skin epidermis 5 in between it and the diffusate chamber 6.

This test uses cells which have a diffusion area of 0.20 cm². The diffusate chamber 6 holds approximately 5 ml. The bottom of the chamber 6 is flattened to allow spinning of a small magnetic stirring bar 10. The bars 10 are spun by a series of permanent magnets attached to individual electric motors. The motors turn at 300 rpm.

The test solution reservoir 2 holds approximately 1 ml of test solution 3. During testing the tops of reservoir 2 are sealed with Parafilm (RTM) to inhibit evaporation and prevent spillage.

Skin Epidermis Preparation

Prior to testing, the skin epidermis is separated from the whole skin. The whole skin is thawed to room temperature and adipose tissue (cellular fat) is removed by surgical scalpel to the adipose-dermal junction. This "defatted" skin is immersed in 60°C water for 80 seconds. This heat treatment makes possible the separation of the epidermis from the dermis at their junction. The epidermis is placed on a sheet of aluminum foil and rinsed with hexane for 5 seconds to remove excess fat globules still clinging to the surface. It is then sprayed with distilled water for 5 seconds to wash away the hexane. The epidermis sample is wrapped in Saran Wrap (RTM) placed in an aluminum pouch, and frozen in a standard freezer until needed.

Test Procedure

The following discussion describes a typical skin barrier penetration test. There are 30 cells per test, 5 replicates per treatment. This permits testing of 5 surfactant solutions plus a water control. Due to large variability in percutaneous transport (site to site and individual to individual), the water control is a mandatory reference point for each experiment.

Dose test solutions are prepared on a weight/weight basis on an analytical ($\pm .0001$ g) balance. Since each static diffusion cell 1 is dosed with 500 μ l, a minimum of 3 ml of each test solution 3 should be prepared (5 doses + standard). Each solution is spiked with $^3\text{H-H}_2\text{O}$ to an approximate activity of 1.8 $\mu\text{Ci/ml}$ dosing solution. This insures a sufficient level of ^3H transport for the sampling interval (2 hours).

During sample collection, the entire volume of the diffusate chamber 6 is emptied through the sampling port 7, rinsed, and refilled with fresh distilled water 11. Samples are collected at 2, 4, 6, 8 and 24 hours after initial dosing. The sample plus rinse is emptied into a scintillation vial, brought to 7 ml total volume and analyzed by liquid scintillation counting.

Standards are included for each sample. One hundred microliters (in duplicate) are placed in a scintillation vial and brought to 7 ml total volume with distilled water. The counts per minute (cpm) for diffusate samples are converted to equivalent mg water using the relation

$$\frac{\text{mg water}}{\text{vial}} = \text{cpm sample} \times \frac{\text{mg water standard}}{\text{cpm standard}}$$

The "water flux" is computed as a total sum penetrant using standard programs on a Tektronix (RTM) 4052A computer. It is important to realize that this "water flux" is a mg water equivalent of the ^3H flux. It does not represent a net water flux, since water is diffusing in both directions in this experiment. The relative flux (vs. water control) is an indication of barrier integrity.

Skin Barrier Penetration Test Values for AGS and cocoyl isethionate are given in Table 1. Water and sodium dodecylsulfate ("SDS") are the mild and harsh controls, respectively. The test

results are expressed in terms of the total amount of water penetrating through the skin in 24 hours. These amounts have also been converted to a relative scale (see Table 1) with water set to zero, SDS to 100, and AGS and isethionate interpolated in between.

TABLE 1
Skin Barrier Penetration

10	<u>Treatment</u>	Total mg Water Penetrating the Skin in 24 Hours	Relative Skin Barrier Penetration Value
	Water Control	12.0 ± 3.6	0
	1% AGS ^a	22.8 ± 7.9	9
	1% Cocoyl Isethionate	60.3 ± 17.7	40
15	1% SDS ^b - Reference	131.7 ± 68.0	100

^a AGS is sodium coco AGS

^b SDS is sodium dodecylsulfate

20 A mild surfactant as defined herein includes those which have a Relative Skin Barrier Penetration Value of as close to zero as possible. Most mild surfactants have Values of from about 7 to about 75, preferably 50 or less, and more preferably 40 or less. Surfactants which have Relative Skin Barrier Penetration Values
25 of greater than 75 can be used along with the mild surfactant at low levels in the compositions of this invention, so long as their use does not significantly change the clinical skin mildness of the total skin cleansing composition.

Moisturizers/Emollients

30 Moisturizers are included to provide the skin conditioning benefits and to improve the mildness of the product. The selection of the levels and types of moisturizers to be incorporated into the product is made without adversely affecting the stability of the product or its in-use characteristics, thereby delivering
35 good moisturization and lather.

The term "moisturizer" is often used within the cosmetic industry without very exact definition. The term is sometimes used as synonymous with emollient, and is then meant to describe a material which imparts a smooth and soft feeling to the skin surface.

There are two ways of reducing water loss from the stratum corneum. One is to deposit on the surface of the skin an occlusive layer which reduces the rate of evaporation. The second method is to add nonocclusive hygroscopic substances to the stratum corneum which will retain water, and make this water available to the stratum corneum to alter its physical properties and produce a cosmetically desirable effect. Nonocclusive moisturizers also function by improving the lubricity of the skin.

Both occlusive and nonocclusive moisturizers can work in the present invention. Some examples of moisturizers are long chain fatty acids, liquid water-soluble polyols, glycerin, propylene glycol, sorbitol, polyethylene glycol, ^{polyethylene glycol and propylene glycol} ~~ethoxylated/propoxylated~~ ethers of methyl glucose (e.g., methyl gluceth-20) and ^{polyethylene glycol and propylene glycol} ~~ethoxylated/propoxylated~~ ethers of lanolin alcohol (e.g., Solulan (RIM)-75).

The moisturizers useful in the present invention are used at a level of 10% to 40% by weight of the composition. The preferred and more preferred levels of moisturizers are, respectively, 10% to 30% and 12% to 25%. The preferred moisturizers are the coco and tallow fatty acids. Some other preferred moisturizers are the nonocclusive liquid water-soluble polyols and the essential amino acid compounds found naturally in the skin. The most preferred moisturizer is a mixture of stearic and lauric acids having a ratio of from 4:1 to 1:1, preferably 2:1 to 1:1, and is present at a level of 15-20%. These moisturizers also aid in solid bar integrity at said levels. The moisturizer, at these levels, provides superior moisturization. In the composition of this invention the high level of moisturizer with soap can also provide an enhanced lather and mildness.

The moisturizer to soap ratios are preferably 1:1 to 4:1.

1:1 to 7:1 and

Other preferred nonocclusive moisturizers are

sodium pyrrolidone carboxylic acid, lactic acid, urea, L-proline, guanidine and pyrrolidone. Examples of other nonocclusive moisturizers include hexadecyl, myristyl, isodecyl or isopropyl esters of adipic, lactic, oleic, stearic, isostearic, myristic or linoleic acids,

sodium isostearoyl-2-lactylate, sodium capryl lactylate, hydrolyzed protein and other collagen-derived proteins, aloe vera gel and acetamide MEA.

Some occlusive moisturizers include petrolatum, mineral oil, beeswax, silicones, lanolin and oil-soluble lanolin derivatives, saturated and unsaturated fatty alcohols such as behenyl alcohol, squalene and squalane, and various animal and vegetable oils such as almond oil, peanut oil, wheat germ oil, linseed oil, jojoba oil, oil of apricot pits, walnuts, palm nuts, pistachio nuts, sesame seeds, rapeseed, cade oil, corn oil, peach pit oil, poppyseed oil, pine oil, castor oil, soybean oil, avocado oil, safflower oil, coconut oil, hazelnut oil, olive oil, grape seed oil and sunflower seed oil.

Other examples of both types of moisturizers are disclosed in "Emollients -- A Critical Evaluation," by J. Mausner, Cosmetics & Toiletries, May 1981, incorporated herein by reference.

The Polymeric Skin Feel and Skin Mildness Aids

The polymeric skin feel and mildness aids used in the present invention are selected from cationic and nonionic polysaccharides; cationic and nonionic homopolymers and copolymers derived from acrylic and/or methacrylic acid; cationic copolymers of dimethyldiallylammonium chloride and acrylic acid; cationic homopolymers of dimethyldiallylammonium chloride; cationic polyalkylene and ethoxypolyalkylene imines, and mixtures thereof; Reduced skin irritation benefits as measured by patch testing of cationic and nonionic types of polymers are set out in "Polymer JR for Skin Care" Bulletin, by Union Carbide, 1977. The cationics are preferred over the others because they provide better skin feel benefits. Examples of the cationic polymers and the nonionic polymers useful in the present invention are set out below.

The amount of polymeric skin feel and mildness aids used in the composition of the present invention is from 0.1% to 5%, preferably from 0.3% to 4%.

The soap to polymer present in the composition of this invention has a stoichiometric ratios of 1:1 to 14:1, preferably 4:1 to 10:1 on a charge basis.

In order to achieve superior skin mildness and moisturization for this composition, it was discovered that a combination of moisturizer (10-40% in the composition) and a selected polymeric ingredient, e.g., Polymer JR-400, made by Union Carbide Corporation, is required.

Other types of high molecular weight polymeric skin feel and skin mildness aids, such as nonionic guar gums, Merquats 100 and 550, made by Merck & Co., Inc; JAGUAR (RTM) C-14-S made by Stein Hall; Mirapol A15 made by Miranol Chemical Company, Inc.; and Galactasol 811[®], made by Henkel, Inc.; plus others, are usable. The polymer also provides enhanced creamy lather benefits.

The nonionic polymers found to be useful include the nonionic polysaccharides, e.g., nonionic hydroxypropyl guar gums, offered by Celanese Water Soluble Polymers, a Division of Celanese Corp. A preferred nonionic hydroxypropyl guar gum material is JAGUAR^R HP-60 having molar substitution of about 0.6. Another class of useful nonionics is the cellulosic nonionic polymers, e.g., HEC and CMC.

The cationic polymers employed in this invention also provide a desirable silky, soft, smooth in-use feeling.

While not being bound to any theory, it is believed that cationic polymers chemically interact with the anionic surfactants (e.g., AGS and sarcosinates) to form complexes which may enhance the mildness to skin characteristics of the already mild surfactants. Also, there is reason to believe that the positively charged cationic polymers can bind with negatively charged sites on the skin to provide a soft skin feel after use. Not to be bound by any theory, it is believed that the greater the charge density of the cationic polymer, the more effective it is for skin feel benefits.

Other suitable cationic polymers are the cationic starches, e.g., StaLok[®] 300 and 400 made by Staley, Inc.

Soaps

Soaps are included in the compositions of this invention for improved lather, particularly under heavy soil loads. It is also surprising that the soap as used in the compositions of this invention provide clinical benefits for less skin dryness at levels up to 25% and less skin irritation (erythema) at levels up to 15%.

Soaps are used at levels of 5.5-25%, preferably 6-15%, and most preferably at a level of 6-8%. The soaps are preferably made in situ via adding a base, e.g., NaOH, to convert free fatty acids in the composition mix. A minor amount of soap, about 0.1-0.5% as shown in Tables 2, 3 and 5, is present as a by-product of the synthetic surfactant. See the process for making an ultra mild bar set out below.

The preferred ultra mild skin cleansing composition is a bar comprising: 20-50% mild synthetic surfactant; 10-40% moisturizer; 0.1-5% polymeric skin feel aid, and 5.5-25% soap; wherein the synthetic surfactant and soap has a ratio of from 2:1 to 12:1; the moisturizer to soap has a ratio of from 1:1 to 7:1; and the soap to polymer has a stoichiometric ratio of from 1:1 to 14:1.

Preferably the synthetic ^{surfactant} to soap ratio is 4:1 to 10:1, and more preferably 6:1 to 9:1. The preferred moisturizer to soap ratio is from 2:1 to 4:1. The preferred soap to polymer stoichiometric ratio is from 4:1 to 10:1.

The preferred level of polymeric skin feel and mildness aid is from 2-5%.



The soap is preferably made in situ from free fatty acids and a base selected from magnesium hydroxide and potassium hydroxide, preferably sodium hydroxide and triethanolamine. The preferred soap level is 6-15% of the composition, particularly in
5 ultra mild skin cleansing compositions which contain from 10-30% fatty acids. Particularly preferred fatty acids are mixtures of stearic and lauric acids having a ratio of from 2:1 to 1:1.

Optionals

The usual optionals can be used in the composition of the
10 present invention.

Perfumes may be used in formulating the skin cleansing products, generally at a level of about 0.1% to about 1.5% of the composition. Colorants, and also fillers such as talc and clay, may also be used. Preservatives, e.g., EDTA, generally at a
15 level of less than 1% of the composition, may be incorporated in the cleansing products to prevent microbiological growth. Anti-bacterials can also be incorporated, usually at levels up to 1.5%.

Process for Making the Ultra Mild Toilet Bar

In order to make the product, an analysis of the surfactant
20 paste is needed. To illustrate the process, an AGS paste with the following nominal analysis will be used.

Cationic Titration for $\text{SO}_3 = 48.5\%$ AGS

NaCl = 1.5%

Moisture = 43%

25 After the composition of the AGS paste is determined, the crutcher mix is calculated using the AGS/sarcosinate ratio of 4:1, 55% moisture in the crutcher mix and about 3.5% NaCl in the final bar.

Crutching

30 Assuming 200 lb (90.8 Kg) total mix and the above AGS analysis.

1. Heat crutcher ~~mix~~ to 200°F (93°C) by adjusting steam and water valves.
2. Add 83.7 lbs. (38 Kg) of AGS paste.
- 35 3. Turn on agitator and recirculation pump.

4. When the AGS temperature is 130°F (54°C) add 12.0 lbs. (5.45 Kg) of stearic acid.
5. When the crutcher mix temperature is back up to 130°F (54°C) add 2.18 lbs. (991g) of 50% NaOH.
- 5 6. Add 2-3 gal. (7.6-11.4 liters) of hot water (160-180°F, 71-82°C) as needed to thin the crutcher mix to obtain good mixing.
7. Allow the crutcher contents to mix for about 20 minutes and maintain the temperature at 130-140°F (54-60°C).
- 10 8. Add 10.15 lbs. (4.61 Kg) of Hamposyl L-95 C₁₂ sarcosinate.
9. Add 2-3 gal. (7.6-11.4 liters) of hot water as needed for good mixing.
10. Maintain 140°F (60°C).
- 15 11. In a suitable container, slurry 2.71 lbs. (1.23 Kg) of JR-400 in hot water.
12. Add the JR-400 slurry to the crutcher; add more hot water as needed to get to 55% H₂O in the crutcher.
13. Add 8.0 lbs. (3.63 Kg) of lauric acid.
- 20 14. Add 0.22 lbs. (101 g) of TiO₂.
15. Add 1.16 lbs. (527 g) of NaCl.
16. Mix for at least 20 min. and bring the crutcher mix temperature up to 175°F (79°C).

Drying

- 25 The crutcher mix is dried on a atmospheric drum dryer. The optimum operating conditons are 3-4 rpm, 80 psi (551.5 kPa) and approximately 310°F(154°C). The nip of the drums are set to give a fairly thin flake. The nip is increased to give thicker flake and to allow the higher moisture in the flake. Typical moisture for
- 30 the flake is 2-3%. The three ways to regulate the moisture, in the order of preference, are (1) opening or closing the drum gap; (2) increasing or decreasing the drum speed; and (3) increasing or decreasing the steam pressure.

Amalgamating

- 35 The flakes are weighed and mixed in a batch amalgamator to obtain a uniform flake sample for moisture analysis. Once the

percent water is known, water and perfume are added to bring the flakes up to final target formula. About $\frac{1}{8}$ extra water is added to allow for losses during processing. The sticky, perfumed flakes are transferred to the mill hopper.

5 Milling

The 3-roll soap mills are set up with the first roll at 100°F (38°C) and the other 2 rolls at about 70°F (21°C). The soap is passed through the mills several times and forms sticky flakes. These flakes are put into a sealed plastic bag and placed in the
10 100°F (38°C) constant temperature room overnight. The flakes are milled one final time prior to plodding.

Plodding and Stamping

The plodder is set up with the barrel temperature at ambient to cold and the nose temperature at 120-130°F (49-54°C). The
15 ideal plodder is a dual stage plodder that allows use of a vacuum of about 15-25 in. Hg. (50.8 - 84.6 KPa). The plugs should be cut in 5" (12.7 cm) sections and stamped with a cold die block and stamp using a die liquor such as alcohol.

CLINICAL ASSESSMENT OF ULTRA MILD TOILET BAR

20 The clinical test procedure used to evaluate several toilet bar formulations for skin mildness is given below.

Clinical Mildness Arm Wash Test Procedure (2 Washes Per Day for Two Weeks for 23 Washes)

The washings were performed by technical personnel. Each
25 panelist was assigned a set of two test products for washings. These bars were all kept under technical personnel control in plastic dishes. The test procedure is:

1. Wet the anterior portion (inside) of the forearm of the panelist with 90-100°F (32-38°C) water (6-8 grains hardness -
30 102.6 - 136.8 mgCaCO₃/litre) from the elbow to the wrist.
2. Pick up and wet the bar.
3. Rub bar from wrist area up to the elbow and back down again in approximately 1 second. Repeat rubbing 5 more up-down cycles for a total of 6 cycles in 6 seconds.
- 35 4. Put bar aside and rub area in the same manner with the hand for 10 additional seconds (approximately 10 additional up-down cycles).

5. Rinse forearm thoroughly with tap water. Pat dry with clean paper towels.
6. Repeat procedure on other arm.

For Panelists on Water Treatment

- 5 1. Wet the anterior portion (inside) of the forearm with 90-100°F (32-38°C) water (6-8 grains hardness - 102.6 - 136.8 mg CaCO₃/litre) from the elbow to the wrist.
2. Rub hand (up and down) on arm for 15 seconds.
3. Rinse forearm thoroughly with tap water. Pat dry with clean paper towels.

Test Product Formulations

The formulations of the products tested clinically are given in the Tables. The molar equivalents of only Example F are given in Table 3 to illustrate the stoichiometric ratios of several key ingredients. The moles of polymer are based on the average molecular weight of monomer containing one positive charge.

TABLE 2

Test Product Formulations

20	Examples: <u>Ingredients</u>	<u>A</u>	<u>C</u>	<u>F</u>	<u>G</u>
		5% JR-400	0.5% JR-400	3% JR-400	1.5% JR-400
	NaAGS	36.0	36.0	48.14	48.74
	NaC ₁₂ Sarcosinate	9.5	9.5	11.07	11.21
	Stearic Acid	24.5	24.5	9.58	10.22
25	Lauric Acid	-	-	6.39	6.81
	JR-400	5.0	0.5	3.08	1.56
	Dextrin	8.7	14.2	-	-
	Unsulfonated				
	Alcohol	3.9	3.9	1.24	1.26
30	Na Laurate	0.4	0.4	0.49	0.50
	Na Soap				
	(60/40 T/C)	-	-	7.69	7.79
	NaCl	3.4	3.4	3.29	4.03
	Na ₂ SO ₄	1.6	1.6	1.24	1.26
35	TiO ₂	0.5	0.5	0.26	0.26
	Perfume	0.5	0.5	1.03	1.04
	Water	6.0	5.0	6.43	5.26

Control H (mildness results are shown in Fig. 3 as Curve -H)- is substantially the same formulation as Example G, except that the formulation contained no JR-400 polymer. Controls E and K are 50/50 tallow/coconut super fatted bar soap.

5 Control D is Dove(RTM) (a commercially available bar) which does not contain polymeric skin feel and polymeric skin mildness aids.

TABLE 3

Molar Equivalents of Example F with 3% JR-400

10	<u>Ingredient</u>	<u>m Moles/100 g Bar</u>
	Sodium Soap (60/40)	27.7 "
	Sodium ACS	116.0
	Unsulfonated Alcohol	15.8
	Stearic Acid	32.8
15	Sodium Lauryl Sarcosinate	36.7
	Sodium Laurate	2.1
	Lauric Acid	31.1
	Polymer JR-400	4.3

20 Note from Table 3 that the stoichiometric ratio of soap (27.7) to polymer (4.3) is about 7:1. It will be appreciated that the total synthetic surfactant to soap weight ratio in Example F reported in Table 2 is about 7.5:1, and the total moisturizer to soap ratio is about 2:1.

25 **Clinical Test Skin Grading Scales**

The forearm skin grading scales reported herein are set out below.

Forearm Grading Scale

Skin Dryness

- 30 0 Perfect skin.
- 1.0 Patches of checking and/or slight powderiness and occasional patches of small scales may be seen. Distribution generalized.
- 2.0 Generalized slight powderiness. Early cracking
- 35 or occasional small lifting scales may be present.

- 3.0 Generalized moderate powderiness and/or moderate cracking and lifting scales.
- 4.0 Generalized heavy powderiness and/or heavy cracking and lifting scales.
- 5 5.0 Generalized high cracking and lifting scales. Eczematous change may be present. Powderiness may be present but not prominent. May see bleeding crack.
- 10 6.0 Generalized severe cracking. Eczematous change may be present. Bleeding cracks may be present. Scales large, may be beginning to disappear.

Forearm Grading Scales

	<u>Skin Erythema</u>	<u>Skin Smoothness</u>
15	0 No redness	0 Very high smoothness/slick
	1.0 Barely detectible redness	1.0 Extreme smoothness
	2.0 Slight redness	2.0 Substantial smoothness
	3.0 Moderate redness	3.0 Moderate smoothness
	4.0 Heavy or substantial redness	4.0 Slight smoothness
20	5.0 Severe redness	5.0 Barely detectible smoothness
	6.0 Extreme redness	6.0 No smoothness

The term "generalized" as used herein means that more than 50% of the surface area of the forearm exhibits the attribute. Whole unit skin grades reflect generalized condition. Half units are used to represent intermediate conditions.

Clinical Test Results

The clinical mildness test results expressed in terms of skin grades for dryness, erythema, and smoothness are shown below in Table 4 and the skin dryness grades are plotted vs. time in Fig. 3. Skin dryness grades from a second clinical test are shown in a similar fashion in Fig. 2. Water treatment results are shown as curves B and I in Figs. 2 and 3, respectively. Curves D and J are the results of DOVE (RTM) (a commercially available bar).

TABLE 4
Two Week Clinical Results
Skin Grades

	<u>Product</u>	<u>Dryness</u>	<u>Erythema</u>	<u>Smoothness</u>
5	F	1.16 H-K	1.04 J,K	2.76 H
	G	1.25 J,K	1.07 J,K	2.75
	H	1.37 F,J,K	1.19 K	2.58 F,I,K
	I	1.44 F,K	1.23 K	2.81 H
	J	1.55 F,G,K	1.28 F,G,K	2.72 K
10	K	2.17 F-J	1.75 F-J	2.87 I,J

In Table 4 the visual skin grades for dryness, erythema and tactile grades for smoothness are all assigned from a six point scale (0-6), the lower the grade the better the skin mildness. Letter codes following the skin grades indicate those products which have statistically significant ($\alpha = 0.05$) differences in skin grades from the specified Example product. Also see Figs. 2 and 3 for a graphic report of these data.

These data show that the toilet skin cleansing bars of this invention (Examples G and F) result in the desired mild skin cleansing action with improved skin feel and moisturization benefits. Note that Control A listed in Table 2 is as mild as "B" water (see Fig. 2). It will be appreciated that bars F and G are actually milder than water (I) as shown in Table 4 and Fig. 3. It will also be appreciated that the toilet skin cleansing bars G and F of the present invention are not only mild but also have excellent bar smear and lather performance.

It will be appreciated that the addition of selected levels of soap surprisingly provide improved skin mildness. Products L-P set out in Table 5 show the selected levels. The effect of varying amounts of soap in this invention on clinical skin condition is demonstrated in Table 6. The skin dryness and skin erythema grades are respectively plotted in Figs. 4 and 5 vs. soap level. The overall conclusion from these data is that selected levels of soap surprisingly provide additional clinical mildness benefit for reduced skin dryness and erythema and enhanced skin smoothness. The reduced skin dryness benefit shown in Fig. 4 is

maximum at the 7-20% soap level, whereas the reduced erythema shown in Fig. 5 (and smoothness benefit in Table 6) is maximum at about the 7% soap level.

5

TABLE 5

Test Product Formulations

<u>Test Product</u> <u>Ingredients</u>		<u>L</u>	<u>M</u>	<u>N</u>	<u>O</u>	<u>P</u>
Na AGS		43.15	43.15	33.15	16.00	9.15
10 NaC ₁₂						
Sarcosinate		10.80	10.80	8.30	-	2.30
Stearic Acid		16.00	11.50	11.84	12.89	12.73
Lauric Acid		10.67	7.67	7.89	8.59	8.49
JR-400		3.00	3.00	3.00	3.00	3.00
15 Unsulfonated						
Alcohol		0.83	0.83	0.64	0.31	0.18
Na Laurate		0.48	0.48	0.37	-	0.10
Na Soap		-	7.50	20.05	45.01	49.99
NaCl		4.04	4.04	4.04	4.00	4.03
20 Na ₂ SO ₄		1.20	1.20	0.92	0.44	0.25
TiO ₂		0.25	0.25	0.25	0.25	0.25
Perfume		1.00	1.00	1.00	1.00	1.00
Water		8.50	8.50	8.50	8.50	8.50
Miscellaneous		0.08	0.08	0.06	-	0.02

25

TABLE 6

Two Week Clinical Results

Skin Grades

<u>Test</u>					
30	<u>Product</u>	<u>% Soap</u>	<u>Dryness</u>	<u>Erythema</u>	<u>Smoothness</u>
	L	-	2.11	1.78	1.79
	M	7.50	1.89	1.64	1.66
	N	20.05	1.91	1.80	1.76
	O	45.01	2.15	1.75	1.73
35	P	49.99	2.09	1.79	1.73

Laboratory Assessment of Bar Performance

The following test procedures are used to evaluate the critical bar performance attributes of lather volume and bar firmness during use.

5

Bar Soap Firmness Test

The bar soap firmness test measures two quantitative parameters of bar firmness after sitting in a wet soap dish for 16 hours: (1) depth of bar surface softening and (2) amount of bar messiness by weight percent of initial bar weight.

10

Equipment

The following equipment is used:

1. "Alathon" rectangular plastic soap dishes (U. S. Pat. No. 2,842,178) with low ridges.
2. Precision Scientific Co. penetrometer with ball shaft, 300 gram weight.
3. Analytical weight balance.
4. Scraper or spatula for removing gelatinous layer.
5. Paper towels.

15

Procedure

20

The following procedure is used:

1. Test bar(s) are weighed to establish an initial dry weight data point.
2. If desired, initial dry bar hardness/softness may be determined by penetrometer with cone shaft and 200 gram weight for penetration measurement.
3. Bar(s) are placed centrally on soap dishes containing 5 ml. of distilled water at room temperature. Rock the dish to break water tension.
4. Store overnight (approximately 18 hours) in standard room conditions of temperature and humidity (80°F/80% RH).
5. Next morning, remove bar(s) with a careful vertical lift and invert for testing.
6. Run ball penetrometer reading on the gelatinous surface, generally three readings in a triangular design; record the average mm depth reading of gelatinous layer. The shaft ball is cleaned with a tissue after each reading.

25

30

35

7. Bar(s) are then carefully scraped to remove the gelatinous layer down to the solid surface. Do not use enough force to remove any of the solid surface area. Slight remaining gelatinous soap may be wiped from bar with paper towels until surface appears firm and somewhat dry.
8. Bar(s) are then allowed to openly dry out on bench surface for no more than one hour.
9. Bar(s) are then weighed to determine the difference from the initial dry untested weight. The weight difference divided by the original weight is calculated into % weight loss or % gelatinous smear weight. Data is recorded.
10. Comparison of any given series of data points obtained will illustrate a formula characteristic that may tend to relate to poorer smear or "messiness" attribute. Averaging of historical data on control type formulas (such as an existing National brand formula) can be used to establish a benchmark for comparison.

Bar Soap Handwash Lather Volume Test

The handwash lather test is used to provide in-use lather volume measurements for the lather performance of skin cleansing bars. The test measures both the ultimate lather volume generated and the volume which is generated after a very short lathering period (to reflect lathering ease). The lather volumes are generated under both soil-loaded and nonloaded conditions.

Synthetic soil is used for the soil-loaded lather volume test reported herein. Its formula and procedure for making it are set out below.

TABLE 9
Synthetic Soil

	<u>Ingredients</u>	<u>Wt. %</u>
	Hyfac 430 ^a	1.87
5	Lauric Acid ^b	1.42
	Neo-fat 14 ^c	5.68
	Neo-fat 16 ^d	11.16
	Neo-fat 18 ^e	5.40
	Neo-fat 90-04 ^f	9.81
10	Industrene 226 ^g	1.26
	Paraffin Wax	7.30
	Squalane ^h	3.70
	Lanolin Anhydrous	19.40
	Coconut Oil	3.30
15	Tallow	<u>29.70</u>
		100.00%

a Emery Industries, Inc., Cincinnati, Ohio

b Emery Industries, Inc., Cincinnati, Ohio

20 c Armour Industrial Chemical Co., Chicago, Illinois

d Armour Industrial Chemical Co., Chicago, Illinois

e Armour Industrial Chemical Co., Chicago, Illinois

f Armour Industrial Chemical Co., Chicago, Illinois

g Humko Products, Memphis, Tennessee

25 h Robeco Chemicals, Inc., New York, New York

Procedure

1. Heat above materials together stirring continuously between 160-175°F.
2. Mix 25 parts of above formula with 25 parts of, a 5% to 80%
30 tallow/20% coconut soap solution and 50 parts of distilled water at 150°F.
3. Cool mixture to room temperature while stirring constantly.
4. Store in covered glass container.

Equipment

- 35 The following equipment is used:

1. Water source and sink with temperature control. The water source should be medium hardness (6-9 grain/gallon) for most testing, although water of lower and higher hardness can be used for special purposes.
- 5 2. Synthetic soil (see Table 9).
3. Paper towels.
4. Test bars.
5. Control bars (i.e., usually marketed brands such as CAMAY (RTM) or the bar of Example D).

10 Procedure

The following procedure is used:

1. Set temperature at 95-100°F.
2. Rub 0.22 cc of soil on hands (if doing soil-loaded test).
3. Wet hands.
- 15 4. Rotate bar 3 times in both hands.
5. Add a little water, rub both hands 5 times.
6. Rotate hands 3 times (without soap), grade for flash volume.
7. Rotate 7 more times, grade for ultimate volume.
8. Collect lather and deposit on sink top.
- 20 9. Compare volume with standard bar target volume and assign grade.

Grading Scale

	<u>Non-Soil Loaded</u>	<u>Soil-Loaded</u>
25	10 - Very much higher than target	
	9 - Higher than target	
	8 - Target volume*	Very much higher than target
30	7 - Lower than target	Higher than target
	6 - Very much lower than target	Target volume*
	5 - Unacceptably lower than target	Slightly lower than target
35	4 - Unacceptably lower than target	Lower than target

Grading Scale - Continued

	<u>Non-Soil Loaded</u>	<u>Soil-Loaded</u>
	3 - Unacceptably lower than target	Very much lower
5	2 - Unacceptably lower than target	Unacceptably low

* Based on 50 tallow/50 coconut - 7% free fatty acid bar.

10

TABLE 7
Bar Formulations

		Na AGS	NaC ₁₂ Sarcosinate	<u>Free Fatty Acid</u>		Soap	JR-400
15	<u>Product</u>	<u>Wt.%</u>	<u>Wt.%</u>	<u>Stearic</u>	<u>Lauric</u>	<u>Wt.%</u>	<u>Wt.%</u>
	1	42	9.0	18	-	-	0.5
	2	43	10.8	15	10	-	3.0
	3	43	11.4	10	6.8	7.0	3.0
	4	Camay (RTM)					
20	5	Dove (RTM)					

20

TABLE 8
Bar Firmness and Hand Lather

25	Pene-	Smear Wt.	Hand Lather				
	trometer	% of	Ultimate Volume		Flash Volume		
	Product	mm	Bar Wt.	w/o Soil	Soil	w/o Soil	Soil
	1	2.9	8.7	8	3	8	2.5
	2	2.6	6.5	8	2.5	8	2.5
	3	2.3	7.2	8	5	8	4
30	4	1.1	3.2	8	6.5	8	6.5
	5	1.8	5.0	8	3	8	3

30

The results listed in Table 8 show the importance of utilizing the preferred ratio of stearic to lauric fatty acids to achieve an acceptable level of firmness and the importance of utilizing soap to achieve an acceptable soil-loaded lather

35

performance. The acceptable levels of bar firmness are expressed as a penetrometer value of equal to or less than 2.7 and smear weight of equal to or less than 7.5. The acceptable levels of hand lather are: ultimate volume, without soil equal to or greater than 7.5; with soil, equal to or greater than 4. Flash volume acceptable levels are: without soil, equal to or greater than 7.5; with soil, equal to or greater than 4.

Control 1 described a sodium AGS bar formula with acceptable mildness but with poor lather and firmness performance. Control 2 demonstrates the advantage of using a 3:2 ratio of stearic:lauric fatty acids. ^{Product} Example 3 demonstrates the benefit of neutralizing the fatty acids in situ forming 7% soap to boost the soil-loaded lather.

Control 4 is a 50/50 tallow/coconut superfatted soap bar which demonstrates very acceptable bar firmness and lather properties. Control 5 is the same as Control D which shows acceptable firmness and lather properties.

Bar Soap Blender Lather Volume Method

Purpose

This quantitative blender lather volume method is used to compliment hand lathering evaluations of different soap formulations.

Equipment

1. Specially designed 500 ml graduate cylinder adapted with blender impeller.
2. Waring Blender Base No. 7011-31BL92 with high/low speed switch.
3. Variac (RTM) Voltage Regulator

Procedure

1. Dissolve 5% by weight of the bar material to be evaluated in distilled water by heating to 150°F (65.7°C).
2. Add 10 mls of soap solution to 90 mls of water both at 38°C in a 100 ml graduate.
3. Immediately pour resultant 0.5% solution into blender graduate.

4. Set blender to run at about 4700 rpm's (high speed) and blend solution for fifteen seconds.
5. After blending read lather height on cylinder.
6. Replicate a minimum of three times and average.

5 The effect of varying the amount of soap in ^{Product} ~~Example~~ L (by converting free fatty acid to soap with NaOH) on lather volume (using the blender method) is shown in Fig. 6. This curve demonstrates that lather volume is increased by the selected levels of soap in the formulation.

10 Thus, it has been demonstrated that an ultra mild skin cleansing composition can be made with selected mild synthetic surfactants; selected moisturizers; polymeric skin feel and mildness aids; and selected levels of soap.

15

20

25

30

35

29

CLAIMS

1. An ultra mild skin cleansing composition comprising:
 - A. 20-70% by weight of mild synthetic surfactant selected from alkyl glyceryl ether sulfonates (AGS); anionic acyl sarcosinates; methyl acyl taurates; N-acyl glutamates; alkyl glucosides; acyl isethionates; alkyl sulfosuccinates; alkyl phosphate esters; alkyl ether sulfates; ethoxylated alkyl phosphate esters; methyl glucose esters; protein condensates; mixtures of ethoxylated alkyl sulfates and alkyl amine oxides; betaines; sultaines; and mixtures thereof;
 - B. 10-40% by weight of moisturizer;
 - C. 0.1-5% by weight of polymeric skin feel and mildness aid selected from cationic and nonionic polysaccharides; cationic and nonionic homopolymers and copolymers derived from acrylic and/or methacrylic acid; cationic copolymers of dimethyldiallylammonium chloride and acrylic acid; cationic homopolymers of dimethyldiallylammonium chloride; cationic polyalkylene and ethoxypolyalkylene imines; and mixtures thereof;
 - D. 5.5-25% by weight of soap;wherein said synthetic surfactant to said soap have a weight ratio of from 2:1 to 12:1; and
wherein said moisturizer to soap have a weight ratio of from 1:1 to 7:1; and
wherein said soap to polymer have a stoichiometric ratio (i.e. ratio of g.ion equivalents) of from 1:1 to 14:1.



2. The ultra mild skin cleansing composition of Claim 1 wherein said synthetic surfactant to soap ratio is 4:1 to 10:1.
3. The ultra mild skin cleansing composition of Claim 1 wherein said synthetic surfactant to soap ratio is 6:1 to 9:1.
4. The ultra mild skin cleansing composition of Claim 1 wherein said moisturizer to soap ratio is from 1:1 to 4:1.
5. The ultra mild skin cleansing composition of Claim 1 wherein said soap to polymer stoichiometric ratio is from 4:1 to 10:1.
6. The ultra mild skin cleansing composition of Claim 1 wherein said mild surfactant is an AGS/sarcosinate mix having ratio of 1:1 to 5:1.
7. The ultra mild skin cleansing composition of Claim 6 wherein said AGS/sarcosinate has a ratio of 2:1 to 4:1.
8. The ultra mild skin cleansing composition of Claim 1 wherein said composition is a toilet bar which contains from 5.5-20% soap and wherein said soap is made in situ from free fatty acids and a base selected from KOH, $Mg(OH)_2$, NaOH and triethanolamine.
9. The ultra mild skin cleansing composition of Claim 8 wherein said soap is 6-15% of said composition; and wherein said composition contains from 10-30% fatty acids; and wherein said base is selected from NaOH and triethanolamine.



10. The ultra mild skin cleansing composition of Claim 9 wherein said fatty acids are stearic and lauric having a ratio of 4:1 to 1:1.

11. The ultra mild skin cleansing composition of Claim 10 wherein said moisturizer is a mixture of stearic and lauric acids having a ratio of from 2:1 to 1:1 and is present in said composition at a level of from 15-20%.

12. The ultra mild skin cleansing composition of Claim 1 wherein the moisturizer is selected from
glycerin, polyethylene glycol, propylene glycol, sorbitol, polyethylene glycol and propylene glycol ethers of methyl glucose, polyethylene glycol and propylene glycol ethers of lanolin alcohol, sodium pyrrolidone carboxylic acid, lactic acid, and L-proline.

13. The ultra mild skin cleansing composition of Claim 1 wherein said moisturizer is a mixture of stearic and lauric acids having a ratio of from 4:1 to 1:1.

14. The ultra mild skin cleansing composition of Claim 1 wherein said moisturizer is selected from
petrolatum, mineral oil, beeswax, silicones, lanolin and oil-soluble lanolin derivatives, saturated and unsaturated fatty alcohols such as behenyl alcohol, squalene and squalane and animal and vegetable oils such as almond oil, peanut oil, wheat germ oil, linseed oil, jojoba oil, oil of apricot pits, walnuts, palm nuts, pistachio nuts, sesame seeds, rapeseed, cade oil, corn oil, peach pit oil, poppyseed oil, pine oil, castor oil, soybean oil, avocado oil, safflower oil, coconut oil, hazelnut oil, olive oil, grape seed oil and sunflower seed oil.



* C L H S *

15. The ultra mild skin cleansing composition of Claim 1 wherein said moisturizer is present at a level of 10-30% of said composition and said moisturizer to soap ratio is from 2:1 to 4:1.

16. The ultra mild skin cleansing composition of Claim 1 wherein said moisturizer is present at a level of 10-30% of said composition.

17. The composition of Claim 1 wherein said polymer is selected from cationic and nonionic cellulosic resins.

18. The ultra mild skin cleansing composition of Claim 12 wherein said polymer is a cationic guar gum.

19. The ultra mild skin cleansing composition of Claim 1 wherein said polymer is present at 0.3-5% by weight; and said moisturizer is free fatty acids present at 10-30% by weight of said composition.

20. The ultra mild skin cleansing composition of Claim 1 wherein said mild synthetic surfactant is present at a level of 40-70% of said composition, and wherein said mild synthetic surfactant has a Relative Skin Barrier Penetration Value of from 7 to 75.

21. The ultra mild skin cleansing composition of Claim 1 wherein said mild synthetic surfactant is present at a level of ~~50~~-65% of said composition and wherein said synthetic surfactant has a Relative Skin Barrier Penetration Value of less than 50.



22. The ultra mild skin cleansing composition of Claim 1 wherein

- A. said mild surfactant is present at 40-70%;
- B. said moisturizer at 10-40%;
- C. said polymer at 0.1-4%; and
- D. said soap at 6-15%.

23. The ultra mild skin cleansing composition of Claim 21 wherein said composition is a toilet bar which contains from 6-8% soap, 12-25% moisturizer, and 2-5% polymeric skin feel and mildness aid.



* C L M S *