





FIGURE 1




FIGURE 2

3/6




FIGURE 3




FIGURE 4

Compound	Sodium Channel Site 2, 3H-Batrachotoxinin Binding					Serotonin 5-HT _{2A} Receptor, 3H-Ketanserin Binding				
	Maximal Concentration	Percentage Inhibition	IC ₅₀	K _i	nH	Maximal Concentration	Percentage Inhibition	IC ₅₀	K _i	nH
Example 1	1 μ M	65	0.39 μ M	0.35 μ M	0.68	10 μ M	88	1.22 μ M	0.35 μ M	0.95
Example 1A	1 μ M	53	0.78 μ M	0.71 μ M	0.89	30 μ M	85	6.59 μ M	1.88 μ M	1.16
Example 1B	10 μ M	84	1.91 μ M	1.75 μ M	0.84	30 μ M	87	5.90 μ M	1.69 μ M	1.11
Example 1C	1 μ M	68	0.39 μ M	0.35 μ M	0.86	1 μ M	60	0.61 μ M	0.17 μ M	0.80
Example 1D	1 μ M	61	0.43 μ M	0.39 μ M	0.66	10 μ M	75	3.03 μ M	0.86 μ M	0.86
Example 1E	1 μ M	66	0.51 μ M	0.46 μ M	0.89	3 μ M	66	1.46 μ M	1.46 μ M	0.89
Dibucaine	nd	nd	0.68 μ M	0.62 μ M	1.23	nd	nd	nd	nd	nd
Ketanserin	nd	nd	nd	nd	nd	nd	nd	0.47 nM	0.14 nM	1.06

FIGURE 5




FIGURE 6