

[1,2,4]TRIAZOLO[1,5-a]PYRIMIDIN-7-YL COMPOUND

FIELD OF THE INVENTION


5

The present invention relates to a novel [1,2,4]triazolo[1,5-a]pyrimidin-yl derivative as inhibitor of phosphodiesterase 2 (PDE2). The invention is also directed to pharmaceutical compositions comprising the compound, to processes for preparing such compound and compositions, and to the use of such compound and compositions for the prevention and treatment of disorders in which PDE2 is involved, such as neurological and psychiatric disorders.

BACKGROUND OF THE INVENTION

15 Phosphodiesterases (PDEs) are a family of enzymes encoded by 21 genes and subdivided into 11 distinct families according to structural and functional properties. These enzymes metabolically inactivate widely occurring intracellular second messengers, 3',5'-cyclic adenosine monophosphate (cAMP) and 3',5'-cyclic guanosine monophosphate (cGMP). These two messengers regulate a wide variety of biological processes, including pro-inflammatory mediator production and action, ion channel function, muscle contraction, learning, differentiation, apoptosis, lipogenesis, glycogenolysis, and gluconeogenesis. They do this by activation of protein kinase A (PKA) and protein kinase G (PKG), which in turn phosphorylate a wide variety of substrates including transcription factors and ion channels that regulate innumerable physiological responses. In neurons, this includes the activation of cAMP and cGMP-dependent kinases and subsequent phosphorylation of proteins involved in acute regulation of synaptic transmission as well as in neuronal differentiation and survival. Intracellular concentrations of cAMP and cGMP are strictly regulated by the rate of biosynthesis by cyclases and by the rate of degradation by PDEs. PDEs are hydrolases that inactivate cAMP and cGMP by catalytic hydrolysis of the 3'-ester bond, forming the inactive 5'-monophosphate (Scheme A).

Scheme A



On the basis of substrate specificity, the PDE families can be divided into three groups:

- 5 i) the cAMP-specific PDEs, which include PDE4, 7 and 8; ii) the cGMP-selective enzymes PDE5, 6 and 9; and iii) the dual-substrate PDEs, PDE1, 2 and 3, as well as PDE10 and 11.

Furthermore, PDEs are expressed differentially throughout the organism, including the central nervous system. Different PDE isozymes therefore may have different physiological functions. Compounds that inhibit selectively PDE families or isozymes may display particular therapeutic activity, fewer side effects, or both.

10 Phosphodiesterase 2A (PDE2A) inactivates intracellular signalling mechanisms reliant on cyclic nucleotide signalling mediated by cAMP and cGMP via their degradation (by hydrolyzing the biologically relevant second messengers cAMP and cGMP into non-signalling AMP and GMP, respectively). Such signalling pathways are known to play a role in the regulation of genes involved in the induction of synaptic plasticity.

The pharmacological inhibition of PDE2 therefore causes increased levels of synaptic plasticity (an underlying correlate of learning and memory), suggesting that PDE2A modulation may be a target for alleviating cognitive deficits seen in people suffering from disorders such as for example, schizophrenia, Alzheimer's disease, Parkinson's disease and other CNS disorders associated with cognitive dysfunction.

20 Phosphodiesterase 2A (PDE2A) is more abundantly expressed in the brain relative to peripheral tissues. The high expression of PDE2 in the limbic system (isocortex, hippocampus, amygdala, habenula, basal ganglia) suggests that PDE2 may modulate neuronal signalling involved in emotion, perception, concentration, learning and memory. Additionally, PDE2 is expressed in the nucleus accumbens, the olfactory bulb, the olfactory tubercle and the amygdala, supporting the suggestion that PDE2 may also be involved in anxiety and depression. (see for instance, Lakics, V. et al. (2010) Quantitative comparison of phosphodiesterase mRNA distribution in human brain and peripheral tissues. *Neuropharmacol.* 59, 367-374).

Additionally, PDE2 inhibitors have been shown to be beneficial in the reduction of oxidative stress-induced anxiety, supporting their use in the treatment of anxiety in neuropsychiatric and neurodegenerative disorders that involve oxidative stress, such as Alzheimer's disease, Parkinson's disease and multiple sclerosis.

- 5 PDE2 inhibitors have been shown to enhance long term potentiation of synaptic transmission and to improve memory acquisition and consolidation in the object recognition and in the social recognition tests in rats. Furthermore, PDE2 inhibitors have been shown to reverse the MK-801 induced working memory deficit in the T-maze in mice. PDE2 inhibitors have also been shown to display activity in forced swim test and light/dark box models; and to show anxiolytic-like effects in elevated plus-maze, hole-board and open-field tests and to prevent stress-induced changes in apoptosis and behaviour.


Thus, PDE2 inhibitors may be useful in the treatment of memory deficiency, cognitive disorders, anxiety, bipolar disorder and depression.

- 15 WO2015/164508 (Dart Neuroscience, LLC) discloses substituted [1,2,4]triazolo[1,5-a]pyrimidin-yl compounds as PDE2 inhibitors.

- There is still a need for PDE2 inhibitor compounds with an advantageous balance of properties, such as for example selectivity for PDE2, good chemical stability and target engagement by occupying PDE2 and increasing cyclic nucleotides levels in relevant brain regions.

- SUMMARY It is the object of the present disclosure to provide a novel inhibitor of PDE2 that may be potentially useful in the treatment of diseases related to PDE2 enzyme activity.

- 25 Thus, in one aspect, the present disclosure is directed to compound 1



or a pharmaceutically acceptable salt or a solvate thereof.

In a particular example, the pharmaceutically acceptable salt is a hydrochloride salt, more in particular, the .2HCl salt. In another aspect, the present disclosure is directed to the hydrochloride salt of the compound of formula (1) as defined herein.

5 Illustrative of the disclosure is a pharmaceutical composition comprising a pharmaceutically acceptable carrier and compound 1 above, or a pharmaceutically acceptable salt or a solvate thereof. In another aspect, the present disclosure is directed to a pharmaceutical composition comprising a therapeutically effective amount of a compound as defined herein and a pharmaceutically acceptable carrier. An illustration
10 of the disclosure is a pharmaceutical composition made by mixing compound 1 above, or a pharmaceutically acceptable salt or a solvate thereof, and a pharmaceutically acceptable carrier. Illustrating the disclosure is a process for making a pharmaceutical composition comprising mixing compound 1 above, or a pharmaceutically acceptable salt or a solvate thereof, and a pharmaceutically acceptable carrier. In another aspect,
15 the present disclosure is directed to a process for preparing a pharmaceutical composition as defined herein, characterized in that a pharmaceutically acceptable carrier is intimately mixed with a therapeutically effective amount of a compound as defined herein.

Further illustrative of the disclosure are methods to enhance neuronal plasticity
20 comprising administering to a subject in need thereof a therapeutically effective amount of compound 1, or a pharmaceutically acceptable salt or a solvate thereof, or pharmaceutical compositions described above.

Exemplifying the disclosure are methods of treating a disorder mediated by the PDE2 enzyme, comprising administering to a subject in need thereof a therapeutically
25 effective amount of compound 1, or a pharmaceutically acceptable salt or a solvate thereof, or pharmaceutical compositions described above.

Further exemplifying the disclosure are methods of inhibiting the PDE2 enzyme, comprising administering to a subject in need thereof a therapeutically effective amount of compound 1, or a pharmaceutically acceptable salt or a solvate
30 thereof, or pharmaceutical compositions described above.

An example of the disclosure is a method of treating a disorder selected from the group consisting of neurological and psychiatric disorders, comprising administering to a subject in need thereof, a therapeutically effective

amount of compound 1, or a pharmaceutically acceptable salt or a solvate thereof, or pharmaceutical compositions described above.

5 An example of the disclosure is a method of treating a disorder selected from the group of neurological and psychiatric disorders selected from psychotic disorders and conditions; anxiety disorders; movement disorders; drug abuse; mood disorders; neurodegenerative disorders; disorders or conditions comprising as a symptom a deficiency in attention and/or cognition; stroke; and autistic disorders, comprising administering to a subject in need thereof, a therapeutically effective amount of compound 1, or a pharmaceutically acceptable salt or a solvate thereof, or a

pharmaceutically acceptable salt or a solvate thereof or pharmaceutical compositions described above.

5 An example of the disclosure is a method of treating a disorder selected from the group consisting of neurological and psychiatric disorders comprising administering to a subject in need thereof, a therapeutically effective amount of compound 1 or a pharmaceutically acceptable salt or a solvate thereof, or pharmaceutical compositions described above.

10 An example of the disclosure is a method of treating a disorder selected from the group of neurological and psychiatric disorders selected from psychotic disorders and conditions; anxiety disorders; movement disorders; drug abuse; mood disorders; neurodegenerative disorders; disorders or conditions comprising as a symptom a deficiency in attention and/or cognition; stroke; and autistic disorders, comprising administering to a subject in need thereof, a therapeutically effective amount of compound 1 or a salt or a solvate thereof, or pharmaceutical compositions described
15 above.

Also exemplifying the disclosure is compound 1 or a salt or a solvate thereof, or a pharmaceutical composition described above, for use as a medicament. In another aspect, the present disclosure is directed to a compound as defined herein, of a pharmaceutical composition as defined herein, for use as a medicament.

20 In another aspect, the present disclosure is directed to a compound as defined herein or a pharmaceutical composition as defined herein for use in treating or preventing a central nervous system disorder selected from the group of psychotic disorders and conditions; anxiety disorders; movement disorders; drug abuse; mood disorders; neurodegenerative disorders; disorders or conditions comprising as a
25 symptom a deficiency in attention and/or cognition; disorders related to memory acquisition and consolidation; stroke; and autistic disorder.

In another aspect, the present disclosure is directed to a compound as defined herein in combination with an additional pharmaceutical agent for use in the treatment or prevention of a condition as defined herein.

30 Further exemplifying the disclosure is compound 1 or a salt or a solvate thereof, or a pharmaceutical composition according to the disclosure for use in the treatment, prevention, amelioration, control or reduction of the risk of various neurological and psychiatric disorders associated with phosphodiesterase 2 dysfunction in a mammal, including a human, the treatment or prevention of which is affected or facilitated by the
35 inhibition of phosphodiesterase 2.

An example of the disclosure is compound 1 or a pharmaceutically acceptable salt or a solvate thereof according to the present disclosure or a pharmaceutical composition according to the disclosure for use in the treatment, prevention, amelioration, control or reduction of the risk of various disorders selected from
5 psychotic disorders and conditions; anxiety disorders; movement disorders; drug abuse; mood disorders; neurodegenerative disorders; disorders or conditions comprising as a symptom a deficiency in attention and/or cognition; stroke; and autistic disorder.

In another aspect, the present disclosure is directed to use of a compound as defined herein or a pharmaceutical composition as defined herein in the manufacture of
10 a medicament for the treatment of a disorder selected from the group of psychotic disorders and conditions; anxiety disorders; movement disorders; drug abuse; mood disorders; neurodegenerative disorders; disorders or conditions comprising as a symptom a deficiency in attention and/or cognition; disorders related to memory acquisition and consolidation; stroke; and autistic disorder.

15 An example of the disclosure is a method of treating a disorder selected from the group consisting of Alzheimer's disease, mild cognitive impairment, senility, dementia, dementia with Lewy bodies, Down's syndrome, dementia associated with stroke, dementia associated with Parkinson's disease and dementia associated with beta-amyloid, preferably Alzheimer's disease, comprising administering to a subject in need

thereof, a therapeutically effective amount of compound 1 or a pharmaceutically acceptable salt of a solvate thereof, or pharmaceutical compositions described above.

Another example of the disclosure is compound 1 or a pharmaceutically acceptable salt or a solvate thereof described above for use in treating: (a) Alzheimer's Disease, (b) mild cognitive impairment, (c) senility, (d) dementia, (e) dementia with Lewy bodies, (f) Down's syndrome, (g) dementia associated with stroke, (h) dementia associated with Parkinson's disease, (i) dementia associated with beta-amyloid, (j) depressive disorders and (k) anxiety disorders, in a subject in need thereof.

In another aspect, the present disclosure is directed to a product comprising (a) a compound as defined herein; and (b) an additional pharmaceutical agent, as a combined preparation for simultaneous, separate or sequential use in the treatment or prevention of a condition.

DESCRIPTION OF THE FIGURES

Figure 1 shows the effect of compound 1 on pGlu1 levels in the hippocampus of Sprague Dawley rats (10 mg/kg and 40 mg/kg Compound 1). Western blots from individual rats are shown (n = 5 per treatment) (Fig 1a). Quantification (pGlu1 normalized against total Glu1 levels) is shown in Fig 1b.

Figure 2 shows occupancy of PDE2 by compound 1.

Figure 3 shows the effect of compound 1 on basal synaptic transmission at the mossy fiber synapse.

Figure 4 shows the dose response effect of compound 1 on basal synaptic transmission at the mossy fiber synapse.

Figures 5a and 5b show the effect of compound 1 on weak HFS-induction of long term potentiation (LTP) at the mossy fiber synapse.

Figure 6 shows the effect of [CAS 1394033-54-5] 1 on basal synaptic transmission at the mossy fiber synapse.

Figure 7 shows measurement of cGMP levels in CSF in Marshall Beagle dogs.

Figure 8a shows Input–output curve for the slope of the field excitatory postsynaptic potential (fEPSP) recorded in the dentate gyrus; sample recordings showing mean population spike slope (PSA) responses at 30 min intervals; figure 8b shows LTP induced by high frequency stimulation (HFS) was enhanced by compound 1 at perforant path synapses as compared to vehicle condition; mean normalized PSA slope before and after HFS is plotted as a function of time; inset bar graphs show mean data over 30 min intervals before and after tetanization procedure; figure 8c shows lasting increases in the fEPSP slope. * p<0.05 compound 1 versus vehicle at each 30 min time intervals.

DETAILED DESCRIPTION OF THE INVENTION

DEFINITIONS

5 The term "subject" as used herein, refers to an animal, preferably a mammal, most preferably a human, who is or has been the object of treatment, observation or experiment.

10 The term "therapeutically effective amount" as used herein, means that amount of active compound or pharmaceutical agent that elicits the biological or medicinal response in a tissue system, animal or human that is being sought by a researcher, veterinarian, medical doctor or other clinician, which includes alleviation of the symptoms of the disease or disorder being treated.

As used herein, the term "composition" is intended to encompass a product comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from combinations of the specified ingredients in the specified amounts.

15 The term "host" refers to a mammal, in particular to humans, mice, dogs and rats.

The term "cell" refers to a cell expressing or incorporating the PDE2 enzyme.

20 The term "compound of the invention" as used herein, is meant to include compound 1, and the salts and solvates thereof.

As used herein, any chemical formula with bonds shown only as solid lines and not as solid wedged or hashed wedged bonds, or otherwise indicated as having a particular configuration (e.g. *R*, *S*) around one or more atoms, contemplates each possible stereoisomer, or mixture of two or more stereoisomers.

25 The absolute configuration is specified according to the Cahn-Ingold-Prelog system. The configuration at an asymmetric atom is specified by either *R* or *S*.

When a specific stereoisomer is identified, this means that said stereoisomer is substantially free, i.e. associated with less than 50%, preferably less than 20%, more preferably less than 10%, even more preferably less than 5%, in particular less than 2% and most preferably less than 1%, of the other stereoisomers.

30 In addition, the compound of the present invention may form solvates with water (i.e., hydrates) or common organic solvents, and such solvates are also intended to be encompassed within the scope of this invention.

For use in medicine, the salts of compound 1 refer to non-toxic
35 "pharmaceutically acceptable salts". Other salts may, however, be useful in the

preparation of compound 1 or of its pharmaceutically acceptable salts. Suitable pharmaceutically acceptable salts of compound 1 include acid addition salts which may, for example, be formed by mixing a solution of the compound with a solution of a pharmaceutically acceptable acid such as hydrochloric acid, sulfuric acid, fumaric acid, maleic acid, succinic acid, acetic acid, benzoic acid, citric acid, tartaric acid, carbonic acid or phosphoric acid. Representative acids which may be used in the preparation of pharmaceutically acceptable salts include, but are not limited to, the following: acetic acid, 2,2-dichloroacetic acid, acylated amino acids, adipic acid, alginic acid, ascorbic acid, L-aspartic acid, benzenesulfonic acid, benzoic acid, 4-acetamidobenzoic acid, (+)-camphoric acid, camphorsulfonic acid, capric acid, caproic acid, caprylic acid, cinnamic acid, citric acid, cyclamic acid, ethane-1,2-disulfonic acid, ethanesulfonic acid, 2-hydroxy-ethanesulfonic acid, formic acid, fumaric acid, galactaric acid, gentisic acid, glucoheptonic acid, D-gluconic acid, D-glucuronic acid, L-glutamic acid, beta-oxo-glutaric acid, glycolic acid, hippuric acid, hydrobromic acid, hydrochloric acid, (+)-L-lactic acid, ($\pm$)-DL-lactic acid, lactobionic acid, maleic acid, (-)-L-malic acid, malonic acid, ($\pm$)-DL-mandelic acid, methanesulfonic acid, naphthalene-2-sulfonic acid, naphthalene-1,5-disulfonic acid, 1-hydroxy-2-naphthoic acid, nicotinic acid, nitric acid, oleic acid, orotic acid, oxalic acid, palmitic acid, pamoic acid, phosphoric acid, L-pyroglutamic acid, salicylic acid, 4-amino-salicylic acid, sebacic acid, stearic acid, succinic acid, sulfuric acid, tannic acid, (+)-L-tartaric acid, thiocyanic acid, p-toluene-sulfonic acid, trifluoromethylsulfonic acid, and undecylenic acid.

PHARMACOLOGY

The compound according to the invention inhibits PDE2 enzyme activity, in particular PDE2A, and hence raises the levels of cAMP or cGMP within cells that express PDE2. Accordingly, inhibition of PDE2 enzyme activity may be useful in the treatment of diseases caused by deficient amounts of cAMP or cGMP in cells. PDE2 inhibitors may also be of benefit in cases in which raising the amount of cAMP or cGMP above normal levels results in a therapeutic effect. Inhibitors of PDE2 may be used to treat neurological and psychiatric disorders.

Hence, the present invention relates to compound 1 or a pharmaceutically acceptable salt or a solvate thereof according to the present invention, for use as a medicine, as well as to the use of compound 1 or a pharmaceutically acceptable salt or a solvate thereof according to the invention or a pharmaceutical composition according to the invention for the manufacture of a medicament. The present invention also relates to compound 1 or a pharmaceutically acceptable salt or a solvate thereof according to the present invention or a pharmaceutical composition according to the

invention for use in the treatment or prevention of, in particular treatment of, a condition in a mammal, including a human, the treatment or prevention of which is affected or facilitated by the inhibition of phosphodiesterase 2 enzyme. The present invention also relates to the use of compound 1 or a pharmaceutically acceptable salt or a solvate thereof according to the present invention or a pharmaceutical composition according to the invention for the manufacture of a medicament for the treatment or prevention of, in particular treatment of, a condition in a mammal, including a human, the treatment or prevention of which is affected or facilitated by the inhibition of phosphodiesterase 2 enzyme.

10 The present invention also relates to compound 1 or a pharmaceutically acceptable salt or a solvate thereof according to the invention, or a pharmaceutical composition according to the invention for use in the treatment, prevention, amelioration, control or reduction of the risk of various neurological and psychiatric disorders associated with phosphodiesterase 2 dysfunction in a mammal, including a human, the treatment or prevention of which is affected or facilitated by the inhibition of phosphodiesterase 2.

Also, the present invention relates to the use of compound 1 or a pharmaceutically acceptable salt or a solvate thereof according to the invention or a pharmaceutical composition according to the invention for the manufacture of a medicament for treating, preventing, ameliorating, controlling or reducing the risk of various neurological and psychiatric disorders associated with phosphodiesterase 2 dysfunction in a mammal, including a human, the treatment or prevention of which is affected or facilitated by the inhibition of phosphodiesterase 2.

Where the invention is said to relate to the use of compound 1 or a pharmaceutically acceptable salt or a solvate thereof or composition according to the invention for the manufacture of a medicament for e.g. the treatment of a subject, e.g. a mammal, it is understood that such use is to be interpreted in certain jurisdictions as a method of e.g. treatment of a subject, comprising administering to a subject in need of such e.g. treatment, an effective amount of compound 1 or a pharmaceutically acceptable salt or a solvate thereof or composition according to the invention.

In particular, the indications that may be treated with PDE2 inhibitors, either alone or in combination with other drugs, include, but are not limited to, those diseases thought to be mediated in part by the basal ganglia, prefrontal cortex and hippocampus.

These indications include neurological and psychiatric disorders selected from psychotic disorders and conditions; anxiety disorders; movement disorders; drug abuse; mood disorders; neurodegenerative disorders; disorders or conditions comprising as a

symptom a deficiency in attention and/or cognition; stroke; and autistic disorder or autism.

In particular, the psychotic disorders and conditions associated with PDE2 dysfunction include one or more of the following conditions or diseases: schizophrenia, for example of the paranoid, disorganized, catatonic, undifferentiated or residual type; 5 schizophreniaform disorder; schizoaffective disorder, such as delusional or depressive type; delusional disorder; substance-induced psychotic disorder such as psychosis induced by alcohol, amphetamine, cannabis, cocaine, hallucinogens, inhalants, opioids, or phencyclidine; personality disorders of the paranoid type; and personality disorder of 10 the schizoid type.

In particular, the anxiety disorders include panic disorder; agoraphobia; specific phobia; social phobia; obsessive-compulsive disorder; post-traumatic stress disorder; acute stress disorder; and generalized anxiety disorder.

In particular, movement disorders include Huntington's disease and dyskinesia; 15 Parkinson's disease; restless leg syndrome and essential tremor. Additionally, Tourette's syndrome and other tic disorders can be included.

In particular, the central nervous system disorder is a substance-related disorder selected from the group of alcohol abuse; alcohol dependence; alcohol withdrawal; alcohol withdrawal delirium; alcohol-induced psychotic disorder; amphetamine 20 dependence; amphetamine withdrawal; cocaine dependence; cocaine withdrawal; nicotine dependence; nicotine withdrawal; opioid dependence and opioid withdrawal.

In particular, mood disorders and mood episodes include depression, mania and bipolar disorders. Preferably, the mood disorder is selected from the group of bipolar disorders (I and II); cyclothymic disorder; depression; dysthymic disorder; major 25 depressive disorder; treatment-resistant depression; and substance-induced mood disorder.

In particular, neurodegenerative disorders include Parkinson's disease; Huntington's disease; dementia such as for example Alzheimer's disease; multi-infarct dementia; AIDS-related dementia or fronto temporal dementia. The neurodegenerative 30 disorder or condition comprises dysfunction of striatal medium spiny neurons responses.

In particular, disorders or conditions comprising as a symptom a deficiency in attention and/or cognition include dementia, such as Alzheimer's disease; multi-infarct dementia; dementia due to Lewy body disease; alcoholic dementia or substance- 35 induced persisting dementia; dementia associated with intracranial tumours or cerebral trauma; dementia associated with Huntington's disease; dementia associated with Parkinson's disease; AIDS-related dementia; dementia due to Pick's disease; dementia

due to Creutzfeldt-Jakob disease; other diseases include delirium; amnestic disorder; post-traumatic stress disorder; stroke; progressive supranuclear palsy; mental retardation; a learning disorder; attention-deficit/hyperactivity disorder (ADHD); mild cognitive disorder; Asperger's syndrome; age-related cognitive impairment; and

5 cognitive impairment related to perception, concentration, learning or memory.

In particular, disorders related to memory acquisition and consolidation include, memory disorders, such as age-associated memory losses, memory deficiency.

Preferably, the psychotic disorder is selected from the group of schizophrenia, delusional disorder, schizoaffective disorder, schizophreniform disorder and

10 substance-induced psychotic disorder.

Preferably, the central nervous system disorder is a personality disorder selected from the group of obsessive-compulsive personality disorder and schizoid, schizotypal disorder.

Preferably, the central nervous system disorder is a mood disorder selected from

15 the group of bipolar disorders (I & II), cyclothymic disorder, depression, dysthymic disorder, major depressive disorder; treatment-resistant depression; and substance-induced mood disorder.

Preferably, the central nervous system disorder is attention-deficit/hyperactivity disorder.

20 Preferably, the central nervous system disorder is a cognitive disorder selected from the group of delirium, substance-induced persisting delirium, dementia, dementia due to HIV disease, dementia due to Huntington's disease, dementia due to Parkinson's disease, dementia of the Alzheimer's type, substance-induced persisting dementia and mild cognitive impairment.

25 Preferably the disorders treated by the compounds of formula (I) or a pharmaceutically acceptable salt or a solvate thereof of the present invention are selected from schizophrenia; obsessive-compulsive disorder; generalized anxiety disorder; Huntington's disease; dyskinesia; Parkinson's disease; depression; bipolar disorders; dementia such as Alzheimer's disease; attention-deficit/hyperactivity

30 disorder; drug abuse; stroke; and autism.

Preferably, the disorders treated by the compounds of formula (I) or a pharmaceutically acceptable salt or a solvate thereof of the present invention are schizophrenia, including positive and negative symptoms thereof, and cognitive deficits, such as impaired attention or memory.

35 Of the disorders mentioned above, the treatment of anxiety, obsessive-compulsive disorder, post-traumatic stress disorder; generalized anxiety disorder, schizophrenia, depression, attention-deficit/hyperactivity disorder, Alzheimer's disease,

dementia due to Huntington's disease, dementia due to Parkinson's disease, dementia of the Alzheimer's type, substance-induced persisting dementia and mild cognitive impairment are of particular importance.

Of the disorders mentioned above, the treatment of anxiety, obsessive-compulsive disorder, schizophrenia, depression, attention-deficit/hyperactivity disorder, and Alzheimer's disease are of particular importance.

Other central nervous system disorders include schizoanxiety disorder, and comorbid depression and anxiety, in particular major depressive disorder with comorbid generalized anxiety disorder, social anxiety disorder, or panic disorder; it is understood that comorbid depression and anxiety may also be referred to by the terms anxious depression, mixed anxiety depression, mixed anxiety-depressive disorder, or major depressive disorder with anxiety symptoms, which are used indistinctively herein.

At present, the fourth edition of the Diagnostic & Statistical Manual of Mental Disorders (DSM-IV) of the American Psychiatric Association provides a diagnostic tool for the identification of the disorders described herein. The person skilled in the art will recognize that alternative nomenclatures, nosologies, and classification systems for neurological and psychiatric disorders described herein exist, and that these evolve with medical and scientific progresses. For example, the "American Psychiatric Association: Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition. Arlington, VA, American Psychiatric Association, 2013" (DSM-5TM) utilizes terms such as depressive disorders, in particular, major depressive disorder, persistent depressive disorder (dysthymia), substance-medication-induced depressive disorder; neurocognitive disorders (NCDs) (both major and mild), in particular, neurocognitive disorders due to Alzheimer's disease, vascular NCD (such as vascular NCD present with multiple infarctions), NCD due to HIV infection, NCD due to traumatic brain injury (TBI), NCD due to Parkinson's disease, NCD due to Huntington's disease, frontotemporal NCD, NCD due to prion disease, and substance/medication-induced NCD; neurodevelopmental disorders, in particular, intellectual disability, specific learning disorder, neurodevelopmental motor disorder, communication disorder, and attention-deficit/hyperactivity disorder (ADHD); substance-related disorders and addictive disorders, in particular, alcohol use disorder, amphetamine use disorder, cannabis use disorder, cocaine use disorder, other hallucinogen use disorder, tobacco use disorder, opioid use disorder, and phencyclidine use disorder; schizophrenia spectrum and other psychotic disorders, in particular, schizophrenia, schizophreniform disorder, schizoaffective disorder, delusional disorder, brief psychotic disorder,

substance/medication-induced psychotic disorder; and cyclothymic disorder (which under DSM-5TM falls under the bipolar and related disorders category). Such terms may be used by the skilled person as an alternative nomenclature for some of the diseases or conditions referred to herein. An additional neurodevelopmental disorder includes autism spectrum disorder (ASD), which encompasses according to the DSM-5TM, disorders previously known by the terms early infantile autism, childhood autism, Kanner's autism, high-functioning autism, atypical autism, pervasive developmental disorder not otherwise specified, childhood disintegrative disorder, and Asperger's disorder.

10 Therefore, the invention also relates to compound 1 or a pharmaceutically acceptable salt or a solvate thereof according to the invention, for use in the treatment of any one of the diseases mentioned hereinbefore.

 The invention also relates to compound 1 or a pharmaceutically acceptable salt or a solvate thereof according to the invention for use in treating any one of the diseases
15 mentioned hereinbefore.

 The invention also relates to compound 1 or a pharmaceutically acceptable salt or a solvate thereof according to the invention, for the treatment or prevention, in particular treatment, of any one of the diseases mentioned hereinbefore.

 The invention also relates to the use of compound 1 or a pharmaceutically acceptable salt or a solvate thereof according to the invention, for the manufacture of a
20 medicament for the treatment or prevention of any one of the disease conditions mentioned hereinbefore.

 The invention also relates to the use of compound 1 or a pharmaceutically acceptable salt or a solvate thereof according to the invention for the manufacture of a
25 medicament for the treatment of any one of the disease conditions mentioned hereinbefore.

 Compound 1 or a pharmaceutically acceptable salt or a solvate thereof of the present invention can be administered to mammals, preferably humans, for the treatment or prevention of any one of the diseases mentioned hereinbefore.

30 In view of the utility of compound 1, or a pharmaceutically acceptable salt or a solvate thereof, according to the invention, there is provided a method of treating a disorder or disease mentioned hereinbefore, comprising administering to a subject in need thereof, a therapeutically effective amount of compound 1 or a pharmaceutically acceptable salt or a solvate thereof or pharmaceutical compositions described herein.

35 Said methods comprise the administration, i.e. the systemic or topical administration, preferably oral administration, of a therapeutically effective amount of

compound 1 or a pharmaceutically acceptable salt or a solvate thereof according to the invention to warm-blooded animals, including humans.

Therefore, the invention also relates to a method for the prevention and/or treatment of any one of the diseases mentioned hereinbefore comprising administering
5 a therapeutically effective amount of compound 1 or a pharmaceutically acceptable salt or a solvate thereof according to the invention to a patient in need thereof.

The PDE2 inhibitor described herein can be used alone, in combination or in combination with other pharmaceutical agents such as other agents used in the treatment of psychoses, such as schizophrenia and bipolar disorder, obsessive-
10 compulsive disorder, Parkinson's disease, cognitive impairment and/or memory loss, e.g. nicotinic α -7 agonists, PDE4 inhibitors (Rolipram, GEBR-7b, GSK356278, GSK256066, Apremilast, MK-0952, Roflumilast, AN2898, AN2728, Ariflo Cilomilast, Dotraverine, Ronomilast Elbimilast, Revamilast, Tetomilast, E6005, GDP-1116, HT0712, MK-0873), PDE5 inhibitors (Sildenafil, Vardenafil, Tadalafil, Udenafil,
15 Avanafil, Mirodenafil, Lodenafil, Dasantafil, PF-00489791), PDE9 (PF-04447943), other PDE2 inhibitors (Bay 60-7550, PF-999, ND-7001), PDE10 inhibitors (PF-02545920, AMG579), PDE2 and 10 inhibitors, calcium channel blockers, muscarinic m1 and m2 modulators, adenosine receptor modulators, ampakines, NMDA-R modulators, mGluR modulators, dopamine modulators, serotonin modulators,
20 cannabinoid modulators, HDAC inhibitors (Vorinostat SAHA, Panobinostat, Quisinostat, Valproic acid) and cholinesterase inhibitors (e.g. donepezil, rivastigmine, and galantamine). In such combinations, the compound 1 or a pharmaceutically acceptable salt or a solvate thereof of the present invention may be utilized in combination with one or more other drugs in the treatment, prevention, control,
25 amelioration, or reduction of risk of diseases or conditions for which compound 1 or the other drugs may have utility, where the combination of the drugs together are safer or more effective than either drug alone.

One skilled in the art will recognize that a therapeutically effective amount of the PDE2 inhibitor of the present invention is the amount sufficient to inhibit the PDE2
30 enzyme and that this amount varies *inter alia*, depending on the type of disease, the concentration of the compound in the therapeutic formulation, and the condition of the patient. Generally, an amount of PDE2 inhibitor to be administered as a therapeutic agent for treating diseases in which inhibition of the PDE2 enzyme is beneficial, such as the disorders described herein, will be determined on a case by case by an attending
35 physician.

Generally, a suitable dose is one that results in a concentration of the PDE2 inhibitor at the treatment site in the range of 0.5 nM to 200 μ M, and more usually 5 nM

to 50 μ M. To obtain these treatment concentrations, a patient in need of treatment likely will be administered between 0.001 mg/kg to 15 mg/kg body weight, in particular from 0.01 mg/kg to 2.50 mg/kg body weight, in particular, from 0.01 to 1.5 mg/kg body weight, in particular from 0.1 mg/kg to 0.50 mg/kg body weight. The amount of a compound according to the present invention, also referred to here as the active ingredient, which is required to achieve a therapeutical effect will, of course vary on case-by-case basis, vary with the particular compound, the route of administration, the age and condition of the recipient, and the particular disorder or disease being treated. A method of treatment may also include administering the active ingredient on a regimen of between one and four intakes per day. In these methods of treatment the compound according to the invention is preferably formulated prior to admission. As described herein below, suitable pharmaceutical formulations are prepared by known procedures using well known and readily available ingredients.

15 PHARMACEUTICAL COMPOSITIONS

The present invention also provides compositions for preventing or treating diseases in which inhibition of PDE2 is beneficial, such as neurological and psychiatric disorders. Said compositions comprising a therapeutically effective amount of compound 1 and a pharmaceutically acceptable carrier or diluent.

While it is possible for the active ingredient to be administered alone, it is preferable to present it as a pharmaceutical composition. Accordingly, the present invention further provides a pharmaceutical composition comprising a compound according to the present invention, together with a pharmaceutically acceptable carrier or diluent. The carrier or diluent must be "acceptable" in the sense of being compatible with the other ingredients of the composition and not deleterious to the recipients thereof.

The pharmaceutical compositions of this invention may be prepared by any methods well known in the art of pharmacy. A therapeutically effective amount of the particular compound, in base form or addition salt form, as the active ingredient is combined in intimate admixture with a pharmaceutically acceptable carrier, which may take a wide variety of forms depending on the form of preparation desired for administration. These pharmaceutical compositions are desirably in unitary dosage form suitable, preferably, for systemic administration such as oral, percutaneous or parenteral administration; or topical administration such as via inhalation, a nose spray, eye drops or via a cream, gel, shampoo or the like. For example, in preparing the compositions in oral dosage form, any of the usual pharmaceutical media may be

employed, such as, for example, water, glycols, oils, alcohols and the like in the case of oral liquid preparations such as suspensions, syrups, elixirs and solutions: or solid carriers such as starches, sugars, kaolin, lubricants, binders, disintegrating agents and the like in the case of powders, pills, capsules and tablets. Because of their ease in administration, tablets and capsules represent the most advantageous oral dosage unit form, in which case solid pharmaceutical carriers are obviously employed. For parenteral compositions, the carrier will usually comprise sterile water, at least in large part, though other ingredients, for example, to aid solubility, may be included.

Injectable solutions, for example, may be prepared in which the carrier comprises saline solution, glucose solution or a mixture of saline and glucose solution. Injectable suspensions may also be prepared in which case appropriate liquid carriers, suspending agents and the like may be employed. In the compositions suitable for percutaneous administration, the carrier optionally comprises a penetration enhancing agent and/or a suitable wettable agent, optionally combined with suitable additives of any nature in minor proportions, which additives do not cause any significant deleterious effects on the skin. Said additives may facilitate the administration to the skin and/or may be helpful for preparing the desired compositions. These compositions may be administered in various ways, e.g., as a transdermal patch, as a spot-on or as an ointment.

It is especially advantageous to formulate the aforementioned pharmaceutical compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used in the specification and claims herein refers to physically discrete units suitable as unitary dosages, each unit containing a predetermined quantity of active ingredient calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. Examples of such dosage unit forms are tablets (including scored or coated tablets), capsules, pills, powder packets, wafers, injectable solutions or suspensions, teaspoonfuls, tablespoonfuls and the like, and segregated multiples thereof.

Depending on the mode of administration, the pharmaceutical composition will comprise from 0.05 to 99 % by weight, preferably from 0.1 to 70 % by weight, more preferably from 0.1 to 50 % by weight of the active ingredient, and, from 1 to 99.95 % by weight, preferably from 30 to 99.9 % by weight, more preferably from 50 to 99.9 % by weight of a pharmaceutically acceptable carrier, all percentages being based on the total weight of the composition.

The present compound can be used for systemic administration such as oral, percutaneous or parenteral administration; or topical administration such as via

inhalation, a nose spray, eye drops or via a cream, gel, shampoo or the like. The compound is preferably orally administered.

The exact dosage and frequency of administration depends on the compound, the particular condition being treated, the severity of the condition being treated, the age, weight, sex, extent of disorder and general physical condition of the particular patient as well as other medication the individual may be taking, as is well known to those skilled in the art. Furthermore, it is evident that said effective daily amount may be lowered or increased depending on the response of the treated subject and/or depending on the evaluation of the physician prescribing the compound of the instant invention.

The amount of compound 1 that can be combined with a carrier material to produce a single dosage form will vary depending upon the disease treated, the mammalian species, and the particular mode of administration. However, as a general guide, suitable unit doses for the compound of the present invention can, for example, preferably contain between 0.1 mg to about 1000 mg of the active compound. A preferred unit dose is between 1 mg to about 500 mg. A more preferred unit dose is between 1 mg to about 300 mg. Even more preferred unit dose is between 1 mg to about 100 mg. Such unit doses can be administered more than once a day, for example, 2, 3, 4, 5 or 6 times a day, but preferably 1 or 2 times per day, so that the total dosage for a 70 kg adult is in the range of 0.001 to about 15 mg per kg weight of subject per administration. A preferred dosage is 0.01 to about 1.5 mg per kg weight of subject per administration, and such therapy can extend for a number of weeks or months, and in some cases, years. It will be understood, however, that the specific dose level for any particular patient will depend on a variety of factors including the activity of the specific compound employed; the age, body weight, general health, sex and diet of the individual being treated; the time and route of administration; the rate of excretion; other drugs that have previously been administered; and the severity of the particular disease undergoing therapy, as is well understood by those of skill in the area.

A typical dosage can be one 1 mg to about 100 mg tablet or 1 mg to about 300 mg taken once a day, or, multiple times per day, or one time-release capsule or tablet taken once a day and containing a proportionally higher content of active ingredient. The time-release effect can be obtained by capsule materials that dissolve at different pH values, by capsules that release slowly by osmotic pressure, or by any other known means of controlled release.

It can be necessary to use dosages outside these ranges in some cases as will be apparent to those skilled in the art. Further, it is noted that the clinician or treating physician will know how and when to start, interrupt, adjust, or terminate therapy in conjunction with individual patient response.

- 5 For the compositions, methods and kits provided above, one of skill in the art will understand that the preferred compound for use in each is the compound noted herein

EXPERIMENTAL PART

- 10 As used herein, the term "ACN" means acetonitrile, "AcOH" means acetic acid, "DMAP" 4-dimethylaminopyridine, "DSC" means differential scanning calorimetry, "LCMS" means liquid chromatography/mass spectrometry, "HPLC" means high-performance liquid chromatography, "RP HPLC" means reverse phase high-performance liquid chromatography, "aq." means aqueous, "DCM" means
15 dichloromethane, "DIPE" means diisopropyl ether, "DIPEA" means diisopropylethyl amine, "DMF" means *N,N*-dimethylformamide, "EtOH" means ethanol, "Et₂O" means diethylether, "EtOAc" means ethyl acetate, "Et₃N" means triethylamine, "HBTU" means O-(benzotriazol-1-yl)-*N,N,N',N'*-tetramethyluroniumhexafluoro-phosphate, "THF" means tetrahydrofuran, "min" means minutes, "h" means hours, "MeOH"
20 means methanol, "iPrOH" means 2-propanol, "RM" means reaction mixture, "RT" means room temperature, "OL" means organic layer, "R_t" means retention time (in minutes), "quant." means quantitative, "sat." means saturated, "sol." means solution, "m.p." means melting point, "q.s." means quantum sufficit.
- 25 Thin layer chromatography (TLC) was carried out on silica gel 60 F254 plates (Merck) using reagent grade solvents. Open column chromatography was performed on silica gel, mesh 230-400 particle size and 60 Å pore size (Merck) under standard techniques. Automated flash column chromatography was performed using ready-to-connect cartridges from Merck, on irregular silica gel, particle size 15-40 µm (normal phase
30 disposable flash columns) on an SPOT or LAFLASH system from Armen Instrument.


- The absolute stereochemical configuration for some of the compounds was determined using vibrational circular dichroism (VCD). They were measured on a Bruker Equinox 55 equipped with a PMA 37, in a KBr liquid cell using CD₂Cl₂ as solvent (PEM: 1350cm⁻¹, LIA: 1mV, resolution: 4 cm⁻¹). A description on the use of VCD for the
35 determination of absolute configuration can be found in Dyatkin A.B. et. al, Chirality, 14:215-219 (2002).

Ab initio calculations: A thorough conformational search is performed at molecular mechanics level using MacroModel to do a mixed torsional/low-mode sampling with the OPLS-2005 force field. The located minima were optimized using Jaguar on the B3LYP/6-31G** level with a Poisson-Boltzmann continuum solvation model to mimic a dichloromethane solvent. All conformations within 10 kJ/mol interval were used to simulate VCD and IR spectrum. Dipole and rotational strengths were calculated at the same B3LYP/6-31G** level, using Jaguar. The calculated VCD spectra, generated after scaling the frequencies with a factor of 0.97, converting to a Lorentzian bandshape, and summing up the contribution of each conformer assuming a Boltzmann ensemble, were visually compared with the experimental spectra for assigning the correct stereochemistry.

The following examples are intended to illustrate but not to limit the scope of the present invention. Unless otherwise noted, all starting materials were obtained from commercial suppliers and used without further purification.

A. Synthesis of intermediates


Intermediate 1



Procedure a: 4-Methyl-3-pyridinecarboxylic acid hydrochloride (1:1) (40 g, 230.4 mmol) was added to a refluxing mixture of sulphuric acid (20 mL) and MeOH (400 mL). The mixture was refluxed overnight, then it was evaporated and the resulting slurry was added to a cold solution of NaHCO₃ (64 g) in water (360 mL). The product was extracted with DCM and the organic layer was dried over MgSO₄, filtered and evaporated, yielding intermediate 1 (28.70 g, 83%).

Procedure b: A metal reactor was charged with 3-bromo-4-methyl-pyridine (200 g, 0.116 mol) and a mixture of DMF/MeOH (1 L/1L). To this was added Et₃N (400 g, 0.395 mol), palladium (II) acetate (8 g, 0.036 mol) and 1,1'-bis(diphenylphosphino)ferrocene (16 g, 0.029 mol). The reactor was closed and pressurized with CO gas (3 MPa) and the reaction mixture was stirred and heated overnight at 140 °C. The RM was cooled, filtered and concentrated *in vacuo*. The residue was purified by flash column chromatography over silica gel (gradient eluent: EtOAc/Petroleum ether from 1/1 to 1/0). The product fractions were collected and the solvent was evaporated to afford the desired intermediate 1 (90 g, 51%).


Intermediate 2



Procedure a: A hydrogenation flask was charged with AcOH (500 mL) and then PtO₂ (15.02 g, 66.2 mmol) was added. Intermediate 1 (50 g, 330.8 mmol) was added and the mixture was hydrogenated at 50 °C for 7 days. The RM was filtered over dicalite® and the filtrate was evaporated to yield intermediate 2 (52 g), which was used in the next step without further purification.

Procedure b: Platinum oxide (5 g, 0.022 mol) was added to a solution of intermediate 1 (90 g, 0.595 mol) and AcOH (1 L). The r.m. was stirred and hydrogenated for 5 days at 50°C under a pressure of 3.5 kPa. The cooled RM was concentrated *in vacuo* to give intermediate 2 as the acetic acid salt (140 g, 97%, 90% purity determined by ¹H-NMR).


Intermediate 3



Procedure a: To a solution of intermediate 2 (52 g, 330.8 mmol) in DCM (869 mL), DIPEA (85.5 g, 661.5 mmol) and DMAP (4.04 g, 33.08 mmol) were added. Then di-tert-butyl dicarbonate (72.19 g, 330.8 mmol) was added to this solution in small portions and the reaction was stirred at RT for 1 h. The RM was washed with water and brine and the organic layer was dried over MgSO₄, filtered and evaporated. The product was purified by column chromatograph (silica gel, eluent: DCM, 1% MeOH in DCM, 2%, 4%). The desired fractions were evaporated, yielding intermediate 3 (64.1 g, 75%).

Procedure b: To a stirred and cooled (0 °C) solution of intermediate 2 (140 g, 0.595 mol) in DCM (1.5 L) was added sequentially di-tert-butyl dicarbonate (130 g, 0.596 mol), Et₃N (225 g, 1.74 mol) and DMAP (10 g, 0.082 mol) and stirring was continued at RT for 2 h. The reaction mixture was poured onto H₂O (500 mL) and extracted with DCM (2x 100 mL). The organic layers were separated, dried (Na₂SO₄), and the solvent was evaporated to give crude intermediate 3 (150 g, 90%, 90% purity determined by ¹H-NMR) which was used as such in the next.


Intermediate 4



Procedure a: Intermediate 3 (64.1 g, 249.1 mmol) was stirred in MeOH (500 mL) at
5 RT. NaOH (2 M, 747.3 mL) was added and the mixture was stirred for 2 h at RT. The
RM was acidified with HCl 1N and the product was extracted with Et₂O. The OL was
washed with brine and dried over MgSO₄, filtered and evaporated, yielding
intermediate 4 (59.70 g) as a white solid.

Procedure b: To a stirred solution of intermediate 3 (150 g, 90% pure, 0.524 mol) in
10 MeOH (0.9L) was added a solution of a 2M NaOH solution (1.8 mol). After 14 h at
RT, the RM was extracted with MTBE (2 x 0.8 L). The aqueous layer was acidified
with 10% citric acid and then extracted with EtOAc (4 x 1 L). The combined organic
layers were dried over Na₂SO₄, filtered and concentrated *in vacuo* to give crude
intermediate 4 (142 g, 90% purity determined by ¹H-NMR, 100%) which was used as
15 such in the next step.

Intermediate 5




Procedure a: To a solution of intermediate 4 (59.7 g, 0.25 mol) in THF (800 mL), was
20 added di-1H-imidazol-1-yl-methanone (54 g, 0.33 mol) and the mixture was stirred at
RT for 1 h. In another flask, to a suspension of N-methoxy-methanamine
hydrochloride (1:1) (32.93 g, 0.34 mol) in ACN (500 mL), was added trimethylamine
(35.75 g, 0.35 mol). Both mixtures were combined and stirred at 50 °C while
monitoring. The intermediate product crystallized out of the RM and did not react with
25 N-methoxy-methanamine to form the desired product. DCM was added until the
intermediate dissolved. The reaction was left stirring for 1 week at 80 °C. The solvents
were evaporated. The residue was dissolved in DCM and washed with water, 20%

AcOH solution and finally with a saturated NaHCO_3 solution. The organic layer was dried over MgSO_4 , filtered and evaporated. The product was purified by column chromatography (silica gel, eluent: 2% MeOH in DCM, 4%). The pure fractions were evaporated, yielding intermediate 5 (70 g, quantitative).


- 5 Procedure b: To a stirred and ice-cooled solution of intermediate 4 (140 g, 0.518 mol) in DCM (2 L) was added N,O-dimethylhydroxylamine (113 g, 1.16 mol) and Et_3N (113 g, 1.79 mol). Then HATU (235 g, 0.618 mol) was added and stirring was continued for 14 h. The solvent was evaporated and a NaHCO_3 solution (0.5 L) was added and then extracted with DCM (3 x 1 L). The combined organic layers were
- 10 separated, dried over Na_2SO_4 , filtered and concentrated *in vacuo*. The residue was purified by silica gel flash chromatography eluting with 1-10% EtOAc in petroleum ether to afford intermediate 5 (152 g, 100%).

Intermediate 6




- 15 Procedure a: Intermediate 5 (70 g, 244.4 mmol) in THF (250 mL) was charged in a flask under N_2 and cooled to -15°C . Methylmagnesium bromide (1.4 M in toluene/THF 75/25, 206 mL) was added dropwise, with the temperature not exceeded 0°C . After addition, the RM was stirred at RT for 1 h. Then the RM was poured on
- 20 ice with 20 mL AcOH. The product was extracted with Et_2O and the organic layer was washed with a 5% NaHCO_3 solution. The organic layer was dried over MgSO_4 , filtered and evaporated to give intermediate 6 (53.35 g, 90%).
- 25 Procedure b: To a stirred and cooled solution (0°C) of intermediate 5 (150 g, 0.524 mol) in THF (2 L) was added dropwise a 3M methylmagnesium bromide solution in THF (0.75 L, 2.25 mol) and stirring was continued at RT for 2 h. The reaction mixture was poured onto aqueous NH_4Cl solution and extracted with DCM. The combined organic layers were dried over Na_2SO_4 , filtered and concentrated *in vacuo*. The residue was purified by silica gel chromatography eluting with 1-5% EtOAc in petroleum ether to afford intermediate 6 (120 g, 95%).

Intermediate 7



Intermediate 6 (53.35 g, 0.22 mol) was stirred in toluene (1500 mL) at 0 °C under N₂. Potassium tert-butoxide (34.14 g) was added at 0-5 °C, 2,2-difluoro-acetic acid ethyl ester (33.01 g, 0.27 mol) was added dropwise at 0-5 °C. The RM was stirred at RT for 2 h, then washed with 10% H₂SO₄ in water and the organic layer was dried on MgSO₄, filtered and evaporated, yielding intermediate 7 (70.50 g, quantitative).


Intermediate 8




Intermediate 7 (70.5 g, 220.8 mmol), 1*H*-1,2,4-triazol-5-amine hydrochloride (1:1) (53.22 g, 441.52 mmol) and DMF (1500 mL) were stirred at 80 °C for 24 h. Et₃N (20 g) and di-tert-butyl dicarbonate (20 g) were added. The mixture was stirred for 30 min, evaporated and then dissolved in EtOAc, washed with water and brine. The organic layer was dried over MgSO₄, filtered and evaporated. Four isomers were observed. The first fraction crystallized from Et₂O. The crystals were filtered off and dried, yielding intermediate 8 (24.60 g, 30%). The mother liquor yielded a second fraction of the compound. The crystals were filtered off and dried, yielding intermediate 8 (2.53 g, 3%).

N.B. "RS" means the intermediate is a racemic mixture of two enantiomers of *trans* relative configuration.


Intermediates 9, 9a and 9b



Intermediate 9



Intermediate 9a




Intermediate 9b

- To a solution of intermediate 8 (24.6 g, 67 mmol) in MeOH (350 mL), was added HCl-iPrOH (350 mL) and the RM was stirred for 2 h at RT. The RM was evaporated and the product was crystallized from EtOH. The crystals were filtered off and dried,
- 5 yielding 20.33 g of a crude, to which water, Na₂CO₃ and DCM were added. The organic layer was dried over MgSO₄, filtered and evaporated, yielding 12.80 g of intermediate 9. This free base was separated into enantiomers 9a and 9b by purification by Prep SFC (Stationary phase: Chiralpak Diacel AD 30 x 250 mm; mobile phase: CO₂, ((MeOH - iPrOH 50/50) with 0.4% iPrNH₂), yielding intermediate 9a (5g, 19%,
- 10 R_t = 7.57 min) and intermediate 9b (5.13 g, 19%, R_t = 9.36 min).

- Intermediates 9a and 9b were isolated as free bases or alternatively, they were dissolved in MeOH, followed by addition of HCl/i-PrOH and the mixture evaporated. The hydrochloride salts (in each instance, .HCl) were crystallized from ACN, filtered
- 15 off and dried.

B-Synthesis of Final Compound

Compound 1



- 20 2,6-Dimethylpyridine-4-carboxylic acid (1.84 g, 12.2 mmol) was stirred in DCM (100 mL), DIPEA (6.31 g, 48.8 mmol) and HBTU (4.63 g, 12.2 mmol) were added, stirring was continued for 0.5 h at RT. Intermediate 9b (3.26 g, 12.2 mmol) was added to the solution and stirring was continued for 5 h at RT. NaOH 1N solution was added and stirred for 5 min. The organic layer was separated, dried over MgSO₄, filtered and

evaporated. The product was purified by column chromatography (silica gel, eluent: 1% MeOH in DCM, 2%, 4%). The pure fractions were evaporated and the product was crystallized from DIPE, filtered off and dried, yielding compound 1 (3.85 g, 79%).

- 5 A separate batch of the compound was crystallized as HCl salt from Et₂O to yield compound 1 as the hydrochloride salt (.2 HCl) (yield: 175 mg, 70%, starting from 175 mg of intermediate 9b.HCl) .

The stereoconfiguration of compound 1 was confirmed by vibrational circular dichroism (VCD).

10 ANALYTICAL PART

Melting Points

Values are either peak values or melt ranges, and are obtained with experimental uncertainties that are commonly associated with this analytical method.

DSC823e (indicated as DSC)

- 15 The melting point was determined with a DSC823e (Mettler-Toledo). The melting point was measured with a temperature gradient of 10°C/min. Maximum temperature was 300°C.

Table 1

Co. No.	MP
1	149.31

20 Optical Rotation

Optical rotation was measured on a Perkin-Elmer 341 polarimeter with a sodium lamp and reported as follows: $[\alpha]^{\circ}(\lambda, c \text{ g/100ml, solvent, } T^{\circ}\text{C})$.

$[\alpha]_{\lambda}^T = (100\alpha) / (l \times c)$: where l is the path length in dm and c is the concentration in g/100 ml for a sample at a temperature T (°C) and a wavelength λ (in nm). If the

- 25 wavelength of light used is 589 nm (the sodium D line), then the symbol D might be used instead. The sign of the rotation (+ or -) should always be given. When using this equation the concentration and solvent are always provided in parentheses after the rotation. The rotation is reported using degrees and no units of concentration are given (it is assumed to be g/100 ml).

30 Table 2

Co. No.	OR
1	+28.91 ° (589 nm, c 0.2975 w/v %, DMF, 20 °C)

SFC-MS methods

The SFC measurement was performed using an Analytical Supercritical fluid chromatography (SFC) system composed by a binary pump for delivering carbon dioxide (CO₂) and modifier, an autosampler, a column oven, a diode array detector equipped with a high-pressure flow cell standing up to 400 bars. If configured with a Mass Spectrometer (MS) the flow from the column was brought to the (MS). It is within the knowledge of the skilled person to set the tune parameters (e.g. scanning range, dwell time...) in order to obtain ions allowing the identification of the compound's nominal monoisotopic molecular weight (MW). Data acquisition was performed with appropriate software.

Table 3a. Analytical SFC-MS Methods (Flow expressed in mL/min; column temperature (T) in °C; Run time in minutes, Backpressure (BPR) in bars.

Method code	column	mobile phase	gradient	Flow	Run time
				----- Col T	----- BPR
1	Daicel (AD, OD, OJ, AS, ID)-H -H column (5.0 µm, 250 x 4.6 mm)	A: CO ₂ B: 5 different solvent for B used: MeOH, EtOH, iPrOH, MeOH-iPrOH (50-50) and EtOH-iPrOH (50-50)	10%-55% B in 4 min, 55-50% in 0.45 min hold 2.55 min	5 ----- 40	7 ----- 110

Table 3b. Analytical SFC data – R_t means retention time (in minutes), [M+H]⁺ means the protonated mass of the compound, method refers to the method used for (SFC)MS analysis of enantiomerically pure compounds.

Co. No.	R _t	[M+H] ⁺	Method	Isomer Elution Order
1	2.93	401	1	Only one enantiomer

LC/MS methods

The High Performance Liquid Chromatography (HPLC) measurement was performed using a LC pump, a diode-array (DAD) or a UV detector and a column as specified in the respective methods. If necessary, additional detectors were included (see table of methods below).

Flow from the column was brought to the Mass Spectrometer (MS) which was configured with an atmospheric pressure ion source. It is within the knowledge of the

skilled person to set the tune parameters (e.g. scanning range, dwell time...) in order to obtain ions allowing the identification of the compound's nominal monoisotopic molecular weight (MW). Data acquisition was performed with appropriate software.

Compounds are described by their experimental retention times (R_t) and ions. If not specified differently in the table of data, the reported molecular ion corresponds to the $[M+H]^+$ (protonated molecule) and/or $[M-H]^-$ (deprotonated molecule). In case the compound was not directly ionizable the type of adduct is specified (i.e. $[M+NH_4]^+$, $[M+HCOO]^-$, etc...). For molecules with multiple isotopic patterns (Br, Cl), the reported value is the one obtained for the lowest isotope mass. All results were obtained with experimental uncertainties that are commonly associated with the method used. Hereinafter, "SQD" means Single Quadrupole Detector, "MSD" Mass Selective Detector, "RT" room temperature, "BEH" bridged ethylsiloxane/silica hybrid, "DAD" Diode Array Detector, "HSS" High Strength silica.

Table 4a. LCMS Method codes (Flow expressed in mL/min; column temperature (T) in °C; Run time in minutes)

Method code	Instrument	Column	Mobile phase	Gradient	Flow ----- Col T	Run time (min)
Method A	Waters: Acquity® UPLC® - DAD-SQD	Waters: BEH C18 (1.7µm, 2.1*50mm)	A: 10mM CH ₃ COONH ₄ in 95% H ₂ O + 5% CH ₃ CN B: CH ₃ CN	From 95% A to 5% A in 1.3 min, held for 0.7 min.	0.8 mL/min ----- 55°C	2

Table 4b. Analytical LCMS data – R_t means retention time (in minutes), $[M+H]^+$ means the protonated mass of the compound, method refers to the method used for (LC)MS analysis.

Co. No.	R_t	$[M+H]^+$	$[M+H]^+$	Method
1	0.76	401.2	399.2	Method A

Nuclear Magnetic Resonance (NMR)

The ^1H NMR spectrum was recorded either on Bruker DPX-400 spectrometer with standard pulse sequences, operating at 400 MHz. Chemical shifts (δ) are reported in parts per million.

Co. No. 1: ^1H NMR (400 MHz, $\text{DMSO}-d_6$, 120°C) δ ppm 0.81 (d, $J=6.6$ Hz, 3 H) 1.43 (qd, $J=12.4$, 4.4 Hz, 1 H) 1.89 (br dq, $J=13.4$, 3.1 Hz, 1 H) 2.44 (s, 6 H) 2.49 - 2.53 (m, 1 H) 3.11 (t, $J=12.7$ Hz, 1 H) 3.35 (dd, $J=12.9$, 11.1 Hz, 1 H) 3.57 (td, $J=10.8$, 4.1 Hz, 1 H) 4.00 - 4.27 (m, 2 H) 6.98 (t, $J=54.2$ Hz, 1 H) 7.00 (s, 2 H) 7.53 (s, 1 H) 8.68 (s, 1 H)

PHARMACOLOGICAL EXAMPLES

The compound provided in the present invention is an inhibitor of PDE2, particularly of PDE2A. The results of testing compound 1 in several pharmacological assays is shown below.

IN VITRO ASSAY PDE2A

Human recombinant PDE2A (hPDE2A) was expressed in Sf9 cells using a recombinant rPDE10A baculovirus construct. Cells were harvested after 48 h of infection and the hPDE2A protein was purified by metal chelate chromatography on Ni-sepharose 6FF . Tested compounds were dissolved and diluted in 100% DMSO to a concentration 100 fold of the final concentration in the assay. Compound dilutions (0.4 μl) were added in 384 well plates to 20 μl of incubation buffer (50 mM Tris pH 7.8, 8.3 mM MgCl_2 , 1.7 mM EGTA). 10 μl of hPDE2A enzyme in incubation buffer was added and the reaction was started by addition of 10 μl substrate to a final concentration of 10 μM cGMP and 0.01 μCi ^3H -cGMP. The reaction was incubated for 45 minutes at room temperature. After incubation, the reaction was stopped with 20 μl of stop solution consisting of 17.8 mg/ml PDE SPA scintillation proximity assay) beads supplemented with 200 mM ZnCl_2 . After sedimentation of the beads during 30 minutes the radioactivity was measured in a Perkin Elmer Topcount scintillation counter and results were expressed as cpm. For blanc values the enzyme was omitted from the reaction and replaced by incubation buffer. Control values were obtained by addition of a final concentration of 1% DMSO instead of compound. A best fit curve is fitted by a minimum sum of squares method to the plot of % of control value subtracted with blanc value versus compound concentration and the half maximal inhibitory concentration (IC_{50}) value is derived from this curve.

IN VITRO ASSAY PDE3A

Human recombinant PDE3A (hPDE3A) was supplied as a partially purified insect cell lysate by Scottish Biomedical, it was cloned from human brain and expressed in Sf9 cells. Tested compounds were dissolved and diluted in 100% DMSO to a concentration

100 fold of the final concentration in the assay. Compound dilutions (0.4 μ l) were added in 384 well plates to 20 μ l of incubation buffer (50 mM Tris pH 7.8, 8.3 mM MgCl_2 , 1.7 mM EGTA). 10 μ l of hPDE3A enzyme in incubation buffer was added and the reaction was started by addition of 10 μ l substrate to a final concentration of

5 0.4 μ M cAMP and 2.4 μ Ci/ml [^3H]-cAMP. The reaction was incubated for 60 min at room temperature. After incubation, the reaction was stopped with 20 μ l of stop solution consisting of 17.8 mg/ml PDE SPA (scintillation proximity assay) beads supplemented with 200 mM ZnCl_2 . After sedimentation of the beads during 30 min the radioactivity was measured in a Perkin Elmer Topcount scintillation counter and results

10 were expressed as cpm. For blanc values the enzyme was omitted from the reaction and replaced by incubation buffer. Control values were obtained by addition of a final concentration of 1% DMSO instead of compound. A best fit curve is fitted by a minimum sum of squares method to the plot of % of control value subtracted with blanc value versus compound concentration and the half maximal inhibitory

15 concentration (IC_{50}) value is derived from this curve.

IN VITRO ASSAY PDE10A

Rat recombinant PDE10A (rPDE10A2) was expressed in Sf9 cells using a recombinant rPDE10A baculovirus construct. Cells were harvested after 48 h of infection and the

20 rPDE10A protein was purified by metal chelate chromatography on Ni-sepharose 6FF. Tested compounds were dissolved and diluted in 100% DMSO to a concentration 100 fold of the final concentration in the assay. Compound dilutions (0.4 μ l) were added in 384 well plates to 20 μ l of incubation buffer (50 mM Tris pH 7.8, 8.3 mM MgCl_2 , 1.7 mM EGTA). 10 μ l of rPDE10A enzyme in incubation buffer was added and the

25 reaction was started by addition of 10 μ l substrate to a final concentration of 60 nM cAMP and 0.008 μ Ci ^3H -cAMP. The reaction was incubated for 60 minutes at room temperature. After incubation, the reaction was stopped with 20 μ l of stop solution consisting of 17.8 mg/ml PDE SPA (scintillation proximity assay) beads. After sedimentation of the beads during 30 minutes the radioactivity was measured in a

30 Perkin Elmer Topcount scintillation counter and results were expressed as cpm. For blanc values the enzyme was omitted from the reaction and replaced by incubation buffer. Control values were obtained by addition of a final concentration of 1% DMSO instead of compound. A best fit curve is fitted by a minimum sum of squares method to the plot of % of control value subtracted with blanc value versus compound

35 concentration and the half maximal inhibitory concentration (IC_{50}) value is derived from this curve.

Table 5a.

Compound	IC ₅₀ PDE2A (nM)
1	0.95
1 . 2HCl	0.7

Table 5b. pIC₅₀ corresponds to the -log IC₅₀ expressed in mol/L.

Compound	pIC ₅₀ (PDE2A)	pIC ₅₀ PDE3B	pIC ₅₀ PDE10A2
1	9.07	5.21	7.06
1 . 2HCl	9.11	5.15	6.93

5

WESTERN BLOT DETECTION OF GLUR1 PHOSPHORYLATION

PDE2 is mainly expressed in hippocampus, cortex and striatum and can hydrolyze cAMP and cGMP. AMPA-R trafficking can be regulated through activation of PKA (via cAMP) or cGKII (via cGMP). Phosphorylation of the Glu1 subunit of the AMPA-R has been shown to be critical for LTD (decrease) and LTP (increase) expression and the retention of memories.

10

Methods

Compound 1 (solved in 10%CD + 1 HCl) was administered p.o.(orally) to Sprague Dawley rats (180-200g; 10 and 40 mg/kg) and 2 hours later, animals were sacrificed by decapitation. Hippocampus was dissected and tissue was snap frozen and stored at -80 °C.

15

After thawing, tissue lysis was performed in Tissue Extraction Reagent supplemented with 5 mM EDTA and a protease and phosphatase inhibitor cocktail. Protein samples were denatured by LDS sample buffer and reducing agent (Life Technologies, Invitrogen, Carlsbad, CA, USA) and finally, 50 µg of protein was loaded and electrophoresed using a 10% Bis-Tris polyacrylamide gel (Bio-Rad, Hercules, CA, USA) at 90-160V. The proteins on the gels were then electroblotted onto a Trans blot turbo 0.2 µm nitrocellulose membrane (Bio-Rad), by using a Trans-blot Turbo transfer system (Bio-Rad). Membranes were blocked for 1 h at RT in Tween-20 Trisbuffered Saline (TBS-T: 10 mM Tris-HCl pH 8.0, 150 mM NaCl, 0.05% Tween-20) containing 5% non-fat dry milk (Santa Cruz Biotechnology, Dallas, TX, USA), and incubated with the primary antibody overnight at 4 °C allowing gentle shaking (Total Glu1 Abcam 31232, Ser845 pGlu1 Abcam 76321, both dilution 1/1000). Blots were washed five times with TBS-T buffer and incubated with the secondary antibody for 1 h at RT (secondary donkey anti-rabbit HRP-conjugated, dilution 1/1000). Immunostaining was revealed after washing with TBST buffer via SuperSignal West Femto Maximum Sensitivity Substrate (Thermo scientific, Cramlington, United Kingdom). Signals were

20

25

30

captured and quantified by chemiluminescence (G-box Syngene, Syngene, Cambridge, United Kingdom).

The results of this test are shown in Figure 1.

5 PDE2 OCCUPANCY BY COMPOUND 1

Methods

Occupancy of PDE2A was evaluated by ex-vivo autoradiography using [³H]B-17a (described in WO2013/000924) as radioligand (compound 12 in Buijnsters et al., (2014). Structure-Based Design of a Potent, Selective, and Brain Penetrating PDE2

10 Inhibitor with Demonstrated Target Engagement. ACS Med Chem Lett. 5(9):1049-53.)

Male Wistar rats (200-250 g) were treated by oral administration of vehicle or increasing doses of [³H]B-17a and killed one hour after. Brains were immediately removed from the skull and rapidly frozen in dry-ice cooled 2-methylbutane (-40°C).

15 Twenty µm-thick striatal sections were cut using a Leica CM 3050 cryostat-microtome (van Hopplynus, Belgium), thaw-mounted on microscope slides (SuperFrost Plus Slides, LaboNord, France) and stored at -20°C until use.

After thawing, sections were dried under a cold stream of air and incubated for one minute with 30 nM [³H]B-17a in Tris-HCl (50 mM, pH7.4) containing 0.3% BSA.

Brain sections from drug-treated and vehicle-treated animals were incubated in parallel.

20 Non-specific binding was measured on cerebellar sections, a brain area which does not contain the PDE2A enzyme. After incubation, the excess of [³H]B-17a was washed off in ice-cold buffer 2 times 10 minutes, followed by a quick dip in distilled water. The sections were then dried under a stream of cold air.

Brain sections were loaded in a β-imager (Biospace, Paris) for 4 hours and

25 radioactivity emerging from delineated brain area was quantified using the Beta vision program (Biospace, Paris). Specific binding was determined as the difference between total binding in the striatum and non-specific binding in the cerebellum. Percentage receptor occupancy of the drug administered to the animal corresponded to 100% minus the percentage receptor labeled in the treated animal. For the determination of

30 ED₅₀-values, the percentage of receptor occupancy was plotted against dose and the sigmoidal log dose-effect curve of best fit was calculated by non-linear regression analysis, using the GraphPad Prism program. ED₅₀s (the drug dose producing 50% receptor occupancy) with 95% confidence limits were calculated from the dose-response curves.

35 The results of this test are shown in Figure 2.

EFFECT OF COMPOUND 1 ON SYNAPTIC TRANSMISSION

Critical Reagents

Sucrose dissection buffer contained (in mM) sucrose (150), NaCl (40), KCl (4), NaH₂PO₄·H₂O (0.3), MgCl₂·6H₂O (7), NaHCO₃ (26), CaCl₂·2H₂O (0.5), D-glucose (10),
5 equilibrated with 95% O₂ and 5% CO₂ gas mixture. Artificial cerebrospinal fluid (ACSF) used during equilibration and recording contained (in mM): NaCl (124), KCl (2.7), NaH₂PO₄·H₂O (1.25), MgSO₄·7H₂O (1.3), NaHCO₃ (26), CaCl₂·2H₂O (2), D-glucose (10), Ascorbic acid (2), equilibrated with 95% O₂ and 5% CO₂ gas mixture. CNQX and Kynurenic acid were prepared in ACSF at a 50 µM and 1mM concentration
10 respectively. Compound 1 was prepared fresh from stock solution (with DMSO) in ACSF and with a final DMSO concentration that did not exceed 0.1%. All reagents were from Sigma-Aldrich, unless otherwise indicated.

Animals (Species, Weight, and Gender)

Animals used were male Sprague-Dawley rats with a weight range between 145 and
15 200g provided by Charles River Germany.

Preparation of hippocampal slices

Horizontal brain slices (300 µm) were obtained from the mid- to ventral hippocampus of male Sprague-Dawley rats anesthetized with isofluorane according to standard protocol. Slices were cut using a vibrating tissue slicer (Leica VT1200S) in cold (4°C)
20 sucrose dissection buffer at a speed of 0.1 mm/s. After cut, slices were placed for equilibration at 35°C for 20 min and then allowed to recover at RT for at least one hour in artificial cerebrospinal fluid (ACSF). Three to four slices were prepared from one brain.

Test System

All data were recorded with a MEA set-up commercially available from MultiChannel Systems MCS GmbH (Reutlingen, Germany) composed of a 4-channel stimulus generator and a 60-channels amplifier head-stage connected to a 60-channels A/D card. Software for stimulation, recordings and analysis are the ones commercially available from Multi Channel Systems: MC Stim (II 2.0.0 release) and MC Rack (3.8.1.0
30 release), respectively. All of the experiments were carried out with 3-dimensional MEA (Ayanda Biosystems, S.A., CH-1015 Lausanne, Switzerland) that consist of 60 tip-shaped and 60-µm-high electrodes spaced by 100 µm. The MEA electrodes are made of platinum with 600 kΩ < impedance < 900 kΩ.

Experimental Design

35 The effect of compound 1 on synaptic transmission was investigated by recording the extracellular field potentials in hippocampal slices. It is well established that synaptic transmission can generate a deflection of the extracellular field potential that reflects

the synchronized synaptic activity in the population of neurons surrounding the recording electrode.

- Extracellular field potential recordings.* After recovery, brain slices were mounted on MEA chip under microscope and locating the 60 recording electrodes on the mossy fiber synapse (Dentate Gyrus - CA3) region of the hippocampus. ACSF solutions were continuously perfused at a rate of 2 mL/min. The temperature of the MEA chamber was maintained at 32 ± 0.1 °C with a Peltier element located in the MEA amplifier headstage. All data were recorded with a MEA set-up commercially available from MultiChannel Systems MCS GmbH (Reutlingen, Germany). Two adjacent electrodes of the chip were selected to stimulate the mossy fibres in the hilar region of the dentate gyrus and the fEPSP was recorded the terminal zone area of the CA3 region of the hippocampus. Field extracellular post-synaptic potentials (fEPSPs) were evoked by stimulation of the mossy fibre input with two consecutive electrical pulses separated by 30 ms and repeated every 60 s (pulse width 100 μ s, and current stimulation strength (μ A) 40% relative maximum amplitude). Control experiments were performed simultaneously from slices that were randomly assigned to be treated with vehicle (DMSO). N represents the number of slices and usually 3-4 slices were used per animal. Evoked-responses at post-synaptic neurons level (fEPSP) are recorded if they satisfy certain quality criteria including: correct location, stable baseline (fluctuation within ± 10 % during ten consecutive minutes, amplitude >100 μ V. The fEPSP from selected electrodes were sampled at 5 kHz and recorded on the hard disk of a PC for offline analysis. In parallel, fEPSP amplitudes of selected electrodes were compiled online (with MC Rack program) to monitor and to follow the quality of the experiment. Data are plotted in a spreadsheet file for off-line analysis.
- Weak Long Term Potentiation (LTP) was evoked by a single high frequency stimulus (HFS) to produce a less than maximal potentiation of the fEPSP.
- The results of this test are shown in Figures 3 and 4 for the effect of compound 1 on basal synaptic transmission and in Figure 5 for the effect of compound 1 on the facilitation on induction of LTP with a weak Long Term Potentiation protocol.
- Interestingly, similar results were obtained with other PDE2 inhibitors such as 4-(1-azetidiny)-7-methyl-5-[1-methyl-5-[5-(trifluoromethyl)-2-pyridinyl]-1H-pyrazol-4-yl]-imidazo[5,1-f][1,2,4]triazine [CAS 1394033-54-5] (WO2012114222, Pfizer) (see Figure 6)

SINGLE DOSE PK/PD PDE2i DOG STUDY

For these studies male and female Marshall Beagle dogs (1-6y) were used: 2 males and 2 females per treatment group. Cerebrospinal fluid (CSF) was sampled from the lateral ventricle via a needle guide cannula in instrumented conscious animals.

- 5 Baseline CSF and blood samples were taken 2 to 5 days before dosing. The dogs are fasted overnight and the next morning dosed on an empty stomach (orally by gavage). At predetermined time points after dosing blood and/or CSF was collected for the measurement of compound levels and cGMP. Analysis of cGMP was done by LC-MS/MS: 25µl CSF was diluted with 125µl artificial CSF (STIL (20 ng/ml)),
10 centrifugated and 25µl was injected. The systems used were: a Shimadzu SIL-30 UPLC-system (Hypersil (50 mmx 1 mm (3 µm)) column, basic (10mM ammonium carbonate) aqueous-acetonitrile gradient (5% to 98% in 5.5 minutes) at a flow-rate of 250 µl/min) and an API Sciex 5500 system equipped with an ESI source (selective MRM transition (m/z 346.1->152.1 (75 msec dwelltime)). The results of this study are
15 summarized in Figure 7. After a single dose of compound 1 the following observations were made: slight to moderate tremors in 3/8 animals at 0.5 mg/kg; sedation and/or tremors in 6 out of 7 animals at 1 mg/kg (one animal was not dosed due to limited stock of the compound). Plasma pharmacokinetics showed a non-dose linearity. There was a dose related increase in cGMP in the CSF. Limited individual data in the vehicle group
20 of PDE2 H-2 (n=2 at 1, 4 and 8 h) due to analytical errors.

PDE2 INHIBITION ENHANCED SYNAPTIC PLASTICITY IN ANESTHETIZED RATS: CASE PILOT STUDY WITH COMPOUND 1

Introduction

- 25 Synaptic plasticity is a fundamental mechanism to many neurobiological functions. Long-term potentiation (LTP), a form of a long-lasting highly localized increase in synaptic strength in the hippocampus as well as in the cortex, is a synaptic substrate for memory and learning (Cooke and Bliss, Curr Opin Investig Drugs. 2005;6(1):
25-34). The increase and decrease of synaptic strength depends on the activity of
30 presynaptic and postsynaptic neurons, on how networks in the brain operate in setting up sensory representation of multiple items in the memory and producing appropriate motor response. Different features of these synaptic modifications, in intact brain, are crucial to the operation of different types of network and operations of several different brain circuit systems. Therefore, LTP is expected to be compromised in in aging
35 psychiatric and neurodegenerative disorders such as Alzheimer's disease (Bergado and Almaguer, Neural Plast. 2002;9(4):217-32; Rowan et al., Biochem Soc Trans. 2005

;33: 563-7). In animals, the procedure carried out under anesthesia in intact highly interconnected brain regions, provides a powerful tool to investigate lasting changes in effective connectivity and plasticity in hippocampal-cortex circuits following a tetanic electrical stimulation with low and high frequency delivered in single pulse or paired pulse (Albensi et al., *Exp Neurol.* 2007; 204:1-13). The studies help expand understanding of the neural circuits underlying development of impaired synaptic strength i.e. determine the direct- circuit path and the role of specific biological target harbored by a specific inter-regional network connections in mediating synaptic weakening. The procedure allows testing pharmacological agents aimed to restore the pathological forms of neuroplasticity e.g. reverse deficits in LTP and network connectivity by increasing synaptic efficacy, which is expected to have beneficial effects on related cognitive and learning ability (Cooke and Bliss, 2005; Albensi et al., 2007).

Phosphodiesterases (PDEs) are a class of enzymes responsible for metabolic inactivation of secondary messengers 3',5'-cyclic adenosine monophosphate (cAMP) and 3',5'-cyclic guanosine monophosphate (cGMP) (Francis et al. *Physiol Rev.* 2011, 9: 651-90). Up to 11 families of PDEs were categorized based on their structural, enzymatic and distribution (Omori and Kotera *Circ Res.* 2007;100:309-27). The role of PDEs in the augmentation of cyclic nucleotide signaling makes these enzymes attractive targets for regulating excitability and enhancing the effects of neuronal communication. In the brain, PDE2 is mainly expressed in cortex, hippocampus and striatum where it controls the hydrolysis of cAMP. Over the last few years, research groups has focused on the development of PDE2 inhibitors as a way to modify intracellular second messengers, cGMP and cAMP to exert action on plasticity and cognitive processes (Duinen et al., *Curr Pharm Des.* 2015;21:3813-28; Gomez and Breitenbucher, *Bioorg Med Chem Lett.* 2013;23: 6522-7; Xu et al., *Neurobiol Aging.* 2015; 36:955-70; Barco et al., *Expert Opin Ther Targets* 2003; 7: 101–114). In the present study, it was investigated whether PDE2 inhibition, using compound 1, leads to alterations in excitability or in ability to express synaptic potentiation in the dentate gyrus of anesthetized adult Sprague Dawley.

Material and Methods

Animals

The present experiments were conducted in strict accordance with the guidelines of the Association for Assessment and Accreditation of Laboratory Animal Care International (AAALAC), and with the European Communities Council Directive of 24th November 1986 (86/609/EEC) and were approved by local ethical committee.

Sprague Dawley rats weighing 170-200 g at the time of surgery) were group-housed in ventilated cages located on a 12-h light/ dark cycle (lights on at 07:00 AM) after their arrival to animal facilities maintained under controlled environmental conditions.

Surgery and electrophysiology

- 5 Rats were anesthetized with an intra-peritoneal injection of urethane 1.5 g/kg body weight. Animals were placed in a stereotactic frame for the insertion of electrodes and their body temperature was constantly monitored through a rectal probe and maintained at 37 °C with a heating pad. Supplementary administration of urethane (0.2–0.5 g/kg) was given when necessary to ensure full anesthesia. Two small holes (1 mm diameter)
- 10 were drilled in the skull at the position of left hippocampus structures for stimulating and recording electrodes. A bipolar stimulating electrode; a pair of twisted stainless steel polyimide-coated wires with tips horizontally separated 0.125 µm apart (MS303/13-B. PlasticsOne), were positioned at the medial perforant pathway (mPP) (AP -7.5, ML -3.8, DV -2.5), and a stainless coated recording electrode (MS303T-2-
- 15 AIU, 0.008-0.005) are positioned at the Dentate Gyrus (DG) area of the dorsal hippocampus (AP -2.8, ML -3.8, DV -3.8). The dura was pierced through both holes, and the stimulating and recording electrodes were lowered very slowly (0.2 mm/min) through the cortex and upper layers of the hippocampus into the mPP and the DG of the dorsal hippocampus. During surgery all efforts were made to minimize animal
- 20 suffering.

- The field excitatory postsynaptic potential (fEPSP) slope is used as a measure of excitatory synaptic transmission. Single monophasic square 0.1 or 0.2 ms wave pulses generated by a constant current unit (MC, Germany) were applied for instance to the mPP and evoked responses are generated in the DG. Extracellular field potentials are
- 25 amplified; band pass filtered between 1 Hz and 2 kHz, digitized and analyzed using custom made software. The electrodes were lowered until a negative deflecting fEPSP with the maximum response is observed. A minimum of 30 min is allowed to ensure stabilization excitability before measurements. Next, monophasic constant current pulses with stimulus intensities ranging from 50 to 500 µA were delivered to generate
 - 30 Input/Output (I/O) curves and determine the maximum PSA and fEPSP slopes, and then stimulus intensity that produced 50% of the maximum response (i.e., test pulse) was used in subsequent experiments.

- LTP induction: Test stimulation was then applied every 5 min before and after tetanic stimulation. Responses were evoked by high frequency stimulation (10 stimulus trains
- 35 of 20 square wave pulses, 0.2 ms duration at 200 Hz, 5 ms inter-stimulus interval, with 2 seconds inter-train interval). Five evoked responses were averaged for each time point measured during the experiments, half hour of baseline recording, immediately

before drug application or tetanic stimulation (control for LTP induction). The magnitude of synaptic potentiation is expressed as the percentage of increase in the amplitude DG population spike (PSA), as well as fEPSP slope at the time interval after tetanic stimulation relative to the slopes averaged over the stable 30-min of the pharmacological period.

Responses at the selected pulse intensity were collected and averaged up to 130 min post-tetanization. The amplitude of the population spike was defined as the average of the amplitude from the first positive peak (a) to the first negative peak (b) and the amplitude from the negative peak (b) to the second positive peak (c): $[(a - b)/(c - b)]/2$. For quantification of the slope of fEPSPs, only very early component of the waveform ($\Delta V/\Delta t$) was measured to avoid contamination by the population spike.

Histology

At the end of the electrophysiological study, electrical stimulation of 500 μ A for 20 sec was delivered to produce a lesion at the end tip of the stimulation and recording electrodes and brains were harvested for histological verification of electrodes placement. Brain sections (20 μ m) were examined using a light microscope. Animals with incorrect electrode placement were excluded from the study.

Drug

Compound 1 was dissolved in 10% Cyclodextrine (CD) + 1HCl + NaCl for subcutaneous (SC) administration.

Statistic

For each animal, the stable baseline (pre-tetanus) responses over 30 min were averaged and the mean was normalized as being 100%, and the post-tetanus response data were expressed relative by the baseline average. Comparison of the effects of vehicle and compound 1 after tetanus was performed on 30 min intervals using one-way repeated-measures analysis of variance (ANOVA) on ranks followed by Dunnett's post hoc comparisons against baseline (100% values). Differences between treatments at discrete time points were examined using a two-tailed Student's t test. All statistical procedures were performed using StatExact Software.

Results

Basal synaptic transmission was not affected by compound 1 as no significant changes were found between vehicle-treated control during baseline pre-tetanus (Fig 8b).

During the LTP induction paradigm, subcutaneous administration of compound 1 (40 mg/kg) enhanced an enduring (>2 h) synaptic potentiation (Fig 8b). At 0-30 min after completion of the tetanization, PSA slopes were $164 \pm 13\%$ as compared to vehicle level $124 \pm 5\%$, $p < 0.05$). At 90-120 min post-tetanization PSA amplitude was still higher ($179 \pm 20\%$ as compared to vehicle level $116 \pm 17\%$, $p < 0.05$). Similarly,

analysis of stimulus–response curves revealed a significant lasting increase in the fEPSP slope as compared to vehicle condition (90-120 min: 137 ± 24 % as compared to vehicle level 94 ± 7 %, $p < 0.05$) (Fig 8c).

Overall, compound 1 facilitates LTP in vivo, but does not affect basal synaptic

5 transmission.

PROPHETIC COMPOSITION EXAMPLES

“Active ingredient” as used throughout these examples relates to compound 1, a pharmaceutically acceptable salt thereof, or a solvate thereof.

10 Typical examples of recipes for the formulation of the invention are as follows:

1. Tablets

Compound 1	5 to 50 mg
Di-calcium phosphate	20 mg
Lactose	30 mg
15 Talcum	10 mg
Magnesium stearate	5 mg
Potato starch	ad 200 mg

2. Suspension

20 An aqueous suspension is prepared for oral administration so that each 1 milliliter contains 1 to 5 mg of one of compound 1, 50 mg of sodium carboxymethyl cellulose, 1 mg of sodium benzoate, 500 mg of sorbitol and water ad 1 ml.

3. Injectable

25 A parenteral composition is prepared by stirring 1.5 % by weight of compound 1 of the invention in 10% by volume propylene glycol in water.


4. Ointment

Compound 1	5 to 1000 mg
30 Stearyl alcohol	3 g
Lanoline	5 g
White petroleum	15 g
Water	ad 100 g

35 Reasonable variations are not to be regarded as a departure from the scope of the invention. It will be obvious that the thus described invention may be varied in many ways by those skilled in the art.

Claims

1. A compound having the formula (1)



- 5 or a pharmaceutically acceptable salt or a solvate thereof.
2. The hydrochloride salt of the compound of formula (1) according to claim 1.
3. A pharmaceutical composition comprising a therapeutically effective amount of a
10 compound according to claim 1 or 2 and a pharmaceutically acceptable carrier.
4. A compound according to claim 1 or 2, of a pharmaceutical composition
according to claim 3, for use as a medicament.
- 15 5. A compound according to claim 1 or 2 or a pharmaceutical composition
according to claim 3 for use in treating or preventing a central nervous system disorder
selected from the group of psychotic disorders and conditions; anxiety disorders;
movement disorders; drug abuse; mood disorders; neurodegenerative disorders;
disorders or conditions comprising as a symptom a deficiency in attention and/or
20 cognition; disorders related to memory acquisition and consolidation; stroke; and
autistic disorder.
6. A compound or a pharmaceutical composition for use according to claim 5,
wherein
25 the psychotic disorders are selected from the group of schizophrenia;
schizophreniform disorder; schizoaffective disorder; delusional disorder; substance-
induced psychotic disorder; personality disorders of the paranoid type; and personality
disorder of the schizoid type;

the anxiety disorders are selected from the group of panic disorder; agoraphobia; specific phobia; social phobia; obsessive-compulsive disorder; post-traumatic stress disorder; acute stress disorder; and generalized anxiety disorder;

the movement disorders are selected from the group of Huntington's disease and dyskinesia; Parkinson's disease; restless leg syndrome and essential tremor; Tourette's syndrome and other tic disorders;

the substance-related disorders are selected from the group of alcohol abuse; alcohol dependence; alcohol withdrawal; alcohol withdrawal delirium; alcohol-induced psychotic disorder; amphetamine dependence; amphetamine withdrawal; cocaine dependence; cocaine withdrawal; nicotine dependence; nicotine withdrawal; opioid dependence and opioid withdrawal;

the mood disorders are selected from depression; mania; bipolar I disorder, bipolar II disorder; cyclothymic disorder; dysthymic disorder; major depressive disorder; treatment-resistant depression; and substance-induced mood disorder;

the neurodegenerative disorders are selected from the group of Parkinson's disease; Huntington's disease; dementia; Alzheimer's disease; multi-infarct dementia; AIDS-related dementia or frontotemporal dementia;

the disorders or conditions comprising as a symptom a deficiency in attention and/or cognition are selected from the group of dementia associated with Alzheimer's disease; multi-infarct dementia; dementia due to Lewy body disease; alcoholic dementia or substance-induced persisting dementia; dementia associated with intracranial tumours or cerebral trauma; dementia associated with Huntington's disease; dementia associated with Parkinson's disease; AIDS-related dementia; dementia due to Pick's disease; dementia due to Creutzfeldt-Jakob disease; delirium; amnesic disorder; post-traumatic stress disorder; stroke; progressive supranuclear palsy; mental retardation; a learning disorder; attention-deficit/hyperactivity disorder (ADHD); mild cognitive impairment; Asperger's syndrome; age-related cognitive impairment; and cognitive impairment related to perception, concentration, learning or memory;

the disorders related to memory acquisition and consolidation are selected from memory disorders.

7. A process for preparing a pharmaceutical composition as defined in claim 3, characterized in that a pharmaceutically acceptable carrier is intimately mixed with a therapeutically effective amount of a compound as defined in claim 1 or 2.

8. A compound according to claim 1 or 2 in combination with an additional pharmaceutical agent for use in the treatment or prevention of a condition as cited in any one of claims 5 to 6.

5 9. A product comprising

(a) a compound as defined in claim 1 or 2; and

(b) an additional pharmaceutical agent,

as a combined preparation for simultaneous, separate or sequential use in the treatment or prevention of a condition as cited in any one of claims 5 to 6.

10

10. Use of a compound according to claim 1 or 2 or a pharmaceutical composition according to claim 3 in the manufacture of a medicament for the treatment of a disorder selected from the group of psychotic disorders and conditions; anxiety disorders; movement disorders; drug abuse; mood disorders; neurodegenerative disorders; disorders or conditions comprising as a symptom a deficiency in attention and/or cognition; disorders related to memory acquisition and consolidation; stroke; and autistic disorder.

15