

NOVEL COMPOSITIONS WITH TISSUE-SPECIFIC TARGETING MOTIFS AND COMPOSITIONS CONTAINING SAME

BACKGROUND OF THE INVENTION

5 The Adeno-Associated Virus (AAV) is currently the gene therapy vector of choice. This is because AAVs can deliver a transgene that is stably expressed for decades from a non-integrating genome, and because the AAV is remarkably safe and non-immunogenic. However, AAV gene therapy is currently limited to a small number of diseases due to challenges in delivery and tropism. This is particularly true for disorders of the central nervous system
10 (CNS). Direct delivery of AAV gene therapy vectors is possible, by injecting the vector directly into the cerebrospinal Fluid (CSF), but this method typically transduces 1% or less of brain cells. Furthermore, most of that transduction is concentrated on the cells that are in direct contact with the CSF. Cells in the “deep brain” are rarely transduced. This has limited the number of CNS disorders treatable by gene therapy.

15 In contrast to the CSF network, the vascular system of the brain reaches nearly every cell in the CNS. This is because of a high demand these tissues have for glucose, oxygen, and other nutrients. However, cells in the brain and spinal cord are protected from the circulatory system by a specialized vascular unit, the Blood Brain Barrier (BBB). The BBB limits the diffusion of large molecules like viral vectors and proteins, and even many small molecule
20 drugs through a complex network of tightly-linked cells that surround the blood vessels of the brain and spinal cord. Thus, a grand challenge in gene therapy delivery to the CNS has been the engineering of an AAV variant capable of crossing the BBB at high efficiency and transducing cells in the deep brain.

 One AAV capsid developed at California Institute of Technology (CalTech) has a seven
25 amino acid peptide inserted into hypervariable loop 8 (HVR8) on the AAV9 capsid to generate a rAAV called AAV9-PHP.B which is reported to mediate interaction with Ly6a, a GPI-anchored receptor on the brain vasculature of some mouse strains. US Patent Published Application No. 2017/0166926A1. This interaction drives transport of AAV9-PHP.B across the BBB, resulting in ~50-fold higher transduction of brain cells than AAV9. However, this finding
30 has not translated to larger animals or humans.

 There remains a need for vectors which can specifically target selected tissue and cell types.

SUMMARY OF THE INVENTION

In certain embodiments, a recombinant adeno-associated virus particle (rAAV) having a capsid comprising an amino acid sequence that comprises the motif N- x- (T/I/V/A)- (K/R) (SEQ ID NO: 47) is provided. Suitably, the amino acid sequence is part of at least the AAV vp3 protein in the capsid and a vector genome packaged in the capsid which comprises a nucleic acid sequence encoding a gene product under control of sequences which direct expression thereof, provided that the capsid is not a mutant AAV2 capsid comprising an NDVRAVS (SEQ ID NO: 48) sequence. In certain embodiments, the amino acid sequence comprising the N- x- (T/I/V/A)- (K/R) motif optionally flanked at the amino terminus and/or the carboxy terminus of the motif by two amino acids to seven amino acids is inserted into the AAV capsid vp3 region. In certain embodiments, the sequence inserted into the capsid comprises: (a) SSNTVKLTSGH (SEQ ID NO: 40); (b) EFSSNTVKLTS (SEQ ID NO: 38); (c) GGVLTNARGEYMRGG (SEQ ID NO: 46); (d) GGIEINATRAGTNLGG (SEQ ID NO: 43); (e) GGSSNTVKLTSGHGG (SEQ ID NO: 39); (f) IEINATRAGTNL (SEQ ID NO: 42); or (g) SANFIKPTSY (SEQ ID NO: 41). In certain embodiments, the amino acid sequence of the motif is NTVK, which is optionally flanked by two to seven amino acids at its carboxy- and/or amino terminus and inserted between amino acids 588 and 589 of an AAV9 capsid protein, based on the numbering of amino acid sequence: SEQ ID NO: 44.

In certain embodiments, a rAAV having an inserted sequence of NTVK in its capsid, which sequence is optionally flanked by two to seven amino acids at its carboxy- and/or amino terminus and inserted between amino acids 588 and 589 of an AAV9 capsid protein, based on the numbering of amino acid sequence: SEQ ID NO: 44.

In certain embodiments, a composition comprises the rAAV having the inserted motif and optional flanking sequences and one or more of a physiologically compatible carrier, excipient, and/or aqueous suspension base.

In certain embodiments, an endothelial cell targeting peptide is provided, the peptide comprising an amino acid sequence of N- x- (T/I/V/A)- (K/R) (SEQ ID NO: 47) optionally flanked at the amino terminus and/or the carboxy terminus of the motif by two amino acids to seven amino acids, and optionally further conjugated to a nanoparticle, a second molecule, or a viral capsid protein. In certain embodiments, the endothelial cell targeting peptide comprises: (a) SSNTVKLTSGH (SEQ ID NO: 40); (b) EFSSNTVKLTS (SEQ ID NO: 38); (c)

GGVLTNIARGEYMRGG (SEQ ID NO: 46); (d) GGIEINATRAGTNLGG (SEQ ID NO: 43);
(e) GGSSNTVKLTSGHG (SEQ ID NO: 39); (f) IEINATRAGTNL (SEQ ID NO: 42); or (g)
SANFIKPTSY (SEQ ID NO: 41). In certain embodiments, the amino acid sequence of the motif
is NTVK. In certain embodiments, a composition is provided which comprises the endothelial
5 cell targeting peptide and one or more of a physiologically compatible carrier, excipient, and/or
aqueous suspension base.

In certain embodiments, a fusion polypeptide or protein comprising a brain endothelial
cell targeting peptide and a fusion partner which comprises at least one polypeptide or protein is
provided herein. In certain embodiments, a composition comprising a fusion polypeptide or
10 protein according to claim 11 and one or more of a physiologically compatible carrier,
excipient, and/or aqueous suspension base.

Provided herein are compositions and methods for using an rAAV, an endothelial cell
targeting peptide, a fusion polypeptide or protein, and/or a composition as described herein of
for delivering a therapeutic to a patient in need thereof. In certain embodiments, the therapeutic
15 is targeted to the brain endothelial cells.

In certain embodiments, a composition and method is provided for treating Allan-
Herndon-Dudley disease by delivering to a subject in need thereof an rAAV as described herein
wherein the encoded gene product is an MCT8 protein.

In certain embodiments, a method is provided for targeting therapy to the lung
20 comprising administering to a patient in need thereof an rAAV as described herein.

In certain embodiments, a method is provided for treating a disease of the lung by
delivering to a subject in need thereof an rAAV having a capsid with the inserted targeting
peptide and encoding a therapeutic gene product, wherein the encoded gene product is a soluble
Ace2 protein, an anti-SARS antibody, an anti-SARS-CoV2 antibody, an anti-influenza
25 antibody, or a cystic fibrosis transmembrane protein.

In certain embodiments, a method is provided for increasing transduction of AAV
production cells in vitro comprising inserting an N- x- (T/I/V/A)- (K/R) motif into an AAV
capsid. In certain embodiments, the production cells are 293 cells.

These and other embodiments and advantages of the invention will be apparent from the
30 specification, including, without limitation, the detailed description of the invention.

BRIEF DESCRIPTION OF THE FIGURES

FIGs 1A to 1B shows the enrichment scores for the top performing peptide hits in mouse brain in the screen, with reference peptides. FIG 1A shows enrichment scores for C57BL/6J mice. FIG 1B shows enrichment scores for Balb/c mice.

5 FIGs 2A and 2B show the enrichment scores for the top performing hits in NHP tissue in the screen. FIG 2A shows enrichment scores for NHP brain. FIG 2B shows enrichment scores for NHP spinal cord tissue.

FIGs 3A to 3D show a secondary validation of the transduction levels of top performing peptide hits in AAV capsid comprising GFP reporter transgene. The results are plotted relative to AAV9 transduction. FIG 3A shows secondary validation screen of selected peptide targeting of brain tissue in Balb/c mice. FIG 3B shows secondary validation screen of selected peptide targeting of brain tissue in C57BL/6 mice. FIG 3C shows secondary validation screen of selected peptide targeting of liver tissue in Balb/c mice. FIG 3D shows secondary validation screen of selected peptide targeting of liver tissue in C57BL/6 mice.

15 FIG. 4 shows region of the alignment of the amino acid sequences of the various AAV capsid proteins of AA9, AAV8, AAV7, AAV6, AAV5, AAV4, AAV3B, AAV2, and AAV1, which is focused on the region HVRVIII in which the targeting peptide may be inserted (based on structure analysis).

FIG 5 shows that “NxTK” motif is the critical motif for brain biodistribution in the SAN insert, and shows average impact of substitution (fold-change from original sequence).

FIG 6 show that “NxTK” motif controls plasmid-to-AAV conversion in the SAN peptide insert, and shows average impact of substitution (fold-change from original sequence).

FIGs 7A to 7D show that “NxTK” motif confers broad transduction advantage across cell lines. FIG 7A shows relative transduction levels when compared to AAV9 capsid in 293 cells. FIG 7B shows relative transduction levels when compared to AAV9 capsid in NIH3T3 cells. FIG 7C shows relative transduction levels when compared to AAV9 capsid in HUH7 cells. FIG 7D shows transduction levels at day 3 post-transduction (3DPT) and day 7 post transduction (7DPT) in macaque primary airway cells. FIG 7E shows microscopic analysis of the macaque primary airway epithelial cells in a control sample treated with carrier (i.e., no vector). FIG. 7F shows microscopic analysis of the macaque primary airway epithelial cells post-transduction with AAV9-GFP vector. FIG. 7G shows microscopic analysis of the macaque primary airway epithelial cells post-transduction with AAV9-GFP vector comprising EFS

peptide insert. FIG. 7H shows microscopic analysis of the macaque primary airway epithelial cells post-transduction with AAV9-GFP comprising SAN peptide inserts.

FIG 8 shows a preliminary transduction test with GFP vectors in cultured human cells (nasal, bronchial and tracheal) plotted as a ratio of mRNA copy number over micro-gram total mRNA.

DETAILED DESCRIPTION OF THE INVENTION

A targeting peptide sequence is provided herein. Also provided herein are fusion proteins, modified proteins, mutant viral capsids and other moieties operably linked to an exogenous targeting peptide motif of N- x- (T/I/V/A)- (K/R) (SEQ ID NO: 47). In certain embodiments, this exogenous motif confers on these compositions a modified native tissue specificity of the source (parental) protein, viral vector, or other moiety. In certain embodiments, targeting peptides in this motif provides enhanced or altered endothelial cell targeting. In certain embodiments, targeting peptides in this motif provide enhanced or altered lung, bronchial, tracheal and/or nasoeptithelial targeting. In certain embodiments, viral vectors having modified capsids with this motif exhibit increased transduction of AAV production cells in vitro.

The targeting peptide may be linked to a recombinant protein (e.g., for enzyme replacement therapy) or polypeptide (e.g., an immunoglobulin) to target to the desired tissue (e.g., CNS or lung) to form a fusion protein or a conjugate. Additionally, the targeting peptide may be linked to a liposome and/or a nanoparticle (a lipid nanoparticle, LNP) forming a peptide-coated liposome and/or LNP to target the desired tissue. Sequences encoding at least one copy of a targeting peptide and optional linking sequences may be fused in frame with the coding sequence for the recombinant protein and co-expressed with the protein or polypeptide to provide fusion proteins or conjugates. Alternatively, other synthetic methods may be used to form a conjugate with a protein, polypeptide or another moiety (e.g., DNA, RNA, or a small molecule). In certain embodiments, multiple copies of a targeting peptide are in the fusion protein/conjugate. Suitable methods for conjugating a targeting peptide to a recombinant protein include modifying the amino (N)-terminus and one or more residues on a recombinant human protein (e.g., an enzyme) using a first crosslinking agent to give rise to a first crosslinking agent modified recombinant human protein, modifying the amino (N)-terminus of a short extension linker region preceding a targeting peptide using a second crosslinking agent to give rise to a

second crosslinking agent modified variant target peptide, and then conjugating the first crosslinking agent modified recombinant human protein to the second crosslinking agent modified variant targeting peptide containing a short extension linker. Other suitable methods of conjugating a targeting peptide to a recombinant protein include conjugating a first

5 crosslinking agent modified recombinant human protein to one or more second crosslinking agent modified variant targeting peptides, wherein the first crosslinking agent modified recombinant protein comprises a recombinant protein characterized as having a chemically modified N-terminus and one or more modified lysine residues and the one or more second crosslinking agent modified variant targeting peptides comprise one or more variant targeting

10 peptides comprising a modified N-terminal amino acid of a short extension linker preceding the targeting peptide. Still other suitable methods for conjugating a targeting peptide to a protein, polypeptide, nanoparticle, or another biologically useful chemical moiety may be selected. See, e.g., US Patent No. US 9,545,450 B2 (NHS-phosphine cross-linking agents; NHS-Azide cross-linking agents); US Published Patent Application No. US 2018/0185503 A1 (aldehyde-

15 hydrazide crosslinking).

In certain embodiments, the targeting peptide may be inserted into a suitable site within a protein or polypeptide (e.g., a viral capsid protein). In certain these embodiments and in certain other embodiments, a targeting peptide may be flanked at its carboxy (COO-) and/or amino (N) terminus by a short extension linker. Such a linker may be 1 to 20 amino acid

20 residues in length, or about 2 to 20 amino acids residues, or about 1 to 15 amino acid residues, or about 2 to 12 amino acid residues, or 2 to 7 amino acid residues in length. The short extension linker can also be about 10 amino acids in length. The presence and length of a linker at the N-terminus is independently selected from a linker at the carboxy-terminus, and the presence and length of a linker at the carboxy terminus is independently selected from a linker

25 at the N-terminus. Suitable short extension linkers can be provided using a 5-amino acid flexible GS extension linker (glycine-glycine-glycine-glycine-serine), a 10-amino acid extension linker comprising 2 flexible GS linkers, a 15-amino acid extension linker comprising 3 flexible GS linkers, a 20-amino acid extension linker comprising 4 flexible GS linkers, or any combination thereof.

30 In certain embodiment, a composition is provided which is useful for targeting an endothelial cell. The composition is a mutant capsid, fusion protein or another conjugate comprising at least one exogenous targeting peptide comprising: an amino acid sequence of N-

x- (T/I/V/A)- (K/R) (SEQ ID NO: 47) optionally flanked at the amino terminus and/or the carboxy terminus of the motif by two amino acids to seven amino acids, and optionally further conjugated to a nanoparticle, a second molecule, or a viral capsid protein. The targeting peptide comprising the following sequence with optional linking sequences:

- 5 (a) SSNTVKLTSGH (SEQ ID NO: 40);
- (b) EFSSNTVKLTS (SEQ ID NO: 38);
- (c) GGVLTNIARGEYMRGG (SEQ ID NO: 46);
- (d) GGIEINATRAGTNLGG (SEQ ID NO: 43);
- (e) GGSSNTVKLTSGHGG (SEQ ID NO: 39);
- 10 (f) IEINATRAGTNL (SEQ ID NO: 42); or
- (g) SANFIKPTSY (SEQ ID NO: 41).

In certain embodiments, the targeting peptide motif is encoded by a nucleic acid sequence selected from:

- (a) agcagcaacaccgtgaagctgaccagcggacac (SEQ ID NO: 54);
- 15 (b) gagttcagcagcaacaccgtgaagctgaccagc (SEQ ID NO: 50);
- (c) ggaggagtgtgaccaacatcgctaggagagtacatgagaggagga (SEQ ID NO: 56);
- (d) ggaggaatcgagatcaacgctaccagagctggaaccaacctgggagga (SEQ ID NO: 52);
- (e) ggaggaagcagcaacaccgtgaagctgaccagcggacacggagga (SEQ ID NO: 55);
- (f) atcgagatcaacgctaccagagctggaaccaacctg (SEQ ID NO: 51); or
- 20 (g) agcgctaacttcataagcctaccagctac (SEQ ID NO: 53).

In certain embodiments, the targeting peptide is encoded by a nucleic acid sequence of SEQ ID NO: 50, or a sequence at least about 70% identical thereto. In certain embodiments, the targeting peptide is encoded by a nucleic acid sequence of SEQ ID NO: 51, or a sequence at least about 70% identical thereto. In certain embodiments, the targeting peptide is encoded by a nucleic acid sequence of SEQ ID NO: 52, or a sequence at least about 70% identical thereto. In certain embodiments, the targeting peptide is encoded by a nucleic acid sequence of SEQ ID NO: 53, or a sequence at least about 70% identical thereto. In certain embodiments, the targeting peptide is encoded by a nucleic acid sequence of SEQ ID NO: 54, or a sequence at least about 70% identical thereto. In certain embodiments, the targeting peptide is encoded by a nucleic acid sequence of SEQ ID NO: 55, or a sequence at least about 70% identical thereto. In certain embodiments, the targeting peptide is encoded by a nucleic acid sequence of SEQ ID

NO: 56, or a sequence at least about 70% identical thereto. In some embodiments, the nucleic acid sequence encoding for the targeting peptide motif is optionally flanked at the 5' and/or 3' ends of the nucleic acid sequence of the motif by six to twenty one nucleotides of an extension linker.

5

In certain embodiments, the targeting peptide NTVK. In certain embodiments, the targeting peptide NTVR. In certain embodiment, more than one copy of a targeting peptide within this motif is provided in a conjugate or modified protein (e.g., a parvovirus capsid). In certain embodiments, two or more different targeting peptides are present

10

In certain embodiment, a composition is provided which is useful for targeting a naso-epithelial and/or lung epithelial cell. The composition is a mutant capsid, fusion protein or another conjugate comprising at least one exogenous targeting peptide comprising: an amino acid sequence of N- x- (T/I/V/A)- (K/R) (SEQ ID NO: 47) optionally flanked at the amino terminus and/or the carboxy terminus of the motif by two amino acids to seven amino acids, and optionally further conjugated to a nanoparticle, a second molecule, or a viral capsid protein. The targeting peptide comprises: (a) SSNTVKLTSGH (SEQ ID NO: 40); (b) EFSSNTVKLTS (SEQ ID NO: 38); (c) GGVLTNARGEYMRGG (SEQ ID NO: 46); (d) GGIEINATRAGTNLGG (SEQ ID NO: 43); (e) GGSSNTVKLTSGHGG (SEQ ID NO: 39); (f) IEINATRAGTNL (SEQ ID NO: 42); or (g) SANFIKPTSY (SEQ ID NO: 41).

15

20

In certain embodiments, the targeting peptide NTVK. In certain embodiments, the targeting peptide NTVR, optionally flanked by spacer amino acids as described herein. In certain embodiment, more than one copy of a targeting peptide within this motif is provided in a conjugate or modified protein (e.g., a parvovirus capsid). In certain embodiments, two or more different targeting peptides are present.

25

Examples of suitable proteins, including enzymes, immunoglobulins, therapeutic proteins, immunogenic polypeptides, nanoparticles, DNA, RNA, and other moieties (e.g., small molecules, etc.) for targeting are described in more detail below. These and other biologic and chemical moieties are suitable for use with the targeting peptide(s) provided herein.

30

In certain embodiments, a composition is a nucleic acid sequence molecule, wherein the nucleic acid sequence is a DNA molecule or RNA molecule, e.g., naked DNA, naked plasmid DNA, messenger RNA (mRNA), containing the targeting peptide sequence motif linked to the nucleic acid molecule. In some embodiments, the nucleic acid molecule is further coupled with various compositions and nano particles, including, e.g., micelles, liposomes, cationic lipid -

nucleic acid compositions, poly-glycan compositions and other polymers, lipid and/or cholesterol-based - nucleic acid conjugates, and other constructs such as are described herein. See, e.g., WO2014/089486, US 2018/035361A1, US2013/0037977A1, WO2015/074085A1, US9670152B2, and US 8,853,377B2, X. Su et al, Mol. Pharmaceutics, 2011, 8 (3), pp 774–787; 5 web publication: March 21, 2011; WO2013/182683, WO 2010/053572 and WO 2012/170930, all of which are incorporated herein by reference. In certain embodiments, the targeting peptide motif is chemically linked to a nanoparticle surface, wherein the nanoparticle encapsulates a nucleic acid molecule. In some embodiments the nanoparticle comprising the targeting peptide linked to the surface is designed for targeted tissue-specific delivery. In some embodiments two 10 or more different targeting peptides are linked to the surface of the nanoparticle. Suitable chemical linking or cross-linking include those known to one skilled in the art.

Capsids

In certain embodiments, a recombinant parvovirus is provided which has a modified 15 parvovirus capsid having at least exogenous peptide from the N- x- (T/I/V/A)- (K/R) targeting motif. Such a recombinant parvovirus may be a hybrid bocavirus/AAV or a recombinant AAV vector. In other embodiments, other viral vectors may be generated having one or more exogenous targeting peptides from the N- x- (T/I/V/A)- (K/R) motif (which may be same or different, or combinations thereof) in an exposed capsid protein to modulate and/or alter the 20 targeting specificity of the viral vector as compared to the parental vector.

The targeting peptide may be inserted into a hypervariable loop (HVR) VIII at any suitable location. For example, based on the numbering of the AAV9 capsid, the peptide is inserted with linkers of various lengths between amino acids 588 and 589 (Q-A) of the AAV9 capsid protein, based on the numbering of the AAV9 VP1 amino acid sequence: SEQ ID NO: 25 44. See, also, WO 2019/168961, published September 6, 2019, including Table G providing the deamidation pattern for AAV9 and WO 2020/160582, filed September 7, 2018. The amino acid residue locations are identical in AAVhu68 (SEQ ID NO: 45). However, another site may be selected within HVRVIII. Alternatively, another exposed loop HVR (e.g., HVRIV) may be selected for the insertion. Comparable HVR regions may be selected in other capsids. In certain 30 embodiments, the location for the HVRVIII and HVRIV is determined using an algorithm and/or alignment technique as described in US Patent No. US 9,737,618 B2 (column 15, lines 3-23), and US Patent No. US 10,308,958 B2 (column 15, line 46 – column 16, line 6), which

are incorporated herein by reference in its entirety. In certain embodiments, AAV1 capsid protein is selected as a parental capsid, wherein the targeting peptide with linkers of various lengths is inserted in a suitable location of the HVRVIII region of amino acid 582 to 585, or HVRIV region of amino acid 456 to 459 based on vp1 numbering (Gurda, BL., et al., Capsid Antibodies to Different Adeno-Associated Virus Serotypes Bind Common Regions, 2012, 5 Journal of Virology, June 12, 2013, 87(16): 9111-91114). In certain embodiments, AAV8 is selected as parental capsid, wherein the targeting peptide with linkers of various length is inserted in a suitable location of HVRVIII region of amino acid 586 to 591 (e.g., 590-591 (N-T)), or HVRIV region of amino acid 456 to 460, based on VP1 numbering (Gurda, BL., et al., 10 Mapping a Neutralizing epitope onto the Capsid of Adeno-Associated Virus Serotype 8, 2012, Journal of Virology, May 16, 2012, 86(15):7739-7751). In certain embodiments, the AAV7 is selected as parental capsid, wherein the targeting peptide with linkers of various length is inserted in a suitable location of amino acid 589 to 590 (N-T). In certain embodiments, the AAV6 is selected as parental capsid, wherein the targeting peptide with linkers of various 15 length is inserted in a suitable location of amino acid 588 to 589 (S-T). In certain embodiments, the AAV5 is selected as parental capsid, wherein the targeting peptide with linkers of various length is inserted in a suitable location of amino acid 577 to 578 (T-T). In certain embodiments, the AAV4 is selected as parental capsid, wherein the targeting peptide with linkers of various length is inserted in a suitable location of amino acid 586 to 587 (S-N). In certain embodiments, 20 the AAV3B is selected as parental capsid, wherein the targeting peptide with linkers of various length is inserted in a suitable location of amino acid 588 to 589 (N-T). In certain embodiments, the AAV2 is selected as parental capsid, wherein the targeting peptide with linkers of various length is inserted in a suitable location of amino acid 587 to 588 (N-R). In certain embodiments, the AAV1 is selected as parental capsid, wherein the targeting peptide with linkers of various 25 length is inserted in a suitable location of amino acid 589 to 589 (S-T). See also, FIG. 4.

In certain embodiments, the parental capsid modified to contain the N- x- (T/I/V/A)- (K/R) motif, with optional flanking sequences, is selected from parvoviruses which natively target the CNS (e.g., Clade F AAV (e.g., AAVhu68 or AAV9), Clade E (e.g., AAV8), or certain Clade A AAV (e.g., AAV1, AAVrh91)) capsids, or non-parvovirus capsids (e.g., herpes 30 simplex virus, etc.) in order enhance expression and/or otherwise modulate the type of CNS targeted cells. In other embodiments, the capsid is selected from parvoviruses which do not natively target the CNS (e.g., Clade F AAV, e.g., AAVhu68 or AAV9, or certain Clade A

AAV, e.g., AAV1, AAVrh91) capsids, or non-parvovirus capsids (e.g., herpes simplex virus (HSV), etc.). See, e.g., WO 2020/223231, published November 5, 2020 (rh91, including table with deamidation pattern), US Provisional Patent Application No. 63/065,616, filed August 14, 2020 and US Provisional Patent Application No. 63/109734, filed November 4, 2020. In certain
5 embodiments, the capsid is selected from AAV Clade F AAVhu95 and AAVhu96 capsids. See, e.g., US Provisional Application No. 63/251,599, filed October 2, 2201.

In certain embodiments, the parental capsid modified to contain the N- x- (T/I/V/A)- (K/R) motif is selected from viruses (e.g., AAV) which natively target nasal epithelial cells, nasopharynx cells, and/or lung cells in order to enhance targeting as compared to the parental
10 AAV (e.g., Clade A AAV, e.g., AAV1, AAVrh32.33, AAV6.2, AAV6, AAVrh91), or AAV5, or certain Clade F AAV, e.g., AAVhu68 or AAV9, capsids, or non-parvovirus capsids (e.g., adenoviruses, HSV, RSV, etc.). See, e.g., WO 2020/223231, published November 5, 2020 (rh91, including table with deamidation pattern), US Provisional Patent Application No. 63/065,616, filed August 14, 2020 and US Provisional Patent Application No. 63/109734, filed
15 November 4, 2020.

In certain embodiments, the AAV capsid is not a mutant AAV2 capsid comprising an NDVRAVS (SEQ ID NO: 48) sequence.

For example, capsids from Clade F AAV such as AAVhu68 or AAV9 may be selected. Methods of generating vectors having the AAV9 capsid or AAVhu68 capsid, and/or chimeric
20 capsids derived from AAV9 have been described. See, e.g., US 7,906,111, which is incorporated by reference herein. Other AAV serotypes which transduce nasal cells or another suitable target (e.g., muscle or lung) may be selected as sources for capsids of AAV viral vectors including, e.g., AAV1, AAV2, AAV3, AAV4, AAV5, AAV6, AAV6.2, AAV7, AAV8, AAV9, rh10, AAVrh64R1, AAVrh64R2, rh8, AAVrh32.33 (See, e.g., US Published Patent
25 Application No. 2007-0036760-A1; US Published Patent Application No. 2009-0197338-A1; and EP 1310571). See also, WO 2003/042397 (AAV7 and other simian AAV), US Patent 7790449 and US Patent 7282199 (AAV8), WO 2005/033321 (AAV9), and WO 2006/110689, or yet to be discovered, or a recombinant AAV based thereon, may be used as a source for the AAV capsid. See, e.g., WO 2020/223232 A1 (AAV rh90), WO 2020/223231 A1 and
30 International Application No. PCT/US21/45945, filed August 13, 2021 (AAV rh91), and WO 2020/223236 A1 (AAV rh92, AAV rh93, AAV rh9193), which are incorporated herein by

reference in its entirety. These documents also describe other AAV which may be selected for generating AAV and are incorporated by reference. In some embodiments, an AAV capsid (cap) for use in the viral vector can be generated by mutagenesis (i.e., by insertions, deletions, or substitutions) of one of the aforementioned AAV caps or its encoding nucleic acid. In some
5 embodiments, the AAV capsid is chimeric, comprising domains from two or three or four or more of the aforementioned AAV capsid proteins. In some embodiments, the AAV capsid is a mosaic of Vp1, Vp2, and Vp3 monomers from two or three different AAVs or recombinant AAVs. In some embodiments, an rAAV composition comprises more than one of the aforementioned caps.

10 As used herein, the term “clade” as it relates to groups of AAV refers to a group of AAV which are phylogenetically related to one another as determined using a Neighbor-Joining algorithm by a bootstrap value of at least 75% (of at least 1000 replicates) and a Poisson correction distance measurement of no more than 0.05, based on alignment of the AAV vp1 amino acid sequence. The Neighbor-Joining algorithm has been described in the literature. See,
15 e.g., M. Nei and S. Kumar, *Molecular Evolution and Phylogenetics* (Oxford University Press, New York (2000)). Computer programs are available that can be used to implement this algorithm. For example, the MEGA v2.1 program implements the modified Nei-Gojobori method. Using these techniques and computer programs, and the sequence of an AAV vp1 capsid protein, one of skill in the art can readily determine whether a selected AAV is contained
20 in one of the clades identified herein, in another clade, or is outside these clades. See, e.g., G Gao, et al, *J Virol*, 2004 Jun; 78(12): 6381-6388, which identifies Clades A, B, C, D, E and F, and provides nucleic acid sequences of novel AAV, GenBank Accession Numbers AY530553 to AY530629. See, also, WO 2005/033321.

 As used herein, an “AAV9 capsid” is a self-assembled AAV capsid composed of
25 multiple AAV9 vp proteins. The AAV9 vp proteins are typically expressed as alternative splice variants encoded by a nucleic acid sequence which encodes the vp1 amino acid sequence of GenBank accession: AAS99264. These splice variants result in proteins of different length. In certain embodiments, “AAV9 capsid” includes an AAV having an amino acid sequence which is 99% identical to AAS99264 or 99% identical thereto. See, also, WO 2019/168961, published
30 September 6, 2019, including Table G providing the deamidation pattern for AAV9. See, also US7906111 and WO 2005/033321. As used herein “AAV9 variants” include those described in, e.g., WO2016/049230, US 8,927,514, US 2015/0344911, and US 8,734,809.

A rAAVhu68 is composed of an AAVhu68 capsid and a vector genome. An AAVhu68 capsid is an assembly of a heterogenous population of vp1, a heterogenous population of vp2, and a heterogenous population of vp3 proteins. As used herein when used to refer to vp capsid proteins, the term “heterogenous” or any grammatical variation thereof, refers to a population
5 consisting of elements that are not the same, for example, having vp1, vp2 or vp3 monomers (proteins) with different modified amino acid sequences. See, also, PCT/US2018/019992, WO 2018/160582, entitled “Adeno-Associated Virus (AAV) Clade F Vector and Uses Therefor”, and which are incorporated herein by reference in its entirety.

For other recombinant viral vectors, suitable exposed portions of the viral capsid or
10 envelope protein which is responsible for targeting specificity are selected for insertion of the targeting peptide. For example, in an adenovirus, it may be desirable to modify the hexon protein. In a lentivirus, an envelope fusion protein may modified comprise one or more copies of the targeting motif. For vaccinia virus, the major glycoprotein may be modified to comprise one or more copies of the targeting motif. Suitably, these recombinant viral vectors are
15 replication-defective for safety purposes.

Expression Cassette and Vectors

Vector genomic sequences which are packaged into an AAV capsid and delivered to a host cell are typically composed of, at a minimum, a transgene and its regulatory sequences, and
20 AAV inverted terminal repeats (ITRs). Both single-stranded AAV and self-complementary (sc) AAV are encompassed with the rAAV. The transgene is a nucleic acid coding sequence, heterologous to the vector sequences, which encodes a polypeptide, protein, functional RNA molecule (e.g., miRNA, miRNA inhibitor) or other gene product, of interest. The nucleic acid coding sequence is operatively linked to regulatory components in a manner which permits
25 transgene transcription, translation, and/or expression in a cell of a target tissue.

The AAV sequences of the vector typically comprise the cis-acting 5' and 3' inverted terminal repeat (ITR) sequences (See, e.g., B. J. Carter, in “Handbook of Parvoviruses”, ed., P. Tijsser, CRC Press, pp. 155 168 (1990)). The ITR sequences are about 145 base pairs (bp) in length. Preferably, substantially the entire sequences encoding the ITRs are used in the
30 molecule, although some degree of minor modification of these sequences is permissible. The ability to modify these ITR sequences is within the skill of the art. (See, e.g., texts such as Sambrook et al, “Molecular Cloning. A Laboratory Manual”, 2d ed., Cold Spring Harbor

Laboratory, New York (1989); and K. Fisher et al., J. Virol., 70:520-532 (1996)). An example of such a molecule employed in the present invention is a “cis-acting” plasmid containing the transgene, in which the selected transgene sequence and associated regulatory elements are flanked by the 5' and 3' AAV ITR sequences. In one embodiment, the ITRs are from an AAV different than that supplying a capsid. In one embodiment, the ITR sequences are from AAV2. A shortened version of the 5' ITR, termed Δ ITR, has been described in which the D-sequence and terminal resolution site (trs) are deleted. In certain embodiments, the vector genome (e.g., of a plasmid) includes a shortened AAV2 ITR of 130 base pairs, wherein the external A elements is deleted. The shortened ITR may revert back to the wild-type length of 145 base pairs during vector DNA amplification using the internal A element as a template and packaging into the capsid to form the viral particle. In other embodiments, the full-length AAV 5' and 3' ITRs are used. However, ITRs from other AAV sources may be selected. Where the source of the ITRs is from AAV2 and the AAV capsid is from another AAV source, the resulting vector may be termed pseudotyped. However, other configurations of these elements may be suitable.

In addition to the major elements identified above for the recombinant AAV vector, the vector also includes conventional control elements necessary which are operably linked to the transgene in a manner which permits its transcription, translation and/or expression in a cell transfected with the plasmid vector or infected with the virus produced by the invention. As used herein, “operably linked” sequences include both expression control sequences that are contiguous with the gene of interest and expression control sequences that act in trans or at a distance to control the gene of interest.

The regulatory control elements typically contain a promoter sequence as part of the expression control sequences, e.g., located between the selected 5' ITR sequence and the coding sequence. Constitutive promoters, regulatable promoters [see, e.g., WO 2011/126808 and WO 2013/04943], tissue specific promoters, or a promoter responsive to physiologic cues may be used may be utilized in the vectors described herein.

Examples of constitutive promoters suitable for controlling expression of the therapeutic products include, but are not limited to chicken β -actin (CB) promoter, CB7 promoter, human cytomegalovirus (CMV) promoter, ubiquitin C promoter (UbC), the early and late promoters of simian virus 40 (SV40), U6 promoter, metallothionein promoters, EFl α promoter, ubiquitin promoter, hypoxanthine phosphoribosyl transferase (HPRT) promoter, dihydrofolate reductase

(DHFR) promoter (Scharfmann et al., Proc. Natl. Acad. Sci. USA 88:4626-4630 (1991), adenosine deaminase promoter, phosphoglycerol kinase (PGK) promoter, pyruvate kinase promoter phosphoglycerol mutase promoter, the β -actin promoter (Lai et al., Proc. Natl. Acad. Sci. USA 86: 10006-10010 (1989)), the long terminal repeats (LTR) of Moloney Leukemia

5 Virus and other retroviruses, the thymidine kinase promoter of Herpes Simplex Virus and other constitutive promoters known to those of skill in the art. Examples of tissue- or cell-specific promoters suitable for use in the present invention include, but are not limited to, endothelin-I (ET -I) and Flt-I, which are specific for endothelial cells, FoxJ1 (that targets ciliated cells). Other examples of tissue specific promoters suitable for use in the present invention include, but

10 are not limited to, liver-specific promoters. Examples of liver-specific promoters may include, e.g., thyroid hormone-binding globulin (TBG), albumin, Miyatake et al., (1997) J. Virol., 71:5124-32; hepatitis B virus core promoter, Sandig et al., (1996) Gene Ther., 3:1002-9; or human alpha 1-antitrypsin, phosphoenolpyruvate carboxykinase (PECK), or alpha fetoprotein (AFP), Arbuthnot et al., (1996) Hum. Gene Ther., 7:1503-14). Preferably, such promoters are

15 of human origin.

Inducible promoters suitable for controlling expression of the therapeutic product include promoters responsive to exogenous agents (e.g., pharmacological agents) or to physiological cues. These response elements include, but are not limited to a hypoxia response element (HRE) that binds HIF-1 α and β , a metal-ion response element such as described by

20 Mayo et al. (1982, Cell 29:99-108); Brinster et al. (1982, Nature 296:39-42) and Searle et al. (1985, Mol. Cell. Biol. 5:1480-1489); or a heat shock response element such as described by Nouer et al. (in: Heat Shock Response, ed. Nouer, L., CRC, Boca Raton, Fla., pp167-220, 1991).

In one embodiment, expression of the gene product is controlled by a regulatable promoter that provides tight control over the transcription of the sequence encoding the gene

25 product, e.g., a pharmacological agent, or transcription factors activated by a pharmacological agent or in alternative embodiments, physiological cues. Promoter systems that are non-leaky and that can be tightly controlled are preferred.

Examples of regulatable promoters which are ligand-dependent transcription factor complexes that may be used in the invention include, without limitation, members of the nuclear

30 receptor superfamily activated by their respective ligands (e.g., glucocorticoid, estrogen, progestin, retinoid, ecdysone, and analogs and mimetics thereof) and rTTA activated by tetracycline. In one aspect of the invention, the gene switch is an EcR-based gene switch.

Examples of such systems include, without limitation, the systems described in US Patent Nos. 6,258,603, 7,045,315, U.S. Published Patent Application Nos. 2006/0014711, 2007/0161086, and International Published Application No. WO 01/70816. Examples of chimeric ecdysone receptor systems are described in U.S. Pat. No. 7,091,038, U.S. Published Patent Application
5 Nos. 2002/0110861, 2004/0033600, 2004/0096942, 2005/0266457, and 2006/0100416, and International Published Application Nos. WO 01/70816, WO 02/066612, WO 02/066613, WO 02/066614, WO 02/066615, WO 02/29075, and WO 2005/108617, each of which is incorporated by reference in its entirety. An example of a non-steroidal ecdysone agonist-regulated system is the RheoSwitch® Mammalian Inducible Expression System (New England
10 Biolabs, Ipswich, MA).

Still other promoter systems may include response elements including but not limited to a tetracycline (tet) response element (such as described by Gossen & Bujard (1992, Proc. Natl. Acad. Sci. USA 89:5547-551); or a hormone response element such as described by Lee et al. (1981, Nature 294:228-232); Hynes et al. (1981, Proc. Natl. Acad. Sci. USA 78:2038-2042);
15 Klock et al. (1987, Nature 329:734-736); and Israel & Kaufman (1989, Nucl. Acids Res. 17:2589-2604) and other inducible promoters known in the art. Using such promoters, expression of the soluble hACE2 construct can be controlled, for example, by the Tet-on/off system (Gossen et al., 1995, Science 268:1766-9; Gossen et al., 1992, Proc. Natl. Acad. Sci. USA., 89(12):5547-51); the TetR-KRAB system (Urrutia R., 2003, Genome Biol., 4(10):231;
20 Deuschle U et al., 1995, Mol Cell Biol. (4):1907-14); the mifepristone (RU486) regulatable system (Geneswitch; Wang Y et al., 1994, Proc. Natl. Acad. Sci. USA., 91(17):8180-4; Schillinger et al., 2005, Proc. Natl. Acad. Sci. U S A. 102(39):13789-94); and the humanized tamoxifen-dep regulatable system (Roscelli et al., 2002, Mol. Ther. 6(5):653-63).

In another aspect, the gene switch is based on heterodimerization of FK506 binding
25 protein (FKBP) with FKBP rapamycin associated protein (FRAP) and is regulated through rapamycin or its non-immunosuppressive analogs. Examples of such systems, include, without limitation, the ARGENT™ Transcriptional Technology (ARIAD Pharmaceuticals, Cambridge, Mass.) and the systems described in U.S. Pat. Nos. 6,015,709, 6,117,680, 6,479,653, 6,187,757, and 6,649,595, U.S. Publication No. 2002/0173474, U.S. Publication No. 200910100535, U.S.
30 Patent No. 5,834,266, U.S. Patent No. 7,109,317, U.S. Patent No. 7,485,441, U.S. Patent No. 5,830,462, U.S. Patent No. 5,869,337, U.S. Patent No. 5,871,753, U.S. Patent No. 6,011,018, U.S. Patent No. 6,043,082, U.S. Patent No. 6,046,047, U.S. Patent No. 6,063,625, U.S. Patent

No. 6,140,120, U.S. Patent No. 6,165,787, U.S. Patent No. 6,972,193, U.S. Patent No. 6,326,166, U.S. Patent No. 7,008,780, U.S. Patent No. 6,133,456, U.S. Patent No. 6,150,527, U.S. Patent No. 6,506,379, U.S. Patent No. 6,258,823, U.S. Patent No. 6,693,189, U.S. Patent No. 6,127,521, U.S. Patent No. 6,150,137, U.S. Patent No. 6,464,974, U.S. Patent No. 5 6,509,152, U.S. Patent No. 6,015,709, U.S. Patent No. 6,117,680, U.S. Patent No. 6,479,653, U.S. Patent No. 6,187,757, U.S. Patent No. 6,649,595, U.S. Patent No. 6,984,635, U.S. Patent No. 7,067,526, U.S. Patent No. 7,196,192, U.S. Patent No. 6,476,200, U.S. Patent No. 6,492,106, WO 94/18347, WO 96/20951, WO 96/06097, WO 97/31898, WO 96/41865, WO 98/02441, WO 95/33052, WO 99/110508, WO 99/110510, WO 99/36553, WO 99/41258, WO 10 01114387, ARGENT™ Regulated Transcription Retrovirus Kit, Version 2.0 (9109102), and ARGENT™ Regulated Transcription Plasmid Kit, Version 2.0 (9109/02), each of which is incorporated herein by reference in its entirety. The Ariad system is designed to be induced by rapamycin and analogs thereof referred to as "rapalogs". Examples of suitable rapamycins are provided in the documents listed above in connection with the description of the ARGENT™

15 system. In one embodiment, the molecule is rapamycin [e.g., marketed as Rapamune™ by Pfizer]. In another embodiment, a rapalog known as AP21967 [ARIAD] is used. Examples of these dimerizer molecules that can be used in the present invention include, but are not limited to rapamycin, FK506, FK1012 (a homodimer of FK506), rapamycin analogs ("rapalogs") which are readily prepared by chemical modifications of the natural product to add a "bump" that

20 reduces or eliminates affinity for endogenous FKBP and/or FRAP. Examples of rapalogs include, but are not limited to such as AP26113 (Ariad), AP1510 (Amara, J.F., et al., 1997, Proc. Natl. Acad. Sci. USA, 94(20): 10618-23) AP22660, AP22594, AP21370, AP22594, AP23054, AP1855, AP1856, AP1701, AP1861, AP1692 and AP1889, with designed 'bumps' that minimize interactions with endogenous FKBP. Still other rapalogs may be selected, e.g.,

25 AP23573 [Merck]. In certain embodiments, rapamycin or a suitable analog may be delivered locally to the AAV-transfected cells of the nasopharynx. This local delivery may be by intranasal injection, topically to the cells via bolus, cream, or gel. See US Patent Application US 2019/0216841 A1, which is incorporated herein by reference.

Other suitable enhancers include those that are appropriate for a desired target tissue

30 indication. In one embodiment, the expression cassette comprises one or more expression enhancers. In one embodiment, the expression cassette contains two or more expression enhancers. These enhancers may be the same or may differ from one another. For example, an

enhancer may include a CMV immediate early enhancer. This enhancer may be present in two copies which are located adjacent to one another. Alternatively, the dual copies of the enhancer may be separated by one or more sequences. In still another embodiment, the expression cassette further contains an intron, e.g., the chicken beta-actin intron. Other suitable introns include those known in the art, e.g., such as are described in WO 2011/126808. Examples of suitable polyadenylation (polyA) sequences include, e.g., rabbit binding globulin (also referenced to as rabbit beta globin, or rBG), SV40, SV50, bovine growth hormone (bGH), human growth hormone, and synthetic polyAs. Optionally, one or more sequences may be selected to stabilize mRNA. An example of such a sequence is a modified WPRE sequence, which may be engineered upstream of the polyA sequence and downstream of the coding sequence (see, e.g., MA Zanta-Boussif, et al, Gene Therapy (2009) 16: 605-619).

An AAV viral vector may include multiple transgenes. In certain situations, a different transgene may be used to encode each subunit of a protein (e.g., an immunoglobulin domain, an immunoglobulin heavy chain, an immunoglobulin light chain). In one embodiment, a cell produces the multi-subunit protein following infected/transfection with the virus containing each of the different subunits. In another embodiment, different subunits of a protein may be encoded by the same transgene. An IRES is desirable when the size of the DNA encoding each of the subunits is small, e.g., the total size of the DNA encoding the subunits and the IRES is less than five kilobases. As an alternative to an IRES, the DNA may be separated by sequences encoding a 2A peptide, which self-cleaves in a post-translational event. See, e.g., ML Donnelly, et al, (Jan 1997) J. Gen. Virol., 78(Pt 1):13-21; S. Furler, S et al, (June 2001) Gene Ther., 8(11):864-873; H. Klump, et al., (May 2001) Gene Ther., 8(10):811-817. This 2A peptide is significantly smaller than IRES, making it well suited for use when space is a limiting factor. More often, when the transgene is large, consists of multi-subunits, or two transgenes are co-delivered, rAAV carrying the desired transgene(s) or subunits are co-administered to allow them to concatamerize in vivo to form a single vector genome. In such an embodiment, a first AAV may carry an expression cassette which expresses a single transgene and a second AAV may carry an expression cassette which expresses a different transgene for co-expression in the host cell. However, the selected transgene may encode any biologically active product or other product, e.g., a product desirable for study.

In addition to the elements identified above for the expression cassette, the vector also includes conventional control elements which are operably linked to the coding sequence in a

manner which permits transcription, translation and/or expression of the encoded product (e.g., soluble hACE2 construct, an anti-influenza antibody, an anti-COVID19 antibody) in a cell transfected with the plasmid vector or infected with the virus produced by the invention.

Examples of other suitable transgenes are provided herein. As used herein, “operably linked”

- 5 sequences include both expression control sequences that are contiguous with the gene of interest and expression control sequences that act in trans or at a distance to control the gene of interest.

- Expression control sequences include appropriate enhancer; transcription factor; transcription terminator; promoter; efficient RNA processing signals such as splicing and polyadenylation (polyA) signals; sequences that stabilize cytoplasmic mRNA, for example Woodchuck Hepatitis Virus (WHP) Posttranscriptional Regulatory Element (WPRE); sequences that enhance translation efficiency (i.e., Kozak consensus sequence); sequences that enhance protein stability; and when desired, sequences that enhance secretion of the encoded product. In one embodiment, the regulatory sequences are selected such that the total rAAV vector genome is about 2.0 to about 5.5 kilobases in size. In one embodiment, it is desirable that the rAAV vector genome approximate the size of the native AAV genome. Thus, in one embodiment, the regulatory sequences are selected such that the total rAAV vector genome is about 4.7 kb in size. In another embodiment, the total rAAV vector genome is less about 5.2kb in size. The size of the vector genome may be manipulated based on the size of the regulatory sequences including the promoter, enhancer, intron, poly A, etc. See, Wu et al, Mol. Ther., Jan 2010 18(1):80-6, which is incorporated herein by reference.
- 10
15
20

- Thus, in one embodiment, an intron is included in the vector. Suitable introns include chicken beta-actin intron, the human beta globin IVS2 (Kelly et al, Nucleic Acids Research, 43(9):4721-32 (2015)); the Promega chimeric intron (Almond, B. and Schenborn, E. T. A Comparison of pCI-neo Vector and pcDNA4/HisMax Vector); and the hFIX intron. Various introns suitable herein are known in the art and include, without limitation, those found at bpg.utoledo.edu/~afedorov/lab/eid.html, which is incorporated herein by reference. See also, Shepelev V., Fedorov A. Advances in the Exon-Intron Database. Briefings in Bioinformatics 2006, 7: 178-185, which is incorporated herein by reference.
- 25

- Several different viral genomes were generated in the studies described herein. However, it will be understood by the skilled artisan that other genomic configurations,
- 30

including other regulatory sequences may be substituted for the promoter, enhancer and other coding sequences may be selected.

rAAV Vector Production

5 For use in producing an AAV viral vector (e.g., a recombinant (r) AAV), the expression cassettes can be carried on any suitable vector, e.g., a plasmid, which is delivered to a packaging host cell. The plasmids useful in this invention may be engineered such that they are suitable for replication and packaging in vitro in prokaryotic cells, insect cells, mammalian cells, among others. Suitable transfection techniques and packaging host cells are known and/or
10 can be readily designed by one of skill in the art.

In certain embodiments, the inclusion of the at least one copy of the N- x- (T/I/V/A)- (K/R) motif into an AAV capsid provides advantages in production as compared to the method without inclusion of at least one copy of motif in AAV capsid, and wherein the production cells are 293 cells.

15 Methods of preparing AAV-based vectors (e.g., having an AAV9 or another AAV capsid) are known. See, e.g., US Published Patent Application No. 2007/0036760 (February 15, 2007), which is incorporated by reference herein. The invention is not limited to the use of AAV9 or other clade F AAV amino acid sequences, but encompasses peptides and/or proteins containing the terminal β -galactose binding generated by other methods known in the art,
20 including, e.g., by chemical synthesis, by other synthetic techniques, or by other methods. The sequences of any of the AAV capsids provided herein can be readily generated using a variety of techniques. Suitable production techniques are well known to those of skill in the art. See, e.g., Sambrook et al, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Press (Cold Spring Harbor, NY). Alternatively, peptides can also be synthesized by the well-known
25 solid phase peptide synthesis methods (Merrifield, (1962) J. Am. Chem. Soc., 85:2149; Stewart and Young, Solid Phase Peptide Synthesis (Freeman, San Francisco, 1969) pp. 27-62). These methods may involve, e.g., culturing a host cell which contains a nucleic acid sequence encoding an AAV capsid; a functional rep gene; a minigene composed of, at a minimum, AAV inverted terminal repeats (ITRs) and a transgene; and sufficient helper functions to permit
30 packaging of the minigene into the AAV capsid protein. These and other suitable production methods are within the knowledge of those of skill in the art and are not a limitation of the present invention.

The components required to be cultured in the host cell to package an AAV minigene in an AAV capsid may be provided to the host cell in trans. Alternatively, any one or more of the required components (e.g., minigene, rep sequences, cap sequences, and/or helper functions) may be provided by a stable host cell which has been engineered to contain one or more of the required components using methods known to those of skill in the art. Most suitably, such a stable host cell will contain the required component(s) under the control of an inducible promoter. However, the required component(s) may be under the control of a constitutive promoter. Examples of suitable inducible and constitutive promoters are provided herein, in the discussion of regulatory elements suitable for use with the transgene. In still another alternative, a selected stable host cell may contain selected component(s) under the control of a constitutive promoter and other selected component(s) under the control of one or more inducible promoters. For example, a stable host cell may be generated which is derived from 293 cells (which contain E1 helper functions under the control of a constitutive promoter), but which contains the rep and/or cap proteins under the control of inducible promoters. Still other stable host cells may be generated by one of skill in the art.

These rAAVs are particularly well suited to gene delivery for therapeutic purposes and for preventing infection. Further, the compositions of the invention may also be used for production of a desired gene product in vitro. For in vitro production, a desired product (e.g., a protein) may be obtained from a desired culture following transfection of host cells with a rAAV containing the molecule encoding the desired product and culturing the cell culture under conditions which permit expression. The expressed product may then be purified and isolated, as desired. Suitable techniques for transfection, cell culturing, purification, and isolation are known to those of skill in the art. Methods for generating and isolating AAVs suitable for use as vectors are known in the art. See generally, e.g., Grieger & Samulski, 2005, "Adeno-associated virus as a gene therapy vector: Vector development, production and clinical applications," *Adv. Biochem. Engin/Biotechnol.* 99: 119-145; Buning et al., 2008, "Recent developments in adeno-associated virus vector technology," *J. Gene Med.* 10:717-733; and the references cited below, each of which is incorporated herein by reference in its entirety. For packaging a transgene into virions, the ITRs are the only AAV components required in cis in the same construct as the nucleic acid molecule containing the expression cassettes. The cap and rep genes can be supplied in trans.

In one embodiment, the expression cassettes described herein are engineered into a genetic element (e.g., a shuttle plasmid) which transfers the immunoglobulin construct sequences carried thereon into a packaging host cell for production a viral vector. In one embodiment, the selected genetic element may be delivered to an AAV packaging cell by any
5 suitable method, including transfection, electroporation, liposome delivery, membrane fusion techniques, high velocity DNA-coated pellets, viral infection and protoplast fusion. Stable AAV packaging cells can also be made. Alternatively, the expression cassettes may be used to generate a viral vector other than AAV, or for production of mixtures of antibodies in vitro. The methods used to make such constructs are known to those with skill in nucleic acid
10 manipulation and include genetic engineering, recombinant engineering, and synthetic techniques. See, e.g., *Molecular Cloning: A Laboratory Manual*, ed. Green and Sambrook, Cold Spring Harbor Press, Cold Spring Harbor, NY (2012).

The term “AAV intermediate” or “AAV vector intermediate” refers to an assembled rAAV capsid which lacks the desired genomic sequences packaged therein. These may also be
15 termed an “empty” capsid. Such a capsid may contain no detectable genomic sequences of an expression cassette, or only partially packaged genomic sequences which are insufficient to achieve expression of the gene product. These empty capsids are non-functional to transfer the gene of interest to a host cell.

The recombinant AAV described herein may be generated using techniques which are
20 known. See, e.g., WO 2003/042397; WO 2005/033321, WO 2006/110689; US 7588772 B2. Such a method involves culturing a host cell which contains a nucleic acid sequence encoding an AAV capsid; a functional rep gene; an expression cassette composed of, at a minimum, AAV inverted terminal repeats (ITRs) and a transgene; and sufficient helper functions to permit packaging of the expression cassette into the AAV capsid protein. Methods of generating the
25 capsid, coding sequences therefore, and methods for production of rAAV viral vectors have been described. See, e.g., Gao, et al, *Proc. Natl. Acad. Sci. U.S.A.* 100 (10), 6081-6086 (2003) and US 2013/0045186A1.

In one embodiment, cells are manufactured in a suitable cell culture (e.g., HEK 293 cells). Methods for manufacturing the gene therapy vectors described herein include methods
30 well known in the art such as generation of plasmid DNA used for production of the gene therapy vectors, generation of the vectors, and purification of the vectors. In some embodiments, the gene therapy vector is an AAV vector and the plasmids generated are an

AAV cis-plasmid encoding the AAV genome and the gene of interest for packaging into the capsid, an AAV trans-plasmid containing AAV rep and cap genes, and an adenovirus helper plasmid. The vector generation process can include method steps such as initiation of cell culture, passage of cells, seeding of cells, transfection of cells with the plasmid DNA, post-
5 transfection medium exchange to serum free medium, and the harvest of vector-containing cells and culture media. The harvested vector-containing cells and culture media are referred to herein as crude cell harvest. In yet another system, the gene therapy vectors are introduced into insect cells by infection with baculovirus-based vectors. For reviews on these production systems, see generally, e.g., Zhang et al., 2009, "Adenovirus-adenovirus hybrid for
10 large-scale recombinant adeno-associated virus production," Human Gene Therapy 20:922-929, which is incorporated herein by reference in its entirety. Methods of making and using these and other AAV production systems are also described in the following U.S. patents, the contents of each of which is incorporated herein by reference in its entirety: 5,139,941; 5,741,683; 6,057,152; 6,204,059; 6,268,213; 6,491,907; 6,660,514; 6,951,753; 7,094,604; 7,172,893;
15 7,201,898; 7,229,823; and 7,439,065. In certain embodiments, the methods of making and using AAV production systems includes that of which uses pseudorabies viruses (rPRV) described in US Patent Application 63/016,894 filed on April 28, 2020, incorporated herein by reference.

The crude cell harvest may thereafter be subject method steps such as concentration of the vector harvest, diafiltration of the vector harvest, microfluidization of the vector harvest,
20 nuclease digestion of the vector harvest, filtration of microfluidized intermediate, crude purification by chromatography, crude purification by ultracentrifugation, buffer exchange by tangential flow filtration, and/or formulation and filtration to prepare bulk vector.

A two-step affinity chromatography purification at high salt concentration followed anion exchange resin chromatography are used to purify the vector drug product and to remove
25 empty capsids. These methods are described in more detail in International Patent Application No. PCT/US2016/065970, filed December 9, 2016, entitled "Scalable Purification Method for AAV9", which is incorporated by reference. Purification methods for AAV8, International Patent Application No. PCT/US2016/065976, filed December 9, 2016, and rh10, International Patent Application No. PCT/US16/66013, filed December 9, 2016, entitled "Scalable
30 Purification Method for AAVrh10", also filed December 11, 2015, and for AAV1, International Patent Application No. PCT/US2016/065974, filed December 9, 2016 for "Scalable Purification Method for AAV1", filed December 11, 2015, are all incorporated by reference herein.

To calculate empty and full particle content, vp3 band volumes for a selected sample (e.g., in examples herein an iodixanol gradient-purified preparation where number of GC = number of particles) are plotted against GC particles loaded. The resulting linear equation ($y = mx + c$) is used to calculate the number of particles in the band volumes of the test article peaks.

- 5 The number of particles (pt) per 20 μ L loaded is then multiplied by 50 to give particles (pt) /mL. Pt/mL divided by GC/mL gives the ratio of particles to genome copies (pt/GC). Pt/mL–GC/mL gives empty pt/mL. Empty pt/mL divided by pt/mL and x 100 gives the percentage of empty particles.

- Generally, methods for assaying for empty capsids and AAV vector particles with packaged genomes have been known in the art. See, e.g., Grimm et al., Gene Therapy (1999) 6:1322-1330; and Sommer et al., Molec. Ther. (2003) 7:122-128. To test for denatured capsid, the methods include subjecting the treated AAV stock to SDS-polyacrylamide gel electrophoresis, consisting of any gel capable of separating the three capsid proteins, for example, a gradient gel containing 3-8% Tris-acetate in the buffer, then running the gel until
- 15 sample material is separated, and blotting the gel onto nylon or nitrocellulose membranes, preferably nylon. Anti-AAV capsid antibodies are then used as the primary antibodies that bind to denatured capsid proteins, preferably an anti-AAV capsid monoclonal antibody, most preferably the B1 anti-AAV-2 monoclonal antibody (Wobus et al., J. Virol. (2000) 74:9281-9293). A secondary antibody is then used, one that binds to the primary antibody and contains a
- 20 means for detecting binding with the primary antibody, more preferably an anti-IgG antibody containing a detection molecule covalently bound to it, most preferably a sheep anti-mouse IgG antibody covalently linked to horseradish peroxidase. A method for detecting binding is used to semi-quantitatively determine binding between the primary and secondary antibodies, preferably a detection method capable of detecting radioactive isotope emissions,
- 25 electromagnetic radiation, or colorimetric changes, most preferably a chemiluminescence detection kit. For example, for SDS-PAGE, samples from column fractions can be taken and heated in SDS-PAGE loading buffer containing reducing agent (e.g., DTT), and capsid proteins were resolved on pre-cast gradient polyacrylamide gels (e.g., Novex). Silver staining may be performed using SilverXpress (Invitrogen, CA) according to the manufacturer's instructions or
- 30 other suitable staining method, i.e., SYPRO ruby or coomassie stains. In one embodiment, the concentration of AAV vector genomes (vg) in column fractions can be measured by quantitative real time PCR (Q-PCR). Samples are diluted and digested with DNase I (or another suitable

nuclease) to remove exogenous DNA. After inactivation of the nuclease, the samples are further diluted and amplified using primers and a TaqManTM fluorogenic probe specific for the DNA sequence between the primers. The number of cycles required to reach a defined level of fluorescence (threshold cycle, Ct) is measured for each sample on an Applied Biosystems Prism 5 7700 Sequence Detection System. Plasmid DNA containing identical sequences to that contained in the AAV vector is employed to generate a standard curve in the Q-PCR reaction. The cycle threshold (Ct) values obtained from the samples are used to determine vector genome titer by normalizing it to the Ct value of the plasmid standard curve. End-point assays based on the digital PCR can also be used.

10 Additionally, another example of measuring empty to full particle ratio is also known in the art. Sedimentation velocity, as measured in an analytical ultracentrifuge (AUC) can detect aggregates, other minor components as well as providing good quantitation of relative amounts of different particle species based upon their different sedimentation coefficients. This is an absolute method based on fundamental units of length and time, requiring no standard
15 molecules as references. Vector samples are loaded into cells with 2-channel charcoal-epon centerpieces with 12mm optical path length. The supplied dilution buffer is loaded into the reference channel of each cell. The loaded cells are then placed into an AN-60Ti analytical rotor and loaded into a Beckman-Coulter ProteomeLab XL-I analytical ultracentrifuge equipped with both absorbance and RI detectors. After full temperature equilibration at 20 °C the rotor is
20 brought to the final run speed of 12,000 rpm. A₂₈₀ scans are recorded approximately every 3 minutes for ~5.5 hours (110 total scans for each sample). The raw data is analyzed using the c(s) method and implemented in the analysis program SEDFIT. The resultant size distributions are graphed and the peaks integrated. The percentage values associated with each peak represent the peak area fraction of the total area under all peaks and are based upon the raw data generated at
25 280nm; many labs use these values to calculate empty: full particle ratios. However, because empty and full particles have different extinction coefficients at this wavelength, the raw data can be adjusted accordingly. The ratio of the empty particle and full monomer peak values both before and after extinction coefficient-adjustment is used to determine the empty-full particle ratio.

30 In one aspect, an optimized q-PCR method is used which utilizes a broad spectrum serine protease, e.g., proteinase K (such as is commercially available from Qiagen). More particularly, the optimized qPCR genome titer assay is similar to a standard assay, except that

after the DNase I digestion, samples are diluted with proteinase K buffer and treated with proteinase K followed by heat inactivation. Suitably samples are diluted with proteinase K buffer in an amount equal to the sample size. The proteinase K buffer may be concentrated to 2-fold or higher. Typically, proteinase K treatment is about 0.2 mg/mL, but may be varied from 0.1 mg/mL to about 1 mg/mL. The treatment step is generally conducted at about 55 °C for about 15 minutes, but may be performed at a lower temperature (e.g., about 37 °C to about 50 °C) over a longer time period (e.g., about 20 minutes to about 30 minutes), or a higher temperature (e.g., up to about 60 °C) for a shorter time period (e.g., about 5 to 10 minutes). Similarly, heat inactivation is generally at about 95 °C for about 15 minutes, but the temperature may be lowered (e.g., about 70 to about 90 °C) and the time extended (e.g., about 20 minutes to about 30 minutes). Samples are then diluted (e.g., 1000-fold) and subjected to TaqMan analysis as described in the standard assay. Quantification also can be done using ViroCyt or flow cytometry.

Additionally, or alternatively, droplet digital PCR (ddPCR) may be used. For example, methods for determining single-stranded and self-complementary AAV vector genome titers by ddPCR have been described. See, e.g., M. Lock et al, *Hu Gene Therapy Methods*, Hum. Gene Ther. Methods. 2014 Apr;25(2):115-25. doi: 10.1089/hgtb.2013.131. Epub 2014 Feb 14.

Therapeutic Proteins and Delivery Systems

Fusion partners, conjugate partners and recombinant vectors containing the targeting motif provided herein, N- x- (T/I/V/A)- (K/R) motif, are useful with a variety of different therapeutic proteins, polypeptides, nanoparticles, and delivery systems. Examples of proteins and compounds useful in compositions provided herein and targeted delivery include the following. It will be understood that the viral vectors, nanoparticles and other delivery systems contain sequences encoding the selected proteins (or conjugates) for expression in vivo.

In certain embodiments, the protein is MCT8 protein (SLC16A2 gene) and other compounds for treating of Allan-Herndon-Dudley disease and the symptoms thereof.

In certain embodiments, the protein is selected from a disease associated with a transport defect such as, e.g., cystic fibrosis (a cystic fibrosis transmembrane regulator), alpha-1-antitrypsin (hereditary emphysema), FE (hereditary hemochromatosis), tyrosinase (oculocutaneous albinism), Protein C (protein C deficiency), Complement C inhibitor (type I hereditary angioedema), alpha-D-galactosidase (Fabry disease), beta hexosaminidase (Tay-

Sachs), sucrase-isomaltase (congenital sucrase-isomaltase deficiency), UDP-glucuronosyl-transferase (Crigler-Najjar type II), insulin receptor (diabetes mellitus), growth hormone receptor (laron syndrome), among others. Examples of other genes and proteins those associated with, e.g. spinal muscular atrophy (SMA, SMN1), Huntingdon's Disease, Rett

5 Syndrome (e.g., methyl-CpG-binding protein 2 (MeCP2); UniProtKB – P51608), Amyotrophic Lateral Sclerosis (ALS), Duchenne Type Muscular dystrophy, Friedrichs Ataxia (e.g., frataxin), ATXN2 associated with spinocerebellar ataxia type 2 (SCA2)/ALS; TDP-43 associated with ALS, progranulin (PRGN) (associated with non-Alzheimer's cerebral degenerations, including, frontotemporal dementia (FTD), progressive non-fluent aphasia (PNFA) and semantic

10 dementia), among others. See, e.g., www.orpha.net/consor/cgi-bin/Disease_Search_List.php; rarediseases.info.nih.gov/diseases. Further illustrative genes which may be delivered via the rAAV include, without limitation, glucose-6-phosphatase, associated with glycogen storage disease or deficiency type 1A (GSD1), phosphoenolpyruvate-carboxykinase (PEPCK), associated with PEPCK deficiency; cyclin-dependent kinase-like 5 (CDKL5), also known as

15 serine/threonine kinase 9 (STK9) associated with seizures and severe neurodevelopmental impairment; galactose-1 phosphate uridyl transferase, associated with galactosemia; phenylalanine hydroxylase (PAH), associated with phenylketonuria (PKU); gene products associated with Primary Hyperoxaluria Type 1 including Hydroxyacid Oxidase 1 (GO/HAO1) and AGXT, branched chain alpha-ketoacid dehydrogenase, including BCKDH, BCKDH-E2,

20 BAKDH-E1a, and BAKDH-E1b, associated with Maple syrup urine disease; fumarylacetoacetate hydrolase, associated with tyrosinemia type 1; methylmalonyl-CoA mutase, associated with methylmalonic acidemia; medium chain acyl CoA dehydrogenase, associated with medium chain acetyl CoA deficiency; ornithine transcarbamylase (OTC), associated with ornithine transcarbamylase deficiency; argininosuccinic acid synthetase (ASS1),

25 associated with citrullinemia; lecithin-cholesterol acyltransferase (LCAT) deficiency; methylmalonic acidemia (MMA); NPC1 associated with Niemann-Pick disease, type C1); propionic academia (PA); TTR associated with Transthyretin (TTR)-related Hereditary Amyloidosis; low density lipoprotein receptor (LDLR) protein, associated with familial hypercholesterolemia (FH), LDLR variant, such as those described in WO 2015/164778;

30 PCSK9; ApoE and ApoC proteins, associated with dementia; UDP-glucouronosyltransferase, associated with Crigler-Najjar disease; adenosine deaminase, associated with severe combined immunodeficiency disease; hypoxanthine guanine phosphoribosyl transferase, associated with

Gout and Lesch-Nyan syndrome; biotinidase, associated with biotinidase deficiency; alpha-galactosidase A (a-Gal A) associated with Fabry disease; beta-galactosidase (GLB1) associated with GM1 gangliosidosis; ATP7B associated with Wilson's Disease; beta-glucocerebrosidase, associated with Gaucher disease type 2 and 3; peroxisome membrane protein 70 kDa, associated with Zellweger syndrome; arylsulfatase A (ARSA) associated with metachromatic leukodystrophy, galactocerebrosidase (*GALC*) enzyme associated with Krabbe disease, alpha-glucosidase (GAA) associated with Pompe disease; sphingomyelinase (SMPD1) gene associated with Nieman Pick disease type A; argininosuccinate synthase associated with adult onset type II citrullinemia (CTLN2); carbamoyl-phosphate synthase 1 (CPS1) associated with urea cycle disorders; survival motor neuron (SMN) protein, associated with spinal muscular atrophy; ceramidase associated with Farber lipogranulomatosis; b-hexosaminidase associated with GM2 gangliosidosis and Tay-Sachs and Sandhoff diseases; aspartylglucosaminidase associated with aspartyl-glucosaminuria; α -fucosidase associated with fucosidosis; α -mannosidase associated with alpha-mannosidosis; porphobilinogen deaminase, associated with acute intermittent porphyria (AIP); alpha-1 antitrypsin for treatment of alpha-1 antitrypsin deficiency (emphysema); erythropoietin for treatment of anemia due to thalassemia or to renal failure; vascular endothelial growth factor, angiopoietin-1, and fibroblast growth factor for the treatment of ischemic diseases; thrombomodulin and tissue factor pathway inhibitor for the treatment of occluded blood vessels as seen in, for example, atherosclerosis, thrombosis, or embolisms; aromatic amino acid decarboxylase (AADC), and tyrosine hydroxylase (TH) for the treatment of Parkinson's disease.

Examples of proteins and compounds useful in compositions provided herein and targeted delivery include therapeutic proteins and other compounds and vaccine protein derivatives of the following respiratory-associated infectious diseases and passive immunoglobulins direct against these infectious disease. Examples of suitable therapeutic proteins include, e.g., alpha-1-antitrypsin, cystic fibrosis transmembrane protein, and variants thereof, surfactant-B, bone morphogenetic protein receptor type II (associated with pulmonary arterial hypertension), and various cancer therapeutics.

Examples of suitable vaccine or passive immunization include proteins derived from airborne pathogens, including the human respiratory coronaviruses, have been associated with severe acute respiratory syndrome (SARS-CoV1), the common cold, and non A, B or C hepatitis. SARS-CoV2 is the causative agent of COVID-19 and antibodies specific for this

virus have been described. Examples of IgG antibodies which have been described as being useful for binding the spike protein of human ACE2 of SARS-CoV2 and having neutralizing activity include, e.g., LY-CoV555 (Eli Lilly), TY027 (Tychon), STI-1499 and STI-2020 (COVI-GUARD; Sorrento), 80R, ADI055689/56046 (Adimab) (Renn et al., Trends in Pharmacological Sciences, 2020); BD-217, BD-218, BD-236 (Cao et al., Cell, 182, 73-84 (2020)). Examples of IgG antibodies which have been described as being useful for binding the receptor binding domain (RBD) of human ACE2 of SARS-COV2 and having neutralizing activity include, e.g., COV2-2196, COV2-2130, COV2-2165 (Zost et al., Nature, 584, 443-465 (2020)); BD-361, BD-368, BD-368-2 (Cao et al., Cell, 182, 73-84 (2020)); B38, H4 (Y. Wu et al., Science 10.1126/science.abc2241 (2020); Jahanshahlu and Rezaei, Biomedicine and Pharmacotherapy 129 (2020)); S309, S315, S304 (Pinto et al., Nature, 583, 290-311 (2020)); CC6.29, CC6.30, CC6.33, CC12.1, CC12.3 (Rogers et al., Science 369, 956-963 (2020)); JS016 (Eli Lilly), CA1, CB6-LALA, P2C-1F11/P2B-2F6/P2A-1A3, 311mab-31B5311/32D4, COVA 2-15, 414-1, (Renn et al., Trends in Pharmacological Sciences, 2020). Examples of IgG antibodies which have been described as being useful for binding the spike protein of human ACE2 of SARS-COV1 and having neutralizing activity include, e.g., m396 and CR3104 (Prabakaran et al., Journal of Biological Chemistry, 281, 15829-15836 (2006); ter Meulen et al., PLoS, 3, 7 (2006)). Examples of IgG antibodies which have been described as being useful for binding either RBD or spike protein of human ACE2 of both SARS-COV1 and SARS-CoV2 and having neutralizing activity include, e.g., CR3022 and 47D11 (Wang et al., Nature Communications, 11, nature.com/naturecommunications (2020)) .

Examples of other target viruses include influenza virus from the orthomyxoviridae family, which includes: Influenza A, Influenza B, and Influenza C. The type A viruses are the most virulent human pathogens. The serotypes of influenza A which have been associated with pandemics include, H1N1, which caused Spanish Flu in 1918, and Swine Flu in 2009; H2N2, which caused Asian Flu in 1957; H3N2, which caused Hong Kong Flu in 1968; H5N1, which caused Bird Flu in 2004; H7N7; H1N2; H9N2; H7N2; H7N3; and H10N7. Broadly neutralizing antibodies against influenza A have been described. As used herein, a "broadly neutralizing antibody" refers to a neutralizing antibody which can neutralize multiple strains from multiple subtypes. For example, CR6261 [The Scripps Institute/ Crucell] has been described as a monoclonal antibody that binds to a broad range of the influenza virus including the 1918 "Spanish flu" (SC1918/H1) and to a virus of the H5N1 class of avian influenza that jumped

from chickens to a human in Vietnam in 2004 (Viet04/H5). CR6261 recognizes a highly conserved helical region in the membrane-proximal stem of hemagglutinin, the predominant protein on the surface of the influenza virus. This antibody is described in WO 2010/130636, incorporated by reference herein. Another neutralizing antibody, F10 [XOMA Ltd] has been
5 described as being useful against H1N1 and H5N1. [Sui et al, Nature Structural and Molecular Biology (Sui, et al. 2009, 16(3):265-73)] Other antibodies against influenza, *e.g.*, Fab28 and Fab49, may be selected. See, *e.g.*, WO 2010/140114 and WO 2009/115972, which are incorporated by reference. Still other antibodies, such as those described in WO 2010/010466, US Published Patent Publication US/2011/076265, and WO 2008/156763, may be readily
10 selected.

Other target pathogenic viruses include, arenaviruses (including funin, machupo, and Lassa), filoviruses (including Marburg and Ebola), hantaviruses, picornoviridae (including rhinoviruses, echovirus), coronaviruses, paramyxovirus, morbillivirus, respiratory syncytial virus, togavirus, coxsackievirus, parvovirus B19, parainfluenza, adenoviruses, reoviruses,
15 variola (*Variola major* (Smallpox)) and *Vaccinia* (Cowpox) from the poxvirus family, and varicella-zoster (pseudorabies). Viral hemorrhagic fevers are caused by members of the arenavirus family (Lassa fever) (which family is also associated with Lymphocytic choriomeningitis (LCM)), filovirus (ebola virus), and hantavirus (puremala). The members of picornavirus (a subfamily of rhinoviruses), are associated with the common cold in humans.
20 The coronavirus family, which includes a number of non-human viruses such as infectious bronchitis virus (poultry), porcine transmissible gastroenteric virus (pig), porcine hemagglutinin encephalomyelitis virus (pig), feline infectious peritonitis virus (cat), feline enteric coronavirus (cat), canine coronavirus (dog). The paramyxovirus family includes parainfluenza Virus Type 1, parainfluenza Virus Type 3, bovine parainfluenza Virus Type 3,
25 rubelavirus (mumps virus, parainfluenza Virus Type 2, parainfluenza virus Type 4, Newcastle disease virus (chickens), rinderpest, morbillivirus, which includes measles and canine distemper, and pneumovirus, which includes respiratory syncytial virus (RSV). The parvovirus family includes feline parvovirus (feline enteritis), feline panleucopeniavirus, canine parvovirus, and porcine parvovirus. The adenovirus family includes viruses (EX, AD7, ARD, O.B.) which
30 cause respiratory disease.

A neutralizing antibody construct against a bacterial pathogen may also be selected for use in the present invention. In one embodiment, the neutralizing antibody construct is directed

against the bacteria itself. In another embodiment, the neutralizing antibody construct is directed against a toxin produced by the bacteria. Examples of airborne bacterial pathogens include, e.g., *Neisseria meningitidis* (meningitis), *Klebsiella pneumonia* (pneumonia), *Pseudomonas aeruginosa* (pneumonia), *Pseudomonas pseudomallei* (pneumonia),

5 *Pseudomonas mallei* (pneumonia), *Acinetobacter* (pneumonia), *Moraxella catarrhalis*, *Moraxella lacunata*, *Alkaligenes*, *Cardiobacterium*, *Haemophilus influenzae* (flu), *Haemophilus parainfluenzae*, *Bordetella pertussis* (whooping cough), *Francisella tularensis* (pneumonia/fever), *Legionella pneumonia* (Legionnaires disease), *Chlamydia psittaci* (pneumonia), *Chlamydia pneumoniae* (pneumonia), *Mycobacterium tuberculosis* (tuberculosis

10 (TB)), *Mycobacterium kansasii* (TB), *Mycobacterium avium* (pneumonia), *Nocardia asteroides* (pneumonia), *Bacillus anthracis* (anthrax), *Staphylococcus aureus* (pneumonia), *Streptococcus pyogenes* (scarlet fever), *Streptococcus pneumoniae* (pneumonia), *Corynebacteria diphtheria* (diphtheria), *Mycoplasma pneumoniae* (pneumonia). The causative agent of anthrax is a toxin produced by *Bacillus anthracis*. Neutralizing antibodies against protective agent (PA), one of

15 the three peptides which form the toxoid, have been described. The other two polypeptides consist of lethal factor (LF) and edema factor (EF). Anti-PA neutralizing antibodies have been described as being effective in passively immunization against anthrax. See, e.g., US Patent number 7,442,373; R. Sawada-Hirai et al, J Immune Based Ther Vaccines. 2004; 2: 5. (on-line 2004 May 12). Still other anti-anthrax toxin neutralizing antibodies have been described and/or

20 may be generated. Similarly, neutralizing antibodies against other bacteria and/or bacterial toxins may be used to generate a non-IgG antibody as described herein.

Other infectious diseases may be caused by airborne fungi including, e.g., *Aspergillus* species, *Absidia corymbifera*, *Rhizopus stolonifer*, *Mucor plumbeus*, *Cryptococcus neoformans*, *Histoplasma capsulatum*, *Blastomyces dermatitidis*, *Coccidioides immitis*, *Penicillium* species,

25 *Micropolyspora faeni*, *Thermoactinomyces vulgaris*, *Alternaria alternate*, *Cladosporium* species, *Helminthosporium*, and *Stachybotrys* species.

In addition to airborne infectious disease conditions which affect humans, many of which are described above, passive immunization according to the invention may be used to prevent conditions associated with direct inoculation of the nasal passages, e.g., conditions

30 which may be transmitted by direct contact of the fingers with the nasal passages. These conditions may include fungal infections (e.g., athlete's foot), ringworm, or viruses, bacteria, parasites, fungi, and other pathogens which can be transmitted by direct contact. In addition, a

variety of conditions which affect household pets, cattle and other livestock, and other animals. For example, in dogs, infection of the upper respiratory tract by canine sinonasal aspergillosis causes significant disease. In cats, upper respiratory disease or feline respiratory disease complex originating in the nose causes morbidity and mortality if left untreated. Cattle are prone to infections by the infectious bovine rhinotracheitis (commonly called IBR or red nose) is an acute, contagious virus disease of cattle. In addition, cattle are prone to Bovine Respiratory Syncytial Virus (BRSV) which causes mild to severe respiratory disease and can impair resistance to other diseases. Still other pathogens and diseases will be apparent to one of skill in the art.

10 An antibody, and particularly, a neutralizing antibody, against a pathogen such as those specifically identified herein (e.g., anti-SARS-CoV2, anti-SARS-CoV1, anti-influenza, anti-ebola, anti-RSV), may be used to generate a class-switched or non-IgG antibody. Monoclonal antibodies (mAbs) with broad neutralizing capacity can be identified using antibody phage display to screen libraries from donors recently vaccinated with the seasonal flu vaccine, from 15 non-immune humans or from survivors of a natural infection. In the case of influenza, antibodies have been identified which neutralize more than one influenza subtype by blocking viral fusion with the host cell. This technique may be utilized with other infections to obtain a neutralizing monoclonal antibody. See, e.g., US 5,811,524, which describes generation of anti-respiratory syncytial virus (RSV) neutralizing antibodies. The techniques described therein are 20 applicable to other pathogens. Such an antibody may be used intact, or its sequences (scaffold) modified to generate an artificial or recombinant neutralizing antibody construct. Such methods have been described [see, e.g., WO 2010/13036; WO 2009/115972; WO 2010/140114]. In one embodiment, mouse, rat, hamster or other host animals, is immunized with an immunizing agent to generate lymphocytes that produce antibodies with binding specificity to the 25 immunizing antigen. In an alternative approach, the lymphocytes may be immunized in vitro. Human antibodies can be produced using techniques such as phage display libraries (Hoogenboom and Winter, J. Mol. Biol. 1991, 227:381, Marks et al., J. Mol. Biol. 1991, 222:581).

30 Compositions and Uses

Provided herein are compositions containing at least one rAAV stock (e.g., an rAAV9 or rAAVhu68 mutant stock) and an optional carrier, excipient and/or preservative. An rAAV

stock refers to a plurality of rAAV vectors which are the same, e.g., such as in the amounts described below in the discussion of concentrations and dosage units.

In certain embodiments, a composition may contain at least a second, different rAAV stock. This second vector stock may vary from the first by having a different AAV capsid
5 and/or a different vector genome. In certain embodiments, a composition as described herein may contain a different vector expressing an expression cassette as described herein, or another active component (e.g., an antibody construct, another biologic, and/or a small molecule drug).

As used herein, "carrier" includes any and all solvents, dispersion media, vehicles, coatings, diluents, antibacterial and antifungal agents, isotonic and absorption delaying agents,
10 buffers, carrier solutions, suspensions, colloids, and the like. The use of such media and agents for pharmaceutical active substances is well known in the art. Supplementary active ingredients can also be incorporated into the compositions. The phrase "pharmaceutically-acceptable" refers to molecular entities and compositions that do not produce an allergic or similar untoward reaction when administered to a host. Delivery vehicles such as liposomes, nanocapsules,
15 microparticles, microspheres, lipid particles, vesicles, and the like, may be used for the introduction of the compositions of the present invention into suitable host cells. In particular, the rAAV vector delivered transgenes may be formulated for delivery either encapsulated in a lipid particle, a liposome, a vesicle, a nanosphere, or a nanoparticle or the like.

In one embodiment, a composition includes a final formulation suitable for delivery to a
20 subject, e.g., is an aqueous liquid suspension buffered to a physiologically compatible pH and salt concentration. Optionally, one or more surfactants are present in the formulation. In another embodiment, the composition may be transported as a concentrate which is diluted for administration to a subject. In other embodiments, the composition may be lyophilized and reconstituted at the time of administration.

25 A suitable surfactant, or combination of surfactants, may be selected from among non-ionic surfactants that are nontoxic. In one embodiment, a difunctional block copolymer surfactant terminating in primary hydroxyl groups is selected, e.g., such as Pluronic® F68 [BASF], also known as Poloxamer 188, which has a neutral pH, has an average molecular weight of 8400. Other surfactants and other Poloxamers may be selected, i.e., nonionic triblock
30 copolymers composed of a central hydrophobic chain of polyoxypropylene (poly(propylene oxide)) flanked by two hydrophilic chains of polyoxyethylene (poly(ethylene oxide)), SUTOL HS 15 (Macrogol-15 Hydroxystearate), LABRASOL (Polyoxy caprylic glyceride),

polyoxy 10 oleyl ether, TWEEN (polyoxyethylene sorbitan fatty acid esters), ethanol and polyethylene glycol. In one embodiment, the formulation contains a poloxamer. These copolymers are commonly named with the letter "P" (for poloxamer) followed by three digits: the first two digits x 100 give the approximate molecular mass of the polyoxypropylene core, and the last digit x 10 gives the percentage polyoxyethylene content. In one embodiment Poloxamer 188 is selected. The surfactant may be present in an amount up to about 0.0005 % to about 0.001% of the suspension.

In one embodiment, the formulation buffer is phosphate-buffered saline (PBS) with total salt concentration of 200 mM, 0.001% (w/v) pluronic F68 (Final Formulation Buffer, FFB).

The vectors are administered in sufficient amounts to transfect the cells and to provide sufficient levels of gene transfer and expression to provide a therapeutic benefit without undue adverse effects, or with medically acceptable physiological effects, which can be determined by those skilled in the medical arts. In certain embodiments, the vectors are formulated for delivery via intranasal delivery devices for targeted delivery to nasal and/or nasopharynx epithelial cells.

In certain embodiments, vectors are formulated for aerosol delivery devices, e.g., via a nebulizer or through other suitable devices. Other conventional and pharmaceutically acceptable routes of administration include, but are not limited to, direct delivery to a desired organ (e.g., lung), oral inhalation, intrathecal, intratracheal, intraarterial, intraocular, intravenous, intramuscular, subcutaneous, intradermal, and other parenteral routes of administration. In one embodiment, the vector is administered intranasally using intranasal mucosal atomization device (LMA[®] MAD Nasal[™]- MAD110). In another embodiment the vector is administered intrapulmonary in nebulized form using Vibrating Mesh Nebulizer (Aerogen[®] Solo) or MADgic[™] Laryngeal Mucosal Atomizer. Routes of administration may be combined, if desired. Routes of administration and utilization of which for delivering rAAV vectors are also described in the following published US Patent Applications, the contents of each of which is incorporated herein by reference in its entirety: US 2018/0155412A1, US 2018/0243416A1, US 2014/0031418 A1, and US 2019/0216841A1.

Dosages of the viral vector will depend primarily on factors such as the condition being treated, the age, weight and health of the patient, and may thus vary among patients. For example, a therapeutically effective human dosage of the viral vector is generally in the range of from about 25 to about 1000 microliters to about 5 mL of aqueous suspending liquid containing doses of from about 10^9 to 4×10^{14} GC of AAV vector. The dosage will be adjusted to balance

the therapeutic benefit against any side effects and such dosages may vary depending upon the therapeutic application for which the recombinant vector is employed. The levels of expression of the transgene can be monitored to determine the frequency of dosage resulting in viral vectors, preferably AAV vectors containing the minigene. Optionally, dosage regimens similar to those described for therapeutic purposes may be utilized for immunization using the compositions of the invention.

The replication-defective virus compositions can be formulated in dosage units to contain an amount of replication-defective virus that is in the range of about 10^9 GC to about 10^{16} GC (to treat an average subject of 70 kg in body weight) including all integers or fractional amounts within the range, and preferably 10^{12} GC to 10^{14} GC for a human patient. In one embodiment, the compositions are formulated to contain at least 10^9 , 2×10^9 , 3×10^9 , 4×10^9 , 5×10^9 , 6×10^9 , 7×10^9 , 8×10^9 , or 9×10^9 GC per dose including all integers or fractional amounts within the range. In another embodiment, the compositions are formulated to contain at least 10^{10} , 2×10^{10} , 3×10^{10} , 4×10^{10} , 5×10^{10} , 6×10^{10} , 7×10^{10} , 8×10^{10} , or 9×10^{10} GC per dose including all integers or fractional amounts within the range. In another embodiment, the compositions are formulated to contain at least 10^{11} , 2×10^{11} , 3×10^{11} , 4×10^{11} , 5×10^{11} , 6×10^{11} , 7×10^{11} , 8×10^{11} , or 9×10^{11} GC per dose including all integers or fractional amounts within the range. In another embodiment, the compositions are formulated to contain at least 10^{12} , 2×10^{12} , 3×10^{12} , 4×10^{12} , 5×10^{12} , 6×10^{12} , 7×10^{12} , 8×10^{12} , or 9×10^{12} GC per dose including all integers or fractional amounts within the range. In another embodiment, the compositions are formulated to contain at least 10^{13} , 2×10^{13} , 3×10^{13} , 4×10^{13} , 5×10^{13} , 6×10^{13} , 7×10^{13} , 8×10^{13} , or 9×10^{13} GC per dose including all integers or fractional amounts within the range. In another embodiment, the compositions are formulated to contain at least 10^{14} , 2×10^{14} , 3×10^{14} , 4×10^{14} , 5×10^{14} , 6×10^{14} , 7×10^{14} , 8×10^{14} , or 9×10^{14} GC per dose including all integers or fractional amounts within the range. In another embodiment, the compositions are formulated to contain at least 10^{15} , 2×10^{15} , 3×10^{15} , 4×10^{15} , 5×10^{15} , 6×10^{15} , 7×10^{15} , 8×10^{15} , or 9×10^{15} GC per dose including all integers or fractional amounts within the range. In one embodiment, for human application the dose can range from 10^{10} to about 10^{12} GC per dose including all integers or fractional amounts within the range. In one embodiment, for human application the dose can range from 10^9 to about 7×10^{13} GC per dose including all integers or fractional amounts within the range. In one embodiment, for human application the dose ranges from 6.25×10^{12} GC to 5.00×10^{13} GC. In a further embodiment, the dose is about 6.25×10^{12} GC, about 1.25×10^{13} GC, about 2.50×10^{13} GC,

or about 5.00×10^{13} GC. In certain embodiment, the dose is divided into one half thereof equally and administered to each nostril. In certain embodiments, for human application the dose ranges from 6.25×10^{12} GC to 5.00×10^{13} GC administered as two aliquots of 0.2 ml per nostril for a total volume delivered in each subject of 0.8ml.

5 These above doses may be administered in a variety of volumes of carrier, excipient or buffer formulation, ranging from about 25 to about 1000 microliters, or higher volumes, including all numbers within the range, depending on the size of the area to be treated, the viral titer used, the route of administration, and the desired effect of the method. In one embodiment, the volume of carrier, excipient or buffer is at least about 25 μ L. In one embodiment, the
10 volume is about 50 μ L. In another embodiment, the volume is about 75 μ L. In another embodiment, the volume is about 100 μ L. In another embodiment, the volume is about 125 μ L. In another embodiment, the volume is about 150 μ L. In another embodiment, the volume is about 175 μ L. In yet another embodiment, the volume is about 200 μ L. In another embodiment, the volume is about 225 μ L. In yet another embodiment, the volume is about 250 μ L. In yet
15 another embodiment, the volume is about 275 μ L. In yet another embodiment, the volume is about 300 μ L. In yet another embodiment, the volume is about 325 μ L. In another embodiment, the volume is about 350 μ L. In another embodiment, the volume is about 375 μ L. In another embodiment, the volume is about 400 μ L. In another embodiment, the volume is about 450 μ L. In another embodiment, the volume is about 500 μ L. In another embodiment, the volume is
20 about 550 μ L. In another embodiment, the volume is about 600 μ L. In another embodiment, the volume is about 650 μ L. In another embodiment, the volume is about 700 μ L. In another embodiment, the volume is between about 700 and 1000 μ L.

 In certain embodiments, the recombinant vectors may be dosed intranasally by using two sprays to each nostril. In one embodiment, the two sprays are administered by alternating to
25 each nostril, e.g., left nostril spray, right nostril spray, then left nostril spray, right nostril spray. In certain embodiments, there may be a delay between alternating sprays. For example, each nostril may receive multiple sprays which are separated by an interval of about 10 to 60 seconds, or 20 to 40 seconds, or about 30 seconds, to a few minutes, or longer. Such sprays may deliver, e.g., about 150 μ L to 300 μ L, or about 250 μ L in each spray, to achieve a total volume
30 dosed of about 200 μ L to about 600 μ L, 400 μ L to 700 μ L, or 450 μ L to 1000 μ L.

 In certain embodiment, the recombinant AAV vector may be dosed intranasally to achieve a concentration of 5-20 ng/ml of the expression product of the transgene as measured in

a nasal wash solution post-dosing, e.g., one week to four weeks, or about two weeks after administration of the vector. Methods of acquiring the nasal wash solution in the subjected as well as methods of quantification of the expression product of the transgene are conventional.

For other routes of administration, e.g., intravenous or intramuscular, dose levels would
5 be higher than for intranasal delivery. For example, such suspensions may be volumes doses of about 1 mL to about 25 mL, with doses of up to about 2.5×10^{15} GC.

In certain embodiments, the intranasal delivery device provides a spray atomizer which delivers a mist of particles having an average size range of about 30 microns to about 100 microns in size. In certain embodiments, the average size range is about 10 microns to about 50
10 microns. Suitable devices have been described in the literature and some are commercially available, e.g., the LMA MAD NASAL™ (Teleflex Medical; Ireland); Teleflex VaxInator™ (Teleflex Medical; Ireland); Controlled Particle Dispersion® (CPD) from Kurve Technologies. See, also, PG Djupesland, Drug Deliv and Transl. Res (2013) 3: 42-62. In certain
15 embodiments, the particle size and volume of delivery is controlled in order to preferentially target nasal epithelial cells and minimize targeting to the lung. In other embodiments, the mist of particles is about 0.1 micron to about 20 microns, or less, in order to deliver to lung cells. Such smaller particle sizes may minimize retention in the nasal epithelium.

One device mists particles at an average diameter of about 16 microns to about 22 microns. The mist may be delivered directly to the tracheobronchial tree inserted through the
20 suction channel of a 3.5-mm flexible fiberoptic bronchoscope (Olympus, Melville, NY). Other suitable delivery devices may include a laryngo-tracheal mucosal atomizer, which provides for administration across the upper airway past the vocal cords. It fits through vocal cords and down a laryngeal mask or into nasal cavity. The droplets are atomized at an average diameter of about 30 microns to about 100 microns. A standard device has a tip diameter of about 0.18
25 in (4.6 mm), a length of about 4.5-8.5 inches, and is inserted through the suction channel and advanced approximately 3 mm beyond the distal tip of the scope. Doses may be administered is 10 aliquots (approximately 150 µl each) of control with saline or rAAV sprayed into right and left main stem bronchi.

In one embodiment, a frozen composition is provided which contains an rAAV in a
30 buffer solution as described herein, in frozen form. Optionally, one or more surfactants (e.g., Pluronic F68), stabilizers or preservatives is present in this composition. Suitably, for use, a

composition is thawed and titrated to the desired dose with a suitable diluent, e.g., sterile saline or a buffered saline.

In one embodiment, a composition comprising one or more exogenous endothelial cell
5 targeting peptide from the motif: N- x- (T/I/V/A)- (K/R) (SEQ ID NO: 47) and optional
flanking linker sequences are provided, together with one or more of a physiologically
compatible carrier, excipient, and/or aqueous suspension base. Further provided are
compositions comprising nucleic acid sequences encoding same. In certain embodiments, the
targeting peptide is of SEQ ID NO: 40 and is encoded by a nucleic acid sequence of SEQ ID
10 NO: 54, or a sequence at least about 70% identical thereto. In certain embodiments, the
targeting peptide is of SEQ ID NO: 38 and is encoded by a nucleic acid sequence of SEQ ID
NO: 50, or a sequence at least about 70% identical thereto. In certain embodiments, the
targeting peptide is of SEQ ID NO: 46 and is encoded by a nucleic acid sequence of SEQ ID
NO: 56, or a sequence at least about 70% identical thereto. In certain embodiments, the
15 targeting peptide is of SEQ ID NO: 43 and is encoded by a nucleic acid sequence of SEQ ID
NO: 52, or a sequence at least about 70% identical thereto. In certain embodiments, the
targeting peptide is of SEQ ID NO: 39 and is encoded by a nucleic acid sequence of SEQ ID
NO: 55, or a sequence at least about 70% identical thereto. In certain embodiments, the
targeting peptide is of SEQ ID NO: 42 and is encoded by a nucleic acid sequence of SEQ ID
20 NO: 51, or a sequence at least about 70% identical thereto. In certain embodiments, the
targeting peptide is of SEQ ID NO: 41 and is encoded by a nucleic acid sequence of SEQ ID
NO: 53, or a sequence at least about 70% identical thereto.

In another embodiment, a fusion polypeptide or protein is provided comprising one or
more exogenous brain endothelial cell targeting peptide from the motif: N- x- (T/I/V/A)- (K/R)
25 (SEQ ID NO: 47) are provided and fusion partner which comprises at least one polypeptide or
protein. Further provided are nucleic acid sequences encoding same.

In certain embodiments, a composition comprising a fusion polypeptide or protein, or a
nucleic acid sequence encoding the fusion polypeptide or protein, or a nanoparticle containing
same are provided. The composition may further comprise one or more of a physiologically
30 compatible carrier, excipient, and/or aqueous suspension base.

In certain embodiments, a nucleic acid sequence encoding the fusion polypeptide
protein is encapsulated in a lipid nanoparticle (LNP). As used herein, the phrase "lipid

nanoparticle" or "nanoparticle" refers to a transfer vehicle comprising one or more lipids (e.g., cationic lipids, non- cationic lipids, and PEG-modified lipids). Preferably, the lipid nanoparticles are formulated to deliver one or more nucleic acid sequences to one or more target cells (e.g., liver and/or muscle). Examples of suitable lipids include, for example, the

5 phosphatidyl compounds (e.g., phosphatidylglycerol, phosphatidylcholine, phosphatidylserine, phosphatidylethanolamine, sphingolipids, cerebrosides, and gangliosides). Also contemplated is the use of polymers as transfer vehicles, whether alone or in combination with other transfer vehicles. Suitable polymers may include, for example, polyacrylates, polyalkycyanoacrylates, polylactide, polylactide- polyglycolide copolymers, polycaprolactones, dextran, albumin,

10 gelatin, alginate, collagen, chitosan, cyclodextrins, dendrimers and polyethylenimine. In one embodiment, the transfer vehicle is selected based upon its ability to facilitate the transfection of a nucleic acid sequence encapsulated therein to a target cell. Useful lipid nanoparticles for nucleic acid sequence comprise a cationic lipid to encapsulate and/or enhance the delivery of such nucleic acid sequence into the target cell that will act as a depot for protein production. As

15 used herein, the phrase "cationic lipid" refers to any of a number of lipid species that carry a net positive charge at a selected pH, such as physiological pH. The contemplated lipid nanoparticles may be prepared by including multi-component lipid mixtures of varying ratios employing one or more cationic lipids, non-cationic lipids and PEG- modified lipids. Several cationic lipids have been described in the literature, many of which are commercially available. See, e.g.,

20 WO2014/089486, US 2018/0353616A1, and US 8,853,377B2, which are incorporated by reference. In certain embodiments, LNP formulation is performed using routine procedures comprising cholesterol, ionizable lipid, helper lipid, PEG-lipid and polymer forming a lipid bilayer around encapsulated nucleic acid sequence (Kowalski et al., 2019, Mol. Ther. 27(4):710-728). In some embodiments, LNP comprises a cationic lipids (i.e. N-[1-(2,3-

25 dioleoyloxy)propyl]-N,N,N-trimethylammonium chloride (DOTMA), or 1,2-dioleoyl-3-trimethylammonium-propane (DOTAP)) with helper lipid DOPE. In some embodiments, LNP comprises an ionizable lipid Dlin-MC3-DMA ionizable lipids, or diketopiperazine-based ionizable lipids (cKK-E12). In some embodiments, polymer comprises a polyethyleneimine (PEI), or a poly(β -amino)esters (PBAEs). See, e.g., WO2014/089486, US 2018/0353616A1,

30 US2013/0037977A1, WO2015/074085A1, US9670152B2, and US 8,853,377B2, which are incorporated by reference.

In certain embodiments, a composition, e.g., an rAAV having a modified capsid with a N- x- (T/I/V/A)- (K/R) (SEQ ID NO: 47) peptide and optional linker sequences, a fusion polypeptide or protein, or a conjugate comprising a nanoparticle or chemical moiety, is useful for delivering a therapeutic to a patient in need thereof. In certain embodiments, the method is for targeting therapy to the brain endothelial cells. In certain embodiments, the method is for treating Allan-Herndon-Dudley disease by delivering an MCT8 protein (e.g., UniProt ID No.: P36021) or a gene which expresses MCT8 in vivo. In other embodiments, the method is for targeting therapy to the lung. In certain embodiments, the delivered product is a soluble Ace2 protein (e.g., hAce2 decoy or hAce2 decoy fusion), an anti-SARS antibody, an anti-SARS-CoV2 antibody, an anti-influenza antibody, or a cystic fibrosis transmembrane protein. See also, [omim.org/entry/300523](https://www.ncbi.nlm.nih.gov/omim.org/entry/300523), the content of which is incorporated herein by reference. See also, US Provisional Application No. 63/143,614, filed January 29, 2021, US Provisional Application No. 63/165,051, filed March 12, 2021, and US Patent Application No. 63/166,686, filed March 26, 2021, US Provisional Patent Application No. 63/215,159, filed June 25, 2021, US Provisional Application No. 63/253,654, filed October 8, 2021 are incorporated herein by reference.

In certain embodiments, a rAAV having a modified capsid as described herein may be delivered in a co-therapeutic regimen which further comprises one or more other active components. In certain embodiments, the regimen may involve co-administration of an immunomodulatory component. Such an immunomodulatory regimen may include, e.g., but are not limited to immunosuppressants such as, a glucocorticoid, steroids, antimetabolites, T-cell inhibitors, a macrolide (e.g., a rapamycin or rapalog), and cytostatic agents including an alkylating agent, an anti-metabolite, a cytotoxic antibiotic, an antibody, or an agent active on immunophilin. The immune suppressant may include a nitrogen mustard, nitrosourea, platinum compound, methotrexate, azathioprine, mercaptopurine, fluorouracil, dactinomycin, an anthracycline, mitomycin C, bleomycin, mithramycin, IL-2 receptor- (CD25-) or CD3-directed antibodies, anti-IL-2 antibodies, cyclosporin, tacrolimus, sirolimus, IFN- β , IFN- γ , an opioid, or TNF- α (tumor necrosis factor-alpha) binding agent. In certain embodiments, the immunosuppressive therapy may be started prior to the gene therapy administration. Such therapy may involve co-administration of two or more drugs, the (e.g., prednisone, micophenolate mofetil (MMF) and/or sirolimus (i.e., rapamycin)) on the same day. One or more of these drugs may be continued after gene therapy administration, at the same dose or an

adjusted dose. Such therapy may be for about 1 week, about 15 days, about 30 days, about 45 days, 60 days, or longer, as needed. Still other co-therapeutics may include, e.g., anti-IgG enzymes, which have been described as being useful for depleting anti-AAV antibodies (and thus may permit administration to patients testing above a threshold level of antibody for the selected AAV capsid), and/or delivery of anti-FcRN antibodies which is described, e.g., in US Provisional Patent Application No. 63/040,381, filed June 17, 2020, entitled "Compositions and Methods for Treatment of Gene Therapy Patients", and/or one or more of a) a steroid or combination of steroids and/or (b) an IgG-cleaving enzyme, (c) an inhibitor of Fc-IgE binding; (d) an inhibitor of Fc-IgM binding; (e) an inhibitor of Fc-IgA binding; and/or (f) gamma interferon.

An antibody "Fc region" refers to the crystallizable fragment which is the region of an antibody which interacts with the cell surface receptors (Fc receptors). In one embodiment, the Fc region is a human IgG1 Fc. In one embodiment, the Fc region is a human IgG2 Fc. In one embodiment, the Fc region is a human IgG4 Fc. In one embodiment, the Fc region is an engineered Fc fragment. See, e.g., Lobner, Elisabeth, et al. "Engineered IgG1-Fc—one fragment to bind them all." *Immunological reviews* 270.1 (2016): 113-131; Saxena, Abhishek, and Donghui Wu. "Advances in therapeutic Fc engineering—modulation of IgG-Associated effector functions and serum half-life." *Frontiers in immunology* 7 (2016); Irani, Vashti, et al. "Molecular properties of human IgG subclasses and their implications for designing therapeutic monoclonal antibodies against infectious diseases." *Molecular immunology* 67.2 (2015): 171-182; Rath, Timo, et al. "Fc-fusion proteins and FcRn: structural insights for longer-lasting and more effective therapeutics." *Critical reviews in biotechnology* 35.2 (2015): 235-254; and Invivogen, IgG-Fc Engineering For Therapeutic Use, invivogen.com/docs/Insight200605.pdf, April 2006; each of which is incorporated by reference herein.

An antibody "hinge region" is a flexible amino acid portion of the heavy chains of IgG and IgA immunoglobulin classes, which links these two chains by disulfide bonds.

An "immunoglobulin molecule" is a protein containing the immunologically-active portions of an immunoglobulin heavy chain and immunoglobulin light chain covalently coupled together and capable of specifically combining with antigen. Immunoglobulin molecules are of any type (e.g., IgG, IgE, IgM, IgD, IgA and IgY), class (e.g., IgG1, IgG2, IgG3, IgG4, IgA1 and IgA2) or subclass. The terms "antibody" and "immunoglobulin" may be used interchangeably herein.

An “immunoglobulin heavy chain” is a polypeptide that contains at least a portion of the antigen binding domain of an immunoglobulin and at least a portion of a variable region of an immunoglobulin heavy chain or at least a portion of a constant region of an immunoglobulin heavy chain. Thus, the immunoglobulin derived heavy chain has significant regions of amino acid sequence homology with a member of the immunoglobulin gene superfamily. For example, the heavy chain in a Fab fragment is an immunoglobulin-derived heavy chain.

An “immunoglobulin light chain” is a polypeptide that contains at least a portion of the antigen binding domain of an immunoglobulin and at least a portion of the variable region or at least a portion of a constant region of an immunoglobulin light chain. Thus, the immunoglobulin-derived light chain has significant regions of amino acid homology with a member of the immunoglobulin gene superfamily.

“Neutralizing antibody titer” (NAb titer) a measurement of how much neutralizing antibody (e.g., anti-AAV NAb) is produced which neutralizes the physiologic effect of its targeted epitope (e.g., an AAV). Anti-AAV NAb titers may be measured as described in, e.g., Calcedo, R., et al., Worldwide Epidemiology of Neutralizing Antibodies to Adeno-Associated Viruses. *Journal of Infectious Diseases*, 2009, 199 (3): p. 381-390, which is incorporated by reference herein.

As used herein, a “subpopulation” of vp proteins refers to a group of vp proteins which has at least one defined characteristic in common and which consists of at least one group member to less than all members of the reference group, unless otherwise specified. For example, a “subpopulation” of vp1 proteins is at least one (1) vp1 protein and less than all vp1 proteins in an assembled AAV capsid, unless otherwise specified. A “subpopulation” of vp3 proteins may be one (1) vp3 protein to less than all vp3 proteins in an assembled AAV capsid, unless otherwise specified. For example, vp1 proteins may be a subpopulation of vp proteins; vp2 proteins may be a separate subpopulation of vp proteins, and vp3 are yet a further subpopulation of vp proteins in an assembled AAV capsid. In another example, vp1, vp2 and vp3 proteins may contain subpopulations having different modifications, e.g., at least one, two, three or four highly deamidated asparagines, e.g., at asparagine - glycine pairs. Unless otherwise specified, highly deamidated refers to at least 45% deamidated, at least 50% deamidated, at least 60% deamidated, at least 65% deamidated, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, 97%, 99%, up to about 100% deamidated at a referenced amino acid position, as compared to the predicted amino acid sequence at the reference amino acid

position. Such percentages may be determined using 2D-gel, mass spectrometry techniques, or other suitable techniques.

As used herein, a “stock” of rAAV refers to a population of rAAV. Despite heterogeneity in their capsid proteins due to deamidation, rAAV in a stock are expected to share an identical vector genome. A stock can include rAAV having capsids with, for example, heterogeneous deamidation patterns characteristic of the selected AAV capsid proteins and a selected production system. The stock may be produced from a single production system or pooled from multiple runs of the production system. A variety of production systems, including but not limited to those described herein, may be selected. See, e.g., WO 2019/168961, published September 6, 2019, including Table G providing the deamidation pattern for AAV9 and WO 2020/160582, filed September 7, 2018. See, also, e.g., WO 2020/223231, published November 5, 2020 (rh91, including table with deamidation pattern), US Provisional Patent Application No. 63/065,616, filed August 14, 2020, and US Provisional Patent Application No. 63/109,734, filed November 4, 2020, and International Patent Application No. PCT/US21/45945, filed August 13, 2021, which are all incorporated herein by reference in its entirety.

The abbreviation “sc” refers to self-complementary. “Self-complementary AAV” refers to a construct in which a coding region carried by a recombinant AAV nucleic acid sequence has been designed to form an intra-molecular double-stranded DNA template. Upon infection, rather than waiting for cell mediated synthesis of the second strand, the two complementary halves of scAAV will associate to form one double stranded DNA (dsDNA) unit that is ready for immediate replication and transcription. See, e.g., D M McCarty et al, “Self-complementary recombinant adeno-associated virus (scAAV) vectors promote efficient transduction independently of DNA synthesis”, *Gene Therapy*, (August 2001), Vol 8, Number 16, Pages 1248-1254. Self-complementary AAVs are described in, e.g., U.S. Patent Nos. 6,596,535; 7,125,717; and 7,456,683, each of which is incorporated herein by reference in its entirety.

As used herein, the term “operably linked” refers to both expression control sequences that are contiguous with the gene of interest and expression control sequences that act in trans or at a distance to control the gene of interest.

The term “heterologous” when used with reference to a protein or a nucleic acid indicates that the protein or the nucleic acid comprises two or more sequences or subsequences which are not found in the same relationship to each other in nature. For instance, the nucleic

acid is typically recombinantly produced, having two or more sequences from unrelated genes arranged to make a new functional nucleic acid. For example, in one embodiment, the nucleic acid has a promoter from one gene arranged to direct the expression of a coding sequence from a different gene. Thus, with reference to the coding sequence, the promoter is heterologous.

5 A "replication-defective virus" or "viral vector" refers to a synthetic or artificial viral particle in which an expression cassette containing a gene of interest is packaged in a viral capsid or envelope, where any viral genomic sequences also packaged within the viral capsid or envelope are replication-deficient; i.e., they cannot generate progeny virions but retain the ability to infect target cells. In one embodiment, the genome of the viral vector does not include
10 genes encoding the enzymes required to replicate (the genome can be engineered to be "gutless" - containing only the transgene of interest flanked by the signals required for amplification and packaging of the artificial genome), but these genes may be supplied during production. Therefore, it is deemed safe for use in gene therapy since replication and infection by progeny virions cannot occur except in the presence of the viral enzyme required for replication.

15 A "recombinant AAV" or "rAAV" is a DNase-resistant viral particle containing two elements, an AAV capsid and a vector genome containing at least non-AAV coding sequences packaged within the AAV capsid. In certain embodiments, the capsid contains about 60 proteins composed of vp1 proteins, vp2 proteins, and vp3 proteins, which self-assemble to form the capsid. Unless otherwise specified, "recombinant AAV" or "rAAV" may be used
20 interchangeably with the phrase "rAAV vector". The rAAV is a "replication-defective virus" or "viral vector", as it lacks any functional AAV rep gene or functional AAV cap gene and cannot generate progeny. In certain embodiments, the only AAV sequences are the AAV inverted terminal repeat sequences (ITRs), typically located at the extreme 5' and 3' ends of the vector genome in order to allow the gene and regulatory sequences located between the ITRs to be
25 packaged within the AAV capsid.

 The term "nuclease-resistant" indicates that the AAV capsid has assembled around the expression cassette which is designed to deliver a transgene to a host cell and protects these packaged genomic sequences from degradation (digestion) during nuclease incubation steps designed to remove contaminating nucleic acids which may be present from the production
30 process.

 As used herein, a "vector genome" refers to the nucleic acid sequence packaged inside a parvovirus (e.g., rAAV) capsid which forms a viral particle. Such a nucleic acid sequence

contains AAV inverted terminal repeat sequences (ITRs). In the examples herein, a vector genome contains, at a minimum, from 5' to 3', an AAV 5' ITR, coding sequence(s) (i.e., transgene(s)), and an AAV 3' ITR. ITRs from AAV2, a different source AAV than the capsid, or other than full-length ITRs may be selected. In certain embodiments, the ITRs are from the same AAV source as the AAV which provides the rep function during production or a transcomplementing AAV. Further, other ITRs, e.g., self-complementary (scAAV) ITRs, may be used. Both single-stranded AAV and self-complementary (sc) AAV are encompassed with the rAAV. The transgene is a nucleic acid coding sequence, heterologous to the vector sequences, which encodes a polypeptide, protein, functional RNA molecule (e.g., miRNA, miRNA inhibitor) or other gene product, of interest. The nucleic acid coding sequence is operatively linked to regulatory components in a manner which permits transgene transcription, translation, and/or expression in a cell of a target tissue. Suitable components of a vector genome are discussed in more detail herein. In one example, a "vector genome" contains, at a minimum, from 5' to 3', a vector-specific sequence, a nucleic acid sequence encoding protein of interest operably linked to regulatory control sequences (which direct their expression in a target cell), where the vector-specific sequence may be a terminal repeat sequence which specifically packages the vector genome into a viral vector capsid or envelope protein. For example, AAV inverted terminal repeats are utilized for packaging into AAV and certain other parvovirus capsids.

As used herein, "operably linked" sequences include both expression control sequences that are contiguous with the gene of interest and expression control sequences that act in trans or at a distance to control the gene of interest.

In certain embodiments, non-viral genetic elements used in manufacture of a rAAV, will be referred to as vectors (e.g., production vectors). In certain embodiments, these vectors are plasmids, but the use of other suitable genetic elements is contemplated. Such production plasmids may encode sequences expressed during rAAV production, e.g., AAV capsid or rep proteins required for production of a rAAV, which are not packaged into the rAAV. Alternatively, such a production plasmid may carry the vector genome which is packaged into the rAAV.

As used herein, a "parental capsid" refers to a non-mutated or a non-modified capsid selected from parvovirus or other viruses (e.g., AAV, adenovirus, HSV, RSV, etc.). In certain embodiments, the parental capsid includes any naturally occurring AAV capsids comprising a

wild-type genome encoding for capsid proteins (i.e., vp proteins), wherein the capsid proteins direct the AAV transduction and/or tissue-specific tropism. In some embodiments, the parent capsid is selected from AAV which natively targets CNS. In other embodiments, the parental capsid is selected from AAV which do not natively target CNS.

5 As used herein, a “variant capsid” or a “variant AAV” or “variant AAV capsid” refers to a modified capsid or a mutated capsid, wherein the capsid protein comprises an insertion of a tissue-specific targeting peptide.

 As used herein, an “expression cassette” refers to a nucleic acid molecule which comprises a biologically useful nucleic acid sequence (e.g., a gene cDNA encoding a protein,
10 enzyme or other useful gene product, mRNA, etc.) and regulatory sequences operably linked thereto which direct or modulate transcription, translation, and/or expression of the nucleic acid sequence and its gene product. As used herein, “operably linked” sequences include both regulatory sequences that are contiguous or non-contiguous with the nucleic acid sequence and regulatory sequences that act in trans or cis nucleic acid sequence. Such regulatory sequences
15 typically include, e.g., one or more of a promoter, an enhancer, an intron, a Kozak sequence, a polyadenylation sequence, and a TATA signal. The expression cassette may contain regulatory sequences upstream (5' to) of the gene sequence, e.g., one or more of a promoter, an enhancer, an intron, etc., and one or more of an enhancer, or regulatory sequences downstream (3' to) a gene sequence, e.g., 3' untranslated region (3' UTR) comprising a polyadenylation site, among
20 other elements. In certain embodiments, the regulatory sequences are operably linked to the nucleic acid sequence of a gene product, wherein the regulatory sequences are separated from nucleic acid sequence of a gene product by an intervening nucleic acid sequences, i.e., 5'-untranslated regions (5'UTR). In certain embodiments, the expression cassette comprises nucleic acid sequence of one or more of gene products. In some embodiments, the expression
25 cassette can be a monocistronic or a bicistronic expression cassette. In other embodiments, the term “transgene” refers to one or more DNA sequences from an exogenous source which are inserted into a target cell.

 The term “translation” in the context of the present invention relates to a process at the ribosome, wherein an mRNA strand controls the assembly of an amino acid sequence to
30 generate a protein or a peptide.

 The term “expression” is used herein in its broadest meaning and comprises the production of RNA or of RNA and protein. Expression may be transient or may be stable.

The term “substantial homology” or “substantial similarity,” when referring to a nucleic acid, or fragment thereof, indicates that, when optimally aligned with appropriate nucleotide insertions or deletions with another nucleic acid (or its complementary strand), there is nucleotide sequence identity in at least about 95 to 99% of the aligned sequences. Preferably, the homology is over full-length sequence, or an open reading frame thereof, or another suitable fragment which is at least 15 nucleotides in length. Examples of suitable fragments are described herein.

The terms “sequence identity” “percent sequence identity” or “percent identical” in the context of nucleic acid sequences refers to the residues in the two sequences which are the same when aligned for maximum correspondence. The length of sequence identity comparison may be over the full-length of the genome, the full-length of a gene coding sequence, or a fragment of at least about 500 to 5000 nucleotides, is desired. However, identity among smaller fragments, e.g., of at least about nine nucleotides, usually at least about 20 to 24 nucleotides, at least about 28 to 32 nucleotides, at least about 36 or more nucleotides, may also be desired. Similarly, “percent sequence identity” may be readily determined for amino acid sequences, over the full-length of a protein, or a fragment thereof. Suitably, a fragment is at least about 8 amino acids in length and may be up to about 700 amino acids. Examples of suitable fragments are described herein.

The term “substantial homology” or “substantial similarity,” when referring to amino acids or fragments thereof, indicates that, when optimally aligned with appropriate amino acid insertions or deletions with another amino acid (or its complementary strand), there is amino acid sequence identity in at least about 95 to 99% of the aligned sequences. Preferably, the homology is over full-length sequence, or a protein thereof, e.g., an immunoglobulin region or domain, an AAV cap protein, or a fragment thereof which is at least 8 amino acids, or more desirably, at least 15 amino acids in length. Examples of suitable fragments are described herein.

By the term “highly conserved” is meant at least 80% identity, preferably at least 90% identity, and more preferably, over 97% identity. Identity is readily determined by one of skill in the art by resort to algorithms and computer programs known by those of skill in the art.

Generally, when referring to “identity”, “homology”, or “similarity” between two different adeno-associated viruses, “identity”, “homology” or “similarity” is determined in reference to “aligned” sequences. “Aligned” sequences or “alignments” refer to multiple nucleic

acid sequences or protein (amino acids) sequences, often containing corrections for missing or additional bases or amino acids as compared to a reference sequence. Alignments are performed using any of a variety of publicly or commercially available Multiple Sequence Alignment Programs. Examples of such programs include, “Clustal Omega”, “Clustal W”, “CAP Sequence Assembly”, “MAP”, and “MEME”, which are accessible through Web Servers on the internet. Other sources for such programs are known to those of skill in the art. Alternatively, Vector NTI utilities are also used. There are also a number of algorithms known in the art that can be used to measure nucleotide sequence identity, including those contained in the programs described above. As another example, polynucleotide sequences can be compared using Fasta™, a program in GCG Version 6.1. Fasta™ provides alignments and percent sequence identity of the regions of the best overlap between the query and search sequences. For instance, percent sequence identity between nucleic acid sequences can be determined using Fasta™ with its default parameters (a word size of 6 and the NOPAM factor for the scoring matrix) as provided in GCG Version 6.1, herein incorporated by reference. Multiple sequence alignment programs are also available for amino acid sequences, e.g., the “Clustal Omega”, “Clustal X”, “MAP”, “PIMA”, “MSA”, “BLOCKMAKER”, “MEME”, and “Match-Box” programs. Generally, any of these programs are used at default settings, although one of skill in the art can alter these settings as needed. Alternatively, one of skill in the art can utilize another algorithm or computer program which provides at least the level of identity or alignment as that provided by the referenced algorithms and programs. See, e.g., J. D. Thomson et al, Nucl. Acids. Res., “A comprehensive comparison of multiple sequence alignments”, 27(13):2682-2690 (1999).

An effective amount may be determined based on an animal model, rather than a human patient.

As described above, the term “about” when used to modify a numerical value means a variation of $\pm 10\%$, ($\pm 10\%$, e.g., ± 1 , ± 2 , ± 3 , ± 4 , ± 5 , ± 6 , ± 7 , ± 8 , ± 9 , ± 10 , or values therebetween) from the reference given, unless otherwise specified.

In certain instances, the term “E+#” or the term “e+#” is used to reference an exponent. For example, “5E10” or “5e10” is 5×10^{10} . These terms may be used interchangeably.

As used throughout this specification and the claims, the terms “comprise” and “contain” and its variants including, “comprises”, “comprising”, “contains” and “containing”, among other variants, is inclusive of other components, elements, integers, steps and the like.

The term “consists of” or “consisting of” are exclusive of other components, elements, integers, steps and the like.

It is to be noted that the term “a” or “an”, refers to one or more, for example, “an enhancer”, is understood to represent one or more enhancer(s). As such, the terms “a” (or “an”),
5 “one or more,” and “at least one” is used interchangeably herein.

With regard to the description of these inventions, it is intended that each of the compositions herein described, is useful, in another embodiment, in the methods of the invention. In addition, it is also intended that each of the compositions herein described as useful in the methods, is, in another embodiment, itself an embodiment of the invention.

10 Unless defined otherwise in this specification, technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs and by reference to published texts, which provide one skilled in the art with a general guide to many of the terms used in the present application.

15 EXAMPLES

The following examples are illustrative only and are not a limitation on the invention described herein.

Example 1. Primary Screen

20 It has been shown that small peptide insertions into flexible loop on the surface of the AAV capsid can mediate interactions with new cellular receptors. In one case discovered at CalTech (AAV9-PHP.B), a seven amino acid peptide inserted into the HVR8 loop on AAV9 mediates interaction with Ly6a, a GPI-anchored receptor on the brain vasculature of some mouse strains. This interaction drives transport of AAV9-PHP.B across the blood-brain barrier
25 (BBB), resulting in ~50-fold higher transduction of brain cells than AAV9. In this work, we search for peptide inserts that can bind cell membrane targets on the BBB and thus have the potential to drive the AAV9 capsid across the BBB.

We sought to solve the AAV-BBB problem by first surveying the available academic and patent literature for peptide sequences that may have the potential to interact with the
30 vascular cells in the brain. We found the following sources of these peptides:

- Published results of phage-display experiments in which phage display libraries were panned against primary brain endothelial cells;

- Natural ligand peptides to known BBB-resident membrane proteins;
- CDRs of antibodies targeted to BBB-resident membrane proteins;
- Viral coat proteins of flaviviruses that cause encephalitis; and
- Bacterial toxins that have cell-binding activities directed at GPI

5 anchorages.

We generated a library of AAV9 insertion mutants containing hundreds of peptides from these sources, all inserted individually at the HVR8 locus (between position 588 and 589, based on the numbering of the amino acid sequence of AAV9 capsid of SEQ ID NO: 44). Each peptide was typically present in the library in multiple forms that differed by 1) length of
10 peptide inserted 2) presence of flexible GSG or GG linker sequences on both sides of the peptide. Peptides were also encoded using multiple synonymous codons so that we could independently observe replicate activities in the screen.

Additionally, we generated a library of insert variants in HVR8 of AAVhu68 capsid with either known or suspected ligand peptides which target the blood-brain barrier (BBB)
15 receptors (between position 588 and 589, based on the numbering of the amino acid sequence of AAVhu68 capsid of SEQ ID NO: 45). Such were:

- Peptides binding mammalian brain endothelium (published phage display data);
- Classical RMT receptor ligands (e.g., Tf);
- CDRs of mAbs against RMT receptors (e.g., anti-TfR); and
- Coat proteins of flaviviruses which cause encephalitis.

As a control a PHP.B peptide was included as well (positive control for C57/BL6 and negative control for Balb/c & NHP). Each peptide was encoded in multiple ways (with and without a linker, and in several synonymous DNA sequences).

25 We injected this library at high-dose intravenously (IV) to 2 mouse strains and to one non-human primate. After a 2-3 week in-life period, the animals were necropsied, and tissues were collected. We extracted the DNA genomes of AAV vectors from CNS and other tissues, and subjected these to next-generation sequencing (NGS). The vector variants encapsidate their own capsid gene variant, allowing us to track capsid activity through the relative abundance of
30 the capsid gene variant in the tissue of interest. We scored the BBB activity (“enrichment score”) of each variant in the library by calculating its abundance in the CNS normalized to its abundance in the injected library mixture.

In mouse study, tope brain enriched HVR8 insertions in C57/BL6 mice were: TLAVPFK (SEQ ID NO: 49) (PHP.B), Positive control PHP.B comes up 3 times independently as the most enriched hit. Three of the PHP.B peptides with synonymous codons are independently enriched. Several other peptides are also enriched in brain. FIGs 1A and 1B show the enrichment scores for the best mouse brain hits in the screen, with reference peptides (FIG 1A for C57BL/6J mice; and FIG 1B for Balb/c mice. FIGs 2A and 2B show the enrichment scores for the top performing NHP brain (FIG 2A) and spinal cord (FIG 2B) tissue.

Table 1. The hit peptide list from the primary screen (as identified in screen in NHPs and mice).

Abbreviation	Peptide Amino Acid Sequence	SEQ ID NO
EFS	EFSSNTVKLTS	38
SSN-L	GGSSNTVKLTSGHGG	39
SSN	SSNTVKLTSGH	40
SAN	SANFIKPTSY	41
VLT-L	GGVLTNIARGEYMRGG	46
IEI	IEINATRAGTNL	42
IEI-L	GGIEINATRAGTNLGG	43

Example 2. Secondary validation

We followed up the primary screen in mice by generating GFP reporter vectors for several of the hit capsids. The vectors were injected at high dose IV to C57BL/6J mice. 2 weeks later, we necropsied the mice and collected GFP images of brain sections (data not shown). All of the hit vectors tested in the GFP study were de-localized from the liver, as evident from liver GFP staining (data not shown).

Higher magnification imaging revealed that the capsids SSN and SAN are significantly brain localized, but restricted to the endothelium. RCA-lectin is a co-stain marker of brain endothelial cells in these sections (data not shown). These results indicate that the identified AAV capsids in the screen are BBB-receptor binding, but do not cross the BBB. The mutant series showed a range of affinities for Ly6a receptor. Brain transduction levels decreased as correlated with observed tighter affinity binding of peptide to Ly6a receptor. The observed tight binding reduced transduction due to genomes likely stuck in the endothelium.

This endothelial localization may be useful for certain diseases. Also, this activity may be optimized to convert these from brain localizing to BBB-crossing vectors. We confirmed these BBB-crossing and brain-localizing activities in a barcoded vector study. Briefly, each capsid was used to individually produce vector containing a GFP reporter gene with a unique DNA barcode included. The barcoded capsid preps were mixed in equal proportions, and injected into C57BL/6J or Balb/c mice (FIG 3A-3D). At the conclusion of in-life, the mouse tissues were subjected to NGS sequencing to count the abundance of each barcode among the vector genomes extracted from the tissue. The results confirm brain localization of vector genomes for all the hit capsids identified in the primary screen. In Balb/c mice, the secondary validation screen showed brain targeting for all hit sequences discovered in primary screen (FIG. 3A). In C57BL/6 mice the secondary validation screen showed brain targeting for all hit sequences discovered in primary screen (FIG 3B). In both, Balb/c and C57BL/6 mice, liver de-targeting for all hit sequences, relative to AVA9 was consistent with affinity for brain vasculature (FIG 3C and 3D).

15

Example 3. Endothelial targeting sequences.

For NHP secondary validation, a barcode study was performed. The study was performed in two NHPs with injected mixture of 27 barcoded vectors, including AAV9 at 4.5×10^{13} GC/kg following 21-day in life. Analysis of a whole-brain tissue homogenate was performed. While some vectors showed a modestly improved vector biodistribution, no vectors showed an improved whole-brain transduction. The accumulation of vector genomes in the brain (DNA) and the expression of the vector-derived transcripts (mRNA) were poorly correlated (Table 2). We observed poor mRNA versus DNA correlation for endothelial-targeting vectors.

25

Table 2.

	DNA		mRNA	
	NHP1	NHP2	NHP1	NHP2
SSN	7.2	4.2	0.1	0.1
EFS	7.0	3.8	0.1	0.1
VLT-L	4.4	1.6	0.1	0
IEI-L	1.7	4.2	0.1	0.1

	DNA		mRNA	
	NHP1	NHP2	NHP1	NHP2
SSN-L	2.9	2.3	0.1	0.1
IEI	2.8	1.9	0.1	0.1
SAN	2.8	1.9	0.1	0.1
AAV9	1.0	1.0	1.0	1.0

Table 3, below, shows the average relative localization score for both of the NHPs in the barcode study, while normalized to AAV9 (equal to 1). In line with the brain-focused library design, most vectors have unremarkable localization in non-brain tissues. An exception to this is the vector PMK, which significantly re-localized to the spleen of both NHPs relative to AAV9.

Table 3.

AAV	Eye	Kidney	Liver	Lungs	Pancreas	Spleen	Testicle	Diaphr.	Quadrcp	Heart
AAV9	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
SSN	1.5	0.3	0.3	0.3	0.2	0.0	0.4	1.1	1.0	0.6
SSN-L	1.3	0.2	0.5	0.2	0.2	0.1	0.6	0.9	0.9	0.6
EFS	2.6	0.4	0.4	0.7	0.2	0.1	0.9	1.6	1.4	1.0
VLT-L	0.9	0.2	0.5	0.3	0.2	0.1	0.4	0.6	0.6	0.4
IEI	0.6	0.2	0.1	0.5	0.1	0.0	0.2	0.3	0.5	0.2
IEI-L	1.0	0.2	0.1	0.4	0.1	0.6	0.8	0.5	0.7	0.5
SAN	1.0	0.3	0.3	0.5	0.1	0.1	0.3	0.7	0.8	0.4

Table 4, below, shows that brain endothelial hits have common in vitro transduction profiles (as measured in 293 transduction) and common production profiles. Importantly, vectors with the endothelial targeting activity all show dramatically increased relative abundance in the AAV vs Plasmid libraries as measured by NGS.

Table 4.

Vector Name	Barcode DNA	Barcode mRNA	Insert Sequence	293 transd	Plasmid to AAV conversion in library	Vector Yield
SSN	5.7	0.1	SSNTVKLTSGH	20	54	0.9
EFS	5.4	0.1	EFSSNTVKLTS	17	35	0.8
VLT-L	3.0	0.0	GGVLTNIARGEYMRGG	21	16	0.6
IEI-L	3.0	0.1	GGIEINATRAGTNLGG	10	20	0.9
SSN-L	2.6	0.1	GGSSNTVKLTSGHGG	24	69	0.8
IEI	2.3	0.1	IEINATRAGTNL	10	14	0.9
SAN	2.3	0.1	SANFIKPTSY	43	26	0.7
AAV9	1.0	1.0		1		1.0

The mutation library of SAN peptide confirms the role of “NxTK” motif in brain targeting. In this study, every possible single amino acid change to SAN insert was made, the optimized library variants were injected in mice, and scores for biodistribution and yield for each variant was measured. “NxTK” motif is the critical motif for brain biodistribution in the SAN insert (Table 5 and FIG 5). The “NxTK” motif controls plasmid-to-AAV conversion in the SAN peptide insert (Table 6 and FIG 6). Furthermore, the “NxTK” motif linked three properties of these endothelial vectors in the “NxTK” class: the endothelial cell transduction, improved 293 cell transduction and ability to propagate during library production. FIGS 7A to 7D show that “NxTK” motif confers broad transduction advantage across cell lines. The relative transduction levels were improved when compared to AAV9 capsid in 293 cells (FIG 7A), NIH3T3 cells (FIG 7B), and HUH7 cells (FIG 7C). FIG 7D shows a significant early improvement of transduction at day 3 post-transduction (3DPT) and approximately a 10-times improvement by day 7 post transduction (7DPT). AAV-GFP vectors with EFS and SAN peptide inserts showed an improved transduction when transduced the primary macaque airway epithelial cells (FIG 7E-7H).

Table 5. Brain biodistribution (average from top to bottom).

	S	A	N	F	I	K	P	T	S	Y
A	0.4	-0.4	-4.8	1.0	0.3	-3.8	-0.3	-0.5	-0.2	0.1
C	-6.0	0.0	-4.0	-1.5	-4.6	-3.1	-4.0	-4.0	-3.1	-1.1
D	0.7	-1.0	-3.7	0.5	-2.6	-8.3	-1.2	-0.8	-0.3	0.7
E	0.0	-0.8	-2.1	-0.5	-4.2	-5.3	-1.8	-0.4	0.2	0.9
F	-1.3	-2.4	-1.6	-0.4	-3.9	-1.9	-1.4	-0.9	-2.2	-0.2
G	0.1	-0.9	-2.7	1.3	-3.1	-4.6	0.5	0.4	-0.2	-0.7
H	-1.8	-1.0	-4.3	0.2	-2.4	-3.0	-1.1	-1.7	-1.9	-1.3
I	-1.0	-0.9	-5.2	0.8	-0.4	-3.5	-0.9	-1.2	-0.8	0.2
K	-4.3	-2.8	-4.9	-0.4	-3.5	-4.0	-1.2	-2.1	-3.2	-2.7
L	-0.6	-0.8	-3.8	0.2	-3.7	-3.3	-1.1	-1.1	-0.5	0.4
M	-0.7	0.2	-4.2	0.6	-3.0	-4.1	-0.8	-0.8	-0.5	0.4
N	-0.3	-0.7	-0.4	1.0	-1.1	-4.4	-0.2	-0.5	-0.7	0.5
P	-1.1	-0.1	-4.5	-2.5	-2.3	-3.6	-0.4	-0.8	-0.2	-0.2
Q	0.0	-0.6	-3.4	<u>1.2</u>	-4.6	-3.6	-0.1	-0.3	-0.1	0.8
R	-2.3	-3.6	-3.9	-0.3	-1.1	-1.1	-2.3	-2.5	-3.1	-2.6
S	-0.4	-1.2	-3.6	0.0	-1.0	-4.6	-0.2	-0.5	-0.4	0.3
T	0.2	-0.9	-3.4	1.2	0.2	-4.2	-0.3	-0.4	-0.3	0.6
V	-1.0	-1.1	-3.9	1.0	-0.1	-4.7	-1.1	-0.6	-0.5	0.6
W	-2.0	-3.0	0.0	-2.0	-2.9	-4.8	0.0	-2.1	0.9	-1.3
Y	-1.6	-4.8	-1.9	0.0	-4.4	-4.1	-1.1	-1.4	-0.8	-0.4

Table 6. Plasmid to AAV yield conversion

	S	A	N	F	I	K	P	T	S	Y
A	0.6	0.4	-1.1	1.1	1.3	-1.4	1.3	-0.1	0.4	1.2
C	-3.1	11.0	-6.2	-2.6	-6.0	-6.8	-4.3	-5.4	-4.4	-1.7
D	-0.1	0.6	-0.7	0.3	-0.7	-1.1	1.0	0.2	0.8	0.9
E	0.3	-1.2	-1.0	0.7	-0.6	-0.6	0.9	0.1	0.8	0.8

	S	A	N	F	I	K	P	T	S	Y
F	-2.8	-5.5	-5.1	0.4	-3.7	-7.2	-5.5	-3.7	-3.2	0.2
G	0.4	0.7	-0.9	1.2	-0.4	-0.5	1.2	0.9	0.7	1.1
H	0.2	-0.9	-1.6	1.3	-0.6	-1.7	1.0	0.0	0.1	0.9
I	-0.9	-1.4	-2.9	1.1	0.4	-3.3	-1.9	-0.7	-0.9	1.0
K	-0.5	-0.5	-1.3	0.3	-1.6	-3	-1.2	-2.5	-1.8	-1.1
L	-0.5	-1.1	-2.6	1.1	-1.1	-4.5	-1.7	-1.2	-1.2	0.6
M	-0.6	-1.5	-2.4	1.3	-0.6	-3.0	-0.4	-0.9	-0.8	1.0
N	1.0	0.5	0.4	1.5	-0.1	-1.0	1.5	0.6	0.7	1.2
P	0.7	-0.2	-1.1	-0.4	-0.4	-0.9	0.4	-0.4	0.4	0.7
Q	0.8	1.2	-0.6	<u>1.6</u>	-0.1	-1.2	1.5	0.0	0.4	1.0
R	-2.1	-2.9	-4.6	-1.0	-5.2	-1.3	-3.2	-4.2	-3.9	-1.3
S	0.4	0.5	-1.5	1.0	1.3	-0.9	1.3	0.3	0.4	1.2
T	0.5	0.4	-1.0	1.3	1.1	-1.0	1.3	0.4	0.5	1.1
V	0.3	-0.3	-2.3	1.1	0.8	-3.7	-0.1	-0.2	0.2	1.1
W	-4.4	-5.6	-11.0	-2.0	-5.5	-5.8	-11.6	-4.9	-7.3	-1.1
Y	-1.8	-2.9	-6.3	0.2	-3.8	-6.8	-4.3	-2.9	-3.1	0.4

In summary, the selected amino acid sequences, as listed in Table 7, all contain the functional motif N- x- (T/I/V/A)- (K/R) motif (SEQ ID NO: 47). Beyond the selected sequences shown in Table 7, others were identified during the screening. We have data that supports that many substitutions to these insert sequences also support or even improve the endothelial targeting activity. Additionally, we have discovered approximately thousands of sequences that fit this motif from large, random insert libraries — these sequences likely all share improved transduction properties.

10

Table 7. Selected Endothelial-Targeting Sequences

Vector name	Insert sequence	SEQ ID NO
SSN	SSNTVKLTSGH	40
EFS	EFSSNTVKLTS	38

Vector name	Insert sequence	SEQ ID NO
VLT-L	GGVLTNIARGEYMRGG	46
IEI-L	GGIEINATRAGTNLGG	43
SSN-L	GGSSNTVKLTSGHGG	39
IEI	IEINATRAGTNL	42
SAN	SANFIKPTSY	41

We completed the barcode evaluation of primary screen hits in NHP. The brain localization is the most prominent feature of hits from this library, while there were no significant enhancements in targeting to peripheral tissues with possible exception of spleen targeting by AAV9-PMK. We defined a sequence motif common to all peptide inserts with brain-endothelial targeting activity. We confirmed the activity of this motif in brain-endothelial targeting as well as in conferring broad in vitro transduction advantage. A single sequence motif “NxTK” defines inserts from 4 unrelated sources that shares three properties:

- (1) dramatically improved in vitro transduction on 293s and other cell lines;
- (2) parasitic expansion during library production – “spreading” phenotype; and
- (3) endothelial biodistribution in vivo in mice and NHP.

The “NxTK” motif was mapped in systemic mutational screen of SAN and EFS vector inserts. In systemic mutational screen, “NxTK” motif is shown to be critical for brain endothelial biodistribution as well as for abundance in library production. The plasmid-to-AAV conversion, meaning the capsid yield during library production, is controlled by 2 factors: presence of parasitic spreading motif (major factor) and intrinsic capsid yield (minor factor). One of the spreading motifs was identified to be “NxTK”, and was likely to interact with 293 cell-surface receptors and confer transduction advantage. The transduction advantage in 293 could lead to propagation of vector genome (Cap) to neighboring cells during production period since library production is done with limiting Cap, so most cells initially have everything except a Cap gene. The minor factor was only evident after digital filtering out of the vectors with parasitic spreading motif.

Example 4. Engineering Strategy for AAV Capsid Development for Airway Delivery

Current AAV vectors poorly transduce cells of the nasal passages and upper airway, limiting the efficacy of prophylactic AAV strategies to protect against upper airway infectious

diseases like influenza or COVID-19. This project aims to engineer AAV capsids for improved transduction of these tissues, specifically a directed evolution and receptor targeting strategies are pursued. An engineering strategy pursued for AAV airway-specific capsid selection comprises of steps including but not limited to generating a diverse library of constructs with

5 inserts (10^3 to 10^9 initial diversity), screening and selection in primary primate or human airway cells, identifying the genomic identity (DNA or RNA) of selected constructs, performing additional screening to converge on hits, validate hit of improved capsids in NHP. The sources for capsid diversity for airway delivery comprise of pursuing two approaches: unbiased and

10 biased approach. In an unbiased approach, random peptides are inserted into AAV capsid surface to generate large libraries of greater than 10^7 variant diversity. In biased approach, peptides with known or suspected airway-cell binding activities inserted into AAV Capsid to generate small libraries with approximately 10^3 variant diversity. The source for such known or suspected airway-cell binding peptides: published phage display results, peptides from viral

15 receptor binding domains, known ligands to airway receptors, and peptides generated in previous in vitro library screening. For screening of the generated library of capsid in vitro, an assay with primary airway cells in air-liquid interface (ALI) cultures is used. The cell used are of human and macaque origin, and comprise of nasal, trachea and bronchi cells origin. A preliminary transduction test with GFP vectors in macaque primary airway epithelial cell

20 cultures showed a significant early improvement in transduction for EFS and SAN inserted peptides into AAV capsid (FIGs 7D and 7E-7H). FIG 7E shows microscopic analysis of the macaque primary airway epithelial cells in a control sample treated with carrier (i.e., no vector). FIG. 7F shows microscopic analysis of the macaque primary airway epithelial cells post-transduction with AAV9-GFP vector. FIG. 7G shows microscopic analysis of the macaque primary airway epithelial cells post-transduction with AAV9-GFP vector comprising EFS

25 peptide insert. FIG. 7H shows microscopic analysis of the macaque primary airway epithelial cells post-transduction with AAV9-GFP comprising SAN peptide inserts. A preliminary transduction test with GFP vectors in cultured human cells showed overall lower transduction, wherein at day 7 the ration of mRNA copy number over micro-gram total mRNA was 1×10^4 for cultured human cells (FIG 8) and 1×10^6 for cultured macaque primary airway epithelial cells

30 (FIG 7D). The SAN motif showed an advantage in transduction at Day 7 when in comparison to AAV9 transduction (FIG 8). The EFS motif showed a poorer transduction in bronchial and tracheal cultured human cells (FIG 8).

- Furthermore, we have developed a number of insert sequences through in vitro selection schemes on other projects that confer dramatically improved cell binding and transduction activity. The generated AAV-insert vectors are tested in barcoded pools on ALI cultures. Results show that all in vitro selected AAV9-insert vectors were better than AAV9 vectors (results not shown). Specifically, AAV9-insert vectors of Spr3L (NxTK) were the best in comparison, the latter showing an approximately 50-fold better transduction than AAV9 capsid.

(Sequence Listing Free Text)

- 10 The following information is provided for sequences containing free text under numeric identifier <223>.

SEQ ID NO:	Free Text under <223>
1	<220> <223> AAV2/9 n.588.EFS nucleic acid sequence expression cassette <220> <221> misc_feature <222> (1)..(36) <223> truncated promoter <220> <221> promoter <222> (1)..(7) <223> p5 promoter <220> <221> CDS <222> (37)..(1899) <223> AAV2-Rep <220>

	<221> CDS <222> (1919)..(4162) <223> AAV9 Cap <220> <221> misc_feature <222> (3683)..(3715) <223> EFS <220> <221> misc_feature <222> (4253)..(4383) <223> p5 promoter <220> <221> misc_feature <222> (4511)..(4725) <223> LacZ promoter
2	<220> <223> Synthetic Construct
3	<220> <223> Synthetic Construct
4	<220> <223> AAV2/9 n.588.IEI nucleic acid sequence expression cassette <220> <221> misc_feature <222> (1)..(36) <223> truncated promoter <220>

	<221> promoter <222> (1)..(7) <223> p5 promoter <220> <221> CDS <222> (37)..(1899) <223> AAV2-Rep <220> <221> CDS <222> (1919)..(4165) <223> AAV9 Cap <220> <221> misc_feature <222> (3683)..(3718) <223> IEI <220> <221> misc_feature <222> (4256)..(4386) <223> p5 promoter <220> <221> misc_feature <222> (4514)..(4728) <223> LacZ promoter
5	<220> <223> Synthetic Construct
6	<220> <223> Synthetic Construct

7	<220> <223> AAV2/9 n.588.IEI-L nucleic acid sequence expression cassette <220> <221> misc_feature <222> (1)..(36) <223> truncated promoter <220> <221> promoter <222> (1)..(7) <223> p5 Promoter <220> <221> CDS <222> (37)..(1899) <223> AAV2 Rep <220> <221> CDS <222> (1919)..(4177) <223> AAV9 Cap <220> <221> misc_feature <222> (3683)..(3739) <223> IEI-L <220> <221> misc_feature <222> (4268)..(4398)
---	---

	<223> p5 promoter <220> <221> misc_feature <222> (4526)..(4740) <223> LacZ promoter
8	<220> <223> Synthetic Construct
9	<220> <223> Synthetic Construct
10	<210> 10 <211> 4722 <212> DNA <213> Artificial Sequence <220> <223> AAV2/9 n.588.SAN nucleic acid sequence expression cassette <220> <221> misc_feature <222> (1)..(36) <223> truncated promoter <220> <221> promoter <222> (1)..(7) <223> p5 promoter <220> <221> CDS <222> (37)..(1899)

	<223> AAV2 Rep <220> <221> CDS <222> (1919)..(4159) <223> AAV 9 Cap <220> <221> misc_feature <222> (3683)..(3712) <223> SAN <220> <221> misc_feature <222> (4250)..(4380) <223> p5 promoter <220> <221> misc_feature <222> (4508)..(4722) <223> LacZ promoter
11	<220> <223> Synthetic Construct
12	<220> <223> Synthetic Construct
13	<220> <223> AAV2/9 n.588.SSN nucleic acid sequence expression cassette <220> <221> misc_feature <222> (1)..(36)

	<223> trunctaed promoter <220> <221> promoter <222> (1)..(7) <223> p5 promoter <220> <221> CDS <222> (37)..(1899) <223> AAV2 Rep <220> <221> CDS <222> (1919)..(4162) <223> AAV9 Cap <220> <221> misc_feature <222> (3683)..(3715) <223> SSN <220> <221> misc_feature <222> (4253)..(4383) <223> p5 promoter <220> <221> misc_feature <222> (4511)..(4725) <223> LacZ promoter
14	<220>

	<223> Synthetic Construct
15	<220> <223> Synthetic Construct
16	<220> <223> AAV2/9 n.588.SSN-L nucleic acid sequence expression cassette <220> <221> misc_feature <222> (1)..(36) <223> truncated promoter <220> <221> promoter <222> (1)..(7) <223> p5 promoter <220> <221> CDS <222> (37)..(1899) <223> AAV2 Rep <220> <221> CDS <222> (1919)..(4174) <223> AAV9 Cap <220> <221> misc_feature <222> (3683)..(3727) <223> SSN-L

	<220> <221> misc_feature <222> (4253)..(4737) <223> LacZ promoter <220> <221> misc_feature <222> (4265)..(4395) <223> p5 promoter
17	<220> <223> Synthetic Construct
18	<220> <223> Synthetic Construct
19	<220> <223> AAV2/9 n.588.VLT-L nucleic acid sequence expression cassette <220> <221> misc_feature <222> (1)..(36) <223> truncated promoter <220> <221> promoter <222> (1)..(7) <223> p5 promoter <220> <221> CDS <222> (37)..(1899) <223> AAV2 Rep

	<220> <221> CDS <222> (1919)..(4177) <223> AAV9 Cap <220> <221> misc_feature <222> (3683)..(3730) <223> VLT-L <220> <221> misc_feature <222> (4268)..(4398) <223> p5 promoter <220> <221> misc_feature <222> (4526)..(4740) <223> LacZ promoter
20	<220> <223> Synthetic Construct
21	<220> <223> Synthetic Construct
22	<220> <223> AAV2 Rep nucleic acid sequence
23	<220> <223> AAV2 Rep amino acid sequence
24	<220> <223> AAV9 Cap n.588.EFS nucleic acid sequence <220>

	<221> misc_feature <222> (1765)..(1797) <223> EFS
25	<220> <223> AAV9 Cap n.588.EFS amino acid sequence <220> <221> MISC_FEATURE <222> (499)..(599) <223> EFS
25	<220> <223> AAV9 Cap n588.IEI nucleic acid sequence <220> <221> misc_feature <222> (1765)..(1800) <223> IEI
27	<220> <223> AAV9 Cap n588.IEI amino acid sequence <220> <221> MISC_FEATURE <222> (499)..(600) <223> IEI
28	<220> <223> AAV9 Cap n.588.IEI-L nucleic acid sequence <220>

	<221> misc_feature <222> (1765)..(1812) <223> IEI-L
29	<220> <223> AAV9 Cap n.588.IEI-L amino acid sequence <220> <221> MISC_FEATURE <222> (499)..(604) <223> IEI-L
30	<220> <223> AAV9 Cap n.588.SAN nucleic acid sequence <220> <221> misc_feature <222> (1765)..(1794) <223> SAN
31	<220> <223> AAV9 Cap n.588.SAN amino acid sequence <220> <221> MISC_FEATURE <222> (499)..(598) <223> SAN
32	<220> <223> AAV9 Cap n.588.SSN nucleic acid sequence <220>

	<221> misc_feature <222> (1765)..(1797) <223> SSN
33	<220> <223> AAV9 Cap n.588.SSN amino acid sequence <220> <221> MISC_FEATURE <222> (499)..(599) <223> SSN
34	<220> <223> AAV9 Cap n.588.SSN-L nucleic acid sequence <220> <221> misc_feature <222> (1765)..(1809) <223> SSN-L
35	<220> <223> AAV9 Cap n.588.SSN-L amino acid sequence <220> <221> MISC_FEATURE <222> (499)..(603) <223> SSN-L
36	<220> <223> AAV9 Cap n588.VLT-L nucleic acid sequence <220>

	<221> misc_feature <222> (1765)..(1812) <223> VLT-L
37	<220> <223> AAV9 Cap n588.VLT-L amino acid sequence <220> <221> MISC_FEATURE <222> (499)..(604) <223> VLT-L
38	<220> <223> EFS peptide sequence
39	<220> <223> SSN-L peptide sequence
40	<220> <223> SSN peptide sequence
41	<220> <223> SAN peptide sequence
42	<220> <223> IEI peptide sequence
43	<220> <223> IEI-L peptide sequence
44	<220> <223> AAV9 capsid
45	<220> <223> AAVhu68 capsid
46	<220> <223> VLT-L peptide sequence
47	<220> <223> N-X-(T/I/V/A)-(K/R) motif

	<220> <221> MISC_FEATURE <222> (2)..(2) <223> any amino acid <220> <221> MISC_FEATURE <222> (3)..(3) <223> Xaa is selected from Threonine (T), Isoleucine (I), Valine (V) or Alanine (A) <220> <221> MISC_FEATURE <222> (4)..(4) <223> Xaa is selected from Lysine (K) or Arginine (R)
48	<220> <223> AAV2 variant peptide NDVRAVS
49	<220> <223> PHP.B peptide insert
50	<220> <223> nucleic acid sequence EFS
51	<220> <223> nucleic acid sequence IEI
52	<220> <223> nucleic acid sequence IEI-L
53	<220> <223> nucleic acid sequence SAN
54	<220> <223> nucleic acid sequence SSN
55	<220> <223> nucleic acid sequence SSN-L

56	<220> <223> nucleic acid sequence VLT-L
----	--

All documents cited in this specification are incorporated herein by reference. US Provisional Application No. 63/119,863, filed December 1, 2020 is incorporated herein by reference in its entirety. The sequence listing filed herewith named “20-9409PCT_ST25” and
5 the sequences and text therein are incorporated by reference. While the invention has been described with reference to particular embodiments, it will be appreciated that modifications can be made without departing from the spirit of the invention. Such modifications are intended to fall within the scope of the appended claims.

CLAIMS:

1. A recombinant adeno-associated virus particle (rAAV) having a capsid comprising an amino acid sequence that comprises the motif N- x- (T/I/V/A)- (K/R) (SEQ ID NO: 47), wherein the amino acid sequence is part of at least the AAV vp3 protein in the capsid and a vector genome packaged in the capsid which comprises a nucleic acid sequence encoding a gene product under control of sequences which direct expression thereof, provided that the capsid is not a mutant AAV2 capsid comprising an NDVRAVS (SEQ ID NO: 48) sequence.
2. The rAAV according to claim 1, wherein the amino acid sequence comprises comprising the N- x- (T/I/V/A)- (K/R) motif optionally flanked at the amino terminus and/or the carboxy terminus of the motif by two amino acids to seven amino acids is inserted into the AAV capsid vp3 region.
3. The rAAV according to claim 1 or 2, wherein the sequence inserted into the capsid comprises:
 - (a) SSNTVKLTSGH (SEQ ID NO: 40);
 - (b) EFSSNTVKLTS (SEQ ID NO: 38);
 - (c) GGVLTNIARGEYMRGG (SEQ ID NO: 46);
 - (d) GGIEINATRAGTNLGG (SEQ ID NO: 43);
 - (e) GGSSNTVKLTSGHGG (SEQ ID NO: 39);
 - (f) IEINATRAGTNL (SEQ ID NO: 42); or
 - (g) SANFIKPTSY (SEQ ID NO: 41).
4. The rAAV according to any one of claims 1 to 3, wherein the amino acid sequence of the motif is NTVK.
5. The rAAV of claim 1, wherein the motif N- x- (T/I/V/A)- (K/R) (SEQ ID NO: 47) is optionally flanked by two to seven amino acids at its carboxy- and/or amino terminus and inserted between amino acids 588 and 589 of an AAV9 capsid protein, based on the numbering of amino acid sequence: SEQ ID NO: 44.

6. A composition comprising a stock of the rAAV according to any of claims 1 to 5 and one or more of a physiologically compatible carrier, excipient, and/or aqueous suspension base.

7. An endothelial cell targeting peptide, wherein the endothelial cell targeting peptide comprises a motif comprising an amino acid sequence of N- x- (T/I/V/A)- (K/R) (SEQ ID NO: 47) optionally flanked at the amino terminus and/or the carboxy terminus of the motif by two amino acids to seven amino acids, and optionally further conjugated to a nanoparticle, a second molecule, or a viral capsid protein.

8. The endothelial cell targeting peptide according to claim 7, wherein the endothelial cell targeting peptide comprises:

- (a) SSNTVKLTSGH (SEQ ID NO: 40);
- (b) EFSSNTVKLTS (SEQ ID NO: 38);
- (c) GGVLTNARGEYMRGG (SEQ ID NO: 46);
- (d) GGIEINATRAGTNLGG (SEQ ID NO: 43);
- (e) GGSSNTVKLTSGHGG (SEQ ID NO: 39);
- (f) IEINATRAGTNL (SEQ ID NO: 42); or
- (g) SANFIKPTSY (SEQ ID NO: 41).

9. The endothelial cell targeting peptide according to claim 7 or 8, wherein the amino acid sequence of the motif is NTVK.

10. A composition comprising the endothelial cell targeting peptide according to any one of claim 7 to 9 and one or more of a physiologically compatible carrier, excipient, and/or aqueous suspension base.

11. A fusion polypeptide or protein comprising a brain endothelial cell targeting peptide according to any of claims 7 to 9 and a fusion partner which comprises at least one polypeptide or protein.

12. A composition comprising a fusion polypeptide or protein according to claim 11 and one or more of a physiologically compatible carrier, excipient, and/or aqueous suspension base.

13. Use of a stock of the rAAV according to any one of claims 1 to 5, an endothelial cell targeting peptide according to any one of claims 7 to 9, or a fusion polypeptide or protein according to claim 11, or a composition according to any one of claims 6, 10 or 12, for delivering a therapeutic to a patient in need thereof.

14. A method for targeting therapy to the endothelial cells in brain, said method comprising administering to a patient in need thereof a stock of rAAV according to any one of claims 1 to 5.

15. A method for treating Allan-Herndon-Dudley disease by delivering to a subject in need thereof a stock of rAAV according to any one of claims 1 to 5, wherein the encoded gene product is an MCT8 protein.

16. A method for targeting therapy to the lung comprising administering to a patient in need thereof a stock of rAAV according to any one of claims 1 to 5.

17. A method for treating a disease of the lung by delivering to a subject in need thereof a stock of rAAV according to any one of claims 1 to 5, wherein the encoded gene product is a soluble Ace2 protein, an anti-SARS antibody, an anti-SARS-CoV2 antibody, an anti-influenza antibody, or a cystic fibrosis transmembrane protein.

18. A method for increasing transduction of AAV production cells in vitro comprising inserting an N- x- (T/I/V/A)- (K/R) (SEQ ID NO: 47) motif into an AAV capsid.

19. The method according to claim 16, wherein the production cells are 293 cells.