

**SUBSTITUTED HETEROCYCLIC ACYL-TRIPEPTIDES USEFUL
AS THROMBIN RECEPTOR MODULATORS**

5

BACKGROUND OF THE INVENTION

Thrombin is an important serine protease in hemostasis and thrombosis. One of the key actions of thrombin is receptor activation. A functional human thrombin receptor (TR), cloned by Coughlin in 1991 (T.-K. Vu, Cell 1991, 64, 1057), was found to be a member of the G-protein coupled receptor (GPCR) superfamily. The receptor activation putatively occurs by N-terminal recognition and proteolytic cleavage at the Arg-41/Ser-42 peptide bond to reveal a truncated N-terminus. This new receptor sequence, which has an SFLLRN (Ser-Phe-Leu-Leu-Arg-Asn) N-terminus acting as a tethered ligand to recognize a site on the receptor, can trigger activation and signal transduction leading to platelet aggregation.

Peptide analogues based on this hexapeptide have also shown good agonist activity leading to platelet aggregation. Since 1991, two other protease-activated receptors with extensive homology to the thrombin receptor, "PAR-2" and "PAR-3," were cloned, and found to be activated by similar N-terminal hexapeptide sequences. Hence, agonists / antagonists of the thrombin receptor, such as those included in this invention, may be useful in activating / antagonizing these protease-activated receptors as well.

- 2 -

Activation of the thrombin receptor by agonist compounds of this invention may mimic thrombin's role in tissue repair. Thrombin can initiate effects related to wound healing, such as: increasing vascular permeability to allow entry of cells and fluid into injured tissue (A. B. Malik, *Semin. Thromb. Haemostasis* 1986, 184); increasing the synthesis of PDGF by endothelial cells (J. M. Harlan, *J. Cell Biol.* 1986, 103, 1129); and increasing the adhesion of platelets, monocytes, and neutrophils to endothelial cells (M. P. Bevilacqua, *Science* 1989, 243, 1160). Tissue repair in rats following surgical incision is accelerated by the use of thrombin (D. H. Carney, *J. Clin. Invest.* 1992, 89, 1469). Thus, agonists at the thrombin receptor may be useful as wound healing agents or in tissue repair.


The peptide-based antagonists of the thrombin receptor in this present invention may show efficacy against myocardial infarction, stroke, restenosis, angina, atherosclerosis, and ischemic attacks by virtue of their ability to prevent platelet aggregation. The thrombin receptor has also been identified on other cell types: endothelial, fibroblast, osteosarcoma, smooth muscle, and neuronal/glia. Thrombin activation of endothelial cells upregulates P-selectin to induce polymorphonuclear leukocyte adhesion - an inflammatory response of the vessel wall (Y. Sugama, *J. Cell Biol.* 1992, 119, 935). In fibroblasts, thrombin receptor activation induces proliferation and transmission of mitogenic signals (D. T. Hung, *J. Cell Biol.* 1992, 116, 827). Thrombin has been implicated in osteoblast proliferation through its activation of osteoblast cells

- 3 -

(D. N. Tatakis, *Biochem. Biophys. Res. Commun.* 1991, 174, 181). Thrombin has been implicated in the regulation and retraction of neurons (K. Jalink, *J. Cell. Biol.* 1992, 118, 411). Therefore, in this context, the antagonist compounds of this invention may also be useful against inflammation, restenosis, cancer, osteoporosis, and neurodegenerative disorders.

SUMMARY OF THE INVENTION

The present invention is directed to compounds represented by the following general formula (I):




wherein A_1 , A_2 , A_3 , X, Y, and Z are defined later.

These compounds are thrombin receptor modulators and may be useful either as agonists in wound healing and tissue repair or as antagonists in myocardial infarction, stroke, restenosis, angina, atherosclerosis, ischemic attacks, inflammation, cancer, osteoporosis, or neurodegenerative disorders.

- 4 -

DETAILED DESCRIPTION OF THE INVENTION

More particularly, the present invention is directed to compounds of the following formula (I):



wherein

10 A_1 is an alkyl amino acid residue selected from Cha, Leu and Ile, an amino alkyl amino acid residue selected from Arg and Lys, or an aryl amino acid residue selected from Phe, substituted Phe, Tyr, or Trp;

15 A_2 is an amino alkyl amino acid residue selected from Lys, Orn, Arg, and homo Arg;

20 A_3 is an aryl amino acid residue selected from Phe, substituted Phe, Tyr, Trp, phenyl-Gly, 2-thienyl-Ala and 3-thienyl-Ala, an alkyl amino acid residue selected from Cha, Leu and Ile, an amido alkyl amino acid selected from Asn and Gln, or an amino alkyl amino acid residue selected from Arg, homo Arg, Orn and Lys;

- 5 -

X is selected from CO, CS, or SO₂;

Y is selected from aryl, substituted aryl, heterocycloalkyl, substituted heterocycloalkyl, heteroaryl, substituted heteroaryl, heteroarylethylenyl, substituted heteroarylethylenyl, arylacrylamidoheteroaryl, substituted arylacrylamidoheteroaryl, heteroarylacrylamidoheteroaryl and substituted heteroarylacrylamidoheteroaryl, preferably, Y is not pyrrolidinyl, substituted pyrrolidinyl, phenyl or 2-aminophenyl; most preferably, Y is selected from heteroaryl, substituted heteroaryl, arylacrylamidoheteroaryl, and substituted arylacrylamidoheteroaryl;

Z is selected from NH₂, NH-alkyl, NH-aralkyl, or an amino alkyl amino acid residue selected from Arg-NH₂; and

wherein all amino acids are of the L configuration; and the pharmaceutically acceptable salts thereof.

In the compounds of formula (I), the amino acid residues comprising the A₁, A₂, and A₃ substituents are attached to the adjacent moiety according to standard nomenclature so that the amino-terminus (N-terminus) of the amino acid is drawn on the left and the carboxy-terminus of the amino acid is drawn on the right. So, for example, in Compound 1, where A₁ is Cha, A₂ is Arg and A₃ is Phe, the N-terminus of the Cha (A₁) is attached to the X substituent and the carboxy-terminus of the Cha (A₁) is attached to the N-terminus of the A₂ substituent

- 6 -

(Arg), similarly, the the N-terminus of the Arg (A₂) is attached to the carboxy-terminus of the A₁ substituent and the carboxy-terminus of the Arg (A₂) is attached to the N-terminus of the A₃ substituent (Phe), similarly, the N-terminus of the Phe (A₃) is attached to the carboxy-terminus of the A₂ substituent and the carboxy-terminus of the Phe (A₃) is attached to the Z substituent.

When a particular group is "substituted" (e.g., Phe, heterocycloalkyl, heteroaryl, acrylamidoheteroaryl), that group may have from 1 to 4 substituents independently selected from: halo (I, Br, Cl, F), C₁-C₈ alkyl, C₁-C₈ alkoxy, amino, amido, carboxyl, cyano, nitro, fluorinated C₁-C₈ alkyl, fluorinated C₁-C₈ alkoxy, or aryl (preferably, phenyl or substituted phenyl).

The term "subject" as used herein, refers to an animal, preferably a mammal, most preferably a human, who has been the object of treatment, observation or experiment.

The term "therapeutically effective amount" as used herein, means that amount of active compound or pharmaceutical agent that elicits the biological or medicinal response in a tissue system, animal or human that is being sought by a researcher, veterinarian, medical doctor or other clinician, which includes alleviation of the symptoms of the disease or disorder being treated.

- 7 -

As used herein, unless otherwise noted alkyl and alkoxy whether used alone or as part of a substituent group, include straight and branched chains having 1 to 8 carbon atoms, or any number within this range. For example, alkyl radicals include methyl, ethyl, propyl, isopropyl, *n*-butyl, isobutyl, *sec*-butyl, *t*-butyl, *n*-pentyl, 3-(2-methyl)butyl, 2-pentyl, 2-methylbutyl, neopentyl, *n*-hexyl, 2-hexyl and 2-methylpentyl. Alkoxy radicals are oxygen ethers formed from the previously described straight or branched chain alkyl groups. Cycloalkyl groups contain 3 to 8 ring carbons and preferably 5 to 7 carbons. Similarly, alkenyl and alkynyl groups include straight and branched chain alkenes and alkynes having 1 to 8 carbon atoms, or any number within this range.

The term "aryl" as used herein, alone or in combination with other terms, represents an aromatic hydrocarbon group. Examples of aryl groups include, but are not limited to, phenyl, naphthyl, phenanthryl, anthryl, biphenylenyl, fluorenyl, or azulenylyl. Preferred aryl groups include phenyl, naphthyl and biphenylenyl.

The term "heterocycloalkyl" as used herein represents an unsubstituted or substituted stable three to seven membered monocyclic saturated ring system which

- 8 -

consists of carbon atoms and from one to three heteroatoms selected from N, O or S, and wherein the nitrogen or sulfur heteroatoms may optionally be oxidized, and the nitrogen heteroatom may optionally be quaternized. The heterocyloalkyl group may be attached at any heteroatom or carbon atom which results in the creation of a stable structure. Examples of such heterocyloalkyl groups include, but are not limited to azetidiny, piperidiny, piperaziny, oxopiperaziny, oxopiperidiny, oxoazepiny, azepiny, tetrahydrofurany, dioxolany, tetrahydroimidazolyl, tetrahydrooxazolyl, tetrahydropyrany, morpholiny, thiomorpholiny, thiomorpholiny sulfoxide, thiomorpholiny sulfone and oxadiazolyl.

The term "heteroaryl" as used herein represents an unsubstituted or substituted stable five or six membered monocyclic aromatic ring system or an unsubstituted or substituted nine or ten membered benzo-fused heteroaromatic ring system which consists of carbon atoms and from one to three heteroatoms selected from N, O or S, and wherein the nitrogen or sulfur heteroatoms may optionally be oxidized, and the nitrogen heteroatom may optionally be quaternized. The heteroaryl group may be attached at any heteroatom or carbon atom which results in the creation of a stable structure. Examples of heteroaryl groups include, but are not limited to pyridyl, pyridaziny, thienyl, furany, imidazolyl,

- 9 -

isoxazolyl, oxazolyl, pyrazolyl, pyrrolyl, thiazolyl, thiadiazolyl, triazolyl, benzimidazolyl, benzofuranyl, benzothienyl, benzisoxazolyl, benzoxazolyl, benzopyrazolyl, indolyl, benzothiazolyl, benzothiadiazolyl, benzotriazolyl or quinolinyl.

Preferred heteroaryl groups include pyridyl, pyrrolyl, pyrazinyl, thiadiazolyl, pyrazolyl, thienyl, triazolyl and quinolinyl.

The term "aralkyl" means an alkyl group substituted with an aryl group (e.g., benzyl, phenylethyl). Similarly, the term "aralkoxy" indicates an alkoxy group substituted with an aryl group (e.g., benzyloxy).

The term "acyl" as used herein means an organic radical having 1 to 6 carbon atoms (branched or straight chain) derived from an organic acid by removal of the hydroxyl group.


The term "amido" refers to the group $C(O)NH$ or $C(O)NH_2$.

The term "carbonyl" refers to the group $C(O)$.

The term "arylacrylamidoheteroaryl" as used herein means an aryl group attached to an ethylene which is attached to an amido group which is attached to a heteroaryl group, where the terms "aryl", "amido" and

- 10 -

"heteroaryl" are as defined above. The term "arylacrylamidoheteroaryl" can therefore refer to a group such as Aryl-C=C-C(O)-NH-heteroaryl, with a specific example of such an "arylacrylamidoheteroaryl" group being 5-(-Me-cinnamamido)triazol-3-yl O having the structure



It is intended that the definition of any substituent or variable at a particular location in a molecule be independent of its definitions elsewhere in that molecule. It is understood that substituents and substitution patterns on the compounds of this invention can be selected by one of ordinary skill in the art to provide compounds that are chemically stable and that can be readily synthesized by techniques known in the art as well as those methods set forth herein.

As used herein, the term "composition" is intended to encompass a product comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from combinations of the specified ingredients in the specified amounts.

- 11 -

The compounds of the present invention may also be present in the form of a pharmaceutically acceptable salt. The pharmaceutically acceptable salt generally takes a form in which the nitrogen of the amino-

5 substituted heterocycle or an amino-acid's basic side-chain is protonated with an inorganic or organic acid. Representative organic or inorganic acids include hydrochloric, hydrobromic, hydriodic, perchloric, sulfuric, nitric, phosphoric, acetic, propionic,

10 glycolic, lactic, succinic, maleic, fumaric, malic, tartaric, citric, benzoic, mandelic, methanesulfonic, hydroxyethanesulfonic, benzenesulfonic, oxalic, pamoic, 2-naphthalenesulfonic, p-toluenesulfonic, cyclohexanesulfamic, salicylic, saccharinic or

15 trifluoroacetic.

The present invention includes within its scope prodrugs of the compounds of this invention. In general, such prodrugs will be functional derivatives of

20 the compounds which are readily convertible *in vivo* into the required compound. Thus, in the methods of treatment of the present invention, the term "administering" shall encompass the treatment of the various disorders described with the compound

25 specifically disclosed or with a compound which may not be specifically disclosed, but which converts to the specified compound *in vivo* after administration to the patient. Conventional procedures for the selection and

- 12 -

preparation of suitable prodrug derivatives are described, for example, in "Design of Prodrugs", ed. H. Bundgaard, Elsevier, 1985.


5 Where the compounds according to this invention have at least one chiral center, they may accordingly exist as enantiomers. Where the compounds possess two or more chiral centers, they may additionally exist as diastereomers. It is to be understood that all such
10 isomers and mixtures thereof are encompassed within the scope of the present invention. Furthermore, some of the crystalline forms for the compounds may exist as polymorphs and as such are intended to be included in the present invention. In addition, some of the
15 compounds may form solvates with water (i.e., hydrates) or common organic solvents, and such solvates are also intended to be encompassed within the scope of this invention.

20 Until the present invention by Applicants, the known thrombin receptor agonists were peptides (i.e., PAR-1 peptide agonists) having a minimum sequence length of five amino acids. Applicants have unexpectedly discovered the instant thrombin receptor agonists which
25 are significantly truncated or that contain a heterocycle within the peptide backbone as a peptidomimetic unit.

- 13 -

Particularly preferred compounds of the present invention that could be useful as thrombin receptor agonists include those compounds shown in Table 1, which shows EC₅₀ values for platelet aggregation and binding IC₅₀ at the thrombin receptor. Amino acids bear the "L" absolute configuration unless denoted otherwise.

TABLE 1



#	Y	A ₁	A ₂	A ₃	EC ₅₀ (M)	
	Bndg (M)					
15	1	5-H ₂ N-1,2,4-triazol-3-yl		Cha	Arg	Phe
	0.75	1.9				
	2	5-Bromopyridin-3-yl	Cha	Arg	Phe	0.46
		1.7				
	3	2-Chromonyl	Cha	Arg	Phe	0.51 1.6
20	4	5-(Me-cinnamamido)triazol-3-yl		Cha	Arg	Phe
	0.76	4.2				
	5	5-Naphthylacrylamidotriazol-3-yl		Cha	Arg	Phe
	0.99	16				
	6	Quinoxalin-2-yl	Cha	Arg	Phe	1.03
25		3.1				
	7	5-(Cl-cinnamamido)triazol-3-yl		Cha	Arg	Phe
	1.14	1.4				
	8	6-Aminopyridin-3-yl		Cha	Arg	Phe
	1.24	2.0				

- 14 -

	9	5-H ₂ N-1,2,4-triazol-3-yl	Cha	Arg	Phe-Arg	
	1.21	0.5				
	10	Thiadiazol-4-yl	Cha	Arg	Phe	
	1.28	9.4				
5	115-	(2,3-diMeO-cinnamamido)triazol-3-yl	Cha	Arg	Phe	1.64
	6.4					
	12	5-(-F-cinnamamido)triazol-3-yl	Cha	Arg	Phe	
	1.72	2.2				
	13	5-(m-NO ₂ -cinnamamido)triazol-3-yl	Cha	Arg	Phe	
10	1.89	8.5				
	14	5-(o-NO ₂ -cinnamamido)triazol-3-yl	Cha	Arg	Phe	
	1.89	1.6				
	15	Pyridin-3-yl	Cha	Arg	Phe	1.97
	1.3					
15	16	5-(m-Cl-cinnamamido)triazol-3-yl	Cha	Arg	Phe	
	2.38	15				
	17	5-H ₂ N-1,2,4-triazol-3-yl	Phe	Arg	Phe	
	2.4	2.1				
	18	5-H ₂ N-1,2,4-triazol-3-yl	Cha	Lys	Phe	
20	2.67	5.4				
	19	5-H ₂ N-1,2,4-triazol-3-yl	Cha	Arg	Cha	
	2.8	2.0				
	20	5-H ₂ N-1,2,4-triazol-3-yl	Cha	Arg		
		Phgly 3.13 7.1				
25	21	5-(thiophen-2-ylacrylamido)triazol-3-yl	Cha	Arg	Phe	
	2.9	4.4				
	22	3-H ₂ N-pyrazin-2-yl	Cha	Arg	Phe	4.2
	8.5					
	23	trans 2-(3-pyridyl)ethylenyl	Cha	Arg	Phe	2.6
30	5.5					
	24	5-(p-MeO-cinnamamido)triazol-3-yl	Cha	Arg	Phe	
	5.6	4				

- 15 -

	25	5-(p-CN-cinnamamido)triazol-3-yl	Cha	Arg	Phe	
	5.8	31				
	26	5-(p-F-cinnamamido)triazol-3-yl	Cha	Arg	Phe	
	7.9	1.7				
5	27	2-H ₂ N-pyridin-3-yl	Cha	Arg	Phe	8.8
		2.3				
	#	X	A ₁	A ₂	A ₃	EC ₅₀ (M)
		Bndg (M)				
	28	5-H ₂ N-1,2,4-triazol-3-yl	Cha	Arg	Tyr	
10	9.7	6				
	29	5-H ₂ N-1,2,4-triazol-3-yl		Cha	Arg	2-
	Thala	11	4			
	30	Pyridin-2-yl	Cha	Arg	Phe	
	14	4				
15	31	5-(p-Phenyl-cinnamamido)triazol-3-yl	Cha	Arg	Phe	
	25	12				
	32	N-(p-F-phenylalanyl)-piperidin-3-yl		Cha	Arg	Phe
		26	2			
	33	5-(Cinnamamido)triazol-3-yl	Cha	Arg	Phe	
20	28	6				
	34	5-(-phenyl-cinnamamido)triazol-3-yl	Cha	Arg	Phe	
	28	3				
	38	3-aminophenyl	Cha	Arg	Phe	2.9
		2				
25	39	1-biphenyl	Cha	Arg	Phe	
	(56%) ¹	13				
	40	2-biphenylenyl		Cha	Arg	Phe
	1.2	2.4				
	41	benzimidazol-5-yl	Cha	Arg	Phe	4
30		4				

1. Percent aggregation induced at 50 M.

- 16 -

Particularly preferred compounds of the present invention that could be useful as thrombin receptor antagonists or mixed agonists / antagonists include those compounds shown in Table 2, which shows IC_{50} values for inhibition of platelet aggregation (gel-filtered platelets aggregation induced by thrombin) and binding IC_{50} at the thrombin receptor. Amino acids bear the "L" absolute configuration unless denoted otherwise.

TABLE 2



#	Y	A ₁	A ₂	X		IC_{50} (M)
	<u>Bndg (M)</u>					
7	5-(Cl-cinnamamido)	triazol-3-yl	Cha	Arg	CO	
	3.6	1.4				
21	5-(Thien-2-ylacrylamido)	triazol-3-yl	Cha	Arg		
CO	5.4	4.4				
33	5-(Cinnamamido)	triazol-3-yl	Cha	Arg	CO	
	9.8	6.4				
4	5-(Me-cinnamamido)	triazol-3-yl	Cha	Arg	CO	
	14	4.2				
34	5-(Ph-cinnamamido)	triazol-3-yl	Cha	Arg	CO	
19		3.1				
35	6-Cinnamamidopyridin-3-yl		Cha	Arg	CO	
	26	2.2				
36	5-Cl, 3-Me-benzothiophen-2-yl		Cha	Arg	SO ₂	
	8.4	4.6				
26	5-(p-F-cinnamamido)	triazol-3-yl	Cha	Arg	CO	
	13	1.7				

- 17 -

37	Benzothiophen-2-yl	Cha	Arg	CO
21	2.3			
42	1-naphthyl	Cha	Arg	SO ₂
	7.8		1.7	
5 43	2-naphthyl	Cha	Arg	SO ₂
	12.52.1			

Particularly preferred compounds of the present invention are:

(5-Bromopyridin-3-yl) carbonyl-cyclohexylalanyl-arginyl-phenylalanineamide;

2-Chromonyl carbonyl-cyclohexylalanyl-arginyl-phenylalanineamide;

(5-Aminotriazol-3-yl) carbonyl-cyclohexylalanyl-arginyl-phenylalanineamide;

[5-(-Methyl) cinnamamidotriazol-3-yl] carbonyl-cyclohexylalanyl-arginyl-phenylalanineamide;

{5-[3-(1-Naphthyl) acrylamido] triazol-3-yl} carbonyl-cyclohexylalanyl-arginyl-phenylalanineamide;

[Quinoxalin-2-yl] carbonyl-cyclohexylalanyl-arginyl-phenylalanineamide;

[5-(o-Chlorocinnamamido) triazol-3-yl] carbonyl-cyclohexylalanyl-arginyl-phenylalanineamide;

(6-Aminopyridin-3-yl) carbonyl-cyclohexylalanyl-arginyl-phenylalanineamide;

(5-Aminotriazol-3-yl) carbonyl-phenylalanyl-arginyl-phenylalanyl-arginineamide;

(5-Aminotriazol-3-yl) carbonyl-cyclohexylalanyl-lysiny-phenylalanineamide;

- 18 -

{5-[3-(2-Thienyl)acrylamido]triazol-3-yl}carbonyl-cyclohexylalanyl-arginyl-phenylalanineamide;


5 [5-cinnamamidotriazol-3-yl]carbonyl-cyclohexylalanyl-arginyl-phenylalanineamide;

(6-Cinnamamidopyridin-3-yl)carbonyl-cyclohexylalanyl-arginyl-phenylalanineamide; or

10 (5-Chloro-3-methyl-benzothiophen-2-yl)carbonyl-cyclohexylalanyl-arginyl-phenylalanineamide;

and pharmaceutically acceptable salts thereof.

15 The modulators of this invention may be prepared using solid phase chemistry as shown in Scheme A.


Scheme A:

Thus, the resin is Fmoc deprotected via agitation
 using piperidine or any dialkyl amine in an appropriate
 solvent such as DMF, washed with fresh solvent and then
 coupled with an Fmoc protected amino-acid A₃ using
 suitable coupling reagents such as
 diisopropylcarbodiimide (DIC) or

- 20 -

dicyclohexylcarbodiimide (DCC) or Bis(2-oxo-3-oxazolidinyl)phosphinic chloride (BOP-Cl) along with hydroxybenzotriazole (HOBT) in DMF or another dipolar aprotic solvent. After washing, the amino-acid loaded resin is Fmoc deprotected as above with a dialkyl amine in DMF and coupled again with the second Fmoc protected amino-acid A₂ with a coupling agent as above. This Fmoc dipeptido loaded resin is further deprotected with dialkylamine in DMF, or appropriate solvent, and coupled again to the Fmoc amino-acid A₁ using a coupling agent as above. The tripeptido resin is deprotected with a dialkyl amine in DMF and coupled to the substituted acid Y-X-OH with an appropriate coupling agent. At this point the product may be cleaved from the resin using a strong acid such as trifluoroacetic (TFA) acid in any inert solvent, such as DCM, to give the peptide amide 1. However, if the Y group possesses a protected amino functionality, such as Fmoc-amino, this may be deprotected and the amino-peptide product cleaved from the resin or the freed-up amine can be reacted further prior to cleavage of the peptide from the resin. Thus, the Fmoc can be removed using a dialkyl amine in DMF as above and the amino-peptide can be 1) cleaved from the resin with TFA or 2) coupled with an aryl acrylic acid, using appropriate coupling agent such as BOP-Cl, and the product can be cleaved from the resin with TFA to afford peptide 11. Following this general route, preparation of compound 1 as outlined in Scheme 1 is prepared, as described in more detail in the experimental section.


Scheme 1:



Alternatively, the compounds may be prepared via normal solution phase chemistry as shown in Scheme B.

5

Scheme B:



10

15

Thus, Fmoc protected amino-acid A₃ can be coupled with an amine ZH using the normal peptide coupling agents such as DIC or DCC and HOBT in dipolar aprotic solvents such as acetonitrile (ACN) or dimethyl formamide (DMF). The isolated product can then be Fmoc deprotected with diethylamine (DEA), or another dialkyl amine, in a dipolar aprotic solvent such as ACN, and the resultant amine coupled to the second Fmoc protected amino-acid A₂. This dipeptide can similarly be deprotected as above and coupled with an appropriate coupling agent with Fmoc protected amino-acid A₁ to give the Fmoc-tripeptide. Deprotection of the Fmoc group

- 24 -


with a dialkyl amine is followed by coupling of this tripeptide to the acid Y-X-OH using a coupling agent such as DIC with HOBT to give the product 111. Using this general route, synthesis of compound 2 was prepared as outlined in Scheme 2, and further described in the experimental section.

10

15

Scheme 2:

- 25 -



5 The utility of the compounds of formula (I) to act as thrombin receptor modulators can be determined according to the procedures described in Examples 4 to 5 herein. The present invention therefore provides a method of treating a condition mediated by modulation of the thrombin receptor in a subject in need thereof which

- 26 -

comprises administering any of the compounds or pharmaceutical compositions as defined herein in a quantity effective to treat the condition. Additionally, the present invention includes the use of a compound of formula (I) for the preparation of a medicament for the treatment of a condition mediated by modulation of the thrombin receptor. The compound may be administered to a patient by any conventional route of administration, including, but not limited to, intravenous, oral, subcutaneous, intramuscular, intradermal and parenteral.

The present invention also provides pharmaceutical compositions comprising one or more compounds of this invention in association with a pharmaceutically acceptable carrier.

To prepare the pharmaceutical compositions of this invention, one or more compounds of formula (I) or salt thereof of the invention as the active ingredient, is intimately admixed with a pharmaceutical carrier according to conventional pharmaceutical compounding techniques, which carrier may take a wide variety of forms depending of the form of preparation desired for administration, e.g., oral or parenteral such as intramuscular. In preparing the compositions in oral dosage form, any of the usual pharmaceutical media may be employed. Thus, for liquid oral preparations, such as for example, suspensions, elixirs and solutions,

- 27 -

suitable carriers and additives include water, glycols, oils, alcohols, flavoring agents, preservatives, coloring agents and the like; for solid oral preparations such as, for example, powders, capsules, caplets, gelcaps and tablets, suitable carriers and additives include starches, sugars, diluents, granulating agents, lubricants, binders, disintegrating agents and the like. Because of their ease in administration, tablets and capsules represent the most advantageous oral dosage unit form, in which case solid pharmaceutical carriers are obviously employed. If desired, tablets may be sugar coated or enteric coated by standard techniques. For parenterals, the carrier will usually comprise sterile water, through other ingredients, for example, for purposes such as aiding solubility or for preservation, may be included. Injectable suspensions may also be prepared, in which case appropriate liquid carriers, suspending agents and the like may be employed. The pharmaceutical compositions herein will contain, per dosage unit, e.g., tablet, capsule, powder, injection, teaspoonful and the like, an amount of the active ingredient necessary to deliver an effective dose as described above. The pharmaceutical compositions herein will contain, per unit dosage unit, e.g., tablet, capsule, powder, injection, suppository, teaspoonful and the like, of from about 0.03 mg to 100 mg/kg (preferred 0.1-30 mg/kg) and may be given at a dosage of from about 0.1-300

- 28 -

mg/kg/day (preferred 1-50 mg/kg/day). The dosages, however, may be varied depending upon the requirement of the patients, the severity of the condition being treated and the compound being employed. The use of either daily administration or post-periodic dosing may be employed.

Preferably these compositions are in unit dosage forms from such as tablets, pills, capsules, powders, granules, sterile parenteral solutions or suspensions, metered aerosol or liquid sprays, drops, ampoules, autoinjector devices or suppositories; for oral parenteral, intranasal, sublingual or rectal administration, or for administration by inhalation or insufflation. Alternatively, the composition may be presented in a form suitable for once-weekly or once-monthly administration; for example, an insoluble salt of the active compound, such as the decanoate salt, may be adapted to provide a depot preparation for intramuscular injection. For preparing solid compositions such as tablets, the principal active ingredient is mixed with a pharmaceutical carrier, e.g. conventional tableting ingredients such as corn starch, lactose, sucrose, sorbitol, talc, stearic acid, magnesium stearate, dicalcium phosphate or gums, and other pharmaceutical diluents, e.g. water, to form a solid preformulation composition containing a homogeneous mixture of a compound of the present invention, or a pharmaceutically acceptable salt thereof. When referring to these

- 29 -

preformulation compositions as homogeneous, it is meant that the active ingredient is dispersed evenly throughout the composition so that the composition may be readily subdivided into equally effective dosage forms such as tablets, pills and capsules. This solid preformulation composition is then subdivided into unit dosage forms of the type described above containing from 0.1 to about 500 mg of the active ingredient of the present invention.

The tablets or pills of the novel composition can be coated or otherwise compounded to provide a dosage form affording the advantage of prolonged action. For example, the tablet or pill can comprise an inner dosage and an outer dosage component, the latter being in the form of an envelope over the former. The two components can be separated by an enteric layer which serves to resist disintegration in the stomach and permits the inner component to pass intact into the duodenum or to be delayed in release. A variety of material can be used for such enteric layers or coatings, such materials including a number of polymeric acids with such materials as shellac, cetyl alcohol and cellulose acetate.

The liquid forms in which the novel compositions of the present invention may be incorporated for administration orally or by injection include, aqueous solutions, suitably flavoured syrups, aqueous or oil suspensions, and flavoured emulsions with edible oils such as cottonseed oil, sesame oil, coconut oil or peanut

- 30 -

oil, as well as elixirs and similar pharmaceutical vehicles. Suitable dispersing or suspending agents for aqueous suspensions, include synthetic and natural gums such as tragacanth, acacia, alginate, dextran, sodium carboxymethylcellulose, methylcellulose, polyvinylpyrrolidone or gelatin.

Where the processes for the preparation of the compounds according to the invention give rise to mixture of stereoisomers, these isomers may be separated by conventional techniques such as preparative chromatography. The compounds may be prepared in racemic form, or individual enantiomers may be prepared either by enantiospecific synthesis or by resolution. The compounds may, for example, be resolved into their components enantiomers by standard techniques, such as the formation of diastereomeric pairs by salt formation with an optically active acid, such as (-)-di-p-toluoyl-d-tartaric acid and/or (+)-di-p-toluoyl-l-tartaric acid followed by fractional crystallization and regeneration of the free base. The compounds may also be resolved by formation of diastereomeric esters or amides, followed by chromatographic separation and removal of the chiral auxiliary. Alternatively, the compounds may be resolved using a chiral HPLC column.

During any of the processes for preparation of the compounds of the present invention, it may be necessary

- 31 -

and/or desirable to protect sensitive or reactive groups on any of the molecules concerned. This may be achieved by means of conventional protecting groups, such as those described in Protective Groups in Organic
5 Chemistry, ed. J.F.W. McOmie, Plenum Press, 1973; and T.W. Greene & P.G.M. Wuts, Protective Groups in Organic Synthesis, John Wiley & Sons, 1991. The protecting groups may be removed at a convenient subsequent stage using methods known from the art.

10

The method of treating conditions modulated by the thrombin receptor described in the present invention may also be carried out using a pharmaceutical composition comprising any of the compounds as defined herein and a
15 pharmaceutically acceptable carrier. The pharmaceutical composition may contain between about 0.01 mg and 100 mg, preferably about 5 to 50 mg, of the compound, and may be constituted into any form suitable for the mode of administration selected. Carriers include necessary and
20 inert pharmaceutical excipients, including, but not limited to, binders, suspending agents, lubricants, flavorants, sweeteners, preservatives, dyes, and coatings. Compositions suitable for oral administration include solid forms, such as pills, tablets, caplets,
25 capsules (each including immediate release, timed release and sustained release formulations), granules, and powders, and liquid forms, such as solutions, syrups, elixers, emulsions, and suspensions. Forms useful for

- 32 -

parenteral administration include sterile solutions, emulsions and suspensions.

Advantageously, compounds of the present invention may be administered in a single daily dose, or the total daily dosage may be administered in divided doses of two, three or four times daily. Furthermore, compounds for the present invention can be administered in intranasal form via topical use of suitable intranasal vehicles, or via transdermal skin patches well known to those of ordinary skill in that art. To be administered in the form of a transdermal delivery system, the dosage administration will, of course, be continuous rather than intermittent throughout the dosage regimen.

For instance, for oral administration in the form of a tablet or capsule, the active drug component can be combined with an oral, non-toxic pharmaceutically acceptable inert carrier such as ethanol, glycerol, water and the like. Moreover, when desired or necessary, suitable binders; lubricants, disintegrating agents and coloring agents can also be incorporated into the mixture. Suitable binders include, without limitation, starch, gelatin, natural sugars such as glucose or beta-lactose, corn sweeteners, natural and synthetic gums such as acacia, tragacanth or sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride and the like. Disintegrators include,

- 33 -

without limitation, starch, methyl cellulose, agar, bentonite, xanthan gum and the like.

5 The liquid forms in suitably flavored suspending or dispersing agents such as the synthetic and natural gums, for example, tragacanth, acacia, methyl-cellulose and the like. For parenteral administration, sterile suspensions and solutions are desired. Isotonic preparations which generally contain suitable preservatives are employed when
10 intravenous administration is desired.

 The compound of the present invention can also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar
15 vesicles, and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine or phosphatidylcholines.

 Compounds of the present invention may also be
20 delivered by the use of monoclonal antibodies as individual carriers to which the compound molecules are coupled. The compounds of the present invention may also be coupled with soluble polymers as targetable drug carriers. Such polymers can include
25 polyvinylpyrrolidone, pyran copolymer, polyhydroxypropylmethacrylamidephenol, polyhydroxy-ethylaspartamidephenol, or polyethyl eneoxydepolylysine substituted with palmitoyl residue. Furthermore, the

- 34 -

compounds of the present invention may be coupled to a class of biodegradable polymers useful in achieving controlled release of a drug, for example, polylactic acid, polyepsilon caprolactone, polyhydroxy butyeric acid, polyorthoesters, polyacetals, polydihydropyrans, polycyanoacrylates and cross-linked or amphipathic block copolymers of hydrogels.

Compounds of this invention may be administered in any of the foregoing compositions and according to dosage regimens established in the art whenever treatment of thrombotic disorders is required.

The daily dosage of the products may be varied over a wide range from 0.01 to 1,000 mg per adult human per day. For oral administration, the compositions are preferably provided in the form of tablets containing, 0.01, 0.05, 0.1, 0.5, 1.0, 2.5, 5.0, 10.0, 15.0, 25.0, 50.0, 100, 150, 200, 250 and 500 milligrams of the active ingredient for the symptomatic adjustment of the dosage to the patient to be treated. An effective amount of the drug is ordinarily supplied at a dosage level of from about 0.1 mg/kg to about 300 mg/kg of body weight per day. Preferably, the range is from about 1 to about 50 mg/kg of body weight per day. The compounds may be administered on a regimen of 1 to 4 times per day.

- 35 -

Optimal dosages to be administered may be readily determined by those skilled in the art, and will vary with the particular compound used, the mode of administration, the strength of the preparation, the mode of administration, and the advancement of the disease condition. In addition, factors associated with the particular patient being treated, including patient age, weight, diet and time of administration, will result in the need to adjust dosages.

To prepare the pharmaceutical compositions of this invention, one or more compounds of formula (I) or salt thereof of the invention as the active ingredient, is intimately admixed with a pharmaceutical carrier according to conventional pharmaceutical compounding techniques, which carrier may take a wide variety of forms depending of the form of preparation desired for administration, e.g., oral or parenteral such as intramuscular. In preparing the compositions in oral dosage form, any of the usual pharmaceutical media may be employed. Thus, for liquid oral preparations, such as for example, suspensions, elixirs and solutions, suitable carriers and additives include water, glycols, oils, alcohols, flavoring agents, preservatives, coloring agents and the like; for solid oral preparations such as, for example, powders, capsules, caplets, gelcaps and tablets, suitable carriers and additives include starches, sugars, diluents, granulating agents, lubricants, binders, disintegrating agents and the like. Because of their ease in

- 36 -

administration, tablets and capsules represent the most advantageous oral dosage unit form, in which case solid pharmaceutical carriers are obviously employed. If desired, tablets may be sugar coated or enteric coated by standard techniques. For parenterals, the carrier will usually comprise sterile water, through other ingredients, for example, for purposes such as aiding solubility or for preservation, may be included. Injectable suspensions may also be prepared, in which case appropriate liquid carriers, suspending agents and the like may be employed. The pharmaceutical compositions herein will contain, per dosage unit, e.g., tablet, capsule, powder, injection, teaspoonful and the like, an amount of the active ingredient necessary to deliver an effective dose as described above. The pharmaceutical compositions herein will contain, per unit dosage unit, e.g., tablet, capsule, powder, injection, suppository, teaspoonful and the like, of from about 0.03 mg to 100 mg/kg (preferred 0.1-30 mg/kg) and may be given at a dosage of from about 0.1-300 mg/kg/day (preferred 1-50 mg/kg/day). The dosages, however, may be varied depending upon the requirement of the patients, the severity of the condition being treated and the compound being employed. The use of either daily administration or post-periodic dosing may be employed.

The following Examples are set forth to aid in the understanding of the invention, and are not intended and should not be construed to limit in any way the invention set forth in the claims which follow thereafter.

- 37 -

Protected amino acids were purchased from Novabiochem, Synthetech or Bachem Bioscience Inc. All other chemicals were purchased from Aldrich Chemical Company, Inc. High field ^1H NMR spectra were recorded on a Bruker AC-300 spectrometer at 300 MHz, and coupling constants are given in Herz. Microanalyses were performed at Robertson Microlit Laboratories, Inc., Madison, New Jersey. In the examples and throughout this application, the following abbreviations have the meanings recited hereinafter:

ACN	Acetonitrile
BOP-Cl	Bis(2-oxo-3-oxazolidinyl)phosphinic chloride
DCM	Dichloromethane
DCC	Dicyclohexylcarbodiimide
DIC	Diisopropylcarbodiimide
DEA	Diethylamine
DMAP	4-Dimethylaminopyridine
DMF	N, N-Dimethylformamide
Et ₂ O	Diethyl ether
Fmoc	Fluorenylmethoxycarbonyl
HOBT	Hydroxybenzotriazole
Me	Methyl
Ph	Phenyl
Pmc	2,2,5,7,8-Pentamethylchroman-6-sulfonyl
RT	room temperature
TFA	Trifluoroacetic acid

Amino acid abbreviations are defined below:


Arg Arginine

- 38 -

	Asn	Asparagine
	Cha	Cyclohexylalanine
	Gln	Glutamine
	hArg	Homoarginine
5	hPhe	Homophenylalanine
	Ile	Isoleucine
	Leu	Leucine
	Lys	Lysine
	Orn	Ornithine
10	Phe	Phenylalanine
	Phgly	Phenylglycine
	2-Thala	2-Thienylalanine
	3-Thala	3-Thienylalanine
	Trp	Tryptophan
15	Tyr	Tyrosine

EXAMPLE 1

20 N-(5-Amino-1,2,4-triazol-3-yl)carbonyl)-
cyclohexylalanyl-argininyl-phenylalanine amide (1)



1

25 Rink amide resin (4.0 g, 3.24 mm) was placed in a solid
phase hour-glass reactor and agitated (nitrogen

- 39 -

bubbling) with 20% piperidine / DMF (25 mL) for 1 hr. The solution was drained off and the resin was washed with DMF (4X), DCM (3X), and DMF (3X). The resin was combined with Fmoc-Phe-OH (3.77 g, 9.75 mm), HOBT (1.5 g, 9.75 mm) and DIC (1.23 g, 9.75 mm) in DMF (25 mL) and agitated in the reactor for 16 hr at ambient temperature. The solution was drained and resin was washed with DMF (5X) and DCM (4X) and dried in vacuo. A portion (1.0 g, 0.80 mm) was agitated in the reactor at ambient temperature with 20% piperidine / DMF (15 mL) for 1 hr, the solution was drained and resin washed with DMF (4X), DCM (3X), and DMF (3X). This was combined with Fmoc-Arg(Pmc)-OH (1.84 g, 2.4 mm), HOBT (0.37 g, 2.4 mm), and DIC (0.31 g, 2.4 mm) in DMF (10 mL) and agitated at ambient temperature for 16 hr. After draining the solution, the resin was washed with DMF (4X), and DCM (3X). Half of this batch was washed with DMF (3X) and then agitated with 20% piperidine / DMF (10 mL) for 1 hr. The solution was drained and the resin was washed with DMF (4X), DCM (3X), and DMF (3X) and then combined with Fmoc-Cha-OH (0.47 g, 1.2 mm), HOBT (0.18 g, 1.2 mm) and DIC (0.15 g, 1.2 mm) in DMF (10 mL) and agitated for 16 hr at ambient temperature. The solution was drained and the resin was washed with DMF (4X), DCM (3X), and DMF (3X) and treated with 20% piperidine / DMF for 1 hr. The solution was drained and the resin was washed with DMF (4X) and then combined with Fmoc-3-amino-1,2,4 triazole-5-carboxylic acid (0.42 g, 1.2 mm), HOBT (0.18 g, 1.2 mm), and DIC (0.15 g, 1.2 mm) in DMF (10 mL) and agitated at ambient temperature for 16 hr. The solution was drained off, and the resin was washed with DMF (4X), DCM (3X), and DMF (3X) and then agitated


- 40 -

with 20% piperidine / DMF (15 mL) for 1 hr. The solution was drained and the resin was washed with DMF (4X), and DCM (3X) and dried under a nitrogen stream. The resin was then agitated with 99% TFA (20 mL) at ambient temperature for 1.5 hr. The TFA solution was collected and evaporated in vacuo to an oil, which was triturated with Et₂O (3X) to give a white solid (152 mg).

Purification was accomplished via reverse phase HPLC using 0.16% TFA in ACN / 0.20% TFA in water (35:65) and upon lyophilization afforded white floccular solid 1 (120 mg). Anal. calcd. for C₂₇H₄₁N₁₁O₄ • 2.25 TFA • 1.0 H₂O (858.26): C, 44.08; H, 5.13; N, 17.95; F, 14.94. Found: C, 43.82; H, 5.27; N, 17.91; F, 14.50. FAB-MS m/e 584.9 (MH⁺). ¹H NMR (DMSO / D₂O) 7.25 (m, 5H), 4.5 (dd, 1H), 4.4 (dd, 1H), 4.28 (dd, 1H), 3.1 - 2.95 (m, 3H), 2.8 (dd, 1H), 1.8 - 1.35 (m, 11H), 1.3 - 1.0 (m, 4H), 0.9 (m, 2H).

EXAMPLE 2

N-(5-Bromopyridin-3-yl-carbonyl)-cyclohexylalanyl-argininyl-phenylalanine amide (2)



- 41 -

Fmoc-phenylalanine amide (3.87 g, 10 mm) was stirred in ACN (100 mL) and DEA (5 mL) was added and stirred at RT for 1 hr. The solution was evaporated in vacuo to an oil, which was triturated 3X with hexane (100 mL) and dissolved in ACN (100 mL); Fmoc-Arg(PMC)-OH (6.63 g, 10 mm) and HOBT (1.53 g, 10 mm) were added, followed by DCC (4.1 g, 20 mm) and solution was stirred at RT. The urea by-product was filtered and the filtrate was evaporated in vacuo to an oil, which was triturated 3X with hexane (100 mL). The crude product was stirred in ACN (100 mL) and DEA (5 mL) was added and stirred at RT for 1 hr. The solution was evaporated in vacuo to an oil, which was triturated 3X with hexane (100 mL) to a solid. This dipeptide was combined in ACN (100 mL) with Fmoc-Cha-OH (3.93 g, 10 mm) and HOBT (1.53 g, 10 mm) and then DIC (2.52 g, 20 mm) was added and reaction stirred at RT for 16 hr. Solid tripeptide was filtered; tripeptide may also be recovered from the filtrate via evaporation and silica gel column chromatography. The tripeptide (5.0 g, 5 mm) was stirred in ACN (100 mL) containing DEA (5 mL) until deprotection is complete and then evaporated in vacuo and triturated with hexane 3X (100 mL). This was combined in ACN (100 mL) and 5-bromonicotinic acid (1.01 g, 5 mm) and HOBT (0.76 g, 5 mm) were added in, followed by DIC (1.26 g, 10 mm) and stirred at RT for 16 hr. The solution was evaporated in vacuo and purified via silica gel column chromatography. This protected product was then stirred with DCM/TFA (1:1; 50 mL) for 1 hr and then evaporated in vacuo to an oil, which was triturated 3X with Et₂O (100 mL) to afford white solid 3 as a trifluoroacetate salt: MS m/e 657.4/659.4 (MH⁺).

- 42 -

¹H NMR (DMSO) 9.0 (s, 1H), 8.9 (s, 1H), 8.8 (d, 1H),
8.5 (s, 1H), 8.2 (d, 1H), 7.8 (d, 1H), 7.4 (s, 2H), 7.1
- 7.3 (m, 6H), 4.5 (m, 2H), 4.2 (q, 1H), 4.0 (s, 1H),
3.0 (m, 1H), 2.8 (q, 1H), 1.3 - 1.8 (m, 12H), 1.1 (m,
4H), 0.9 (m, 3H).

EXAMPLE 3

As a specific embodiment of an oral composition, 100
mg of the compound 1 of Example 1 is formulated with
sufficient finely divided lactose to provide a total
amount of 580 to 590 mg to fill a size 0 hard gel
capsule.

BIOLOGY

The compounds of the present invention modulate
platelet activation induced by thrombin's proteolytic
cleavage of its platelet surface receptor, and thereby
activate / inhibit platelet aggregation. Compounds that
exhibit agonist activity may be expected to aid in wound
healing and tissue repair, while antagonist compounds
may be useful in treating platelet-mediated thrombotic
disorders such as arterial and venous thrombosis, acute
myocardial infarction, reocclusion following
thrombolytic therapy and angioplasty, and a variety of
vaso-occlusive disorders.

EXAMPLE 4

IN VITRO THROMBIN RECEPTOR BINDING ASSAY.

- 43 -

CHRF membranes (Jones, *Biochim. Biophys. Acta* 1992, 1136, 272) are thawed from -70°C, centrifuged at maximum speed for 5 min, washed twice with binding buffer (50 mM HEPES containing 5 mM MgCl₂ and 0.1% BSA), and re-suspended in binding buffer (25 g/100 mL). 100 l membranes are added to the 24-Wallac plates and delivered to the Tomtech apparatus. In a typical experiment, 6 l of samples (from a 125 g/mL intermediary plate, 20%DMSO) and 44 l buffer are delivered to the plates (final conc. of compounds is 3.7 g/mL, 0.6% DMSO). Similarly, 6 l 20%DMSO and 44 l buffer are delivered to both column 1 (NSB) and column 12 (TB). 10 l Ser-pFPhe-Har-Leu-Har-Lys-Tyr-NH₂ (721-40; 500 M in deionized water) is added to column 1. 50 l tritiated 721-40 (specific activity 46 Ci/mmol) is added to all the wells. The plates are mixed well for 20 seconds, incubated for 30 min, and then harvested with 10 mM HEPES/138 mM NaCl using the Skatron harvester. The filters (GF/C Brandel FPXLR 296 filters are presoaked 3 h in 0.5% polyethylenimine in HEPES/0.1 M N-acetylglucosamine) are set in saran wrap and dried for 3 min in the microwave, and placed in sample bags (Wallac 1450-432). 4.5 mL scintillation fluid (Wallac, Betaplate Scint 1205-440) is added. The bags are sealed, placed in filter cassettes (Wallac 1450-104), and analyzed on the microbeta counter.

EXAMPLE 5

30

- 44 -

IN VITRO AGONIST PLATELET AGGREGATION ASSAY /
INHIBITION OF THROMBIN-INDUCED GEL-FILTERED PLATELET
AGGREGATION ASSAY.


5 The percentage of platelet aggregation is calculated
as an increase in light transmission of compound-treated
platelet concentrate vs. control-treated platelet
concentrate. Human blood is obtained from drug free,
normal donors into tubes containing 0.13 M sodium
10 citrate. Platelet rich plasma (PRP) is collected by
centrifugation of whole blood at 200 x g for 10 min at
25°C. The PRP (5 mL) is gel filtered through Sepharose
2B (bed volume 50 mL), and the platelet count is
adjusted to 2×10^7 platelets per sample. The following
15 constituents are added to a siliconized cuvette:
concentrated platelet filtrate and Tyrode's buffer (0.14
M NaCl, 0.0027 M KCl, 0.012 M NaHCO₃, 0.76 mM Na₂HPO₄,
0.0055 M glucose, 2 mg/mL BSA and 5.0 mM HEPES @ pH 7.4)
in an amount equal to 350 μ l, 50 μ l of 20 mM calcium and
20 50 μ l of the test compound. Aggregation is monitored in
a BIODATA aggregometer for the 3 min following the
addition of agonist (thrombin 50 μ l of 1 unit/mL).

25 While the foregoing specification teaches the
principles of the present invention, with examples
provided for the purpose of illustration, it will be
understood that the practice of the invention encompasses
all of the usual variations, adaptations and/or
modifications as come within the scope of the following
30 claims and their equivalents.



WE CLAIM:

1. A compound represented by the general formula (I):



10 wherein A_1 is an alkyl amino acid residue selected from Cha, Leu and Ile, an amino alkyl amino acid residue selected from Arg and Lys, or an aryl amino acid residue selected from Phe, substituted Phe, Tyr, or Trp;

15 A_2 is an amino alkyl amino acid residue selected from Lys, Orn, Arg, and homo Arg;

20 A_3 is an aryl amino acid residue selected from Phe, substituted Phe, homo Phe, Tyr, Trp, phgly, 2-Thala and 3-Thala, an alkyl amino acid residue selected from Cha, Leu and Ile, an amido alkyl amino acid selected from Asn and Gln, or an amino alkyl amino acid residue selected from Arg, homo Arg, Orn and Lys;

25 X is selected from CO, CS, or SO_2 ;

Y is selected from aryl, substituted aryl, heterocycloalkyl, substituted heterocycloalkyl, heteroaryl, substituted heteroaryl, heteroarylethylenyl,



- 46 -

substituted heteroarylethylenyl,
arylacrylamidoheteroaryl, substituted
arylacrylamidoheteroaryl, heteroarylacrylamidoheteroaryl
and substituted heteroarylacrylamidoheteroaryl, provided
that Y is not pyrrolidinyl, phenyl or 2-aminophenyl;

Z is selected from NH₂, NH-alkyl, NH-aralkyl, or an
amino alkyl amino acid residue selected from Arg-NH₂; and

wherein all amino acids are of the L configuration;
and pharmaceutically acceptable salts thereof.

2. The compound of Claim 1, wherein

A₃ is an aryl amino acid residue selected from Phe,
substituted Phe, Tyr, Trp, phgly, 2-Thala and 3-Thala,
an alkyl amino acid residue selected from Cha, Leu and
Ile, an amido alkyl amino acid selected from Asn and
Gln, or an amino alkyl amino acid residue selected
from Arg, homo Arg, Orn and Lys; and

Y is selected from heteroaryl, substituted
heteroaryl, arylacrylamidoheteroaryl, and substituted
arylacrylamidoheteroaryl;

and pharmaceutically acceptable salts thereof.



- 47 -

3. The compound of Claim 2, wherein

A₁ is an alkyl amino acid residue selected from Cha, Leu and Ile, or an aryl amino acid residue selected from Phe, substituted Phe, Tyr, or Trp;

A₂ is an amino alkyl amino acid residue selected from Lys or Arg;

A₃ is an aryl amino acid residue selected from Phe, substituted Phe, Tyr, Trp, Phgly and 2-Thala;

X is selected from CO or SO₂; and pharmaceutically acceptable salts thereof.

4. The compound of Claim 3, wherein

X is CO;

Y is selected from benzothiophenyl, substituted benzothiophenyl, pyridinyl, substituted pyridinyl, triazolyl, substituted triazolyl, chromonyl, quinoxaliny, thiadiazolyl, substituted thiadiazolyl, pyrazinyl, substituted pyrazinyl, pyridylethylenyl, substituted pyridylethylenyl, cinnamamido-triazolyl, substituted cinnamamido-triazolyl, thiophenylacrylamido-triazolyl, or naphthylacrylamido-triazolyl;



- 48 -

Z is selected from NH_2 or Arg-NH_2 ;

and pharmaceutically acceptable salts thereof.

5

5. The compound of Claim 4, wherein

A_1 is selected from Cha or Phe;

10

A_2 is selected from Arg or Lys;

A_3 is selected from Phe;

and pharmaceutically acceptable salts thereof.

15

6. The compound of Claim 5, selected from:

(5-Bromopyridin-3-yl) carbonyl-cyclohexylalanyl-arginyl-phenylalanineamide;

20

2-Chromonyl carbonyl-cyclohexylalanyl-arginyl-phenylalanineamide;

(5-Aminotriazol-3-yl) carbonyl-cyclohexylalanyl-arginyl-phenylalanineamide;

25

[5-(-Methyl) cinnamamidotriazol-3-yl] carbonyl-cyclohexylalanyl-arginyl-phenylalanineamide;

30

{5-[3-(1-Naphthyl) acrylamido] triazol-3-yl} carbonyl-cyclohexylalanyl-arginyl-phenylalanineamide;

[Quinoxalin-2-yl] carbonyl-cyclohexylalanyl-arginyl-phenylalanineamide;



- 49 -

[5-(*o*-Chlorocinnamamido)triazol-3-yl]carbonyl-cyclohexylalanyl-arginyl-phenylalanineamide;

5 (6-Aminopyridin-3-yl)carbonyl-cyclohexylalanyl-arginyl-phenylalanineamide;

(5-Aminotriazol-3-yl)carbonyl-phenylalanyl-arginyl-phenylalanyl-arginineamide;

10 (5-Aminotriazol-3-yl)carbonyl-cyclohexylalanyl-lysiny-phenylalanineamide;

15 {5-[3-(2-Thienyl)acrylamido]triazol-3-yl}carbonyl-cyclohexylalanyl-arginyl-phenylalanineamide;

[5-cinnamamidotriazol-3-yl]carbonyl-cyclohexylalanyl-arginyl-phenylalanineamide;

20 (6-Cinnamamidopyridin-3-yl)carbonyl-cyclohexylalanyl-arginyl-phenylalanineamide; or

(5-Chloro-3-methyl-benzothiophen-2-yl)carbonyl-cyclohexylalanyl-arginyl-phenylalanineamide;

25 and pharmaceutically acceptable salts thereof.

7. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a compound of Claim 1.

30

8. A pharmaceutical composition made by mixing a compound of Claim 1 and a pharmaceutically acceptable carrier.

35



- 50 -

9. A process for making a pharmaceutical composition comprising mixing a compound of Claim 1 and a pharmaceutically acceptable carrier.

5 10. A method of treating a condition mediated by modulation of the thrombin receptor in a subject in need thereof comprising administering to the subject a therapeutically effective amount of the compound of Claim 1.

10 11. The method of Claim 10, wherein the condition is selected from wound healing, tissue repair, myocardial infarction, stroke, restenosis, angina, atherosclerosis, ischemic attacks, inflammation, cancer, 15 osteoporosis, or neurodegenerative disorders.

12. The method of Claim 11, wherein the therapeutically effective amount of the compound is about 0.1 to about 300 mg/kg/day.

20 13. The method of Claim 12, wherein the therapeutically effective amount of the compound is about 1 to about 50 mg/kg/day.

25 14. A method of treating a condition modulated by the thrombin receptor in a subject in need thereof comprising administering to the subject a therapeutically effective amount of the composition of Claim 7.



- 51 -

15. The method of Claim 14, wherein the condition
is selected from wound healing, tissue repair,
myocardial infarction, stroke, restenosis, angina,
5 atherosclerosis, ischemic attacks, inflammation, cancer,
osteoporosis, or neurodegenerative disorders.

16. The method of Claim 14, wherein the
therapeutically effective amount of the compound is
10 about 0.1 to about 300 mg/kg/day.

17. The method of Claim 16, wherein the
therapeutically effective amount of the compound is
about 1 to about 50 mg/kg/day.

15