

METHODS AND COMPOSITIONS FOR THE TREATMENT OF CANCER

FIELD OF THE INVENTION

5 The invention relates to the field of cancer and, more particularly, to methods and compositions for the treatment of cancer using the Omomyc polypeptide and to compositions comprising Omomyc and one or more anticancer drugs and their use for the treatment of cancer.

10 BACKGROUND OF THE INVENTION

 The ideal cancer drug should target a non-redundant function continuously necessary for tumor maintenance, but dispensable for maintenance and function of any normal tissues. Hence, the most common logic is to target gene products that are specifically mutated in cancer, on the basis that these mutant molecules would be the likely "drivers" of the cancer and, perhaps, less critical for normal tissues. For these reasons, much attention has focused on cataloguing recurring lesions in specific cancer types. Unfortunately, there are several problems to this approach. First, most solid human cancers pass through episodes of genomic instability and exhibit a mutational noise that can obscure the "driver" mutations and their attendant effector pathways. Second, cancers are the end result of a process that involves transitions through multiple evolutionary bottlenecks. Each bottleneck may require a specific type of mutation whose function is thereafter dispensable for tumor maintenance and, consequently, not a good therapeutic target after that point in the tumor's evolution.

25 Myc is a basic helix-loop-helix leucine zipper (b-HLH-LZ) protein involved in growth control and cancer, which operates in a network with the structurally related proteins Max, Mad and Mnt. Myc/Max dimers activate gene transcription and induce cell proliferation or apoptosis. Mad/Max and Mnt/Max complexes act as repressors and cause cell growth arrest and differentiation. All dimers recognize the same DNA consensus site, the CACGTG E-box.

 Myc is tightly regulated in normal cells, where its levels are higher in proliferating and lower in non-proliferating. Aberrantly high and/or deregulated Myc

activity is causally implicated in most cancers and often associated with aggressive, poorly differentiated and angiogenic tumors. The deregulation of Myc expression is due to overexpression through gene amplifications, loss of transcriptional control, impaired degradation or increased stabilization. This results in aberrant proliferation, increased survival, changes in metabolism, angiogenesis and inflammation, all of which represent major hallmarks of cancer. Multiple studies substantiated the crucial role of Myc in governing intracellular and extracellular aspects of tumorigenesis suggesting that targeting its function would be therapeutically valuable.

It is known that down-regulation of myc by a BET bromodomain inhibitor results in the regression of multiple tumor types (Delmore, J. E., et al., 2011, *Cell*, 146: 904–917). While this approach displays good potential, it presents some limitations such as toxicity and numerous off targets effects.

Many small molecules disrupting the Myc/Max interaction have displayed low specificity *in cellulo* (Prochownik, E. V. and Vogt, P. K., 2010, *Genes Cancer* 1, 650-659).

A Myc inhibitor, however, has yet to become clinically available and its design presents various caveats: first, Myc is a nuclear transcription factor, which is consequently more difficult to reach than membrane or cytoplasmic molecules; second, Myc does not have an enzymatic "active site" that could be targeted; third, the Myc family comprises 3 different proteins, c-, N and L- Myc, which in certain conditions are functionally redundant, so all of them require simultaneous inhibition. Furthermore, there have been concerns that Myc inhibition would induce serious side effects by inhibiting proliferation of normal tissues. For all these reasons, making a Myc inhibitor drug is challenging.

Omomyc is a dominant-negative MYC mutant comprising the b-HLH-LZ domain of Myc and harboring four amino acid substitutions in the leucine zipper of Myc (Soucek, L. et al., 1998, *Oncogene* 17, 2463-2472; Soucek, L. et al. (2002), *Cancer Res* 62: 3507-3510). The amino acid substitutions E61T, E68I, R74Q, and R75N confer altered dimerization specificity to the protein, which retains the ability to bind its natural partner Max and to form homodimers and heterodimers with wild type c-, N- and L-Myc.

Because of these properties, Omomyc is able to prevent Myc-dependent gene transactivation functions both *in vitro* and *in vivo* by negating the ability of Myc to bind its DNA recognition binding site, the E box (Savino, M. et al. ,2011, *PLoS One* 6, e22284; Soucek, L. et al. (2004), *Cell Death Differ* 11, 1038 1045). At the same time,

5 Omomyc strongly potentiates Myc-induced apoptosis in a manner dependent on Myc expression level and thereby strengthens Myc transrepression activity. Omomyc thus prevents Myc binding to promoter E-boxes and transactivation of target genes while retaining Miz-1-dependent binding to promoters and transrepression. In the presence of Omomyc, the Myc interactome is channelled to repression and its activity switches

10 from a pro-oncogenic to a tumor-suppressive one.

TRE-Omomyc;CMVrtTA mice, in which Omomyc expression is controlled by a tetracycline-responsive promoter element and the widely expressed rtTA transactivator is driven by a CMV promoter, exhibit high Omomyc expression in most tissues following administration of doxycycline (Soucek et al., 2008, *Nature*, 455: 679-683).

15 These mice were crossed with the well-established LSL-Kras^{G12D} murine model of lung tumorigenesis. Just 3 days of Omomyc expression were sufficient to cause dramatic shrinkage of the tumors and one week renders the animals essentially tumor free. Importantly, although other dividing tissues, such as skin, testis and intestine, exhibited significantly decreased proliferation rates during the treatment, and displayed a certain

20 degree of atrophy, the mice exhibited no obvious signs of distress or disease. Moreover, the side effects of Myc inhibition resulting from Omomyc expression are completely reversible and disappear upon discontinuation of the treatment.

To date, despite the fact that the expression of Omomyc has proven to be an efficacious Myc inhibiting strategy *in vivo*, it has been applied solely using a gene

25 therapy approach. Indeed, Omomyc is a peptide considered too bulky and unfit for delivery to the desired cellular compartment (Montagne M. et al., *PLoS One*. 2012;7:e32172. doi: 10.1371/journal.pone.0032172), Savino M. et al., *PLoS One*. 2011; 6:e22284. doi: 10.1371/journal.pone.0022284) and *Genes Dev.*, 2011, 25: 895-7. doi: 10.1101/gad.2053311.)

30 Moreover, Omomyc is predicted to display poor ability to cross physiological barriers because of its intrinsic physico-chemical properties (e.g. hydrophobicity, as predicted using Kyte & Doolittle hydropathy plot, Kyte J., Doolittle R.F. (1982) *J. Mol.*

Biol. 157:105-132). In addition, despite the presence of several arginine residues within the basic region of Omomyc, the most recent algorithms predicting spontaneous cell-penetration capacity of peptides do not predict Omomyc to possess such property (Gautam et al. Journal of Translational Medicine 2013, 11:74).

5 Therefore, providing therapeutic approaches for the treatment of cancer based on b-HLH-LZ domains capable of transducing across the cellular membrane of eukaryotic cells and inhibiting Myc-dependent gene transactivation would be advantageous.

BRIEF SUMMARY OF THE INVENTION

10

In a first aspect, the invention refers to a polypeptide of SEQ ID NO: 1 or a functionally equivalent variant thereof for use in medicine as well as for use in the prevention and/or treatment of cancer.

In another aspect the invention refers to a fusion protein comprising:

- 15 (i) the polypeptide of SEQ ID NO: 1 or a functionally equivalent variant thereof and
- (ii) a cell-penetrating peptide sequence and/or a nuclear localization signal.

20 In another aspect, the invention refers to a pharmaceutical composition comprising the fusion protein according to the invention as well as to the use of the fusion protein in medicine and, in particular, for the treatment of cancer.

In another aspect, the invention relates to a composition comprising together or separately:

- 25 (i) a polypeptide of SEQ ID NO: 1, a functionally equivalent variant thereof or a fusion protein according to the invention and
- (ii) an antitumoral agent.

In another aspect, the invention refers to a pharmaceutical composition comprising the composition of the invention as well as the use of the composition of the invention in medicine and, in particular, for the treatment of cancer.

30

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1. (A) Fluorescence imaging of A549 cells incubated for 2h with Omomyc-FITC at 37°C shows that Omomyc-FITC is localized in the nucleus and cytoplasm (B) Hoescht staining of the nuclei. (C) Phase contrast.

Figure 2. Cells incubated with 10 μ M of either Omomyc or Max* were counted and stained with AnnexinV and PI. (A) Total cell number of PBS, Omomyc- or Max-treated A549. (B) Percentage of dead cells stained with PI. (C) Percentage of live (non-stained) cells. (D) Percentage of AnnexinV positive cells.

Figure 3. (A) Circular dichroism (CD) spectra recorded at 20°C for c-Myc*, Max* and Omomyc purified proteins (32 μ M) showing that Omomyc has a folded structure more similar to that of Max* than Myc*. (mdeg, millidegrees) (B) Thermal denaturation studied by circular dichroism at 1°C/min for c-Myc*, Max* and Omomyc purified proteins showing that Omomyc has a folded structure more thermally stable than that of Max*. ($^{\circ}$ m, millidegrees at the specified wavelength of 222 nm).

Figure 4. Quantification of fluorescence from confocal microscopy images of A549 cells fixed with 4% PFA after 2 hours of incubation at 37°C with Omomyc or Max* at different peptide concentrations (5, 10 and 25 μ M) (30-80 cells counted per image). AU, arbitrary units.

Figure 5. Quantification of fluorescence from confocal microscopy images of A549 live cells after 20 minutes of incubation at 37°C with Omomyc or Max* (20 μ M). AU, arbitrary units.

Figure 6. Crystal violet staining of A549 and H1650 lung adenocarcinoma cells treated with 25 μ M Omomyc or Max* peptide for the indicated times.

Figure 7. Quantification of the inhibition of proliferation by crystal violet staining of A549 and H1650 lung adenocarcinoma cells treated with 25 μ M Omomyc or Max* peptides for the indicated times.

5 **Figure 8.** Dose response of A549 cells to Omomyc and Max* by crystal violet staining quantification.

Figure 9. Quantification of proliferation of U87 glioma cells treated with Omomyc or Max* peptides at 25 μ M.

10

Figure 10. (A) Lung of untreated (left) and treated (right) animals 10 minutes after intranasal administration of the fluorescent peptide Omomyc (single dose of 37.5 mg/kg). (B) Brain of untreated (left) and treated (right) animals 10 minutes after intranasal administration of the fluorescent peptide Omomyc (single dose of 37.5

15 mg/kg).

Figure 11. Treatment of lung adenocarcinoma with PBS or with Omomyc by intranasal administration. (A) Proliferative rate of tumors. (B) Cellular density.

20 **Figure 12.** Measurement of the % tumor area after treatment of lung adenocarcinoma with PBS or with Omomyc by intranasal administration.

DETAILED DESCRIPTION OF THE INVENTION

25 The authors of the present invention have found that, surprisingly, Omomyc is capable of efficiently transducing across the cellular membrane and translocate to the nucleus, wherein it exerts its tumor-suppressive effect. Therefore, Omomyc is a bona fide Protein Transduction Domain (PTD). Hence, for the first time Omomyc can be used itself as an anti-Myc drug without the need of either using vehicles for delivery of

30 the polypeptide to the cytoplasm of the cell or using gene therapy approaches for delivery the nucleic acid encoding Omomyc to the cell. This allows the use of the Omomyc polypeptide for the treatment of diseases associated with deregulated cell

proliferation such as cancer. The authors of the present invention have surprisingly found that Omomyc has several advantages in comparison to the bHLHZ domain of Max (Max*):

- Omomyc shows greater cell penetrating ability in comparison to Max* in different cell types and at different concentrations (Example 6)
- Omomyc is more thermally stable than Max* and this is a clear advantage for drug design (Example 5)
- Omomyc is more efficient than Max* in both preventing the growth of cells (Example 8) and increasing death of cancer cells (Example 4).

In addition, Omomyc is capable of crossing the blood-brain barrier (Example 9 and Figure 10B) and exerting its therapeutic effect *in vivo* (Example 10).

Therapeutic methods using Omomyc

The present invention provides methods for the treatment of cancer based on the use of a polypeptide having the sequence of SEQ ID NO: 1, which corresponds to Omomyc, or of a functionally equivalent variant thereof.

In a first aspect, the invention refers to a polypeptide of SEQ ID NO: 1 or a functionally equivalent variant thereof for use in medicine.

In another aspect, the invention refers to a polypeptide of SEQ ID NO: 1 or a functionally equivalent variant thereof for use in the prevention and/or treatment of cancer.

In another aspect, the invention also refers to a method for the prevention and/or treatment of cancer that comprises administering to a subject in need thereof a therapeutically effective amount of the polypeptide of SEQ ID NO: 1 or a functionally equivalent variant thereof.

In another aspect, the invention also refers to the polypeptide of SEQ ID NO: 1 or a functionally equivalent variant thereof for the preparation of a medicament for the prevention and/or treatment of cancer.

The polypeptide of sequence SEQ ID NO: 1 corresponds to the Omomyc protein sequence. The term “Omomyc” as used herein, refers to a polypeptide which consists of a mutated version of the bHLHZip domain of the Myc carrying the E61T, E68I,

R74Q and R75N mutations (wherein the numbering of the mutated positions is given with respect to the sequence of Myc region corresponding to amino acids 365-454 of the polypeptide as defined under accession number NP_002458 in the NCBI database, release of June 27, 2012). The sequence of c-Myc provided in the NCBI database under the accession number NP_002458 is shown below, wherein the region from which Omomyc derives is shown underlined:

```

1  mdffrvvenq qppatmplnv sftnrnydld ydsvqpyfyc deenfyqqq qqselsppap
61  sediwwkfel lptpplspsr rsglcspsyv avtpfslrgd ndggggsfst adqlemvtel
10 121 lggdmvnqsf icdpddetfi kniiiqdcmw sgfsaaaklv seklasyqaa rkdsgspnpa
181 rghsvctstss lylqdlstaa secidpsvfv pypIndsssp kscasqdssa fspssdsls
241 stesspggsp eplvlheetp pttssdseee qedeeidvv svekrqapgk rsesgspasg
301 ghskpphspl vlkrchvsth qhnyaappst rkdypaakrv kldsvrvlrq isnnrkctsp
361 rssdteenvk rrthnvlerq rrnelkrsff alrdqipele nnekapkvvi lkkatayils
15 421 vgaeeqklis eedllrkrre qlkhkleqlr nsca (SEQ ID NO:2)

```

The polynucleotide encoding Omomyc (SEQ ID NO: 3) and the corresponding polypeptide sequence (SEQ ID NO: 1) is shown below, wherein the underlined and bold triplets correspond to those positions which are mutated with respect to Myc:

```

20
1ACC GAG GAG AAT GTC AAG AGG CGA ACA CAC AAC GTC TTG GAG CGC CAG
1Thr Glu Glu Asn Val Lys Arg Arg Thr His Asn Val Leu Glu Arg Gln

49AGG AGG AAC GAG CTA AAA CGG AGC TTT TTT GCC CTG CGT GAC CAG ATC
25 17Arg Arg Asn Glu Leu Lys Arg Ser Phe Phe Ala Leu Arg Asp Gln Ile

97CCG GAG TTG GAA AAC AAT GAA AAG GCC CCC AAG GTA GTT ATC CTT AAA
33Pro Glu Leu Glu Asn Asn Glu Lys Ala Pro Lys Val Val Ile Leu Lys

30 145AAA GCC ACA GCA TAC ATC CTG TCC GTC CAA GCA GAG ACG CAA AAG CTC
49Lys Ala Thr Ala Tyr Ile Leu Ser Val Gln Ala Glu Thr Gln Lys Leu

193ATT TCT GAA ATC GAC TTG TTG CGG AAA CAA AAC GAA CAG TTG AAA CAC

```

⁶⁵ *Ile Ser Glu Ile Asp Leu Leu Arg Lys Gln Asn Glu Gln Leu Lys His*

²⁴¹AAA CTT GAA CAG CTA CGG AAC TCT TGT GCG TAA

⁸¹*Lys Leu Glu Gln Leu Arg Asn Ser Cys Ala End*

5

Omomyc also contains the M2 domain of c-Myc, having the sequence RQRRNELKRSF (SEQ ID NO: 49) (see Dang and Lee, Mol.Cell. Biol., 1988, 8:4048-4054) (double underlined above), and which corresponds to a nuclear localization signal.

10 Omomyc is characterized in that it shows increased dimerization capacity with all three oncogenic Myc proteins (c-Myc, N-Myc and L-Myc). Omomyc can derive from the bHLHZip domain of any Myc protein known in the art, provided that the mutations which result in the tumor suppressor effect are preserved. Thus, the Omomyc that can be used in the present invention may derive from any mammal species,
15 including but not being limited to domestic and farm animals (cows, horses, pigs, sheep, goats, dog, cats or rodents), primates and humans. Preferably, the Omomyc protein is derived from human Myc protein (accession number NP_002458, release of June 27, 2012).

The term “Myc”, as used, herein, refers to a family of transcription factors which
20 includes c-Myc, N-Myc and L-Myc. Myc protein activates expression of many genes through binding on consensus sequence CACGTG (Enhancer Box sequences or E-boxes and recruiting histone acetyl-transferases or HATs). However, Myc can also act as a transcriptional repressor. By binding the Miz-1 transcription factor and displacing p300 co-activator, it inhibits expression of Miz-1 target genes. Myc also has a direct
25 role in the control of DNA replication.

The Myc b-HLH-LZ or Myc basic region helix-loop-helix leucine zipper domain refers to a region which determines Myc dimerization with Max protein and binding to Myc-target genes. This region corresponds to amino acids 365-454 of human Myc and is characterized by two alpha helices connected by a loop (Nair, S. K., & Burley, S. K.,
30 2003, Cell, 112: 193–205).

The term “functionally equivalent variant”, when referring to Omomyc, refers to any polypeptide which results from the deletion, insertion or addition of one or more amino acids with respect to the polypeptide of SEQ ID NO:1 or which results from the

chemical modification of the polypeptide of SEQ ID NO: 1 and which substantially preserves the tumor suppressor activity of the Omomyc polypeptide. The skilled person will understand that the preservation of the tumor suppressor activity of Omomyc requires that the variant can dimerize with Myc and inhibit its activity once found in the
 5 nucleus, that it is capable of translocating across the cell membrane and that it is capable of translocating across the nuclear envelope.

Suitable functionally equivalent variants of Omomyc include polypeptides consisting essentially of the polypeptide of SEQ ID NO:1. In this context, "consisting essentially of" means that the specified molecule would not contain any additional
 10 sequences that would alter the activity of Omomyc.

Suitable functional variants of the targeting peptide are those showing a degree of identity with respect to the peptide of SEQ ID NO:1 of about greater than 25% amino acid sequence identity, such as 25% 40%, 60%, 70%, 80%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99%. The degree of identity between two polypeptides is
 15 determined using computer algorithms and methods that are widely known for the persons skilled in the art. The identity between two amino acid sequences is preferably determined by using the BLASTP algorithm as described previously [BLAST Manual, Altschul, S., et al., NCBI NLM NIH Bethesda, Md. 20894, Altschul, S., et al., J. Mol. Biol. 1990;215: 403-410]. In a preferred embodiment, the sequence identity is
 20 determined throughout the whole length of the polypeptide of SEQ ID NO:1 or throughout the whole length of the variant or of both.

The functionally equivalent variants of the Omomyc polypeptide may also include post-translational modifications, such as glycosylation, acetylation, isoprenylation, myristoylation, proteolytic processing, etc.

25 Alternatively, suitable functional variants of the targeting peptide are those wherein one or more positions within the Omomyc polypeptide contain an amino acid which is a conservative substitution of the amino acid present in the Omomyc protein mentioned above. "Conservative amino acid substitutions" result from replacing one amino acid with another having similar structural and/or chemical properties. For
 30 example, the following six groups each contain amino acids that are conservative substitutions for one another: 1) Alanine (A), Serine (S), Threonine (T); 2) Aspartic acid (D), Glutamic acid (E); 3) Asparagine (N), Glutamine (Q); 4) Arginine (R), Lysine

(K); 5) Isoleucine (I), Leucine (L), Methionine (M), Valine (V); and 6) Phenylalanine (F), Tyrosine (Y), Tryptophan (W). Selection of such conservative amino acid substitutions is within the skill of one of ordinary skill in the art and is described, for example by Dordo et al. et al., (J. Mol. Biol, 1999, 217:721-739) and Taylor et al., (J. Theor. Biol., 1986, 119:205-218).

It will be understood that the functionally equivalent variants of Omomyc contain mutations at positions corresponding to the mutations E61T, E68I, R74Q and R75N found in Omomyc derived from human c-Myc. The position wherein said mutations have to occur in the functionally equivalent variant can be determined by a multiple sequence alignment of different Myc sequences and identified by the alignment of those positions corresponding to positions 61, 68, 74 and 75 within the sequence of Omomyc derived from human c-Myc.

A multiple sequence alignment is an extension of pairwise alignment to incorporate more than two sequences at a time. Multiple alignment methods align all of the sequences in a given query set. A preferred multiple sequence alignment program (and its algorithm) is ClustalW, Clusal2W or ClustalW XXL (see Thompson et al. (1994) Nucleic Acids Res 22:4673-4680). Once the sequences of c-Myc from different organisms and of the variant are compared (aligned) as described herein, the skilled artisan can readily identify the positions within each of the sequence corresponding to positions and introduce within the Omomyc variant mutations corresponding to the E61T, E68I, R74Q and R75N mutations found in Omomyc derived from human c-Myc.

Suitable assays for determining whether a polypeptide can be considered as a functionally equivalent variant of Omomyc include, without limitation:

- Assays which measure the capacity of the polypeptide to form dimeric complexes with Max and Myc, such as the assays based on the expression of a reporter gene as described in Soucek et al. (Oncogene, 1998, 17: 2463 - 2472) as well as PLA (protein Ligation assay) or Co-immunoprecipitation.
- Assays which measure the capacity of the polypeptide to bind to the Myc/Max recognition site within DNA (the CACGTG site), such as the electrophoretic mobility shift assay (EMSA) described in Soucek et al. (*supra.*)

- Assays which measure the capacity to repress Myc-induced transactivation, such as the assay based on the expression of a reporter gene under the control of the DNA binding sites specific for Myc/Max as described by Soucek et al. (supra.).
- Assays based on the capacity of the polypeptide to inhibit growth of cells expressing the myc oncogene, as described by Soucek et al. (supra.).
- Assays which measure the ability of the polypeptide to enhance myc-induced apoptosis, such as the assays described by Soucek et al. (Oncogene, 1998: 17, 2463 - 2472). Moreover, any assay commonly known in the art for assessing apoptosis in a cell can be used, such as the Hoechst staining, Propidium Iodide (PI) or Annexin V staining), trypan blue, DNA laddering/fragmentation and TUNEL.

In a preferred embodiment, a polypeptide is considered a functionally equivalent variant of Omomyc if it shows an activity in one or more of the above assays which is at least 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or 100% of the native Omomyc.

Additionally, functionally equivalent variants of Omomyc are also capable of transducing cells after the variant is contacted with said cell. It will be understood that functionally equivalent variants of Omomyc contain the protein transducing domain found in native Omomyc or another functional protein transducing domain.

The term “cell penetrating peptide sequence” is used in the present specification interchangeably with “CPP”, “protein transducing domain” or “PTD”. It refers to a peptide chain of variable length that directs the transport of a protein inside a cell. The delivering process into cell commonly occurs by endocytosis but the peptide can also be internalized into cell by means of direct membrane translocation. CPPs typically have an amino acid composition that either contains a high relative abundance of positively charged amino acids such as lysine or arginine or has sequences that contain an alternating pattern of polar/charged amino acid and non-polar, hydrophobic amino acids. Examples of CPPs which can be used in the present invention include, without limitation, the CPP found in *Drosophila antennapedia* protein (RQIKIWFQNRRMKWKK, SEQ ID NO:4), the CPP found in the herpesvirus simplex 1 (HSV-1) VP22 DNA-binding protein (DAATATRGRSAASRPTERPRAPARSASRPRRPVE, SEQ ID NO:5), the CPP of

Bac-7 (RRIRPRPPRLPRPRPLPFPRPG; SEQ ID NO: 6), the CPPs of the HIV-1 TAT protein consisting of amino acids 49-57 (RKKRRQRRR; SEQ ID NO: 7), amino acids 48-60 (GRKKRRQRRRTPQ; SEQ ID NO: 8), amino acids 47-57 (YGRKKRRQRRR; SEQ ID NO: 9); the CPP of S413-PV peptide
 5 (ALWKTLLKKVLKAPKKKRKV; SEQ ID NO: 10), the CPP of penetratin (RQIKWFAQNRMMKWKK; SEQ ID NO: 11), the CPP of SynB1 (RGGRLSYSRRRFSTSTGR; SEQ ID NO: 12), the CPP of SynB3 (RRLSYSRRRF; SEQ ID NO: 13), the CPP of PTD-4 (PIRRRKCLRRLK; SEQ ID NO: 14), the CPP of PTD-5 (RRQRRTSKLMKR; SEQ ID NO: 15), the CPP of the FHV Coat-(35-49)
 10 (RRRRNRTRNRNRVR; SEQ ID NO: 16), the CPP of BMV Gag-(7-25) (KMTRAQRRAAARRNRWTAR; SEQ ID NO: 17), the CPP of HTLV-II Rex-(4-16) (TRRQRTRRARRNR; SEQ ID NO: 18), the CPP of D-Tat (GRKKRRQRRRPPQ; SEQ ID NO: 19), the CPP R9-Tat (GRRRRRRRRRPPQ; SEQ ID NO: 20), the CPP of MAP (KLALKLALKLALKLA; SEQ ID NO: 21), the CPP of SBP
 15 (MGLGLHLLVLAAALQGAWSQPKKKRKV; SEQ ID NO: 22), the CPP of FBP (GALFLGWLGAAGSTMGAWSQPKKKRKV; SEQ ID NO: 23), the CPP of MPG (ac-GALFLGLGAAGSTMGAWSQPKKKRKV-cya; SEQ ID NO: 24), the CPP of MPG(ENLS) (ac-GALFLGLGAAGSTMGAWSQPKSKRKV-cya; SEQ ID NO: 25), the CPP of Pep-1 (ac-KETWWETWWTEWSQPKKKRKV-cya; SEQ ID NO: 26), the
 20 CPP of Pep-2 (ac-KETWFETWFTEWSQPKKKRKV-cya; SEQ ID NO: 27), a polyarginine sequence having the structure R_N (wherein N is between 4 and 17), the GRKKRRQRRR sequence (SEQ ID NO: 28), the RRRRRRLR sequence (SEQ ID NO: 29), the RRQRRTS KLMKR sequence (SEQ ID NO: 30); Transportan GWTLNSAGYLLGKINLKALAALAKKIL (SEQ ID NO: 31);
 25 KALAWKALAKALAKALAKHLAKALAKALKCEA (SEQ ID NO: 32); RQIKWFAQNRMMKWKK (SEQ ID NO: 33), the YGRKKRRQRRR sequence (SEQ ID NO: 34); the RKKRRQRR sequence (SEQ ID NO: 35); the YARAAARQARA sequence (SEQ ID NO: 36); the THRLPRRRRRR sequence (SEQ ID NO: 37); the GRRRRRRRRR sequence (SEQ ID NO: 38).

30 Suitable assays for determining whether a polypeptide preserves the cell membrane translocation capacity of Omomyc include, without limitation, assays which measure the capacity of the polypeptide to transduce cells in culture, such as the assay

shown in example 3 of the present invention. This assay is based on contacting the polypeptide with culture cells and detecting the presence of the polypeptide in an intracellular location. In a preferred embodiment, the detection of the polypeptide of the invention is performed by fluorescence microscopy.

5 In a preferred embodiment, a polypeptide is considered as a functionally equivalent variant of Omomyc if it is capable of transducing a target cell at least 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or 100% as efficiently as native Omomyc.

Additionally, functionally equivalent variants of Omomyc are also capable of reaching the nuclei of the transduced cells after the variant is contacted with said cell. It
10 will be understood that functionally equivalent variants of Omomyc contain the NLS found in native Omomyc or another functional NLS.

The term "nuclear localization signal", as used herein, refers to an amino acid sequence of about 4-20 amino acid residues in length, which serves to direct a protein to the nucleus. Typically, the nuclear localization sequence is rich in basic amino acids and
15 exemplary sequences are well known in the art (Gorlich D. (1998) EMBO 5.17:2721-7). In some embodiments, the NLS is selected from the group consisting of the SV40 large T Antigen NLS (PKKKRKV, SEQ ID NO: 39); the Nucleoplasmin NLS (KRPAATKKAGQ AKKKK, SEQ ID NO: 40); the CBP80 NLS (RRRHSDENDGGQPHKRRK, SEQ ID NO: 41); the HIV-I Rev protein NLS
20 (RQARRNRRRWE, SEQ ID NO: 42); the HTLV-I Rex (MPKTRRRPRRSQRKRPT, SEQ ID NO: 43); the hnRNP A NLS (NQSSNFGPMKGGNFGGRSSGPYGGGGQYFKPRNQGGY, SEQ ID NO: 44); the rpL23a NLS (VSHKKKKIRTSPFTTPKTLRLRRQPKYPRKSAPRRNKLDHY, SEQ ID NO: 45). In one embodiment of the invention, the nuclear localization signal
25 comprises the motif K (K/ R) X (K/ R) (SEQ ID NO: 46).

Suitable assays for determining whether a polypeptide is a functionally equivalent variant of Omomyc in terms of its ability to translocate across the cellular membrane include double labelling of a cell with a reagent specific for the polypeptide and with a dye which specifically labels the nucleus of the cell (such as DAPI or
30 Hoechst dye). Such assays are shown in Example 6 of the present invention. In a preferred embodiment, the detection of the polypeptide of the invention is performed by confocal microscopy or by fluorescence microscopy.

In a preferred embodiment, a polypeptide is considered as a functionally equivalent variant of Omomyc if it is capable of translocating to the nucleus of the target tumor cells at least 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or 100% as efficiently as native Omomyc.

- 5 Suitable functionally equivalent variants include the polypeptides Omomyc*TAT and Omomyc*LZArg as defined below:

Name	SEQ ID NO:	Sequence
Omomyc*TAT	47	MTEENVKRRTHNVLERQRRNELKRSFFALRDQIPELENNEKAPKVVI LKKATAYILSVQAETQKLI SEIDLLRKQNEQLKHKLEQLRNSCAGRK KRRQRRR
Omomyc*LZArg	48	MTEENVKRRTHNVLERQRRNELKRSFFALRDQIPELENNEKAPKVVI LKKATAYILSVQAETQKLI SEIDLLRKQNEQLKHKLEQLRNSCARRR RRRLR

- According to the invention, the Omomyc polypeptide or the functionally
10 equivalent variant thereof are used in a method for the prevention or treatment of cancer in a subject. It will be understood that the preventive or therapeutic method according to the invention involves the direct use of the Omomyc polypeptide or of the functionally equivalent variant thereof. Thus, the preventive or therapeutic methods according to the invention do not involve the administration of the nucleic acid encoding Omomyc or the
15 functionally equivalent variant thereof.

“Prevention” is understood as the administration of a polypeptide of SEQ ID NO: 1 or a functionally equivalent variant thereof according to the first aspect of the invention, or of a medicament containing it in an initial or early stage of the disease, or to also prevent its onset.

- 20 The term “treatment” is used to designate the administration of a polypeptide of SEQ ID NO: 1 or a functionally equivalent variant thereof according to the first aspect of the invention, or of a medicament containing it to control the progression of the disease before or after the clinical signs have appeared. Control of the progression of the disease is understood as the beneficial or desired clinical results which include but are
25 not limited to reduction of the symptoms, reduction of the duration of the disease, stabilization of pathological conditions (specifically avoiding additional impairment), delaying the progression of the disease, improving the pathological condition and remission (both partial and complete). The control of the progression of the disease also

involves a prolongation of survival in comparison to the expected survival if the treatment was not applied.

The term “cancer” is referred to a disease characterized by uncontrolled cell division (or by an increase of survival or apoptosis resistance), by the ability of said
 5 cells to invade other neighbouring tissues (invasion) or by the spread to other areas of the body where the cells are not normally located (metastasis) through the lymphatic and blood vessels. Depending on whether or not tumours can spread by invasion and metastasis, they are classified as being either benign or malignant: benign tumours are tumours that cannot spread by invasion or metastasis, i.e., they only grow locally;
 10 whereas malignant tumours are tumours that are capable of spreading by invasion and metastasis. The methods according to the present invention are useful for the treatment of local and malignant tumours. As used herein, the term cancer includes, but is not limited to, the following types of cancer: breast cancer; biliary tract cancer; bladder cancer; brain cancer including glioblastomas and medulloblastomas; cervical cancer;
 15 choriocarcinoma; colon cancer; endometrial cancer; esophageal cancer; gastric cancer; hematological neoplasms including acute lymphocytic and myelogenous leukemia; T-cell acute lymphoblastic leukemia/lymphoma; hairy cell leukemia; chronic myelogenous leukemia, multiple myeloma; AIDS-associated leukemias and adult T-cell leukemia/lymphoma; intraepithelial neoplasms including Bowen's disease and Paget's
 20 disease; liver cancer; lung cancer; lymphomas including Hodgkin's disease and lymphocytic lymphomas; neuroblastomas; oral cancer including squamous cell carcinoma; ovarian cancer including those arising from epithelial cells, stromal cells, germ cells and mesenchymal cells; pancreatic cancer; prostate cancer; rectal cancer; sarcomas including leiomyosarcoma, rhabdomyosarcoma, liposarcoma, fibrosarcoma, and osteosarcoma;
 25 skin cancer including melanoma, Merkel cell carcinoma, Kaposi's sarcoma, basal cell carcinoma, and squamous cell cancer; testicular cancer including germinal tumors such as seminoma, non-seminoma (teratomas, choriocarcinomas), stromal tumors, and germ cell tumors; thyroid cancer including thyroid adenocarcinoma and medullar carcinoma; and renal cancer including adenocarcinoma and Wilms tumor.
 30 Other cancers will be known to one of ordinary skill in the art. In a preferred embodiment, the cancer treated is lung cancer, preferably lung adenocarcinoma, more preferably a KRas-driven lung adenocarcinoma.

The authors of the present invention have also observed that Omomyc or the functionally equivalent variant thereof is capable of decreasing cell proliferation irrespective of whether the cancer shows increased expression or activity of the Myc protein.

5 A "subject," as used herein, includes any animal that has a cancer or exhibits a symptom or cancer, or is at risk for having a cancer or exhibiting a symptom of cancer. Suitable subjects (patients) include laboratory animals (such as mouse, rat, rabbit, or guinea pig), farm animals, and domestic animals or pets (such as a cat or dog). Non-human primates and, preferably, human patients, are included.

10 The appropriate dosage of Omomyc or of the functionally equivalent variant thereof to be used in the methods according to the invention will depend on different factors such as the type of cancer to be treated, the severity and course of the disease, whether the composition is administered for preventive or therapeutic purposes, previous therapy, the patient's clinical history and response to the peptide or
15 polypeptide, and the discretion of the attending physician.

 The amount of polypeptide of SEQ ID NO:1, of the functionally equivalent variant thereof is suitably administered to the patient at one time or over a series of treatments. Depending on the type and severity of the disease, an appropriate dosage level will generally be about 0.01 to 500 mg per kg patient body weight per day which
20 can be administered in single or multiple doses. Preferably, the dosage level will be about 0.1 to about 250 mg/kg per day; more preferably about 0.5 to about 100 mg/kg per day. A suitable dosage level may be about 0.01 to 250 mg/kg per day, about 0.05 to 100 mg/kg per day, or about 0.1 to 50 mg/kg per day. Within this range the dosage may be 0.05 to 0.5, 0.5 to 5 or 5 to 50 mg/kg per day. For oral administration, the
25 compositions are preferably provided in the form of tablets containing 1.0 to 1000 milligrams of the active ingredient, particularly 1.0, 5.0, 10.0, 15.0, 20.0, 25.0, 50.0, 75.0, 100.0, 150.0, 200.0, 250.0, 300.0, 400.0, 500.0, 600.0, 750.0, 800.0, 900.0, and 1000.0 milligrams of the active ingredient for the symptomatic adjustment of the dosage to the patient to be treated. The compounds may be administered on a regimen of 1 to 4
30 times per day, preferably once or twice per day.

 The polypeptide of SEQ ID NO:1, or of the functionally equivalent variant thereof may be administered by any type of suitable route, such as by oral route, topical

route, by inhalation or parenteral route so that the pharmaceutically acceptable excipients necessary for the formulation of the desired dosage form will be included. The preferred route of administration of said pharmaceutical compositions is the endovenous route. In another embodiment, the route of administration is the intranasal
5 route.

In one embodiment, the Omomyc or the functionally equivalent variant thereof is prepared with carriers which will protect said polypeptide from a rapid elimination from the body, such as a controlled release formulation, including implants and microencapsulated administration systems. Biodegradable biocompatible polymers such as
10 ethylene vinylacetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters and polylactic acid can be used. The processes for preparing said formulations will be clear for persons skilled in the art. The materials can also be commercially obtained in Alza Corporation and Nova Pharmaceuticals, Inc.

Despite the fact that Omomyc and functionally equivalent variants thereof are
15 capable of translocating across biological membranes, it is possible to formulate Omomyc or any of its functionally equivalent variants in nanoparticles. The nanoparticles may contribute to preserve the integrity of the polypeptide in the biological fluids until it reaches the target organ. In addition, nanoparticles can also be modified so as to include moieties which allow the targeting of the nanoparticle to an
20 organ of interest. In this way, Omomyc or the functionally equivalent variant thereof will be delivered in the proximity of the target organ, facilitating access of Omomyc to the interior of the cells where its biological activity is required.

Thus, in another embodiment, Omomyc or any of its functionally equivalent variants are provided forming part of a nanoparticle.

25 As used herein, the term "nanoparticle" refers to any material having dimensions in the 1-1,000 nm range. In some embodiments, nanoparticles have dimensions in the 2-200 nm range, preferably in the 2-150 nm range, and even more preferably in the 2-100 nm range. Nanoparticles that can be used in the present invention include such nanoscale materials as a lipid-based nanoparticle, a superparamagnetic nanoparticle, a
30 nanoshell, a semiconductor nanocrystal, a quantum dot, a polymer-based nanoparticle, a silicon-based nanoparticle, a silica-based nanoparticle, a metal-based nanoparticle, a fullerene and a nanotube.

Targeted delivery can be achieved by the addition of ligands without compromising the ability of nanoparticles to deliver their polypeptide payloads. It is contemplated that this will enable delivery to specific cells, tissues and organs. The targeting specificity of the ligand-based delivery systems are based on the distribution
 5 of the ligand receptors on different cell types. The targeting ligand may either be non-covalently or covalently associated with a nanoparticle, and can be conjugated to the nanoparticles by a variety of methods as discussed herein.

Examples of proteins or peptides that can be used to target nanoparticles include transferrin, lactoferrin, TGF- β , nerve growth factor, albumin, HIV Tat peptide, RGD
 10 peptide, and insulin, as well as others.

It will be understood that the formulation of Omomyc or of the functionally equivalent variant thereof in a nanoparticle is not intended or is not solely intended for facilitating the access of the Omomyc to the interior of the cell but to protect Omomyc from degradation and/or for facilitating targeting of the nanoparticle to the organ of
 15 interest.

Omomyc conjugates and fusion proteins comprising Omomyc

The present invention also provides conjugates which includes a first region comprising the polypeptide of SEQ ID NO:1 or a functionally equivalent variant thereof
 20 and a second region comprising a chemical moiety that facilitates cellular uptake of the polypeptide. Examples of moieties for enhancing cellular uptake include but are not limited to: a hydrophobic group (e.g., a lipid or fatty acid), a protein transducing domain, and certain metal chelates.

The presence of additional chemical moieties in the Omomyc molecule results in
 25 conjugates showing increased capability for being translocated across biological membranes with respect to unmodified Omomyc, thereby resulting in increased tumor suppressor activity.

Thus, in another aspect, the invention relates to a conjugate comprising:

- (i) the polypeptide of SEQ ID NO: 1 or a functionally equivalent variant thereof,
 30 and
- (ii) a chemical moiety that facilitates cellular uptake of the polypeptide.

The term “conjugate”, as used herein, refers to two or more compounds which are covalently linked together so that the function of each compound is retained in the conjugate.

In preferred embodiments, the conjugates according to the invention comprise at least 1, at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, at least 10 or more chemical moieties that facilitate cellular uptake of the polypeptide or of the functionally equivalent variant thereof.

In one embodiment, the chemical moiety that facilitates cellular uptake of the polypeptide is a lipid or a fatty acid.

A fatty acid generally is a molecule comprising a carbon chain with an acidic moiety (e.g., carboxylic acid) at an end of the chain. The carbon chain may of a fatty acid may be of any length, however, it is preferred that the length of the carbon chain be of at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 or more carbon atoms, and any range derivable therein. In certain embodiments, the length of the carbon chain is from 4 to 18 carbon atoms in the chain portion of the fatty acid. In certain embodiments the fatty acid carbon chain may comprise an odd number of carbon atoms, however, an even number of carbon atoms in the chain may be preferred in certain embodiments. A fatty acid comprising only single bonds in its carbon chain is called saturated, while a fatty acid comprising at least one double bond in its chain is called unsaturated. The fatty acid may be branched, though in preferable embodiments of the present invention, it is unbranched. Specific fatty acids include, but are not limited to, linoleic acid, oleic acid, palmitic acid, linolenic acid, stearic acid, lauric acid, myristic acid, arachidic acid, palmitoleic acid, arachidonic acid.

In a preferred embodiment, the chemical moiety that facilitates cellular uptake of the polypeptide is a cell penetrating peptide sequence, in which case, the conjugate is a fusion protein comprising Omomyc or the functionally equivalent variant thereof and the cell penetrating peptide sequence.

The term “fusion protein” relates to proteins generated by gene technology which consist of two or more functional domains derived from different proteins. A fusion protein may be obtained by conventional means, e.g., by means of gene expression of the nucleotide sequence encoding for said fusion protein in a suitable cell. It will be understood that the cell penetrating peptide refers to a cell penetrating peptide

which is different from the cell penetrating peptide which forms part of the polypeptide of SEQ ID NO: 1 or of the a functionally equivalent variant thereof.

The terms “polypeptide of SEQ ID NO:1”, “functionally equivalent variant of the polypeptide of SEQ ID NO:1” and “cell penetrating peptide” have been described in
5 detail in the context of the medical uses of the invention and are equally applicable in the context of the fusion protein.

In a preferred embodiment, said cell-penetrating peptide is not the endogenous Omomyc cell penetrating peptide.

In one embodiment, the cell-penetrating peptide sequence is fused at the N-
10 terminus of the polypeptide of SEQ ID NO: 1 or of the functionally equivalent variant thereof. In another embodiment, the cell-penetrating peptide is fused at the C-terminus of the polypeptide of SEQ ID NO: 1 or of the functionally equivalent variant thereof.

In preferred embodiments, the fusion proteins according to the invention comprise, in addition to the own cell penetrating peptide found in the polypeptide of
15 SEQ ID NO:1 or of the functionally equivalent variant thereof, at least 1, at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, at least 10 or more additional cell penetrating peptides.

In another preferred embodiment, the conjugates or fusion proteins of the invention comprise the polypeptide of SEQ ID NO: 1 or a functionally equivalent
20 variant thereof and further comprise an N-terminal or C-terminal nuclear localization signal. The term “nuclear localization signal (NLS)” has been described in the context of the therapeutic uses of the invention and is equally applicable to the fusion proteins of the invention. It will be appreciated that the additional NLS refers to an NLS which is different to the endogenous NLS found in Omomyc or in the functionally equivalent
25 variant thereof. The additional NLS may be the same or different to the endogenous NLS found in Omomyc or in the functionally equivalent variant thereof.

In one embodiment, the NLS is one of the NLS which appears endogenously in the Myc sequence, such as the M1 peptide (PAAKRVKLD, SEQ ID NO: 50) or the M2 peptide (RQRRNELKRSF, SEQ ID NO: 49) (see Dang and Lee, *supra.*).

30 In preferred embodiments, the conjugates or fusion proteins according to the invention comprise, in addition to the endogenous NLS found in the polypeptide of

SEQ ID NO:1 or in the functionally equivalent variant thereof, at least 1, at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, at least 10 or NLS.

The skilled person will understand that it may be desirable that the fusion protein further comprises one or more flexible peptides that connect the polypeptide of SEQ ID NO:1 or the functionally equivalent variant thereof, the cell penetrating peptide sequence and/or the NLS. Thus, in a particular embodiment the polypeptide of the invention is directly connected to the cell penetrating peptide sequence. In another particular embodiment, the polypeptide of the invention is connected to the cell penetrating peptide sequence through a flexible peptide. In a particular embodiment the polypeptide of the invention is directly connected to the cell penetrating peptide sequence and to the NLS. In another particular embodiment, the polypeptide of the invention is connected to the cell penetrating peptide sequence through a first flexible peptide linker and to the NLS through a second flexible peptide linker.

As used herein, the term "flexible peptide", "spacer peptide" or "linker peptide" refers to a peptide that covalently binds two proteins or moieties but which is not part of either polypeptide, allowing movement of one with respect to the other, without causing a substantial detrimental effect on the function of either the protein or the moiety. Thus, the flexible linker does not affect the tumour suppressor activity of the Omomyc sequence, the cell penetrating activity of the cell penetrating peptide or the nuclear localization capacity of the NLS.

The flexible peptide comprises at least one amino acid, at least two amino acids, at least three amino acids, at least four amino acids, at least five amino acids, at least six amino acids, at least seven amino acids, at least eight amino acids, at least nine amino acids, the least 10 amino acids, at least 12 amino acids, at least 14 amino acids, at least 16 amino acids, at least 18 amino acids, at least 20 amino acids, at least 25 amino acids, at least 30 amino acids, at least 35 amino acids, at least 40 amino acids, the least 45 amino acids, at least 50 amino acids, at least 60 amino acids, at least 70 amino acids, at least 80 amino acids, at least 90 amino acids, or about 100 amino acids. In some embodiments the flexible peptide will permit the movement of one protein with respect to the other in order to increase solubility of the protein and/or to improve its activity. Suitable linker regions include a poly-glycine region, the GPRRRR sequence (SEQ ID NO: 51) of combinations of glycine, proline and alanine residues.

In some embodiments the fusion protein of the invention can comprise an additional chemical moiety including, among others, fluorescence groups, biotin, polyethylene glycol (PEG), amino acid analogs, unnatural amino acids, phosphate groups, glycosyl groups, radioisotope labels, and pharmaceutical molecules. In other
 5 embodiments, the heterologous polypeptide can comprise one or more chemically reactive groups including, among others, ketone, aldehyde, Cys residues and Lys residues.

In a particular embodiment, the conjugates or fusion proteins of the invention comprise a tag bound to the conjugate or to the C-terminal or N-terminal domain of said
 10 fusion protein or variant thereof. Said tag is generally a peptide or amino acid sequence which can be used in the isolation or purification of said fusion protein. Thus, said tag is capable of binding to one or more ligands, for example, one or more ligands of an affinity matrix such as a chromatography support or bead with high affinity. An example of said tag is a histidine tag (His-tag or HT), such as a tag comprising 6
 15 residues of histidine (His6 or H6), which can bind to a column of nickel (Ni²⁺) or cobalt (Co²⁺) with high affinity. His-tag has the desirable feature that it can bind its ligands under conditions that are denaturing to most proteins and disruptive to most protein-protein interactions. Thus, it can be used to remove the bait protein tagged with H6 following the disruption of protein-protein interactions with which the bait has
 20 participated.

Additional illustrative, non-limitative, examples of tags useful for isolating or purifying a fusion protein include Arg-tag, FLAG-tag (DYKDDDDK; SEQ ID NO:52), Strep-tag (WSHPQFEK, SEQ ID NO:53), an epitope capable of being recognized by an antibody, such as c-myc-tag (recognized by an anti-c-myc antibody), HA tag
 25 (YPYDVPDYA, SEQ ID NO:54), V5 tag (GKPIPNPLLGLDST, SEQ ID NO:55), SBP-tag, S-tag, calmodulin binding peptide, cellulose binding domain, chitin binding domain, glutathione S-transferase-tag, maltose binding protein, NusA, TrxA, DsbA, Avi-tag, etc. (Terpe K., Appl. Microbiol. Biotechnol. 2003, 60:523-525), an amino acid sequence such as AHGHRP (SEQ ID NO:56) or PIHDHDHPLVHSGMTCXXC
 30 (SEQ ID NO:57), β -galactosidase and the like.

The tag can be used, if desired, for the isolation or purification of said fusion protein.

In another aspect, the invention relates to the conjugate or fusion protein according to the invention for use in medicine.

In another aspect, the invention relates to the conjugate or fusion protein according to the invention for use in the prevention and/or treatment of cancer.

5 In another aspect, the invention relates to the use of the conjugate or the fusion protein according to the invention for the preparation of a medicament for the prevention and/or treatment of cancer.

In another aspect, the invention relates to a method of treatment and/or prevention of cancer in a subject which comprises the administration to said subject of
10 the conjugate or of the fusion protein according to the invention.

In a preferred embodiment the cancer to be prevented or treated is Myc-induced cancer. In another preferred embodiment, the cancer to be prevented or treated is a cancer associated with a mutation in the *KRAS* gene. In one embodiment, the mutation in the *KRAS* gene is a mutation at the glycine at position 12, at the glycine at position 13
15 or at the glutamine at position 61. In a more preferred embodiment, the mutation is selected from the group consisting of the G12S mutation, the G12V mutation, the G13D mutation, the G12C mutation, the G12R mutation, the G12F mutation, the G12I mutation, the G13C mutation, the G13R mutation, or the Q61L mutation.

The terms “medicament”, “prevention”, “treatment”, “cancer” and “Myc-
20 induced cancer” have been defined previously and equally apply to this aspect of the invention.

Antitumor compositions

25 The unexpected finding that the Omomyc polypeptide is capable of translocating across biological membranes and exerting its tumor suppressor activity when provided as a polypeptide opens the possibility of formulating Omomyc with other antitumor drugs. Thus, in another aspect, the present invention also provides compositions containing, together or separately, a first component selected from the group consisting
30 of:

- (i) Omomyc,
- (ii) a functionally equivalent variant thereof,

(iii) a conjugate or fusion protein according to the invention
and, as a second component, an antitumor compound.

The first component of the compositions according to the invention includes a polypeptide according to SEQ ID NO:1, a functionally equivalent variant thereof or a
5 fusion protein according to the invention. Suitable polypeptides, functionally equivalent variants of the polypeptide of SEQ ID NO:1 and fusion proteins have been described above in the context of the therapeutic methods according to the invention and are equally applicable to the compositions according to the invention.

As used herein, “antitumoral agent” is understood as said biological or chemical
10 compound which treat tumors or prevent the formation thereof. In a preferred embodiment said antitumoral agent is selected from the group consisting of a cytotoxic agent, an antiangiogenic agent, an antimetastatic agent and an antiproliferative agent.

As used in the present invention, the term “cytotoxic agent” relates to an agent which is capable of promoting cell death and which has capacity for reducing the
15 growth, stopping the growth or destroying cells and, particularly, rapidly proliferating cells and, yet more particularly, tumor cells. Cell death can be caused by any mechanism, such as for example apoptosis, although it is not limited to this cause, by the metabolism inhibition, the interference with the organization of the cytoskeleton or the chemical modification of the DNA. The term cytotoxic agent comprises any
20 chemotherapy agent including small organic molecules, peptides, oligonucleotides and the like; toxins; enzymes; cytokines; radioisotopes or radiotherapy agents.

“Chemotherapy agents” are understood as chemical compounds such as, without limitation, anthracycline antibiotics such as doxorubicin and daunorubicin, taxanes such as Taxol™ and docetaxel, vinca alkaloids such as vincristine and vinblastine, 5-
25 fluorouracil (5-FU), leucovorin, irinotecan, idarubicin, mitomycin C, oxaliplatin, raltitrexed, tamoxifen, cisplatin, carboplatin, methotrexate, actinomycin D, mitoxantrone, blenoxane or mithramycin.

“Toxin” is understood as a toxic agent which conjugates with the polypeptide according to the first aspect of the invention or the fusion protein according to the
30 second aspect of the invention forming an immunotoxin. The conjugation of determined toxins with said polypeptide or with said fusion protein reduces the toxicity of the former, enabling their use as therapeutic agents, because otherwise they would be too

toxic. The binding between the toxin and the polypeptide according to the first aspect of the invention or the fusion protein according to the second aspect of the invention is performed chemically, conserving its biological activity. Their separation generally occurs in the lysosomes of the target cells such that the mentioned chemical binding is

5 only broken in the enclosed acidic cellular environment provided by the lysosomes. Toxins useful in the context of the present invention are plant toxins, bacterial toxins, toxins of fungal or animal origin and fragments thereof, such as, without limitation, the ricin A-chain, saponin, the diphtheria A-chain, active non-binding fragments of the diphtheria toxin, *Pseudomonas aeruginosa* exotoxin A-chain, abrin A-chain, modecin A-

10 chain, α -sarcin, *Leurites fordii* A-proteins, dianthin proteins, *Phytolaca americana* (PAPI, PAPII and PAP-S) proteins, *Momordica charantia* inhibitor, curcine, crotin, *Saponaria officinalis* inhibitor, gelonin, mitogelin, restrictocin, phenomycin, enomycin and trichothecenes.

“Enzymes” are understood in the context of the present invention as toxin or

15 drug activating enzymes, such as, without limitation, alkaline phosphatase which activates etoposide and doxorubicin; carboxypeptidase G2 which activates nitrogen mustards; beta-lactamase which activates doxorubicin, paclitaxel and mitomycin.

“Cytokines” are understood as peptides of different sizes and molecular weights which synthesize the cells of the immune system for the purpose of regulating the

20 immune response, and they can be hormones, growth factors, necrosis factors, etc. They can be of natural origin or from recombinant cell cultures and biologically active equivalents of natural sequence cytokines. Their conjugation with antibodies gives rise to immunocytokines. Cytokines useful in the present invention are, without limitation, TNF factor alpha, INF-gamma, GM-CSF factor or IL-2.

25 “Radioisotopes” is understood as radioactive isotopes such as, without limitation, ^{131}I , ^{90}Y , ^{177}Lu , ^{188}Re , ^{67}Cu , ^{211}At , ^{213}Bi , ^{125}I , ^{111}In .

“Antiangiogenic agent” is understood as a chemical or biological substance which inhibits or reduces the formation of new blood vessels, i.e., angiogenesis.

Antiangiogenic agents that can be conjugated with the polypeptide according to

30 the first aspect of the invention or with the fusion protein according to the second aspect of the invention include, without limitation, an antiangiogenic agent selected from the group of paclitaxel, 2-methoxyestradiol, prinomastat, batimastat, BAY 12-9566,

carboxyamidotriazole, CC-1088, dextromethorphan acetic acid, dimethylxanthenone acetic acid, endostatin, IM-862, marimastat, penicillamine, PTK787/ZK 222584, RPI.4610, squalamine lactate, SU5416, thalidomide, combretastatin, tamoxifen, COL-3, neovastat, BMS-275291, SU6668, anti-VEGF antibodies, Medi-522 (Vitaxin II), CAI, 5 interleukin 12, IM862, amiloride, angiostatin, K1-3 angiostatin, K1-5 angiostatin, Captopril, DL-alpha-difluoromethylornithine, DL-alpha-difluoromethylornithine HCl, endostatin, fumagillin, herbimycin A, 4-Hydroxyphenylretinamide, juglone, laminin, laminin hexapeptide, laminin pentapeptide, lavendustin A, medroxyprogesterone, minocycline, placenta ribonuclease inhibitor, suramin, thrombospondin, antibodies 10 directed against proangiogenic factors (for example, Avastin, Erbitux, Vectibix, Herceptin); low molecular weight tyrosine kinase inhibitors of proangiogenic growth factors (for example Tarceva, Nexavar, Sutent, Iressa); mTOR inhibitors (for example Torisel); interferon alpha, beta and gamma, IL-12, matrix metalloproteinase inhibitors (for example, COL3, marimastat, batimastat); ZD6474, SU11248, vitaxin; PDGFR 15 inhibitors (for example Gleevec); NM3 and 2- ME2; cyclopeptides such as cilengitide.

“Antimetastatic agent” is understood as a chemical or biological substance which inhibits or reduces metastasis, i.e., the distance propagation, fundamentally by the lymphatic or blood stream, of the cancer causing cells, and the growth of new tumors in the destination sites of said metastasis.

20 “Antiproliferative agent” is understood as a chemical or biological substance which is capable of preventing or inhibiting the formation or growth of tumors. Antiproliferative agents include but are not limited to (i) antimetabolites such as folic acid antimetabolites (aminopterin, denopterin, methotrexate, edatrexate, trimetrexate, nolatrexed, lometrexol, pemetrexed, raltitrexed, piritrexim, pteropterin, leucovorin, 10-propargyl-5,8- 25 dideazafolate (PDDF, CB3717)), purine analogs (cladribine, clofarabine, fludarabine, mercaptopurine, pentostatin, thioguanine) and pyrimidine analogs (capecitabine, cytarabine or ara-C, decitabine, fluorouracil, 5-fluorouracil, doxifluridine, floxuridine and gemcitabine) (ii) natural products, such as antitumor antibiotics and mitotic inhibitors such vinca alkaloids such as vindesine, vincristine, vinblastine, vinorelbine; 30 taxanes such as paclitaxel (Taxol™), docetaxel (Taxotere™); colchicine (NSC 757), thiocolchicine (NSC 361792), colchicine derivatives (e. g., NSC 33410), and allocolchicine (NSC 406042); halichondrin B (NSC 609395); dolastatin 10 (NSC

376128); maytansine (NSC 153858); rhizoxin (NSC 332598); epothilone A, epothilone B; discodermolide; estramustine; nocodazole; (iii) hormones and antagonist thereof, such tamoxifen, toremifene, anastrozole, arzoxifene, lasofoxifene, raloxifene, nafoxidine, fulvestrant, aminoglutethimide, testolactone, atamestane, exemestane, fadrozole, formestane, letrozole, goserelin, leuprorelin or leuprolide, buserelin, histrelin, megestrol and fluoxymesterone; (iv) biological agents, such as viral vectors, interferon alpha and interleukines; (v) platinum based compounds such as carboplatin, cisplatin [cis-diamminedichloroplatinum, (CDDP)], oxaliplatin, iproplatin, nedaplatin, triplatin tetranitrate, tetraplatin, satraplatin (JM216), JM118 [cis ammine dichloro (II)], JM149 [cis ammine dichloro (cyclohexylamine) trans dihydroxo platinum (IV)], JM335 [trans ammine dichloro dihydroxo platinum (IV)], transplatin, ZD0473, cis, trans, cis-Pt(NH₃)(C₆H₁₁NH₂)(O₂CC₃H₇)₂Cl, malanate-1,2-diaminocyclohexanoplatin(II), 5-sulphosalicylate-trans-(1,2- diaminocyclohexane)platin (II) (SSP), poly-[(trans-1,2-diaminocyclohexane)platin]- carboxyamyllose (POLY-PLAT) and 4-hydroxy-sulphonylphenylacetate (trans- 1,2- diaminocyclohexane) platinum (II) (SAP) and the like and (vi) DNA-alkylating drugs such as nitrogen mustards, nitrosoureas, ethylenimine derivatives, alkyl sulfonates and triazenes, including, but not limited to, cyclophosphamide (CytosanTM), busulfan, improsulfan, piposulfan, pipobroman, melphalan (L-sarcolysin), chlorambucil, mechlorethamine or mustine, uramustine or uracil mustard, novembichin, phenesterine, trofosfamide, ifosfamide, carmustine (BCNU), lomustine (CCNU), chlorozotocin, fotemustine, nimustine, ranimustine, semustine (methyl-CCNU), streptozocin, thiotepa, triethylenemelamine, triethylenethiophosphoramine, procarbazine, altretamine, dacarbazine, mitozolomide and temozolomide.

25

Pharmaceutical compositions

The invention also provides pharmaceutical compositions comprising the conjugates and fusion proteins of the invention or the compositions according to the invention. Thus, in another aspect, the invention provides a pharmaceutical composition comprising a pharmaceutically active amount of the conjugate or fusion protein according to the invention and a pharmaceutically active carrier (first pharmaceutical

30

composition of the invention). In another aspect, the invention provides a pharmaceutical composition comprising a pharmaceutically active amount of the composition according to the invention and a pharmaceutically active carrier (second pharmaceutical composition according to the invention).

5 As it is used in the present invention, the expression “pharmaceutical composition” relates to a formulation that has been adapted for administering a predetermined dose of one or several therapeutic useful agents to a cell, a group of cells, an organ, a tissue or an animal in which cell division is uncontrolled, such cancer.

 The first pharmaceutical composition of the invention contains a pharmaceutical
10 effective amount of a conjugate or fusion protein according to the invention. Suitable conjugates and fusion proteins for use in the pharmaceutical compositions according to the present invention include any of the fusion proteins mentioned above under the paragraph Omomyc conjugates and fusion proteins of the invention.

 The second pharmaceutical composition of the invention contains a
15 pharmaceutical effective amount of a composition according to the invention and a pharmaceutically active carrier. The second pharmaceutical composition of the invention comprise the polypeptide of SEQ ID NO:1, a functionally equivalent variant thereof or a fusion protein according to the invention. Suitable functionally equivalent variants of the polypeptide of SEQ ID NO:1 or suitable fusion proteins for use in the
20 second pharmaceutical compositions according to the invention are as defined above under therapeutic uses of the invention or under fusion proteins of the invention, respectively.

 The expression “Pharmaceutical effective amount”, as used herein, is understood as an amount capable of providing a therapeutic effect, and which can be determined by
25 the person skilled in the art by commonly used means. The amount of the Omomyc polypeptide, of the functionally equivalent variant thereof or of the fusion protein or of the antitumoral compound that may be combined in the pharmaceutical compositions according to the invention will vary depending upon the subject and the particular mode of administration. Those skilled in the art will appreciate that dosages may also be
30 determined with guidance from Goodman and Goldman's The Pharmacological Basis of Therapeutics, Ninth Edition (1996), Appendix II, pp. 1707-1711 and from Goodman

and Goldman's The Pharmacological Basis of Therapeutics, Tenth Edition (2001), Appendix II, pp. 475-493.

The appropriate dosage of the active principle or principles within the pharmaceutical composition will depend on the type of cancer to be treated, the severity and course of the disease, whether the composition is administered for preventive or therapeutic purposes, previous therapy, the patient's clinical history and response to the peptide or polypeptide, and the discretion of the attending physician. The amount of polypeptide of SEQ ID NO:1, of the functionally equivalent variant thereof, of the fusion protein is suitably administered to the patient at one time or over a series of treatments. Depending on the type and severity of the disease, an appropriate dosage level will generally be about 0.01 to 500 mg per kg patient body weight per day which can be administered in single or multiple doses. Preferably, the dosage level will be about 0.1 to about 250 mg/kg per day; more preferably about 0.5 to about 100 mg/kg per day. A suitable dosage level may be about 0.01 to 250 mg/kg per day, about 0.05 to 100 mg/kg per day, or about 0.1 to 50 mg/kg per day. Within this range the dosage may be 0.05 to 0.5, 0.5 to 5 or 5 to 50 mg/kg per day. For oral administration, the compositions are preferably provided in the form of tablets containing 1.0 to 1000 milligrams of the active ingredient, particularly 1.0, 5.0, 10.0, 15.0, 20.0, 25.0, 50.0, 75.0, 100.0, 150.0, 200.0, 250.0, 300.0, 400.0, 500.0, 600.0, 750.0, 800.0, 900.0, and 1000.0 milligrams of the active ingredient for the symptomatic adjustment of the dosage to the patient to be treated. The compounds may be administered on a regimen of 1 to 4 times per day, preferably once or twice per day.

In the case of the second pharmaceutical compositions according to the invention, which contain a first component selected from the polypeptide of SEQ ID NO:1, a functionally equivalent variant thereof and a fusion protein according to the invention and a second component which is an antitumoral agent, the composition may be presented as a single formulation (for example, as a tablet or a capsule comprising a fixed quantity of each one of the components) or can, on the other hand, be presented as separate formulations to be later combined for joint, sequential, or separate administration. The compositions of the invention also include the formulation as a kit-of-parts wherein the components are formulated separately but are packaged in the same container. Those skilled in the art will appreciate that the formulation of the different

components in the case of the second pharmaceutical composition according to the invention may be similar, in other words, similarly formulated (in tablets or pills), which allows their administration by the same route. In the case where the different components of the invention are formulated separately, the two components can be presented in a blister. Each blister contains the drugs that must be consumed during the day. If the drugs must be administered several times a day, the drugs corresponding to each administration can be placed in different sections of the blister, preferably recording in each section of the blister the time of day when they should be administered. Alternatively, the components of the composition of the invention can be formulated differently so that the different components are differently administered. Thus, it is possible that the first component is formulated as a tablet or capsule for its oral administration and the second component is formulated for its intravenous administration or vice versa. The ratio between the components that are part of the compositions used in the second pharmaceutical composition according to the invention can be adjusted by the skilled person depending on the antitumor agent used in each particular case, as well as of the desired indication. Thus, the invention envisages compositions wherein the ratio between the quantities of the two components can range from 50:1 to 1:50, in particular from 20:1 to 1:20, from 1:10 to 10:1, or from 5:1 to 1:5.

The first and the second pharmaceutical compositions of the invention can also contain one or several additional compounds for the prevention and/or treatment of pathologies in which there is an uncontrolled cell division, such as cancer. Said additional compounds, such as antitumoral agents can form part of the pharmaceutical composition as independent entities.

The first and the second pharmaceutical compositions of the invention also contain one or several additional pharmaceutically acceptable excipients. "Pharmaceutically acceptable excipient" is understood a therapeutically inactive substance said to be used for incorporating the active ingredient and which is acceptable for the patient from a pharmacological/toxicological point of view and for the pharmaceutical chemist who manufactures it from a physical/chemical point of view with respect to the composition, formulation, stability, acceptance of the patient and bioavailability.

The number and the nature of the pharmaceutically acceptable excipients

depend on the desired dosage form. The pharmaceutically acceptable excipients are known by the person skilled in the art (Faulí y Trillo C. (1993) “Tratado de Farmacia Galénica”, Luzán 5, S.A. Ediciones, Madrid). Said compositions can be prepared by means of the conventional methods known in the state of the art (“Remington: The
 5 Science and Practice of Pharmacy”, 20th edition (2003) Genaro A.R., ed., Lippincott Williams & Wilkins, Philadelphia, US).

The first and the second pharmaceutical compositions of the invention or can be administered by any type of suitable route, such as by oral route, topical route, by inhalation or parenteral route so that the pharmaceutically acceptable excipients
 10 necessary for the formulation of the desired dosage form will be included. The preferred route of administration of said pharmaceutical compositions is the endovenous route.

“Oral route” is understood as the pharmaceutical composition incorporated into the organism after deglutition. In a particular embodiment, the pharmaceutical composition of the invention can be in a dosage form suitable for its administration by oral
 15 route, whether it is solid or liquid. The dosage forms suitable for their administration by oral route can be tablets, capsules, syrups or solutions, and can contain any conventional excipient known in the art, such as binders, for example syrup, acacia, gelatin, sorbitol or polyvinylpyrrolidone; filling agents, for example lactose, sugar, corn starch, calcium phosphate, sorbitol or glycine; lubricants for compression, for example, magnesium
 20 stearate; disintegrating agents, for example starch, polyvinylpyrrolidone, sodium glycolate of starch or microcrystalline cellulose; or pharmaceutically acceptable wetting agents such as sodium lauryl sulfate. The solid oral compositions can be prepared by means of conventional processes of mixing, filling or compressing. Repetitive mixing operations can be used to completely distribute the active agent in those compositions that use high
 25 amounts of filling agents. Said operations are conventional in the art. The tablets can be prepared, for example, by means of wet or dry granulation, and optionally coating them according to the processes known in the common pharmaceutical practice, particularly with an enteric coating.

On the other hand, “topical route” is understood as an administration by non-
 30 systemic route, and includes the application of a pharmaceutical composition of the invention externally on the epidermis, in the oral cavity and the instillation of said composition into ears, eyes and nose, and in which it does not significantly enter the blood

stream. "Systemic route" is understood as the administration by oral route, intravenous route, intraperitoneal route and intramuscular route. The amount of antibody required for the therapeutic or prophylactic effect will naturally vary according to the elected antibody, the nature and the severity of the illness that is going to be treated, and the patient.

5 "Inhalation" is understood as the administration by intranasal route and by oral inhalation. The dosage forms suitable for said administration, such as a formulation in aerosol or a meter dosed inhaler can be prepared by means of conventional techniques. In an embodiment the route of administration is the intranasal route.

 As it is used herein, the term "parenteral", includes administration by intravenous
10 route, intraperitoneal route, intramuscular route or subcutaneous route. Subcutaneous, intramuscular and intravenous dosage forms of parenteral administration are generally preferred.

 In one embodiment, the first and the second pharmaceutical compositions of the invention can be adapted for their parenteral administration, such as sterile solutions,
15 suspensions or lyophilized products in the appropriate dosage unit form. The pharmaceutical compositions suitable for its injectable use include sterile aqueous solutions (when they are soluble in water), or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. For its administration by intravenous route, some suitable carriers include saline solution buffered
20 with phosphate (PBS). In all the cases, the composition must be sterile, and must be fluid to the point which that there exists easy ability for being injected. It must be stable in the preparation and storage conditions, and must be protected from the contamination action of microorganisms such as bacteria and fungi. The carrier can be a solvent or a dispersion medium which contains, for example, water, ethanol, a pharmaceutically acceptable polyol
25 such as glycerol, propylene glycol, liquid polyethylene glycol and suitable mixtures thereof. Suitable fluidity can be maintained, for example, by means of using a coating such as lecithin, by means of maintaining the particle size required in the case of dispersion and by means of using surfactants. The prevention of the action of the microorganisms can be achieved by means of various antibacterial and antifungal agents, for example, parabens,
30 chlorobutanol, phenol, ascorbic acid, thiomersal, and the like. In most cases, it will be preferable to include isotonic agents, for example, sugars; polyalcohols such as mannitol, sorbitol; or sodium chloride in the composition. The prolonged absorption of the injectable

compositions may be caused by the inclusion of an agent which delays the absorption, for example, aluminum and gelatin monostearate.

The injectable sterile solutions can be prepared by incorporating the active compound in the required amount in a suitable solvent with one or a combination of the
5 aforementioned ingredients, as needed, followed by sterilization by filtration through sterile membranes. Generally, the dispersions are prepared by incorporating the active compound in a sterile vehicle containing a basic dispersion medium and the rest of the ingredients required from among those previously listed. In the case of sterile powders for the preparation of injectable sterile solutions, the preferred preparation processes are
10 vacuum drying and lyophilization which give rise to a powder with the active ingredient plus any desired additional ingredient from a previously filtered sterile solution thereof.

The pharmaceutical compositions of the invention can be suitably administered by means of pulse infusion, for example, with decreasing doses of the composition. Preferably, the dose is administered by means of injections, more preferably intravenous or
15 subcutaneous injections, partly depending if the administration is acute or chronic.

In one embodiment, the first or second pharmaceutical compositions of the invention are prepared with carriers which will protect said polypeptide from a rapid elimination from the body, such as a controlled release formulation, including implants and microencapsulated administration systems. Biodegradable biocompatible polymers
20 such as ethylene vinylacetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters and polylactic acid can be used. The processes for preparing said formulations will be clear for persons skilled in the art. The materials can also be commercially obtained in Alza Corporation and Nova Pharmaceuticals, Inc.

The sustained release compositions also include preparations of antibody
25 crystals suspended in suitable formulations which can maintain the crystals in suspension. These preparations, when they are injected by subcutaneous or intraperitoneal route may produce a sustained release effect. Other compositions also include antibodies trapped in liposomes. The liposomes containing such antibodies are prepared by means of known methods such as Epstein *et al.*, Proc. Natl. Acad. Sci.
30 USA, (1985) 82:3688-3692; Hwang *et al.*, Proc. Natl. Acad. Sci. USA, (1980) 77:4030-4034; EP 52,322; EP 36,676; EP 88,046; EP 143,949.

Despite the fact that Omomyc and the conjugates and fusion proteins containing

Omomyc are capable of translocating across biological membranes, the skilled person will understand that it may also be convenient to formulate the conjugates or fusion proteins comprising Omomyc in nanoparticles. The nanoparticles may contribute to preserve the integrity of the polypeptide in the biological fluids until it reaches the target organ. Moreover, in the case of compositions comprising an antitumor agent, encapsulation of the composition may decrease secondary effects caused by the antitumor agent. Lastly, nanoparticles can also be modified so as to include moieties which allow the targeting of the nanoparticule to an organ of interest.

Thus, in another embodiment, the pharmaceutical compositions of the invention comprise the conjugates, fusion proteins and compositions according to the invention forming part of a nanoparticle.

Suitable nanoparticles that can be used in the context of the present invention include such nanoscale materials as a lipid-based nanoparticle, a superparamagnetic nanoparticle, a nanoshell, a semiconductor nanocrystal, a quantum dot, a polymer-based nanoparticle, a silicon-based nanoparticle, a silica-based nanoparticle, a metal-based nanoparticle, a fullerene and a nanotube.

Targeted delivery can be achieved by the addition of ligands without compromising the ability of nanoparticles to deliver their polypeptide payloads. It is contemplated that this will enable delivery to specific cells, tissues and organs. The targeting specificity of the ligand-based delivery systems are based on the distribution of the ligand receptors on different cell types. The targeting ligand may either be non-covalently or covalently associated with a nanoparticle, and can be conjugated to the nanoparticles by a variety of methods as discussed herein.

Examples of proteins or peptides that can be used to target nanoparticles include transferin, lactoferrin, TGF- β , nerve growth factor, albumin, HIV Tat peptide, RGD peptide, and insulin, as well as others.

The first and the second pharmaceutical compositions of the invention may be formulated with a pharmaceutically acceptable carrier. In a preferred embodiment, the carrier does not allow direct delivery of the fusion protein or of the composition to the cytoplasm of the cells, i.e. the carrier is not capable of fusing with the plasmatic membrane of the target cells. As used herein “carrier” is meant any substance that serves to improve the delivery and the effectiveness of the active principle within the

pharmaceutical composition. Examples of pharmaceutically acceptable carriers include one or more of water, saline, phosphate buffered saline, dextrose, glycerol, ethanol and the like, as well as combinations thereof. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as mannitol, sorbitol, or sodium chloride in the composition. Pharmaceutically acceptable carriers may further comprise minor amounts of auxiliary substances such as wetting or emulsifying agents, preservatives or buffers, which enhance the shelf life or effectiveness of the fusion protein or of the compositions forming part of the pharmaceutical compositions. Examples of proper carriers are well known in the literature (see for example Remington's Pharmaceutical Sciences, 19th ed., Mack Publishing Company, Easton, PA, 1995). Examples of carriers without limitation are a series of saccharide such as lactose, dextrose, sucrose, sorbitol, mannitol, xylitol, erythritol, and maltitol; a series of starch such as corn starch, wheat starch, rice starch, and potato starch; a series of cellulose such as cellulose, methyl cellulose, sodium carboxy methyl cellulose, and hydroxyl propylmethyl cellulose; and a series of filler such as gelatin and polyvinyl pyrrolidone. In some cases, a disintegrants such as cross-linked polyvinyl pyrrolidone, agar, alginic acid, or sodium alginate may be added.

The first and the second pharmaceutical compositions of the invention are suitable for the administration into any type of mammal, preferably a human being.

The invention is detailed below by means of the following examples which are merely illustrative and by no means limiting for the scope of the invention.

EXAMPLES

Materials and Methods

Example 1

Expression of the Omomyc construct in rich medium or in minimal M9 medium

For Omomyc production in rich medium, 1 L of 2YT medium containing ampicillin with BL21-AI™ One Shot® Chemically competent *Escherichia coli* (Life technologies) transformed with the Omomyc-pET-3a construct was inoculated, grown at 37°C with agitation at 250 rpm until OD₆₀₀ reached 0.8. The protein expression was

induced with 10 mL of Arabinose solution, incubating at 37°C for 12 h. Alternatively, for Omomyc production in M9 minimal medium, *E. coli* transformed with the Omomyc-pET3a construct was inoculated in 1L of M9 minimal medium containing chloramphenicol and ampicillin with BL21-CodonPlus (Stratagene) and grown at 37°C
 5 with agitation at 250 rpm until OD₆₀₀ reached 0.8. Protein expression was induced with 1 mL of IPTG 1000X solution. The cell culture was incubated with agitation at 250 rpm at 37°C for 12h.

The bacterial culture was centrifuged at 10,000 rpm, 4°C, 5 minutes in a SLA-1500 rotor using 250 mL bottles. The culture was centrifuged sequentially to combine
 10 the equivalent of 1L of culture per bottle.

In order to verify the successful protein expression, 1 mL aliquot from the cultures was collected and its OD₆₀₀ was measured. 800 µL of said aliquot was spined at 16,060 xg for 1 min and the supernatant was discarded. The pellet was suspended in a volume of IB Laemmli buffer equivalent to (0.1 x OD₆₀₀ value) µL in order to ensure
 15 equal quantities of lysed cell extract in each sample and facilitate comparison between cells before and after protein expression. Then the samples were mixed by vortex, sonicated for 5 seconds at maximum power, frozen in liquid nitrogen and finally, boiled for 2 minutes (repeated 3 times) prior to loading on a denaturing 16.5% acrylamide SDS-PAGE gel. Electrophoresis was followed by Coomassie blue staining or Western
 20 blot. The IB Laemmli buffer was optimized to allow solubilization of the inclusion bodies and to ensure adequate migration and protein separation to visualize the expression of Omomyc.

Example 2

25 Purification of the Omomyc b-HLH-LZ

Cell pellets were suspended in 3 mL of Lysis buffer per gram of pellet by vortexing. 150 µL of Triton solution per gram of culture pellet was added. The viscosity of the suspension was reduced by sonicating on ice, at power 15 for 6 x 15 seconds using an ultrasonic homogenizer.

30 Then, the equivalent of 100 µL/g of culture pellet of bovine pancreatic DNase I was added followed by an incubation of 60 minutes at 37°C with agitation at 50 rpm.

Samples were centrifuged 20 min at $12,000 \times g$ (13,000 rpm in a SS34 rotor in a Sorvall RC 5B Plus, use 35 mL centrifuge bottles), 4°C to pellet the inclusion bodies as well as high molecular weight complexes such as cell walls, ribosomes and non-degraded genomic DNA. Lipids, soluble proteins, amino acids, sugars and nucleic acids
5 constitute the supernatant that was discarded after centrifugation.

The pelleted inclusion bodies were solubilized with 15 mL of Bull cracker buffer part A by vortexing. This acidic, high ionic strength and denaturant buffer enables complete solubilization of the Omomyc construct from the inclusion bodies and allows elimination of high molecular weight complexes and residual DNA by centrifugation.

10 The inclusion bodies solution 1:1 was diluted with Bull cracker part B. After that, samples were centrifuged 30 min at $30,000 \times g$ (19,000 rpm in a SS34 rotor in a Sorvall RC 5B Plus) at 4°C .

The supernatant was purified by cation exchange chromatography on 5, cationic exchange columns of 5 mL (HiTrapTM SP Sepharose HP columns from GE Healthcare or the equivalent) mounted in series on a FPLC system. First, column was washed with
15 2 column volumes (CV) of FPLC buffer B at a flow rate of 5 mL/min. Then, the column was equilibrated with 5 CV (125 mL) of FPLC buffer A at a flow rate of 2.5 mL/min. The supernatant (equivalent to a maximum of 9 g of culture pellet per load) was loaded at 2.5 mL/min. Immediately after loading the supernatant, column was washed with 75
20 mL (3 CV) of U8 buffer, at a flow rate of 2.5 mL/min. Wash with 1 CV of buffer A. Then, 3 CV of 10% Buffer B was added to the column. Omomyc elution was done with a gradient of Buffer B of 10-35% in 50 mL (0.5%/mL) at 2.5 mL/min in 2.5 mL fractions. In these conditions, Omomyc eluted at ~ 1.5 M NaCl (i.e. $\sim 30\%$ v/v of the gradient) with a purity of 90%.

25 The fractions containing the pure construct were pooled and desalted on 5 consecutive HiTrapTM Desalting columns from GE Healthcare (or the equivalent) equilibrated with Desalting TFA buffer, at a 3.0 mL/min flow rate. The pooled fractions were injected (maximum volume of 7.5 mL per injection) and the elution was followed with OD₂₈₀ detector. The protein eluted within the first 15 mL after injection. The
30 following fractions showing low OD₂₈₀ values and high conductivity were discarded. This method allowed $\sim 95\%$ recovery of 98-99% desalted protein.

The purified protein can concentrate either by centrifugation in Amicon®Ultra centrifugal filters Ultracel®-3K (Millipore™) or the equivalent, or by lyophilization. For concentration using the Amicon®Ultra, follow manufacturer's indication. Alternatively, for lyophilization, add 50 % v/v acetonitrile to the desalted fractions,
 5 freeze in liquid nitrogen and lyophilize. The foamy lyophilized protein obtained can be weighted and resolubilized to 100 µl/mg in 50 % v/v acetonitrile containing 0.05 % v/v TFA, aliquoted 1 to 10 mg per eppendorf and lyophilized again.

Yields around 25 mg/L of culture in rich medium and 15 mg/L of culture in minimal M9 medium.

10

Example 3

Omomyc transduces into A549 cells

The authors of the invention aimed to demonstrate that cells treated with Omomyc show that, even after such a short time, Omomyc already reaches the nucleus.

15 Expression and purification of Omomyc was done as it is shown in Examples 1 and 2.

A549 cells (ATCC) were maintained in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% fetal bovine serum at 37°C in a humidified atmosphere containing 5% CO₂. For fluorescence microscopy, 20,000 A549 cells were
 20 seeded on 0.5 inch glass cover slips and grown for 16 hours. Fresh media supplemented with Omomyc peptide (35 µM) was added and incubated for 2 hours at 37°C. Cells were washed 3 times with PBS and fixed in 3% paraformaldehyde, and observed under the microscope.

Figure 1 shows that Omomyc can work as a protein transduction domain
 25 transducing across the cellular membrane and translocating to the nucleus.

Example 4

Omomyc reduces the number of viable A549 cells and induces apoptosis

A549 cells were incubated in 24-well plates over a period of 72 hours with 10
 30 µM Max* or Omomyc. Max* was obtained as described in Montagne M. et al. (Montagne M. et al. 2012. PLoS One, 7(2):e32172). Cells were harvested and stained with Annexin V and PI according to the manufacturer protocol (Tali® Apoptosis Kit

A10788, Life technologies) and the fluorescence quantified using Tali® image-based cytometer.

In this study the authors of the present invention show that Omomyc is more efficient at reducing the number of viable A549 cells than the bHLHZ domain of Max (Figure 2 A and C). Consistently, the percentage of dead/apoptotic cells is higher upon treatment with the Omomyc peptide than with the Max* peptide (Figure 2B and D).

Example 5

Omomyc is more folded than c-Myc* and more thermally stable than both Max* and c-Myc* in solution.

Thermally stable polypeptides are advantageous in those circumstances when the polypeptide has to be formulated as a pharmaceutical composition. Hence, the comparative thermal stability of Omomyc and Max* were determined.

The Omomyc gene was expressed as described in **Example 1** and purified as in **Example 2** above. The purification of Max* was carried out as described in detail in Beaulieu M.E. et al., 2013 (Methods Mol Biol, 1012:7-20). Briefly, BL21-DE3-pLys bacteria were transformed with a pET-3a expression vector containing the Max* insert, and cultured as described above. Induction was carried out with IPTG (0.6mM) for 4 hours, after which cells were harvested. After lysis with the Lysis buffer and centrifugation at 30 000 x g, the solubilized protein extract was purified using cation-exchange chromatography and desalted as described above. The purified and lyophilized proteins were resuspended in 50 mM KH₂PO₄, 50 mM KCl, 5 mM DTT (pH was adjusted using 1N KOH or 1N HCl).

Circular dichroism measurements were performed as described in Beaulieu M.E. et al., 2012 (J Mol Recognit, 25(7):414-26), using a protein concentration of 32 µM. The CD spectra were recorded at 20°C and the thermal denaturation was performed at 1°C/min (Figure 3A and 3B), monitoring at a wavelength of 222 nm to measure the helical content (reflecting the folded, structured state) of the protein.

Example 6

Omomyc penetrates nuclei of cells more efficiently than Max*

Briefly, the purified Omomyc and Max* peptides were labelled with fluorescein-maleimide (FITC) through an engineered C-terminal cysteine residue. After purification and assessment of the presence of a single fluorescent label per protein molecule by mass spectrometry, different concentrations of protein (5 μ M, 10 μ M or 25 μ M) were added to A549 cells (K-Ras mutant lung adenocarcinoma) cultured in RPMI containing 0.5% serum and incubated for 2h at 37°C before being washed thrice with PBS and fixed in 4% PFA for 10 min, stained with 1.5 μ g/mL Hoechst for 10 min and mounted on microscopy slides. The confocal microscopy images showed that Omomyc penetrates the nuclei of A549 cells more efficiently than Max* at concentrations of 5, 10 and 25 μ M (data not shown). Nuclear fluorescence intensity was quantified using ImageJ, with 30 to 80 cells counted per image. The quantification of the cell penetration observed for Omomyc and Max* in A549 fixed cells is shown in Figure 4.

The superior cell penetrating ability of Omomyc over Max* was confirmed at a concentration of 20 μ M in live cells (to avoid potential artifacts of fixation) after 20 minutes of incubation. Briefly, the FITC-labelled Omomyc and Max* were added to A549 culture cells in RPMI containing 0.5% serum and incubated for 20 minutes at 37°C before being washed thrice in PBS and mounted with Hoechst-containing mounting media. The confocal microscopy images showed that Omomyc penetrates the nuclei of A549 live cells more efficiently than Max* at 20 μ M (data not shown). Nuclear fluorescence intensity was quantified using ImageJ, with 30 to 80 cells counted per image. The quantification of the cell penetration observed for Omomyc and Max* in A549 live cells is shown in Figure 5.

Example 7

Omomyc entry occurs through an ATP-dependent process

Max* was shown to penetrate cells through an ATP-dependent process, which can be blocked lowering the temperature at 4°C. A similar mechanism is involved in Omomyc's cell penetrating ability. A549 cells were incubated for 2 hours with 20 μ M FITC-labelled peptide at 4°C and fixed with 4% PFA, before being washed thrice with PBS and mounted on microscopy slides using Hoechst-containing mounting media. Confocal microscopy images showed that Omomyc entry can be blocked lowering the temperature at 4°C .

Example 8

Omomyc reduces the total number of cells more efficiently than Max*

Briefly, A549 and H1650 lung adenocarcinoma cells were incubated with 25
 5 μ M Omomyc or Max* peptides in RPMI containing 5% serum and fixed at the
 indicated time (2 days or 4 days) with 4% PFA followed by crystal violet staining (0.5%
 crystal violet in 25% methanol for 10 minutes) and washing at least three times with
 water. Both Omomyc and Max* prevent growth of cells, but Omomyc is more efficient
 in doing so (Figure 6). The crystal violet stained cells were dissolved in 10% acetic acid
 10 and the resulting solution was diluted 1:4 in H₂O and quantified by measuring
 absorbance at OD₅₉₅. Quantification of the inhibition of proliferation by crystal violet
 showed that Omomyc is clearly more efficient than Max* to reduce proliferation of
 H1650 cells (EGFR-mutant lung adenocarcinoma cells) when used at 25 μ M
 concentration over 6 days (Figure 7).

15 In order to calculate the IC₅₀ of Omomyc and Max, A549 cells were treated with
 the indicated peptide concentration in culture media containing 5% serum and stained
 with crystal violet as described above. Quantification was performed as described
 above. Figure 8 shows dose response of A549 cells to Omomyc versus Max* showing
 that Omomyc has a lower IC₅₀ than Max* (i.e. Omomyc needs a lower dose compared
 20 to Max* in order to reduce the number of cells to a half). Specifically, IC₅₀ for
 Omomyc is 1.2 μ M and IC₅₀ for Max* is 2.6 μ M.

The effect of Omomyc on proliferation of cells was assayed in U87 glioma cells.
 U87 glioma cells were incubated with Omomyc or Max* peptides (25 μ M final
 concentration) in DMEM containing 5% serum for 48 h. Cells were trypsinized and
 25 counted using Tali cell counter (Life technologies Inc.). Figure 9 shows that Omomyc
 reduces the proliferation of U87 glioma cells more efficiently than Max*.

Example 9

Omomyc efficiently reaches lung and brain tissue after intranasal administration

30 Animals were treated with a single dose of 37.5 mg/kg of Omomyc or left
 untreated (treated with PBS) by intranasal administration. 10 minutes after intranasal
 administration the fluorescent peptide was detected. The lung of the treated animals

appears fluorescent as a consequence of Omomyc localization (Figure 10A). This demonstrates that the Omomyc peptide can be administered to animals by intranasal instillation and it efficiently reaches the lung. The brain of the same animals described in the previous figure also appears fluorescent (Figure 10B). It is apparent that the
5 fluorescent Omomyc peptide can pass the blood-brain barrier (BBB) and reach the brain.

Example 10

Omomyc peptide has a therapeutic effect *in vivo*

10 Administered as a peptide intranasally, Omomyc reduces proliferation in a KRas-driven lung adenocarcinoma mouse model. The inventors made use of a mouse model of KRas driven tumorigenesis (the LSLKRasG12D model). Briefly, 8 weeks old mice were instilled with an AdenoCre virus to induce expression of the KRas-G12D mutant protein specifically in the lungs. When the mice developed lung adenocarcinoma
15 (18 weeks after infection), the animals were treated intranasally with the Omomyc peptide (15 mg/kg in 35 μ L volume) daily for 3 days. Omomyc reduced the proliferative rate of tumors (Ki67 staining; Figure 11A) and reduced cellular density (Figure 11B).

The Omomyc peptide also reduces tumor burden *in vivo* after 1 week treatment.
20 Using the same mouse model and experimental conditions as described previously, the animals were treated daily for 1 week with 15 mg/kg of peptide. The % tumor area was measured using ImageJ. Figure 12 shows that Omomyc reduces tumor burden in a KRas-driven lung adenocarcinoma mouse model.

CLAIMS

1. A polypeptide of SEQ ID NO: 1 or a functionally equivalent variant thereof having a degree of identity with respect to SEQ ID NO: 1 greater than 70% for use in medicine, wherein the polypeptide or the functionally equivalent variant thereof is administered and wherein the functionally equivalent variant is a polypeptide which results from the insertion or addition of one amino acid and/or from the deletion of one or more amino acids and/or from the conservative substitution of one or more amino acids with respect to the polypeptide of SEQ ID NO: 1.
2. A polypeptide of SEQ ID NO: 1 or a functionally equivalent variant thereof having a degree of identity with respect to SEQ ID NO: 1 greater than 70% for use in the prevention and/or treatment of cancer, wherein the polypeptide or the functionally equivalent variant thereof is administered and wherein the functionally equivalent variant is a polypeptide which results from the insertion or addition of one amino acid and/or from the deletion of one or more amino acids and/or from the conservative substitution of one or more amino acids with respect to the polypeptide of SEQ ID NO: 1.
3. A conjugate comprising:
 - (i) the polypeptide of SEQ ID NO: 1 or a functionally equivalent variant thereof having a degree of identity with respect to SEQ ID NO: 1 greater than 70% and
 - (ii) a chemical moiety that facilitates cellular uptake of the polypeptide or of the functionally equivalent variant thereof.
4. The conjugate according to claim 3 wherein the chemical moiety that facilitates cellular uptake of the polypeptide or of the functionally equivalent variant thereof is a cell penetrating peptide and wherein said cell penetrating peptide and said polypeptide or the functionally equivalent variant thereof form a fusion protein.

5. The conjugate according to claim 4 wherein the cell-penetrating peptide sequence is selected from the group consisting of GRKKRRQRRR (SEQ ID NO: 28) and RRRRRRLR (SEQ ID NO: 29).
- 5 6. The conjugate according to any of claims 3 to 5 further comprising a nuclear-localization signal.
7. The conjugate according to claim 6 wherein the nuclear localization signal is selected from the group consisting of PKKKRKV (SEQ ID NO: 39),
10 PAAKRVKLD (SEQ ID NO: 50) and KRPAATKKAGQAKKKK (SEQ ID NO: 40).
8. The conjugate according to any of claims 3 to 6, wherein the functionally equivalent variant of the polypeptide of SEQ ID NO: 1 is selected from the group consisting of
15 SEQ ID NO: 47 and SEQ ID NO: 48.
9. A pharmaceutical composition comprising a pharmaceutically active amount of a conjugate according to any of claims 3 to 8 and a pharmaceutically acceptable excipient.
20
10. A conjugate according to any of claims 3 to 8 for use in medicine.
11. A conjugate according to any of claims 3 to 8 for use in the prevention and/or treatment of cancer.
25
12. The conjugate for use according to any of claims 10 or 11, wherein the conjugate is administered.
13. A composition comprising, together or separately:

- (i) a polypeptide of SEQ ID NO: 1, a functionally equivalent variant thereof having a degree of identity with respect to SEQ ID NO: 1 greater than 70% or a conjugate according to any of claims 3 to 8 and
- (ii) an antitumoral agent

5

14. The composition according to claim 13 wherein the antitumoral agent is selected from the group consisting of a cytotoxic agent, an antiangiogenic agent and an antimetastatic agent.

10 15. The composition according to any of claims 13 or 14, wherein the functionally equivalent variant of the polypeptide of SEQ ID NO: 1 is selected from the group consisting of SEQ ID NO: 47 and SEQ ID NO: 48.

15 16. A pharmaceutical composition comprising a pharmaceutically active amount of a composition according to any of claims 13 to 15 and a pharmaceutically acceptable excipient.

17. A composition according to any of claims 13 to 15 for use in medicine.

20 18. A composition according to any of claims 13 to 15 for use in the prevention and/or treatment of cancer.

25 19. The composition for use according to claims 17 or 18 wherein the polypeptide of SEQ ID NO: 1, the functionally equivalent variant thereof or the conjugate is administered.