RTMPLS copy

REPLACED SHEETS SG

- ☑ Description : paged ...1... to paged 101
- ☑ Claim(s) : paged 102 to paged 121
- ☐ Drawing(s) : paged ..... to paged .....
- ☑ Abstract : paged 122 to paged .....

138/CSQ55
139/CSQ56

5

10 TITLE OF THE INVENTION

HIV PROTEASE INHIBITORS USEFUL FOR THE TREATMENT OF AIDS

15 The present invention is concerned with compounds which inhibit the protease encoded by human immunodeficiency virus (HIV) or pharmaceutically acceptable salts thereof and are of value in the prevention of infection by HIV, the treatment of 20 infection by HIV and the treatment of the resulting acquired immune deficiency syndrome (AIDS). It also relates to pharmaceutical compositions containing the compounds and to a method of use of the present compounds and other agents for the 25 treatment of AIDS and viral infection by HIV.

30

This Sheet Has Been Replaced

BACKGROUND OF THE INVENTION

A retrovirus designated human immunodeficiency virus (HIV) is the etiological agent of the complex disease that includes progressive destruction of the immune system (acquired immune deficiency syndrome; AIDS) and degeneration of the central and peripheral nervous system. This virus was previously known as LAV, HTLV-III, or ARV. A common feature of retrovirus replication is the extensive post-translational processing of precursor polyproteins by a virally encoded protease to generate mature viral proteins required for virus assembly and function. Inhibition of this processing prevents the production of normally infectious virus. For example, Kohl, N.E. et al., Proc. Nat'l Acad. Sci. 85, 4686 (1988) demonstrated that genetic inactivation of the HIV encoded protease resulted in the production of immature, non-infectious virus particles. These results indicate that inhibition of the HIV protease represents a viable method for the treatment of AIDS and the prevention or treatment of infection by HIV.

The nucleotide sequence of HIV shows the presence of a pol gene in one open reading frame [Ratner, L. et al., Nature, 313, 277(1985)]. Amino acid sequence homology provides evidence that the pol sequence encodes reverse transcriptase, an endonuclease and an HIV protease [Toh, H. et al., EMBO J. 4, 1267 (1985); Power, M.D. et al., Science, 231, 1567 (1986); Pearl, L.H. et al., Nature 329, 351 (1987)]. Applicants demonstrate that the compounds of this invention are inhibitors of HIV protease.

This Sheet Has Been Replaced



BRIEF DESCRIPTION OF THE INVENTION

Compounds of formula I, as herein defined, are disclosed. These compounds are useful in the inhibition of HIV protease, the prevention of infection by HIV, the treatment of infection by HIV and in the treatment of AIDS, either as compounds, pharmaceutically acceptable salts, pharmaceutical composition ingredients, whether or not in combination with other antivirals, immunomodulators, antibiotics or vaccines. Methods of treating AIDS, methods of preventing infection by HIV, and methods of treating infection by HIV are also disclosed.

Some abbreviations that may appear in this application are as follows.

ABBREVIATIONS

Designation	Protecting Group
BOC (Boc)	t-butyloxycarbonyl
CBZ (Cbz)	benzyloxycarbonyl(carbobenzoxy)
TBS (TBDMS)	t-butyl-dimethylsilyl
	Activating Group
HBT(HOBT or HOBt)	1-hydroxybenzotriazole hydrate


This Sheet Has Been Replaced

Designation	Coupling Reagent
BOP reagent	benzotriazol-1-yloxytris-(dimethylamino)phosphonium hexafluorophosphate
BOP-Cl	bis(2-oxo-3-oxazolidinyl) phosphinic chloride
EDC	1-ethyl-3-(3-dimethylaminopropyl) carbodiimide hydrochloride
	Other
$(BOC)_2O$ (BOC_2O)	di-t-butyl dicarbonate
$n\text{-}Bu_4N^+F^-$	tetrabutyl ammonium fluoride
nBuLi (n-Buli)	n-butyllithium
DMF	dimethylformamide
Et_3N	triethylamine
EtOAc	ethyl acetate
TFA	trifluoroacetic acid
DMAP	dimethylaminopyridine
DME	dimethoxyethane
LDA	lithium diisopropylamide
THF	tetrahydrofuran
	Amino Acid
Ile	L-isoleucine
Val	L-valine

This Sheet Has Been Replaced

DETAILED DESCRIPTION OF THE INVENTION AND PREFERRED EMBODIMENTS

This invention is concerned with compounds of formula I, combinations thereof, or pharmaceutically acceptable salts thereof, in the inhibition of HIV protease, the prevention or treatment of infection by HIV and in the treatment of the resulting acquired immune deficiency syndrome (AIDS). Compounds of formula I are defined as follows:



I

wherein

X is -OH or $-NH_2$;

Z is -O, -S, or -NH;

R is hydrogen or C_{1-4} alkyl;

R^1 and R^2 are independently:

1) hydrogen,
2) $-C_{1-4}$ alkyl unsubstituted or substituted with one or more of
   a) halo,
   b) hydroxy,
   c) C_{1-3} alkoxy,
   d) aryl unsubstituted or substituted with one or more of C_{1-4}alkyl, hydroxy or aryl,

This Sheet Has Been Replaced

e) -W-aryl or -W-benzyl, wherein W is -O-, -S-, or -NH-,

f) a 5-7 membered cycloalkyl group unsubstituted or substituted with one or more of
   i) halo,
   ii) hydroxy,
   iii) C_{1-3} alkoxy, or
   iv) aryl,

g) heterocycle unsubstituted or substituted with one or more of hydroxy, C_{1-4}alkyl optionally substituted with hydroxy, or Boc,

h) $-NH-\overset{O}{\overset{\|}{C}}OC_{1-3}alkyl$,

i) $-NH-\overset{O}{\overset{\|}{C}}-C_{1-3}alkyl$,

j) $-NH-SO_2C_{1-3}alkyl$,

k) $-NR_2$,

l) -COOR, or

m) $-((CH_2)_mO)_nR$ wherein m is 2-5 and n is zero, 1, 2 or 3, or

3) aryl, unsubstituted or substituted with one or more of
   a) halo,
   b) hydroxy,
   c) $-NO_2$ or $-NR_2$,
   d) C_{1-4}alkyl,
   e) C_{1-3} alkoxy, unsubstituted or substituted with one or more of -OH or C_{1-3} alkoxy,

This Sheet Has Been Replaced

f) -COOR,

g) $-\overset{O}{\overset{\|}{C}}NR_2$,

h) $-CH_2NR_2$,

i) $-CH_2NH\overset{O}{\overset{\|}{C}}R$,

j) -CN,

k) $-CF_3$,

l) $-NH\overset{O}{\overset{\|}{C}}R$,

m) aryl C_{1-3} alkoxy,

n) aryl,

o) $-NRSO_2R$,

p) $-OP(O)(OR_x)_2$, or

q) $-R^5$, as defined below; or

R^1 and R^2 can be joined together to form with the nitrogen to which R^1 is attached a 3 to 10 membered monocyclic or bicyclic saturated ring system which consists of the nitrogen to which R^1 is attached and from 2 to 9 carbon atoms, and is unsubstituted or substituted with

1) hydroxy,

2) C_{1-4} alkyl unsubstituted or substituted with one or more of

a) halo,

b) hydroxy,

c) C_{1-3} alkoxy,

d) aryl,

This Sheet Has Been Replaced

e) a 5-7 membered cycloalkyl group unsubstituted or substituted with one or more of
i) halo,
ii) hydroxy,
iii) C_{1-3} alkoxy, or
iv) aryl,
f) heterocycle, or
g) $-NR_2$,

3) C_{1-3} alkoxy,

4) $-NH-\overset{O}{\overset{\|}{C}}OC_{1-3}alkyl$,

5) $-NH-\overset{O}{\overset{\|}{C}}-C_{1-3}alkyl$,

6) $-NH-SO_2C_{1-3}alkyl$,

7) heterocycle,

8) -W-aryl, or

9) $-W-\overset{O}{\overset{\|}{C}}-aryl$,

wherein W is defined above; or

R^1 and R^2 can be joined together to form with the nitrogen to which R^1 is attached a 3 to 10 membered monocyclic or bicyclic saturated ring system which consists of the nitrogen to which R^1 is attached, from 1 to 8 carbon atoms and one or more unsubstituted or substituted heteroatom selected from

1) $-\overset{|}{N}-V-R^1$,

wherein V is absent or $-\overset{O}{\overset{\|}{C}}-Q-$ or $-SO_2-Q-$,

This Sheet Has Been Replaced

R^1 is defined as above for when R^1 is independent from and not joined to R^2, and wherein Q is absent or -O-, -NR-, or heterocycle optionally substituted with

$-C_{1-4}$alkyl,

2) $-\underset{|}{N}-$ heterocycle,

3) $-\underset{|}{N}-$ C_{1-4} alkenyl, unsubstituted or substituted with aryl,

4) $-\underset{|}{N}-$ SO_2-C_{1-4}alkenyl, unsubstituted or substituted with aryl,

5) $-S(O)_p-$, wherein p is zero, 1 or 2, or

6) -O-; or

R^1 and R^2 can be joined together to form with the nitrogen to which R^1 is attached a 3 to 10 membered monocyclic or bicyclic saturated ring system, which consists of the nitrogen to which R^1 is attached and from 2 to 9 carbon atoms, in which the saturated ring system is fused to a phenyl ring and the phenyl ring is unsubstituted or substituted with one or more of

1) halo,
2) C_{1-3} alkoxy,
3) hydroxy,
4) C_{1-4} alkyl,
5) $-NHR^1$, wherein R^1 is defined as above for when R^1 is independent from and not joined to R^2, or
6) -NH-heterocycle;

This Sheet Has Been Replaced

R^3 is

1) $-(CH_2)_r-R^4$,
   wherein r is zero through 5,
2) C_{1-4}alkenyl-R^4, or
3) C_{1-4}alkynyl-R^4;

R^4 is

1) hydrogen,
2) C_{1-4} alkyl,
3) C_5-C_{10} cycloalkyl, optionally substituted with hydroxy,
4) C_6-C_{10} aryl, unsubstituted or substituted with one or more of
   a) halo,
   b) hydroxy,
   c) $-NO_2$ or $-NR_2$,
   d) C_{1-4}alkyl,
   e) C_{1-3} alkoxy, unsubstituted or substituted with one or more of -OH or C_{1-3} alkoxy,
   f) -COOR,
   g) $-\overset{\overset{O}{\|}}{C}NR_2$,
   h) $-CH_2NR_2$,
   i) $-CH_2NH\overset{\overset{O}{\|}}{C}R$,
   j) -CN,
   k) $-CF_3$,
   l) $-NH\overset{\overset{O}{\|}}{C}R$,
   m) aryl C_{1-3} alkoxy,
   n) aryl,

This Sheet Has Been Replaced

o) $-NRSO_2R$,
p) $-OP(O)(OR_x)_2$, or
q) $-R^5$, as defined below, or

5) monocyclic or bicyclic heterocyle containing from 1 to 3 heteroatoms chosen from the group consisting of N, O, and S and which is unsubstituted or substituted with R^5 and optionally with one or more of
a) halo,
b) C_{1-4} alkyl, or
c) C_{1-3} alkoxy;

R_x is H or aryl;

R^5 is

1) $-W-(CH_2)_m-NR^6R^7$
wherein W is as defined above,
m is 2-5, and
R^6 and R^7 are independently
a) hydrogen,
b) C_{1-6} alkyl, unsubstituted or substituted with one or more of
i) C_{1-3} alkoxy,
ii) -OH, or
iii) $-NR_2$,
c) the same or different and joined together to form a 5-7 member heterocycle, such as morpholino, containing up to two additional heteroatoms selected from

$$-\overset{R}{\underset{|}{N}}-, \quad -O-, \quad -\overset{O}{\underset{\|}{S}}-, \quad -S-, \quad \text{or} \quad -SO_2-,$$


the heterocycle optionally substituted with C_{1-4} alkyl, or

This Sheet Has Been Replaced

d) aromatic heterocycle unsubstituted or substituted with one or more of
i) C_{1-4} alkyl, or
ii) $-NR_2$,

2) $-(CH_2)_q-NR^6R^7$ wherein q is 1-5, and R^6 and R^7 are defined above, except that R^6 or R^7 are not H or unsubstituted C_{1-6} alkyl, or

3) benzofuryl, indolyl, azacycloalkyl, azabicyclo C_{7-11} cycloalkyl, or benzopiperidinyl, unsubstituted or substituted with C_{1-4} alkyl;

B is absent, or ,

wherein R^8 is
1) $-CH(CH_3)_2$,
2) $-CH(CH_3)(CH_2CH_3)$, or
3) -phenyl;

J^1 and J^2 are independently
1) $-YR^9$ wherein
Y is -O- or -NH-, and
R^9 is
a) hydrogen,
b) C_{1-6} alkyl, unsubstituted or substituted with one or more of
i) $-NR_2$,
ii) -OR,
iii) $-NHSO_2C_{1-4}$ alkyl,

This Sheet Has Been Replaced

iv) $-NHSO_2$ aryl, or $-NHSO_2$(dialkyl-aminoaryl),

v) $-CH_2OR$,

vi) $-C_{1-4}$ alkyl,

vii) $-\overset{O}{\overset{\|}{C}}OR$,

viii) $-\overset{O}{\overset{\|}{C}}NR_2$,

ix) $-NH\!-\!\overset{NH}{\overset{\|}{C}}\!-\!NR_2$ or $-NH\!-\!\overset{N-CN}{\overset{\|}{C}}\!-\!NR_2$,

x) $-NH\overset{O}{\overset{\|}{C}}R^{13}$, wherein R^{13} is

A) -H,

B) $-C_{1-4}$ alkyl,

C) -aryl,

D) -heterocycle, or

E) -NH-, -O- or $-(CH_2)_n-$ wherein n is zero, 1, 2 or 3, substituted with

I) $-C_{1-4}$ alkyl, unsubstituted or substituted with one or more of aryl or heterocycle, or

II) aryl, unsubstituted or substituted with heterocycle,

xi) $-NR_3^{\oplus}$ $A^{\ominus}$ wherein $A^{\ominus}$ is a counterion,

This Sheet Has Been Replaced

xii) $-NR^{10}R^{11}$ wherein R^{10} and R^{11} are the same or different and are C_{1-5} alkyl joined together directly to form a 5-7 membered heterocycle containing up to one additional heteroatom selected from -O-, -S-, or -NR-,

xiii) aryl,

xiv) -CHO,

xv) $-OP(O)(OR_x)_2$,

xvi) $-O-\overset{O}{\overset{\|}{C}}-C_{1-4}$alkyl substituted with one or more of amine or quaternary amine, or $-O-((CH_2)_mO)_n-R$, or $-OP(O)(OR_x)_2$,


xvii) $-O\overset{O}{\overset{\|}{C}}-R$, or

xviii) $-O\overset{O}{\overset{\|}{C}}-NH-CH_2$-heterocycle, or

c) $-((CH_2)_mO)_nCH_3$ or $-((CH_2)_mO)_n$ H, wherein m and n are defined above,

2) $-N(R^9)_2$,

3) $-NR^{10}R^{11}$ wherein R^{10} and R^{11} are defined above, or

4) 

wherein Y, R^9 and n are defined above; and

This Sheet Has Been Replaced

R^{12} is

1) hydrogen,
2) aryl, unsubstituted or substituted with one or more of
   a) R^{14}, wherein R^{14} is
      i) halo,
      ii) -OR,
      iii) $-\overset{O}{\overset{\|}{C}}NR_2$,
      iv) $-CH_2NR_2$,
      v) $-SO_2NR_2$,
      vi) $-NR_2$,
      vii) $-NH\overset{O}{\overset{\|}{C}}R$,
      viii) C_{1-4} alkyl,
      ix) phenyl
      x) $-CF_3$,
      xi) $-\overset{R}{\overset{|}{N}}-SO_2R$,
      xii) $-OP(O)(OR_x)_2$, or
      xiii) $-\overset{O}{\overset{\|}{C}}OR$,
   b) $-C_{1-4}$ alkyl-NR_2, or
   c) $-O-\overset{O}{\overset{\|}{C}}-C_{1-4}$alkyl substituted with one or more of amine or quaternary amine or $-OP(O)(OR_x)_2$,

This Sheet Has Been Replaced

3) heterocycle, such as isochroman, chroman, isothiochroman, thiochroman, benzimidazole, benzothiopyran, oxobenzothiopyran, benzopyran, benzothiopyranylsulfone, benzothiopyranylsulfoxide, the ring or rings being unsubstituted or substituted with one or more of
a) R^{14}, as defined above,
b) $-OC_{1-4}$ alkenyl,
c) phenyl-C_{1-4} alkyl,
d) $-O-\overset{O}{\overset{\|}{C}}-C_{1-4}$alkyl substituted with one or more of amine or quaternary amine, or $-OP(O)(OR_x)_2$, or $-O((CH_2)_mO)_n-R$, or
e) $-O-\overset{O}{\overset{\|}{C}}-O-((CH_2)_mO)_n-R$, or

4) A 5 to 7 membered carbocyclic or 7-10 membered bicyclic carbocyclic ring, such as cyclopentane, cyclohexane, indane, norbornane, naphthalene, thiopyran, isothiopyran, or benzopyran, the carbocyclic ring being unsubstituted or substituted with one or more of
a) R^{14}, as defined above,
b) $-CH_2OR$,
c) $-(CH_2)_n-NR_2$, C_{5-16}alkyl, pyridine, $-(CH_2)_nNR-(CH_2)_n-NR_2$, $-(CH_2)_n-\overset{O}{\overset{\|}{C}}-OR$,

This Sheet Has Been Replaced

$-((CH_2)_mO)_n-R$, quinuclidiniumyl substituted with R, piperazine-C_{1-4}alkyl-benzyl substituted once or more with R, or morpholino-C_{1-4}alkyl-benzyl,

d) $-O-\overset{O}{\overset{\|}{C}}-C_{1-4}$alkyl substituted with one or more of amine or quaternary amine, $-OP(O)(OR_x)_2$, or $-O-((CH_2)_mO)_n-R$,

e) $-O-\overset{O}{\overset{\|}{C}}-O-((CH_2)_mO)_n-R$, or

f) $-C_{1-4}$alkyl-phenyl;

or a pharmaceutically acceptable salt thereof.

In a preferred embodiment of this invention, R^1 and R^2 are joined together to form with the nitrogen to which R^1 is attached a 3 to 10 membered monocyclic or bicyclic saturated ring system which consists of the nitrogen to which R^1 is attached and from 2 to 9 carbon atoms, and is unsubstituted or substituted with

1) hydroxy,
2) C_{1-4} alkyl unsubstituted or substituted with one or more of
   a) hydroxy,
   b) C_{1-3} alkoxy,
   c) aryl,
   d) a 5-7 membered cycloalkyl group unsubstituted or substituted with one or more of

This Sheet Has Been Replaced

i) halo,
ii) hydroxy,
iii) C_{1-3} alkoxy, or
iv) aryl,

e) heterocycle, or

f) $-NR_2$,

3) C_{1-3} alkoxy,

4) $-NH-\overset{O}{\overset{\|}{C}}OC_{1-3}alkyl$,

5) $-NH-\overset{O}{\overset{\|}{C}}-C_{1-3}alkyl$,

6) $-NH-SO_2C_{1-3}alkyl$,

7) -W-aryl, or

8) $-W-\overset{O}{\overset{\|}{C}}-aryl$,

wherein W is -O-, -S-, or -NH-; or

R^1 and R^2 are joined together to form with the nitrogen to which R^1 is attached a 3 to 10 membered monocyclic or bicyclic saturated ring system which consists of the nitrogen to which R^1 is attached, from 1 to 8 carbon atoms and one or more unsubstituted or substituted heteroatom selected from

1) $-\underset{\underset{V-R^1}{|}}{N}-$,

wherein V is absent or $-\overset{O}{\overset{\|}{C}}-Q-$ or $-SO_2-Q-$, R^1 is defined as above for when R^1 is independent from and not joined to R^2, and wherein Q is absent or -O-, -NR-, or heterocycle optionally substituted with $-C_{1-4}alkyl$,

This Sheet Has Been Replaced

2) $-\underset{\underset{C_{1-4}\text{ alkenyl}}{|}}{N}-$, unsubstituted or substituted with aryl,

3) $-S(O)_p-$, wherein p is zero, 1 or 2, or

4) $-O-$; or

R^1 and R^2 are joined together to form with the nitrogen to which R^1 is attached a 3 to 10 membered monocyclic or bicyclic saturated ring system, which consists of the nitrogen to which R^1 is attached and from 2 to 9 carbon atoms, in which the saturated ring system is fused to a phenyl ring and the phenyl ring is unsubstituted or substituted with one or more of

1) C_{1-3} alkoxy,

2) hydroxy,

3) C_{1-4} alkyl, or

4) $-NHR^1$,

wherein R^1 is defined as above for when R^1 is independent from and not joined to R^2.

A second, more preferred embodiment of this invention is further limited to compounds where:

R^1 and R^2 are joined together to form with the nitrogen to which R^1 is attached a 3 to 10 membered monocyclic or bicyclic saturated ring system which consists of the nitrogen to which R^1 is attached and from 2 to 9 carbon atoms, and is unsubstituted or substituted with

This Sheet Has Been Replaced

1) hydroxy,
2) C_{1-4} alkyl unsubstituted or substituted with one or more of
   a) hydroxy,
   b) C_{1-3} alkoxy,
   c) aryl,
   d) a 5-7 membered cycloalkyl group unsubstituted or substituted with one or more of
      i) halo,
      ii) hydroxy,
      iii) C_{1-3} alkoxy, or
      iv) aryl,
   e) heterocycle, or
   f) $-NR_2$,
3) C_{1-3} alkoxy,
4) $-NH-\overset{O}{\overset{\|}{C}}OC_{1-3}$alkyl,
5) $-NH-\overset{O}{\overset{\|}{C}}-C_{1-3}$alkyl,
6) $-NH-SO_2C_{1-3}$alkyl,
7) -W-aryl, or
8) $-W-\overset{O}{\overset{\|}{C}}$-aryl,

wherein W is -O-, -S-, or -NH-; or

R^1 and R^2 are joined together to form with the nitrogen to which R^1 is attached a 3 to 10 membered monocyclic or bicyclic saturated ring system which consists of the nitrogen to which R^1 is attached, from 1 to 8 carbon atoms and one or more unsubstituted or substituted heteroatom selected from

This Sheet Has Been Replaced

1) $-N(V-R^1)-$,

wherein V is absent or $-\overset{O}{\overset{\|}{C}}-Q-$ or $-SO_2-Q-$, R^1 is defined as above for when R^1 is independent from and not joined to R^2, and wherein Q is absent or -O-, -NR-, or heterocycle optionally substituted with $-C_{1-4}$alkyl,

2) $-S(O)_p-$,

wherein p is zero, 1 or 2, or

3) -O-;

R^3 is benzyl, unsubstituted or substituted with one or more of

a) hydroxy,

b) $-NO_2$, or $-NR_2$,

c) C_{1-4}alkyl,

d) C_{1-3} alkoxy, unsubstituted or substituted with one or more of -OH or C_{1-3} alkoxy,

e) $-\overset{O}{\overset{\|}{C}}NR_2$,

f) $-CH_2NR_2$,

g) $-CH_2NH\overset{O}{\overset{\|}{C}}R$,

h) $-CF_3$,

i) $-NH\overset{O}{\overset{\|}{C}}R$,

j) $-NRSO_2R$,

k) $-OP(O)(OR_x)_2$, or

l) $-R^5$;

This Sheet Has Been Replaced

and B is absent.

A third, most preferred embodiment of this invention is further limited to compounds where:

X is -OH;


Z is -O;

R^1 and R^2 are joined together to form with the nitrogen to which R^1 is attached a 3 to 10 membered monocyclic or bicyclic saturated ring system which consists of the nitrogen to which R^1 is attached and from 2 to 9 carbon atoms, and is unsubstituted or substituted with -W-aryl or -W-C(=O)-aryl; or

R^1 and R^2 are joined together to form with the nitrogen to which R^1 is attached a 3 to 10 membered monocyclic or bicyclic saturated ring system which consists of the nitrogen to which R^1 is attached, from 1 to 8 carbon atoms and one of -N(V-R^1)- ,

wherein V is absent or -C(=O)-Q- or -SO_2-Q-,
R^1 is defined as above for when R^1 is independent from and not joined to R^2,
and wherein Q is absent or -O-, -NR- or heterocycle optionally substituted with -C_{1-4}alkyl;


R^3 is benzyl, unsubstituted or substituted with one or more of (1) hydroxy, (2) C_{1-3} alkoxy substituted with one or more of -OH or (3)



;


This Sheet Has Been Replaced

J^1 is $-NH-C_{1-4}$alkyl; and
J^2 is



The most preferred compounds of this invention are compounds A through H and J, shown below.


Compound A:



N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(2-(3(S)-N'-(t-butylcarboxamido)-(4aS,8aS)-decahydroisoquinoline)yl)-pentaneamide,


This Sheet Has Been Replaced

Compound B:



N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(1-(4-carbobenzyloxy-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide,


Compound C:



N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-(2-(4-morpholinyl)ethoxy)phenyl)methyl)-4(S)-hydroxy-5-(2-(3(S)-N'-(t-butylcarboxamido)-(4aS,8aS)-decahydroisoquinoline)yl)-pentaneamide,


This Sheet Has Been Replaced

Compound D:



N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-(2-(4-morpholinyl)ethoxy)phenyl)methyl)-4(S)-hydroxy-5-(1-(4-carbobenzyloxy-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide,


Compound E:



N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-((2-hydroxy)-ethoxy)phenyl)methyl)-4(S)-hydroxy-5-(2-(3(S)-N'-(t-butylcarboxamido)-(4aS,8aS)-decahydroisoquinoline)-yl)-pentaneamide,


This Sheet Has Been Replaced

Compound F:



N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-((2-hydroxy)-ethoxy)phenyl)methyl)-4(S)-hydroxy-5-(1-(4-carbobenzyloxy-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide,


Compound G:



N-(4(S)-3,4-dihydro-1H-2,2-dioxobenzothiopyranyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(2-(3(S)-N'-(t-butyl-carboxamido)-(4aS,8aS)-decahydroisoquinoline)yl)-pentaneamide,


This Sheet Has Been Replaced

Compound H:



N-(4(S)-3,4-dihydro-1H-2,2-dioxobenzothiopyranyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(1-(4-carbobenzyloxy-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide,

Compound J:



N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(1-(4-(3-pyridylmethyl)-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide.

Novel compounds of the present invention also include but are not limited to the following compounds:

This Sheet Has Been Replaced

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4-(S)-hydroxy-5-(1-(N'-(t-butyl)-4(S)-phenoxyproline-amid)yl)-pentaneamide,

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4-(S)-hydroxy-5-(1-(N'-t-butyl-4(S)-2-naphthyloxy-prolineamid)yl)-pentaneamide,

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4-(S)-hydroxy-5-(1-(N'-t-butyl-4(S)-1-naphthyloxy-prolineamid)yl)-pentaneamide,

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4-(S)-amino-5-(2-(3(S)-N'-(t-butylcarboxamido)-(4aS,8aS)-decahydroisoquinoline)yl)-pentaneamide,

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4-(S)-hydroxy-5-(1-(4-(3-phenylpropionyl)-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide,

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4-(S)-hydroxy-5-(1-(4-benzoyl-2(S)-N'-(t-butylcarbox-amido)-piperazinyl))-pentaneamide,

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4-(S)-hydroxy-5-(1-(4-(3-phenylpropyl)-2(S)-N'-(t-butyl-carboxamido)-piperazinyl))-pentaneamide,

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4-(S)-amino-5-(1-(4-carbobenzyloxy-2(S)-N'-(t-butyl-carboxamido)-piperazinyl))-pentaneamide,

This Sheet Has Been Replaced

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-(2-(4-morpholinyl)ethoxy)phenyl)methyl)-4(S)-hydroxy-5-(1-(N'-(t-butyl)-4(S)-phenoxyprolineamid)yl)-pentaneamide,

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-(2-(4-morpholinyl)ethoxy)phenyl)methyl)-4(S)-hydroxy-5-(1-(N'-t-butyl-4(S)-2-naphthyloxy-prolineamid)yl)-pentaneamide,

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-(2-(4-morpholinyl)ethoxy)phenyl)methyl)-4(S)-hydroxy-5-(1-(N'-t-butyl-4(S)-1-naphthyloxy-prolineamid)yl)-pentaneamide,

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-(2-(4-morpholinyl)ethoxy)phenyl)methyl)-4(S)-amino-5-(2-(3(S)-N'-(t-butylcarboxamido)-(4aS,8aS)-decahydroisoquinoline)yl)-pentaneamide,

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-(2-(4-morpholinyl)ethoxy)phenyl)methyl)-4(S)-hydroxy-5-(1-(4-(3-phenylpropionyl)-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide,

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-(2-(4-morpholinyl)ethoxy)phenyl)methyl)-4(S)-hydroxy-5-(1-(4-benzoyl-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide,

This Sheet Has Been Replaced

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-(2-(4-morpholinyl)ethoxy)phenyl)methyl)-4(S)-hydroxy-5-(1-(4-(3-phenylpropyl)-2(S)-N'-(t-butylcarboxamido))-piperazinyl)-pentaneamide,

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-(2-(4-morpholinyl)ethoxy)phenyl)methyl)-4(S)-amino-5-(1-(4-carbobenzyloxy-2(S)-N'-(t-butylcarboxamido)-piperazinyl)pentaneamide,

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-((2-hydroxy)-ethoxy)phenyl)methyl)-4(S)-hydroxy-5-(1-(N'-(t-butyl)-4(S)-phenoxyprolineamid)yl)-pentaneamide,

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-((2-hydroxy)-ethoxy)phenyl)methyl)-4(S)-hydroxy-5-(1-(N'-t-butyl-4(S)-2-naphthyloxy-prolineamid)yl)-pentaneamide,

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-((2-hydroxy)-ethoxy)phenyl)methyl)-4(S)-hydroxy-5-(1-(N'-t-butyl-4(S)-1-naphthyloxy-prolineamid)yl)-pentaneamide,

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-((2-hydroxy)-ethoxy)phenyl)methyl)-4(S)-amino-5-(2-(3(S)-N'-(t-butylcarboxamido)-(4aS,8aS)-decahydroisoquinoline)-yl)pentaneamide,

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-((2-hydroxy)-ethoxy)phenyl)methyl)-4(S)-hydroxy-5-(1-(4-(3-phenyl-propionyl)-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide,

This Sheet Has Been Replaced

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-((2-hydroxy)-ethoxy)phenyl)methyl)-4(S)-hydroxy-5-(1-(4-benzoyl-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide,

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-((2-hydroxy)-ethoxy)phenyl)methyl)-4(S)-hydroxy-5-(1-(4-(3-phenyl-propyl)-2(S)-N'-(t-butylcarboxamido))-piperazinyl)-pentaneamide,

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-((2-hydroxy)-ethoxy)phenyl)methyl)-4(S)-amino-5-(1-(4-carbobenzyl-oxy-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide,

N-(4(S)-3,4-dihydro-1H-2,2-dioxobenzothiopyranyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(1-(N'-(t-butyl)-4(S)-phenoxyprolineamid)yl)-pentaneamide,

N-(4(S)-3,4-dihydro-1H-2,2-dioxobenzothiopyranyl)-2-(R)-phenylmethyl-4(S)-hydroxy-5-(1-(N'-t-butyl-4(S)-2-naphthyloxy-prolineamid)yl)-pentaneamide,

N-(4(S)-3,4-dihydro-1H-2,2-dioxobenzothiopyranyl)-2-(R)-phenylmethyl-4(S)-hydroxy-5-(1-(N'-t-butyl-4(S)-1-naphthyloxy-prolineamid)yl)-pentaneamide,

N-(4(S)-3,4-dihydro-1H-2,2-dioxobenzothiopyranyl)-2-(R)-phenylmethyl-4(S)-amino-5-(2-(3(S)-N'-(t-butyl-carboxamido)-(4aS,8aS)-decahydroisoquinoline)yl)-pentaneamide,

This Sheet Has Been Replaced

N-(4(S)-3,4-dihydro-1H-2,2-dioxobenzothiopyranyl)-2-(R)-phenylmethyl-4(S)-hydroxy-5-(1-(4-(3-phenylpropionyl)-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide,

N-(4(S)-3,4-dihydro-1H-2,2-dioxobenzothiopyranyl)-2-(R)-phenylmethyl-4(S)-hydroxy-5-(1-(4-benzoyl-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide,

N-(4(S)-3,4-dihydro-1H-2,2-dioxobenzothiopyranyl)-2-(R)-phenylmethyl-4(S)-hydroxy-5-(1-(4-(3-phenylpropyl)-2(S)-N'-(t-butylcarboxamido))-piperazinyl)-pentaneamide, or

(4(S)-3,4-dihydro-1H-2,2-dioxobenzothiopyranyl)-2(R)-phenylmethyl-4(S)-amino-5-(1-(4-carbobenzyloxy-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide.

The compounds of the present invention, may have asymmetric centers and occur as racemates, racemic mixtures and as individual diastereomers, or enantiomers with all isomeric forms being included in the present invention.


When any variable (e.g., aryl, heterocycle, R, R^1, R^2, A^-, n, Z, etc.) occurs more than one time in any constituent or in formula I, its definition on each occurrence is independent of its definition at every other occurrence. Also, combinations of substituents and/or variables are permissible only if such combinations result in stable compounds.

This Sheet Has Been Replaced

As used herein except where noted, "alkyl" is intended to include both branched- and straight-chain saturated aliphatic hydrocarbon groups having the specified number of carbon atoms (Me is methyl, Et is ethyl, Pr is propyl, Bu is butyl); "alkoxy" represents an alkyl group of indicated number of carbon atoms attached through an oxygen bridge; and "cycloalkyl" is intended to include saturated ring groups, such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl (Cyh) and cycloheptyl. "Alkenyl" is intended to include hydrocarbon groups of either a straight or branched configuration with one or more carbon-carbon double bonds which may occur in any stable point along the chain, such as ethenyl, propenyl, butenyl, pentenyl, and the like. "Alkynyl" is intended to include hydrocarbon groups of either a straight or branched configuration with one or more carbon-carbon triple bonds which may occur in any stable point along the chain, such as ethynyl, propynyl, butynyl, pentynyl, and the like. "Halo", as used herein, means fluoro, chloro, bromo and iodo; and "counterion" is used to represent a small, single negatively-charged species, such as chloride, bromide, hydroxide, acetate, trifluroacetate, perchlorate, nitrate, benzoate, maleate, tartrate, hemitartrate, benzene sulfonate, and the like.

As used herein, with exceptions as noted, "aryl" is intended to mean phenyl (Ph) or naphthyl. "Carbocyclic" is intended to mean any stable 5- to 7-membered carbon ring or 7- to 10-membered bicyclic carbon ring any ring of which may be saturated or unsaturated.

This Sheet Has Been Replaced



The term heterocycle or heterocyclic, as used herein except where noted, represents a stable 5- to 7-membered mono- or bicyclic or stable 7- to 10-membered bicyclic heterocyclic ring system any ring of which may be saturated or unsaturated, and which consists of carbon atoms and from one to three heteroatoms selected from the group consisting of N, O and S, and wherein the nitrogen and sulfur heteroatoms may optionally be oxidized, and the nitrogen heteroatom may optionally be quaternized, and including any bicyclic group in which any of the above-defined heterocyclic rings is fused to a benzene ring. The heterocyclic ring may be attached at any heteroatom or carbon atom which results in the creation of a stable structure. Examples of such heterocyclic elements include piperidinyl, piperazinyl, 2-oxopiperazinyl, 2-oxopiperidinyl, 2-oxopyrrolodinyl, 2-oxoazepinyl, azepinyl, pyrrolyl, 4-piperidonyl, pyrrolidinyl, pyrazolyl, pyrazolidinyl, imidazolyl, imidazolinyl, imidazolidinyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, oxazolyl, oxazolidinyl, isoxazolyl, isoxazolidinyl, morpholinyl, thiazolyl, thiazolidinyl, isothiazolyl, quinuclidinyl, isothiazolidinyl, indolyl, quinolinyl, isoquinolinyl, benzimidazolyl, thiadiazoyl, benzopyranyl, benzothiazolyl, benzoxazolyl, furyl, tetrahydrofuryl, tetrahydropyranyl, thienyl, benzothienyl, thiamorpholinyl, thiamorpholinyl sulfoxide, thiamorpholinyl sulfone, and oxadiazolyl. Morpholino is the same as morpholinyl.

This Sheet Has Been Replaced

The pharmaceutically-acceptable salts of the compounds of Formula I (in the form of water- or oil-soluble or dispersible products) include the conventional non-toxic salts or the quaternary ammonium salts which are formed, e.g., from inorganic or organic acids or bases. Examples of such acid addition salts include acetate, adipate, alginate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, citrate, camphorate, camphorsulfonate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, fumarate, glucoheptanoate, glycerophosphate, hemisulfate, heptanoate, hexanoate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethanesulfonate, lactate, maleate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, oxalate, pamoate, pectinate, persulfate, 3-phenyl-propionate, picrate, pivalate, propionate, succinate, tartrate, thiocyanate, tosylate, and undecanoate. Base salts include ammonium salts, alkali metal salts such as sodium and potassium salts, alkaline earth metal salts such as calcium and magnesium salts, salts with organic bases such as dicyclohexylamine salts, N-methyl-D-glucamine, and salts with amino acids such as arginine, lysine, and so forth. Also, the basic nitrogen-containing groups may be quaternized with such agents as lower alkyl halides, such as methyl, ethyl, propyl, and butyl chloride, bromides and iodides; dialkyl sulfates like dimethyl, diethyl, dibutyl; and diamyl sulfates, long chain halides such as decyl, lauryl, myristyl and stearyl chlorides, bromides and iodides, aralkyl halides like benzyl and phenethyl bromides and others.

This Sheet Has Been Replaced



Schemes I-III for preparing the novel compounds of this invention are presented below. Tables I and II which follow the schemes illustrate the compounds that can be synthesized by Schemes I-III, but Schemes I-III are not limited by the compounds in the tables nor by any particular substituents employed in the schemes for illustrative purposes. The examples specifically illustrate the application of the following schemes to specific compounds.

Amide couplings used to form the compounds of this invention are typically performed by the carbodiimide method with reagents such as dicyclohexylcarbodiimide, or 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide. Other methods of forming the amide or peptide bond include, but are not limited to the synthetic routes via an acid chloride, azide, mixed anhydride or activated ester. Typically, solution phase amide coupling are performed, but solid-phase synthesis by classical Merrifield techniques may be employed instead. The addition and removal of one or more protecting groups is also typical practice.

Additional related information on synthetic background is contained in EPO 0337714.

One method for producing formula I compounds is provided by Scheme I. Dihydro-5(S)-(tert-butyldimethylsilyloxymethyl)-3(2H)-furanone (compound 1 below) is prepared by standard methods known in the art from commercially available dihydro-5(S)-(hydroxymethyl)-2(3H)-furanone. After alkylation of compound 1 to form compound 2, the protecting group of lactone 2 is removed with aqueous HF to afford compound 3.

This Sheet Has Been Replaced


The alcohol group of 3 is activated by conversion into a leaving group such as mesylate, tosylate or trifylate by treating the alcohol with a sulfonyl chloride or sulfonic anhydride, such as trifluoromethanesulfonic anhydride, in the presence of a hindered amine base such as triethylamine, diethyl isopropylamine or 2,6 lutidine, to afford a compound such as compound 4. The leaving group of compound 4 is displaced by an amine 5, such as N'-t-butyl-(4aS,8aS)-(decahydroisoquinoline)-3(S)-carboxamide, in a high boiling solvent such as DMF or xylene to produce a compound such as 6. A trifluoromethanesulfonyloxy group can be displaced by an amine at room temperature in a solvent such as isopropanol by treatment with N,N-diisopropylethylamine.

Compound 6 is hydrolyzed with aqueous lithium or sodium hydroxide and the resultant hydroxy acid 7 is converted into a protected hydroxy acid 8. The hydroxyl group is conveniently protected with a standard silyl protecting group such as t-butyldimethyl silyl or t-butyldiphenyl silyl.

The protected hydroxy-acid 8 is then coupled to the desired R^{12} amine to produce compound 9, and the silyl protecting group is removed with fluoride ion to arrive at compound 10.

This Sheet Has Been Replaced


SCHEME I




1 2 3 4 5 6 7 8

This Sheet Has Been Replaced

SCHEME I (CONT'D)



9



10

A second method for forming products of general formula I is shown in Scheme II. In Scheme II, alkylation of 11 is performed by a first step of deprotonation of 11 with n-butyllithium or lithium diisopropylamide (LDA) followed by a second step of adding an alkenyl halide (such as allyl bromide) to afford 12.

Dihydroxylation of the olefin of 12 with osmium tetroxide and N-methylmorpholine-N-oxide (NMO) produces a diasteriomeric mixture of diols, 13. Selective mesylation of the primary alcohol of 13 with methanesulfonyl chloride and either triethylamine or pyridine gives a mesylate 14.


This Sheet Has Been Replaced

Heating mesylate 14 with an amine in a refluxing alcoholic solvent such as methanol or isopropanol which contains an excess of potassium carbonate produces an amino alcohol such as compound 15. The diasteriomers can be separated at this step by standard techniques well known to those of skill in the art. Alternatively, the separation can be done after removal of the ketal.

Removal of the ketal in compound 15 is accomplished by treatment with acid in the presence of methanol, or by aqueous acid or by 1N HCl in THF, to form compound 16.

This Sheet Has Been Replaced

SCHEME II




This Sheet Has Been Replaced

A third method for forming products of general formula I is shown in Scheme III. Protection of the pyrrolidine -NH- group of compound 17 is carried out with BOC-anhydride and dimethylaminopyridine to give the protected compound 18. Alkylation of 18 is performed by a first step of deprotonation of 18 with a strong base such as lithium hexamethyldisilamide (LHMDS) or lithium diisopropylamide (LDA) followed by a second step of adding an alkyl halide (such as benzyl bromide) to afford compound 19.


The TBS protecting and BOC protecting group of 19 are removed by treatment with aqueous HF in acetonitrile to give alcohol 20. Mesylation of the primary alcohol of 20 with methanesulfonyl chloride and either triethylamine or pyridine gives mesylate 21 which is heated with an amine in a refluxing alcoholic solvent such as methanol or isopropanol which contains an excess of potassium carbonate to produce an amino pyrrolidinone such as compound 22. The pyrrolidine -NH- group of 22 is reprotected as a BOC group as before and the resultant compound 23 is hydrolized open with a base such as lithium or sodium hydroxide to afford the acid 24. Compound 24 is then coupled to an NH_2R^{12} amine in a standard manner and the BOC is removed with gaseous HCl or trifluoro-acetic acid to give the desired product, exemplified by compound 25.

This Sheet Has Been Replaced


SCHEME III




17 18




19




20



21




22




23

This Sheet Has Been Replaced

SCHEME III (CONT'D)




24



25


A compound of formula 26



26

wherein P is a nitrogen protecting group such as -BOC or -CBZ, is preferably prepared according to the method described in Scheme I, preferably employing the 5-trifluoromethanesulfonyloxymethyl analog of lactone 4 therein (see Example 15, Step 1).


Compounds of formula 27



27

This Sheet Has Been Replaced

can be obtained by a variety of routes from compound 28



28


which is obtained after removal of the nitrogen protecting group in 26 using methods well known in the art, e.g., catalytic hydrogenation to remove a CBZ group, or treatment with trimethylsilyltriflate and 2,6 lutidine at about 0°C in a solvent such as CH_2Cl_2 to remove a BOC group.






For example, the 4-position piperazinyl nitrogen of compound 28 can be alkylated with a compound of formula R^1-X in a solvent such as DMF in the presence of Et_3N at room temperature, wherein X is -Cl, Br or -I, or a sulfonamide group can be formed by treatment of 28 with a sulfonyl chloride compound of formula R^1SO_2Cl under similar conditions. Also, standard amide coupling techniques can be used to form an amide group at the piperazinyl 4-position. Techniques for these procedures are well known to those skilled in the art. The R^1 group of R^1-X or R^1SO_2Cl is defined above in the definition of compounds of formula I wherein R^1 is independent from and not joined to R^2, except that R^1 can not be hydrogen or a group with a free hydroxy substituent, such as -C_{1-4}alkyl substitued with hydroxy, with the further exception that R^1 can be aryl substituted with a hydroxy group.

The compounds of this invention are also illustrated by Tables I-IV, which follow.

This Sheet Has Been Replaced






TABLE I



R^3	X	D
$-CH_2-Ph$	-OH	
$-CH_2-Ph$	-OH	
$-CH_2-Ph$	-OH	
$-CH_2-Ph$	-OH	
$-CH_2-Ph$	-OH	






This Sheet Has Been Replaced

TABLE I. CONT'D

R^3	X	D
$-CH_2-Ph$	-OH	
$-CH_2-Ph$	-OH	
$-CH_2-Ph$	-OH	
$-CH_2-Ph$	-OH	
$-CH_2-Ph$	-OH	





This Sheet Has Been Replaced

TABLE I, CONT'D

R^3	X	D
$-CH_2-Ph$	-OH	
$-CH_2-Ph$	-OH	
$-CH_2-Ph$	-OH	
$-CH_2-Ph$	-OH	
$-CH_2-Ph$	-OH	











This Sheet Has Been Replaced

TABLE I. CONT'D

R^3	X	D
$-CH_2-Ph$	-OH	
$-CH_2-Ph$	$-NH_2$	
$-CH_2-Ph$	-OH	
$-CH_2-Ph$	-OH	















This Sheet Has Been Replaced

TABLE I. CONT'D

R^3	X	D
$-CH_2-Ph$	$-NH_2$	
$-CH_2-Ph$	$-OH$	
$-CH_2-Ph$	$-OH$	
$-CH_2-Ph$	$-OH$	
	$-OH$	
	$-OH$	
	$-OH$	











This Sheet Has Been Replaced

TABLE I. CONT'D

R^3	X	D
	$-NH_2$	
	$-NH_2$	
	-OH	
	-OH	
	$-NH_2$	
	-OH	
	-OH	















This Sheet Has Been Replaced

TABLE I. CONT'D

R^3	X	D
	$-NH_2$	
	-OH	
	-OH	
	-OH	
	$-NH_2$	
	-OH	

This Sheet Has Been Replaced







TABLE I, CONT'D

R^3	X	D
	-OH	
	$-NH_2$	
	-OH	
	-OH	
	$-NH_2$	
	$-NH_2$	
	-OH	

This Sheet Has Been Replaced









TABLE I. CONT'D

R^3	X	D
$-CH_2CH=CH-Ph$	-OH	
$-CH_2CH=CH-Ph$	-OH	
	-OH	
	-OH	

This Sheet Has Been Replaced








TABLE II



A	X	D
	-OH	
	-OH	
	$-NH_2$	















This Sheet Has Been Replaced

TABLE II Cont'd

A	X	D
	$-NH_2$	
	-OH	
	-OH	
	-OH	









This Sheet Has Been Replaced

TABLE II Cont'd

A	X	D
	-OH	
	$-NH_2$	
	-OH	
	-OH	
	-OH	
	-OH	
	-OH	







This Sheet Has Been Replaced

TABLE II CONT'D

A	X	D
	-OH	
	-OH	
	-OH	
	-OH	











This Sheet Has Been Replaced

TABLE II CONT'D

A	X	D
	-OH	
	-OH	
	-OH	





This Sheet Has Been Replaced

TABLE II, CONT'D

A	X	D
	-OH	
	-OH	
	-OH	
	-OH	
	-OH	


This Sheet Has Been Replaced










TABLE II CONT'D

A	X	D
	-OH	
	-OH	

This Sheet Has Been Replaced


TABLE III




A	A, con't
	$CH_3CH_2CH_2CH_2-$
CH_3-	$(CH_3)_2CH-$
	
	
	
	


This Sheet Has Been Replaced

TABLE IV




A





$(CH_3)_3C-O-$



$Ph-CH_2CH_2-$

This Sheet Has Been Replaced

The compounds of the present invention are useful in the inhibition of HIV protease the prevention or treatment of infection by the human immunodeficiency virus (HIV) and the treatment of consequent pathological conditions such as AIDS. Treating AIDS or preventing or treating infection by HIV is defined as including, but not limited to, treating a wide range of states of HIV infection: AIDS, ARC (AIDS related complex), both symptomatic and asymptomatic, and actual or potential exposure to HIV. For example, the compounds of this invention are useful in treating infection by HIV after suspected past exposure to HIV by, e.g., blood transfusion, organ transplant, exchange of body fluids, bites, accidental needle stick, or exposure to patient blood during surgery.

For these purposes, the compounds of the present invention may be administered orally, parenterally (including subcutaneous injections, intravenous, intramuscular, intrasternal injection or infusion techniques), by inhalation spray, or rectally, in dosage unit formulations containing conventional non-toxic pharmaceutically-acceptable carriers, adjuvants and vehicles.

Thus, in accordance with the present invention there is further provided a method of treating and a pharmaceutical composition for treating HIV infection and AIDS. The treatment involves administering to a patient in need of such treatment a pharmaceutical composition comprising a pharmaceutical carrier and a therapeutically effective amount of a compound of the present invention, or a pharmaceutically acceptable salt thereof.

This Sheet Has Been Replaced



These pharmaceutical compositions may be in the form of orally-administrable suspensions or tablets; nasal sprays; sterile injectable preparations, for example, as sterile injectable aqueous or oleagenous suspensions or suppositories.

When administered orally as a suspension, these compositions are prepared according to techniques well-known in the art of pharmaceutical formulation and may contain microcrystalline cellulose for imparting bulk, alginic acid or sodium alginate as a suspending agent, methylcellulose as a viscosity enhancer, and sweetners/flavoring agents known in the art. As immediate release tablets, these compositions may contain microcrystalline cellulose, dicalcium phosphate, starch, magnesium stearate and lactose and/or other excipients, binders, extenders, disintegrants, diluents and lubricants known in the art.

When administered by nasal aerosol or inhalation, these compositions are prepared according to techniques well-known in the art of pharmaceutical formulation and may be prepared as solutions in saline, employing benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability, fluorocarbons, and/or other solubilizing or dispersing agents known in the art.

The injectable solutions or suspensions may be formulated according to known art, using suitable non-toxic, parenterally-acceptable diluents or solvents, such as mannitol, 1,3-butanediol, water, Ringer's solution or isotonic sodium chloride solution, or suitable dispersing or wetting and suspending agents, such as sterile, bland, fixed oils, including

This Sheet Has Been Replaced



synthetic mono- or diglycerides, and fatty acids, including oleic acid.

When rectally administered in the form of suppositories, these compositions may be prepared by mixing the drug with a suitable non-irritating excipient, such as cocoa butter, synthetic glyceride esters or polyethylene glycols, which are solid at ordinary temperatures, but liquidify and/or dissolve in the rectal cavity to release the drug.

Dosage levels of the order of 0.02 to 5.0 or 10.0 grams-per-day are useful in the treatment or prevention of the above-indicated conditions, with oral doses two-to-five times higher. For example, infection by HIV is effectively treated by the administration of from 10 to 50 milligrams of the compound per kilogram of body weight from one to three times per day. It will be understood, however, that the specific dose level and frequency of dosage for any particular patient may be varied and will depend upon a variety of factors including the activity of the specific compound employed, the metabolic stability and length of action of that compound, the age, body weight, general health, sex, diet, mode and time of administration, rate of excretion, drug combination, the severity of the particular condition, and the host undergoing therapy.

The present invention is also directed to combinations of the HIV protease inhibitory compounds with one or more agents useful in the treatment of AIDS. For example, the compounds of this invention may be effectively administered, whether at periods of pre-exposure and/or post-exposure, in combination with effective amounts of the AIDS antivirals,

This Sheet Has Been Replaced



immunomodulators, anti-infectives, or vaccines known to those of ordinary skill in the art.

It will be understood that the scope of combinations of the compounds of this invention with AIDS antivirals, immunomodulators, anti-infectives or vaccines include in principle any combination with any pharmaceutical composition useful for the treatment of AIDS.

Assay for Inhibition of Microbial Expressed Viral Protease

Inhibition studies of the reaction of the protease expressed in Eschericia coli with a peptide substrate [Val-Ser-Gln-Asn-(betanapthyl)Ala-Pro-Ile-Val, 0.5 mg/mL at the time the reaction is initiated] were in 50 mM Na acetate, pH 5.5, at 30°C for 1 hour. Various concentrations of inhibitor in 1.0 ul DMSO were added to 25 ul of the peptide solution in water. The reaction is initiated by the addition of 15 ul of 0.33 nM protease (0.11 ng) in a solution of 0.133 M Na acetate pH 5.5 and 0.1% bovine serum albumin. The reaction was quenched with 160 ul of 5% phosphoric acid. Products of the reaction were separated by HPLC (VYDAC wide pore 5 cm C-18 reverse phase, acetonitrile gradient, 0.1% phosphoric acid). The extent of inhibition of the reaction was determined from the peak heights of the products. HPLC of the products, independently synthesized, proved quantitation standards and confirmation of the product composition. The products of synthesis in Examples 1-7 inclusive showed IC_{50} values in the range of 1-100 nM. Compounds A, B and J showed IC_{50} values of between about 0.3 and about 6 nM.

This Sheet Has Been Replaced



EXAMPLE 1

Preparation of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(1-(N'-(t-butyl)-4(S)-phenoxyprolineamide)yl)-pentaneamide


Step 1: Preparation of N-(2(R)-hydroxy-1(S)-indanyl)-3-phenylpropaneamide

To a cold (0°C) solution of methylene chloride (30 ml) containing 2(R)-hydroxy-1(S)-aminoindane (750 mg, 5.0 mmol) and triethylamine (606 mg, 6.0 mmol) was added a solution of hydrocinnamoyl chloride (843 mg, 5.0 mmol) in 5 ml of methylene chloride. After 2 hr the reaction was poured into a separatory funnel containing 50 ml of methylene chloride and washed with 10% citric acid solution (2 x 30 ml). The organic layer was dried, filtered and concentrated to afford a white solid.

Step 2: Preparation of N-(2(R)-hydroxy-1(S)-indan-N,O-isopropylidene-yl)-3-phenyl-propaneamide

The crude white solid from step 1 above was dissolved in 50 ml of methylene chloride and 5 ml of dimethoxypropane was added followed by the addition of 100 mg of p-toluenesulfonic acid. The reaction was stirred at room temperature for 18 hr and then poured into a separatory funnel and washed with saturated $NaHCO_3$ solution (2 x 30 ml). The organic layer was dried, filtered and concentrated to afford an oil which was chromatographed (SiO_2, 40% EtOAc/Hexane) to give an oil which eventually crystallized.

This Sheet Has Been Replaced



Step 3: Preparation of N-(2(R)-hydroxy-1(S)-indan-N,O-isopropylidene-yl)-2(S)-phenylmethyl-pent-4-eneamide

To a solution of N-(2(R)-hydroxy-1(S)-indan-N,O-isopropylidene-yl)-3-phenyl-propaneamide (1.03 gm, 2.9 mmol) in 20 ml of THF cooled to -78°C was added n-BuLi (2.5M, 1.40 ml, 3.5 mmol). After 20 min, allyl bromide (0.48 gm, 3.9 mmol) was added, the reaction was stirred at -78°C for 1 hr and then 10 ml of saturated NH_4Cl solution was added to quench the reaction. The reaction was diluted with 50 ml of water, extracted with ethyl acetate (2 x 50 ml), the organic phase was washed with saturated NaCl solution (50 ml), dried filtered and concentrated to afford the crude product. The crude product was purified on silica gel to afford the title compound.

Step 4: Preparation of N-(2(R)-hydroxy-1(S)-indan-N,O-isopropylidene-yl)-2(S)-phenylmethyl-(4(RS),5-dihydroxy)-pentaneamide

To 800 mg (2.2 mmol) of N-(2(R)-hydroxy-1(S)-indan-N,O-isopropylidene-yl)-2(S)-phenylmethyl-pent-4-eneamide dissolved in 40 ml of a 9:1 mixture of acetone/water was added 0.8 ml of a 60% solution of N-methylmorpholine-N-oxide in water followed by 4 ml of a 2.5% solution of osmium tetroxide in t-BuOH. After 18 hr, excess solid sodium bisulfate was added, the reaction was stirred for 2 hr and then filtered through a pad of celite. The filtrate was concentrated, diluted with 50 ml of water, extracted with methylene chloride (2 X 50 ml), the organic phase was dried, filtered and concentrated to give the product as a foam.

This Sheet Has Been Replaced



Step 5: Preparation of N-(2(R)-hydroxy-1(S)-indan-N,O-isopropylidene-yl)-2(S)-phenylmethyl-4(RS)-hydroxy-5-methanesulfonyloxy-pentaneamide

To 200 mg (0.527 mmol) of N-(2(R)-hydroxy-1(S)-indan-N,O-isopropylidene-yl)-2(S)-phenylmethyl-(4(RS),5-dihydroxy)-pentaneamide dissolved in 7 ml of methylene chloride at 0°C was added triethylamine (59 mg, 0.58 mmol), followed by methanesulfonyl chloride (66 mg, 0.579 mmol). After 4 hr the reaction was worked up by washing with 10% citric acid solution (2 X 50 ml) and the organic phase was dried, filtered and concentrated to afford the monomesylate as a mixture of alcohols.

Step 6: Preparation of N'-t-butyl-N-Boc-4(R)-hydroxy-L-prolineamide

To a solution of N-Boc-4(R)-hydroxyproline (2.00 g) in DMF (20 mL) cooled to 0°C was added EDC (1.987 g), HOBt (1.401 g), tert butyl amine (1.09 mL) and triethylamine (2.41 mL). After 18 h the reaction mixture was diluted with ethyl acetate (150 mL) and washed with 10% HCl, saturated $NaHCO_3$, water and brine. The solution was then dried over $MgSO_4$ and concentrated to afford a white solid.

Step 7: Preparation of N'-t-butyl-N-Boc-4(S)-phenoxy-L-prolineamide

To a solution of N'-t-butyl-N-Boc-4(R)-hydroxy-L-prolineamide (0.6 g) in THF (5 mL) was added phenol (0.295 g), triphenylphosphine (0.824 g) and then diethylazo-dicarboxylate (0.495 mL) dropwise. The reaction mixture stirred for 24 h at ambient

This Sheet Has Been Replaced

temperature and was diluted with ethyl acetate (200 mL) and washed with saturated $NaHCO_3$, water, brine and dried over $MgSO_4$. Concentration in vacuo afforded a yellow oil which was purified by flash chromatography (elution hexane: EtOAc 1:1, 30 mm column).

Step 8: Preparation of N-t-butyl-4(S)-phenoxy-L-prolineamide trifluoroacetic acid salt

To a solution of N'-t-butyl-N-Boc-4(S)-phenoxy-L-prolineamide (0.596 g) in methylene chloride (4 mL) at 0°C was added trifluoroacetic acid (2 mL). After 30 min the reaction was warmed to room temperature and stirred for two hours. The solvent was removed in vacuo and a slightly yellow oil was obtained.

Step 9: Preparation of N-(2(R)-hydroxy-1(S)-indan-N,O-isopropylidene-yl)-2-(R)-phenylmethyl-4-(S)-hydroxy-5-(1-(N'-(t-butyl)-4(S)-phenoxy-prolineamide)yl)-pentaneamide

To a solution of N-t-butyl-4(S)-phenoxy-L-prolineamide trifloroacetic acid salt (0.36 g) and N-(2(R)-hydroxy-1(S)-indan-N,O-isopropylidene-yl)-2(S)-phenylmethyl-4(RS)-hydroxy-5-methanesulfonyloxy-pentaneamide (0.226 g) in 3 mL of isopropanol was added potassium carbonate (0.441 g) and the reaction was warmed to 80°C. After 18 h the reaction was cooled to room temperature, filtered through celite which was washed with further portions of EtOAc. The filtrate was concentrated, the residue was dissolved in EtOAc (100mL) and washed with water, brine and dried over $MgSO_4$. The solvent was removed in vacuo and the resulting oil was purified by flash chromatography to

This Sheet Has Been Replaced

afford the product as a mixture of diastereomers.

Step 10: Prep of N-(2(R)-hydroxy-1(S)-indanyl)-2-(R)-phenylmethyl-4-(S)-hydroxy-5-(1-(N'-t-butyl-4(S)-phenoxyprolineamid)yl)-pentaneamide

To a solution of N-(2(R)-hydroxy-1(S)-indan-N,O-isopropylidene-yl)-2-(R)-phenylmethyl-4-(S)-hydroxy-5-(1-(N'-(t-butyl)-4(S)-phenoxyprolineamide)-yl)-pentaneamide (0.13 g) in MeOH (5 mL) was added camphorsulfonic acid (CSA) (0.070 g) at ambient temperature. After 5 hours more CSA (0.025 g) was added and the reaction was stirred for total of 18 hours. The reaction was quenched with saturated $NaHCO_3$ (5 mL) and the solvent was removed to a volume of 4 mL. The aqueous layer was thoroughly extracted with EtOAc and the organic layer was washed with water, brine and dried. After removal of the solvent in vacuo the resulting oil was purified via flash chromatography to provide the title compound as a white foam. The foam was dissolved in EtOAc : hexanes and the mother liquor was decanted away from the oil. The oil was then dried in a high vacuum desiccator to afford a white foam.

EXAMPLE 2

Preparation of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(1-(N'-t-butyl-4(S)-2-naphthyloxy-prolineamid)yl)-pentaneamide

Step 1: Preparation of N-t-butyl-4(S)-2-naphthyloxy-L-prolineamide trifluoroacetic acid salt

Following substantially the same procedure

This Sheet Has Been Replaced

for synthesizing N-t-butyl-4(S)-phenoxy-L-proline-amide trifluoroacetic acid salt as outlined in Example 1, Steps 6 through 8, but substituting 2-naphthol for the phenol used therein, the 2-naphthyloxy proline amide was produced.

Step 2: Preparation of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(1-(N'-t-butyl-4(S)-2-naphthyloxy-prolineamid)yl)-pentaneamide

The title compound was produced by following substantially the same procedure outlined in Example 1, Steps 9 and 10, but substituting N-t-butyl-4(S)-2-naphthyloxy-L-prolineamide trifluoroacetic acid salt for the N-t-butyl-4(S)- phenoxy-L-prolineamide trifloroacetic acid salt used in step 9 therein.

EXAMPLE 3

Preparation of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(1-(N'-t-butyl-4(S)-1-naphthyloxy-prolineamid)yl)-pentaneamide

Step 1: Preparation of N-t-butyl-4(S)-1-naphthyloxy-L-prolineamide trifluoroacetic acid salt

Following substantially the same procedure for synthesizing N-t-butyl-4(S)-phenoxy-L-proline-amide trifluoroacetic acid salt as outlined in Example 1, Steps 6 through 8, but substituting 1-naphthol for the phenol used therein, the 1-naphthyloxy proline amide was produced.

This Sheet Has Been Replaced

Step 2: Preparation of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(1-(N'-t-butyl-4(S)-2-naphthyloxy-prolineamid)yl)-pentaneamide

The title compound.was produced by following the procedure outlined in Example 1, Steps 9 and 10, but substituting N-t-butyl-4(S)-1-naphthyloxy-L-prolineamide trifluoroacetic acid salt for the N-t-butyl-4(S)-phenoxy-L-prolineamide trifluoroacetic acid salt used in Step 9.

EXAMPLE 4

Preparation of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(2-(3(S)-N'-(t-butyl-carboxamido)-(4aS,8aS)-decahydroisoquinoline)yl)-pentaneamide

Step 1: Preparation of dihydro-5(S)-((t-butyldi-phenylsilyl)oxymethyl)-3(R)phenylmethyl-3(2H)-furanone

A solution of lithium diisopropylamide (LDA) was generated by the addition 1.55 ml of n-BuLi (2.5M in hexane) to 0.55 ml (3.9 mmol) of diisopropylamine in 10 ml of THF at -78°C. After 30 minutes a solution of dihydro-5-(S)-((t-butyldiphenylsilyl)-oxymethyl)-3(2H)-furanone (1.38 g, 3.89 mmol) in 5 ml of THF was added. After an additional 30 minutes of stirring, benzyl bromide (0.68 g, 3.9 mmol) was added and stirring was continued for 3 h after which time the reaction was quenched with the addition of a 10% aqueous citric acid solution. The solution was

This Sheet Has Been Replaced

extracted with ethyl acetate (2 x 50 ml) which was backwashed with brine, dried, filtered and concentrated to afford an oil. The product was purified by chromatography (SiO_2, 20% EtOAc/Hexane) to afford the title compound.

Step 2: Preparation of dihydro-5(S)-(hydroxymethyl)-3(R)-phenylmethyl-3(2H)-furanone

To 5.26 g of dihydro-5(S)-((t-butyldiphenylsilyl)oxymethyl)-3(R)phenylmethyl-3(2H)-furanone in 40 ml of acetonitrile was added 1.34 ml of a 49% aqueous HF solution. After 18 hr at room temperature the reaction was concentrated to dryness and the residue was partitioned between water (50 ml) and ethyl acetate (50 ml). The organic layer was washed with brine, dried filtered and concentrated to afford the product as a tan solid (mp 69-72°C).

Step 3: Preparation of dihydro-5(S)-((methanesulfonyl)oxymethyl)-3(R)phenylmethyl-3(2H)-furanone

To a solution of 2.93 g (14 mmol) of dihydro-5(S)-(hydroxymethyl)-3(R)-phenylmethyl-3(2H)-furanone in methylene chloride cooled to 0°C was added triethylamine (1.98ml, 15.6 mmol) followed by the addition of methanesulfonyl chloride (1.20 ml, 15.6 mmol). After 1 hour at 0°C, the reaction was poured into 10% aqueous citric acid solution, washed with ethyl acetate (2 x 100 ml) which was backwashed with water (100 ml), brine (100 ml), dried, filtered and concentrated to give the product as a waxy brown solid.

This Sheet Has Been Replaced



Step 4: Preparation of dihydro-5(S)-(2-(3(S)-N-(t-butylcarboxamido)-(4aS, 8aS)-(decahydroisoquinoline)yl)methyl)-3(R)-phenylmethyl-3(2H)-furanone

To 70 mg of dihydro-5(S)-((methanesulfonyl)-oxymethyl)-3(R)phenylmethyl-3(2H)-furanone (0.25 mmol) in 10 ml of xylene containing 100 mg of potassium carbonate was added 65 mg (0.27 mmol) of N-t-butyl-(4aS,8aS)-(decahydroisoquinoline)-3(S)-carboxamide and the reaction was heated to 140°C. After 6 hours, the reaction was cooled, poured into 30 ml of water which was washed with ethyl acetate (2 x 30 ml). The organic phase was dried, filtered and concentrated to afford a residue which was chromatographed (50/50 EtOAc/Hexane) to give the product.

Step 5: Preparation of 2(R)-phenylmethyl-4(S)-(t-butyldimethylsilyloxy)-5-(2-(3(S)-N-(t-butylcarboxamido)-(4aS,8aS)-decahydroisoquinoline)yl)-pentanoic acid

To 130 mg (0.305 mmol) of dihydro-5(S)-(2-(3(S)-N-(t-butylcarboxamido)-(4aS, 8aS)-(decahydroisoquinoline)yl)methyl)-3(R)-phenylmethyl-3-(2H)furanone in 2 ml of DME was added 1 ml lithium hydroxide solution. After 4 hours at room temperature, the reaction was concentrated to dryness and azeotroped with toluene (3X) to remove excess water. The residue was dissolved in 5 ml of DMF and 414 mg (6.10 mmol) of imidazole and 465 mg (3.05 mmol) of t-butyldimethylsilyl chloride was added. After two days at room

This Sheet Has Been Replaced

temperature, 1 ml of methanol was added to the reaction and after 1 hour the solution was evaporated to dryness. The residue was partitioned between saturated NH_4Cl solution (aq) and washed with ethyl acetate which was dried, filtered and concentrated to give an oil which was a mixture of product and the furanone starting material. This material was carried on crude into the next reaction.

Step 6: Preparation of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-(t-butyldimethyl-silyloxy-5-(2-(3(S)-N'-(t-butylcarboxamido)-(4aS,8aS)-decahydroisoquinoline)yl)-pentaneamide

The crude product of step 5, above, was dissolved in 3 ml of DMF along with 47 mg (0.246 mmol) of EDC, 33 mg (0.246 mmol) of HOBT and 37 mg of 2(R)-hydroxy-1(S)-aminoindane. The pH of the solution was adjusted to 8.5-9.0 with triethylamine and after 18 hours it was worked up by concentrating to dryness, dissolving the residue in 10% aq. citric acid solution and washing the aqueous layer with ethyl acetate. The organic layer was dried, filtered and concentrated and the resultant oil was chromatographed (SiO_2, 30% EtOAc/Hexane) to yield the title compound.

Step 7: Preparation of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(2-(3(S)-N'-(t-butylcarboxamido)-(4aS,8aS)-decahydro-isoquinoline)yl)-pentaneamide

The product from step 6, above, was dissolved

This Sheet Has Been Replaced

in 1 ml of THF and 1 ml of a 1M solution of tetrabutylammonium fluoride in THF was added. After 18 hr at room temperature the reaction was diluted with 20 ml of saturated $NaHCO_3$ solution (aq) and the product was extracted into ethyl acetate which was dried, filtered and concentrated to give a foam. The resultant material was chromatographed on a prep plate (0.5 mm, 5% MeOH/$CHCl_3$) and the title product isolated in the usual manner as a solid with mp 105-107°C.

EXAMPLE 5

Preparation of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-amino-5-(2-(3(S)-N'-(t-butylcarboxamido)-(4aS,8aS)-decahydroisoquinoline)yl)-pentaneamide

Step 1: Preparation of 5(S)-((t-butyl-dimethyl-silyloxy)methyl)-3(R)-phenylmethyl-N-BOC-2-pyrrolidinone

A solution of 5(S)-((t-butyl-dimethyl-silyloxy)methyl)-N-BOC-2-pyrrolidinone (400 mg, 1.26 mmol) in 2 ml of THF was added to a precooled (-78°C) 1M solution of lithium hexamethyldisilazide (1.3 ml) in 5 ml of THF. After 45 min, 0.15 ml of benzyl bromide (1.3 mmol) was added and the stirring was continued. After 5 h the reaction was worked up by pouring into a separatory funnel containing 30 ml of an aqueous 10% citric solution. The aqueous layer was extracted (2 x 30 ml EtOAc) which was backwashed with brine (50 ml) dried, filtered and concentrated to an oil. The residue was chromatographed (SiO_2, 20% EtOAc/Hexane) to afford the product as an oil.

This Sheet Has Been Replaced

Step 2: Preparation of 5(S)-hydroxymethyl-3(R)-phenylmethyl-2-pyrrolidinone

To 130 mg (0.34 mmol) of 5(S)-((t-butyldimethylsilyloxy)methyl)-3(R)-phenylmethyl-N-BOC-2-pyrrolidinone in 5 ml of acetonitrile was added 0.1 ml of a solution of 48% HF in water. After 3 hr at room temperature the reaction was concentrated to dryness and diluted with 30 ml of an aqueous 10% $NaHCO_3$ solution. This was extracted with EtOAc (2 X 30 ml), dried filtered and concentrated to afford the crude product.

Step 3: Preparation of 5(S)-(methanesulfonyloxy)-methyl-3(R)-phenylmethyl-2-pyrrolidinone

To a solution of the crude product from Step 2, in 5 ml of methylene chloride cooled to 0°C was added triethylamine (42 mg, 0.41 mmol) and methanesulfonyl chloride (47 mg, 0.41 mmol). The reaction was slowly allowed to warm to room temperature and was stirred for 18 hr after which time it was diluted with 30 ml of methylene chloride, washed with 30 ml of 10% citric acid solution, dried filtered and concentrated to afford the product as an oil.

Step 4: Preparation of 5(S)-(2-(3(S)-N-(t-butylcarboxamido)-(4aS,8aS)-(decahydroisoquinoline)-yl)-methyl)-3(R)-phenylmethyl-2-pyrrolidinone

To a solution of 380 mg (1.34 mmol) of 5(S)-(methanesulfonyloxy)methyl-3(R)-phenylmethyl-2-pyrrolidinone in 20 ml of isopropanol was added 350 mg of potassium carbonate and 360 mg of N-t-butyl-(4aS,8aS)-(decahydroisoquinoline)-3(S)-carboxamide and the reaction was heated to 85°C. After 18 hr the

This Sheet Has Been Replaced

cooled reaction was filtered through celite, evaporated to dryness and the residue was dissolved in water which was extracted with EtOAc (2 X 50 ml). The organics were dried, filtered and concentrated, and the residue was chromatographed (SiO_2, 50/50 EtOAc/Hexane) to afford the product as an oil.

Step 5: Preparation of 5(S)-(2-(3(S)-N'-(t-butylcarboxamido)-(4aS,8aS)-(decahydroisoquinoline)-yl)-methyl)-3(R)-phenylmethyl-N-BOC-2-pyrrolidinone

To a solution of the product from step 4, above, (260 mg, 0.611 mmol) in 10 ml of methylene chloride was added dimethylaminopyridine (74 mg, 0.6 mmol) and 133 mg (0.61 mmol) of BOC-anhydride. After 18 hr at room temperature the reaction was worked up by diluting with 30 ml of methylene chloride and the organics washed with 30 ml of 10% citric acid solution, brine (30 ml) dried, filtered and concentrated to afford an oil. Chromatography (SiO_2, 40% EtOAc/Hexane) gave the title compound.

Step 6: Preparation of 5-(2-(3(S)-N'-(t-butylcarboxamido)-(4aS,8aS)-decahydroisoquinoline)-yl)-4(S)-[(1',1')-(dimethylethoxycarbonyl)-amino]-2(R)-phenylmethyl-pentanoic acid

To a solution of the product of step 5, above, (260 mg, 0.495 mmol) dissolved in 3 ml of dimethoxyethane was added 1.5 ml of a 1M solution of aqueous lithium hydroxide (1.5 mmol). The reaction was worked up after 2 hr by concentrating to dryness, dissolving the residue in saturated aqueous ammonium chloride solution and the aqueous phase was washed with

This Sheet Has Been Replaced

ethyl acetate (2 x 50 ml) which was dried, filtered and concentrated to afford the crude acid.

Step 7: Preparation of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-[(1',1')-(dimethyl-ethoxycarbonyl)amino]-5-(2-(3(S)-N'-(t-butyl-carboxamido)-(4aS,8aS)-decahydroisoquinol-ine)yl)-pentaneamide

To a solution of the product of step 6, above, (260 mg, 0.49 mmol) in methylene chloride was added EDC (94 mg, 0.49 mmol), HOBT (66 mg, 0.49 mmol), 2(R)-hydroxy-1(S)-aminoindane (73 mg, 0.49 mmol) and the pH of the reaction was adjusted to 8.5-9.0 using triethylamine. After 5 hr at room temperature the reaction was worked up by diluting with 50 ml of methylene chloride and washing the organics with saturated aqueous ammonium chloride solution. The organic phase was dried, filtered and concentrated and the residue was chromatographed to afford the title compound as a foam.

Step 8: Preparation of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(2-(3(S)-N'-(t-butylcarboxamido)-(4aS,8aS)-decahydro-isoquinoline)yl)-pentaneamide

To a solution of the product of step 7, above, (180 mg, 0.28 mmol) in 5 ml of methylene chloride cooled to 0°C was added 1 ml of trifluoro-acetic acid. After 4 hr the reaction was worked up by concentrating to dryness and the residue was dissolved in 50 ml of methylene chloride and washed with 10% aqueous $NaHCO_3$ solution. The organic layer was dried, filtered and concentrated to give the product as a

This Sheet Has Been Replaced



solid which was chromatographed (SiO_2, 7% MeOH/CH_2Cl_2) to afford the title compound, mp = 92-95°C.

EXAMPLE 6

Preparation of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(1-(4-carbobenzyloxy-2-(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide

Employing substantially the same procedure used in Example 1, but substituting N-t-butyl-4-CBZ-piperazine-2(S)-carboxamide for N-t-butyl-4(S)-phenoxy-L-prolineamide used in step 9 therein, the title compound was obtained.

EXAMPLE 7

Preparation of N''-(N-(2-pyridyl)-valyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(2-(3(S)-(N'-t-butylcarboxamido)-(4aS,8aS)-decahydroisoquinoline)yl)pentaneamide

Employing substantially the same procedure used in Example 4, but substituting N-2-pyridyl-valine for the 2(R)-hydroxy-1(S)aminoindane used in step 6 therein, the title compound was obtained.

EXAMPLE 8

Preparation of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(2(S)-(N'-t-butyl-3-phenyl-propionamide)amino)-pentaneamide

Employing substantially the same procedure used in Example 1, but substituting N-t-butyl-phenylalanine amide for the N'-t-butyl-4(S)-phenoxy-

This Sheet Has Been Replaced

L-prolineamide used in step 9 therein, the title compound is obtained.

EXAMPLE 9

Preparation of N-(4(S)-3,4-dihydro-1H-2,2-dioxobenzothiopyranyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(2-(3(S)-N'-(t-butylcarboxamido)-(4aS,8aS)-decahydroisoquinoline)yl)-pentaneamide

Step 1: Preparation of N-(4(S)-3,4-dihydro-1H-benzothiopyranyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(2-(3(S)-t-butylcarboxamido)-(4aS,8aS)-decahydroisoquinoline)yl)-pentaneamide

Employing substantially the same procedure used in Example 4 but substituting 4(S)-amino-3,4-dihydro-1H-benzothiopyran for the 2(R)-hydroxy-1(S)-aminoindane used in step 6 therein, the title compound is obtained.

Step 2: Preparation of N-(4(S)-3,4-dihydro-1H-2,2-dioxobenzothiopyranyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(2-(3(S)-t-butylcarboxamido)-(4aS,8aS)-decahydroisoquinoline)yl)-pentaneamide

The compound from step 1 above is dissolved in a 1:1 mixture of methanol and water. To this is added 10 eq. of OXONE and the reaction is stirred at room temperature. When the reaction is complete, it is concentrated to dryness, water is added and extracted with ethyl acetate which is dried, filtered and concentrated to give the title compound.

This Sheet Has Been Replaced

EXAMPLE 10

Preparation of N-(4(S)-3,4-dihydro-1H-2,2-dioxobenzothiopyranyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(1-(4-carbobenzyloxy-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide

Step 1: Preparation of dihydro-5(S)-(1-(4-carbobenzyloxy-2(S)-N'-(t-butylcarboxamido)-piperazinyl)methyl)-3(R)-phenylmethyl-3(2H)-furanone

Employing substantially the same procedure used in Example 4, step 4 but substituting 4-carbobenzyloxy-2(S)-N'-(t-butylcarboxamido)-piperazine for the N'-t-butyl-(4aS,8aS)-(decahydroisoquinoline)-3(S)-carboxamide used therein, the title compound is produced.

Step 2: Preparation of 2(R)-phenylmethyl-4(S)-(t-butyldimethylsilyloxy)-5-(1-(4-carbobenzyloxy-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentanoic acid

Employing substantially the same procedure used in Example 4, step 5 but substituting dihydro-5(S)-(1-(4-carbobenzyloxy-2(S)-N'-(t-butylcarboxamido)-piperazinyl)methyl)-3(R)-phenylmethyl-3(2H)-furanone for the dihydro-5(S)-(2-(3(S)-N'-(t-butylcarboxamido)-(4aS,8aS)-(decahydroisoquinoline)yl)-methyl)-3(R)-phenylmethyl-3(2H) furanone used therein, the title compound is produced.

This Sheet Has Been Replaced

Step 3: Preparation of N-(4(S)-3,4-dihydro-1H-benzothiopyranyl)-2(R)-phenylmethyl-4(S)-(t-butyldimethylsilyloxy)-5-(1-(4-carbobenzyloxy-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide

The crude 2(R)-phenylmethyl-4(S)-(t-butyldimethylsilyloxy)-5-(1-(4-carbobenzyloxy-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentanoic acid is dissolved in 3ml of DMF along with 1 eq of EDC, 1 eq of HOBT and 1 eq of 4(S)-amino-3,4-dihydro-1H- benzothiopyran. The pH of the solution is adjusted to 8.5-9.0 with triethylamine and after 18 hours it is worked up by concentrating to dryness, dissolving the residue in 10% aq citric acid solution and washing the aqueous layer with ethyl acetate. The organic layer is dried, filtered and concentrated and the resultant residue is chromatographed to yield the title product.

Step 4: Preparation of N-(4(S)-3,4-dihydro-1H-benzothiopyranyl)-2(R)-phenylmethyl-4(S)-hydroxy)-5-(1-(4-carbobenzyloxy-2(S)-(t-butylcarboxamido)-piperazinyl))-pentaneamide

The product from step 3 above is dissolved in 1 ml of THF and 1 ml of a 1M solution of tetra-butylammonium fluoride in THF is added. After 18 hr at room temperature the reaction is diluted with 20 ml of saturated $NaHCO_3$ solution (aq) and the product is extracted into ethyl acetate which is dried, filtered and concentrated to give a residue. The residue is chromatographed to afford the product.

This Sheet Has Been Replaced



Step 5: Preparation of N-(4(S)-3,4-dihydro-1H-2,2-dioxobenzothiopyranyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(1-(4-carbobenzyloxy-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide

The compound from step 4 above is dissolved in a 1:1 mixture of methanol and water. To this is added 10 eq of OXONE and the reaction is stirred at room temperature. When the reaction is complete, it is concentrated to dryness, water is added and extracted with ethyl acetate which is dried, filtered and concentrated to give the title compound.

EXAMPLE 11

Preparation of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-((2-hydroxy)ethoxy)phenyl)methyl)-4(S)-hydroxy-5-(2-(3-(S)-N'-(t-butylcarboxamido)-(4aS,8aS)-decahydroisoquinoline)yl)pentaneamide

Step 1: Preparation of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-(2-allyloxy)phenyl)methyl)-4(S)-hydroxy-5-(2-(3(S)-t-butylcarboxamido)-(4aS,8aS)-decahydroisoquinoline)yl)-pentaneamide

To a solution of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-hydroxyphenyl)methyl)-4(S)-hydroxy-5-(2-(3(S)-t-butylcarboxamido)-(4aS,8aS)-decahydroisoquinoline)yl)-pentaneamide in dioxane is added 6 eq of allyl bromide and 6 eq of cesium carbonate. The reaction is heated to 90°C. When the reaction is complete, the precipitate is filtered off, the dioxane is concentrated to dryness and the residue is diluted with

This Sheet Has Been Replaced

water which is washed with ethyl acetate. The organic phase is dried, filtered and concentrated to afford the product.

Step 2: Preparation of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-((2-hydroxy)ethoxy)phenyl)methyl)-4(S)-hydroxy-5-(2-(3(S)-N'-(t-butylcarboxamido)-(4aS,8aS)-decahydroisoquinoline)yl)-pentaneamide

The product from step 1 above is dissolved in methanol, 1 eq of p-toluenesulfonic acid is added and the reaction is cooled to -78°C. Excess ozone is bubbled through the reaction until a blue color persists. The flask is purged with nitrogen to remove any ozone and excess sodium borohydride solution is added. The reaction is warmed to room temperature and then saturated $NaHCO_3$ solution is added. The methanol is concentrated off on the rotoevaporater and the aqueous residue is washed with ethyl acetate which is dried, filtered and concentrated to afford the title compound.

EXAMPLE 12

Preparation of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-((2-hydroxy)ethoxy)phenyl)methyl)-4(S)-hydroxy-5-(1-(4-carbobenzyloxy-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide

Employing substantially the same prodecure used in Example 11 but substituting N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-hydroxyphenyl)methyl)-4(S)-hydroxy-5-(1-(4-carbobenzyloxy-2(S)-(t-butylcarbox-

This Sheet Has Been Replaced

amido)-piperazinyl)-pentaneamide for the N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-hydroxyphenyl)methyl)-4(S)-hydroxy-5-(2-(3(S)-t-butylcarboxamido)-(4aS,8aS)-decahydroisoquinoline)yl)-pentaneamide used therein, the title compound is obtained.

EXAMPLE 13

Preparation of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-(2-(4-morpholinyl)ethoxy)phenyl)methyl)-4(S)-hydroxy-5-(2-(3(S)-N'-(t-butylcarboxamido)-(4aS,8aS)-decahydroisoquinoline)yl)-pentaneamide

To a solution of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-hydroxyphenyl)methyl)-4(S)-hydroxy-5-(2-(3(S)-N'-(t-butylcarboxamido)-(4aS,8aS)-decahydroisoquinoline)yl)-pentaneamide in dioxane is added 6 eq of chloroethyl morpholine and 6 eq of cesium carbonate. The reaction is heated to 90°C. When the reaction is complete, the precipitate is filtered off, the dioxane is concentrated to dryness and the residue is diluted with water which is washed with ethyl acetate. The organic phase is dried, filtered and concentrated to afford the title compound.

EXAMPLE 14

Preparation of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-(2-(4-morpholinyl)ethoxy)phenyl)methyl)-4(S)-hydroxy-5-(1-(4-carbobenzyloxy-2(S)-N'-(t-butyl-carboxamido)-piperazinyl))-pentaneamide

To a solution of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-hydroxyphenyl)methyl)-4(S)-hydroxy-5-(1-(4-carbobenzyloxy-2(S)-(t-butylcarboxamido)-

This Sheet Has Been Replaced

piperazinyl)-pentaneamide in dioxane is added 6 eq of chloroethyl morpholine and 6 eq of cesium carbonate. The reaction is heated to 90°C. When the reaction is complete, the precipitate is filtered off, the dioxane is concentrated to dryness and the residue is diluted with water which is washed with ethyl acetate. The organic phase is dried, filtered and concentrated to afford the title compound.

EXAMPLE 15

Preparation of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(1-(4-(3-pyridylmethyl)-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide

Step 1: Preparation of dihydro-5(S)-((trifluoromethanesulfonyl)oxymethyl)-3(R)-phenylmethyl-3(2H)-furanone

To a solution of 18.4g (89.2 mmol) of dihydro-5(S)-(hydroxymethyl)-3(R)-phenylmethyl-3(2H)-furanone in 350 mL of methylene chloride cooled to 0°C was added 13.51 mL 2,6-lutidine (115.98 mmol) followed by a dropwise addition of 16.51 mL of trifluoromethanesulfonic anhydride (98.1 mmol). After 1.5 hours at 0°C, the reaction was poured into a mixture of 300 mL ice/brine and stirred for 0.5 hours. The aqueous layer was then extracted with methylene chloride (3 x 150 mL), the organic layers were washed with 10% HCl (2 x 75 mL), saturated $NaHCO_3$ (100mL), water (100mL), dried over $MgSO_4$, filtered and concentrated to give a solid residue. Purification via flash column chromatography (120 x 150 mm column, gradient elution of hexanes:EtOAc, 4:1 to 3:1) afforded the title product; mp 53-54°C.

This Sheet Has Been Replaced

Step 2: Preparation of 4-(1,1-dimethylethyl)-1-(phenylmethyl)-1,2(S),4-piperazinetricarboxylate

The title compound was prepared following the procedure of Bigge, C.F.; Hays, S.J.; Novak, P.M.; Drummond, J.T.; Johnson, G.; Bobovski, T.P. Tetrahedron Lett. 1989, 30, 5193; starting with 2(S)-piperazinecarboxylic acid. (see Felder, E.; Maffei, S.; Pietra, S.; Pitre, D.; Helv. Chim. Acta 1960, 117, 888.

Step 3: Preparation of N-t-butyl-4-(1,1-dimethylethoxycarbonylamino)-1-(phenylmethylcarbonylamino)piperazine-2(S)-carboxamide

To 9.90g (27.16 mmol) of 4-(1,1-dimethylethyl)-1-(phenylmethyl)-1,2(S),4-piperazinetricarboxylate dissolved in 75 mL of DMF and cooled to 0°C was added 5.73g (29.88 mmol) of EDC, 4.03g (29.88 mmol) of HOBt, 3.14 mL (29.88 mmol) of t-butylamine, and finally 4.16 mL (29.88 mmol) of triethylamine. The reaction mixture was stirred for 18 hours and the reaction volume was concentrated by half. The mixture was then diluted with 600 mL of EtOAc and washed with 10% HCl (2 x 75 mL), saturated $NaHCO_3$ (1 x 75 mL), water (3 x 75 mL) and brine (1 x 50 mL), dried over $MgSO_4$ and concentrated to a solid. This solid was triturated with EtOAc: hexane (1:2) and filtered to provide the title product as a white solid;
mp 134-135°C.

This Sheet Has Been Replaced

Step 4: Preparation of N-t-butyl-4-(1,1-dimethylethoxycarbonylamino)piperazine-2(S)-carboxamide

To 1.20g (2.86 mmol) of N-t-butyl-4-(1,1-dimethylethoxycarbonylamino)-1-(phenylmethylcarbonylamino)piperazine-2(S)-carboxamide and 1.1g (0.086 mmol) of 10% Pd/C was added 15 mL of methanol. The vessel was charged with hydrogen and the reaction stirred for 2 hours, filtered through celite and washed with ethanol. The solvents were removed in vacuo to provide the title product as a foam.
^{1}H NMR (300 MHz, $CDCl_3$) δ 6.65 (br, 1H), 4.10 (m, 1H), 3.81 (br, 1H), 3.21 (dd, J=18 and 7 Hz, 1H), 3.02-2.70 (m, 4H), 2.10-2.0 (br, 1H), 1.50 (s, 9H), 1.41(s, 9H).

Step 5: Preparation of dihydro-5(S)-(4-(1,1-dimethylethoxycarbonylamino))-2(S)-N-(t-butylcarboxamido)-piperazinyl)methyl)-3(R)-phenylmethyl-3(2H)-furanone

To a solution of 22.40g (0.0662 mol) dihydro-5(S)-((trifluoromethanesulfonyl)oxymethyl)-3(R)-phenylmethyl-3(2H)-furanone (prep in step 1) and 18.0g (0.063mol) of n-t-butyl-4-(1,1-dimethylethoxy-carbonylamino)piperazine-2(S)-carboxamide dissolved in 180 mL of isopropanol was added 11.53 mL (0.0662 mol) of N,N-diisopropylethylamine. After 2.5 hours another 1.2g of dihydro-5(S)-((trifluoromethanesulfonyl)oxy-methyl)-3(R)-phenylmethyl-3(2H)-furanone was added. The reaction was complete by thin layer chromatography (tlc) after 3.5 hours and was concentrated to a thick oil. Trituration with EtOAc:hexanes (1:2, 200mL) provided a white solid which was filtered and discarded. The oil was purified by flash column

This Sheet Has Been Replaced

chromatography (120 x 150 mm column, EtOAc:hexanes gradient elution 1:1, 2:1, 3:1 to all EtOAc) to afford the title compound.

^{1}H NMR (400 MHz, $CDCl_3$) δ 7.34-7.17 (m, 5H), 6.31 (br s, 1H), 4.38 (br m, 1H), 3.96-3.92 (m, 1H), 3.79 (br m, 1H), 3.16 (dd, J=13.6 and 4.4 Hz, 1H), 3.08-2.99 (m, 3H), 2.90-2.82 (m, 1H), 2.80 (dd, J=13.5 and 8.9 Hz, 1H), 2.78 (m, 1H), 2.67-2.61 (m,1H), 2.58-2.49 (m, 1H), 2.38-2.32 (m,1H), 2.32-2.04 (m, 1H), 1.99-1.92 (m, 1H,) 1.45 (s, 9H), 1.29 (s, 9H).

Step 6: Preparation of 2(R)-phenylmethyl-4(S)-(t-butyldimethylsilyloxy)-5-(1-(4-(1,1-dimethylethoxycarbonylamino)))-2(S)-N-(t-butylcarboxamido)-piperazinyl))-pentaneamide

To 25.50g (52.50 mmol) of dihydro-5(S)-(4-(1,1-dimethylethoxycarbonylamino))-2(S)-N-(t-butyl-carboxamido)-piperazinyl)methyl)-3(R)-phenylmethyl-3(2H)-furanone dissolved in 120 mL DME cooled to 0°C was added a solution of 60 mL of water and 1.512g (63.01 mmol) of lithium hydroxide. After 0.5 hours the reaction was quenched with the addition of 10% HCl until pH 6 and the solution was concentrated in vacuo. The residue was dissolved in 50 mL water and extracted with EtOAc (4 x 75 mL) and the organic layers were washed with water (1 x 20 mL), brine (1 x 20 mL). The aqueous was back extracted with EtOAc (2 x 75 mL) and the combined organic layers were dried over $MgSO_4$ and concentrated to provide a yellow solid. This crude product was dissolved in 100 mL of DMF and 17.87g (0.262 mol) of imidazole was added, cooled to 0°C and then 31.50g (0.21 mol) of t-butyldimethylsilyl chloride was added. This stirred 1 hour at 0°C and was then

This Sheet Has Been Replaced

warmed to room temperature. After 20 hours the reaction was quenched with 10 mL methanol and concentrated to half the volume. 100 mL of pH 7 buffered water was added and the aqueous was extracted with EtOAc (4 x 100 mL), the combined organic layers were washed with 10% HCl (2 x 50 mL), water (3 x 75 mL), and brine (1 x 50 mL), dried over $MgSO_4$ and concentrated to obtain the title compound. This material was used directly in the next step.

Step 7: Preparation of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-(t-butyldimethylsilyl-oxy)-5-(1-(4-(1,1-dimethylethoxycarbonyl-amino)))-2(S)-N-(t-butylcarboxamido)-piper-azinyl))-pentaneamide

To 27.0g (0.0446mol) of the crude material from step 6 dissolved in 180 mL of DMF and cooled to 0°C was added 8.98g (0.0468 mol) of EDC, 6.32g (0.0468 mol) of HOBt, and 7.31g (0.049 mol) aminohydroxy indane. Triethylamine (6.52 mL, 0.0468 mol) was added and the reaction stirred at 0°C for 2 hours, room temperature for 16 hours and was quenched by diluting with 500 mL of EtOAc. The organic layer was washed with 10% HCl (2 x 100 mL), saturated $NaHCO_3$ (1 x 100 mL), water (3 x 150 mL), brine (1 x 75 mL), dried over $MgSO_4$ and concentrated to yield the title compound as a white foam.

^1H NMR (400 MHz, $CDCl_3$) δ 7.4-7.17 (m, 9H), 6.51 (br s, 1H), 5.79 (br s, 1H), 5.23 (m, 1H), 4.23 (br s, 1H), 4.06 (m, 1H), 3.96-3.84 (m, 2H), 3.07-2.78 (m, 8H), 3.65 (dd, J=9.6 and 4.1 Hz, 1H), 2.56-2.44 (m, 2H), 2.29 (dd, J=12.0 and 4.5 Hz, 1H), 2.17-2.09 (m, 1H), 1.79 (br s, 1H), 1.44 (s, 9H), 1.35 (s, 9H), 1.10 (s, 1H), 0.84 (s, 9H), 0.12 (s, 3H), 0.08 (s, 3H).

This Sheet Has Been Replaced

Step 8: Preparation of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-(hydroxy)-5-(1-(4-(1,1-dimethylethoxycarbonylamino)))-2(S)-N-(t-butylcarboxamido)-piperazinyl))-pentaneamide

To 32.20g (0.0437 mol) of N-(2(R)-hydroxy-1-(S)-indanyl)-2(R)-phenylmethyl-4(S)-(t-butyldimethylsilyloxy)-5-(1-(4-(1,1-dimethylethoxycarbonylamino)))-2(S)-N-(t-butylcarboxamido)-piperazinyl))-pentaneamide was added 437 mL (0.437 mol) of tetrabutylammonium fluoride (1.0M solution in THF, Aldrich). The reaction stirred for 18 hours and was then concentrated to 200 mL and diluted with 700 mL of EtOAc. This was washed with water (2 x 100 mL), brine (1 x 50 mL) and the aqueous layers were back extracted with EtOAc (2 x 200 mL). The combined organic layers were dried over $MgSO_4$ and concentrated to an oil. Purification via flash column chromatography (120 x 150 mm column, gradient elution CH_2Cl_2: $CHCl_3$/saturated with NH_3: methanol, increasing methanol from 1%, 1.5%, 2%) afforded the title compound as a white foam.
^1H NMR (400 MHz, $CDCl_3$) δ 7.31-7.11 (m, 9H), 6.41 (br s, 1H), 6.23 (d, J=8.6 Hz, 1H), 5.25 (dd, J=8.6 and 4.7Hz, 1H), 4.21 (m, 1H), 3.83-3.82 (m, 2H), 3.78-3.61 (m, 2H), 3.22-3.19 (m, 2H), 3.03-2.78 (m, 8H), 2.62-2.58 (m, 1H), 2.41-2.35 (m, 2H), 2.04-2.02 (m, 1H), 1.57-1.50 (m, 1H), 1.45 (s, 9H), 1.32 (s, 9H).

This Sheet Has Been Replaced

Step 9: Preparation of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-(hydroxy)-5-(1-(2(S)-N-(t-butylcarboxamido)-piperazinyl)-pentaneamide

To 21.15g (0.034 mol) of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-(hydroxy)-5-(1-(4-(1,1-dimethylethoxycarbonylamino)))-2(S)-N-(t-butylcarboxamido)-piperazinyl))-pentaneamide dissolved in 350 mL of methylene chloride and cooled to 0°C was added 22.43 mL (0.204 mol) 2,6-lutidine and then 32.85 mL (0.170 mol) of trimethylsilyltriflate over 5 minutes. After 0.5 hours the reaction was quenched with 10% HCl (80 mL) and this stirred 0.5 hours. To this was added 100 mL of saturated $NaHCO_3$ and then solid $NaHCO_3$ until pH 8. The aqueous layer was then extracted with EtOAc (4 x 100 mL) and the combined organic layers were washed with water (1 x 50 mL), brine (1 x 75 mL), dried over $MgSO_4$ and concentrated. The residue was purified via column chromatography (120 x 150 mm column, gradient elution CH_2Cl_2:$CHCl_3$ saturated with NH_3: MeOH, slowly increasing methanol 2%, 3%, 4%, 5%, 6%, to 10%). This provided the title product as a white foam.

^1H NMR (400 MHz, $CDCl_3$) δ 7.53 (s, 1H), 7.29-7.09 (m, 9H), 6.52 (d, J=8.3 Hz, 1H), 5.24 (dd, J=8.2 and 4.9 Hz, 1H), 4.23 (dd, J=4.7 and 4.03 Hz, 1H), 4.25-4.00 (br s, 1H), 3.83-3.81 (m, 1H), 3.03-2.88 (m, 4H), 2.82-2.73 (m, 7H), 2.50-1.60 (br s, 2H), 2.45 (d, J=6.2 Hz, 2H), 2.32-2.29 (m, 1H), 1.98 (m, 1H), 1.51 (m, 1H), 1.33 (s, 9H).

This Sheet Has Been Replaced


Step 10: Preparation of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(1-(4-(3-pyridylmethyl)-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide

To 10.0g (0.019 mol) of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-hydroxy)-5-(1(-2(S)-N-(t-butylcarboxamido)-piperazinyl)-pentaneamide and 3.45g (0.021 mol) of 3-picolyl chloride dissolved in 40 mL of DMF was added 5.85 mL (0.042 mol) of triethylamine. After 3 hours an additional 0.313g of 3-picolyl chloride was added. After an additional 2 hours the reaction was diluted with 400 mL of EtOAc and washed with water (3 x 75 mL), brine (1 x 100 mL), dried over $MgSO_4$ and concentrated. The residue was triturated with 30 mL of EtOAc and the resulting white precipitate was collected. Further recrystallization from EtOAc provided the title product (mp 167.5-168°C).

EXAMPLE 16


Employing substantially the same procedure as described in Example 15, but treating the N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(1-(2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide used therein (compound (i) below) with the alkylating agent (ii) indicated below in place of the 3-picolyl chloride used in Step 10 therein, the following products defined by formula (iii) were made:

This Sheet Has Been Replaced







i







ii











iii

R^1	X
	I
	Cl
	Cl
	I






This Sheet Has Been Replaced

R^1	X
CH_3CH_2-	I
	Br
	Cl
	Cl
	Cl
	I
	Cl

This Sheet Has Been Replaced

R^1	X
	Cl
	I
	Cl
	Cl
	Cl
$CH_3O(CH_2CH_2O)_2-CH_2CH_2-$	I
	I
	I
	Cl

This Sheet Has Been Replaced

R^1	X
	I
	Cl
	Cl
	Cl
	Cl

EXAMPLE 17

Preparation of dihydro-5(S)-(tert-butyldimethylsilyl-oxymethyl)-3(2H)-furanone

To a solution of 3.00g (25.8 mmol) of dihydro-5(S)-(hydroxymethyl)-2(3H)-furanone dissolved in 25 mL of dichloromethane was added 3.51g (51.6 mmol) of imidazole and then 4.67g (31.0 mmol) of tert-butyl-

This Sheet Has Been Replaced


dimethylsilyl chloride. The reaction stirred at room temperature for 8 hours and was quenched with 2 mL of methanol. The mixture was concentrated to an oil and then diluted with 150 mL of ether and washed with 5% HCl (2 x 10 mL), saturated $NaHCO_3$ (1 x 10 mL), water (1 x 10 mL), and brine (1 x 10 mL), dried over $MgSO_4$ and concentrated. The residue was purified by flash chromatography (40 x 150 mm column, gradient elution, hexanes:ethyl/acetate 5:1 to 4:1) to afford the product as a clear oil.
^1H NMR (300 MHz, $CDCl_3$) δ 4.68-4.60 (m, 1H), 3.89 (dd, J=3.3 and 11.3 Hz, 1H), 3.71 (dd, J=3.2 and 5411.3 Hz,1H), 2.71-2.45 (m, 2H), 2.35-2.16 (m, 2H), 0.91 (s, 9H), 0.10 (s, 3H), 0.09 (s, 3H).

While the foregoing specification teaches the principles of the present invention, with examples provided for the purpose of illustration, it will be understood that the practice of the invention emcompasses all of the usual variations, adaptations, or modifications, as come within the scope of the following claims and its equivalents.

This Sheet Has Been Replaced

WHAT IS CLAIMED IS:

1. A compound of the formula:



I

wherein

X is -OH or $-NH_2$;

Z is -O, -S, or -NH;

R is hydrogen or C_{1-4} alkyl;

R^1 and R^2 are independently:

1) hydrogen,

2) $-C_{1-4}$ alkyl unsubstituted or substituted with one or more of

a) halo,

b) hydroxy,

c) C_{1-3} alkoxy,

d) aryl unsubstituted or substituted with one or more of C_{1-4}alkyl, hydroxy or aryl,

e) -W-aryl or -W-benzyl, wherein W is -O-, -S-, or -NH-,

f) a 5-7 membered cycloalkyl group unsubstituted or substituted with one or more of

This Sheet Has Been Replaced

i) halo,
ii) hydroxy,
iii) C_{1-3} alkoxy, or
iv) aryl,

g) heterocycle unsubstituted or substituted with one or more of hydroxy, C_{1-4}alkyl optionally substituted with hydroxy, or Boc,

h) $-NH-\overset{O}{\overset{\|}{C}}OC_{1-3}alkyl$,

i) $-NH-\overset{O}{\overset{\|}{C}}-C_{1-3}alkyl$,

j) $-NH-SO_2C_{1-3}alkyl$,

k) $-NR_2$,

l) -COOR, or

m) $-((CH_2)_mO)_nR$ wherein m is 2-5 and n is zero, 1, 2 or 3, or

3) aryl, unsubstituted or substituted with one or more of

a) halo,

b) hydroxy,

c) $-NO_2$ or $-NR_2$,

d) C_{1-4}alkyl,

e) C_{1-3} alkoxy, unsubstituted or substituted with one or more of -OH or C_{1-3} alkoxy,

f) -COOR,

g) $-\overset{O}{\overset{\|}{C}}NR_2$,

h) $-CH_2NR_2$,

This Sheet Has Been Replaced

i) $-CH_2NHC(=O)R$,
j) $-CN$,
k) $-CF_3$,
l) $-NHC(=O)R$,
m) aryl C_{1-3} alkoxy,
n) aryl,
o) $-NRSO_2R$,
p) $-OP(O)(OR_x)_2$, or
q) $-R^5$, as defined below; or

R^1 and R^2 can be joined together to form with the nitrogen to which R^1 is attached a 3 to 10 membered monocyclic or bicyclic saturated ring system which consists of the nitrogen to which R^1 is attached and from 2 to 9 carbon atoms, and is unsubstituted or substituted with

1) hydroxy,
2) C_{1-4} alkyl unsubstituted or substituted with one or more of
   a) halo,
   b) hydroxy,
   c) C_{1-3} alkoxy,
   d) aryl,
   e) a 5-7 membered cycloalkyl group unsubstituted or substituted with one or more of
      i) halo,
      ii) hydroxy,
      iii) C_{1-3} alkoxy, or
      iv) aryl,
   f) heterocycle, or
   g) $-NR_2$,

This Sheet Has Been Replaced

3) C_{1-3} alkoxy,

4) $-NH-\overset{O}{\overset{\|}{C}}OC_{1-3}alkyl$,

5) $-NH-\overset{O}{\overset{\|}{C}}-C_{1-3}alkyl$,

6) $-NH-SO_2C_{1-3}alkyl$,

7) heterocycle,

8) -W-aryl, or

9) $-W-\overset{O}{\overset{\|}{C}}-aryl$,

wherein W is defined above; or

R^1 and R^2 can be joined together to form with the nitrogen to which R^1 is attached a 3 to 10 membered monocyclic or bicyclic saturated ring system which consists of the nitrogen to which R^1 is attached, from 1 to 8 carbon atoms and one or more unsubstituted or substituted heteroatom selected from

1) $-\overset{|}{N}-V-R^1$,

wherein V is absent or $-\overset{O}{\overset{\|}{C}}-Q-$ or $-SO_2-Q-$, R^1 is defined as above for when R^1 is independent from and not joined to R^2, and wherein Q is absent or -O-, -NR-, or heterocycle optionally substituted with $-C_{1-4}alkyl$,

2) $-\overset{|}{N}-$heterocycle,

3) $-\overset{|}{N}-C_{1-4}$ alkenyl, unsubstituted or substituted with aryl,

This Sheet Has Been Replaced

4) $-N(SO_2-C_{1-4}alkenyl)-$, unsubstituted or substituted with aryl,

5) $-S(O)_p-$, wherein p is zero, 1 or 2, or

6) $-O-$; or

R^1 and R^2 can be joined together to form with the nitrogen to which R^1 is attached a 3 to 10 membered monocyclic or bicyclic saturated ring system, which consists of the nitrogen to which R^1 is attached and from 2 to 9 carbon atoms, in which the saturated ring system is fused to a phenyl ring and the phenyl ring is unsubstituted or substituted with one or more of

1) halo,
2) C_{1-3} alkoxy,
3) hydroxy,
4) C_{1-4} alkyl,
5) $-NHR^1$, wherein R^1 is defined as above for when R^1 is independent from and not joined to R^2, or
6) -NH-heterocycle;

R^3 is

1) $-(CH_2)_r-R^4$, wherein r is zero through 5,
2) $C_{1-4}alkenyl-R^4$, or
3) $C_{1-4}alkynyl-R^4$;

R^4 is

1) hydrogen,
2) C_{1-4} alkyl,

This Sheet Has Been Replaced

3) C_5-C_{10} cycloalkyl, optionally substituted with hydroxy,

4) C_6-C_{10} aryl, unsubstituted or substituted with one or more of

a) halo,

b) hydroxy,

c) $-NO_2$ or $-NR_2$,

d) C_{1-4}alkyl,

e) C_{1-3} alkoxy, unsubstituted or substituted with one or more of -OH or C_{1-3} alkoxy,

f) -COOR,

g) $-\overset{\overset{O}{\|}}{C}NR_2$,

h) $-CH_2NR_2$,

i) $-CH_2NH\overset{\overset{O}{\|}}{C}R$,

j) -CN,

k) $-CF_3$,

l) $-NH\overset{\overset{O}{\|}}{C}R$,

m) aryl C_{1-3} alkoxy,

n) aryl,

o) $-NRSO_2R$,

p) $-OP(O)(OR_x)_2$, or

q) $-R^5$, as defined below, or

5) monocyclic or bicyclic heterocyle containing from 1 to 3 heteroatoms chosen from the group consisting of N, O, and S and which is unsubstituted or substituted with R^5 and optionally with one or more of

a) halo,

b) C_{1-4} alkyl, or

c) C_{1-3} alkoxy;

This Sheet Has Been Replaced

R_x is H or aryl;


R^5 is

1) $-W-(CH_2)_m-NR^6R^7$
wherein W is as defined above,
m is 2-5, and
R^6 and R^7 are independently
a) hydrogen,
b) C_{1-6} alkyl, unsubstituted or substituted with one or more of
i) C_{1-3} alkoxy,
ii) -OH, or
iii) $-NR_2$,
c) the same or different and joined together to form a 5-7 member heterocycle, such as morpholino, containing up to two additional heteroatoms selected from
$-\overset{R}{\underset{|}{N}}-$, -O-, $-\overset{O}{\overset{\|}{S}}-$, -S-, or $-SO_2-$, the heterocycle optionally substituted with C_{1-4} alkyl, or
d) aromatic heterocycle unsubstituted or substituted with one or more of
i) C_{1-4} alkyl, or
ii) $-NR_2$,

2) $-(CH_2)_q-NR^6R^7$ wherein q is 1-5, and R^6 and R^7 are defined above, except that R^6 or R^7 are not H or unsubstituted C_{1-6} alkyl, or

3) benzofuryl, indolyl, azacycloalkyl, azabicyclo C_{7-11} cycloalkyl, or benzopiperidinyl, unsubstituted or substituted with C_{1-4} alkyl;

This Sheet Has Been Replaced

B is absent, or ,

wherein R^8 is

1) $-CH(CH_3)_2$,
2) $-CH(CH_3)(CH_2CH_3)$, or
3) -phenyl;

J^1 and J^2 are independently

1) $-YR^9$ wherein

Y is -O- or -NH-, and

R^9 is

a) hydrogen,

b) C_{1-6} alkyl, unsubstituted or substituted with one or more of

i) $-NR_2$,

ii) -OR,

iii) $-NHSO_2C_{1-4}$ alkyl,

iv) $-NHSO_2$ aryl, or $-NHSO_2$(dialkyl-aminoaryl),

v) $-CH_2OR$,

vi) $-C_{1-4}$ alkyl,

vii) $-\overset{O}{\overset{\|}{C}}OR$,

viii) $-\overset{O}{\overset{\|}{C}}NR_2$,

ix) $-NH(\overset{NH}{\overset{\|}{C}}NR_2)$ or $-NH(\overset{N-CN}{\overset{\|}{C}}NR_2)$,

x) $-NH\overset{O}{\overset{\|}{C}}R^{13}$, wherein R^{13} is

This Sheet Has Been Replaced

A) -H,
B) -C_{1-4} alkyl,
C) -aryl,
D) -heterocycle, or
E) -NH-, -O- or -$(CH_2)_n$-
wherein n is zero, 1, 2 or 3,
substituted with
I) -C_{1-4} alkyl,
unsubstituted or
substituted with one or
more of aryl or
heterocycle, or
II) aryl, unsubstituted or
substituted with
heterocycle,

xi) -$NR_3^{\oplus}$ $A^{\ominus}$ wherein $A^{\ominus}$ is a counterion,

xii) -$NR^{10}R^{11}$ wherein R^{10} and R^{11} are the same or different and are C_{1-5} alkyl joined together directly to form a 5-7 membered heterocycle containing up to one additional heteroatom selected from -O-, -S-, or -NR-,

xiii)-aryl,

xiv) -CHO,

xv) -$OP(O)(OR_x)_2$,

xvi) -O-$\overset{O}{\overset{\|}{C}}$-$C_{1-4}$alkyl substituted with one or more of amine or quaternary amine, or -O-$((CH_2)_mO)_n$-R, or -$OP(O)(OR_x)_2$,

This Sheet Has Been Replaced


xvii) $-O\overset{O}{\overset{\|}{C}}-R$, or

xviii) $-O\overset{O}{\overset{\|}{C}}-NH-CH_2$-heterocycle, or

c) $-((CH_2)_mO)_nCH_3$ or $-((CH_2)_mO)_n$ H, wherein m and n are defined above,

2) $-N(R^9)_2$,

3) $-NR^{10}R^{11}$ wherein R^{10} and R^{11} are defined above, or

4) 

wherein Y, R^9 and n are defined above; and

R^{12} is

1) hydrogen,

2) aryl, unsubstituted or substituted with one or more of

a) R^{14}, wherein R^{14} is

i) halo,

ii) $-OR$,

iii) $-\overset{O}{\overset{\|}{C}}NR_2$,

iv) $-CH_2NR_2$,

v) $-SO_2NR_2$,

vi) $-NR_2$,

vii) $-NH\overset{O}{\overset{\|}{C}}R$,

viii) $-C_{1-4}$ alkyl,

ix) phenyl

This Sheet Has Been Replaced

x) $-CF_3$,

xi) $-\overset{R}{\underset{|}{N}}-SO_2R$,

xii) $-OP(O)(OR_x)_2$, or

xiii) $-\overset{O}{\overset{\|}{C}}OR$,

b) $-C_{1-4}$ alkyl-NR_2, or

c) $-O-\overset{O}{\overset{\|}{C}}-C_{1-4}$alkyl substituted with one or more of amine or quaternary amine or $-OP(O)(OR_x)_2$,

3) heterocycle, the ring or rings being unsubstituted or substituted with one or more of

a) R^{14}, as defined above,

b) $-OC_{1-4}$ alkenyl,

c) phenyl-C_{1-4} alkyl,

d) $-O-\overset{O}{\overset{\|}{C}}-C_{1-4}$alkyl substituted with one or more of amine or quaternary amine, or $-OP(O)(OR_x)_2$, or $-O((CH_2)_mO)_n-R$, or

e) $-O-\overset{O}{\overset{\|}{C}}-O-((CH_2)_mO)_n-R$, or

4) A 5 to 7 membered carbocyclic or 7-10 membered bicyclic carbocyclic ring, the carbocyclic ring being unsubstituted or substituted with one or more of

a) R^{14}, as defined above,

b) $-CH_2OR$,

This Sheet Has Been Replaced

c) $-(CH_2)_n-NR_2$, C_{5-16}alkyl, pyridine, $-(CH_2)_nNR-(CH_2)_n-NR_2$, $-(CH_2)_n-\overset{O}{\overset{\|}{C}}-OR$, $-((CH_2)_mO)_n-R$, quinuclidiniumyl substituted with R, piperazine-C_{1-4}alkyl-benzyl substituted once or more with R, or morpholino-C_{1-4}alkyl-benzyl,

d) $-O-\overset{O}{\overset{\|}{C}}-C_{1-4}$alkyl substituted with one or more of amine or quaternary amine, $-OP(O)(OR_x)_2$, or $-O-((CH_2)_mO)_n-R$,

e) $-O-\overset{O}{\overset{\|}{C}}-O-((CH_2)_mO)_n-R$, or

f) $-C_{1-4}$alkyl-phenyl;

or a pharmaceutically acceptable salt thereof.

2. A compound according to Claim 1 wherein:

R^1 and R^2 are joined together to form with the nitrogen to which R^1 is attached a 3 to 10 membered monocyclic or bicyclic saturated ring system which consists of the nitrogen to which R^1 is attached and from 2 to 9 carbon atoms, and is unsubstituted or substituted with

1) hydroxy,
2) C_{1-4} alkyl unsubstituted or substituted with one or more of
   a) hydroxy,
   b) C_{1-3} alkoxy,
   c) aryl,

This Sheet Has Been Replaced

d) a 5-7 membered cycloalkyl group unsubstituted or substituted with one or more of
   i) halo,
   ii) hydroxy,
   iii) C_{1-3} alkoxy, or
   iv) aryl,
e) heterocycle, or
f) $-NR_2$,

3) C_{1-3} alkoxy,

4) $-NH-\overset{O}{\overset{\|}{C}}OC_{1-3}alkyl$,

5) $-NH-\overset{O}{\overset{\|}{C}}-C_{1-3}alkyl$,

6) $-NH-SO_2C_{1-3}alkyl$,

7) -W-aryl, or

8) $-W-\overset{O}{\overset{\|}{C}}-aryl$,

wherein W is -O-, -S-, or -NH-; or

R^1 and R^2 are joined together to form with the nitrogen to which R^1 is attached a 3 to 10 membered monocyclic or bicyclic saturated ring system which consists of the nitrogen to which R^1 is attached, from 1 to 8 carbon atoms and one or more unsubstituted or substituted heteroatom selected from

1) $-\underset{\underset{V-R^1}{|}}{N}-$,

wherein V is absent or $-\overset{O}{\overset{\|}{C}}-Q-$ or $-SO_2-Q-$, R^1 is defined as above for when R^1 is independent from and not joined to R^2, and wherein Q is absent or -O-, -NR-, or

This Sheet Has Been Replaced

heterocycle optionally substituted with $-C_{1-4}$alkyl,

2) $-\overset{|}{N}-$
C_{1-4} alkenyl, unsubstituted or substituted with aryl,

3) $-S(O)_p-$,
wherein p is zero, 1 or 2, or

4) $-O-$; or

R^1 and R^2 are joined together to form with the nitrogen to which R^1 is attached a 3 to 10 membered monocyclic or bicyclic saturated ring system, which consists of the nitrogen to which R^1 is attached and from 2 to 9 carbon atoms, in which the saturated ring system is fused to a phenyl ring and the phenyl ring is unsubstituted or substituted with one or more of

1) C_{1-3} alkoxy,
2) hydroxy,
3) C_{1-4} alkyl, or
4) $-NHR^1$,
wherein R^1 is defined as above for when R^1 is independent from and not joined to R^2.

3. A compound according to Claim 2 wherein:

R^1 and R^2 are joined together to form with the nitrogen to which R^1 is attached a 3 to 10 membered monocyclic or bicyclic saturated ring system which consists of the nitrogen to which R^1 is attached and from 2 to 9 carbon atoms, and is unsubstituted or substituted with

1) hydroxy,
2) C_{1-4} alkyl unsubstituted or substituted with one or more of

This Sheet Has Been Replaced

a) hydroxy,
b) C_{1-3} alkoxy,
c) aryl,
d) a 5-7 membered cycloalkyl group unsubstituted or substituted with one or more of
i) halo,
ii) hydroxy,
iii) C_{1-3} alkoxy, or
iv) aryl,
e) heterocycle, or
f) $-NR_2$,

3) C_{1-3} alkoxy,

4) $-NH-\overset{O}{\overset{\|}{C}}OC_{1-3}alkyl$,

5) $-NH-\overset{O}{\overset{\|}{C}}-C_{1-3}alkyl$,

6) $-NH-SO_2C_{1-3}alkyl$,

7) -W-aryl, or

8) $-W-\overset{O}{\overset{\|}{C}}-aryl$,

wherein W is -O-, -S-, or -NH-; or

R^1 and R^2 are joined together to form with the nitrogen to which R^1 is attached a 3 to 10 membered monocyclic or bicyclic saturated ring system which consists of the nitrogen to which R^1 is attached, from 1 to 8 carbon atoms and one or more unsubstituted or substituted heteroatom selected from

1) $-\underset{\underset{V-R^1}{|}}{N}-$,

wherein V is absent or $-\overset{O}{\overset{\|}{C}}-Q-$ or $-SO_2-Q-$,

This Sheet Has Been Replaced

R^1 is defined as above for when R^1 is independent from and not joined to R^2, and wherein Q is absent or -O-, -NR-, or heterocycle optionally substituted with

$-C_{1-4}$alkyl,

2) $-S(O)_p-$,
   wherein p is zero, 1 or 2, or

3) -O-;

R^3 is benzyl, unsubstituted or substituted with one or more of

a) hydroxy,
b) $-NO_2$, or $-NR_2$,
c) C_{1-4}alkyl,
d) C_{1-3} alkoxy, unsubstituted or substituted with one or more of -OH or C_{1-3} alkoxy,
e) $-\overset{\overset{O}{\|}}{C}NR_2$,
f) $-CH_2NR_2$,
g) $-CH_2NH\overset{\overset{O}{\|}}{C}R$,
h) $-CF_3$,
i) $-NH\overset{\overset{O}{\|}}{C}R$,
j) $-NRSO_2R$,
k) $-OP(O)(OR_x)_2$, or
l) $-R^5$;

and B is absent.

This Sheet Has Been Replaced

4. A compound according to Claim 3 wherein:

X is -OH;


Z is -O;

R^1 and R^2 are joined together to form with the nitrogen to which R^1 is attached a 3 to 10 membered monocyclic or bicyclic saturated ring system which consists of the nitrogen to which R^1 is attached and from 2 to 9 carbon atoms, and is unsubstituted or substituted with -W-aryl or -W-C(=O)-aryl; or

R^1 and R^2 are joined together to form with the nitrogen to which R^1 is attached a 3 to 10 membered monocyclic or bicyclic saturated ring system which consists of the nitrogen to which R^1 is attached, from 1 to 8 carbon atoms and one of -N(V-R^1)- ,


wherein V is absent or -C(=O)-Q- or -SO_2-Q-,
R^1 is defined as above for when R^1 is independent from and not joined to R^2,
and wherein Q is absent or -O-, -NR- or heterocycle optionally substituted with -C_{1-4}alkyl;

R^3 is benzyl, unsubstituted or substituted with one or more of (1) hydroxy, (2) C_{1-3} alkoxy substituted with one or more of -OH or (3)


;

This Sheet Has Been Replaced

J^1 is $-NH-C_{1-4}$alkyl; and
J^2 is



5. The compound

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(2-(3(S)-N'-(t-butylcarboxamido)-(4aS,8aS)-decahydroisoquinoline)yl)-pentaneamide,

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(1-(4-carbobenzyloxy-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide,

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-(2-(4-morpholinyl)ethoxy)phenyl)methyl)-4(S)-hydroxy-5-(2-(3(S)-N'-(t-butylcarboxamido)-(4aS,8aS)-decahydroisoquinoline)yl)-pentaneamide,

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-(2-(4-morpholinyl)ethoxy)phenyl)methyl)-4(S)-hydroxy-5-(1-(4-carbobenzyloxy-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide,

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-((2-hydroxy)-ethoxy)phenyl)methyl)-4(S)-hydroxy-5-(2-(3(S)-N'-(t-butylcarboxamido)-(4aS,8aS)-decahydroisoquinoline)-yl)-pentaneamide,

This Sheet Has Been Replaced

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-((2-hydroxy)-ethoxy)phenyl)methyl)-4(S)-hydroxy-5-(1-(4-carbobenzyloxy-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide,

N-(4(S)-3,4-dihydro-1H-2,2-dioxobenzothiopyranyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(2-(3(S)-N'-(t-butyl-carboxamido)-(4aS,8aS)-decahydroisoquinoline)yl)-pentaneamide,

N-(4(S)-3,4-dihydro-1H-2,2-dioxobenzothiopyranyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(1-(4-carbobenzyl-oxy-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide,

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(1-(4-(3-pyridylmethyl)-2(S)-N'-(t-butyl-carboxamido)-piperazinyl))-pentaneamide, or

a pharmaceutically acceptable salt thereof.

6. A pharmaceutical composition comprising a compound of Claim 1 and a pharmaceutically acceptable carrier.

7. The use of a compound of Claim 1 for the preparation of a medicament useful for treating AIDS in a patient receiving AIDS therapy.

This Sheet Has Been Replaced

8. The use of a compound of Claim 1 for the preparation of a medicament for preventing or treating infection by HIV in a patient.

9.. The use of a compound of Claim 1 for the preparation of a medicament useful for inhibiting HIV protease in a patient receiving HIV protease therapy.

This Sheet Has Been Replaced

138/CSQ55
139/CSQ56


18597IA

TITLE OF THE INVENTION

HIV PROTEASE INHIBITORS USEFUL FOR THE TREATMENT OF AIDS

ABSTRACT OF THE DISCLOSURE

Compounds of formula



where R^1 and R^2 are independently hydrogen or optionally-substituted C_{1-4}alkyl or aryl, or R^1 and R^2 are joined together to form a monocyclic or bicyclic ring system, are HIV protease inhibitors.

These compounds are useful in the prevention or treatment of infection by HIV and in the treatment of AIDS, either as compounds, pharmaceutically acceptable salts, pharmaceutical composition ingredients, whether or not in combination with other antivirals, immunomodulators, antibiotics or vaccines. Methods of treating AIDS and methods of preventing or treating infection by HIV are also described.

This Sheet Has Been Replaced

REPLACEMENT SHEET
04 FEB 1997

REPLACEMENT SHEETS SG
☒ Description : paged 1 to paged 101
☒ Claim(s) : paged 102 to paged 121
☐ Drawing(s) : paged ..... to paged .....
☒ Abstract : paged 122 to paged .....
DATE OF RECEIPT 4/2/97

TITLE OF THE INVENTION

HIV PROTEASE INHIBITORS USEFUL FOR THE TREATMENT OF AIDS

The present invention is concerned with compounds which inhibit the protease encoded by human immunodeficiency virus (HIV) or pharmaceutically acceptable salts thereof and are of value in the prevention of infection by HIV, the treatment of infection by HIV and the treatment of the resulting acquired immune deficiency syndrome (AIDS). It also relates to pharmaceutical compositions containing the compounds and to a method of use of the present compounds and other agents for the treatment of AIDS and viral infection by HIV.

REPLACEMENT SHEET
04 FEB 1997

BACKGROUND OF THE INVENTION

A retrovirus designated human immunodeficiency virus (HIV) is the etiological agent of the complex disease that includes progressive destruction of the immune system (acquired immune deficiency syndrome; AIDS) and degeneration of the central and peripheral nervous system. This virus was previously known as LAV, HTLV-III, or ARV. A common feature of retrovirus replication is the extensive post-translational processing of precursor polyproteins by a virally encoded protease to generate mature viral proteins required for virus assembly and function. Inhibition of this processing prevents the production of normally infectious virus. For example, Kohl, N.E. et al., Proc. Nat'l Acad. Sci. 85, 4686 (1988) demonstrated that genetic inactivation of the HIV encoded protease resulted in the production of immature, non-infectious virus particles. These results indicate that inhibition of the HIV protease represents a viable method for the treatment of AIDS and the prevention or treatment of infection by HIV.

The nucleotide sequence of HIV shows the presence of a pol gene in one open reading frame [Ratner, L. et al., Nature, 313, 277(1985)]. Amino acid sequence homology provides evidence that the pol sequence encodes reverse transcriptase, an endonuclease and an HIV protease [Toh, H. et al., EMBO J. 4, 1267 (1985); Power, M.D. et al., Science, 231, 1567 (1986); Pearl, L.H. et al., Nature 329, 351 (1987)]. Applicants demonstrate that the compounds of this invention are inhibitors of HIV protease.

REPLACEMENT SHEET
04 FEB 1997

BRIEF DESCRIPTION OF THE INVENTION

Compounds of formula I, as herein defined, are disclosed. These compounds are useful in the inhibition of HIV protease, the prevention of infection by HIV, the treatment of infection by HIV and in the treatment of AIDS, either as compounds, pharmaceutically acceptable salts, pharmaceutical composition ingredients, whether or not in combination with other antivirals, immunomodulators, antibiotics or vaccines. Methods of treating AIDS, methods of preventing infection by HIV, and methods of treating infection by HIV are also disclosed.

Some abbreviations that may appear in this application are as follows.

ABBREVIATIONS

Designation	Protecting Group
BOC (Boc)	t-butyloxycarbonyl
CBZ (Cbz)	benzyloxycarbonyl(carbo-benzoxy)
TBS (TBDMS)	t-butyl-dimethylsilyl
	Activating Group
HBT(HOBT or HOBt)	1-hydroxybenzotriazole hydrate


REPLACEMENT SHEET
04 FEB 1997

Designation	Coupling Reagent
BOP reagent	benzotriazol-1-yloxytris-(dimethylamino)phospho-nium hexafluorophosphate
BOP-Cl	bis(2-oxo-3-oxazolidinyl) phosphinic chloride
EDC	1-ethyl-3-(3-dimethyl-aminopropyl) carbodiimide hydrochloride
	Other
$(BOC)_2O$ (BOC_2O)	di-t-butyl dicarbonate
$n\text{-}Bu_4N^+F^-$	tetrabutyl ammonium fluoride
nBuLi (n-Buli)	n-butyllithium
DMF	dimethylformamide
Et_3N	triethylamine
EtOAc	ethyl acetate
TFA	trifluoroacetic acid
DMAP	dimethylaminopyridine
DME	dimethoxyethane
LDA	lithium diisopropylamide
THF	tetrahydrofuran
	Amino Acid
Ile	L-isoleucine
Val	L-valine

REPLACEMENT SHEET
04 FEB 1997

DETAILED DESCRIPTION OF THE INVENTION AND PREFERRED EMBODIMENTS

This invention is concerned with compounds of formula I, combinations thereof, or pharmaceutically acceptable salts thereof, in the inhibition of HIV protease, the prevention or treatment of infection by HIV and in the treatment of the resulting acquired immune deficiency syndrome (AIDS). Compounds of formula I are defined as follows:



I

wherein

X is -OH or $-NH_2$;

Z is -O, -S, or -NH;

R is hydrogen or C_{1-4} alkyl;

R^1 and R^2 are independently:

1) hydrogen,
2) $-C_{1-4}$ alkyl unsubstituted or substituted with one or more of
   a) halo,
   b) hydroxy,
   c) C_{1-3} alkoxy,
   d) aryl unsubstituted or substituted with one or more of C_{1-4}alkyl, hydroxy or aryl,

REPLACEMENT SHEET
04 FEB 1997

e) -W-aryl or -W-benzyl,
wherein W is -O-, -S-, or -NH-,

f) a 5-7 membered cycloalkyl group unsubstituted or substituted with one or more of
i) halo,
ii) hydroxy,
iii) C_{1-3} alkoxy, or
iv) aryl,

g) heterocycle unsubstituted or substituted with one or more of hydroxy, C_{1-4}alkyl optionally substituted with hydroxy, or Boc,

h) $-NH-\overset{O}{\overset{\|}{C}}OC_{1-3}alkyl$,

i) $-NH-\overset{O}{\overset{\|}{C}}-C_{1-3}alkyl$,

j) $-NH-SO_2C_{1-3}alkyl$,

k) $-NR_2$,

l) -COOR, or

m) $-((CH_2)_mO)_nR$ wherein m is 2-5 and n is zero, 1, 2 or 3, or

3) aryl, unsubstituted or substituted with one or more of
a) halo,
b) hydroxy,
c) $-NO_2$ or $-NR_2$,
d) C_{1-4}alkyl,
e) C_{1-3} alkoxy, unsubstituted or substituted with one or more of -OH or C_{1-3} alkoxy,

REPLACEMENT SHEET
04 FEB 1997

f) -COOR,

g) $-\overset{O}{\overset{\|}{C}}NR_2$,

h) $-CH_2NR_2$,

i) $-CH_2NH\overset{O}{\overset{\|}{C}}R$,

j) -CN,

k) $-CF_3$,

l) $-NH\overset{O}{\overset{\|}{C}}R$,

m) aryl C_{1-3} alkoxy,

n) aryl,

o) $-NRSO_2R$,

p) $-OP(O)(OR_x)_2$, or

q) $-R^5$, as defined below; or

R^1 and R^2 can be joined together to form with the nitrogen to which R^1 is attached a 3 to 10 membered monocyclic or bicyclic saturated ring system which consists of the nitrogen to which R^1 is attached and from 2 to 9 carbon atoms, and is unsubstituted or substituted with

1) hydroxy,

2) C_{1-4} alkyl unsubstituted or substituted with one or more of

a) halo,

b) hydroxy,

c) C_{1-3} alkoxy,

d) aryl,

REPLACEMENT SHEET
04 FEB 1997

e) a 5-7 membered cycloalkyl group unsubstituted or substituted with one or more of
   i) halo,
   ii) hydroxy,
   iii) C_{1-3} alkoxy, or
   iv) aryl,
f) heterocycle, or
g) $-NR_2$,

3) C_{1-3} alkoxy,

4) $-NH-\overset{O}{\overset{\|}{C}}OC_{1-3}alkyl$,

5) $-NH-\overset{O}{\overset{\|}{C}}-C_{1-3}alkyl$,

6) $-NH-SO_2C_{1-3}alkyl$,

7) heterocycle,

8) -W-aryl, or

9) $-W-\overset{O}{\overset{\|}{C}}-aryl$,

wherein W is defined above; or

R^1 and R^2 can be joined together to form with the nitrogen to which R^1 is attached a 3 to 10 membered monocyclic or bicyclic saturated ring system which consists of the nitrogen to which R^1 is attached, from 1 to 8 carbon atoms and one or more unsubstituted or substituted heteroatom selected from

1) $-\overset{|}{\underset{V-R^1}{N}}-$,

wherein V is absent or $-\overset{O}{\overset{\|}{C}}-Q-$ or $-SO_2-Q-$,

REPLACEMENT SHEET
04 FEB 1997

R^1 is defined as above for when R^1 is independent from and not joined to R^2, and wherein Q is absent or -O-, -NR-, or heterocycle optionally substituted with

$-C_{1-4}$alkyl,

2) -N- (heterocycle),

3) -N- (C_{1-4} alkenyl, unsubstituted or substituted with aryl,

4) -N- (SO_2-C_{1-4}alkenyl, unsubstituted or substituted with aryl,

5) $-S(O)_p-$, wherein p is zero, 1 or 2, or

6) -O-; or

R^1 and R^2 can be joined together to form with the nitrogen to which R^1 is attached a 3 to 10 membered monocyclic or bicyclic saturated ring system, which consists of the nitrogen to which R^1 is attached and from 2 to 9 carbon atoms, in which the saturated ring system is fused to a phenyl ring and the phenyl ring is unsubstituted or substituted with one or more of

1) halo,

2) C_{1-3} alkoxy,

3) hydroxy,

4) C_{1-4} alkyl,

5) $-NHR^1$, wherein R^1 is defined as above for when R^1 is independent from and not joined to R^2, or

6) -NH-heterocycle;

REPLACEMENT SHEET
04 FEB 1997

R^3 is

1) $-(CH_2)_r-R^4$,
   wherein r is zero through 5,
2) C_{1-4}alkenyl-R^4, or
3) C_{1-4}alkynyl-R^4;

R^4 is

1) hydrogen,
2) C_{1-4} alkyl,
3) C_5-C_{10} cycloalkyl, optionally substituted with hydroxy,
4) C_6-C_{10} aryl, unsubstituted or substituted with one or more of
   a) halo,
   b) hydroxy,
   c) $-NO_2$ or $-NR_2$,
   d) C_{1-4}alkyl,
   e) C_{1-3} alkoxy, unsubstituted or substituted with one or more of -OH or C_{1-3} alkoxy,
   f) -COOR,
   g) $-\overset{\overset{O}{\|}}{C}NR_2$,
   h) $-CH_2NR_2$,
   i) $-CH_2NH\overset{\overset{O}{\|}}{C}R$,
   j) -CN,
   k) $-CF_3$,
   l) $-NH\overset{\overset{O}{\|}}{C}R$,
   m) aryl C_{1-3} alkoxy,
   n) aryl,

REPLACEMENT SHEET
04 FEB 1997

o) $-NRSO_2R$,
p) $-OP(O)(OR_x)_2$, or
q) $-R^5$, as defined below, or

5) monocyclic or bicyclic heterocyle containing from 1 to 3 heteroatoms chosen from the group consisting of N, O, and S and which is unsubstituted or substituted with R^5 and optionally with one or more of
   a) halo,
   b) C_{1-4} alkyl, or
   c) C_{1-3} alkoxy;

R_x is H or aryl;

R^5 is


1) $-W-(CH_2)_m-NR^6R^7$
wherein W is as defined above,
m is 2-5, and
R^6 and R^7 are independently
   a) hydrogen,
   b) C_{1-6} alkyl, unsubstituted or substituted with one or more of
      i) C_{1-3} alkoxy,
      ii) -OH, or
      iii) $-NR_2$,
   c) the same or different and joined together to form a 5-7 member heterocycle, such as morpholino, containing up to two additional heteroatoms selected from
      $-\overset{R}{\underset{|}{N}}-$, -O-, $-\overset{O}{\underset{\|}{S}}-$, -S-, or $-SO_2-$, the heterocycle optionally substituted with C_{1-4} alkyl, or

REPLACEMENT SHEET
04 FEB 1997

d) aromatic heterocycle unsubstituted or substituted with one or more of
i) C_{1-4} alkyl, or
ii) $-NR_2$,

2) $-(CH_2)_q-NR^6R^7$ wherein q is 1-5, and R^6 and R^7 are defined above, except that R^6 or R^7 are not H or unsubstituted C_{1-6} alkyl, or

3) benzofuryl, indolyl, azacycloalkyl, azabicyclo C_{7-11} cycloalkyl, or benzopiperidinyl, unsubstituted or substituted with C_{1-4} alkyl;

B is absent, or ,

wherein R^8 is
1) $-CH(CH_3)_2$,
2) $-CH(CH_3)(CH_2CH_3)$, or
3) -phenyl;

J^1 and J^2 are independently
1) $-YR^9$ wherein
Y is -O- or -NH-, and
R^9 is
a) hydrogen,
b) C_{1-6} alkyl, unsubstituted or substituted with one or more of
i) $-NR_2$,
ii) -OR,
iii) $-NHSO_2C_{1-4}$ alkyl,

REPLACEMENT SHEET
04 FEB 1997

iv) $-NHSO_2$ aryl, or $-NHSO_2$(dialkyl-aminoaryl),

v) $-CH_2OR$,

vi) $-C_{1-4}$ alkyl,

vii) $-\overset{O}{\overset{\|}{C}}OR$,

viii) $-\overset{O}{\overset{\|}{C}}NR_2$,

ix) $-NH\!-\!\overset{NH}{\overset{\|}{C}}\!-\!NR_2$ or $-NH\!-\!\overset{N\text{-}CN}{\overset{\|}{C}}\!-\!NR_2$,

x) $-NH\overset{O}{\overset{\|}{C}}R^{13}$, wherein R^{13} is

A) -H,

B) $-C_{1-4}$ alkyl,

C) -aryl,

D) -heterocycle, or

E) -NH-, -O- or $-(CH_2)_n-$ wherein n is zero, 1, 2 or 3, substituted with

I) $-C_{1-4}$ alkyl, unsubstituted or substituted with one or more of aryl or heterocycle, or

II) aryl, unsubstituted or substituted with heterocycle,

xi) $-NR_3^{\oplus}$ $A^{\ominus}$ wherein $A^{\ominus}$ is a counterion,

REPLACEMENT SHEET
04 FEB 1997

xii) $-NR^{10}R^{11}$ wherein R^{10} and R^{11} are the same or different and are C_{1-5} alkyl joined together directly to form a 5-7 membered heterocycle containing up to one additional heteroatom selected from -O-, -S-, or -NR-,

xiii) aryl,

xiv) -CHO,

xv) $-OP(O)(OR_x)_2$,

xvi) $-O-\overset{O}{\overset{\|}{C}}-C_{1-4}$alkyl substituted with one or more of amine or quaternary amine, or $-O-((CH_2)_mO)_n-R$, or $-OP(O)(OR_x)_2$,


xvii) $-O\overset{O}{\overset{\|}{C}}-R$, or

xviii) $-O\overset{O}{\overset{\|}{C}}-NH-CH_2$-heterocycle, or

c) $-((CH_2)_mO)_nCH_3$ or $-((CH_2)_mO)_n$ H, wherein m and n are defined above,

2) $-N(R^9)_2$,

3) $-NR^{10}R^{11}$ wherein R^{10} and R^{11} are defined above, or

4) 

wherein Y, R^9 and n are defined above; and

REPLACEMENT SHEET
04 FEB 1997

R^{12} is

1) hydrogen,

2) aryl, unsubstituted or substituted with one or more of

a) R^{14}, wherein R^{14} is

i) halo,

ii) -OR,

iii) $-\overset{O}{\overset{\|}{C}}NR_2$,

iv) $-CH_2NR_2$,

v) $-SO_2NR_2$,

vi) $-NR_2$,

vii) $-NH\overset{O}{\overset{\|}{C}}R$,

viii) C_{1-4} alkyl,

ix) phenyl

x) $-CF_3$,

xi) $-\overset{R}{\overset{|}{N}}-SO_2R$,

xii) $-OP(O)(OR_x)_2$, or

xiii) $-C\overset{O}{\overset{\|}{O}}R$,

b) $-C_{1-4}$ alkyl$-NR_2$, or

c) $-O-\overset{O}{\overset{\|}{C}}-C_{1-4}$alkyl substituted with one or more of amine or quaternary amine or $-OP(O)(OR_x)_2$,

REPLACEMENT SHEET
04 FEB 1997

3) heterocycle, such as isochroman, chroman, isothiochroman, thiochroman, benzimidazole, benzothiopyran, oxobenzothiopyran, benzopyran, benzothiopyranylsulfone, benzothiopyranylsulfoxide, the ring or rings being unsubstituted or substituted with one or more of
a) R^{14}, as defined above,
b) $-OC_{1-4}$ alkenyl,
c) phenyl-C_{1-4} alkyl,
d) $-O-\overset{O}{\overset{\|}{C}}-C_{1-4}$alkyl substituted with one or more of amine or quaternary amine, or $-OP(O)(OR_x)_2$, or $-O((CH_2)_mO)_n-R$, or
e) $-O-\overset{O}{\overset{\|}{C}}-O-((CH_2)_mO)_n-R$, or

4) A 5 to 7 membered carbocyclic or 7-10 membered bicyclic carbocyclic ring, such as cyclopentane, cyclohexane, indane, norbornane, naphthalene, thiopyran, isothiopyran, or benzopyran, the carbocyclic ring being unsubstituted or substituted with one or more of
a) R^{14}, as defined above,
b) $-CH_2OR$,
c) $-(CH_2)_n-NR_2$, C_{5-16}alkyl, pyridine, $-(CH_2)_nNR-(CH_2)_n-NR_2$, $-(CH_2)_n-\overset{O}{\overset{\|}{C}}-OR$,

REPLACEMENT SHEET
04 FEB 1997

$-((CH_2)_mO)_n-R$, quinuclidiniumyl substituted with R, piperazine-C_{1-4}alkyl-benzyl substituted once or more with R, or morpholino-C_{1-4}alkyl-benzyl,

d) $-O-\overset{O}{\overset{\|}{C}}-C_{1-4}$alkyl substituted with one or more of amine or quaternary amine, $-OP(O)(OR_x)_2$, or $-O-((CH_2)_mO)_n-R$,

e) $-O-\overset{O}{\overset{\|}{C}}-O-((CH_2)_mO)_n-R$, or

f) $-C_{1-4}$alkyl-phenyl;

or a pharmaceutically acceptable salt thereof.

In a preferred embodiment of this invention, R^1 and R^2 are joined together to form with the nitrogen to which R^1 is attached a 3 to 10 membered monocyclic or bicyclic saturated ring system which consists of the nitrogen to which R^1 is attached and from 2 to 9 carbon atoms, and is unsubstituted or substituted with

1) hydroxy,
2) C_{1-4} alkyl unsubstituted or substituted with one or more of
   a) hydroxy,
   b) C_{1-3} alkoxy,
   c) aryl,
   d) a 5-7 membered cycloalkyl group unsubstituted or substituted with one or more of

REPLACEMENT SHEET
04 FEB 1997

i) halo,
ii) hydroxy,
iii) C_{1-3} alkoxy, or
iv) aryl,

e) heterocycle, or

f) $-NR_2$,

3) C_{1-3} alkoxy,

4) $-NH-\overset{O}{\overset{\|}{C}}OC_{1-3}alkyl$,

5) $-NH-\overset{O}{\overset{\|}{C}}-C_{1-3}alkyl$,

6) $-NH-SO_2C_{1-3}alkyl$,

7) -W-aryl, or

8) $-W-\overset{O}{\overset{\|}{C}}-aryl$,

wherein W is -O-, -S-, or -NH-; or

R^1 and R^2 are joined together to form with the nitrogen to which R^1 is attached a 3 to 10 membered monocyclic or bicyclic saturated ring system which consists of the nitrogen to which R^1 is attached, from 1 to 8 carbon atoms and one or more unsubstituted or substituted heteroatom selected from

1) $-\overset{}{\underset{|}{N}}-$
$V-R^1$,

wherein V is absent or $-\overset{O}{\overset{\|}{C}}-Q-$ or $-SO_2-Q-$, R^1 is defined as above for when R^1 is independent from and not joined to R^2, and wherein Q is absent or -O-, -NR-, or heterocycle optionally substituted with $-C_{1-4}alkyl$,

REPLACEMENT SHEET
04 FEB 1997

2) $-\underset{\underset{1}{}}{N}-$
C_{1-4} alkenyl, unsubstituted or substituted with aryl,

3) $-S(O)_p-$,
wherein p is zero, 1 or 2, or

4) $-O-$; or

R^1 and R^2 are joined together to form with the nitrogen to which R^1 is attached a 3 to 10 membered monocyclic or bicyclic saturated ring system, which consists of the nitrogen to which R^1 is attached and from 2 to 9 carbon atoms, in which the saturated ring system is fused to a phenyl ring and the phenyl ring is unsubstituted or substituted with one or more of

1) C_{1-3} alkoxy,
2) hydroxy,
3) C_{1-4} alkyl, or
4) $-NHR^1$,
wherein R^1 is defined as above for when R^1 is independent from and not joined to R^2.

A second, more preferred embodiment of this invention is further limited to compounds where:

R^1 and R^2 are joined together to form with the nitrogen to which R^1 is attached a 3 to 10 membered monocyclic or bicyclic saturated ring system which consists of the nitrogen to which R^1 is attached and from 2 to 9 carbon atoms, and is unsubstituted or substituted with

REPLACEMENT SHEET
04 FEB 1997

1) hydroxy,
2) C_{1-4} alkyl unsubstituted or substituted with one or more of
   a) hydroxy,
   b) C_{1-3} alkoxy,
   c) aryl,
   d) a 5-7 membered cycloalkyl group unsubstituted or substituted with one or more of
      i) halo,
      ii) hydroxy,
      iii) C_{1-3} alkoxy, or
      iv) aryl,
   e) heterocycle, or
   f) $-NR_2$,
3) C_{1-3} alkoxy,
4) $-NH-\overset{O}{\overset{\|}{C}}OC_{1-3}alkyl$,
5) $-NH-\overset{O}{\overset{\|}{C}}-C_{1-3}alkyl$,
6) $-NH-SO_2C_{1-3}alkyl$,
7) -W-aryl, or
8) $-W-\overset{O}{\overset{\|}{C}}-aryl$,

wherein W is -O-, -S-, or -NH-; or

R^1 and R^2 are joined together to form with the nitrogen to which R^1 is attached a 3 to 10 membered monocyclic or bicyclic saturated ring system which consists of the nitrogen to which R^1 is attached, from 1 to 8 carbon atoms and one or more unsubstituted or substituted heteroatom selected from

REPLACEMENT SHEET
04 FEB 1997

1) $-N(V\text{-}R^1)-$,

wherein V is absent or $-\overset{O}{\overset{\|}{C}}-Q-$ or $-SO_2-Q-$, R^1 is defined as above for when R^1 is independent from and not joined to R^2, and wherein Q is absent or -O-, -NR-, or heterocycle optionally substituted with $-C_{1-4}$alkyl,

2) $-S(O)_p-$,
wherein p is zero, 1 or 2, or

3) -O-;

R^3 is benzyl, unsubstituted or substituted with one or more of

a) hydroxy,
b) $-NO_2$, or $-NR_2$,
c) C_{1-4}alkyl,
d) C_{1-3} alkoxy, unsubstituted or substituted with one or more of -OH or C_{1-3} alkoxy,
e) $-\overset{O}{\overset{\|}{C}}NR_2$,
f) $-CH_2NR_2$,
g) $-CH_2NH\overset{O}{\overset{\|}{C}}R$,
h) $-CF_3$,
i) $-NH\overset{O}{\overset{\|}{C}}R$,
j) $-NRSO_2R$,
k) $-OP(O)(OR_x)_2$, or
l) $-R^5$;

REPLACEMENT SHEET
04 FEB 1997

and B is absent.

A third, most preferred embodiment of this invention is further limited to compounds where:

X is -OH;

Z is -O;

R^1 and R^2 are joined together to form with the nitrogen to which R^1 is attached a 3 to 10 membered monocyclic or bicyclic saturated ring system which consists of the nitrogen to which R^1 is attached and from 2 to 9 carbon atoms, and is unsubstituted or substituted with -W-aryl or -W-C(=O)-aryl; or


R^1 and R^2 are joined together to form with the nitrogen to which R^1 is attached a 3 to 10 membered monocyclic or bicyclic saturated ring system which consists of the nitrogen to which R^1 is attached, from 1 to 8 carbon atoms and one of -N(V-R^1)- ,

wherein V is absent or -C(=O)-Q- or -SO_2-Q-,

R^1 is defined as above for when R^1 is independent from and not joined to R^2,


and wherein Q is absent or -O-, -NR- or heterocycle optionally substituted with -C_{1-4}alkyl;

R^3 is benzyl, unsubstituted or substituted with one or more of (1) hydroxy, (2) C_{1-3} alkoxy substituted with one or more of -OH or (3)




REPLACEMENT SHEET
04 FEB 1997

J^1 is $-NH-C_{1-4}$alkyl; and
J^2 is



The most preferred compounds of this invention are compounds A through H and J, shown below.


Compound A:



N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(2-(3(S)-N'-(t-butylcarboxamido)-(4aS,8aS)-decahydroisoquinoline)yl)-pentaneamide,


REPLACEMENT SHEET
04 FEB 1997

Compound B:



N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(1-(4-carbobenzyloxy-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide,


Compound C:



N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-(2-(4-morpholinyl)ethoxy)phenyl)methyl)-4(S)-hydroxy-5-(2-(3(S)-N'-(t-butylcarboxamido)-(4aS,8aS)-decahydroisoquinoline)yl)-pentaneamide,


REPLACEMENT SHEET
04 FEB 1997

Compound D:



N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-(2-(4-morpholinyl)ethoxy)phenyl)methyl)-4(S)-hydroxy-5-(1-(4-carbobenzyloxy-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide,


Compound E:



N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-((2-hydroxy)-ethoxy)phenyl)methyl)-4(S)-hydroxy-5-(2-(3(S)-N'-(t-butylcarboxamido)-(4aS,8aS)-decahydroisoquinoline)-yl)-pentaneamide,


REPLACEMENT SHEET
04 FEB 1997

Compound F:



N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-((2-hydroxy)-ethoxy)phenyl)methyl)-4(S)-hydroxy-5-(1-(4-carbobenzyloxy-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide,


Compound G:



N-(4(S)-3,4-dihydro-1H-2,2-dioxobenzothiopyranyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(2-(3(S)-N'-(t-butyl-carboxamido)-(4aS,8aS)-decahydroisoquinoline)yl)-pentaneamide,


REPLACEMENT SHEET
04 FEB 1997

Compound H:



N-(4(S)-3,4-dihydro-1H-2,2-dioxobenzothiopyranyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(1-(4-carbobenzyloxy-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide,

Compound J:



N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(1-(4-(3-pyridylmethyl)-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide.

Novel compounds of the present invention also include but are not limited to the following compounds:

REPLACEMENT SHEET
04 FEB 1997

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4-(S)-hydroxy-5-(1-(N'-(t-butyl)-4(S)-phenoxyproline-amid)yl)-pentaneamide,

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4-(S)-hydroxy-5-(1-(N'-t-butyl-4(S)-2-naphthyloxy-prolineamid)yl)-pentaneamide,

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4-(S)-hydroxy-5-(1-(N'-t-butyl-4(S)-1-naphthyloxy-prolineamid)yl)-pentaneamide,

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4-(S)-amino-5-(2-(3(S)-N'-(t-butylcarboxamido)-(4aS,8aS)-decahydroisoquinoline)yl)-pentaneamide,

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4-(S)-hydroxy-5-(1-(4-(3-phenylpropionyl)-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide,

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4-(S)-hydroxy-5-(1-(4-benzoyl-2(S)-N'-(t-butylcarbox-amido)-piperazinyl))-pentaneamide,

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4-(S)-hydroxy-5-(1-(4-(3-phenylpropyl)-2(S)-N'-(t-butyl-carboxamido)-piperazinyl))-pentaneamide,

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4-(S)-amino-5-(1-(4-carbobenzyloxy-2(S)-N'-(t-butyl-carboxamido)-piperazinyl))-pentaneamide,

REPLACEMENT SHEET
04 FEB 1997

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-(2-(4-morpholinyl)ethoxy)phenyl)methyl)-4(S)-hydroxy-5-(1-(N'-(t-butyl)-4(S)-phenoxyprolineamid)yl)-pentaneamide,

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-(2-(4-morpholinyl)ethoxy)phenyl)methyl)-4(S)-hydroxy-5-(1-(N'-t-butyl-4(S)-2-naphthyloxy-prolineamid)yl)-pentaneamide,

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-(2-(4-morpholinyl)ethoxy)phenyl)methyl)-4(S)-hydroxy-5-(1-(N'-t-butyl-4(S)-1-naphthyloxy-prolineamid)yl)-pentaneamide,

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-(2-(4-morpholinyl)ethoxy)phenyl)methyl)-4(S)-amino-5-(2-(3(S)-N'-(t-butylcarboxamido)-(4aS,8aS)-decahydroisoquinoline)yl)-pentaneamide,

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-(2-(4-morpholinyl)ethoxy)phenyl)methyl)-4(S)-hydroxy-5-(1-(4-(3-phenylpropionyl)-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide,

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-(2-(4-morpholinyl)ethoxy)phenyl)methyl)-4(S)-hydroxy-5-(1-(4-benzoyl-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide,

REPLACEMENT SHEET
04 FEB 1997

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-(2-(4-morpholinyl)ethoxy)phenyl)methyl)-4(S)-hydroxy-5-(1-(4-(3-phenylpropyl)-2(S)-N'-(t-butylcarboxamido))-piperazinyl)-pentaneamide,

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-(2-(4-morpholinyl)ethoxy)phenyl)methyl)-4(S)-amino-5-(1-(4-carbobenzyloxy-2(S)-N'-(t-butylcarboxamido)-piperazinyl)pentaneamide,

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-((2-hydroxy)-ethoxy)phenyl)methyl)-4(S)-hydroxy-5-(1-(N'-(t-butyl)-4(S)-phenoxyprolineamid)yl)-pentaneamide,

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-((2-hydroxy)-ethoxy)phenyl)methyl)-4(S)-hydroxy-5-(1-(N'-t-butyl-4(S)-2-naphthyloxy-prolineamid)yl)-pentaneamide,

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-((2-hydroxy)-ethoxy)phenyl)methyl)-4(S)-hydroxy-5-(1-(N'-t-butyl-4(S)-1-naphthyloxy-prolineamid)yl)-pentaneamide,

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-((2-hydroxy)-ethoxy)phenyl)methyl)-4(S)-amino-5-(2-(3(S)-N'-(t-butylcarboxamido)-(4aS,8aS)-decahydroisoquinoline)-yl)pentaneamide,

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-((2-hydroxy)-ethoxy)phenyl)methyl)-4(S)-hydroxy-5-(1-(4-(3-phenyl-propionyl)-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide,

REPLACEMENT SHEET
04 FEB 1997

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-((2-hydroxy)-ethoxy)phenyl)methyl)-4(S)-hydroxy-5-(1-(4-benzoyl-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide,

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-((2-hydroxy)-ethoxy)phenyl)methyl)-4(S)-hydroxy-5-(1-(4-(3-phenyl-propyl)-2(S)-N'-(t-butylcarboxamido))-piperazinyl)-pentaneamide,

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-((2-hydroxy)-ethoxy)phenyl)methyl)-4(S)-amino-5-(1-(4-carbobenzyl-oxy-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide,

N-(4(S)-3,4-dihydro-1H-2,2-dioxobenzothiopyranyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(1-(N'-(t-butyl)-4(S)-phenoxyprolineamid)yl)-pentaneamide,

N-(4(S)-3,4-dihydro-1H-2,2-dioxobenzothiopyranyl)-2-(R)-phenylmethyl-4(S)-hydroxy-5-(1-(N'-t-butyl-4(S)-2-naphthyloxy-prolineamid)yl)-pentaneamide,

N-(4(S)-3,4-dihydro-1H-2,2-dioxobenzothiopyranyl)-2-(R)-phenylmethyl-4(S)-hydroxy-5-(1-(N'-t-butyl-4(S)-1-naphthyloxy-prolineamid)yl)-pentaneamide,

N-(4(S)-3,4-dihydro-1H-2,2-dioxobenzothiopyranyl)-2-(R)-phenylmethyl-4(S)-amino-5-(2-(3(S)-N'-(t-butyl-carboxamido)-(4aS,8aS)-decahydroisoquinoline)yl)-pentaneamide,

REPLACEMENT SHEET
04 FEB 1997

N-(4(S)-3,4-dihydro-1H-2,2-dioxobenzothiopyranyl)-2-(R)-phenylmethyl-4(S)-hydroxy-5-(1-(4-(3-phenylpropionyl)-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide,

N-(4(S)-3,4-dihydro-1H-2,2-dioxobenzothiopyranyl)-2-(R)-phenylmethyl-4(S)-hydroxy-5-(1-(4-benzoyl-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide,

N-(4(S)-3,4-dihydro-1H-2,2-dioxobenzothiopyranyl)-2-(R)-phenylmethyl-4(S)-hydroxy-5-(1-(4-(3-phenylpropyl)-2(S)-N'-(t-butylcarboxamido))-piperazinyl)-pentaneamide, or

(4(S)-3,4-dihydro-1H-2,2-dioxobenzothiopyranyl)-2(R)-phenylmethyl-4(S)-amino-5-(1-(4-carbobenzyloxy-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide.

The compounds of the present invention, may have asymmetric centers and occur as racemates, racemic mixtures and as individual diastereomers, or enantiomers with all isomeric forms being included in the present invention.

When any variable (e.g., aryl, heterocycle, R, R^1, R^2, A^-, n, Z, etc.) occurs more than one time in any constituent or in formula I, its definition on each occurrence is independent of its definition at every other occurrence. Also, combinations of substituents and/or variables are permissible only if such combinations result in stable compounds.

REPLACEMENT SHEET
04 FEB 1997

As used herein except where noted, "alkyl" is intended to include both branched- and straight-chain saturated aliphatic hydrocarbon groups having the specified number of carbon atoms (Me is methyl, Et is ethyl, Pr is propyl, Bu is butyl); "alkoxy" represents an alkyl group of indicated number of carbon atoms attached through an oxygen bridge; and "cycloalkyl" is intended to include saturated ring groups, such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl (Cyh) and cycloheptyl. "Alkenyl" is intended to include hydrocarbon groups of either a straight or branched configuration with one or more carbon-carbon double bonds which may occur in any stable point along the chain, such as ethenyl, propenyl, butenyl, pentenyl, and the like. "Alkynyl" is intended to include hydrocarbon groups of either a straight or branched configuration with one or more carbon-carbon triple bonds which may occur in any stable point along the chain, such as ethynyl, propynyl, butynyl, pentynyl, and the like. "Halo", as used herein, means fluoro, chloro, bromo and iodo; and "counterion" is used to represent a small, single negatively-charged species, such as chloride, bromide, hydroxide, acetate, trifluroacetate, perchlorate, nitrate, benzoate, maleate, tartrate, hemitartrate, benzene sulfonate, and the like.

As used herein, with exceptions as noted, "aryl" is intended to mean phenyl (Ph) or naphthyl. "Carbocyclic" is intended to mean any stable 5- to 7-membered carbon ring or 7- to 10-membered bicyclic carbon ring any ring of which may be saturated or unsaturated.

REPLACEMENT SHEET
04 FEB 1997

The term heterocycle or heterocyclic, as used herein except where noted, represents a stable 5- to 7-membered mono- or bicyclic or stable 7- to 10-membered bicyclic heterocyclic ring system any ring of which may be saturated or unsaturated, and which consists of carbon atoms and from one to three heteroatoms selected from the group consisting of N, O and S, and wherein the nitrogen and sulfur heteroatoms may optionally be oxidized, and the nitrogen heteroatom may optionally be quaternized, and including any bicyclic group in which any of the above-defined heterocyclic rings is fused to a benzene ring. The heterocyclic ring may be attached at any heteroatom or carbon atom which results in the creation of a stable structure. Examples of such heterocyclic elements include piperidinyl, piperazinyl, 2-oxopiperazinyl, 2-oxopiperidinyl, 2-oxopyrrolodinyl, 2-oxoazepinyl, azepinyl, pyrrolyl, 4-piperidonyl, pyrrolidinyl, pyrazolyl, pyrazolidinyl, imidazolyl, imidazolinyl, imidazolidinyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, oxazolyl, oxazolidinyl, isoxazolyl, isoxazolidinyl, morpholinyl, thiazolyl, thiazolidinyl, isothiazolyl, quinuclidinyl, isothiazolidinyl, indolyl, quinolinyl, isoquinolinyl, benzimidazolyl, thiadiazoyl, benzopyranyl, benzothiazolyl, benzoxazolyl, furyl, tetrahydrofuryl, tetrahydropyranyl, thienyl, benzothienyl, thiamorpholinyl, thiamorpholinyl sulfoxide, thiamorpholinyl sulfone, and oxadiazolyl. Morpholino is the same as morpholinyl.

REPLACEMENT SHEET
04 FEB 1997

The pharmaceutically-acceptable salts of the compounds of Formula I (in the form of water- or oil-soluble or dispersible products) include the conventional non-toxic salts or the quaternary ammonium salts which are formed, e.g., from inorganic or organic acids or bases. Examples of such acid addition salts include acetate, adipate, alginate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, citrate, camphorate, camphorsulfonate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, fumarate, glucoheptanoate, glycerophosphate, hemisulfate, heptanoate, hexanoate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethanesulfonate, lactate, maleate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, oxalate, pamoate, pectinate, persulfate, 3-phenyl-propionate, picrate, pivalate, propionate, succinate, tartrate, thiocyanate, tosylate, and undecanoate. Base salts include ammonium salts, alkali metal salts such as sodium and potassium salts, alkaline earth metal salts such as calcium and magnesium salts, salts with organic bases such as dicyclohexylamine salts, N-methyl-D-glucamine, and salts with amino acids such as arginine, lysine, and so forth. Also, the basic nitrogen-containing groups may be quaternized with such agents as lower alkyl halides, such as methyl, ethyl, propyl, and butyl chloride, bromides and iodides; dialkyl sulfates like dimethyl, diethyl, dibutyl; and diamyl sulfates, long chain halides such as decyl, lauryl, myristyl and stearyl chlorides, bromides and iodides, aralkyl halides like benzyl and phenethyl bromides and others.

REPLACEMENT SHEET
04 FEB 1997

Schemes I-III for preparing the novel compounds of this invention are presented below. Tables I and II which follow the schemes illustrate the compounds that can be synthesized by Schemes I-III, but Schemes I-III are not limited by the compounds in the tables nor by any particular substituents employed in the schemes for illustrative purposes. The examples specifically illustrate the application of the following schemes to specific compounds.

Amide couplings used to form the compounds of this invention are typically performed by the carbodiimide method with reagents such as dicyclohexylcarbodiimide, or 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide. Other methods of forming the amide or peptide bond include, but are not limited to the synthetic routes via an acid chloride, azide, mixed anhydride or activated ester. Typically, solution phase amide coupling are performed, but solid-phase synthesis by classical Merrifield techniques may be employed instead. The addition and removal of one or more protecting groups is also typical practice.

Additional related information on synthetic background is contained in EPO 0337714.

One method for producing formula I compounds is provided by Scheme I. Dihydro-5(S)-(tert-butyldimethylsilyloxymethyl)-3(2H)-furanone (compound 1 below) is prepared by standard methods known in the art from commercially available dihydro-5(S)-(hydroxymethyl)-2(3H)-furanone. After alkylation of compound 1 to form compound 2, the protecting group of lactone 2 is removed with aqueous HF to afford compound 3.

REPLACEMENT SHEET
04 FEB 1997


The alcohol group of 3 is activated by conversion into a leaving group such as mesylate, tosylate or trifylate by treating the alcohol with a sulfonyl chloride or sulfonic anhydride, such as trifluoromethanesulfonic anhydride, in the presence of a hindered amine base such as triethylamine, diethyl isopropylamine or 2,6 lutidine, to afford a compound such as compound 4. The leaving group of compound 4 is displaced by an amine 5, such as N'-t-butyl-(4aS,8aS)-(decahydroisoquinoline)-3(S)-carboxamide, in a high boiling solvent such as DMF or xylene to produce a compound such as 6. A trifluoromethanesulfonyloxy group can be displaced by an amine at room temperature in a solvent such as isopropanol by treatment with N,N-diisopropylethylamine.

Compound 6 is hydrolyzed with aqueous lithium or sodium hydroxide and the resultant hydroxy acid 7 is converted into a protected hydroxy acid 8. The hydroxyl group is conveniently protected with a standard silyl protecting group such as t-butyldimethyl silyl or t-butyldiphenyl silyl.

The protected hydroxy-acid 8 is then coupled to the desired R^{12} amine to produce compound 9, and the silyl protecting group is removed with fluoride ion to arrive at compound 10.


REPLACEMENT SHEET
04 FEB 1997

SCHEME I




REPLACEMENT SHEET
04 FEB 1997

SCHEME I (CONT'D)



9



10

A second method for forming products of general formula I is shown in Scheme II. In Scheme II, alkylation of 11 is performed by a first step of deprotonation of 11 with n-butyllithium or lithium diisopropylamide (LDA) followed by a second step of adding an alkenyl halide (such as allyl bromide) to afford 12.

Dihydroxylation of the olefin of 12 with osmium tetroxide and N-methylmorpholine-N-oxide (NMO) produces a diasteriomeric mixture of diols, 13. Selective mesylation of the primary alcohol of 13 with methanesulfonyl chloride and either triethylamine or pyridine gives a mesylate 14.


REPLACEMENT SHEET
04 FEB 1997

Heating mesylate 14 with an amine in a refluxing alcoholic solvent such as methanol or isopropanol which contains an excess of potassium carbonate produces an amino alcohol such as compound 15. The diasteriomers can be separated at this step by standard techniques well known to those of skill in the art. Alternatively, the separation can be done after removal of the ketal.

Removal of the ketal in compound 15 is accomplished by treatment with acid in the presence of methanol, or by aqueous acid or by 1N HCl in THF, to form compound 16.

REPLACEMENT SHEET
04 FEB 1997

SCHEME II



11 12 13 14 15 16


REPLACEMENT SHEET
04 FEB 1997

A third method for forming products of general formula I is shown in Scheme III. Protection of the pyrrolidine -NH- group of compound 17 is carried out with BOC-anhydride and dimethylaminopyridine to give the protected compound 18. Alkylation of 18 is performed by a first step of deprotonation of 18 with a strong base such as lithium hexamethyldisilamide (LHMDS) or lithium diisopropylamide (LDA) followed by a second step of adding an alkyl halide (such as benzyl bromide) to afford compound 19.

The TBS protecting and BOC protecting group of 19 are removed by treatment with aqueous HF in acetonitrile to give alcohol 20. Mesylation of the primary alcohol of 20 with methanesulfonyl chloride and either triethylamine or pyridine gives mesylate 21 which is heated with an amine in a refluxing alcoholic solvent such as methanol or isopropanol which contains an excess of potassium carbonate to produce an amino pyrrolidinone such as compound 22. The pyrrolidine -NH- group of 22 is reprotected as a BOC group as before and the resultant compound 23 is hydrolized open with a base such as lithium or sodium hydroxide to afford the acid 24. Compound 24 is then coupled to an NH_2R^{12} amine in a standard manner and the BOC is removed with gaseous HCl or trifluoro-acetic acid to give the desired product, exemplified by compound 25.


REPLACEMENT SHEET
04 FEB 1997

SCHEME III




REPLACEMENT SHEET
04 FEB 1997

SCHEME III (CONT'D)




24



25


A compound of formula 26



26

wherein P is a nitrogen protecting group such as -BOC or -CBZ, is preferably prepared according to the method described in Scheme I, preferably employing the 5-trifluoromethanesulfonyloxymethyl analog of lactone 4 therein (see Example 15, Step 1).


Compounds of formula 27



27

REPLACEMENT SHEET
04 FEB 1997

can be obtained by a variety of routes from compound 28



28


which is obtained after removal of the nitrogen protecting group in 26 using methods well known in the art, e.g., catalytic hydrogenation to remove a CBZ group, or treatment with trimethylsilyltriflate and 2,6 lutidine at about 0°C in a solvent such as CH_2Cl_2 to remove a BOC group.






For example, the 4-position piperazinyl nitrogen of compound 28 can be alkylated with a compound of formula R^1-X in a solvent such as DMF in the presence of Et_3N at room temperature, wherein X is -Cl, Br or -I, or a sulfonamide group can be formed by treatment of 28 with a sulfonyl chloride compound of formula R^1SO_2Cl under similar conditions. Also, standard amide coupling techniques can be used to form an amide group at the piperazinyl 4-position. Techniques for these procedures are well known to those skilled in the art. The R^1 group of R^1-X or R^1SO_2Cl is defined above in the definition of compounds of formula I wherein R^1 is independent from and not joined to R^2, except that R^1 can not be hydrogen or a group with a free hydroxy substituent, such as -C_{1-4}alkyl substitued with hydroxy, with the further exception that R^1 can be aryl substituted with a hydroxy group.

The compounds of this invention are also illustrated by Tables I-IV, which follow.

REPLACEMENT SHEET
04 FEB 1997






TABLE I



R^3	X	D
$-CH_2-Ph$	-OH	
$-CH_2-Ph$	-OH	
$-CH_2-Ph$	-OH	
$-CH_2-Ph$	-OH	
$-CH_2-Ph$	-OH	






REPLACEMENT SHEET
04 FEB 1997

TABLE I. CONT'D

R^3	X	D
$-CH_2-Ph$	-OH	
$-CH_2-Ph$	-OH	
$-CH_2-Ph$	-OH	
$-CH_2-Ph$	-OH	
$-CH_2-Ph$	-OH	





REPLACEMENT SHEET
04 FEB 1997

TABLE I. CONT'D

R^3	X	D
$-CH_2-Ph$	-OH	
$-CH_2-Ph$	-OH	
$-CH_2-Ph$	-OH	
$-CH_2-Ph$	-OH	
$-CH_2-Ph$	-OH	











REPLACEMENT SHEET
04 FEB 1997

TABLE I, CONT'D

R^3	X	D
$-CH_2-Ph$	-OH	
$-CH_2-Ph$	$-NH_2$	
$-CH_2-Ph$	-OH	
$-CH_2-Ph$	-OH	















REPLACEMENT SHEET
04 FEB 1997

TABLE I. CONT'D

R^3	X	D
$-CH_2-Ph$	$-NH_2$	
$-CH_2-Ph$	$-OH$	
$-CH_2-Ph$	$-OH$	
$-CH_2-Ph$	$-OH$	
	$-OH$	
	$-OH$	
	$-OH$	











REPLACEMENT SHEET
04 FEB 1997

TABLE I. CONT'D

R^3	X	D
	$-NH_2$	
	$-NH_2$	
	-OH	
	-OH	
	$-NH_2$	
	-OH	
	-OH	















REPLACEMENT SHEET
04 FEB 1997

TABLE I. CONT'D

R^3	X	D
	$-NH_2$	
	-OH	
	-OH	
	-OH	
	$-NH_2$	
	-OH	







REPLACEMENT SHEET
04 FEB 1997

TABLE I. CONT'D

	R^3	X	D
5		-OH	
10		$-NH_2$	
15		-OH	
20		-OH	
		$-NH_2$	
25		$-NH_2$	
30		-OH	


REPLACEMENT SHEET
04 FEB 1997





TABLE I. CONT'D

R^3	X	D
$-CH_2CH=CH-Ph$	-OH	
$-CH_2CH=CH-Ph$	-OH	
$-CH_2CH=CH$ 	-OH	
$-CH_2CH=CH$ 	-OH	

REPLACEMENT SHEET
04 FEB 1997









TABLE II



A	X	D
	-OH	
	-OH	
	$-NH_2$	















REPLACEMENT SHEET
04 FEB 1997

TABLE II Cont'd

A	X	D
	$-NH_2$	
	-OH	
	-OH	
	-OH	









REPLACEMENT SHEET
04 FEB 1997

TABLE II Cont'd

A	X	D
	-OH	
	$-NH_2$	
	-OH	
	-OH	
	-OH	
	-OH	
	-OH	







REPLACEMENT SHEET
04 FEB 1997

TABLE II CONT'D

A	X	D
	-OH	
	-OH	
	-OH	
	-OH	











REPLACED
04 FEB 1997

TABLE II CONT'D

A	X	D
	-OH	
	-OH	
	-OH	





REPLACEMENT SHEET
04 FEB 1997

TABLE II, CONT'D

A	X	D
	-OH	
	-OH	
	-OH	
	-OH	
	-OH	


REPLACEMENT SHEET
04 FEB 1997









TABLE II CONT'D

A	X	D
	-OH	
	-OH	

REPLACEMENT SHEET
04 FEB 1997


TABLE III






A	A, con't
	$CH_3CH_2CH_2CH_2-$
CH_3-	$(CH_3)_2CH-$
	
	
	
	

REPLACEMENT SHEET
04 FEB 1997

TABLE IV



A


$(CH_3)_3C-O-$

$Ph-CH_2CH_2-$

REPLACEMENT SHEET
04 FEB 1997

The compounds of the present invention are useful in the inhibition of HIV protease the prevention or treatment of infection by the human immunodeficiency virus (HIV) and the treatment of consequent pathological conditions such as AIDS. Treating AIDS or preventing or treating infection by HIV is defined as including, but not limited to, treating a wide range of states of HIV infection: AIDS, ARC (AIDS related complex), both symptomatic and asymptomatic, and actual or potential exposure to HIV. For example, the compounds of this invention are useful in treating infection by HIV after suspected past exposure to HIV by, e.g., blood transfusion, organ transplant, exchange of body fluids, bites, accidental needle stick, or exposure to patient blood during surgery.

For these purposes, the compounds of the present invention may be administered orally, parenterally (including subcutaneous injections, intravenous, intramuscular, intrasternal injection or infusion techniques), by inhalation spray, or rectally, in dosage unit formulations containing conventional non-toxic pharmaceutically-acceptable carriers, adjuvants and vehicles.

Thus, in accordance with the present invention there is further provided a method of treating and a pharmaceutical composition for treating HIV infection and AIDS. The treatment involves administering to a patient in need of such treatment a pharmaceutical composition comprising a pharmaceutical carrier and a therapeutically effective amount of a compound of the present invention, or a pharmaceutically acceptable salt thereof.

REPLACEMENT SHEET
04 FEB 1997

These pharmaceutical compositions may be in the form of orally-administrable suspensions or tablets; nasal sprays; sterile injectable preparations, for example, as sterile injectable aqueous or oleagenous suspensions or suppositories.

When administered orally as a suspension, these compositions are prepared according to techniques well-known in the art of pharmaceutical formulation and may contain microcrystalline cellulose for imparting bulk, alginic acid or sodium alginate as a suspending agent, methylcellulose as a viscosity enhancer, and sweetners/flavoring agents known in the art. As immediate release tablets, these compositions may contain microcrystalline cellulose, dicalcium phosphate, starch, magnesium stearate and lactose and/or other excipients, binders, extenders, disintegrants, diluents and lubricants known in the art.

When administered by nasal aerosol or inhalation, these compositions are prepared according to techniques well-known in the art of pharmaceutical formulation and may be prepared as solutions in saline, employing benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability, fluorocarbons, and/or other solubilizing or dispersing agents known in the art.

The injectable solutions or suspensions may be formulated according to known art, using suitable non-toxic, parenterally-acceptable diluents or solvents, such as mannitol, 1,3-butanediol, water, Ringer's solution or isotonic sodium chloride solution, or suitable dispersing or wetting and suspending agents, such as sterile, bland, fixed oils, including

REPLACEMENT SHEET
04 FEB 1997

synthetic mono- or diglycerides, and fatty acids, including oleic acid.

When rectally administered in the form of suppositories, these compositions may be prepared by mixing the drug with a suitable non-irritating excipient, such as cocoa butter, synthetic glyceride esters or polyethylene glycols, which are solid at ordinary temperatures, but liquidify and/or dissolve in the rectal cavity to release the drug.

Dosage levels of the order of 0.02 to 5.0 or 10.0 grams-per-day are useful in the treatment or prevention of the above-indicated conditions, with oral doses two-to-five times higher. For example, infection by HIV is effectively treated by the administration of from 10 to 50 milligrams of the compound per kilogram of body weight from one to three times per day. It will be understood, however, that the specific dose level and frequency of dosage for any particular patient may be varied and will depend upon a variety of factors including the activity of the specific compound employed, the metabolic stability and length of action of that compound, the age, body weight, general health, sex, diet, mode and time of administration, rate of excretion, drug combination, the severity of the particular condition, and the host undergoing therapy.

The present invention is also directed to combinations of the HIV protease inhibitory compounds with one or more agents useful in the treatment of AIDS. For example, the compounds of this invention may be effectively administered, whether at periods of pre-exposure and/or post-exposure, in combination with effective amounts of the AIDS antivirals,

REPLACEMENT SHEET
04 FEB 1997

immunomodulators, anti-infectives, or vaccines known to those of ordinary skill in the art.

It will be understood that the scope of combinations of the compounds of this invention with AIDS antivirals, immunomodulators, anti-infectives or vaccines include in principle any combination with any pharmaceutical composition useful for the treatment of AIDS.

Assay for Inhibition of Microbial Expressed Viral Protease

Inhibition studies of the reaction of the protease expressed in Eschericia coli with a peptide substrate [Val-Ser-Gln-Asn-(betanapthyl)Ala-Pro-Ile-Val, 0.5 mg/mL at the time the reaction is initiated] were in 50 mM Na acetate, pH 5.5, at 30°C for 1 hour. Various concentrations of inhibitor in 1.0 ul DMSO were added to 25 ul of the peptide solution in water. The reaction is initiated by the addition of 15 ul of 0.33 nM protease (0.11 ng) in a solution of 0.133 M Na acetate pH 5.5 and 0.1% bovine serum albumin. The reaction was quenched with 160 ul of 5% phosphoric acid. Products of the reaction were separated by HPLC (VYDAC wide pore 5 cm C-18 reverse phase, acetonitrile gradient, 0.1% phosphoric acid). The extent of inhibition of the reaction was determined from the peak heights of the products. HPLC of the products, independently synthesized, proved quantitation standards and confirmation of the product composition. The products of synthesis in Examples 1-7 inclusive showed IC_{50} values in the range of 1-100 nM. Compounds A, B and J showed IC_{50} values of between about 0.3 and about 6 nM.

REPLACEMENT SHEET
04 FEB 1997

EXAMPLE 1

Preparation of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(1-(N'-(t-butyl)-4(S)-phenoxyprolineamide)yl)-pentaneamide

Step 1: Preparation of N-(2(R)-hydroxy-1(S)-indanyl)-3-phenylpropaneamide

To a cold (0°C) solution of methylene chloride (30 ml) containing 2(R)-hydroxy-1(S)-aminoindane (750 mg, 5.0 mmol) and triethylamine (606 mg, 6.0 mmol) was added a solution of hydrocinnamoyl chloride (843 mg, 5.0 mmol) in 5 ml of methylene chloride. After 2 hr the reaction was poured into a separatory funnel containing 50 ml of methylene chloride and washed with 10% citric acid solution (2 x 30 ml). The organic layer was dried, filtered and concentrated to afford a white solid.

Step 2: Preparation of N-(2(R)-hydroxy-1(S)-indan-N,O-isopropylidene-yl)-3-phenyl-propaneamide

The crude white solid from step 1 above was dissolved in 50 ml of methylene chloride and 5 ml of dimethoxypropane was added followed by the addition of 100 mg of p-toluenesulfonic acid. The reaction was stirred at room temperature for 18 hr and then poured into a separatory funnel and washed with saturated $NaHCO_3$ solution (2 x 30 ml). The organic layer was dried, filtered and concentrated to afford an oil which was chromatographed (SiO_2, 40% EtOAc/Hexane) to give an oil which eventually crystallized.

REPLACEMENT SHEET
04 FEB 1997

Step 3: Preparation of N-(2(R)-hydroxy-1(S)-indan-N,O-isopropylidene-yl)-2(S)-phenylmethyl-pent-4-eneamide

To a solution of N-(2(R)-hydroxy-1(S)-indan-N,O-isopropylidene-yl)-3-phenyl-propaneamide (1.03 gm, 2.9 mmol) in 20 ml of THF cooled to -78°C was added n-BuLi (2.5M, 1.40 ml, 3.5 mmol). After 20 min, allyl bromide (0.48 gm, 3.9 mmol) was added, the reaction was stirred at -78°C for 1 hr and then 10 ml of saturated NH_4Cl solution was added to quench the reaction. The reaction was diluted with 50 ml of water, extracted with ethyl acetate (2 x 50 ml), the organic phase was washed with saturated NaCl solution (50 ml), dried filtered and concentrated to afford the crude product. The crude product was purified on silica gel to afford the title compound.

Step 4: Preparation of N-(2(R)-hydroxy-1(S)-indan-N,O-isopropylidene-yl)-2(S)-phenylmethyl-(4(RS),5-dihydroxy)-pentaneamide

To 800 mg (2.2 mmol) of N-(2(R)-hydroxy-1(S)-indan-N,O-isopropylidene-yl)-2(S)-phenylmethyl-pent-4-eneamide dissolved in 40 ml of a 9:1 mixture of acetone/water was added 0.8 ml of a 60% solution of N-methylmorpholine-N-oxide in water followed by 4 ml of a 2.5% solution of osmium tetroxide in t-BuOH. After 18 hr, excess solid sodium bisulfate was added, the reaction was stirred for 2 hr and then filtered through a pad of celite. The filtrate was concentrated, diluted with 50 ml of water, extracted with methylene chloride (2 X 50 ml), the organic phase was dried, filtered and concentrated to give the product as a foam.

REPLACEMENT SHEET
04 FEB 1997

Step 5: Preparation of N-(2(R)-hydroxy-1(S)-indan-N,O-isopropylidene-yl)-2(S)-phenylmethyl-4(RS)-hydroxy-5-methanesulfonyloxy-pentaneamide

To 200 mg (0.527 mmol) of N-(2(R)-hydroxy-1(S)-indan-N,O-isopropylidene-yl)-2(S)-phenylmethyl-(4(RS),5-dihydroxy)-pentaneamide dissolved in 7 ml of methylene chloride at 0°C was added triethylamine (59 mg, 0.58 mmol), followed by methanesulfonyl chloride (66 mg, 0.579 mmol). After 4 hr the reaction was worked up by washing with 10% citric acid solution (2 X 50 ml) and the organic phase was dried, filtered and concentrated to afford the monomesylate as a mixture of alcohols.

Step 6: Preparation of N'-t-butyl-N-Boc-4(R)-hydroxy-L-prolineamide

To a solution of N-Boc-4(R)-hydroxyproline (2.00 g) in DMF (20 mL) cooled to 0°C was added EDC (1.987 g), HOBt (1.401 g), tert butyl amine (1.09 mL) and triethylamine (2.41 mL). After 18 h the reaction mixture was diluted with ethyl acetate (150 mL) and washed with 10% HCl, saturated $NaHCO_3$, water and brine. The solution was then dried over $MgSO_4$ and concentrated to afford a white solid.

Step 7: Preparation of N'-t-butyl-N-Boc-4(S)-phenoxy-L-prolineamide

To a solution of N'-t-butyl-N-Boc-4(R)-hydroxy-L-prolineamide (0.6 g) in THF (5 mL) was added phenol (0.295 g), triphenylphosphine (0.824 g) and then diethylazo-dicarboxylate (0.495 mL) dropwise. The reaction mixture stirred for 24 h at ambient

REPLACEMENT SHEET
04 FEB 1997

temperature and was diluted with ethyl acetate (200 mL) and washed with saturated $NaHCO_3$, water, brine and dried over $MgSO_4$. Concentration in vacuo afforded a yellow oil which was purified by flash chromatography (elution hexane: EtOAc 1:1, 30 mm column).

Step 8: Preparation of N-t-butyl-4(S)-phenoxy-L-prolineamide trifluoroacetic acid salt

To a solution of N'-t-butyl-N-Boc-4(S)-phenoxy-L-prolineamide (0.596 g) in methylene chloride (4 mL) at 0°C was added trifluoroacetic acid (2 mL). After 30 min the reaction was warmed to room temperature and stirred for two hours. The solvent was removed in vacuo and a slightly yellow oil was obtained.

Step 9: Preparation of N-(2(R)-hydroxy-1(S)-indan-N,O-isopropylidene-yl)-2-(R)-phenylmethyl-4-(S)-hydroxy-5-(1-(N'-(t-butyl)-4(S)-phenoxy-prolineamide)yl)-pentaneamide

To a solution of N-t-butyl-4(S)-phenoxy-L-prolineamide trifloroacetic acid salt (0.36 g) and N-(2(R)-hydroxy-1(S)-indan-N,O-isopropylidene-yl)-2(S)-phenylmethyl-4(RS)-hydroxy-5-methanesulfonyloxy-pentaneamide (0.226 g) in 3 mL of isopropanol was added potassium carbonate (0.441 g) and the reaction was warmed to 80°C. After 18 h the reaction was cooled to room temperature, filtered through celite which was washed with further portions of EtOAc. The filtrate was concentrated, the residue was dissolved in EtOAc (100mL) and washed with water, brine and dried over $MgSO_4$. The solvent was removed in vacuo and the resulting oil was purified by flash chromatography to

REPLACEMENT SHEET
04 FEB 1997

afford the product as a mixture of diastereomers.

Step 10: Prep of N-(2(R)-hydroxy-1(S)-indanyl)-2-(R)-phenylmethyl-4-(S)-hydroxy-5-(1-(N'-t-butyl-4(S)-phenoxyprolineamid)yl)-pentaneamide

To a solution of N-(2(R)-hydroxy-1(S)-indan-N,O-isopropylidene-yl)-2-(R)-phenylmethyl-4-(S)-hydroxy-5-(1-(N'-(t-butyl)-4(S)-phenoxyprolineamide)-yl)-pentaneamide (0.13 g) in MeOH (5 mL) was added camphorsulfonic acid (CSA) (0.070 g) at ambient temperature. After 5 hours more CSA (0.025 g) was added and the reaction was stirred for total of 18 hours. The reaction was quenched with saturated $NaHCO_3$ (5 mL) and the solvent was removed to a volume of 4 mL. The aqueous layer was thoroughly extracted with EtOAc and the organic layer was washed with water, brine and dried. After removal of the solvent in vacuo the resulting oil was purified via flash chromatography to provide the title compound as a white foam. The foam was dissolved in EtOAc : hexanes and the mother liquor was decanted away from the oil. The oil was then dried in a high vacuum desiccator to afford a white foam.

EXAMPLE 2

Preparation of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(1-(N'-t-butyl-4(S)-2-naphthyloxy-prolineamid)yl)-pentaneamide

Step 1: Preparation of N-t-butyl-4(S)-2-naphthyloxy-L-prolineamide trifluoroacetic acid salt

Following substantially the same procedure

REPLACEMENT SHEET
04 FEB 1997

for synthesizing N-t-butyl-4(S)-phenoxy-L-proline-amide trifluoroacetic acid salt as outlined in Example 1, Steps 6 through 8, but substituting 2-naphthol for the phenol used therein, the 2-naphthyloxy proline amide was produced.

Step 2: Preparation of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(1-(N'-t-butyl-4(S)-2-naphthyloxy-prolineamid)yl)-pentaneamide

The title compound was produced by following substantially the same procedure outlined in Example 1, Steps 9 and 10, but substituting N-t-butyl-4(S)-2-naphthyloxy-L-prolineamide trifluoroacetic acid salt for the N-t-butyl-4(S)- phenoxy-L-prolineamide trifloroacetic acid salt used in step 9 therein.

EXAMPLE 3

Preparation of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(1-(N'-t-butyl-4(S)-1-naphthyloxy-prolineamid)yl)-pentaneamide

Step 1: Preparation of N-t-butyl-4(S)-1-naphthyloxy-L-prolineamide trifluoroacetic acid salt

Following substantially the same procedure for synthesizing N-t-butyl-4(S)-phenoxy-L-proline-amide trifluoroacetic acid salt as outlined in Example 1, Steps 6 through 8, but substituting 1-naphthol for the phenol used therein, the 1-naphthyloxy proline amide was produced.

REPLACEMENT SHEET
04 FEB 1997

Step 2: Preparation of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(1-(N'-t-butyl-4(S)-2-naphthyloxy-prolineamid)yl)-pentaneamide

The title compound was produced by following the procedure outlined in Example 1, Steps 9 and 10, but substituting N-t-butyl-4(S)-1-naphthyloxy-L-prolineamide trifluoroacetic acid salt for the N-t-butyl-4(S)-phenoxy-L-prolineamide trifluoroacetic acid salt used in Step 9.

EXAMPLE 4

Preparation of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(2-(3(S)-N'-(t-butyl-carboxamido)-(4aS,8aS)-decahydroisoquinoline)yl)-pentaneamide

Step 1: Preparation of dihydro-5(S)-((t-butyldiphenylsilyl)oxymethyl)-3(R)phenylmethyl-3(2H)-furanone

A solution of lithium diisopropylamide (LDA) was generated by the addition 1.55 ml of n-BuLi (2.5M in hexane) to 0.55 ml (3.9 mmol) of diisopropylamine in 10 ml of THF at -78°C. After 30 minutes a solution of dihydro-5-(S)-((t-butyldiphenylsilyl)-oxymethyl)-3(2H)-furanone (1.38 g, 3.89 mmol) in 5 ml of THF was added. After an additional 30 minutes of stirring, benzyl bromide (0.68 g, 3.9 mmol) was added and stirring was continued for 3 h after which time the reaction was quenched with the addition of a 10% aqueous citric acid solution. The solution was

REPLACEMENT SHEET
04 FEB 1997

extracted with ethyl acetate (2 x 50 ml) which was backwashed with brine, dried, filtered and concentrated to afford an oil. The product was purified by chromatography (SiO_2, 20% EtOAc/Hexane) to afford the title compound.

Step 2: Preparation of dihydro-5(S)-(hydroxymethyl)-3(R)-phenylmethyl-3(2H)-furanone

To 5.26 g of dihydro-5(S)-((t-butyldiphenylsilyl)oxymethyl)-3(R)phenylmethyl-3(2H)-furanone in 40 ml of acetonitrile was added 1.34 ml of a 49% aqueous HF solution. After 18 hr at room temperature the reaction was concentrated to dryness and the residue was partitioned between water (50 ml) and ethyl acetate (50 ml). The organic layer was washed with brine, dried filtered and concentrated to afford the product as a tan solid (mp 69-72°C).

Step 3: Preparation of dihydro-5(S)-((methanesulfonyl)oxymethyl)-3(R)phenylmethyl-3(2H)-furanone

To a solution of 2.93 g (14 mmol) of dihydro-5(S)-(hydroxymethyl)-3(R)-phenylmethyl-3(2H)-furanone in methylene chloride cooled to 0°C was added triethylamine (1.98ml, 15.6 mmol) followed by the addition of methanesulfonyl chloride (1.20 ml, 15.6 mmol). After 1 hour at 0°C, the reaction was poured into 10% aqueous citric acid solution, washed with ethyl acetate (2 x 100 ml) which was backwashed with water (100 ml), brine (100 ml), dried, filtered and concentrated to give the product as a waxy brown solid.

REPLACEMENT SHEET
04 FEB 1997

Step 4: Preparation of dihydro-5(S)-(2-(3(S)-N-(t-butylcarboxamido)-(4aS, 8aS)-(decahydroisoquinoline)yl)methyl)-3(R)-phenylmethyl-3(2H)-furanone

To 70 mg of dihydro-5(S)-((methanesulfonyl)-oxymethyl)-3(R)phenylmethyl-3(2H)-furanone (0.25 mmol) in 10 ml of xylene containing 100 mg of potassium carbonate was added 65 mg (0.27 mmol) of N-t-butyl-(4aS,8aS)-(decahydroisoquinoline)-3(S)-carboxamide and the reaction was heated to 140°C. After 6 hours, the reaction was cooled, poured into 30 ml of water which was washed with ethyl acetate (2 x 30 ml). The organic phase was dried, filtered and concentrated to afford a residue which was chromatographed (50/50 EtOAc/Hexane) to give the product.

Step 5: Preparation of 2(R)-phenylmethyl-4(S)-(t-butyldimethylsilyloxy)-5-(2-(3(S)-N-(t-butylcarboxamido)-(4aS,8aS)-decahydroisoquinoline)yl)-pentanoic acid

To 130 mg (0.305 mmol) of dihydro-5(S)-(2-(3(S)-N-(t-butylcarboxamido)-(4aS, 8aS)-(decahydroisoquinoline)yl)methyl)-3(R)-phenylmethyl-3-(2H)furanone in 2 ml of DME was added 1 ml lithium hydroxide solution. After 4 hours at room temperature, the reaction was concentrated to dryness and azeotroped with toluene (3X) to remove excess water. The residue was dissolved in 5 ml of DMF and 414 mg (6.10 mmol) of imidazole and 465 mg (3.05 mmol) of t-butyldimethylsilyl chloride was added. After two days at room

REPLACEMENT SHEET
04 FEB 1997

temperature, 1 ml of methanol was added to the reaction and after 1 hour the solution was evaporated to dryness. The residue was partitioned between saturated NH_4Cl solution (aq) and washed with ethyl acetate which was dried, filtered and concentrated to give an oil which was a mixture of product and the furanone starting material. This material was carried on crude into the next reaction.

Step 6: Preparation of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-(t-butyldimethyl-silyloxy-5-(2-(3(S)-N'-(t-butylcarboxamido)-(4aS,8aS)-decahydroisoquinoline)yl)-pentaneamide

The crude product of step 5, above, was dissolved in 3 ml of DMF along with 47 mg (0.246 mmol) of EDC, 33 mg (0.246 mmol) of HOBT and 37 mg of 2(R)-hydroxy-1(S)-aminoindane. The pH of the solution was adjusted to 8.5-9.0 with triethylamine and after 18 hours it was worked up by concentrating to dryness, dissolving the residue in 10% aq. citric acid solution and washing the aqueous layer with ethyl acetate. The organic layer was dried, filtered and concentrated and the resultant oil was chromatographed (SiO_2, 30% EtOAc/Hexane) to yield the title compound.

Step 7: Preparation of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(2-(3(S)-N'-(t-butylcarboxamido)-(4aS,8aS)-decahydro-isoquinoline)yl)-pentaneamide

The product from step 6, above, was dissolved

REPLACEMENT SHEET
04 FEB 1997

in 1 ml of THF and 1 ml of a 1M solution of tetrabutylammonium fluoride in THF was added. After 18 hr at room temperature the reaction was diluted with 20 ml of saturated $NaHCO_3$ solution (aq) and the product was extracted into ethyl acetate which was dried, filtered and concentrated to give a foam. The resultant material was chromatographed on a prep plate (0.5 mm, 5% MeOH/$CHCl_3$) and the title product isolated in the usual manner as a solid with mp 105-107°C.

EXAMPLE 5

Preparation of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-amino-5-(2-(3(S)-N'-(t-butylcarboxamido)-(4aS,8aS)-decahydroisoquinoline)yl)-pentaneamide

Step 1: Preparation of 5(S)-((t-butyl-dimethyl-silyloxy)methyl)-3(R)-phenylmethyl-N-BOC-2-pyrrolidinone

A solution of 5(S)-((t-butyl-dimethyl-silyloxy)methyl)-N-BOC-2-pyrrolidinone (400 mg, 1.26 mmol) in 2 ml of THF was added to a precooled (-78°C) 1M solution of lithium hexamethyldisilazide (1.3 ml) in 5 ml of THF. After 45 min, 0.15 ml of benzyl bromide (1.3 mmol) was added and the stirring was continued. After 5 h the reaction was worked up by pouring into a separatory funnel containing 30 ml of an aqueous 10% citric solution. The aqueous layer was extracted (2 x 30 ml EtOAc) which was backwashed with brine (50 ml) dried, filtered and concentrated to an oil. The residue was chromatographed (SiO_2, 20% EtOAc/Hexane) to afford the product as an oil.

REPLACEMENT SHEET
04 FEB 1997

Step 2: Preparation of 5(S)-hydroxymethyl-3(R)-phenylmethyl-2-pyrrolidinone

To 130 mg (0.34 mmol) of 5(S)-((t-butyldimethylsilyloxy)methyl)-3(R)-phenylmethyl-N-BOC-2-pyrrolidinone in 5 ml of acetonitrile was added 0.1 ml of a solution of 48% HF in water. After 3 hr at room temperature the reaction was concentrated to dryness and diluted with 30 ml of an aqueous 10% $NaHCO_3$ solution. This was extracted with EtOAc (2 X 30 ml), dried filtered and concentrated to afford the crude product.

Step 3: Preparation of 5(S)-(methanesulfonyloxy)-methyl-3(R)-phenylmethyl-2-pyrrolidinone

To a solution of the crude product from Step 2, in 5 ml of methylene chloride cooled to 0°C was added triethylamine (42 mg, 0.41 mmol) and methanesulfonyl chloride (47 mg, 0.41 mmol). The reaction was slowly allowed to warm to room temperature and was stirred for 18 hr after which time it was diluted with 30 ml of methylene chloride, washed with 30 ml of 10% citric acid solution, dried filtered and concentrated to afford the product as an oil.

Step 4: Preparation of 5(S)-(2-(3(S)-N-(t-butylcarboxamido)-(4aS,8aS)-(decahydroisoquinoline)-yl)-methyl)-3(R)-phenylmethyl-2-pyrrolidinone

To a solution of 380 mg (1.34 mmol) of 5(S)-(methanesulfonyloxy)methyl-3(R)-phenylmethyl-2-pyrrolidinone in 20 ml of isopropanol was added 350 mg of potassium carbonate and 360 mg of N-t-butyl-(4aS,8aS)-(decahydroisoquinoline)-3(S)-carboxamide and the reaction was heated to 85°C. After 18 hr the

REPLACEMENT SHEET
04 FEB 1997

cooled reaction was filtered through celite, evaporated to dryness and the residue was dissolved in water which was extracted with EtOAc (2 X 50 ml). The organics were dried, filtered and concentrated, and the residue was chromatographed (SiO_2, 50/50 EtOAc/Hexane) to afford the product as an oil.

Step 5: Preparation of 5(S)-(2-(3(S)-N'-(t-butylcarboxamido)-(4aS,8aS)-(decahydroisoquinoline)-yl)-methyl)-3(R)-phenylmethyl-N-BOC-2-pyrrolidinone

To a solution of the product from step 4, above, (260 mg, 0.611 mmol) in 10 ml of methylene chloride was added dimethylaminopyridine (74 mg, 0.6 mmol) and 133 mg (0.61 mmol) of BOC-anhydride. After 18 hr at room temperature the reaction was worked up by diluting with 30 ml of methylene chloride and the organics washed with 30 ml of 10% citric acid solution, brine (30 ml) dried, filtered and concentrated to afford an oil. Chromatography (SiO_2, 40% EtOAc/Hexane) gave the title compound.

Step 6: Preparation of 5-(2-(3(S)-N'-(t-butylcarboxamido)-(4aS,8aS)-decahydroisoquinoline)-yl)-4(S)-[(1',1')-(dimethylethoxycarbonyl)-amino]-2(R)-phenylmethyl-pentanoic acid

To a solution of the product of step 5, above, (260 mg, 0.495 mmol) dissolved in 3 ml of dimethoxyethane was added 1.5 ml of a 1M solution of aqueous lithium hydroxide (1.5 mmol). The reaction was worked up after 2 hr by concentrating to dryness, dissolving the residue in saturated aqueous ammonium chloride solution and the aqueous phase was washed with

REPLACEMENT SHEET
04 FEB 1997

ethyl acetate (2 x 50 ml) which was dried, filtered and concentrated to afford the crude acid.

Step 7: Preparation of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-[(1',1')-(dimethyl-ethoxycarbonyl)amino]-5-(2-(3(S)-N'-(t-butyl-carboxamido)-(4aS,8aS)-decahydroisoquinol-ine)yl)-pentaneamide

To a solution of the product of step 6, above, (260 mg, 0.49 mmol) in methylene chloride was added EDC (94 mg, 0.49 mmol), HOBT (66 mg, 0.49 mmol), 2(R)-hydroxy-1(S)-aminoindane (73 mg, 0.49 mmol) and the pH of the reaction was adjusted to 8.5-9.0 using triethylamine. After 5 hr at room temperature the reaction was worked up by diluting with 50 ml of methylene chloride and washing the organics with saturated aqueous ammonium chloride solution. The organic phase was dried, filtered and concentrated and the residue was chromatographed to afford the title compound as a foam.

Step 8: Preparation of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(2-(3(S)-N'-(t-butylcarboxamido)-(4aS,8aS)-decahydro-isoquinoline)yl)-pentaneamide

To a solution of the product of step 7, above, (180 mg, 0.28 mmol) in 5 ml of methylene chloride cooled to 0°C was added 1 ml of trifluoro-acetic acid. After 4 hr the reaction was worked up by concentrating to dryness and the residue was dissolved in 50 ml of methylene chloride and washed with 10% aqueous $NaHCO_3$ solution. The organic layer was dried, filtered and concentrated to give the product as a

REPLACEMENT SHEET
04 FEB 1997

solid which was chromatographed (SiO_2, 7% MeOH/CH_2Cl_2) to afford the title compound, mp = 92-95°C.

EXAMPLE 6

Preparation of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(1-(4-carbobenzyloxy-2-(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide

Employing substantially the same procedure used in Example 1, but substituting N-t-butyl-4-CBZ-piperazine-2(S)-carboxamide for N-t-butyl-4(S)-phenoxy-L-prolineamide used in step 9 therein, the title compound was obtained.

EXAMPLE 7

Preparation of N''-(N-(2-pyridyl)-valyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(2-(3(S)-(N'-t-butylcarboxamido)-(4aS,8aS)-decahydroisoquinoline)yl)pentaneamide

Employing substantially the same procedure used in Example 4, but substituting N-2-pyridyl-valine for the 2(R)-hydroxy-1(S)aminoindane used in step 6 therein, the title compound was obtained.

EXAMPLE 8

Preparation of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(2(S)-(N'-t-butyl-3-phenyl-propionamide)amino)-pentaneamide

Employing substantially the same procedure used in Example 1, but substituting N-t-butyl-phenylalanine amide for the N'-t-butyl-4(S)-phenoxy-

REPLACEMENT SHEET
04 FEB 1997

L-prolineamide used in step 9 therein, the title compound is obtained.

EXAMPLE 9

Preparation of N-(4(S)-3,4-dihydro-1H-2,2-dioxobenzothiopyranyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(2-(3(S)-N'-(t-butylcarboxamido)-(4aS,8aS)-decahydroisoquinoline)yl)-pentaneamide

Step 1: Preparation of N-(4(S)-3,4-dihydro-1H-benzothiopyranyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(2-(3(S)-t-butylcarboxamido)-(4aS,8aS)-decahydroisoquinoline)yl)-pentaneamide

Employing substantially the same procedure used in Example 4 but substituting 4(S)-amino-3,4-dihydro-1H-benzothiopyran for the 2(R)-hydroxy-1(S)-aminoindane used in step 6 therein, the title compound is obtained.

Step 2: Preparation of N-(4(S)-3,4-dihydro-1H-2,2-dioxobenzothiopyranyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(2-(3(S)-t-butylcarboxamido)-(4aS,8aS)-decahydroisoquinoline)yl)-pentaneamide

The compound from step 1 above is dissolved in a 1:1 mixture of methanol and water. To this is added 10 eq. of OXONE and the reaction is stirred at room temperature. When the reaction is complete, it is concentrated to dryness, water is added and extracted with ethyl acetate which is dried, filtered and concentrated to give the title compound.

REPLACEMENT SHEET
04 FEB 1997

EXAMPLE 10

Preparation of N-(4(S)-3,4-dihydro-1H-2,2-dioxobenzothiopyranyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(1-(4-carbobenzyloxy-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide

Step 1: Preparation of dihydro-5(S)-(1-(4-carbobenzyloxy-2(S)-N'-(t-butylcarboxamido)-piperazinyl)methyl)-3(R)-phenylmethyl-3(2H)-furanone

Employing substantially the same procedure used in Example 4, step 4 but substituting 4-carbobenzyloxy-2(S)-N'-(t-butylcarboxamido)-piperazine for the N'-t-butyl-(4aS,8aS)-(decahydroisoquinoline)-3(S)-carboxamide used therein, the title compound is produced.

Step 2: Preparation of 2(R)-phenylmethyl-4(S)-(t-butyldimethylsilyloxy)-5-(1-(4-carbobenzyloxy-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentanoic acid

Employing substantially the same procedure used in Example 4, step 5 but substituting dihydro-5(S)-(1-(4-carbobenzyloxy-2(S)-N'-(t-butylcarboxamido)-piperazinyl)methyl)-3(R)-phenylmethyl-3(2H)-furanone for the dihydro-5(S)-(2-(3(S)-N'-(t-butylcarboxamido)-(4aS,8aS)-(decahydroisoquinoline)yl)-methyl)-3(R)-phenylmethyl-3(2H) furanone used therein, the title compound is produced.

REPLACEMENT SHEET
04 FEB 1997

Step 3: Preparation of N-(4(S)-3,4-dihydro-1H-benzothiopyranyl)-2(R)-phenylmethyl-4(S)-(t-butyldimethylsilyloxy)-5-(1-(4-carbobenzyloxy-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide

The crude 2(R)-phenylmethyl-4(S)-(t-butyldimethylsilyloxy)-5-(1-(4-carbobenzyloxy-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentanoic acid is dissolved in 3ml of DMF along with 1 eq of EDC, 1 eq of HOBT and 1 eq of 4(S)-amino-3,4-dihydro-1H- benzothiopyran. The pH of the solution is adjusted to 8.5-9.0 with triethylamine and after 18 hours it is worked up by concentrating to dryness, dissolving the residue in 10% aq citric acid solution and washing the aqueous layer with ethyl acetate. The organic layer is dried, filtered and concentrated and the resultant residue is chromatographed to yield the title product.

Step 4: Preparation of N-(4(S)-3,4-dihydro-1H-benzothiopyranyl)-2(R)-phenylmethyl-4(S)-hydroxy)-5-(1-(4-carbobenzyloxy-2(S)-(t-butylcarboxamido)-piperazinyl))-pentaneamide

The product from step 3 above is dissolved in 1 ml of THF and 1 ml of a 1M solution of tetra-butylammonium fluoride in THF is added. After 18 hr at room temperature the reaction is diluted with 20 ml of saturated $NaHCO_3$ solution (aq) and the product is extracted into ethyl acetate which is dried, filtered and concentrated to give a residue. The residue is chromatographed to afford the product.

REPLACEMENT SHEET
04 FEB 1997

Step 5: Preparation of N-(4(S)-3,4-dihydro-1H-2,2-dioxobenzothiopyranyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(1-(4-carbobenzyloxy-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide

The compound from step 4 above is dissolved in a 1:1 mixture of methanol and water. To this is added 10 eq of OXONE and the reaction is stirred at room temperature. When the reaction is complete, it is concentrated to dryness, water is added and extracted with ethyl acetate which is dried, filtered and concentrated to give the title compound.

EXAMPLE 11

Preparation of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-((2-hydroxy)ethoxy)phenyl)methyl)-4(S)-hydroxy-5-(2-(3-(S)-N'-(t-butylcarboxamido)-(4aS,8aS)-decahydroisoquinoline)yl)pentaneamide

Step 1: Preparation of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-(2-allyloxy)phenyl)methyl)-4(S)-hydroxy-5-(2-(3(S)-t-butylcarboxamido)-(4aS,8aS)-decahydroisoquinoline)yl)-pentaneamide

To a solution of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-hydroxyphenyl)methyl)-4(S)-hydroxy-5-(2-(3(S)-t-butylcarboxamido)-(4aS,8aS)-decahydroisoquinoline)yl)-pentaneamide in dioxane is added 6 eq of allyl bromide and 6 eq of cesium carbonate. The reaction is heated to 90°C. When the reaction is complete, the precipitate is filtered off, the dioxane is concentrated to dryness and the residue is diluted with

REPLACEMENT SHEET
04 FEB 1997

water which is washed with ethyl acetate. The organic phase is dried, filtered and concentrated to afford the product.

Step 2: Preparation of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-((2-hydroxy)ethoxy)phenyl)methyl)-4(S)-hydroxy-5-(2-(3(S)-N'-(t-butylcarboxamido)-(4aS,8aS)-decahydroisoquinoline)yl)-pentaneamide

The product from step 1 above is dissolved in methanol, 1 eq of p-toluenesulfonic acid is added and the reaction is cooled to -78°C. Excess ozone is bubbled through the reaction until a blue color persists. The flask is purged with nitrogen to remove any ozone and excess sodium borohydride solution is added. The reaction is warmed to room temperature and then saturated $NaHCO_3$ solution is added. The methanol is concentrated off on the rotoevaporater and the aqueous residue is washed with ethyl acetate which is dried, filtered and concentrated to afford the title compound.

EXAMPLE 12

Preparation of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-((2-hydroxy)ethoxy)phenyl)methyl)-4(S)-hydroxy-5-(1-(4-carbobenzyloxy-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide

Employing substantially the same prodecure used in Example 11 but substituting N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-hydroxyphenyl)methyl)-4(S)-hydroxy-5-(1-(4-carbobenzyloxy-2(S)-(t-butylcarbox-

REPLACEMENT SHEET
04 FEB 1997

amido)-piperazinyl)-pentaneamide for the N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-hydroxyphenyl)methyl)-4(S)-hydroxy-5-(2-(3(S)-t-butylcarboxamido)-(4aS,8aS)-decahydroisoquinoline)yl)-pentaneamide used therein, the title compound is obtained.

EXAMPLE 13

Preparation of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-(2-(4-morpholinyl)ethoxy)phenyl)methyl)-4(S)-hydroxy-5-(2-(3(S)-N'-(t-butylcarboxamido)-(4aS,8aS)-decahydroisoquinoline)yl)-pentaneamide

To a solution of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-hydroxyphenyl)methyl)-4(S)-hydroxy-5-(2-(3(S)-N'-(t-butylcarboxamido)-(4aS,8aS)-decahydroisoquinoline)yl)-pentaneamide in dioxane is added 6 eq of chloroethyl morpholine and 6 eq of cesium carbonate. The reaction is heated to 90°C. When the reaction is complete, the precipitate is filtered off, the dioxane is concentrated to dryness and the residue is diluted with water which is washed with ethyl acetate. The organic phase is dried, filtered and concentrated to afford the title compound.

EXAMPLE 14

Preparation of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-(2-(4-morpholinyl)ethoxy)phenyl)methyl)-4(S)-hydroxy-5-(1-(4-carbobenzyloxy-2(S)-N'-(t-butyl-carboxamido)-piperazinyl))-pentaneamide

To a solution of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-hydroxyphenyl)methyl)-4(S)-hydroxy-5-(1-(4-carbobenzyloxy-2(S)-(t-butylcarboxamido)-

REPLACEMENT SHEET
04 FEB 1997

piperazinyl)-pentaneamide in dioxane is added 6 eq of chloroethyl morpholine and 6 eq of cesium carbonate. The reaction is heated to 90°C. When the reaction is complete, the precipitate is filtered off, the dioxane is concentrated to dryness and the residue is diluted with water which is washed with ethyl acetate. The organic phase is dried, filtered and concentrated to afford the title compound.

EXAMPLE 15

Preparation of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(1-(4-(3-pyridylmethyl)-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide

Step 1: Preparation of dihydro-5(S)-((trifluoromethanesulfonyl)oxymethyl)-3(R)-phenylmethyl-3(2H)-furanone

To a solution of 18.4g (89.2 mmol) of dihydro-5(S)-(hydroxymethyl)-3(R)-phenylmethyl-3(2H)-furanone in 350 mL of methylene chloride cooled to 0°C was added 13.51 mL 2,6-lutidine (115.98 mmol) followed by a dropwise addition of 16.51 mL of trifluoromethanesulfonic anhydride (98.1 mmol). After 1.5 hours at 0°C, the reaction was poured into a mixture of 300 mL ice/brine and stirred for 0.5 hours. The aqueous layer was then extracted with methylene chloride (3 x 150 mL), the organic layers were washed with 10% HCl (2 x 75 mL), saturated $NaHCO_3$ (100mL), water (100mL), dried over $MgSO_4$, filtered and concentrated to give a solid residue. Purification via flash column chromatography (120 x 150 mm column, gradient elution of hexanes:EtOAc, 4:1 to 3:1) afforded the title product; mp 53-54°C.

REPLACEMENT SHEET
04 FEB 1997

Step 2: Preparation of 4-(1,1-dimethylethyl)-1-(phenylmethyl)-1,2(S),4-piperazinetricarboxylate

The title compound was prepared following the procedure of Bigge, C.F.; Hays, S.J.; Novak, P.M.; Drummond, J.T.; Johnson, G.; Bobovski, T.P. Tetrahedron Lett. 1989, 30, 5193; starting with 2(S)-piperazinecarboxylic acid. (see Felder, E.; Maffei, S.; Pietra, S.; Pitre, D.; Helv. Chim. Acta 1960, 117, 888.

Step 3: Preparation of N-t-butyl-4-(1,1-dimethylethoxycarbonylamino)-1-(phenylmethylcarbonylamino)piperazine-2(S)-carboxamide

To 9.90g (27.16 mmol) of 4-(1,1-dimethylethyl)-1-(phenylmethyl)-1,2(S),4-piperazinetricarboxylate dissolved in 75 mL of DMF and cooled to 0°C was added 5.73g (29.88 mmol) of EDC, 4.03g (29.88 mmol) of HOBt, 3.14 mL (29.88 mmol) of t-butylamine, and finally 4.16 mL (29.88 mmol) of triethylamine. The reaction mixture was stirred for 18 hours and the reaction volume was concentrated by half. The mixture was then diluted with 600 mL of EtOAc and washed with 10% HCl (2 x 75 mL), saturated $NaHCO_3$ (1 x 75 mL), water (3 x 75 mL) and brine (1 x 50 mL), dried over $MgSO_4$ and concentrated to a solid. This solid was triturated with EtOAc: hexane (1:2) and filtered to provide the title product as a white solid;
mp 134-135°C.

REPLACEMENT SHEET
04 FEB 1997

Step 4: Preparation of N-t-butyl-4-(1,1-dimethyl-ethoxycarbonylamino)piperazine-2(S)-carboxamide

To 1.20g (2.86 mmol) of N-t-butyl-4-(1,1-dimethylethoxycarbonylamino)-1-(phenylmethylcarbonyl-amino)piperazine-2(S)-carboxamide and 1.1g (0.086 mmol) of 10% Pd/C was added 15 mL of methanol. The vessel was charged with hydrogen and the reaction stirred for 2 hours, filtered through celite and washed with ethanol. The solvents were removed in vacuo to provide the title product as a foam.
^{1}H NMR (300 MHz, $CDCl_3$) δ 6.65 (br, 1H), 4.10 (m, 1H), 3.81 (br, 1H), 3.21 (dd, J=18 and 7 Hz, 1H), 3.02-2.70 (m, 4H), 2.10-2.0 (br, 1H), 1.50 (s, 9H), 1.41(s, 9H).

Step 5: Preparation of dihydro-5(S)-(4-(1,1-dimethylethoxycarbonylamino))-2(S)-N-(t-butylcarboxamido)-piperazinyl)methyl)-3(R)-phenylmethyl-3(2H)-furanone

To a solution of 22.40g (0.0662 mol) dihydro-5(S)-((trifluoromethanesulfonyl)oxymethyl)-3(R)-phenylmethyl-3(2H)-furanone (prep in step 1) and 18.0g (0.063mol) of n-t-butyl-4-(1,1-dimethylethoxy-carbonylamino)piperazine-2(S)-carboxamide dissolved in 180 mL of isopropanol was added 11.53 mL (0.0662 mol) of N,N-diisopropylethylamine. After 2.5 hours another 1.2g of dihydro-5(S)-((trifluoromethanesulfonyl)oxy-methyl)-3(R)-phenylmethyl-3(2H)-furanone was added. The reaction was complete by thin layer chromatography (tlc) after 3.5 hours and was concentrated to a thick oil. Trituration with EtOAc:hexanes (1:2, 200mL) provided a white solid which was filtered and discarded. The oil was purified by flash column

REPLACEMENT SHEET
04 FEB 1997

chromatography (120 x 150 mm column, EtOAc:hexanes gradient elution 1:1, 2:1, 3:1 to all EtOAc) to afford the title compound.

^{1}H NMR (400 MHz, $CDCl_3$) δ 7.34-7.17 (m, 5H), 6.31 (br s, 1H), 4.38 (br m, 1H), 3.96-3.92 (m, 1H), 3.79 (br m, 1H), 3.16 (dd, J=13.6 and 4.4 Hz, 1H), 3.08-2.99 (m, 3H), 2.90-2.82 (m, 1H), 2.80 (dd, J=13.5 and 8.9 Hz, 1H), 2.78 (m, 1H), 2.67-2.61 (m,1H), 2.58-2.49 (m, 1H), 2.38-2.32 (m,1H), 2.32-2.04 (m, 1H), 1.99-1.92 (m, 1H,) 1.45 (s, 9H), 1.29 (s, 9H).

Step 6: Preparation of 2(R)-phenylmethyl-4(S)-(t-butyldimethylsilyloxy)-5-(1-(4-(1,1-dimethylethoxycarbonylamino)))-2(S)-N-(t-butylcarboxamido)-piperazinyl))-pentaneamide

To 25.50g (52.50 mmol) of dihydro-5(S)-(4-(1,1-dimethylethoxycarbonylamino))-2(S)-N-(t-butyl-carboxamido)-piperazinyl)methyl)-3(R)-phenylmethyl-3(2H)-furanone dissolved in 120 mL DME cooled to 0°C was added a solution of 60 mL of water and 1.512g (63.01 mmol) of lithium hydroxide. After 0.5 hours the reaction was quenched with the addition of 10% HCl until pH 6 and the solution was concentrated in vacuo. The residue was dissolved in 50 mL water and extracted with EtOAc (4 x 75 mL) and the organic layers were washed with water (1 x 20 mL), brine (1 x 20 mL). The aqueous was back extracted with EtOAc (2 x 75 mL) and the combined organic layers were dried over $MgSO_4$ and concentrated to provide a yellow solid. This crude product was dissolved in 100 mL of DMF and 17.87g (0.262 mol) of imidazole was added, cooled to 0°C and then 31.50g (0.21 mol) of t-butyldimethylsilyl chloride was added. This stirred 1 hour at 0°C and was then

REPLACEMENT SHEET
04 FEB 1997

warmed to room temperature. After 20 hours the reaction was quenched with 10 mL methanol and concentrated to half the volume. 100 mL of pH 7 buffered water was added and the aqueous was extracted with EtOAc (4 x 100 mL), the combined organic layers were washed with 10% HCl (2 x 50 mL), water (3 x 75 mL), and brine (1 x 50 mL), dried over $MgSO_4$ and concentrated to obtain the title compound. This material was used directly in the next step.

Step 7: Preparation of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-(t-butyldimethylsilyl-oxy)-5-(1-(4-(1,1-dimethylethoxycarbonyl-amino)))-2(S)-N-(t-butylcarboxamido)-piper-azinyl))-pentaneamide

To 27.0g (0.0446mol) of the crude material from step 6 dissolved in 180 mL of DMF and cooled to 0°C was added 8.98g (0.0468 mol) of EDC, 6.32g (0.0468 mol) of HOBt, and 7.31g (0.049 mol) aminohydroxy indane. Triethylamine (6.52 mL, 0.0468 mol) was added and the reaction stirred at 0°C for 2 hours, room temperature for 16 hours and was quenched by diluting with 500 mL of EtOAc. The organic layer was washed with 10% HCl (2 x 100 mL), saturated $NaHCO_3$ (1 x 100 mL), water (3 x 150 mL), brine (1 x 75 mL), dried over $MgSO_4$ and concentrated to yield the title compound as a white foam.

^1H NMR (400 MHz, $CDCl_3$) δ 7.4-7.17 (m, 9H), 6.51 (br s, 1H), 5.79 (br s, 1H), 5.23 (m, 1H), 4.23 (br s, 1H), 4.06 (m, 1H), 3.96-3.84 (m, 2H), 3.07-2.78 (m, 8H), 3.65 (dd, J=9.6 and 4.1 Hz, 1H), 2.56-2.44 (m, 2H), 2.29 (dd, J=12.0 and 4.5 Hz, 1H), 2.17-2.09 (m, 1H), 1.79 (br s, 1H), 1.44 (s, 9H), 1.35 (s, 9H), 1.10 (s, 1H), 0.84 (s, 9H), 0.12 (s, 3H), 0.08 (s, 3H).

REPLACEMENT SHEET
04 FEB 1997

Step 8: Preparation of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-(hydroxy)-5-(1-(4-(1,1-dimethylethoxycarbonylamino)))-2(S)-N-(t-butylcarboxamido)-piperazinyl))-pentaneamide

To 32.20g (0.0437 mol) of N-(2(R)-hydroxy-1-(S)-indanyl)-2(R)-phenylmethyl-4(S)-(t-butyldimethylsilyloxy)-5-(1-(4-(1,1-dimethylethoxycarbonylamino)))-2(S)-N-(t-butylcarboxamido)-piperazinyl))-pentaneamide was added 437 mL (0.437 mol) of tetrabutylammonium fluoride (1.0M solution in THF, Aldrich). The reaction stirred for 18 hours and was then concentrated to 200 mL and diluted with 700 mL of EtOAc. This was washed with water (2 x 100 mL), brine (1 x 50 mL) and the aqueous layers were back extracted with EtOAc (2 x 200 mL). The combined organic layers were dried over $MgSO_4$ and concentrated to an oil. Purification via flash column chromatography (120 x 150 mm column, gradient elution CH_2Cl_2: $CHCl_3$/saturated with NH_3: methanol, increasing methanol from 1%, 1.5%, 2%) afforded the title compound as a white foam.
^1H NMR (400 MHz, $CDCl_3$) δ 7.31-7.11 (m, 9H), 6.41 (br s, 1H), 6.23 (d, J=8.6 Hz, 1H), 5.25 (dd, J=8.6 and 4.7Hz, 1H), 4.21 (m, 1H), 3.83-3.82 (m, 2H), 3.78-3.61 (m, 2H), 3.22-3.19 (m, 2H), 3.03-2.78 (m, 8H), 2.62-2.58 (m, 1H), 2.41-2.35 (m, 2H), 2.04-2.02 (m, 1H), 1.57-1.50 (m, 1H), 1.45 (s, 9H), 1.32 (s, 9H).

REPLACEMENT SHEET
04 FEB 1997

Step 9: Preparation of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-(hydroxy)-5-(1-(2(S)-N-(t-butylcarboxamido)-piperazinyl)-pentaneamide

To 21.15g (0.034 mol) of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-(hydroxy)-5-(1-(4-(1,1-dimethylethoxycarbonylamino)))-2(S)-N-(t-butylcarboxamido)-piperazinyl))-pentaneamide dissolved in 350 mL of methylene chloride and cooled to 0°C was added 22.43 mL (0.204 mol) 2,6-lutidine and then 32.85 mL (0.170 mol) of trimethylsilyltriflate over 5 minutes. After 0.5 hours the reaction was quenched with 10% HCl (80 mL) and this stirred 0.5 hours. To this was added 100 mL of saturated $NaHCO_3$ and then solid $NaHCO_3$ until pH 8. The aqueous layer was then extracted with EtOAc (4 x 100 mL) and the combined organic layers were washed with water (1 x 50 mL), brine (1 x 75 mL), dried over $MgSO_4$ and concentrated. The residue was purified via column chromatography (120 x 150 mm column, gradient elution CH_2Cl_2:$CHCl_3$ saturated with NH_3: MeOH, slowly increasing methanol 2%, 3%, 4%, 5%, 6%, to 10%). This provided the title product as a white foam.

^1H NMR (400 MHz, $CDCl_3$) δ 7.53 (s, 1H), 7.29-7.09 (m, 9H), 6.52 (d, J=8.3 Hz, 1H), 5.24 (dd, J=8.2 and 4.9 Hz, 1H), 4.23 (dd, J=4.7 and 4.03 Hz, 1H), 4.25-4.00 (br s, 1H), 3.83-3.81 (m, 1H), 3.03-2.88 (m, 4H), 2.82-2.73 (m, 7H), 2.50-1.60 (br s, 2H), 2.45 (d, J=6.2 Hz, 2H), 2.32-2.29 (m, 1H), 1.98 (m, 1H), 1.51 (m, 1H), 1.33 (s, 9H).

REPLACEMENT SHEET
04 FEB 1997


Step 10: Preparation of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(1-(4-(3-pyridylmethyl)-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide

To 10.0g (0.019 mol) of N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-hydroxy)-5-(1(-2(S)-N-(t-butylcarboxamido)-piperazinyl)-pentaneamide and 3.45g (0.021 mol) of 3-picolyl chloride dissolved in 40 mL of DMF was added 5.85 mL (0.042 mol) of triethylamine. After 3 hours an additional 0.313g of 3-picolyl chloride was added. After an additional 2 hours the reaction was diluted with 400 mL of EtOAc and washed with water (3 x 75 mL), brine (1 x 100 mL), dried over $MgSO_4$ and concentrated. The residue was triturated with 30 mL of EtOAc and the resulting white precipitate was collected. Further recrystallization from EtOAc provided the title product (mp 167.5-168°C).

EXAMPLE 16

Employing substantially the same procedure as described in Example 15, but treating the N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(1-(2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide used therein (compound (i) below) with the alkylating agent (ii) indicated below in place of the 3-picolyl chloride used in Step 10 therein, the following products defined by formula (iii) were made:


REPLACEMENT SHEET
04 FEB 1997







i

R^1-X







ii











iii

R^1	X
	I
	Cl
	Cl
	I






REPLACEMENT SHEET
04 FEB 1997

	R^1	X
	CH_3CH_2-	I
		Br
		Cl
		Cl
		Cl
		I
		Cl

REPLACEMENT SHEET
04 FEB 1997

	R^1	X
5		Cl
		I
10		Cl
		Cl
15		Cl
20	$CH_3O(CH_2CH_2O)_2-CH_2CH_2-$	I
		I
25		I
30		Cl

REPLACEMENT SHEET
04 FEB 1997

R^1	X
	I
	Cl
	Cl
	Cl
	Cl

EXAMPLE 17

Preparation of dihydro-5(S)-(tert-butyldimethylsilyl-oxymethyl)-3(2H)-furanone

To a solution of 3.00g (25.8 mmol) of dihydro-5(S)-(hydroxymethyl)-2(3H)-furanone dissolved in 25 mL of dichloromethane was added 3.51g (51.6 mmol) of imidazole and then 4.67g (31.0 mmol) of tert-butyl-

REPLACEMENT SHEET
04 FEB 1997

dimethylsilyl chloride. The reaction stirred at room temperature for 8 hours and was quenched with 2 mL of methanol. The mixture was concentrated to an oil and then diluted with 150 mL of ether and washed with 5% HCl (2 x 10 mL), saturated $NaHCO_3$ (1 x 10 mL), water (1 x 10 mL), and brine (1 x 10 mL), dried over $MgSO_4$ and concentrated. The residue was purified by flash chromatography (40 x 150 mm column, gradient elution, hexanes:ethyl/acetate 5:1 to 4:1) to afford the product as a clear oil.

^1H NMR (300 MHz, $CDCl_3$) δ 4.68-4.60 (m, 1H), 3.89 (dd, J=3.3 and 11.3 Hz, 1H), 3.71 (dd, J=3.2 and 5411.3 Hz,1H), 2.71-2.45 (m, 2H), 2.35-2.16 (m, 2H), 0.91 (s, 9H), 0.10 (s, 3H), 0.09 (s, 3H).


While the foregoing specification teaches the principles of the present invention, with examples provided for the purpose of illustration, it will be understood that the practice of the invention emcompasses all of the usual variations, adaptations, or modifications, as come within the scope of the following claims and its equivalents.

THIS SHEET HAS BEEN REPLACED

REPLACEMENT SHEET
04 FEB 1997

WHAT IS CLAIMED IS:

1. A compound of the formula:



I

wherein
X is -OH or $-NH_2$;
Z is -O, -S, or -NH;
R is hydrogen or C_{1-4} alkyl;
R^1 and R^2 are independently:
1) hydrogen,
2) $-C_{1-4}$ alkyl unsubstituted or substituted with one or more of
a) halo,
b) hydroxy,
c) C_{1-3} alkoxy,
d) aryl unsubstituted or substituted with one or more of C_{1-4}alkyl, hydroxy or aryl,
e) -W-aryl or -W-benzyl,
wherein W is -O-, -S-, or -NH-,
f) a 5-7 membered cycloalkyl group unsubstituted or substituted with one or more of

REPLACEMENT SHEET
04 FEB 1997
THIS SHEET HAS BEEN REPLACED

* C L M S *

i) halo,
ii) hydroxy,
iii) C_{1-3} alkoxy, or
iv) aryl,

g) heterocycle unsubstituted or substituted with one or more of hydroxy, C_{1-4}alkyl optionally substituted with hydroxy, or Boc,

h) $-NH-\overset{O}{\overset{\|}{C}}OC_{1-3}alkyl$,

i) $-NH-\overset{O}{\overset{\|}{C}}-C_{1-3}alkyl$,

j) $-NH-SO_2C_{1-3}alkyl$,

k) $-NR_2$,

l) -COOR, or

m) $-((CH_2)_mO)_nR$ wherein m is 2-5 and n is zero, 1, 2 or 3, or

3) aryl, unsubstituted or substituted with one or more of

a) halo,
b) hydroxy,
c) $-NO_2$ or $-NR_2$,
d) C_{1-4}alkyl,
e) C_{1-3} alkoxy, unsubstituted or substituted with one or more of -OH or C_{1-3} alkoxy,
f) -COOR,
g) $-\overset{O}{\overset{\|}{C}}NR_2$,
h) $-CH_2NR_2$,

REPLACEMENT SHEET
04 FEB 1997
THIS SHEET HAS BEEN REPLACED

* C L M S *

i) $-CH_2NHC(=O)R$,
j) $-CN$,
k) $-CF_3$,
l) $-NHC(=O)R$,
m) aryl C_{1-3} alkoxy,
n) aryl,
o) $-NRSO_2R$,
p) $-OP(O)(OR_x)_2$, or
q) $-R^5$, as defined below; or

R^1 and R^2 can be joined together to form with the nitrogen to which R^1 is attached a 3 to 10 membered monocyclic or bicyclic saturated ring system which consists of the nitrogen to which R^1 is attached and from 2 to 9 carbon atoms, and is unsubstituted or substituted with

1) hydroxy,
2) C_{1-4} alkyl unsubstituted or substituted with one or more of
   a) halo,
   b) hydroxy,
   c) C_{1-3} alkoxy,
   d) aryl,
   e) a 5-7 membered cycloalkyl group unsubstituted or substituted with one or more of
      i) halo,
      ii) hydroxy,
      iii) C_{1-3} alkoxy, or
      iv) aryl,
   f) heterocycle, or
   g) $-NR_2$,

REPLACEMENT SHEET
04 FEB 1997
THIS SHEET HAS BEEN REPLACED

* C L M S *

3) C_{1-3} alkoxy,

4) $-NH-\overset{O}{\overset{\|}{C}}OC_{1-3}alkyl$,

5) $-NH-\overset{O}{\overset{\|}{C}}-C_{1-3}alkyl$,

6) $-NH-SO_2C_{1-3}alkyl$,

7) heterocycle,

8) -W-aryl, or

9) $-W-\overset{O}{\overset{\|}{C}}-aryl$,

wherein W is defined above; or

R^1 and R^2 can be joined together to form with the nitrogen to which R^1 is attached a 3 to 10 membered monocyclic or bicyclic saturated ring system which consists of the nitrogen to which R^1 is attached, from 1 to 8 carbon atoms and one or more unsubstituted or substituted heteroatom selected from

1) $-\overset{|}{N}-V-R^1$,

wherein V is absent or $-\overset{O}{\overset{\|}{C}}-Q-$ or $-SO_2-Q-$, R^1 is defined as above for when R^1 is independent from and not joined to R^2, and wherein Q is absent or -O-, -NR-, or heterocycle optionally substituted with $-C_{1-4}alkyl$,

2) $-\overset{|}{N}-$heterocycle,

3) $-\overset{|}{N}-C_{1-4}$ alkenyl, unsubstituted or substituted with aryl,

REPLACEMENT SHEET
04 FEB 1997
THIS SHEET HAS BEEN REPLACED

* C L M S *

4) $-N(SO_2-C_{1-4}alkenyl)-$, unsubstituted or substituted with aryl,

5) $-S(O)_p-$, wherein p is zero, 1 or 2, or

6) $-O-$; or

R^1 and R^2 can be joined together to form with the nitrogen to which R^1 is attached a 3 to 10 membered monocyclic or bicyclic saturated ring system, which consists of the nitrogen to which R^1 is attached and from 2 to 9 carbon atoms, in which the saturated ring system is fused to a phenyl ring and the phenyl ring is unsubstituted or substituted with one or more of

1) halo,
2) C_{1-3} alkoxy,
3) hydroxy,
4) C_{1-4} alkyl,
5) $-NHR^1$, wherein R^1 is defined as above for when R^1 is independent from and not joined to R^2, or
6) -NH-heterocycle;

R^3 is

1) $-(CH_2)_r-R^4$, wherein r is zero through 5,
2) $C_{1-4}alkenyl-R^4$, or
3) $C_{1-4}alkynyl-R^4$;

R^4 is

1) hydrogen,
2) C_{1-4} alkyl,

REPLACEMENT SHEET
04 FEB 1997
THIS SHEET HAS BEEN REPLACED

* C L M S *

3) C_5-C_{10} cycloalkyl, optionally substituted with hydroxy,

4) C_6-C_{10} aryl, unsubstituted or substituted with one or more of

a) halo,

b) hydroxy,

c) $-NO_2$ or $-NR_2$,

d) C_{1-4}alkyl,

e) C_{1-3} alkoxy, unsubstituted or substituted with one or more of -OH or C_{1-3} alkoxy,

f) -COOR,

g) $-\overset{\overset{O}{\|}}{C}NR_2$,

h) $-CH_2NR_2$,

i) $-CH_2NH\overset{\overset{O}{\|}}{C}R$,

j) -CN,

k) $-CF_3$,

l) $-NH\overset{\overset{O}{\|}}{C}R$,

m) aryl C_{1-3} alkoxy,

n) aryl,

o) $-NRSO_2R$,

p) $-OP(O)(OR_x)_2$, or

q) $-R^5$, as defined below, or

5) monocyclic or bicyclic heterocyle containing from 1 to 3 heteroatoms chosen from the group consisting of N, O, and S and which is unsubstituted or substituted with R^5 and optionally with one or more of

a) halo,

b) C_{1-4} alkyl, or

c) C_{1-3} alkoxy;

REPLACEMENT SHEET
04 FEB 1997
THIS SHEET HAS BEEN REPLACED

* C L M S *

R_x is H or aryl;

R^5 is

1) $-W-(CH_2)_m-NR^6R^7$
wherein W is as defined above,
m is 2-5, and
R^6 and R^7 are independently
a) hydrogen,
b) C_{1-6} alkyl, unsubstituted or substituted with one or more of
i) C_{1-3} alkoxy,
ii) -OH, or
iii) $-NR_2$,
c) the same or different and joined together to form a 5-7 member heterocycle, such as morpholino, containing up to two additional heteroatoms selected from
$-\overset{R}{\underset{|}{N}}-$, -O-, $-\overset{O}{\overset{\|}{S}}-$, -S-, or $-SO_2-$, the heterocycle optionally substituted with C_{1-4} alkyl, or
d) aromatic heterocycle unsubstituted or substituted with one or more of
i) C_{1-4} alkyl, or
ii) $-NR_2$,

2) $-(CH_2)_q-NR^6R^7$ wherein q is 1-5, and R^6 and R^7 are defined above, except that R^6 or R^7 are not H or unsubstituted C_{1-6} alkyl, or

3) benzofuryl, indolyl, azacycloalkyl, azabicyclo C_{7-11} cycloalkyl, or benzopiperidinyl, unsubstituted or substituted with C_{1-4} alkyl;

REPLACEMENT SHEET
04 FEB 1997
THIS SHEET HAS BEEN REPLACED

* C L M S *

B is absent, or -NH-C(=Z)-R^8,

wherein R^8 is

1) -CH(CH$_3$)$_2$,
2) -CH(CH$_3$)(CH$_2$CH$_3$), or
3) -phenyl;

J^1 and J^2 are independently

1) -YR^9 wherein

Y is -O- or -NH-, and

R^9 is

a) hydrogen,

b) C$_{1-6}$ alkyl, unsubstituted or substituted with one or more of

i) -NR$_2$,

ii) -OR,

iii) -NHSO$_2$C$_{1-4}$ alkyl,

iv) -NHSO$_2$ aryl, or -NHSO$_2$(dialkylaminoaryl),

v) -CH$_2$OR,

vi) -C$_{1-4}$ alkyl,

vii) -C(=O)-OR,

viii) -C(=O)-NR$_2$,

ix) -NH-C(=NH)-NR$_2$ or -NH-C(=N-CN)-NR$_2$,

x) -NH-C(=O)-R^{13}, wherein R^{13} is

REPLACEMENT SHEET
04 FEB 1997
THIS SHEET HAS BEEN REPLACED

* C L M S *

A) -H,
B) $-C_{1-4}$ alkyl,
C) -aryl,
D) -heterocycle, or
E) -NH-, -O- or $-(CH_2)_n-$ wherein n is zero, 1, 2 or 3, substituted with
I) $-C_{1-4}$ alkyl, unsubstituted or substituted with one or more of aryl or heterocycle, or
II) aryl, unsubstituted or substituted with heterocycle,

xi) $-NR_3^{\oplus}$ $A^{\ominus}$ wherein $A^{\ominus}$ is a counterion,

xii) $-NR^{10}R^{11}$ wherein R^{10} and R^{11} are the same or different and are C_{1-5} alkyl joined together directly to form a 5-7 membered heterocycle containing up to one additional heteroatom selected from -O-, -S-, or -NR-,

xiii) -aryl,

xiv) -CHO,

xv) $-OP(O)(OR_x)_2$,

xvi) $-O-\overset{O}{\overset{\|}{C}}-C_{1-4}$alkyl substituted with one or more of amine or quaternary amine, or $-O-((CH_2)_mO)_n-R$, or $-OP(O)(OR_x)_2$,

REPLACEMENT SHEET
04 FEB 1997
THIS SHEET HAS BEEN REPLACED

* C L M S *


xvii) $-\overset{O}{\overset{\|}{C}}-R$, or

xviii) $-\overset{O}{\overset{\|}{C}}-NH-CH_2$-heterocycle, or

c) $-((CH_2)_mO)_nCH_3$ or $-((CH_2)_mO)_n H$, wherein m and n are defined above,

2) $-N(R^9)_2$,

3) $-NR^{10}R^{11}$ wherein R^{10} and R^{11} are defined above, or

4) 

wherein Y, R^9 and n are defined above; and

R^{12} is

1) hydrogen,

2) aryl, unsubstituted or substituted with one or more of

a) R^{14}, wherein R^{14} is

i) halo,

ii) $-OR$,

iii) $-\overset{O}{\overset{\|}{C}}NR_2$,

iv) $-CH_2NR_2$,

v) $-SO_2NR_2$,

vi) $-NR_2$,

vii) $-NH\overset{O}{\overset{\|}{C}}R$,

viii) $-C_{1-4}$ alkyl,

ix) phenyl

REPLACEMENT SHEET
04 FEB 1997
THIS SHEET HAS BEEN REPLACED

* C L M S *

x) $-CF_3$,

xi) $-\overset{R}{\underset{|}{N}}-SO_2R$,

xii) $-OP(O)(OR_x)_2$, or

xiii) $-\overset{O}{\overset{\|}{C}}OR$,

b) $-C_{1-4}$ alkyl-NR_2, or

c) $-O-\overset{O}{\overset{\|}{C}}-C_{1-4}$alkyl substituted with one or more of amine or quaternary amine or $-OP(O)(OR_x)_2$,

3) heterocycle, the ring or rings being unsubstituted or substituted with one or more of

a) R^{14}, as defined above,

b) $-OC_{1-4}$ alkenyl,

c) phenyl-C_{1-4} alkyl,

d) $-O-\overset{O}{\overset{\|}{C}}-C_{1-4}$alkyl substituted with one or more of amine or quaternary amine, or $-OP(O)(OR_x)_2$, or $-O((CH_2)_mO)_n-R$, or

e) $-O-\overset{O}{\overset{\|}{C}}-O-((CH_2)_mO)_n-R$, or

4) A 5 to 7 membered carbocyclic or 7-10 membered bicyclic carbocyclic ring, the carbocyclic ring being unsubstituted or substituted with one or more of

a) R^{14}, as defined above,

b) $-CH_2OR$,

REPLACEMENT SHEET
04 FEB 1997
THIS SHEET HAS BEEN REPLACED

c) $-(CH_2)_n-NR_2$, C_{5-16}alkyl, pyridine, $-(CH_2)_nNR-(CH_2)_n-NR_2$, $-(CH_2)_n-\overset{O}{\overset{\|}{C}}-OR$, $-((CH_2)_mO)_n-R$, quinuclidiniumyl substituted with R, piperazine-C_{1-4}alkyl-benzyl substituted once or more with R, or morpholino-C_{1-4}alkyl-benzyl,

d) $-O-\overset{O}{\overset{\|}{C}}-C_{1-4}$alkyl substituted with one or more of amine or quaternary amine, $-OP(O)(OR_x)_2$, or $-O-((CH_2)_mO)_n-R$,

e) $-O-\overset{O}{\overset{\|}{C}}-O-((CH_2)_mO)_n-R$, or

f) $-C_{1-4}$alkyl-phenyl;

or a pharmaceutically acceptable salt thereof.

2. A compound according to Claim 1 wherein:

R^1 and R^2 are joined together to form with the nitrogen to which R^1 is attached a 3 to 10 membered monocyclic or bicyclic saturated ring system which consists of the nitrogen to which R^1 is attached and from 2 to 9 carbon atoms, and is unsubstituted or substituted with

1) hydroxy,
2) C_{1-4} alkyl unsubstituted or substituted with one or more of
   a) hydroxy,
   b) C_{1-3} alkoxy,
   c) aryl,

REPLACEMENT SHEET
04 FEB 1997
THIS SHEET HAS BEEN REPLACED

d) a 5-7 membered cycloalkyl group unsubstituted or substituted with one or more of
   i) halo,
   ii) hydroxy,
   iii) C_{1-3} alkoxy, or
   iv) aryl,
e) heterocycle, or
f) $-NR_2$,

3) C_{1-3} alkoxy,

4) $-NH-\overset{O}{\overset{\|}{C}}OC_{1-3}alkyl$,

5) $-NH-\overset{O}{\overset{\|}{C}}-C_{1-3}alkyl$,

6) $-NH-SO_2C_{1-3}alkyl$,

7) -W-aryl, or

8) $-W-\overset{O}{\overset{\|}{C}}-aryl$,

wherein W is -O-, -S-, or -NH-; or

R^1 and R^2 are joined together to form with the nitrogen to which R^1 is attached a 3 to 10 membered monocyclic or bicyclic saturated ring system which consists of the nitrogen to which R^1 is attached, from 1 to 8 carbon atoms and one or more unsubstituted or substituted heteroatom selected from

1) $-\overset{}{\underset{|}{N}}-$
   $V-R^1$,

wherein V is absent or $-\overset{O}{\overset{\|}{C}}-Q-$ or $-SO_2-Q-$, R^1 is defined as above for when R^1 is independent from and not joined to R^2, and wherein Q is absent or -O-, -NR-, or

* C L M S *

REPLACEMENT SHEET
04 FEB 1997
THIS SHEET HAS BEEN REPLACED

heterocycle optionally substituted with $-C_{1-4}$alkyl,

2) $-\overset{|}{N}-$
C_{1-4} alkenyl, unsubstituted or substituted with aryl,

3) $-S(O)_p-$,
wherein p is zero, 1 or 2, or

4) $-O-$; or

R^1 and R^2 are joined together to form with the nitrogen to which R^1 is attached a 3 to 10 membered monocyclic or bicyclic saturated ring system, which consists of the nitrogen to which R^1 is attached and from 2 to 9 carbon atoms, in which the saturated ring system is fused to a phenyl ring and the phenyl ring is unsubstituted or substituted with one or more of

1) C_{1-3} alkoxy,
2) hydroxy,
3) C_{1-4} alkyl, or
4) $-NHR^1$,
wherein R^1 is defined as above for when R^1 is independent from and not joined to R^2.

3. A compound according to Claim 2 wherein:

R^1 and R^2 are joined together to form with the nitrogen to which R^1 is attached a 3 to 10 membered monocyclic or bicyclic saturated ring system which consists of the nitrogen to which R^1 is attached and from 2 to 9 carbon atoms, and is unsubstituted or substituted with

1) hydroxy,
2) C_{1-4} alkyl unsubstituted or substituted with one or more of

* C L M S *

REPLACEMENT SHEET
04 FEB 1997
THIS SHEET HAS BEEN REPLACED

a) hydroxy,
b) C_{1-3} alkoxy,
c) aryl,
d) a 5-7 membered cycloalkyl group unsubstituted or substituted with one or more of
i) halo,
ii) hydroxy,
iii) C_{1-3} alkoxy, or
iv) aryl,
e) heterocycle, or
f) $-NR_2$,

3) C_{1-3} alkoxy,

4) $-NH-\overset{O}{\overset{\|}{C}}OC_{1-3}alkyl$,

5) $-NH-\overset{O}{\overset{\|}{C}}-C_{1-3}alkyl$,

6) $-NH-SO_2C_{1-3}alkyl$,

7) -W-aryl, or

8) $-W-\overset{O}{\overset{\|}{C}}-aryl$,

wherein W is -O-, -S-, or -NH-; or

R^1 and R^2 are joined together to form with the nitrogen to which R^1 is attached a 3 to 10 membered monocyclic or bicyclic saturated ring system which consists of the nitrogen to which R^1 is attached, from 1 to 8 carbon atoms and one or more unsubstituted or substituted heteroatom selected from

1) $-\underset{\underset{V-R^1}{|}}{N}-$,

wherein V is absent or $-\overset{O}{\overset{\|}{C}}-Q-$ or $-SO_2-Q-$,

REPLACEMENT SHEET
04 FEB 1997
THIS SHEET HAS BEEN REPLACED

R^1 is defined as above for when R^1 is independent from and not joined to R^2, and wherein Q is absent or -O-, -NR-, or heterocycle optionally substituted with $-C_{1-4}$alkyl,

2) $-S(O)_p-$, wherein p is zero, 1 or 2, or

3) -O-;

R^3 is benzyl, unsubstituted or substituted with one or more of

a) hydroxy,

b) $-NO_2$, or $-NR_2$,

c) C_{1-4}alkyl,

d) C_{1-3} alkoxy, unsubstituted or substituted with one or more of -OH or C_{1-3} alkoxy,

e) $-\overset{\overset{O}{\|}}{C}NR_2$,

f) $-CH_2NR_2$,

g) $-CH_2NH\overset{\overset{O}{\|}}{C}R$,

h) $-CF_3$,

i) $-NH\overset{\overset{O}{\|}}{C}R$,

j) $-NRSO_2R$,

k) $-OP(O)(OR_x)_2$, or

l) $-R^5$;

and B is absent.

* C L M S *

REPLACEMENT SHEET
04 FEB 1997
THIS SHEET HAS BEEN REPLACED

4. A compound according to Claim 3 wherein:

X is -OH;


Z is -O;

R^1 and R^2 are joined together to form with the nitrogen to which R^1 is attached a 3 to 10 membered monocyclic or bicyclic saturated ring system which consists of the nitrogen to which R^1 is attached and from 2 to 9 carbon atoms, and is unsubstituted or substituted with -W-aryl or -W-C(=O)-aryl; or

R^1 and R^2 are joined together to form with the nitrogen to which R^1 is attached a 3 to 10 membered monocyclic or bicyclic saturated ring system which consists of the nitrogen to which R^1 is attached, from 1 to 8 carbon atoms and one of -N(V-R^1)- ,

wherein V is absent or -C(=O)-Q- or -SO_2-Q-,
R^1 is defined as above for when R^1 is independent from and not joined to R^2,
and wherein Q is absent or -O-, -NR- or heterocycle optionally substituted with -C_{1-4}alkyl;


R^3 is benzyl, unsubstituted or substituted with one or more of (1) hydroxy, (2) C_{1-3} alkoxy substituted with one or more of -OH or (3)


;



REPLACEMENT SHEET
04 FEB 1997
THIS SHEET HAS BEEN REPLACED

J^1 is $-NH-C_{1-4}$alkyl; and
J^2 is



5. The compound

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(2-(3(S)-N'-(t-butylcarboxamido)-(4aS,8aS)-decahydroisoquinoline)yl)-pentaneamide,

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(1-(4-carbobenzyloxy-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide,

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-(2-(4-morpholinyl)ethoxy)phenyl)methyl)-4(S)-hydroxy-5-(2-(3(S)-N'-(t-butylcarboxamido)-(4aS,8aS)-decahydroisoquinoline)yl)-pentaneamide,

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-(2-(4-morpholinyl)ethoxy)phenyl)methyl)-4(S)-hydroxy-5-(1-(4-carbobenzyloxy-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide,

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-((2-hydroxy)-ethoxy)phenyl)methyl)-4(S)-hydroxy-5-(2-(3(S)-N'-(t-butylcarboxamido)-(4aS,8aS)-decahydroisoquinoline)-yl)-pentaneamide,

CLMS

REPLACEMENT SHEET
04 FEB 1997
THIS SHEET HAS BEEN REPLACED

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-((4-((2-hydroxy)-ethoxy)phenyl)methyl)-4(S)-hydroxy-5-(1-(4-carbobenzyloxy-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide,

N-(4(S)-3,4-dihydro-1H-2,2-dioxobenzothiopyranyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(2-(3(S)-N'-(t-butyl-carboxamido)-(4aS,8aS)-decahydroisoquinoline)yl)-pentaneamide,

N-(4(S)-3,4-dihydro-1H-2,2-dioxobenzothiopyranyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(1-(4-carbobenzyl-oxy-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide,

N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(1-(4-(3-pyridylmethyl)-2(S)-N'-(t-butyl-carboxamido)-piperazinyl))-pentaneamide, or

a pharmaceutically acceptable salt thereof.

6. A pharmaceutical composition comprising a compound of Claim 1 and a pharmaceutically acceptable carrier.

7. The use of a compound of Claim 1 for the preparation of a medicament useful for treating AIDS in a patient receiving AIDS therapy.

* C L M S *

THIS SHEET HAS BEEN REPLACED

8. The use of a compound of Claim 1 for the preparation of a medicament for preventing or treating infection by HIV in a patient.

9.. The use of a compound of Claim 1 for the preparation of a medicament useful for inhibiting HIV protease in a patient receiving HIV protease therapy.

REPLACEMENT SHEET
04 FEB 1997

This Sheet Has Been Replaced

THIS SHEET HAS BEEN REPLACED


REPLACEMENT SHEET
04 FEB 1997

TITLE OF THE INVENTION

HIV PROTEASE INHIBITORS USEFUL FOR THE TREATMENT OF AIDS

ABSTRACT OF THE DISCLOSURE

Compounds of formula



where R^1 and R^2 are independently hydrogen or optionally-substituted C_{1-4}alkyl or aryl, or R^1 and R^2 are joined together to form a monocyclic or bicyclic ring system, are HIV protease inhibitors.

These compounds are useful in the prevention or treatment of infection by HIV and in the treatment of AIDS, either as compounds, pharmaceutically acceptable salts, pharmaceutical composition ingredients, whether or not in combination with other antivirals, immunomodulators, antibiotics or vaccines. Methods of treating AIDS and methods of preventing or treating infection by HIV are also described.

This Sheet Has Been Replaced


* C L M S *

REPLACEMENT SHEET
20 DEC 1997

8. The use of a compound of Claim 1 for the preparation of a medicament for preventing or treating infection by HIV in a patient.


9. The use of a compound of Claim 1 for the preparation of a medicament useful for inhibiting HIV protease in a patient receiving HIV protease therapy.

10. A process for making a compound of the formula I



I

comprising reacting a compound (i) of the formula



i

with R^1-X; wherein R^1 is C_{1-4} alkyl unsubstituted or substituted with one or more of

(1) aryl unsubstituted or substituted with one or more of C_{1-4} alkyl, amino, hydroxy or aryl;

(2) a 5-7 membered cycloalkyl group unsubstituted or substituted with one

REPLACEMENT SHEET
20 DEC 1997

or more of halo, C_{1-3} alkoxy or aryl; or

(3) heterocycle unsubstituted or substituted with one or more of oxo, halo, C_{1-4} alkoxy, C_{1-4} alkyl, $-C(O)OC_{1-3}$ alkyl, $-NHC(O)C_{1-3}$ alkyl, or Boc; and

X is selected from Cl, Br or I;

to give the compound of formula I.


11. The process of Claim 10, wherein

R^1 is C_{1-4} alkyl unsubstituted or substituted with a heterocycle wherein the heterocycle is unsubstituted or substituted with one or more of oxo, halo, C_{1-4} alkoxy, C_{1-4} alkyl, $-C(O)OC_{1-3}$ alkyl, $-NHC(O)C_{1-3}$ alkyl, or Boc.

12. The process of Claim 11, wherein

R^1 is C_{1-4} alkyl substituted with a heterocycle, wherein the heterocycle is selected from piperidinyl, pyridyl, thienyl, pyrrolyl, thiazolyl, imidazolyl, furyl, benzimidazolyl, pyrazinyl, isoxazolyl, pyridazinyl, or quinolinyl, and wherein the heterocycle is unsubstituted or substituted with one or more of oxo, halo, C_{1-4} alkoxy, C_{1-4} alkyl, $-C(O)OC_{1-3}$ alkyl, $-NHC(O)C_{1-3}$ alkyl, or Boc.


13. The process of Claim 12 for making a compound J of the formula



J


REPLACEMENT SHEET
20 DEC 1997

comprising reacting a compound (i) of the formula




i

with a compound of the formula




to give the compound J.

14. The process of Claim 13 for making a compound J of the formula



J

comprising reacting a compound (i) of the formula




i

with 3-picolyl chloride to give the compound J.

REPLACEMENT SHEET
20 DEC 1997


15. A compound (i) of the formula:



i

or a salt thereof.

16. A compound J of the formula:



J

and the pharmaceutically acceptable salts thereof.


REPLACEMENT SHEET
20 DEC 1997

TITLE OF THE INVENTION

HIV PROTEASE INHIBITORS USEFUL FOR THE TREATMENT OF AIDS

ABSTRACT OF THE DISCLOSURE

Compounds of formula



where R^1 and R^2 are independently hydrogen or optionally-substituted C_{1-4}alkyl or aryl, or R^1 and R^2 are joined together to form a monocyclic or bicyclic ring system, are HIV protease inhibitors.

These compounds are useful in the prevention or treatment of infection by HIV and in the treatment of AIDS, either as compounds, pharmaceutically acceptable salts, pharmaceutical composition ingredients, whether or not in combination with other antivirals, immunomodulators, antibiotics or vaccines. Methods of treating AIDS and methods of preventing or treating infection by HIV are also described.