


Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6



Difference between the posology 100mg/kg and the posology based on alcohol consumption (in % of the posology 100mg/kg) - Patients without hepatic impairment
80%
70%
60%
50%
40%
30%
20%
10%
0%
-10%
-20%
LOW
MEDIUM
HIGH
VERY HIGH
Daily alcohol consumption
100
90
80
70
60
50
40
Weight (kg)
-20%--10%
-10%-0%
0%-10%
10%-20%
20%-30%
30%-40%
40%-50%
50%-60%
60%-70%
70%-80%

Figure 7



Difference between the posology 50mg/kg and the posology based on alcohol consumption (in % of the posology 50mg/kg) - Patients with hepatic impairment
80%
70%
60%
50%
40%
30%
20%
10%
0%
-10%
-20%
LOW
MEDIUM
HIGH
VERY HIGH
Daily alcohol consumption
100
90
80
70
60
50
40
Weight (kg)
-20%--10%
-10%-0%
0%-10%
10%-20%
20%-30%
30%-40%
40%-50%
50%-60%
60%-70%
70%-80%

Figure 8



Difference between the posology 100mg/kg and the posology based on alcohol consumption (in % of the posology 100mg/kg) - Patients with hepatic impairment
90%
80%
70%
60%
50%
40%
30%
20%
LOW
MEDIUM
HIGH
VERY HIGH
Daily alcohol consumption
100
90
80
70
60
50
40
Weight (kg)
40%-50%
50%-60%
60%-70%
70%-80%
80%-90%

Figure 9