

**COMBINATIONS OF EXTRACTS OF *SERENOA REPENS* AND
LIPOPHILIC EXTRACTS OF *ZINGIBER OFFICINALIS* AND
ECHINACEA ANGUSTIFOLIA, THE USE THEREOF, AND
FORMULATIONS CONTAINING THEM**

The present invention relates to the combination of an extract of *Serenoa repens* with a lipophilic extract of *Zingiber officinalis* and a lipophilic extract of *Echinacea angustifolia* or *Zanthoxylum bungeanum* which is useful in the treatment of urinary incontinence, the symptoms of benign prostate hypertrophy, prostatitis and prostatism, and pelvic pain.

Prior art

Prostate hypertrophy, an age-dependent physiological event caused by a modified hormone ratio between androgens and oestrogens, is often characterised by a series of unpleasant events such as dysuria, pollakiuria and incontinence, associated with inflammatory states and pelvic pain.

Many of the supposed symptoms associated with increased prostate volume also appear to be associated with bladder dysrhythmia, which is common in both men and women, and characterised by similar symptoms.

Serenoa repens extracts are widely used in the clinical and nutritional fields to treat the symptoms of benign prostate hypertrophy in men and urinary incontinence in women (Hale E.M. (1898). Saw palmetto (*Sabal serrulata*, *Serenoa Serrulata*): its History, Botany, Chemistry, Pharmacology, Provings, Clinical Experience and Therapeutical Applications. Philadelphia: Boericke and Taffel).

Zingiber officinalis extract is traditionally used to treat dyspepsia, nausea and vomiting and to a lesser extent indigestion, flatulence, diarrhoea, cough and inflammation. Gingerols, the active ingredients of the plant, interact with NFkB and TNF- α . They also control gastric voiding, with the related consequences on the digestion (Rhode J. Et al., BMC Complement. Altern. Med. 7, 44, 2007; El-

Sharaki et al., Food Chem. Toxicol 47, 1584, 2009; Micklefield G.H. et al., Int. J. Clin. Pharmacol. Ther. 37, 341, 1999).

The extracts of this plant are particularly useful in the treatment of nausea.

Description of the invention

5 It has now surprisingly been found that by combining *Serenoa repens* extract with a lipophilic extract of *Zingiber officinalis* and a lipophilic extract of *Echinacea angustifolia* or *Zanthoxylum bungeanum*, the urogenital symptoms of both men and woman regress completely, with a rapid reduction of inflammation in prostatism and pelvic pain in general which has never been observed with the
10 sole use of the extracts administered individually.

The present invention therefore relates to compositions containing a *Serenoa repens* extract, a lipophilic extract of *Zingiber officinalis* and a lipophilic extract of *Echinacea angustifolia* or *Zanthoxylum bungeanum*.

The invention also relates to said compositions for use in the treatment of
15 urinary incontinence, benign prostate hypertrophy, prostatitis and prostatism, and pelvic pain.

Serenoa repens extract, which is rich in fatty acids, lipophilic extract of *Zingiber officinalis* and lipophilic extract of *Echinacea angustifolia* or *Zanthoxylum bungeanum* are preferably obtained by extraction with carbon
20 dioxide under supercritical conditions.

The *Zingiber officinalis* extract used in the present invention, prepared by extraction from the rhizomes with carbon dioxide under supercritical conditions, is characterised by the fact that it contains 20 to 35% gingerols, preferably 25%.

The *Serenoa repens* extracts according to the invention contain between 60
25 and 90% fatty acids, preferably about 80%. The extracts can also be prepared with solvents selected from hexane, ethanol and acetone. Alternatively, the extract obtained by pressing the fresh fruit can be used. An example of a lipophilic extract of *Serenoa* is disclosed in EP 0250953.

The lipophilic extracts of *Echinacea angustifolia* or *Zanthoxylum bungeanum* contain between 15 and 35%, preferably 25%, of isobutylamides, and are prepared with carbon dioxide under supercritical conditions as reported in EP 0464298 and WO 00/02570. Isobutylamides are ligands of the cannabinoid
5 receptors, particularly CB2, and agonists of TRPV1, the vanilloid receptor with a painkilling function.

The extracts of *Serenoa repens*, *Zingiber officinalis* and *Echinacea angustifolia* or *Zanthoxylum bungeanum* are preferably present in the compositions according to the invention in the ratios of 1:0.25-0.5:0.05-0.1, which minimise the
10 side effects and consequently allow safe chronic treatment.

The therapeutic effect was evaluated on the basis of internationally accepted scores according to the NIH-CPSI (National Institute of Health - Chronic Prostatitis Symptoms Index) which monitors CP-CPPS (chronic prostatitis/chronic pelvic pain syndrome), which is associated with urinary symptoms such as
15 pollakiuria, dysuria, nocturia and sexual dysfunctions which have lasted for at least three to six months and are present in the absence of urogenital infections, tumours or anatomical abnormalities. The evaluations are functional in the absence of specific markers, and require an adequate number of patients. The scale comprises an NIH-CPSI range of 0 to 43 (43 for the most serious cases). A population with
20 variations of 0 to 30 in the inclusion criteria was evaluated.

The trial was carried out on 80 patients aged between 55 and 70 years (63 ± 7) with category IIIB CPPS. The patients were divided into 4 groups after randomisation; the first group was treated with capsules of *Serenoa repens* lipophilic extract (120 mg, 3 capsules a day), the second with the combination
25 described in example 1 (3 capsules a day), the third with the mixture of *Echinacea angustifolia*/*Zingiber officinalis* lipophilic extract (3 capsules a day) prepared as described in EP 2379095 and administered in groundnut oil, and the fourth with a placebo. The data are reported in Table 1.

Table 1

Variations in NIH-CPSI for pelvic pain and quality of life after treatment with Combination 1 and the ingredients thereof

		NIH-CPSI TOTAL	NIH-CPSI PAIN	NIH-CPSI QUALITY OF LIFE
<i>Serenoa repens</i>	Start	23.1 (1.8)	9.8 (1.6)	8.0 (0.9)
	3 months	14.3 (1.3)	7.3 (1.1)	3.8 (1.1)
	6 months	10.2 (1.1)	6.5 (1.2)	4.5 (0.8)
Combination 1	Start	22.4 (1.5)	9.7 (1.5)	7.8 (0.9)
	3 months	11.3 (1.1)	3.2 (1.0)	2.2 (0.9)
	6 months	8.2 (1.2)	1.5 (1.1)	2.1 (0.8)
Echinacea/ginger	Start	22.7(11.4)	9.1 (1.1)	7.9 (1.2)
	3 months	17.4 (1.5)	8.2 (1.2)	6.2 (1.1)
	6 months	12.6 (1.4)	7.3 (1.1)	6.2 (1.1)
Placebo	Start	24.2 (1.7)	9.2 (1.3)	7.6 (1.1)
	3 months	15.1 (1.6)	7.3 (1.2)	6.9 (0.8)
	6 months	11.2 (1.5)	6.5 (1.1)	5.9 (1.0)

5 A second group of 60 patients was treated similarly to the first, evaluating the changes in the IPSS (International Prostate Symptom Score) and quality of life; the *Serenoa repens* extract, the combination and the placebo were compared for six months. The results are reported in Table 2.

Table 2

	Start	2 months	4 months	Final	MODIFICATIONS
SYMPTOMS (score)					
<i>Serenoa repens</i>	16.5 ±4.7	12.9 ± 4.4	11.9 ±5.0	12.1 ± 5.3	-4.4 ±5.0 (p<0.036)
Combination 1	17.1 ±4.6	9.5 ±4.2	8.6 ±5.2	7.6 ±5.8	-9.5 ±5.2 (p<0.001)
Placebo	15.9 ±4.5	12.3 ±5.2	13.2 ±5.3	13.5 ± 6.4	-2.4 ±5.4
QUALITY OF LIFE (score)					
<i>Serenoa repens</i>	3.3±1.2	3.1 ±1.3	2.7 ±1.1	2.6 ±1.4	-0.7 ±1.3
Combination 1	3.2 ±1.2	2.8 ±1.1	2.0 ±1.2	1.8±1.1	-1.4 ±1.1 (p<0.05)
Placebo	3.1 ±1.2	2.9 ±1.0	2.8 ±1.2	2.8 ±1.1	-0.3 ±1.1

Values expressed as mean ± S.D. p<0.05 (significance vs. placebo)

The results of the studies demonstrate the synergy of the composition compared with the individual ingredients.

The pharmaceutical compositions mainly take the form of soft gelatin or cellulose capsules suitable for oils that disperse rapidly in the stomach, or capsules or tablets after microencapsulation of the oils.

A preferred lipophilic formulation uses as dispersing agent oils rich in ω -3 fatty acids, which promote rapid absorption of the active ingredient. Examples of said oils include *Oenothera biennis* oil, *Linum usitatissimum* oil and *Ribes nigrum* oil, and natural or semi-synthetic fish oils. Said compositions are mainly used in curative and maintenance treatments, and to reduce pelvic pain.

According to a preferred aspect, the compositions according to the invention will also contain antioxidants, such as palmitoyl ascorbate, catechin and gallic

polyphenols, or lycopene and extracts containing it, and trace elements such as zinc, selenium and manganese.

These compositions can be prepared according to conventional methods, as described in “Remington’s Pharmaceutical Handbook”, Mack Publishing Co.,
5 N.Y., USA, using suitable excipients. Some formulation examples are reported below.

EXAMPLE 1 - SOFT GELATIN CAPSULES

<i>Serenoa repens</i> extract	120 mg
<i>Zingiber officinalis</i> lipophilic extract	25 mg
<i>Echinacea angustifolia</i> lipophilic extract	5 mg
Soya lecithin	10 mg
Palmitoyl ascorbate	20 mg
<i>Oenothera biennis</i> lipophilic extract	150 mg

10 EXAMPLE 2 - CELLULOSE CAPSULES FOR OILS

<i>Serenoa repens</i> extract	120 mg
<i>Zingiber officinalis</i> lipophilic extract	50 mg
<i>Echinacea angustifolia</i> lipophilic extract	10 mg
Soya lecithin	10 mg
<i>Oenothera biennis</i> lipophilic extract	150 mg

EXAMPLE 3 - TABLETS

<i>Serenoa repens</i> extract	120 mg
<i>Zingiber officinalis</i> lipophilic extract	50 mg
<i>Echinacea angustifolia</i> lipophilic extract	10 mg
Soya lecithin	10 mg
Palmitoyl ascorbate	20 mg
<i>Oenothera biennis</i> lipophilic extract	150 mg

EXAMPLE 4 - SOFT GELATIN CAPSULES

5

<i>Serenoa repens</i> extract	130 mg
<i>Zingiber officinalis</i> lipophilic extract	25 mg
<i>Echinacea angustifolia</i> lipophilic extract	5 mg
Soya lecithin	10 mg
Palmitoyl ascorbate	20 mg
<i>Lycopersicon esculentum</i> extract (10% lycopene)	50 mg
<i>Oenothera biennis</i> lipophilic extract	150 mg

EXAMPLE 5 - SOFT GELATIN CAPSULES

<i>Serenoa repens</i> extract	120 mg
<i>Zingiber officinalis</i> lipophilic extract	25 mg
<i>Zanthoxylum bungeanum</i> lipophilic extract	5 mg
Soya lecithin	10 mg
Palmitoyl ascorbate	20 mg
<i>Oenothera biennis</i> lipophilic extract	150 mg

CLAIMS

1. Compositions comprising a *Serenoa repens* extract, a *Zingiber officinalis* lipophilic extract and an *Echinacea angustifolia* or *Zanthoxylum bungeanum*
5 lipophilic extract.
2. Compositions according to claim 1 wherein the extracts of *Serenoa repens*, *Zingiber officinalis*, *Echinacea angustifolia* and *Zanthoxylum bungeanum* are prepared by extraction with supercritical carbon dioxide.
3. Compositions according to claim 1 or 2 wherein the *Serenoa repens* extract:
10 *Zingiber officinalis* lipophilic extract: *Echinacea angustifolia* or *Zanthoxylum bungeanum* lipophilic extract ratios are 1:0.25-0.5:0.05-0.1.
4. Compositions according to one or more of claims 1 to 3 further comprising oils rich in ω -3 fatty acids.
5. Compositions according to claim 4 wherein the oils are selected from
15 *Oenothera biennis*, *Linum usitatissimum* or *Ribes nigrum* oils and natural or semi-synthetic fish oils.
6. Compositions according to one or more of claims 1 to 5 further containing antioxidants and trace elements.
7. The compositions of claims 1-6 for use in the treatment of urinary
20 incontinence, benign prostate hypertrophy, prostatitis and prostatic, and pelvic pain.