

Novel Heterocyclic Carboxamides as Modulators of Kinase Activity

5 ***Field of the invention***

The invention relates to a series of heterocyclic carboxamide compounds that are useful in the treatment of hyperproliferative diseases, such as cancer, in mammals. Also encompassed by the present invention is the use of such compounds in the treatment of hyperproliferative diseases in mammals, especially humans, and pharmaceutical
10 compositions containing such compounds.

Summary of the related art

Protein kinases constitute a large family of structurally related enzymes that are responsible for the control of a wide variety of signal transduction processes within the
15 cell (Hardie, G. and Hanks, S. (1995) The Protein Kinase Facts Book. I and II, Academic Press, San Diego, CA). The kinases may be categorized into families by the substrates they phosphorylate (e.g., protein-tyrosine, protein-serine/threonine, lipids, etc.).

Sequence motifs have been identified that generally correspond to each of these kinase families (e.g., Hanks, S.K., Hunter, T., FASEB J., 9:576-596 (1995); Knighton, et al.,
20 Science, 253:407-414 (1991); Hiles, et al., Cell, 70:419-429 (1992); Kunz, et al., Cell, 73:585-596 (1993); Garcia-Bustos, et al., EMBO J., 13:2352-2361 (1994)).

Protein kinases may be characterized by their regulation mechanisms. These mechanisms include, for example, autophosphorylation, transphosphorylation by other kinases, protein-protein interactions, protein-lipid interactions, and protein-polynucleotide
25 interactions. An individual protein kinase may be regulated by more than one mechanism.

Kinases regulate many different cell processes including, but not limited to, proliferation, differentiation, apoptosis, motility, transcription, translation and other signalling
30 processes, by adding phosphate groups to target proteins. These phosphorylation events act as molecular on/off switches that can modulate or regulate the target protein biological function. Phosphorylation of target proteins occurs in response to a variety of extracellular signals (hormones, neurotransmitters, growth and differentiation factors, etc.), cell cycle events, environmental or nutritional stresses, etc. The appropriate protein
35 kinase functions in signalling pathways to activate or inactivate (either directly or

indirectly), for example, a metabolic enzyme, regulatory protein, receptor, cytoskeletal protein, ion channel or pump, or transcription factor. Uncontrolled signalling due to defective control of protein phosphorylation has been implicated in a number of diseases, including, for example, inflammation, cancer, allergy/asthma, diseases and conditions of the immune system, diseases and conditions of the central nervous system, and angiogenesis.

Protein kinase 70S6K, the 70 kDa ribosomal protein kinase p70S6K (also known as SK6, p70/p85 S6 kinase, p70/p85 ribosomal S6 kinase and p70S6K), is a member of the AGC subfamily of protein kinases. p70S6K is a serine-threonine kinase that is a component of the phosphatidylinositol 3 kinase (PI3K)/AKT pathway. p70S6K is downstream of PI3K, and activation occurs through phosphorylation at a number of sites in response to numerous mitogens, hormones and growth factors. p70S6K activity is also under the control of a mTOR-containing complex (TORC1) since rapamycin acts to inhibit p70S6K activity. p70S6K is regulated by PI3K downstream targets AKT and PKC. AKT directly phosphorylates and inactivates TSC2, thereby activating mTOR. In addition, studies with mutant alleles of p70S6K that inhibited by Wortmannin but not by rapamycin suggest that the PI3K pathway can exhibit effects on p70S6K independent of the regulation of mTOR activity.

The enzyme p70S6K modulates protein synthesis by phosphorylation of the S6 ribosomal protein. S6 phosphorylation correlates with increased translation of mRNAs encoding components of the translational apparatus, including ribosomal proteins and translational elongation factors whose increased expression is essential for cell growth and proliferation. These mRNAs contain an oligopyrimidine tract at their 5' transcriptional start (termed 5'TOP), which has been shown to be essential for their regulation at the translational level.

In addition to its involvement in translation, p70S6K activation has also been implicated in cell cycle control, neuronal cell differentiation, regulation of cell motility and a cellular response that is important in tumor metastases, the immune response and tissue repair. Antibodies to p70S6K abolish the mitogenic response driven entry of rat fibroblasts into S phase, indication that p70S6K function is essential for the progression from G1 to S phase in the cell cycle. Furthermore, inhibition of cell cycle proliferation at the G1 to S

phase of the cell cycle by rapamycin has been identified as a consequence of inhibition of the production of the hyperphosphorylated, activated form of p70S6K.

5 A role for p70S6K in tumor cell proliferation and protection of cells from apoptosis is supported based on its participation in growth factor receptor signal transduction, overexpression and activation in tumor tissues. For example, Northern and Western analyses revealed that amplification of the PS6K gene was accompanied by corresponding increases in mRNA and protein expression, respectively (Cancer Res. (1999) 59: 1408-11-Localization of PS6K to Chromosomal Region 17q23 and
10 Determination of Its Amplification in Breast Cancer).

Chromosome 17q23 is amplified in up to 20% of primary breast tumors, in 87% of breast tumors containing BRCA2 mutations and in 50% of tumors containing BRCA1 mutations, as well as other cancer types such as pancreatic, bladder and neuroblastoma (see M.
15 Barlund, O. Monni, J. Kononen, R. Cornelison, J. Torhorst, G. Sauter, O.-P. Kallioniemi and Kallioniemi A., Cancer Res., 2000, 60:5340-5346). It has been shown that 17q23 amplifications in breast cancer involve the PAT1, RAD51C, PS6K, and SIGMA1B genes (Cancer Res. (2000): 60, pp. 5371-5375).

20 The p70S6K gene has been identified as a target of amplification and overexpression in this region, and statistically significant association between amplification and poor prognosis has been observed. Clinical inhibition of p70S6K activation was observed in renal carcinoma patients treated with CCI-779 (rapamycin ester), an inhibitor of the upstream kinase mTOR. A significant linear association between disease progression
25 and inhibition of p70S6K activity was reported. In response to energy stress, the tumor suppressor LKB1 activates AMPK which phosphorylates the TSC1/2 complex and enables it to inactivate the mTOR/p70S6K pathway. Mutations in LKB1 cause Peutz-Jeghers syndrome (PJS), where patients with PJS are 15 times more likely to develop cancer than the general population. In addition, 1/3 of lung adenocarcinomas harbor
30 inactivating LKB1 mutations. P70S6K has been implicated in metabolic diseases and disorders. It was reported that the absence of p70S6K protects against age- and diet-induced obesity while enhancing insulin sensitivity. A role for p70S6K in metabolic and disorders such as obesity, diabetes, metabolic syndrome, insulin resistance, hyperglycemia, hyperaminoacidemia, and hyperlipidemia is supported based upon the
35 findings.


Compounds described as suitable for p70S6K inhibition are disclosed in, i.a.
WO 03/064397, WO 04/092154, WO 05/054237, WO 05/056014, WO 05/033086, WO
05/117909, WO 05/039506, WO 06/120573, WO 06/136821, WO 06/071819, WO
5 06/131835, WO 08/140947 and PCT/US10/000313.

Description of the invention

It is the object of the present invention to provide novel compounds that modulate kinase
activity. This protein kinase modulation includes, but is not limited to, p70S6K inhibition
10 and AKT inhibition useful in the treatment of hyperproliferative diseases, especially
those related to the hyperactivity of the above mentioned protein kinases, such as
cancer in mammals, with superior pharmacological properties both with respect to their
activities as well as their solubility, metabolic clearance and bioavailability
characteristics.

15 As a result, this invention provides novel, heterocyclic carboxamides and
pharmaceutically acceptable salts, solvates or prodrugs thereof, that are kinase
inhibitors and useful in the treatment of the above mentioned diseases.

20 The compounds are defined by Formula (I) and pharmaceutically acceptable salts,
solvates, solvates of salts, or prodrugs thereof, wherein:



(I)

- 25
- X is N, or C-R⁴;
Y is N-R⁵, O, or absent;

- R¹ is L¹-R⁶, L¹-R⁶-L²-R⁷;
- R² is H, Hal, OH, OA, CN, NH₂, or NHA;
- R³ is H, CH₃, or C(Hal)₃;
- R⁴ is H, Hal, OH, COOH, NH₂, or CN;
- 5 R⁵ is H, LA or monocyclic alkyl having 3, 4, 5, 6, or 7 ring atoms, in which one or two CH₂ groups may be replaced by a -NH- group, or
- R⁵, R¹ together form monocyclic alkyl having 3, 4, 5, 6, or 7 ring atoms, in which one or two CH₂ groups may be replaced by an O atom and/or an -NH-, -NA-, -N(L¹-R⁶)-, -CHA-, -CA₂-, CH(L¹-R⁶)- or -CO- group, and which monocyclic alkyl may be
- 10 substituted by NH₂;
- L¹, L² are independently a single bond, or unbranched or branched alkyl having 1, 2, 3, 4 or 5 C atoms, which may be unsubstituted, or mono- or disubstituted with Hal, OH, NH₂, NH(LA), N(LA)₂, and wherein one or two CH₂ groups may be replaced by an O atom or by a -CO-, -NH-, -N(LA)-, -CONH-, -N(LA)COO- or -NHCO-
- 15 group;
- R⁶ is Ar or monocyclic alkyl having 3, 4, 5, 6, or 7 ring atoms, in which one or two CH₂ groups may be replaced by an O atom and/or an -NH-, -NA-, -CHA-, -CO- or -CONHA- group;
- Ar is a mono- or bicyclic aromatic homo- or heterocycle having 0, 1, 2, 3 or 4 N, O
- 20 and/or S atoms and 5, 6, 8, 9, or 10 skeleton atoms, which may be unsubstituted or, indenpendently of one another, mono-, di- or trisubstituted by Hal, A, OA, OH, NH₂, or NHA;
- A is an unbranched or branched linear or cyclic alkyl having 1, 2, 3, 4, 5 or 6 C atoms, in which one or two CH₂ groups may be replaced by an O atom and/or an
- 25 -NH-, -NHCOAr or -CONHAr group in which 1-3 H atoms may be replaced by Hal, and in which one or two CH₃ groups may be replaced by NH₂, OH, NH(LA) or N(LA)₂ group;
- LA is unbranched or branched, linear alkyl having 1, 2, 3, or 4 C atoms wherein 1, 2 or 3 H atoms may be replaced by Hal (*e.g.*, methyl, ethyl, and trifluoromethyl);
- 30 and
- Hal is F, Cl, Br or I.

In general, all residues which occur more than once may be identical or different, i.e. are independent of one another. Above and below, the residues and parameters have the

35 meanings indicated for the Formula (I), unless expressly indicated otherwise.

Accordingly, the invention relates, in particular, to the compounds of the Formula (I) in which at least one of the said residues has one of the preferred meanings indicated below.

5 Hal denotes fluorine, chlorine, bromine or iodine, in particular fluorine or chlorine.

“A” denotes, for example, methyl, furthermore ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl or tert-butyl, furthermore also pentyl, 1-, 2- or 3-methylbutyl, 1,1-, 1,2- or 2,2-dimethylpropyl, 1-ethylpropyl, hexyl, 1-, 2-, 3- or 4-methylpentyl, 1,1-, 1,2-, 1,3-, 2,2-,
10 2,3- or 3,3-dimethylbutyl, 1- or 2-ethylbutyl, 1-ethyl-1-methylpropyl, 1-ethyl-2-methylpropyl, 1,1,2- or 1,2,2-trimethylpropyl.

“A” further denotes alkyl as defined above, in which one or two CH₂ groups may be replaced by O or S atoms and/or by NH, N(LA), CONH, NHCO or -CH=CH-groups and/or in addition 1-3 H atoms may be replaced by F and/or Cl, such as, for example,
15 trifluoromethyl, pentafluoroethyl, 1,1-difluoromethyl, 1,1,1-trifluoroethyl, methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, isobutoxy, sec-butoxy or tert-butoxy.

In other examples of “A”, one or two CH₃ groups is replaced by OH, SH, NH₂, N(LA)H, N(LA)₂ or CN, such as, for example, N,N'-dimethylaminoalkyl, 2-aminoethyl, 3-amino-propyl, 4-aminobutyl, 5-aminopentyl, 3-aminomethylcyclobutyl or cyanoalkyl.
20 Cyclic A preferably denotes cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or cycloheptyl.

“LA” denotes unbranched or branched, linear alkyl having 1, 2, 3 or 4 C atoms, i.e. methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl or tert-butyl.

25 “Ar” denotes, for example, unsubstituted phenyl, naphthyl or biphenyl, furthermore preferably, for example, phenyl, naphthyl or biphenyl, each of which is mono-, di- or trisubstituted by A, fluorine, chlorine, bromine, iodine, hydroxyl, methoxy, ethoxy, propoxy, butoxy, pentyloxy, hexyloxy, nitro, cyano, formyl, acetyl, propionyl, trifluoro-
30 methyl, amino, methylamino, ethylamino, dimethylamino, diethylamino, benzyloxy, sulfonamido, methylsulfonamido, ethylsulfonamido, propylsulfonamido, butylsulfonamido, dimethylsulfonamido, phenylsulfonamido, carboxyl, methoxycarbonyl, ethoxycarbonyl, aminocarbonyl.

5 “Ar” furthermore denotes o-, m- or p-tolyl, o-, m- or p-ethylphenyl, o-, m- or p-propylphenyl, o-, m- or p-isopropylphenyl, o-, m- or p-tert-butylphenyl, o-, m- or p-hydroxyphenyl, o-, m- or p-nitrophenyl, o-, m- or p-aminophenyl, o-, m- or p-(N-methylamino)phenyl, o-, m- or p-(N-methylaminocarbonyl)phenyl, o-, m- or p-acetamidophenyl, o-,
 10 m- or p-methoxyphenyl, o-, m- or p-ethoxyphenyl, o-, m- or p-ethoxycarbonylphenyl, o-, m- or p-(N,N-dimethylamino)phenyl, o-, m- or p-(N,N-dimethylaminocarbonyl)phenyl, o-, m- or p-(N-ethylamino)phenyl, o-, m- or p-(N,N-diethylamino)phenyl, o-, m- or p-fluorophenyl, o-, m- or p-bromophenyl, o-, m- or p-chlorophenyl, o-, m- or p-(methylsulfonamido)phenyl, o-, m- or p-(methylsulfonyl)phenyl, further preferably 2,3-, 2,4-, 2,5-, 2,6-,
 15 3,4- or 3,5-difluorophenyl, 2,3-, 2,4-, 2,5-, 2,6-, 3,4- or 3,5-dichlorophenyl, 2,3-, 2,4-, 2,5-, 2,6-, 3,4- or 3,5-dibromophenyl, 2,4- or 2,5-dinitrophenyl, 2,5- or 3,4-dimethoxyphenyl, 3-nitro-4-chlorophenyl, 3-amino-4-chloro-, 2-amino-3-chloro-, 2-amino-4-chloro-, 2-amino-5-chloro- or 2-amino-6-chlorophenyl, 2-nitro-4-N,N-dimethylamino- or 3-nitro-4-N,N-dimethylaminophenyl, 2,3-diaminophenyl, 2,3,4-, 2,3,5-, 2,3,6-, 2,4,6- or 3,4,5-tri-
 20 chlorophenyl, 2,4,6-trimethoxyphenyl, 2-hydroxy-3,5-dichlorophenyl, p-iodophenyl, 3,6-dichloro-4-aminophenyl, 4-fluoro-3-chlorophenyl, 2-fluoro-4-bromophenyl, 2,5-difluoro-4-bromophenyl, 3-bromo-6-methoxyphenyl, 3-chloro-6-methoxyphenyl, 3-chloro-4-acetamidophenyl, 3-fluoro-4-methoxyphenyl, 3-amino-6-methylphenyl, 3-chloro-4-acetamidophenyl or 2,5-dimethyl-4-chlorophenyl, (4-methoxyphenyl)methyl, (3-methoxyphenyl)methyl, (4-methoxyphenyl)ethyl, (3-methoxyphenyl)ethyl.

25 “Ar” furthermore preferably denotes 2-, 3- or 4-phenyl, 2-, 3- or 4-phenylmethyl, 2-, 3- or 4-phenylethyl, 2- or 3-furyl, 2- or 3-thienyl, 1-, 2- or 3-pyrrolyl, 1-, 2, 4- or 5-imidazolyl, 1-, 3-, 4- or 5-pyrazolyl, 2-, 4- or 5-oxazolyl, 3-, 4- or 5-isoxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-isothiazolyl, 2-, 3- or 4-pyridyl, 2-, 3- or 4-pyridylmethyl, 2-, 3- or 4-pyridylethyl, 2-, 4-, 5- or 6-pyrimidinyl, 2-, 3-, 5-, or 6-pyrazin-1- or 4-yl, furthermore preferably 1,2,3-triazol-1-, -4- or -5-yl, 1,2,4-triazol-1-, -3- or 5-yl, 1- or 5-tetrazolyl, 1,2,3-oxadiazol-4- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-oxadiazol-2-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,2,4-thiadiazol-3- or -5-yl, 1,2,3-thiadiazol-4- or -5-yl, 3- or 4-pyridazinyl, 1-, 2-, 3-, 4-, 5-, 6- or
 30 7-indolyl, 2-, 3-, 4- or 5-isoindolyl, 2-, 6-, -or 8-purinyl, 1-, 2-, 4- or 5-benzimidazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzopyrazolyl, 2-, 4-, 5-, 6- or 7-benzoxazolyl, 3-, 4-, 5-, 6- or 7-benzisoxazolyl, 2-, 4-, 5-, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 4-, 5-, 6- or 7-benz-2,1,3-oxadiazolyl, 1-, 3-, 4-, 5-, 6-, 7- or 8-isoquinolinyl, 3-, 4-, 5-, 6-, 7- or 8-quinolinyl, 2-, 4-, 5-, 6-, 7- or 8-quinazolinyl, quinoxalin-2-, 3-, 4- or 5-yl, 4-, 5-, or
 35 6-phthalazinyl, 2-, 3-, 5-, 6-, 7- or 8-2H-benzo-1,4-oxazinyl,

further preferably 1,3-benzodioxol-2-, 4- or 5-yl, thiophen-2- or 3-yl, 1,4-benzodioxan-6-yl, 2,1,3-benzothiadiazol-4- or -5-yl or 2,1,3-benzoxadiazol-5-yl, furan-2- or 3-yl, 2,3-dihydro-benzofuran-2-, 3-, 4- or 5-yl,

each of which is unsubstituted or may be mono-, di- or trisubstituted, for example, by
5 carbonyl oxygen, F, Cl, Br, methyl, ethyl, propyl, phenyl, benzyl, -CH₂-cyclohexyl, hydroxyl, methoxy, ethoxy, amino, methylamino, dimethylamino, nitro, cyano, carboxyl, methoxycarbonyl, aminocarbonyl, methylaminocarbonyl, dimethylaminocarbonyl, acetamino, ureido, methylsulfonylamino, formyl, acetyl, aminosulfonyl and/or methyl-sulfonyl.

10

Particularly preferred are compounds of Subformulae 1 to 10 of Formula (I), and pharmaceutically acceptable salts, solvates or prodrugs thereof, wherein

in Subformula 1

15 X is N,
Y is N-R⁵,

in Subformula 2

X is N,
20 Y is N-R⁵,

R⁵, R¹ together form monocyclic alkyl having 3, 4, 5, 6, or 7 ring atoms, in which one or two CH₂ groups may be replaced by an -NH-, -NA-, -N(L¹-R⁶)-, -CHA-, -CA₂-, CH(L¹-R⁶)- or -CO- group, and which monocyclic alkyl may be substituted by NH₂,

25

in Subformula 3

X is N,
Y is N-R⁵,

R⁵, R¹ together form monocyclic alkyl having 4, 5 or 6 ring atoms, in which one CH₂
30 group is replaced by an -N(L¹-R⁶)- group, and which monocyclic alkyl may be substituted by NH₂,

in Subformula 4

X is N,
35 Y is N-R⁵,

R⁵, R¹ together form monocyclic alkyl having 4, 5 or 6 ring atoms, in which one CH₂ group is replaced by an -N(L¹-R⁶)- group, and which monocyclic alkyl may be substituted by NH₂,

L¹ is a bond, -CONH-, -NHCO-, -CONHCH₂-, CH₂CONH-,

5

in Subformula 5

X is N,

Y is N-R⁵,

R⁵, R¹ together form monocyclic alkyl having 4, 5 or 6 ring atoms, in which one CH₂ group is replaced by an -N(L¹-R⁶)- group, and which monocyclic alkyl may be substituted by NH₂,

10

R⁶ is phenyl, which is unsubstituted, or independently mono-, di- or trisubstituted by Hal, C(Hal)₃, CH₃, C(Hal)₃O,

15 in Subformula 6

X is N,

Y is N-R⁵,

X is N,

Y is N-R⁵,

20 R⁵, R¹ together form monocyclic alkyl having 4, 5 or 6 ring atoms, in which one CH₂ group is replaced by an -N(L¹-R⁶)- group, and which monocyclic alkyl may be substituted by NH₂,

L¹ is a bond, -CONH-, -NHCO-, -CONHCH₂-, CH₂CONH-,

R⁶ is phenyl, which is unsubstituted, or independently mono-, di- or trisubstituted by Hal, C(Hal)₃, CH₃, CH₃O, C(Hal)₃O,

25

in Subformula 7

X is N,

Y is NH,

30 R¹ is L¹-R⁶-L²-R⁷,

L¹, L² are a bond,

R⁶ monocyclic alkyl having 4, 5 or 6 ring atoms, in which one CH₂ group is replaced by an -NH-, group,

R⁷ is phenyl, which is unsubstituted, or independently mono-, di- or trisubstituted by Hal, C(Hal)₃, CH₃, CH₃O, C(Hal)₃O,

35

in Subformula 8

X is N,

Y is NH,

5 R¹ is L¹-R⁶-L²-R⁷,

L¹, L² are a bond,

R⁶ is piperidyl or pyrrolidinyl,

R⁷ is phenyl, which is mono- or disubstituted by Hal, C(Hal)₃, CH₃, CH₃O, C(Hal)₃O,



10 in Subformula 9

X is N,

Y is NH,

R¹ is L¹-R⁶-L²-R⁷,

L¹, L² are a bond,

15 R⁶ is  or ,

R⁷ is phenyl, which is mono- or disubstituted by Hal, C(Hal)₃, CH₃, CH₃O, C(Hal)₃O,

in Subformula 10

X is N,

20 Y is NH,

R¹ is L¹-R⁶,

R⁶ is phenyl, which is unsubstituted, or independently mono-, di- or trisubstituted by Hal, C(Hal)₃, CH₃, CH₃O, C(Hal)₃O,

25 and the remaining residues have the meaning as indicated for Formula (I) above.

The compounds of the present invention can be in the form of a prodrug compound.

"Prodrug compound" means a derivative that is converted into a biologically active compound according to the present invention under physiological conditions in the living
30 body, e.g., by oxidation, reduction, hydrolysis or the like, each of which is carried out enzymatically, or without enzyme involvement. Examples of prodrugs are compounds, wherein the amino group in a compound of the present invention is acylated, alkylated or phosphorylated, e.g., eicosanoylamino, alanylamino, pivaloyloxymethylamino or wherein

the hydroxyl group is acylated, alkylated, phosphorylated or converted into the borate, e.g. acetyloxy, palmitoyloxy, pivaloyloxy, succinyloxy, fumaryloxy, alanyloxy or wherein the carboxyl group is esterified or amidated, or wherein a sulfhydryl group forms a disulfide bridge with a carrier molecule, e.g. a peptide, that delivers the drug selectively to a target and/or to the cytosol of a cell. These compounds can be produced from compounds of the present invention according to well-known methods. Other examples of prodrugs are compounds, wherein the carboxylate in a compound of the present invention is for example converted into an alkyl-, aryl-, choline-, amino, acyloxymethylester, linolenoyl-ester.

Metabolites of compounds of the present invention are also within the scope of the present invention.

Where tautomerism, e.g., keto-enol tautomerism, of compounds of the present invention or their prodrugs may occur, the individual forms, e.g., the keto or the enol form, are claimed separately and together as mixtures in any ratio. The same applies for stereoisomers, e.g., enantiomers, cis/trans isomers, conformers and the like.

If desired, isomers can be separated by methods well known in the art, e.g. by liquid chromatography. The same applies for enantiomers, e.g., by using chiral stationary phases. Additionally, enantiomers may be isolated by converting them into diastereomers, i.e., coupling with an enantiomerically pure auxiliary compound, subsequent separation of the resulting diastereomers and cleavage of the auxiliary residue. Alternatively, any enantiomer of a compound of the present invention may be obtained from stereoselective synthesis using optically pure starting materials.

The compounds of the present invention can be in the form of a pharmaceutically acceptable salt or a solvate. The term "pharmaceutically acceptable salts" refers to salts prepared from pharmaceutically acceptable non-toxic bases or acids, including inorganic bases or acids and organic bases or acids. In cases where the compounds of the present invention contain one or more acidic or basic groups, the invention also comprises their corresponding pharmaceutically or toxicologically acceptable salts, in particular their pharmaceutically utilizable salts. Thus, the compounds of the present invention which contain acidic groups can be present in salt form, and can be used according to the invention, for example, as alkali metal salts, alkaline earth metal salts or as ammonium salts. More precise examples of such salts include sodium salts,

potassium salts, calcium salts, magnesium salts or salts with ammonia or organic amines such as, for example, ethylamine, ethanolamine, triethanolamine or amino acids. Compounds of the present invention which contain one or more basic groups, i.e. groups which can be protonated, can be present in salt form, and can be used according to the invention in the form of their addition salts with inorganic or organic acids.

Examples of suitable acids include hydrogen chloride, hydrogen bromide, phosphoric acid, sulfuric acid, nitric acid, methanesulfonic acid, p-toluenesulfonic acid, naphthalenedisulfonic acids, oxalic acid, acetic acid, tartaric acid, lactic acid, salicylic acid, benzoic acid, formic acid, propionic acid, pivalic acid, diethylacetic acid, malonic acid, succinic acid, pimelic acid, fumaric acid, maleic acid, malic acid, sulfaminic acid, phenylpropionic acid, gluconic acid, ascorbic acid, isonicotinic acid, citric acid, adipic acid, and other acids known to the person skilled in the art. If the compounds of the present invention simultaneously contain acidic and basic groups in the molecule, the invention also includes, in addition to the salt forms mentioned, inner salts or betaines (zwitterions). The respective salts can be obtained by customary methods which are known to a person skilled in the art, for example by contacting these with an organic or inorganic acid or base in a solvent or dispersant, or by anion exchange or cation exchange with other salts. The present invention also includes all salts of the compounds of the present invention which, owing to low physiological compatibility, are not directly suitable for use in pharmaceuticals but which can be used, for example, as intermediates for chemical reactions or for the preparation of pharmaceutically acceptable salts.

The term "substituted" preferably relates to the substitution by the above-mentioned substituents, where a plurality of different degrees of substitution are possible, unless indicated otherwise.

All physiologically acceptable salts, derivatives, solvates, solvates of salts, and stereoisomers of these compounds, including mixtures thereof in all ratios, are also in accordance with the invention.

The compounds of Formula (I) may have one or more centres of chirality. They may accordingly occur in various enantiomeric forms and be in racemic or optically active form. The invention therefore also relates to the optically active forms (stereoisomers),

the enantiomers, the racemates, the diastereomers and hydrates and solvates of these compounds.

Since the pharmaceutical activity of the racemates or stereoisomers of the compounds according to the invention may differ, it may be desirable to use the enantiomers. In these cases, the end product or even the intermediates can be separated into enantiomeric compounds by chemical or physical measures known to the person skilled in the art or even employed as such in the synthesis.

In the case of racemic amines, diastereomers are formed from the mixture by reaction with an optically active resolving agent. Examples of suitable resolving agents are optically active acids, such as the R and S forms of tartaric acid, diacetyltartaric acid, dibenzoyltartaric acid, mandelic acid, malic acid, lactic acid, suitably N-protected amino acids (for example N-benzoylproline or N-benzenesulfonylproline), or the various optically active camphorsulfonic acids. Also advantageous is chromatographic enantiomer resolution with the aid of an optically active resolving agent (for example dinitrobenzoylphenylglycine, cellulose triacetate or other derivatives of carbohydrates or chirally derivatised methacrylate polymers immobilised on silica gel). Suitable eluents for this purpose are aqueous or alcoholic solvent mixtures, such as, for example, hexane/isopropanol/ acetonitrile, for example in the ratio 82:15:3. A method for the resolution of racemates containing ester groups (for example acetyl esters) is the use of enzymes, in particular esterases.

Furthermore, the present invention relates to pharmaceutical compositions comprising a compound of the present invention, or a prodrug compound thereof, or a pharmaceutically acceptable salt or solvate thereof as an active ingredient together with a pharmaceutically acceptable carrier.

"Pharmaceutical composition" means one or more active ingredients, and one or more inert ingredients that make up the carrier, as well as any product which results, directly or indirectly, from combination, complexation or aggregation of any two or more of the ingredients, or from dissociation of one or more of the ingredients, or from other types of reactions or interactions of one or more of the ingredients. Accordingly, the pharmaceutical compositions of the present invention encompass any composition

made by admixing a compound of the present invention and a pharmaceutically acceptable carrier.

A pharmaceutical composition of the present invention may additionally comprise one or more other compounds as active ingredients, such as one or more additional compounds of the present invention, or a prodrug compound or other p70S6K inhibitors.

The pharmaceutical compositions include compositions suitable for oral, rectal, topical, parenteral (including subcutaneous, intramuscular, and intravenous), ocular (ophthalmic), pulmonary (nasal or buccal inhalation), or nasal administration, although the most suitable route in any given case will depend on the nature and severity of the conditions being treated and on the nature of the active ingredient. They may be conveniently presented in unit dosage form and prepared by any of the methods well-known in the art of pharmacy.

In one embodiment, said compounds and pharmaceutical composition are for the treatment of cancer such as brain, lung, colon, epidermoid, squamous cell, bladder, gastric, pancreatic, breast, head, neck, renal, kidney, liver, ovarian, prostate, colorectal, uterine, rectal, oesophageal, testicular, gynecological, thyroid cancer, melanoma, hematologic malignancies such as acute myelogenous leukemia, multiple myeloma, chronic myelogenous leukemia, myeloid cell leukemia, glioma, Kaposi's sarcoma, or any other type of solid or liquid tumors. Preferably, the cancer to be treated is chosen from breast, colorectal, lung, prostate or pancreatic cancer or glioblastoma.

The invention also relates to the use of compounds according to the invention for the preparation of a medicament for the treatment of hyperproliferative diseases related to the hyperactivity of p70S6K as well as diseases modulated by the p70S6K cascade in mammals, or disorders mediated by aberrant proliferation, such as cancer and inflammation.

The invention also relates to a compound or pharmaceutical composition for treating a disease related to vasculogenesis or angiogenesis in a mammal which comprises a therapeutically effective amount of a compound of the present invention, or a pharmaceutically acceptable salt, prodrug or hydrate thereof, and a pharmaceutically acceptable carrier.

In one embodiment, said compound or pharmaceutical composition is for treating a disease selected from the group consisting of tumor angiogenesis, chronic inflammatory disease such as rheumatoid arthritis, inflammatory bowel disease, atherosclerosis, skin diseases such as psoriasis, eczema, and sclerodema, diabetes, diabetic retinopathy, retinopathy of prematurity and age-related macular degeneration.

This invention also relates to a compound or pharmaceutical composition for inhibiting abnormal cell growth in a mammal which comprises an amount of a compound of the present invention, or a pharmaceutically acceptable salt or solvate or prodrug thereof, in combination with an amount of another anti-cancer therapeutic, wherein the amounts of the compound, salt, solvate, or prodrug, and of the chemotherapeutic are together effective in inhibiting abnormal cell growth. Many anti-cancer therapeutics are presently known in the art. In one embodiment, the anti-cancer therapeutic is a chemotherapeutic selected from the group consisting of mitotic inhibitors, alkylating agents, anti-metabolites, intercalating antibiotics, growth factor inhibitors, cell cycle inhibitors, enzymes, topoisomerase inhibitors, biological response modifiers, anti-hormones, angiogenesis inhibitors such as cilengitide, vaccines such as BLP-25, and anti-androgens. In another embodiment the anti-cancer therapeutic is an antibody selected from the group consisting of bevacizumab, CD40-specific antibodies, chTNT-1/B, denosumab, zanolimumab, IGF1R-specific antibodies, lintuzumab, edrecolomab, WX G250, rituximab, ticilimumab, trastuzumab and cetuximab. In yet another embodiment the anti-cancer therapeutic is an inhibitor of another protein kinase, such as Axl, Aurora A, Aurora B, dyrk2, epha2, fgfr3, igf1r, IKK2, JNK3, Vegfr1, Vegfr2, Vegfr3 (also known as Flt-4), KDR, MEK, MET, Plk1, RSK1, Src, TrkA, Zap70, cKit, bRaf, EGFR, Jak2, PI3K, NPM-Alk, c-Abl, BTK, FAK, PDGFR, TAK1, LimK, Flt-3, PDK1 and Erk.

This invention further relates to a method for inhibiting abnormal cell growth in a mammal or treating a hyperproliferative disorder that comprises administering to the mammal an amount of a compound of the present invention, or a pharmaceutically acceptable salt or solvate or prodrug thereof, in combination with surgery or radiation therapy, wherein the amounts of the compound, salt, solvate, or prodrug, is in combination with the radiation therapy effective in inhibiting abnormal cell growth or treating the hyperproliferative disorder in the mammal. Techniques for administering radiation therapy are known in the art, and these techniques can be used in the

combination therapy described herein. The administration of a compound of the invention in this combination therapy can be determined as described herein. It is believed that the compounds of the present invention can render abnormal cells more sensitive to treatment with radiation for purposes of killing and/or inhibiting the growth of such cells.

Accordingly, this invention further relates to a method for sensitizing abnormal cells in a mammal to treatment with radiation which comprises administering to the mammal an amount of a compound of the present invention or pharmaceutically acceptable salt or solvate or prodrug thereof, which amount is effective in sensitizing abnormal cells to treatment with radiation. The amount of the compound, salt, or solvate in this method can be determined according to the means for ascertaining effective amounts of such compounds described herein. The invention also relates to a method for inhibiting abnormal cell growth in a mammal that comprises an amount of a compound of the present invention, or a pharmaceutically acceptable salt or solvate thereof, a prodrug thereof, or an isotopically-labeled derivative thereof, and an amount of one or more substances selected from anti-angiogenesis agents, signal transduction inhibitors, and antiproliferative agents.

In practical use, the compounds of the present invention can be combined as the active ingredient in intimate admixture with a pharmaceutical carrier according to conventional pharmaceutical compounding techniques. The carrier may take a wide variety of forms depending on the form of preparation desired for administration, *e.g.*, oral or parenteral (including intravenous). In preparing the compositions for oral dosage form, any of the usual pharmaceutical media may be employed, such as, for example, water, glycols, oils, alcohols, flavoring agents, preservatives, coloring agents and the like. In the case of oral liquid preparations, any of the usual pharmaceutical media may be employed, such as, for example, suspensions, elixirs and solutions; or carriers such as starches, sugars, microcrystalline cellulose, diluents, granulating agents, lubricants, binders, disintegrating agents and the like. In the case of oral solid preparations the composition may take forms such as, for example, powders, hard and soft capsules and tablets, with the solid oral preparations being preferred over the liquid preparations.

Because of their ease of administration, tablets and capsules represent the most advantageous oral dosage unit form in which case solid pharmaceutical carriers are

obviously employed. If desired, tablets may be coated by standard aqueous or nonaqueous techniques. Such compositions and preparations should contain at least 0.1 percent of active compound. The percentage of active compound in these compositions may, of course, be varied and may conveniently be between about 2 percent to about 60 percent of the weight of the unit. The amount of active compound in such therapeutically useful compositions is such that an effective dosage will be obtained. The active compounds can also be administered intranasally as, for example, liquid drops or spray.

10 The tablets, pills, capsules, and the like may also contain a binder such as gum tragacanth, acacia, corn starch or gelatin; excipients such as dicalcium phosphate; a disintegrating agent such as corn starch, potato starch, alginic acid; a lubricant such as magnesium stearate; and a sweetening agent such as sucrose, lactose or saccharin. When a dosage unit form is a capsule, it may contain, in addition to materials of the
15 above type, a liquid carrier such as a fatty oil.

Various other materials may be present as coatings or to modify the physical form of the dosage unit. For instance, tablets may be coated with shellac, sugar or both. A syrup or elixir may contain, in addition to the active ingredient, sucrose as a sweetening agent,
20 methyl and propylparabens as preservatives, a dye and a flavoring such as cherry or orange flavor.

Compounds of the present invention may also be administered parenterally. Solutions or suspensions of these active compounds can be prepared in water suitably mixed with a
25 surfactant such as hydroxy-propylcellulose. Dispersions can also be prepared in glycerol, liquid polyethylene glycols and mixtures thereof in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.

30 The pharmaceutical forms suitable for injectable use include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. In all cases, the form must be sterile and must be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of
35 microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion

medium containing, for example, water, ethanol, polyol (e.g., glycerol, propylene glycol and liquid polyethylene glycol), suitable mixtures thereof, and vegetable oils.

Any suitable route of administration may be employed for providing a mammal, especially a human, with an effective dose of a compound of the present invention. For example, oral, rectal, topical, parenteral, ocular, pulmonary, nasal, and the like may be employed. Dosage forms include tablets, troches, dispersions, suspensions, solutions, capsules, creams, ointments, aerosols, and the like. Preferably compounds of the present invention are administered orally.

The effective dosage of active ingredient employed may vary depending on the particular compound employed, the mode of administration, the condition being treated and the severity of the condition being treated. Such dosage may be ascertained readily by a person skilled in the art.

When treating or preventing cancer, inflammation or other proliferative diseases for which compounds of the present invention are indicated, generally satisfactory results are obtained when the compounds of the present invention are administered at a daily dosage of from about 0.01 milligram to about 100 milligram per kilogram of animal body weight, preferably given as a single daily dose. For most large mammals, the total daily dosage is from about 0.1 milligrams to about 1000 milligrams, preferably from about 0.2 milligram to about 50 milligrams. In the case of a 70 kg adult human, the total daily dose will generally be from about 0.2 milligrams to about 200 milligrams. This dosage regimen may be adjusted to provide the optimal therapeutic response.

The invention also relates to a set (kit) consisting of separate packs of

a) an effective amount of a compound according to the invention or a physiologically acceptable salt, solvate or prodrug thereof, and

b) an effective amount of a further medicament active ingredient.

The set comprises suitable containers, such as boxes, individual bottles, bags or ampoules. The set may, for example, comprise separate ampoules, each containing an effective amount of a compound according to the invention and/or pharmaceutically usable derivatives, solvates and stereoisomers thereof, including mixtures thereof in all ratios, and an effective amount of a further medicament active ingredient in dissolved or lyophilised form.

Experimental Section

Some abbreviations that may appear in this application are as follows:

5

Abbreviations

<i>Designation</i>	
ACN	acetonitrile
AcOH	Acetic acid
AIBN	Azobisisobutyronitrile
ATP	Adenosine triphosphate
b	Broad peak
Bop-Cl	Bis(2-oxo-3-oxazolidinyl)phosphinic chloride
Conc.	concentrated
d	Doublet
DCM	Dichloromethane
DCE	dichloroethane
DMAP	dimethylaminopyridine
DMF	dimethylformamide
DMSO	dimethylsulfoxide
DIEA/DIPEA	N,N-Diisopropylethylamine
DTT	dithiothreitol
EDTA	Ethylenediaminetetraacetic acid
equiv./eq.	equivalents
Et	ethyl
h	hour
HEPES	4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid
HPLC	High pressure liquid chromatography
LC/MS	Liquid chromatography coupled to mass spectrometry
LiOH	Lithium hydroxide
m	multiplet
M	Molecular ion
m/z	Mass-to-charge ratio

Me	methyl
MeOH	methanol
min	minute
MS	Mass spectrometry
N	Normal (unit of concentration)
NaOH	Sodium hydroxide
NBS	N-bromosuccinimide
NMO	4-methylmorpholine <i>N</i> -oxide
NMP	N-Methyl-2-pyrrolidone
NMR	Nuclear Magnetic Resonance
PG	Protecting group
psi	Pounds per square inch
q	Quartette (or quartet)
R _f	Retention factor
RT/rt	Room temperature
R _t .	Retention time
s	Singlet
T3P	Propylphosphonic anhydride
TBAF	Tetrabutylammonium fluoride
Tert	Tertiary
TEA	Triethylamine
TFA	Trifluoroacetic acid
THAB	Tetrahexylammonium bromide
THF	Tetrahydrofuran
UV	ultraviolet
VIS	visible

The compounds of the present invention can be prepared according to the procedures of the following Schemes and Examples, using appropriate materials and are further exemplified by the following specific examples.

5

Moreover, by utilizing the procedures described herein, in conjunction with ordinary skills in the art, additional compounds of the present invention claimed herein can be readily prepared. The compounds illustrated in the examples are not, however, to be construed

as forming the only genus that is considered as the invention. The examples further illustrate details for the preparation of the compounds of the present invention. Those skilled in the art will readily understand that known variations of the conditions and processes of the following preparative procedures can be used to prepare these compounds.

The instant compounds are generally isolated in the form of their pharmaceutically acceptable salts, such as those described above. The amine-free bases corresponding to the isolated salts can be generated by neutralization with a suitable base, such as aqueous sodium hydrogencarbonate, sodium carbonate, sodium hydroxide and potassium hydroxide, and extraction of the liberated amine-free base into an organic solvent, followed by evaporation. The amine-free base, isolated in this manner, can be further converted into another pharmaceutically acceptable salt by dissolution in an organic solvent, followed by addition of the appropriate acid and subsequent evaporation, precipitation or crystallization.

The invention will be illustrated, but not limited, by reference to the specific embodiments described in the following schemes and examples. Unless otherwise indicated in the schemes, the variables have the same meaning as described above.


Unless otherwise specified, all starting materials are obtained from commercially suppliers and used without further purifications. Unless otherwise specified, all temperatures are expressed in °C and all reactions are conducted at room temperature. Compounds were purified by either silica chromatography or preparative HPLC.

The present invention also relates to processes for manufacturing the compounds of Formula (I) according to the hereinafter described schemes and working examples.

Synthetic Schemes Describing Intermediate and End Product Compounds

Quinazoline bromide intermediate **1a** is obtained from commercially available 2-Amino-3-methylbenzoic acid by an 8-step synthesis outlined in Scheme 1.


Scheme 1



Quinoline intermediates **9a**, **10a** are commercially available and **11a** is prepared according to the 5-step synthesis outlined in Scheme 2.


5

Scheme 2




10 Amine intermediates are either commercially available or prepared by the following synthetic routes (Scheme 3, Scheme 4, Scheme 5 and Scheme 6).

Scheme 3




Protection of the amino group of compound **3a** with Toc yields **3b**. **3b** was hydrolyzed to release carboxylic acid **3c**. **3c** is coupled with amine, followed by removing Toc protecting group to generate the amine intermediate **e**.

Scheme 4




Arylaldehyde **4a** was reacted with nitromethane under basic conditions to provide the hydroxyl derivative **4b**, which was converted to alkene **4c** under the elimination condition promoted by acetic anhydride. Cyclization of **4c** with N-benzyl-1-methoxy-N-((trimethylsilyl)methyl)methanamine gave pyrrolidine derivative **4d**. Reduction of the nitro group in **4d** using Raney nickel as a catalyst, followed by the Teoc protection of the resulting amino moiety with N-[2-(trimethylsilyl)ethoxycarbonyloxy]succinimide gave **4f**. The N-benzyl group of **4f** was removed under hydrogenation conditions, and protected with *di-tert*-butyl dicarbonate to provide **4h**. The Teoc protected primary amine in **4h** was released upon treatment with *tetra-n*-butylammonium fluoride to provide amine intermediate **4i**.

Scheme 5




Amino acid **5a** was protected with Boc to give **5b**. **5b** was then reacted with amine to give amide **5c**. Cbz group was removed under hydrogenation condition to generate the desired amine intermediate **5d**.

Scheme 6




Amino group of amino acid **6a** was selectively protected with Teoc, following the acid moiety coupling with amine gave amide **6c**. Teoc was removed by TBAF to afford the amine intermediate **6d**.

Scheme 7




Under reductive amination conditions, **7a** and **7b** are reacted to give **7c**, which was then
 5 treated with acid to give the deprotected pyrrolidine intermediate, **7d**, as the trifluoroacetate salt.

Scheme 8




Bromide intermediate **1a** is coupled with amine **8b**, where P is a protecting group when it
 10 is needed, to yield **8c**. Hydrolysis of nitrile intermediate **8c** affords amide **8d** under basic condition assistant by hydrogen peroxide, and followed by removing protecting group, gave the desired compound **8e**.

Scheme 9




9a was oxidized hydrolysis with hot concentrated sulfuric acid to convert the nitrile to carboxylic amide **9b**. **9b** was coupled with carboxylic acid provided **9c**. De-protection of **9c** to afford **9d** as the desired compound.

Scheme 10




Benzyl bromides **10a** were reacted with aniline **9b** to yield the desired compounds **10b**.

Scheme 11



5-bromoquinoline-8-carbonitrile **11a** was reacted with aminopyrrolidine derivative under Buchwald Harting cross coupling condition to give **11b**. The nitrile moiety was converted to carboxylic amide using hot concentrated sulfuric acid to afford the compound **11c**.

Scheme 12



Under Buchwald-Hartwig cross coupling condition, methyl 5-bromoquinoline-8-carboxylate **12a** is reacted with the amine intermediate **8b** gave **12b**. **12b** was treated with ammonia in methanol to yield the carboxylic amide **12c**, which was then removed the protecting group to generate **12d**.

Analytical Methodology

Analytical LC/MS was performed using the following three methods:

Method A: A Discovery C¹⁸, 5 μm , 3 x 30 mm column was used at a flow rate of 400 $\mu\text{L}/\text{min}$, sample loop 5 μL , mobile phase: (A) water with 0.1% formic acid, mobile phase, (B) methanol with 0.1% formic acid; retention times are given in minutes. Method details: (I) runs on a Quaternary Pump G1311A (Agilent) with UV/VIS diode array detector G1315B (Agilent) and Finnigan LCQ Duo MS detector in ESI + mode with UV-detection at 254 and 280 nm with a gradient of 15-95% (B) in a 3.2 min linear gradient (II) hold for 1.4 min at 95% (B) (III) decrease from 95-15% (B) in a 0.1 min linear gradient (IV) hold for 2.3 min at 15% (B).

Method B: A Waters Symmetry C¹⁸, 3.5 μm , 4.6 x 75 mm column at a flow rate of 1 mL/min, sample loop 10 μL , mobile phase (A) is water with 0.05% TFA, mobile phase (B) is ACN with 0.05% TFA; retention times are given in minutes. Method details: (I) runs on a Binary Pump G1312A (Agilent) with UV/Vis diode array detector G1315B (Agilent) and Agilent G1956B (SL) MS detector in ESI + mode with UV-detection at 254 and 280 nm with a gradient of 20-85% (B) in a 10 min linear gradient (II) hold for 1 min at 85% (B) (III) decrease from 20-85% (B) in a 0.2 min linear gradient (IV) hold for 3.8 min at 20% (B).

Method C: Gradient: 4.2 min/ Flow: 2 ml/min 99:01 - 0:100 Water + 0.1%(Vol.) TFA; Acetonitril + 0.1%(Vol.) TFA; 0.0 to 0.2 min: 99:01; 0.2 to 3.8 min: 99:01→ 0:100; 3.8 to 4.2 min: 0:100; Column: Chromolith Performance RP18e; 100 mm long, 3 mm diameter; Wavelength: 220nm.

5

Analytical Chiral HPLC

Analytical chiral HPLC was performed using a ChiralPak AD-H column (250 X 4.6 mm) from Daicel Chemical Industries, Ltd. on an Agilent 1100 Series system. The method used a 5.0 µL injection volume, with a flow rate of 1 mL/min of 100% methanol for 15 min at 25 °C, and UV-detection at 254 and 280 nm.

10

Preparative HPLC

Preparative HPLC was performed using either a Waters Atlantis dC₁₈ OBD™ 10 µM (30 X 250 mm) column or a Waters Sunfire Prep C₁₈ OBD 10 µM (30 X 250 mm) column.

15

The columns were used at a flow rate of 60 mL/min on a Waters Prep LC 4000 System equipped with a sample loop (10 mL) and an ISCO UA-6 UV/Vis detector. The mobile phase was drawn from two solvent reservoirs containing (A) water and (B) HPLC-grade acetonitrile. A typical preparative run used a linear gradient (e.g., 0-60 % solvent B over 60 min).


20

Examples

The working examples presented below are intended to illustrate particular embodiments of the invention and are not intended to limit the scope of the specification or the claims in any way. These working examples describe both the synthesis of reagents used to prepare the heterocyclic carboxamide compounds useful in the treatment of hyperproliferative diseases and these same heterocyclic quinazoline and quinoline carboxamide compounds.

25

Preparation of the intermediates



30

Intermediate A: 5-bromoquinazoline-8-carbonitrile

Step1: 8-Methylquinazolin-4(3H)-one

2-Amino-3-methylbenzoic acid (100 g, 0.66 mol), formamidine acetate (206 g, 1.98 mol) and formamide (26 mL, 0.6600 mol) were mixed in a 2L R.B fitted with Mechanical stirrer. The reaction mixture was heated at 160 °C for 16h. The reaction completion was monitored by LCMS. After completion, the reaction mixture was cooled to was RT and diluted with 2N NaOH solution (300 mL). After stirring at the same temperature for 15min, the reaction mixture neutralised with 1.5N HCl solution. The solid precipitated was filtered off, washed with ice cold water and dried under vacuum to yield (90 g, 86% yield) of the titled compound as an off white solid. ¹H NMR (DMSO-d₆, 400MHz) δ 12.21 (bs, 1H), 8.10 (s, 1H), 7.95-7.93 (dd, J = 8.8, 7.9 Hz, 1H), 7.65-7.63 (d, J = 7.9 Hz, 1H), 7.39-7.35 (t, J = 15.2 Hz, 1H), 2.51 (s, 3H).

Step 2: 4-Chloro-8-methylquinazoline

POCl₃ (300 mL) was taken in a 2 L round bottom flask under nitrogen. To this was added 8-Methylquinazolin-4(3H)-one (45 g) in portions. The reaction mixture refluxed at 120 °C for 12h. Reaction completion was monitored by TLC and LCMS. After completion, the reaction mixture was cooled to RT and evaporated to dryness under reduced pressure. The resulted residue was dissolved in DCM (500 mL) and quenched slowly into an ice cold solution of saturated K₂CO₃ with constant stirring. Then the organic layer was separated and washed with brine solution, dried over sodium sulphate and concentrated under vacuum to afford (45g, 90% yield) of the titled compound as yellow solid. This was taken for next step without further purification. ¹H NMR (CDCl₃, 400MHz) δ 9.03 (s, 1H), 8.08-8.06 (dd, J = 8.9, 8.4 Hz, 1H), 7.77-7.76 (d, J = 7.1 Hz, 1H), 7.59-7.56 (d, J = 15.5 Hz, 1H), 2.75 (s, 3H).

Step 3: 8-Methylquinazoline

To a stirred solution of 4-Chloro-8-methylquinazoline (45 g, 0.252 mol) in DCM (700 mL) under N₂ was added p-toluenesulfonylhydrazide (65.9 g, 0.353 mol) in portions. The reaction mixture was heated at 45°C for 12h. The reaction completion was monitored by LCMS and TLC. After completion, the reaction mixture was cooled to RT, the solvent evaporated to dryness and the resulted residue redissolved in EtOH (500 mL) and added 2N NaOH solution (300 mL) and refluxed for 6h. After confirming by LCMS, the reaction mixture was cooled to RT and extracted with MTBE (3 x 600 mL). The combined organic layers were washed with brine solution, dried over sodium sulfate and concentrated under vacuum. The resulted residue was filtered through column chromatography using neutralized silica gel (60-120mesh) and pet ether/ ethyl acetate as an eluent to yield (15 g, 27% yield) of the titled compound as a low melting yellow

solid. ¹H NMR (DMSO-d₆, 400MHz) δ 9.54 (s, 1H), 9.31 (s, 1H), 7.97-7.94 (dd, J = 8.8, 8.1 Hz, 1H), 7.87-7.84 (m, 1H), 7.65-7.62 (d, J = 15.2 Hz, 1H), 2.67 (s, 3H).

Step 4: 5-Bromo-8-methylquinazoline

To a solution of 8-Methylquinazoline (10 g, 0.0694 mol) in conc.H₂SO₄ (100 mL), was added silver sulphate (34.64 g, 0.1111mol) in portions at 0°C. To this was added Bromine (4.4 mL, 0.0832mol) in drops. The reaction mixture was stirred at RT for 16h. Reaction was monitored by LCMS at regular intervals. At the end of 16h, LCMS showed 42% starting material and 51% of product. The reaction mixture was quenched with ice and basified with NH₄OH solution. The aqueous layer was extracted with EtOAc (4 x 500 mL), washed with water and brine solution. The organic layer was dried over Na₂SO₄ and concentrated under vacuum. The crude was purified by column chromatography using neutralized silica gel (60-120 mesh) and pet ether/ ethyl acetate as an eluent to yield the titled compound as pale yellow liquid (51% yield). ¹H NMR (DMSO-d₆, 400MHz) δ 9.60-9.58 (s, 1H), 9.40-9.38 (s, 1H), 7.93-7.91 (d, J = 7.72 Hz, 1H), 7.77-7.75 (d, J = 7.72 Hz, 1H), 2.62 (s, 3H).

Step 5: 5-Bromo-8-(dibromomethyl) quinazoline

To a stirred solution of 5-Bromo-8-methylquinazoline (46 g, 0.206 mol) in CCl₄ (800 mL) under N₂, was added N-Bromosuccinimide (80.4 g, 0.451mol) followed by AIBN (6.74 g, 0.041 mol) at RT. Then the reaction mixture was heated at 90°C for 12h. After completion, the reaction mixture cooled to RT, filtered off and washed with CCl₄. The filtrate was concentrated to dryness to yield (61g) of the titled compound as yellow liquid. The crude product was taken as such for next step without purification. ¹H NMR (DMSO-d₆, 400MHz) δ 9.73 (s, 1H), 9.53 (s, 1H), 8.45-8.43 (d, J = 8.0 4Hz, 1H), 8.22-8.20 (d, J = 8.04 Hz, 1H), 8.02 (s, 1H).

Step 6: 5-Bromoquinazoline-8-carbaldehyde

To a stirred solution of 5-Bromo-8-(dibromomethyl) quinazoline (61g, crude mixture) in Acetone (500 mL) and water (100 mL), was added silver nitrate (61 g) in portions at 0°C. The reaction mixture was stirred at RT for 6h. The reaction completion was confirmed by TLC. The reaction mixture was filtered off and filtrate was extracted with ethyl acetate (3 x 500 mL). The organic layer was washed with 10% NaHCO₃ solution, water and brine solution. The solvent was dried over Na₂SO₄ and concentrated under vacuum to afford the crude product with HPLC (56%) (25 g, 65%) was subjected to prep.HPLC condition: COLUMN: Xterra, C18 (19 X 300 mm), 10 micron, Mobile: 0.1% TFA; B: MeOH to isolate the compound. The resulted solid was basified with NH₄OH and extracted with EtOAc. The organic layer was washed water. The solvent was dried over Na₂SO₄ and


concentrated under vacuum to yield (6.2 g) of the titled compound as pale yellow solid. ¹H NMR (DMSO-d₆, 400MHz) 11.14 (s, 1H), 9.80 (s, 1H), 9.58 (s, 1H), 8.30-8.27 (d, J = 12.3 Hz, 2H).

Step 7: 5-Bromo-quinazoline-8-carbaldehyde oxime

- 5 To a suspension solution of 5-Bromo-quinazoline-8-carbaldehyde (2400.00 mg; 10.12 mmol; 1.00 eq.) in acetonitrile (10ml), hydroxylamine hydrogen chloride (773.90 mg; 11.14 mmol; 1.10 eq.) was added, followed by triethylamine (1.57 ml; 11.14 mmol; 1.10 eq.). The reaction mixture was stirred at 100°C for 1hr, LCMS showed MS 270/272 (M+18). The reaction was cooled, filtrate and washed with ether to collect product 5-Bromo-quinazoline-8-carbaldehyde oxime 3000mg, yield 117% (1HNMR showed contained 1eq triethylamine HCl salt).
- 10

Step 8: 5-bromoquinazoline-8-carbonitrile

- To a solution of 5-Bromo-quinazoline-8-carbaldehyde oxime (3000.00 mg; 11.90 mmol; 1.00 eq.) in DMF (15ml) was added 2,4,6-Tripropyl-[1,3,5,2,4,6] trioxatriphosphinane 2,4,6-trioxide (10.42 ml; 17.85 mmol; 1.50 eq.) 50% in DMF. The resulting mixture was stirred at 100°C for 45min, LCMS showed major peak (MS: 252/254, 234/236) and small (MS: 208/209). The reaction mixture was cooled, poured into water, extracted with EtOAc, dried and concentrated to yield sticky solid, which was washed with ether to collected the title product (1665mg. yield 59.6%). LC-MS (232/234 and 250/252).
- 15
- 20



Intermediate B: Methyl 5-bromoquinoline-8-carboxylate

Step1: 5-Bromo-8-methylquinoline

- To a solution of 8-Methyl quinoline (30g, 0.209mol) in Conc.H₂SO₄ (300ml) was added Silver sulphate(97.98g, 0.314mol) in lots at 0°C and followed by Bromine(10.74ml, 0.209mol) drop wise for 10 min. After addition of Bromine, the reaction mixture was stirred at RT for 4h and the reaction completion was confirmed by TLC. After completion of reaction the reaction mixture was quenched with ice and basified with NH₄OH solution and extracted with ethyl acetate. The organic layer was washed with water, brine solution and dried over anhydrous sodium sulphate and concentrated to afford (43g, 92.4% yield) as dark brown liquid. The crude product was as such taken for next step
- 25
- 30

without further purification. **¹H NMR** (DMSO-d₆, 400MHz): δ 8.97-8.96 (dd, J=4.16, 5.8Hz, 1H), 8.45-8.42 (dd, J=8.56, 10.2Hz, 1H), 7.81-7.79 (d, J=7.68Hz, 1H), 7.67-7.64 (dd, J=8.52, 12.64Hz, 1H), 7.53-7.51 (dd, J=7.64, 8.52Hz, 1H), 2.66(s, 3H).

Step2: 5-Bromo-8-(dibromomethyl)quinoline

- 5 To a solution of 5-Bromo-8-methylquinoline (86g, 0.387mol) in CCl₄ (700ml) was added N-Bromosuccinamide(144g, 0.813mol), Followed by Benzoylperoxide(8.6g) at RT and the reaction mixture was heated for 90°C for 12h. The reaction completion was confirmed by TLC. After completion of reaction, the reaction mixture was filtered and concentrated to afford (140g, 95% yield) as pale orange solid. The crude product was as such taken
- 10 for next step without further purification.. **¹H NMR** (DMSO-d₆, 400MHz): δ 9.1-9.08 (dd, J=4.16, 5.8Hz, 1H), 8.59-8.57 (dd, J=8.56, 10.2Hz, 1H), 8.25-8.23 (d, J=8.04Hz, 1H), 8.16-8.11 (t, J=19.08Hz, 2H), 7.82-7.79 (dd, J=8.6, 12.8Hz, 1H).

Step 3: 5-Bromoquinoline-8-carbaldehyde

- To a solution of 5-Bromo-8-(dibromomethyl)quinoline (75g, 0.197mol) in Acetone
- 15 (400ml), Water(100ml) was added Silvernitrate(75g) at 0°C in lots for 10min and this reaction mixture was stirred at RT for 6h. The reaction completion was confirmed by TLC. After completion of reaction the reaction mixture was filtered and the filtrate was extracted with MTBE (1000ml). The organic layer was washed with 10% NaHCO₃ solution water, brine and the organic layer was dried over sodium sulphate and
- 20 concentrated to afford the title compound as off white solid (15g, 32% yield). **¹H NMR** (DMSO-d₆, 400MHz): δ 11.272-11.27 (s, 1H), 9.17-9.15 (dd, J = 4.2, 5.8 Hz, 1H), 8.64-8.62 (dd, J = 8.6, 10.3 Hz, 1H), 8.17-8.15 (d, J=7.84, 1H), 8.09-8.07 (d, J=7.88Hz, 1H), 7.85-7.82 (dd, J= 8.64, 12.84Hz, 1H).


Step 4: 5-Bromoquinoline-8-carboxylic acid

- 25 To a solution of 5-Bromoquinoline-8-carbaldehyde (10 g, 0.039mol) in THF (300ml) was added aqueous NaOH (30 g) and followed by silver nitrate (10.79 g, 0.0635 mol) lot wise at RT for 10 min and this reaction mixture was stirred at RT for 30min. The reaction completion was confirmed by TLC. After completion of reaction the reaction mixture was filtered. The black solid was washed with THF, MeOH and DMF (50ml each). (Note:
- 30 Product was not soluble in any of these solvents!! The filtrate and washings were discarded). The black solid was dried to afford the title compound (crude -30 g). The crude product was as such taken for the next step. **¹H NMR** (DMSO-d₆, 400MHz): δ 8.92-8.91 (dd, J=4.12, 5.8Hz, 1H), 8.49 (s, 1H), 8.46-8.44 (dd, J=8.56, 10.2Hz, 1H), 7.83-7.81 (dd, J=7.6, Hz, 1H), 7.63-7.6 (dd, J=8.52, 12.68Hz, 1H), 7.46-7.44 (d, J=7.6Hz, 1H).

- 35 *Step 5: Methyl 5-bromoquinoline-8-carboxylate*

To a solution of 5-Bromoquinoline-8-carboxylic acid (30g, 0.119mol, crude) in DMF (400ml) was added potassium carbonate (41.12g, 0.297mol) and MeI (22.3ml, 0.357mol) at RT. The reaction mixture was stirred at RT for 12h and the reaction completion was confirmed by TLC. After completion of reaction the reaction mixture was filtered and evaporated. The reaction mixture was basified with 10%NaHCO₃ and extracted with ethyl acetate. The organic layer was washed with water, brine and dried over sodium sulphate and concentrated to afford the title compound as brown liquid (4.6g). LCMS: Mass found (M+, 268). ¹H NMR (DMSO-d₆, 400MHz): δ 9.04-9.02 (dd, J=3.92, 5.68, Hz, 1H), 8.58-8.56 (m, 1H), 8.05-8.03 (d, J=7.72Hz, 1H), 7.87-7.86 (d, J=7.76Hz, 1H), 7.78-7.75 (dd, J=8.6, 12.8Hz, 1H), 3.91 (s, 3H).

Representative synthesis of Intermediate C (Scheme 4)



tert-butyl 3-amino-4-(4-(trifluoromethoxy)phenyl)pyrrolidine-1-carboxylate (trans, racemic)

Step 1: 1-(3-trifluoromethoxy-phenyl)-2-nitro-ethanol

A solution of 3-fluoromethoxybenzaldehyde (21.37 ml; 201.43 mmol; 1.00 eq.) and nitromethane (13.06 ml; 241.71 mmol; 1.20 eq.) in MeOH (40ml) was cooled to -10 °C. A solution of NaOH (8.46 g; 211.50 mmol; 1.05 eq.) in H₂O (20 ml) was added over 10 min, keeping the temperature below -5 °C. The reaction mixture was stirred at -5 °C for 15 min, during which the reaction solution solidified as a white solid. The reaction mixture was warmed to 0 °C, and diluted with H₂O (150 ml). Upon dissolution of all of the solids, HCl (4M, 100 ml) was added. The reaction mixture was extracted with DCM (300ml x 2). The combined extracts were washed with brine and concentrated to provide the desired 1-(3-trifluoromethoxy-phenyl)-2-nitro-ethanol 34.8g, yield 93%.

Step 2: 1-trifluoromethoxy-3-((E)-2-nitro-vinyl)-benzene

N,N-Dimethylpyridin-4-amine (2.30 g; 18.80 mmol) was added to a solution of 1-(3-fluoromethoxyphenyl)-2-nitroethanol (34.80 g; 187.95 mmol) in acetic anhydride (35.53 ml; 375.90 mmol) at 0°C, and stirred at RT for 72 h. The reaction mixture was quenched by pouring into a vigorously stirred satd. NaHCO₃ solution (400 mL). The desired intermediate was extracted with ethyl acetate (3 x 100 mL). The organic extracts were

washed with satd. NaHCO₃, brine, dried over MgSO₄, filtered, and concentrated to provide the desired 1-trifluoromethoxy-3-((E)-2-nitro-vinyl)-benzene 26.0 g, yield 83%.

Step 3: Trans-1-Benzyl-3-(3-trifluoromethoxy-phenyl)-4-nitro-pyrrolidine

N-benzyl-1-methoxy-N-[(trimethylsilyl)methyl]methanamine was added to a solution of 1-trifluoromethoxy-3-((E)-2-nitro-vinyl)-benzene (6.00 g; 35.90 mmol) in DCM (50ml). The reaction solution was cooled to 0°C, TFA (0.30 ml; 3.95 mmol) was added dropwise, and stirred overnight at RT. The reaction solution was washed with H₂O and brine, dried over MgSO₄, filtered, and concentrated. The crude material was purified via Biotage (340 g column) eluting with 5% EtOAc in hexane to provide the desired product 5.5g, yield

51%.

Step 4: (Trans-1-Benzyl-4-(3-trifluoromethoxy-phenyl)-pyrrolidin-3-ylamine

Trans-1-benzyl-3-(3-trifluoromethoxyphenyl)-4-nitropyrrolidine (5.50 g; 18.31 mmol) was dissolved in MeOH (300 mL). NH₃ (30ml, 2.0M in MeOH) was added, and the solution was passed through the H cube (flow 1.5min/min, full H₂, at 50°C). The reaction solution was concentrated to provide (Trans-1-Benzyl-4-(3-trifluoromethoxy-phenyl)-pyrrolidin-3-ylamine 4.6 g, yield 92%.

Step 5: 2-(trimethylsilyl)ethyl [trans-1-Benzyl-4-(3-trifluoromethoxy-phenyl)-pyrrolidin-3-yl]-carbamate

1-({[2-(Trimethylsilyl)ethoxy]carbonyl}oxy)pyrrolidine-2,5-dione (4.5 g; 17.37 mmol) was added to a solution of trans-1-benzyl-4-(3-trifluoromethoxyphenyl) pyrrolidinyl-3-amine (4.5 g; 16.87 mmol) and DIEA (4.5 ml; 25.30 mmol) in DCM (50 ml) at 0°C. The reaction mixture was then warmed to RT and stirred for 1hr at RT. The reaction solution was washed with brine, dried over MgSO₄, filtered, and concentrated. The crude material was purified by Biotage eluting with a gradient of 30 to 60% EtOAc in hexane to provide the title compound 6.0 g, yield 99%.

Step 6: 2-(trimethylsilyl)ethyl [trans-4-(3-trifluoromethoxy-phenyl)-pyrrolidin-3-yl]-carbamate

AcOH (2 mL) was added to a solution of 2-(trimethylsilyl)ethyl [trans-benzyl-4-(3-trifluoromethoxyphenyl)pyrrolidin-3-yl]carbamate (2.50 g; 6.03 mmol) in EtOH (150 ml). Pd/C (1.25 g, wet, 10% Pd) was then added, and the reaction mixture was put on a par shaker (60Psi), and reacted for 2 h. The reaction mixture was filtered, and the filtrate was concentrated to provide the title compound (1.96 g, quantitative yield).


Step 7: Tert-butyl trans-3-(3-trifluoromethoxyphenyl)-4-({[2-(trimethylsilyl)ethoxy]carbonyl}amino)-pyrrolidine-1-carboxylate

Di-tert-butyl dicarbonate (1.27 g; 5.82 mmol) was added to a solution of 2-(trimethylsilyl)ethyl [trans-4-(3-fluoromethoxyphenyl)pyrrolidin-3-yl]carbamate (1.80 g; 5.55 mmol) and DIEA (2.2 ml; 12.26 mmol) in DCM (100 ml) and stirred overnight at RT. The reaction mixture was concentrated, and the crude product was purified via Biotage eluting with a gradient of 20 to 60% EtOAc in hexanes to provide the title compound 2.0 g, yield 85%.

Step 8: tert-butyl 3-amino-4-(4-(trifluoromethoxy)phenyl)pyrrolidine-1-carboxylate (trans, racemic)

Tert-butyl trans-3-(3-(trifluoromethoxyphenyl)-4-([2-(trimethylsilyl)ethoxy]carbonyl)amino)-pyrrolidine-1-carboxylate (2.4 g; 5.76 mmol) and N,N,N-tributylbutan-1-aminium fluoride (20.00 ml; 1.00 M; 20.00 mmol) was dissolved in MeOH, and stirred overnight at RT. The crude product was purified via Biotage eluting with a gradient of 5 to 10% MeOH in DCM to provide title compound (1.61 g, yield 79%). LC-MS (M + H = 346, obsd = 347). ¹HNMR (DMSO-*d*₆) δ 1.39(s, 9H), 1.55 (s, 1H), 2.90-2.99 (m, 2H), 3.24-3.26(m, 1H), 3.33-3.37 (m, 1H), 3.55-3.57 (m, 1H), 3.60-3.68 (m, 1H), 3.70-3.72 (m, 1H), 7.05-7.06 (m, 1H), 7.13-7.15 (m, 2H), 7.35-7.36 (m, 1H).

Representative synthesis of Intermediate D (Scheme 6)



Benzyl ((3-(phenylcarbamoyl)piperidin-3-yl)methyl)carbamate (racemic)

Step 1: 3-(((benzyloxy)carbonyl)amino)methyl)-1-((2-(trimethylsilyl)ethoxy)-carbonyl)piperidine-3-carboxylic acid

To 3-(Benzyloxycarbonylamino-methyl)-piperidine-3-carboxylic acid hydrochloride (2000.00 mg; 6.08 mmol; 1.00 eq.) in DME (60ml) was added DIEA (3.82 ml; 21.29 mmol; 3.50 eq.). After stirring for 15mins, 3-trimethylsilanyl-propionic acid 2,5-dioxo-pyrrolidin-1-yl ester was added (1628.18 mg; 6.69 mmol; 1.10 eq.). The reaction mixture was stirred overnight at RT. Added another 50ml of DME to the reaction solution, then washed with 1% citric acid, brine, dried and concentrated to yield the residue showed clean ms as the title compound (2462mg, yield 85%), which was directly used for the next step reaction.


Step 2: 2-(trimethylsilyl)ethyl 3-(((benzyloxy)carbonyl)amino)methyl)-3-(phenylcarbamoyl)piperidine-1-carboxylate

To a solution of 3-((((benzyloxy)carbonyl)amino)methyl)-1-((2-(trimethylsilyl)ethoxy)carbonyl)piperidine-3-carboxylic acid (2462.00 mg; 5.64 mmol; 1.00 eq.) in DCE (4.0ml) bis(2-oxo-1,3-oxazolidin-3-yl)phosphinic chloride (1435.59 mg; 5.64 mmol; 1.00 eq.) was added. After stirring for 15mins at RT, DIEA (1.52 ml; 8.46 mmol; 1.50 eq.) and Phenylamine (525.18 mg; 5.64 mmol; 1.00 eq.) were added. The reaction mixture was stirred overnight at RT. The reaction mixture was added 50ml of DCM, washed with brine, dried and concentrated. The residue was subjected to SNAP column (100g) for purification, eluted with 20-50 % EtOAc in hexane to yield the title compound (2600mg, yield 90.1%)

Step 3: benzyl ((3-(phenylcarbamoyl)piperidin-3-yl)methyl)carbamate

The reaction mixture of 2-(trimethylsilyl)ethyl 3-((((benzyloxy)carbonyl)amino)methyl)-3-(phenylcarbamoyl)piperidine-1-carboxylate (1200.00 mg; 2.35 mmol; 1.00 eq.) and TBAF (10.00 ml; 9.38 mmol; 4.00 eq.) in THF was stirred at RT overnight. The crude was purified by prep HPLC to afford the title compound (600mg, 69.6%)

Representative synthesis of Intermediate E (Scheme 5)



tert-Butyl 3-aminomethyl-3-phenylcarbamoyl-pyrrolidine-1-carboxylate (racemic)

Step 1: 3-(Benzyloxycarbonylamino-methyl)-pyrrolidine-1,3-dicarboxylic acid 1-tert-butyl ester

A reaction mixture of 3-(benzyloxycarbonylamino-methyl)-pyrrolidine-3-carboxylic acid, hydrochloride (2000.00 mg; 6.35 mmol; 1.00 eq.) and di-tert-butyl dicarbonate (1.77 ml; 8.26 mmol; 1.30 eq.) was suspended in t-BuOH (25.00 ml) and then treated with aqueous 2N NaOH (3.97 ml; 7.94 mmol; 1.25 eq.). The reaction mixture was warmed to 75 °C (immediate effervescence was observed) for 2 h. The reaction mixture was concentrated. The residue was dissolved in EtOAc (100ml) and washed with water and brine, dried over MgSO₄, concentrated and dried to afford 3-(benzyloxycarbonylamino-methyl)-pyrrolidine-1,3-dicarboxylic acid 1-tert-butyl ester (2200mg, 89.4%).


Step 2: 3-(Benzyloxycarbonylamino-methyl)-3-phenylcarbamoyl-pyrrolidine-1-carboxylic acid tert-butyl ester

To a solution of 3-(benzyloxycarbonylamino-methyl)-pyrrolidine-1,3-dicarboxylic acid 1-tert-butyl ester (2200.00 mg; 5.81 mmol; 1.00 eq.) in DCE (40.0ml), bis(2-oxo-1,3-oxazolidin-3-yl)phosphinic chloride (1627.95 mg; 6.40 mmol; 1.10 eq.) was added. After stirring for 15mins. DIEA (1.25 ml; 6.98 mmol; 1.20 eq.) and phenylamine (595.55 mg; 6.40 mmol; 1.10 eq) were added. The reaction mixture was stirred overnight at RT. DCM (100ml) was added. The solution was washed with brine, dried and concentrated to give the crude product, which was subjected for biotage for purification (20-50 % EtOAc in hexane) to afford the title compound (1800mg, 68.3%).

Step 3: tert-Butyl 3-aminomethyl-3-phenylcarbamoyl-pyrrolidine-1-carboxylate

To 3-(benzyloxycarbonylamino-methyl)-3-phenylcarbamoyl-pyrrolidine-1-carboxylic acid tert-butyl ester (1200.00 mg; 2.65 mmol; 1.00 eq.) in 30ml of methanol was added of ammonium formate (1668.40 mg; 26.46 mmol; 10.00 eq.) and 10%Pd/C (wet) 1.2g. The mixture was stirred at 65°C for 1hr and then filtered. The filtrate was concentrated to give the crude, which was dissolved in DCM and washed with 5% NaHCO₃, brine, dried and concentrated to afford the title compound (820mg, 97%).

Representative synthesis of Intermediate F (Scheme 3)



tert-butyl 3-amino-4-((3-fluorophenyl)carbamoyl)pyrrolidine-1-carboxylate (racemic)

Step 1: 4-(2-Trimethylsilyl-ethoxycarbonylamino)-pyrrolidine-1,3-dicarboxylic acid 1-tert-butyl ester 3-ethyl ester

A reaction mixture of 4-(2-Trimethylsilyl-ethoxycarbonylamino)-pyrrolidine-1,3-dicarboxylic acid 1-tert-butyl ester 3-ethyl ester (4443.00 mg; 11.04 mmol; 1.00 eq.), LiOH.H₂O (1389.45 mg; 33.11 mmol; 3.00 eq.) in water (16.50 ml) and MeOH (16.50 ml) were stirred at 45°C overnight. After removing partial of solvents, the mixture was extracted with DCM, washed with 5% citric acid, then brine, dried and concentrated to yield light yellow oil as the title compound (3200mg, 77.4%).

Step 2: 3-(3-Fluoro-phenylcarbamoyl)-4-(2-trimethylsilyl-ethoxycarbonylamino)-pyrrolidine-1-carboxylic acid tert-butyl ester


To a solution of 4-(2-trimethylsilyl-ethoxycarbonylamino)-pyrrolidine-1,3-dicarboxylic acid 1-tert-butyl ester (3200.00 mg; 8.54 mmol; 1.00 eq.) in DCE (40.0ml), bis(2-oxo-1,3-oxazolidin-3-yl)phosphinic chloride (2392.68 mg; 9.40 mmol; 1.10 eq.) was added. After

stirring for 15mins. DIEA (1.69 ml; 9.40 mmol; 1.10 eq.) and 3-fluoro-phenylamine (1044.40 mg; 9.40 mmol; 1.10 eq) were added. The reaction mixture was stirred overnight at 45°C. The reaction solution was diluted with EtOAc, washed, dried and concentrated to give the crude product, which was subjected for SNAP (100g) column for purification (eluted with 20-50 % EtOAc in hexane) to afford the title compound (1700mg, 42.5%).

Step 3: tert-Butyl 3-amino-4-(3-fluoro-phenylcarbamoyl)-pyrrolidine-1-carboxylate

To 3-(3-Fluoro-phenylcarbamoyl)-4-(2-trimethylsilanyl-ethoxycarbonylamino)-pyrrolidine-1-carboxylic acid tert-butyl ester (1700.00 mg; 3.64 mmol; 1.00 eq.) was added of TBAF (2851.67 mg; 10.91 mmol; 3.00 eq.) (1.0M in THF 11ml). The resulting mixture was stirred overnight. The reaction mixture was poured into water, washed with brine, then 5% NaHCO₃, brine, dried and concentrated to give the crude, which was added of ether (5ml). The precipitated white solid was filtered to afford the title compound (710mg, 60.4%).

Representative synthesis of Intermediate G (Scheme 7)



N-(3,4-dichlorobenzyl)pyrrolidin-3-amine (racemic)

Step 1: tert-Butyl 3-((3,4-dichlorobenzyl)amino)pyrrolidine-1-carboxylate


A solution of tert-butyl 3-aminopyrrolidine-1-carboxylate (1.5 g; 8.05 mmol; 1.00 eq and 3,4-dichlorobenzaldehyde (1339.00 mg; 7.65 mmol; 0.95 eq.) in methanol (10.00 ml) and acetic acid (1.00 ml) was stirred for 30 minutes at ambient temperature, then sodium cyanoborohydride (9.66 ml; 9.66 mmol; 1.20 eq.) was added. The reaction mixture was stirred for an additional 12 hours at ambient temperature at which time the reaction was quenched by pouring into aqueous ammonia (30 mL). The resulting inorganic precipitate was filtered off, and washed with dichloromethane (50 mL). The organic layer was separated and the remaining aqueous layer was extracted with dichloromethane (50 mL x 3). The combined organic extracts were washed with brine (50 mL), dried over sodium sulfate, filtered and concentrated in vacuo. The resultant oil is purified under flash chromatography using silicon column with gradient of ethyl acetate in hexanes to afford the desired intermediate, tert-Butyl 3-((3,4-dichlorobenzyl)amino)pyrrolidine-1-

carboxylate (2.47g; 7.16mmol; 88.9%), as a clear viscous oil. LC-MS (M+H =346, obsd. =288 (M-57))

Step 2: N-(3,4-dichlorobenzyl)pyrrolidin-3-amine

To a solution of tert-Butyl 3-((3,4-dichlorobenzyl)amino)pyrrolidine-1-carboxylate (100 mg; 0.29 mmol; 1.00 eq) in dichloromethane (3 mL) is added trifluoroacetic acid (0.04 mL; 0.58 mmol; 2.00 eq). The reaction flask is equipped with argon inlet and stirred until deemed complete as determined by TLC (10% methanol in dichloromethane). Upon completion, the solution is concentrated en vacuo to a dry residue, then taken up with dichloromethane and concentrated (x 3) to give the desired **Intermediate G** as the TFA salt in quantitative yield. LC-MS (M+H =246, obsd. =246).

Example Compounds according to Formula (I)



Example 1 5-((4-(4-(trifluoromethoxy)phenyl)pyrrolidin-3-yl)amino)quinazoline-8-carboxamide (trans_racemic)

Step 1: Tert-butyl 3-((8-cyano-quinazolin-5-yl)amino)-4-(4-(trifluoromethoxy)phenyl)pyrrolidine-1-carboxylate (trans_racemic)

A reaction mixture of Intermediate **A** (100.00 mg; 0.43 mmol; 1.00 eq.), tert-butyl 3-amino-4-(4-(trifluoromethoxy)phenyl)pyrrolidine-1-carboxylate (trans_racemic) (155.38 mg; 0.45 mmol; 1.05 eq.) and DIEA (0.15 ml; 0.85 mmol; 2.00 eq.) in NMP (2ml) was stirred at 120°C for overnight. The reaction mixture was purified by pre-HPLC to give the title compound (120 mg, 56.8%).

Step 2: Tert-butyl 3-((8-carbamoylquinazolin-5-yl)amino)-4-(4-(trifluoromethoxy)phenyl)pyrrolidine-1-carboxylate (trans_racemic)

Tert-butyl 3-(8-Cyano-quinazolin-5-ylamino)-4-(4-trifluoromethoxy-phenyl)-pyrrolidine-1-carboxylate (100.00 mg; 0.20 mmol; 1.00 eq.) was stirred with aqueous 2.0M NaOH (1ml; 2.00 mmol; 10.00 eq.) and aqueous 35% H₂O₂ (0.19 ml; 2.00 mmol; 10.00 eq.) in DMSO (8ml) at 40°C overnight. The crude was purified by pre-HPLC to yield the title compound (58mg, 56.0%).

Step 3: 5-((4-(4-(trifluoromethoxy)phenyl)pyrrolidin-3-yl)amino)quinazoline-8-carboxamide (trans_racemic)

To Tert-butyl 3-((8-carbamoylquinazolin-5-yl)amino)-4-(4-


(trifluoromethoxy)phenyl)pyrrolidine-1-carboxylate (trans_racemic) 58.0mg was added

1ml of methanol, and 1ml of 4.0M HCl in dioxane. The resulting mixture was stirred at RT for 3hrs. The solvents were removed to give the crude product, which was treated

with acetonitrile to yield **Example 1** (42mg) as HCl salt. LC-MS (M+H = 418, obsd. = 418). ¹HNMR: (DMSO) δ 10.12(s, 1H), 9.80 (s, 1H), 9.67 (s, 1H), 9.32 (1s, 1H), 8.49-8.51

(d, 2H), 8.05-8.07 (d, 1H), 7.67-7.70(d, 2H), 7.38-7.40(d, 2H), 6.72-6.7 (d, 1H), 4.65 (t, 1H), 3.77-3.81 (m, 3H), 3.35-3.40(m, 1H), 3.15-3.25(m, 1H), 3.20(s, 1H).


P70S6K IC₅₀: 23 nM



Example 2 5-((4-(3-chloro-5-fluorophenyl)pyrrolidin-3-yl)amino)quinazoline-8-carboxamide (Trans_racemic)

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling with tert-butyl 3-amino-4-(3-chloro-5-fluorophenyl)pyrrolidine-1-carboxylate(trans_racemic). LC-MS (M+H = 386,obsd. = 386/388).


P70S6K IC₅₀: 160 nM



Example 3 5-((4-phenylpiperidin-3-yl)amino)quinazoline-8-carboxamide (trans_racemic)

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling with tert-butyl 3-amino-4-phenylpiperidine-1-carboxylate (trans_racemic). LC-MS. (M+H = 348, obsd = 348).


P70S6K IC₅₀: 4.1 nM



Example 4 5-(((4-(3-fluoro-4-(trifluoromethyl)phenyl)piperidin-3-yl)amino)quinazoline-8-carboxamide(trans_racemic)

- 5 The title compound was synthesized according to the procedure described for the preparation of Example 1 by coupling with tert-butyl 3-amino-4-(3-fluoro-4-(trifluoromethyl)phenyl)piperidine-1-carboxylate (trans_racemic). LC-MS. (M+H = 434, obsd.=434). ¹HNMR: (DMSO) δ 9.85 (s, 1H), 9.20 (s, 1H), 9.24-9.26 (d, 2H), 8.50-8.52 (d, 1H), 7.73-7.75 (m, 1H), 7.47-7.49(d, 1H), 7.29-7.30 (d, 1H), 6.88-6.90 (d, 1H), 4.50-4.55 (d, 1H), 3.58-3.60 (m, 1H), 3.46-3.49 (m, 1H), 3.35-3.40 (m, 2H), 3.12-3.14(m, 1H), 2.80-2.84 (m, 1H), 2.30-2.35 (m, 1H).
- 10


P70S6K IC₅₀: 5.9 nM



- 15 **Example 5** 5-(((3R)-4-(3-fluoro-4-(trifluoromethyl)phenyl)piperidin-3-yl)amino)quinazoline-8-carboxamide (chiral, one of enantiomer of **Example 4** with unknown absolute configuration)

The title compound was isolated via chiral chromatography of **Example 4**. LC-MS (M+H = 434, obsd.= 434).


- 20 P70S6K IC₅₀: 9.9 nM



Example 6 5-(((3S,4S)-4-(3-fluoro-4-(trifluoromethyl)phenyl)piperidin-3-yl)amino)quinazoline-8-carboxamide (chiral, second enantiomer of **Example 4** with unknown absolute configuration)

The title compound was isolated via chiral chromatography of **Example 4**. LC-MS (M+H = 434, obsd.= 434).

P70S6K IC₅₀: 54 nM




Example 7 (S)-5-((1-(3-fluorophenyl)-2-(methylamino)ethyl)amino)quinazoline-8-carboxamide (chiral)

A reaction mixture of **A** (100.00 mg; 0.43 mmol; 1.00 eq.), N-[(S)-2-Amino-2-(3-fluorophenyl)-ethyl]-4-nitro-benzenesulfonamide (152.24 mg; 0.45 mmol; 1.05 eq.) and DIEA (0.15 ml; 0.85 mmol; 2.00 eq.) in NMP (1ml) was stirred at 120°C for overnight. The reaction solution was cooled and then poured into water, extracted with EtOAc and washed with brine, dried, and concentrated to give the crude product, which was subjected to SNAP column (25g, eluted with 20-80% EtOAc in hexane) to afford N-[(S)-2-(8-Cyano-quinazolin-5-ylamino)-2-(3-fluoro-phenyl)-ethyl]-N-methyl-4-nitro-benzenesulfonamide (120mg).

To a stirred solution of N-[(S)-2-(8-Cyano-quinazolin-5-ylamino)-2-(3-fluoro-phenyl)-ethyl]-N-methyl-4-nitro-benzenesulfonamide (120.00 mg; 0.24 mmol; 1.00 eq.) in DMSO (8ml) was added aqueous 2.0M NaOH (0.59ml; 1.18 mmol; 5.00 eq.) and aqueous 35% hydrogen peroxide (0.14 ml; 1.42 mmol; 6.00 eq.) and continued stirring at 40°C overnight. The crude product was purified by prep-HPLC to yield the desired product, which was then added of acetonitrile (2ml), thiophenol (0.1ml), CsCO₃ (200mg), and

stirred at 40°C overnight. The crude was purified by prepHPLC to afford the title product.
LC-MS (M+H = 434, obsd.= 434).

P70S6K IC₅₀: >1000 nM




5

Example 8 5-(4-(2-amino-1-(4-(trifluoromethyl)phenyl)ethyl)piperazin-1-yl)quinazoline-8-carboxamide (racemic)

The title compound was synthesized according to the procedure described for the preparation of **Example 7** by coupling with 4-nitro-N-(2-(piperazin-1-yl)-2-(4-(trifluoromethyl)phenyl)ethyl)benzenesulfonamide, followed by hydrolysis and deprotection. LC-MS. (M+H = 445, obsd= 445).

10

P70S6K IC₅₀: 210 nM




15

Example 9 5-(3-amino-3-((2,4-difluorobenzamido)methyl)pyrrolidin-1-yl)quinazoline-8-carboxamide (racemic)

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling with N-((3-aminopyrrolidin-3-yl)methyl)-2,4-difluorobenzamide, followed by nitrile hydrolysis. LC-MS (M+H = 427, obsd = 427).

20


p70S6K IC₅₀: >1000 nM



Example 10 5-(3-amino-3-((4-fluoro-2-hydroxybenzamido)methyl)pyrrolidin-1-yl)quinazoline-8-carboxamide(racemic)

The title compound was isolated as a side product from the synthesis of **Example 9**. LC-


- 5 MS (M+H = 425, obsd = 425).
p70S6K IC₅₀: >1000 nM



Example 11 5-(3-(aminomethyl)-3-(phenylcarbamoyl)piperidin-1-yl)quinazoline-8-carboxamide (racemic)

- 10 The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling with benzyl ((3-(phenylcarbamoyl)piperidin-3-yl)methyl)carbamate, followed by hydrolysis and de-protection. LC-MS (M+H = 404, obsd = 405).


- 15 p70S6K IC₅₀: >1000 nM



Example 12 5-(((3-(phenylcarbamoyl)pyrrolidin-3-yl)methyl)amino)quinazoline-8-carboxamide (racemic)

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling with tert-butyl 3-(aminomethyl)-3-(phenylcarbamoyl)pyrrolidine-1-carboxylate (racemic. LC-MS (M+H =390, obsd.= 391).


p70S6K IC₅₀: 650 nM



Example 13 5-((-4-((3-fluorophenyl)carbamoyl)pyrrolidin-3-yl)amino)quinazoline-8-carboxamide (trans_racemic)


The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling with tert-butyl 3-amino-4-((3-fluorophenyl)carbamoyl)pyrrolidine-1-carboxylate (trans_racemic). LC-MS (M+H =395, obsd. = 395).


p70S6K IC₅₀: 290 nM



Example 14 5-(((4-(phenylcarbamoyl)piperidin-4-yl)methyl)amino)quinazoline-8-carboxamide (racemic)


The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling with 2-(trimethylsilyl)ethyl 4-(aminomethyl)-4-(phenylcarbamoyl)piperidine-1-carboxylate (racemic). LC-MS (M+H =404, obsd. = 405). p70S6K IC₅₀: >1000 nM





Example 15 5-(3-amino-3-(phenylcarbamoyl)pyrrolidin-1-yl)quinazoline-8-carboxamide (racemic)

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling with tert-butyl (3-(phenylcarbamoyl)pyrrolidin-3-yl)carbamate (racemic). LC-MS (M+H =377,obsd. = 377).
p70S6K IC₅₀: 2300 nM




Example 16 (S)-5-((1-(3-fluorophenyl)-2-(methylamino)ethyl)amino)quinoline-8-carboxamide (chiral)

A reaction mixture of 5-Bromo-quinoline-8-carboxylic acid methyl ester (600.00 mg; 2.25 mmol; 1.00 eq.), N-[(S)-2-Amino-2-(3-fluoro-phenyl)-ethyl]-N-methyl-4-nitro-benzenesulfonamide (876.48 mg; 2.48 mmol; 1.10 eq.), potassium phosphate, tribasic (957.26 mg; 4.51 mmol; 2.00 eq.) , dicyclohexyl-(2',4',6'-triisopropyl-biphenyl-2-yl)-phosphane (214.99 mg; 0.45 mmol; 0.20 eq.), palladium (+2) acetate (50.62 mg; 0.23 mmol; 0.10 eq.) and toluene (5ml) in microwave tube were stirred at 100°C overnight. The crude was purified by prep HPLC (Basic condition) to afford (S)-5-{1-(3-Fluoro-phenyl)-2-[methyl-(4-nitro-benzenesulfonyl)-amino]-ethylamino}-quinoline-8-carboxylic acid methyl ester (250mg). LC-MS (M+H =339,obsd. = 339).

The above methyl ester (250.00 mg; 0.46 mmol; 1.00 eq.) was added 7.0M ammonia in MeOH (15.00 ml) , stirred at 50°C for 5 days. The reaction mixture was concentrated and purified by prep HPLC to yield (S)-5-{1-(3-Fluoro-phenyl)-2-[methyl-(4-nitro-benzenesulfonyl)-amino]-ethylamino}-quinoline-8- carboxamide, which was dissolved in acetonitrile (5ml) and added of benzenethiol (204.58 mg; 1.86 mmol; 4.00 eq.) and Cs₂CO₃ (605.00 mg; 1.86 mmol; 4.00 eq.). The reaction mixture ewas stirred at RT

overnight. After workup, the residue was purified by prep HPLC to afford **Example 16** (30mg, 19%). LC-MS (M+H =339, obsd=339).

p70S6K IC₅₀: >1000 nM




5

Example 17 5-((-4-(quinolin-7-yl)pyrrolidin-3-yl)amino)quinoline-8-carboxamide (trans_racemic)

The title compound was synthesized according to the procedure described for the preparation of **Example 16** by using methyl 5-bromoquinazoline-8-carboxylate coupled with tert-butyl 3-amino-4-(quinolin-7-yl)pyrrolidine-1-carboxylate (trans_racemic), conversion of the methyl ester to the amide with NH₃ in MeOH, followed by N-Boc deprotection to afford **Example 17**. LC-MS (M+H =384, obsd= 384).

10

p70S6K IC₅₀: 190 nM




15

Example 18 5-(3-phenylpiperazin-1-yl)quinoline-8-carboxamide (racemic)

The title compound was synthesized according to the procedure described for the preparation of **Example 16** by using methyl 5-bromoquinazoline-8-carboxylate coupled with tert-butyl 2-phenylpiperazine-1-carboxylate, conversion of the methyl ester to the amide with NH₃ in MeOH, followed by N-Boc deprotection to afford **Example 18**. LC-MS. (M+H = 333, obsd = 333).

20


p70S6K IC₅₀: 670 nM



Example 19 5-((3-phenylazetidin-3-yl)amino)quinoline-8-carboxamide

The title compound was synthesized according to the procedure described for the preparation of **Example 16** by using methyl 5-bromoquinazoline-8-carboxylate coupled with tert-butyl 3-amino-3-phenylazetidine-1-carboxylate (racemic), conversion of the methyl ester to the amide with NH_3 in MeOH, followed by N-Boc deprotection to afford **Example 19**. LC-MS ($\text{M}+\text{H} = 319$, obsd = 319).


p70S6K IC_{50} : >1000 nM



Example 20 5-(((3-(phenylcarbamoyl)pyrrolidin-3-yl)methyl)amino)quinoline-8-carboxamide (racemic)

The title compound was synthesized according to the procedure described for the preparation of **Example 16** by using methyl 5-bromoquinazoline-8-carboxylate coupled with tert-butyl 3-(aminomethyl)-3-(phenylcarbamoyl)pyrrolidine-1-carboxylate (racemic), conversion of the methyl ester to the amide with NH_3 in MeOH, followed by N-Boc deprotection to afford **Example 20**. LC-MS ($\text{M}+\text{H} = 390$, obsd. = 390).

p70S6K IC_{50} : >1000 nM




Example 21 5-((-4-(3-(trifluoromethoxy)phenyl)pyrrolidin-3-yl)amino)quinoline-8-carboxamide (racemic_trans)

A reaction mixture of 5-Bromo-quinoline-8-carbonitrile (200.00 mg; 0.86 mmol; 1.00 eq), (3-amino-4-(3-trifluoromethoxy-phenyl)-pyrrolidine-1-carboxylic acid tert-butyl ester (0.41 ml; 1.03 mmol; 1.20 eq.) (trans_racemic), sodium; 2-methyl-propan-2-olate (181.43 mg; 1.89 mmol; 2.20 eq.), and dicyclohexyl-(2',4',6'-triisopropyl-biphenyl-2-yl)-phosphane (122.73 mg; 0.26 mmol; 0.30 eq.) in toluene (5ml) in microwave tube was degas, followed by adding Pd₂(dba)₃ (74.02 mg; 0.13 mmol; 0.15 eq.). The resulting mixture was stirred for 10min at RT, placed in microwave at 100°C for 20min. The reaction mixture was concentrated and re-dissolved in DMSO, purified by prep HPLC (Basic condition, 70-75% acetonitrile in water) to give 3-(8-cyano-quinolin-5-ylamino)-4-(3-trifluoromethoxy-phenyl)-pyrrolidine-1-carboxylic acid tert-butyl ester (trans_racemic) (150mg, 35%).

A reaction mixture of 3-(8-Cyano-quinolin-5-ylamino)-4-(3-trifluoromethoxy-phenyl)-pyrrolidine-1-carboxylic acid tert-butyl ester (trans_racemic) (136.00 mg; 0.27 mmol; 1.00 eq.) in concentrate sulfate acid (2.00 ml; 37.52 mmol; 137.53 eq.) was heated at 100°C for 1hr. The reaction solution was cooled and poured to crashed ice. Solid sodium hydroxide was added to adjust PH=9. The separated oil was purified by prep HPLC to afford **Example 21** (15mg). LC-MS(M+H =417, obsd= 417).


p70S6K IC₅₀: 22 nM



Example 22 5-((((4S)-4-(3-(trifluoromethoxy)phenyl)pyrrolidin-3-yl)amino)quinoline-8-carboxamide (chiral, 1st enantiomer of **Example 21** with unknown absolute configuration)

The title compound was isolated via chiral chromatography of racemic **Example 21**. LC-MS (M+H =417, obsd= 417).


p70S6K IC₅₀: 31 nM



Example 23 5-((((3S,4R)-4-(3-(trifluoromethoxy)phenyl)pyrrolidin-3-yl)amino)quinoline-8-carboxamide (chiral, 2nd enantiomer of **Example 21** with unknown absolute configuration)

- 5 The title compound was isolated via chiral chromatography of racemic **Example 21**. LC-MS (M+H =417, obsd= 417). ¹HNMR: (DMSO) δ 9.92 (s, 1H), 9.76 (s, 1H), 8.20-8.23 (d, 1H), 7.98-8.00(d, 1H), 7.30-7.32 (d, 1H), 7.25 (s, 1H), 7.15 (t, 1H), 6.80-6.82 (d, 1H), 4.60-4.63 (m, 1H), 3.80-3.90 (m, 2H), 3.65 (t, 1H), 3.30(t, 1H), 3.20-3.25 (m, 1H).
p70S6K IC₅₀: 690 nM


10



Example 24 5-((-4-(3-(trifluoromethoxy)phenyl)pyrrolidin-3-yl)amino)quinoline-8-carboxamide (racemic_cis)

The title compound was isolated as a side product from the synthesis of **Example 21**.

- 15 LC-MS. (M+H =417, obsd. = 417).
p70S6K IC₅₀: 23 nM




Example 25 5-((4-(3-fluorophenyl)piperidin-3-yl)amino)quinoline-8-carboxamide
(racemic_trans)

The title compound was synthesized according to the procedure described for the

preparation of **Example 16** by using methyl 5-bromoquinazoline-8-carboxylate coupled with tert-butyl 3-amino-4-(3-fluorophenyl)piperidine-1-carboxylate (trans_racemic), conversion of the methyl ester to the amide with NH₃ in MeOH, followed by N-Boc deprotection to afford **Example 25**. LC-MS (M+H =365, obsd.= 365).


5 p70S6K IC₅₀: 520 nM



Example 26 5-(((4R)-4-(3-chloro-5-fluorophenyl)pyrrolidin-3-yl)amino)quinoline-8-carboxamide (racemic_trans)

10 The title compound was synthesized according to the procedure described for the preparation of **Example 16** by using methyl 5-bromoquinazoline-8-carboxylate coupled with tert-butyl 3-amino-4-(3-chloro-5-fluorophenyl)pyrrolidine-1-carboxylate (trans_racemic), conversion of the methyl ester to the amide with NH₃ in MeOH, followed by N-Boc deprotection to afford **Example 26**. LC-MS. (M+H =385, obsd= 384/386).

15 p70S6K IC₅₀: 350 nM




Example 27 5-(((4S)-4-(3-fluorophenyl)pyrrolidine-3-carboxamido)quinoline-8-carboxamide (chiral, absolutely)

20 A reaction mixture of 5-amino-quinoline-8-carbonitrile (1000.00 mg; 4.29 mmol; 1.00 eq.) in sulfuric acid (5.00 ml; 93.80 mmol; 21.86 eq.) was stirred at 100°C for 1h. The reaction mixture was then cooled, poured into ice and neutralized with 2N sodium hydroxide to PH=9. The precipitate was filtered, washed with water and dried to afford 5-aminoquinoline-8-carboxamide (800mg, yield 74.3%).

25 To a solution of tert-butyl (3R,4S)-4-(3-fluoro-phenyl)-pyrrolidine-1,3-dicarboxylate (125.00 mg; 0.40 mmol; 1.00 eq.) in DCE (4.0ml) bis(2-oxo-1,3-oxazolidin-3-

yl)phosphinic chloride (102.87 mg; 0.40 mmol; 1.00 eq.) was added. After stirring for 15mins at RT, DIEA (0.15 ml; 0.81 mmol; 2.00 eq.) and 5-aminoquinoline-8-carboxamide (75.65 mg; 0.40 mmol; 1.00 eq) were added. The reaction mixture was stirred overnight at 60°C. The crude was purified by prep HPLC to yield (4S)-tert-butyl 3-((8-carbamoylquinolin-5-yl)carbamoyl)-4-(3-fluorophenyl)pyrrolidine-1-carboxylate, which was added 1ml 4.0M HCl and 1ml methanol, stirred at RT for 3h. The reaction mixture was concentrated, neutralized to Ph 7 and purified by prep HPLC to afford **Example 27**. (M+H =479, obsd= 479).


p70S6K IC₅₀: >1000 nM



Example 28 5-((4S)-4-phenylpyrrolidine-2-carboxamido)quinoline-8-carboxamide
(chiral, absolutely)

The title compound was synthesized according to the procedure described for the preparation of **Example 27** by using 5-aminoquinoline-8-carboxamide coupled with (4S)-1-(tert-butoxycarbonyl)-4-phenylpyrrolidine-2-carboxylic acid, followed by removal of the protecting group to afford **Example 28**. LC-MS. (M+H =361, obsd= 361).


p70S6K IC₅₀: 578 nM



Example 29 5-(2-amino-2-phenylacetamido)quinoline-8-carboxamide (racemic)

The title compound was synthesized according to the procedure described for the preparation of **Example 27** by using 5-aminoquinoline-8-carboxamide coupled with 2-((tert-butoxycarbonyl)amino)-2-phenylacetic acid, followed by removal of the protecting group to afford **Example 29**. LC-MS (M+H =321, obsd= 321).

p70S6K IC₅₀: >1000 nM




Example 30 5-((3S)-3-phenylazetidine-2-carboxamido)quinoline-8-carboxamide (chiral, absolutely)

- 5 The title compound was synthesized according to the procedure described for the preparation of Example 27 by using 5-aminoquinoline-8-carboxamide coupled with (3S)-1-(tert-butoxycarbonyl)-3-phenylazetidine-2-carboxylic acid, followed by de-Boc to afford **Example 30**. LC-MS (M+H =347, obsd= 347).

p70S6K IC₅₀: >1000 nM

10




Example 31 5-(4-Fluorobenzylamino)-quinoline-8-carboxamide

- 15 To a solution of 5-aminoquinoline-8-carboxamide (120 mg, 0.64 mmol) in anhydrous DMF (2 mL) were added 4-fluorobenzyl bromide (242.34 mg, 1.28 mmol) and potassium carbonate (531.56 mg, 3.85 mmol). The suspension was heated at 50 °C overnight. The mixture was diluted with MeOH (4 mL) and the solid was filtered. The crude product was purified through reverse phase HPLC to afford **Example 31** (90 mg). LC-MS (M+H =296, obsd= 296).

p70S6K IC₅₀: 875 nM


20



Example 32 5-(2-Fluorobenzylamino)-quinoline-8-carboxamide

The title compound was synthesized according to the procedure described for the preparation of **Example 31**. LC-MS ($M+H = 296$, obsd= 296).

p70S6K IC_{50} : >1000 nM




5

Example 33 5-(3-Fluorobenzylamino)-quinoline-8-carboxamide

The title compound was synthesized according to the procedure described for the preparation of **Example 31**. LC-MS ($M+H = 296$, obsd= 296).

p70S6K IC_{50} : 369 nM


10



Example 34 5-(3-Chlorobenzylamino)-quinoline-8-carboxamide

The title compound was synthesized according to the procedure described for the preparation of **Example 31**. LC-MS ($M+H = 312$, obsd= 312).

15 p70S6K IC_{50} : 140 nM




Example 35 5-(3,4-Difluorobenzylamino)-quinoline-8-carboxamide

The title compound was synthesized according to the procedure described for the preparation of **Example 31**. LC-MS ($M+H = 314$, obsd= 314).

20


p70S6K IC_{50} : >1000 nM



Example 36 5-(3,4,5-trifluorobenzylamino)-quinoline-8-carboxamide

The title compound was synthesized according to the procedure described for the preparation of **Example 31**. LC-MS ($M+H = 332$, obsd. = 332).

5 p70S6K IC_{50} : >1000 nM




Example 37 5-(4-Fluoro-3-trifluoromethylbenzylamino)-quinoline-8-carboxamide

The title compound was synthesized according to the procedure described for the preparation of **Example 31**. LC-MS ($M+H = 364$, obsd. = 364).

10

p70S6K IC_{50} : 695 nM




Example 38 5-(4-Chloro-3-trifluoromethylbenzylamino)-quinoline-8-carboxamide

The title compound was synthesized according to the procedure described for the preparation of **Example 31**. LC-MS ($M+H = 380$, obsd. = 380).

15


p70S6K IC_{50} : >1000 nM



Example 39 5-((1-phenyl-2(pyrrolidin-1-yl)ethyl)amino)-quinoline-8-carboxamide

The title compound was synthesized according to the procedure described for the preparation of **Example 31**. LC-MS ($M+H = 361$, obsd. = 361).


5 p70S6K IC_{50} : >1000 nM



Example 40 5-(Phenylethylamino)-quinoline-8-carboxamide

The title compound was synthesized according to the procedure described for the preparation of **Example 31**. LC-MS ($M+H = 292$, obsd. = 292).


10 p70S6K IC_{50} : >1000 nM



Example 41 5-(Phenylpropylamino)-quinoline-8-carboxamide

The title compound was synthesized according to the procedure described for the preparation of **Example 31**. LC-MS ($M+H = 306$, obsd. = 306).


15 p70S6K IC_{50} : >1000 nM



Example 42 5-(Benzylamino)quinoline-8-carboxamide

The title compound was synthesized according to the procedure described for the preparation of **Example 31**. LC-MS ($M+H = 278$, obsd = 278).


5 p70S6K IC_{50} : 680 nM



Example 43 5-[3-amino-3-[(4-fluorophenyl)carbamoyl]pyrrolidin-1-yl]quinazoline-8-carboxamide (racemic)

10 The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling with 3-amino-N-(4-fluorophenyl)pyrrolidine-3-carboxamide (racemic), followed by hydrolysis of the nitrile intermediate. LC-MS ($M+H = 395$, obsd. = 395). 1H NMR (400 MHz, DMSO) δ 13.03 (s, 0H), 9.76 (d, $J = 13.4$ Hz, 2H), 9.24 – 9.17 (m, 1H), 8.54 (t, $J = 9.2$ Hz, 1H), 7.75 (dd, $J = 9.0, 5.0$ Hz, 2H), 7.55 (d, $J = 4.1$ Hz, 1H), 7.17 (dd, $J = 16.2, 7.4$ Hz, 2H), 6.88 (t, $J = 8.2$ Hz, 1H), 4.36 – 4.26 (m, 1H), 4.11 (dd, $J = 16.7, 9.7$ Hz, 1H), 3.68 (d, $J = 10.4$ Hz, 1H), 2.08 (t, $J = 8.5$ Hz, 1H), 1.45 – 1.18 (m, 1H).

15 p70S6K IC_{50} : 1200 nM




Example 44 5-(3-amino-3-((3,4-difluorophenyl)carbamoyl)pyrrolidin-1-yl)quinazoline-8-carboxamide (racemic)

The title compound was synthesized according to the procedure described for the

- 5 preparation of **Example 1** by coupling with 3-amino-N-(3,4-difluorophenyl)pyrrolidine-3-carboxamide (racemic), followed by hydrolysis of the nitrile intermediate. LC-MS (M+H = 413, obsd. = 413). ¹H NMR (400 MHz, DMSO) δ 9.79 – 9.71 (m, 2H), 9.21 (s, 1H), 8.56 (d, J = 8.9 Hz, 1H), 7.93 (ddd, J = 13.3, 7.5, 2.5 Hz, 1H), 7.59 – 7.49 (m, 2H), 7.41 (dd, J = 19.7, 9.1 Hz, 1H), 6.89 (d, J = 9.0 Hz, 1H), 4.31 (d, J = 10.3 Hz, 1H), 4.10 (dd, J = 16.5, 9.7 Hz, 1H), 3.78 (t, J = 8.0 Hz, 1H), 3.68 (d, J = 10.3 Hz, 1H), 2.08 (t, J = 8.5 Hz, 1H).

p70S6K IC₅₀: 1300 nM




- 15 **Example 45** 5-(3-amino-3-((3-fluorophenyl)carbamoyl)pyrrolidin-1-yl)quinazoline-8-carboxamide (racemic)

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling with 3-amino-N-(3-fluorophenyl)pyrrolidine-3-

- 20 carboxamide (racemic), followed by hydrolysis of the nitrile intermediate. LC-MS (M+H = 395, obsd. = 395). ¹H NMR (400 MHz, DMSO) δ 13.48 (s, 1H), 9.76 (d, J = 13.6 Hz, 2H), 9.21 (d, J = 6.6 Hz, 1H), 8.55 (t, J = 9.0 Hz, 1H), 7.74 (d, J = 11.4 Hz, 1H), 7.54 (dd, J = 28.9, 10.7 Hz, 2H), 7.42 – 7.26 (m, 1H), 7.19 (d, J = 8.8 Hz, 1H), 6.96 – 6.76 (m,

2H), 5.22 (s, 1H), 4.31 (dd, $J = 19.8, 11.2$ Hz, 2H), 4.11 (s, 1H), 3.93 – 3.73 (m, 2H), 3.69 (d, $J = 10.3$ Hz, 1H).

p70S6K IC₅₀: 2000 nM



5


Example 46 5-(3-amino-3-((4-fluoro-3-(trifluoromethyl)phenyl)carbamoyl)pyrrolidin-1-yl)quinazoline-8-carboxamide (racemic)

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling with 3-amino-N-(4-fluoro-3-(trifluoromethyl)-phenyl)pyrrolidine-3-carboxamide (racemic), followed by hydrolysis of the nitrile intermediate. LC-MS ($M+H = 395$, obsd. = 395). ¹H NMR (400 MHz, DMSO) δ 9.81 – 9.68 (m, 2H), 9.21 (d, $J = 7.7$ Hz, 1H), 8.56 (d, $J = 8.9$ Hz, 1H), 8.27 (dd, $J = 6.4, 2.6$ Hz, 1H), 8.10 – 8.00 (m, 1H), 7.60 – 7.46 (m, 2H), 6.88 (t, $J = 8.2$ Hz, 1H), 4.32 (d, $J = 10.4$ Hz, 1H), 4.10 (dd, $J = 16.7, 9.7$ Hz, 1H), 3.79 (t, $J = 8.0$ Hz, 1H), 3.69 (d, $J = 10.1$ Hz, 1H), 2.11 (d, $J = 5.1$ Hz, 1H).

10

15

p70S6K IC₅₀: 270 nM



Example 47 5-(3-(2-phenylacetamido)pyrrolidin-1-yl)quinazoline-8-carboxamide

20 (racemic)


The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling with 2-phenyl-N-(pyrrolidin-3-yl)acetamide (racemic), followed by hydrolysis of the nitrile intermediate. LC-MS ($M+H = 376$, obsd = 376). ¹H NMR (500 MHz, dmso) δ 9.75 (d, $J = 9.8$ Hz, 2H), 9.21 (d, $J = 1.3$ Hz, 1H), 8.59 – 8.52 (m, 1H), 8.43 (d, $J = 6.3$ Hz, 1H), 7.57 (s, 1H), 7.23 (td, $J = 15.7, 8.0$ Hz, 5H),

25

6.88 (d, $J = 8.9$ Hz, 1H), 4.37 (s, 1H), 4.03 (dd, $J = 10.2, 6.0$ Hz, 1H), 3.85 (t, $J = 8.5$ Hz, 1H), 3.76 (s, 1H), 3.59 – 3.51 (m, 1H), 3.40 (s, 2H), 2.21 (dd, $J = 12.6, 6.0$ Hz, 1H), 2.00 (d, $J = 5.3$ Hz, 1H).

p70S6K IC₅₀: 7900 nM

5




Example 48 5-(3-(3-(trifluoromethyl)benzamido)pyrrolidin-1-yl)quinazoline-8-carboxamide (racemic)

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling with N-(pyrrolidin-3-yl)-3-

(trifluoromethyl)benzamide (racemic), followed by hydrolysis of the nitrile intermediate.

LC-MS ($M+H = 430$, obsd = 430). ¹H NMR (500 MHz, dmsd) δ 9.81 (s, 1H), 9.74 (s, 1H), 9.22 (s, 1H), 8.90 (d, $J = 6.0$ Hz, 1H), 8.56 (d, $J = 8.9$ Hz, 1H), 8.22 – 8.13 (m, 2H), 7.92 (d, $J = 8.0$ Hz, 1H), 7.73 (t, $J = 8.0$ Hz, 1H), 7.57 (s, 1H), 6.91 (d, $J = 8.9$ Hz, 1H), 4.66 (d, $J = 5.4$ Hz, 1H), 4.17 – 4.08 (m, 1H), 3.93 (s, 1H), 3.87 – 3.74 (m, 2H), 2.99 (s, 0H), 2.54 (s, 2H), 2.33 (d, $J = 6.1$ Hz, 1H), 2.20 (s, 1H).

p70S6K IC₅₀: 2200 nM



Example 49 5-(3-(benzyl(2-(dimethylamino)ethyl)carbamoyl)pyrrolidin-1-yl)quinazoline-8-carboxamide (racemic)


The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling with N-benzyl-N-(2-

(dimethylamino)ethyl)pyrrolidine-3-carboxamide (racemic), followed by hydrolysis of the

nitrile intermediate. LC-MS ($M+H = 447$, obsd = 447). ¹H NMR (400 MHz, DMSO) δ 9.74

(s, 2H), 9.21 (d, $J = 4.5$ Hz, 1H), 8.54 (t, $J = 9.1$ Hz, 1H), 7.55 (s, 1H), 7.41 – 7.20 (m, 5H), 6.88 (dd, $J = 23.8, 8.7$ Hz, 1H), 4.74 (s, 1H), 4.65 (d, $J = 15.0$ Hz, 1H), 4.55 (d, $J = 14.6$ Hz, 1H), 3.91 (ddd, $J = 23.2, 16.7, 8.5$ Hz, 3H), 3.73 (s, 1H), 3.57 (d, $J = 44.4$ Hz, 2H), 3.38 (dd, $J = 14.9, 7.8$ Hz, 2H), 3.00 (s, 1H), 2.54 (d, $J = 5.9$ Hz, 1H), 2.43 – 2.28 (m, 3H), 2.12 (d, $J = 14.3$ Hz, 5H).


p70S6K IC₅₀: 1600 nM



Example 50 5-(3-(3,4-difluorobenzamido)pyrrolidin-1-yl)quinazoline-8-carboxamide
(racemic)

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling with 3,4-difluoro-N-(pyrrolidin-3-yl)benzamide (racemic), followed by hydrolysis of the nitrile intermediate. LC-MS ($M+H = 398$, obsd = 398). ¹H NMR (500 MHz, dmsO) δ 9.81 (s, 1H), 9.74 (s, 1H), 9.22 (s, 1H), 8.71 (d, $J = 6.2$ Hz, 1H), 8.56 (d, $J = 8.7$ Hz, 1H), 7.97 – 7.88 (m, 1H), 7.76 (s, 1H), 7.56 (dt, $J = 16.9, 8.6$ Hz, 2H), 6.90 (d, $J = 8.9$ Hz, 1H), 4.64 – 4.58 (m, 1H), 4.10 (dd, $J = 10.4, 6.1$ Hz, 1H), 3.97 – 3.88 (m, 1H), 3.82 (s, 1H), 3.74 (dd, $J = 10.7, 4.3$ Hz, 1H), 2.30 (dd, $J = 12.9, 5.9$ Hz, 1H), 2.17 (d, $J = 5.2$ Hz, 1H).

p70S6K IC₅₀: 380 nM




Example 51 5-(3-(2-(4-fluorophenyl)acetamido)pyrrolidin-1-yl)quinazoline-8-carboxamide (racemic)

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling with 2-(4-fluorophenyl)-N-(pyrrolidin-3-yl)-acetamide (racemic), followed by hydrolysis of the nitrile intermediate. LC-MS ($M+H =$

394, obsd = 394). ¹H NMR (400 MHz, DMSO) δ 9.76 (s, 2H), 9.21 (s, 1H), 8.55 (d, *J* = 8.5 Hz, 1H), 8.44 (d, *J* = 6.0 Hz, 1H), 7.59 (s, 1H), 7.25 (s, 2H), 7.08 (t, *J* = 8.0 Hz, 2H), 6.88 (d, *J* = 9.0 Hz, 1H), 4.36 (s, 1H), 4.04 (d, *J* = 6.6 Hz, 1H), 3.85 (s, 1H), 3.77 (s, 1H), 3.54 (d, *J* = 10.1 Hz, 1H), 3.38 (d, *J* = 14.2 Hz, 2H), 2.20 (s, 1H), 1.99 (s, 1H).


5 p70S6K IC₅₀: 5400 nM



Example 52 5-(3-((3,4-dichlorobenzyl)amino)pyrrolidin-1-yl)quinazoline-8-carboxamide (racemic)

10 The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling with N-(3,4-dichlorobenzyl)pyrrolidin-3-amine (racemic), followed by hydrolysis of the nitrile intermediate. LC-MS (*M*+*H* = 417, obsd = 417). ¹H NMR (500 MHz, dmso) δ 9.73 (s, 2H), 9.18 (s, 1H), 8.52 (d, *J* = 8.9 Hz, 1H), 7.59 (d, *J* = 1.8 Hz, 1H), 7.55 – 7.47 (m, 2H), 7.34 – 7.28 (m, 1H), 6.82 (d, *J* = 9.0 Hz, 1H), 3.91 (dd, *J* = 10.4, 5.5 Hz, 1H), 3.88 – 3.81 (m, 1H), 3.74 (s, 2H), 3.68 (s, 1H), 3.54 (dd, *J* = 10.3, 3.9 Hz, 1H), 3.38 (s, 1H), 2.63 (s, 1H), 2.13 – 2.04 (m, 1H), 1.94 (d, *J* = 5.3 Hz, 1H).

p70S6K IC₅₀: 490 nM




20

Example 53 5-(3-((3,4-dichlorobenzyl)(methyl)amino)pyrrolidin-1-yl)quinazoline-8-carboxamide (racemic)

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling with N-(3,4-dichlorobenzyl)-N-methylpyrrolidin-3-

amine (racemic), followed by hydrolysis of the nitrile intermediate. LC-MS ($M+H = 431$, obsd = 431). ^1H NMR (400 MHz, DMSO) δ 9.78 (d, $J = 17.6$ Hz, 2H), 9.21 (s, 1H), 8.54 (d, $J = 8.7$ Hz, 1H), 7.63 – 7.50 (m, 3H), 7.34 (d, $J = 8.1$ Hz, 1H), 6.90 (d, $J = 8.9$ Hz, 1H), 3.94 – 3.73 (m, 4H), 3.61 (s, 2H), 3.20 (s, 2H), 2.27 (s, 2H), 2.18 (s, 3H), 2.06 – 1.93 (m, 2H), 0.08 (s, 1H).


p70S6K IC_{50} : 570 nM



Example 54 5-((3-((3,4-difluorobenzamido)methyl)pyrrolidin-1-yl)quinazoline-8-carboxamide (racemic)

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling 5-bromo-quinazoline-8-carbonitrile with 3,4-difluoro-N-(pyrrolidin-3-ylmethyl)-benzamide (racemic), followed by hydrolysis of the nitrile intermediate. LC-MS ($M+H = 412$, obsd = 412). ^1H NMR (400 MHz, DMSO) δ 9.81 (s, 1H), 9.75 (d, $J = 4.0$ Hz, 1H), 9.20 (s, 1H), 8.72 (t, $J = 5.6$ Hz, 1H), 8.53 (d, $J = 8.9$ Hz, 1H), 7.89 (ddd, $J = 11.5, 7.8, 2.0$ Hz, 1H), 7.74 (dd, $J = 5.1, 3.3$ Hz, 1H), 7.62 – 7.50 (m, 2H), 6.85 (d, $J = 9.0$ Hz, 1H), 3.92 – 3.74 (m, 3H), 3.65 (dd, $J = 10.2, 7.0$ Hz, 1H), 3.41 (dtd, $J = 19.7, 13.3, 6.1$ Hz, 2H), 2.63 (dt, $J = 13.7, 6.8$ Hz, 1H), 2.16 (td, $J = 12.0, 5.9$ Hz, 1H), 1.85 (dq, $J = 15.2, 7.5$ Hz, 1H).

p70S6K IC_{50} : 233 nM




Example 55 5-(3-((2-fluoro-4-(trifluoromethyl)benzamido)methyl)pyrrolidin-1-yl)quinazoline-8-carboxamide (racemic)

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling with 2-fluoro-N-(pyrrolidin-3-ylmethyl)-4-

(trifluoromethyl)benzamide (racemic), followed by hydrolysis of the nitrile intermediate. LC-MS (M+H = 462, obsd = 462). ¹H NMR (400 MHz, DMSO) δ 10.87 (s, 1H), 9.80 (s, 1H), 9.75 (d, J = 3.9 Hz, 1H), 9.19 (s, 1H), 8.53 (d, J = 8.9 Hz, 1H), 7.89 (d, J = 8.1 Hz, 1H), 7.52 (d, J = 3.8 Hz, 1H), 6.90 – 6.80 (m, 2H), 6.72 (d, J = 8.0 Hz, 1H), 3.87 – 3.77 (m, 3H), 3.61 (dd, J = 10.1, 7.3 Hz, 1H), 3.45 (s, 2H), 2.71 – 2.57 (m, 1H), 2.14 (dt, J = 19.7, 7.0 Hz, 1H), 1.93 – 1.79 (m, 1H).


p70S6K IC₅₀: 290 nM



Example 56 5-(3-(aminomethyl)piperidin-1-yl)quinazoline-8-carboxamide (racemic)

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling 5-bromo-quinazoline-8-carbonitrile with 3-N-boc-aminomethyl piperidine (racemic), followed by hydrolysis of the nitrile intermediate. LC-MS (M+H = 287, obsd. = 286.2/287.2).

P70S6K IC₅₀: 56000 nM AKT IC₅₀: >100000 nM




Example 57 5-((1-(m-tolyl)piperidin-4-yl)amino)quinazoline-8-carboxamide

The title compound was synthesized according to the procedure described for the

preparation of **Example 1** by coupling 5-Bromo-quinazoline-8-carbonitrile with 1-(m-tolyl)piperidin-4-amine, followed by hydrolysis of the nitrile intermediate. LC-MS (M+H = 363, obsd. = 362.2/363.3).

P70S6K IC₅₀: 22000 nM AKT IC₅₀: >100000 nM




Example 58 5-((1-methylpyrrolidin-3-yl)amino)quinazoline-8-carboxamide (racemic)

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling 5-Bromo-quinazoline-8-carbonitrile with 1-methylpyrrolidin-3-amine (racemic), followed by hydrolysis of the nitrile intermediate. LC-MS (M+H = 273, obsd. = 272.1/273.1).

P70S6K IC₅₀: 20000 nM

AKT IC₅₀: >100000 nM




Example 59 5-(3-((methylamino)methyl)piperidin-1-yl)quinazoline-8-carboxamide (racemic)

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling 5-Bromo-quinazoline-8-carbonitrile with tert-butyl methyl(piperidin-3-ylmethyl)carbamate (racemic), followed by hydrolysis of the nitrile intermediate. LC-MS (M+H = 300, obsd. = 300.1/301.1).

P70S6K IC₅₀: 5900 nM

AKT IC₅₀: 18000 nM




Example 60 (R)-5-((pyrrolidin-3-ylmethyl)amino)quinazoline-8-carboxamide (chiral)

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling 5-Bromo-quinazoline-8-carbonitrile with (S)-tert-butyl 3-(aminomethyl)pyrrolidine-1-carboxylate, followed by hydrolysis of the nitrile intermediate. LC-MS (M+H = 273, obsd. = 272.1/273.1).

P70S6K IC₅₀: 4800 nM

AKT IC₅₀: >100000 nM




Example 61 (S)-5-((pyrrolidin-3-ylmethyl)amino)quinazoline-8-carboxamide (chiral)

- 5 The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling 5-Bromo-quinazoline-8-carbonitrile with (R)-tert-butyl 3-(aminomethyl)pyrrolidine-1-carboxylate, followed by hydrolysis of the nitrile intermediate. LC-MS (M+H = 273, obsd. = 272.1/273.1).

P70S6K IC₅₀: 1700 nM

AKT IC₅₀: >100000 nM

10




Example 62 5-((pyridin-3-ylmethyl)amino)quinazoline-8-carboxamide

- 15 The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling 5-Bromo-quinazoline-8-carbonitrile with pyridin-3-ylmethanamine, followed by hydrolysis of the nitrile intermediate. LC-MS (M+H = 279, obsd. = 280.1/281.1).

P70S6K IC₅₀: 1600 nM

AKT IC₅₀: 70000 nM




- 20 **Example 63** 5-((3-chlorobenzyl)amino)quinazoline-8-carboxamide

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling 5-Bromo-quinazoline-8-carbonitrile with (3-

chlorophenyl)methanamine, followed by hydrolysis of the nitrile intermediate. LC-MS (M+H = 313, obsd. = 313.1/314.1).

P70S6K IC₅₀: 59.2 nM

AKT IC₅₀: 12000 nM



5


Example 64 5-(3-phenethylpyrrolidin-1-yl)quinazoline-8-carboxamide (racemic)

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling 5-Bromo-quinazoline-8-carbonitrile with 3-phenethylpyrrolidine (racemic), followed by hydrolysis of the nitrile intermediate. LC-MS (M+H = 347, obsd. = 347.2/348.2).

10

P70S6K IC₅₀: 240 nM

AKT IC₅₀: 3100 nM



Example 65 5-(3-(benzyloxy)pyrrolidin-1-yl)quinazoline-8-carboxamide (racemic)


The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling 5-Bromo-quinazoline-8-carbonitrile with 3-(benzyloxy)pyrrolidine (racemic), followed by hydrolysis of the nitrile intermediate. LC-MS (M+H = 349, obsd. = 349.1/350.1).

15

P70S6K IC₅₀: 710 nM

AKT IC₅₀: >100000 nM


20



Example 66 5-(3-hydroxy-3-phenylpyrrolidin-1-yl)quinazoline-8-carboxamide (racemic)

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling 5-Bromo-quinazoline-8-carbonitrile with 3-phenylpyrrolidin-3-ol (racemic), followed by hydrolysis of the nitrile intermediate. LC-MS (M+H = 335, obsd. = 335.1/336.1).

5 P70S6K IC₅₀: 1800 nM AKT IC₅₀: >100000 nM




Example 67 5-((3S,4R)-3-(hydroxymethyl)-4-(m-tolyl)pyrrolidin-1-yl)quinazoline-8-carboxamide (chiral)

10 The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling 5-Bromo-quinazoline-8-carbonitrile with ((3S,4R)-4-(m-tolyl)pyrrolidin-3-yl)methanol, followed by hydrolysis of the nitrile intermediate. LC-MS (M+H = 363, obsd. = 363.2/364.1).

P70S6K IC₅₀: 190 nM AKT IC₅₀: 8200 nM


15



Example 68 5-((2-(3-chlorophenyl)-2-(dimethylamino)ethyl)amino)quinazoline-8-carboxamide (racemic)

20 The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling 5-Bromo-quinazoline-8-carbonitrile with 1-(3-chlorophenyl)-N1,N1-dimethylethane-1,2-diamine (racemic), followed by hydrolysis of the nitrile intermediate. LC-MS (M+H = 370, obsd. = 370.1/371.1).

P70S6K IC₅₀: 2600 nM AKT IC₅₀: >100000 nM




Example 69 5-((2-phenyl-2-(piperazin-1-yl)ethyl)amino)quinazoline-8-carboxamide
(racemic)

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling 5-Bromo-quinazoline-8-carbonitrile with tert-butyl 4-(2-amino-1-phenylethyl)piperazine-1-carboxylate (racemic), followed by hydrolysis of the nitrile intermediate. LC-MS ($M+H = 363$, obsd. = 363.2/364.1).

P70S6K IC_{50} : 170 nM

AKT IC_{50} : 10000 nM




Example 70 5-((3-(4-fluorophenyl)piperidin-4-yl)amino)quinazoline-8-carboxamide
(diastereomeric racemic mixture)

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling 5-Bromo-quinazoline-8-carbonitrile with tert-butyl 4-amino-3-(4-fluorophenyl)piperidine-1-carboxylate (diastereomeric racemic mixtures), followed by hydrolysis of the nitrile intermediate. LC-MS ($M+H = 366$, obsd. = 366.1/367.2).

P70S6K IC_{50} : 550 nM

AKT IC_{50} : 2500 nM




Example 71 5-((4-(3-(trifluoromethyl)phenyl)pyrrolidin-3-yl)amino)quinazoline-8-carboxamide
(diastereomeric racemic mixtures)

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling 5-Bromo-quinazoline-8-carbonitrile with tert-butyl

3-amino-4-(3-(trifluoromethyl)phenyl)pyrrolidine-1-carboxylate (diastereomeric racemic mixtures), followed by hydrolysis of the nitrile intermediate. LC-MS (M+H = 402, obsd. = 402.2).

P70S6K IC₅₀: 21 nM AKT IC₅₀: 4500 nM

5




Example 72 5-((4-(3-(trifluoromethoxy)phenyl)pyrrolidin-3-yl)amino)quinazoline-8-carboxamide (diastereomeric racemic mixtures)

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling 5-Bromo-quinazoline-8-carbonitrile with tert-butyl 3-amino-4-(3-(trifluoromethoxy)phenyl)pyrrolidine-1-carboxylate (diastereomeric racemic mixtures), followed by hydrolysis of the nitrile intermediate. LC-MS (M+H = 418, obsd. = 418.2/418.9).

P70S6K IC₅₀: 100 nM AKT IC₅₀: >100000 nM


15



Example 73 5-((4-(m-tolyl)piperidin-3-yl)amino)quinazoline-8-carboxamide (diastereomeric racemic mixtures)

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling 5-Bromo-quinazoline-8-carbonitrile with tert-butyl 3-amino-4-(m-tolyl)piperidine-1-carboxylate (diastereomeric racemic mixtures), followed by hydrolysis of the nitrile intermediate. LC-MS (M+H = 362, obsd. = 362.2/363.2).

P70S6K IC₅₀: 390 nM AKT IC₅₀: 9000 nM




Example 74 5-((4-(3-(trifluoromethyl)phenyl)piperidin-3-yl)amino)quinazoline-8-carboxamide (diastereomeric racemic mixtures)

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling 5-bromo-quinazoline-8-carbonitrile with tert-butyl 3-amino-4-(3-(trifluoromethyl)phenyl)piperidine-1-carboxylate (diastereomeric racemic mixtures), followed by hydrolysis of the nitrile intermediate. LC-MS (M+H = 416, obsd. = 416.2/417.2).

P70S6K IC₅₀: 140 nM

AKT IC₅₀: 2100 nM




Example 75 5-((4-(4-chloro-3-fluorophenyl)piperidin-3-yl)amino)quinazoline-8-carboxamide (diastereomeric racemic mixtures)

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling 5-Bromo-quinazoline-8-carbonitrile with tert-butyl 3-amino-4-(4-chloro-3-fluorophenyl)piperidine-1-carboxylate (diastereomeric racemic mixtures), followed by hydrolysis of the nitrile intermediate. LC-MS (M+H = 400.8, obsd. = 400.2).

P70S6K IC₅₀: 8.1 nM


AKT IC₅₀: 130 nM



Example 76 5-((4-(3-chloro-4-fluorophenyl)piperidin-3-yl)amino)quinazoline-8-carboxamide (diastereomeric racemic mixtures)

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling 5-bromo-quinazoline-8-carbonitrile with *tert*-butyl 3-amino-4-(3-chloro-4-fluorophenyl)piperidine-1-carboxylate (diastereomeric racemic mixtures), followed by hydrolysis of the nitrile intermediate. LC-MS (M+H = 400.8, obsd. = 400.1/401.1).


P70S6K IC₅₀: 33 nM AKT IC₅₀: 570 nM



Example 77 5-((((3R,4R)-4-(3-chloro-4-fluorophenyl)piperidin-3-yl)amino)quinazoline-8-carboxamide (chiral, one of the enantiomers of **Example 76** with unknown absolute configuration)

The title compound was separated from the diastereomeric mixture (**Example 76**) via chiral SFC. LC-MS (M+H = 400.8, obsd. = 400.1/401.1). ¹HNMR (MeOH-*d*₄) δ 1.16(m, 1H), 1.31 (m, 1H), 1.78 (m, 1H), 2.34 (m, 1H), 2.89 (m, 1H), 3.20-3.32 (m, 3H), 6.29 (m, 1H), 6.97 (m, 1H), 7.25 (m, 1H), 7.39 (m, 1H), 8.33 (m, 1H), 9.21 (s, 1H), 9.93 (s, 1H).


P70S6K IC₅₀: 50 nM AKT IC₅₀: 850 nM



Example 78 5-((((3S,4S)-4-(3-chloro-4-fluorophenyl)piperidin-3-yl)amino)quinazoline-8-carboxamide (chiral, one of the enantiomers of **Example 76** with unknown absolute configuration)

The title compound was separated from the diastereomeric mixture (**Example 76**) via chiral SFC. LC-MS (M+H = 400.8, obsd. = 400.2).

P70S6K IC₅₀: 350 nM AKT IC₅₀: 5700 nM




Example 79 5-((4-(4-fluorophenyl)piperidin-3-yl)amino)quinazoline-8-carboxamide
(diastereomeric racemic mixtures)

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling 5-bromo-quinazoline-8-carbonitrile with *tert*-butyl 3-amino-4-(4-fluorophenyl)piperidine-1-carboxylate (diastereomeric racemic mixtures), followed by hydrolysis of the nitrile intermediate. LC-MS ($M+H = 366$, obsd. = 366.1/367.1).

P70S6K IC_{50} : 290 nM

AKT IC_{50} : 2000 nM




Example 80 5-(((3R,4R)-4-(4-fluorophenyl)piperidin-3-yl)amino)quinazoline-8-carboxamide (chiral, one of the enantiomers of **Example 79** with unknown absolute configuration)

The title compound was separated from the diastereomeric mixture (**Example 79**) via chiral SFC. LC-MS ($M+H = 366$, obsd. = 366.2).

P70S6K IC_{50} : 290 nM


AKT IC_{50} : 2200 nM



Example 81 5-(((3S,4S)-4-(4-fluorophenyl)piperidin-3-yl)amino)quinazoline-8-carboxamide (chiral, one of the enantiomers of **Example 79** with unknown absolute configuration)

The title compound was separated from the diastomeric mixture (**Example 79**) via chiral SFC. LC-MS (M+H = 366, obsd. = 366.1).

P70S6K IC₅₀: 52 nM AKT IC₅₀: 900 nM




5

Example 82 5-(3-amino-4-phenylpiperidin-1-yl)quinazoline-8-carboxamide (racemic)

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling 5-bromo-quinazoline-8-carbonitrile with 4-phenylpiperidin-3-ylamine, followed by hydrolysis of the nitrile intermediate. LC-MS (M + H = 347, obsd = 348).

10

P70S6K IC₅₀: 120 nM




Example 83 5-(3-amino-4-(4-(trifluoromethyl)phenyl)piperidin-1-yl)quinazoline-8-carboxamide (racemic)

15

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling 5-bromo-quinazoline-8-carbonitrile with 4-(4-trifluoromethyl-phenyl)-piperidin-3-ylamine, followed by hydrolysis of the nitrile intermediate. LC-MS (M + H = 415, obsd = 416).

20


P70S6K IC₅₀: 350 nM



Example 84 (S)-5-(3-amino-3-(phenylcarbamoyl)pyrrolidin-1-yl)quinazoline-8-carboxamide (pure enantiomer, absolute structure unknown).

The title compound was isolated by chiral HPLC from **Example 15**. LC-MS (M + H = 376, obsd = 377).


P70S6K IC₅₀: 2150 nM



Example 85 (R)-5-(3-amino-3-(phenylcarbamoyl)pyrrolidin-1-yl)quinazoline-8-carboxamide (pure enantiomer, absolute structure unknown).

The title compound was isolated by chiral HPLC from **Example 15**. LC-MS (M + H = 376, obsd = 377).


P70S6K IC₅₀: 11500 nM



Example 86 5-[(3S,4R)-3-Amino-4-(4-fluoro-phenyl)-piperidin-1-yl]-quinazoline-8-carboxylic acid amide (racemic_cis)

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling 5-bromo-quinazoline-8-carbonitrile with *cis*-4-(4-fluorophenyl)piperidin-3-amine, followed by hydrolysis of the nitrile intermediate. LC-MS (M + H = 366, obsd = 366).


P70S6K IC₅₀: 250 nM



Example 87 5-[(3R,4R)-3-Amino-4-(4-fluoro-phenyl)-piperidin-1-yl]-quinazoline-8-carboxylic acid amide. (racemic_trans)

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling 5-bromo-quinazoline-8-carbonitrile with *trans*-4-(4-fluorophenyl)piperidin-3-amine, followed by hydrolysis of the nitrile intermediate. LC-MS (M + H = 366, obsd = 366).


P70S6K IC₅₀: 7100 nM



Example 88 5-[(3S,4R)-3-Amino-4-(3-fluoro-phenyl)-piperidin-1-yl]-quinazoline-8-carboxylic acid amide (racemic_cis)

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling 5-bromo-quinazoline-8-carbonitrile with *cis*-4-(3-fluorophenyl)piperidin-3-amine, followed by hydrolysis of the nitrile intermediate. LC-MS (M + H = 366, obsd = 366).


P70S6K IC₅₀: **17100** nM



Example 89 5-[(3R,4R)-3-Amino-4-(3-fluoro-phenyl)-piperidin-1-yl]-quinazoline-8-carboxylic acid amide (racemic_trans)

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling 5-bromo-quinazoline-8-carbonitrile with *trans*-4-(3-fluorophenyl)piperidin-3-amine, followed by hydrolysis of the nitrile intermediate. LC-MS (M + H = 366, obsd = 366).

P70S6K IC₅₀: **2400** nM




Example 90 5-[4-(4-Trifluoromethyl-phenyl)-piperazin-1-yl]-quinazoline-8-carboxylic acid amide

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling with 5-bromo-quinazoline-8-carbonitrile and 1-(4-trifluoromethyl-phenyl)-piperazine, followed by hydrolysis of the nitrile intermediate. LC-MS. (M+H = 402, obsd= 402).

P70S6K IC₅₀: 24000 nM

AKT IC₅₀: >100000 nM




Example 91 5-[4-phenyl-piperazin-1-yl]-quinazoline-8-carboxylic acid amide

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling with 5-bromo-quinazoline-8-carbonitrile and 1-

5 phenylpiperazine, followed by hydrolysis of the nitrile intermediate. LC-MS. (M+H = 334, obsd= 334).

P70S6K IC₅₀: 3000 nM

AKT IC₅₀: 15000 nM




10 **Example 92** 5-[4-(3,4-difluoro-phenyl)-piperazin-1-yl]-quinazoline-8-carboxylic acid amide

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling with 5-bromo-quinazoline-8-carbonitrile and 1-(3,4-difluoro-phenyl)-piperazine, followed by hydrolysis of the nitrile intermediate. LC-

15 MS. (M+H = 370, obsd= 370).

P70S6K IC₅₀: 1000 nM


AKT IC₅₀: >100000 nM



Example 93 5-[4-(4-chloro-phenyl)-piperazin-1-yl]-quinazoline-8-carboxylic acid amide

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling with 5-bromo-quinazoline-8-carbonitrile and 1-(4-chlorophenyl)piperazine, followed by hydrolysis of the nitrile intermediate. LC-MS. (M+H = 368, obsd= 368).

P70S6K IC₅₀: 1000 nM




Example 94 5-[4-(3-chloro-4-fluorophenyl)-piperazin-1-yl]-quinazoline-8-carboxylic acid amide

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling with 5-bromo-quinazoline-8-carbonitrile and 1-(3-chloro-4-fluoro-phenyl)-piperazine, followed by hydrolysis of the nitrile intermediate. LC-MS. (M+H = 386, obsd= 386).

P70S6K IC₅₀: 12000 nM


AKT IC₅₀: >100000 nM



Example 95 5-[4-(3-fluoro-phenyl)-piperazin-1-yl]-quinazoline-8-carboxylic acid amide

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling with 5-bromo-quinazoline-8-carbonitrile and 1-(3-fluoro-phenyl)-piperazine , followed by hydrolysis of the nitrile intermediate. LC-MS. (M+H = 352, obsd= 352).


5 P70S6K IC₅₀: 770 nM AKT IC₅₀: 6400 nM



Example 96 5-[4-(3-chloro-phenyl)-piperazin-1-yl]-quinazoline-8-carboxylic acid amide

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling with 5-bromo-quinazoline-8-carbonitrile and 1-(3-chloro-phenyl)-piperazine , followed by hydrolysis of the nitrile intermediate. LC-MS. (M+H = 368, obsd= 368).


P70S6K IC₅₀: 10000 nM AKT IC₅₀: >100000 nM



Example 97 5-(benzylamino)quinazoline-8-carboxamide

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling with benzylamine, followed by hydrolysis of the nitrile intermediate. LC-MS (M+H = 261, obsd = 261). ¹H NMR (400 MHz, DMSO) δ 9.96 (s, 1H), 9.30 (s, 1H), 8.83 (t, J = 5.6 Hz, 1H), 8.11 (d, J = 8.6 Hz, 1H), 7.42 (d, J = 7.7 Hz, 2H), 7.35 (t, J = 7.3 Hz, 2H), 7.27 (t, J = 6.9 Hz, 1H), 6.62 (d, J = 8.7 Hz, 1H), 4.64 (d, J = 5.8 Hz, 2H).

p70S6K IC₅₀: = 690 nM AKT IC₅₀: = 57000 nM




Example 98 5-(phenethylamino)quinazoline-8-carboxamide

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling with 2-phenethylamine, followed by hydrolysis of the nitrile intermediate. LC-MS ($M+H = 293$, obsd = 293). 1H NMR (500 MHz, dmsO) δ 9.84 (s, 1H), 9.69 (d, $J = 3.9$ Hz, 1H), 9.26 (s, 1H), 8.56 (d, $J = 8.7$ Hz, 1H), 7.86 (t, $J = 5.3$ Hz, 1H), 7.52 (d, $J = 3.9$ Hz, 1H), 7.39 – 7.27 (m, 4H), 7.26 – 7.19 (m, 1H), 6.85 (d, $J = 8.8$ Hz, 1H), 3.62 – 3.52 (m, 2H), 3.04 – 2.97 (m, 2H).

p70S6K IC_{50} : = 89 nM; AKT IC_{50} : = 5600 nM

10




Example 99 5-[2-(3-fluorophenyl)ethylamino]quinazoline-8-carboxamide

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling with 2-(3-fluorophenyl)ethanamine, followed by hydrolysis of the nitrile intermediate. LC-MS ($M+H = 311$, obsd = 311). 1H NMR (500 MHz, dmsO) δ 9.83 (s, 1H), 9.69 (d, $J = 3.9$ Hz, 1H), 9.26 (s, 1H), 8.56 (d, $J = 8.7$ Hz, 1H), 7.84 (t, $J = 5.5$ Hz, 1H), 7.52 (d, $J = 3.9$ Hz, 1H), 7.38 – 7.30 (m, 1H), 7.24 – 7.18 (m, 1H), 7.16 (d, $J = 7.6$ Hz, 1H), 7.04 (td, $J = 8.6, 2.5$ Hz, 1H), 6.87 (d, $J = 8.8$ Hz, 1H), 3.59 (dd, $J = 13.9, 6.4$ Hz, 2H), 3.06 – 2.98 (m, 2H).

p70S6K IC_{50} : = 120 nM

AKT IC_{50} : = 2400 nM

20




Example 100 5-[2-(4-fluorophenyl)ethylamino]quinazoline-8-carboxamide

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling with 2-(4-fluorophenyl)ethanamine, followed by hydrolysis of the nitrile intermediate. LC-MS (M+H = 311, obsd = 311). ¹H NMR (500 MHz, dmso) δ 9.83 (s, 1H), 9.69 (d, J = 3.9 Hz, 1H), 9.26 (s, 1H), 8.55 (d, J = 8.7 Hz, 1H), 7.84 (t, J = 5.4 Hz, 1H), 7.52 (d, J = 3.8 Hz, 1H), 7.39 – 7.31 (m, 2H), 7.16 – 7.07 (m, 2H), 6.85 (d, J = 8.8 Hz, 1H), 3.55 (dd, J = 14.0, 6.4 Hz, 2H), 3.03 – 2.95 (m, 2H).

p70S6K IC₅₀: = 290 nM

AKT IC₅₀: = 4800 nM

10




Example 101 5-[2-(3,4-difluorophenyl)ethylamino]quinazoline-8-carboxamide

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling with 2-(3,4-difluorophenyl)ethanamine, followed by hydrolysis of the nitrile intermediate. LC-MS (M+H = 329, obsd = 329). ¹H NMR (500 MHz, dmso) δ 9.82 (s, 1H), 9.69 (d, J = 3.9 Hz, 1H), 9.26 (s, 1H), 8.56 (d, J = 8.7 Hz, 1H), 7.81 (t, J = 5.3 Hz, 1H), 7.53 (d, J = 3.7 Hz, 1H), 7.45 (ddd, J = 12.0, 7.9, 2.0 Hz, 1H), 7.35 (dt, J = 10.9, 8.5 Hz, 1H), 7.15 (s, 1H), 6.87 (d, J = 8.8 Hz, 1H), 3.57 (dd, J = 13.7, 6.4 Hz, 2H), 2.99 (t, J = 7.3 Hz, 2H).

p70S6K IC₅₀: = 360 nM

AKT IC₅₀: = 41000 nM

20




Example 102 5-(4-phenylbutylamino)quinazoline-8-carboxamide

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling with 4-phenylbutan-1-amine, followed by

- 5 hydrolysis of the nitrile intermediate. LC-MS ($M+H = 321$, obsd = 321). 1H NMR (500 MHz, dmsO) δ 9.86 (s, 1H), 9.68 (d, $J = 3.9$ Hz, 1H), 9.24 (s, 1H), 8.53 (d, $J = 8.7$ Hz, 1H), 7.74 (t, $J = 5.2$ Hz, 1H), 7.50 (d, $J = 3.9$ Hz, 1H), 7.27 (t, $J = 7.6$ Hz, 2H), 7.22 (d, $J = 7.0$ Hz, 2H), 7.17 (t, $J = 7.2$ Hz, 1H), 6.75 (d, $J = 8.8$ Hz, 1H), 3.34 (dd, $J = 12.1, 6.8$ Hz, 2H), 2.66 (t, $J = 7.1$ Hz, 2H), 1.79 – 1.62 (m, 4H).

10 p70S6K IC_{50} : = 87 nM AKT IC_{50} : > 100000 nM




Example 103 5-[methyl(phenethyl)amino]quinazoline-8-carboxamide

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling with *N*-methyl-2-phenylethanamine, followed by

- 15 hydrolysis of the nitrile intermediate. LC-MS ($M+H = 307$, obsd = 307). 1H NMR (500 MHz, dmsO) δ 9.71 (d, $J = 2.5$ Hz, 1H), 9.48 (s, 1H), 9.29 (s, 1H), 8.59 (d, $J = 8.4$ Hz, 1H), 7.76 (d, $J = 2.4$ Hz, 1H), 7.32 – 7.19 (m, 5H), 7.18 (M, 1H), 3.60 (t, 2H), 3.13 (s, 3H), 2.99 (t, 2H).

20 p70S6K IC_{50} : = 220 nM AKT IC_{50} : = 14000 nM




Example 104 5-[2-(3,4-dichlorophenyl)ethylamino]quinazoline-8-carboxamide

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling with 2-(3,4-dichlorophenyl)ethanamine, followed by hydrolysis of the nitrile intermediate. LC-MS ($M+H = 362$, obsd = 362). 1H NMR (500 MHz, cd_3od) δ 9.65 (s, 1H), 9.22 (s, 1H), 8.69 (d, $J = 8.8$ Hz, 1H), 7.48 (d, $J = 1.9$ Hz, 1H), 7.44 (d, $J = 8.2$ Hz, 1H), 7.22 (dd, $J = 8.3, 2.0$ Hz, 1H), 6.89 (d, $J = 8.8$ Hz, 1H), 3.71 – 3.64 (m, 2H), 3.05 (t, $J = 7.3$ Hz, 2H).

p70S6K IC_{50} : = 98 nM

AKT IC_{50} : > 100000 nM




Example 105 5-[phenethyl-[(3R)-3-piperidyl]amino]quinazoline-8-carboxamide (chiral)

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling with (*R*)-*tert*-butyl 3-(phenethylamino)piperidine-1-carboxylate (**Scheme 7**), followed by hydrolysis of the nitrile intermediate and deprotection. LC-MS ($M+H = 376$, obsd = 376). 1H NMR (500 MHz, $dmsO$) δ 10.34 (s, 1H), 8.39 (s, 1H), 7.95 (d, $J = 8.4$ Hz, 1H), 7.41 (s, 1H), 7.32 – 7.06 (m, 5H), 6.88 (d, $J = 8.7$ Hz, 1H), 5.56 (s, 1H), 3.57 – 3.47 (m, 1H), 3.26 – 3.10 (m, 2H), 2.93 – 2.74 (m, 3H), 2.68 – 2.60 (m, 1H), 2.42 – 2.33 (m, 1H), 2.07 – 1.99 (m, 1H), 1.93 – 1.75 (m, 2H), 1.74 – 1.56 (m, 2H), 1.27 – 1.19 (m, 1H).

p70S6K IC_{50} : = 11000 nM

AKT IC_{50} : > 100000 nM




Example 106 5-(3-((2,5-difluorobenzamido)methyl)pyrrolidin-1-yl)quinazoline-8-carboxamide (racemic)

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling with 2,5-difluoro-N-(pyrrolidin-3-

ylmethyl)benzamide (racemic), followed by hydrolysis of the nitrile intermediate. LC-MS (M+H = 412, obsd = 412). ¹H NMR (400 MHz, DMSO) δ 9.81 (s, 1H), 9.75 (d, J = 4.2 Hz, 1H), 9.21 (s, 1H), 8.62 (s, 1H), 8.54 (d, J = 8.8 Hz, 1H), 7.53 (s, 1H), 7.49 – 7.32 (m, 4H), 6.85 (d, J = 8.9 Hz, 1H), 3.90 – 3.82 (m, 1H), 3.82 – 3.75 (m, 2H), 3.69 – 3.61 (m, 1H), 3.41 (ddd, J = 34.0, 13.6, 7.3 Hz, 3H), 2.67 – 2.58 (m, 1H), 2.16 (dd, J = 12.1, 6.2 Hz, 1H), 1.90 – 1.79 (m, 1H).

p70S6K IC₅₀: 510 nM




Example 107 5-[3-Amino-4-(3-trifluoromethyl-phenyl)-piperidin-1-yl]-quinazoline-8-carboxylic acid amide (racemic)

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling with 5-bromo-quinazoline-8-carbonitrile and tert-butyl (4-(3-(trifluoromethyl)phenyl)piperidin-3-yl)carbamate, followed by hydrolysis of the nitrile intermediate and deprotection. LC-MS (M + H = 416, obsd = 416).

P70S6K IC₅₀: **380** nM

AKt IC₅₀: 8100 nM




Example 108 5-[4-(4-fluorophenyl)-piperazin-1-yl]-quinazoline-8-carboxylic acid amide

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling with 5-bromo-quinazoline-8-carbonitrile and 1-(4-fluorophenyl)piperazine, followed by hydrolysis of the nitrile intermediate. LC-MS. (M+H = 352, obsd= 352).

P70S6K IC₅₀: **2300** nM

AKt IC₅₀: >100000 nM




Example 109 5-[4-(3-trifluoromethylphenyl)-piperazin-1-yl]-quinazoline-8-carboxylic acid amide

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling with 5-bromo-quinazoline-8-carbonitrile and 1-(3-trifluoromethylphenyl)piperazine, followed by hydrolysis of the nitrile intermediate.

LC-MS. (M+H = 402, obsd= 402).

P70S6K IC₅₀: 11000 nM

AKt IC₅₀: 23000 nM




Example 110 5-[4-(3,4-dichlorophenyl)-piperazin-1-yl]-quinazoline-8-carboxylic acid amide

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling with 5-bromo-quinazoline-8-carbonitrile and 1-(3,4-dichlorophenyl)-piperazine, followed by hydrolysis of the nitrile intermediate. LC-MS. (M+H = 403, obsd= 403).

P70S6K IC₅₀: >100000 nM

AKt IC₅₀: >100000 nM




Example 111 5-[4-(5-chloro-2-methylphenyl)-piperazin-1-yl]-quinazoline-8-carboxylic acid amide

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling with 5-Bromo-quinazoline-8-carbonitrile and 1-(5-chloro-2-methylphenyl)piperazine, followed by hydrolysis of the nitrile intermediate. LC-MS. (M+H = 382, obsd= 382).

P70S6K IC₅₀: >100000 nM

AKt IC₅₀: >100000 nM




Example 112 5-{3-[(2,4-Difluoro-benzoylamino)-methyl]-pyrrolidin-1-yl}-quinazoline-8-carboxylic acid amide (racemic)

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling with 2,4-difluoro-N-(pyrrolidin-3-

ylmethyl)benzamide (racemic), followed by hydrolysis of the nitrile intermediate. LC-MS (M+H = 412, obsd = 412).

p70S6K IC50: 260 nM




5

Example 113 5-{3-[(3-Trifluoromethyl-benzoylamino)-methyl]-pyrrolidin-1-yl}-quinazoline-8-carboxylic acid amide (racemic)

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling with N-(pyrrolidin-3-ylmethyl)-3-

10 (trifluoromethyl)benzamide (racemic), followed by hydrolysis of the nitrile intermediate. LC-MS (M+H = 444, obsd = 444).

p70S6K IC50: nd nM



15 **Example 114**


5-{(R)-3-[(3,4-Difluoro-benzoylamino)-methyl]-pyrrolidin-1-yl}-quinazoline-8-carboxylic acid amide (pure enantiomer, absolute structure unknown)

The title compound was isolated by chiral HPLC from **Example 54**. LC-MS (M+H = 412, obsd = 412). ¹H NMR (400 MHz, DMSO) δ 9.77 (d, J = 26.9 Hz, 2H), 9.20 (s, 1H), 8.71

20 (s, 1H), 8.53 (d, J = 7.2 Hz, 1H), 7.95 – 7.83 (m, 1H), 7.70 (d, J = 13.6 Hz, 4H), 7.61 – 7.45 (m, 2H), 6.84 (d, J = 8.5 Hz, 1H), 4.30 – 3.97 (m, 4H), 3.84 (dd, J = 23.0, 14.6 Hz,

3H), 3.68 – 3.59 (m, 1H), 3.44 (s, 1H), 2.63 (dd, $J = 17.2, 8.6$ Hz, 1H), 2.14 (s, 1H), 1.82 (s, 2H), 1.63 (s, 3H), 1.26 (d, $J = 18.5$ Hz, 19H), 0.95 – 0.66 (m, 15H).

p70S6K IC₅₀: 470 nM




5

Example 115

5-[(S)-3-[(3,4-Difluoro-benzoylamino)-methyl]-pyrrolidin-1-yl]-quinazoline-8-carboxylic acid amide (pure enantiomer, absolute structure unknown)

The title compound was isolated by chiral HPLC from **Example 54**. LC-MS ($M+H = 412$, obsd = 412). ¹H NMR (400 MHz, DMSO) δ 9.82 (s, 1H), 9.75 (s, 1H), 9.21 (s, 1H), 8.79 (s, 1H), 8.54 (d, $J = 8.8$ Hz, 2H), 7.91 (s, 1H), 7.76 (s, 1H), 7.56 (d, $J = 10.0$ Hz, 2H), 6.86 (d, $J = 9.2$ Hz, 1H), 4.37 (s, 4H), 3.83 (d, $J = 28.4$ Hz, 3H), 3.64 (s, 2H), 3.45 (d, $J = 6.2$ Hz, 5H), 3.15 (d, $J = 29.4$ Hz, 2H), 2.94 (s, 1H), 2.15 (s, 2H), 1.88 (s, 1H), 1.59 (s, 2H), 1.25 (s, 4H), 1.05 (dd, $J = 18.1, 12.1$ Hz, 9H), 0.84 (s, 2H).

15 p70S6K IC₅₀: 450 nM



Example 116

5-[(S)-3-Amino-3-(4-fluoro-3-trifluoromethyl-phenylcarbamoyl)-pyrrolidin-1-yl]-quinazoline-8-carboxylic acid amide (pure enantiomer, absolute structure unknown)


The title compound was isolated by chiral HPLC from **Example 46**. LC-MS ($M+H = 463$, obsd = 463). ¹H NMR (400 MHz, DMSO) δ 9.77 (d, $J = 12.8$ Hz, 2H), 9.22 (s, 1H), 8.56

20

(d, $J = 9.0$ Hz, 1H), 8.28 (s, 1H), 8.06 (s, 1H), 7.61 – 7.46 (m, 2H), 6.90 (d, $J = 8.7$ Hz, 1H), 4.33 (d, $J = 10.7$ Hz, 1H), 4.11 (s, 2H), 3.79 (s, 1H), 3.70 (d, $J = 9.1$ Hz, 1H), 2.10 (s, 2H).

p70S6K IC₅₀: 520 nM

5



Example 117


5-[(R)-3-Amino-3-(4-fluoro-3-trifluoromethyl-phenylcarbamoyl)-pyrrolidin-1-yl]-

10 quinazoline-8-carboxylic acid amide (pure enantiomer, absolute structure unknown)

The title compound was isolated by chiral HPLC from **Example 46**. LC-MS ($M+H = 463$, obsd = 463). ¹H NMR (400 MHz, DMSO) δ 9.76 (d, $J = 12.2$ Hz, 2H), 9.22 (s, 1H), 8.56 (d, $J = 7.4$ Hz, 1H), 8.27 (s, 1H), 8.05 (s, 1H), 7.62 – 7.44 (m, 2H), 6.89 (d, $J = 8.8$ Hz, 1H), 4.32 (d, $J = 9.8$ Hz, 1H), 4.10 (d, $J = 8.5$ Hz, 1H), 3.79 (s, 1H), 3.69 (d, $J = 10.3$ Hz, 1H), 2.10 (s, 2H).

15

p70S6K IC₅₀: 380 nM




20

Example 118

5-{3-[(2,4-Difluoro-benzoylamino)-methyl]-pyrrolidin-1-yl}-quinazoline-8-carboxylic acid amide (racemic)

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling with 2-phenyl-N-(pyrrolidin-3-ylmethyl)acetamide (racemic), followed by hydrolysis of the nitrile intermediate. LC-MS (M+H = 390, obsd. = 390). ¹H NMR (400 MHz, DMSO) δ 9.81 – 9.70 (m, 2H), 9.21 (s, 1H), 8.53 (d, J = 8.9 Hz, 1H), 8.22 (t, J = 5.5 Hz, 1H), 7.54 (d, J = 3.7 Hz, 1H), 7.33 – 7.24 (m, 4H), 7.24 – 7.16 (m, 1H), 6.81 (d, J = 8.9 Hz, 1H), 3.76 (dd, J = 14.9, 8.1 Hz, 3H), 3.54 (dd, J = 10.2, 7.2 Hz, 1H), 3.43 (s, 2H), 3.21 (dtd, J = 19.7, 13.3, 6.2 Hz, 2H), 2.08 (dd, J = 11.8, 5.9 Hz, 1H), 1.83 – 1.67 (m, 1H).

p70S6K IC₅₀: 260 nM




Example 119

5-[4-(2-chloro-phenyl)-piperazin-1-yl]-quinazoline-8-carboxylic acid amide

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling with 5-Bromo-quinazoline-8-carbonitrile and 1-(2-chloro-phenyl)-piperazine, followed by hydrolysis. LC-MS. (M+H = 368, obsd= 368).

P70S6K IC₅₀: 400 nM




Example 120

5-[4-(2-methyl-phenyl)-piperazin-1-yl]-quinazoline-8-carboxylic acid amide

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling with 5-Bromo-quinazoline-8-carbonitrile and 1-(2-methyl-phenyl)-piperazine , followed by hydrolysis. LC-MS. (M+H = 348, obsd= 348). P70S6K IC₅₀: 980 nM


5



Example 121

5-[4-(2-fluorophenyl)-piperazin-1-yl]-quinazoline-8-carboxylic acid amide

10 The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling with 5-Bromo-quinazoline-8-carbonitrile and 1-(2-fluorophenyl)-piperazine , followed by hydrolysis. LC-MS. (M+H = 352, obsd= 352). P70S6K IC₅₀: 12000 nM



15

Example 122

5-[4-(2-methoxyphenyl)-piperazin-1-yl]-quinazoline-8-carboxylic acid amide

The title compound was synthesized according to the procedure described for the preparation of **Example 1** by coupling with 5-Bromo-quinazoline-8-carbonitrile and 1-(2-methoxyphenyl)-piperazine , followed by hydrolysis. LC-MS. (M+H = 364, obsd= 364). P70S6K IC₅₀: 11000 nM

20

Biological Activity

5 **P70S6K enzyme assay**

P70S6K inhibitor compounds are diluted and plated in 96 well plates. A reaction mixture including the following components is then added to the compound plate to initiate the enzyme reaction; P70S6K (3 nM, T412E mutant, Millipore) is mixed with 24 μ M ATP in an assay buffer containing 100 mM Hepes (pH 7.5), 5 mM MgCl₂, 1mM DTT, 0.015% Brij and 1 μ M of the substrate peptide FITC-AHA-AKRRRLSSLRA-OH (derived from the S6 ribosomal protein sequence, FITC = fluorescein isothiocyanate, AHA = 6-aminohexanoic acid). The reaction is incubated for 90 min at 25° C, before the addition of 10 mM EDTA to stop the reaction. The proportion of substrate and product (phosphorylated) peptide is analysed on a Caliper Life Sciences Lab Chip 3000, using a pressure of - 1.4 psi, and upstream and downstream voltages of - 3000 and - 700 respectively. Product peaks are resolved before substrate peaks on the resulting chromatograms.

20

AKT enzyme assay

A TTP Mosquito liquid handling instrument is used to place 125nl of the appropriate concentration of inhibitor in 100% DMSO (for a dose response curve calculation) into each well of a 384-well plate. To this reaction components are added to a final volume of 12.5 μ l:

0.1 ng/ μ l His-AKT (Full Length), (Invitrogen, Part # P2999, Lot # 641228C).
160 μ M ATP (Fluka, 02055)
1 mM DTT (Sigma, D0632)
1 mM MgCl₂ (Sigma, M1028)
1 μ M substrate peptide (sequence FITC-AHA-GRPRTSSFAEG-NH₂), synthesized by Tufts Peptide Synthesis service.
100 mM HEPES pH 7.5 (Calbiochem, 391338)
0.015% Brij-35 (Sigma, B4184)

35

The reaction is incubated for 90 min at 25 C, and then stopped by the addition of 70 μ l of Stop buffer (100 mM HEPES pH 7.5, 0.015% Brij-35, 10 mM EDTA (Sigma, E7889)).


5 The plate is read on a Caliper LC 3000 in an Off-Chip mobility shift assay format, using the following parameters for a 12-sipper chip: screening pressure -2.3 psi, upstream voltage -500, and downstream voltage -3000. These conditions cause unphosphorylated substrate and phosphorylated product peptide to resolve as separate peaks allowing direct measurement of percentage of conversion of substrate to product. The percent conversion can be plotted against concentration of inhibitor to produce a sigmoidal dose response
10 curve, from which an IC₅₀ can be calculated.

Claims

We claim:

1. A compound of Formula (I),

5



(I)

or its stereoisomers or tautomers, or pharmaceutically acceptable salts of each of the foregoing, including mixtures thereof in all ratios, wherein

X is N;

10 Y is N-R⁵, O, or absent;

R¹ is L¹-R⁶, or L¹-R⁶-L²-R⁷;

R² is H, Hal, OH, OA, CN, NH₂, or NHA;

R³ is H, CH₃, or C(Hal)₃;

R⁴ is H, Hal, OH, COOH, NH₂, or CN;

15 R⁵ is H, LA or monocyclic alkyl having 3, 4, 5, 6, or 7 ring atoms, wherein one or two CH₂ groups may be replaced by a -NH- group, or

R⁵, R¹ together form a monocyclic alkyl ring having 3, 4, 5, 6, or 7 ring atoms, wherein one or two CH₂ groups may be replaced by an O atom and/or an -NH-, -NA-, -N(L¹-R⁶)-, -CHA-, -CA₂-, CH(L¹-R⁶)- or -CO- group, and wherein monocyclic alkyl ring may be substituted by NH₂;

20

L¹, L² are each independently a single bond, or unbranched or branched alkyl having 1, 2, 3, 4 or 5 C atoms, which may be unsubstituted, or mono- or disubstituted with Hal, OH, NH₂, NH(LA), N(LA)₂, and wherein one or two CH₂ groups may be replaced by an O atom or by a -CO-, -NH-, -N(LA)-, -CONH-, -N(LA)COO- or -NHCO- group;

25

R⁶, R⁷ are each independently Ar or a monocyclic alkyl ring having 3, 4, 5, 6, or 7 ring atoms, wherein one or two CH₂ groups may be replaced by an O atom and/or an –NH-, -NA-, -CHA-, -CO- or -CONHA- group;

each Ar is independently a mono- or bicyclic aromatic homo- or heterocycle having 0, 1, 2, 3 or 4 N, O and/or S atoms and 5, 6, 8, 9, or 10 skeleton atoms, which may be unsubstituted or, independently of one another, mono-, di- or trisubstituted by Hal, A, OA, OH, NH₂, or NHA;

each A is independently an unbranched or branched linear or cyclic alkyl having 1, 2, 3, 4, 5 or 6 C atoms, wherein one or two CH₂ groups may be replaced by an O atom and/or an –NH-, -NHCOAr or –CONHAr group wherein 1-3 H atoms may be replaced by Hal, and wherein one or two CH₃ groups may be replaced by NH₂, OH, NH(LA) or N(LA)₂ group;

each LA is independently an unbranched or branched, linear alkyl having 1, 2, 3, or 4 C atoms wherein 1, 2 or 3 H atoms may be replaced by Hal; and

each Hal is independently F, Cl, Br or I.

2. The compound according to Claim 1, or its stereoisomers or tautomers, or pharmaceutically acceptable salts of each of the foregoing, including mixtures thereof in all ratios, X is N and Y is N-R⁵.

3. The compound according to claim 1. or its stereoisomers or tautomers, or pharmaceutically acceptable salts of each of the foregoing, including mixtures thereof in all ratios, wherein X is N; Y is N-R⁵; R⁵ and R¹ together form a monocyclic alkyl ring having 3, 4, 5, 6, or 7 ring atoms, wherein one or two CH₂ groups may be replaced by an –NH-, -NA-, -N(L¹-R⁶)-, -CHA-, -CA₂-, -CH(L¹-R⁶)-, or –CO- group, and wherein the monocyclic alkyl ring is optionally substituted by NH₂.

4. The compound according to Claim 1, wherein the compound is selected from the group consisting of:

5-((4-(4-(trifluoromethoxy)phenyl)pyrrolidin-3-yl)amino)quinazoline-8-carboxamide (trans_racemic),

5-((4-phenylpiperidin-3-yl)amino)quinazoline-8-carboxamide (trans_racemic),

5-((4-(3-fluoro-4-(trifluoromethyl)phenyl)piperidin-3-yl)amino)quinazoline-8-carboxamide(trans_racemic),

5-((-4-(3-(trifluoromethoxy)phenyl)pyrrolidin-3-yl)amino)quinoline-8-carboxamide
(trans_racemic)

5-((4-(3-(trifluoromethyl)phenyl)pyrrolidin-3-yl)amino)quinazoline-8-carboxamide,

5-((4-(4-chloro-3-fluorophenyl)piperidin-3-yl)amino)quinazoline-8-carboxamide,

5 5-((4-(3-chloro-4-fluorophenyl)piperidin-3-yl)amino)quinazoline-8-carboxamide,

5-(((3R,4R)-4-(3-chloro-4-fluorophenyl)piperidin-3-yl)amino)quinazoline-8-carboxamide,

and

5-((4-(4-fluorophenyl)piperidin-3-yl)amino)quinazoline-8-carboxamide,

or its stereoisomers or tautomers, or pharmaceutically acceptable salts of each of the

10 foregoing, including mixtures thereof in all ratios.



5. A pharmaceutical composition comprising a compound of claim 1, and/or its
stereoisomers or tautomers, or pharmaceutically acceptable salts of each of the
foregoing, including mixtures thereof in all ratios, as active ingredient, together with a
15 pharmaceutically acceptable carrier.

6. A kit consisting of separate packs of an effective amount of a compound of claim 1
and/or its stereoisomers or tautomers, or pharmaceutically acceptable salts of each of
the foregoing, including mixtures thereof in all ratios, and an effective amount of a further
20 medicament active ingredient.

7. The compound according to claim 1, or its stereoisomers or tautomers, or
pharmaceutically acceptable salts of each of the foregoing, including mixtures thereof in
all ratios, wherein, X is N; Y is N-R⁵; and R⁵ and R¹ together form a monocyclic alkyl ring
25 having 4, 5 or 6 ring atoms, wherein one CH₂ group is replaced by an -N(L¹-R⁶)- group,
and wherein the monocyclic alkyl ring is optionally substituted by NH₂.

8. The compound according to claim 1, or its stereoisomers or tautomers, or
pharmaceutically acceptable salts of each of the foregoing, including mixtures thereof in
30 all ratios, wherein, X is N; Y is N-R⁵; R⁵ and R¹ together form a monocyclic alkyl ring
having 4, 5 or 6 ring atoms, wherein one CH₂ group is replaced by an -N(L¹-R⁶)- group,
and wherein the monocyclic alkyl ring is optionally substituted by NH₂; and L¹ is a bond,
-CONH-, -NHCO-, -CONHCH₂-, or CH₂CONH-.

9. The compound according to claim 1, or its stereoisomers or tautomers, or pharmaceutically acceptable salts of each of the foregoing, including mixtures thereof in all ratios, wherein, X is N; Y is N-R⁵; R⁵ and R¹ together form a monocyclic alkyl ring having 4, 5 or 6 ring atoms, wherein one CH₂ group is replaced by an -N(L¹-R⁶)- group, and wherein the monocyclic alkyl is optionally substituted by NH₂; and R⁶ is phenyl, which is unsubstituted, or independently mono-, di- or trisubstituted by Hal, C(Hal)₃, CH₃, or C(Hal)₃O.
10. The compound according to claim 1, or its stereoisomers or tautomers, or pharmaceutically acceptable salts of each of the foregoing, including mixtures thereof in all ratios, wherein, X is N; Y is N-R⁵; R⁵ and R¹ together form a monocyclic alkyl ring having 4, 5 or 6 ring atoms, wherein one CH₂ group is replaced by an -N(L¹-R⁶)- group, and wherein the monocyclic alkyl ring is optionally substituted by NH₂; L¹ is a bond, -CONH-, -NHCO-, -CONHCH₂-, or CH₂CONH-; and R⁶ is phenyl, which is unsubstituted, or independently mono-, di- or trisubstituted by Hal, C(Hal)₃, CH₃, CH₃O, or C(Hal)₃O.
11. The compound according to claim 1, or its stereoisomers or tautomers, or pharmaceutically acceptable salts of each of the foregoing, including mixtures thereof in all ratios, wherein, X is N; Y is NH; R¹ is L¹-R⁶-L²-R⁷; L¹ and L² are a bond; R⁶ is a monocyclic alkyl ring having 4, 5 or 6 ring atoms, wherein one CH₂ group is replaced by an -NH-, group; and R⁷ is phenyl, which is unsubstituted, or independently mono-, di- or trisubstituted by Hal, C(Hal)₃, CH₃, CH₃O, or C(Hal)₃O.
12. The compound according to claim 1, or its stereoisomers or tautomers, or pharmaceutically acceptable salts of each of the foregoing, including mixtures thereof in all ratios, wherein, X is N; Y is NH; R¹ is L¹-R⁶-L²-R⁷; L¹ and L² are a bond; R⁶ is piperidyl or pyrrolidinyl; and R⁷ is phenyl, which is mono- or disubstituted by Hal, C(Hal)₃, CH₃, CH₃O, or C(Hal)₃O.
13. The compound according to claim 1, or its stereoisomers or tautomers, or pharmaceutically acceptable salts of each of the foregoing, including mixtures thereof in all ratios, wherein, X is N; Y is NH; R¹ is L¹-R⁶-L²-R⁷; L¹ and L² are a bond; R⁶ is


 or
 
 , and R⁷ is phenyl, which is mono- or disubstituted by Hal, C(Hal)₃, CH₃, CH₃O, or C(Hal)₃O.