

Claims

1. A method for assessing the risk for development of a cardiovascular disorder in a subject, comprising:

- 5 (a) determining in a sample from said subject
 - (i) the amount of a first marker Trefoil Factor 3, alpha-2-Macroglobulin, and alpha-Glutathione-S-Transferase; and
 - (ii) the amount of a further marker, wherein said further marker is Nt-pro BNP, Peroxiredoxin-4, Osteoprotegerin, Apolipoprotein B, and Angiopoietin-2; and
- 10 (b) correlating that said subject is at risk for development of a cardiovascular disorder when said amount is altered compared to a reference amount for the at least first marker.

2. A method for assessing the risk for development of a cardiovascular disorder in a subject, comprising:

- 15 (a) determining in a sample from said subject
 - (i) the amount of a first marker Trefoil Factor 3, alpha-2-Macroglobulin, alpha-Glutathione-S-Transferase and Hepatocyte Growth Factor Receptor; and
 - (ii) the amount of a further marker Nt-pro BNP, Angiopoietin-2, Apolipoprotein B, Osteoprotegerin, Growth-Factor-Differentiation-Factor-15 and Chromogranin A; and
- 20 (b) correlating that said subject is at risk for development of a cardiovascular disorder when said amount is altered compared to a reference amount for the at least first marker.