# Combination Therapies for the Treatment of Hepatocellular Carcinoma

## CROSS-REFERENCE TO RELATED APPLICATIONS


This application claims the benefit of priority to United States Provisional Patent Application No. 62/378,455, filed on August 23, 2016. That application is incorporated by reference as if fully rewritten herein.

## BACKGROUND

Liver cancer is the second greatest cause of mortality from any type of cancer, and the 16th most common cause of death worldwide (Llovet JM, *et al.*, 2015 "Advances in targeted therapies for hepatocellular carcinoma in the genomic era." *Nat. Rev. Clin Oncology* 12, 408–424). Hepatocellular carcinoma (HCC) accounts for up to 90% of all primary liver cancers (Llovet JM *et al.* 2015).

Various signaling pathways have been implicated in HCC, including fibroblast growth factors (FGF) (particularly FGF19/FGFR4), epidermal growth factor (EGF), vascular endothelial growth factor (VEGF), ERK/MAPK, and mechanistic target of rapamycin (mTOR), among others (Llovet JM *et al.*, 2015). FGF19 is overexpressed in about a third of all HCC, and this overexpression is hypothesized to hyperactivate FGFR4 and its downstream signaling pathway leading to enhanced tumor growth (Xie MH *et al.*, 1999 "FGF-19, a novel fibroblast growth factor with unique specificity for FGFR4." *Cytokine*. 1999 Oct;11(10):729-35; Sawey, *et al.*, 2011 "Identification of a therapeutic strategy targeting amplified FGF19 in liver cancer by Oncogenomic screening." *Cancer Cell.* 2011 Mar 8;19(3):347-58.). Similar to FGF19/FGFR4 pathway, the CDK4/6 pathway activation is also involved in HCC pathogenesis (Rivadeneira *et al.*, 2010 "Proliferative Suppression by CDK4/6 Inhibition: Complex Function of the Retinoblastoma Pathway in Liver Tissue and Hepatoma Cells." *Gastroenterology* 138:1920–1930).


Compound 1 is a selective, orally bioavailable small molecule FGFR4 inhibitor with the structure shown in Formula I, and the chemical name *N*-(2-((6-(3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-methylureido)pyrimidin-4-yl)amino)-5-(4-ethylpiperazin-1-yl)phenyl)acrylamide:



(I).


Compound 1 and its synthesis are reported in PCT International Application Publication No. WO2015/057938, published on April 23, 2015. That document is incorporated by reference herein.

Palbociclib (6-acetyl-8-cyclopentyl-5-methyl-2-{[5-(piperazin-1-yl)pyridin-2-yl]amino}pyrido[2,3-*d*]pyrimidin-7(8*H*)-one) is an FDA-approved inhibitor of cyclin-dependent kinase (CDK) 4 and 6. Palbociclib has the following structure:




See U.S. Patent Nos. 6,936,612; 7,208,489, and 7,456,168, which are incorporated by reference herein.

Ribociclib (7-Cyclopentyl-2-(5-piperazin-1-yl-pyridin-2-ylamino)-7H-pyrrolo[2,3-d]pyrimidine-6-carboxylic acid dimethylamide) is an FDA-approved inhibitor of cyclin-dependent kinase (CDK) 4 and 6. Ribociclib has the following structure:



See U.S. Patent App. Pub. No. US20120115878, PCT Publication No. WO2007140222, PCT Publication No. WO2012061156; PCT Publication No. WO2011130232; PCT Publication No. WO2011101417; and PCT Publication No. WO2010020675, all of which are incorporated by reference herein.


Abemaciclib is an inhibitor of CDK 4/6 with the name *N*-(5-((4-ethylpiperazin-1-yl)methyl)pyridin-2-yl)-5-fluoro-4-(4-fluoro-1-isopropyl-2-methyl-*1H*-benzo[d]imidazol-6-yl)pyrimidin-2-amine. Abemaciclib has the following structure:



See O'Leary, *et al.*, "Treating Cancer with Selective CDK 4/6 Inhibitors" *Nat. Rev.* (Published Online March. 31, 2016); PCT Publication No. WO2016110224, United States Patent App. Pub. No. 20100160340; and PCT Publication No. WO2016025650, all of which are incorporated by reference herein.


G1T-38 (also referred to as GZ-38-1 or G1T38-1) is a reported inhibitor of CDK 4/6. G1T-38, which is studied by G1 Therapeutics, Inc., of Research Triangle Park, North Carolina, is reported in Abstract #2824 of the 2016 AACR Annual Meeting, held April 16-20 in New Orleans, Louisiana, entitled "G1T38, A Novel, Oral, Potent and Selective CDK 4/6 Inhibitor for the Treatment of RB Competent Tumors," by J. Sorrentino, J. Bisi, P. Roberts, and J. Strum. that document is incorporated by reference herein. G1T38 has the chemical

name 2'-((5-(4-isopropylpiperazin-1-yl)pyridin-2-yl)amino)-7', 8'dihydro-6'Hspiro[cyclohexane1,9' pyrazino[1',2':1,5] pyrrolo[2,3-d]pyrimidin]-6'-one dihydrochloride, and the structure set forth below:




See Bisi, *et al.*, "Preclinical development of G1T38: A novel, potent and selective inhibitor of cyclin dependent kinases 4/6 for use as an oral antineoplastic in patients with CDK4/6 sensitive tumors," *Oncotarget*, Advance Publications 2017 (March 15, 2017), incorporated by reference herein.

G1T-28 is an inhibitor of CDK 4/6 with the name 2'-((5-(4-methylpiperazin-1-yl)pyridin-2-yl)amino)-7',8'-dihydro-6'H-spiro[cyclohexane-1,9'-pyrazino[1',2':1,5]pyrrolo[2,3-d]pyrimidin]-6'-one. G1T-28 has the following structure:




See, for example, Bisi, *et al.*, "Preclinical Characterization of G1T28: A Novel CDK4/6 Inhibitor for Reduction of Chemotherapy-induced Myelosuppression" *Mol. Cancer Ther.*; 15(5) 783-93, May 2016; U.S. Patent Application Publication No. US20160220569; PCT International Patent Application Publication Nos. WO2014144326; WO2014144847; and WO2016040848, all of which are incorporated by reference herein.

AT-7519 is an inhibitor of CDK 4/6 with the name N-(4-piperidinyl)-4-(2,6-dichlorobenzoylamino)-1*H*-pyrazole-3 carboxamide. AT-7519 has the following structure:




See, for example, PCT International Patent Application Publication Nos. WO 2005012256; WO 2006077424; WO 2006077426; WO 2008001101; WO 2006077425; WO 2006077428; WO 2008007113; WO 2008007122; and WO 2008009954, which are incorporated by reference herein.

FLX-925 (also known as AMG-925) is an inhibitor of CDK 4/6 with the name 2-Hydroxy-1-[2-[[9-(trans-4-methylcyclohexyl)-9*H*-pyrido[4',3':4,5]pyrrolo[2,3-d]pyrimidin-2-yl]amino]-7,8-dihydro-1,6-naphthyridin-6(5*H*)-yl]ethanone. FLX-925 has the following structure:



See, for example, U.S. Patent Application Pub. No. 2014163052 and PCT International Patent Application Publication No. WO 2012129344, both of which are incorporated by reference herein.

Alvocidib is an inhibitor of CDK 4/6 with the name 2-(2-chlorophenyl)-5,7-dihydroxy-8-((3*S*,4*R*)-3-hydroxy-1-methylpiperidin-4-yl)-4*H*-chromen-4-one. Alvocidib has the following structure:




.

See, for example, U.S. Patent Application Publication No. US2011189175 and US2011189175; PCT International Patent Application Publication Nos. WO 2000044362; WO 2001041747; WO 2001053293; WO 2001053294; WO 2002022133; WO 2007010946, all of which are incorporated by reference herein.

Despite advances in the treatment of HCC, there is a need to provide improved treatment for HCC. Despite advances in the treatment of IHCC, there is a need to provide improved treatment for IHCC.

## SUMMARY

In one aspect, the present invention refers to a pharmaceutical formulation comprising Compound 1 or a pharmaceutically acceptable salt thereof and a CDK 4/6 inhibitor or a pharmaceutically acceptable salt thereof. In a second aspect, the present invention refers to use of N-(2-((6-(3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-methylureido)pyrimidin-4-yl)amino)-5-(4-ethylpiperazin-1-yl)phenyl)acrylamide or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for the treatment of hepatocellular carcinoma, wherein N-(2-((6-(3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-methylureido)pyrimidin-4-yl)amino)-5-(4-ethylpiperazin-1-yl)phenyl)acrylamide or a pharmaceutically acceptable salt thereof is to be administered simultaneously or sequentially with a CDK 4/6 inhibitor or a pharmaceutically acceptable salt thereof to a patient in need thereof.

Examples provide a combination therapy, comprising an effective amount of Compound 1 and an effective amount of a CDK4/6 inhibitor. In certain examples the CDK 4/6 inhibitor is palbociclib. In other examples the CDK 4/6 inhibitor is ribociclib. In other examples the CDK 4/6 inhibitor is abemaciclib. In other examples the CDK 4/6 inhibitor is G1T38. In other examples the CDK 4/6 inhibitor is G1T28. Combination therapy provided herein may lead to an enhanced reduction in the viability of HCC cells and may lead to tumor growth inhibition of HCC in patients in need of treatment. Combination therapy provided

herein may also be effective in treatment of hepatic cholangiocarcinoma, including for example intrahepatic cholangiocarcinoma (IHCC).

Examples may provide a method of treating hepatocellular carcinoma in a patient in need thereof, including administering to the patient combination of *N*-(2-((6-(3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-methylureido)pyrimidin-4-yl)amino)-5-(4-ethylpiperazin-1-yl)phenyl)acrylamide or a pharmaceutically acceptable salt thereof and a CDK 4/6 inhibitor or a pharmaceutically acceptable salt thereof. In some examples the *N*-(2-((6-(3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-methylureido)pyrimidin-4-yl)amino)-5-(4-ethylpiperazin-1-yl)phenyl)acrylamide or a pharmaceutically acceptable salt thereof is administered in a daily dosage between 50 mg to 600 mg. In some examples the *N*-(2-((6-(3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-methylureido)pyrimidin-4-yl)amino)-5-(4-ethylpiperazin-1-yl)phenyl)acrylamide or a pharmaceutically acceptable salt thereof is administered in a daily dosage between 200 mg to 400 mg. In some examples the *N*-(2-((6-(3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-methylureido)pyrimidin-4-yl)amino)-5-(4-ethylpiperazin-1-yl)phenyl)acrylamide or a pharmaceutically acceptable salt thereof is administered in a daily dosage of 300 mg.

In some examples the CDK 4/6 inhibitor is selected from, for example, 6-acetyl-8-cyclopentyl-5-methyl-2-{[5-(piperazin-1-yl)pyridin-2yl]amino}pyrido[2,3-*d*]pyrimidin-7(8*H*)-one (palbociclib); 7-Cyclopentyl-2-(5-piperazin-1-yl-pyridin-2-ylamino)-7H-pyrrolo[2,3-d]pyrimidine-6-carboxylic acid dimethylamide (ribociclib); and *N*-(5-((4-ethylpiperazin-1-yl)methyl)pyridin-2-yl)-5-fluoro-4-(4-fluoro-1-isopropyl-2-methyl-*1H*-benzo[d]imidazol-6-yl)pyrimidin-2-amine (abemaciclib).

In some examples the CDK 4/6 inhibitor is palbociclib. Palbociclib may be administered, for example in a dosage of 75, 100, or 125 mg/day. Typically a dosage is administered orally as a single capsule for 21 consecutive days followed by a 7 day off-treatment period

In some examples the CDK 4/6 inhibitor is ribociclib. Ribociclib may be administered, for example, in a dosage of 200, 400, or 600 mg/day. Typically ribociclib is administered orally as 200 mg capsules or tables, for 21 consecutive days, followed by a 7 day off-treatment period.

In some examples the CDK 4/6 inhibitor is abemaciclib. Abemaciclib may be administered, for example, in a dosage of 200, 300, or 400 mg/day. Typically abemaciclib is administered twice-daily in dosages of 100, 150, or 200 mg/dose. Abemaciclib is typically administered for 21 consecutive days or 28 consecutive days, followed by a 7 day off-

treatment period.

In some examples the CDK 4/6 inhibitor is G1T-38. G1T-38 may be administered, for example, in dosages of 10, 50, or 100 mg/kg. In some examples the CDK 4/6 inhibitor is G1T-28. G1T-28 may be administered, for example, in dosages of between 190 and 200 $mg/m^2$.

In some examples the CDK 4/6 inhibitor is AT-7519. AT-7519 may be administered, for example, in dosages of 14.4 to 32.4 $mg/m^2$. AT-7519 may be dosed every three weeks, with drug given on days 1, 4, 8, and 11. In one example the dose is 27 $mg/m^2$, given at the above frequencies.

In some examples the CDK 4/6 inhibitor is FLX-925. In some examples the CDK 4/6 inhibitor is alvocidib. Alvocidib may be administered, for example, in amounts between 8 and 122 mg/m2. Alvocidib may be administered as a 72 hour infusion. Maximum tolerated dosages of aovocidib have been reported as 40, 50, or 78 $mg/m^2$.

In some examples the *N*-(2-((6-(3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-methylureido)pyrimidin-4-yl)amino)-5-(4-ethylpiperazin-1-yl)phenyl)acrylamide or a pharmaceutically acceptable salt thereof and the CDK 4/6 inhibitor or a pharmaceutically acceptable salt thereof are administered as separate formulations. Typically the time between administration of each formulation does not exceed 12 hours. In some examples the *N*-(2-((6-(3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-methylureido)pyrimidin-4-yl)amino)-5-(4-ethylpiperazin-1-yl)phenyl)acrylamide or a pharmaceutically acceptable salt thereof and the CDK 4/6 inhibitor or a pharmaceutically acceptable salt thereof are administered as a single formulation. In some examples the *N*-(2-((6-(3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-methylureido)pyrimidin-4-yl)amino)-5-(4-ethylpiperazin-1-yl)phenyl)acrylamide or a pharmaceutically acceptable salt thereof and the CDK 4/6 inhibitor or a pharmaceutically acceptable salt thereof are administered sequentially with other treatments. In some examples the *N*-(2-((6-(3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-methylureido)pyrimidin-4-yl)amino)-5-(4-ethylpiperazin-1-yl)phenyl)acrylamide or a pharmaceutically acceptable salt thereof and the CDK 4/6 inhibitor or a pharmaceutically acceptable salt thereof are administered simultaneously.

In some examples the form of *N*-(2-((6-(3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-methylureido)pyrimidin-4-yl)amino)-5-(4-ethylpiperazin-1-yl)phenyl)acrylamide that is administered is the free base form. In some examples the form of *N*-(2-((6-(3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-methylureido)pyrimidin-4-yl)amino)-5-(4-ethylpiperazin-1-yl)phenyl)acrylamide that is administered is a hydrochloride salt form.

Further examples may provide a pharmaceutical formulation including *N*-(2-((6-(3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-methylureido)pyrimidin-4-yl)amino)-5-(4-ethylpiperazin-1-yl)phenyl)acrylamide or a pharmaceutically acceptable salt thereof and a CDK 4/6 inhibitor or a pharmaceutically acceptable salt thereof. In some examples the *N*-(2-((6-(3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-methylureido)pyrimidin-4-yl)amino)-5-(4-ethylpiperazin-1-yl)phenyl)acrylamide is a free base form. In some examples the *N*-(2-((6-(3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-methylureido)pyrimidin-4-yl)amino)-5-(4-ethylpiperazin-1-yl)phenyl)acrylamide is a hydrochloride salt form.

Further examples may provide use of a combination of *N*-(2-((6-(3-(2,6-dichloro-3,5-

dimethoxyphenyl)-1-methylureido)pyrimidin-4-yl)amino)-5-(4-ethylpiperazin-1-yl)phenyl)acrylamide or a pharmaceutically acceptable salt thereof and a CDK 4/6 inhibitor in the treatment of hepatocellular carcinoma. Further examples may provide use of a combination of *N*-(2-((6-(3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-methylureido)pyrimidin-4-yl)amino)-5-(4-ethylpiperazin-1-yl)phenyl)acrylamide or a pharmaceutically acceptable salt thereof and a CDK 4/6 inhibitor in the preparation of a medicament for treatment of hepatocellular carcinoma.

## BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1A through FIG. 1C show antitumor effects of Compound 1 and CDK 4/6 inhibitor ribociclib in the JHH7 xenograft model of hepatocellular carcinoma. Both Compound 1 and ribociclib, each as a free base,were given orally (PO) once daily (QD) for 8 days. Data represent the mean ±SEM for tumor volume.

FIG. 1A shows Ribociclib as single agent (****$P \leq 0.0001$ compared to vehicle controls using two way ANOVA followed by Sidak post hoc test).

FIG. 1B shows two dose levels of Ribociclib in combination with 300mg/kg Compound 1 (****$P \leq 0.0001$ compared to 300mg/kg Compound 1 single agent group using two way ANOVA followed by Sidak post hoc test).

FIG. 1C shows two dose levels of Ribociclib in combination with 500mg/kg Compound 1 (****$P \leq 0.0001$ for 500mg/kg Compound 1 single agent group in comparison to the vehicle controls and ****$P \leq 0.0001$ for combinations compared to 500mg/kg Compound 1 single agent group using two way ANOVA followed by Sidak post hoc test).

FIG. 2A through FIG. 2C show antitumor effects of Compound 1 and CDK 4/6 inhibitor palbociclib in the JHH xenograft model of hepatocellular carcinoma. Both Compound 1 and Palbociclib, each as a free base, were given orally (PO) once daily (QD) for 8 days. Data represent the mean ±SEM for Tumor Volume.

FIG. 2A shows palbociclib as single agent (*$P \leq 0.05$ compared to vehicle controls using two way ANOVA followed by Sidak post hoc test).

FIG. 2B shows two dose levels of palbociclib in combination with 300mg/kg Compound 1 (**$P \leq 0.01$ compared to vehicle control and ****$P \leq 0.0001$ compared to

300mg/kg Compound 1 single agent group using two way ANOVA followed by Sidak post hoc test).

FIG. 2C shows two dose levels of palbociclib in combination with 500mg/kg Compound 1 (****P ≤ 0.0001 for 500mg/kg Compound 1 single agent group in comparison to the vehicle controls and ****P ≤ 0.0001 for combinations compared to 500mg/kg Compound 1 single agent group using two way ANOVA followed by Sidak post hoc test).

FIG. 3A and FIG. 3B show antitumour effects of Compound 1 and CDK 4/6 inhibitor palbociclib in the LIX066 patient derived xenograft model of hepatocellular carcinoma. Both Compound 1 and Palbociclib were given orally (PO) once daily (QD) for 8 days. Data represent the mean ±SEM for Tumor Volume.

FIG. 3A shows Compound 1 300mg/kg either as single agent or in combination with 100mg/kg palbociclib (***P ≤ 0.001 for palbociclib 100mg/kg single agent compared to vehicle controls, ****P ≤ 0.0001 for 300mg/kg Compound 1 single agent group in comparison to the vehicle controls and ****P ≤ 0.0001 for combinations compared to 300mg/kg Compound 1 single agent group using two way ANOVA followed by Sidak post hoc test).

FIG. 3B shows Compound 1 500mg/kg either as single agent or in combination with 100mg/kg palbociclib (***P ≤ 0.001 for palbociclib 100mg/kg single agent compared to vehicle controls, ****P ≤ 0.0001 for 500mg/kg Compound 1 single agent group in comparison to the vehicle controls and ****P ≤ 0.0001 for combinations compared to 500mg/kg Compound 1 single agent group using two way ANOVA followed by Sidak post hoc test).


## DETAILED DESCRIPTION OF EMBODIMENTS

Provided herein are combination therapies useful in treating hepatocellular carcinoma (HCC) and intrahepatic cholangiocarcinoma (IHCC). In some embodiments, the combination therapies include administration of Compound 1 in combination with a CDK 4/6 inhibitor. In certain embodiments the CDK 4/6 inhibitor is palbociclib. In other embodiments, the CDK 4/6 inhibitor is ribociclib. In still other embodiments the CDK 4/6 inhibitor is abemaciclib.

Provided herein are combinations of therapeutic agents and methods for administration of the combination of agents to treat hepatocellular carcinoma. As used herein, a "combination of therapeutic agents" and similar terms refer to a combination of two types

of therapeutic agents: (1) Compound 1 and/or pharmacologically active salts thereof and (2) a CDK 4/6 inhibitor, and/or pharmacologically active salts thereof. "Combination" as used herein (including in the term "combination of therapeutic agents") refers to these types of therapeutic agents co-formulated in a single dosage form, individually formulated and co-administered, or individually formulated and sequentially administered.

Compound 1 is a selective, orally bioavailable small molecule FGFR4 inhibitor with the structure shown in Formula I:




(I).

Compound 1 and its synthesis are reported in PCT International Application Publication No. WO2015/057938, published on April 23, 2015. That document is incorporated by reference herein. Compound 1 may also be used alone or in combinations described herein as treatment for HCC or cholangiocarcinoma, including intrahepatic cholangiocarcinoma (IHCC). When used alone or in combinations as described herein, Compound 1 may be administered to patients in any of the following daily dosage amounts: 150 mg, 300 mg, 600 mg, 1000 mg, 1500 mg or 2000 mg. The daily dosage amount may be from 50 mg to 3000 mg, from 50 mg to 600 mg, or from 200 mg to 400 mg. The daily dosage may be part of a cyclic regimen lasting 14 days or 21 days. The daily dosage amount may be administered as a single dosage or as multiple dosages.

CDK 4/6 inhibitors suitable for use herein may include, for example, ribociclib, palbociclib, and abemaciclib, and their pharmaceutically acceptable salts and hydrates.

Administration of a combination of therapeutic agents comprises administration of the individual therapeutic agents in combination in a single formulation or unit dosage form, administration of the individual therapeutic agents of the combination concurrently but separately, or administration of the individual agents of the combination sequentially by any suitable route. The dosage of the individual therapeutic agents of the combination may require more frequent administration of one of the agents as compared to the other agent in the combination. Therefore, to permit appropriate dosing, packaged pharmaceutical products may contain one or more dosage forms that contain the combination of agents, and one or

more dosage forms that contain one of the combinations of agents, but not the other agent(s) of the combination.

Combinations as reported herein may include embodiments wherein one or more of Compound 1 and a CDK 4/6 inhibitor are administered as a pharmaceutically acceptable salt or as a free base. There is no requirement that both compounds be administered as the same pharmaceutically acceptable salt, but they may be. In particular embodiments combinations comprise a free base form of Compound 1 and a free base form of CDK 4/6 inhibitor. In other embodiments combinations comprise an HCl form of Compound 1 and a CDK 4/6 inhibitor. In some embodiments the CDK 4/6 inhibitor may be a free base. In some embodiments the CDK 4/6 inhibitor may be a pharmaceutically acceptable salt. In some embodiments the CDK 4/6 inhibitor may be a hydrate.

"Pharmaceutically acceptable salt" as used herein refers to acid addition salts or base addition salts of the compounds in the present disclosure. A pharmaceutically acceptable salt is any salt which retains the activity of the parent compound and does not impart any unduly deleterious or undesirable effect on a subject to whom it is administered and in the context in which it is administered. Pharmaceutically acceptable salts include, but are not limited to, metal complexes and salts of both inorganic and carboxylic acids. Pharmaceutically acceptable salts also include metal salts such as aluminum, calcium, iron, magnesium, manganese and complex salts. In addition, pharmaceutically acceptable salts include, but are not limited to, acid salts such as acetic, aspartic, alkylsulfonic, arylsulfonic, axetil, benzenesulfonic, benzoic, bicarbonic, bisulfuric, bitartaric, butyric, calcium edetate, camsylic, carbonic, chlorobenzoic, citric, edetic, edisylic, estolic, esyl, esylic, formic, fumaric, gluceptic, gluconic, glutamic, glycolic, glycolylarsanilic, hexamic, hexylresorcinoic, hydrabamic, hydrobromic, hydrochloric, hydroiodic, hydroxynaphthoic, isethionic, lactic, lactobionic, maleic, malic, malonic, mandelic, methanesulfonic, methylnitric, methylsulfuric, mucic, muconic, napsylic, nitric, oxalic, p-nitromethanesulfonic, pamoic, pantothenic, phosphoric, monohydrogen phosphoric, dihydrogen phosphoric, phthalic, polygalactouronic, propionic, salicylic, stearic, succinic, sulfamic, sulfanlic, sulfonic, sulfuric, tannic, tartaric, teoclic, toluenesulfonic, and the like.

Embodiments may be hydrochloride salts. Pharmaceutically acceptable salts may be derived from amino acids including, but not limited to, cysteine. Methods for producing compounds as salts are known to those of skill in the art (*see, e.g.*, Stahl et al., Handbook of Pharmaceutical Salts: Properties, Selection, and Use, Wiley-VCH; Verlag Helvetica Chimica

Acta, Zurich, 2002; Berge et al., J. Pharm. Sci. 66: 1, 1977).

An "effective amount" of a combination of therapeutic agents (e.g., Compound 1 and a CDK 4/6 inhibitor) is an amount sufficient to provide an observable therapeutic benefit compared to HCC or IHCC left untreated in a subject or patient.

Active agents as reported herein can be combined with a pharmaceutically acceptable carrier to provide pharmaceutical formulations thereof. The particular choice of carrier and formulation will depend upon the particular route of administration for which the composition is intended.

"Pharmaceutically acceptable carrier" as used herein refers to a nontoxic carrier, adjuvant, or vehicle that does not destroy the pharmacological activity of the compound with which it is formulated. Pharmaceutically acceptable carriers, adjuvants or vehicles that may be used in the compositions of this invention include, but are not limited to, sorbic acid, potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts or electrolytes, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, cellulose-based substances, polyethylene glycol, sodium carboxymethylcellulose, polyacrylates, waxes, polyethylene glycol and wool fat.

The compositions of the present invention may be suitable for parenteral, oral, inhalation spray, topical, rectal, nasal, buccal, vaginal or implanted reservoir administration, etc. In some embodiments, the formulation comprises ingredients that are from natural or non-natural sources. In some embodiments, the formulation or carrier may be provided in a sterile form. Non-limiting examples of a sterile carrier include endotoxin-free water or pyrogen-free water.

The term "parenteral" as used herein includes subcutaneous, intravenous, intramuscular, intra-articular, intra-synovial, intrasternal, intrathecal, intrahepatic, intralesional and intracranial injection or infusion techniques. In particular embodiments, the compounds are administered intravenously, orally, subcutaneously, or via intramuscular administration. Sterile injectable forms of the compositions of this invention may be aqueous or oleaginous suspension. These suspensions may be formulated according to techniques known in the art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a nontoxic parenterally acceptable diluent or solvent. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride

solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium.

For this purpose, any bland fixed oil may be employed including synthetic mono- or di-glycerides. Fatty acids and their glyceride derivatives are useful in the preparation of injectables, as are natural pharmaceutically acceptable oils, such as olive oil or castor oil, especially in their polyoxyethylated versions. These oil solutions or suspensions may also contain a long-chain alcohol diluent or dispersant, such as carboxymethyl cellulose or similar dispersing agents that are commonly used in the formulation of pharmaceutically acceptable dosage forms including emulsions and suspensions. Other commonly used surfactants, such as Tweens, Spans and other emulsifying agents that are commonly used in the manufacture of pharmaceutically acceptable solid, liquid, or other dosage forms may also be used for the purposes of formulation.

For oral administration, a compound or salt may be provided in an acceptable oral dosage form, including, but not limited to, capsules, tablets, aqueous suspensions or solutions. In the case of tablets for oral use, carriers commonly used include lactose and corn starch. Lubricating agents, such as magnesium stearate, may also be added. For oral administration in a capsule form, useful diluents include lactose and dried cornstarch. When aqueous suspensions are required for oral use, the active ingredient may be combined with emulsifying and suspending agents. If desired, certain sweetening, flavoring or coloring agents may also be added. In addition preservatives may also be added. Suitable examples of pharmaceutically acceptable preservatives include, but are not limited to, various antibacterial and antifungal agents such as solvents, for example ethanol, propylene glycol, benzyl alcohol, chlorobutanol, quaternary ammonium salts, and parabens (such as methyl paraben, ethyl paraben, propyl paraben, etc.).

"Immediate-release" is meant to include a conventional release, in which release of the drug starts immediately after administration. As used herein, the term "immediate release" includes dosage forms that allow the drug to dissolve in the gastrointestinal contents, with no intention of delaying or prolonging the dissolution or absorption of the drug. The objective is for the drug to be released rapidly after administration, for example for it to be possible to release at least 80% of the drug within approximately 30 minutes after commencement of dissolution in a dissolution test.

"Sustained-release" or "extended-release" includes dosage forms whose drug-release characteristics of time course and/or location are chosen to accomplish therapeutic or

convenience objectives not offered by conventional dosage forms such as a solution or an immediate release dosage form.

The term "steady-state" means that a plasma level for a given active agent or combination of active agents, has been achieved and which is maintained with subsequent doses of the active agent(s) at a level which is at or above the minimum effective therapeutic level and is below the minimum toxic plasma level for a given active agent(s).

The term "single formulation" as used herein refers to a single carrier or vehicle formulated to deliver effective amounts of both therapeutic agents to a patient. The single vehicle is designed to deliver an effective amount of each of the agents along with any pharmaceutically acceptable carriers or excipients. In some embodiments, the vehicle is a tablet, capsule, pill, or a patch.

The term "unit dose" is used herein to mean simultaneous administration of both agents together, in one dosage form, to the patient being treated. In some embodiments, the unit dose is a single formulation. In certain embodiments, the unit dose includes one or more vehicles such that each vehicle includes an effective amount of at least one of the agents (Compound 1 or a CDK 4/6 inhibitor) along with pharmaceutically acceptable carriers and excipients. In some embodiments, the unit dose is one or more tablets, capsules, pills, or patches administered to the patient at the same time.

The term "dose range" as used herein refers to an upper and a lower limit of an acceptable variation of the amount of agent specified. Typically, a dose of an agent in any amount within the specified range can be administered to patients undergoing treatment.

The term "treat" is used herein to mean to relieve, reduce or alleviate at least one symptom of a disease in a subject. For example, in relation to HCC, the term "treat" may mean to arrest, delay the onset (i.e., the period prior to clinical manifestation of a disease or symptom of a disease) and/or reduce the risk of developing or worsening a symptom of a disease. The term "protect" is used herein to mean prevent delay or treat, or all, as appropriate, development or continuance or aggravation of symptoms of the disease in a subject.

The term "subject" or “patient” is intended to include animals, which are capable of suffering from or afflicted with HCC or IHCC. Examples of subjects or patients include mammals, e.g., humans, dogs, cows, horses, pigs, sheep, goats, cats, mice, rabbits, rats, and transgenic non-human animals. In certain embodiments, the subject is a human, e.g., a human suffering from, at risk of suffering from, or potentially capable of suffering from HCC or

IHCC.

The term "about" or "approximately" usually means within 20%, more preferably within 10%, and most preferably still within 5% of a given value or range. Alternatively, especially in biological systems, the term "about" means approximately within a log (i.e., an order of magnitude) preferably within a factor of two of a given value.

The use of the terms "a" and "an" and "the" and similar referents in the context of describing the invention (especially in the context of the following claims) are to be construed to cover both the singular and the plural, unless otherwise indicated herein or clearly contradicted by context. The terms "comprising," "having," "including," and "containing" are to be construed as open-ended terms (i.e., meaning "including, but not limited to") unless otherwise noted. Recitation of ranges of values herein are merely intended to serve as a shorthand method of referring individually to each separate value falling within the range, unless otherwise indicated herein, and each separate value is incorporated into the specification as if it were individually recited herein.

The term "enhanced effect" as used herein, refers to action of two agents that administered together provide a greater or improved result than when the individual agents are administered alone without co-administration of the other agent. Administration of the agents together may provide an enhanced effect when they are administered simultaneously or sequentially. Sequential administration of the agents includes administrations separated by several seconds, minutes, hours or days. Administration of the agents together may provide an enhanced effect when the agents are administered either as part of a single formulation, or when administered in separate formulations. Examples of agents that may be administered together include Compound 1 and CDK4/6 inhibtors. Additional examples of agents that may be administered together include i) Compound 1 and ribociclib; ii) Compound 1 and palbociclib; and iii) Compound 1 and abemaciclib.

The enhanced effect's greater or improved result may include, for example, one or more of the following: i) improved quality of tumor response, ii) improved speed of the tumor response, and iii) a tumor response that is more than additive of the response that might otherwise be achieved had the individual agents been administered alone. Examples of improved quality of tumor response may include complete regregression (CR) instead of partial regression (PR), stable disease (SD) or progressive disease (PD). Another example of improved quality of tumor response may include partial regression (PR) instead of stable disease (SD) or progressive disease (PD). Another example of improved quality of tumor

response may include stable disease (SD) instead of progressive disease (PD). Controlled studies to determine whether administration of the agents together resulted with an enhanced effect of a tumor response more than additive of the corresponding responses achieved when the individual agents are respectively administered alone may be done, for example, in mice, rats, dogs, monkeys or other animals. Such controlled studies may evaluate, for example, the resulting tumor volume or metastatic or other status. Likewise, controlled studies may be used to determine an enhanced effect resulting in a faster tumor response.

In some embodiments, treatment is provided to a subject having hepatocellular carcinoma with altered FGFR4 and /or FGF19 (fibroblast growth factor 19) status.

In some embodiments, treatment may include or be performed in conjunction with analyzing FGFR4 and/or FGF19 status in a biological sample containing cells of said hepatocellular carcinoma, and if said hepatocellular carcinoma exhibits an FGFR4 and/or FGF19 alteration, treating a subject with a treatment effective amount of a therapeutic combination as described herein.

Methods of Treatment

Provided herein is a combination therapy useful for the treatment of HCC or IHCC. As discussed below, combinations provided herein may have a number of advantages.

One advantage of the combination disclosed herein is the unexpected enhanced effect of a combination of Compound 1 and a CDK 4/6 inhibitor on treatment of tumor grown inhibition and treatment of HCC or IHCC.

In some embodiments, provided herein is a single pharmaceutical formulation containing a combination of Compound 1 and a CDK 4/6 inhibitor. An advantage provided herein is the enhanced effect that results in the treatment of HCC compared to treatment with a single dose of either drug. When the drugs are provided in a single unit dose or single formulation, the "pill burden" on a patient suffering from HCC is not increased.

As specified above, in one aspect, provided herein is a drug combination useful for treating, preventing, arresting, delaying the onset of and/or reducing the risk of developing, or reversing HCC in a mammal comprising administering to said mammal a combination therapy, comprising an effective amount of Compound 1 and an effective amount of a CDK 4/6 inhibitor.

In some embodiments, the subject to be treated (e.g., patient) is determined to be non-responsive or resistant to one or more HCC therapies, e.g., Compound 1. In other embodiments, the individual to be treated is responsive to Compound 1 therapy, but the

therapy was improved with the administration of a CDK 4/6 inhibitor. For example, the patient is administered Compound 1 (e.g., 50 mg to 600 mg per day, 200 mg to 400 mg per day, or 300 mg per day for some period of time, e.g., more than one day, more than two days, more than three days, more than one week, for 21 days, more than one month, etc. After that time, a CDK 4/6 inhibitor could be administered to that patient in combination with Compound 1.

Amounts of CDK 4/6 inhibitor may vary depending on the CDK 4/6 inhibitor that is used. For example, palbociclib may be administered, for example in a dosage of 75, 100, or 125 mg/day; ribociclib may be administered, for example, in a dosage of 200, 400, or 600 mg/day. Typically a dosage is administered orally as a single capsule for 21 consecutive days followed by a 7 day off-treatment period.

The daily dosage may be part of a cyclic regimen lasting 14 to 21 days or longer. The daily dosage amount may be administered as a single dosage or as multiple dosages.

One skilled in the art appreciates that the effective dose of the active drug may be lower than the actual amount administered. As such, provided herein are doses necessary to achieve a therapeutic dose.

In various embodiments, provided herein are methods of treating HCC by administering an effective amount of Compound 1 and a CDK 4/6 inhibitor, to an individual having HCC. The amount of the combination of agents is effective to treat the HCC. In one embodiment, the combination of agents has an enhanced effect. In one embodiment, even though one or more of the agents administered alone at a particular dosage may be effective, when administered in combination, at the same dosage of each agent, the treatment is more effective. For example, in one embodiment a combination of Compound 1 and palbociclib is more effective than is administration of either agent alone. In another embodiment a combination of Compound 1 and ribociclib is more effective than is administration of either agent alone.

Dosages

The optimal dose of the combination of agents for treatment of HCC can be determined empirically for each individual using known methods and will depend upon a variety of factors, including the activity of the agents; the age, body weight, general health, gender and diet of the individual; the time and route of administration; and other medications the individual is taking. Optimal dosages may be established using routine testing and procedures that are well known in the art.

For the combination therapy of the instant invention, the daily dose of Compound 1 is in the range of 50 mg to 600 mg. In some embodiments, the daily dose of Compound 1 is up to 600 mg. In certain embodiments, the daily dose of Compound 1 is up to 400 mg. In various embodiments, the daily dose of Compound 1 is up to 300 mg. In certain embodiments, the daily dose of Compound 1 is 200 mg to 400 mg. In one embodiment, the daily dose is 300 mg.

The time of administration can be chosen such that both the drugs are administered simultaneously, separately or sequentially, either in the morning or at night. Alternatively, one drug can be administered in the morning and the other at night. In certain embodiments, both the drugs can be administered as a single tablet, capsule, pill, patch or jelly formulation, once daily, either in the morning or at night.

The amount of combination of agents that may be combined with the carrier materials to produce a single dosage form will vary depending upon the individual treated and the particular mode of administration. In some embodiments the unit dosage forms containing the combination of agents as described herein will contain the amounts of each agent of the combination that are typically administered when the agents are administered alone.

Pharmaceutical Formulations and Routes of Administration

Provided herein are pharmaceutical formulations comprising a combination of agents for the treatment of HCC. The pharmaceutical formulations may additionally comprise a carrier or excipient, stabilizer, flavoring agent, and/or coloring agent.

A combination of agents may be administered using a variety of routes of administration known to those skilled in the art. Routes of administration include oral administration. In certain embodiments, a pharmaceutical formulation comprising a combination of agents may be taken orally in the form of liquid, syrup, tablet, capsule, powder, sprinkle, chewtab, or dissolvable disk. Alternatively, pharmaceutical formulations of the present invention can be administered intravenously or transdermally. Additional routes of administration are known to those skilled in the art (see, e.g., Remington's Pharmaceutical Sciences, Gennaro A. R., Ed., 20.sup.th Edition, Mack Publishing Co., Easton, Pa.).

In some embodiments, the Compound 1 and CDK 4/6 inhibitor are formulated as a paste, jelly, or suspension. For example, the drugs are dissolved, entrapped or suspended in the form of drug particles, microencapsulated particles, or drug-polymer particles in a

gelatinous solution or semi-solid. An advantage of an oral jelly formulation is that it is easier to administer the drugs to patients who have difficulty swallowing tablets, capsules or pills. In certain embodiments, both agents are thoroughly mixed and suspended in an appropriate medium to form a paste or a gel. Additional agents can optionally be mixed to provide flavor during oral administration. Peanut butter or alginate, flavored with raspberry and a sweetener are examples of the many suitable taste masking agents. In various embodiments, the paste or jelly can also be formulated with suitable binders or excipients known in the art for topical administration.

Methods of preparing sustained release formulations in the form of tablets, capsules or pills are known in the art. In some embodiments, the sustained release formulation is prepared by coating the active ingredient of the drug with a polymer, preferably a water-insoluble polymer. For example, a water-insoluble polymer used in the pharmaceutical field as a sustained release coating agent, an enteric coating agent, or a gastric coating agent. The water-insoluble polymer can include, for example, ethyl cellulose, purified shellac, white shellac, aminoalkyl methacrylate copolymer RS, hydroxypropyl methylcellulose phthalate, hydroxypropyl methylcellulose acetate succinate, carboxymethylethyl-cellulose, cellulose acetate phthalate, methacrylic acid copolymer L, methacrylic acid copolymer LD, methacrylic acid copolymer S, aminoalkyl methacrylate copolymer E, or polyvinyl acetal diethylaminoacetate.

The type, degree of substitution and molecular weight of the water-insoluble polymers can depend on solubility of the active ingredient in water or an alcohol, the desired sustained release level and the like. The water-insoluble polymers can be used either alone or in combination. There can be further incorporated a hydrogenated oil, stearic acid, or cetanol as a coating auxiliary agent, and a middle-chain triglyceride, triacetin, triethyl citrate, or cetanol as a plasticizer.

In some embodiments, the sustained release formulation is a matrix-type tablet or granule. The active ingredient can be coated with up to 3 different types of polymers. These three different types of polymers can include: 1) a water insoluble polymer, such as ethylcellulose; 2) a pH independent gelling polymer, such as hydroxypropyl methylcellulose; and 3) a pH dependent gelling polymer, such as sodium alginate. These three different types of polymers can be used together to attenuate the release rate of the drugs.

Dosage Forms: Release Properties

Sustained-release formulations can achieve a degree of sustained effect. However, the exposure and/or the bioavailability of the active ingredient may vary based on a variety of factors, such as for example, the absorption window, the carriers or excipients used in the formulation, the mode of delivery of the formulation, and/or the transit time of the active ingredient through the gastrointestinal tract of the patient.

A combination therapy can contain at least one sustained-release portion for performing a sustained-release function and one immediate release portion for performing an immediate release function. In certain embodiments, when the combination therapy is in a single dosage form, it can be in the form of tablets formed from a mixture of sustained-release granules constituting a sustained-release portion and immediate-release granules constituting an immediate-release portion, a capsule preparation obtained by filling a capsule with sustained-release granules and immediate-release granules, or press-coated tablets in which an outer layer constituting an immediate-release portion is formed on an inner core constituting a sustained-release portion. There is, however, no limitation to the above embodiments.

Moreover, there are no particular limitations on the state of containment of each drug in the composition or in an immediate-release portion or a sustained-release portion; the Compound 1 may be dispersed uniformly in the composition, immediate release portion or sustained release portion, or may be contained in only one part of the composition, immediate-release portion or sustained-release portion, or may be contained such that there is a concentration gradient.

A sustained-release portion in the composition according to the present invention can contain at least one non-pH-dependent polymeric substance or pH-dependent polymeric substance for controlling drug release.

A non-pH-dependent polymeric substance used herein can comprise a polymeric substance whose charge state hardly changes under pH conditions generally found in the gastrointestinal tract, specifically from pH 1 to pH 8. This means, for example, a polymeric substance that does not have functional groups whose charge state changes depending on the pH such as basic functional groups such as amino groups or acidic functional groups such as carboxylic acid groups. Note that the non-pH-dependent polymeric substance can be included for giving the composition according to the present invention a sustained-release function, but may also be included for another purpose. Moreover, the non-pH-dependent polymeric

substance used in the present invention may be water-insoluble, or may swell in water or dissolve in water to form a gel.

Examples of water-insoluble non-pH-dependent polymeric substances include, but are not limited to, cellulose ethers, cellulose esters, and methacrylic acid-acrylic acid copolymers (trade name Eudragit, manufactured by Rohm GmbH & Co. KG, Darmstadt, Germany). Examples include, but are not limited to, cellulose alkyl ethers such as ethylcellulose (trade name Ethocel, manufactured by Dow Chemical Company, USA), ethyl methylcellulose, ethyl propylcellulose or isopropylcellulose, and butylcellulose, cellulose aralkyl ethers such as benzyl cellulose, cellulose cyanoalkyl ethers such as cyanoethylcellulose, cellulose organic acid esters such as cellulose acetate butyrate, cellulose acetate, cellulose propionate or cellulose butyrate, and cellulose acetate propionate, ethyl acrylate-methyl methacrylate copolymers (trade name Eudragit NE, manufactured by Rohm GmbH & Co. KG, Darmstadt, Germany), and aminoalkyl methacrylate copolymer RS (trade names Eudragit RL, Eudragit RS). There are no particular limitations on the mean particle diameter of a water-insoluble polymer used in the present invention, but usually the lower this mean particle diameter the better the performance, with the mean particle diameter preferably being from 0.1 to 100 μm, more preferably from 1 to 50 μm, particularly preferably from 3 to 15 μm, most preferably from 5 to 15 μm. Moreover, examples of water-soluble or water-swelling non-pH-dependent polymeric substances include, but are not limited to, polyethylene oxide (trade name Polyox, manufactured by Dow Chemical Company, molecular weight 100,000 to 7,000,000), low-substituted hydroxypropyl cellulose (trade name L-HPC, manufactured by Shin-Etsu Chemical, Japan), hydroxypropyl cellulose (trade name HPC, manufactured by Nippon Soda, Co., Ltd, Japan), hydroxypropyl methylcellulose (trade names Metolose 60SH, 65SH, 90SH, manufactured by Shin-Etsu Chemical, Japan), and methylcellulose (trade name Metolose SM, manufactured by Shin-Etsu Chemical, Japan).

In some embodiments a single non-pH-dependent polymeric substance may be contained in the composition, or a plurality of the non-pH-dependent polymeric substances may be contained. The non-pH-dependent polymeric substance, if used in embodiments reported herein, may be a water-insoluble polymeric substance, more preferably ethylcellulose, an ethyl acrylate-methyl methacrylate copolymer (trade name Eudragit NE), or an aminoalkyl methacrylate copolymer RS (trade name Eudragit RL, Eudragit RS). Particularly preferable is at least one of ethylcellulose and an aminoalkyl methacrylate copolymer RS. Most preferable is ethylcellulose. There are no particular limitations on the

amount of the non-pH-dependent polymeric substance contained in the composition; this amount can be adjusted as appropriate in accordance with the purpose such as controlling sustained drug release.

A pH-dependent polymeric substance that can be used in embodiments reported herein may be a polymeric substance whose charge state changes under pH conditions generally found in the gastrointestinal tract, specifically from pH 1 to pH 8. This means, for example, a polymeric substance having functional groups whose charge state changes depending on the pH such as basic functional groups such as amino groups or acidic functional groups such as carboxylic acid groups. The pH-dependent functional groups of the pH-dependent polymeric substance are preferably acidic functional groups, with the pH-dependent polymeric substance most preferably having carboxylic acid groups.

A pH-dependent polymeric substance used in the present invention may be water-insoluble, or may swell in water or dissolve in water to form a gel. Examples of pH-dependent polymeric substances used in the present invention include, but are not limited to, enteric polymeric substances. Examples of enteric polymeric substances include, but are not limited to, methacrylic acid-methyl methacrylate copolymers (Eudragit L100, Eudragit S100, manufactured by Rohm GmbH & Co. KG, Darmstadt, Germany), methacrylic acid-ethyl acrylate copolymers (Eudragit L100-55, Eudragit L30D-55, manufactured by Rohm GmbH & Co. KG, Darmstadt, Germany), hydroxypropyl methylcellulose phthalate (HP-55, HP-50, manufactured by Shin-Etsu Chemical, Japan), hydroxypropyl methylcellulose acetate succinate (AQOAT, manufactured by Shin-Etsu Chemical, Japan), carboxymethyl ethylcellulose (CMEC, manufactured by Freund Corporation, Japan), and cellulose acetate phthalate.

Examples of pH-dependent polymeric substances that swell in water or dissolve in water to form a gel include, but are not limited to, alginic acid, pectin, carboxyvinyl polymer, and carboxymethyl cellulose. In the present invention, a single pH-dependent polymeric substance may be contained in the composition, or a plurality of pH-dependent polymeric substances may be contained. The pH-dependent polymeric substance used in the present invention is preferably an enteric polymeric substance, more preferably a methacrylic acid-ethyl acrylate copolymer, a methacrylic acid-methyl methacrylate copolymer, hydroxypropyl methylcellulose phthalate, or hydroxypropyl methylcellulose acetate succinate, particularly preferably a methacrylic acid-ethyl acrylate copolymer.

When using a pH-dependent polymeric substance in the manufacturing process of a

composition according to the present invention, a commercially available product of a powder type or a granular type, or a suspension type in which the pH-dependent polymeric substance has been dispersed in a solvent in advance can be used as is, or such a commercially available product can be used dispersed in water or an organic solvent. The lower the particle diameter of the pH-dependent polymeric substance the better the performance, with the pH-dependent polymeric substance preferably being of the powder type. In the case of a methacrylic acid-ethyl acrylate copolymer, an example is Eudragit L100-55. There are no particular limitations on the mean particle diameter of a pH-dependent polymeric substance used in the present invention, but the mean particle diameter is preferably from 0.05 to 100 μm, more preferably from 0.05 to 70 μm, most preferably from 0.05 to 50 μm. Moreover, there are no particular limitations on the amount of the pH-dependent polymeric substance, for example, in the case of an enteric polymeric substance, the amount is generally from 0.1 to 90 parts by weight, preferably from 1 to 70 parts by weight, more preferably from 5 to 60 parts by weight, particularly preferably from 10 to 50 parts by weight, based on 100 parts by weight of the composition.

A combination therapy according to embodiments reported herein may further contain any of various additives, such as any of various pharmacologically acceptable carriers such as diluents, lubricants, binders and disintegrants, as well as preservatives, colorants, sweeteners, plasticizers, film coating agents and so on, as necessary. Examples of diluents include, but are not limited to, lactose, mannitol, dibasic calcium phosphate, starch, pregelatinized starch, crystalline cellulose, light silicic anhydride, synthetic aluminum silicate, magnesium aluminate metasilicate or the like. Examples of lubricants include, but are not limited to, magnesium stearate, calcium stearate, talc, sodium stearyl fumarate or the like. Examples of binders include, but are not limited to, hydroxypropyl cellulose, methylcellulose, sodium carboxymethyl cellulose, hydroxypropyl methylcellulose, polyvinylpyrrolidone or the like. Examples of disintegrants include, but are not limited to, carboxymethyl cellulose, calcium carboxymethyl cellulose, croscarmellose sodium, sodium carboxymethyl starch, low-substituted hydroxypropyl cellulose or the like. Examples of preservatives include, but are not limited to, paraoxybenzoic acid esters, chlorobutanol, benzyl alcohol, phenethyl alcohol, dehydroacetic acid, sorbic acid or the like. Preferable examples of colorants include, but are not limited to, water-insoluble lake pigments, natural pigments (e.g., .beta.-carotene, chlorophyll, red ferric oxide), yellow ferric oxide, red ferric oxide, black ferric oxide or the like. Preferable examples of sweeteners include, but are not limited to, sodium saccharin,

dipotassium glycyrrhizate, aspartame, stevia or the like. Examples of plasticizers include, but are not limited to, glycerol fatty acid esters, triethyl citrate, propylene glycol, polyethylene glycol or the like. Examples of film coating agents include, but are not limited to, hydroxypropyl methylcellulose, hydroxypropyl cellulose or the like.

Manufacturing Methods

To manufacture embodiments as reported herein, a single conventional method, or a combination of conventional methods, can be used. For example, when manufacturing drug-containing granules as a sustained-release portion or an immediate-release portion, granulation is the main operation, but this may be combined with other operations such as mixing, drying, sieving, and classification. As the granulation method, for example, a wet granulation method in which a binder and a solvent are added to the powder and granulation is carried out, a dry granulation method in which the powder is compressed and granulation is carried out, a molten granulation method in which a binder that melts on heating is added and heating and granulation are carried out, or the like can be used.

Furthermore, in accordance with the granulation method, an operating method such as a mixing granulation method using a planetary mixer, a screw mixer or the like, a high-speed mixing granulation method using a Henschel mixer, a Super mixer or the like, an extruding granulation method using a cylindrical granulator, a rotary granulator, a screw extruding granulator, a pellet mill type granulator or the like, a wet high-shear granulation method, a fluidized-bed granulation method, a compression granulation method, a crushing granulation method, or a spraying granulation method can be used. After the granulation, drying using a dryer, a fluidized bed or the like, cracking, and sieving can be carried out to obtain the granules or fine granules for use. Moreover, a granulation solvent may be used when preparing the composition according to the present invention. There are no particular limitations on such a granulation solvent, which may be water or any of various organic solvents, for example, water, a lower alcohol such as methanol or ethanol, a ketone such as acetone or methyl ethyl ketone, methylene chloride, or a mixture thereof.

For sustained-release granules contained in embodiments, at least one drug and at least one selected from non-pH-dependent polymeric substances and pH-dependent polymeric substances are mixed together, a diluent and a binder are added as necessary, and granulation is carried out to obtain granular matter. The granular matter obtained is dried

using a tray dryer, a fluidized bed dryer or the like, and sieving is carried out using a mill or an oscillator, whereby the sustained-release granules can be obtained. Alternatively, as a method of manufacturing sustained-release granules in the present invention, it is possible to add at least one drug, at least one selected from non-pH-dependent polymeric substances and pH-dependent polymeric substances, and as necessary a diluent and a binder using a dry compactor such as a roller compactor or a slug tabletting machine, and carry out compression-molding while mixing, and then carry out granulation by cracking down to a suitable size. The granular matter prepared using such a granulator may be used as is as granules or fine granules according to the present invention, or may be further cracked using a power mill, a roll granulator, a rotor speed mill or the like, and sieved to obtain sustained-release granules. Note that immediate-release granules can also be manufactured as for the sustained-release granules.

A compression-molded product can be manufactured as a drug-containing sustained-release portion or immediate-release portion, or as a composition reported herein using a single conventional method, or a combination of conventional methods. For example, at least one drug, at least one selected from non-pH-dependent polymeric substances and pH-dependent polymeric substances, a diluent such as mannitol or lactose, a binder such as polyvinylpyrrolidone or crystalline cellulose, a disintegrant such as carmellose sodium or crospovidone, and a lubricant such as magnesium stearate or talc are used, and tableting is carried out using an ordinary method, whereby the compression-molded product can be obtained. In this case, tabletting is the main operation in the method of manufacturing the compression-molded product, but this may be combined with other operations such as mixing, drying, sugar coating formation, and coating.

Examples of the method for the tabletting include, but are not limited to, direct compression molding in which at least one drug and pharmacologically acceptable additives are mixed together and then the mixture is directly compression-molded into tablets using a tabletting machine, and dry granule compression or wet granule compression in which sustained-release granules or immediate-release granules according to the present invention are subjected to compression-molding after adding a lubricant or a disintegrant as necessary. There are no particular limitations on the tabletting machine used in the compression molding; for example, a single-punch tabletting machine, a rotary tabletting machine, or a press-coated tabletting machine can be used.

Drug-containing sustained-release granules or immediate-release granules, or

compression-molded product according to embodiments herein can be used as is in the form of granules or a tablet as the composition, but may also be subjected to further processing to manufacture the composition. For example, the compression-molded product or granules can be given a film coating using a film base material such as ethylcellulose, casein, methylcellulose, hydroxypropyl methylcellulose, methacrylic acid copolymer L, cellulose acetate phthalate, shellac or the like, or given a sugar coating using a sugar coating liquid containing saccharose, sugar alcohol, gum arabic powder, talc or the like, thus producing film-coated tablets or sugar-coated tablets. One solvent in this coating technique may be purified water, but an organic solvent such as an alcohol, a ketone, an ether or a chlorinated hydrocarbon, or a mixture thereof can also be used. For example, ethanol, acetone, methylene chloride or the like can be used as an organic solvent. Moreover, as the coating apparatus, an apparatus ordinarily used in coating techniques for manufacturing medicines can be used, with examples including a spray coating apparatus in which the coating is carried out by spraying a coating liquid or the like, and a rotor fluidized bed granulator for layering.

In the case of manufacturing capsule preparations, capsule preparations can be manufactured by filling sustained-release granules or immediate-release granules as above, or mini-tablets into hard gelatin capsules or HPMC capsules using an automatic capsule filling machine. Alternatively, in the case of the preparations for per-tube administration or a dry syrup that is used mixed with water or the like when taken, sustained-release granules or immediate-release granules as above can be mixed with a thickener or a dispersant so as to disperse these granules, the mixture then being made into granules or tablets. Furthermore, a liquid or jelly can be made using water, and substances selected from dispersants, emulsifiers, thickeners, preservatives, pH adjustors, sweeteners, flavorings, fragrances and so on. However, with respect to other manufacturing methods, there are no limitations to the above.

So that embodiments described herein may be more fully understood, the following examples are set forth. It should be understood that these examples are for illustrative purposes only and are not to be construed as limiting.

## EXAMPLES

Materials and methods

Cell Lines Tested

The cell line used, JHH7, was sourced from Japanese Collection of Research Bioresources Cell Bank (JCRB), verified free of mycobacterium contamination and verified for identity by short tandem repeat analysis of 9 markers.

Cell Line Maintenance and Study Conditions

JHH7 cells were maintained in William's E Medium (Thermo Fisher Scientific, 12551-032) with 10% fetal bovine serum. Cells were maintained prior to and during experiments at 37 °C, 5% $CO_2$, and at 95% relative humidity. Cell passage number was limited to between 12 and 20.

Xenograft Generation, Dosing and Measurement of Antitumor Activity

The human hepatocellular cancer cell line JHH7 was cultured in William's E Medium (Thermo Fisher Scientific, 12551-032) containing 10% fetal bovine serum at 37 °C in a 5% $CO_2$ atmosphere and kept in exponential growth phase. For harvesting, the cells were washed with phosphate buffered saline, incubated with 0.25% trypsin-EDTA, and suspended in a 1:1 mixture of William's E Medium and Matrigel (Corning) at a final concentration of 5 x $10^7$cells/mL. To generate xenografts, 0.1-mL of the inoculum was injected subcutaneously into the right flank region of NU/NU immuno compromised 6-8 week old female mice, giving a final concentration of 5 x $10^6$ cells/mouse. When the mean tumor volume (TV) reached approximately 170 $mm^3$ (10 days after implantation), 144 mice were selected based on their TVs, and were sorted into 18 treatment groups with 8 animals per group. Per os (PO) treatment with Compound 1 (300 and 500 mg/kg) alone or in combination with vehicle (control) or palbociclib (50 and 100mg/kg) or ribociclib (75 and 150mg/kg) administered once daily (QD) continued for 8 days. The administration volume (0.1 mL/10 g body weight) was calculated from the individual mouse body weight (BW) before administration. Body weights were measured daily and tumor measurements were performed twice weekly.

The human primary hepatocellular carcinoma model LIX066 model from ChemPartner (shangpharma.com) were implanted into the female severe combined immunodeficiency (SCID) mice. When the tumors developed in mice, the mice were sacrificed and the tumors were resected and implanted into female nude mice for tumor preservation, histopathology diagnosis and in vivo efficacy study. Solid tumor tissues were depleted of necrotic components, cut into 10-15 mg pieces, and mixed. Three to five pieces were mixed with 15-30 mL Matrigel and implanted into a flank of 6-8-week-old female

NU/NU immune compromised mice weighing 18-20 g nude mice. 30 mice with average tumor volume of 160 $mm^3$ selected and randomized into 6 groups of 5 mice each. All of the primary human tumors utilized in this study had undergone 3-5 passages in vivo and the tumor histology of each was maintained over the serial transplantation process. Per os (PO) treatment with Compound 1 (300 and 500 mg/kg) alone or in combination with vehicle (control) or palbociclib (100mg/kg) administered once daily (QD) continued for 17 days. The administration volume (0.1 mL/10 g body weight) was calculated from the individual mouse body weight (BW) before administration. Body weights were measured daily and tumor measurements were performed twice weekly.

The TV in $mm^3$ was calculated according to the following formula: TV = length × $width^2$ × 0.5 length: largest diameter of tumor (mm) width: diameter perpendicular to length (mm) The Tumor Growth Inhibition% (TGI) was calculated according to the following formula:

$$\frac{\text{Average Control TV Day X - Treatment TV Day X}}{\text{Average Control TV Day X}} \times 100$$

where Day X is any day of measurement

The anti-tumor effects of the treatment, Partial (PR) and complete regression (CR), stable (SD) and progressive (PD) disease were defined by the Xenograft Model Response Criteria (see below). Mice with > 20% body weight loss compared to their Day 1 body weight or bearing tumors with the longest diameter > 2000 mm were immediately euthanized to prevent any pain or suffering in the animal as according to IACUC guidelines.

Statistical Analysis

Data are expressed as the mean ±SEM for Tumor Volume (TV). The differences in TV between groups were analyzed by two way ANOVA followed by Sidak post hoc test. Statistical analyses were performed using the GraphPad Prism version 6 (GraphPad Software, La Jolla, CA).

Xenograft Model Response Criteria

Progressive disease (PD): 3 consecutive measurements >120% of starting volume or 3 consecutive increasing measurements from best response, Stable disease (SD): 3

consecutive measurements >50% and <120% of starting volume, Partial regression (PR): 3 consecutive measurements <50% of starting volume, Complete regression (CR): 3 consecutive measurements <30mm$^3$.

Formulation of CDK 4/6 Inhibitors

In the examples reported below, ribociclib and palbociclib were formulated as follows. This type of formulation is exemplary and not required in particular embodiments of the invention. In these examples both ribociclib and palbociclib were presented as free bases.

Palbociclib was formulated in a 25mM Sodium Bicarbonate, 15mM Lactic Acid solution with 2% Cremaphor. Add cremaphor first and sonicate until a fine, even suspension is formed. This compound should be made fresh each time.

Ribociclib was formulated in a 0.5% Methylcellulose in distilled water with 1% Cremaphor. Add Cremaphor first and sonicate until a fine even suspension is formed. This compound should be made fresh each time

Example 1 - Compound 1 and Ribociclib

The JHH7 cell line was grown as a xenograft in female nude immunocompromised mice and tumor bearing mice were orally treated daily for 8 days with 300 or 500mg/kg Compound 1 as single agent or in combination with 75 and 150mg/kg ribociclib. Ribociclib, as single agent at 75mg/kg did not significantly inhibit tumor growth with 13% TGI whereas 150mg/kg significantly inhibit tumor growth ($P \leq 0.0001$) with 23% TGI in comparison to the vehicle controls. The single agent Compound 1 at 300 and 500mg/kg resulted in significant inhibition of tumor growth ($P \leq 0.0001$) with TGI of 14% and 35%, respectively in comparison to the vehicle controls. The combination of 300mg/kg Compound 1 and 75mg/kg ribociclib resulted in significant enhancement of tumor growth inhibition ($P \leq 0.0001$) in comparison to 300mg/kg Compound 1 single agent with TGI of 44%. The combination of 300mg/kg Compound 1 and 150mg/kg ribociclib also resulted in significant enhancement of tumor growth inhibition ($P \leq 0.0001$) in comparison to the 300mg/kg Compound 1 alone with TGI of 64%. The combination of 500mg/kg Compound 1 and 75mg/kg ribociclib resulted in significant enhancement of tumor growth inhibition ($P \leq 0.0001$ in comparison to 500mg/kg Compound 1 alone with TGI of 61%. The combination of 500mg/kg Compound 1 and 150mg/kg Ribociclib also resulted in significant enhancement of tumor growth inhibition ($P \leq 0.0001$) in comparison to the 500mg/kg Compound 1 alone with

TGI of 73%.

These data demonstrate that ribociclib can significantly enhance the antitumor effects of Compounr 1. Treatment with Compound 1 as single agent or in combination with Ribociclib did not cause any CR, PR or SD and all groups had PD. All combination dosing groups were well tolerated based on body weight measurements and routine clinical observation.

Results are shown in FIG. 1A through FIG. 1C.

Example 2 – Compound 1 and Palbociclib

The JHH7 cell line was grown as a xenograft in female nude immunocompromised mice and tumor bearing mice were orally treated daily for 8 days with 300 or 500mg/kg Compound 1 as single agent or in combination with 50 and 100mg/kg palbociclib. Palbociclib, as single agent at 50mg/kg did not significantly inhibit tumor growth with 11% TGI whereas 100mg/kg significantly inhibit tumor growth ($P \leq 0.05$) with 26% TGI in comparison to the vehicle controls. The single agent Compound 1 at 300mg/kg resulted in significant inhibition of tumor growth ($P \leq 0.01$) with TGI of 12% and 500mg/kg also resulted in significant tumor growth inhibition ($P \leq 0.0001$) with 32% in comparison to the vehicle controls. The combination of 300mg/kg Compound 1 and 50mg/kg Palbociclib resulted in significant enhancement of tumor growth inhibition ($P \leq 0.0001$) in comparison to 300mg/kg Compound 1 single agent with TGI of 59%.

The combination of 300mg/kg Compound 1 and 100mg/kg palbociclib also resulted in significant enhancement of tumor growth inhibition ($P \leq 0.0001$) in comparison to the 300mg/kg Compound 1 alone with TGI of 77%. The combination of 500mg/kg Compound 1 and 50mg/kg palbociclib resulted in significant enhancement of tumor growth inhibition ($P \leq 0.0001$) in comparison to 500mg/kg Compound 1 alone with TGI of 62%. The combination of 500mg/kg Compound 1 and 100mg/kg palbociclib also resulted in significant enhancement of tumor growth inhibition ($P \leq 0.0001$) in comparison to the 500mg/kg Compound 1 alone with TGI of 77%.

These data demonstrate that palbociclib can significantly enhance the antitumor effects of Compound 1. Treatment with Compound 1 as single agent or in combination with palbociclib did not cause any CR, PR or SD and all groups had PD. All combination dosing groups were well tolerated based on body weight measurements and routine clinical observation.

Results are shown in FIG. 2A through FIG. 2C.

Example 3 – Compound 1 and Palbociclib

In a patient derived xenograft model (PDX), LIX066 PDX fragments were inoculated in female nude immunocompromised mice and tumor bearing mice were orally treated daily for 17 days with 300 or 500mg/kg Compound 1 as single agent or in combination with 100mg/kg palbociclib. Palbociclib, as single agent at 100mg/kg significantly inhibited tumor growth ($P \leq 0.001$), in comparison to the vehicle controls, with 62% TGI. All the enrolled animals in this group showed PD. The single agent Compound 1 at 300mg/kg and 500mg/kg significantly inhibited tumor growth ($P \leq 0.0001$) in comparison to the vehicle controls, with 59% TGI and 70% TGI, respectively. All animals in the 300mg/kg and the 500mg/kg Compound 1 groups showed PD.

The combination of 300mg/kg Compound 1 and 100mg/kg palbociclib resulted in significant enhancement of tumor growth inhibition ($P \leq 0.0001$) in comparison to 300mg/kg Compound 1 alone with TGI of 96%. This combination led to SD in 3/5 animals and PR in 2/5 animals. The combination of 500mg/kg Compound 1 and 100mg/kg palbociclib also resulted in significant enhancement of tumor growth inhibition ($P \leq 0.0001$) in comparison to the 300mg/kg Compound 1 alone with TGI of 97%. This combination led to SD in 1/5 animals and PR in 4/5 animals. All combination dosing groups were well tolerated based on body weight measurements and routine clinical observation.

Results are shown in FIG. 3A and FIG. 3B. Consistent with the discovery that Palbociclib can enhance Compound 1 antitumor effects in HCC models, additional testing in 7 more PDX models showed similar enhancement in 3 out of 7 models.

Claims

1. A pharmaceutical formulation comprising *N*-(2-((6-(3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-methylureido)pyrimidin-4-yl)amino)-5-(4-ethylpiperazin-1-yl)phenyl)acrylamide or a pharmaceutically acceptable salt thereof and a CDK 4/6 inhibitor or a pharmaceutically acceptable salt thereof.

2. The pharmaceutical formulation of claim 1, comprising a free base form of *N*-(2-((6-(3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-methylureido)pyrimidin-4-yl)amino)-5-(4-ethylpiperazin-1-yl)phenyl)acrylamide.

3. The pharmaceutical formulation of claim 1, wherein the pharmaceutically acceptable salt of *N*-(2-((6-(3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-methylureido)pyrimidin-4-yl)amino)-5-(4-ethylpiperazin-1-yl)phenyl)acrylamide is a hydrochloride salt form of *N*-(2-((6-(3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-methylureido)pyrimidin-4-yl)amino)-5-(4-ethylpiperazin-1-yl)phenyl)acrylamide.

4. The pharmaceutical formulation as defined in claim 1 for use in therapy.

5. Use of N-(2-((6-(3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-methylureido)pyrimidin-4-yl)amino)-5-(4-ethylpiperazin-1-yl)phenyl)acrylamide or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for the treatment of hepatocellular carcinoma, wherein N-(2-((6-(3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-methylureido)pyrimidin-4-yl)amino)-5-(4-ethylpiperazin-1-yl)phenyl)acrylamide or a pharmaceutically acceptable salt thereof is to be administered simultaneously or sequentially with a CDK 4/6 inhibitor or a pharmaceutically acceptable salt thereof to a patient in need thereof.

6. The use of claim 5, wherein the *N*-(2-((6-(3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-methylureido)pyrimidin-4-yl)amino)-5-(4-ethylpiperazin-1-yl)phenyl)acrylamide or a pharmaceutically acceptable salt thereof is to be administered in a daily dosage selected from the group consisting of from 50 mg to 600 mg/day, from 200 mg to 400 mg/day, from 50 mg/day to 3000 mg/day, 150 mg/day, 300 mg/day, 600 mg/day, 1000 mg/day, 1500 mg/day, and 2000 mg/day.

7. The use of claim 5 or 6, wherein the *N*-(2-((6-(3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-methylureido)pyrimidin-4-yl)amino)-5-(4-ethylpiperazin-1-yl)phenyl)acrylamide or a pharmaceutically acceptable salt thereof is to be administered as a single dosage or in multiple dosages.

8. The use of any one of claims 5 to 7, wherein the CDK 4/6 inhibitor is selected

from the group consisting of 6-acetyl-8-cyclopentyl-5-methyl-2-{[5-(piperazin-1-yl)pyridin-2yl]amino}pyrido[2,3-*d*]pyrimidin-7(8*H*)-one (palbociclib) and pharmaceutically acceptable salts thereof; *N*-(5-((4-ethylpiperazin-1-yl)methyl)pyridin-2-yl)-5-fluoro-4-(4-fluoro-1-isopropyl-2-methyl-*1H*-benzo[d]imidazol-6-yl)pyrimidin-2-amine (abemaciclib) and pharmaceutically acceptable salts thereof; and 7-cyclopentyl-2-(5-piperazin-1-yl-pyridin-2-ylamino)-7H-pyrrolo[2,3-d]pyrimidine-6-carboxylic acid dimethylamide (ribociclib); G1T-38; 2'-((5-(4-methylpiperazin-1-yl)pyridin-2-yl)amino)-7',8'-dihydro-6'H-spiro[cyclohexane-1,9'-pyrazino[1',2':1,5]pyrrolo[2,3-d]pyrimidin]-6'-one (G1T-28); N-(4-piperidinyl)-4-(2,6-dichlorobenzoylamino)-1*H*-pyrazole-3 carboxamide (AT-7519); 2-Hydroxy-1-[2-[[9-(trans-4-methylcyclohexyl)-9*H*-pyrido[4',3':4,5]pyrrolo[2,3-d]pyrimidin-2-yl]amino]-7,8-dihydro-1,6-naphthyridin-6(5*H*)-yl]ethanone (FLX-925); 2-(2-chlorophenyl)-5,7-dihydroxy-8-((3*S*,4*R*)-3-hydroxy-1-methylpiperidin-4-yl)-4*H*-chromen-4-one (alvocidib) and pharmaceutically acceptable salts thereof.

9. The use of any one of claims 5 to 8, wherein the CDK 4/6 inhibitor is palbociclib.

10. The use of claim 9, wherein the palbociclib is to be administered in a daily dosage selected from the group consisting of 75 mg/day, 100 mg/day, and 125 mg/day.

11. The use of any one of claims 5 to 8, wherein the CDK 4/6 inhibitor is ribociclib.

12. The use of claim 11, wherein the ribociclib is to be administered in a daily dosage selecting from the group consisting of 200 mg/day, 400 mg/day, and 600 mg/day.

13. The use of any one of claims 5 to 8, wherein the CDK 4/6 inhibitor is abemaciclib.

14. The use of claim 13, wherein the abemaciclib is to be administered in a daily dosage selected from the group consisting of 200 mg/day, 300 mg/day, and 400 mg/day.

15. The use of any one of claims 5 to 14, wherein the *N*-(2-((6-(3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-methylureido)pyrimidin-4-yl)amino)-5-(4-ethylpiperazin-1-yl)phenyl)acrylamide or a pharmaceutically acceptable salt thereof and the CDK 4/6 inhibitor or pharmaceutically acceptable salt thereof are to be administered as separate formulations.

16. The use of any one of claims 5 to 14, wherein the *N*-(2-((6-(3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-methylureido)pyrimidin-4-yl)amino)-5-(4-ethylpiperazin-1-yl)phenyl)acrylamide or a pharmaceutically acceptable salt thereof and the CDK 4/6 inhibitor or pharmaceutically acceptable salt thereof are to be administered simultaneously as a single

formulation.

17. The use of any one of claims 5 to 14, wherein the *N*-(2-((6-(3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-methylureido)pyrimidin-4-yl)amino)-5-(4-ethylpiperazin-1-yl)phenyl)acrylamide or a pharmaceutically acceptable salt thereof and the CDK 4/6 inhibitor or pharmaceutically acceptable salt thereof are to be administered sequentially.

18. The use of any one of claims 5 to 14, wherein the *N*-(2-((6-(3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-methylureido)pyrimidin-4-yl)amino)-5-(4-ethylpiperazin-1-yl)phenyl)acrylamide or a pharmaceutically acceptable salt thereof and the CDK 4/6 inhibitor or pharmaceutically acceptable salt thereof are to be administered simultaneously.

19. The use of any one of claims 5 to 18, wherein the *N*-(2-((6-(3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-methylureido)pyrimidin-4-yl)amino)-5-(4-ethylpiperazin-1-yl)phenyl)acrylamide is the free base form of *N*-(2-((6-(3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-methylureido)pyrimidin-4-yl)amino)-5-(4-ethylpiperazin-1-yl)phenyl)acrylamide.

20. The use of any one of claims 5 to 18, wherein the pharmaceutically acceptable salt of *N*-(2-((6-(3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-methylureido)pyrimidin-4-yl)amino)-5-(4-ethylpiperazin-1-yl)phenyl)acrylamide is a hydrochloride salt form of *N*-(2-((6-(3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-methylureido)pyrimidin-4-yl)amino)-5-(4-ethylpiperazin-1-yl)phenyl)acrylamide.