

METHODS FOR TREATING HUNTINGTON'S DISEASE

The present description relates to compounds, forms, and pharmaceutical compositions thereof and methods of using such compounds, forms, or compositions thereof for treating or ameliorating Huntington's disease. In particular, the present description relates to substituted
5 monocyclic heteroaryl compounds, forms and pharmaceutical compositions thereof and methods of using such compounds, forms, or compositions thereof for treating or ameliorating Huntington's disease.


BACKGROUND

Huntington's disease (HD) is a progressive, autosomal dominant neurodegenerative
10 disorder of the brain, having symptoms characterized by involuntary movements, cognitive impairment, and mental deterioration. Death, typically caused by pneumonia or coronary artery disease, usually occurs 13 to 15 years after the onset of symptoms. The prevalence of HD is between three and seven individuals per 100,000 in populations of western European descent. In North America, an estimated 30,000 people have HD, while an additional 200,000 people are at
15 risk of inheriting the disease from an affected parent. The disease is caused by an expansion of uninterrupted trinucleotide CAG repeats in the "mutant" huntingtin (Htt) gene, leading to production of HTT (Htt protein) with an expanded poly-glutamine (polyQ) stretch, also known as a "CAG repeat" sequence. There are no current small molecule therapies targeting the underlying cause of the disease, leaving a high unmet need for medications that can be used for
20 treating or ameliorating HD. Consequently, there remains a need to identify and provide small molecule compounds for treating or ameliorating HD.

All other documents referred to herein are incorporated by reference into the present application as though fully set forth herein.

SUMMARY

25 The present description relates to methods for treating or ameliorating HD in a subject in need thereof comprising administering to the subject an effective amount of a compound of Formula (I):




wherein W, X, A and B are as defined herein, or forms and compositions thereof.

In particular, the present description relates to a use of a compound of Formula (I) or a
 5 form or composition thereof in a method for treating or ameliorating HD in a subject in need
 thereof comprising, administering an effective amount of the compound or a form or
 composition thereof, to the subject.

The present description further relates to the use of a compound of Formula (I) or a form
 thereof in combination with agents having additive or synergistic activity, thus providing a
 10 combination product for the treatment of HD.

DETAILED DESCRIPTION

The present description relates to a method or use of a compound for treating or
 ameliorating HD in a subject in need thereof comprising administering to the subject an effective
 amount of a compound of Formula (I):



15

or a form thereof, wherein

W is CH=CH or S;

X is CH₂, CH(C₁₋₄alkyl), C(C₁₋₄alkyl)₂, CH=CH, O, NR₅, or a bond;

20 A is aryl, heteroaryl, heterocyclyl, or C₉₋₁₀cycloalkyl,

wherein aryl is selected from phenyl and naphthyl, each optionally substituted with 1, 2, 3, or 4
 substituents each selected from R₁,

wherein heteroaryl is a saturated monocyclic, bicyclic or tricyclic ring system having 1, 2, or 3
 heteroatom ring members independently selected from N, O, or S, each optionally

25 substituted with 1, 2, 3, 4, or 5 substituents each selected from R₁,

wherein heterocyclyl is a saturated or partially unsaturated monocyclic, bicyclic or tricyclic ring system having 1, 2, or 3 heteroatom ring members independently selected from N, O, or S, each optionally substituted with 1, 2, 3, 4, or 5 substituents each selected from R₂, and wherein C₉₋₁₀cycloalkyl is a saturated or partially unsaturated bicyclic ring system optionally substituted with 1, 2, 3, 4, or 5 substituents each selected from R₂;

B is heterocyclyl,

wherein heterocyclyl is a saturated or partially unsaturated monocyclic, bicyclic or polycyclic ring system having 1, 2, or 3 heteroatom ring members independently selected from N, O, or S, each optionally substituted with 1, 2, 3, 4, or 5 substituents each selected from R₄;

R₁ is halogen, hydroxyl, cyano, C₁₋₄alkyl, halo-C₁₋₄alkyl, amino, C₁₋₄alkyl-amino, (C₁₋₄alkyl)₂-amino, amino-C₁₋₄alkyl, C₁₋₄alkyl-amino-C₁₋₄alkyl, (C₁₋₄alkyl)₂-amino-C₁₋₄alkyl, amino-carbonyl, C₁₋₄alkyl-amino-carbonyl, (C₁₋₄alkyl)₂-amino-carbonyl, C₁₋₄alkyl-amino-carbonyl-C₁₋₄alkyl, (C₁₋₄alkyl)₂-amino-carbonyl-C₁₋₄alkyl, C₁₋₄alkyl-carbonyl-amino, C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkyl, hydroxyl-C₁₋₄alkyl, C₁₋₄alkyl-carbonyl, C₁₋₄alkoxy, halo-C₁₋₄alkoxy, amino-C₁₋₄alkoxy, hydroxyl-C₁₋₄alkoxy, C₁₋₄alkyl-C₁₋₄alkoxy, C₁₋₄alkyl-amino-C₁₋₄alkoxy, (C₁₋₄alkyl)₂-amino-C₁₋₄alkoxy, C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkoxy, C₁₋₄alkoxy-C₁₋₄alkoxy, C₁₋₄alkoxy-carbonyl, C₁₋₄alkoxy-carbonyl-amino, C₁₋₄alkoxy-carbonyl-amino-C₁₋₄alkoxy, C₂₋₄alkenyl, C₂₋₄alkenyl-amino-carbonyl, C₃₋₇cycloalkyl, C₃₋₇cycloalkyl-C₁₋₄alkoxy, C₃₋₇cycloalkenyl, heteroaryl, heteroaryl-C₁₋₄alkyl, heteroaryl-C₁₋₄alkyl-amino, heteroaryl-C₁₋₄alkyl-amino-carbonyl, heteroaryl-C₁₋₄alkyl-carbonyl-amino, heteroaryl-C₁₋₄alkyl-amino-carbonyl-C₁₋₄alkyl, heteroaryl-C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkyl, heterocyclyl, heterocyclyl-C₁₋₄alkyl, heterocyclyl-C₁₋₄alkoxy, phenyl, or phenyl-C₁₋₄alkoxy,

wherein heteroaryl is a saturated monocyclic or bicyclic ring system having 1, 2, or 3 heteroatom ring members selected from N, O, and S,

wherein heterocyclyl is a saturated or partially unsaturated monocyclic or bicyclic ring system having 1, 2, or 3 heteroatom ring members selected from N, O, and S, and

wherein each instance of phenyl, heteroaryl or heterocyclyl is optionally substituted with 1, or 2 substituents each selected from R₃;

R₂ is halogen, hydroxyl, cyano, oxo, hydroxyl-imino, C₁₋₄alkyl, halo-C₁₋₄alkyl, amino, C₁₋₄alkyl-amino, (C₁₋₄alkyl)₂-amino, amino-C₁₋₄alkyl, C₁₋₄alkyl-amino-C₁₋₄alkyl, (C₁₋₄alkyl)₂-amino-C₁₋₄alkyl, amino-carbonyl, hydroxyl-C₁₋₄alkyl, C₁₋₄alkoxy, C₁₋₄alkoxy-carbonyl, C₂₋₄alkenyl, C₃₋₇cycloalkyl, or heterocyclyl-C₁₋₄alkyl,

- 5 wherein heterocyclyl is a saturated or partially unsaturated monocyclic or bicyclic ring system having 1, 2, or 3 heteroatom ring members selected from N, O, and S, and wherein each instance of heterocyclyl is optionally substituted with 1, or 2 substituents each selected from R₃;


- R₃ is halogen, hydroxyl, nitro, oxo, hydroxyl-imino, C₁₋₄alkyl, halo-C₁₋₄alkyl, amino, C₁₋₄alkyl-amino, (C₁₋₄alkyl)₂-amino, amino-C₁₋₄alkyl, C₁₋₄alkyl-amino-C₁₋₄alkyl, (C₁₋₄alkyl)₂-amino-C₁₋₄alkyl, amino-carbonyl, C₁₋₄alkyl-amino-carbonyl, (C₁₋₄alkyl)₂-amino-carbonyl, C₁₋₄alkyl-amino-carbonyl-C₁₋₄alkyl, (C₁₋₄alkyl)₂-amino-carbonyl-C₁₋₄alkyl, C₁₋₄alkyl-carbonyl-amino, C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkyl, hydroxyl-C₁₋₄alkyl, C₁₋₄alkyl-carbonyl, C₁₋₄alkoxy, halo-C₁₋₄alkoxy, amino-C₁₋₄alkoxy, hydroxyl-C₁₋₄alkoxy, C₁₋₄alkyl-C₁₋₄alkoxy, C₁₋₄alkyl-amino-C₁₋₄alkoxy, (C₁₋₄alkyl)₂-amino-C₁₋₄alkoxy, C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkoxy, C₁₋₄alkoxy-C₁₋₄alkoxy, C₁₋₄alkoxy-carbonyl, C₁₋₄alkoxy-carbonyl-amino, C₁₋₄alkoxy-carbonyl-amino-C₁₋₄alkoxy, C₂₋₄alkenyl, C₂₋₄alkenyl-amino-carbonyl, C₃₋₇cycloalkyl, C₃₋₇cycloalkyl-C₁₋₄alkoxy, C₃₋₇cycloalkenyl, heteroaryl, heteroaryl-C₁₋₄alkyl, heteroaryl-C₁₋₄alkyl-amino, heteroaryl-C₁₋₄alkyl-amino-carbonyl, heteroaryl-C₁₋₄alkyl-carbonyl-amino, heteroaryl-C₁₋₄alkyl-amino-carbonyl-C₁₋₄alkyl, heteroaryl-C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkyl, heterocyclyl, heterocyclyl-C₁₋₄alkyl, phenyl, or phenyl-C₁₋₄alkoxy;

- R₄ is independently selected from halogen, C₁₋₄alkyl, hydroxyl-C₁₋₄alkyl, amino, C₁₋₄alkyl-amino, (C₁₋₄alkyl)₂-amino or hydroxyl-C₁₋₄alkyl-amino; and

R₅ is hydrogen, C₁₋₄alkyl, or hydroxyl-C₁₋₄alkyl;

wherein a form of the compound is selected from the group consisting of a prodrug, salt, hydrate, solvate, clathrate, isotopologue, racemate, enantiomer, diastereomer, stereoisomer, polymorph and tautomer form thereof.

An embodiment of the present description further relates to a method or use of a compound for treating or ameliorating HD in a subject in need thereof comprising administering to the subject an effective amount of a compound of Formula (I) selected from a compound of Formula (Ia) and Formula (Ib):



5 or a form thereof, wherein

X is CH₂, CH(C₁₋₄alkyl), C(C₁₋₄alkyl)₂, CH=CH, O, NR₅, or a bond;

A is aryl, heteroaryl, heterocyclyl, or C₉₋₁₀cycloalkyl,

wherein aryl is selected from phenyl and naphthyl, each optionally substituted with 1, 2, 3, or 4 substituents each selected from R₁,

10 wherein heteroaryl is a saturated monocyclic, bicyclic or tricyclic ring system having 1, 2, or 3 heteroatom ring members independently selected from N, O, or S, each optionally substituted with 1, 2, 3, 4, or 5 substituents each selected from R₁,

wherein heterocyclyl is a saturated or partially unsaturated monocyclic, bicyclic or tricyclic ring system having 1, 2, or 3 heteroatom ring members independently selected from N, O, or

15 S, each optionally substituted with 1, 2, 3, 4, or 5 substituents each selected from R₂, and wherein C₉₋₁₀cycloalkyl is a saturated or partially unsaturated bicyclic ring system optionally substituted with 1, 2, 3, 4, or 5 substituents each selected from R₂;

B is heterocyclyl,

20 wherein heterocyclyl is a saturated or partially unsaturated monocyclic, bicyclic or polycyclic ring system having 1, 2, or 3 heteroatom ring members independently selected from N, O, or S, each optionally substituted with 1, 2, 3, 4, or 5 substituents each selected from R₄;

R₁ is halogen, hydroxyl, cyano, C₁₋₄alkyl, halo-C₁₋₄alkyl, amino, C₁₋₄alkyl-amino,

(C₁₋₄alkyl)₂-amino, amino-C₁₋₄alkyl, C₁₋₄alkyl-amino-C₁₋₄alkyl,

(C₁₋₄alkyl)₂-amino-C₁₋₄alkyl, amino-carbonyl, C₁₋₄alkyl-amino-carbonyl,

25 C₁₋₄alkyl-amino-carbonyl-C₁₋₄alkyl, C₁₋₄alkyl-carbonyl-amino, C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkyl, hydroxyl-C₁₋₄alkyl, C₁₋₄alkyl-carbonyl, C₁₋₄alkoxy, halo-C₁₋₄alkoxy, amino-C₁₋₄alkoxy, hydroxyl-C₁₋₄alkoxy, C₁₋₄alkyl-C₁₋₄alkoxy, C₁₋₄alkyl-amino-C₁₋₄alkoxy, (C₁₋₄alkyl)₂-amino-C₁₋₄alkoxy, C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkoxy, C₁₋₄alkoxy-

- C₁₋₄alkoxy, C₁₋₄alkoxy-carbonyl, C₁₋₄alkoxy-carbonyl-amino, C₁₋₄alkoxy-carbonyl-
 amino-C₁₋₄alkoxy, C₂₋₄alkenyl, C₂₋₄alkenyl-amino-carbonyl, C₃₋₇cycloalkyl,
 C₃₋₇cycloalkyl-C₁₋₄alkoxy, C₃₋₇cycloalkenyl, heteroaryl, heteroaryl-C₁₋₄alkyl,
 heteroaryl-C₁₋₄alkyl-amino, heteroaryl-C₁₋₄alkyl-amino-carbonyl,
 5 heteroaryl-C₁₋₄alkyl-carbonyl-amino, heteroaryl-C₁₋₄alkyl-amino-carbonyl-C₁₋₄alkyl,
 heteroaryl-C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkyl, heterocyclyl, heterocyclyl-C₁₋₄alkyl,
 heterocyclyl-C₁₋₄alkoxy, phenyl, or phenyl-C₁₋₄alkoxy,
 wherein heteroaryl is a saturated monocyclic or bicyclic ring system having 1, 2, or 3 heteroatom
 ring members selected from N, O, and S,
 10 wherein heterocyclyl is a saturated or partially unsaturated monocyclic or bicyclic ring system
 having 1, 2, or 3 heteroatom ring members selected from N, O, and S, and
 wherein each instance of phenyl, heteroaryl or heterocyclyl is optionally substituted with 1, or 2
 substituents each selected from R₃;
 R₂ is halogen, hydroxyl, cyano, oxo, hydroxyl-imino, C₁₋₄alkyl, halo-C₁₋₄alkyl, amino,
 15 C₁₋₄alkyl-amino, (C₁₋₄alkyl)₂-amino, amino-C₁₋₄alkyl, amino-carbonyl,
 hydroxyl-C₁₋₄alkyl, C₁₋₄alkoxy, C₁₋₄alkoxy-carbonyl, C₂₋₄alkenyl, C₃₋₇cycloalkyl, or
 heterocyclyl-C₁₋₄alkyl;
 R₃ is halogen, hydroxyl, nitro, oxo, hydroxyl-imino, C₁₋₄alkyl, amino, C₁₋₄alkyl-amino,
 (C₁₋₄alkyl)₂-amino, amino-C₁₋₄alkyl, C₁₋₄alkyl-amino-C₁₋₄alkyl,
 20 (C₁₋₄alkyl)₂-amino-C₁₋₄alkyl, amino-carbonyl, C₁₋₄alkyl-amino-carbonyl,
 C₁₋₄alkyl-amino-carbonyl-C₁₋₄alkyl, C₁₋₄alkyl-carbonyl-amino, C₁₋₄alkyl-carbonyl-amino-
 C₁₋₄alkyl, hydroxyl-C₁₋₄alkyl, C₁₋₄alkyl-carbonyl, C₁₋₄alkoxy, halo-C₁₋₄alkoxy, amino-
 C₁₋₄alkoxy, hydroxyl-C₁₋₄alkoxy, C₁₋₄alkyl-C₁₋₄alkoxy, C₁₋₄alkyl-amino-C₁₋₄alkoxy,
 (C₁₋₄alkyl)₂-amino-C₁₋₄alkoxy, C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkoxy, C₁₋₄alkoxy-
 25 C₁₋₄alkoxy, C₁₋₄alkoxy-carbonyl, C₁₋₄alkoxy-carbonyl-amino, C₁₋₄alkoxy-carbonyl-
 amino-C₁₋₄alkoxy, C₂₋₄alkenyl, C₂₋₄alkenyl-amino-carbonyl, C₃₋₇cycloalkyl,
 C₃₋₇cycloalkyl-C₁₋₄alkoxy, C₃₋₇cycloalkenyl, heteroaryl, heteroaryl-C₁₋₄alkyl,
 heteroaryl-C₁₋₄alkyl-amino, heteroaryl-C₁₋₄alkyl-amino-carbonyl,
 heteroaryl-C₁₋₄alkyl-carbonyl-amino, heteroaryl-C₁₋₄alkyl-amino-carbonyl-C₁₋₄alkyl,
 30 heteroaryl-C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkyl, heterocyclyl, heterocyclyl-C₁₋₄alkyl,
 phenyl, or phenyl-C₁₋₄alkoxy;


R₄ is independently selected from halogen, C₁₋₄alkyl, hydroxyl-C₁₋₄alkyl, amino, C₁₋₄alkyl-amino, (C₁₋₄alkyl)₂-amino or hydroxyl-C₁₋₄alkyl-amino; and

R₅ is hydrogen, C₁₋₄alkyl, or hydroxyl-C₁₋₄alkyl;

wherein a form of the compound is selected from the group consisting of a prodrug, salt, hydrate, solvate, clathrate, isotopologue, racemate, enantiomer, diastereomer, stereoisomer, polymorph and tautomer form thereof.

Another embodiment of the present description further relates to methods for treating or ameliorating HD in a subject in need thereof comprising administering to the subject an effective amount of a compound of Formula (I) selected from a compound of Formula (Ia) and Formula

(Ib):




or a form thereof, wherein


X is O, NH, N(CH₃) or a bond;


A is aryl, heteroaryl or heterocyclyl,


wherein aryl is selected from the group consisting of




wherein heteroaryl is selected from the group consisting of









wherein heterocyclyl is selected from the group consisting of



B is heterocyclyl selected from the group consisting of





R_{1a} , R_{1b} and R_{1c} are each, where allowed by available valences, one or more substituents each selected from halogen, hydroxyl, cyano, C_{1-4} alkyl, halo- C_{1-4} alkyl, amino, C_{1-4} alkyl-amino, (C_{1-4} alkyl)₂-amino, amino- C_{1-4} alkyl, C_{1-4} alkyl-amino- C_{1-4} alkyl, (C_{1-4} alkyl)₂-amino- C_{1-4} alkyl, amino-carbonyl, C_{1-4} alkyl-amino-carbonyl, (C_{1-4} alkyl)₂-amino-carbonyl, C_{1-4} alkyl-amino-carbonyl- C_{1-4} alkyl, (C_{1-4} alkyl)₂-amino-carbonyl- C_{1-4} alkyl, C_{1-4} alkyl-carbonyl-amino, C_{1-4} alkyl-carbonyl-amino- C_{1-4} alkyl, hydroxyl- C_{1-4} alkyl, C_{1-4} alkyl-carbonyl, C_{1-4} alkoxy, halo- C_{1-4} alkoxy, amino- C_{1-4} alkoxy, hydroxyl- C_{1-4} alkoxy, C_{1-4} alkyl- C_{1-4} alkoxy, C_{1-4} alkyl-amino- C_{1-4} alkoxy, (C_{1-4} alkyl)₂-amino- C_{1-4} alkoxy, C_{1-4} alkyl-carbonyl-amino- C_{1-4} alkoxy, C_{1-4} alkoxy- C_{1-4} alkoxy, C_{1-4} alkoxy-carbonyl, C_{1-4} alkoxy-carbonyl-amino, C_{1-4} alkoxy-carbonyl-amino- C_{1-4} alkoxy, C_{2-4} alkenyl, C_{2-4} alkenyl-amino-carbonyl, C_{3-7} cycloalkyl, C_{3-7} cycloalkyl- C_{1-4} alkoxy, C_{3-7} cycloalkenyl, heteroaryl, heteroaryl- C_{1-4} alkyl, heteroaryl- C_{1-4} alkyl-amino, heteroaryl- C_{1-4} alkyl-amino-carbonyl, heteroaryl- C_{1-4} alkyl-carbonyl-amino, heteroaryl- C_{1-4} alkyl-amino-carbonyl- C_{1-4} alkyl, heteroaryl- C_{1-4} alkyl-carbonyl-amino- C_{1-4} alkyl, heterocyclyl, heterocyclyl- C_{1-4} alkyl, heterocyclyl- C_{1-4} alkoxy, phenyl, or phenyl- C_{1-4} alkoxy,

wherein heteroaryl is a saturated monocyclic or bicyclic ring system having 1, 2, or 3 heteroatom ring members selected from N, O, and S,





wherein heterocyclyl is a saturated or partially unsaturated monocyclic or bicyclic ring system having 1, 2, or 3 heteroatom ring members selected from N, O, and S, and

wherein each instance of phenyl, heteroaryl or heterocyclyl is optionally substituted with 1, or 2 substituents each selected from R_3 ;

R_{2a} , R_{2b} and R_{2c} are each, where allowed by available valences, one or more substituents each selected from halogen, hydroxyl, cyano, oxo, hydroxyl-imino, C_{1-4} alkyl, halo- C_{1-4} alkyl, amino, C_{1-4} alkyl-amino, (C_{1-4} alkyl)₂-amino, amino- C_{1-4} alkyl, C_{1-4} alkyl-amino- C_{1-4} alkyl,

- (C₁₋₄alkyl)₂-amino-C₁₋₄alkyl, amino-carbonyl, hydroxyl-C₁₋₄alkyl, C₁₋₄alkoxy, C₁₋₄alkoxy-carbonyl, C₂₋₄alkenyl, C₃₋₇cycloalkyl, or heterocyclyl-C₁₋₄alkyl, wherein heterocyclyl is a saturated or partially unsaturated monocyclic or bicyclic ring system having 1, 2, or 3 heteroatom ring members selected from N, O, and S, and
- 5 wherein each instance of heterocyclyl is optionally substituted with 1, or 2 substituents each selected from R₃;
- R₃ is halogen, hydroxyl, nitro, oxo, hydroxyl-imino, C₁₋₄alkyl, amino, C₁₋₄alkyl-amino, (C₁₋₄alkyl)₂-amino, amino-C₁₋₄alkyl, C₁₋₄alkyl-amino-C₁₋₄alkyl, (C₁₋₄alkyl)₂-amino-C₁₋₄alkyl, amino-carbonyl, C₁₋₄alkyl-amino-carbonyl,
- 10 (C₁₋₄alkyl)₂-amino-carbonyl, C₁₋₄alkyl-amino-carbonyl-C₁₋₄alkyl, (C₁₋₄alkyl)₂-amino-carbonyl-C₁₋₄alkyl, C₁₋₄alkyl-carbonyl-amino, C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkyl, hydroxyl-C₁₋₄alkyl, C₁₋₄alkyl-carbonyl, C₁₋₄alkoxy, halo-C₁₋₄alkoxy, amino-C₁₋₄alkoxy, hydroxyl-C₁₋₄alkoxy, C₁₋₄alkyl-C₁₋₄alkoxy, C₁₋₄alkyl-amino-C₁₋₄alkoxy, (C₁₋₄alkyl)₂-amino-C₁₋₄alkoxy, C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkoxy, C₁₋₄alkoxy-
- 15 C₁₋₄alkoxy, C₁₋₄alkoxy-carbonyl, C₁₋₄alkoxy-carbonyl-amino, C₁₋₄alkoxy-carbonyl-amino-C₁₋₄alkoxy, C₂₋₄alkenyl, C₂₋₄alkenyl-amino-carbonyl, C₃₋₇cycloalkyl, C₃₋₇cycloalkyl-C₁₋₄alkoxy, C₃₋₇cycloalkenyl, heteroaryl, heteroaryl-C₁₋₄alkyl, heteroaryl-C₁₋₄alkyl-amino, heteroaryl-C₁₋₄alkyl-amino-carbonyl, heteroaryl-C₁₋₄alkyl-carbonyl-amino, heteroaryl-C₁₋₄alkyl-amino-carbonyl-C₁₋₄alkyl,
- 20 heteroaryl-C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkyl, heterocyclyl, heterocyclyl-C₁₋₄alkyl, phenyl, or phenyl-C₁₋₄alkoxy; and
- R_{4a}, R_{4b}, R_{4c}, R_{4d}, R_{4e}, R_{4f} and R_{4g} are independently selected from halogen, C₁₋₄alkyl, hydroxyl-C₁₋₄alkyl, amino, C₁₋₄alkyl-amino, (C₁₋₄alkyl)₂-amino or hydroxyl-C₁₋₄alkyl-amino;
- 25 wherein a form of the compound is selected from the group consisting of a prodrug, salt, hydrate, solvate, clathrate, isotopologue, racemate, enantiomer, diastereomer, stereoisomer, polymorph and tautomer form thereof.

Another aspect of the present description relates to a compound of Formula (I) selected from a compound of Formula (Ia11), Formula (Ia15), Formula (Ia18) or Formula (Ib1):



or a form thereof, wherein (when present),

X is selected from O, NR₅, or a bond;

- 5 A is selected from phenyl, thiophenyl, indazolyl, pyridinyl, pyrimidinyl or phenoxy, wherein phenyl and phenoxy are each optionally substituted with 1, 2 or 3 substituents each selected from R_{1a}, wherein thiophenyl, indazolyl, pyridinyl, pyrimidinyl are each optionally substituted with 1 or 2 substituents each selected from R_{1a},
- 10 B is selected from 1H-pyrazolyl, piperidinyl, 1,2,3,6-tetrahydropyridinyl, (1R,5S)-8-azabicyclo[3.2.1]octyl, 8-azabicyclo[3.2.1]oct-2-enyl, 2,6-diazaspiro[3.4]octyl or 2,7-diazaspiro[3.5]nonyl, each optionally substituted with 1 or 2 substituents each selected from R_{4a};

R_{1a} is selected from halogen, hydroxyl, C₁₋₄alkyl, halo-C₁₋₄alkyl, amino, C₁₋₄alkoxy, or

- 15 heteroaryl,

wherein heteroaryl is a saturated monocyclic or bicyclic ring system having 1, 2, or 3 heteroatom ring members selected from N, O, and S optionally substituted with 1 or 2 substituents each selected from R_{3a};

R_{3a} is selected from nitro or C₁₋₄alkyl; and,


- 20 R_{4a} is C₁₋₄alkyl;

R_{5a} is hydrogen, C₁₋₄alkyl, or hydroxyl-C₁₋₄alkyl;

wherein a form of the compound is selected from the group consisting of a prodrug, salt, hydrate, solvate, clathrate, isotopologue, racemate, enantiomer, diastereomer, stereoisomer, polymorph and tautomer form thereof.

- Another aspect of the present description relates to a compound of Formula (I) selected from a compound of Formula (Ia11), Formula (Ia15), Formula (Ia18) or Formula (Ib1):
- 5 or a form thereof, wherein (when present),
- R_{1a} is selected from fluoro, chloro, hydroxyl, methyl, difluoromethyl, amino, methoxy or 1H-pyrazolyl or 1H-imidazol-1-yl,
- wherein 1H-pyrazolyl is optionally substituted with 1 or 2 substituents each selected from R_{3a} ;
- 10 R_{3a} is selected from nitro or methyl or amino; and,
- R_{4a} is methyl or ethyl;
- R_{5a} is hydrogen or methyl;
- wherein a form of the compound is selected from the group consisting of a prodrug, salt, hydrate, solvate, clathrate, isotopologue, racemate, enantiomer, diastereomer, stereoisomer,
- 15 polymorph and tautomer form thereof.

- Another embodiment of the method of the present description includes the use of a compound of Formula (Ia) or a form thereof selected from a compound of Formula (Ia1) or a form thereof, wherein substituents R_{1a} , R_{1b} , and X, when present, are indicated in the table below with multiple substituents separated by a comma; and, "--" indicates that one or more R_{1a} , R_{1b} ,
- 20 and X substituents are not present:




(Ia1)

Cpd	R_{1a}	R_{1b}	X
1	--	--	NH
8	2-OH	--	N(CH ₃)
40	1-CH ₂ CH=CH ₂ , 2-OH	--	N(CH ₃)

Cpd	R _{1a}	R _{1b}	X
106	1-Br, 2-OH	7-OH	N(CH ₃)
107	1-Cl, 2-OH	7-OH	N(CH ₃)

Another embodiment of the method of the present description includes the use of a compound of Formula (Ia) or a form thereof selected from a compound of Formula (Ia2) or a form thereof, wherein substituents R_{1a}, R_{1b}, and R_{4a}, when present, are indicated in the table below with multiple substituents separated by a comma; and, "--" indicates that one or more R_{1a},

5 R_{1b}, and R_{4a} substituents are not present:




(Ia2)

Cpd	R _{1a}	R _{1b}	R _{4a}
13	--	--	--
207	--	7-OH	--
208	--	7-OH	CH ₃
210	2-CH ₃	7-OH	--
222	3-Cl	7-OH	--
223	3-Br	7-OH	--
224	3-CN	7-OH	--
225	3-(1-CH ₃ -1H-imidazol-4-yl)	7-OH	--
226	3-(1H-imidazol-1-yl)	7-OH	--
227	3-OH	7-OH	--
228	3-CH ₂ CH ₃	7-OH	--
229	3-CH(CH ₃) ₂	7-OH	--
232	2-CH ₃ , 4-OCH ₃ ,	7-OH	--
233	2-CH ₃ , 4-(pyrrolidin-1-yl)	7-OH	--

Cpd	R _{1a}	R _{1b}	R _{4a}
234	2-CH ₃ , 4-(morpholin-4-yl)	7-OH	--
235	2-CH ₃ , 4-N(CH ₃) ₂	7-OH	--
236	2-CH ₃ , 4-OCH ₂ CH ₃	7-OH	--
237	2-CH ₃ , 4-(1-CH ₃ -1H-pyrazol-4-yl)	7-OH	--
240	3-(tetrahydro-2H-pyran-4-yl)	7-OH	--
249	4-OCH ₃	7-OH	--
250	2-CH ₃ , 4-(azetidin-1-yl)	7-OH	--
251	2-CH ₃ , 4-CN	7-OH	--
252	2-CH ₃ , 4-cyclopropyl	7-OH	--
253	2-CH ₃ , 4-(3,6-dihydro-2H-pyran-4-yl)	7-OH	--
254	2-CH ₃ , 4-(tetrahydro-2H-pyran-4-yl)	7-OH	--
255	2-CH ₃ , 4-(oxetan-3-yl)	7-OH	--
256	4-N(CH ₃) ₂	7-OH	--
262	2-CN	7-OH	--
265	2-C(O)NH ₂	7-OH	--
293	3-Cl	7-OH	--
294	3-CH(CH ₃) ₂	7-OH	--
296	2-CH ₃ , 4-Cl	7-OH	--


Another embodiment of the method of the present description includes the use of a compound of Formula (Ia) or a form thereof selected from a compound of Formula (Ia3) or a form thereof, wherein substituents R_{1a} , R_{1b} and X, when present, are indicated in the table below with multiple substituents separated by a comma; and, "--" indicates that one or more R_{1a} , R_{1b} and X substituents are not present:



(Ia3)

Cpd	R_{1a}	R_{1b}	X
11	--	--	O
15	--	--	N(CH ₃)
218	--	7-OH	N(CH ₃)
261	1-CN	7-OH	N(CH ₃)
272	1-CH ₃	7-OH	N(CH ₃)
275	1-CN, 3-CH ₃	7-OH	N(CH ₃)
292	3-(OCH ₂ -phenyl)	--	N(CH ₃)

Another embodiment of the method of the present description includes the use of a compound of Formula (Ia) or a form thereof selected from a compound of Formula (Ia4) or a form thereof, wherein substituents X, R_{1a} , R_{1b} and R_{4a} , when present, are indicated in the table below; and, "--" indicates that one or more X, R_{1a} , R_{1b} and R_{4a} substituents are not present:




(Ia4)

Cpd	R_{1a}	R_{1b}	X	R_{4a}
10	--	--	O	--

Cpd	R _{1a}	R _{1b}	X	R _{4a}
14	--	--	N(CH ₃)	--
159	1-(OCH ₂ -phenyl)	--	N(CH ₃)	--
211	--	6-OH	N(CH ₃)	CH ₃
212	--	6-OH	N(CH ₃)	--
213	--	6-OH	O	--
215	1-cyclopropyl	6-OH	N(CH ₃)	--
216	1-OH	6-OH	N(CH ₃)	--
217	1-CN	6-OH	N(CH ₃)	--
264	1-C(O)NH ₂	6-OH	N(CH ₃)	--
273	1-CH ₃	6-OH	N(CH ₃)	--
274	1,3-(CH ₃) ₂	6-OH	N(CH ₃)	--
276	1-NH ₂	6-OH	N(CH ₃)	--
283	1-OCH ₂ CH ₃	6-OH	N(CH ₃)	--
284	1-OH	6-OH	O	--
285	3-phenyl	6-OH	N(CH ₃)	--
286	3-CH ₃	6-OH	N(CH ₃)	--
287	3-cyclopropyl	6-OH	N(CH ₃)	--
288	3-CH(CH ₃) ₂	6-OH	N(CH ₃)	--
289	3-(CH ₂) ₂ CH ₃	6-OH	N(CH ₃)	--
290	3-CH(CH ₃) ₂	6-OH	O	--


Another embodiment of the method of the present description includes the use of a compound of Formula (Ia) or a form thereof selected from a compound of Formula (Ia5) or a form thereof, wherein substituents R_{1a} and R_{1b}, when present, are indicated in the table below with multiple substituents separated by a comma; and, "--" indicates that one or more R_{1a} and R_{1b} substituents are not present:



(Ia5)

Cpd	R _{1a}	R _{1b}
12	--	--
220	--	6-OH
221	2-CH ₃	6-OH
238	4-OCH ₃	6-OH
241	3-Cl	6-OH
242	3-Br	6-OH
243	3-CH ₃	6-OH
244	3-CH ₃	5-Br, 6-OH
263	2-CN	6-OH
266	2-C(O)NH ₂	6-OH
267	2-CO ₂ CH ₃	6-OH
297	4-Cl	6-OH
300	--	6-OH


Another embodiment of the method of the present description includes the use of a compound of Formula (Ia) or a form thereof selected from a compound of Formula (Ia6) or a form thereof, wherein substituents R_{1a} , when present, are indicated in the table below; and, "--" indicates that one or more R_{1a} substituents are not present:



(Ia6)

Cpd	R_{1a}
239	--
246	2,3-(CH ₃) ₂
247	2-CH ₃
248	3-CH ₃


Another embodiment of the method of the present description includes the use of a compound of Formula (Ia) or a form thereof selected from a compound of Formula (Ia7) or a form thereof, wherein substituents R_{1a} , when present, are indicated in the table below; and, "--" indicates that one or more R_{1a} substituents are not present:



(Ia7)


Cpd	R_{1a}
258	--
260	2-CH ₃

Another embodiment of the method of the present description includes the use of a compound of Formula (Ia) or a form thereof selected from a compound of Formula (Ia8) or a form thereof, wherein substituents R_{1a} and B, when present, are indicated in the table below; and, "--" indicates that one or more R_{1a} and B substituents are not present:




Cpd	R_{1a}	B
209	--	6-((3aR,6aS)-5-CH ₃ -hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)
269	2-CN	piperazin-1-yl

Another embodiment of the method of the present description includes the use of a compound of Formula (Ia) or a form thereof selected from a compound of Formula (Ia9) or a form thereof, wherein substituents R_{1a} and B, when present, are indicated in the table below; and, "--" indicates that one or more R_{1a} and B substituents are not present:



Cpd	R_{1a}	B
214	--	6-((3aR,6aS)-5-CH ₃ -hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)
270	--	piperazin-1-yl
291	3-CH ₃	piperazin-1-yl


Another embodiment of the method of the present description includes the use of a compound of Formula (Ia) or a form thereof selected from a compound of Formula (Ia10) or a form thereof, wherein substituents R_{1a} and B, when present, are indicated in the table below; and, "--" indicates that one or more R_{1a} and B substituents are not present:



(Ia10)

Cpd	R_{1a}	B
268	2-CN	piperazin-1-yl
271	--	1,2,3,6-tetrahydropyridin-4-yl

Another embodiment of the method of the present description includes the use of a compound of Formula (Ia) or a form thereof selected from a compound of Formula (Ia11) or a form thereof, wherein substituents A, X and R_{4a} , when present, are indicated in the table below; and, "--" indicates that one or more A, X and R_{4a} substituents are not present:



(Ia11)


Cpd	A	X	R_{4a}
2	benzo[b]thiophen-2-yl	N(CH ₃)	--
4	5-CN-benzo[b]thiophen-2-yl	N(CH ₃)	--
5	quinolin-3-yl	NH	--
6	benzo[b]thiophen-2-yl	O	--
9	benzo[b]thiophen-2-yl	NH	--
16	imidazo[1,2-a]pyridin-6-yl	N(CH ₃)	--
17	6-phenyl-pyridin-3-yl	N(CH ₃)	--
18	6-(1H-pyrrol-1-yl)-pyridin-3-yl	N(CH ₃)	--
19	6-(1H-pyrazol-1-yl)-pyridin-3-yl	N(CH ₃)	--

Cpd	A	X	R _{4a}
20	quinoxalin-2-yl	N(CH ₃)	--
21	quinolin-3-yl	N(CH ₃)	--
22	phthalazin-6-yl	N(CH ₃)	--
23	benzo[c][1,2,5]oxadiazol-5-yl	NH	--
24	benzo[d]thiazol-5-yl	NH	--
25	2-CH ₃ -benzo[d]oxazol-6-yl	NH	--
30	2-(4-CN-phenol)	N(CH ₃)	--
32	2-(4-CF ₃ -phenol)	N(CH ₃)	--
33	6-(2-F-phenol)	N(CH ₃)	--
34	2-[3,5-(OCH ₃) ₂ -phenol]	N(CH ₃)	--
35	2-[4,5-(OCH ₃) ₂ -phenol]	N(CH ₃)	--
37	2-(4,5-F ₂ -phenol)	N(CH ₃)	--
41	benzo[b]thiophen-2-yl	NH	CH ₃
53	2-[4-(1H-pyrazol-1-yl)-phenol]	N(CH ₃)	--
115	2-[3-OH-5-(1H-pyrazol-4-yl)-phenol]	N(CH ₃)	--
116	2-[3-OCH ₃ -5-(1H-pyrazol-4-yl)-phenol]	N(CH ₃)	--
117	2-[5-(1H-pyrazol-4-yl)-3-OCF ₃ -phenol]	NH	--
118	2-[5-(1-CH ₃ -1H-pyrazol-4-yl)-3-OCF ₃ -phenol]	N(CH ₃)	--
119	2-[5-(1H-pyrazol-4-yl)-3-OCF ₃ -phenol]	N(CH ₃)	--
120	2-[5-(1-CH ₃ -pyridin-2(1H)-one)-3-OCF ₃ -phenol]	N(CH ₃)	--
121	2-[3-OCH ₃ -5-(1-CH ₃ -1H-pyrazol-4-yl)-phenol]	N(CH ₃)	--
122	2-[3-OCH ₃ -5-(5,6,7,8-tetrahydroimidazo-[1,2-a]pyridin-3-yl)-phenol]	N(CH ₃)	--
123	2-[3-OCH ₃ -5-(pyridin-3-yl)-phenol]	N(CH ₃)	--
124	2-[3-OCH ₃ -5-(1-cyclopentyl-1H-pyrazol-4-yl)-phenol]	N(CH ₃)	--
125	2-[5-(3-OCH ₃ -phenyl)-3-OCH ₃ -phenol]	N(CH ₃)	--
126	2-[3-benzyloxy-5-(5-CH ₃ -oxazol-2-yl)-phenol]	N(CH ₃)	--
127	2-[3-OCH ₂ CH ₃ -5-(5-CH ₃ -oxazol-2-yl)-phenol]	N(CH ₃)	--
128	2-[3-(OCH ₂ -cyclopropyl)-5-(5-CH ₃ -oxazol-2-yl)-phenol]	N(CH ₃)	--
129	5-(2-CH ₃ -1H-benzo[d]imidazol-6-ol)	N(CH ₃)	--
134	2-[4-(1H-pyrazol-4-yl)-phenol]	N(CH ₃)	--
135	2-[4-(4,5,6,7-tetrahydropyrazolo[1,5-a]pyridin-3-yl)-phenol]	N(CH ₃)	--

Cpd	A	X	R _{4a}
136	2-[4-(4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-3-yl)-phenol]	N(CH ₃)	--
137	2-[4-(1H-indol-2-yl)-phenol]	N(CH ₃)	--
138	2-[4-(cyclopent-1-en-1-yl)-phenol]	N(CH ₃)	--
139	2-[4-(1H-pyrazol-3-yl)-phenol]	N(CH ₃)	--
140	2-[4-(2-OH-pyridin-4-yl)-phenol]	N(CH ₃)	--
141	2[4-(1-CH ₃ -pyridin-2(1H)-one)-phenol]	O	--
142	2-[4-(2-OH-pyridin-4-yl)-phenol]	O	--
144	2-[4-Cl-5-(1H-pyrazol-4-yl)-phenol]	N(CH ₃)	--
145	2-[4-F-5-(1H-pyrazol-4-yl)-phenol]	N(CH ₃)	--
146	2-[5-F-4-(1H-imidazol-4-yl)-phenol]	N(CH ₃)	--
147	2-[5-F-4-(1H-pyrazol-4-yl)-phenol]	N(CH ₃)	--
148	2-[5-F-(1H-pyrazol-5-yl)-phenol]	N(CH ₃)	--
149	6-OH-1-oxo-2,3-dihydro-1H-inden-5-yl	N(CH ₃)	--
150	6-(1,4-dihydroindeno[1,2-c]-1H-pyrazol-7-ol)	N(CH ₃)	--
151	6-OH-1-OH-imino-2,3-dihydro-1H-inden-5-yl	N(CH ₃)	--
152	6-OH-1-OH-2,3-dihydro-1H-inden-5-yl	N(CH ₃)	--
153	6-(2-NH ₂ -8H-indeno[1,2-d]thiazol-5-ol)	N(CH ₃)	--
154	9-(5,6-dihydroimidazo[5,1-a]isoquinolin-8-ol)	N(CH ₃)	--
155	2-{4-[C(O)NHCH ₂ -(1-CH ₃ -1H-pyrazol-4-yl)]-phenol}	N(CH ₃)	--
156	2-[4-(4-CH ₂ OH-1H-pyrazol-1-yl)-phenol]	N(CH ₃)	--
158	3-(OCH ₂ -phenyl)-isoquinolin-6-yl	N(CH ₃)	--
160	2-[3-F-5-(2-OCH ₃ -pyridin-4-yl)-phenol]	N(CH ₃)	--
161	4-[1-(4-pyridin-2(1H)-one)-3-F-5-OH-phenyl]	N(CH ₃)	--
162	4-{1-[4-(1-CH ₃ -pyridin-2(1H)-one)]-3-F-5-OH-phenyl}	N(CH ₃)	--
163	4-{1-[5-(1-CH ₃ -pyridin-2(1H)-one)]-3-F-5-OH-phenyl}	N(CH ₃)	--
164	2-[3-F-5-(1H-pyrazol-4-yl)-phenol]	O	--
165	2-(5-Cl-3-F-phenol)	N(CH ₃)	--
166	2-[3-F-5-(1H-pyrazol-4-yl)-phenol]	N(CH ₃)	--
167	2-[3-F-5-(1-CH ₃ -1H-pyrazol-4-yl)-phenol]	N(CH ₃)	--
219	8-(quinolin-7-ol)	N(CH ₃)	--
230	6-(7-OH-quinolin-2(1H)-one)	N(CH ₃)	--

Cpd	A	X	R _{4a}
231	6-(7-OH-1-CH ₃ -quinolin-2(1H)-one)	N(CH ₃)	--
245	7-(6-OH-1-CH ₃ -quinolin-4(1H)-one)	N(CH ₃)	--
257	6-(7-OH-quinazolin-4(1H)-one)	N(CH ₃)	--
259	6-(7-OH-1-CH ₃ -3,4-dihydroquinolin-2(1H)-one)	N(CH ₃)	--
277	7-OH-1,3-(CH ₃) ₂ -quinazolin-6-yl-2,4(1H,3H)-dione	N(CH ₃)	--
278	6-OH-benzo[d]oxazol-5-yl-2(3H)-one	N(CH ₃)	--
279	2-CH ₃ -6-OH-2H-indazol-5-yl	N(CH ₃)	--
280	1-CH ₃ -6-OH-1H-indazol-5-yl	N(CH ₃)	--
281	7-(6-OH-2-CH ₃ -isoquinolin-1(2H)-one)	N(CH ₃)	--
282	7-(6-OH-2-CH ₂ CH ₃ -isoquinolin-1(2H)-one)	O	--

Another embodiment of the present description includes a compound of Formula (Ia) or a form thereof selected from a compound of Formula (Ia11) or a form thereof, wherein substituents A, X and R_{4a}, when present, are indicated in the table below; and, "--" indicates that one or more A, X and R_{4a} substituents are not present:




(Ia11)

Cpd	A	X
420	2-OCH ₃ -4-(4-NO ₂ -1H-pyrazol-1-yl)phenyl	N(CH ₃)
428	2,5-F ₂ -4-(1H-pyrazol-4-yl)phenyl	N(CH ₃)
430	2,3-F ₂ -4-(1H-pyrazol-4-yl)phenyl	N(CH ₃)
431	2,5-F ₂ -4-(1H-pyrazol-4-yl)phenyl	O
434	2-OCH ₃ -4-(1H-pyrazol-4-yl)phenyl	N(CH ₃)
435	4-(1H-pyrazol-4-yl)phenyl	O
437	2-F-4-(1H-pyrazol-4-yl)phenyl	O
438	4-(1-CH ₃ -1H-pyrazol-4-yl)thiophen-2-yl	O
440	2-F-4-OH-phenyl	N(CH ₃)

Cpd	A	X
442	2-CH ₃ -2H-indazol-5-yl	N(CH ₃)
443	2-CH ₃ -2H-indazol-5-yl	O
444	4-Cl-2-OCH ₃ -phenyl	O
445	2-CH ₃ -pyrazolo[1,5-a]pyridin-3-yl	N(CH ₃)
446	imidazo[1,2-a]pyridin-6-yl	O
447	2-OCH ₃ -4-(1H-pyrazol-1-yl)phenyl	O
448	5-(1H-pyrazol-4-yl)thiophen-2-yl	O
449	5-(1-CH ₃ -1H-pyrazol-4-yl)thiophen-2-yl	O
450	4-(1H-pyrazol-4-yl)thiophen-2-yl	O
451	2-OH-4-[3,5-(CH ₃) ₂ -1H-pyrazol-4-yl]phenyl	O
452	2-F-4-(1H-pyrazol-4-yl)phenyl	N(CH ₃)
453	2-OCH ₃ -4-OH-phenyl	O
454	2-OCH ₃ -4-(4-NO ₂ -1H-pyrazol-1-yl)phenyl	O
455	2,4-(OH) ₂ -phenyl	O
456	2-Cl-4-(1H-pyrazol-4-yl)phenyl	N(CH ₃)
457	5-amino-2-(1H-pyrazol-4-yl)pyrimidin-4-yl	O
458	2,6-F ₂ -4-(1H-pyrazol-4-yl)phenyl	O
464	2-(CHF ₂)-4-(1H-pyrazol-4-yl)phenyl	O
465	2-(CHF ₂)-4-(1H-pyrazol-4-yl)phenyl	N(CH ₃)

Another embodiment of the method of the present description includes the use of a compound of Formula (Ia) or a form thereof selected from a compound of Formula (Ia11) or a form thereof, wherein substituents A, X and R_{4a}, when present, are indicated in the table below; and, "--" indicates that one or more A, X and R_{4a} substituents are not present:




(Ia11)

Cpd	A	X
420	2-OCH ₃ -4-(4-NO ₂ -1H-pyrazol-1-yl)phenyl	N(CH ₃)
428	2,5-F ₂ -4-(1H-pyrazol-4-yl)phenyl	N(CH ₃)
430	2,3-F ₂ -4-(1H-pyrazol-4-yl)phenyl	N(CH ₃)
431	2,5-F ₂ -4-(1H-pyrazol-4-yl)phenyl	O
434	2-OCH ₃ -4-(1H-pyrazol-4-yl)phenyl	N(CH ₃)
435	4-(1H-pyrazol-4-yl)phenyl	O
437	2-F-4-(1H-pyrazol-4-yl)phenyl	O
438	4-(1-CH ₃ -1H-pyrazol-4-yl)thiophen-2-yl	O
440	2-F-4-OH-phenyl	N(CH ₃)
442	2-CH ₃ -2H-indazol-5-yl	N(CH ₃)
443	2-CH ₃ -2H-indazol-5-yl	O
444	4-Cl-2-OCH ₃ -phenyl	O
445	2-CH ₃ -pyrazolo[1,5-a]pyridin-3-yl	N(CH ₃)
446	imidazo[1,2-a]pyridin-6-yl	O
447	2-OCH ₃ -4-(1H-pyrazol-1-yl)phenyl	O
448	5-(1H-pyrazol-4-yl)thiophen-2-yl	O
449	5-(1-CH ₃ -1H-pyrazol-4-yl)thiophen-2-yl	O
450	4-(1H-pyrazol-4-yl)thiophen-2-yl	O
451	2-OH-4-[3,5-(CH ₃) ₂ -1H-pyrazol-4-yl]phenyl	O

Cpd	A	X
452	2-F-4-(1H-pyrazol-4-yl)phenyl	N(CH ₃)
453	2-OCH ₃ -4-OH-phenyl	O
454	2-OCH ₃ -4-(4-NO ₂ -1H-pyrazol-1-yl)phenyl	O
455	2,4-(OH) ₂ -phenyl	O
456	2-Cl-4-(1H-pyrazol-4-yl)phenyl	N(CH ₃)
457	5-amino-2-(1H-pyrazol-4-yl)pyrimidin-4-yl	O
458	2,6-F ₂ -4-(1H-pyrazol-4-yl)phenyl	O
464	2-(CHF ₂)-4-(1H-pyrazol-4-yl)phenyl	O
465	2-(CHF ₂)-4-(1H-pyrazol-4-yl)phenyl	N(CH ₃)


Another embodiment of the method of the present description includes the use of a compound of Formula (Ia) or a form thereof selected from a compound of Formula (Ia12) or a form thereof, wherein substituents X, R_{1a} and B, when present, are indicated in the table below; and, "--" indicates that one or more X, R_{1a} and B substituents are not present:



(Ia12)


Cpd	R _{1a}	X	B
66	H	NH	azetidin-3-yl
82	OH	--	piperazin-1-yl
85	H	--	1,2,3,6-tetrahydropyridin-4-yl
86	OH	--	1,2,3,6-tetrahydropyridin-4-yl
87	OH	--	2,2,6,6-tetramethyl-(1,2,3,6-tetrahydropyridin-4-yl)
88	OH	--	1-CH ₃ -(1,2,3,6-tetrahydropyridin-4-yl)
89	OH	--	piperidin-4-yl
99	H	CH ₂	piperidin-4-yl

Another embodiment of the method of the present description includes the use of a compound of Formula (Ia) or a form thereof selected from a compound of Formula (Ia13) or a form thereof, wherein substituents X, R_{1a} and R_{4a}, when present, are indicated in the table below; and, "--" indicates that one or more X, R_{1a} and R_{4a} substituents are not present:



Cpd	X	R _{1a}	R _{4a}
26	N(CH ₃)	H	--
28	NH	H	--
31	O	H	--
90	O	OH	--
91	N(CH ₃)	OH	--
92	NH	OH	--
93	N(CH ₃)	O(CH ₂) ₃ NHCO ₂ C(CH ₃) ₃	--
94	N(CH ₃)	O(CH ₂) ₃ NH ₂	--
95	N(CH ₃)	O(CH ₂) ₃ NHCO ₂ CH ₃	--
96	N(CH ₃)	O(CH ₂) ₃ OH	--
97	N(CH ₃)	O(CH ₂) ₃ OCH ₃	--
98	O	O(CH ₂) ₃ -morpholin-4-yl	--
103	N(CH ₃)	CN	--
104	N(CH ₃)	CH ₂ -1-piperidiny1	--
105	N(CH ₃)	CH ₂ -pyrrolidin-1-yl	--
108	N(CH ₃)	OCH ₃	--
109	N(CH ₃)	OCH ₃	CH ₃
110	N(CH ₃)	3,6-dihydro-2H-pyran-4-yl	--
111	N(CH ₃)	tetrahydro-2H-pyran-4-yl	--
112	N(CH ₃)	CHF ₂	--
113	N(CH ₃)	OC(CH ₃) ₂ (CH ₂) ₂ OH	--
114	N(CH ₃)	O(CH ₂) ₂ C(CH ₃) ₂ OH	--

Another embodiment of the method of the present description includes the use of a compound of Formula (Ia) or a form thereof selected from a compound of Formula (Ia14) or a form thereof, wherein substituents X and B, when present, are indicated in the table below; and, "--" indicates that one or more X and B substituents are not present:




(Ia14)

Cpd	X	B
55	O	piperidin-4-yl
56	O	(2S,4R,6R)-2,6-(CH ₃) ₂ -piperidin-4-yl
57	O	2,6-(CH ₃) ₂ -piperidin-4-yl
58	O	pyrrolidin-3-yl
59	O	2-CH ₃ -piperidin-4-yl
60	OCH ₂	1H-pyrrolidin-3-yl
61	O	3-F-piperidin-4-yl
65	--	piperazin-1-yl
67	NH	azetidin-3-yl
68	--	3,5-(CH ₃) ₂ -piperazin-1-yl
69	--	7-CH ₃ -2,7-diazaspiro[4.4]non-2-yl
70	--	[1,4]diazepan-1-yl
71	--	4-CH ₂ CH ₂ OH-piperazin-1-yl
72	--	2,7-diazaspiro[3.5]non-7-yl
73	--	2,7-diazaspiro[3.5]non-7-yl
74	--	3-CH ₂ OH-piperazin-1-yl
75	--	1,7-diazaspiro[4.4]non-7-yl
76	--	4-NH ₂ -4-CH ₃ -piperidin-1-yl
77	--	3-N(CH ₃) ₂ -piperidin-1-yl
79	--	3,3-(CH ₃) ₂ -piperazin-1-yl
80	--	7-CH ₂ CH ₂ OH-2,7-diazaspiro[4.4]-nonan-2-yl
83	--	1,2,3,6-tetrahydropyridin-4-yl
84	--	piperidin-4-yl

Cpd	X	B
102	O	(6S)-6-[(S)-CH(OH)CH ₃]-2,2-(CH ₃) ₂ -piperidin-4-yl
133	O	2,2-(CH ₃) ₂ -piperidin-4-yl

Another embodiment of the method of the present description includes the use of a compound of Formula (Ia) or a form thereof selected from a compound of Formula (Ia15) or a form thereof, wherein substituents X, R_{1a} and R_{4a}, when present, are indicated in the table below; and, "--" indicates that one or more X, R_{1a} and R_{4a} substituents are not present:




(Ia15)

Cpd	X	R _{1a}	R _{4a}
3	NH	H	--
7	N(CH ₃)	H	--
27	N(CH ₃)	Cl	CH ₃
29	NH	Cl	CH ₃
36	N(CH ₃)	OCH ₃	--
38	N(CH ₃)	F	--
39	N(CH ₃)	CN	--
42	N(CH ₃)	C(O)NHCH ₂ CH=CH ₂	--
43	N(CH ₃)	1H-pyrazol-1-yl	--
44	N(CH ₃)	5-CH ₃ -oxazol-2-yl	--
45	N(CH ₃)	4-CH ₂ OH-1H-pyrazole-1-yl	--
46	N(CH ₃)	1H-imidazole-1-yl	--
47	N(CH ₃)	4-NH ₂ -1H-pyrazol-1-yl	--
48	N(CH ₃)	1H-pyrazol-4-yl	--
49	N(CH ₃)	3-NH ₂ -1H-pyrazol-1-yl	--
50	N(CH ₃)	1-(CH ₂ CH ₂ -morpholin-4-yl)-1H-pyrazol-4-yl	--
51	N(CH ₃)	1-CH ₃ -1H-pyrazol-4-yl	--
52	N(CH ₃)	5-NH ₂ -1H-pyrazol-1-yl	--
54	N(CH ₂ CH ₂ OH)	1H-pyrazol-1-yl	--

Cpd	X	R _{1a}	R _{4a}
62	O	1H-pyrazol-1-yl	CH ₃
63	O	1H-pyrazol-1-yl	--
64	O	1H-pyrazol-4-yl	--
78	NH	1H-pyrazol-1-yl	CH ₃
100	CH ₂	1H-pyrazol-1-yl	--
130	N(CH ₃)	Cl	--
131	NH	1H-pyrazol-1-yl	--
132	NH	CN	--
143	N(CH ₃)	1H-indazol-7-yl	--
157	CH ₂	1H-pyrazol-4-yl	--
168	N(CH ₃)	5-OCH ₃ -pyridin-3-yl	--
169	N(CH ₃)	5-pyridin-2-ol	--
170	N(CH ₃)	4-pyridin-2-ol	--
171	N(CH ₃)	6-OCH ₃ -pyridin-3-yl	--
172	N(CH ₃)	5-(3-CF ₃ -pyridin-2-ol)	--
173	N(CH ₃)	5-(1-CH ₃ -pyridin-2(1H)-one)	--
174	N(CH ₃)	4-(1-CH ₃ -pyridin-2(1H)-one)	--
175	N(CH ₃)	2-OCH ₃ -pyridin-4-yl	--
176	O	4-pyridin-2-ol	--
177	N(CH ₃)	6-N(CH ₃) ₂ -pyridin-3-yl	--
178	O	4-(1-CH ₃ -pyridin-2(1H)-one)	--
179	N(CH ₃)	pyrimidin-5-yl	--
180	N(CH ₃)	5-pyridin-3-ol	--
181	N(CH ₃)	4-(1-cyclopropyl-pyridin-2(1H)-one)	--
182	N(CH ₃)	1,2,3,6-tetrahydropyridin-4-yl	--
183	N(CH ₃)	cyclopent-1-en-1-yl	--
184	N(CH ₃)	3,6-dihydro-2H-pyran-4-yl	--
185	N(CH ₃)	imidazo[1,5-a]pyridin-7-yl	--
186	N(CH ₃)	imidazo[1,2-a]pyridin-7-yl	--
187	N(CH ₃)	2-CH ₃ -pyridin-4-yl	--
188	N(CH ₃)	1H-imidazol-2-yl	--
189	N(CH ₃)	1H-imidazol-4-yl	--
190	N(CH ₃)	imidazo[1,2-a]pyrazin-3-yl	--


Cpd	X	R _{1a}	R _{4a}
191	N(CH ₃)	5,6,7,8-tetrahydroimidazo[1,2-a]pyrazin-3-yl	--
192	N(CH ₃)	4-CH ₃ -1H-imidazol-2-yl	--
193	N(CH ₃)	1-CH ₃ -1H-imidazol-4-yl	--
194	N(CH ₃)	1-CH ₃ -1H-imidazol-5-yl	--
195	N(CH ₃)	4-NO ₂ -1H-imidazol-2-yl	--
196	N(CH ₃)	2-CH ₃ -1H-imidazol-4-yl	--
197	N(CH ₃)	1,2-(CH ₃) ₂ -1H-imidazol-4-yl	--
198	N(CH ₃)	4-C(O)NH ₂ -1H-pyrazol-1-yl	--
206	N(CH ₃)	H	--

Another embodiment of the present description includes a compound of Formula (Ia) or a form thereof selected from a compound of Formula (Ia15) or a form thereof, wherein substituents X, R_{1a} and R_{4a}, when present, are indicated in the table below; and, "--" indicates that one or more X, R_{1a} and R_{4a} substituents are not present:



Cpd	X	R _{1a}	R _{4a}
413	NH	1H-pyrazol-4-yl	--
414	O	1-CH ₃ -1H-pyrazol-4-yl	--
416	N(CH ₃)	5-CH ₃ -1H-pyrazol-4-yl	--
417	O	1H-imidazol-1-yl	--
418	O	5-CH ₃ -1H-pyrazol-4-yl	--
419	N(CH ₃)	4-NO ₂ -1H-pyrazol-1-yl	--
421	O	4-NH ₂ -1H-pyrazol-1-yl	--
423	O	4-NO ₂ -1H-pyrazol-1-yl	--
460	N(CH ₃)	1H-pyrazol-4-yl	--
461	O	1H-pyrazol-4-yl	--


Another embodiment of the method of the present description includes the use of a compound of Formula (Ia) or a form thereof selected from a compound of Formula (Ia15) or a form thereof, wherein substituents X, R_{1a} and R_{4a}, when present, are indicated in the table below; and, "--" indicates that one or more X, R_{1a} and R_{4a} substituents are not present:



(Ia15)

Cpd	X	R _{1a}
413	NH	1H-pyrazol-4-yl
414	O	1-CH ₃ -1H-pyrazol-4-yl
416	N(CH ₃)	5-CH ₃ -1H-pyrazol-4-yl
417	O	1H-imidazol-1-yl
418	O	5-CH ₃ -1H-pyrazol-4-yl
419	N(CH ₃)	4-NO ₂ -1H-pyrazol-1-yl
421	O	4-NH ₂ -1H-pyrazol-1-yl
423	O	4-NO ₂ -1H-pyrazol-1-yl
460	N(CH ₃)	1H-pyrazol-4-yl
461	O	1H-pyrazol-4-yl


Another embodiment of the method of the present description includes the use of a compound of Formula (Ia) or a form thereof selected from a compound of Formula (Ia16) or a form thereof, wherein substituents R_{1a} and R_{4a} , when present, are indicated in the table below; and, "--" indicates that one or more R_{1a} and R_{4a} substituents are not present:



(Ia16)

Cpd	R_{1a}	R_{4a}
81	1H-pyrazol-1-yl	--
199	1H-pyrazol-4-yl	$(CH_2)_2OH$
200	1H-pyrazol-4-yl	--
201	1H-pyrazol-4-yl	CH_3
202	4-(1- CH_3 -pyridin-2(1H)-one)	CH_3
203	4-(1- CH_3 -pyridin-2(1H)-one)	CH_3


Another embodiment of the method of the present description includes the use of a compound of Formula (Ia) or a form thereof selected from a compound of Formula (Ia17) or a form thereof, wherein substituent R_{1a} , when present, is indicated in the table below; and, "--" indicates that one or more R_{1a} substituents are not present:



(Ia17)

Cpd	R_{1a}
204	1H-pyrazol-4-yl
205	4-(1- CH_3 -pyridin-2(1H)-one)


Another embodiment of the present description includes a compound of Formula (Ia) or a form thereof selected from a compound of Formula (Ia18) or a form thereof, wherein substituents X and B, when present, are indicated in the table below; and, "--" indicates that one or more X and B substituents are not present:



(Ia18)

Cpd	X	B
411	N(CH ₃)	(1 <i>R</i> ,5 <i>S</i>)-8-azabicyclo[3.2.1]oct-3-yl
412	NH	(1 <i>R</i> ,5 <i>S</i>)-8-azabicyclo[3.2.1]oct-3-yl
415	O	(1 <i>R</i> ,5 <i>S</i>)-8-azabicyclo[3.2.1]oct-3-yl
422	--	1-CH ₃ -(1,2,3,6-tetrahydropyridin-4-yl)
424	--	1,2,3,6-tetrahydropyridin-4-yl
425	--	1-CH ₃ CH ₂ -(1,2,3,6-tetrahydropyridin-4-yl)
426	N(CH ₃)	piperidin-4-yl
427	NH	piperidin-4-yl
429	--	8-azabicyclo[3.2.1]oct-2-en-3-yl
432	O	piperidin-4-yl
433	NH	(1 <i>R</i> ,5 <i>S</i>)-8-azabicyclo[3.2.1]oct-3-yl
436	O	2,6-(CH ₃) ₂ -piperidin-4-yl
439	--	2,7-diazaspiro[3.5]non-2-yl
441	O	2,6-(CH ₃) ₂ -piperidin-4-yl
459	--	2,6-diazaspiro[3.4]oct-2-yl


Another embodiment of the method of the present description includes the use of a compound of Formula (Ia) or a form thereof selected from a compound of Formula (Ia18) or a form thereof, wherein substituents X, R_{1a} and B, when present, are indicated in the table below; and, "--" indicates that one or more X, R_{1a} and B substituents are not present:



(Ia18)

Cpd	X	B
411	N(CH ₃)	(1 <i>R</i> ,5 <i>S</i>)-8-azabicyclo[3.2.1]oct-3-yl
412	NH	(1 <i>R</i> ,5 <i>S</i>)-8-azabicyclo[3.2.1]oct-3-yl
415	O	(1 <i>R</i> ,5 <i>S</i>)-8-azabicyclo[3.2.1]oct-3-yl
422	--	1-CH ₃ -(1,2,3,6-tetrahydropyridin-4-yl)
424	--	1,2,3,6-tetrahydropyridin-4-yl
425	--	1-CH ₃ CH ₂ -(1,2,3,6-tetrahydropyridin-4-yl)
426	N(CH ₃)	piperidin-4-yl
427	NH	piperidin-4-yl
429	--	8-azabicyclo[3.2.1]oct-2-en-3-yl
432	O	piperidin-4-yl
433	NH	(1 <i>R</i> ,5 <i>S</i>)-8-azabicyclo[3.2.1]oct-3-yl
436	O	2,6-(CH ₃) ₂ -piperidin-4-yl
439	--	2,7-diazaspiro[3.5]non-2-yl
441	O	2,6-(CH ₃) ₂ -piperidin-4-yl
459	--	2,6-diazaspiro[3.4]oct-2-yl

Another embodiment of the method of the present description includes the use of a compound of Formula (Ib) or a form thereof selected from a compound of Formula (Ib1) or a form thereof, wherein substituent A is indicated in the table below:



(Ib1)


5

Cpd	A
302	6-(naphthalen-2-yl)
320	6-(naphthalen-2,7-diyl)
331	7-OCH ₃ -quinolin-6-yl
332	7-OH-quinolin-6-yl
337	2-CN-7-OCH ₃ -quinolin-6-yl
355	3-F-5-(1H-pyrazol-4-yl)-pyridin-2-yl
364	2-(6-OCH ₃ -3,4-dihydroisoquinolin-1(2H)-one)
392	6-OH-1-oxo-2,3-dihydro-1H-inden-5-yl
401	3-(4-OCH ₃ -1-CH ₃ -quinolin-2(1H)-one)
402	3-(4-OH-1-CH ₃ -quinolin-2(1H)-one)
403	3-(quinolin-2(1H)-one)
404	3-(1-OCH ₃ -quinolin-2(1H)-one)
408	5-CN-benzo[b]thiophen-2-yl
409	3-Cl-benzo[b]thiophen-2-yl

Another embodiment of the present description includes a compound of Formula (Ib) or a form thereof selected from a compound of Formula (Ib1) or a form thereof, wherein substituent A is indicated in the table below:

Cpd	A
462	3-(1H-pyrazol-4-yl)phenoxy
463	4-(1H-pyrazol-4-yl)phenoxy


Another embodiment of the method of the present description includes the use of a compound of Formula (Ib) or a form thereof selected from a compound of Formula (Ib1) or a form thereof, wherein substituent A is indicated in the table below:



(Ib1)

Cpd	A
462	3-(1H-pyrazol-4-yl)phenoxy
463	4-(1H-pyrazol-4-yl)phenoxy


Another embodiment of the method of the present description includes the use of a compound of Formula (Ib) or a form thereof selected from a compound of Formula (Ib2) or a form thereof, wherein substituent A is indicated in the table below:



(Ib2)

Cpd	A
321	6-naphthalen-2,7-diol


Another embodiment of the method of the present description includes the use of a compound of Formula (Ib) or a form thereof selected from a compound of Formula (Ib3) or a form thereof, wherein substituents R_{1a}, R_{1b} and B, when present, are indicated in the table below; and, "--" indicates that one or more R_{1a}, R_{1b} and B substituents are not present:



(Ib3)

Cpd	R _{1a}	R _{1b}	B
329	1H-pyrazol-1-yl	OCH ₃	1,2,3,6-tetrahydropyridin-4-yl
330	1H-pyrazol-1-yl	OH	piperazin-1-yl
381	1H-pyrazol-1-yl	Cl	5-((3aR,6aR)-1-CH ₃ -hexahydropyrrolo[3,4-b]pyrrol-5(1H)-yl)
382	1H-pyrazol-1-yl	Cl	2-NHCH(CH ₃) ₂ -morpholin-4-yl
383	1H-pyrazol-1-yl	Cl	2-OCH ₃ -2,7-diazaspiro[4.5]decan-7-yl
385	1-CH ₃ -1H-pyrazol-4-yl	OCH ₃	5-((3aR,6aS)-5-CH ₃ -hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)
394	1-CH ₃ -1H-pyrazol-4-yl	OH	5-((3aR,6aS)-5-CH ₃ -hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)
406	1H-pyrazol-1-yl	Cl	2,7-diazaspiro[4.5]decan-2-yl
407	1H-pyrazol-1-yl	Cl	(3R)-(3-(R)-CH ₂ OH)-piperazin-1-yl

Another embodiment of the present description includes the use of a compound of Formula (Ib) or a form thereof selected from a compound of Formula (Ib4) or a form thereof, wherein substituents R_{1a}, R_{1b}, R_{1c}, R_{1d} (each representative of the scope of R₁) and X, when present, are indicated in the table below; and, "--" indicates that one or more R_{1a}, R_{1b}, R_{1c}, R_{1d} and X substituents are not present:



(Ib4)


Cpd	R _{1a}	R _{1b}	R _{1c}	R _{1d}	X
301	1H-pyrazol-1-yl	OCH ₃	H	H	N(CH ₃)
305	1H-pyrazol-1-yl	OCH ₃	H	H	N(CH ₃)
306	1-CH ₃ -1H-pyrazol-4-yl	OCH ₃	H	H	N(CH ₃)
307	1H-pyrazol-4-yl	OCH ₃	H	H	N(CH ₃)
308	4-(1-CH ₃ -pyridin-2(1H)-one)	OCH ₃	H	H	N(CH ₃)
309	5-pyridin-2-ol	OCH ₃	H	H	N(CH ₃)
310	5-(1-CH ₃ -pyridin-2(1H)-one)	OCH ₃	H	H	N(CH ₃)
311	1-CH ₃ -1H-pyrazol-4-yl	CH ₃	H	H	N(CH ₃)

Cpd	R _{1a}	R _{1b}	R _{1c}	R _{1d}	X
312	4-(1-CH ₃ -pyridin-2(1H)-one)	OCF ₃	H	H	N(CH ₃)
313	3,5-(CH ₃) ₂ -1H-pyrazol-4-yl	OCH ₃	H	H	N(CH ₃)
314	1-CH ₃ -1H-pyrazol-4-yl	CF ₃	H	H	N(CH ₃)
315	1-CH ₃ -1H-pyrazol-4-yl	OH	H	H	N(CH ₃)
316	1H-pyrazol-1-yl	OH	H	H	N(CH ₃)
317	5-(1-CH ₃ -pyridin-2(1H)-one)	OH	H	H	N(CH ₃)
318	4-(1-CH ₃ -pyridin-2(1H)-one)	OH	H	H	N(CH ₃)
319	5-pyridin-2-ol	OH	H	H	N(CH ₃)
324	H	OH	1H-pyrazol-1-yl	H	N(CH ₃)
325	1-CH ₃ -1H-pyrazol-4-yl	H	H	Cl	N(CH ₃)
326	1-CH ₃ -1H-pyrazol-4-yl	OH	H	Cl	N(CH ₃)
327	1-CH ₃ -1H-pyrazol-4-yl	H	H	Cl	N(CH ₃)
328	5-CH ₃ -oxazol-2-yl	OH	H	OCH ₃	N(CH ₃)
333	CN	OCH ₃	H	H	N(CH ₃)
334	CN	F	H	H	N(CH ₃)
335	CO ₂ CH ₃	F	H	H	N(CH ₃)
336	3-NHCH ₃ -1H-pyrazol-1-yl	OCH ₃	H	H	N(CH ₃)
338	4-(1-CH ₃ -pyridin-2(1H)-one)	OCH ₃	H	H	N(CH ₃)
339	4-(1-CH ₃ -pyridin-2(1H)-one)	Cl	H	H	N(CH ₃)
340	1H-pyrazol-4-yl	Cl	H	H	N(CH ₃)
341	4,5,6,7-tetrahydropyrazolo[1,5-a]pyridin-3-yl	Cl	H	H	N(CH ₃)
343	1-CH ₃ -1H-pyrazol-4-yl	Cl	H	H	O
344	6-OCH ₃ -pyridin-3-yl	Cl	H	H	N(CH ₃)
345	6-NH ₂ -pyridin-3-yl	F	H	H	N(CH ₃)
346	3-CH ₃ -1H-pyrazol-5-yl	F	H	H	N(CH ₃)
347	1H-pyrazol-5-yl	F	H	H	N(CH ₃)
348	1H-pyrazol-4-yl	H	F	F	N(CH ₃)
349	1H-pyrazol-5-yl	H	F	F	N(CH ₃)
350	1H-pyrazol-4-yl	F	F	H	N(CH ₃)
351	1H-pyrazol-5-yl	F	F	H	N(CH ₃)
352	1H-pyrazol-4-yl	F	H	F	N(CH ₃)
354	1H-pyrazol-4-yl	Cl	F	H	N(CH ₃)

Cpd	R _{1a}	R _{1b}	R _{1c}	R _{1d}	X
356	2-NH ₂ -pyrimidin-4-yl	Cl	H	H	N(CH ₃)
357	H	Cl	2-NH ₂ -pyrimidin-4-yl	H	N(CH ₃)
358	2,4-(CH ₃) ₂ -thiazol-5-yl	F	F	H	N(CH ₃)
359	2,4-(CH ₃) ₂ -thiazol-5-yl	H	F	F	N(CH ₃)
360	4-(1-CH ₃ -pyridin-2(1H)-one)	OH	H	OCF ₃	N(CH ₃)
361	1H-pyrazol-4-yl	OCH ₃	H	F	N(CH ₃)
363	1H-pyrazol-4-yl	OCH ₃	F	F	N(CH ₃)
365	1H-pyrazol-1-yl	Cl	H	H	N(CH ₃)
366	1H-1,2,3-triazol-1-yl	Cl	H	H	N(CH ₃)
367	2H-1,2,3-triazol-2-yl	Cl	H	H	N(CH ₃)
368	1H-1,2,4-triazol-1-yl	Cl	H	H	N(CH ₃)
369	3-NH ₂ -1H-pyrazol-1-yl	Cl	H	H	N(CH ₃)
371	1H-imidazol-1-yl	Cl	H	H	N(CH ₃)
372	1H-imidazol-1-yl	F	H	H	N(CH ₃)
373	1H-pyrazol-5-yl	OCH ₃	H	H	N(CH ₃)
374	2,4-(CH ₃) ₂ -thiazol-5-yl	OCH ₃	H	H	N(CH ₃)
375	pyridin-3-yl	OCH ₃	H	H	N(CH ₃)
376	1H-pyrazol-4-yl	F	H	H	N(CH ₃)
377	2-OCH ₃ -pyridin-4-yl	OCH ₃	H	H	N(CH ₃)
378	6-OCH ₃ -pyridin-3-yl	OCH ₃	H	H	N(CH ₃)
387	1H-pyrazol-1-yl	OH	H	H	N(CH ₃)
388	5-(pyridin-2(1H)-one)	Cl	H	H	N(CH ₃)
389	3-NHCH ₃ -1H-pyrazol-1-yl	OH	H	H	N(CH ₃)
390	1H-pyrazol-4-yl	OH	H	F	N(CH ₃)
391	1H-pyrazol-4-yl	OH	F	F	N(CH ₃)
393	1H-pyrazol-4-yl	OH	H	H	N(CH ₃)
397	1H-pyrazol-4-yl	OH	H	Cl	N(CH ₃)
398	1H-pyrazol-1-yl	OCH ₃	H	H	CH ₂
410	1H-pyrazol-4-yl	OCH ₃	H	H	N(CH ₃)

Another embodiment of the method of the present description includes the use of a compound of Formula (Ib) or a form thereof selected from a compound of Formula (Ib5) or a form thereof, wherein substituents R_{1a} , R_{1b} , R_{1c} , R_{1d} (each representative of the scope of R_1) and R_{4a} , when present, are indicated in the table below; and, "--" indicates that one or more R_{1a} , R_{1b} ,


5 R_{1c} , R_{1d} and R_{4a} substituents are not present:



(Ib5)

Cpd	R_{1a}	R_{1b}	R_{1c}	R_{1d}	R_{4a}
353	1H-pyrazol-4-yl	F	F	H	--
362	1H-pyrazol-4-yl	OCH ₃	H	F	CH ₃
370	1H-imidazol-1-yl	Cl	H	H	CH ₃
379	1-CH ₃ -1H-pyrazol-4-yl	Cl	H	H	CH ₃
380	1H-pyrazol-4-yl	Cl	H	H	CH ₃
384	1H-pyrazol-4-yl	F	H	H	CH ₃
396	1H-pyrazol-4-yl	F	H	OH	--
405	1H-pyrazol-4-yl	Cl	H	H	--


Another embodiment of the method of the present description includes the use of a compound of Formula (Ib) or a form thereof selected from a compound of Formula (Ib6) or a form thereof, wherein substituents R_{1a} , R_{1b} , R_{1c} and R_{1d} (each representative of the scope of R_1), when present, are indicated in the table below; and, "--" indicates that one or more R_{1a} , R_{1b} , R_{1c} and R_{1d} substituents are not present:



(Ib6)

Cpd	R_{1a}	R_{1b}	R_{1c}	R_{1d}
386	1-CH ₃ -1H-pyrazol-4-yl	OCH ₃	H	H
395	1-CH ₃ -1H-pyrazol-4-yl	OH	H	H
399	1H-pyrazol-4-yl	H	F	F
400	1H-pyrazol-4-yl	OH	H	F


Another embodiment of the method of the present description includes the use of a compound of Formula (Ib) or a form thereof selected from a compound of Formula (Ib7) or a form thereof, wherein substituent R_{1b} , when present, is indicated in the table below:



(Ib7)

Cpd	R_{1b}
304	OCH ₃
322	OH


Another embodiment of the method of the present description includes the use of a compound of Formula (Ib) or a form thereof selected from a compound of Formula (Ib8) or a form thereof, wherein substituent R_{1b} , when present, is indicated in the table below:




(Ib8)

Cpd	R_{1b}
303	OCH ₃
323	OH


An embodiment of the use of a compound of Formula (I) or a form thereof includes a method of use of a compound of Formula (I) or a form thereof for treating or ameliorating HD in a subject in need thereof, comprising administering an effective amount of the compound of Formula (I) or a form thereof to the subject, selected from the group consisting of:




1




2




3




4




5




6




7




8




9




10




11




12




13




14




15




16




17

















18

























19
















20

**21****23****25****27****29****31****33****22****24****26****28****30****32****34**












69




70




71




72




73




74




75















76

























77

























78









































**101****102****103****104****105****106****107****108****109****110****111****112**











**113****114****115****116****117****118****119****120****121****122**













**123****124****125****126****127****128****129****130****131****132****133****134**

























**149****150****151****152****153****154****155****156****157****158****159****160**















**161****162****163****164****165****166****167****168****169****170**





























**181****182****183****184****185****186****187****188****189****190****191****192**













**193****194****195****196****197****198****199****200****201****202**



























































305




306




307




308




309




310




311




312




313




314




315




316




317











318




319




320







329




330




331




332




333




334




335




336




337




338




339




340




341




342




343




344




345




346




347




348




349




350




351




352




353




354




355




356




357




358




359




360




361




362




363




364




365




366




367




368




369




370




371




372




373




374




375




376




377




378




379




380




381




382




383




384




385




386




387











388












389




390

**391****392****393****394****395****396****397****398**


**399****400****401****402****403****404****405****406**




407




408




409




410




411




412




413




414




415




416




417




418




419




420




421




422




423




424




425




426




427




428




429




430




431




432




433




434




435




436




437




438




439




440




441




442




443




444




445




446




447




448




449




450




451




453




455




457




459




461




452




454




456






458



460










462




wherein a form of the compound is selected from the group consisting of a prodrug, salt, hydrate, solvate, clathrate, isotopologue, racemate, enantiomer, diastereomer, stereoisomer, polymorph and tautomer form thereof.

In another aspect, the compound of Formula (I) or a form thereof includes a compound
5 selected from the group consisting of:







417




418




419




420




421




422




423




424




425




426




427




428




429




430




431




432




433




434




435




436




437




438




439




440




441




442




443




444




445




446




447




448




449




450




451




452




453




454




455




456




457




458




459




460



461



462



wherein a form of the compound is selected from the group consisting of a prodrug, salt, hydrate, solvate, clathrate, isotopologue, racemate, enantiomer, diastereomer, stereoisomer, polymorph and tautomer form thereof.

Another embodiment of the use of a compound of Formula (I) or a form thereof includes
 5 a method of use of a compound of Formula (I) or a form thereof for treating or ameliorating HD in a subject in need thereof, comprising administering an effective amount of the compound of Formula (I) or a form thereof (wherein compound number (#¹) indicates that the salt form was isolated) to the subject, selected from the group consisting of:

Cpd	Name
1	6-(naphthalen-2-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
2	6-(benzo[b]thiophen-2-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
3	2-(6-(2,2,6,6-tetramethylpiperidin-4-yl-amino)-pyridazin-3-yl)phenol
4	2-(6-(methyl-(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)benzo[b]-thiophene-5-carbonitrile
5	6-(quinolin-3-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
6	3-(benzo[b]-thiophen-2-yl)-6-(2,2,6,6-tetramethylpiperidin-4-yl-oxy)pyridazine
7	2-(6-(methyl-(2,2,6,6-tetramethylpiperidin-4-yl)amino)-pyridazin-3-yl)phenol
8	6-(6-(methyl-(2,2,6,6-tetramethylpiperidin-4-yl)amino)-pyridazin-3-yl)naphthalen-2-ol
9	6-(benzo[b]-thiophen-2-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
10	7-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)isoquinoline
11	6-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)isoquinoline
12	N-methyl-6-(quinolin-7-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine

Cpd	Name
13	N-methyl-6-(quinolin-6-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
14	6-(isoquinolin-7-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
15	6-(isoquinolin-6-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
16	6-(imidazo[1,2-a]pyridin-6-yl-pyridazin-3-yl)-methyl-(2,2,6,6-tetramethylpiperidin-4-yl)-amine
17	N-methyl-6-(6-phenylpyridin-3-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
18	6-(6-(1H-pyrrol-1-yl)pyridin-3-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
19	6-(6-(1H-pyrazol-1-yl)pyridin-3-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
20	methyl-(6-quinoxalin-2-yl-pyridazin-3-yl)-(2,2,6,6-tetramethylpiperidin-4-yl)-amine
21	methyl-(6-quinolin-3-yl-pyridazin-3-yl)-(2,2,6,6-tetramethylpiperidin-4-yl)-amine
22	N-methyl-6-(phthalazin-6-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
23	6-(benzo[c][1,2,5]oxa-diazol-5-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
24	6-(benzo[d]thiazol-5-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
25	6-(2-methylbenzo-[d]oxazol-6-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
26	3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalen-2-ol
27	5-chloro-2-(6-(methyl(1,2,2,6,6-pentamethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
28	3-(6-(2,2,6,6-tetramethylpiperidin-4-yl-amino)pyridazin-3-yl)naphthalen-2-ol
29	5-chloro-2-(6-(1,2,2,6,6-pentamethylpiperidin-4-ylamino)pyridazin-3-yl)phenol
30	4-hydroxy-3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)benzonitrile
31	3-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)naphthalen-2-ol
32 ¹	2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(trifluoromethyl)phenol
33	2-fluoro-6-(6-(methyl-(2,2,6,6-tetramethylpiperidin-4-yl)-amino)-pyridazin-3-yl)phenol
34	3,5-dimethoxy-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
35	4,5-dimethoxy-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
36	5-methoxy-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
37	4,5-difluoro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
38	5-fluoro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol

Cpd	Name
39	3-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)benzonitrile
40	1-allyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalen-2-ol
41	6-(benzo[b]thiophen-2-yl)-N-(1,2,2,6,6-pentamethylpiperidin-4-yl)pyridazin-3-amine
42	N-allyl-3-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)benzamide
43	2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
44	5-(5-methyl-oxazol-2-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)-amino)-pyridazin-3-yl)phenol
45	5-(4-hydroxymethyl)-1H-pyrazole-1-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
46	5-(1H-imidazol-1-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
47	5-(4-amino-1H-pyrazole-1-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
48	2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1H-pyrazol-4-yl)phenol
49	5-(3-amino-1H-pyrazol-1-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
50	2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1-(2-morpholino-ethyl)-1H-pyrazol-4-yl)phenol
51	2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1-methyl-1H-pyrazol-4-yl)phenol
52	5-(5-amino-1H-pyrazol-1-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
53 ¹	2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(1H-pyrazol-1-yl)phenol
54	2-(((6-((2-hydroxy-ethyl)-(2,2,6,6-tetramethylpiperidin-4-yl)-amino)-pyridazin-3-yl)-5-pyrazol-1-yl)phenol
55	2-(6-(piperidin-4-yloxy)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
56	2-(6-(((2S,4R,6R)-2,6-dimethylpiperidin-4-yl)oxy)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
57	2-(6-((-2,6-dimethylpiperidin-4-yl)oxy)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
58	5-(1H-pyrazol-1-yl)-2-(6-(pyrrolidin-3-yl-oxy)pyridazin-3-yl)phenol
59	2-(6-(((2S,4S)-2-methylpiperidin-4-yl)oxy)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
60	(5-(1H-pyrazol-1-yl)-2-(6-(pyrrolidin-3-ylmethoxy)pyridazin-3-yl)phenol

Cpd	Name
61	2-(6-((3-fluoropiperidin-4-yl)oxy)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
62	2-(6-(1,2,2,6,6-pentamethyl-piperidin-4-yl-oxy)-pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
63	5-1H-pyrazol-1-yl-2-(6-(2,2,6,6-tetramethylpiperidin-4-yl-oxy)-pyridazin-3-yl)phenol
64	5-(1H-pyrazol-4-yl)-2-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)phenol
65 ¹	2-(6-piperazin-1-yl-pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
66	3-(6-(azetidin-3-ylamino)-pyridazin-3-yl)naphthalen-2-ol
67	2-(6-(azetidin-3-ylamino)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
68	2-(6-(3,5-dimethylpiperazin-1-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
69	2-(6-(7-methyl-2,7-diazaspiro[4.4]nonan-2-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
70	2-(6-(1,4-diazepan-1-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
71	2-(6-(4-(2-hydroxyethyl)piperazin-1-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
72	2-(6-(3,6-diazabicyclo[3.2.1]octan-3-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
73	2-(6-(2,7-diazaspiro[3.5]nonan-7-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
74	2-(6-(3-(hydroxymethyl)piperazin-1-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
75	2-(6-(1,7-diazaspiro[4.4]nonan-7-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
76	2-(6-(4-amino-4-methylpiperidin-1-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
77	2-(6-(3-(dimethylamino)piperidin-1-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
78	2-(6-(1,2,2,6,6-pentamethylpiperidin-4-ylamino)-pyridazin-3-yl)-5-1H-pyrazol-1-yl-phenol
79	2-(6-(3,3-dimethylpiperazin-1-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
80	2-(6-(7-(2-hydroxyethyl)-2,7-diazaspiro[4.4]-nonan-2-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
81	2-(6-((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
82 ¹	3-(6-(piperazin-1-yl)pyridazin-3-yl)naphthalene-2,7-diol
83	5-(1H-pyrazol-1-yl)-2-(6-(1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl)phenol
84	2-(6-piperidin-4-yl-pyridazin-3-yl)-5-1H-pyrazol-1-yl-phenol
85	3-(6-(1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl)naphthalen-2-ol
86 ¹	3-(6-(1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl)naphthalene-2,7-diol
87	3-(6-(2,2,6,6-tetramethyl-1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl)naphthalene-2,7-diol
88 ¹	3-(6-(1-methyl-1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl)naphthalene-2,7-diol
89 ¹	3-(6-(piperidin-4-yl)pyridazin-3-yl)naphthalene-2,7-diol
90	3-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)naphthalene-2,7-diol

Cpd	Name
91	3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalene-2,7-diol
92	3-(6-((2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalene-2,7-diol
93	tert-butyl 3-((7-hydroxy-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalen-2-yl)oxy)propyl)carbamate
94	7-(3-amino-propoxy)-3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)-amino)-pyridazin-3-yl)naphthalen-2-ol
95	N-(3-((7-hydroxy-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalen-2-yl)oxy)propyl)acetamide
96	7-(3-hydroxypropoxy)-3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalen-2-ol
97	7-(3-methoxypropoxy)-3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalen-2-ol
98	7-(2-morpholinoethoxy)-3-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)naphthalen-2-ol
99	3-(6-(piperidin-4-ylmethyl)pyridazin-3-yl)naphthalen-2-ol
100	5-(1H-pyrazol-1-yl)-2-(6-((2,2,6,6-tetramethylpiperidin-4-yl)methyl)pyridazin-3-yl)phenol
101	3-methoxy-2-(6-(methyl(2,2,6-trimethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(5-methyloxazol-2-yl)phenol
102	2-(6-(((6S)-6-((S)-1-hydroxyethyl)-2,2-dimethylpiperidin-4-yloxy)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
103	7-hydroxy-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-2-naphthonitrile
104	3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-7-(piperidinylmethyl)naphthalen-2-ol
105	3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-7-(pyrrolidinylmethyl)naphthalen-2-ol
106	1-bromo-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalene-2,7-diol
107	1-chloro-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalene-2,7-diol
108	7-methoxy-3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalene-2-ol
109	7-methoxy-3-(6-(methyl(1,2,2,6,6-pentamethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalen-2-ol
110	7-(3,6-dihydro-2H-pyran-4-yl)-3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalen-2-ol

Cpd	Name
111	3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-7-(tetrahydro-2H-pyran-4-yl)naphthalene-2-ol
112	7-(difluoromethyl)-3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalen-2-ol
113	7-((4-hydroxy-2-methylbutan-2-yl)oxy)-3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalen-2-ol
114	7-(3-hydroxy-3-methylbutoxy)-3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalen-2-ol
115	2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1H-pyrazol-4-yl)benzene-1,3-diol
116	3-methoxy-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1H-pyrazol-4-yl)phenol
117	5-(1H-pyrazol-4-yl)-2-(6-((2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-3-(trifluoromethoxy)phenol
118	2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1-methyl-1H-pyrazol-4-yl)-3-(trifluoromethoxy)phenol
119	2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1H-pyrazol-4-yl)-3-(trifluoromethoxy)phenol
120	4-(3-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(trifluoromethoxy)phenyl)-1-methylpyridin-2(1H)-one
121	3-methoxy-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1-methyl-1H-pyrazol-4-yl)phenol
122	3-methoxy-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(5,6,7,8-tetrahydroimidazo[1,2-a]pyridin-3-yl)phenol
123	3-methoxy-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(pyridine-3-yl)phenol
124	5-(1-cyclopentyl-1H-pyrazol-4-yl)-3-methoxy-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
125	3',5-dimethoxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-(1,1'-biphenyl)-3-ol
126	3-(benzyloxy)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(5-methyloxazol-2-yl)phenol
127	3-ethoxy-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(5-methyloxazol-2-yl)phenol
128	3-(cyclopropylmethoxy)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(5-methyloxazol-2-yl)phenol
129	2-methyl-5-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-1H-benzo[d]imidazol-6-ol
130	5-chloro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol

Cpd	Name
131	5-(1H-pyrazol-1-yl)-2-(6-((2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
132	3-hydroxy-4-(6-((2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)benzonitrile
133	2-(6-((2,2-dimethylpiperidin-4-yl)oxy)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
134	2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(1H-pyrazol-4-yl)phenol
135	2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridin-3-yl)-4-(4,5,6,7-tetrahydropyrazolo[1,5-a]pyridin-3-yl)phenol
136	2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-3-yl)phenol
137	4-(1H-indol-2-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
138	4-(cyclopent-1-en-1-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
139	2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(1H-pyrazol-3-yl)phenol
140	4-(4-hydroxy-3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)pyridin-2-ol
141	4-(4-hydroxy-3-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)phenyl)-1-methylpyridin-2(1H)-one
142	4-(4-hydroxy-3-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)phenyl)pyridin-2-ol
143	5-(1H-indazol-7-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
144	4-chloro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1H-pyrazol-4-yl)phenol
145	4-fluoro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1H-pyrazol-4-yl)phenol
146	5-fluoro-4-(1H-imidazol-4-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
147	5-fluoro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(1H-pyrazol-4-yl)phenol
148	5-fluoro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(1H-pyrazol-5-yl)phenol
149	6-hydroxy-5-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-2,3-dihydro-1H-inden-1-one
150	6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-1,4-dihydroindeno[1,2-c]-1H-pyrazol-7-ol

Cpd	Name
151 ¹	6-hydroxy-5-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-2,3-dihydro-1H-inden-1-one oxime
152	5-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-2,3-dihydro-1H-indene-1,6-diol
153 ¹	2-amino-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-8H-indeno[1,2-d]thiazol-5-ol
154 ¹	9-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5,6-dihydroimidazo[5,1-a]isoquinolin-8-ol
155	4-hydroxy-3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-N-((1-methyl-1H-pyrazol-4-yl)methyl)benzamide
156	4-(4-(hydroxymethyl)-1H-pyrazol-1-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
157	5-(1H-pyrazol-4-yl)-2-(6-((2,2,6,6-tetramethylpiperidin-4-yl)methyl)pyridazin-3-yl)phenol
158	6-(3-(benzyloxy)isoquinolin-6-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
159	6-(1-(benzyloxy)isoquinolin-7-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
160 ¹	3-fluoro-5-(2-methoxypyridin-4-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
161 ¹	4-(3-fluoro-5-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)pyridin-2(1H)-one
162 ¹	4-(3-fluoro-5-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)-1-methylpyridin-2(1H)-one
163 ¹	5-(3-fluoro-5-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)-1-methylpyridin-2(1H)-one
164 ¹	3-fluoro-5-(1H-pyrazol-4-yl)-2-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)phenol
165 ¹	5-chloro-3-fluoro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
166 ¹	3-fluoro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1H-pyrazol-4-yl)phenol
167 ¹	3-fluoro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1-methyl-1H-pyrazol-4-yl)phenol
168	5-(5-methoxypyridin-3-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
169	5-(3-hydroxy-4-(6-methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)pyridin-2-ol

Cpd	Name
170	4-(3-hydroxy-4-(6-methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)pyridin-2-ol
171	5-(6-methoxypyridin-3-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
172	5-(3-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)-3-(trifluoromethyl)pyridin-2-ol
173	5-(3-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)-1-methylpyridin-2(1H)-one
174	4-(3-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)-1-methylpyridin-2(1H)-one
175	5-(2-methoxypyridin-4-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
176	4-(3-hydroxy-4-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)phenyl)pyridin-2-ol
177	5-(6-(dimethylamino)pyridin-3-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
178	4-(3-hydroxy-4-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)phenyl)-1-methylpyridin-2(1H)-one
179	2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(pyrimidin-5-yl)phenol
180	5-(3-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)pyridin-3-ol
181	1-cyclopropyl-4-(3-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)pyridin-2(1H)-one
182	2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1,2,3,6-tetrahydropyridin-4-yl)phenol
183	5-(cyclopent-1-en-1-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
184	5-(3,6-dihydro-2H-pyran-4-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
185	5-(imidazo[1,5-a]pyridin-7-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
186	5-(imidazo[1,2-a]pyridin-7-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
187	2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(2-methylpyridin-4-yl)phenol
188	5-(1H-imidazol-2-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol

Cpd	Name
189	5-(1H-imidazol-4-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
190	5-(imidazo[1,2-a]pyrazin-3-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
191	2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(5,6,7,8-tetrahydroimidazo[1,2-a]pyrazin-3-yl)phenol
192	2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(4-methyl-1H-imidazol-2-yl)phenol
193	2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1-methyl-1H-imidazol-4-yl)phenol
194	2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1-methyl-1H-imidazol-5-yl)phenol
195	2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(4-nitro-1H-imidazol-2-yl)phenol
196	2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(2-methyl-1H-imidazol-4-yl)phenol
197	5-(1,2-dimethyl-1H-imidazol-4-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
198	1-(3-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)-1H-pyrazole-4-carboxamide
199	2-(6-((3aR,6aS)-5-(2-hydroxyethyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)pyridazin-3-yl)-5-(1H-pyrazol-4-yl)phenol
200	2-(6-((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)pyridazin-3-yl)-5-(1H-pyrazol-4-yl)phenol
201	2-(6-((3aR,6aS)-5-methylhexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)pyridazin-3-yl)-5-(1H-pyrazol-4-yl)phenol
202	4-(3-hydroxy-4-(6-(5-methylhexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)pyridazin-3-yl)phenyl)-1-methylpyridin-2(1H)-one
203	4-(3-hydroxy-4-(6-((3aR,6aR)-1-methylhexahydropyrrolo[3,4-b]pyrrol-5(1H)-yl)pyridazin-3-yl)phenyl)-1-methylpyridin-2(1H)-one
204	2-(6-(2,7-diazaspiro[4.5]decan-2-yl)pyridazin-3-yl)-5-(1H-pyrazol-4-yl)phenol
205	4-(4-(6-(2,7-diazaspiro[4.5]decan-2-yl)pyridazin-3-yl)-3-hydroxyphenyl)-1-methylpyridin-2(1H)-one
206	2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)-amino)-pyridazin-3-yl)phenol
207	6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
208	6-(6-(methyl(1,2,2,6,6-pentamethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
209	6-(6-((3aR,6aS)-5-methylhexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)pyridazin-3-yl)quinolin-7-ol

Cpd	Name
210	2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
211	7-(6-(methyl(1,2,2,6,6-pentamethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-6-ol
212	7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-6-ol
213	7-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)isoquinoline-6-ol
214	7-(6-((3aR,6aS)-5-methylhexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)pyridazin-3-yl)isoquinolin-6-ol
215	1-cyclopropyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-6-ol
216	7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinoline-1,6-diol
217	6-hydroxy-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinoline-1-carbonitrile
218	6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-7-ol
219	8-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
220	6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-6-ol
221	2-methyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-6-ol
222	3-chloro-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
223	3-bromo-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
224	7-hydroxy-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinoline-3-carbonitrile
225	6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-3-(1-methyl-1H-imidazol-4-yl)quinolin-7-ol
226 ¹	3-(1H-imidazol-1-yl)-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
227	6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinoline-3,7-diol
228	3-ethyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
229	3-isopropyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
230	7-hydroxy-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-2(1H)-one
231 ¹	7-hydroxy-1-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-2(1H)-one

Cpd	Name
232	4-methoxy-2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
233	2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(pyrrolidin-1-yl)quinolin-7-ol
234	2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-morpholinoquinolin-7-ol
235	4-(dimethylamino)-2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
236	4-ethoxy-2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
237	2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(1-methyl-1H-pyrazol-4-yl)quinolin-7-ol
238 ¹	4-methoxy-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-6-ol
239 ¹	7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinoxalin-6-ol
240	6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-3-(tetrahydro-2H-pyran-4-yl)quinolin-7-ol
241	3-chloro-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-6-ol
242	3-bromo-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-6-ol
243	3-methyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-6-ol
244	5-bromo-3-methyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-6-ol
245	6-hydroxy-1-methyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-4(1H)-one
246	2,3-dimethyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-6-ol
247	2-methyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinoxalin-6-ol
248	3-methyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinoxalin-6-ol
249	4-methoxy-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
250	4-(azetidin-1-yl)-2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
251	7-hydroxy-2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolone-4-carbonitrile

Cpd	Name
252	4-cyclopropyl-2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-quinolin-7-ol
253	4-(3,6-dihydro-2H-pyran-4-yl)-2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
254 ¹	2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(tetrahydro-2H-pyran-4-yl)quinolin-7-ol
255	2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(oxetan-3-yl)quinolin-7-ol
256 ¹	4-(dimethylamino)-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-quinolin-7-ol
257	7-hydroxy-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinazolin-4(1H)-one
258	6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinazolin-7-ol
259	7-hydroxy-1-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-3,4-dihydroquinolin-2(1H)-one
260	2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinazolin-7-ol
261	7-hydroxy-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinoline-1-carbonitrile
262	7-hydroxy-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinoline-2-carbonitrile
263	6-hydroxy-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-2-carbonitrile
264	6-hydroxy-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinoline-1-carboxamide
265	7-hydroxy-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-2-carboxamide
266	6-hydroxy-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinoline-2-carboxamide
267	methyl 6-hydroxy-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinoline-2-carboxylate
268	6-hydroxy-7-(6-(piperazin-1-yl)pyridazin-3-yl)quinoline-2-carbonitrile
269	7-hydroxy-6-(6-(piperazin-1-yl)pyridazin-3-yl)quinoline-2-carbonitrile
270	7-(6-(piperazin-1-yl)pyridazin-3-yl)isoquinolin-6-ol
271	7-(6-(1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl)quinolin-6-ol
272	1-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-7-ol
273	1-methyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-6-ol

Cpd	Name
274	1,3-dimethyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-6-ol
275	7-hydroxy-3-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinoline-1-carbonitrile
276	1-amino-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-6-ol
277	7-hydroxy-1,3-dimethyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinazoline-2,4(1H,3H)-dione
278	6-hydroxy-5-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)benzo[d]oxazol-2(3H)-one
279	2-methyl-5-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-2H-indazol-6-ol
280	1-methyl-5-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-1H-indazol-6-ol
281 ¹	6-hydroxy-2-methyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-1(2H)-one
282	2-ethyl-6-hydroxy-7-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)isoquinolin-1(2H)-one
283	1-ethoxy-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-6-ol
284	7-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)isoquinoline-1,6-diol
285	7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-pyridazin-3-yl)-3-phenylisoquinolin-6-ol
286	3-methyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-6-ol
287	3-cyclopropyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-6-ol
288	3-isopropyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-6-ol
289	3-propyl-7-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)-pyridazin-3-yl)isoquinolin-6-ol
290	3-isopropyl-7-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)-pyridazin-3-yl)isoquinolin-6-ol
291	3-methyl-7-(6-(piperazin-1-yl)pyridazin-3-yl)isoquinolin-6-ol
292	6-(3-(benzyloxy)isoquinolin-6-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
293	3-chloro-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
294	3-isopropyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol

Cpd	Name
295	3-methyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinoxalin-6-ol
296	4-chloro-2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
297	4-chloro-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-6-ol
300	7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-6-ol
301	5-(2-methoxy-4-(1H-pyrazol-1-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
302	6-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)naphthalen-2-ol
303	5-(2-methoxyquinolin-3-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
304	5-(3-methoxy-naphthalen-2-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
305	5-(2-methoxy-4-(1H-pyrazol-1-yl)phenyl)-N-(1,2,2,6,6-pentamethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
306	5-(2-methoxy-4-(1-methyl-1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
307	5-(2-methoxy-4-(1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
308	4-(3-methoxy-4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)phenyl)-1-methylpyridin-2(1H)-one
309	5-(3-methoxy-4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)phenyl)pyridin-2-ol
310	5-(3-methoxy-4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)phenyl)-1-methylpyridin-2(1H)-one
311	N-methyl-5-(2-methyl-4-(1-methyl-1H-pyrazol-4-yl)phenyl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
312	1-methyl-4-(4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-3-(trifluoromethoxy)phenyl)pyridin-2(1H)-one
313	5-(4-(3,5-dimethyl-1H-pyrazol-4-yl)-2-methoxyphenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
314	5-(2-methoxy-4-(1-methyl-1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
315	2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-5-(1-methyl-1H-pyrazol-4-yl)phenol
316	2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-5-(1H-pyrazol-1-yl)phenol

Cpd	Name
317	5-(3-hydroxy-4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)phenyl)-1-methylpyridin-2(1H)-one
318	4-(3-hydroxy-4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)phenyl)-1-methylpyridin-2(1H)-one
319	5-(3-hydroxy-4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)phenyl)pyridin-2-ol
320	3-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)naphthalene-2,7-diol
321	3-(5-((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-1,3,4-thiadiazol-2-yl)naphthalene-2,7-diol
322 ¹	3-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)naphthalen-2-ol
323	3-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)quinolin-2-ol
324	2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-4-(1H-pyrazol-1-yl)phenol
325	5-(2-chloro-4-(1-methyl-1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
326	3-chloro-2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-5-(1-methyl-1H-pyrazol-4-yl)phenol
327	5-(2-chloro-4-(1-methyl-1H-pyrazol-4-yl)phenyl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
328	3-methoxy-2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-5-methyloxazol-2-yl)phenol
329	2-(2-methoxy-4-(1H-pyrazol-1-yl)phenyl)-5-(1,2,3,6-tetrahydropyridin-4-yl)-1,3,4-thiadiazole
330	2-(5-(piperazin-1-yl)-1,3,4-thiadiazol-2-yl)-5-(1H-pyrazol-1-yl)phenol
331	5-(7-methoxyquinolin-6-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazole-2-amine
332	6-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)quinolin-7-ol
333	3-methoxy-4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)benzonitrile
334	3-fluoro-4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)benzonitrile
335	methyl-3-fluoro-4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)benzoate
336	5-(2-methoxy-4-(3-(methylamino)-1H-pyrazol-1-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

Cpd	Name
337	7-methoxy-6-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)quinoline-2-carbonitrile
338	4-(3-methoxy-4-(5-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)-1,3,4-thiadiazol-2-yl)phenyl)-1-methylpyridin-2(1H)-one
339	4-(3-chloro-4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)phenyl)-1-methylpyridin-2(1H)-one
340	5-(2-chloro-4-(1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
341	5-(2-chloro-4-(4,5,6,7-tetrahydropyrazolo[1,5-a]pyridin-3-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
342 ¹	N-methyl-5-(5-(1-methyl-1H-pyrazol-4-yl)pyridin-2-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
343	2-(2-chloro-4-(1-methyl-1H-pyrazol-4-yl)phenyl)-5-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)-1,3,4-thiadiazole
344	5-(2-chloro-4-(6-methoxypyridin-3-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
345	5-(4-(6-aminopyridin-3-yl)-2-fluorophenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
346	5-(2-fluoro-4-(3-methyl-1H-pyrazol-5-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
347	5-(2-fluoro-4-(1H-pyrazol-5-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
348	5-(2,3-difluoro-4-(1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
349	5-(2,3-difluoro-4-(1H-pyrazol-5-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
350	5-(2,5-difluoro-4-(1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
351	5-(2,5-difluoro-4-(1H-pyrazol-5-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
352	5-(2,6-difluoro-4-(1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
353	2-(2,5-difluoro-4-(1H-pyrazol-4-yl)phenyl)-5-((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-1,3,4-thiadiazole
354	5-(2-chloro-5-fluoro-4-(1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
355	5-(3-fluoro-5-(1H-pyrazol-4-yl)pyridin-2-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

Cpd	Name
356	5-(4-(2-aminopyrimidin-4-yl)-2-chlorophenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
357	5-(5-(2-aminopyrimidin-4-yl)-2-chlorophenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
358	5-(4-(2,4-dimethylthiazol-5-yl)-2,5-difluorophenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
359	5-(4-(2,4-dimethylthiazol-5-yl)-2,3-difluorophenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
360	4-(3-hydroxy-4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-5-(trifluoromethoxy)phenyl)-1-methylpyridin-2(1H)-one
361	5-(2-fluoro-6-methoxy-4-(1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
362	2-(2-fluoro-6-methoxy-4-(1H-pyrazol-4-yl)phenyl)-5-((3aR,6aS)-5-methylhexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-1,3,4-thiadiazole
363	5-(2,3-difluoro-6-methoxy-4-(1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
364	6-methoxy-2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-3,4-dihydroisoquinolin-1-(2H)-one
365	5-(2-chloro-4-(1H-pyrazol-1-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
366	5-(2-chloro-4-(1H-1,2,3-triazol-1-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
367	5-(2-chloro-4-(2H-1,2,3-triazol-2-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
368	5-(2-chloro-4-(1H-1,2,4-triazol-1-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
369	5-(4-(3-amino-1H-pyrazol-1-yl)-2-chlorophenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
370	2-(2-chloro-4-(1H-imidazol-1-yl)phenyl)-5-((3aR,6aS)-5-methylhexahydropyrrolo[3,4c]pyrrol-2(1H)-yl)-1,3,4-thiadiazole
371	5-(2-chloro-4-(1H-imidazol-1-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
372	5-(2-fluoro-4-(1H-imidazol-1-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
373	5-(2-methoxy-4-(1H-pyrazol-5-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
374	5-(4-(2,4-dimethylthiazol-5-yl)-2-methoxyphenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

Cpd	Name
375	5-(2-methoxy-4-(pyridin-3-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
376	5-(2-fluoro-4-(1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
377	5-(2-methoxy-4-(2-methoxypyridin-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
378	5-(2-methoxy-4-(6-methoxypyridin-3-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
379	2-(2-chloro-4-(1-methyl-1H-pyrazol-4-yl)phenyl)-5-((3aR,6aS)-5-methylhexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-1,3,4-thiadiazole
380	2-(2-chloro-4-(1H-pyrazol-4-yl)phenyl)-5-((3aR,6aS)-5-methylhexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-1,3,4-thiadiazole
381	2-(2-chloro-4-(1H-pyrazol-4-yl)phenyl)-5-((3aR,6aR)-1-methylhexahydropyrrolo[3,4-b]pyrrol-5(1H)-yl)-1,3,4-thiadiazole
382	1-(4-(5-(2-chloro-4-(1H-pyrazol-4-yl)phenyl)-1,3,4-thiadiazol-2-yl)morpholin-2-yl)-N,N-dimethylmethanamine
383	2-(2-chloro-4-(1H-pyrazol-4-yl)phenyl)-5-(2-methyl-2,7-diazaspiro[4.5]decan-7-yl)-1,3,4-thiadiazole
384	2-(2-fluoro-4-(1H-pyrazol-4-yl)phenyl)-5-((3aR,6aS)-5-methylhexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-1,3,4-thiadiazole
385	2-(2-methoxy-4-(1-methyl-1H-pyrazol-4-yl)phenyl)-5-(2,6-diazaspiro[3.5]nonan-2-yl)-1,3,4-thiadiazole
386	2-(2-methoxy-4-(1-methyl-1H-pyrazol-4-yl)phenyl)-5-(2,7-diazaspiro[3.5]nonan-2-yl)-1,3,4-thiadiazole
387	2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-5-(1H-pyrazol-1-yl)phenol
388	5-(3-chloro-4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)phenyl)pyridin-2(1H)-one
389	2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-5-(3-(methylamino)-1H-pyrazol-1-yl)phenol
390	3-fluoro-2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-5-(1H-pyrazol-4-yl)phenol
391	3,4-difluoro-2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-5-(1H-pyrazol-4-yl)phenol
392	6-hydroxy-5-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-2,3-dihydro-1H-inden-1-one
393	2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-5-(1H-pyrazol-4-yl)phenol

Cpd	Name
394	2-(5-(2,6-diazaspiro[3.5]nonan-2-yl)-1,3,4-thiadiazol-2-yl)-5-(1-methyl-1H-pyrazol-4-yl)phenol
395	2-(5-(2,7-diazaspiro[3.5]nonan-2-yl)-1,3,4-thiadiazol-2-yl)-5-(1-methyl-1H-pyrazol-4-yl)phenol
396 ¹	3-fluoro-2-(5-((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-1,3,4-thiadiazol-2-yl)-5-(1H-pyrazol-4-yl)phenol
397	3-chloro-2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-5-(1H-pyrazol-4-yl)phenol
398	2-(2-methoxy-4-(1H-pyrazol-1-yl)phenyl)-5-((2,2,6,6-tetramethylpiperidin-4-yl)methyl)-1,3,4-thiadiazole
399	2-(2,3-difluoro-4-(1H-pyrazol-4-yl)phenyl)-5-(2,7-diazaspiro[3.5]nonan-2-yl)-1,3,4-thiadiazole
400	2-(5-(2,7-diazaspiro[3.5]nonan-2-yl)-1,3,4-thiadiazol-2-yl)-3-fluoro-5-(1H-pyrazol-4-yl)phenol
401	4-methoxy-1-methyl-3-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)quinolin-2(1H)-one
402	4-hydroxy-1-methyl-3-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)quinolin-2(1H)-one
403	3-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)quinolin-2(1H)-one
404	1-methyl-3-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)quinolin-2(1H)-one
405 ¹	2-(2-chloro-4-(1H-pyrazol-4-yl)phenyl)-5-((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-1,3,4-thiadiazole
406 ¹	2-(2-chloro-4-(1H-pyrazol-4-yl)phenyl)-5-(2,7-diazaspiro[4.5]decan-2-yl)-1,3,4-thiadiazole
407 ¹	(R)-(4-(5-(2-chloro-4-(1H-pyrazol-4-yl)phenyl)-1,3,4-thiadiazol-2-yl)piperazin-2-yl)methanol
408	2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)benzo[b]thiophene-5-carbonitrile
409	5-(3-chlorobenzo[b]thiophen-2-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
410	5-(2-methoxy-4-(1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
411 ¹	2-{6-[(1R,5S)-8-azabicyclo[3.2.1]oct-3-yl(methyl)amino]pyridazin-3-yl}-5-(1H-pyrazol-4-yl)phenol
412 ¹	2-[6-((1R,5S)-8-azabicyclo[3.2.1]oct-3-ylamino)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol

Cpd	Name
413 ¹	5-(1H-pyrazol-4-yl)-2-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl}phenol
414	5-(1-methyl-1H-pyrazol-4-yl)-2-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}phenol
415 ¹	2-[6-((1 <i>R</i> ,5 <i>S</i>)-8-azabicyclo[3.2.1]oct-3-yloxy)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
416	5-(5-methyl-1H-pyrazol-4-yl)-2-{6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl}phenol
417	5-(1H-imidazol-1-yl)-2-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}phenol
418	5-(5-methyl-1H-pyrazol-4-yl)-2-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}phenol
419 ¹	2-{6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl}-5-(4-nitro-1H-pyrazol-1-yl)phenol
420	6-[2-methoxy-4-(4-nitro-1H-pyrazol-1-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
421	5-(4-amino-1H-pyrazol-1-yl)-2-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}phenol
422 ¹	2-[6-(1-methyl-1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
423	5-(4-nitro-1H-pyrazol-1-yl)-2-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}phenol
424 ¹	5-(1H-pyrazol-4-yl)-2-[6-(1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]phenol
425 ¹	2-[6-(1-ethyl-1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
426 ¹	2-{6-[methyl(piperidin-4-yl)amino]pyridazin-3-yl}-5-(1H-pyrazol-4-yl)phenol
427 ¹	2-[6-(piperidin-4-ylamino)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
428 ¹	6-[2,5-difluoro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
429 ¹	2-[6-(8-azabicyclo[3.2.1]oct-2-en-3-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
430 ¹	6-[2,3-difluoro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
431 ¹	3-[2,5-difluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
432	2-[6-(piperidin-4-yloxy)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
433 ¹	2-{6-[(1 <i>R</i> ,5 <i>S</i>)-8-azabicyclo[3.2.1]oct-3-ylamino]pyridazin-3-yl}-5-(1H-pyrazol-4-yl)phenol
434 ¹	6-[2-methoxy-6-(1H-pyrazol-4-yl)pyridin-3-yl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine

Cpd	Name
435	3-[4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
436 ¹	2-{6-[(2,6-dimethylpiperidin-4-yl)oxy]pyridazin-3-yl}-5-(1H-pyrazol-4-yl)phenol
437 ¹	3-[2-fluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
438	3-[4-(1-methyl-1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
439 ¹	2-[6-(2,7-diazaspiro[3.5]non-2-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
440	3-fluoro-4-{6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl}phenol
441	2-{6-[(2,6-dimethylpiperidin-4-yl)oxy]pyridazin-3-yl}-5-(1H-pyrazol-1-yl)phenol
442	N-methyl-6-(2-methyl-2H-indazol-5-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
443	2-methyl-5-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}-2H-indazole
444	3-(4-chloro-2-methoxyphenyl)-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
445	N-methyl-6-(2-methylpyrazolo[1,5-a]pyridin-3-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
446	6-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}imidazo[1,2-a]pyridine
447	3-[2-methoxy-4-(1H-pyrazol-1-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
448 ¹	3-[5-(1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
449	3-[5-(1-methyl-1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
450 ¹	3-[4-(1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
451	5-(3,5-dimethyl-1H-pyrazol-4-yl)-2-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}phenol
452	6-[2-fluoro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
453	3-methoxy-4-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}phenol
454	3-[2-methoxy-4-(4-nitro-1H-pyrazol-1-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
455	4-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}benzene-1,3-diol
456 ¹	6-[2-chloro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
457	2-(1H-pyrazol-4-yl)-4-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}pyrimidin-5-amine
458 ¹	3-[2,6-difluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine

Cpd	Name
459	2-[6-(2,6-diazaspiro[3.4]oct-2-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
460¹	3-{6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl}-6-(1H-pyrazol-4-yl)pyridin-2-ol
461	6-(1H-pyrazol-4-yl)-3-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}pyridin-2-ol
462¹	N,2,2,6,6-pentamethyl-N-{5-[3-(1H-pyrazol-4-yl)phenoxy]-1,3,4-thiadiazol-2-yl}piperidin-4-amine
463¹	N,2,2,6,6-pentamethyl-N-{5-[4-(1H-pyrazol-4-yl)phenoxy]-1,3,4-thiadiazol-2-yl}piperidin-4-amine
464¹	3-[2-(difluoromethyl)-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine and
465¹	6-[2-(difluoromethyl)-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine

wherein a form of the compound is selected from the group consisting of a prodrug, salt, hydrate, solvate, clathrate, isotopologue, racemate, enantiomer, diastereomer, stereoisomer, polymorph and tautomer form thereof.

In one aspect, the compound of Formula (I) or a form thereof includes a compound
5 selected from the group consisting of:

411¹	2-{6-[(1 <i>R</i> ,5 <i>S</i>)-8-azabicyclo[3.2.1]oct-3-yl(methyl)amino]pyridazin-3-yl}-5-(1H-pyrazol-4-yl)phenol
412¹	2-[6-((1 <i>R</i> ,5 <i>S</i>)-8-azabicyclo[3.2.1]oct-3-ylamino)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
413¹	5-(1H-pyrazol-4-yl)-2-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl}phenol
414	5-(1-methyl-1H-pyrazol-4-yl)-2-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}phenol
415¹	2-[6-((1 <i>R</i> ,5 <i>S</i>)-8-azabicyclo[3.2.1]oct-3-yloxy)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
416	5-(5-methyl-1H-pyrazol-4-yl)-2-{6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl}phenol
417	5-(1H-imidazol-1-yl)-2-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}phenol
418	5-(5-methyl-1H-pyrazol-4-yl)-2-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}phenol
419¹	2-{6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl}-5-(4-nitro-1H-pyrazol-1-yl)phenol

- 420 6-[2-methoxy-4-(4-nitro-1H-pyrazol-1-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
- 421 5-(4-amino-1H-pyrazol-1-yl)-2-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}phenol
- 422¹ 2-[6-(1-methyl-1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
- 423 5-(4-nitro-1H-pyrazol-1-yl)-2-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}phenol
- 424¹ 5-(1H-pyrazol-4-yl)-2-[6-(1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]phenol
- 425¹ 2-[6-(1-ethyl-1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
- 426¹ 2-{6-[methyl(piperidin-4-yl)amino]pyridazin-3-yl}-5-(1H-pyrazol-4-yl)phenol
- 427¹ 2-[6-(piperidin-4-ylamino)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
- 428¹ 6-[2,5-difluoro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
- 429¹ 2-[6-(8-azabicyclo[3.2.1]oct-2-en-3-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
- 430¹ 6-[2,3-difluoro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
- 431¹ 3-[2,5-difluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
- 432 2-[6-(piperidin-4-yloxy)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
- 433¹ 2-{6-[(1R,5S)-8-azabicyclo[3.2.1]oct-3-ylamino]pyridazin-3-yl}-5-(1H-pyrazol-4-yl)phenol
- 434¹ 6-[2-methoxy-6-(1H-pyrazol-4-yl)pyridin-3-yl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
- 435 3-[4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
- 436¹ 2-{6-[(2,6-dimethylpiperidin-4-yl)oxy]pyridazin-3-yl}-5-(1H-pyrazol-4-yl)phenol
- 437¹ 3-[2-fluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
- 438 3-[4-(1-methyl-1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
- 439¹ 2-[6-(2,7-diazaspiro[3.5]non-2-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
- 440 3-fluoro-4-{6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl}phenol
- 441 2-{6-[(2,6-dimethylpiperidin-4-yl)oxy]pyridazin-3-yl}-5-(1H-pyrazol-1-yl)phenol
- 442 N-methyl-6-(2-methyl-2H-indazol-5-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
- 443 2-methyl-5-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}-2H-indazole
- 444 3-(4-chloro-2-methoxyphenyl)-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
- 445 N-methyl-6-(2-methylpyrazolo[1,5-a]pyridin-3-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine

- 446 6-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}imidazo[1,2-a]pyridine
 3-[2-methoxy-4-(1H-pyrazol-1-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
 447 3-[5-(1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
 448¹ 3-[5-(1-methyl-1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
 449 3-[4-(1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
 450¹ 5-(3,5-dimethyl-1H-pyrazol-4-yl)-2-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}phenol
 451 6-[2-fluoro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
 452 3-methoxy-4-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}phenol
 453 3-[2-methoxy-4-(4-nitro-1H-pyrazol-1-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
 454 4-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}benzene-1,3-diol
 455 6-[2-chloro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
 456¹ 2-(1H-pyrazol-4-yl)-4-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}pyrimidin-5-amine
 457 3-[2,6-difluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
 458¹ 2-[6-(2,6-diazaspiro[3.4]oct-2-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
 459 3-{6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl}-6-(1H-pyrazol-4-yl)pyridin-2-ol
 460¹ 6-(1H-pyrazol-4-yl)-3-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}pyridin-2-ol
 461 N,2,2,6,6-pentamethyl-N-{5-[3-(1H-pyrazol-4-yl)phenoxy]-1,3,4-thiadiazol-2-yl}piperidin-4-amine
 462¹ N,2,2,6,6-pentamethyl-N-{5-[4-(1H-pyrazol-4-yl)phenoxy]-1,3,4-thiadiazol-2-yl}piperidin-4-amine
 463¹ 3-[2-(difluoromethyl)-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine and
 464¹ 6-[2-(difluoromethyl)-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
 465¹

wherein a form of the compound is selected from the group consisting of a prodrug, salt, hydrate, solvate, clathrate, isotopologue, racemate, enantiomer, diastereomer, stereoisomer, polymorph and tautomer form thereof.

Another embodiment of the use of a compound salt of Formula (I) or a form thereof includes a method of use of a compound salt of Formula (I) or a form thereof for treating or ameliorating HD in a subject in need thereof, comprising administering an effective amount of the compound salt of Formula (I) or a form thereof to the subject, selected from the group

5 consisting of:

Cpd	Name
32	2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(trifluoromethyl)phenol hydrochloride
53	2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(1H-pyrazol-1-yl)phenol hydrochloride
65	2-(6-piperazin-1-yl-pyridazin-3-yl)-5-1H-pyrazol-1-yl-phenol hydrochloride
82	3-(6-(piperazin-1-yl)pyridazin-3-yl)naphthalene-2,7-diol trifluoroacetate
86	3-(6-(1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl)naphthalene-2,7-diol trifluoroacetate
88	3-(6-(1-methyl-1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl)naphthalene-2,7-diol trifluoroacetate
89	3-(6-(piperidin-4-yl)pyridazin-3-yl)naphthalene-2,7-diol trifluoroacetate
151	6-hydroxy-5-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-2,3-dihydro-1H-inden-1-one oxime hydrochloride
153	2-amino-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-8H-indeno[1,2-d]thiazol-5-ol hydrochloride
154	9-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5,6-dihydroimidazo[5,1-a]isoquinolin-8-ol hydrochloride
160	3-fluoro-5-(2-methoxypyridin-4-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol hydrochloride
161	4-(3-fluoro-5-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)pyridin-2(1H)-one hydrochloride
162	4-(3-fluoro-5-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)-1-methylpyridin-2(1H)-one hydrochloride
163	5-(3-fluoro-5-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)-1-methylpyridin-2(1H)-one hydrochloride
164	3-fluoro-5-(1H-pyrazol-4-yl)-2-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)phenol hydrochloride
165	5-chloro-3-fluoro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol hydrochloride
166	3-fluoro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1H-pyrazol-4-yl)phenol hydrochloride
167	3-fluoro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1-methyl-1H-pyrazol-4-yl)phenol hydrochloride

Cpd	Name
226	3-(1H-imidazol-1-yl)-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol hydrochloride
227	6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinoline-3,7-diol formate
231	7-hydroxy-1-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-2(1H)-one hydrochloride
238	4-methoxy-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-6-ol formate
239	7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinoxalin-6-ol hydrochloride
254	2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(tetrahydro-2H-pyran-4-yl)quinolin-7-ol formate
256	4-(dimethylamino)-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-quinolin-7-ol formate
281	6-hydroxy-2-methyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-1(2H)-one hydrochloride
322	3-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)naphthalen-2-ol hydrobromide
342	N-methyl-5-(5-(1-methyl-1H-pyrazol-4-yl)pyridin-2-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine hydrochloride
396	3-fluoro-2-(5-((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-1,3,4-thiadiazol-2-yl)-5-(1H-pyrazol-4-yl)phenol dihydrochloride
405	2-(2-chloro-4-(1H-pyrazol-4-yl)phenyl)-5-((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-1,3,4-thiadiazole hydrochloride
406	2-(2-chloro-4-(1H-pyrazol-4-yl)phenyl)-5-(2,7-diazaspiro[4.5]decan-2-yl)-1,3,4-thiadiazole hydrochloride
407	(R)-(4-(5-(2-chloro-4-(1H-pyrazol-4-yl)phenyl)-1,3,4-thiadiazol-2-yl)piperazin-2-yl)methanol hydrochloride
411	2-{6-[8-azabicyclo[3.2.1]oct-3-yl(methyl)amino]pyridazin-3-yl}-5-(1H-pyrazol-4-yl)phenol hydrochloride
412	2-[6-(8-azabicyclo[3.2.1]oct-3-ylamino)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol hydrochloride
413	5-(1H-pyrazol-4-yl)-2-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl}phenol hydrochloride
415	2-[6-(8-azabicyclo[3.2.1]oct-3-yloxy)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol hydrochloride
419	2-{6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl}-5-(4-nitro-1H-pyrazol-1-yl)phenol dihydrochloride

Cpd	Name
422	2-[6-(1-methyl-1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol trihydrochloride
424	5-(1H-pyrazol-4-yl)-2-[6-(1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]phenol trihydrochloride
425	2-[6-(1-ethyl-1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol trihydrochloride
426	2-{6-[methyl(piperidin-4-yl)amino]pyridazin-3-yl}-5-(1H-pyrazol-4-yl)phenol tetrahydrochloride
427	2-[6-(piperidin-4-ylamino)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol tetrahydrochloride
428	6-[2,5-difluoro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine tetrahydrochloride
429	2-[6-(8-azabicyclo[3.2.1]oct-2-en-3-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol hydrochloride
430	6-[2,3-difluoro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine hydrochloride
431	3-[2,5-difluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine trihydrochloride
433	2-{6-[(1R,5S)-8-azabicyclo[3.2.1]oct-3-ylamino]pyridazin-3-yl}-5-(1H-pyrazol-4-yl)phenol hydrochloride
434	6-[2-methoxy-6-(1H-pyrazol-4-yl)pyridin-3-yl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine hydrochloride
436	2-{6-[(2,6-dimethylpiperidin-4-yl)oxy]pyridazin-3-yl}-5-(1H-pyrazol-4-yl)phenol trihydrochloride
437	3-[2-fluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine hydrochloride
439	2-[6-(2,7-diazaspiro[3.5]non-2-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol tetrahydrochloride
448	3-[5-(1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine hydrochloride
450	3-[4-(1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine hydrochloride
456	6-[2-chloro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine trihydrochloride
458	3-[2,6-difluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine trihydrochloride
460	3-{6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl}-6-(1H-pyrazol-4-yl)pyridin-2-ol hydrochloride
462	N,2,2,6,6-pentamethyl-N-{5-[3-(1H-pyrazol-4-yl)phenoxy]-1,3,4-thiadiazol-2-yl}piperidin-4-amine hydrochloride

Cpd	Name
463	N,2,2,6,6-pentamethyl-N-{5-[4-(1H-pyrazol-4-yl)phenoxy]-1,3,4-thiadiazol-2-yl}piperidin-4-amine hydrochloride
464	3-[2-(difluoromethyl)-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine hydrochloride and
465	6-[2-(difluoromethyl)-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine hydrochloride

wherein a form of the compound salt is selected from the group consisting of a prodrug, hydrate, solvate, clathrate, isotopologue, racemate, enantiomer, diastereomer, stereoisomer, polymorph and tautomer form thereof.

In one aspect, the compound salt of Formula (I) or a form thereof includes a compound
5 salt selected from the group consisting of:

- 411 2-{6-[8-azabicyclo[3.2.1]oct-3-yl(methyl)amino]pyridazin-3-yl}-5-(1H-pyrazol-4-yl)phenol hydrochloride
- 412 2-[6-(8-azabicyclo[3.2.1]oct-3-ylamino)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol hydrochloride
- 413 5-(1H-pyrazol-4-yl)-2-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl]phenol hydrochloride
- 415 2-[6-(8-azabicyclo[3.2.1]oct-3-yloxy)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol hydrochloride
- 419 2-{6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl}-5-(4-nitro-1H-pyrazol-1-yl)phenol dihydrochloride
- 422 2-[6-(1-methyl-1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol trihydrochloride
- 424 5-(1H-pyrazol-4-yl)-2-[6-(1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]phenol trihydrochloride
- 425 2-[6-(1-ethyl-1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol trihydrochloride
- 426 2-{6-[methyl(piperidin-4-yl)amino]pyridazin-3-yl}-5-(1H-pyrazol-4-yl)phenol tetrahydrochloride
- 427 2-[6-(piperidin-4-ylamino)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol tetrahydrochloride
- 428 6-[2,5-difluoro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine tetrahydrochloride
- 429 2-[6-(8-azabicyclo[3.2.1]oct-2-en-3-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol hydrochloride
- 430 6-[2,3-difluoro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine hydrochloride

- 431 3-[2,5-difluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine trihydrochloride
- 433 2-{6-[(1R,5S)-8-azabicyclo[3.2.1]oct-3-ylamino]pyridazin-3-yl}-5-(1H-pyrazol-4-yl)phenol hydrochloride
- 434 6-[2-methoxy-6-(1H-pyrazol-4-yl)pyridin-3-yl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine hydrochloride
- 436 2-{6-[(2,6-dimethylpiperidin-4-yl)oxy]pyridazin-3-yl}-5-(1H-pyrazol-4-yl)phenol trihydrochloride
- 437 3-[2-fluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine hydrochloride
- 439 2-[6-(2,7-diazaspiro[3.5]non-2-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol tetrahydrochloride
- 448 3-[5-(1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine hydrochloride
- 450 3-[4-(1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine hydrochloride
- 456 6-[2-chloro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine trihydrochloride
- 458 3-[2,6-difluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine trihydrochloride
- 460 3-{6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl}-6-(1H-pyrazol-4-yl)pyridin-2-ol hydrochloride
- 462 N,2,2,6,6-pentamethyl-N-{5-[3-(1H-pyrazol-4-yl)phenoxy]-1,3,4-thiadiazol-2-yl}piperidin-4-amine hydrochloride
- 463 N,2,2,6,6-pentamethyl-N-{5-[4-(1H-pyrazol-4-yl)phenoxy]-1,3,4-thiadiazol-2-yl}piperidin-4-amine hydrochloride
- 464 3-[2-(difluoromethyl)-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine hydrochloride and
- 465 6-[2-(difluoromethyl)-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine hydrochloride

wherein a form of the compound salt is selected from the group consisting of a prodrug, hydrate, solvate, clathrate, isotopologue, racemate, enantiomer, diastereomer, stereoisomer, polymorph and tautomer form thereof.

- Another embodiment of the present description includes a method of use of a compound of Formula (I) or a form thereof for treating or ameliorating HD in a subject in need thereof, comprising administering an effective amount of the compound of Formula (I) or a form thereof to the subject.

Another embodiment of the present description includes a method of use of a compound of Formula (I) or a form thereof for treating or ameliorating HD in a subject in need thereof, comprising administering an effective amount of the compound salt of Formula (I) or a form thereof to the subject.

5 Another embodiment of the present description includes a use of the compound of Formula (I) or a form thereof for treating or ameliorating HD in a subject in need thereof, comprising administering an effective amount of the compound of Formula (I) or a form thereof to the subject.

10 Another embodiment of the present description includes a use of the compound salt of Formula (I) or a form thereof for treating or ameliorating HD in a subject in need thereof, comprising administering an effective amount of the compound salt of Formula (I) or a form thereof to the subject.

Chemical Definitions

As used herein, the term “C₁₋₄alkyl” generally refers to saturated hydrocarbon radicals having from one to four carbon atoms in a straight or branched chain configuration, including, without limitation, methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, and the like. In some embodiments, C₁₋₄alkyl includes C₁₋₃alkyl, C₁₋₂alkyl, and the like. A C₁₋₄alkyl radical may be optionally substituted where allowed by available valences.

As used herein, the term “C₂₋₆alkenyl” generally refers to partially unsaturated hydrocarbon radicals having from two to five carbon atoms in a straight or branched chain configuration and one or more carbon-carbon double bonds therein, including, without limitation, ethenyl, allyl, propenyl and the like. In some embodiments, C₂₋₆alkenyl includes C₂₋₄alkenyl, C₂₋₃alkenyl, and the like. A C₂₋₆alkenyl radical may be optionally substituted where allowed by available valences.

25 As used herein, the term “C₁₋₄alkoxy” generally refers to saturated hydrocarbon radicals having from one to four carbon atoms in a straight or branched chain configuration of the formula: -O-C₁₋₄alkyl, including, without limitation, methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, isobutoxy, sec-butoxy, tert-butoxy, and the like. In some embodiments, C₁₋₄alkoxy includes C₁₋₃alkoxy, C₁₋₂alkoxy and the like. A C₁₋₄alkoxy radical may be optionally substituted
30 where allowed by available valences.

As used herein, the term “C₃₋₁₄cycloalkyl” generally refers to a saturated monocyclic, bicyclic or polycyclic hydrocarbon radical, including, without limitation, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, 1H-indanyl, indenyl, tetrahydro-naphthalenyl and the like. In some embodiments, C₃₋₁₄cycloalkyl includes

5 C₃₋₁₀cycloalkyl, C₃₋₈cycloalkyl, C₃₋₇cycloalkyl, C₅₋₈cycloalkyl, C₉₋₁₀cycloalkyl and the like. A C₃₋₁₄cycloalkyl radical may be optionally substituted where allowed by available valences.

As used herein, the term “C₃₋₁₄cycloalkenyl” generally refers to a partially unsaturated monocyclic, bicyclic or polycyclic hydrocarbon radical having one or more chemically stable carbon-carbon double bonds therein, including, without limitation, cyclopropenyl, cyclobutenyl, 10 cyclopentenyl, cyclohexenyl, cycloheptenyl, cyclooctenyl and the like. In some embodiments, C₃₋₁₄cycloalkenyl includes C₃₋₇cycloalkenyl, C₃₋₈cycloalkenyl, C₅₋₈cycloalkenyl, C₃₋₁₀cycloalkenyl and the like. A C₃₋₁₄cycloalkenyl radical may be optionally substituted where allowed by available valences.

As used herein, the term “aryl” generally refers to a monocyclic, bicyclic or polycyclic aromatic carbon atom ring structure radical, including, without limitation, phenyl, naphthyl, 15 anthracenyl, fluorenyl, azulenyl, phenanthrenyl and the like. An aryl radical may be optionally substituted where allowed by available valences.

As used herein, the term “heteroaryl” generally refers to a monocyclic, bicyclic or polycyclic aromatic carbon atom ring structure radical in which one or more carbon atom ring 20 members have been replaced, where allowed by structural stability, with one or more heteroatoms, such as an O, S or N atom, including, without limitation, furanyl, thienyl (also referred to as thiophenyl), pyrrolyl, pyrazolyl, imidazolyl, isoxazolyl, isothiazolyl, oxazolyl, thiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyranyl, thiopyranyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, triazinyl, indolyl, indazolyl, indoliziny, benzofuranyl, 25 benzothienyl, benzimidazolyl, benzothiazolyl, benzooxazolyl, 9H-purinyl, quinoxaliny, isoindolyl, quinoliny, isoquinoliny, quinazoliny, acridiny, phthalaziny, imidazo[1,2-a]pyridiny, imidazo[1,5-a]pyridiny, imidazo[5,1-a]isoquinoliny, 1,4-dihydroindeno[1,2-c]-1H-pyrazolyl, 2,3-dihydro-1H-inden-1-one, 2,3-dihydro-1H-indenyl, 3,4-dihydroquinolin-2(1H)-one, 5,6-dihydroimidazo[5,1-a]isoquinoliny, 8H-indeno[1,2-d]thiazolyl, 30 benzo[c][1,2,5]oxadiazolyl, benzo[d]oxazol-2(3H)-one, quinolin-2(1H)-one, quinazolin-4(1H)-one, quinazoline-2,4(1H,3H)-dione, benzo-[d]oxazolyl, pyrazolo[1,5-a]pyridiny, and the like.

A heteroaryl radical may be optionally substituted on a carbon or nitrogen atom ring member where allowed by available valences.

As used herein, the term “heterocyclyl” generally refers to a saturated or partially unsaturated monocyclic, bicyclic or polycyclic carbon atom ring structure radical in which one or more carbon atom ring members have been replaced, where allowed by structural stability, with a heteroatom, such as an O, S or N atom, including, without limitation, oxiranyl, oxetanyl, azetidiny, dihydrofuranyl, tetrahydrofuranyl, dihydrothienyl, tetrahydrothienyl, pyrrolinyl, pyrrolidinyl, dihydropyrazolyl, pyrazolinyl, pyrazolidinyl, dihydroimidazolyl, imidazoliny, imidazolidinyl, isoxazoliny, isoxazolidinyl, isothiazoliny, isothiazolidinyl, oxazoliny, oxazolidinyl, thiazoliny, thiazolidinyl, triazoliny, triazolidinyl, oxadiazoliny, oxadiazolidinyl, thiadiazoliny, thiadiazolidinyl, tetrazoliny, tetrazolidinyl, dihydro-2H-pyranyl, dihydro-pyridiny, tetrahydro-pyridiny, 1,2,3,6-tetrahydropyridiny, hexahydro-pyridiny, dihydro-pyrimidiny, tetrahydro-pyrimidiny, 1,4,5,6-tetrahydropyrimidiny, dihydro-pyraziny, tetrahydro-pyraziny, dihydro-pyridaziny, tetrahydro-pyridaziny, piperaziny, piperidiny, morpholiny, thiomorpholiny, dihydro-triaziny, tetrahydro-triaziny, hexahydro-triaziny, 1,4-diazepanyl, dihydro-indolyl, indoliny, tetrahydro-indolyl, dihydro-indazolyl, tetrahydro-indazolyl, dihydro-isoindolyl, dihydro-benzofuranyl, tetrahydro-benzofuranyl, dihydro-benzothienyl, tetrahydro-benzothienyl, dihydro-benzimidazolyl, tetrahydro-benzimidazolyl, dihydro-benzooxazolyl, 2,3-dihydrobenzo[d]oxazolyl, tetrahydro-benzooxazolyl, dihydro-benzooxaziny, 3,4-dihydro-2H-benzo[b][1,4]oxaziny, tetrahydro-benzooxaziny, benzo[1,3]dioxolyl, benzo[1,4]dioxanyl, dihydro-puriny, tetrahydro-puriny, dihydro-quinoliny, tetrahydro-quinoliny, 1,2,3,4-tetrahydroquinoliny, dihydro-isoquinoliny, 3,4-dihydroisoquinolin-(1H)-yl, tetrahydro-isoquinoliny, 1,2,3,4-tetrahydroisoquinoliny, dihydro-quinazoliny, tetrahydro-quinazoliny, dihydro-quinoxaliny, tetrahydro-quinoxaliny, 1,2,3,4-tetrahydroquinoxaliny, 1,3-dioxolanyl, 2,5-dihydro-1H-pyrroly, 4,5-dihydro-1H-imidazolyl, tetrahydro-2H-pyranyl, hexahydropyrrolo[3,4-b][1,4]oxazin-(2H)-yl, (4aR,7aS)-hexahydropyrrolo[3,4-b][1,4]oxazin-(4aH)-yl, 3,4-dihydro-2H-pyrido[3,2-b][1,4]oxaziny, (cis)-octahydrocyclopenta[c]pyrroly, hexahydropyrrolo[3,4-b]pyrrol-(1H)-yl, (3aR,6aR)-hexahydropyrrolo[3,4-b]pyrrol-(1H)-yl, (3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-(1H)-yl, 5H-pyrrolo[3,4-b]pyridin-(7H)-yl, 5,7-dihydro-6H-pyrrolo[3,4-b]pyridiny, tetrahydro-1H-pyrrolo[3,4-b]pyridin-(2H,7H,7aH)-yl, hexahydro-1H-

pyrrolo[3,4-b]pyridin-(2H)-yl, (4aR,7aR)-hexahydro-1H-pyrrolo[3,4-b]pyridin-(2H)-yl, octahydro-6H-pyrrolo[3,4-b]pyridinyl, 2,3,4,9-tetrahydro-1H-carbazolyl, 1,2,3,4-tetrahydropyrazino[1,2-a]indolyl, 2,3-dihydro-1H-pyrrolo[1,2-a]indolyl, (3aR,6aR)-hexahydrocyclopenta[c]pyrrol-(1H)-yl, (3aR,4R,6aS)-hexahydrocyclopenta[c]pyrrol-(1H)-yl, 5 (3aR,4S,6aS)-hexahydrocyclopenta[c]pyrrol-(1H)-yl, (3aR,5r,6aS)-hexahydrocyclopenta[c]pyrrol-(1H)-yl, 1,3-dihydro-2H-isoindolyl, octahydro-2H-isoindolyl, (3aS)-1,3,3a,4,5,6-hexahydro-2H-isoindolyl, (3aR,4R,7aS)-1H-isoindolyl, (3H,3aH,4H,5H,6H,7H,7aH)-yl, (3aR,7aS)-octahydro-2H-isoindolyl, (3aR,4R,7aS)-octahydro-2H-isoindolyl, (3aR,4S,7aS)-octahydro-2H-isoindolyl, 2,5-diazabicyclo[2.2.1]heptanyl, 2-azabicyclo[2.2.1]heptenyl, 3-azabicyclo[3.1.0]hexanyl, 3,6-diazabicyclo[3.1.0]hexanyl, (1R,5S)- 10 3-azabicyclo[3.1.0]hexanyl, (1S,5R)-3-azabicyclo[3.2.0]heptanyl, 5-azaspiro[2.4]heptanyl, 2,6-diazaspiro[3.3]heptanyl, 2,5-diazaspiro[3.4]octanyl, 2,6-diazaspiro[3.4]octanyl, 2,7-diazaspiro[3.5]nonanyl, 2,7-diazaspiro[4.4]nonanyl, 2-azaspiro[4.5]decanyl, 2,8-diazaspiro[4.5]decanyl, 3,6-diazabicyclo[3.2.1]octyl, 1,4-dihydroindeno[1,2-c]pyrazolyl, 15 dihydropyranyl, dihydropyridinyl, dihydroquinolinyl, 8H-indeno[1,2-d]thiazolyl, tetrahydroimidazo[1,2-a]pyridinyl, pyridin-2(1H)-one, (1R,5S)-8-azabicyclo[3.2.1]octyl, 8-azabicyclo[3.2.1]oct-2-enyl and the like. A heterocyclyl radical may be optionally substituted on a carbon or nitrogen atom ring member where allowed by available valences.

As used herein, the term “C₂₋₄alkenyl-amino-carbonyl” refers to a radical of the
20 formula: -C(=O)-NH-C₂₋₄alkenyl.

As used herein, the term “C₁₋₄alkoxy-C₁₋₄alkoxy” refers to a radical of the
formula: -O-C₁₋₄alkyl-O-C₁₋₄alkyl.

As used herein, the term “C₁₋₄alkoxy-carbonyl” refers to a radical of the
formula: -C(=O)-O-C₁₋₄alkyl.

As used herein, the term “C₁₋₄alkoxy-carbonyl-amino” refers to a radical of the
25 formula: -NH-C(=O)-O-C₁₋₄alkyl.

As used herein, the term “C₁₋₄alkoxy-carbonyl-amino-C₁₋₄alkoxy” refers to a radical of
the formula: -O-C₁₋₄alkyl-NH-C(=O)-O-C₁₋₄alkyl.

As used herein, the term “C₁₋₄alkyl-C₁₋₄alkoxy” refers to a radical of the
30 formula: -O-C₁₋₄alkyl-C₁₋₄alkyl.

As used herein, the term “C₁₋₄alkyl-amino” refers to a radical of the formula: -NH-C₁₋₄alkyl.

As used herein, the term “(C₁₋₄alkyl)₂-amino” refers to a radical of the formula: -N(C₁₋₄alkyl)₂.

5 As used herein, the term “C₁₋₄alkyl-amino-C₁₋₄alkoxy” refers to a radical of the formula: -O-C₁₋₄alkyl-NH-C₁₋₄alkyl.

As used herein, the term “(C₁₋₄alkyl)₂-amino-C₁₋₄alkoxy” refers to a radical of the formula: -O-C₁₋₄alkyl-N(C₁₋₄alkyl)₂.

10 As used herein, the term “C₁₋₄alkyl-amino-C₁₋₄alkyl” refers to a radical of the formula: -C₁₋₄alkyl-NH-C₁₋₄alkyl.

As used herein, the term “(C₁₋₄alkyl)₂-amino-C₁₋₄alkyl” refers to a radical of the formula: -C₁₋₄alkyl-N(C₁₋₄alkyl)₂.

As used herein, the term “C₁₋₄alkyl-amino-carbonyl” refers to a radical of the formula: -C(=O)-NH-C₁₋₄alkyl.

15 As used herein, the term “(C₁₋₄alkyl)₂-amino-carbonyl” refers to a radical of the formula: -C(=O)-N(C₁₋₄alkyl)₂.

As used herein, the term “C₁₋₄alkyl-amino-carbonyl-C₁₋₄alkyl” refers to a radical of the formula: -C₁₋₄alkyl-C(=O)-NH-C₁₋₄alkyl.

20 As used herein, the term “(C₁₋₄alkyl)₂-amino-carbonyl-C₁₋₄alkyl” refers to a radical of the formula: -C₁₋₄alkyl-C(=O)-N(C₁₋₄alkyl)₂.

As used herein, the term “C₁₋₄alkyl-carbonyl” refers to a radical of the formula: -C(=O)-C₁₋₄alkyl.

As used herein, the term “C₁₋₄alkyl-carbonyl-amino” refers to a radical of the formula: -NH-C(=O)-C₁₋₄alkyl.

25 As used herein, the term “C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkoxy” refers to a radical of the formula: -O-C₁₋₄alkyl-NH-C(=O)-C₁₋₄alkyl.

As used herein, the term “C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkyl” refers to a radical of the formula: -C₁₋₄alkyl-NH-C(=O)-C₁₋₄alkyl.

As used herein, the term “amino” refers to a radical of the formula: -NH₂.

30 As used herein, the term “amino-C₁₋₄alkoxy” refers to a radical of the formula: -O-C₁₋₄alkyl-NH₂.

As used herein, the term “amino-C₁₋₄alkyl” refers to a radical of the formula: -C₁₋₄alkyl-NH₂.

As used herein, the term “amino-carbonyl” refers to a radical of the formula: -C(=O)-NH₂.

5 As used herein, the term “cyano” refers to a radical of the formula: -CN.

As used herein, the term “C₃₋₇cycloalkyl-C₁₋₄alkoxy” refers to a radical of the formula: -O-C₁₋₄alkyl-C₃₋₇cycloalkyl.

As used herein, the term “halo-C₁₋₄alkoxy” refers to a radical of the formula: -O-C₁₋₄alkyl-halo, wherein C₁₋₄alkyl may be partially or completely substituted where
10 allowed by available valences with one or more halogen atoms. In some embodiments, halo-C₁₋₄alkoxy includes halo-C₁₋₆alkoxy, halo-C₁₋₄alkoxy and the like.

As used herein, the term “halo-C₁₋₄alkyl” refers to a radical of the formula: -C₁₋₄alkyl-halo, wherein C₁₋₄alkyl may be partially or completely substituted where
allowed by available valences with one or more halogen atoms. In some embodiments,
15 halo-C₁₋₄alkyl includes halo-C₁₋₆alkyl, halo-C₁₋₄alkyl and the like.

As used herein, the term “heteroaryl-C₁₋₄alkyl” refers to a radical of the formula: -C₁₋₄alkyl-heteroaryl.

As used herein, the term “heteroaryl-C₁₋₄alkyl-amino” refers to a radical of the formula: -NH-C₁₋₄alkyl-heteroaryl.

20 As used herein, the term “heteroaryl-C₁₋₄alkyl-amino-carbonyl” refers to a radical of the formula: -C(=O)-NH-C₁₋₄alkyl-heteroaryl.

As used herein, the term “heteroaryl-C₁₋₄alkyl-amino-carbonyl-C₁₋₄alkyl” refers to a radical of the formula: -C₁₋₄alkyl-C(=O)-NH-C₁₋₄alkyl-heteroaryl.

As used herein, the term “heteroaryl-C₁₋₄alkyl-carbonyl-amino” refers to a radical of the
25 formula: -NH-C(=O)-C₁₋₄alkyl-heteroaryl.

As used herein, the term “heteroaryl-C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkyl” refers to a radical of the formula: -C₁₋₄alkyl-NH-C(=O)-C₁₋₄alkyl-heteroaryl.

As used herein, the term “heterocyclyl-C₁₋₄alkoxy” refers to a radical of the formula: -C₁₋₄alkoxy-heterocyclyl.

30 As used herein, the term “heterocyclyl-C₁₋₄alkyl” refers to a radical of the formula: -C₁₋₄alkyl-heterocyclyl.

As used herein, the term “hydroxyl” refers to a radical of the formula: -OH.

As used herein, the term “hydroxyl-C₁₋₄alkoxy” refers to a radical of the formula: -O-C₁₋₄alkyl-OH, wherein C₁₋₄alkyl may be partially or completely substituted where allowed by available valences with one or more hydroxy radicals.

5 As used herein, the term “hydroxyl-C₁₋₄alkyl” refers to a radical of the formula: -C₁₋₄alkyl-OH, wherein C₁₋₄alkyl may be partially or completely substituted where allowed by available valences with one or more hydroxy radicals.

As used herein, the term “hydroxyl-C₁₋₄alkyl-amino” refers to a radical of the formula: -NH-C₁₋₄alkyl-OH, wherein C₁₋₄alkyl may be partially or completely substituted where
10 allowed by available valences with one or more hydroxyl radicals.

As used herein, the term “hydroxyl-imino” refers to the =NOH radical of the formula: C(=NOH).

As used herein, the term “oxo” refers to the radical of the formula: C=O.

As used herein, the term “phenyl-C₁₋₄alkoxy” refers to a radical of the
15 formula: -C₁₋₄alkoxy-phenyl.

As used herein, the term “substituent” means positional variables on the atoms of a core molecule that are substituted at a designated atom position, replacing one or more hydrogens on the designated atom, provided that the designated atom’s normal valency is not exceeded, and that the substitution results in a stable compound. Combinations of substituents and/or variables
20 are permissible only if such combinations result in stable compounds. A person of ordinary skill in the art should note that any carbon as well as heteroatom with valences that appear to be unsatisfied as described or shown herein is assumed to have a sufficient number of hydrogen atom(s) to satisfy the valences described or shown. In certain instances one or more substituents having a double bond (e.g., “oxo” or “=O”) as the point of attachment may be described, shown
25 or listed herein within a substituent group, wherein the structure may only show a single bond as the point of attachment to the core structure of Formula (I). A person of ordinary skill in the art would understand that, while only a single bond is shown, a double bond is intended for those substituents.

As used herein, the term “and the like,” with reference to the definitions of chemical
30 terms provided herein, means that variations in chemical structures that could be expected by one skilled in the art include, without limitation, isomers (including chain, branching or positional

structural isomers), hydration of ring systems (including saturation or partial unsaturation of monocyclic, bicyclic or polycyclic ring structures) and all other variations where allowed by available valences which result in a stable compound.

For the purposes of this description, where one or more substituent variables for a compound of Formula (I) or a form thereof encompass functionalities incorporated into a compound of Formula (I), each functionality appearing at any location within the disclosed compound may be independently selected, and as appropriate, independently and/or optionally substituted.

As used herein, the terms “independently selected,” or “each selected” refer to functional variables in a substituent list that may occur more than once on the structure of Formula (I), the pattern of substitution at each occurrence is independent of the pattern at any other occurrence. Further, the use of a generic substituent variable on any formula or structure for a compound described herein is understood to include the replacement of the generic substituent with species substituents that are included within the particular genus, *e.g.*, aryl may be replaced with phenyl or naphthalenyl and the like, and that the resulting compound is to be included within the scope of the compounds described herein.

As used herein, the terms “each instance of” or “in each instance, when present,” when used preceding a phrase such as “...C₃₋₁₄cycloalkyl, C₃₋₁₄cycloalkyl-C₁₋₄alkyl, aryl, aryl-C₁₋₄alkyl, heteroaryl, heteroaryl-C₁₋₄alkyl, heterocyclyl and heterocyclyl-C₁₋₄alkyl,” are intended to refer to the C₃₋₁₄cycloalkyl, aryl, heteroaryl and heterocyclyl ring systems when each are present either alone or as a substituent.

As used herein, the term “optionally substituted” means optional substitution with the specified substituent variables, groups, radicals or moieties.

Compound Forms

As used herein, the term “form” means a compound of Formula (I) having a form selected from the group consisting of a free acid, free base, prodrug, salt, hydrate, solvate, clathrate, isotopologue, racemate, enantiomer, diastereomer, stereoisomer, polymorph and tautomer form thereof.

In certain embodiments described herein, the form of the compound of Formula (I) is a free acid, free base or salt thereof.

In certain embodiments described herein, the form of the compound of Formula (I) is a salt thereof.

In certain embodiments described herein, the form of the compound of Formula (I) is an isotopologue thereof.

5 In certain embodiments described herein, the form of the compound of Formula (I) is a stereoisomer, racemate, enantiomer or diastereomer thereof.

In certain embodiments described herein, the form of the compound of Formula (I) is a tautomer thereof.

10 In certain embodiments described herein, the form of the compound of Formula (I) is a pharmaceutically acceptable form.

In certain embodiments described herein, the compound of Formula (I) or a form thereof is isolated for use.

As used herein, the term “isolated” means the physical state of a compound of Formula (I) or a form thereof after being isolated and/or purified from a synthetic process (*e.g.*, from a reaction mixture) or natural source or combination thereof according to an isolation or purification process or processes described herein or which are well known to the skilled artisan (*e.g.*, chromatography, recrystallization and the like) in sufficient purity to be characterized by standard analytical techniques described herein or well known to the skilled artisan.

As used herein, the term “protected” means that a functional group in a compound of Formula (I) or a form thereof is in a form modified to preclude undesired side reactions at the protected site when the compound is subjected to a reaction. Suitable protecting groups will be recognized by those with ordinary skill in the art as well as by reference to standard textbooks such as, for example, T.W. Greene *et al*, *Protective Groups in organic Synthesis* (1991), Wiley, New York. Such functional groups include hydroxy, phenol, amino and carboxylic acid.

25 Suitable protecting groups for hydroxy or phenol include trialkylsilyl or diarylalkylsilyl (*e.g.*, *t*-butyldimethylsilyl, *t*-butyldiphenylsilyl or trimethylsilyl), tetrahydropyranyl, benzyl, substituted benzyl, methyl, methoxymethanol, and the like. Suitable protecting groups for amino, amidino and guanidino include *t*-butoxycarbonyl, benzyloxycarbonyl, and the like. Suitable protecting groups for carboxylic acid include alkyl, aryl or arylalkyl esters. In certain instances,

30 the protecting group may also be a polymer resin, such as a Wang resin or a 2-chlorotrityl-chloride resin. Protecting groups may be added or removed in accordance with standard

techniques, which are well-known to those skilled in the art and as described herein. It will also be appreciated by those skilled in the art, although such protected derivatives of compounds described herein may not possess pharmacological activity as such, they may be administered to a subject and thereafter metabolized in the body to form compounds described herein which are pharmacologically active. Such derivatives may therefore be described as "prodrugs". All prodrugs of compounds described herein are included within the scope of the use described herein.

As used herein, the term "prodrug" means a form of an instant compound (*e.g.*, a drug precursor) that is transformed *in vivo* to yield an active compound of Formula (I) or a form thereof. The transformation may occur by various mechanisms (*e.g.*, by metabolic and/or non-metabolic chemical processes), such as, for example, by hydrolysis and/or metabolism in blood, liver and/or other organs and tissues. A discussion of the use of prodrugs is provided by T. Higuchi and W. Stella, "Pro-drugs as Novel Delivery Systems," Vol. 14 of the A.C.S. Symposium Series, and in *Bioreversible Carriers in Drug Design*, ed. Edward B. Roche, American Pharmaceutical Association and Pergamon Press, 1987.

In one example, when a compound of Formula (I) or a form thereof contains a carboxylic acid functional group, a prodrug can comprise an ester formed by the replacement of the hydrogen atom of the acid group with a functional group such as alkyl and the like. In another example, when a compound of Formula (I) or a form thereof contains a hydroxyl functional group, a prodrug form can be prepared by replacing the hydrogen atom of the hydroxyl with another functional group such as alkyl, alkylcarbonyl or a phosphonate ester and the like. In another example, when a compound of Formula (I) or a form thereof contains an amine functional group, a prodrug form can be prepared by replacing one or more amine hydrogen atoms with a functional group such as alkyl or substituted carbonyl. Pharmaceutically acceptable prodrugs of compounds of Formula (I) or a form thereof include those compounds substituted with one or more of the following groups: carboxylic acid esters, sulfonate esters, amino acid esters, phosphonate esters and mono-, di- or triphosphate esters or alkyl substituents, where appropriate. As described herein, it is understood by a person of ordinary skill in the art that one or more of such substituents may be used to provide a compound of Formula (I) or a form thereof as a prodrug.

One or more compounds described herein may exist in unsolvated as well as solvated forms with pharmaceutically acceptable solvents such as water, ethanol, and the like, and the description herein is intended to embrace both solvated and unsolvated forms.

As used herein, the term "solvate" means a physical association of a compound described herein with one or more solvent molecules. This physical association involves varying degrees of ionic and covalent bonding, including hydrogen bonding. In certain instances the solvate will be capable of isolation, for example when one or more solvent molecules are incorporated in the crystal lattice of the crystalline solid. As used herein, "solvate" encompasses both solution-phase and isolatable solvates. Non-limiting examples of suitable solvates include ethanolates, methanolates, and the like.

As used herein, the term "hydrate" means a solvate wherein the solvent molecule is water.

The compounds of Formula (I) can form salts, which are intended to be included within the scope of this description. Reference to a compound of Formula (I) or a form thereof herein is understood to include reference to salt forms thereof, unless otherwise indicated. The term "salt(s)", as employed herein, denotes acidic salts formed with inorganic and/or organic acids, as well as basic salts formed with inorganic and/or organic bases. In addition, when a compound of Formula (I) or a form thereof contains both a basic moiety, such as, without limitation an amine moiety, and an acidic moiety, such as, but not limited to a carboxylic acid, zwitterions ("inner salts") may be formed and are included within the term "salt(s)" as used herein.

The term "pharmaceutically acceptable salt(s)", as used herein, means those salts of compounds described herein that are safe and effective (*i.e.*, non-toxic, physiologically acceptable) for use in mammals and that possess biological activity, although other salts are also useful. Salts of the compounds of the Formula (I) may be formed, for example, by reacting a compound of Formula (I) or a form thereof with an amount of acid or base, such as an equivalent amount, in a medium such as one in which the salt precipitates or in an aqueous medium followed by lyophilization.

Pharmaceutically acceptable salts include one or more salts of acidic or basic groups present in compounds described herein. Embodiments of acid addition salts include, and are not limited to, acetate, ascorbate, benzoate, benzenesulfonate, bisulfate, bitartrate, borate, bromide, butyrate, chloride, citrate, camphorate, camphorsulfonate, ethanesulfonate, formate, fumarate,

gentisinate, gluconate, glucaronate, glutamate, iodide, isonicotinate, lactate, maleate, methanesulfonate, naphthalenesulfonate, nitrate, oxalate, pamoate, pantothenate, phosphate, propionate, saccharate, salicylate, succinate, sulfate, tartrate, thiocyanate, toluenesulfonate (also known as tosylate), trifluoroacetate salts and the like. Certain embodiments of acid addition salts
5 include chloride, dichloride, trichloride, bromide, acetate, formate or trifluoroacetate salts.

Additionally, acids which are generally considered suitable for the formation of pharmaceutically useful salts from basic pharmaceutical compounds are discussed, for example, by P. Stahl *et al*, Camille G. (eds.) *Handbook of Pharmaceutical Salts. Properties, Selection and Use*. (2002) Zurich: Wiley-VCH; S. Berge *et al*, *Journal of Pharmaceutical Sciences* (1977)
10 66(1) 1-19; P. Gould, *International J. of Pharmaceutics* (1986) 33, 201-217; Anderson *et al*, *The Practice of Medicinal Chemistry* (1996), Academic Press, New York; and in *The Orange Book* (Food & Drug Administration, Washington, D.C. on their website). These disclosures are incorporated herein by reference thereto.

Suitable basic salts include, but are not limited to, aluminum, ammonium, calcium,
15 lithium, magnesium, potassium, sodium and zinc salts.

All such acid salts and base salts are intended to be included within the scope of pharmaceutically acceptable salts as described herein. In addition, all such acid and base salts are considered equivalent to the free forms of the corresponding compounds for purposes of this description.

20 Compounds of Formula (I) and forms thereof, may further exist in a tautomeric form. All such tautomeric forms are contemplated and intended to be included within the scope of the compounds of Formula (I) or a form thereof as described herein.

The compounds of Formula (I) or a form thereof may contain asymmetric or chiral centers, and, therefore, exist in different stereoisomeric forms. The present description is
25 intended to include all stereoisomeric forms of the compounds of Formula (I) as well as mixtures thereof, including racemic mixtures.

The compounds described herein may include one or more chiral centers, and as such may exist as racemic mixtures (*R/S*) or as substantially pure enantiomers and diastereomers. The compounds may also exist as substantially pure (*R*) or (*S*) enantiomers (when one chiral center is
30 present). In one embodiment, the compounds described herein are (*S*) isomers and may exist as enantiomerically pure compositions substantially comprising only the (*S*) isomer. In another

embodiment, the compounds described herein are (*R*) isomers and may exist as enantiomerically pure compositions substantially comprising only the (*R*) isomer. As one of skill in the art will recognize, when more than one chiral center is present, the compounds described herein may also exist as a (*R,R*), (*R,S*), (*S,R*) or (*S,S*) isomer, as defined by *IUPAC* Nomenclature

5 Recommendations.

As used herein, the term “substantially pure” refers to compounds consisting substantially of a single isomer in an amount greater than or equal to 90%, in an amount greater than or equal to 92%, in an amount greater than or equal to 95%, in an amount greater than or equal to 98%, in an amount greater than or equal to 99%, or in an amount equal to 100% of the single isomer.

10 In one aspect of the description, a compound of Formula (I) or a form thereof is a substantially pure (*S*) enantiomer form present in an amount greater than or equal to 90%, in an amount greater than or equal to 92%, in an amount greater than or equal to 95%, in an amount greater than or equal to 98%, in an amount greater than or equal to 99%, or in an amount equal to 100%.

15 In one aspect of the description, a compound of Formula (I) or a form thereof is a substantially pure (*R*) enantiomer form present in an amount greater than or equal to 90%, in an amount greater than or equal to 92%, in an amount greater than or equal to 95%, in an amount greater than or equal to 98%, in an amount greater than or equal to 99%, or in an amount equal to 100%.

20 As used herein, a “racemate” is any mixture of isometric forms that are not “enantiomerically pure”, including mixtures such as, without limitation, in a ratio of about 50/50, about 60/40, about 70/30, or about 80/20.

In addition, the present description embraces all geometric and positional isomers. For example, if a compound of Formula (I) or a form thereof incorporates a double bond or a fused
25 ring, both the *cis*- and *trans*-forms, as well as mixtures, are embraced within the scope of the description. Diastereomeric mixtures can be separated into their individual diastereomers on the basis of their physical chemical differences by methods well known to those skilled in the art, such as, for example, by chromatography and/or fractional crystallization. Enantiomers can be separated by use of chiral HPLC column or other chromatographic methods known to those
30 skilled in the art. Enantiomers can also be separated by converting the enantiomeric mixture into a diastereomeric mixture by reaction with an appropriate optically active compound (*e.g.*, chiral

auxiliary such as a chiral alcohol or Mosher's acid chloride), separating the diastereomers and converting (*e.g.*, hydrolyzing) the individual diastereomers to the corresponding pure enantiomers. Also, some of the compounds of Formula (I) may be atropisomers (*e.g.*, substituted biaryls) and are considered as part of this description.

5 All stereoisomers (for example, geometric isomers, optical isomers and the like) of the present compounds (including those of the salts, solvates, esters and prodrugs of the compounds as well as the salts, solvates and esters of the prodrugs), such as those which may exist due to asymmetric carbons on various substituents, including enantiomeric forms (which may exist even in the absence of asymmetric carbons), rotameric forms, atropisomers, and diastereomeric
10 forms, are contemplated within the scope of this description, as are positional isomers (such as, for example, 4-pyridyl and 3-pyridyl). Individual stereoisomers of the compounds described herein may, for example, be substantially free of other isomers, or may be present in a racemic mixture, as described *supra*.

The use of the terms "salt", "solvate", "ester", "prodrug" and the like, is intended to
15 equally apply to the salt, solvate, ester and prodrug of enantiomers, stereoisomers, rotamers, tautomers, positional isomers, racemates or isotopologues of the instant compounds.

The term "isotopologue" refers to isotopically-enriched compounds described herein which are identical to those recited herein, but for the fact that one or more atoms are replaced by an atom having an atomic mass or mass number different from the atomic mass or mass number
20 usually found in nature. Examples of isotopes that can be incorporated into compounds described herein include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorus, fluorine and chlorine, such as ^2H , ^3H , ^{13}C , ^{14}C , ^{15}N , ^{18}O , ^{17}O , ^{31}P , ^{32}P , ^{35}S , ^{18}F , ^{35}Cl and ^{36}Cl , respectively, each of which are also within the scope of this description.

Certain isotopically-enriched compounds described herein (*e.g.*, those labeled with ^3H
25 and ^{14}C) are useful in compound and/or substrate tissue distribution assays. Tritiated (*i.e.*, ^3H) and carbon-14 (*i.e.*, ^{14}C) isotopes are particularly preferred for their ease of preparation and detectability. Further, substitution with heavier isotopes such as deuterium (*i.e.*, ^2H) may afford certain therapeutic advantages resulting from greater metabolic stability (*e.g.*, increased *in vivo* half-life or reduced dosage requirements) and hence may be preferred in some circumstances.

Polymorphic crystalline and amorphous forms of the compounds of Formula (I) and of the salts, solvates, hydrates, esters and prodrugs of the compounds of Formula (I) are further intended to be included in the present description.

Compound Uses

5 The present description relates to a method of use of a compound of Formula (I) or a form thereof for treating or ameliorating HD in a subject in need thereof, comprising administering an effective amount of the compound or a form thereof to the subject.

 The present description further relates to use of the compound of Formula (I) or a form thereof for treating or ameliorating HD in a subject in need thereof.

10 The present description further relates to use of the compound of Formula (I) or a form thereof having activity toward HD.

 The present description further relates to use of the compound of Formula (I) or a form thereof in a combination therapy to provide additive or synergistic activity, thus enabling the development of a combination product for treating or ameliorating HD.

15 In addition to monotherapeutic use, the instant compounds are useful in a combination therapy with current standard of agents, having additive or synergistic activity with one or more known agents.

 A combination therapy comprising compounds described herein in combination with one or more known drugs may be used to treat HD regardless of whether HD is responsive to the
20 known drug.

 Embodiments of the present description include the use of a compound of Formula (I) or a form thereof in a combination therapy for treating or ameliorating HD in a subject in need thereof, comprising administering an effective amount of the compound of Formula (I) or a form thereof and an effective amount of one or more agent(s).

25 Embodiments of the present description include the use of a compound of Formula (I) or a form thereof in a combination therapy for treating or ameliorating HD in a subject in need thereof, comprising administering an effective amount of the compound of Formula (I) or a form thereof and an effective amount of one or more agent(s).

 In an embodiment of a use or method provided herein, compounds of Formula (I) or a
30 form thereof used in combination with one or more additional agents can be administered to a subject or contacted with a subject or patient cell(s) prior to, concurrently with, or subsequent to

administering to the subject or patient or contacting the cell with an additional agent(s). A compound(s) of Formula (I) or a form thereof and an additional agent(s) can be administered to a subject or contacted with a cell in single composition or different compositions. In a specific embodiment, a compound(s) of Formula (I) or a form thereof is used in combination with gene therapy to inhibit HTT expression (using, *e.g.*, viral delivery vectors) or the administration of another small molecule HTT inhibitor. In another specific embodiment, a compound(s) of Formula (I) or a form thereof are used in combination with cell replacement using differentiated non-mutant HTT stem cells. In another specific embodiment, a compound(s) of Formula (I) or a form thereof are used in combination with cell replacement using differentiated HTT stem cells.

In one embodiment, provided herein is the use of compounds of Formula (I) or a form thereof in combination with supportive standard of care therapies, including palliative care.

An embodiment of the present description includes the use of a compound of Formula (I) or a form thereof in the preparation of a kit comprising the compound of Formula (I) or a form thereof and instructions for administering an effective amount of the compound of Formula (I) or a form thereof and an effective amount of one or more agent(s) in a combination therapy for treating or ameliorating HD in a subject in need thereof.

Accordingly, the present description relates to use of a compound of Formula (I) or a form thereof for treating or ameliorating HD. In accordance with the use of the present description, compounds that are useful in selectively treating or ameliorating HD, have been identified and use of these compounds for treating or ameliorating HD has been provided.

One embodiment of the use of the present description relates to use of a compound of Formula (I) or a form thereof for treating or ameliorating HD in a subject in need thereof, comprising administering an effective amount of the compound of Formula (I) or a form thereof to the subject.

One embodiment of the use of the present description relates to a method of use of a compound of Formula (I) or a form thereof for treating or ameliorating HD in a subject in need thereof, comprising administering an effective amount of the compound to the subject.

An embodiment of the use of the present description relates to a method of use of a compound of Formula (I) or a form thereof for treating or ameliorating HD in a subject in need thereof, comprising administering an effective amount of the compound to the subject.

An embodiment of the use of the present description relates to use of a compound of Formula (I) or a form thereof in the manufacture of a medicament for treating or ameliorating HD in a subject in need thereof, comprising administering an effective amount of the medicament to the subject.

5 An embodiment of the use of the present description relates to use of a compound of Formula (I) or a form thereof in the preparation of a kit comprising the compound of Formula (I) or a form thereof and instructions for administering the compound for treating or ameliorating HD in a subject in need thereof.

10 In one respect, for each of such embodiments, the subject is treatment naive. In another respect, for each of such embodiments, the subject is not treatment naive.

As used herein, the term “treating” refers to: (i) preventing a disease, disorder or condition from occurring in a subject that may be predisposed to the disease, disorder and/or condition but has not yet been diagnosed as having the disease, disorder and/or condition; (ii) inhibiting a disease, disorder or condition, *i.e.*, arresting the development thereof; and/or (iii) relieving a disease, disorder or condition, *i.e.*, causing regression of the disease, disorder and/or condition.

As used herein, the term “subject” refers to an animal or any living organism having sensation and the power of voluntary movement, and which requires oxygen and organic food. Nonlimiting examples include members of the human, primate, equine, porcine, bovine, murine, 20 rattus, canine and feline specie. In some embodiments, the subject is a mammal or a warm-blooded vertebrate animal. In other embodiments, the subject is a human. As used herein, the term “patient” may be used interchangeably with “subject” and “human”.

As used herein, the terms “effective amount” or “therapeutically effective amount” mean an amount of compound of Formula (I) or a form, composition or medicament thereof effective 25 in inhibiting the above-noted diseases and thus producing the desired therapeutic, ameliorative, inhibitory or preventative effect in a subject in need thereof.

The dose administered to achieve an effective target plasma concentration may also be administered based upon the weight of the subject or patient. Doses administered on a weight basis may be in the range of about 0.001 mg/kg/day to about 3500 mg/kg/day, or about 0.001 30 mg/kg/day to about 3000 mg/kg/day, or about 0.001 mg/kg/day to about 2500 mg/kg/day, or about 0.001 mg/kg/day to about 2000 mg/kg/day, or about 0.001 mg/kg/day to about 1500

mg/kg/day, or about 0.001 mg/kg/day to about 1000 mg/kg/day, or about 0.001 mg/kg/day to about 500 mg/kg/day, or about 0.001 mg/kg/day to about 250 mg/kg/day, or about 0.001 mg/kg/day to about 100 mg/kg/day, or about 0.001 mg/kg/day to about 75 mg/kg/day, or about 0.001 mg/kg/day to about 50 mg/kg/day, or about 0.001 mg/kg/day to about 40 mg/kg/day, or about 0.001 mg/kg/day to about 30 mg/kg/day, or about 0.001 mg/kg/day to about 20 mg/kg/day, or about 0.001 mg/kg/day to about 10 mg/kg/day, or about 0.01 mg/kg/day to about 2000 mg/kg/day, or about 0.01 mg/kg/day to about 1500 mg/kg/day, or about 0.01 mg/kg/day to about 1000 mg/kg/day, or about 0.01 mg/kg/day to about 600 mg/kg/day, or about 0.01 mg/kg/day to about 500 mg/kg/day, or about 0.01 mg/kg/day to about 300 mg/kg/day, or about 0.015 mg/kg/day to about 200 mg/kg/day, or about 0.02 mg/kg/day to about 100 mg/kg/day, or about 0.025 mg/kg/day to about 100 mg/kg/day, or about 0.03 mg/kg/day to about 100 mg/kg/day, wherein said amount is orally administered once (once in approximately a 24 hour period), twice (once in approximately a 12 hour period) or thrice (once in approximately an 8 hour period) daily according to subject weight.

In certain embodiments, the effective amount will be in a range of from about 0.001 mg/kg/day to about 500 mg/kg/day, or about 0.01 mg/kg/day to about 500 mg/kg/day, or about 0.1 mg to about 500 mg/kg/day, or about 1.0 mg/day to about 500 mg/kg/day, in single, divided, or a continuous dose for a patient or subject having a weight in a range of between about 40 to about 200 kg (which dose may be adjusted for patients or subjects above or below this range, particularly children under 40 kg). The typical adult subject is expected to have a median weight in a range of about 70 kg.

In another embodiment, where daily doses are adjusted based upon the weight of the subject or patient, compounds described herein may be formulated for delivery at about 0.02, 0.025, 0.03, 0.05, 0.06, 0.075, 0.08, 0.09, 0.10, 0.20, 0.25, 0.30, 0.50, 0.60, 0.75, 0.80, 0.90, 1.0, 1.10, 1.20, 1.25, 1.50, 1.75, 2.0, 3.0, 5.0, 10, 20, 30, 40, 50, 100, 150, 200, 250, 300, 400 or 500 mg/kg/day. Daily doses adjusted based upon the weight of the subject or patient may be administered as a single, divided, or continuous dose. In embodiments where a dose of compound is given more than once per day, the dose may be administered twice, thrice, or more times per day.

Within the scope of the present description, the “effective amount” of a compound of Formula (I) or a form thereof for use in the manufacture of a medicament, for use in the

preparation of a pharmaceutical kit or in a method of use for treating or ameliorating HD in a subject in need thereof is intended to include an amount in a range of from about 0.001 mg/kg/day to about 3500 mg/kg/day, or about 0.001 mg/kg/day to about 3000 mg/kg/day, or about 0.001 mg/kg/day to about 2500 mg/kg/day, or about 0.001 mg/kg/day to about 2000 mg/kg/day, or about 0.001 mg/kg/day to about 1500 mg/kg/day, or about 0.001 mg/kg/day to about 1000 mg/kg/day, or about 0.001 mg/kg/day to about 500 mg/kg/day, or about 0.001 mg/kg/day to about 250 mg/kg/day, or about 0.001 mg/kg/day to about 100 mg/kg/day, or about 0.001 mg/kg/day to about 75 mg/kg/day, or about 0.001 mg/kg/day to about 50 mg/kg/day, or about 0.001 mg/kg/day to about 40 mg/kg/day, or about 0.001 mg/kg/day to about 30 mg/kg/day, or about 0.001 mg/kg/day to about 20 mg/kg/day, or about 0.001 mg/kg/day to about 10 mg/kg/day, or about 0.01 mg/kg/day to about 2000 mg/kg/day, or about 0.01 mg/kg/day to about 1500 mg/kg/day, or about 0.01 mg/kg/day to about 1000 mg/kg/day, or about 0.01 mg/kg/day to about 600 mg/kg/day, or about 0.01 mg/kg/day to about 500 mg/kg/day, or about 0.01 mg/kg/day to about 300 mg/kg/day, or about 0.015 mg/kg/day to about 200 mg/kg/day, or about 0.02 mg/kg/day to about 100 mg/kg/day, or about 0.025 mg/kg/day to about 100 mg/kg/day, or about 0.03 mg/kg/day to about 100 mg/kg/day, wherein said amount is administered once (once in approximately a 24 hour period; i.e., "q.d."), twice (once in approximately a 12 hour period; i.e., "b.i.d." or "q.12h"), thrice (once in approximately an 8 hour period; i.e., "t.i.d." or "q.8h"), or four times (once in approximately a 6 hour period; i.e., "q.d.s.", "q.i.d." or "q.6h") daily according to subject weight.

Such amounts may further include an amount in a range of from about 0.001 mg to about 3500 mg administered daily; 0.001 mg to about 3000 mg administered daily; 0.001 mg to about 2500 mg administered daily; 0.001 mg to about 2000 mg administered daily; 0.001 mg to about 1500 mg administered daily; 0.001 mg to about 1000 mg administered daily; 0.001 mg to about 500 mg administered daily; 0.001 mg to about 250 mg administered daily; 1.0 mg to about 3500 mg administered daily; 1.0 mg to about 1500 mg administered daily; 1.0 mg to about 1000 mg administered daily; 10.0 mg to about 600 mg administered daily; 0.5 mg to about 2000 mg administered daily; or, an amount in a range of from about 5.0 mg to about 300 mg administered daily.

For example, the effective amount may be the amount required to treat HD in a subject or, more specifically, in a human. The effective amount for a subject will depend upon various

factors, including the subject's body weight, size and health. Effective amounts for a given patient can be determined by routine experimentation that is within the skill and judgment of the clinician.

For any compound, the effective amount can be estimated initially either in cell culture
5 assays or in relevant animal models, such as a mouse, chimpanzee, marmoset or tamarin animal model. Relevant animal models may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans. Therapeutic efficacy and toxicity may be determined by standard pharmaceutical procedures in cell cultures or experimental animals, *e.g.*, ED₅₀ (the
10 dose therapeutically effective in 50% of the population) and LD₅₀ (the dose lethal to 50% of the population). The dose ratio between therapeutic and toxic effects is therapeutic index, and can be expressed as the ratio, LD₅₀/ED₅₀. In some embodiments, the effective amount is such that a large therapeutic index is achieved. In further embodiments, the dosage is within a range of circulating concentrations that include an ED₅₀ with little or no toxicity. The dosage may vary
15 within this range depending upon the dosage form employed, sensitivity of the patient, and the route of administration.

More specifically, the concentration-biological effect relationships observed with regard to a compound of Formula (I) or a form thereof indicate a target plasma concentration ranging from approximately 0.001 µg/mL to approximately 50 µg/mL, from approximately 0.01 µg/mL
20 to approximately 20 µg/mL, from approximately 0.05 µg/mL to approximately 10 µg/mL, or from approximately 0.1 µg/mL to approximately 5 µg/mL. To achieve such plasma concentrations, the compounds described herein may be administered at doses that vary, such as, for example, without limitation, from 0.1 ng to 10,000 mg, depending upon the route of administration in single, divided, or continuous doses for a patient weighing between about 10 to
25 about 100 kg (which dose may be adjusted for patients within this weight range, particularly for children under 40 kg).

The exact dosage will be determined by the practitioner, in light of factors related to the subject. Dosage and administration may be adjusted to provide sufficient levels of the active agent(s) or to maintain the desired effect. Factors which may be taken into account include the
30 severity of the disease state, general health of the subject, ethnicity, age, weight, gender, diet, time of day and frequency of administration, drug combination(s), reaction sensitivities,

experience with other therapies, and tolerance/response to therapy. Long-acting pharmaceutical compositions may be administered every 2, 3 or 4 days, once every week, or once every two weeks depending on half-life and clearance rate of the particular formulation.

5 The compounds and compositions described herein may be administered to the subject via any drug delivery route known in the art. Nonlimiting examples include oral, ocular, rectal, buccal, topical, nasal, sublingual, transdermal, subcutaneous, intramuscular, intravenous (bolus and infusion), intracerebral, and pulmonary routes of administration.

10 In one aspect, provided herein are methods for modulating the amount of HTT (huntingtin protein), comprising contacting a human cell with a compound of Formula (I) or a form thereof. In a specific embodiment, provided herein are methods for modulating the amount of HTT, comprising contacting a human cell with a compound of Formula (I) or a form thereof that modulates the expression of HTT. The human cell can be contacted with a compound of Formula (I) or a form thereof *in vitro*, or *in vivo*, *e.g.*, in a non-human animal or in a human. In a specific embodiment, the human cell is from or in a human. In another specific embodiment, the human cell is from or in a human with HD. In another specific embodiment, the human cell is from or in a human with HD, caused by a CAG repeat in the *Htt* gene, resulting in a loss of HTT expression and/or function. In another embodiment, the human cell is from a human with HD. In another embodiment, the human cell is in a human with HD. In one embodiment, the compound is a form of the compound of Formula (I).

20 In a specific embodiment, provided herein is a method for enhancing the inhibition of mutant HTT transcribed from the *Htt* gene, comprising contacting a human cell with a compound of Formula (I) or a form thereof. The human cell can be contacted with a compound of Formula (I) or a form thereof *in vitro*, or *in vivo*, *e.g.*, in a non-human animal or in a human. In a specific embodiment, the human cell is from or in a human. In another specific embodiment, the human cell is from or in a human with HD. In another specific embodiment, the human cell is from or in a human with HD, caused by a CAG repeat in the *Htt* gene, resulting in a loss of wild-type “normal” HTT expression and/or function. In another embodiment, the human cell is from a human with HD. In another embodiment, the human cell is in a human with HD. In one embodiment, the compound is a form of the compound of Formula (I).

30 In another aspect, provided herein is a method for modulating the inhibition of mutant HTT transcribed from the *Htt* gene, comprising administering to a non-human animal model for

HD a compound of Formula (I) or a form thereof. In a specific embodiment, provided herein is a method for modulating the inhibition of mutant HTT transcribed from the *Htt* gene, comprising administering to a non-human animal model for HD a compound of Formula (I) or a form thereof. In a specific embodiment, the compound is a form of the compound of Formula (I).

5 In another aspect, provided herein is a method for decreasing the amount of mutant HTT, comprising contacting a human cell with a compound of Formula (I) or a form thereof. In a specific embodiment, provided herein is a method for decreasing the amount of mutant HTT, comprising contacting a human cell with a compound of Formula (I) that inhibits the transcription of mutant *HTT* (huntingtin mRNA) from the *Htt* gene. In another specific
10 embodiment, provided herein is a method for decreasing the amount of HTT, comprising contacting a human cell with a compound of Formula (I) that inhibits the expression of mutant HTT transcribed from the *Htt* gene. The human cell can be contacted with a compound of Formula (I) or a form thereof *in vitro*, or *in vivo*, e.g., in a non-human animal or in a human. In a specific embodiment, the human cell is from or in a human. In another specific embodiment, the
15 human cell is from or in a human with HD. In another specific embodiment, the human cell is from or in a human with HD, caused by a CAG repeat in the *Htt* gene, resulting in a loss of HTT expression and/or function. In another embodiment, the human cell is from a human with HD. In another embodiment, the human cell is in a human with HD. In one embodiment, the compound is a form of the compound of Formula (I).

20 In certain embodiments, treating or ameliorating HD with a compound of Formula (I) or a form thereof (alone or in combination with an additional agent) has a therapeutic effect and/or beneficial effect. In a specific embodiment, treating HD with a compound of Formula (I) or a form thereof (alone or in combination with an additional agent) results in one, two or more of the following effects: (i) reduces or ameliorates the severity of HD; (ii) delays onset of HD; (iii)
25 inhibits the progression of HD; (iv) reduces hospitalization of a subject; (v) reduces hospitalization length for a subject; (vi) increases the survival of a subject; (vii) improves the quality of life for a subject; (viii) reduces the number of symptoms associated with HD; (ix) reduces or ameliorates the severity of a symptom(s) associated with HD; (x) reduces the duration of a symptom associated with HD; (xi) prevents the recurrence of a symptom associated with
30 HD; (xii) inhibits the development or onset of a symptom of HD; and/or (xiii) inhibits of the progression of a symptom associated with HD.

Metabolites of the Compounds

Also included within the scope of the present description are the use of *in vivo* metabolic products of the compounds described herein. Such products may result, for example, from the oxidation, reduction, hydrolysis, amidation, esterification and the like of the administered compound, primarily due to enzymatic processes. Accordingly, the description includes the use of compounds produced by a process comprising contacting a compound described herein with a mammalian tissue or a mammal for a period of time sufficient to yield a metabolic product thereof.

Such products typically are identified by preparing a radio-labeled isotopologue (*e.g.*, ^{14}C or ^3H) of a compound described herein, administering the radio-labeled compound in a detectable dose (*e.g.*, greater than about 0.5 mg/kg) to a mammal such as a rat, mouse, guinea pig, dog, monkey or human, allowing sufficient time for metabolism to occur (typically about 30 seconds to about 30 hours), and identifying the metabolic conversion products from urine, bile, blood or other biological samples. The conversion products are easily isolated since they are “radiolabeled” by virtue of being isotopically-enriched (others are isolated by the use of antibodies capable of binding epitopes surviving in the metabolite). The metabolite structures are determined in conventional fashion, *e.g.*, by MS or NMR analysis. In general, analysis of metabolites may be done in the same way as conventional drug metabolism studies well-known to those skilled in the art. The conversion products, so long as they are not otherwise found *in vivo*, are useful in diagnostic assays for therapeutic dosing of the compounds described herein even if they possess no biological activity of their own.

Pharmaceutical Compositions

Embodiments of the present description include the use of a compound of Formula (I) or a form thereof in a pharmaceutical composition for treating or ameliorating HD in a subject in need thereof, comprising administering an effective amount of the compound of Formula (I) or a form thereof in admixture with one or more pharmaceutically acceptable excipient(s).

An embodiment of the present description includes the use of a pharmaceutical composition of the compound of Formula (I) or a form thereof in the preparation of a kit comprising the pharmaceutical composition of the compound of Formula (I) or a form thereof

and instructions for administering the compound for treating or ameliorating HD in a subject in need thereof.

As used herein, the term “composition” means a product comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from combination of the specified ingredients in the specified amounts.

The pharmaceutical composition may be formulated to achieve a physiologically compatible pH, ranging from about pH 3 to about pH 11. In some embodiments, the pharmaceutical composition is formulated to achieve a pH of from about pH 3 to about pH 7. In other embodiments, the pharmaceutical composition is formulated to achieve a pH of from about pH 5 to about pH 8.

The term “pharmaceutically acceptable excipient” refers to an excipient for administration of a pharmaceutical agent, such as the compounds described herein. The term refers to any pharmaceutical excipient that may be administered without undue toxicity. Pharmaceutically acceptable excipients may be determined in part by the particular composition being administered, as well as by the particular mode of administration and/or dosage form. Nonlimiting examples of pharmaceutically acceptable excipients include carriers, solvents, stabilizers, adjuvants, diluents, *etc.* Accordingly, there exists a wide variety of suitable formulations of pharmaceutical compositions for the instant compounds described herein (*see, e.g.,* Remington’s Pharmaceutical Sciences).

Suitable excipients may be carrier molecules that include large, slowly metabolized macromolecules such as proteins, polysaccharides, polylactic acids, polyglycolic acids, polymeric amino acids, amino acid copolymers, and inactive antibodies. Other exemplary excipients include antioxidants such as ascorbic acid; chelating agents such as EDTA; carbohydrates such as dextrin, hydroxyalkylcellulose, hydroxyalkylmethylcellulose (*e.g.,* hydroxypropylmethylcellulose, also known as HPMC), stearic acid; liquids such as oils, water, saline, glycerol and ethanol; wetting or emulsifying agents; pH buffering substances; and the like. Liposomes are also included within the definition of pharmaceutically acceptable excipients.

The pharmaceutical compositions described herein may be formulated in any form suitable for the intended use described herein. Suitable formulations for oral administration include solids, liquid solutions, emulsions and suspensions, while suitable inhalable formulations

for pulmonary administration include liquids and powders. Alternative formulations include syrups, creams, ointments, tablets, and lyophilized solids which can be reconstituted with a physiologically compatible solvent prior to administration.

When intended for oral use for example, tablets, troches, lozenges, aqueous or oil
5 suspensions, non-aqueous solutions, dispersible powders or granules (including micronized particles or nanoparticles), emulsions, hard or soft capsules, syrups or elixirs may be prepared. Compositions intended for oral use may be prepared according to any method known to the art for the manufacture of pharmaceutical compositions, and such compositions may contain one or more agents including sweetening agents, flavoring agents, coloring agents, and preserving
10 agents, in order to provide a palatable preparation.

Pharmaceutically acceptable excipients suitable for use in conjunction with tablets include, for example, inert diluents, such as celluloses, calcium or sodium carbonate, lactose, calcium or sodium phosphate; disintegrating agents, such as croscarmellose sodium, cross-linked povidone, maize starch, or alginic acid; binding agents, such as povidone, starch, gelatin or
15 acacia; and lubricating agents, such as magnesium stearate, stearic acid, or talc. Tablets may be uncoated or may be coated by known techniques including microencapsulation to delay disintegration and adsorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate alone or with a wax may be employed.

Formulations for oral use may be also presented as hard gelatin capsules where the active
20 ingredient is mixed with an inert solid diluent, for example celluloses, lactose, calcium phosphate, or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with non-aqueous or oil medium, such as glycerin, propylene glycol, polyethylene glycol, peanut oil, liquid paraffin, or olive oil.

In other embodiments, pharmaceutical compositions described herein may be formulated as suspensions comprising a compound of Formula (I) or a form thereof in admixture with one or more pharmaceutically acceptable excipient(s) suitable for the manufacture of a suspension. In yet other embodiments, pharmaceutical compositions described herein may be formulated as dispersible powders and granules suitable for preparation of a suspension by the addition of one
30 or more excipient(s).

Excipients suitable for use in connection with suspensions include suspending agents, such as sodium carboxymethylcellulose, methylcellulose, hydroxypropyl methylcellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth, gum acacia, dispersing or wetting agents such as a naturally occurring phosphatide (*e.g.*, lecithin), a condensation product of an alkylene oxide with a fatty acid (*e.g.*, polyoxyethylene stearate), a condensation product of ethylene oxide with a long chain aliphatic alcohol (*e.g.*, heptadecaethyleneoxycethanol), a condensation product of ethylene oxide with a partial ester derived from a fatty acid and a hexitol anhydride (*e.g.*, polyoxyethylene sorbitan monooleate); and thickening agents, such as carbomer, beeswax, hard paraffin, or cetyl alcohol. The suspensions may also contain one or more preservatives such as acetic acid, methyl and/or n-propyl p-hydroxy-benzoate; one or more coloring agents; one or more flavoring agents; and one or more sweetening agents such as sucrose or saccharin.

The pharmaceutical compositions described herein may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, such as olive oil or arachis oil, a mineral oil, such as liquid paraffin, or a mixture of these. Suitable emulsifying agents include naturally-occurring gums, such as gum acacia and gum tragacanth; naturally occurring phosphatides, such as soybean lecithin, esters or partial esters derived from fatty acids; hexitol anhydrides, such as sorbitan monooleate; and condensation products of these partial esters with ethylene oxide, such as polyoxyethylene sorbitan monooleate. The emulsion may also contain sweetening and flavoring agents. Syrups and elixirs may be formulated with sweetening agents, such as glycerol, sorbitol or sucrose. Such formulations may also contain a demulcent, a preservative, a flavoring or a coloring agent.

Additionally, the pharmaceutical compositions described herein may be in the form of a sterile injectable preparation, such as a sterile injectable aqueous emulsion or oleaginous suspension. Such emulsion or suspension may be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents which have been mentioned above. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally acceptable diluent or solvent, such as a solution in 1,2-propanediol. The sterile injectable preparation may also be prepared as a lyophilized powder. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile fixed oils may be employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic

mono- or di-glycerides. In addition, fatty acids such as oleic acid may likewise be used in the preparation of injectables.

The compounds described herein may be substantially insoluble in water and sparingly soluble in most pharmaceutically acceptable protic solvents and vegetable oils, but generally
5 soluble in medium-chain fatty acids (*e.g.*, caprylic and capric acids) or triglycerides and in propylene glycol esters of medium-chain fatty acids. Thus, contemplated in the description are compounds which have been modified by substitutions or additions of chemical or biochemical moieties which make them more suitable for delivery (*e.g.*, increase solubility, bioactivity, palatability, decrease adverse reactions, *etc.*), for example by esterification, glycosylation,
10 PEGylation, *etc.*

In some embodiments, the compound described herein is formulated for oral administration in a lipid-based composition suitable for low solubility compounds. Lipid-based formulations can generally enhance the oral bioavailability of such compounds. As such, pharmaceutical compositions described herein may comprise an effective amount of a compound
15 of Formula (I) or a form thereof, together with at least one pharmaceutically acceptable excipient selected from medium chain fatty acids or propylene glycol esters thereof (*e.g.*, propylene glycol esters of edible fatty acids such as caprylic and capric fatty acids) and pharmaceutically acceptable surfactants, such as polysorbate 20 or 80 (also referred to as Tween® 20 or Tween® 80, respectively) or polyoxyl 40 hydrogenated castor oil.

20 In other embodiments, the bioavailability of low solubility compounds may be enhanced using particle size optimization techniques including the preparation of nanoparticles or nanosuspensions using techniques known to those skilled in the art. The compound forms present in such preparations include amorphous, partially amorphous, partially crystalline or crystalline forms.

25 In alternative embodiments, the pharmaceutical composition may further comprise one or more aqueous solubility enhancer(s), such as a cyclodextrin. Nonlimiting examples of cyclodextrin include hydroxypropyl, hydroxyethyl, glucosyl, maltosyl and maltotriosyl derivatives of α -, β -, and γ -cyclodextrin, and hydroxypropyl- β -cyclodextrin (HPBC). In some embodiments, the pharmaceutical composition further comprises HPBC in a range of from about
30 0.1% to about 20%, from about 1% to about 15%, or from about 2.5% to about 10%. The

amount of solubility enhancer employed may depend on the amount of the compound in the composition.

Preparation of Compounds

Compounds provided herein can be prepared by those skilled in the art, such as, by the synthetic methods set forth in International Application Publication Number WO2014/028459 A1, published February 20, 2014, International Application Publication Number WO2014/116845 A1, published July 31, 2014, and International Application Publication Number WO2015/017589 A1, published February 5, 2015, each of which are herein incorporated by reference.


General Schemes

The following reaction schemes illustrate methods to make compounds described herein. It is understood that one skilled in the art would be able to make these compounds by similar methods or by methods known to one skilled in the art.

In general, starting components and reagents may be commercially obtained from various sources such as commercial vendors, synthesized according to methodology known to those skilled in the art, or prepared as described herein. It is understood that in the following description, combinations of substituents and/or variables of the depicted formulae are permissible only if such contributions result in stable compounds.

In general, compounds of Formula (I) described herein can be synthesized following the general procedure for Scheme 1.

Scheme 1



General Procedure for Scheme 1

The starting materials for the above reaction scheme are commercially available or can be prepared according to methods known to one skilled in the art or by methods disclosed herein.

In general, the compounds described herein are prepared in the above reaction Scheme 1 as follows: a halogenated Compound 1a (wherein Hal is a halogen selected from bromine or

chlorine) is reacted with a Compound 1b (wherein H is a reactive hydrogen atom in an amine or alcohol X functional group, wherein X and B for a compound of Formula (I) are as described herein) using a displacement or a metal-mediated cross coupling reaction, such as a Buchwald reaction, to provide an intermediate Compound 1c. Compound 1c is carried forward and reacted
5 with a boronate acid or boronate ester Compound 1d (where B represents a boron atom, R represents a an acid or ester functional group or, when taken together with boron, form a ring system and A for a compound of Formula (I) is as described herein) using a transition metal-mediated cross coupling reaction, such as a Suzuki reaction, to provide a Compound of Formula (I).

10 In a complementary manner, compounds of Formula (I) may also be synthesized in reverse order wherein the boronate Compound 1d is reacted first with Compound 1a followed by reaction of the intermediate with amine Compound 1b. Other variations representing different combinations for substituting either or both boronate Compound 1d and amine Compound 1b on Compound 1a are intended to be included within the scope of the synthetic methodologies described herein.

15 Specific Examples

To assist in understanding the present description, the following specific examples are included. The experiments relating to this description should not, of course, be construed as specifically limiting the description and such variations of the description, now known or later developed, which would be within the purview of one skilled in the art are considered to fall
20 within the scope of the description as described herein and hereinafter claimed.

Other than in the working examples, unless indicated to the contrary, all numbers expressing quantities of materials, reagents, reaction conditions, experimental data, and so forth used in the specification and claims are to be understood as being modified by the term “about”. Accordingly, all such numbers represent approximations that may vary depending upon the
25 desired properties sought to be obtained by a reaction or as a result of variable experimental conditions. Therefore, within an expected range of experimental reproducibility, the term “about” in the context of the resulting data, refers to a range for data provided that may vary according to a standard deviation from the mean. As well, for experimental results provided, the resulting data may be rounded up or down to present data consistently, without loss of significant
30 figures. At the very least, and not as an attempt to limit the application of the doctrine of

equivalents to the scope of the claims, each numerical parameter should be construed in light of the number of significant digits and ordinary rounding techniques.

While the numerical ranges and parameters setting forth the broad scope of the description are approximations, the numerical values set forth in the working examples are reported as precisely as possible. Any numerical value, however, inherently contains certain errors necessarily resulting from the standard deviation found in their respective testing measurements.

The column chromatography system used for product purification was an ISCO CombiFlash™ brand chromatography system (manufactured by Teledyne Isco).

10 Synthetic Examples

Greater details of the present description are provided with reference to the following non-limiting examples, which are offered to more fully illustrate the description, but are not to be construed as limiting the scope thereof. The examples illustrate the preparation of certain compounds described herein, and the testing of these compounds *in vitro* and/or *in vivo*. Those of skill in the art will understand that the techniques described in these examples represent techniques described by the inventors to function well in the practice of the description, and as such constitute preferred modes for the practice thereof. However, those of skill in the art should appreciate in light of the present disclosure that many changes can be made to the specific methods that are disclosed and still obtain a like or similar result without departing from the spirit and scope of the description.

As used above, and throughout this description, the following abbreviations, unless otherwise indicated, shall be understood to have the following meanings:

Abbreviation	Meaning
AcOH or HOAc	acetic acid
ACN or MeCN	acetonitrile
Bn	benzyl
BnBr	benzyl bromide
BnO or OBn	benzyloxy
Boc	<i>tert</i> -butoxycarbonyl
Boc ₂ O or (Boc) ₂ O	di- <i>tert</i> -butyl dicarbonate
<i>t</i> BuXPhos-Pd-G3	[(2-di- <i>tert</i> -butylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)-2-(2'-amino-1,1'-biphenyl)] palladium(II) methanesulfonate
Cbz	benzyloxycarbonyl
DCE	dichloroethane
DCM	dichloromethane (CH ₂ Cl ₂)
DIBAL-H	diisobutylaluminium hydride

Abbreviation	Meaning
DIPEA	N,N-diisopropylethylamine
DMF	dimethyl formamide
DMA	dimethylacetamide
DMAP	4-dimethylaminopyridine
DMB	2,4-dimethoxybenzyl
DMSO	dimethylsulfoxide
EA or EtOAc	ethyl acetate
EtOH	ethanol
Et ₂ O	diethyl ether
HCl	hydrochloric acid
HPLC	high performance liquid chromatography
h/hr/hrs/min/s	hour(s)(h, hr or hrs)/minute(s)(min/mins)/second(s)
K ₂ CO ₃	potassium carbonate
KOAc	potassium acetate
K ₃ PO ₄	potassium phosphate
LAH	lithium aluminium hydride
LC/MS, LCMS or LC-MS	liquid chromatographic mass spectroscopy
LDA	lithium diisopropylamide
LiHMDS	lithium bis(trimethylsilyl)amide
Me ₄ Bu-XPhos	di-tert-butyl(2',4',6'-triisopropyl-3,4,5,6-tetramethyl-[1,1'-biphenyl]-2-yl)phosphine
MeI	methyl iodide
MeOH	methanol
Me ₂ NH or NHMe ₂	dimethyl amine
MS	mass spectroscopy
N ₂	nitrogen
Na ₂ SO ₄	sodium sulfate
NaBH(OAc) ₃	sodium triacetoxymethylborohydride
NaHMDS	sodium bis(trimethylsilyl)amide
NBS	N-bromosuccinimide
NH ₄ OH	ammonium hydroxide
NMO	N-methylmorpholine-N-oxide
NMP	N-methyl-2-pyrrolidone
n-BuLi	n-butyl lithium
NMR	nuclear magnetic resonance
Pd/C	palladium on carbon
Pd ₂ (dba) ₃	tris(dibenzylideneacetone)dipalladium(0)
Pd(dppf)Cl ₂	[1,1'-bis(diphenylphosphino)ferrocene] dichloropalladium(II)
Pd(PPh ₃) ₄	tetrakis(triphenylphosphine)palladium
Pin	pinacol
psi	pounds per square inch pressure
S-Phos	2-dicyclohexylphosphino-2',6'-dimethoxybiphenyl
TBDPSCI	tert-butyldiphenylchlorosilane
TBS	tert-butyldimethylsilyl
TBSCl	tert-butyldimethylsilyl chloride

Abbreviation	Meaning
<i>t</i> -BuOK	potassium tert-butoxide
TEA or NEt ₃	triethylamine
TFA	trifluoroacetic acid
THF	tetrahydrofuran
THP	tetrahydro-2H-pyran-2-yl
THPO or OTHP	tetrahydro-2H-pyran-2-yl-oxy
TIPS-H or TIPSH	triisopropyl silane
TLC	thin layer chromatography
TMP	2,2,6,6-tetramethylpiperidinyl
TMPMgCl-LiCl	2,2,6,6-tetramethylpiperidinylmagnesium chloride lithium chloride
TMSI	trimethylsilyl iodide
TMSOK	potassium trimethylsilanolate

Example 1 (Compound 411)

2-{6-[8-Azabicyclo[3.2.1]oct-3-yl(methyl)amino]pyridazin-3-yl}-5-(1H-pyrazol-4-yl)phenol hydrochloride

- Step 1:** 3,6-Dibromopyridazine (133.7 mg, 0.56 mmol), tert-butyl (1*R*,5*S*)-3-(methylamino)-8-azabicyclo[3.2.1]octane-8-carboxylate (85 mg, 0.28 mmol) and DIPEA (0.15 mL, 0.84 mmol) were mixed in 1 mL of ACN and heated to 100 °C for 1 h until LC-MS showed complete consumption of the starting material. The reaction mixture was concentrated and purified via column chromatography: eluting with gradient CH₂Cl₂/MeOH (0% to 30% MeOH), column: silica 4g to afford tert-butyl (1*S*,5*R*)-3-[(6-bromopyridazin-3-yl)-methyl-amino]-8-azabicyclo[3.2.1]octane-8-carboxylate (55 mg, 49.25% yield).
- Step 2:** tert-Butyl 3-[(6-bromopyridazin-3-yl)-methyl-amino]-8-azabicyclo[3.2.1]octane-8-carboxylate (55 mg, 0.14 mmol), 4-[3-(methoxymethoxy)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]-1-tetrahydropyran-2-yl-pyrazole (68.82 mg, 0.17 mmol), Pd(dppf)Cl₂ (10.34 mg, 0.01 mmol), K₂CO₃ (57.98 mg, 0.42 mmol) were mixed in a Schlenk tube. The reaction was degassed with N₂ for 15 min and dioxane (2 mL) and water (0.5 mL) were added and the reaction was heated to 90 °C for 16 h. The reaction was cooled to room temperature, partitioned between EtOAc and water. The organic layers were dried over Na₂SO₄, concentrated under vacuum, purified via column chromatography: eluting with gradient CH₂Cl₂/EtOAc (0% to 80%), column: silica 4g, to provide tert-butyl (1*S*,5*R*)-3-[[6-[2-(methoxymethoxy)-4-(1-tetrahydropyran-2-yl)pyrazol-4-yl]phenyl]pyridazin-3-yl]-methyl-amino]-8-azabicyclo[3.2.1]octane-8-carboxylate (52 mg, 62%) as a tan solid.

Step 3: To a solution of tert-butyl (1*S*,5*R*)-3-[[6-[2-(methoxymethoxy)-4-(1-tetrahydropyran-2-

ylpyrazol-4-yl)phenyl]pyridazin-3-yl]-methyl-amino]-8-azabicyclo[3.2.1]octane-8-carboxylate (52 mg, 0.086 mmol) in a 1 mL mixture of CH₂Cl₂ and 0.5 mL of MeOH was added HCl (4 mol/L) in 1,4-dioxane (0.06 mL). The reaction was stirred overnight. The precipitate was filtered, then dried under vacuum to provide 2-[6-[8-azabicyclo[3.2.1]octan-3-

- 5 yl(methyl)amino]pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol; hydrochloride (15 mg, 42.25% yield) as a yellow solid.

LC-MS: 377 [M+H]⁺. ¹H NMR (500 MHz, DMSO-*d*₆) δ: 9.37-9.51 (m, 1H), 8.87-8.96 (m, 1H), 8.31 (d, *J*=9.8 Hz, 1H), 8.12 (s, 2H), 7.74 (d, *J*=8.2 Hz, 2H), 7.19-7.27 (m, 2H), 4.80-4.92 (m, 1H), 4.08-4.14 (m, 3H), 3.07 (s, 3H), 2.30 (td, *J*=12.6, 2.8 Hz, 2H), 1.99-2.15 (m, 4H), 1.74-1.87
10 (m, 2H)

Using the procedure described for Example 1 above, additional compounds described herein may be prepared by substituting the appropriate starting materials reagents, and reaction conditions, obtaining compounds such as those selected from:

Cpd	Data
412	LC-MS: 363 [M+H] ⁺ . ¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ: 9.24-9.34 (m, 1H), 9.09-9.21 (m, 1H), 8.30 (d, <i>J</i> =9.8 Hz, 1H), 8.09 (s, 2H), 7.64-7.67 (m, 1H), 7.63 (d, <i>J</i> =8.2 Hz, 1H), 7.25-7.27 (m, 1H), 7.24 (s, 1H), 5.42 (br s, 2H), 4.23-4.31 (m, 1H), 4.00-4.13 (m, 2H), 2.17 (s, 4H), 1.97-2.02 (m, 2H), 1.94 (td, <i>J</i> =12.6, 2.5 Hz, 2H)
413	LC-MS: 393 [M+H] ⁺ . ¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ: 9.34-9.49 (m, 1H), 8.43-8.56 (m, 1H), 8.30 (d, <i>J</i> =9.8 Hz, 1H), 8.10 (s, 2H), 7.63 (s, 2H), 7.19-7.30 (m, 2H), 5.77 (br s, 2H), 4.41-4.62 (m, 1H), 2.08 (dd, <i>J</i> =11.7, 2.8 Hz, 2H), 1.72 (td, <i>J</i> =12.1, 2.8 Hz, 2H), 1.53 (s, 6H), 1.49 (s, 6H)
415	LC-MS: 364 [M+H] ⁺ . ¹ H NMR (500 MHz, methanol- <i>d</i> ₄) δ: 8.61 (d, <i>J</i> =9.5 Hz, 1H), 8.24 (s, 2H), 7.85 (d, <i>J</i> =8.2 Hz, 1H), 7.72 (d, <i>J</i> =9.5 Hz, 1H), 7.38 (dd, <i>J</i> =8.2, 1.6 Hz, 1H), 7.32 (d, <i>J</i> =1.6 Hz, 1H), 5.65 (spt, <i>J</i> =5.7 Hz, 1H), 4.25 (dd, <i>J</i> =3.6, 2.8 Hz, 2H), 4.17-4.32 (m, 2H), 2.24 (s, 4H), 2.11 (tt, <i>J</i> =11.2, 2.8 Hz, 2H) (3 hydrogens corresponding to OH and NH unobserved)
433	LC-MS: 363 [M+H] ⁺ . ¹ H NMR (500 MHz, methanol- <i>d</i> ₄) δ: 8.42 (d, <i>J</i> =9.5 Hz, 1H), 8.34 (s, 2H), 7.80 (d, <i>J</i> =9.8 Hz, 1H), 7.74 (d, <i>J</i> =8.5 Hz, 1H), 7.40 (dd, <i>J</i> =8.2, 2.5 Hz, 1H), 7.32 (d, <i>J</i> =1.6 Hz, 1H), 4.18 (td, <i>J</i> =6.9, 0.9 Hz, 1H), 4.11-4.16 (m, 2H), 2.42-2.52 (m, 4H), 2.32-2.35 (m, 1H), 2.28-2.32 (m, 1H), 2.15-2.25 (m, 2H), (4 hydrogens corresponding to OH and NH unobserved)

Example 2 (Compound 416)

5-(5-Methyl-1H-pyrazol-4-yl)-2-{6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl}phenol

- Step 1: 3-Methoxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl trifluoromethanesulfonate (80 mg, 0.16 mmol), (tert-butyl 3-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole-1-carboxylate (59 mg, 0.19 mmol), Pd(dppf)Cl₂ (7 mg, 0.01 mmol), K₂CO₃ (66 mg, 0.48 mmol) were mixed in a Schlenk tube. The reaction was degassed with N₂ for 15 min and dioxane (2 mL) and water (0.5 mL) were added and the reaction was heated to 90 °C for 16 h. The reaction was cooled to room temperature, partitioned between EtOAc and water. The organic layers were dried over Na₂SO₄, concentrated under vacuum, then purified via column chromatography: eluting with gradient CH₂Cl₂/MeOH (0% to 30% MeOH), column: silica 4g to provide tert-butyl 4-(3-methoxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)-3-methyl-1H-pyrazole-1-carboxylate (46 mg, 54%) as light brown solid.
- Step 2: A solution of tert-butyl 4-(3-methoxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)-3-methyl-1H-pyrazole-1-carboxylate (46 mg, 0.09mmol) in 2 mL of dry CH₂Cl₂ was cooled in ice-water bath. Boron tribromide 1.0 M in CH₂Cl₂ (0.45 mL, 0.45 mmol) was added and the reaction mixture was stirred at room temperature for 16 hours. The reaction was quenched with 2 mL of MeOH, stirred for 30 min, then concentrated and purified via column chromatography: eluting with gradient CH₂Cl₂/MeOH (2.5% NH₄OH) (0% to 30% MeOH/NH₄OH), column: silica 4g, to provide tert-butyl 2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(3-methyl-1H-pyrazol-4-yl)phenol (20 mg, 55%) as a yellow solid.

LC-MS: 421 [M+H]⁺. ¹H NMR (500 MHz, DMSO-*d*₆) δ: 13.74 (s, 1H), 12.63-12.71 (m, 1H), 8.20 (d, *J*=9.5Hz, 1H), 8.03-8.04 (m, 1H), 7.86 (d, *J*=8.0Hz, 1H), 7.73-7.75 (m, 1H), 7.36 (d, *J*=9.5Hz, 1H), 7.03-7.06 (m, 2H), 4.90-5.05 (m, 1H), 2.92 (s, 3H), 2.43 (s, 3H), 1.43-1.54 (m, 2H), 1.15-1.43 (m, 2H), 1.14 (s, 6H), 1.09 (s, 6H)

Using the procedure described for Example 2 above, additional compounds described herein may be prepared by substituting the appropriate starting materials reagents, and reaction conditions, obtaining compounds such as those selected from:

Cpd	Data
414	LC-MS: 408 [M+H] ⁺ . ¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ: 13.84 (s, 1H), 8.10-8.29 (m, 2H), 7.92 (s, 1H), 7.84 (d, <i>J</i> =8.2 Hz, 1H), 7.36 (d, <i>J</i> =9.8 Hz, 1H), 7.08-7.19 (m, 2H), 4.87-5.03 (m, 1H), 3.87 (s, 3H), 2.96 (s, 3H), 1.48-1.59 (m, 2H), 1.38-1.48 (m, 2H), 1.26 (s, 6H), 1.09 (s, 6H)
418	LC-MS: 422 [M+H] ⁺ . ¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ: 12.89 (s, 1H), 12.29-12.48 (m, 1H), 8.41 (dd, <i>J</i> =8.8, 0.9 Hz, 1H), 7.95 (dd, <i>J</i> =7.9, 0.9 Hz, 1H), 7.37 (dd, <i>J</i> =9.1, 1.3 Hz, 1H), 6.85-6.94 (m, 2H), 5.61-5.70 (m, 1H), 2.25 (br s., 6H), 2.09 (dd, <i>J</i> =11.8, 3.9 Hz, 2H), 1.25-1.33 (m, 2H), 1.24 (s, 6H), 1.11 (s, 6H), (1 hydrogen corresponding to OH or NH unobserved)
451	LC-MS: 408 [M+H] ⁺ . ¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ: 13.00 (s, 1H), 12.59-12.80 (m, 1H), 8.42 (d, <i>J</i> =9.5 Hz, 1H), 7.94 (d, <i>J</i> =8.8 Hz, 1H), 7.38 (d, <i>J</i> =9.5 Hz, 1H), 7.03-7.18 (m, 2H), 5.55-5.78 (m, 1H), 2.39 (br s., 3H), 2.03-2.13 (m, 2H), 1.26-1.41 (m, 2H), 1.24 (s, 6H), 1.11 (s, 6H), (1 hydrogen corresponding to OH or NH unobserved)

Example 3 (Compound 419)

- 5 2-{6-[Methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl}-5-(4-nitro-1H-pyrazol-1-yl)phenol dihydrochloride

Step 1: 3-Methoxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl trifluoromethanesulfonate (335 mg, 0.67 mmol), 4-nitro-1H-pyrazole (340 mg, 4.5 mmol), Pd₂(dba)₃ (67 mg, 0.07 mmol), Me₄tBu-XPhos (35 mg, 0.07 mmol), K₃PO₄ (368 mg, 1.73 mmol) were mixed in a Schlenk tube. The reaction was degassed with N₂ for 15 min and dioxane (8 mL) was added and the reaction was heated to 90 °C for 16 h. The reaction was cooled to room temperature, filtered through celite, concentrated, purified via column chromatography: eluting with gradient CH₂Cl₂/MeOH (0% to 30% MeOH), column: silica 12g, to provide 6-(2-methoxy-4-(4-nitro-1H-pyrazol-1-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine (202 mg, 65%).

LC-MS: 466 [M+H]⁺. ¹H NMR (500 MHz, DMSO-*d*₆) δ: 9.81-9.85 (m, 1H), 8.61 (d, *J*=0.6 Hz, 1H), 7.91 (d, *J*=8.5 Hz, 1H), 7.81 (d, *J*=9.8 Hz, 1H), 7.73 (d, *J*=1.9 Hz, 1H), 7.68 (dd, *J*=8.2, 2.2 Hz, 1H), 7.10-7.21 (m, 1H), 5.08-5.26 (m, 1H), 3.96 (s, 3H), 2.95 (s, 3H), 1.59-1.75 (m, 4H), 1.40 (s, 6H), 1.33, (s, 6H), (1 hydrogen corresponding to OH or NH unobserved)

Step 2: 6-(2-Methoxy-4-(4-nitro-1H-pyrazol-1-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine (150 mg, 0.32 mmol), benzenethiol (33 μ L, 0.32 mmol) and K_2CO_3 (44 mg, 0.32 mmol) were mixed in a microwave tube. The reaction was degassed with N_2 for 15 min and dry NMP (1.5 mL) was added. The reaction was heated in a Biotage microwave at 190 $^{\circ}C$ for 20 min, then diluted with 10 mL EtOAc, and washed with water and brine. The product was dried over Na_2SO_4 , concentrated under vacuum, then purified via column chromatography: eluting with gradient $CH_2Cl_2/MeOH$ (0% to 30% MeOH), column: silica 4g, to provide 2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(4-nitro-1H-pyrazol-1-yl)phenol (60 mg, 42%) as a brown solid.

- LC-MS: 452 $[M+H]^+$. ^1H NMR (500 MHz, $DMSO-d_6$) δ : 9.77 (s, 1H), 8.68 (d, $J=9.8$ Hz, 1H), 8.62 (s, 1H), 8.27 (d, $J=8.8$ Hz, 1H), 7.87-7.94 (m, 1H), 7.66 (dd, $J=8.5, 2.6$ Hz, 1H), 7.64 (d, $J=2.6$ Hz, 1H), 5.03-5.22 (m, 1H), 3.01 (s, 3H), 1.84-1.94 (m, 2H), 1.70-1.77 (m, 2H), 1.53 (s, 6H), 1.43 (s, 6H), (2 hydrogens corresponding to OH and NH unobserved)

- Using the procedure described for Example 3 above, additional compounds described herein may be prepared by substituting the appropriate starting materials reagents, and reaction conditions, obtaining compounds such as those selected from:

Cpd	Data
417	LC-MS: 394 $[M+H]^+$. ^1H NMR (500 MHz, $DMSO-d_6$) δ : 13.49 (br s, 1H), 8.49 (d, $J=9.8$ Hz, 1H), 8.39 (s, 1H), 8.10 (d, $J=8.8$ Hz, 1H), 7.79-7.92 (m, 1H), 7.43 (d, $J=9.5$ Hz, 1H), 7.34 (d, $J=2.5$ Hz, 1H), 7.30 (dd, $J=8.5, 2.5$ Hz, 1H), 7.12 (s, 1H), 5.61-5.71 (m, 1H), 2.09 (dd, $J=12.0, 3.8$ Hz, 2H), 1.25-1.32 (m, 2H), 1.24 (s, 6H), 1.11 (s, 6H), (1 hydrogen corresponding to OH or NH unobserved)
421	LC-MS: 409 $[M+H]^+$. ^1H NMR (500 MHz, methanol- d_4) δ : 8.31 (d, $J=9.5$ Hz, 1H), 7.92 (d, $J=9.1$ Hz, 1H), 7.79 (d, $J=0.6$ Hz, 1H), 7.44 (d, $J=0.9$ Hz, 1H), 7.24-7.33 (m, 3H), 5.77 (tt, $J=11.2, 4.1$ Hz, 1H), 2.28 (dd, $J=12.3, 4.2$ Hz, 2H), 1.49 (t, $J=12.3$ Hz, 2H), 1.42 (s, 6H), 1.30 (s, 6H), (4 hydrogens corresponding to OH and NH unobserved)
423	LC-MS: 439 $[M+H]^+$. ^1H NMR (500 MHz, $DMSO-d_6$) δ : 9.71 (d, $J=0.6$ Hz, 1H), 8.58 (d, $J=0.6$ Hz, 1H), 8.47 (d, $J=9.5$ Hz, 1H), 8.14 (d, $J=8.5$ Hz, 1H), 7.59-7.64 (m, 2H), 7.44 (d, $J=9.5$ Hz, 1H), 5.67-5.73 (m, 1H), 2.18-2.28 (m, 2H), 1.45-1.56 (m, 2H), 1.37 (s, 6H), 1.27 (s, 6H), (2 hydrogens corresponding to OH and NH unobserved)

Example 4 (Compound 424)

5-(1H-Pyrazol-4-yl)-2-[6-(1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]phenol trihydrochloride

Step 1: 3-Bromo-6-(2-methoxy-4-(1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazol-4-

yl)phenyl)pyridazine (100 mg, 0.24 mmol), (1-(tert-butoxycarbonyl)-1,2,3,6-tetrahydropyridin-

5 4-yl)boronic acid (66 mg, 0.29 mmol), PddppfCl₂ (8 mg, 0.012 mmol), K₂CO₃ (99 mg, 0.72 mmol) were mixed in a Schlenk tube. The reaction was degassed with N₂ for 15 min and dioxane (2 mL) and water (0.5 mL) were added and the reaction was heated to 90 °C for 16 h. The reaction was cooled to room temperature, and partitioned between EtOAc and water. The organic layers were dried over Na₂SO₄, then concentrated under vacuum, and purified via column chromatography: eluting with gradient CH₂Cl₂/MeOH (0% to 20% MeOH), column: silica 4g, to provide 5-(1H-pyrazol-4-yl)-2-(6-(1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl)phenol (102 mg, 82%) as an off-white solid.

Step 2: A solution of 5-(1H-pyrazol-4-yl)-2-(6-(1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl)phenol (102 mg, 0.196 mmol) in 2 mL of dry CH₂Cl₂ was cooled in ice-water bath. Boron

15 tribromide 1.0 M in CH₂Cl₂ (0.98 mL, 0.98 mmol) was added and the reaction mixture was stirred at room temperature for 16 hours. The reaction was quenched with 2 mL of MeOH, stirred for 30 min, then concentrated and purified using preparative HPLC to provide 5-(1H-pyrazol-4-yl)-2-(6-(1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl)phenol (41 mg, 66%) as an off-white solid.

20 LC-MS: 320 [M+H]⁺. ¹H NMR (500 MHz, DMSO-*d*₆) δ: 9.19 (br s, 2H), 8.54 (d, *J*=9.6Hz, 1H), 8.22, (d, *J*=9.6Hz, 1H), 8.18 (s, 1H), 8.05 (d, *J*=8.8Hz, 1H), 7.25-7.29 (m, 2H), 6.92 (s, 1H), 3.85-3.92 (m, 2H), 3.38-3.41 (m, 2H), 2.88-2.98 (m, 2H), (1 hydrogen corresponding to OH or NH unobserved)

Using the procedure described for Example 4 above, additional compounds described herein may be prepared by substituting the appropriate starting materials reagents, and reaction conditions, obtaining compounds such as those selected from:

Cpd	Data
422	LC-MS: 334 [M+H] ⁺ . ¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ: 11.06 (br s, 1H), 8.56 (d, <i>J</i> =9.2Hz, 1H), 8.25 (d, <i>J</i> =9.2Hz, 1H), 8.20 (s, 2H), 8.05 (d, <i>J</i> =8.8Hz, 1H), 7.22-7.31 (m, 2H), 6.92 (s, 1H), 4.06-4.12 (m, 1H), 3.87-3.94 (m, 1H), 3.62-3.70 (m, 1H), 3.21-3.30 (m, 1H), 3.07-3.12 (m, 2H), 3.03 (s, 3H), (1 hydrogen corresponding to OH or NH unobserved)

Cpd	Data
425	LC-MS: 348 [M+H] ⁺ . ¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ: 10.94 (br s, 1H), 8.56 (d, <i>J</i> =9.6Hz, 1H), 8.26 (d, <i>J</i> =9.6Hz, 1H), 8.19-8.23 (m, 2H), 8.05 (d, <i>J</i> =8.4Hz, 1H), 7.26-7.31 (m, 2H), 6.93 (s, 1H), 4.07-4.13 (m, 1H), 3.85-3.91 (m, 1H), 3.70-3.74 (m, 1H), 3.22-3.30 (m, 3H), 3.08-3.14 (m, 2H), 1.33 (t, <i>J</i> =7.2Hz, 3H), (1 hydrogen corresponding to OH or NH unobserved)
429	LC-MS: 346 [M+H] ⁺ . ¹ H NMR (500 MHz, methanol- <i>d</i> ₄) δ: 1.79 - 1.89 (m, 1H) 1.98 - 2.11 (m, 1H) 2.11 - 2.33 (m, 2H) 2.65 - 2.80 (m, 1H) 3.09 - 3.21 (m, 1H) 4.02 - 4.16 (m, 2H) 7.02 - 7.14 (m, 1H) 7.24 (s, 2 H) 7.85 - 7.97 (m, 1H) 8.04 (d, <i>J</i> =9.46 Hz, 3H) 8.33 (d, <i>J</i> =9.46 Hz, 1H)

Example 5 (Compound 427)

2-[6-(Piperidin-4-ylamino)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol tetrahydrochloride

Step 1: 3,6-Dibromopyridazine (500 mg, 2.1 mmol), 4-(3-methoxy-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazole (970 mg, 2.52 mmol),

5 Pd(dppf)Cl₂ (77 mg, 0.11 mmol), K₂CO₃ (870 mg, 6.3 mmol) were mixed in a Schlenk tube. The reaction was degassed with N₂ for 15 min and dioxane (8 mL) and water (1 mL) were added and the reaction was heated to 90 °C for 16 h. The reaction was cooled to room temperature, partitioned between EtOAc and water. The organic layers were dried over Na₂SO₄, concentrated under vacuum, purified via column chromatography: eluting with gradient hexanes/EtOAc (0% to 40% EtOAc), column: silica 4g, to provide 3-bromo-6-(2-methoxy-4-(1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazol-4-yl)phenyl)pyridazine (690 mg, 79%) as an off-white fluffy solid.

Step 2: 3-Bromo-6-(2-methoxy-4-(1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazol-4-yl)phenyl)pyridazine (50 mg, 0.12 mmol), tert-butyl 4-aminopiperidine-1-carboxylate (36 mg, 0.18 mmol), potassium t-butoxide (41 mg, 0.36 mmol), *t*BuXPhos-Pd-G3 (10 mg, 0.012 mmol)

15 were mixed in a Schlenk tube. The reaction was degassed with Ar then dry THF (2 mL) was added and the reaction was heated to 80 °C for 12 h. The reaction was cooled to room temperature, then partitioned between EtOAc and water. The organic layers were dried over Na₂SO₄, concentrated under vacuum, and purified via column chromatography: eluting with gradient CH₂Cl₂/MeOH (0% to 15% MeOH), column: silica 4g, to provide tert-butyl 4-((6-(2-methoxy-4-(1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazol-4-yl)phenyl)pyridazin-3-yl)amino)piperidine-1-carboxylate (45 mg, 70%) as a tan solid.

Step 3: A solution of 4-((6-(2-methoxy-4-(1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazol-4-

yl)phenyl)pyridazin-3-yl)amino)piperidine-1-carboxylate (45 mg, 0.084 mmol) in 2 mL of dry CH₂Cl₂ was cooled in an ice-water bath. Boron tribromide 1.0 M in CH₂Cl₂ (0.42 mL, 0.42 mmol) was added and the reaction mixture was stirred at room temperature for 16 hours. The reaction was quenched with 2 mL of MeOH, stirred for 30 min, then concentrated and purified using preparative HPLC to provide 2-(6-(piperidin-4-ylamino)pyridazin-3-yl)-5-(1H-pyrazol-4-yl)phenol (21 mg, 73%) as a yellowish solid.

LC-MS: 377 [M+H]⁺. ¹H NMR (500 MHz, DMSO-*d*₆) δ: 8.95 (s, 2H), 8.25 (d, *J*=9.6 Hz, 1H), 8.09 (s, 2H), 7.69 (d, *J*=8.8 Hz, 1H), 7.49-7.41 (m, 1H), 7.21-7.24 (m, 2H), 4.11-4.15 (m, 1H), 3.34-3.39 (m, 4H), 3.01-3.08 (m, 2H), 2.11-2.17 (m, 2H), 1.79 (q, *J*=9.2 Hz, 2H)

- 10 Using the procedure described for Example 5 above, additional compounds described herein may be prepared by substituting the appropriate starting materials reagents, and reaction conditions, obtaining compounds such as those selected from:

Cpd	Data
426	LC-MS: 351 [M+H] ⁺ . ¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ: 9.22-9.28 (m, 2H), 8.34 (d, <i>J</i> =9.6Hz, 1H), 8.06-8.11 (m, 2H), 7.85-7.89 (m, 1H), 7.74 (d, <i>J</i> =8.2 Hz, 1H), 7.20-7.27 (m, 2H), 4.67-4.75 (m, 2H), 3.35-3.39 (m, 2H), 3.07-3.09 (m, 2H), 3.05 (s, 3H), 2.21 (q, <i>J</i> =10.2Hz, 2H), 1.83-1.88 (m, 2H)
432	LC-MS: 338 [M+H] ⁺ . ¹ H NMR (500 MHz, methanol- <i>d</i> ₄) δ: 8.72 (d, <i>J</i> =9.6Hz, 1H), 8.41-8.54 (m, 2H), 7.91-8.08 (m, 1H), 7.82 (d, <i>J</i> =8.0Hz, 1H), 7.49 (d, <i>J</i> =8.2 Hz, 1H), 7.38 (s, 1H), 5.51 (quint, <i>J</i> =2.8Hz, 1H), 3.30-3.46 (m, 2H), 3.28-3.31 (m, 2H), 2.33-2.38 (m, 2H), 2.25-2.32 (m, 2H), (3 hydrogens corresponding to OH and NH unobserved)
436	LC-MS: 366 [M+H] ⁺ . ¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ: 9.17-9.30 (m, 1H), 8.81-8.95 (m, 1H), 8.49 (d, <i>J</i> =9.8 Hz, 1H), 8.17 (s, 2H), 7.94 (d, <i>J</i> =8.8 Hz, 1H), 7.46 (d, <i>J</i> =9.8 Hz, 1H), 7.18-7.29 (m, 2H), 5.47 (spt, <i>J</i> =5.7 Hz, 1H), 3.31-3.52 (m, 2H), 2.41 (dd, <i>J</i> =12.9, 5.7 Hz, 2H), 1.68 (q, <i>J</i> =12.0 Hz, 2H), 1.34 (d, <i>J</i> =6.3 Hz, 6H), (1 hydrogen corresponding to OH or NH unobserved)
439	LC-MS: 363 [M+H] ⁺ . ¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ: 9.07 (s, 2H), 8.28 (d, <i>J</i> =10Hz, 1H), 1.7 (s, 2H), 7.81 (d, <i>J</i> =8.8Hz, 1H), 7.73 (d, <i>J</i> =9.6Hz, 1H), 7.19-7.22 (m, 2H), 3.75-3.81 (m, 4H), 3.58-3.69 (m, 4H), 1.85-1.89 (m, 4H), (1 hydrogen corresponding to OH or NH unobserved)

Cpd	Data
459	LC-MS: 349 [M+H] ⁺ . ¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ: 10.53 (s, 1H), 10.40 (s, 1H), 9.79 (br s, 1H), 9.67 (br s, 1H), 8.28 (d, <i>J</i> =9.6Hz, 1H), 8.06 (s, 2H), 7.65 (d, <i>J</i> =9.6Hz, 1H), 7.58 (d, <i>J</i> =8.4Hz, 1H), 7.21-7.25 (m, 1H), 4.53 (q, <i>J</i> =14Hz, 2H), 3.64 (d, <i>J</i> =12.8Hz, 1H), 3.52 (d, <i>J</i> =12.8Hz, 1H), 3.37-3.40 (m, 2H), 3.15-3.24 (m, 2H), 1.92-2.02 (m, 2H)

Example 6 (Compound 430)

6-[2,3-Difluoro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine hydrochloride

- Step 1:** To a RBF, equipped with a N₂ inlet, were added: 6-bromo-N-methyl-N-(2,2,6,6-tetramethyl-4-piperidyl)pyridazin-3-amine (90 mg, 0.28 mmol), 4-[2,3-difluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]-1-tetrahydropyran-2-yl-pyrazole (128.8 mg, 0.33 mmol), Pd(PPh₃)₄ (31.8 mg, 0.03 mmol), Na₂CO₃ (87.43 mg, 0.83 mmol). The reaction was degassed with N₂ for 15 min then dioxane (8 mL) and water (2 mL) were added. The reaction was heated to 90 °C for 16 h, then cooled to room temperature, and partitioned between EtOAc and water. The organic layers were dried over Na₂SO₄, concentrated on a rotavap, and purified via column chromatography: eluting with gradient CH₂Cl₂/MeOH (0% to 30%), column: silica 4g to provide 6-(2,3-Difluoro-4-(1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine (88 mg, 63%) as a grey solid.
- Step 2:** 6-[2,3-Difluoro-4-(1-tetrahydropyran-2-ylpyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethyl-4-piperidyl)pyridazin-3-amine (88 mg, 0.17 mmol) was dissolved in MeOH (2 mL) and 4M HCl in dioxane (90 μL, 0.34 mmol) was added dropwise. The reaction mixture was stirred at room temperature for 30 min until LC-MS showed complete consumption of starting material. The reaction mixture was concentrated under reduced pressure to provide 6-(2,3-difluoro-4-(1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine hydrochloride salt.

LC-MS: 427 [M+H]⁺. ¹H NMR (500 MHz, DMSO-*d*₆) δ: 9.21-9.33 (m, 1H), 8.27-8.43 (m, 1H), 8.20 (s, 2H), 8.01 (d, *J*=9.8 Hz, 1H), 7.60-7.79 (m, 3H), 5.01-5.14 (m, 1H), 3.05 (s, 3H), 2.10 (td, *J*=13.2, 1.6 Hz, 2H), 1.79 (dd, *J*=12.9, 4.1 Hz, 2H), 1.55 (s, 6H), 1.50 (s, 6H)

Using the procedure described for Example 6 above, additional compounds described herein may be prepared by substituting the appropriate starting materials reagents, and reaction conditions, obtaining compounds such as those selected from:

Cpd	Data
426	LC-MS: 427 [M+H] ⁺ . ¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ: 9.22-9.38 (m, 1H), 8.33-8.45 (m, 1H), 8.21 (d, J=1.9 Hz, 2H), 7.97 (dd, J=10.4, 2.8 Hz, 1H), 7.86 (d, J=5.7 Hz, 1H), 7.81 (dd, J=11.7, 6.6 Hz, 1H), 7.54-7.63 (m, 1H), 5.09-5.16 (m, 1H), 3.03 (s, 3H), 2.09 (t, J=12.9 Hz, 2H), 1.78 (dd, J=12.9, 3.8 Hz, 2H), 1.55 (s, 6H), 1.51 (s, 6H)
431	LC-MS: 414 [M+H] ⁺ . ¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ: 9.24-9.35 (m, 1H), 8.43-8.58 (m, 1H), 8.22 (d, J=1.6 Hz, 2H), 8.06 (dd, J=9.1, 1.9 Hz, 1H), 7.87 (dd, J=12.1, 6.5 Hz, 1H), 7.83 (dd, J=11.5, 6.5 Hz, 1H), 7.37 (d, J=9.5 Hz, 1H), 5.77 (tt, J=10.7, 4.1 Hz, 1H), 2.33 (dd, J=13.1, 3.9 Hz, 2H), 1.84 (dd, J=12.9, 11.3 Hz, 2H), 1.54 (s, 6H), 1.52 (s, 6H)
435	LC-MS: 378 [M+H] ⁺ . ¹ H NMR (500 MHz, methanol- <i>d</i> ₄) δ: 8.57-8.58 (m, 1H), 8.55 (d, J=5.0 Hz, 1H), 8.45 (s, 2H), 8.05-8.10 (m, 1H), 7.95-7.99 (m, 1H), 7.86 (s, 1H), 7.82 (d, J=9.5 Hz, 1H), 5.79 (tt, J=10.4, 4.4 Hz, 1H), 2.53 (dd, J=14.5, 4.5 Hz, 2H), 1.86-2.01 (m, 2H), 1.66 (s, 6H), 1.60 (s, 6H), (2 hydrogens corresponding to NH unobserved)
437	LC-MS: 396 [M+H] ⁺ . ¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ: 9.28-9.44 (m, 1H), 8.47-8.61 (m, 1H), 8.25 (s, 2H), 8.01 (dd, J=9.3, 2.0 Hz, 1H), 7.92 (t, J=8.2 Hz, 1H), 7.69 (dd, J=12.8, 1.7 Hz, 1H), 7.65 (dd, J=8.0, 1.7 Hz, 1H), 7.34 (d, J=9.1 Hz, 1H), 5.76 (tt, J=10.7, 4.1 Hz, 1H), 2.33 (dd, J=13.2, 3.9 Hz, 2H), 1.85 (dd, J=13.2, 10.1 Hz, 2H), 1.54 (s, 6H), 1.53 (s, 6H)
440	LC-MS: 359 [M+H] ⁺ . ¹ H NMR (500 MHz, methanol- <i>d</i> ₄) δ: .06 (d, J=10.1 Hz, 1H), 7.78 (dd, J=9.5, 6.3 Hz, 1H), 7.32 (d, J=9.8 Hz, 1H), 6.58-6.76 (m, 2H), 5.11 (tt, J=12.0, 4.1 Hz, 1H), 3.02 (s, 3H), 1.73 (dd, J=12.9, 3.5 Hz, 2H), 1.63 (t, J=12.5 Hz, 2H), 1.43 (s, 6H), 1.27 (s, 6H), (1 hydrogen corresponding to NH unobserved)
441	LC-MS: 364 [M+H] ⁺ . ¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ: 1.33 (d, J=6.62 Hz, 6H) 1.65 (q, J=12.40 Hz, 2H) 2.42 (d, J=12.61 Hz, 2H) 3.38 - 3.50 (m, 2H) 5.48 (s, 1H) 6.51 - 6.63 (m, 1H) 7.42 - 7.57 (m, 3H) 7.80 (d, J=1.58 Hz, 1H) 8.09 (d, J=8.83 Hz, 1H) 8.50 (d, J=9.46 Hz, 1H) 8.61 (d, J=2.52 Hz, 1H) 8.76 (br s., 1H) 9.12 (br s., 1H) 13.18 (s, 1H)

Cpd	Data
442	LC-MS: 379 [M+H] ⁺ . ¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ: 8.42 (s, 1H), 8.28 (dd, <i>J</i> =1.6, 0.9 Hz, 1H), 8.03 (dd, <i>J</i> =9.1, 1.6 Hz, 1H), 7.95 (s, 1H), 7.67 (d, <i>J</i> =9.1 Hz, 1H), 7.14 (d, <i>J</i> =9.1 Hz, 1H), 5.03-5.17 (m, 1H), 4.19 (s, 3H), 2.94 (s, 3H), 1.49-1.58 (m, 2H), 1.37-1.48 (m, 2H), 1.27 (s, 6H), 1.10 (s, 6H), (1 hydrogen corresponding to NH unobserved)
443	LC-MS: 366 [M+H] ⁺ . ¹ H NMR (500 MHz, methanol- <i>d</i> ₄) δ: 8.35 (s, 1H), 8.31 (dd, <i>J</i> =1.6, 0.9 Hz, 1H), 8.11 (d, <i>J</i> =9.1 Hz, 1H), 7.99 (dd, <i>J</i> =9.1, 1.9 Hz, 1H), 7.75 (dt, <i>J</i> =9.1, 0.9 Hz, 1H), 7.21 (d, <i>J</i> =9.5 Hz, 1H), 5.81 (tt, <i>J</i> =11.2, 4.1 Hz, 1H), 4.27 (s, 3H), 2.28 (dd, <i>J</i> =12.6, 4.1 Hz, 2H), 1.48 (t, <i>J</i> =12.3 Hz, 2H), 1.41 (s, 6H), 1.30 (s, 6H), (1 hydrogen corresponding to NH unobserved)
445	LC-MS: 379 [M+H] ⁺ . ¹ H NMR (500 MHz, acetone- <i>d</i> ₆) δ: 8.52 (dt, <i>J</i> =7.0, 1.1 Hz, 1H), 8.13 (dt, <i>J</i> =8.8, 1.3 Hz, 1H), 7.66 (d, <i>J</i> =9.8 Hz, 1H), 7.28 (ddd, <i>J</i> =9.0, 6.8, 1.3 Hz, 1H), 7.13 (d, <i>J</i> =9.5 Hz, 1H), 6.89 (td, <i>J</i> =6.9, 1.4 Hz, 1H), 5.22 (tt, <i>J</i> =12.3, 3.5 Hz, 1H), 3.04 (s, 3H), 2.63 (s, 3H), 1.67 (dd, <i>J</i> =12.3, 3.5 Hz, 2H), 1.56 (t, <i>J</i> =12.3 Hz, 2H), 1.36 (s, 6H), 1.18 (s, 6H), (1 hydrogen corresponding to NH unobserved)
446	LC-MS: 352 [M+H] ⁺ . ¹ H NMR (500 MHz, acetone- <i>d</i> ₆) δ: 9.14 (dd, <i>J</i> =1.6, 0.9 Hz, 1H), 7.98 (dd, <i>J</i> =9.5, 1.9 Hz, 1H), 7.94 (t, <i>J</i> =0.9 Hz, 1H), 7.89 (d, <i>J</i> =9.5 Hz, 1H), 7.63 (dt, <i>J</i> =9.5, 0.8 Hz, 1H), 7.60 (d, <i>J</i> =1.3 Hz, 1H), 7.14 (d, <i>J</i> =9.5 Hz, 1H), 5.34 (tt, <i>J</i> =12.6, 3.5 Hz, 1H), 1.67 (dd, <i>J</i> =12.3, 3.8 Hz, 2H), 1.55 (t, <i>J</i> =12.1 Hz, 2H), 1.36 (s, 6H), 1.17 (s, 6H), (1 hydrogen corresponding to NH unobserved)
447	LC-MS: 394 [M+H] ⁺ . ¹ H NMR (500 MHz, acetone- <i>d</i> ₆) δ: 8.47 (d, <i>J</i> =2.5 Hz, 1H), 8.05 (d, <i>J</i> =9.1 Hz, 1H), 7.98 (d, <i>J</i> =8.5 Hz, 1H), 7.76 (d, <i>J</i> =1.9 Hz, 1H), 7.70 (d, <i>J</i> =2.2 Hz, 1H), 7.59 (dd, <i>J</i> =8.5, 2.2 Hz, 1H), 7.06 (d, <i>J</i> =9.1 Hz, 1H), 6.57 (dd, <i>J</i> =2.4, 1.7 Hz, 1H), 5.85 (tt, <i>J</i> =11.3, 4.1 Hz, 1H), 2.21 (dd, <i>J</i> =12.1, 3.9 Hz, 2H), 2.04-2.08 (m, 2H), 1.35 (s, 6H), 1.18 (s, 6H), (2 hydrogens corresponding to OH and NH unobserved)
452	LC-MS: 409 [M+H] ⁺ . ¹ H NMR (500 MHz, acetone- <i>d</i> ₆) δ: 7.59 (dd, <i>J</i> =8.5, 6.0 Hz, 1H), 7.46 (s, 2H), 7.31 (dd, <i>J</i> =10.1, 2.5 Hz, 1H), 7.15 (td, <i>J</i> =8.5, 2.5 Hz, 1H), 7.04 (d, <i>J</i> =9.5 Hz, 1H), 6.91 (d, <i>J</i> =9.5 Hz, 1H), 5.23 (tt, <i>J</i> =12.3, 2.8 Hz, 1H), 3.00 (s, 3H), 1.65 (dd, <i>J</i> =12.3, 3.5 Hz, 2H), 1.51 (t, <i>J</i> =12.1 Hz, 2H), 1.33 (s, 6H), 1.15 (s, 6H), (2 hydrogens corresponding to NH unobserved)
455	LC-MS: 344 [M+H] ⁺ . ¹ H NMR (500 MHz, methanol- <i>d</i> ₄) δ: 8.17 (d, <i>J</i> =9.5 Hz, 1H), 7.66 (d, <i>J</i> =8.8 Hz, 1H), 7.19 (d, <i>J</i> =9.5 Hz, 1H), 6.44 (dd, <i>J</i> =8.7, 2.4 Hz, 1H), 6.41 (d, <i>J</i> =2.5 Hz, 1H), 5.71 (tt, <i>J</i> =11.0, 3.8 Hz, 1H), 2.23 (dd, <i>J</i> =12.6, 4.1 Hz, 2H), 1.40-1.47 (m, 2H), 1.39 (s, 6H), 1.27 (s, 6H), (3 hydrogens corresponding to OH and NH unobserved)

Cpd	Data
456	LC-MS: 425 [M+H] ⁺ . ¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ: 9.23-9.39 (m, 1H), 8.39-8.52 (m, 1H), 8.26 (s, 2H), 8.00-8.08 (m, 1H), 7.93 (s, 1H), 7.81-7.85 (m, 1H), 7.78 (d, <i>J</i> =7.9 Hz, 1H), 7.63 (d, <i>J</i> =8.2 Hz, 1H), 4.95-5.03 (m, 1H), 3.08 (s, 3H), 2.13 (t, <i>J</i> =12.9 Hz, 2H), 1.81 (dd, <i>J</i> =12.9, 3.5 Hz, 2H), 1.56 (s, 6H), 1.52 (s, 6H)
458	LC-MS: 414 [M+H] ⁺ . ¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ: 8.70-9.36 (m, 1H), 8.29 (s, 2H), 7.85 (d, <i>J</i> =9.1 Hz, 1H), 7.60 (d, <i>J</i> =9.5 Hz, 1H), 7.33-7.39 (m, 1H), 5.59-5.68 (m, 1H), 2.24-2.33 (m, 2H), 1.72-1.82 (m, 2H), 1.50 (s, 6H), 1.48 (s, 6H), (2 hydrogens corresponding to NH unobserved)

Example 7 (Compound 434)

6-[2-Methoxy-6-(1H-pyrazol-4-yl)pyridin-3-yl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine hydrochloride

- 6-(2-Methoxy-6-(1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazol-4-yl)pyridin-3-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine (6 mg) was dissolved in 1 mL of 4N HCl/dioxane and the resulting mixture was stirred at RT for 20 minutes. The mixture was evaporated and triturated with ether to provide 6-(2-methoxy-6-(1H-pyrazol-4-yl)pyridin-3-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine 4 mg, 75%).
- LC-MS: 422 [M+H]⁺. ¹H NMR (500 MHz, methanol-*d*₄) δ: 8.42 - 8.47 (m, 1H), 8.33 - 8.42 (m, 2 H), 8.09 - 8.17 (m, 1H), 7.99 - 8.08 (m, 1H), 7.50 - 7.56 (m, 1H), 5.0 (m, 1H), 4.16 (s, 3H), 3.20 (s, 3H), 1.94 - 2.15 (m, 4H), 1.65 (s, 6H), 1.57 (s, 6H)

Example 8 (Compound 460)

3-{6-[Methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl}-6-(1H-pyrazol-4-yl)pyridin-2-ol hydrochloride

- Step 1:** 6-Chloro-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine (280 mg, 1.0 mmol) was dissolved in 10 mL, a mixture of dioxane and water (4:1). To the solution was added sodium carbonate (190 mg, 3 eq), (6-chloro-2-methoxypyridin-3-yl)boronic acid (190 mg, 1 eq) and Pd(Ph₃P)₄ (70 mg, 0.1 eq). The reaction mixture was stirred at 90 °C for 16 h, then applied to a column and purified using DCM/MeOH to yield (133 mg, 55%) 6-(6-chloro-2-methoxypyridin-3-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine.
- LC-MS: 390 [M+H]⁺. ¹H NMR (500 MHz, DMSO-*d*₆) δ: 8.25 (d, *J*=7.88 Hz, 1H), 7.83 (d, *J*=9.77 Hz, 1H), 7.24 (d, *J*=7.88 Hz, 1H), 7.11 (d, *J*=9.77 Hz, 1H), 5.00 - 5.24 (m, 1H), 3.93 (s, 3 H), 2.87 - 2.96 (m, 3H), 1.40 - 1.57 (m, 4H), 1.21 - 1.29 (m, 7H), 1.10 (br s, 6H)

Step 2: 6-(6-Chloro-2-methoxypyridin-3-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-

yl)pyridazin-3-amine (90 mg) was dissolved in a mixture of dioxane and water (4:1, 5 mL). To the solution was added sodium carbonate (100 mg, 3 eq), 1-(tetrahydro-2H-pyran-2-yl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole (70 mg, 1.1 eq) and Pd(Ph₃P)₄ (27 mg, 0.1 eq). The reaction mixture was stirred at 90 °C for 16 h, then purified by column chromatography using DCM/MeOH to yield 80 mg (68%) of 6-(2-methoxy-6-(1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazol-4-yl)pyridin-3-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine.

Step 3: 6-(2-Methoxy-6-(1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazol-4-yl)pyridin-3-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine (45 mg) was dissolved in 4 mL of 4N HCl/dioxane and the resulting mixture was heated for 2 hours at 60 °C. The resulting mixture was evaporated and triturated with ether to provide 3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-6-(1H-pyrazol-4-yl)pyridin-2-ol (24 mg, 61%).

LC-MS: 408 [M+H]⁺; ¹H NMR (500 MHz, methanol-*d*₄) δ: 8.62 - 8.69 (m, 1H), 8.55 - 8.61 (m, 1H), 8.39 (s, 2H), 7.95 - 8.06 (m, 1H), 7.04 - 7.14 (m, 1H), 5.02 - 5.24 (m, 1H), 3.18 (s, 3H), 2.03 (s, 4H), 1.67 (s, 6H), 1.57 (s, 6H)

Using the procedure described for Example 8 above, an additional compound described herein may be prepared by substituting the appropriate starting materials reagents, and reaction conditions, obtaining compounds such as those selected from:

Cpd	Data
461	LC-MS: 395 [M+H] ⁺ . ¹ H NMR (500 MHz, methanol- <i>d</i> ₄) δ: 8.46 (d, J=9.5 Hz, 1H), 8.33-8.42 (m, 1H), 8.29 (d, J=7.6 Hz, 1H), 8.10-8.23 (m, 1H), 7.20 (d, J=9.1 Hz, 1H), 6.86 (d, J=7.3 Hz, 1H), 5.84 (tt, J=10.4, 3.8 Hz, 1H), 2.49 (dd, J=13.9, 4.1 Hz, 2H), 1.86 (dd, J=13.9, 10.4 Hz, 2H), 1.65 (s, 6H), 1.56 (s, 6H), (3 hydrogens corresponding to OH and NH unobserved)

Biological Examples

The following *in vitro* biological examples demonstrate the usefulness of the compounds of the present description for treating Huntington's disease.

To describe in more detail and assist in understanding the present description, the following non-limiting biological examples are offered to more fully illustrate the scope of the description and are not to be construed as specifically limiting the scope thereof. Such variations of the present description that may be now known or later developed, which would be within the

purview of one skilled in the art to ascertain, are considered to fall within the scope of the present description and as hereinafter claimed.

Example 1

5 **Development and Validation of Meso Scale Discovery (MSD) Assay to Quantify Mutant and Total Huntingtin Protein (HTT) – Best Pair Analysis for Endogenous HTT Detection**

To identify orally bioavailable small molecules useful for HTT reduction, high throughput ELISA methods to detect and measure both total HTT (wild-type and mutant)(tHTT) and mutant HTT (mHTT) in cells and tissues from HD patients and animal models are needed. The method described herein permits relatively high throughput protein measurement of tHTT and mHTT levels in cells and tissues from HD patients and in HD animal models. The ELISA method described herein enables screening of molecules for their ability to decrease the level of HTT and to identify and optimize small molecules for use in treating HD.

To determine the combination of antibodies that detect HTT (wild type and mutant) in a sensitive, high throughput format, multiple commercially available antibodies (Table 1) were screened in capture/detection combinations.

Protocol for Best Pair Analysis for Antibody Comparison in Detecting Total and Mutant HTT in Cells

1. Cells were cultured in a cell culture-compatible 96-well plate.
2. The medium was removed and 50-100 μ L of lysis buffer per well (composition described below) was added to cells to provide a “cell lysate”. The plate was placed on a shaker at 4° C for 30 minutes, then stored at -20° C.
3. A Capture Antibody Diluent was prepared by diluting the capture antibody in PBS for standard plates as described in the plate layout, then 30 μ L of the Capture Antibody Diluent was added per well. The plate was sealed and incubated at 4° C overnight.
- 25 4. The plate was washed 3X with 200 μ L of Wash Buffer. Blocking Buffer (150 μ L per well) was added, then the plate was sealed and incubated on a plate shaker for 3-4 hours at room temperature (RT).
5. The Blocking Buffer was decanted and the plate was washed 3X with 200 μ L of Wash Buffer. The cell lysate (25 μ L per well) was then added. The plate was sealed and incubated at 4° C overnight.
- 30

6. A First Detection Antibody Diluent was prepared by diluting a first detection antibody for pairing in an Assay Antibody Diluent. The plate was washed 3X with 200 μ L of Wash Buffer, then 25 μ L of the First Detection Antibody Diluent was added per well. The plate was sealed and incubated on a plate shaker for 1 hour at RT.
- 5 7. A Second Detection Antibody Diluent was prepared by diluting a second detection antibody for pairing in an Assay Antibody Diluent. The plate was washed 3X with 200 μ L of Wash Buffer, then 25 μ L of the Second Detection Antibody Diluent was added per well. The plate was sealed and incubated on a plate shaker for 1 hour at RT.
8. The Read Buffer was diluted 4X. The plate was washed 3X with 200 μ L of Wash Buffer, then 150 μ L of the Read Buffer was added per well. The plate were read immediately following.
- 10

The various commercially available antibodies screened in the Best Pair Analysis, providing different capture/detection combinations are shown in Table 1. The capture/detection antibody pairs evaluated are shown in Table 2. The two antibody pairs that gave the maximum spread of values (higher signal to noise ratio) were selected for polyglutamine-expanded (mHTT) and total human HTT detection.

15

Table 1

Species	ID	Antibody Epitope
monoclonal mouse	MAB2166	Huntingtin fragment from aa 181-810 as a fusion protein
monoclonal rat	MAB2174	Huntingtin fusion protein 549-679, fused to Glutathione S-transferase
monoclonal mouse	MAB5374	GST fusion protein from the first 256 amino acids from human huntingtin with the deletion of the polyglutamine tract
goat polyclonal	sc-8767	Peptide mapping near the N-terminus of Huntingtin of human origin (N18)
monoclonal mouse	P1874	GST human Huntingtin (N-terminal fragment of 171 amino acids containing 65Q)
monoclonal mouse	MAB1574	Homopolymeric glutamine stretch
monoclonal rabbit	#5656S	Synthetic peptide corresponding to residues surrounding Pro1220 of human huntingtin protein
monoclonal mouse	MW1	DRPLA-19Q

Table 2

Pair	HTT MSD	ID	Signal to Noise ratio
Capture/Detection	Total	MAB2166/MAB2174	95
Capture/Detection	Total	MAB2166/sc-8767	6
Capture/Detection	Total	sc-8767/#5656	5
Capture/Detection	Total	MAB2166/#5656S	483
Capture/Detection	Mutant	MAB5374/#5656S	8
Capture/Detection	Mutant	MAB1574/#5656S	25
Capture/Detection	Mutant	P1874/#5656S	20
Capture/Detection	Mutant	MW1/#5656S	45

Characteristics for Best Pair capture/detection antibody pairs are those combinations that give minimum background and maximum signal. Such highly sensitive ELISA assays would be useful to detect HTT in the context of drug discovery for screening agents (small and large molecules and gene therapies) and for use in detecting HTT as a biomarker in clinical studies.

- 5 Such capture/detection antibody pairs for quantifying total HTT are selected from MAB2166/MAB2174 and MAB2166/#5656S. Such capture/detection antibody pairs for quantifying mutant HTT are selected from MAB1574/#5656S, P1874/#5656S and MW1/#5656S. The antibody pair for mHTT (MW1/#5656S) and the other for total HTT (MAB2166/#5656S) from this analysis were further optimized and validated to provide high-
- 10 throughput assays that enabled HTT detection in patient fibroblasts and lymphocytes.

- The Endogenous Huntingtin Protein assay used in Example 2 was developed using the Best Pair antibodies identified in Table 2, providing highly sensitive, high-throughput detection assays for polyglutamine-expanded (mHTT) and total human HTT in multiple cell systems. The platform used for the assay is the ELISA-based Meso Scale Discovery (MSD)
- 15 electrochemiluminescence assay platform.

Example 2

Endogenous Huntingtin Protein Assay

- Meso Scale Discovery (MSD) 96-well or 384-well plates were coated overnight at 4°C with MW1 (expanded polyglutamine) or MAB2166 monoclonal antibody (for capture) at a
- 20 concentration of 1 µg/mL in PBS (30 µL per well). Plates were then washed three times with 300 µL wash buffer (0.05% Tween-20 in PBS) and blocked (100 µL blocking buffer; 5% BSA in PBS) for 4-5 hours at room temperature with rotational shaking and then washed three times with wash buffer.

Samples (25 μ L) were transferred to the antibody-coated MSD plate and incubated overnight at 4°C. After removal of the lysates, the plate was washed three times with wash buffer, and 25 μ L of #5656S (Cell signaling; rabbit monoclonal) secondary antibody (diluted to 0.25 μ g/mL in 0.05% Tween-20 in blocking buffer) was added to each well and incubated with shaking for 1 Hour at room temperature. Following incubation with the secondary antibody, the wells were rinsed with wash buffer after which 25 μ L of goat anti-rabbit SULFO TAG secondary detection antibody (required aspect of the MSD system) (diluted to 0.25 μ g/mL in 0.05% Tween-20 in blocking buffer) was added to each well and incubated with shaking for 1 hour at room temperature. After rinsing three times with wash buffer, 150 μ L of read buffer T with surfactant (MSD) were added to each empty well, and the plate was imaged on a SI 6000 imager (MSD) according to manufacturers' instructions provided for 96- or 384-well plates. The resulting IC₅₀ values (μ M) for compounds tested are shown in Table 3.

As shown in Table 3, test compounds described herein had the following IC₅₀ values, an IC₅₀ value between $> 3 \mu\text{M}$ and $\leq 9 \mu\text{M}$ is indicated by a single star (*), an IC₅₀ value between $> 1 \mu\text{M}$ and $\leq 3 \mu\text{M}$ is indicated by two stars (**), an IC₅₀ value between $> 0.5 \mu\text{M}$ and $\leq 1 \mu\text{M}$ is indicated by three stars (***), an IC₅₀ value between $> 0.1 \mu\text{M}$ and $\leq 0.5 \mu\text{M}$ is indicated by four stars (****) and an IC₅₀ value of $\leq 0.1 \mu\text{M}$ is indicated by five stars (*****).

Table 3

Cpd	IC ₅₀	Cpd	IC ₅₀	Cpd	IC ₅₀
16	*	413	*****	441	*
26	****	414	*****	442	*
31	****	415	*****	443	*
460	*	416	*****	444	*
39	*	417	*****	445	*
44	*****	418	*****	446	*
46	*****	419	*****	447	*
47	****	420	*****	448	*
48	*****	421	*****	449	*
51	*****	422	*****	450	*
63	*****	423	*****	451	*
64	*****	424	****	451	*
170	*****	425	****	452	*
176	*****	426	****	454	*


Cpd	IC₅₀	Cpd	IC₅₀	Cpd	IC₅₀
200	*****	427	****	455	*
207	****	428	****	456	*
212	*****	429	****	456	*
218	*****	430	****	456	*
258	****	431	****	457	*
307	****	432	****	458	*
315	*****	433	****	459	*
318	*****	434	***	460	*
348	*****	435	***	461	*
350	*****	436	***	462	*
352	****	437	**	463	*
393	*****	438	**	464	*
411	*****	439	**	465	*
412	*****	440	*		

Without regard to whether a document cited herein was specifically and individually indicated as being incorporated by reference, all documents referred to herein are incorporated by reference into the present application for any and all purposes to the same extent as if each individual reference was fully set forth herein.

- 5 Having now fully described the subject matter of the claims, it will be understood by those having ordinary skill in the art that the same can be performed within a wide range of equivalents without affecting the scope of the subject matter or embodiments described herein. It is intended that the appended claims be interpreted to include all such equivalents.

What is claimed is:

1. A method of use of a compound for treating or ameliorating HD in a subject in need thereof comprising administering to the subject an effective amount of a compound of Formula (I):



(I)

or a form thereof, wherein

W is CH=CH or S;

X is CH₂, CH(C₁₋₄alkyl), C(C₁₋₄alkyl)₂, CH=CH, O, NR₅, or a bond;

10 A is aryl, heteroaryl, heterocyclyl, or C₉₋₁₀cycloalkyl,

wherein aryl is selected from phenyl and naphthyl, each optionally substituted with 1, 2, 3, or 4 substituents each selected from R₁,

wherein heteroaryl is a saturated monocyclic, bicyclic or tricyclic ring system having 1, 2, or 3 heteroatom ring members independently selected from N, O, or S, each optionally

15 substituted with 1, 2, 3, 4, or 5 substituents each selected from R₁,

wherein heterocyclyl is a saturated or partially unsaturated monocyclic, bicyclic or tricyclic ring system having 1, 2, or 3 heteroatom ring members independently selected from N, O, or S, each optionally substituted with 1, 2, 3, 4, or 5 substituents each selected from R₂, and

20 wherein C₉₋₁₀cycloalkyl is a saturated or partially unsaturated bicyclic ring system optionally substituted with 1, 2, 3, 4, or 5 substituents each selected from R₂;

B is heterocyclyl,

wherein heterocyclyl is a saturated or partially unsaturated monocyclic, bicyclic or polycyclic ring system having 1, 2, or 3 heteroatom ring members independently selected from N, O, or S, each optionally substituted with 1, 2, 3, 4, or 5 substituents each selected from R₄;

25 R₁ is halogen, hydroxyl, cyano, C₁₋₄alkyl, halo-C₁₋₄alkyl, amino, C₁₋₄alkyl-amino,

(C₁₋₄alkyl)₂-amino, amino-C₁₋₄alkyl, C₁₋₄alkyl-amino-C₁₋₄alkyl,

(C₁₋₄alkyl)₂-amino-C₁₋₄alkyl, amino-carbonyl, C₁₋₄alkyl-amino-carbonyl,


(C₁₋₄alkyl)₂-amino-carbonyl, C₁₋₄alkyl-amino-carbonyl-C₁₋₄alkyl, (C₁₋₄alkyl)₂-amino-

carbonyl-C₁₋₄alkyl, C₁₋₄alkyl-carbonyl-amino, C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkyl,

- hydroxyl-C₁₋₄alkyl, C₁₋₄alkyl-carbonyl, C₁₋₄alkoxy, halo-C₁₋₄alkoxy, amino-C₁₋₄alkoxy, hydroxyl-C₁₋₄alkoxy, C₁₋₄alkyl-C₁₋₄alkoxy, C₁₋₄alkyl-amino-C₁₋₄alkoxy, (C₁₋₄alkyl)₂-amino-C₁₋₄alkoxy, C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkoxy, C₁₋₄alkoxy-C₁₋₄alkoxy, C₁₋₄alkoxy-carbonyl, C₁₋₄alkoxy-carbonyl-amino, C₁₋₄alkoxy-carbonyl-amino-C₁₋₄alkoxy, C₂₋₄alkenyl, C₂₋₄alkenyl-amino-carbonyl, C₃₋₇cycloalkyl, C₃₋₇cycloalkyl-C₁₋₄alkoxy, C₃₋₇cycloalkenyl, heteroaryl, heteroaryl-C₁₋₄alkyl, heteroaryl-C₁₋₄alkyl-amino, heteroaryl-C₁₋₄alkyl-amino-carbonyl, heteroaryl-C₁₋₄alkyl-carbonyl-amino, heteroaryl-C₁₋₄alkyl-amino-carbonyl-C₁₋₄alkyl, heteroaryl-C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkyl, heterocyclyl, heterocyclyl-C₁₋₄alkyl, heterocyclyl-C₁₋₄alkoxy, phenyl, or phenyl-C₁₋₄alkoxy,
- wherein heteroaryl is a saturated monocyclic or bicyclic ring system having 1, 2, or 3 heteroatom ring members selected from N, O, and S,
- wherein heterocyclyl is a saturated or partially unsaturated monocyclic or bicyclic ring system having 1, 2, or 3 heteroatom ring members selected from N, O, and S, and
- wherein each instance of phenyl, heteroaryl or heterocyclyl is optionally substituted with 1, or 2 substituents each selected from R₃;
- R₂ is halogen, hydroxyl, cyano, oxo, hydroxyl-imino, C₁₋₄alkyl, halo-C₁₋₄alkyl, amino, C₁₋₄alkyl-amino, (C₁₋₄alkyl)₂-amino, amino-C₁₋₄alkyl, C₁₋₄alkyl-amino-C₁₋₄alkyl, (C₁₋₄alkyl)₂-amino-C₁₋₄alkyl, amino-carbonyl, hydroxyl-C₁₋₄alkyl, C₁₋₄alkoxy, C₁₋₄alkoxy-carbonyl, C₂₋₄alkenyl, C₃₋₇cycloalkyl, or heterocyclyl-C₁₋₄alkyl,
- wherein heterocyclyl is a saturated or partially unsaturated monocyclic or bicyclic ring system having 1, 2, or 3 heteroatom ring members selected from N, O, and S, and
- wherein each instance of heterocyclyl is optionally substituted with 1, or 2 substituents each selected from R₃;
- R₃ is halogen, hydroxyl, nitro, oxo, hydroxyl-imino, C₁₋₄alkyl, amino, C₁₋₄alkyl-amino, (C₁₋₄alkyl)₂-amino, amino-C₁₋₄alkyl, C₁₋₄alkyl-amino-C₁₋₄alkyl, (C₁₋₄alkyl)₂-amino-C₁₋₄alkyl, amino-carbonyl, C₁₋₄alkyl-amino-carbonyl, (C₁₋₄alkyl)₂-amino-carbonyl, C₁₋₄alkyl-amino-carbonyl-C₁₋₄alkyl, (C₁₋₄alkyl)₂-amino-carbonyl-C₁₋₄alkyl, C₁₋₄alkyl-carbonyl-amino, C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkyl, hydroxyl-C₁₋₄alkyl, C₁₋₄alkyl-carbonyl, C₁₋₄alkoxy, halo-C₁₋₄alkoxy, amino-C₁₋₄alkoxy, hydroxyl-C₁₋₄alkoxy, C₁₋₄alkyl-C₁₋₄alkoxy, C₁₋₄alkyl-amino-C₁₋₄alkoxy,

(C₁₋₄alkyl)₂-amino-C₁₋₄alkoxy, C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkoxy, C₁₋₄alkoxy-C₁₋₄alkoxy, C₁₋₄alkoxy-carbonyl, C₁₋₄alkoxy-carbonyl-amino, C₁₋₄alkoxy-carbonyl-amino-C₁₋₄alkoxy, C₂₋₄alkenyl, C₂₋₄alkenyl-amino-carbonyl, C₃₋₇cycloalkyl, C₃₋₇cycloalkyl-C₁₋₄alkoxy, C₃₋₇cycloalkenyl, heteroaryl, heteroaryl-C₁₋₄alkyl, heteroaryl-C₁₋₄alkyl-amino, heteroaryl-C₁₋₄alkyl-amino-carbonyl, heteroaryl-C₁₋₄alkyl-carbonyl-amino, heteroaryl-C₁₋₄alkyl-amino-carbonyl-C₁₋₄alkyl, heteroaryl-C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkyl, heterocyclyl, heterocyclyl-C₁₋₄alkyl, phenyl, or phenyl-C₁₋₄alkoxy;

- R₄ is independently selected from halogen, C₁₋₄alkyl, hydroxyl-C₁₋₄alkyl, amino, C₁₋₄alkyl-amino, (C₁₋₄alkyl)₂-amino or hydroxyl-C₁₋₄alkyl-amino; and
- R₅ is hydrogen, C₁₋₄alkyl, or hydroxyl-C₁₋₄alkyl;
- wherein a form of the compound is selected from the group consisting of a prodrug, salt, hydrate, solvate, clathrate, isotopologue, racemate, enantiomer, diastereomer, stereoisomer, polymorph and tautomer form thereof.
2. The method of claim 1, wherein the compound of Formula (I) is selected from a compound of Formula (Ia) and Formula (Ib):



- or a form thereof, wherein
- X is CH₂, CH(C₁₋₄alkyl), C(C₁₋₄alkyl)₂, CH=CH, O, NR₅, or a bond;
- A is aryl, heteroaryl, heterocyclyl, or C₉₋₁₀cycloalkyl,
- wherein aryl is selected from phenyl and naphthyl, each optionally substituted with 1, 2, 3, or 4 substituents each selected from R₁,
- wherein heteroaryl is a saturated monocyclic, bicyclic or tricyclic ring system having 1, 2, or 3 heteroatom ring members independently selected from N, O, or S, each optionally substituted with 1, 2, 3, 4, or 5 substituents each selected from R₁,
- wherein heterocyclyl is a saturated or partially unsaturated monocyclic, bicyclic or tricyclic ring system having 1, 2, or 3 heteroatom ring members independently selected from N, O, or S, each optionally substituted with 1, 2, 3, 4, or 5 substituents each selected from R₂, and

wherein C₉₋₁₀cycloalkyl is a saturated or partially unsaturated bicyclic ring system optionally substituted with 1, 2, 3, 4, or 5 substituents each selected from R₂;

B is heterocyclyl,

wherein heterocyclyl is a saturated or partially unsaturated monocyclic, bicyclic or polycyclic

5 ring system having 1, 2, or 3 heteroatom ring members independently selected from N, O, or S, each optionally substituted with 1, 2, 3, 4, or 5 substituents each selected from R₄;

R₁ is halogen, hydroxyl, cyano, C₁₋₄alkyl, halo-C₁₋₄alkyl, amino, C₁₋₄alkyl-amino,

(C₁₋₄alkyl)₂-amino, amino-C₁₋₄alkyl, C₁₋₄alkyl-amino-C₁₋₄alkyl,

(C₁₋₄alkyl)₂-amino-C₁₋₄alkyl, amino-carbonyl, C₁₋₄alkyl-amino-carbonyl,

10 C₁₋₄alkyl-amino-carbonyl-C₁₋₄alkyl, C₁₋₄alkyl-carbonyl-amino, C₁₋₄alkyl-carbonyl-amino-

C₁₋₄alkyl, hydroxyl-C₁₋₄alkyl, C₁₋₄alkyl-carbonyl, C₁₋₄alkoxy, halo-C₁₋₄alkoxy, amino-

C₁₋₄alkoxy, hydroxyl-C₁₋₄alkoxy, C₁₋₄alkyl-C₁₋₄alkoxy, C₁₋₄alkyl-amino-C₁₋₄alkoxy,

(C₁₋₄alkyl)₂-amino-C₁₋₄alkoxy, C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkoxy, C₁₋₄alkoxy-

C₁₋₄alkoxy, C₁₋₄alkoxy-carbonyl, C₁₋₄alkoxy-carbonyl-amino, C₁₋₄alkoxy-carbonyl-

15 amino-C₁₋₄alkoxy, C₂₋₄alkenyl, C₂₋₄alkenyl-amino-carbonyl, C₃₋₇cycloalkyl,

C₃₋₇cycloalkyl-C₁₋₄alkoxy, C₃₋₇cycloalkenyl, heteroaryl, heteroaryl-C₁₋₄alkyl,

heteroaryl-C₁₋₄alkyl-amino, heteroaryl-C₁₋₄alkyl-amino-carbonyl,

heteroaryl-C₁₋₄alkyl-carbonyl-amino, heteroaryl-C₁₋₄alkyl-amino-carbonyl-C₁₋₄alkyl,

heteroaryl-C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkyl, heterocyclyl, heterocyclyl-C₁₋₄alkyl,

20 heterocyclyl-C₁₋₄alkoxy, phenyl, or phenyl-C₁₋₄alkoxy,

wherein heteroaryl is a saturated monocyclic or bicyclic ring system having 1, 2, or 3 heteroatom ring members selected from N, O, and S,

wherein heterocyclyl is a saturated or partially unsaturated monocyclic or bicyclic ring system having 1, 2, or 3 heteroatom ring members selected from N, O, and S, and

25 wherein each instance of phenyl, heteroaryl or heterocyclyl is optionally substituted with 1, or 2 substituents each selected from R₃;

R₂ is halogen, hydroxyl, cyano, oxo, hydroxyl-imino, C₁₋₄alkyl, halo-C₁₋₄alkyl, amino,

C₁₋₄alkyl-amino, (C₁₋₄alkyl)₂-amino, amino-C₁₋₄alkyl, amino-carbonyl,

hydroxyl-C₁₋₄alkyl, C₁₋₄alkoxy, C₁₋₄alkoxy-carbonyl, C₂₋₄alkenyl, C₃₋₇cycloalkyl, or

30 heterocyclyl-C₁₋₄alkyl;


- R_3 is halogen, hydroxyl, nitro, oxo, hydroxyl-imino, C_{1-4} alkyl, amino, C_{1-4} alkyl-amino, $(C_{1-4}alkyl)_2$ -amino, amino- C_{1-4} alkyl, C_{1-4} alkyl-amino- C_{1-4} alkyl, $(C_{1-4}alkyl)_2$ -amino- C_{1-4} alkyl, amino-carbonyl, C_{1-4} alkyl-amino-carbonyl, C_{1-4} alkyl-amino-carbonyl- C_{1-4} alkyl, C_{1-4} alkyl-carbonyl-amino, C_{1-4} alkyl-carbonyl-amino- C_{1-4} alkyl, hydroxyl- C_{1-4} alkyl, C_{1-4} alkyl-carbonyl, C_{1-4} alkoxy, halo- C_{1-4} alkoxy, amino- C_{1-4} alkoxy, hydroxyl- C_{1-4} alkoxy, C_{1-4} alkyl- C_{1-4} alkoxy, C_{1-4} alkyl-amino- C_{1-4} alkoxy, $(C_{1-4}alkyl)_2$ -amino- C_{1-4} alkoxy, C_{1-4} alkyl-carbonyl-amino- C_{1-4} alkoxy, C_{1-4} alkoxy- C_{1-4} alkoxy, C_{1-4} alkoxy-carbonyl, C_{1-4} alkoxy-carbonyl-amino, C_{1-4} alkoxy-carbonyl-amino- C_{1-4} alkoxy, $C_{2-4}alkenyl$, $C_{2-4}alkenyl$ -amino-carbonyl, $C_{3-7}cycloalkyl$, $C_{3-7}cycloalkyl$ - C_{1-4} alkoxy, $C_{3-7}cycloalkenyl$, heteroaryl, heteroaryl- C_{1-4} alkyl, heteroaryl- C_{1-4} alkyl-amino, heteroaryl- C_{1-4} alkyl-amino-carbonyl, heteroaryl- C_{1-4} alkyl-carbonyl-amino, heteroaryl- C_{1-4} alkyl-amino-carbonyl- C_{1-4} alkyl, heteroaryl- C_{1-4} alkyl-carbonyl-amino- C_{1-4} alkyl, heterocyclyl, heterocyclyl- C_{1-4} alkyl, phenyl, or phenyl- C_{1-4} alkoxy;
- R_4 is independently selected from halogen, C_{1-4} alkyl, hydroxyl- C_{1-4} alkyl, amino, C_{1-4} alkyl-amino, $(C_{1-4}alkyl)_2$ -amino or hydroxyl- C_{1-4} alkyl-amino; and
- R_5 is hydrogen, C_{1-4} alkyl, or hydroxyl- C_{1-4} alkyl;
- wherein a form of the compound is selected from the group consisting of a prodrug, salt, hydrate, solvate, clathrate, isotopologue, racemate, enantiomer, diastereomer, stereoisomer, polymorph and tautomer form thereof.

3. The method of claim 1, wherein:


X is O, NH, $N(CH_3)$ or a bond;


A is aryl, heteroaryl or heterocyclyl,

wherein, aryl is selected from the group consisting of:




wherein, heteroaryl is selected from the group consisting of:







wherein, heterocyclyl is selected from the group consisting of:



B is heterocyclyl selected from the group consisting of:





R_{1a} and R_{1b} are each, where allowed by available valences, one or more substituents each selected from halogen, hydroxyl, cyano, C_{1-4} alkyl, halo- C_{1-4} alkyl, amino, C_{1-4} alkyl-amino, $(C_{1-4}alkyl)_2$ -amino, amino- C_{1-4} alkyl, C_{1-4} alkyl-amino- C_{1-4} alkyl, $(C_{1-4}alkyl)_2$ -amino- C_{1-4} alkyl, amino-carbonyl, C_{1-4} alkyl-amino-carbonyl, $(C_{1-4}alkyl)_2$ -amino-carbonyl, C_{1-4} alkyl-amino-carbonyl- C_{1-4} alkyl, $(C_{1-4}alkyl)_2$ -amino-carbonyl- C_{1-4} alkyl, C_{1-4} alkyl-carbonyl-amino, C_{1-4} alkyl-carbonyl-amino- C_{1-4} alkyl, hydroxyl- C_{1-4} alkyl, C_{1-4} alkyl-carbonyl, C_{1-4} alkoxy, halo- C_{1-4} alkoxy, amino- C_{1-4} alkoxy, hydroxyl- C_{1-4} alkoxy, C_{1-4} alkyl- C_{1-4} alkoxy, C_{1-4} alkyl-amino- C_{1-4} alkoxy, $(C_{1-4}alkyl)_2$ -amino- C_{1-4} alkoxy, C_{1-4} alkyl-carbonyl-amino- C_{1-4} alkoxy, C_{1-4} alkoxy- C_{1-4} alkoxy, C_{1-4} alkoxy-carbonyl, C_{1-4} alkoxy-carbonyl-amino, C_{1-4} alkoxy-carbonyl-amino- C_{1-4} alkoxy, C_{2-4} alkenyl, C_{2-4} alkenyl-amino-carbonyl, C_{3-7} cycloalkyl, C_{3-7} cycloalkyl- C_{1-4} alkoxy, C_{3-7} cycloalkenyl, heteroaryl, heteroaryl- C_{1-4} alkyl, heteroaryl- C_{1-4} alkyl-amino, heteroaryl- C_{1-4} alkyl-amino-carbonyl, heteroaryl- C_{1-4} alkyl-carbonyl-amino, heteroaryl- C_{1-4} alkyl-amino-carbonyl- C_{1-4} alkyl, heteroaryl- C_{1-4} alkyl-carbonyl-amino- C_{1-4} alkyl, heterocyclyl, heterocyclyl- C_{1-4} alkyl, heterocyclyl- C_{1-4} alkoxy, phenyl, or phenyl- C_{1-4} alkoxy,

wherein heteroaryl is a saturated monocyclic or bicyclic ring system having 1, 2, or 3 heteroatom ring members selected from N, O, and S,





wherein heterocyclyl is a saturated or partially unsaturated monocyclic or bicyclic ring system having 1, 2, or 3 heteroatom ring members selected from N, O, and S, and

wherein each instance of phenyl, heteroaryl or heterocyclyl is optionally substituted with 1, or 2 substituents each selected from R_3 ;

R_{2a} , R_{2b} and R_{2c} are each, where allowed by available valences, one or more substituents each selected from halogen, hydroxyl, cyano, oxo, hydroxyl-imino, C_{1-4} alkyl, halo- C_{1-4} alkyl, amino, C_{1-4} alkyl-amino, $(C_{1-4}alkyl)_2$ -amino, amino- C_{1-4} alkyl, C_{1-4} alkyl-amino- C_{1-4} alkyl,

- (C₁₋₄alkyl)₂-amino-C₁₋₄alkyl, amino-carbonyl, hydroxyl-C₁₋₄alkyl, C₁₋₄alkoxy, C₁₋₄alkoxy-carbonyl, C₂₋₄alkenyl, C₃₋₇cycloalkyl, or heterocyclyl-C₁₋₄alkyl, wherein heterocyclyl is a saturated or partially unsaturated monocyclic or bicyclic ring system having 1, 2, or 3 heteroatom ring members selected from N, O, and S, and
- 5 wherein each instance of heterocyclyl is optionally substituted with 1, or 2 substituents each selected from R₃;
- R₃ is halogen, hydroxyl, nitro, oxo, hydroxyl-imino, C₁₋₄alkyl, amino, C₁₋₄alkyl-amino, (C₁₋₄alkyl)₂-amino, amino-C₁₋₄alkyl, C₁₋₄alkyl-amino-C₁₋₄alkyl, (C₁₋₄alkyl)₂-amino-C₁₋₄alkyl, amino-carbonyl, C₁₋₄alkyl-amino-carbonyl,
- 10 (C₁₋₄alkyl)₂-amino-carbonyl, C₁₋₄alkyl-amino-carbonyl-C₁₋₄alkyl, (C₁₋₄alkyl)₂-amino-carbonyl-C₁₋₄alkyl, C₁₋₄alkyl-carbonyl-amino, C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkyl, hydroxyl-C₁₋₄alkyl, C₁₋₄alkyl-carbonyl, C₁₋₄alkoxy, halo-C₁₋₄alkoxy, amino-C₁₋₄alkoxy, hydroxyl-C₁₋₄alkoxy, C₁₋₄alkyl-C₁₋₄alkoxy, C₁₋₄alkyl-amino-C₁₋₄alkoxy, (C₁₋₄alkyl)₂-amino-C₁₋₄alkoxy, C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkoxy, C₁₋₄alkoxy-
- 15 C₁₋₄alkoxy, C₁₋₄alkoxy-carbonyl, C₁₋₄alkoxy-carbonyl-amino, C₁₋₄alkoxy-carbonyl-amino-C₁₋₄alkoxy, C₂₋₄alkenyl, C₂₋₄alkenyl-amino-carbonyl, C₃₋₇cycloalkyl, C₃₋₇cycloalkyl-C₁₋₄alkoxy, C₃₋₇cycloalkenyl, heteroaryl, heteroaryl-C₁₋₄alkyl, heteroaryl-C₁₋₄alkyl-amino, heteroaryl-C₁₋₄alkyl-amino-carbonyl, heteroaryl-C₁₋₄alkyl-carbonyl-amino, heteroaryl-C₁₋₄alkyl-amino-carbonyl-C₁₋₄alkyl,
- 20 heteroaryl-C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkyl, heterocyclyl, heterocyclyl-C₁₋₄alkyl, phenyl, or phenyl-C₁₋₄alkoxy; and
- R_{4a}, R_{4b}, R_{4c}, R_{4d}, R_{4e}, R_{4f} and R_{4g} are independently selected from halogen, C₁₋₄alkyl, hydroxyl-C₁₋₄alkyl, amino, C₁₋₄alkyl-amino, (C₁₋₄alkyl)₂-amino or hydroxyl-C₁₋₄alkyl-amino;
- 25 wherein a form of the compound is selected from the group consisting of a prodrug, salt, hydrate, solvate, clathrate, isotopologue, racemate, enantiomer, diastereomer, stereoisomer, polymorph and tautomer form thereof.

4. A compound selected from a compound of Formula (Ia11), Formula (Ia15), Formula (Ia18) or Formula (Ib1):



or a form thereof, wherein (when present),

X is selected from O, NR₅, or a bond;

- 5 A is selected from phenyl, thiophenyl, indazolyl, pyridinyl, pyrimidinyl or phenoxy, wherein phenyl and phenoxy are each optionally substituted with 1, 2 or 3 substituents each selected from R_{1a}, wherein thiophenyl, indazolyl, pyridinyl, pyrimidinyl are each optionally substituted with 1 or 2 substituents each selected from R_{1a},
- 10 B is selected from 1H-pyrazolyl, piperidinyl, 1,2,3,6-tetrahydropyridinyl, (1*R*,5*S*)-8-azabicyclo[3.2.1]octyl, 8-azabicyclo[3.2.1]oct-2-enyl, 2,6-diazaspiro[3.4]octyl or 2,7-diazaspiro[3.5]nonyl, each optionally substituted with 1 or 2 substituents each selected from R_{4a};
- R_{1a} is selected from halogen, hydroxyl, C₁₋₄alkyl, halo-C₁₋₄alkyl, amino, C₁₋₄alkoxy, or
- 15 heteroaryl,
- wherein heteroaryl is a saturated monocyclic or bicyclic ring system having 1, 2, or 3 heteroatom ring members selected from N, O, and S optionally substituted with 1 or 2 substituents each selected from R_{3a};
- R_{3a} is selected from nitro or C₁₋₄alkyl; and,
- 20 R_{4a} is C₁₋₄alkyl;
- R_{5a} is hydrogen, C₁₋₄alkyl, or hydroxyl-C₁₋₄alkyl;

wherein a form of the compound is selected from the group consisting of a prodrug, salt, hydrate, solvate, clathrate, isotopologue, racemate, enantiomer, diastereomer, stereoisomer, polymorph and tautomer form thereof.

5. The compound of claim 4 wherein (when present),
- 5 R_{1a} is selected from fluoro, chloro, hydroxyl, methyl, difluoromethyl, amino, methoxy or 1H-pyrazolyl or 1H-imidazol-1-yl, wherein 1H-pyrazolyl is optionally substituted with 1 or 2 substituents each selected from R_{3a}; R_{3a} is selected from nitro or methyl or amino; and, R_{4a} is methyl or ethyl;
- 10 R_{5a} is hydrogen or methyl;
- wherein a form of the compound is selected from the group consisting of a prodrug, salt, hydrate, solvate, clathrate, isotopologue, racemate, enantiomer, diastereomer, stereoisomer, polymorph and tautomer form thereof.
6. A method of use of a compound or a form thereof for treating or ameliorating HD in a
- 15 subject in need thereof, comprising administering to the subject an effective amount of a compound selected from the group consisting of:

6-(naphthalen-2-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
 6-(benzo[b]thiophen-2-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
 2-(6-(2,2,6,6-tetramethylpiperidin-4-yl-amino)-pyridazin-3-yl)phenol
 2-(6-(methyl-(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)benzo[b]-thiophene-5-carbonitrile
 6-(quinolin-3-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
 3-(benzo[b]-thiophen-2-yl)-6-(2,2,6,6-tetramethylpiperidin-4-yl-oxy)pyridazine
 2-(6-(methyl-(2,2,6,6-tetramethylpiperidin-4-yl)amino)-pyridazin-3-yl)phenol
 6-(6-(methyl-(2,2,6,6-tetramethylpiperidin-4-yl)amino)-pyridazin-3-yl)naphthalen-2-ol
 6-(benzo[b]-thiophen-2-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
 7-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)isoquinoline
 6-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)isoquinoline
 N-methyl-6-(quinolin-7-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
 N-methyl-6-(quinolin-6-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
 6-(isoquinolin-7-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
 6-(isoquinolin-6-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine

6-(imidazo[1,2-a]pyridin-6-yl-pyridazin-3-yl)-methyl-(2,2,6,6-tetramethylpiperidin-4-yl)-amine
N-methyl-6-(6-phenylpyridin-3-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
6-(6-(1H-pyrrol-1-yl)pyridin-3-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
6-(6-(1H-pyrazol-1-yl)pyridin-3-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
methyl-(6-quinoxalin-2-yl-pyridazin-3-yl)-(2,2,6,6-tetramethylpiperidin-4-yl)-amine
methyl-(6-quinolin-3-yl-pyridazin-3-yl)-(2,2,6,6-tetramethylpiperidin-4-yl)-amine
N-methyl-6-(phthalazin-6-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
6-(benzo[c][1,2,5]oxa-diazol-5-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
6-(benzo[d]thiazol-5-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
6-(2-methylbenzo-[d]oxazol-6-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalen-2-ol
5-chloro-2-(6-(methyl(1,2,2,6,6-pentamethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
3-(6-(2,2,6,6-tetramethylpiperidin-4-yl-amino)pyridazin-3-yl)naphthalen-2-ol
5-chloro-2-(6-(1,2,2,6,6-pentamethylpiperidin-4-ylamino)pyridazin-3-yl)phenol
4-hydroxy-3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)benzonitrile
3-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)naphthalen-2-ol
2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(trifluoromethyl)phenol
2-fluoro-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)-amino)-pyridazin-3-yl)phenol
3,5-dimethoxy-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
4,5-dimethoxy-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
5-methoxy-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
4,5-difluoro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
5-fluoro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
3-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)benzonitrile
1-allyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalen-2-ol
6-(benzo[b]thiophen-2-yl)-N-(1,2,2,6,6-pentamethylpiperidin-4-yl)pyridazin-3-amine
N-allyl-3-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)benzamide
2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol

5-(5-methyl-oxazol-2-yl)-2-(6-(methyl-(2,2,6,6-tetramethylpiperidin-4-yl)-amino)-pyridazin-3-yl)phenol
5-(4-hydroxymethyl)-1H-pyrazole-1-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
5-(1H-imidazol-1-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
5-(4-amino-1H-pyrazole-1-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1H-pyrazol-4-yl)phenol
5-(3-amino-1H-pyrazol-1-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1-(2-morpholinoethyl)-1H-pyrazol-4-yl)phenol
2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1-methyl-1H-pyrazol-4-yl)phenol
5-(5-amino-1H-pyrazol-1-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(1H-pyrazol-1-yl)phenol
2-(((6-((2-hydroxy-ethyl)-(2,2,6,6-tetramethylpiperidin-4-yl)-amino)-pyridazin-3-yl)-5-pyrazol-1-yl)phenol
2-(6-(piperidin-4-yloxy)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
2-(6-(((2S,4R,6R)-2,6-dimethylpiperidin-4-yl)oxy)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
2-(6-((-2,6-dimethylpiperidin-4-yl)oxy)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
5-(1H-pyrazol-1-yl)-2-(6-(pyrrolidin-3-yl-oxy)pyridazin-3-yl)phenol
2-(6-(((2S,4S)-2-methylpiperidin-4-yl)oxy)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
(5-(1H-pyrazol-1-yl)-2-(6-(pyrrolidin-3-ylmethoxy)pyridazin-3-yl)phenol
2-(6-((3-fluoropiperidin-4-yl)oxy)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
2-(6-(1,2,2,6,6-pentamethyl-piperidin-4-yl-oxy)-pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
5-1H-pyrazol-1-yl)-2-(6-(2,2,6,6-tetramethylpiperidin-4-yl-oxy)-pyridazin-3-yl)phenol
5-(1H-pyrazol-4-yl)-2-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)phenol
2-(6-piperazin-1-yl-pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
3-(6-(azetidin-3-ylamino)-pyridazin-3-yl)naphthalen-2-ol
2-(6-(azetidin-3-ylamino)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
2-(6-(3,5-dimethylpiperazin-1-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
2-(6-(7-methyl-2,7-diazaspiro[4.4]nonan-2-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
2-(6-(1,4-diazepan-1-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol

2-(6-(4-(2-hydroxyethyl)piperazin-1-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
2-(6-(3,6-diazabicyclo[3.2.1]octan-3-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
2-(6-(2,7-diazaspiro[3.5]nonan-7-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
2-(6-(3-(hydroxymethyl)piperazin-1-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
2-(6-(1,7-diazaspiro[4.4]nonan-7-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
2-(6-(4-amino-4-methylpiperidin-1-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
2-(6-(3-(dimethylamino)piperidin-1-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
2-(6-(1,2,2,6,6-pentamethylpiperidin-4-ylamino)-pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
2-(6-(3,3-dimethylpiperazin-1-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
2-(6-(7-(2-hydroxyethyl)-2,7-diazaspiro[4.4]nonan-2-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
2-(6-((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
3-(6-(piperazin-1-yl)pyridazin-3-yl)naphthalene-2,7-diol
5-(1H-pyrazol-1-yl)-2-(6-(1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl)phenol
2-(6-piperidin-4-yl-pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
3-(6-(1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl)naphthalene-2-ol
3-(6-(1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl)naphthalene-2,7-diol
3-(6-(2,2,6,6-tetramethyl-1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl)naphthalene-2,7-diol
3-(6-(1-methyl-1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl)naphthalene-2,7-diol
3-(6-(piperidin-4-yl)pyridazin-3-yl)naphthalene-2,7-diol
3-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)naphthalene-2,7-diol
3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalene-2,7-diol
3-(6-((2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalene-2,7-diol
tert-butyl (3-((7-hydroxy-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalen-2-yl)oxy)propyl)carbamate
7-(3-amino-propoxy)-3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)-amino)-pyridazin-3-yl)naphthalen-2-ol
N-(3-((7-hydroxy-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalen-2-yl)oxy)propyl)acetamide
7-(3-hydroxypropoxy)-3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalen-2-ol
7-(3-methoxypropoxy)-3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalen-2-ol
7-(2-morpholinoethoxy)-3-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)naphthalen-2-ol
3-(6-(piperidin-4-ylmethyl)pyridazin-3-yl)naphthalen-2-ol

5-(1H-pyrazol-1-yl)-2-(6-((2,2,6,6-tetramethylpiperidin-4-yl)methyl)pyridazin-3-yl)phenol
3-methoxy-2-(6-(methyl(2,2,6-trimethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(5-methyloxazol-2-yl)phenol
2-(6-((6S)-6-((S)-1-hydroxyethyl)-2,2-dimethylpiperidin-4-yl)oxy)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
7-hydroxy-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-2-naphthonitrile
3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-7-(piperidinylmethyl)naphthalen-2-ol
3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-7-(pyrrolidinylmethyl)naphthalen-2-ol
1-bromo-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalene-2,7-diol
1-chloro-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalene-2,7-diol
7-methoxy-3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalene-2-ol
7-methoxy-3-(6-(methyl(1,2,2,6,6-pentamethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalen-2-ol
7-(3,6-dihydro-2H-pyran-4-yl)-3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalen-2-ol
3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-7-(tetrahydro-2H-pyran-4-yl)naphthalene-2-ol
7-(difluoromethyl)-3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalen-2-ol
7-((4-hydroxy-2-methylbutan-2-yl)oxy)-3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalen-2-ol
7-(3-hydroxy-3-methylbutoxy)-3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalen-2-ol
2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1H-pyrazol-4-yl)benzene-1,3-diol
3-methoxy-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1H-pyrazol-4-yl)phenol
5-(1H-pyrazol-4-yl)-2-(6-((2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-3-(trifluoromethoxy)phenol
2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1-methyl-1H-pyrazol-4-yl)-3-(trifluoromethoxy)phenol
2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1H-pyrazol-4-yl)-3-(trifluoromethoxy)phenol

4-(3-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(trifluoromethoxy)phenyl)-1-methylpyridin-2(1H)-one

3-methoxy-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1-methyl-1H-pyrazol-4-yl)phenol

3-methoxy-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(5,6,7,8-tetrahydroimidazo[1,2-a]pyridin-3-yl)phenol

3-methoxy-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(pyridine-3-yl)phenol

5-(1-cyclopentyl-1H-pyrazol-4-yl)-3-methoxy-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol

3',5-dimethoxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-(1,1'-biphenyl)-3-ol

3-(benzyloxy)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(5-methyloxazol-2-yl)phenol

3-ethoxy-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(5-methyloxazol-2-yl)phenol

3-(cyclopropylmethoxy)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-pyridazin-3-yl)-5-(5-methyloxazol-2-yl)phenol

2-methyl-5-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-1H-benzo[d]imidazol-6-ol

5-chloro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol

5-(1H-pyrazol-1-yl)-2-(6-((2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol

3-hydroxy-4-(6-((2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)benzonitrile

2-(6-((2,2-dimethylpiperidin-4-yl)oxy)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol

2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(1H-pyrazol-4-yl)phenol

2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridin-3-yl)-4-(4,5,6,7-tetrahydropyrazolo[1,5-a]pyridin-3-yl)phenol

2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-3-yl)phenol

4-(1H-indol-2-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol

4-(cyclopent-1-en-1-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol

2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(1H-pyrazol-3-yl)phenol

4-(4-hydroxy-3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)pyridin-2-ol

4-(4-hydroxy-3-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)phenyl)-1-methylpyridin-2(1H)-one

4-(4-hydroxy-3-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)phenyl)pyridin-2-ol

5-(1H-indazol-7-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol

4-chloro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1H-pyrazol-4-yl)phenol

4-fluoro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1H-pyrazol-4-yl)phenol

5-fluoro-4-(1H-imidazol-4-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol

5-fluoro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(1H-pyrazol-4-yl)phenol

5-fluoro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(1H-pyrazol-5-yl)phenol

6-hydroxy-5-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-2,3-dihydro-1H-inden-1-one

6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-1,4-dihydroindeno[1,2-c]-1H-pyrazol-7-ol

6-hydroxy-5-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-2,3-dihydro-1H-inden-1-one oxime

5-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-2,3-dihydro-1H-indene-1,6-diol

2-amino-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-8H-indeno[1,2-d]thiazol-5-ol

9-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5,6-dihydroimidazo[5,1-a]isoquinolin-8-ol

4-hydroxy-3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-N-((1-methyl-1H-pyrazol-4-yl)methyl)benzamide

4-(4-(hydroxymethyl)-1H-pyrazol-1-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol

5-(1H-pyrazol-4-yl)-2-(6-((2,2,6,6-tetramethylpiperidin-4-yl)methyl)pyridazin-3-yl)phenol

6-(3-(benzyloxy)isoquinolin-6-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine

6-(1-(benzyloxy)isoquinolin-7-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine

3-fluoro-5-(2-methoxypyridin-4-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol

4-(3-fluoro-5-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)pyridin-2(1H)-one

4-(3-fluoro-5-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)-1-methylpyridin-2(1H)-one

5-(3-fluoro-5-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)-1-methylpyridin-2(1H)-one

3-fluoro-5-(1H-pyrazol-4-yl)-2-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)phenol

5-chloro-3-fluoro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol

3-fluoro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1H-pyrazol-4-yl)phenol

3-fluoro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1-methyl-1H-pyrazol-4-yl)phenol

5-(5-methoxypyridin-3-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol

5-(3-hydroxy-4-(6-methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)pyridin-2-ol

4-(3-hydroxy-4-(6-methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)pyridin-2-ol

5-(6-methoxypyridin-3-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol

5-(3-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)-3-(trifluoromethyl)pyridin-2-ol

5-(3-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)-1-methylpyridin-2(1H)-one

4-(3-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)-1-methylpyridin-2(1H)-one

5-(2-methoxypyridin-4-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol

4-(3-hydroxy-4-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)phenyl)pyridin-2-ol

5-(6-(dimethylamino)pyridin-3-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol

4-(3-hydroxy-4-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)phenyl)-1-methylpyridin-2(1H)-one

2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(pyrimidin-5-yl)phenol

5-(3-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)pyridin-3-ol

1-cyclopropyl-4-(3-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)pyridin-2(1H)-one

2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1,2,3,6-tetrahydropyridin-4-yl)phenol

5-(cyclopent-1-en-1-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol

5-(3,6-dihydro-2H-pyran-4-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol

5-(imidazo[1,5-a]pyridin-7-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol

5-(imidazo[1,2-a]pyridin-7-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol

2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(2-methylpyridin-4-yl)phenol

5-(1H-imidazol-2-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol

5-(1H-imidazol-4-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol

5-(imidazo[1,2-a]pyrazin-3-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol

2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(5,6,7,8-tetrahydroimidazo[1,2-a]pyrazin-3-yl)phenol

2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(4-methyl-1H-imidazol-2-yl)phenol

2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1-methyl-1H-imidazol-4-yl)phenol

2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1-methyl-1H-imidazol-5-yl)phenol

2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(4-nitro-1H-imidazol-2-yl)phenol

2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(2-methyl-1H-imidazol-4-yl)phenol

5-(1,2-dimethyl-1H-imidazol-4-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol

1-(3-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)-1H-pyrazole-4-carboxamide

2-(6-((3aR,6aS)-5-(2-hydroxyethyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)pyridazin-3-yl)-5-(1H-pyrazol-4-yl)phenol

2-(6-((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)pyridazin-3-yl)-5-(1H-pyrazol-4-yl)phenol

2-(6-((3aR,6aS)-5-methylhexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)pyridazin-3-yl)-5-(1H-pyrazol-4-yl)phenol

4-(3-hydroxy-4-(6-(5-methylhexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)pyridazin-3-yl)phenyl)-1-methylpyridin-2(1H)-one
4-(3-hydroxy-4-(6-((3aR,6aR)-1-methylhexahydropyrrolo[3,4-b]pyrrol-5(1H)-yl)pyridazin-3-yl)phenyl)-1-methylpyridin-2(1H)-one
2-(6-(2,7-diazaspiro[4.5]decan-2-yl)pyridazin-3-yl)-5-(1H-pyrazol-4-yl)phenol
4-(4-(6-(2,7-diazaspiro[4.5]decan-2-yl)pyridazin-3-yl)-3-hydroxyphenyl)-1-methylpyridin-2(1H)-one
2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)-amino)-pyridazin-3-yl)phenol
6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
6-(6-(methyl(1,2,2,6,6-pentamethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
6-(6-((3aR,6aS)-5-methylhexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)pyridazin-3-yl)quinolin-7-ol
2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
7-(6-(methyl(1,2,2,6,6-pentamethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-6-ol
7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-6-ol
7-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)isoquinoline-6-ol
7-(6-((3aR,6aS)-5-methylhexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)pyridazin-3-yl)isoquinolin-6-ol
1-cyclopropyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-6-ol
7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinoline-1,6-diol
6-hydroxy-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinoline-1-carbonitrile
6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-7-ol
8-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-6-ol
2-methyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-6-ol
3-chloro-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
3-bromo-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
7-hydroxy-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinoline-3-carbonitrile
6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-3-(1-methyl-1H-imidazol-4-yl)quinolin-7-ol
3-(1H-imidazol-1-yl)-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinoline-3,7-diol
3-ethyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol

3-isopropyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol

7-hydroxy-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-2(1H)-one

7-hydroxy-1-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-2(1H)-one

4-methoxy-2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol

2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(pyrrolidin-1-yl)quinolin-7-ol

2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-morpholinoquinolin-7-ol

4-(dimethylamino)-2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol

4-ethoxy-2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol

2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(1-methyl-1H-pyrazol-4-yl)quinolin-7-ol

4-methoxy-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-6-ol

7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinoxalin-6-ol

6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-3-(tetrahydro-2H-pyran-4-yl)quinolin-7-ol

3-chloro-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-6-ol

3-bromo-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-6-ol

3-methyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-6-ol

5-bromo-3-methyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-6-ol

6-hydroxy-1-methyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-4(1H)-one

2,3-dimethyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-6-ol

2-methyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinoxalin-6-ol

3-methyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinoxalin-6-ol

4-methoxy-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol

4-(azetidin-1-yl)-2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
7-hydroxy-2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolone-4-carbonitrile
4-cyclopropyl-2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-quinolin-7-ol
4-(3,6-dihydro-2H-pyran-4-yl)-2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(tetrahydro-2H-pyran-4-yl)quinolin-7-ol
2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(oxetan-3-yl)quinolin-7-ol
4-(dimethylamino)-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-quinolin-7-ol
7-hydroxy-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinazolin-4(1H)-one
6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinazolin-7-ol
7-hydroxy-1-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-3,4-dihydroquinolin-2(1H)-one
2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinazolin-7-ol
7-hydroxy-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinoline-1-carbonitrile
7-hydroxy-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinoline-2-carbonitrile
6-hydroxy-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-2-carbonitrile
6-hydroxy-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinoline-1-carboxamide
7-hydroxy-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-2-carboxamide
6-hydroxy-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinoline-2-carboxamide
methyl 6-hydroxy-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinoline-2-carboxylate
6-hydroxy-7-(6-(piperazin-1-yl)pyridazin-3-yl)quinoline-2-carbonitrile
7-hydroxy-6-(6-(piperazin-1-yl)pyridazin-3-yl)quinoline-2-carbonitrile
7-(6-(piperazin-1-yl)pyridazin-3-yl)isoquinolin-6-ol
7-(6-(1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl)quinolin-6-ol

1-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-7-ol

1-methyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-6-ol

1,3-dimethyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-6-ol

7-hydroxy-3-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinoline-1-carbonitrile

1-amino-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-6-ol

7-hydroxy-1,3-dimethyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinazoline-2,4(1H,3H)-dione

6-hydroxy-5-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)benzo[d]oxazol-2(3H)-one

2-methyl-5-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-2H-indazol-6-ol

1-methyl-5-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-1H-indazol-6-ol

6-hydroxy-2-methyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-1(2H)-one

2-ethyl-6-hydroxy-7-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)isoquinolin-1(2H)-one

1-ethoxy-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-6-ol

7-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)isoquinoline-1,6-diol

7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-pyridazin-3-yl)-3-phenylisoquinolin-6-ol

3-methyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-6-ol

3-cyclopropyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-6-ol

3-isopropyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-6-ol

3-propyl-7-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)-pyridazin-3-yl)isoquinolin-6-ol

3-isopropyl-7-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)-pyridazin-3-yl)isoquinolin-6-ol

3-methyl-7-(6-(piperazin-1-yl)pyridazin-3-yl)isoquinolin-6-ol

6-(3-(benzyloxy)isoquinolin-6-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine

3-chloro-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol

3-isopropyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol

3-methyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinoxalin-6-ol

4-chloro-2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol

4-chloro-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-6-ol

7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-6-ol

5-(2-methoxy-4-(1H-pyrazol-1-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

6-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)naphthalen-2-ol

5-(2-methoxyquinolin-3-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

5-(3-methoxy-naphthalen-2-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

5-(2-methoxy-4-(1H-pyrazol-1-yl)phenyl)-N-(1,2,2,6,6-pentamethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

5-(2-methoxy-4-(1-methyl-1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

5-(2-methoxy-4-(1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

4-(3-methoxy-4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)phenyl)-1-methylpyridin-2(1H)-one

5-(3-methoxy-4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)phenyl)pyridin-2-ol

5-(3-methoxy-4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)phenyl)-1-methylpyridin-2(1H)-one

N-methyl-5-(2-methyl-4-(1-methyl-1H-pyrazol-4-yl)phenyl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

1-methyl-4-(4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-3-(trifluoromethoxy)phenyl)pyridin-2(1H)-one

5-(4-(3,5-dimethyl-1H-pyrazol-4-yl)-2-methoxyphenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

5-(2-methoxy-4-(1-methyl-1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-5-(1-methyl-1H-pyrazol-4-yl)phenol

2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-5-(1H-pyrazol-1-yl)phenol

5-(3-hydroxy-4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)phenyl)-1-methylpyridin-2(1H)-one

4-(3-hydroxy-4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)phenyl)-1-methylpyridin-2(1H)-one

5-(3-hydroxy-4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)phenyl)pyridin-2-ol

3-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)naphthalene-2,7-diol

3-(5-((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-1,3,4-thiadiazol-2-yl)naphthalene-2,7-diol

3-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)naphthalen-2-ol

3-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)quinolin-2-ol

2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-4-(1H-pyrazol-1-yl)phenol

5-(2-chloro-4-(1-methyl-1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

3-chloro-2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-5-(1-methyl-1H-pyrazol-4-yl)phenol

5-(2-chloro-4-(1-methyl-1H-pyrazol-4-yl)phenyl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

3-methoxy-2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-5-methyloxazol-2-yl)phenol

2-(2-methoxy-4-(1H-pyrazol-1-yl)phenyl)-5-(1,2,3,6-tetrahydropyridin-4-yl)-1,3,4-thiadiazole

2-(5-(piperazin-1-yl)-1,3,4-thiadiazol-2-yl)-5-(1H-pyrazol-1-yl)phenol

5-(7-methoxyquinolin-6-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazole-2-amine

6-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)quinolin-7-ol

3-methoxy-4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)benzonitrile

3-fluoro-4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)benzonitrile

methyl-3-fluoro-4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)benzoate

5-(2-methoxy-4-(3-(methylamino)-1H-pyrazol-1-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

7-methoxy-6-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)quinoline-2-carbonitrile

4-(3-methoxy-4-(5-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)-1,3,4-thiadiazol-2-yl)phenyl)-1-methylpyridin-2(1H)-one

4-(3-chloro-4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)phenyl)-1-methylpyridin-2(1H)-one

5-(2-chloro-4-(1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

5-(2-chloro-4-(4,5,6,7-tetrahydropyrazolo[1,5-a]pyridin-3-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

N-methyl-5-(5-(1-methyl-1H-pyrazol-4-yl)pyridin-2-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

2-(2-chloro-4-(1-methyl-1H-pyrazol-4-yl)phenyl)-5-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)-1,3,4-thiadiazole

5-(2-chloro-4-(6-methoxypyridin-3-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

5-(4-(6-aminopyridin-3-yl)-2-fluorophenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

5-(2-fluoro-4-(3-methyl-1H-pyrazol-5-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

5-(2-fluoro-4-(1H-pyrazol-5-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

5-(2,3-difluoro-4-(1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

5-(2,3-difluoro-4-(1H-pyrazol-5-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

5-(2,5-difluoro-4-(1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

5-(2,5-difluoro-4-(1H-pyrazol-5-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

5-(2,6-difluoro-4-(1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

2-(2,5-difluoro-4-(1H-pyrazol-4-yl)phenyl)-5-((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-1,3,4-thiadiazole

5-(2-chloro-5-fluoro-4-(1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

5-(3-fluoro-5-(1H-pyrazol-4-yl)pyridin-2-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

5-(4-(2-aminopyrimidin-4-yl)-2-chlorophenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

5-(5-(2-aminopyrimidin-4-yl)-2-chlorophenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

5-(4-(2,4-dimethylthiazol-5-yl)-2,5-difluorophenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

5-(4-(2,4-dimethylthiazol-5-yl)-2,3-difluorophenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

4-(3-hydroxy-4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-5-(trifluoromethoxy)phenyl)-1-methylpyridin-2(1H)-one

5-(2-fluoro-6-methoxy-4-(1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

2-(2-fluoro-6-methoxy-4-(1H-pyrazol-4-yl)phenyl)-5-((3aR,6aS)-5-methylhexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-1,3,4-thiadiazole

5-(2,3-difluoro-6-methoxy-4-(1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

6-methoxy-2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-3,4-dihydroisoquinolin-1-(2H)-one

5-(2-chloro-4-(1H-pyrazol-1-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

5-(2-chloro-4-(1H-1,2,3-triazol-1-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

5-(2-chloro-4-(2H-1,2,3-triazol-2-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

5-(2-chloro-4-(1H-1,2,4-triazol-1-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

5-(4-(3-amino-1H-pyrazol-1-yl)-2-chlorophenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

2-(2-chloro-4-(1H-imidazol-1-yl)phenyl)-5-((3aR,6aS)-5-methylhexahydropyrrolo[3,4c]pyrrol-2(1H)-yl)-1,3,4-thiadiazole

5-(2-chloro-4-(1H-imidazol-1-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

5-(2-fluoro-4-(1H-imidazol-1-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

5-(2-methoxy-4-(1H-pyrazol-5-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

5-(4-(2,4-dimethylthiazol-5-yl)-2-methoxyphenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

5-(2-methoxy-4-(pyridin-3-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

5-(2-fluoro-4-(1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

5-(2-methoxy-4-(2-methoxypyridin-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

5-(2-methoxy-4-(6-methoxypyridin-3-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

2-(2-chloro-4-(1-methyl-1H-pyrazol-4-yl)phenyl)-5-((3aR,6aS)-5-methylhexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-1,3,4-thiadiazole

2-(2-chloro-4-(1H-pyrazol-4-yl)phenyl)-5-((3aR,6aS)-5-methylhexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-1,3,4-thiadiazole

2-(2-chloro-4-(1H-pyrazol-4-yl)phenyl)-5-((3aR,6aR)-1-methylhexahydropyrrolo[3,4-b]pyrrol-5(1H)-yl)-1,3,4-thiadiazole

1-(4-(5-(2-chloro-4-(1H-pyrazol-4-yl)phenyl)-1,3,4-thiadiazol-2-yl)morpholin-2-yl)-N,N-dimethylmethanamine

2-(2-chloro-4-(1H-pyrazol-4-yl)phenyl)-5-(2-methyl-2,7-diazaspiro[4.5]decan-7-yl)-1,3,4-thiadiazole

2-(2-fluoro-4-(1H-pyrazol-4-yl)phenyl)-5-((3aR,6aS)-5-methylhexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-1,3,4-thiadiazole

2-(2-methoxy-4-(1-methyl-1H-pyrazol-4-yl)phenyl)-5-(2,6-diazaspiro[3.5]nonan-2-yl)-1,3,4-thiadiazole

2-(2-methoxy-4-(1-methyl-1H-pyrazol-4-yl)phenyl)-5-(2,7-diazaspiro[3.5]nonan-2-yl)-1,3,4-thiadiazole

2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-5-(1H-pyrazol-1-yl)phenol

5-(3-chloro-4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)phenyl)pyridin-2(1H)-one

2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-5-(3-(methylamino)-1H-pyrazol-1-yl)phenol

3-fluoro-2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-5-(1H-pyrazol-4-yl)phenol

3,4-difluoro-2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-5-(1H-pyrazol-4-yl)phenol

6-hydroxy-5-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-2,3-dihydro-1H-inden-1-one

2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-5-(1H-pyrazol-4-yl)phenol

2-(5-(2,6-diazaspiro[3.5]nonan-2-yl)-1,3,4-thiadiazol-2-yl)-5-(1-methyl-1H-pyrazol-4-yl)phenol

2-(5-(2,7-diazaspiro[3.5]nonan-2-yl)-1,3,4-thiadiazol-2-yl)-5-(1-methyl-1H-pyrazol-4-yl)phenol

3-fluoro-2-(5-((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-1,3,4-thiadiazol-2-yl)-5-(1H-pyrazol-4-yl)phenol

3-chloro-2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-5-(1H-pyrazol-4-yl)phenol

2-(2-methoxy-4-(1H-pyrazol-1-yl)phenyl)-5-((2,2,6,6-tetramethylpiperidin-4-yl)methyl)-1,3,4-thiadiazole

2-(2,3-difluoro-4-(1H-pyrazol-4-yl)phenyl)-5-(2,7-diazaspiro[3.5]nonan-2-yl)-1,3,4-thiadiazole

2-(5-(2,7-diazaspiro[3.5]nonan-2-yl)-1,3,4-thiadiazol-2-yl)-3-fluoro-5-(1H-pyrazol-4-yl)phenol

4-methoxy-1-methyl-3-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)quinolin-2(1H)-one

4-hydroxy-1-methyl-3-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)quinolin-2(1H)-one

3-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)quinolin-2(1H)-one

1-methyl-3-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)quinolin-2(1H)-one

2-(2-chloro-4-(1H-pyrazol-4-yl)phenyl)-5-((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-1,3,4-thiadiazole

2-(2-chloro-4-(1H-pyrazol-4-yl)phenyl)-5-(2,7-diazaspiro[4.5]decan-2-yl)-1,3,4-thiadiazole

(R)-(4-(5-(2-chloro-4-(1H-pyrazol-4-yl)phenyl)-1,3,4-thiadiazol-2-yl)piperazin-2-yl)methanol

2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)benzo[b]thiophene-5-carbonitrile

5-(3-chlorobenzo[b]thiophen-2-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

5-(2-methoxy-4-(1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

2-{6-[8-azabicyclo[3.2.1]oct-3-yl(methyl)amino]pyridazin-3-yl}-5-(1H-pyrazol-4-yl)phenol

2-[6-(8-azabicyclo[3.2.1]oct-3-ylamino)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol

5-(1H-pyrazol-4-yl)-2-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl}phenol

5-(1-methyl-1H-pyrazol-4-yl)-2-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}phenol

2-[6-(8-azabicyclo[3.2.1]oct-3-yloxy)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol

5-(5-methyl-1H-pyrazol-4-yl)-2-{6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl}phenol

5-(1H-imidazol-1-yl)-2-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}phenol

5-(5-methyl-1H-pyrazol-4-yl)-2-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}phenol

2-{6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl}-5-(4-nitro-1H-pyrazol-1-yl)phenol

6-[2-methoxy-4-(4-nitro-1H-pyrazol-1-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine

5-(4-amino-1H-pyrazol-1-yl)-2-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}phenol
2-[6-(1-methyl-1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
5-(4-nitro-1H-pyrazol-1-yl)-2-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}phenol
5-(1H-pyrazol-4-yl)-2-[6-(1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]phenol
2-[6-(1-ethyl-1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
2-{6-[methyl(piperidin-4-yl)amino]pyridazin-3-yl}-5-(1H-pyrazol-4-yl)phenol
2-[6-(piperidin-4-ylamino)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
6-[2,5-difluoro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
2-[6-(8-azabicyclo[3.2.1]oct-2-en-3-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
6-[2,3-difluoro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
3-[2,5-difluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
2-[6-(piperidin-4-yloxy)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
2-{6-[(1R,5S)-8-azabicyclo[3.2.1]oct-3-ylamino]pyridazin-3-yl}-5-(1H-pyrazol-4-yl)phenol
6-[2-methoxy-6-(1H-pyrazol-4-yl)pyridin-3-yl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
3-[4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
2-{6-[(2,6-dimethylpiperidin-4-yl)oxy]pyridazin-3-yl}-5-(1H-pyrazol-4-yl)phenol
3-[2-fluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
3-[4-(1-methyl-1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
2-[6-(2,7-diazaspiro[3.5]non-2-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
3-fluoro-4-{6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl}phenol
2-{6-[(2,6-dimethylpiperidin-4-yl)oxy]pyridazin-3-yl}-5-(1H-pyrazol-1-yl)phenol
N-methyl-6-(2-methyl-2H-indazol-5-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
2-methyl-5-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}-2H-indazole
3-(4-chloro-2-methoxyphenyl)-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
N-methyl-6-(2-methylpyrazolo[1,5-a]pyridin-3-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
6-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}imidazo[1,2-a]pyridine
3-[2-methoxy-4-(1H-pyrazol-1-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine

3-[5-(1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
 3-[5-(1-methyl-1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
 3-[4-(1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
 5-(3,5-dimethyl-1H-pyrazol-4-yl)-2-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}phenol
 6-[2-fluoro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
 3-methoxy-4-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}phenol
 3-[2-methoxy-4-(4-nitro-1H-pyrazol-1-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
 4-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}benzene-1,3-diol
 6-[2-chloro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
 2-(1H-pyrazol-4-yl)-4-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}pyrimidin-5-amine
 3-[2,6-difluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
 2-[6-(2,6-diazaspiro[3.4]oct-2-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
 3-{6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl}-6-(1H-pyrazol-4-yl)pyridin-2-ol
 6-(1H-pyrazol-4-yl)-3-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}pyridin-2-ol
 N,2,2,6,6-pentamethyl-N-{5-[3-(1H-pyrazol-4-yl)phenoxy]-1,3,4-thiadiazol-2-yl}piperidin-4-amine
 N,2,2,6,6-pentamethyl-N-{5-[4-(1H-pyrazol-4-yl)phenoxy]-1,3,4-thiadiazol-2-yl}piperidin-4-amine
 3-[2-(difluoromethyl)-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine and
 6-[2-(difluoromethyl)-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine;

wherein a form of the compound is selected from the group consisting of a prodrug, salt, hydrate, solvate, clathrate, isotopologue, racemate, enantiomer, diastereomer, stereoisomer, polymorph and tautomer form thereof.

7. A compound selected from the group consisting of:

2-{6-[8-azabicyclo[3.2.1]oct-3-yl(methyl)amino]pyridazin-3-yl}-5-(1H-pyrazol-4-yl)phenol
 2-[6-(8-azabicyclo[3.2.1]oct-3-ylamino)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
 5-(1H-pyrazol-4-yl)-2-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl}phenol

5-(1-methyl-1H-pyrazol-4-yl)-2-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}phenol
2-[6-(8-azabicyclo[3.2.1]oct-3-yloxy)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
5-(5-methyl-1H-pyrazol-4-yl)-2-{6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl}phenol
5-(1H-imidazol-1-yl)-2-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}phenol
5-(5-methyl-1H-pyrazol-4-yl)-2-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}phenol
2-{6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl}-5-(4-nitro-1H-pyrazol-1-yl)phenol
6-[2-methoxy-4-(4-nitro-1H-pyrazol-1-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
5-(4-amino-1H-pyrazol-1-yl)-2-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}phenol
2-[6-(1-methyl-1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
5-(4-nitro-1H-pyrazol-1-yl)-2-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}phenol
5-(1H-pyrazol-4-yl)-2-[6-(1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]phenol
2-[6-(1-ethyl-1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
2-{6-[methyl(piperidin-4-yl)amino]pyridazin-3-yl}-5-(1H-pyrazol-4-yl)phenol
2-[6-(piperidin-4-ylamino)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
6-[2,5-difluoro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
2-[6-(8-azabicyclo[3.2.1]oct-2-en-3-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
6-[2,3-difluoro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
3-[2,5-difluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
2-[6-(piperidin-4-yloxy)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
2-{6-[(1R,5S)-8-azabicyclo[3.2.1]oct-3-ylamino]pyridazin-3-yl}-5-(1H-pyrazol-4-yl)phenol
6-[2-methoxy-6-(1H-pyrazol-4-yl)pyridin-3-yl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
3-[4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
2-{6-[(2,6-dimethylpiperidin-4-yl)oxy]pyridazin-3-yl}-5-(1H-pyrazol-4-yl)phenol
3-[2-fluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
3-[4-(1-methyl-1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
2-[6-(2,7-diazaspiro[3.5]non-2-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol

3-fluoro-4-{6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl}phenol
2-{6-[(2,6-dimethylpiperidin-4-yl)oxy]pyridazin-3-yl}-5-(1H-pyrazol-1-yl)phenol
N-methyl-6-(2-methyl-2H-indazol-5-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
2-methyl-5-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}-2H-indazole
3-(4-chloro-2-methoxyphenyl)-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
N-methyl-6-(2-methylpyrazolo[1,5-a]pyridin-3-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
6-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}imidazo[1,2-a]pyridine
3-[2-methoxy-4-(1H-pyrazol-1-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
3-[5-(1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
3-[5-(1-methyl-1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
3-[4-(1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
5-(3,5-dimethyl-1H-pyrazol-4-yl)-2-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}phenol
6-[2-fluoro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
3-methoxy-4-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}phenol
3-[2-methoxy-4-(4-nitro-1H-pyrazol-1-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
4-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}benzene-1,3-diol
6-[2-chloro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
2-(1H-pyrazol-4-yl)-4-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}pyrimidin-5-amine
3-[2,6-difluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
2-[6-(2,6-diazaspiro[3.4]oct-2-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
3-{6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl}-6-(1H-pyrazol-4-yl)pyridin-2-ol
6-(1H-pyrazol-4-yl)-3-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}pyridin-2-ol
N,2,2,6,6-pentamethyl-N-{5-[3-(1H-pyrazol-4-yl)phenoxy]-1,3,4-thiadiazol-2-yl}piperidin-4-amine
N,2,2,6,6-pentamethyl-N-{5-[4-(1H-pyrazol-4-yl)phenoxy]-1,3,4-thiadiazol-2-yl}piperidin-4-amine

3-[2-(difluoromethyl)-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine and

6-[2-(difluoromethyl)-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine;

wherein a form of the compound is selected from the group consisting of a prodrug, salt, hydrate, solvate, clathrate, isotopologue, racemate, enantiomer, diastereomer, stereoisomer, polymorph and tautomer form thereof.

8. A method of use of a compound salt or a form thereof for treating or ameliorating HD in
5 a subject in need thereof, comprising administering to the subject an effective amount of a compound salt selected from the group consisting of:

2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(trifluoromethyl)phenol hydrochloride

2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(1H-pyrazol-1-yl)phenol hydrochloride

2-(6-piperazin-1-yl-pyridazin-3-yl)-5-1H-pyrazol-1-yl-phenol hydrochloride

3-(6-(piperazin-1-yl)pyridazin-3-yl)naphthalene-2,7-diol trifluoroacetate

3-(6-(1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl)naphthalene-2,7-diol trifluoroacetate

3-(6-(1-methyl-1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl)naphthalene-2,7-diol trifluoroacetate

3-(6-(piperidin-4-yl)pyridazin-3-yl)naphthalene-2,7-diol trifluoroacetate

6-hydroxy-5-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-2,3-dihydro-1H-inden-1-one oxime hydrochloride

2-amino-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-8H-indeno[1,2-d]thiazol-5-ol hydrochloride

9-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5,6-dihydroimidazo[5,1-a]isoquinolin-8-ol hydrochloride

3-fluoro-5-(2-methoxypyridin-4-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol hydrochloride

4-(3-fluoro-5-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)pyridin-2(1H)-one hydrochloride

4-(3-fluoro-5-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)-1-methylpyridin-2(1H)-one hydrochloride

5-(3-fluoro-5-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)-1-methylpyridin-2(1H)-one hydrochloride

3-fluoro-5-(1H-pyrazol-4-yl)-2-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)phenol hydrochloride

5-chloro-3-fluoro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol hydrochloride

3-fluoro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1H-pyrazol-4-yl)phenol hydrochloride

3-fluoro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1-methyl-1H-pyrazol-4-yl)phenol hydrochloride

3-(1H-imidazol-1-yl)-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol hydrochloride

6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinoline-3,7-diol formate

7-hydroxy-1-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-2(1H)-one hydrochloride

4-methoxy-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-6-ol formate

7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinoxalin-6-ol hydrochloride

2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(tetrahydro-2H-pyran-4-yl)quinolin-7-ol formate

4-(dimethylamino)-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-quinolin-7-ol formate

6-hydroxy-2-methyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-1(2H)-one hydrochloride

3-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)naphthalen-2-ol hydrobromide

N-methyl-5-(5-(1-methyl-1H-pyrazol-4-yl)pyridin-2-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine hydrochloride

3-fluoro-2-(5-((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-1,3,4-thiadiazol-2-yl)-5-(1H-pyrazol-4-yl)phenol dihydrochloride

2-(2-chloro-4-(1H-pyrazol-4-yl)phenyl)-5-((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-1,3,4-thiadiazole hydrochloride

2-(2-chloro-4-(1H-pyrazol-4-yl)phenyl)-5-(2,7-diazaspiro[4.5]decan-2-yl)-1,3,4-thiadiazole hydrochloride

(R)-(4-(5-(2-chloro-4-(1H-pyrazol-4-yl)phenyl)-1,3,4-thiadiazol-2-yl)piperazin-2-yl)methanol hydrochloride

2-{6-[8-azabicyclo[3.2.1]oct-3-yl(methyl)amino]pyridazin-3-yl}-5-(1H-pyrazol-4-yl)phenol hydrochloride

2-[6-(8-azabicyclo[3.2.1]oct-3-ylamino)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol hydrochloride

5-(1H-pyrazol-4-yl)-2-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl}phenol hydrochloride

2-[6-(8-azabicyclo[3.2.1]oct-3-yloxy)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol hydrochloride

2-{6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl}-5-(4-nitro-1H-pyrazol-1-yl)phenol dihydrochloride

2-[6-(1-methyl-1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol trihydrochloride

5-(1H-pyrazol-4-yl)-2-[6-(1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]phenol trihydrochloride

2-[6-(1-ethyl-1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol trihydrochloride

2-{6-[methyl(piperidin-4-yl)amino]pyridazin-3-yl}-5-(1H-pyrazol-4-yl)phenol tetrahydrochloride

2-[6-(piperidin-4-ylamino)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol tetrahydrochloride

6-[2,5-difluoro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine tetrahydrochloride

2-[6-(8-azabicyclo[3.2.1]oct-2-en-3-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol hydrochloride

6-[2,3-difluoro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine hydrochloride

3-[2,5-difluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine trihydrochloride

2-{6-[(1R,5S)-8-azabicyclo[3.2.1]oct-3-ylamino]pyridazin-3-yl}-5-(1H-pyrazol-4-yl)phenol hydrochloride

6-[2-methoxy-6-(1H-pyrazol-4-yl)pyridin-3-yl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine hydrochloride

2-{6-[(2,6-dimethylpiperidin-4-yl)oxy]pyridazin-3-yl}-5-(1H-pyrazol-4-yl)phenol trihydrochloride

3-[2-fluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine hydrochloride

2-[6-(2,7-diazaspiro[3.5]non-2-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol tetrahydrochloride

3-[5-(1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine hydrochloride

3-[4-(1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine hydrochloride

6-[2-chloro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine trihydrochloride

3-[2,6-difluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine trihydrochloride

3-{6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl}-6-(1H-pyrazol-4-yl)pyridin-2-ol hydrochloride
 N,2,2,6,6-pentamethyl-N-{5-[3-(1H-pyrazol-4-yl)phenoxy]-1,3,4-thiadiazol-2-yl}piperidin-4-amine hydrochloride
 N,2,2,6,6-pentamethyl-N-{5-[4-(1H-pyrazol-4-yl)phenoxy]-1,3,4-thiadiazol-2-yl}piperidin-4-amine hydrochloride
 3-[2-(difluoromethyl)-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine hydrochloride and
 6-[2-(difluoromethyl)-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine hydrochloride;

wherein a form of the compound salt is selected from the group consisting of a prodrug, hydrate, solvate, clathrate, isotopologue, racemate, enantiomer, diastereomer, stereoisomer, polymorph and tautomer form thereof.

9. A compound salt selected from the group consisting of:

2-{6-[8-azabicyclo[3.2.1]oct-3-yl(methyl)amino]pyridazin-3-yl}-5-(1H-pyrazol-4-yl)phenol hydrochloride
 2-[6-(8-azabicyclo[3.2.1]oct-3-ylamino)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol hydrochloride
 5-(1H-pyrazol-4-yl)-2-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl}phenol hydrochloride
 2-[6-(8-azabicyclo[3.2.1]oct-3-yloxy)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol hydrochloride
 2-{6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl}-5-(4-nitro-1H-pyrazol-1-yl)phenol dihydrochloride
 2-[6-(1-methyl-1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol trihydrochloride
 5-(1H-pyrazol-4-yl)-2-[6-(1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]phenol trihydrochloride
 2-[6-(1-ethyl-1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol trihydrochloride
 2-{6-[methyl(piperidin-4-yl)amino]pyridazin-3-yl}-5-(1H-pyrazol-4-yl)phenol tetrahydrochloride
 2-[6-(piperidin-4-ylamino)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol tetrahydrochloride
 6-[2,5-difluoro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine tetrahydrochloride
 2-[6-(8-azabicyclo[3.2.1]oct-2-en-3-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol hydrochloride

6-[2,3-difluoro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine hydrochloride
 3-[2,5-difluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine trihydrochloride
 2-{6-[(1R,5S)-8-azabicyclo[3.2.1]oct-3-ylamino]pyridazin-3-yl}-5-(1H-pyrazol-4-yl)phenol hydrochloride
 6-[2-methoxy-6-(1H-pyrazol-4-yl)pyridin-3-yl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine hydrochloride
 2-{6-[(2,6-dimethylpiperidin-4-yl)oxy]pyridazin-3-yl}-5-(1H-pyrazol-4-yl)phenol trihydrochloride
 3-[2-fluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine hydrochloride
 2-[6-(2,7-diazaspiro[3.5]non-2-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol tetrahydrochloride
 3-[5-(1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine hydrochloride
 3-[4-(1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine hydrochloride
 6-[2-chloro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine trihydrochloride
 3-[2,6-difluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine trihydrochloride
 3-{6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl}-6-(1H-pyrazol-4-yl)pyridin-2-ol hydrochloride
 N,2,2,6,6-pentamethyl-N-{5-[3-(1H-pyrazol-4-yl)phenoxy]-1,3,4-thiadiazol-2-yl}piperidin-4-amine hydrochloride
 N,2,2,6,6-pentamethyl-N-{5-[4-(1H-pyrazol-4-yl)phenoxy]-1,3,4-thiadiazol-2-yl}piperidin-4-amine hydrochloride
 3-[2-(difluoromethyl)-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine hydrochloride and
 6-[2-(difluoromethyl)-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine hydrochloride;

wherein a form of the compound salt is selected from the group consisting of a prodrug, hydrate, solvate, clathrate, isotopologue, racemate, enantiomer, diastereomer, stereoisomer, polymorph and tautomer form thereof.

10. An assay for detecting HTT (Huntingtin protein) using capture/detection antibody pairs, wherein HTT is total HTT and the capture/detection antibody pairs are selected from MAB2166/MAB2174 and MAB2166/#5656S; and, wherein HTT is mutant HTT and the

capture/detection antibody pairs are selected from MAB1574/#5656S, P1874/#5656S and MW1/#5656S.

11. The assay of claim 10, wherein HTT is total HTT and the capture/detection antibody pair is MAB2166/#5656S; and, wherein HTT is mutant HTT and the capture/detection antibody pair is MW1/#5656S.
12. The method of any of claims 1, 6 or 8, wherein the effective amount of the compound is in a range of from about 0.001 mg/kg/day to about 500 mg/kg/day.
13. A use of the compound of any of claims 1, 6 or 8 for treating or ameliorating HD in a subject in need thereof, comprising administering to the subject an effective amount of the compound.
14. The use of claim 13, wherein the effective amount of the compound is in a range of from about 0.001 mg/kg/day to about 500 mg/kg/day.
15. A use of the compound of any of claims 1, 6 or 8 in the manufacture of a medicament for treating or ameliorating HD in a subject in need thereof, comprising administering an effective amount of the medicament to the subject.
16. The use of claim 15, wherein the effective amount of the compound in the medicament is in a range of from about 0.001 mg/kg/day to about 500 mg/kg/day.
17. A use of the compound of any of claims 1, 6 or 8 in admixture with one or more pharmaceutically acceptable excipient(s) in a pharmaceutical composition for treating or ameliorating HD in a subject in need thereof, comprising administering an effective amount of the pharmaceutical composition to the subject.
18. The use of claim 17, wherein the effective amount of the compound in the pharmaceutical composition is in a range of from about 0.001 mg/kg/day to about 500 mg/kg/day.