

TitleDerivatives of Dihydronepetalactone and
Method for Preparation

5

This application claims the benefit of U.S. Provisional Application No. 60/525,304, filed on November 26, 2003.

10


Field of the Invention

The present invention is directed to derivatives of dihydronepetalactone, which are useful
15 as fragrance compounds, to impart, for example, a desirable or attractive fragrance to objects, and are also useful as insect repellents. The present invention is also directed to a method for preparing derivatives of dihydronepetalactone, which method is
20 particularly well-suited for transforming naturally occurring nepetalactone to the desired derivative.

Background of the Invention

25 Dihydronepetalactone (DHN) is an iridoid monoterpenoid that may be represented schematically by the structure of Formula II,


30



II

and has been described by Hallahan in US 2003/0191047 (which is incorporated in its entirety as a part hereof for all purposes) as possessing a fragrance that many humans find attractive, and thus as having usefulness
5 as a fragrance compound. Hallahan, in WO 03/079786 (which is incorporated in its entirety as a part hereof for all purposes), has also disclosed that DHN compares favorably as an insect repellent with N,N-diethyl-m-toluamide (DEET), which has been the insect repellent
10 of choice in most commercially available insect repellents. DHN may be prepared by the catalytic hydrogenation of nepetalactone.

Various derivatives of DHN are known, among
15 which, for example, is nepetalic acid, which may be represented schematically by the structure of Formula III,



Nepetalic acid may be isolated from the essential oil
20 of *nepeta cataria* [see, for example, Eisenbraun et al, *J. Org. Chem.*, 53(17), 3968-3972 (1988)]. Eisenbraun also teaches the treatment of nepetalic acid with an oxidizer and base to produce a dicarboxylic acid ring-opened structure (nepetalinic acid).


25

Nepetalic acid (Formula III) may also be beneficially synthesized by treatment of nepetalactone with strong base according to the method of McElvain et al in *J. Am. Chem. Soc.*, 63, 1558-1563 (1941). The

structure of Formula III was confirmed by Eisenbraun et al in *J. Org. Chem.* 46, 3303-3305 (1981).

Another derivative of DHN, as disclosed in
5 Dawson et al, *Bioorganic and Medicinal Chemistry*, 4
(3), 351-361 (1996), is the ethoxy substituted
variation of the structure of Formula III, represented
by the structure of Formula IV

10.




IV

According to Dawson, *op. cit.*, the compound of Formula
IV is prepared as an intermediate in a synthetic
pathway presented therein for preparing nepetalactone
15 from R-pulegone.

In the references noted above, the respective
derivatives of DHN disclosed therein are not described
as being useful as insect repellents or for purposes
20 related to fragrance, aroma or odor.

Nepetalactone, represented schematically in
structure of Formula I,

25



I

is a major component of an essential oil secreted by plants of the genus *Nepeta*, and is the active ingredient in catnip. Although the aroma of nepetalactone is generally considered unpleasant by humans, cats find it attractive. Nepetalactone may also play a role as an insect repellent as studies of the repellency of catnip oil (predominantly nepetalactone) have shown that it was repellent towards a number of insect species on short-term exposure, but not to a number of other species [Eisner, T., *Science* 146:1318-1320 (1964)].

While nepetalactone and DHN are both compounds that have exhibited some degree of usefulness as an insect repellent, only DHN of the two presents a pleasing odor to humans. It is thus difficult to predict which member of the iridoid monoterpene family of compounds will or will not have any particular property or utility.


A need nevertheless remains to develop new compounds that will be useful as insect repellents but that will have other uses as well, some of which uses may be based on fragrance, aroma and odor. Applicants have discovered previously unknown compounds that are derivatives of DHN, and that have usefulness as insect repellents in addition to other uses based on fragrance, aroma and/or odor.

30

Summary of the Invention

The present invention provides for a compound represented schematically by the structure of Formula (V)

35



V

where R is alkyl or aryl.

5 The present invention further provides for a process to prepare a compound of Formula V comprising contacting nepetalic acid with a base, and with a Grignard reagent; or with only a Grignard reagent.

10 This invention also provides for a variety of compositions that include the compound of Formula V such as a composition that includes a mixture of different compounds according to Formula V, or a composition that includes a compound of Formula V and
15 one or both of an adjuvant and a carrier.

 This invention further provides for a perfumed or insect-repellent composition of matter or article of manufacture that includes a compound of
20 Formula V; and thus correspondingly provides a method of making a perfumed or insect-repellent composition of matter or article of manufacture by forming the composition from, or incorporating into the article, a compound of Formula V.

25

 In particular, this invention further provides a topical treatment for skin, such as a

fragrance or perfume composition, or an insect repellent, that includes a compound of Formula V.

5 This invention further provides a method of treating skin or fur by applying to the skin or fur a compound of Formula V, or a composition including a compound of Formula V. The invention thus also provides a method of repelling one or more insects by exposing the insect(s) to a compound of Formula V.

10

Brief Description of the Drawings

15 Figures 1-5 represent the results of testing the indicated compounds of the present invention against the indicated controls for their effect on the probing behavior of *Aedes aegypti* mosquitoes in the *in vitro* Gupta box landing assay procedure, described hereinbelow. The horizontal scale shows time in minutes, and the vertical scale shows mean number of landings of mosquitoes.

20

Detailed Description of Various Preferred Embodiments


25 The present invention provides an entirely new class of highly odorous bicyclic lactones useful variously as insect repellents and for other purposes related to fragrance, aroma and/or odor of the compound.

30 The compounds described herein will be recognized as exhibiting stereoisomerism, both enantiomerism and diastereomerism as the case may be. Unless a specific stereoisomer is indicated, the discussion will be understood to refer to all possible isomers, whether the structures are shown in the

35

stereochemically ambiguous form of the structures of Formulae I-V, or are shown as a specific stereoisomer (as in various locations hereinbelow) when other stereoisomers are also possible. There are numerous instances in which the discussion may focus on a specific stereoisomer of a diastereomeric pair (such as a case where, of several stereoisomers of identical chemical content, certain members of the group have differing chemical properties or biological activity), and this will also be specifically noted.

The present invention provides for the compounds that may be represented schematically by the structure of Formula V,



V

where R is alkyl or aryl.

While there is in principle no limitation on the type of alkyl or aryl groups that are useful in the practice of the invention, there will be practical considerations as to the size of the R substituent that would have practical use in commerce. Furthermore, it may be desirable to avoid incorporating highly reactive functionality in the R substituents to avoid side reactions.

In addition to practical considerations associated with synthesis, it will be recognized that, for end-uses where odor is a key factor, the higher molecular weight members of the class of compounds of this invention will generally exhibit lower respective vapor pressures under the same conditions than the lower molecular weight members of the class, and therefore may be less preferred in those kinds of circumstances.

10


In view of those considerations, representative values for R may include normal or branched C₁-C₂₀ alkyl such as sec-butyl or neo-pentyl, and is preferably normal C₁-C₁₂ alkyl such as methyl, ethyl, propyl, butyl, pentyl or hexyl; or R may be substituted or unsubstituted C₆-C₂₀ phenyl, and is preferably unsubstituted or meta- or para-substituted phenyl, where the substituent on phenyl may be normal C₁-C₁₂ alkyl or alkoxy, or halogen. Most preferably, R is methyl or ethyl, phenyl, or para- or meta-tolyl.

A compound according to this invention includes a compound that is a single stereoisomer as well as a compound that is a mixture of stereoisomers. A composition may be formed from a mixture of the compounds of this invention in which R, as described above, differs among the various compounds from which the composition is formed.

In the process of this invention, the compound represented schematically by the structure of Formula V is prepared from nepetalic acid by deprotonation, followed by treatment with Grignard reagent.

35

The nepetalic acid starting material is preferably prepared from nepetalactone according to the teachings of McElvain *supra*. The nepetalactone bicyclic structure can exist in any of four stereoisomeric forms, as shown in the structures of Formulae VIIa~VIIId.



VII

Nepetalactone extracted from the essential oil of the *nepeta* (catmint) plant leaves is a preferred source of raw material as nepetalactone is present in large quantity therein and may be readily purified therefrom. This produces a highly desirable route from a natural product to the compounds of the invention. Fractional distillation has been found to be an effective method for both purifying nepetalactone from the essential oils, and for separating the several stereoisomers from one another. Chromatographic separations are also suitable.


20

Only the first three listed stereoisomers of nepetalactone exist in the essential oil of the *nepeta cataria* plant. Cis, trans nepetalactone is the predominant isomer that may be isolated from the *nepeta cataria* plant and is therefore the most useful because of availability. Other plant species have been

25

identified that have essential oils enriched with the trans, cis and cis, cis nepetalactone isomers.

Following the method of McElvain et al, *op. cit.*, the starting nepetalic acid suitable for use in the process of this invention is prepared by hydration of nepetalactone with strong base to produce nepetalic acid. In the hydration of cis, trans nepetalactone, there exists the possibility of obtaining 4 diastereomers of nepetalic acid as shown schematically in structures of Formulae VIa~VI d.




15

The structures depicted in Formulae VI are diastereomeric pairs at C3 (the lactol) and C4 (the methyl proximate to the lactol). Eisenbraun et al, *J. Org. Chem.* 46, 3303-3305 (1981), showed that the absolute stereochemistry of the major nepetalic acid diastereomer that is crystallized from this product mixture is structure VIa. Spectral evidence and the solved single crystal x-ray structure indicate that the cyclic lactol tautomer VIa exists as a single diastereomer with respect to the methyl-bearing asymmetric carbon (C4) adjacent to the lactol carbon. Crystallization unambiguously sets the relative (and

25

thus absolute) stereochemical configuration of the methyl group at C4 such that the methyl occupies an equatorial configuration.

5 When *trans, cis* nepetalactone is subjected to hydration, the ring-opened tautomer is obtained as a diastereomeric pair:



10

Cis, cis-nepetalactone and trans, trans nepetalactone are also suitable for use in preparing nepetalic acid by hydration.


15

After the preparation of nepetalic acid, it is treated with a non-aqueous base in a deprotonation step to form a carboxylate salt. In one embodiment, for example, the nepetalic acid is treated with an alkali metal hydride, preferably KH. In this embodiment, the deprotonation step is followed by treatment with Grignard reagent to form a product represented schematically by the structure Formula V. Typical Grignard reagents include, but are not limited to, those that are prepared by the union of metallic magnesium with an organic chloride, bromide or iodide

usually in the presence of ether and in the complete absence of water.

In a preferred embodiment, the nepetalic acid is derived from *cis*, *trans* nepetalactone with the result that the product created by the above described steps may be represented schematically by the structure Formula VIII, with the absolute stereochemistry of the anti product shown.

10




VIII

Both the anti- and syn- addition products may be formed, and the relative amounts in which the two products are formed is governed primarily by steric factors resulting from the choice of Grignard reagent, as discussed in more detail below.

As previously discussed, nepetalic acid may, in an alternative embodiment, be derived from *trans*, *cis* nepetalactone, thereby forming the nepetalic acid tautomer consisting of the diastereomeric pair shown above. As a consequence, in this embodiment, treatment of that tautomer pair with an alkali metal hydride, preferably KH, followed by treatment with Grignard reagent results in production of the diastereomeric mixture represented schematically by the structures of Formulae IXa-IXb.

30



Although this particular embodiment of the process started with *trans*, *cis* nepetalactone, the final product is in the *cis*, *trans* configuration. Shown in IX are only the anti products. Both the anti- and syn-addition products may be formed, however, and the relative amounts in which the two products are formed is governed primarily by steric factors resulting from the choice of Grignard reagent, as discussed in more detail below.

Very reactive metal hydrides such as lithium aluminum hydride may be too reactive for use in the deprotonation step. Such species may reduce the lactone, which would not be desirable, and are therefore less preferred.

In a further, preferred, embodiment, the nepetalic acid is treated with more than one equivalent of Grignard reagent, and preferably at least two equivalents of Grignard reagent to effect formation of the products, such as those depicted respectively in the structures of Formulae VIII and IX. This embodiment of the process of the invention eliminates the step of first treating nepetalic acid with a separate non-aqueous base such as an alkali metal hydride. Use of an amount of Grignard reagent in slight excess of two equivalents ensures high conversion to the desired product.

Suitable for use in the process herein disclosed are any Grignard reagents that are reactive

with aldehydes including but not limited to alkyl
magnesium chlorides, alkyl magnesium bromides, aryl
magnesium chlorides, and aryl magnesium bromides. Also
included are dialkylzincs, diarylzincs and
5 alkyllithiums and aryllithiums. Representative
substituents that may be present on suitable Grignard
reagents include methyl, ethyl, n-propyl, butyl,
pentyl, hexyl phenyl, para- and meta-substituted phenyl,
including para- and meta-tolyl, and para-methoxy-
10 phenyl.

While there is no specific limitation on the
choice of Grignard reagents suitable for the practice
of the invention, some Grignard reactions are more
15 effective than others at generating products that are
high in purity in terms of stereochemistry. While not
wishing to be bound by any theory, it is believed that
steric factors govern the relative rates of production
of the anti- and syn-configured isomers produced by the
20 treatment of nepetalic acid with base and/or Grignard
reagent, such as those shown in the structures of
Formulae VIII and IX. The larger the organic radical
of the Grignard reagent, the greater the tendency is to
form predominantly anti-configured product. Small
25 organic radicals on the other hand are not sterically
hindered, and the anti-configured product is thus not
energetically favored over the syn-configured product.

It has been found that the smaller the alkyl
30 group on a Grignard reagent, the less
diastereoselective the addition reaction is. Methyl,
ethyl, and n-propyl Grignard show no
diastereoselectivity, forming a 1:1 mixture of anti-
addition product and syn-addition product. Branched
35 alkyls such as isopropyl and sec-butyl, or cyclic alkyl

such as cyclohexyl provide lower yields of the final products than do the comparable linear alkyls.

By contrast, phenyl Grignard is completely
5 diastereoselective for the anti-addition product. In particular, para-substituted phenyl Grignard reagents like p-tolyl Grignard, give exclusively the anti products with excellent 1,2-asymmetric induction from the established methyl-bearing C4. Conversely, ortho-
10 substituted phenyl Grignard reagents are less diastereoselective in this reaction, resulting in some syn product and thus a diastereomeric mixture. The governing mechanism may be related to the extent to which the given reactants follow the Felkin-Anh rules,
15 such rules being known in the art.

Aryl Grignard reagents are diastereoselective and provide 3-substituted dihydronepetalactones, whose absolute stereochemistry is consistent with the
20 structural representation of the embodiments depicted in Formula VIII. This is illustrated in the examples where it is observed that p-tolylmagnesium bromide gave only one diastereomer (as verified by single crystal x-ray and NMR analysis in Example 2) while
25 methylmagnesium bromide gave a 1:1 inseparable mixture of diastereomers (in Examples 3 and 4). Many 3-substituted dihydronepetalactones can be prepared by the process of this invention.

30 In one aspect of the invention, it is found that meta-substituted phenyls such as meta-tolyl give lactone products that are oils. The oils so formed performed better in the insect repellency tests described hereinbelow than did the lactones obtained

from phenyl and para-tolyl which gave more crystalline products such as para-tolyl.

5 In the processes described above, the steps
of adding base and then Grignard reagent, or of adding
Grignard reagent in one or more increments, may be
performed at a temperature in the range of about 25C to
about to -78C, with a temperature in the range of 5C to
-5C being preferred to prevent decomposition during
10 this step

The compounds of this invention are directed
to use for a multiplicity of purposes, such as use in
an effective amount for the repellency of various
15 insect species, or in a perfume composition or in a
topical treatment for skin. For example, the
compounds hereof may be applied in a topical manner to
human or animal skin, fur or feathers, or to growing
plants or crops, to impart insect repellency or a
20 pleasant odor or aroma.

Desirable properties of an insect repellent
include low toxicity, resistance to loss by water
immersion or sweating, low or no odor or at least a
25 pleasant odor, ease of application, and rapid formation
of a dry tack-free surface film on the surface of the
host to which applied. In order to obtain these
properties, the formulation for a topical insect
repellent should permit insect-infested animals (e.g.,
30 dogs with fleas, poultry with lice, cows with ticks,
and humans) to be treated with an insect repellent by
contacting the skin, fur or feathers of such an animal
with an effective amount of the repellent for repelling
the insect from the animal host. Thus, dispersing the
35 repellent into the air or dispersing the liquid as a

liquid mist or fine dust will permit the repellent to fall on or be collected on the desired surface of the host. Likewise, directly spreading of a liquid/semi-solid/solid repellent on the host is an effective
5 method of contacting the surface of the host with an effective amount of the repellent.

In formulating an insect repellent in accordance with this invention, it may be desirable to
10 blend more than one of the compounds hereof together in a composition, which includes the possibility of blending together more than one isomer of a particular compound. It may also be desirable to formulate an insect repellent by combining one or more compounds of
15 this invention to form a composition with a fugitive vehicle for application in the form of a spray.

A carrier useful for forming a composition in this invention can be any one of a variety of
20 commercially available organic and inorganic liquid, solid, or semi- solid carriers or carrier formulations usable in formulating skin or insect repellent products. When formulating a skin product or topical insect repellent, it is preferred to select a
25 dermatologically acceptable carrier. For example the carrier may include silicone, petrolatum, lanolin or many of several other well known carrier components. Examples of organic liquid carriers include liquid aliphatic hydrocarbons (e.g., pentane, hexane, heptane,
30 nonane, decane and their analogs) and liquid aromatic hydrocarbons. Examples of other liquid hydrocarbons include oils produced by the distillation of coal and the distillation of various types and grades of petrochemical stocks, including kerosene oils which are
35 obtained by fractional distillation of petroleum. Other petroleum oils include those generally referred

to as agricultural spray oils (e.g., the so-called light and medium spray oils, consisting of middle fractions in the distillation of petroleum and which are only slightly volatile). Such oils are usually
5 highly refined and may contain only minute amounts of unsaturated compounds. Such oils, moreover, are generally paraffin oils and accordingly can be emulsified with water and an emulsifier, diluted to lower concentrations, and used as sprays. Tall oils,
10 obtained from sulfate digestion of wood pulp, like the paraffin oils, can similarly be used. Other organic liquid carriers can include liquid terpene hydrocarbons and terpene alcohols such as alpha-pinene, dipentene, terpineol, and the like.

15

Other carriers include silicone, petrolatum, lanolin, liquid hydrocarbons, agricultural spray oils, paraffin oil, tall oils, liquid terpene hydrocarbons and terpene alcohols, aliphatic and aromatic alcohols,
20 esters, aldehydes, ketones, mineral oil, higher alcohols, finely divided organic and inorganic solid materials. In addition to the above-mentioned liquid hydrocarbons, the carrier can contain conventional emulsifying agents which can be used for causing the
25 dihydronepetalactone compounds to be dispersed in, and diluted with, water for end-use application. Still other liquid carriers can include organic solvents such as aliphatic and aromatic alcohols, esters, aldehydes, and ketones. Aliphatic monohydric alcohols include
30 methyl, ethyl, normal-propyl, isopropyl, normal-butyl, sec-butyl, and tert-butyl alcohols. Suitable alcohols include glycols (such as ethylene and propylene glycol) and pinacols. Suitable polyhydroxy alcohols include glycerol, arabitol, erythritol, sorbitol, and the like.
35 Finally, suitable cyclic alcohols include cyclopentyl and cyclohexyl alcohols.

Conventional aromatic and aliphatic esters, aldehydes and ketones can be used as carriers, and occasionally are used in combination with the above-mentioned alcohols. Still other liquid carriers include relatively high-boiling petroleum products such as mineral oil and higher alcohols (such as cetyl alcohol). Additionally, conventional or so-called "stabilizers" (e.g., tert-butyl sulfinyl dimethyl dithiocarbonate) can be used in conjunction with, or as a component of, the carrier or carriers comprising the compositions of the present invention.

Further, it may also be desirable to combine one or more compounds of this invention with one or more other types of insect repellents in a composition to achieve some particular beneficial effect as may result from such a combination. Suitable other insect repellents combinable for such purpose include but are not limited to DHN, benzil, benzyl benzoate, 2,3,4,5-bis(butyl-2-ene) tetrahydrofurfural, butoxypolypropylene glycol, N-butylacetanilide, normal-butyl-6,6-dimethyl-5,6-dihydro-1,4-pyrone-2-carboxylate, dibutyl adipate, dibutyl phthalate, di-normal-butyl succinate, N,N-diethyl-meta-toluamide, dimethyl carbate, dimethyl phthalate, 2-ethyl-2-butyl-1,3-propanediol, 2-ethyl-1,3-hexanediol, di-normal-propyl isocinchomeronate, 2-phenylcyclohexanol, p-methane-3,8-diol, and normal-propyl N, N-diethylsuccinamate.

In addition to the compound of Formula V, an insect repellent composition may also include one or more essential oils and/or active ingredients of essential oils. "Essential oils" are defined as any class of volatile oils obtained from plants possessing

the odor and other characteristic properties of the plant. Examples of useful essential oils include: almond bitter oil, anise oil, basil oil, bay oil, caraway oil, cardamom oil, cedar oil, celery oil, chamomile oil, cinnamon oil, citronella oil, clove oil, coriander oil, cumin oil, dill oil, eucalyptus oil, fennel oil, ginger oil, grapefruit oil, lemon oil, lime oil, mint oil, parsley oil, peppermint oil, pepper oil, rose oil, spearmint oil (menthol), sweet orange oil, thyme oil, turmeric oil, and oil of wintergreen. Examples of active ingredients in essential oils are: citronellal, methyl salicylate, ethyl salicylate, propyl salicylate, citronellol, safrole, and limonene.

The insects that may be repelled by the compounds of this invention may include any member of a large group of invertebrate animals characterized, in the adult state (non-adult insect states include larva and pupa) by division of the body into head, thorax, and abdomen, three pairs of legs, and, often (but not always) two pairs of membranous wings. This definition therefore includes a variety of biting insects [e.g. ants, bees, chiggers, fleas, black flies, green head flies, stable flies, horn flies (*haematobia irritans*), mosquitoes, ticks, wasps], wood-boring insects (e.g. termites), noxious insects (e.g. houseflies, cockroaches, lice, roaches, wood lice), and household pests (e.g. flour and bean beetles, dust mites, moths, silverfish, weevils). A host from which it may be desired to repel an insect may include any plant or animal (including humans) affected by insects. Typically, hosts are considered to be insect-acceptable food sources or insect-acceptable habitats.

In another embodiment, a compound may be used as a fragrance compound or in a fragrance composition, and be applied in a topical manner to human or animal

skin or hair to impart a pleasing fragrance, as in skin lotions and perfumes.

Particularly because of the pleasant aroma
5 associated with the compounds hereof, a further
embodiment of this invention is one in which one or
more compounds are formulated into a composition for
use as a product that is directed to other fundamental
purposes. The fragrance and/or insect repellency of
10 these products will be enhanced by the presence therein
of compound(s) of this invention. Included among such
products (but not thereto limited) are colognes,
lotions, sprays, creams, gels, ointments, bath and
shower gels, foam products (e.g., shaving foams),
15 makeup, deodorants, shampoo, hair lacquers/hair rinses,
and personal soap compositions (e.g., hand soaps and
bath/shower soaps). The compound(s) may of course be
incorporated into such products simply to impart a
pleasing aroma. Any means of incorporation such as is
20 practiced in the art is satisfactory.

A corresponding aspect of the wide variety of
products discussed above is a further alternative
embodiment of this invention, which is a process for
25 fabricating a composition of matter, a topical
treatment for skin, or an article of manufacture, by
providing as the composition, or incorporating into the
composition, skin treatment or article, a compound of
Formula V, or a mixture of the stereoisomers thereof.
30 Such products, and the method and process described
above, illustrate the use of a 3-substituted DHN as a
fragrance compound or perfume, or in a fragrance
composition or formulation, or in an article of
manufacture.

35

A composition containing compound(s) of this invention prepared as an insect repellent, fragrance product, or other personal care product may also contain other therapeutically or cosmetically active
5 adjuvants or ingredients as are typical in the personal care industry. Examples of these include fungicides, sunscreens agents, sunblocking agents, vitamins, tanning agents, plant extracts, anti-inflammatory agents, anti-oxidants, radical scavenging agents,
10 retinoids, alpha-hydroxy acids, antiseptics, antibiotics, antibacterial agents, antihistamines; adjuvants such as thickeners, buffering agents, chelating agents, preservatives, gelling agents, stabilizers, surfactants, emollients, coloring agents,
15 aloe vera, waxes, and penetration enhancers; and mixtures of any two or more thereof. The composition may also contain one or more carriers such as disclosed in US 2003/0191047 and/or WO 03/079786.

20 The amount of a compound of this invention contained in a composition will generally not exceed about 80% by weight based on the weight of the final product, however, greater amounts may be utilized in certain applications and this amount is not limiting.
25 More preferably, a suitable amount of a compound will be at least about 0.001% by weight and preferably about 0.01% up to about 50% by weight; and more preferably, from about 0.01% to about 20% weight percent, based on the weight of the composition or article. Specific
30 compositions will depend on the intended use.

In a further embodiment of this invention, compound(s) hereof is incorporated into an article to produce an insect repellent effect. Articles
35 contemplated to fall within this embodiment include manufactured goods, including textile goods such as clothing, outdoor or military equipment, natural

products such as lumber, or the leaves of insect vulnerable plants.

In another embodiment of this invention,
5 compound(s) hereof is incorporated into an article to produce a fragrance pleasing to some humans, or compound(s) hereof is applied to the surface of an object to impart an odor thereto. The particular manner of application will depend upon the surface in
10 question and the concentration required to impart the necessary intensity of odor.

Articles contemplated to fall within these embodiments include manufactured goods, including
15 textile goods, air fresheners, candles, various scented articles, fibers, sheets, paper, paint, ink, clay, wood, furniture (e.g., for patios and decks), carpets, sanitary goods, plastics, polymers, and the like.

20 In another embodiment, the composition of the invention may serve as an attractant for pests, and be highly suitable for use in a trap.

The present invention is further described
25 in, but not limited by, the following specific embodiments.

Examples 1-12: General Procedures

All reactions and manipulations were carried
30 out in a standard laboratory fume hood open in standard laboratory glassware. Nepetalactones were obtained by steam distillation of commercially-available catnip oil from catmint, obtained from Berjé, (Bloomfield, NJ). All inorganic salts and organic solvents, with the


exception for anhydrous THF, were obtained from VWR Scientific. All other reagents used in the examples were obtained from Sigma-Aldrich Chemical (Milwaukee, WI) and used as received. Determination of pH was done
5 with pHDrion paper from Micro Essential Laboratory. The 3-substituted dihydronepetalactone products were purified by column chromatography and characterized by NMR spectroscopy. NMR spectra were obtained on a Bruker DRX Advance (500 MHz ^1H , 125 MHz ^{13}C) using
10 deuterated solvents obtained from Cambridge Isotope Laboratories.

The batches of nepetalic acid employed in Examples 1-12 were prepared according to the following
15 procedures, although there was some variation in the amounts employed from batch to batch. A solution of cis-trans nepetalactone in 5mL of tetrahydrofuran (THF) was treated with sodium hydroxide (1.0 g in 5 (mL of water) resulting in initially a two-phase mixture and
20 then yellow solution. After 1 hour, the basic solution was extracted twice with fresh 20 mL aliquots of ethyl acetate. The aqueous layer from this extraction was acidified with 1N HCl drop-wise to pH = 3 at which point it turned into a white heterogeneous mixture.
25 The thus formed aqueous mixture was extracted twice with ethyl acetate and dried over anhydrous sodium sulfate. Removal of the solvent under vacuum resulted in a yellow oil which was triturated with petroleum ether (100 mL) and allowed to crystallize to a white
30 solid on standing. The white solid was filtered, washed with cold petroleum ether (20 mL) and dried under high vacuum to afford nepetalic acid (1.9 g, 69%) with a melting point of 67° C (lit. 71° C, J. Org. Chem. Vol. 46, No. 16, 1981 3302-3305). The absolute
35 stereochemistry of the product was verified by single crystal analysis and was consistent with the structural representation depicted in structure of Formula VIa.

Example 1

Preparation of 3-phenyl-DHN

((3*R*,4*R*,4a*R*,7*S*,7a*R*)-4,7-dimethyl-3-phenylhexahydrocyclopenta[*c*]pyran-1(3*H*)-one)



$C_{16}H_{20}O_2$
Mol. Wt.: 244.33

A


10

An oven-dried 100 mL three-necked round-bottomed flask was cooled to room temperature ("RT") under a steady stream of nitrogen. A solution of 500 mg of nepetalic acid in 10 mL anhydrous of THF was added to the flask and then cooled to 0°C. 363mg of a suspension of 30% potassium hydride in mineral oil was weighed into another flask and washed with two 10 mL aliquots of hexane and removed by decantation once the solid settled. A further 10 mL aliquot of hexane was added to the solid and the resulting mixture was added to the three-necked round-bottomed flask in one portion. After twenty minutes, the resulting mixture was treated at 0°C with phenylmagnesium bromide solution (4.1 mL of a 1M solution in THF) added drop-wise via syringe. After 30 minutes, the reaction warmed to room temperature and was stirred for 30 minutes. The mixture was acidified with 1N HCl drop-wise to pH = 3.0 and extracted with three 10 mL

aliquots of dichloromethane. The combined organic extracts were dried over anhydrous sodium sulfate. Removal of the solvent under vacuum resulted in a white solid (520 mg) with a melting point of 122-124° C. The
5 product was recrystallized with ether and hexane to >95% purity according to NMR analysis. The absolute stereochemistry of product was verified by single crystal x-ray analysis and consistent with
(3*R*,4*R*,4*aR*,7*S*,7*aR*)-4,7-dimethyl-3-
10 phenylhexahydrocyclopenta[*c*]pyran-1(3*H*)-one represented by structure A.

Example 2Preparation of 3-*p*-tolyl-DHN

((3*R*,4*R*,4*aR*,7*S*,7*aR*)-4,7-dimethyl-3-(4-methylphenyl)hexahydrocyclopenta[*c*]pyran-1(3*H*)-one)



B

$C_{17}H_{22}O_2$
Mol. Wt.: 258.36


10

An oven-dried 100 mL three-necked round-bottomed flask was cooled to RT under a steady stream of nitrogen. A solution of 1.9g of nepetalic acid in 40mL of anhydrous THF was added to the flask and then cooled to 0°C. 407mg of a suspension of 30% potassium hydride in mineral oil was weighed into another flask and washed with two 10 mL aliquots of hexane and removed by decantation once the solid settled. A further 10 mL aliquot of hexane was added to the solid and the resulting mixture was added to the three-necked round-bottomed flask in one portion. After twenty minutes, the resulting mixture was treated at 0°C with *p*-tolylmagnesium bromide solution (15 mL of a 1M solution in THF) drop-wise via syringe. After 30 minutes, the reaction warmed to RT and was stirred for 30 minutes. The mixture was acidified with 1N HCl drop-wise to pH = 3.0 and extracted with three 20mL aliquots of dichloromethane. The combined organic

extracts were dried over anhydrous sodium sulfate. Removal of the solvent under vacuum resulted in a white solid (2.8 g, > 100%) with a melting point of 127-128° C. The product was recrystallized with ether and
5 hexane to >95% purity according to NMR analysis. The absolute stereochemistry of product was verified by single crystal x-ray analysis and consistent with (3*R*,4*R*,4*aR*,7*S*,7*aR*)-4,7-dimethyl-3-(4-methylphenyl)hexahydrocyclopenta[*c*]pyran-1(3*H*)-one
10 represented by structure B.

Example 3

Preparation of 3-*m*-tolyl-DHN
(3*R*,4*R*,4*aR*,7*S*,7*aR*)-4,7-dimethyl-3-(3-methylphenyl)hexahydrocyclopenta[*c*]pyran-1(3*H*)-one)
15



C₁₇H₂₂O₂
Mol. Wt.: 258.36

C

20



An oven-dried 100 mL three-necked round-bottomed flask was cooled to RT under a steady stream of nitrogen. A solution of 1.4g of nepetalic acid in 30mL of anhydrous THF was added to the flask and then
25 cooled to 0°C. 304mg of a suspension of 30% potassium hydride in mineral oil was weighed into another flask and washed with two 10 mL aliquots of hexane and removed by decantation once the solid settled. A fresh

10mL aliquot of hexane was added to the solid and the resulting mixture was added to the three-necked round-bottomed flask in one portion. After twenty minutes, the resulting mixture was treated at 0°C with *m*-
5 tolylmagnesium bromide solution (15 mL of a 1M solution in THF) drop-wise via syringe. After 30 minutes, the reaction warmed to RT and was stirred for 30 minutes. The mixture was acidified with 1N HCl drop-wise to pH = 3.0 and extracted with three 20mL aliquots of
10 dichloromethane (20 mL). The combined organic extracts were dried over anhydrous sodium sulfate. Removal of the solvent under vacuum resulted in a yellow oil (1.7 g). The product was purified by column chromatography using silica gel eluting with hexane followed by ethyl
15 acetate. 0.66 g of purified products of (3*R*,4*R*,4*aR*,7*S*,7*aR*)-4,7-dimethyl-3-(3-methylphenyl)hexahydrocyclopenta[*c*]pyran-1(3*H*)-one was obtained as represented predominantly by the structure C.

20 Example 4

Preparation of 3-*p*-chlorophenyl-DHN
((3*R*,4*R*,4*aR*,7*S*,7*aR*)-3-(4-chlorophenyl)-4,7-dimethylhexahydrocyclopenta[*c*]pyran-1(3*H*)-one)

25



Mol. Wt.: 278.77

D


An oven-dried 100 mL three-necked round-bottomed flask was cooled to RT under a steady stream of nitrogen. A solution of 1.9g of nepetalic acid in 40mL of anhydrous THF was added to the flask and then cooled to 0°C. 407 mg of a suspension of 30% potassium hydride in mineral oil was weighed into another flask and washed with two 10 mL aliquots of hexane and removed by decantation once the solid settled. A fresh 10mL aliquot of hexane was added to the solid and the resulting mixture was added to the three-necked round-bottomed flask in one portion. After twenty minutes, the resulting mixture was treated at 0°C with *p*-chlorophenylmagnesium bromide solution (15 mL of a 1M solution in THF) drop-wise via syringe. After 30 minutes, the reaction warmed to RT and was stirred for 30 minutes. The mixture was acidified with 1N HCl drop-wise to pH = 3.0 and extracted with three 20mL aliquots of dichloromethane. The combined organic extracts were dried over anhydrous sodium sulfate. Removal of the solvent under vacuum resulted in a white solid (1.82 g). The product was purified by column chromatography using silica gel eluting with hexane followed by ethyl acetate. 290 mg of purified products obtained as a crystalline solid with a melting point of 137-139° C and NMR analysis of the product was consistent with the product (3*R*,4*R*,4*aR*,7*S*,7*aR*)-3-(4-chlorophenyl)-4,7-dimethylhexahydrocyclopenta[*c*]pyran-1(3*H*)-one represented in structure D..

30

Example 5Preparation of 3-*p*-methoxyphenyl-DHN

((3*R*,4*R*,4*aR*,7*S*,7*aR*)-3-(4-methoxyphenyl)-4,7-dimethylhexahydrocyclopenta[*c*]pyran-1(3*H*)-one)

5



E


$C_{17}H_{22}O_3$
Mol. Wt.: 274.35

10 An oven-dried 100 mL three-necked round-bottomed flask was cooled to RT under a steady stream of nitrogen. A solution of 1.9g of nepetalic acid (1.9 g) in anhydrous THF (40 mL) was added to the flask and then cooled to 0°C. 407 mg of a suspension of 30%
15 potassium hydride in mineral oil was weighed into another flask and washed with two 10 mL aliquots of hexane and removed by decantation once the solid settled. A fresh 10mL aliquot of hexane was added to the solid and the resulting mixture was added to the
20 three-necked round-bottomed flask in one portion. After twenty minutes, the resulting mixture was treated at 0°C with *p*-methoxyphenylmagnesium bromide solution (30 mL of a 0.5 M solution in THF) drop-wise via syringe. After 30 minutes, the reaction warmed to RT
25 and was stirred for 30 minutes. The mixture was acidified with 1N HCl drop-wise to pH = 3.0 and extracted with three 20mL aliquots of dichloromethane. The combined organic extracts were dried over anhydrous

sodium sulfate. Removal of the solvent under vacuum resulted in a white solid (0.89 g). The product was purified by column chromatography using silica gel eluting with hexane followed by ethyl acetate. 214 mg of purified products obtained as a crystalline solid with a melting point of 125-126° C and NMR analysis of the product was consistent with (3*R*,4*R*,4a*R*,7*S*,7a*R*)-3-(4-methoxyphenyl)-4,7-dimethylhexahydrocyclopenta[*c*]pyran-1(3*H*)-one as represented by the structure E.

Example 6

Preparation of 3-*m*-methoxyphenyl-DHN
((3*R*,4*R*,4a*R*,7*S*,7a*R*)-3-(3-methoxyphenyl)-4,7-dimethylhexahydrocyclopenta[*c*]pyran-1(3*H*)-one)




C₁₇H₂₂O₃
Mol. Wt.: 274.35

F

An oven-dried 100 mL three-necked round-bottomed flask was cooled to RT under a steady stream of nitrogen. A solution of 1.9g of nepetalic acid in 40mL of anhydrous THF was added to the flask and then cooled to 0°C. 407mg of a suspension of 30% potassium hydride in mineral oil was weighed into another flask and washed with two 10 mL aliquots of hexane and removed by decantation once the solid settled. A fresh 10mL aliquot of hexane was added to the solid and the

resulting mixture was added to the three-necked round-bottomed flask in one portion. After twenty minutes, the resulting mixture was treated at 0°C with *m*-methoxyphenylmagnesium bromide solution (15 mL of a 1M solution in THF) drop-wise via syringe. After 30 minutes, the reaction warmed to RT and was stirred for 30 minutes. The mixture was acidified with 1N HCl drop-wise to pH = 3.0 and extracted with three 20mL aliquots of dichloromethane (20 mL). The combined organic extracts were dried over anhydrous sodium sulfate. Removal of the solvent under vacuum resulted in a yellow oil (2.0 g). The product was purified by column chromatography using silica gel eluting with hexane followed by ethyl acetate. 0.466 g of purified products was obtained and NMR analysis of the product was consistent with (3*R*,4*R*,4*aR*,7*S*,7*aR*)-3-(3-methoxyphenyl)-4,7-dimethylhexahydrocyclopenta[*c*]pyran-1(3*H*)-one and represented by structure F.

Example 7
Preparation of 3-*p*-fluorophenyl-DHN
((3*R*,4*R*,4*aR*,7*S*,7*aR*)-3-(4-fluorophenyl)-4,7-dimethylhexahydrocyclopenta[*c*]pyran-1(3*H*)-one)



C₁₆H₁₉FO₂
Mol. Wt.: 262.32


G

An oven-dried 100 mL three-necked round-bottomed flask was cooled to RT under a steady stream of nitrogen. A solution of 1.9g of nepetalic acid in 40mL of anhydrous THF was added to the flask and then
5 cooled to 0°C. 407mg of a suspension of 30% potassium hydride in mineral oil was weighed into another flask and washed with two 10 mL aliquots of hexane and removed by decantation once the solid settled. A fresh 10mL aliquot of hexane was added to the solid and the
10 resulting mixture was added to the three-necked round-bottomed flask in one portion. After twenty minutes, the resulting mixture was treated at 0°C with *p*-methoxyphenylmagnesium bromide solution (15 mL of a 1.0 M solution in THF) drop-wise via syringe. After 30
15 minutes, the reaction warmed to RT and was stirred for 30 minutes. The mixture was acidified with 1N HCl drop-wise to pH = 3.0 and extracted with three 20mL aliquots of dichloromethane. The combined organic extracts were dried over anhydrous sodium sulfate.
20 Removal of the solvent under vacuum resulted in a white solid (2.04 g). The product was purified by column chromatography using silica gel eluting with hexane followed by ethyl acetate. 405 mg of purified products were obtained as a crystalline solid with a melting
25 point of 154-156° C and NMR analysis of the product was consistent with (3*R*,4*R*,4*aR*,7*S*,7*aR*)-3-(4-fluorophenyl)-4,7-dimethylhexahydrocyclopenta[*c*]pyran-1(3*H*)-one as represented by structure G.

Example 8

Preparation of 3-(*p*-fluoro-*m*-methyl)-phenyl-DHN ((3*R*,4*R*,4*aR*,7*S*,7*aR*)-3-(4-fluoro-3-methylphenyl)-4,7-dimethylhexahydrocyclopenta[*c*]pyran-1(3*H*)-one)

5



10

An oven-dried 100 mL three-necked round-bottomed flask was cooled to RT under a steady stream of nitrogen. A solution of 1.9g of nepetalic acid in 40mL of anhydrous THF was added to the flask and then

15 cooled to 0°C. 407mg of a suspension of 30% potassium hydride in mineral oil was weighed into another flask and washed with two 10 mL aliquots of hexane and removed by decantation once the solid settled. A fresh

20 10mL aliquot of hexane was added to the solid and the resulting mixture was added to the three-necked round-bottomed flask in one portion. After twenty minutes, the resulting mixture was treated at 0°C with *m*-methoxyphenylmagnesium bromide solution (15 mL of a 1M solution in THF) drop-wise via syringe. After 30


25 minutes, the reaction warmed to RT and was stirred for 30 minutes. The mixture was acidified with 1N HCl drop-wise to pH = 3.0 and extracted with three 20mL aliquots of dichloromethane. The combined organic

extracts were dried over anhydrous sodium sulfate. Removal of the solvent under vacuum resulted in a yellow oil (3.2 g). The product was purified by column chromatography using silica gel eluting with hexane followed by ethyl acetate. 0.616 g of purified product was obtained and NMR analysis of the product was consistent with (3*R*,4*R*,4*aR*,7*S*,7*aR*)-3-(4-fluoro-3-methylphenyl)-4,7-dimethylhexahydrocyclopenta[*c*]pyran-1(3*H*)-one as represented by structure H.

10 Example 9

Preparation of 3-methyl-DHN

((3*S*,4*R*,4*aR*,7*S*,7*aR*)-3,4,7-trimethylhexahydrocyclopenta[*c*]pyran-1(3*H*)-one, and (3*R*,4*R*,4*aR*,7*S*,7*aR*)-3,4,7-trimethylhexahydrocyclopenta[*c*]pyran-1(3*H*)-one)



$C_{11}H_{18}O_2$
Mol. Wt.: 182.26

20

An oven-dried 100 mL three-necked round-bottomed flask was cooled to RT under a steady stream of nitrogen. A solution of 2.13g of nepetalic acid in 40mL of anhydrous THF (40 mL) was added to the flask and then cooled to 0°C. 1.40g of a suspension of 30% potassium hydride in mineral oil was weighed into another flask and washed with two 20 mL aliquots of


hexane and removed by decantation once the solid settled. A further 20mL aliquot of hexane was added to the solid and the resulting mixture was added to the three-necked round-bottomed flask in one portion.

- 5 After twenty minutes, the resulting mixture was treated at 0°C with methylmagnesium chloride solution (5.8 mL of 3M solution in THF) drop-wise via syringe. After 30 minutes, the reaction warmed to RT and was stirred for 30 minutes. The mixture was acidified with 1N HCl
- 10 drop-wise to pH = 3.0 and extracted with three 40mL aliquots of dichloromethane. The combined organic extracts were dried over anhydrous sodium sulfate. Removal of the solvent under vacuum resulted in a pale oil (1.88 g). Purification of the crude reaction
- 15 product was carried out by column chromatography using silica gel eluting with hexane followed by ethyl acetate. The 3-methyl-dihydronepetalactone diastereomers were not separable by this method. 0.682 g of purified product was obtained and NMR analysis of
- 20 the product was consistent with the (3*S*,4*R*,4*aR*,7*S*,7*aR*)-3,4,7-trimethylhexahydrocyclopenta[*c*]pyran-1(3*H*)-one and (3*R*,4*R*,4*aR*,7*S*,7*aR*)-3,4,7-trimethylhexahydrocyclopenta[*c*]pyran-1(3*H*)-one as represented by the diastereomeric structure I.

25 Example 10

Preparation of 3-ethyl-DHN

- ((3*R*,4*R*,4*aR*,7*S*,7*aR*)-3-ethyl-4,7-dimethylhexahydrocyclopenta[*c*]pyran-1(3*H*)-one, and (3*S*,4*R*,4*aR*,7*S*,7*aR*)-3-ethyl-4,7-
- 30 dimethylhexahydrocyclopenta[*c*]pyran-1(3*H*)-one)



J

$C_{12}H_{20}O_2$
Mol. Wt.: 196.29

An oven-dried 100 mL three-necked round-bottomed flask was cooled to RT under a steady stream of nitrogen. A solution of 1.84g of nepetalic acid in 40mL of anhydrous THF was added to the flask and then cooled to 0°C. 441mg of a suspension of 30% potassium hydride in mineral oil was weighed into another flask and washed with two 20 mL aliquots of hexane and removed by decantation once the solid settled. A further 20 mL aliquot of hexane was added to the solid and the resulting mixture was added to the three-necked round-bottomed flask in one portion. After twenty minutes, the resulting mixture was treated at 0°C with ethylmagnesium chloride solution (15 mL of 1.0 M solution in THF) drop-wise via syringe. After 30 minutes, the reaction warmed to RT and was stirred for 30 minutes. The mixture was acidified with 1N HCl drop-wise to pH = 3.0 and extracted with three 20mL aliquots of dichloromethane. The combined organic extracts were dried over anhydrous sodium sulfate. Removal of the solvent under vacuum resulted in a pale oil (1.21 g). Purification of the crude reaction product was carried out by column chromatography using silica gel eluting with hexane followed by ethyl acetate. The 3-ethyl-dihydronepetalactone diastereomers were not separable by this method. 0.253

g of purified product was obtained and NMR analysis of the product was consistent with (3*R*,4*R*,4*aR*,7*S*,7*aR*)-3-ethyl-4,7-dimethylhexahydrocyclopenta[*c*]pyran-1(3*H*)-one and (3*S*,4*R*,4*aR*,7*S*,7*aR*)-3-ethyl-4,7-


5 dimethylhexahydrocyclopenta[*c*]pyran-1(3*H*)-one as represented by the diastereomeric structure J.

Example 11

Preparation of 3-*n*-propyl-DHN

((3*S*,4*R*,4*aR*,7*S*,7*aR*)-4,7-dimethyl-3-
10 propylhexahydrocyclopenta[*c*]pyran-1(3*H*)-one, and
(3*R*,4*R*,4*aR*,7*S*,7*aR*)-4,7-dimethyl-3-
propylhexahydrocyclopenta[*c*]pyran-1(3*H*)-one)

15



K

$C_{13}H_{22}O_2$
Mol. Wt.: 210.31

An oven-dried 100 mL three-necked round-
20 bottomed flask was cooled to RT under a steady stream
of nitrogen. A solution of 1.84g of nepetalic acid in
40mL of anhydrous THF was added to the flask and then
cooled to 0°C. 441mg of a suspension of 30% potassium
hydride in mineral oil was weighed into another flask
25 and washed with two 20 mL aliquots of hexane and
removed by decantation once the solid settled. A
further 20mL aliquot of hexane was added to the solid
and the resulting mixture was added to the three-necked

round-bottomed flask in one portion. After twenty minutes, the resulting mixture was treated at 0°C with *n*-propylmagnesium chloride solution (15 mL of 1.0 M solution in THF) drop-wise via syringe. After 30 minutes, the reaction warmed to RT and was stirred for 30 minutes. The mixture was acidified with 1N HCl drop-wise to pH = 3.0 and extracted with three 20mL aliquots of dichloromethane. The combined organic extracts were dried over anhydrous sodium sulfate.


Removal of the solvent under vacuum resulted in a pale oil (1.26 g). Purification of the crude reaction product was carried out by column chromatography using silica gel eluting with hexane followed by ethyl acetate. The 4-*n*-propyl-dihydronepetalactone diastereomers were not separable by this method. 0.713 g of purified product was obtained and NMR analysis of the product was consistent with (3*S*,4*R*,4*aR*,7*S*,7*aR*)-4,7-dimethyl-3-propylhexahydrocyclopenta[*c*]pyran-1(3*H*)-one and (3*R*,4*R*,4*aR*,7*S*,7*aR*)-4,7-dimethyl-3-propylhexahydrocyclopenta[*c*]pyran-1(3*H*)-one as represented by the diastereomeric structure K.

Example 12

Preparation of 3-methyl-DHN

((3*S*,4*R*,4*aR*,7*S*,7*aR*)-3,4,7-trimethylhexahydrocyclopenta[*c*]pyran-1(3*H*)-one, and (3*R*,4*R*,4*aR*,7*S*,7*aR*)-3,4,7-trimethylhexahydrocyclopenta[*c*]pyran-1(3*H*)-one)

5



L

$C_{11}H_{18}O_2$
Mol. Wt.: 182.26

An oven-dried 3 liter three-necked round-bottomed flask was cooled to RT under a steady stream of nitrogen. A solution of 55.3g of nepetalic acid in 800mL of anhydrous THF was added to the flask and then cooled to 0°C. The solution was treated with methylmagnesium bromide solution (200 mL 3.0M solution in ether) drop-wise via syringe. After 30 minutes, the reaction warmed to RT and was stirred for 30 minutes. The mixture was acidified with 1N HCl drop-wise to pH = 3.0 and extracted with three 200mL aliquots of dichloromethane. The combined organic extracts were dried over anhydrous sodium sulfate. Removal of the solvent under vacuum resulted in a pale oil (51.0 g). Purification of the crude reaction product was carried out by column chromatography using silica gel eluting with hexane followed by ethyl acetate. The 3-methyl-dihydronepetalactone diastereomers were not separable by this method. 21.77 g of purified product was obtained and NMR analysis of the product was consistent with (3*S*,4*R*,4*aR*,7*S*,7*aR*)-3,4,7-trimethylhexahydrocyclopenta[*c*]pyran-1(3*H*)-one and (3*R*,4*R*,4*aR*,7*S*,7*aR*)-3,4,7-

trimethylhexahydrocyclopenta[c]pyran-1(3H)-one as represented by the diastereomeric structure L.

Examples 13-19

- 5 The products of Examples 1-12 were evaluated for the insect repellency against *Aedes aegypti* mosquitoes in the *in vitro* Gupta box landing assay. In this method a chamber contained 5 wells, each covered by a Baudruche (animal intestine) membrane. Each well was
- 10 filled with bovine blood, containing sodium citrate (to prevent clotting) and ATP (72 mg ATP disodium salt per 26 ml of blood), and heated to 37°C. A volume of 25 µl of isopropyl alcohol (IPA) containing one test specimen or control was applied to each membrane. The
- 15 concentrations were all 1% in IPA except where otherwise indicated. Controls were either neat IPA, an untreated membrane surface, or a membrane surface treated with a 1% solution of DEET..
- 20 After 5 min, approximately 250 4-day-old female *Aedes aegypti* mosquitoes were introduced into the chamber. The number of mosquitoes probing the membranes for each treatment was recorded at 2 min intervals over 20 min. Each datum represents the mean
- 25 of three replicate experiments.


Example 13

- Specimens prepared in Example 1, Example 2, and Example 5 were evaluated according to the test
- 30 method above described. Results are shown in Figure 1.

Claims

1. A compound represented schematically by the structure of Formula V wherein R is alkyl or aryl:

5



2. A compound according to Claim 1 wherein R
10 is C₁ to C₂₀ alkyl.

3. A compound according to Claim 1 wherein R
is C₆ to C₂₀ aryl.

15 4. A compound according to Claim 1 wherein R
is selected from the group consisting of methyl, ethyl,
propyl, phenyl, tolyl, alkoxy-phenyl and halo-phenyl.

20 5. A compound according to Claim 1 which is a
single stereoisomer of a single compound.

6. A compound according to Claim 1 which is a
mixture of stereoisomers of a single compound.

25 7. A compound according to Claim 1 which is
selected from one or more members of the group
consisting of
(3*R*,4*R*,4*aR*,7*S*,7*aR*)-4,7-dimethyl-3-(4-
methylphenyl)hexahydrocyclopenta[*c*]pyran-1(3*H*)-one;

(3*R*,4*R*,4*aR*,7*S*,7*aR*)-3-(3-methoxyphenyl)-4,7-dimethylhexahydrocyclopenta[*c*]pyran-1(3*H*)-one;
(3*S*,4*R*,4*aR*,7*S*,7*aR*)-3,4,7-trimethylhexahydrocyclopenta[*c*]pyran-1(3*H*)-one; and
5 (3*R*,4*R*,4*aR*,7*S*,7*aR*)-3,4,7-trimethylhexahydrocyclopenta[*c*]pyran-1(3*H*)-one.

8. A process for preparing a compound according to Claim 1 comprising contacting nepetalic
10 acid with a base, and with a Grignard reagent.

9. A process according to Claim 8 wherein the base is a non-aqueous base.

15 10. A process according to Claim 8 wherein the base is a Grignard reagent.

11. A process according to Claim 8 wherein the Grignard reagent is an alkyl Grignard reagent.
20

12. A process according to Claim 8 wherein the Grignard reagent is an aryl Grignard reagent.

13. A process according to Claim 8 further
25 comprising a step of providing nepetalic acid by the hydration of nepetalactone.

14. A process for preparing a compound according to Claim 1 comprising contacting nepetalic
30 acid with a Grignard reagent, wherein the amount of Grignard reagent per mole of nepetalic acid is greater than 1.

15. A process according to Claim 14 wherein
35 the Grignard reagent is an alkyl Grignard reagent.

16. A process according to Claim 14 wherein the Grignard reagent is an aryl Grignard reagent.

17. A process according to Claim 14 further
5 comprising a step of providing nepetalic acid by the hydration of nepetalactone.

18. A composition of matter comprising the compound of Claim 1.

10

19. A method of repelling one or more insects comprising exposing the insect(s) to a compound according to Claim 1.

15

20. A method according to Claim 19 wherein the insect(s) are selected from one or more members of the group consisting of biting flies, chiggers, fleas, mosquitoes, ticks and lice.