


6-ALKOXYALKYL ESTRADIOL DERIVATIVES AND METHODS OF USE

CROSS REFERENCE TO RELATED APPLICATIONS

This application claims the benefit of U.S. Provisional
 5 Application No. 60/867,980 filed November 30, 2006. This
 application is also a continuation-in-part of U.S.
 Application No. 11/541,987 filed on October 02, 2006,
 currently pending, which claims the priority benefit from
 U.S. Provisional Application No. 60/722,204 filed September
 10 30, 2005. The teachings of all such applications are
 incorporated herein by their entirety.

FIELD OF THE INVENTION

The present invention relates to compositions and
 methods of making 6-alkoxyalkyl Estradiol compounds with
 15 the general formula (I) and in particular of compounds (R or
 S) 6 hydroxymethyl- or (R or S) 6-methyloxymethyl- (8 R or
 S, 9S, 13 R or S, 14S, 17 R or S) -13-methyl-
 7,8,9,11,12,14,15,16,17-decehydrocyclopenta [a]
 phenantherene-3, 17-diol and their pharmaceutically
 20 acceptable salts, or prodrugs thereof. The present invention
 also pertains to pharmaceutical compositions comprising such
 compounds, both in vitro and in vivo, for both diagnostic
 applications and treatment of proliferative conditions, such
 as cancer.



Formula (I)

BACKGROUND OF THE INVENTION

Proliferative cell disorders such as tumors and primary malignant tumors {herein, cancer(s)} in particular are problematic given their tendency to invade surrounding
5 tissues and metastasize to distant organs in the body. To date, the most frequently used methods for treating neoplasia, especially solid tumor forms of neoplasia, include surgical procedures, radiation therapy, drug therapies, and combinations of the foregoing.

10 With over million cases of cancer being diagnosed annually, and cancer claiming more than half a million lives in the United States each year, there is increased need in new therapeutic modalities against such condition. Prostate, lung and colorectal remains the most common cancer among
15 men; while breast, colorectal and lung cancers are the most common cancers among women.

In recent years, there have been significant gains in the management of these conditions. At least one of the success stories in the clinical management of a cancer is
20 the early diagnosis and treatment options now available for primary breast cancer. The introduction of effective and nontoxic anti-estrogen agents that block the actions of estrogen has been shown effective.

Tamoxifen is primarily one of the first selective
25 estrogen receptor modulators that have become first-line therapy for hormonal treatment of breast cancer, both for adjuvant treatment and for therapy of metastatic disease. Tamoxifen is a competitive inhibitor of estradiol binding to the estrogen receptor inhibiting its estrogen binding to the
30 estrogen binding element on DNA. In addition, there has been an increased interest in the use of aromatase inhibitors to block specifically the local production of


estrogens that may contribute substantially to hormone responsive disease such as breast cancer.

Aromatase(CYP19) is described as the principal enzyme that converts androgens to estrogens both in pre- and
5 postmenopausal women. Estrogen deprivation through aromatase inhibition is described as an effective and selective treatment for some postmenopausal patients with hormone-dependent breast cancer. Exemestane (which is sold as Aromasin, is chemically described as 6-methylenandrosta-1,4-
10 diene-3,17-dione)and acts as an irreversible, steroidal aromatase inactivator. It is believed to act as a false substrate for the aromatase enzyme, and processed to an intermediate that binds irreversibly to the active site of the enzyme causing its inactivation. U.S. Patent Nos.
15 4,808,616, and 4,904,650, the teachings of which are incorporated herein in their entirety, disclose 6-alkylidenandrosta-1,4-diene-3,17-dione derivatives, such as exemestane, and methods of making them. U.S. Patent No. 4,876,045 discloses a method of preparing 6-methylene
20 derivatives of androsta-1,4-diene-3,17-diones. U.S. Patent No. 4,990,635 discloses a process for making 6-methylene derivatives of androsta-1,4-diene-3,17-diones.


The preparation of intermediates that may be useful in preparing exemestane is disclosed in U.S. Patent No.
25 3,274,176. In German patent DD 258820, 6-hydroxymethyl-androsta-1,4-diene-3,17-dione is prepared from androsta-1,4-diene-3,17-dione via 1,3-dipyrrolidinoandrosta-3,5-dien-17-one.

Co-pending international application no.
30 PCT/US2005/001248 filed January 14, 2005 (PCT Publication Number WO 2005/070951) also describes the preparation of intermediates that are useful in preparing exemestane, such

application is incorporated herein by reference, in its entirety. The structure of Exemestane is shown below.



Schneider et. al, in "Course of the reaction of
 5 steroidal 3,5-dienamines with formaldehyde", Helvetica
 Chimica Acta (1973), 56(7), 2396-2404, discloses the
 following compounds:



where the ---- symbol represents a double bond, it means a
 10 keto group and no R₆ is present; and where the ---- symbol
 represents a single bond R₆ is hydrogen (i.e. an alcohol
 group). Unlike the compounds of the present invention,
 Schneider's compounds do not embrace estrodial, testosterone
 or dihydrotestosterone variations.

15 A tri-hydroxyl substituted derivative of estranes is
 disclosed in U.S. Patent No. 3,377,363 to Tadanier et. al,
 and the 3 hydroxy substituent on the aromatic ring of the
 present compounds is not disclosed.

U.S. Patent No. 5,914,324 to De Funari et.al, discloses
 20 6 hydroxy and oxy androstane derivatives for hypertension
 and heart failure. U.S. Patent 6,384,250 to Gobbini, et al.,
 discloses the hydroxyl and ketone substituents at the 6


position in the preparation of (E,Z) 3-(2-aminoethoxyimino)-androstane-6,17-dione. These compounds were directed towards the treatment of heart failure. The effects of alkyl hydroxyl substitution at the 6 position is not disclosed.

5 Tanenbaum, et. al, "Crystallographic comparison of the estrogen and progesterone receptor's ligand binding domains", Proc. Natl. Acad. Sci. USA, Biochemistry, Vol. 95, pp 5998-6003, discloses the mechanism of ER receptors and notes that estradiol containing an aromatic ring with a 3-hydroxy
10 substituent binds well with the ER ligand binding region. It is disclosed that a flat aromatic group without the 19 methyl substituent is favored. Given the above, a need still exists to identify new and effective agents for treating cancer.


15 SUMMARY OF THE INVENTION

In light of the foregoing, the present invention is directed towards chemotherapeutic compounds, compositions and methods for their use and preparation, thereby overcoming various deficiencies and shortcomings of the
20 prior art, including those outlined above.

An aspect of the present invention pertains to a compound of Formula I and II.



25 Formula (I)



Formula (II)

wherein R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , R_7 , R_8 and R_9 are independently hydrogen, C_1 to C_6 alkyl or substituted alkyl, halogen, sulfate, or glucuronide moieties; and the ---- symbol represents either a single or a double bond and when the ---
5 _ symbol is a double bond and forms a keto group at position 3 or 17, then no R_7 or R_6 is present, respectively; the symbol --- represent the presence or absence of a bond at position 10; and the ~~~~~ symbol represents any type of bond regardless of the stereochemistry. The compounds also
10 embrace the enantiomers, other stereochemical isomers, hydrates, solvates, tautomers and pharmaceutically acceptable salts thereof.

The present invention relates to a method of therapeutically treating cancer in a mammalian subject
15 (e.g., a human patient). In this aspect of the invention, methods are provided for inhibiting tumor or cancerous cell growth within the mammalian subject. In such a method, the cells are exposed to or contacted with a compound of Formula (I) or (II) or pharmaceutically acceptable enantiomers,
20 other stereochemical isomers, hydrates, solvates, tautomers, or salts thereof, as shown herein. In a specific, non-limiting embodiment of the methods of the present invention, a compound of Formula (I) or (II) is used to therapeutically treat an identified cancer state as described herein. In
25 another specific non-limiting embodiment of the methods of the present invention, a composition comprising a compound of Formula (I) or (II) is used to therapeutically treat an identified cancer state as described herein.


The compounds of the present invention may be used to
30 treat any tumor which may be either directly or indirectly effected by hormonal and/or estrogen-related activity, including but not in any way limited to solid tumors

associated with breast, pancreatic, lung, colon, prostate, ovarian cancers, as well as brain, liver, spleen, kidney, lymph node, small intestine, blood cells, bone, stomach, endometrium, testicular, ovary, central nervous system, skin, head and neck, esophagus, or bone marrow cancer; as well as hematological cancers, such as leukemia, acute promyelocytic leukemia, lymphoma, multiple myeloma, myelodysplasia, myeloproliferative disease, or refractory anemia.

The compounds of the present invention may also be used in combination-based therapeutic cancer treatments in a mammalian subject. Such methods may comprise administration of a compound of Formula (I) or (II) in combination with other adjunct cancer therapies, such as chemotherapy, radiotherapy, gene therapy, hormone therapy and other cancer therapies known in the art.


Any of the compounds of the present invention may be contemplated for administration to the mammalian subject in the form of a drug, prodrug or even active metabolite. In the methods of treatment of the present invention, the term "administering" shall encompass the treatment of the various conditions described with the compound specifically disclosed or with a compound which may not be specifically disclosed, but which converts to the specified compound *in vivo* after administration to the patient and exhibits therapeutic activity.

Another aspect of the present invention pertains to a method of inhibiting growth of cancer cells comprising providing to a patient a prodrug of Formula (III) wherein R₅ is a methyl or hydrogen;




Formula(III)


and forming metabolites of formula (IV), (V), (VI), (VII),
and(VIII) wherein any of R_3 , R_4 , R_5 , R_7 , R_8 of their Formula II
5 counterparts may be methyl or hydrogen. Such metabolites
could include for example the structures shown below:




(IV) A - NDC-1088




(IV) B - NDC-1099




10 (V) A - NDC-1033




(V) B - NDC-1022




(VI) A -NDC-1110




(VI) B - NDC-1044




(VII) A - NDC-1066



(VII) B - NDC-1055



Formula (VIII)-A



Formula (VIII)-B

Other objects, features, benefits and advantages of the present invention will be apparent from this summary and the following descriptions of certain embodiments, and will be readily apparent to those skilled in the art having knowledge of various chemotherapeutic compounds, methods and/or modes of operation.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1- shows the Estradiol biosynthetic pathway.

Figure 2- shows a predicted metabolic pathway for the present compounds.

5 Figure 3- shows the effect of NDC-1011, NDC-1022, NDC-1033, NDC-1044, NDC-1055 and NDC-1066 on estrogen receptor beta (ER- β) activity as measured by luciferase expression (RLU = relative light units). CV-1 cells were transfected with two plasmid constructs, the reporter construct ERE-tk-luciferase and a CMV-ER- β construct. Transfected control
10 (Ctrl) CV-1 cells received no treatment while estradiol treated cells (E2) received estradiol added alone at 10^{-9} M (1 nM). In the case of NDC compounds, each compound respectively was either added alone at 10^{-8} M (10 nM) (as
15 evident in the left column for each test compound) or at 10^{-8} M plus 10^{-9} M estradiol (E2) (as evident in the right column for each test compound).

Figure 4 shows the effect of NDC-1011, NDC-1033, NDC-1055 and NDC-1066 on estrogen receptor alpha (ER- α) activity
20 as measured by luciferase expression (RLU = relative light units). CV-1 cells were transfected with two plasmid constructs, the reporter construct ERE-tk-luciferase and a CMV-ER- α construct. Transfected control (Ctrl) CV-1 cells received no treatment while estradiol (E2) was added alone
25 at 10^{-9} M (1 nM). In the case of NDC compounds, each compound respectively was either added alone at 10^{-8} M (10 nM) (as evident in the left column for each test compound) or at 10^{-8} M plus 10^{-9} M estradiol (E2) (as evident in the right column for each test compound).

30 Figure 5 shows IC₅₀ growth inhibition data (in μ M) for NDC-1022 (left columns), NDC-1033 (middle columns) and NDC-1044 (right columns) as determined in each of the cell lines

HT-29, SK-OV-3, NCI-H23, MCF-7, MDA-MB-231, OVCAR-3, CAPAN-1, CAPAN-2, SH-SY5Y, A-549 and PC-3.

Figure 6 shows numerical IC₅₀ growth inhibition data (in μ M) for NDC-1022, NDC-1033 and NDC-1044 as determined in each of the cell lines HT-29, SK-OV-3, NCI-H23, MCF-7, MDA-MB-231, OVCAR-3, CAPAN-1, CAPAN-2, SH-SY5Y, A-549 and PC-3.

DETAILED DESCRIPTION OF THE INVENTION

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as is commonly understood by one of skill in the art to which this invention belongs and shall be understood to have the meanings described below. All publications and patents referred to herein are incorporated by reference in their entirety. Unless otherwise specified, a reference to a particular compound includes all such isomeric forms, including racemic and other mixtures thereof. Unless otherwise specified, a reference to a particular compound also includes ionic, salt, solvate (e.g., hydrate), protected forms, prodrugs, and other stereoisomers thereof, for example, as discussed herein.

It may be convenient or desirable to prepare, purify, and/or handle a corresponding salt of the active compound, for example, a pharmaceutically-acceptable salt. Examples of pharmaceutically acceptable salts are discussed in Berge et al., 1977, "Pharmaceutically Acceptable Salts," J. Pharm. Sci., Vol. 66, pp. 1-19, and discussed herein.

Anti-proliferative compounds of the present invention have application in the treatment of cancer, and so the present invention further provides anti-cancer agents. The term "anti-cancer agent" as used herein, pertains to a compound which treats a cancer (i.e., a compound which is useful in the treatment of a cancer). The anti-cancer effect

may arise through one or more mechanisms, including but not limited to, the regulation of cell proliferation, the inhibition of angiogenesis (the formation of new blood vessels), the inhibition of metastasis (the spread of a tumor from its origin), the inhibition of invasion (the spread of tumor cells into neighboring normal structures), or the promotion of apoptosis (programmed cell death).

The invention further provides active compounds for use in a method of treatment of the human or animal body by therapy. Such a method may comprise administering to such a subject a therapeutically-effective amount of an active compound, preferably in the form of a pharmaceutical composition as discussed further herein.

The term "treatment," or "therapy" as used herein in the context of treating a condition, pertains generally to treatment and therapy of a mammalian subject, whether of a human or a non-human animal (e.g., in veterinary applications), in which some desired therapeutic effect is achieved, for example, the inhibition of the progress of the condition, and includes a reduction in the rate of progress, a halt in the rate of progress, amelioration of the condition, and/or cure of the condition. Treatment as a prophylactic measure is also included. Treatment includes combination treatments and therapies, in which two or more treatments or therapies are combined, for example, sequentially or simultaneously. Examples of treatments and therapies include, but are not limited to, chemotherapy (the administration of active agents, including, e.g., drugs, antibodies (e.g., as in immunotherapy), prodrugs (e.g., employing protecting groups including phosphoric acid derivatives and phosphinates at suitable positions such as position 3 or 17, other compounds used for photodynamic

therapy, GDEPT, ADEPT, etc.); surgery; radiation therapy; and gene therapy.

The term "stereochemical isomer" as used herein, refers to isomers that differ from each other only in the way the
5 atoms are oriented in space. The two stereoisomers particularly of importance in the instant invention are enantiomers and diastereomers depending on whether or not the two isomers are mirror images of each other. In the preferred embodiment, the claimed formulations comprise such
10 compounds that isolated, resolved and are "substantially free of other isomers."

The term "therapeutically-effective amount," as used herein, pertains to that amount of an active compound, or a material, composition or dosage form comprising an active
15 compound, which is effective for producing some desired therapeutic effect, commensurate with a reasonable benefit/risk ratio.

The term "patient" refers to animals, including mammals, preferably humans.


20 The term "region of a patient" refers to a particular area or portion of the patient afflicted with a proliferative disorder, cancer or tumor and in some instances to regions throughout the entire patient. Exemplary of such regions are the pulmonary region, the
25 gastrointestinal region, the breast region, the renal region as well as other bodily regions, tissues, lymphocytes, receptors, organs and the like, including the vasculature and circulatory system, and cancerous tissue. "region of a patient" includes, for example, regions to be treated with
30 the disclosed compounds and compositions. The "region of a patient" is preferably internal, although it may be external.

The term "tissue" refers generally to specialized cells which may perform a particular function. The term "tissue" may refer to an individual cell or a plurality or aggregate of cells, for example, membranes, blood or organs. The term


5 "tissue" also includes reference to an abnormal cell or a plurality of abnormal cells. Exemplary tissues include breast tissue, including breast cells, membranous tissues, including endothelium and epithelium, laminae, connective tissue, including interstitial tissue, and tumors.

10 Compounds

The present invention is directed to a chemotherapeutic compound of a Formulas (I)-(II):



Formula (I)



Formula (II)

- 15 wherein R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈ and R₉ are independently hydrogen, C₁ to C₆ alkyl or substituted alkyl, halogen, sulfate, or glucuronide moieties; and the --- symbol represents either a single or a double bond and when the --- symbol is a double bond and forms a keto group at position
- 20 3 or 17, then no R₇ or R₆ is present, respectively; the symbol --- represent the presence or absence of a bond at position 10; and the ~~~~ symbol represents any type of bond regardless of the stereochemistry. The compounds also embrace the enantiomers, other stereochemical isomers,
- 25 hydrates, solvates, tautomers and pharmaceutically acceptable salts thereof.

In an embodiment of Formula I, R₅, R₆, R₇, R₉ are hydrogen atoms, methyl or Cl; R₁, R₂ is hydrogen or methyl; R₃, R₄ is C₁ to about C₆ alkyl or substituted alkyl, halogen and ---- is a single bond corresponding to the alcohol group. In an
 5 another embodiment the stereochemistry at the C-6 carbon comprises a S or R enantiomer or diastereomers.

Embodiment compounds of the present invention can be used in a pharmaceutical composition. Such a composition can comprise one or more compounds selected from those discussed
 10 above, illustrated below or otherwise inferred herein, and combinations thereof. In certain embodiments, such a composition can comprise a pharmaceutically-acceptable carrier component. Without limitation, such a composition can comprise a racemic mixture of compounds. In certain
 15 embodiments, such a compound can be present as the S and R enantiomer, preferably their isolated and purified form which is substantially free of other isomers, and R₅, or R₇ can be selected from H, C₁ to C₆ alkyl or substituted alkyl, and a halogen.

20 The compounds of the present invention may have asymmetric centers and may occur as racemate, racemic mixture or as individual diastereomers or enantiomers such as (S)6-methyloxymethyl(8S, 9S, 13S, 14S, 17S) -13-methyl-7,8,9,11,12,14,15,16,17 -decehydrocyclopenta [a]
 25 phenantherene-3, 17-diol; (R)6-methyloxymethyl(8S, 9S, 13S, 14S, 17R) -13-methyl-7,8,9,11,12,14,15,16,17 -decehydrocyclopenta [a] phenantherene-3, 17-diol; (R)6-methyloxymethyl(8R, 9S, 13R, 14S, 17R) -13-methyl-7,8,9,11,12,14,15,16,17-decehydrocyclopenta [a]
 30 phenantherene-3, 17-diol (NDC-1022); or (S)6-methyloxymethyl(8R, 9S, 13R, 14S, 17R) -13-methyl-

7,8,9,11,12,14,15,16,17-decehydrocyclopenta [a]
phenantherene-3, 17-diol (NDC-1033).

An embodiment of the present invention pertains to the preparation of the R or S enantiomers, R or S diastereomers
5 of 6 substituted estradiols. Methods for the preparation (e.g., asymmetric synthesis) and separation (e.g., fractional crystallization and chromatographic means) of such isomeric forms are either generally known in the art or are readily obtained by adapting the methods taught herein.
10 One such methodologies are described in the co-pending U.S. application SN 11/541,987, the teachings of which are herein incorporated in its entirety.


Another embodiment of present invention pertains to a method for preparing a 6-hydroxymethyl or 6-methyloxymethyl
15 derivative of estradiol. A reaction scheme for preparing estradiol derivatives is given below, Scheme 1. Such a method can comprise reaction of t-butyldimethylsilyl derivative of estradiol with LIDAKOR/THF/formaldehyde to obtain a 6-hydroxylated compound followed by such steps as:
20 (i) hydrolysis to obtain 6-hydroxymethyl derivative of estradiol, NDC-1066; and/or (ii) treatment with dimethylsulfate followed by hydrolysis to obtain 6-methyloxymethyl derivative of estradiol, NDC-1033. NDC-1088 can be obtained by further oxidation of NDC-1033 at the C-17
25 hydroxyl position.

In an alternative approach, the compounds of the present invention can also be prepared by a method comprising such steps as: (i) protecting an estrodial compound, (ii) acylating the protected estradiol compound at
30 the benzylic 6-position with LIDAKOR/Butyl-Lithium/Diisopropylamine/potassium tert-amylate, (iii) reducing the position 6 aldehyde with lithium aluminum


hydride, (iv) deprotecting the protected regions of the estrodial compound. A reaction scheme for preparing estradiol derivatives is given below in Scheme 2.

5 The compounds of the present invention can be synthesized by the following methods as depicted in the schemes below:

Scheme 1 - preparing compounds of the instant invention.



Scheme 2- Alternative methods for preparing the compounds of the invention.



5

Various methoxyalkyl derivatives, in accordance with this invention, are limited only by choice of alkylating agent, such derivatives as would be understood by those


skilled in art made aware of this invention, as available through synthetic procedures of the sort described herein or straight-forward modifications thereof, such modifications as would also be understood by those skilled in the art.

5 Accordingly, without limitation, various C₁ to C₆ alkyl and substituted alkyl (e.g., C₁ to C₆ linear, substituted linear, branched and substituted branched alkyl, such substituents as would be understood in the art) reagents can be used as described herein to prepare the corresponding methyloxyalkyl

10 derivatives.

The present invention relates to a method of treating cancer in a mammalian subject (e.g., a human patient). In this aspect of the invention, methods are provided for inhibiting tumor or cancerous cell growth. In such a

15 method, the cells are exposed to or contacted with a compound of Formula (I) or pharmaceutically acceptable salts or hydrates thereof:



Formula(I)

20 wherein R₁, R₂, R₃, R₄, R₅, R₆, R₇, and R₉ are independently hydrogen, C₁ to C₆ alkyl or substituted alkyl, halogen, sulfate, or glucuronide moieties; and the ---- symbol represents either a single or a double bond and when the --- symbol is a double bond and forms a keto group at position

25 3 or 17, then no R₇ or R₆ is present, respectively; the symbol --- represent the presence or absence of a bond at

position 10; and the  symbol represents any type of bond regardless of the stereochemistry. The compounds also embrace the enantiomers, other stereochemical isomers, hydrates, solvates, tautomers and pharmaceutically acceptable salts thereof.


These methods may be used to treat any tumor which may be either directly or indirectly effected by hormonal and/or estrogen-related activity, including but not in any way limited to solid tumors associated with breast, pancreatic, lung, colon, prostate, ovarian cancers, as well as brain, liver, spleen, kidney, lymph node, small intestine, blood cells, bone, stomach, endometrium, testicular, ovary, central nervous system, skin, head and neck, esophagus, or bone marrow cancer; as well as hematological cancers, such as leukemia, acute promyelocytic leukemia, lymphoma, multiple myeloma, myelodysplasia, myeloproliferative disease, or refractory anemia.

In addition, administration of the compounds of the present invention for treatment of various cancer states may comprise administration of a compound of Formula (I) in combination with other adjunct cancer therapies, such as chemotherapy, radiotherapy, gene therapy, hormone therapy and other cancer therapies known in the art. Combinations of the presently disclosed compounds with other anti-cancer or chemotherapeutic agents are within the scope of the invention. Examples of such agents can be found in *Cancer Principles and Practice of Oncology* by V. T. Devita and S. Hellman (editors), 6th edition (Feb. 15, 2001), Lippincott Williams & Wilkins Publishers. A physician, veterinarian or clinician of ordinary skill in the art would be able to discern which combinations of agents would be useful based on the particular characteristics of the drugs and the

cancer involved. Such anti-cancer agents include the following: estrogen receptor modulators, androgen receptor modulators, retinoid receptor modulators, cytotoxic agents, anti-proliferative agents, prenyl-protein transferase inhibitors, HMG-CoA reductase inhibitors, EHV protease inhibitors, reverse transcriptase inhibitors, aromatase inhibitors, and angiogenesis inhibitors.

In at least one aspect of the invention, inventors illustrate the compounds of the present invention in table I below:

Table I




R_1, R_2, R_3, R_4 : independently H, C₁-C₆ alkyl, substituted alkyl, or halogen
 R_5, R_7 : H, C₁-C₆ alkyl, substituted alkyl, sulfate, or glucuronide
 R_6 : H, C₁-C₆ alkyl, or substituted alkyl, sulfate, or glucuronide, when --- is a single bond; not present, when = is a double bond

Entry	Substituents			Spatial Configuration					
	R_5	R_6	R_7	C-6	C-8	C-9	C-13	C-14	C-17
1	H	H	H	S	S	S	S	S	S
2	H	H	H	S	R	R	R	R	R
3	H	H	H	S	S	S	S	S	R
4	H	H	H	S	R	R	R	R	S
5	H	—	H	S	S	S	S	S	C=O
6	H	—	H	S	R	R	R	R	C=O
7	H	H	H	R	R	R	R	R	R
8	H	H	H	R	S	S	S	S	S
9	H	H	H	R	S	S	S	S	R
10	H	H	H	R	R	R	R	R	S
11	H	—	H	R	S	S	S	S	C=O
12	H	—	H	R	R	R	R	R	C=O
13	Me	H	H	S	S	S	S	S	S
14	Me	H	H	S	R	R	R	R	R
15*	Me	H	H	S	R	S	R	S	R
16	Me	H	H	S	R	R	R	R	S
17	Me	—	H	S	S	S	S	S	C=O

18	Me	—	H	S	R	R	R	R	C=O
19	Me	H	H	R	R	R	R	R	R

Table I-continued




5

R_1, R_2, R_3, R_4 : independently H, C_1 - C_6 alkyl, substituted alkyl, or halogen
 R_5, R_7 : H, C_1 - C_6 alkyl, substituted alkyl, sulfate, or glucuronide
 R_6 : H, C_1 - C_6 alkyl, or substituted alkyl, sulfate, or glucuronide, when $---$ is a single bond; not present, when $---$ is a double bond

10

Entry	Substituents			Spatial Configuration					
	R_5	R_6	R_7	C-6	C-8	C-9	C-13	C-14	C-17
20**	Me	H	H	R	R	S	R	S	R
21	Me	H	H	R	S	S	S	S	R
22	Me	H	H	R	R	R	R	R	S
23	Me	—	H	R	S	S	S	S	C=O
24	Me	—	H	R	R	R	R	R	C=O
25	H	H	SO ₃ H	S	S	S	S	S	S
26	H	H	SO ₃ H	S	R	R	R	R	R
27	H	H	SO ₃ H	S	S	S	S	S	R
28	H	H	SO ₃ H	S	R	R	R	R	S
29	H	—	SO ₃ H	S	S	S	S	S	C=O
30	H	—	SO ₃ H	S	R	R	R	R	C=O
31	H	H	SO ₃ H	R	R	R	R	R	R
32	H	H	SO ₃ H	R	S	S	S	S	S
33	H	H	SO ₃ H	R	S	S	S	S	R
34	H	H	SO ₃ H	R	R	R	R	R	S
35	H	—	SO ₃ H	R	S	S	S	S	C=O
36	H	—	SO ₃ H	R	R	R	R	R	C=O
37	Me	H	SO ₃ H	S	S	S	S	S	S
38	Me	H	SO ₃ H	S	R	R	R	R	R
39	Me	H	SO ₃ H	S	S	S	S	S	R
40	Me	H	SO ₃ H	S	R	R	R	R	S
41	Me	—	SO ₃ H	S	S	S	S	S	C=O
42	Me	—	SO ₃ H	S	R	R	R	R	C=O
43	Me	H	SO ₃ H	R	R	R	R	R	R
44	Me	H	SO ₃ H	R	S	S	S	S	S
45	Me	H	SO ₃ H	R	S	S	S	S	R

Table I-continued



5

R₁, R₂, R₃, R₄: independently H, C₁-C₆ alkyl, substituted alkyl, or halogen
 R₅, R₇: H, C₁-C₆ alkyl, substituted alkyl, sulfate, or glucuronide
 R₆: H, C₁-C₆ alkyl, or substituted alkyl, sulfate, or glucuronide, when
 --- is a single bond; not present, when --- is a double bond

	Substituents			Spatial Configuration					
Entry	R ₅	R ₆	R ₇	C-6	C-8	C-9	C-13	C-14	C-17
46	Me	H	SO ₃ H	R	R	R	R	R	S
47	Me	–	SO ₃ H	R	S	S	S	S	C=O
48	Me	–	SO ₃ H	R	R	R	R	R	C=O
49	H	H	glucuronide	S	S	S	S	S	S
50	H	H	glucuronide	S	R	R	R	R	R
51	H	H	glucuronide	S	S	S	S	S	R
52	H	H	glucuronide	S	R	R	R	R	S
53	H	–	glucuronide	S	S	S	S	S	C=O
54	H	–	glucuronide	S	R	R	R	R	C=O
55	H	H	glucuronide	R	R	R	R	R	R
56	H	H	glucuronide	R	S	S	S	S	S
57	H	H	glucuronide	R	S	S	S	S	R
58	H	H	glucuronide	R	R	R	R	R	S
59	H	–	glucuronide	R	S	S	S	S	C=O
60	H	–	glucuronide	R	R	R	R	R	C=O
61	Me	H	glucuronide	S	S	S	S	S	S
62	Me	H	glucuronide	S	R	R	R	R	R
63	Me	H	glucuronide	S	S	S	S	S	R
64	Me	H	glucuronide	S	R	R	R	R	S
65	Me	–	glucuronide	S	S	S	S	S	C=O
66	Me	–	glucuronide	S	R	R	R	R	C=O
67	Me	H	glucuronide	R	R	R	R	R	R
68	Me	H	glucuronide	R	S	S	S	S	S
69	Me	H	glucuronide	R	S	S	S	S	R
70	Me	H	glucuronide	R	R	R	R	R	S
71	Me	–	glucuronide	R	S	S	S	S	C=O
72	Me	–	glucuronide	R	R	R	R	R	C=O


10

* NDC-1033; ** NDC-1022

The preferred compounds in Table I include compounds 15 and 20. At least one aspect of the instant invention is directed to these preferred compound, their method of use and making.

In at least another aspect of the invention, inventors
5 illustrate the compounds of the present invention in table II below:


Table II



R₁, R₂, R₃, R₄: independently H, C₁-C₆ alkyl, substituted alkyl, or halogen
 R₅: H, C₁-C₆ alkyl, substituted alkyl, sulfate, or glucuronide
 R₆: H, C₁-C₆ alkyl, or substituted alkyl, sulfate, or glucuronide,
 R₈: H, C₁-C₆ alkyl, or substituted alkyl,
 when --- is a single bond; not present, when = is a double bond

Entry	Substituent		Spatial Configuration						
	R ₅	R ₆	C-6	C-8	C-9	C-10	C-13	C-14	C-17
73	H	H	S	S	S	R	S	S	S
74	H	H	S	R	R	R	R	R	R
75	H	H	S	S	S	R	S	S	R
76	H	H	S	R	R	R	R	R	S
77	H	—	S	S	S	R	S	S	C=O
78	H	—	S	R	R	R	R	R	C=O
79	H	H	R	R	R	R	R	R	R
80	H	H	R	S	S	R	S	S	S
81	H	H	R	S	S	R	S	S	R
82	H	H	R	R	R	R	R	R	S
83	H	—	R	S	S	R	S	S	C=O
84	H	—	R	R	R	R	R	R	C=O
85	Me	H	S	S	S	R	S	S	S
86	Me	H	S	R	R	R	R	R	R
87	Me	H	S	S	S	R	S	S	R
88	Me	H	S	R	R	R	R	R	S
89*	Me	—	S	R	S	R	R	S	C=O
90**	Me	—	R	R	S	R	R	S	C=O
91	Me	H	R	R	R	R	R	R	R

Table II



5


R_1, R_2, R_3, R_4 : independently H, C_1 - C_6 alkyl, substituted alkyl, or halogen
 R_5 : H, C_1 - C_6 alkyl, substituted alkyl, sulfate, or glucuronide
 R_6 : H, C_1 - C_6 alkyl, or substituted alkyl, sulfate, or glucuronide,
 R_8 : H, C_1 - C_6 alkyl, or substituted alkyl,
 when --- is a single bond; not present, when --- is a double bond

Entry	Substituent		Spatial Configuration						
	R_5	R_6	C-6	C-8	C-9	C-10	C-13	C-14	C-17
92	Me	H	S	S	S	R	S	S	S
93	Me	H	S	R	R	R	R	R	R
94	Me	H	S	S	S	R	S	S	R
95	Me	H	S	R	R	R	R	R	S
96	Me	—	R	R	S	—	R	S	C=O
97	Me	—	S	R	R	R	R	R	C=O
98***	Me	H	S	R	S	R	R	S	R
99	Me	H	R	S	S	R	S	S	S
100	Me	H	R	S	S	S	S	S	R
101	Me	H	R	R	R	S	R	R	S
102	Me	—	R	S	S	R	S	S	C=O
103	Me	—	R	R	R	R	R	R	C=O
104	H	H	S	S	S	R	S	S	S
105	H	H	S	R	R	R	R	R	R
106	H	H	S	S	S	R	S	S	R
107	H	H	S	R	R	R	R	R	S
108	H	—	S	S	S	R	S	S	C=O
109	H	—	S	R	R	R	R	R	C=O
110	H	H	R	R	R	R	R	R	R
111	H	H	R	S	S	R	S	S	S
112	H	H	R	S	S	R	S	S	R
113	H	H	R	R	R	R	R	R	S
114	H	—	R	S	S	R	S	S	C=O
115	H	—	R	R	R	R	R	R	C=O

10

15

Table II-Continued



5

R_1, R_2, R_3, R_4 : independently H, C_1-C_6 alkyl, substituted alkyl, or halogen
 R_5 : H, C_1-C_6 alkyl, substituted alkyl, sulfate, or glucuronide
 R_6 : H, C_1-C_6 alkyl, or substituted alkyl, sulfate, or glucuronide,
 R_8 : H, C_1-C_6 alkyl, or substituted alkyl,
 when --- is a single bond; not present, when --- is a double bond


Entry	Substituent		Spatial Configuration						
	R_5	R_6	C-6	C-8	C-9	C-10	C-13	C-14	C-17
116	Me	H	S	S	S	R	S	S	S
117	Me	H	S	R	R	R	R	R	R
118	Me	H	S	S	S	R	S	S	R
119	Me	H	S	R	R	R	R	R	S
120	Me	—	S	S	S	S	S	S	C=O
121	Me	—	S	R	R	R	R	R	C=O
122	Me	H	R	R	R	R	R	R	R
123	Me	H	R	S	S	R	S	S	S
124	Me	H	R	S	S	S	S	S	R
125	Me	H	R	R	R	S	R	R	S
126	Me	—	R	S	S	R	S	S	C=O
127	Me	—	R	R	R	R	R	R	C=O
128****	Me	H	R	R	S	R	R	S	R

10


* NDC-1077, when R_8 is a methyl; ** NDC-1011, when R_8 is a methyl; *** NDC-1110,
 when R_8 is a methyl; **** NDC-1044, when R_8 is a methyl.

The preferred compounds in Table II include compounds
 89, 90, 98, and 128. At least one aspect of the instant
 invention is directed to these preferred compound, their
 method of use and making.

One specific non-limiting example for treatment of an
 identified cancer state as described herein includes use a
 compound of Formula (I), which has the Formula (IX) below:




(IX) A- NDC -1033




(IX) B- NDC - 1022

Another specific non-limiting example for treatment of an identified cancer state as described herein includes use
 5 a compound of Formula (I), which has the Formula (X), below:



(X) A- NDC -1066



(X) B- NDC -1055

The above active compounds may also be used as part of an *in vitro* assay, for example, in order to determine
 10 whether a candidate host is likely to benefit from treatment with the compound in question. Any active compound of the present invention may also be used as a standard, for example, in an assay, in order to identify other active compounds, other anti-proliferative agents, other anti-
 15 inflammatory agents, etc.

At least in one aspect of the instant invention, the candidate compounds were evaluated for their estrogen receptor antagonistic activity. The evaluation as to whether a compound is an estrogen receptor antagonist may be carried
 20 out by various methodologies known in the art. In the instant application, such capacity was determined by

conducting the Luciferase binding assay according to the screening methods described herein.

In a more preferred embodiment of this aspect of the invention, the estrogen receptor binding capacity were
5 assessed by transiently transfecting CV-1 cells with expression constructs for either ER(α) or ER (β) plus an ERE-tk-luciferase reporter construct. The cells were then divided into controls and candidate groups wherein the controls received no treatment, or were treated with
10 estradiol alone (1 nM) and the candidate groups received estradiol plus an Endece compound at varying concentrations. After 16-24 hours the cells were harvested and assayed for luciferase activity using a commercially available assay kit.

15 In yet another aspect of the instant invention, the IC_{50} or the half maximal inhibitory concentration of the candidate compounds were determined to assess drug potency and potential dosing regimens for in vivo use. One of ordinary skill in the art is readily able to ascertain such
20 information using commonly known methodologies. As it has been well described in the art, IC_{50} represents and measures how much of a particular substance/molecule is needed to inhibit some biological process by 50%. In the instant case, the IC_{50} of the candidate compounds were determined as the
25 concentration that led to a response of 50% compared to the vehicle control cells.

As noted herein, the salts of the compounds of this invention refer to non-toxic "pharmaceutically acceptable salts." Other salts may, however, be useful in the
30 preparation of the compounds according to the invention or of their pharmaceutically acceptable salts. When the compounds of the present invention contain a basic group,

salts encompassed within the term "pharmaceutically acceptable salts" refer to non-toxic salts which are generally prepared by reacting the free base with a suitable organic or inorganic acid. Representative salts include any
5 such salt known in the art. Where compounds of the present invention carry an acidic moiety, suitable pharmaceutically acceptable salts thereof may include alkali metal salts, e.g., sodium or potassium salts; alkaline earth metal salts, e.g., calcium or magnesium salts; and salts formed with
10 suitable organic ligands, e.g., quaternary ammonium salts.

To treat a mammalian subject, such as a human patient, an effective amount of one or more compounds of the present invention, or a pharmaceutically-acceptable salt thereof, is administered to the mammalian subject so as to promote
15 exposure to or contact of cancer cells or the targeted tumor growth. Effective dosage forms, modes of administration and dosage amounts may be determined empirically, and making such determinations is within the skill of the art. It is understood by the physician, veterinarian or clinician of
20 ordinary skill in the art that the dosage amount will vary with the activity of the particular compound employed, course and/or progression of the disease state, the route of administration, the rate of excretion of the compound, renal and hepatic function of the patient, the duration of the
25 treatment, the identity of any other drugs being administered to the subject, age, size and like factors well known in the medical arts. As discussed herein, the compounds of the present invention can be administered in such oral dosage forms as tablets, capsules (each of which
30 includes sustained release or timed release formulations), pills, powders, micronized compositions, granules, elixirs, tinctures, suspensions, syrups and emulsions. Likewise, they

may also be administered in intravenous (bolus or infusion), intraperitoneal, topical (e.g., ocular eyedrop), subcutaneous, intramuscular or transdermal (e.g., patch) form, all using forms well known to those of ordinary skill
5 in the pharmaceutical arts. Again, the ordinarily skilled physician, veterinarian or clinician can readily determine and prescribe the effective amount of the drug required to prevent, counter or arrest the progress of the condition.

Oral dosages of the present invention, when used for
10 the indicated effects, will range between about 0.01 mg per kg of body weight per day (mg/kg/day) to about 100 mg/kg/day, preferably 0.01 to 10 mg/kg/day, and most preferably 0.1 to 5.0 mg/kg/day. For oral administration, the compositions are preferably provided in the form of
15 tablets containing 0.01, 0.05, 0.1, 0.5, 1.0, 2.5, 5.0, 10.0, 15.0, 25.0, 50.0, 100 and 500 milligrams of the active ingredient for the symptomatic adjustment of the dosage to the patient to be treated. A medicament typically contains from about 0.01 mg to about 500 mg of the active ingredient,
20 preferably, from about 1 mg to about 100 mg of active ingredient. Intravenously, the most preferred doses will range from about 0.1 to about 10 mg/kg/minute during a constant rate infusion. Compounds of the present invention may be administered in a single daily dose, or the total
25 daily dosage may be administered in divided doses of two, three or four times daily.

As noted herein, the compounds of the present invention can be used in combination with other anti-cancer agents or other agents which will enhance the treatment
30 regime for the mammalian subject. The individual components of such combinations can be administered separately at different times during the course of therapy or concurrently


in divided or single combination forms to patients or regions of such patients in need of such therapy. The instant invention is therefore to be understood as embracing all such regimes of simultaneous or alternating treatment and the term "administering" is to be interpreted accordingly. It will be understood that the scope of combinations of the compounds of this invention with other agents useful to treat the targeted cancer condition includes in principle any combination with any pharmaceutical composition useful for treating disorders related to estrogen functioning.

As used herein, the term "composition" is intended to encompass a product comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from combination of the specified ingredients in the specified amounts.


It may be convenient or desirable to prepare, purify, and/or handle the active compound in the form of a prodrug. The term "prodrug" as used herein, pertains to a compound which, when metabolized, yields the desired active compound or in itself is the active compound. This includes for example adding a phosphoric acid ester moiety in suitable positions such as positions 3, 6, 10 or 17. Typically, the prodrug is inactive, or less active than the active compound, but may provide advantageous handling, administration, or metabolic properties. For example, some prodrugs are ethers of the active compound; during metabolism the ether group is cleaved to yield the active drug. Also, some prodrugs are activated enzymatically to yield the active compound, or a compound which, upon further chemical reaction, yields the active compound. Thus, in the methods of treatment of the present invention disclosed

herein, the term "administering" shall encompass the treatment of the various conditions described with the compound specifically disclosed or with a compound which may not be specifically disclosed, but which converts to the specified compound *in vivo* after administration to the patient. Metabolites of these compounds include active species produced upon introduction of compounds of this invention into the mammalian subject.

Compounds of the present invention may be prodrugs for potent anti-proliferative agents. Compounds which exhibit low or moderate intrinsic activity may act as prodrugs, and be metabolically activated (e.g., *in vivo*) to generate more potent compounds. This is especially useful in cancer therapy where metabolic activation can be achieved by an enzyme that is expressed in tumors. Prodrugs, acting as a substrate, may be metabolized by CYP19, 17 β -HSD, HS-demethylase or another steroidal linked enzyme to generate a potent anti-cancer agent. The (R) or (S)-6-methyloxoalkyl derivatives of exemestane suggest that it may be active against numerous forms of cancer beyond breast cancer. Activity in inhibiting tumor cell growth in cells lines derived from breast, lung, colon, prostate, endometrial and ovarian cancers was observed for the NDC-1011 enantiomer. For example, *in vitro* studies of tumor cell growth is the highest in cell lines that are CYP19 positive (MDA-MB-213 and SK-OV-3) and is reduced in those cell lines that are CYP19 negative (MCF-7 and NIH:OVCA-3) indicates that NDC-1011 may act as a pro-drug. See Formula (XI) and (XII) below for the structure of NDC-1077 and NDC-1011.



(XI) NDC-1077



(XII) NDC-1011

While not bound by any theory, for example, if NDC-1011 is a pro-drug, then any number of the body's normal steroidogenic enzymes should be active towards the NDC-1011 compound thereby converting NDC-1011 into the active metabolite(s). This aspect of the invention can apply in the same manner to both the S and the R diastereomers.


The prodrug compounds of the instant invention act in a manner analogous to that observed for endogenous androstenedione, NDC-1011 is converted to an aromatic ring by hydroxylation at the C-3 carbon of NDC-1011 via CYP19 to give rise to the metabolite NDC-1099. see Figure 1 in the Estradiol biosynthetic pathway wherein Androstendione is converted to the Estrone via CYP19. NDC-1099 could undergo further hydroxylation at the C-17 carbon via the reversible action of 17β -hydroxysteroid dehydrogenase (17β -HSD) resulting in the diol compound NDC-1022.

As with Estradiol, the diol compound NDC-1022 has an aromatic ring, but differs from estradiol with respect to the methoxyalkyl substituent at the C-6 carbon. The metabolism of NDC-1011 into the diol compound NDC-1022 could occur in any order as shown in Figure 2. For example, NDC-1044 formed by 17β -HSD is converted to the NDC-1022 diol by CYP19 aromatization activity.

Without being bound to any theories, it has been reported that Estradiol binds to the receptor ligand pocket

of estrogen receptors (both ER α and ER β), via the C17-OH (via His 524); and the C3-OH (via Arg 394 and Glu 353). As with Estradiol, binding of NDC-1022 diol in the same ligand pocket of ER α and ER β via similar amino acid bindings may occur. Additionally, the presence of the methyloxyalkyl substituent at the C-6 carbon of compound NDC-1022 may alter the conformation of the normal ligand-bound receptor resulting in modified activity accounting for the observed anti-tumor activity.

In addition, demethylase enzyme activity directed at the C-6 methyl group of NDC-1011 (or one of the metabolites of NDC-1011), may indicate the formation of triol metabolite NDC-1055. With alcohol groups at the C-3, C-6 and C-17 carbons, such an NDC-1055 triol metabolite may bind to a broad spectrum of steroid receptors in a range of tissues involving various combinations of the C-3, C-6 and C-17 alcohols. One example of such metabolites includes the compound of Formula (VII)B as shown below:



(VII)B -NDC-1055

Pharmaceutical formulations of the present invention include those suitable for oral, nasal, topical (including buccal and sublingual), rectal, vaginal and/or parenteral administration. Regardless of the route of administration selected, the active ingredient(s) are formulated into

pharmaceutically-acceptable dosage forms by conventional methods known to those of skill in the art.

The amount of the active ingredient(s) which will be combined with a carrier material to produce a single dosage
5 form will vary depending upon the host being treated, the particular mode of administration and all of the other factors described above. The amount of the active ingredient(s) which will be combined with a carrier material to produce a single dosage form will generally be that
10 amount of the active ingredient(s) which is the lowest dose effective to produce a therapeutic effect.

Methods of preparing pharmaceutical formulations or compositions include the step of bringing the active ingredient(s) into association with the carrier and,
15 optionally, one or more accessory ingredients. In general, the formulations are prepared by uniformly and intimately bringing the active ingredient(s) into association with liquid carriers, or finely divided solid carriers, or both, and then, if necessary, shaping the product.

20 Formulations of the invention suitable for oral administration may be in the form of capsules, cachets, pills, tablets, lozenges (using a flavored basis, usually sucrose and acacia or tragacanth), powders, granules, or as a solution or a suspension in an aqueous or nonaqueous
25 liquid, or as an oil-in-water or water-in-oil liquid emulsion, or as an elixir or syrup, or as pastilles (using an inert base, such as gelatin and glycerin, or sucrose and acacia) and/or as mouth washes and the like, each containing a predetermined amount of the active ingredient(s). The
30 active ingredient(s) may also be administered as a bolus, electuary or paste.

In solid dosage forms of the invention for oral administration (capsules, tablets, pills, dragees, powders, granules and the like), the prodrug(s), active ingredient(s) (in their micronized form) is/are mixed with one or more
5 pharmaceutically-acceptable carriers, such as sodium citrate or dicalcium phosphate, and/or any of the following: (1) fillers or extenders, such as starches, lactose, sucrose, glucose, mannitol, and/or silicic acid; (2) binders, such as, for example, carboxymethyl-cellulose, alginates, gelatin, polyvinyl pyrrolidone, sucrose and/or acacia; (3)
10 humectants, such as glycerol; (4) disintegrating agents, such as agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate; (5) solution retarding agents, such as paraffin;
15 (6) absorption accelerators, such as quaternary ammonium compounds; (7) wetting agents, such as, for example, cetyl alcohol and glycerol monostearate; (8) absorbents, such as kaolin and bentonite clay; (9) lubricants, such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, and mixtures thereof; and
20 (10) coloring agents. In the case of capsules, tablets and pills, the pharmaceutical compositions may also comprise buffering agents. Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugars, as
25 well as high molecular weight polyethylene glycols and the like.

A tablet may be made by compression or molding, optionally with one or more accessory ingredients.
30 Compressed tablets may be prepared using binder (for example, gelatin or hydroxypropylmethyl cellulose), lubricant, inert diluent, preservative, disintegrant (for

example, sodium starch glycolate or cross-linked sodium carboxymethyl cellulose), surface-active or dispersing agent. Molded tablets may be made by molding in a suitable machine a mixture of the powdered active ingredient(s) moistened with an inert liquid diluent.

The tablets, and other solid dosage forms of the pharmaceutical compositions of the present invention, such as dragees, capsules, pills and granules, may optionally be scored or prepared with coatings and shells, such as enteric coatings and other coatings well known in the pharmaceutical-formulating art. They may also be formulated so as to provide slow or controlled release of the active ingredient(s) therein using, for example, hydroxypropylmethyl cellulose in varying proportions to provide the desired release profile, other polymer matrices, liposomes and/or microspheres. They may be sterilized by, for example, filtration through a bacteria-retaining filter. These compositions may also optionally contain opacifying agents and may be of a composition that they release the active ingredient(s) only, or preferentially, in a certain portion of the gastrointestinal tract, optionally, in a delayed manner. Examples of embedding compositions which can be used include polymeric substances and waxes. The active ingredient(s) can also be in microencapsulated form.

Liquid dosage forms for oral administration of the active ingredient(s) include pharmaceutically-acceptable emulsions, microemulsions, solutions, suspensions, syrups and elixirs. In addition to the active ingredient(s), the liquid dosage forms may contain inert diluents commonly used in the art, such as, for example, water or other solvents, solubilizing agents and emulsifiers, such as ethyl alcohol, isopropyl alcohol, ethylacetate, butyl alcohol, benzyl

benzoate, propylene glycol, glycol, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor and sesame oils), glycerol, amyl alcohol, tetrahydrofuryl polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof.

Besides inert diluents the oral compositions can also include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, coloring, perfuming and preservative agents. Suspensions, in addition to the active ingredient(s), may contain suspending agents as, for example, ethoxylated alcohols, polyoxyethylene sorbitol and sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar and tragacanth, and mixtures thereof.

Formulations of the pharmaceutical compositions of the invention for rectal or vaginal administration may be presented as a suppository, which may be prepared by mixing the active ingredient(s) with one or more suitable nonirritating excipients or carriers comprising, for example, cocoa butter, polyethylene glycol, wax or salicylate and which is solid at room temperature, but liquid at body temperature and, therefore, will melt in the rectum or vaginal cavity and release the active ingredient(s). Formulations of the present invention which are suitable for vaginal administration also include pessaries, tampons, creams, gels, pastes, foams or spray formulations containing such carriers as are known in the art to be appropriate.

Dosage forms for the topical or transdermal administration of the active ingredient(s) include powders, sprays, ointments, pastes, creams, lotions, gels, solutions, patches and inhalants. The active ingredient(s) may be mixed

under sterile conditions with pharmaceutically-acceptable carrier, and with any buffers, or propellants which may be required.

The ointments, pastes, creams and gels may contain, in
5 addition to the active ingredient(s), excipients, such as animal and vegetable fats, oils, waxes, paraffins, starch, tragacanth, cellulose derivatives, polyethylene glycols, silicones, bentonites, silicic acid, talc and zinc oxide, or mixtures thereof. Powders and sprays can contain, in
10 addition to the active ingredient(s), excipients such as lactose, talc, silicic acid, aluminum hydroxide, calcium silicates and polyamide powder, or mixtures of these substances. Sprays can additionally contain customary propellants such as chlorofluorohydrocarbons and volatile
15 unsubstituted hydrocarbons, such as butane and propane.

Compounds of the present invention may be administered in intranasal form via topical use of suitable intranasal vehicles, or via transdermal routes, using those forms of transdermal skin patches well known to those of ordinary
20 skill in the art. A transdermal delivery system provides for continuous administration throughout the dosage regimen. Transdermal patches have the added advantage of providing controlled delivery of the active ingredient(s) to the body. Such dosage forms can be made by dissolving, dispersing or
25 otherwise incorporating the active ingredient(s) in a proper medium, such as an elastomeric matrix material. Absorption enhancers can also be used to increase the flux of the active ingredient(s) across the skin. The rate of such flux can be controlled by either providing a rate-controlling
30 membrane or dispersing the active ingredient(s) in a polymer matrix or gel.

The compounds of the present invention can also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine or phosphatidylcholines.

Another mode of delivery for the compounds of the present invention may be delivery via the use of monoclonal antibodies as individual carriers to which the compound molecules are coupled. The compounds of the present invention may also be coupled with soluble polymers as targetable drug carriers. Such polymers can include polyvinylpyrrolidone, pyran copolymer, polyhydroxypropylmethacrylamide-phenol, polyhydroxyethylaspartamide-phenol, or polyethyleneoxide-polylysine substituted with palmitoyl residues. Furthermore, the compounds of the present invention may be coupled to a class of biodegradable polymers useful in achieving controlled release of a drug, for example, polylactic acid, polyglycolic acid, copolymers of polyactic and polyglycolic acid, polyepsilon caprolactone, polyhydroxy butyric acid, polyorthoesters, polyacetals, polydihydropyrans, polycyanoacrylates and crosslinked or amphipathic block copolymers of hydrogels.

Pharmaceutical compositions of this invention suitable for parenteral administration comprise the active ingredient(s) in combination with one or more pharmaceutically-acceptable sterile isotonic aqueous or nonaqueous solutions, suspensions or emulsions, or sterile powders which may be reconstituted into sterile injectable solutions or dispersions just prior to use, which may contain antioxidants, buffers, solutes which render the

formulation isotonic with the blood of the intended recipient or suspending or thickening agents.

Examples of suitable aqueous and nonaqueous carriers which may be employed in the pharmaceutical compositions of the invention include water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), and suitable mixtures thereof, vegetable oils, such as olive oil, and injectable organic esters, such as ethyl oleate. Proper fluidity can be maintained, for example, by the use of coating materials, such as lecithin, by the maintenance of the required particle size, and by the use of surfactants.

These compositions may also contain adjuvants such as wetting agents, emulsifying agents and dispersing agents. It may also be desirable to include isotonic agents, such as sugars, sodium chloride, and the like in the compositions. In addition, prolonged absorption of the injectable pharmaceutical form may be brought about by the inclusion of agents which delay absorption such as aluminum monostearate and gelatin.

In some cases, in order to prolong the effect of the active ingredient(s), it is desirable to slow the absorption of the drug from subcutaneous or intramuscular injection. This may be accomplished by the use of a liquid suspension of crystalline or amorphous material having poor water solubility. The rate of absorption of the active ingredient(s) then depends upon its/their rate of dissolution which, in turn, may depend upon crystal size and crystalline form. Alternatively, delayed absorption of parenterally-administered active ingredient(s) is accomplished by dissolving or suspending the active ingredient(s) in an oil vehicle.

Injectable depot forms are made by forming microencapsule matrices of the active ingredient(s) in biodegradable polymers such as polylactide-polyglycolide. Depending on the ratio of the active ingredient(s) to polymer, and the nature of the particular polymer employed, the rate of release of the active ingredient(s) can be controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides). Depot injectable formulations are also prepared by entrapping the active ingredient(s) in liposomes or microemulsions which are compatible with body tissue. The injectable materials can be sterilized for example, by filtration through a bacterial-retaining filter.

The formulations may be presented in unit-dose or multi-dose sealed containers, for example, ampoules and vials, and may be stored in a lyophilized condition requiring only the addition of the sterile liquid carrier, for example water for injection, immediately prior to use. Extemporaneous injection solutions and suspensions may be prepared from sterile powders, granules and tablets of the type described above.

The pharmaceutical compositions of the present invention may also be used in the form of veterinary formulations, including those adapted for the following: (1) oral administration, for example, drenches (aqueous or nonaqueous solutions or suspensions), tablets, boluses, powders, granules or pellets for admixture with feed stuffs, pastes for application to the tongue; (2) parenteral administration, for "ampule, by subcutaneous, intramuscular or intravenous injection as, for example, a sterile solution or suspension or, when appropriate, by intramammary injection where a suspension or solution is introduced into

the udder of the animal via its teat; (3) topical application, for example, as a cream, ointment or spray applied to the skin; or (4) intravaginally, for example, as a pessary, cream or foam or any other methods fit to by
5 those of ordinary skill in the art for administration to a region of interest.

Although the present invention has been described with reference to certain embodiments, one skilled in the art will appreciate that the present invention can be practiced
10 by other than the described embodiments, which have been presented for purposes of illustration and not of limitation. Therefore, the scope of the appended claims should not be limited to the description of the embodiments contained herein.

15 The general methods given in Schemes 1 and 2 for the preparation of compounds exemplified in Tables I and II above of the present invention are further illustrated by the following examples. Specifically, the methods given in Schemes 1 and 2 for the preparation of 6-alkoxyalkyl
20 Estradiol compounds are illustrated by Examples 1-5 shown below. An example of assessing the estrogen receptor binding capacity is articulated in example 4, and finally assessing the IC_{50} of the preferred compounds of the instant invention and their comparative efficacy is given in example 5. Unless
25 otherwise specified all starting materials and reagents are of standard commercial grade, and are used without further purification, or are readily prepared from such materials by routine methods. Those skilled in the art of organic synthesis will recognize that starting materials and
30 reaction conditions may be varied to achieve the desired end product.

Example 1

Methods of Preparing 6-hydroxymethyl-androsta-1,4-diene-3,17 dione.

In a reaction system, sufficient amounts of (+)androsta-1,4-diene-3,17-dione (ADD), 12.2 equivalents
5 pyrrolidine, catalytic acetic acid, denatured ethanol (95/5 ethanol/methanol) and 6-7% tetrahydrofuran (THF) were heated to 30 to 40° C for a minimum of 16 hours to form 1,3 - dipyrrolidinoandrosta-3,5-diene-17one. Once the ADD content reaches to a less than 3% by HPLC area, or it becomes static
10 or the resulting dipyrrolidinoandrostadiene begins to revert to ADD, the reaction mixture is cooled to $5 \pm 5^\circ$ C. The resulting compound is then collected and washed with cold denatured ethanol. Yields are typically 70-80% on a dry basis with purities typically 90-95% by HPLC area percent.

15 The resulting 1,3-dipyrrolidinoandrosta-3,5-diene-17one is then mixed in amount of 1 equivalent, with 2.6 equivalents formalin (formaldehyde) in 10 ml dichloromethane/g at room temperature. The reaction mixture is then acidified to a pH of about 2 with 2% sulfuric acid
20 solution. Accordingly, an organic layer is formed, which is washed with 2% sulfuric acid and 1:1 water/brine. Solvent exchange into toluene (approximately 10ml/g) is then carried out wherein the product crystallizes as toluene exchange transpires. Said product is collected washed and dried to
25 provide 6-hydroxymethyl-androsta-1,4-diene-3,17 dione. One of ordinary skill in the art can further modify the stereochemistry at position 6, if so desired by employing known techniques in the art.

Example 2

30 Methods of preparing compound NDC-1022 and NDC-1033

As outlined in Scheme 2, estradiol derivatives NDC-1022, NDC-1033 were synthesized in the following manner.

The protected estradiol compound 2 is prepared by reaction of compound 1 with dihydropyran in THF, using toluenesulfonic acid or camphorsulfonic acid as catalyst. As one of ordinary skill in the art can appreciate this
5 reaction is an equilibrium reaction and would not go to completion under such conditions. Thus, both the mono-protected estradiols can be found in the reaction mixture. Such crude reaction mixture would undergo a trituration step with acetonitrile causing the desired bis-THP estradiol to
10 crystallize in approximately 70 % yield.

As shown in Scheme 2, the key intermediate compound 3 is obtained via acylation at the benzylic 6-position with the strong base mixture referred to as LiDAKOR: butyl lithium, diisopropylamine, and potassium tert-amylate.
15 Under such conditions at -70 °C, one of ordinary skill in the art can appreciate the abstraction of a proton at a benzylic position. The intermediate compound 3 is then purified by column chromatography to give a syrup in approximately 50 % yield, still containing minor impurities
20 and column solvents. Reduction of the aldehyde with an excess of lithium aluminum hydride results in high yields of the racemic hydroxymethyl estradiol compound 4 as a glassy foam.

For purposes of preparing NDC-1022 and NDC 1033, the
25 methoxymethyl intermediate compound 7 was prepared by methylation of compound 4 with sodium hydride and methyl iodide. Compound 7 was purified by column chromatography to give a glassy foam. Deprotecting the protected groups would give racemic 6-methoxymethyl estradiol compound 8.
30 Separation of the enantiomers was performed using chiral preparative HPLC to give the compounds NDC-1022 and NDC-1033. For compound NDC-1022, a chiral purity of >95:5 R:S

was realized. For compound NDC-1033, a chiral purity of 86:14 *S*:*R* was realized. It is well within the level of one of ordinary skill in the art to employ NMR for determination of the absolute stereochemistry of the 6-position, where the
5 4-and 6-protons are diagnostic.

Example 3

Methods of preparing NDC-1055 and NDC-1066

Using the same methodologies described in Example 2,
10 compound 4 is synthesized. Deprotection of compound 4 was then achieved with catalytic hydrogen chloride in methanol, and racemic compound 5 was separated on chiral preparative HPLC to give two fractions, one enriched for NDC-1055 and the other enriched for NDC-1066. For each compound, chiral
15 purity of >95:5 *R*:*S* and *S*:*R* was realized respectively. Absolute stereochemistry of the 6-position was established by NMR, where the 4-and 6-protons are diagnostic.

Example 4

Methods of determining estrogen receptor binding
20 capacity using Luciferase activity.

Estrogen receptor-negative CV-1 kidney cells were maintained in Dulbecco's modified Eagle's medium with 4.5 g/L glucose supplemented with 10% fetal bovine serum and 100 units / ml penicillin-streptomycin at 37° C in a humidified
25 5% CO₂ atmosphere.

The cells were then plated in 6-well dishes at a density of 2×10^5 cells per well in phenol-red free Dulbecco's modified Eagle's medium containing 10% charcoal-dextran-stripped fetal bovine serum. CV-1 cells were
30 transfected using LipofectAMINE reagent according to the manufacturer's protocol. Transfections containing 1.5 ug of reporter plasmid (containing ERE-tk-luciferase containing a

single ERE cloned upstream of the thymidine kinase promoter and luciferase gene) and 0.5 ug of either ER α or ER β expression vector (containing CMV-ER α or CMV-ER β full length coding sequence respectively).

5 The next day, cells received no treatment (controls) or were treated with estradiol alone (1 nM) or estradiol plus an Endece compound (at varying concentrations). After 16-24 hours, cells were harvested and assayed for luciferase activity.

10 At the outset, cell monolayers were washed twice with ice-cold phosphate-buffered saline and incubated for 15 minutes in 250 ul of 1X cell culture lysis reagent (Promega, Madison, WI). Cell extracts were transferred to a fresh tube and assayed using the luciferase assay system
15 (Promega). For each assay, 10 ul of extract was diluted with 90 ul of 1X cell culture lysis reagent. Luminescence was read using an AutoLumat LB953 luminometer.

A compound or a salt thereof, which is identified by the binding assay described herein is a compound that
20 inhibits the binding of estrodial at the ligand binding site of the estrogen receptors. Specifically, it is a compound or a salt thereof that is envisioned to cause cell proliferation stasis and accordingly exerts its pharmacological activity. As indicated in Figure 3 and 4,
25 lead compounds NDC-1022, NDC-1033 exhibit strong competitive behavior against estradiol in binding to either of the estrogen receptors thereby causing stasis of cell proliferation activity.

Example 5

30 Method of determining the IC₅₀ values of the candidate compounds. The cell lines listed were maintained at approximately 5% CO₂, 37°C, 95% relative humidity in the


media appropriate for that cell line. The cells were sub-cultured every two to three days and plated in clear bottom 96-well plates at a density of 1×10^4 cells/well and incubated at ca. 5% CO₂, 37°C overnight prior to initiation of the assay. To begin cell viability assays, the media in the cell plate (100 µL) was replaced with fresh media (100 µL). The test articles were serially diluted 1:2 in fresh media in duplicate and added to the cells (100 µL) at final sample concentrations of 0.46, 1.37, 4.12, 12.35, 37.04, 111.1, 333.3 and 1000 µM ($\leq$ 1% DMSO) in a total volume of 200 µL. Wells containing no cells and wells containing cells lysed with 0.1% Triton-X were used for baseline controls. Tamoxifen was used as a known control for each assay and DMSO only will be run as vehicle control. The samples were incubated at ca. 37 °C in humidified 5% CO₂ atmosphere for 72 hours. The plate was monitored once a day during the incubation period, paying special attention to the level of confluence. If the cells approach confluence prior to the end of the 72 hour incubation period the experiment was terminated at that time and cell viability measured as described below.

Cell Viability was determined using a commercially available kit to determine ATP levels by luminescence. Briefly, the cell plate had the media removed and replaced with 100 µL of fresh media, and the buffer and lyophilized substrate were equilibrated to room temperature. The buffer was used to reconstitute the substrate just prior to addition to the wells of the cell plate (100 µL per well). The plate was placed into the Infinite M200 plate reader, allowed to shake for 10 minutes followed by a 10 minute wait period, the plate was read using an integration time of 0.5 sec with no attenuation.

The mean baseline controls (wells with Triton X-100 or no cells) were subtracted from the total luminescence to give the net luminescence for that well. This total was compared to the control of DMSO only. An IC_{50} was calculated
5 as the concentration that led to a response of 50% compared to the vehicle control cells. Figures 5 and 6 depict the results of the tests. Accordingly, those of ordinary skill in the art can appreciate that the R configuration of the instantly claimed composition are superior to other
10 stereoisomers.

We claim:

1. A compound having the structure:



wherein R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , R_7 and R_9 are independently
 5 hydrogen, C_1 to C_6 alkyl or substituted alkyl, halogen, sulfate or glucuronide moieties; the ---- symbol represents either a single or a double bond and when the ---- symbol is a double bond and forms a keto group at position 3 or 17, then no R_7 or R_6 is present, respectively; the symbol ---
 10 represent the presence or absence of a bond at position 10; and the ~~~~ symbol represents any type of bond regardless of the stereochemistry; and the enantiomers, other stereochemical isomers, hydrates, solvates, tautomers and pharmaceutically acceptable salts of said compounds.

15 2. The compound of claim 1 wherein r_5 is C_1 to C_6 alkyl or substituted alkyl.

3. The compound of claim 1 wherein r_5 is methyl or hydrogen and r_1 , r_2 , r_3 , and r_4 are hydrogen and ---- is a single bond at positions 3 and 17.


20 4. The compound of claim 3, wherein r_5 is hydrogen.

5. The compound of claim 3 or 4, wherein said compound is the r and s stereoisomer.

6. The Compound Of Claim 2, Wherein R_1 , R_2 , R_3 , R_4 , R_6 , And R_7 Are Independently Hydrogen, Methyl, Or Ethyl.

7. The compound of claim 6, wherein the $\sim$ symbol represents a bond forming a R or a S stereoisomer.

5 8. A compound having the structure:



wherein:

R_1 , R_2 , R_3 , R_4 are independently selected from the group consisting of H, C_1 - C_6 alkyl, substituted alkyl, and halogen;

10 R_5 , R_6 , R_7 are independently selected from the group consisting of H, C_1 - C_6 alkyl, substituted alkyl, sulfate, and glucuronide, and


the --- symbol represents either a single or a double bond with the proviso that when the --- symbol is a double bond and forms a keto group at position 3 or 17, then no R_7 or R_6 is respectively present.

9. The compound of claim 8 wherein R_5 is a C_1 to C_6 alkyl or a substituted alkyl.

10. The compound of claim 9 wherein R_6 and R_7 are
20 independently selected from a group consisting of methyl, hydrogen, or HSO_3^- group.

11. The compound of claim 9 wherein R_5 is methyl or hydrogen and R_1, R_2, R_3, R_4 , is hydrogen.

12. A compound having the structure:



5 wherein:

$R_1, R_2, R_3, R_4, R_8, R_9$ are independently selected from the group consisting of H, C_1 - C_6 alkyl, substituted alkyl, and halogen,


R_5, R_6, R_7 are independently selected from the group
10 consisting of H, C_1 - C_6 alkyl, substituted alkyl, sulfate, or glucuronide, and

the ---- symbol represents either a single or a double bond with the proviso that when the ---- symbol is a double bond and forms a keto group at position 3 or 17, then no R_7 or R_6 is respectively present; the ~~~~~ symbol represents any
15 type of bond regardless of the stereochemistry; and the symbol --- represents the presence or absence of a bond at position 10.

13. The compound of claim 12, wherein R_5 is C_1 to C_6 alkyl or
20 substituted alkyl.

14. The compound of claim 13, wherein R_5 is methyl or hydrogen and R_1, R_2, R_3, R_4 , is hydrogen.

15. A compound having the structure:



wherein:

R₁, R₃, R₄, R₈ are independently selected from the group
 5 consisting of H, C₁-C₆ alkyl, substituted alkyl, and halogen,

R₅, R₇ are independently selected from the group
 consisting of H, C₁-C₆ alkyl, substituted alkyl, sulfate, or
 glucuronide, and == is a single or a double bond.

16. A pharmaceutical composition comprising the compound
 10 of claim 1 and a pharmaceutically-acceptable carrier,
 wherein said compound comprises at least one S enantiomer, R
 enantiomer, S diastereomers or R diastereomers.

17. A pharmaceutical composition comprising the compound
 of claim 8 and a pharmaceutically-acceptable carrier,
 15 wherein said compound comprises at least one S enantiomer or
 R enantiomer, S diastereomer or R diastereomer.

18. A pharmaceutical composition comprising the compound of
 claim 12 and a pharmaceutically-acceptable carrier, wherein
 said compound comprises at least one S enantiomer, R
 20 enantiomer, S diastereomer or R diastereomer.


19. A pharmaceutical composition comprising the compound of
 claim 15 and a pharmaceutically-acceptable carrier, wherein

said compound comprises at least one S enantiomer, R enantiomer, S diastereomer or R diastereomer.

20. A pharmaceutical composition according to any of the claims 16, 17, 18, and 19 wherein said compound is an
5 isolated S or R diastereomer at positions 3 and 17 and the composition is substantially free of the other diastereomer for said positions.

21. A method of inhibiting growth of cancer cells comprising: providing a growth of cancer cells; and
10 contacting said cancer cells with a compound of claim 1, 8, 12, and 15, said compound in an amount at least partially sufficient to inhibit said growth.

22. The method of claim 21 wherein said compound has the following structure:



15

wherein:

R₁, R₂, R₃, R₄ are independently selected from the group consisting of H, C₁-C₆ alkyl, substituted alkyl, and halogen,


R₅, R₆ are independently selected from the group
20 consisting of H, C₁-C₆ alkyl, substituted alkyl, sulfate, or glucuronide, and

the ---- symbol represents either a single or a double bond with the proviso that when the ---- symbol is a double bond and when forms a keto group at position 17, then no R₆ is present.


5 23. The method of claim 22, wherein said cancer growth
cells comprises a solid tumor selected from the group of
solid tumors consisting of breast, pancreatic, lung, colon,
prostate, ovarian, brain, liver, spleen, kidney, lymph node,
small intestine, blood cells, bone, stomach, endometrium,
10 testicular, ovary, central nervous system, skin, head and
neck, esophagus, and bone marrow cancer.

24. The method of claim 23, wherein said compound is in a
pharmaceutical suitable carrier.


25. A method of inhibiting growth of cancer cells
15 comprising: administering to a region of patient in need
thereof a compound having the formula selected from the
group consisting of:




NDC-1099




NDC-1022



NDC-1033



NDC-1044, and




NDC-1055.

5

26. A method of inhibiting solid tumor growth in a mammalian host which comprises administering to the host a therapeutically effective amount of a compound having the structure:

10



wherein R₁, R₂, R₃, R₄, R₅ and R₆ is as described above.

27. The method of claim 26 wherein the mammalian host is a human.

- 5 28. The method of claim 27 wherein the solid tumor growth represents a solid tumor associated with a cancer selected from the group consisting of breast, pancreatic, lung, colon, prostate, ovarian, brain, liver, spleen, kidney, lymph node, small intestine, blood cells, bone, stomach,
- 10 endometrium, testicular, ovary, central nervous system, skin, head and neck, esophagus, and bone marrow cancer.